



# *A Remark on the Uniqueness of Solutions to Hyperbolic Conservation Laws*

ALBERTO BRESSAN & CAMILLO DE LELLIS

*Communicated by T.-P. LIU*

## Abstract

Given a strictly hyperbolic  $n \times n$  system of conservation laws, it is well known that there exists a unique Lipschitz semigroup of weak solutions, defined on a domain of functions with small total variation, which are limits of vanishing viscosity approximations. The aim of this note is to prove that every weak solution taking values in the domain of the semigroup, and whose shocks satisfy the Liu admissibility conditions, actually coincides with a semigroup trajectory.

## 1. Introduction

We consider the Cauchy problem for a strictly hyperbolic  $n \times n$  system of conservation laws in one space dimension

$$u_t + f(u)_x = 0, \quad (1.1)$$

$$u(0, x) = \bar{u}(x), \quad (1.2)$$

with  $f \in \mathcal{C}^2(\mathbb{R}^n, \mathbb{R}^n)$ . In this setting, it is well known that there exists a Lipschitz continuous semigroup  $S : \mathcal{D} \times [0, +\infty[ \mapsto \mathcal{D}$  of weak solutions, defined on a domain  $\mathcal{D} \subset \mathbf{L}^1(\mathbb{R}; \mathbb{R}^n)$  of functions with sufficiently small total variation. The trajectories of this semigroup are the unique limits of vanishing viscosity approximations [3]. All of their shocks satisfy the Liu admissibility conditions [2, 17, 18]. We recall that the semigroup is globally Lipschitz continuous w.r.t. the  $\mathbf{L}^1$  distance. Namely, there exists a constant  $L$  such that

$$\|S_t \bar{u} - S_s \bar{v}\|_{\mathbf{L}^1} \leq L(|t - s| + \|\bar{u} - \bar{v}\|_{\mathbf{L}^1}) \quad \text{for all } s, t \geq 0, \bar{u}, \bar{v} \in \mathcal{D}. \quad (1.3)$$

Given any weak solution  $u = u(t, x)$  of (1.1), (1.2), various conditions have been derived in [7, 9, 10] which guarantee the identity

$$u(t) = S_t \bar{u} \quad \text{for all } t \geq 0. \quad (1.4)$$

Since the semigroup  $S$  is unique, the identity (1.4) yields the uniqueness of solutions to the Cauchy problem (1.1), (1.2). In addition to the standard assumption that each characteristic field is either linearly degenerate or genuinely nonlinear, earlier results required some additional regularity conditions, such as “Tame Variation” or “Tame Oscillation”, controlling the behavior of the solution near a point where the variation is small.

The aim of the present note is to show that uniqueness is guaranteed in a fully general setting: without any assumption about genuine nonlinearity, and without any of the above regularity conditions. Moreover, no assumption is made about the existence of a convex entropy. Our only requirement is that all points of approximate jump satisfy the Liu admissibility conditions.

As in Refs. [7, 9, 10], the proof relies on the elementary error estimate

$$\|u(t) - S_t \bar{u}\|_{\mathbf{L}^1} \leq L \cdot \int_0^t \liminf_{h \rightarrow 0+} \frac{\|u(\tau + h) - S_h u(\tau)\|_{\mathbf{L}^1}}{h} d\tau. \quad (1.5)$$

Indeed, we will prove that the integrand is zero for a.e. time  $\tau \geq 0$ . Following an argument introduced in BRESSAN [4], this is achieved by two estimates:

- (i) In a neighborhood of a point  $(\tau, y)$  where  $u(\tau, \cdot)$  has a large jump, the weak solution  $u$  is compared with the solution to a Riemann problem.
- (ii) In a region where the total variation is small, the weak solution  $u$  is compared with the solution to a linear system with constant coefficients.

To fix our ideas, let

$$M \doteq \sup \left\{ \text{Tot.Var.}\{\bar{u}; \mathbb{R}\}; \bar{u} \in \mathcal{D} \right\} \quad (1.6)$$

be an upper bound for the total variation of all functions in the domain of the semigroup. Notice that this implies

$$\|\bar{u}\|_{\mathbf{L}^\infty} \leq M \quad \text{for all } \bar{u} \in \mathcal{D}. \quad (1.7)$$

Moreover, for each BV function  $\bar{u} \in \mathcal{D}$ , we shall take its right-continuous representative, so that  $\bar{u}(x) = \lim_{y \rightarrow x+} u(y)$ .

To state our result, we first describe the basic setting.

**(A1) (Conservation equations)** The function  $u = u(t, x)$  is a weak solution of the Cauchy problem (1.1), (1.2) taking values within the domain of the semigroup.

More precisely,  $u : [0, T] \mapsto \mathcal{D}$  is continuous w.r.t. the  $\mathbf{L}^1$  distance. The identity  $u(0, \cdot) = \bar{u}$  holds in  $\mathbf{L}^1$ , and, moreover,

$$\iint (u \varphi_t + f(u) \varphi_x) dx dt = 0 \quad (1.8)$$

for every  $\mathcal{C}^1$  function  $\varphi$  with compact support contained inside the open strip  $[0, T[ \times \mathbb{R}$ .

To introduce the Liu admissibility condition on the shocks [17, 18], we first recall that **(A1)** implies that  $u = u(t, x)$  is a function of bounded variation in time and space (cf. [15, Section 5.1] for the definition). Indeed, by [13, Theorem 4.3.1], we have the Lipschitz bound

$$\|u(t_2, \cdot) - u(t_1, \cdot)\|_{L^1(\mathbb{R})} \leq C_M (t_2 - t_1) \quad \text{for all } 0 \leq t_1 \leq t_2, \quad (1.9)$$

for some constant  $C_M > 0$  depending only on the flux  $f$  and on the upper bound  $M$  for the total variation.

By the structure theorem for BV functions of two variables (see e.g. [15, Section 5.9] or [1]), there is a Borel subset  $\mathbf{J} \subset [0, 1] \times \mathbb{R}$  with the following three properties:

- (i) Every point  $(\tau, \xi) \notin \mathbf{J}$  is a point of approximate continuity.
- (ii)  $\mathbf{J}$  is countably 1-rectifiable, i.e. it can be covered by countably many Lipschitz curves, possibly leaving out a subset of zero  $\mathcal{H}^1$  measure ( $\mathcal{H}^1$  denotes the Hausdorff 1-dimensional measure, cf. [15, Section 2.1]).
- (iii)  $\mathcal{H}^1$ -almost every point  $(\tau, \xi) \in \mathbf{J}$  is an approximate jump of the function  $u$ . More precisely there exist states  $u^-, u^+ \in \mathbb{R}^n$  and a speed  $\lambda \in \mathbb{R}$  such that, calling

$$U(t, x) \doteq \begin{cases} u^- & \text{if } x < \lambda t, \\ u^+ & \text{if } x > \lambda t, \end{cases} \quad (1.10)$$

if holds that

$$\lim_{r \rightarrow 0+} \frac{1}{r^2} \int_{-r}^r \int_{-r}^r |u(\tau + t, \xi + x) - U(t, x)| \, dx \, dt = 0. \quad (1.11)$$

Defining the rescaled functions

$$u_r(t, x) \doteq u(\tau + rt, \xi + rx) \quad (1.12)$$

by (iii) and **(A1)**, it follows that  $u_r$  converges to  $U$  in  $L^1_{loc}(\mathbb{R}^2)$ . In particular the conservation equations (1.8) must hold for the piecewise constant function  $U$ , and the triple  $(u^+, u^-, \lambda)$  must therefore satisfy the Rankine-Hugoniot equations

$$f(u^+) - f(u^-) = \lambda(u^+ - u^-). \quad (1.13)$$

Now let a left state  $u^- \in \mathbb{R}^n$  be given. Since the system is strictly hyperbolic, there exist  $n$  shock curves

$$s \mapsto S_i(s)(u^-), \quad i = 1, \dots, n,$$

parameterizing the sets of right states  $u^+$  connected to the left state  $u^-$  by a shock of the  $i$ -th family [5, 13, 16]. As in (1.13), denote by  $\lambda = \lambda_i(s)$  the Rankine-Hugoniot speed of a shock with left and right states  $u^-$  and  $u^+ = S_i(s)(u^-)$ ,

**(A2) (Liu admissibility condition)** *In the above setting, a shock with left and right states  $u^-$  and  $u^+ = S_i(\sigma)(u^-)$  is Liu-admissible if  $\lambda_i(\sigma) \leq \lambda_i(s)$  for all  $s \in [0, \sigma]$ .*

Our result can be simply stated as follows:

**Theorem 1.1.** Let (1.1) be a strictly hyperbolic  $n \times n$  system. Then every weak solution  $u : [0, T] \mapsto \mathcal{D}$ , taking values within the domain of the semigroup and whose shocks satisfy the Liu admissibility condition, coincides with a semigroup trajectory.

Under the additional assumptions that each characteristic family is either linearly degenerate or genuinely nonlinear, and that the  $n \times n$  system (1.1) admits a strictly convex entropy selecting the admissible shocks, this uniqueness result was recently proved in [8]. Restricted to a class of  $2 \times 2$  systems, an earlier proof can also be found in [12].

## 2. Proof of the Theorem

**1.** Let  $\mathbf{J}$  be the set introduced in the previous section and let  $\mathbf{S} \subset \mathbf{J}$  be the subset of all points which are not approximate jumps. Since  $\mathcal{H}^1(\mathbf{S}) = 0$ , its projection on the time axis is a subset  $\mathcal{N} \subset [0, T]$  which is null for the Lebesgue measure. Every point  $(t, x) \in [0, T] \times \mathbb{R}$  with  $t \notin \mathcal{N}$  is therefore either a point of approximate jump, or a point of approximate continuity.

Let us denote by  $\mathcal{J}$  the set

$$\mathcal{J} \doteq \{(\tau, \xi) : \tau \notin \mathcal{N} \text{ and } (\tau, \xi) \text{ is an approximate jump}\}.$$

While it follows immediately from the aforementioned BV structure theorem that  $\mathcal{J}$  is rectifiable, we claim here a stronger property:  $\mathcal{J}$  can be covered by the graphs of countably many Lipschitz functions

$$x = \phi_\ell(t), \quad \ell \in \mathbb{N}, \quad (2.1)$$

and moreover the Lipschitz constant of each  $\phi_\ell$  is bounded by a number  $\Lambda$  which depends only on  $f$  and on the constant  $M$  in (1.6). More precisely, recalling (1.7) we can set

$$\Lambda \doteq 2 \operatorname{Lip}(f, B_M), \quad (2.2)$$

where the right hand side denotes the Lipschitz constant of the function  $f$  over the ball  $B_M \subset \mathbb{R}^n$  centered at the origin with radius  $M$ .

Since by (1.7) it follows

$$|u^\pm(\tau, \xi)| \leq M \quad \text{for all } (\tau, \xi) \in \mathcal{J},$$

the above definition implies that the shock speed  $\lambda = \lambda(\tau, \xi)$  at (1.10), (1.13) satisfies the bound

$$|\lambda(\tau, \xi)| \leq \operatorname{Lip}(f, B_M) \leq \frac{\Lambda}{2}. \quad (2.3)$$

In order to prove our claim, we decompose  $\mathcal{J}$  in the countable union of suitable pieces. First of all, for every integer  $k \geq 1$  we define

$$\mathcal{J}_k \doteq \left\{ (\tau, \xi) \in \mathcal{J} : |u^+(\tau, \xi) - u^-(\tau, \xi)| \geq \frac{1}{k} \right\}.$$

Obviously,  $\mathcal{J} = \bigcup_k \mathcal{J}_k$ . Next, given any pairs  $(\tau_1, \xi_1)$  and  $(\tau_2, \xi_2)$  in  $\mathcal{J}_k$ , consider the two piecewise constant functions

$$U_j(t, x) \doteq \begin{cases} u^-(\tau_j, \xi_j) & \text{if } x < \lambda(\tau_j, \xi_j)t, \\ u^+(\tau_j, \xi_j) & \text{if } x > \lambda(\tau_j, \xi_j)t, \end{cases} \quad j = 1, 2. \quad (2.4)$$

By (2.3) there is a positive number  $\varepsilon(k, \Lambda)$  depending on  $k$  and  $\Lambda$  such that the following holds. If  $(\tau_s, \xi_s)$  is yet a third point in the plane with the properties that  $\tau_s^2 + \xi_s^2 = 1$  and  $|\xi_s| \geq \Lambda|\tau_s|$ , then if we “shift”  $U_2$  by this vector we get the inequality

$$\int_{B_1} |U_1(t, x) - U_2(t + \tau_s, x + \xi_s)| dx dt \geq 6\varepsilon, \quad (2.5)$$

where  $B_1$  denotes the unit disk centered at the origin in  $\mathbb{R}^2$ . We subdivide further  $\mathcal{J}_k$  as a union of sets  $\mathcal{J}_{k,j}$ ,  $j \geq 1$ , where  $(\tau, \xi)$  belongs to  $\mathcal{J}_{k,j}$  if

$$\int_{B_r(\tau, \xi)} |u(t, x) - U(t - \tau, x - \xi)| dx dt \leq \varepsilon r^2 \quad \text{for all } r < \frac{1}{j}. \quad (2.6)$$

Here  $U$  is defined as in (1.10) and  $B_r(\tau, \xi)$  denotes the disk centered at  $(\tau, \xi)$  with radius  $r$ . Clearly,

$$\mathcal{J} = \bigcup_k \mathcal{J}_k = \bigcup_{k,j} \mathcal{J}_{k,j}.$$

Next, consider two points  $(\tau_1, \xi_1), (\tau_2, \xi_2) \in \mathcal{J}_{k,j}$  such that  $r \doteq |(\tau_1, \xi_1) - (\tau_2, \xi_2)| < \frac{1}{2j}$ . Let  $U_j$  be as in (2.4) and set the shift  $(\tau_s, \xi_s)$  to be

$$\tau_s = \frac{\tau_1 - \tau_2}{r}, \quad \xi_s = \frac{\xi_1 - \xi_2}{r}.$$

We claim that (2.5) cannot hold with this shift. This will enable us to conclude  $|\xi_s| \leq \Lambda|\tau_s|$ . To prove the claim, observe first that

$$\begin{aligned} & \int_{B_1} |U_1(t, x) - U_2(t + \tau_s, x + \xi_s)| dx dt \\ & \leq \int_{B_1} |U_1(t, x) - u(\tau_1 + rt, \xi_1 + rx)| dx dt \\ & \quad + \int_{B_1} |u(\tau_1 + rt, \xi_1 + rx) - U_2(t + \tau_s, x + \xi_s)| dx dt \end{aligned}$$

We then change variables in the integrals to  $(\sigma, y) = (\tau_1 + rt, \xi_1 + rx)$ . Observe that

$$\begin{aligned} t + \tau_s &= t + r^{-1}(\tau_1 - \tau_2) = r^{-1}(t + \tau_1 - \tau_2) = r^{-1}(\sigma - \tau_2), \\ x + \xi_s &= x + r^{-1}(\xi_1 - \xi_2) = r^{-1}(y - \xi_2), \end{aligned}$$

while  $U_j(r^{-1}(\sigma - \tau_j), r^{-1}(y - \xi_j)) = U_j(\sigma - \tau_j, y - \xi_j)$  because of the 0-homogeneity of the functions  $U_j$ . Hence the change of variables yields

$$\begin{aligned} \int_{B_1} |U_1(t, x) - U_2(t + \tau_s, x + \xi_s)| \, dx \, dt \\ \leq \frac{1}{r^2} \int_{B_r(\tau_1, \xi_1)} |U_1(\sigma - \tau_1, y - \xi_1) - u(\sigma, y)| \, dy \, d\sigma \\ + \frac{1}{r^2} \int_{B_r(\tau_1, \xi_1)} |u(\sigma, y) - U_2(\sigma - \tau_2, y - \xi_2)| \, dy \, d\sigma \\ \leq \frac{1}{r^2} \int_{B_r(\tau_1, \xi_1)} |U_1(\sigma - \tau_1, y - \xi_1) - u(\sigma, y)| \, dy \, d\sigma \\ + \frac{1}{r^2} \int_{B_{2r}(\tau_2, \xi_2)} |u(\sigma, y) - U_2(\sigma - \tau_2, y - \xi_2)| \, dy \, d\sigma, \end{aligned}$$

where we have used the inclusion  $B_r(\tau_1, \xi_1) \subset B_{2r}(\tau_2, \xi_2)$  to get the last inequality. Note next that  $r < 2r < \frac{1}{j}$  and, since both  $(\tau_1, \xi_1)$  and  $(\tau_2, \xi_2)$  belong to  $\mathcal{J}_{k,j}$ , we can use (2.6) to bound the first summand by  $\varepsilon$  and the second summand by  $4\varepsilon$ . In particular we conclude

$$\int_{B_1} |U_1(t, x) - U_2(t + \tau_s, x + \xi_s)| \, dx \, dt \leq 5\varepsilon.$$

As already pointed out, since the latter inequality contradicts (2.5), we conclude that  $|\xi_s| \leq \Lambda |\tau_s|$ , which in turn gives  $|\xi_2 - \xi_1| \leq \Lambda |\tau_2 - \tau_1|$ .

We have thus proved the following fact:

**(L)** *If  $(\tau_1, \xi_1), (\tau_2, \xi_2) \in \mathcal{J}_{k,j}$  and  $|(\tau_1, \xi_1) - (\tau_2, \xi_2)| < \frac{1}{2j}$ , then  $|\xi_2 - \xi_1| \leq \Lambda |\tau_2 - \tau_1|$ .*

It is well known that from (L) it follows that  $\mathcal{J}_{k,j}$  can be covered by countably many Lipschitz graphs of functions  $x = \phi_\ell(t)$ ,  $\ell \in \mathbb{N}$ , with Lipschitz constant at most  $\Lambda$ . See for instance [14, Lemma 4.7].

For readers' convenience, we include here a proof. If  $B = B_{1/4j}(x_0, t_0)$  is any disk of radius  $1/(4j)$  and we set  $F \doteq \mathcal{J}_{k,j} \cap B$ , then (L) implies

$$|x - y| \leq \Lambda |t - s| \quad \text{for all } (t, x), (s, y) \in F. \quad (2.7)$$

This obviously implies that there are no points of  $F$  which lie on the same line  $\{t = \text{const}\}$ . Hence, if  $G$  is the projection of  $F$  on the time axis, then there is a function  $\phi : G \rightarrow \mathbb{R}$  such that  $F = \{(t, \phi(t)) : t \in G\}$ . On the other hand (2.7) is equivalent to the statement that the Lipschitz constant of  $G$  is at most  $\Lambda$ . By the classical Lipschitz extension theorem we can simply extend  $\phi$  to a Lipschitz function defined on the whole time axis. Since  $\mathcal{J}_{k,j}$  can be covered by a countable collection of disks with radius  $1/(4j)$ , the existence of the desired covering by means of countably many Lipschitz graphs follows immediately.

**2.** Next, we wish to show that, if  $t \notin \mathcal{N}$  and  $(t, x) \notin \mathbf{J}$ , then  $y \mapsto u(t, y)$  is continuous at  $x$ . We start by noticing that, since  $(t, x) \notin \mathbf{J}$ ,  $u$  is approximately

continuous at  $(t, x)$  as a function of two variables. Therefore there is a  $u_0 \in \mathbb{R}^n$  such that

$$\lim_{r \downarrow 0} \frac{1}{r^2} \int_{t-r}^{t+r} \int_{x-r}^{x+r} |u(s, y) - u_0| \, ds \, dy = 0.$$

In particular, for every fixed  $\varepsilon > 0$  there is an  $r_0(\varepsilon) > 0$  such that

$$\int_{t-r}^{t+r} \int_{x-r}^{x+r} |u(s, y) - u_0| \, ds \, dy \leq \varepsilon r^2 \quad \text{for all } r \leq r_0(\varepsilon).$$

An elementary application of Chebyshev's inequality and Fubini's theorem yields then the existence of a  $t(r) \in [t - r\sqrt{\varepsilon}, t + r\sqrt{\varepsilon}]$  such that

$$\int_{x-r}^{x+r} |u(t(r), y) - u_0| \, dy \leq r\sqrt{\varepsilon}.$$

Furthermore we can use the Lipschitz estimate (1.9) to bound

$$\frac{1}{r} \int_{x-r}^{x+r} |u(t, y) - u_0| \, dy \leq \sqrt{\varepsilon} + C_M \frac{|t - t(r)|}{r} \leq (1 + C_M)\sqrt{\varepsilon} \quad \text{for all } r \leq r_0(\varepsilon). \quad (2.8)$$

On the other hand recall that  $y \mapsto u(t, y)$  is a function of bounded variation on the real line. As such, every point is either a classical jump point, or a point of continuity. Since  $\varepsilon$  in (2.8) can be chosen arbitrarily small,  $x$  cannot be a classical jump point of  $u(t, \cdot)$  and must therefore be a point of continuity, which was in fact our initial claim.

**3.** Together with the functions  $\phi_\ell$  in (2.1), we consider functions of the form

$$\phi^{y+}(t) = y + \Delta t, \quad \phi^{y-}(t) = y - \Delta t, \quad y \in \mathbb{Q}. \quad (2.9)$$

Since here  $y$  is a rational point, there are countably many of these functions. For convenience, the countable set of all functions in (2.1) together with those in (2.9) will be relabeled as

$$\{\psi_n ; n \geq 1\}. \quad (2.10)$$

Next, we observe that, for every  $j, k \geq 1$ , the scalar function

$$W_{jk}(t) \doteq \begin{cases} \text{Tot.Var.} \{u(t) ; [\psi_j(t), \psi_k(t)]\} & \text{if } \psi_j(t) < \psi_k(t), \\ 0 & \text{otherwise,} \end{cases} \quad (2.11)$$

is bounded and measurable (indeed, it is lower semicontinuous). Therefore a.e.  $t \in [0, T]$  is a Lebesgue point. We denote by  $\mathcal{N}' \subset [0, T]$  the set of all times  $t$  which are NOT Lebesgue for at least one of the countably many functions  $W_{jk}$ . Of course,  $\mathcal{N}'$  has zero Lebesgue measure.

In view of (1.5), we will prove the theorem by establishing the following claim.

**(C)** For every  $\tau \in [0, T] \setminus (\mathcal{N} \cup \mathcal{N}')$  and  $\varepsilon > 0$ , one has

$$\limsup_{h \rightarrow 0+} \frac{1}{h} \|u(\tau + h) - S_h u(\tau)\|_{\mathbf{L}^1} \leq \varepsilon. \quad (2.12)$$

**4.** Assume  $\tau \notin \mathcal{N} \cup \mathcal{N}'$ . By induction on  $k = 0, 1, 2, \dots, N$ , we construct points

$$z_0 < z_1 < \dots < z_n$$

such that

$$\text{Tot.Var.}\{u(\tau, \cdot); \]z_0, \infty[\} < \varepsilon, \quad (2.13)$$

$$\text{Tot.Var.}\{u(\tau, \cdot); \]z_N, +\infty[\} < \varepsilon, \quad (2.14)$$

and moreover, for  $k = 1, 2, \dots, N$ ,

$$z_k = \sup \left\{ x > z_{k-1}; \text{Tot.Var.}\{u(\tau, \cdot); \]y_{k-1}, x[\} < \varepsilon \right\}.$$

Notice that this implies

$$N \leq \frac{\text{Tot.Var.}\{u(\tau, \cdot)\}}{\varepsilon}. \quad (2.15)$$

We then choose additional points

$$y_0 \leqq z_0 \leqq y'_0 < y_1 \leqq z_1 \leqq y'_1 < \dots < y_N \leqq z_N \leqq y'_N$$

according to the following rules.

- (i) If  $u(\tau, \cdot)$  has a jump at  $x = z_k$ , then  $y_k = y'_k = z_k$ .
- (ii) If  $u(\tau, \cdot)$  is continuous at  $x = z_k$ , then the points  $y_k < z_k < y'_k$  are chosen so that

$$y_k - \tau \in \mathbb{Q}, \quad y'_k + \tau \in \mathbb{Q}, \quad (2.16)$$

$$\text{Tot.Var.}\{u(\tau, \cdot); \]y_{k-1}, y'_k[\} < \varepsilon/2. \quad (2.17)$$

Notice that, for every  $k = 1, \dots, N$ , this implies

$$\text{Tot.Var.}\{u(\tau, \cdot); \]y_{k-1}, y'_k[\} \leqq 2\varepsilon. \quad (2.18)$$


**5.** The remainder of the proof is very similar to the one in [8]. For any given  $y \in \mathbb{R}$ , we denote by  $U^\sharp = U_{(u, \tau, y)}^\sharp(t, x)$  the solution to the Riemann problem for (1.1), with initial data

$$u(\tau, x) = \begin{cases} u(\tau, y-) & \text{if } x < y, \\ u(\tau, y+) & \text{if } x > y. \end{cases} \quad (2.19)$$

Moreover, for every given  $k \geqq 1$  we denote by  $U^\flat = U_{(u, \tau, k)}^\flat(t, x)$  the solution to the linear Cauchy problem with constant coefficients

$$v_t + Av_x = 0, \quad v(\tau, x) = u(\tau, x). \quad (2.20)$$

Here the  $n \times n$  matrix  $A$  is the Jacobian matrix of  $f$  computed at the midpoint of the interval  $[y_{k-1}, y'_k]$ . Namely,



**Fig. 1.** The points  $z_k, y_k, y'_k$  constructed in the proof of the theorem

$$A = Df \left( u \left( \tau, \frac{y_{k-1} + y'_k}{2} \right) \right).$$

With reference to Fig. 1, to estimate the lim-sup in (2.12), we need to estimate two types of integrals.

(I) The integral of  $|u(t, x) - U_{(u, \tau, y_k)}^\sharp(t, x)|$  over the intervals

$$J_y(t) \doteq [y_k - \Lambda(t - \tau), y_k + \Lambda(t - \tau)],$$

for every  $k$  such that  $y_k = y'_k$ .

(II) The integral of  $|u(t, x) - U_{(u, \tau, k)}^\flat(t, x)|$  over the intervals

$$J_k(t) = [y_{k-1} + \Lambda(t - \tau), y'_k - \Lambda(t - \tau)].$$

**6.** To estimate integrals of type (I) we observe that, since  $\tau \notin \mathcal{N}$ ,  $(\tau, y_k)$  is either a Lebesgue point or a point of approximate jump of the function  $u$ . Therefore

$$\lim_{t \rightarrow \tau^+} \frac{1}{t - \tau} \int_{y_k - \Lambda(t - \tau)}^{y_k + \Lambda(t - \tau)} |u(t, x) - U_{(u, \tau, y_k)}^\sharp(t, x)| dx = 0. \quad (2.21)$$

Indeed, this follows from (1.11) and the Lipschitz continuity of the map  $t \mapsto u(t, \cdot) \in \mathbf{L}^1(\mathbb{R})$ ; see Theorem 2.6 in [5] for details.

**7.** To estimate integrals of type (II), two main cases will be considered.

CASE 1:  $y_{k-1} - \Lambda\tau \in \mathbb{Q}$  and  $y'_k + \Lambda\tau \in \mathbb{Q}$ . In this case, since  $\tau \notin \mathcal{N}'$ , the function

$$V(t) \doteq \text{Tot.Var.} \left\{ u(t, \cdot) ; [y_{k-1} + \Lambda(t - \tau), y'_k - \Lambda(t - \tau)] \right\} \quad (2.22)$$

has a Lebesgue point at  $t = \tau$ . Hence

$$\lim_{h \rightarrow 0^+} \frac{1}{h} \int_{\tau}^{\tau+h} |V(t) - V(\tau)| dt = 0.$$

By (2.18) it follows  $V(\tau) \leq 2\varepsilon$ . Therefore

$$\limsup_{h \rightarrow 0+} \frac{1}{h} \int_{\tau}^{\tau+h} V(t) dt \leq 2\varepsilon. \quad (2.23)$$

CASE 2:  $y_{k-1} = \phi_{\ell}(\tau)$ ,  $y'_k = \phi_{\ell'}(\tau)$  for some indices  $\ell, \ell' \in \mathbb{N}$ . In this case, since  $\tau \notin \mathcal{N}'$ , the function

$$W(t) \doteq \text{Tot.Var.} \left\{ u(t, \cdot); [\phi_{\ell}(t), \phi_{\ell'}(t)] \right\}$$

has a Lebesgue point at  $t = \tau$ . Recalling that the functions  $\phi_{\ell}, \phi_{\ell'}$  have Lipschitz constant  $\leq \Lambda$ , a comparison with (2.22) immediately yields  $V(t) \leq W(t)$  for all  $t \geq \tau$ . Since our construction implies  $W(\tau) \leq 2\varepsilon$ , we thus conclude that (2.23) again holds.

The remaining two cases, where  $y_{k-1} = \phi_{\ell}(\tau)$  and  $y'_k + \Lambda\tau \in \mathbb{Q}$ , or where  $y_{k-1} - \Lambda\tau \in \mathbb{Q}$  and  $y'_k = \phi_{\ell'}(\tau)$  for some  $\ell, \ell' \in \mathbb{N}$ , can be handled in the same way. Namely, (2.23) always holds.

Using again the fact that the map  $t \mapsto u(t, \cdot)$  is Lipschitz continuous from  $[0, T]$  into  $\mathbf{L}^1(\mathbb{R})$ , the same argument used in [8] now yields

$$\limsup_{t \rightarrow \tau+} \frac{1}{t - \tau} \int_{y_{k-1} + \Lambda(t - \tau)}^{y'_k - \Lambda(t - \tau)} \left| u(t, x) - U_{(u, \tau, k)}^{\flat}(t, x) \right| dx = \mathcal{O}(1) \cdot \varepsilon^2. \quad (2.24)$$

Indeed, this corresponds to formula (3.20) in [8]. Based on (2.23), the proof is identical and will not be repeated here.

**8.** On the other hand, as showed in [3] all trajectories of the semigroup are weak solutions of (1.1) which satisfy the Liu admissibility conditions. Therefore they satisfy the same bounds as in (2.21) and (2.24). More precisely,

$$\lim_{t \rightarrow \tau+} \frac{1}{t - \tau} \int_{y_{k-1} - \Lambda(t - \tau)}^{y_k + \Lambda(t - \tau)} \left| (S_{t-\tau} u(\tau))(x) - U_{(u, \tau, y_k)}^{\sharp}(t, x) \right| dx = 0, \quad (2.25)$$

$$\limsup_{t \rightarrow \tau+} \frac{1}{t - \tau} \int_{y_{k-1} + \Lambda(t - \tau)}^{y'_k - \Lambda(t - \tau)} \left| (S_{t-\tau} u(\tau))(x) - U_{(u, \tau, k)}^{\flat}(t, x) \right| dx = \mathcal{O}(1) \cdot \varepsilon^2. \quad (2.26)$$

**9.** Combining the previous estimates (2.21), (2.24), (2.25), (2.26), and recalling that the total number of intervals is  $N \leq 2M\varepsilon^{-1}$ , we establish the limit (2.12), proving the theorem.  $\square$

**Acknowledgements.** The research by A. Bressan was partially supported by NSF with grant DMS-2006884, “Singularities and error bounds for hyperbolic equations”. C. De Lellis has been supported by NSF with grant DMS-1946175.

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other right-holder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

## References

1. AMBROSIO, L., FUSCO, N., PALLARA, D.: Functions of Bounded Variation and Free Discontinuity Problems. Clarendon Press, Oxford (2000)
2. BIANCHINI, S.: On the Riemann problem for non-conservative hyperbolic systems. *Arch. Rational Mech. Anal.* **166**, 1–26, 2003
3. BIANCHINI, S., BRESSAN, A.: Vanishing viscosity solutions of nonlinear hyperbolic systems. *Ann. Math.* **161**, 223–342, 2005
4. BRESSAN, A.: The unique limit of the Glimm scheme. *Arch. Rational Mech. Anal.* **130**, 205–230, 1995
5. BRESSAN, A.: Hyperbolic Systems of Conservation Laws. The One Dimensional Cauchy Problem. Oxford University Press, Oxford (2000)
6. BRESSAN, A., CRASTA, G., PICCOLI, B.: Well posedness of the Cauchy problem for  $n \times n$  systems of conservation laws. *Am. Math. Soc. Memoir* **694**, 2000
7. BRESSAN, A., GOATIN, P.: Oleinik type estimates and uniqueness for  $n \times n$  conservation laws. *J. Differ. Equ.* **156**, 26–49, 1999
8. BRESSAN, A., GUERRA, G.: Unique solutions to hyperbolic conservation laws with a strictly convex entropy, preprint, [arxiv:2305.10737](https://arxiv.org/abs/2305.10737)
9. BRESSAN, A., LEFLOCH, P.: Uniqueness of weak solutions to systems of conservation laws. *Arch. Rational Mech. Anal.* **140**, 301–317, 1997
10. BRESSAN, A., LEWICKA, M.: A uniqueness condition for hyperbolic systems of conservation laws. *Discr. Cont. Dyn. Syst.* **6**, 673–682, 2000
11. BRESSAN, A., LIU, T.P., YANG, T.:  $L^1$  stability estimates for  $n \times n$  conservation laws. *Arch. Rational Mech. Anal.* **149**, 1–22, 1999
12. CHEN, G., KRUPA, S., VASSEUR, A.: Uniqueness and weak-BV stability for  $2 \times 2$  conservation laws. *Arch. Rational Mech. Anal.* **246**, 299–332, 2022
13. DAFERMOS, C.: Hyperbolic Conservation Laws in Continuum Physics, 4th edn. Springer, Berlin (2016)
14. DE LELLIS, C.: Rectifiable sets, Densities, and Tangent Measures. Zurich Lectures in Advanced Mathematics. European Mathematical Society, Zurich (2008)
15. EVANS, L.C., GARIEPY, R.F.: Measure Theory and Fine Properties of Functions. CRC Press, Boca Raton (1991)
16. HOLDEN, H., RISEBRO, N.H.: Front Tracking for Hyperbolic Conservation Laws. Springer, Berlin (2015)
17. LIU, T.P.: Riemann problem for general  $2 \times 2$  conservation laws. *Trans. Am. Math. Soc.* **199**, 89–112, 1974
18. LIU, T.P.: The entropy condition and the admissibility of shocks. *J. Math. Anal. Appl.* **53**, 78–88, 1976

ALBERTO BRESSAN  
Department of Mathematics,  
Penn State University,  
Pa.16803 University Park  
USA.  
e-mail: axb62@psu.edu

and

CAMILLO DE LELLIS  
Institute for Advanced Study,  
Princeton  
NJ 08540  
USA.  
e-mail: camillo.delellis@ias.edu

(Received June 24, 2023 / Accepted October 3, 2023)  
Published online October 30, 2023

© The Author(s), under exclusive licence to Springer-Verlag GmbH, DE, part of Springer Nature (2023)