



INVITED PAPER

Effect of Temperature on Mosquito Olfaction

Chloé Lahondère^{*†‡§¶||}, Clément Vinauger^{*†¶}, Jessica E. Liaw^{||}, Kennedy K. S. Tobin^{||}, Jillian M. Joiner^{||} and Jeffrey A. Riffell^{||}

^{*}Department of Biochemistry, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA; [†]The Fralin Life Science Institute, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA; [‡]The Global Change Center, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA; [§]Department of Entomology, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA; [¶]Center of Emerging, Zoonotic and Arthropod-borne Pathogens, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA; ^{||}Department of Biology, University of Washington, Seattle, WA 98195, USA

¹E-mail: lahonder@vt.edu

Synopsis Mosquitoes use a wide range of cues to find a host to feed on, eventually leading to the transmission of pathogens. Among them, olfactory cues (e.g., host-emitted odors, including CO₂, and skin volatiles) play a central role in mediating host-seeking behaviors. While mosquito olfaction can be impacted by many factors, such as the physiological state of the insect (e.g., age, reproductive state), the impact of environmental temperature on the olfactory system remains unknown. In this study, we quantified the behavioral responses of *Aedes aegypti* mosquitoes, vectors of dengue, yellow fever, and Zika viruses, among other pathogens, to host and plant-related odors under different environmental temperatures.

Introduction

Temperature is one of the most important physical factors affecting the life of insects. Because of their poikilothermic nature (i.e., their inability to maintain a constant body temperature, independently of the temperature of the environment), excessive heat or cold can have deleterious effects on insects' physiology and behavior. For each insect species, its behavioral performance, such as feeding or flying, is optimal at a given temperature and possible within a range of temperatures (i.e., between a critical minimum and maximum temperature). Although these responses and behaviors are possible within a range of temperatures (i.e., tolerance range), beyond a critical minimum and maximum point it is impossible for the insect to perform (Huey and Stevenson 1979). Nevertheless, insects have developed strategies to avoid thermal stress and some species can rely on thermoregulation processes (Heinrich 1993). For example, they can adjust their body position toward the sun (May 1976), seek shade (Herrera 1997), use evaporative cooling of droplets of nectar (Heinrich 1979) or urine and blood (Lahondère and Lazzari 2012; Lazzari et al. 2021; Reinhold et al. 2021), or actively pro-

duce or release heat via counter-current heat exchangers (Heinrich 1976; Lahondère et al. 2017).

In blood-feeding insects, the physiological processes regulating host-seeking behavior have been well-described (e.g., nutritional, and reproductive states, age, time of the day, etc. reviewed by Lazzari 2009). By contrast, comparatively less is known about how the temperature of the environment could affect their sensory-mediated behaviors and systems, and by extension, their host-seeking behavior and the transmission of the pathogens responsible for diseases. In mosquitoes, the deadliest animals on earth (WHO, 2019), the temperature of the environment has been shown to affect their behavior (Thomson 1938; Kirby and Lindsay 2004), metabolism (Clements 1992), development (Rueda et al. 1990; Lyimo et al. 1992; Lanciani and Le 1995), and reproduction and blood-feeding (Eldridge 1968). Moreover, by affecting the development of the pathogens that they might carry, temperature can also affect the vectorial capacity of mosquitoes (Paaijmans et al. 2011; Reinhold et al. 2018).

In the context of global warming and the ability of mosquitoes to invade new areas, there is an urgent need

Advance Access publication June 12, 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of the Society for Integrative and Comparative Biology. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

to identify the physical processes influencing mosquito behavior and biology (Lahondère and Bonizzoni 2022). In the present study, we focused on the mosquito olfactory system—essential for host and plant seeking—and how the temperature of the environment affects mosquito responses. More specifically, we examined: (1) how temperature affects flight dynamics of tethered and free-flying mosquitoes; (2) how *Aedes aegypti*, a major vector of pathogens such as Zika, dengue, or yellow fever viruses, responds to host and plant-related odors under different, yet ecologically relevant, thermal conditions; and (3) the possible modulation of antennal responses to odors by the environmental temperature. Specifically, given the effect of temperature on olfactory reception in *Drosophila melanogaster* (Riveron et al. 2009; Martin et al. 2011), we hypothesize that at higher ambient temperatures, *Ae. aegypti* females will be more sensitive and behaviorally responsive to olfactory cues.

Material and methods

Insects

Female *Aedes aegypti* mosquitoes (wild type, line F21 MRA-726, MR4, ATCC[®], Manassas, VA, USA) were used for all experiments. The colony was kept in a climatic chamber maintained at $25 \pm 1^\circ\text{C}$, $60 \pm 10\%$ relative humidity (RH) and under a 12–12 h light–dark cycle. Groups of 160–200 larvae were placed in $26 \times 35 \times 4$ cm covered pans containing Milli-Q[®] water and were fed daily on fish food (Hikari Tropic 382 First Bites—Petco, San Diego, CA, USA). Groups of same day pupae (both males and females) were then isolated in 16 Oz (i.e., 473 mL) containers (Mosquito Breeder Jar, Bioquip Products, Rancho Dominguez, CA, USA) until emergence. Adults were then transferred into mating cages (BioQuip Products, Rancho Dominguez, CA, USA) and maintained on 10% sucrose. To maintain our laboratory colonies, an artificial feeder (D.E. Lillie Glassblowers, Atlanta, GA, USA; 2.5 cm internal diameter) filled with heparinized bovine blood (Lampire Biological Laboratories, Pipersville, PA, USA) placed on the top of the cage and heated at 37°C using a water-bath circulation was used to feed mosquitoes from the main colony weekly.

For the experiments, groups of 120 pupae were isolated from the main colony and maintained in a separate container for 6 days after their emergence. This gave the mosquitoes time to mate in the containers before the experiments (random dissection of females revealed that 95% of them had oocytes). Mosquitoes had access to 10% sucrose *ad libitum* but were not blood-fed before the experiments. The day the experiments were conducted, females were selected manually with forceps af-

ter placing the container for a few minutes in a climatic chamber at 10°C . Depending on the type of experiment, they were then either placed individually or in groups of 20 in 50 mL conical Falcon[™] tubes (Thermo Fisher Scientific, Pittsburgh, PA, USA) and covered with a piece of mesh. We gave them at least 1 h to recover before testing them in the olfactometer. All experiments were conducted when *Ae. aegypti* mosquitoes are the most active and responsive to host related cues, either 2 h before their subjective night or 2 h after their subjective dawn (Trpis et al. 1973; Vinauger et al. 2014).

Tethered flight under different thermal conditions

Flight arena

The in-flight response to CO_2 of individual female mosquitoes was quantified in an LED-based arena (*sensu* Reiser and Dickinson 2008; Vinauger et al. 2018, 2019). Mosquitoes were cold anesthetized on ice and tethered by the thorax with a tungsten wire using UV-activated glue (Loctite 3104 Light Cure Adhesive, Loctite, Düsseldorf, Germany), positioning their main body axis at a 30° angle from the tether. Mosquitoes were then kept at room temperature in a closed container for an approximate 30-min recovery period. A tethered mosquito was centered in a hovering position within an arena composed of 12 columns of two panels each, which were arranged into a regular dodecagon and produced a display resolution of 96×16 pixels (Fig. 1A; Reiser and Dickinson 2008; Vinauger et al. 2018, 2019). Mosquitoes were placed directly under an infrared (IR) diode and above an optical sensor coupled to a wing-beat analyzer (JFI Electronics, University of Chicago; Götz 1987; Lehmann and Dickinson 1997; Reiser and Dickinson 2008). The beating wings cast a shadow onto the sensor, allowing the analyzer to track their motion and measure the amplitude and frequency of each wing stroke (Götz 1987; Lehmann and Dickinson 1997; Reiser and Dickinson 2008; Vinauger et al. 2018, 2019). Measurements were sampled at 5 kHz and acquired with a National Instrument Acquisition board (BNC-2090A, National Instruments, Austin, TX, USA). The mosquito was centered between an air inlet and a vacuum line aligned diagonally with one another, 30° from the vertical axis. The air inlet was positioned 12 mm in front of and slightly above the mosquito's head, targeting the antennae from an angle of 15° . The vacuum line was positioned behind the mosquito 25 mm away from the tip of the abdomen. Two different airlines independently controlled by a solenoid valve (The Lee Company, Westbrook, CT, USA) intersected this main air inlet, either delivering clean air or CO_2 diluted with nitrogen (N_2) to obtain the desired concentration. A visual




Fig. 1 (A) Schematic of the LED arena. The odor delivery line was connected to a thermostat-controlled conductive tubing, maintaining the main air flow at a constant temperature set at either 20, 25, or 30°C. (B) Trigger-averaged and baseline-subtracted wingbeat frequency (WBF), in response to a 1s long pulse of nitrogen (N₂, gray bar, $n = 11$), or CO₂ diluted into N₂ at 20°C (blue bar, $n = 16$), 25°C (purple bar, $n = 14$), or 30°C (red bar, $n = 13$). (C) Proportion of active mosquitoes (i.e., the number of mosquitoes which showed a change in their WBF) at different temperatures. Asterisks indicate distributions that are significantly different from random (Binomial tests; $P < 0.05$). Vertical bars represent S.E.M. (i.e., standard errors of the mean). (D) Trigger-averaged and baseline-subtracted wingbeat amplitude (WBA) in response to the same stimuli.

pattern of alternating stripes of either inactive or fully lit LEDs, each 16×6 pixels in size (22.5°) was used in conjunction with the odor stimulus and set in closed-loop with the mosquito behavior to maintain their motivation to fly. Odor stimuli consisted of a 1sec pulse followed by a 60-s inter-trial-interval, after which the sequence was repeated for a total of five pulses. The responses of the mosquitoes were tested at 20 ($n = 16$), 25 ($n = 14$), and 30°C ($n = 13$). A N₂ control ($n = 11$) at 25°C was also performed.

Data analysis

Data were analyzed in R version 4.0.0. For each pulse (either CO₂ or N₂ control), the wingbeat frequency

and amplitude were normalized by averaging the signal across a 1-s time window preceding the stimulus delivery and then subtracting this baseline value from the signal values following the pulse. Trials were discarded in which mosquitoes had frequency fluctuations greater than 5 Hz in this 1-s window or frequency changes greater than 30 Hz that did not begin within the 4-s following stimulation (as they presumably were not in response to the stimulus). The mean response for each individual was calculated from the saved trials and used as a replicate to calculate the mean response for each treatment group. This latter was calculated using the difference between the mean frequency—and amplitude—before (200 ms before stimulation) and the maxima

of the signal after stimulus delivery (within a window starting at 1 s and ending 3 s after the pulse). One-tailed Student's *t*-tests for paired samples were used to test for differences from baseline and Student's *t*-tests for independent samples were used to test for differences between groups. *P*-values were adjusted for multiple comparisons using the Bonferroni correction (Jafari et al. 2019).

Free flight orientation to the olfactory cue under different thermal conditions

Olfactometer

To test if the ambient temperature impacts mosquito's olfaction, we measured and compared the mosquito behavioral responses to different odors under various thermal conditions using a Y-maze Plexiglas® olfactometer. Three different temperature conditions were tested: 20, 25, and 30°C, which are in the range of activity of *Ae. aegypti* [i.e., 17 to 34°C, Christophers (1960)]. The olfactometer was composed of a releasing chamber connected to a tube of 30 cm long and 10 cm of diameter attached to a central box where two choice arms (39 cm long, 10 cm diameter) were placed. White cardboard was placed under and on the sides of the olfactometer to prevent the mosquitoes from being distracted by surroundings. Fans (Rosewill, Los Angeles, CA, USA) were placed at the end of the two arms of the olfactometer to generate a constant airflow (air speed approximately 40 cm.s⁻¹). To create a laminar flow, the air first entered a charcoal filter [to remove odor contaminants (C16 × 48, Complete Filtration Services, Greenville, NC, USA)] and a series of mesh screens and honeycomb metal sheet (10 cm thick). The odor delivery system to the two choice arms was made using a charcoal-filtered air stream adjusted with flow meters equipped with needle valves. Teflon® tubing (3 mm internal diameter) conducted the air flow through each of the two 20 mL scintillation vials (Thermo Fisher Scientific, Waltham, MA, USA) containing 12 mL of either the tested odor or the control solution (i.e., MilliQ water or ethanol 100%). Each tubing was connected to the corresponding choice arm of the olfactometer and placed at about 4 cm from the fans and its end located in the center of the arm. All the olfactometer experiments were conducted in a well-ventilated climatic chamber (Environmental Structures, Colorado Springs, CO, USA), 50% RH. After each experiment, the olfactometer, tubing, and vials were cleaned with water and then ethanol 100% to avoid any odorant contamination between experiments. Finally, to avoid any side biases in the arms of the olfactometer, both stimulus and control vials were randomly tested on both sides for all groups.

Experimental procedure

Response to carbon dioxide in groups

The first experiment consisted in measuring the response of groups of mosquitoes to carbon dioxide (CO₂), which is a strong activator and attractant (i.e., host-seeking cue). Carbon dioxide was chemically generated using sulfuric acid (1 M) and sodium carbonate (0.3 M) via the reaction: Na₂CO₃ + H₂SO₄ → CO₂ + H₂O + Na₂SO₄. Briefly, a solution of Na₂CO₃ was injected at a constant rate (10 mL.h⁻¹) in a jar containing 100 mL of H₂SO₄ using a drip-feed device (Barrozo and Lazzari 2004). The production of CO₂ was monitored along the experiment with a carbon dioxide detector and calibrated so the level of CO₂ in the olfactometer was of 2000 ppm. The jar containing sulfuric acid was placed on a stir plate to help mixing the two chemicals and ensure a constant CO₂ production. To send the odor to the olfactometer, an air pump gently injected air to the jar through tubing from the jar to the arm of the olfactometer. As a control, 100 mL of sulfuric acid was placed in a jar and the same air delivery system was used to connect to the olfactometer. Twenty female mosquitoes were released in the olfactometer and were given 30 min total to choose between the two arms carrying either the tested odor or the control. The total number of females in each arm (i.e., females who made a choice) was then counted as well as the total number of females that stayed in the central box and main arm of the olfactometer, which were considered as inactive.

Individual response to ecologically relevant odors

The individual response to different odors was then tested using the olfactometer. This allowed us to observe the behavior of each mosquito and record their flight behavior and analyze their flight trajectories. We tested seven different biologically relevant odors, both host- and plant-related: lactic acid [L-(+)-lactic, Sigma Aldrich, ≥98% purity—concentration: 50 mM in milliQ water]; Octenol [1-octen-3-ol, Sigma Aldrich, ≥98% purity; enantiomeric ratio: ≥99:1 (GC)—concentration: 140 mM in milliQ water]; Nonanol [1-nonanol, Sigma Aldrich, purum, ≥98.0% (GC)—concentration: 1.58 mM in milliQ water], Hexanol [1-hexanol; Sigma Aldrich, >98% purity; 92% of the Z isomer—concentration: 85 mM in milliQ water], Benzaldehyde (Sigma Aldrich, purified by redistillation, ≥99.5%—concentration: 0.98 M in milliQ water), DEET (N, N-diethyl-meta-toluamide, Supelco, ≥95% purity—concentration: 40% diluted in 100% ethanol to match commercial version of the repellent), and CO₂ (chemically produced as previously described—concentration: 2000 ppm above ambient air levels). Hexanol and DEET are known olfactory repellents and

feeding deterrents of *Ae. aegypti*, while octenol and lactic acid [here, used at a concentration similar to human skin emissions (Eiras and Jepson 1991; Cork and Park 1996; Geier et al. 1996)] evoke a slight attraction preference, and nonanol is neutral. Carbon dioxide was used as a positive control and we also ran experiments with two clean air currents as a negative control and to assess that no bias (i.e., innate preference for one side of the olfactometer) could be evinced. We also performed each set of experiments in duplicate, switching the odor side to control for any possible side preference. The results from the two sets of experiments were then grouped and summed. After releasing the mosquito in the olfactometer, each female was given 5 min to choose between the two arms. We considered that the mosquito made a choice when it crossed the entry of one of the two decision arms. If after 5 min the mosquito did not make any choice and stayed either in the central box or the main arm, then this individual was considered as inactive. A camera placed above the olfactometer allowed us to record the flight of each of the tested insects. Then, flight trajectories were recreated with video tracking using MATLAB (toolbox: DLT digitizing tools) and a mean flight velocity of the mosquitoes was calculated (as in Vinauger et al. 2018).

Data analyses

Comparisons of mosquito choices in the olfactometer Binary data collected in the olfactometer were analyzed (i.e., the number of mosquitoes in each choice arm) and all statistical tests were computed using R software (R Development Core Team 2022). Comparisons were performed by means of the Exact Binomial test ($\alpha = 0.05$). For each treatment, the distribution of mosquitoes in the olfactometer was either compared to a random distribution of 50% on each arm of the maze or to the distribution of the corresponding control when appropriate. For binary data, the standard errors (SE) were calculated as in Le (2003):

$$SEM = \left(\frac{p(1-p)}{n} \right)^{\frac{1}{2}}$$

A preference index (PI) was calculated as follows: $PI = [(number\ of\ females\ in\ the\ odor\ arm - number\ of\ females\ in\ the\ control\ arm) / (total\ number\ of\ active\ mosquitoes)]$. A PI with a positive value indicates that the mosquitoes were attracted to the odor while a PI with a negative value indicates that mosquitoes were repelled. A PI of 0 indicates that 50% of insects chose the odor arm and 50% the control arm (adapted from Schwaerzel et al. 2003).

In addition, the total number of mosquitoes making a decision (i.e., entering one of the two decision arms)

as well as the number of mosquitoes that remained in the first sections of the olfactometer were used to calculate the proportion of “active” mosquitoes at each temperature. These numbers were compared by means of a Chi-square test for experiments conducted in groups and by means of Binomial Exact tests for experiments conducted individually.

Comparisons of mosquito velocity

Flight patterns were recreated with video tracking using MATLAB (toolbox: DLT digitizing tools) and the mean flight velocity of the mosquitoes was calculated. A two-way ANOVA was performed to evaluate the impact of both temperature and odor on the mosquitoes’ flight velocity.

Electroantennography (EAG)

Mosquito head preparation

Electroantennograms were performed following the procedure by Lahondère (2021). Briefly, the head of the mosquitoes was excised and one segment of the tip of each antenna was cut off with fine scissors under a binocular microscope (Carl Zeiss, Oberkochen, Germany). The head was then mounted on an electrode made of a silver wire 0.01 (A-M Systems, Carlsborg, WA, USA) and a borosilicate pulled capillary (0.78 mm I.D. Sutter Instrument Company, Novato, CA, USA) filled with a 1:3 mix of saline solution (Beyenbach and Masia 2002) and electrode gel (Parker Laboratories, Fairfield, NJ, USA) to limit desiccation during the experiment. The head was mounted by the neck on the reference electrode. The preparation was then moved to the EAG setup and using a micromanipulator (Narishige, Japan) the tips of the antennae were inserted, under the microscope (Optiphot-2, Nikon, Tokyo, Japan) into a recording electrode, identical to the reference electrode. The mounted head was oriented at 90° from the main airline, which was carrying clean air (Praxair, Danbury, CT, USA) at either 20, 25, or 30°C and volatiles from the syringe to the preparation for the whole duration of the experiment. Two pulses of a duration of 2.5 s and separated by 10 s were delivered to the mosquito head preparation by switching a solenoid valve controlled by a custom MATLAB script. All the chemicals tested in the olfactometer (except for CO₂, which is detected by the mosquito palps) were used for these experiments. Pulses of clean air were used as a control as well as 100% ethanol (solvent of DEET).

Data acquisition and analyses

Electrophysiological signals were amplified 100× and filtered (0.1–500 Hz) (A-M Systems Model 1800,

Sequim, WA, USA), recorded and digitized at 20 Hz using WinEDR software (Strathclyde Electrophysiology Software, Glasgow, UK) and a BNC-2090A analog-to-digital board (National Instruments, Austin, TX, USA) on a personal computer. A Humbug noise eliminator (Quest Scientific, Vancouver, Canada) was used to eliminate electrical noise. The voltage pulse delivered to the solenoid valve was recorded simultaneously, on a separate channel, and a custom-written R script was used to detect the onset of the odor pulses to trigger averaging responses to each stimulus. The electrophysiological signal collected was further filtered (Butterworth digital filter set as a low pass with a Nyquist frequency of 0.99; R package *signal*). Amplitude of the EAG response was determined as the difference between the mean baseline signal in a 2-s window immediately preceding the onset of the odor pulse, and largest voltage deflection in a 5-s window following the onset of the odor pulse. A generalized linear model (glm) was fitted to the data, as: $EAG\ Response \sim Temperature * odor\ type$. As a post-hoc analysis, general linear hypotheses and multiple comparisons were tested with the *glht* of the *multcomp* package in R.

Results

Flight arena

To examine how temperature modulates flight kinematics, mosquitoes were tethered by the thorax and maintained in an air flow at different temperatures in an LED arena (Fig. 1A). First, we found that the average baseline wingbeat frequency was lower at 20°C (408.8 ± 11.2 Hz; mean \pm SEM) than at 25°C (449.3 ± 11 Hz) and 30°C (501.2 ± 11.5 Hz) (Student's *t*-tests, $P < 0.01$) (Fig. 1B). Regarding the amplitude, we found that the baseline was higher at 25°C (4.39 ± 0.45 a.u.) than 20°C (3.91 ± 0.38 a.u.) and 30°C (3.14 ± 0.50 a.u.) (Student's *t*-tests, all $P < 0.01$) (Fig. 1B). We also found that with increasing temperature, higher the numbers of mosquitoes responded (20°C: 43%; 25°C: 64%, and 30°C: 84%; Binomial Exact tests: 20–25: $P = 0.08$; 25–30: $P = 0.05$; 20–30: $p = 0.0002$) (Fig. 1C). As a control for a potential mechanical perturbation induced by the onset of the odor delivery, we stimulated mosquitoes with a pulse of the carrier gas (N₂) at the same flow rate. Mosquitoes did not change their wingbeat frequency (Student's *t*-test, $t = -1.1013$, $df = 10$, $P = 0.2966$) or wingbeat amplitude ($t = 0.18055$, $df = 10$, $P = 0.8603$) in response to the N₂ pulse (Fig. 1D).

Next, changes in wingbeat frequency and amplitude in response to pulses of CO₂ were measured. We found that the temperature affected mosquitoes' response to CO₂ and that the average increase in wingbeat fre-

quency was lower at 20°C (4.62 ± 1.28 Hz) than at 25°C (12.2 ± 4.22 Hz) and 30°C (22.22 ± 5.06 Hz) (20–25: $t = -1.7178$, $df = 15.396$, $P = 0.05$; 25–30: $t = -1.5206$, $df = 23.841$, $P = 0.07$; 20–30: $t = -3.3693$, $df = 13.543$, $P = 0.002$) (Fig. 1D). Moreover, the average changes in amplitude in response to a pulse of CO₂ gradually increased with increasing temperatures (20°C: 0.28 ± 0.11 a.u., 25°C: 0.73 ± 0.19 a.u., and 30°C: 0.98 ± 0.29 a.u.; 20–25: $t = -2.0295$, $df = 20.799$, $P = 0.03$; 25–30: $t = -0.7196$, $df = 21.229$, $P = 0.24$; 20–30: $t = -2.2695$, $df = 15.431$, $P = 0.02$) (Fig. 1D). The intensity of the responses to the CO₂ pulse was higher at 30°C than at 20°C (Student's *t*-tests, $P < 0.01$).

Olfactometer

Response to carbon dioxide in groups

To test whether the temperature of the environment could affect the mosquito's olfactory behaviors, we first measured their response to carbon dioxide, an important cue used by the insects for host-seeking, in the olfactometer under different thermal conditions. We found that mosquito groups were more attracted by CO₂ when tested at 30°C compared to 20 and 25°C (Binomial tests, $P < 0.05$) (Fig. 2A). Moreover, the proportion of mosquitoes that made a choice in the olfactometer was also higher at higher temperatures (5.4% at 20°C, 45.3% at 25°C, and 64.8% at 30°C; Chi₂, $P < 0.05$) (Fig. 2B).

Individual response to ecologically relevant odors

To determine whether releasing the mosquitoes in groups had any effect, and to easily track the behavior of the mosquitoes, we conducted the same experiment with individual female mosquitoes. We confirmed the results we obtained for the response to CO₂ in groups and found that higher temperatures elicited a stronger level of attraction for the odor (Binomial tests, $P < 0.05$) (Fig. 3A). We also found that hexen-1-ol, benzaldehyde, and DEET significantly repelled more mosquitoes at higher temperatures (Binomial tests, $P < 0.05$) (Fig. 3A). Interestingly, for some other odors such as L-(+)-lactic acid, 1-octen-3-ol, and nonanol, no effect of the temperature was evinced on the mosquito choice (Fig. 3A, top panel). However, the overall level of activity was generally higher at higher temperature, independently of the odor tested (Binomial tests, $P < 0.05$) (Fig. 3B).

Comparisons of mosquito velocity

We then compared the flight velocity of the mosquitoes and found that temperature affected it for some odors [e.g., benzaldehyde, hexen-1-ol], i.e., mosquitoes




Fig. 2 (A) Responses of groups of female mosquitoes to carbon dioxide tested in the y-maze olfactometer at different ambient temperatures (20, 25, and 30°C). Mosquitoes were released in groups and given a choice between CO₂ (2000 ppm) (gray bars) and clean air (white bars). Number of groups of mosquitoes tested at 20°C ($N = 12, n = 240$), 25°C ($N = 14, n = 267$), and 30°C ($N = 9, n = 179$). **(B)** Proportion of active mosquitoes (i.e., the number of mosquitoes that entered one of the two decision arms over the total number of individuals tested) at different temperatures. Asterisks indicate distributions that are significantly different from random (Binomial tests; $P < 0.05$). Vertical bars represent S.E.M.

Table 1 Results of the analysis of variance (two-way ANOVA with interaction) on the mean flight velocity of female mosquitoes under different thermal conditions and tested for their responses to different odors.

Source of variation	Df	Sum sq	Mean sq	F-value	Pr(>F)
Temperature	2	185	92.50	6.423	0.001772**
Odor	6	1524	254.05	17.641	< 2e-16***
Temperature:Odor	12	573	47.75	3.316	0.000124***
Residuals	461	6639	14.40		

Df: degree of freedom, Sum sq: sum square, Mean sq: mean square, F-value: Mean square/residual, Pr(>F): P-value. Statistically significant differences: $P < 0$ "***," $P < 0.001$ "**."

flew faster at higher temperatures (Benzaldehyde: 20°C = 13 ± 1.07 cm.s⁻¹; 25°C = 19.1 ± 0.72 cm.s⁻¹; 30°C = 18.7 ± 1.22 cm.s⁻¹; Hexen-1-ol: 20°C = 14.5 ± 1.21 cm.s⁻¹; 25°C = 14.5 ± 0.57 cm.s⁻¹; 30°C = 21.9 ± 4.20 cm.s⁻¹], but not for others (e.g., DEET, nonanol) (DEET: 20°C = 19.3 ± 2.30 cm.s⁻¹; 25°C = 16.1 ± 0.94 cm.s⁻¹; 30°C = 18.7 ± 0.79 cm.s⁻¹; Nonanol: 20°C = 15.4 ± 1.29 cm.s⁻¹; 25°C = 18.1 ± 1.55 cm.s⁻¹; 30°C = 16.1 ± 0.92 cm.s⁻¹) (mean \pm S.E.M., see ANOVA results, Table 1) (Fig. 4A and B).

Electroantennography

Recording from the antennae of mosquitoes allowed us to quantify the peripheral olfactory sensitivity under different temperatures. Fitting a generalized linear model on the EAG data showed a significant effect of temperature (Analysis of deviance: $P < 0.001$) and of the odor identity ($P < 0.0001$) on the electrophysiological responses. Overall, responses at 25°C were significantly different from responses at 20 and 30°C (Simultaneous tests for general linear hypotheses; Multiple comparisons of means: Tukey Contrasts; 20 vs 25°C: $P < 0.001$; 25 vs 30°C: $P = 0.003$). However, responses at 20 and 30°C were overall not significantly different from each other (Simultaneous tests for general linear hypotheses; 20 vs 30°C: $P = 0.73$) (Fig. 5).

No significant effect of the interaction between temperature and the odor identity was found (Analysis of deviance: $P = 0.078$), although we observed a trend for some odors (e.g., octenol, nonanol, hexanol, benzaldehyde, ethanol), but not others (e.g., lactic acid, DEET), to display an optimum in EAG response at 25°C (Fig. 5).

Discussion

The present study provides clear evidence that temperature modulates mosquito olfactory behavior in both tethered preparations and free flight. In addition, our




Fig. 3 (A) Responses of individual female mosquitoes to different odors, at different environmental temperatures (20, 25, and 30°C). Mosquitoes were tested individually and given a choice between the test odor and a control (i.e., the solvent of the odor). Mosquito choices are represented as the preference index calculated from the distribution of insects in the olfactometer. Each bar represents an experimental group tested at either 20 (blue bars), 25 (purple bars), or 30°C (red bars). Odor tested: L-(+) lactic acid (20°C: $n = 19$, 25°C: $n = 23$, and 30°C: $n = 29$), octenol (20°C: $n = 12$, 25°C: $n = 24$, and 30°C: $n = 29$), nonanol (20°C: $n = 16$, 25°C: $n = 34$, and 30°C: $n = 31$), hexanol (20°C: $n = 19$, 25°C: $n = 29$, and 30°C: $n = 29$), DEET (20°C: $n = 21$, 25°C: $n = 40$, and 30°C: $n = 32$), and CO₂ (20°C: $n = 11$, 25°C: $n = 18$, and 30°C: $n = 13$). As a control, mosquitoes were also exposed to two clean air currents (20°C: $n = 5$, 25°C: $n = 30$, and 30°C: $n = 22$). **(B)** Proportion of active mosquitoes (i.e., the number of mosquitoes that entered one of the two decision arms over the total number of individuals tested) at different temperatures. Odor tested: L-(+) lactic acid (20°C: $N = 40$, 25°C: $N = 38$, and 30°C: $N = 33$), octenol (20°C: $N = 40$, 25°C: $N = 32$, and 30°C: $N = 34$), nonanol (20°C: $N = 39$, 25°C: $N = 44$, and 30°C: $N = 33$), hexanol (20°C: $N = 36$, 25°C: $N = 44$, and 30°C: $N = 34$), DEET (20°C: $N = 40$, 25°C: $N = 46$, and 30°C: $N = 38$), and CO₂ (20°C: $N = 19$, 25°C: $N = 20$, and 30°C: $N = 13$). A clean air control where mosquitoes are exposed to two clean air currents was also performed (20°C: $N = 27$, 25°C: $N = 38$, and 30°C: $N = 31$). Asterisks indicate distributions that are significantly different from random (Binomial tests; $P < 0.05$). Vertical bars represent S.E.M.

electrophysiological results show that the effect of ambient temperature on mosquito behavior is mediated by a modulation of the antennal sensitivity to odors. The approach employed here allowed us to discriminate between effects on the overall activity of mosquitoes and effects on the oriented response to odorants. Indeed, the proportion of responsive mosquitoes tended to be higher when the temperature increased in both the flight arena and the olfactometer, but the response of the mosquitoes was modulated in an odor-dependent manner. While a larger proportion of mosquitoes made a choice at higher temperature in free flight, this response remained neutral for several chemicals at the three tested temperatures. Interestingly, a stronger level of attraction (i.e., CO₂) or aversion (i.e., benzaldehyde, DEET, and hexen-1-ol) were noted for several chem-

icals, indicating that temperature affects mosquito olfactory choices as well as their flight kinematics. In the arena, we noted that baseline wing beat frequency proportionally increased with temperature, while the amplitude of wing motion was maximal at 25°C. It is well known that ambient temperature (T_a) affects insect flight (e.g., Heinrich 1993; Reinhold et al. 2018) and that the flight force generated by insect wings is affected by changes in stroke amplitude and frequency (Lehmann and Dickinson 1997). Using a mosquito flight mill, Rowley and Graham (1968) found that 21°C is the optimal flight temperature in terms of flight distance and duration but indicated that the maximum flight speed occurred between 27 and 32°C. Christophers (1960) also highlighted that flight frequency increases with T_a , reporting 367 beat s⁻¹ at 18°C vs 427 beat s⁻¹ at 25°C in




Fig. 4 (A) Examples of reconstructed flight trajectories obtained via video-tracking of mosquitoes in the olfactometer at 25°C. Multiple individual tracks were overlapped to obtain the present figures for mosquitoes tested for their response to CO₂ (left; $N = 18$), or two control clean air currents (right; $N = 30$). Tracks are color coded as a function of the instantaneous flight velocity of the insect. The dark gray circle represents the arm delivering CO₂, while the white circle represents the control side. **(B)** Mean flight velocities (cm s^{-1}) of female mosquitoes in the olfactometer. Each boxplot represents an experimental group tested with the different odors at either 20°C (blue box plots), 25°C (purple box plots), or 30°C (red boxplots). Dots represent the mean flight velocity of individual mosquitoes; the central bar represents the median and the upper and lower bars of the plot represent the first and third quartiles, respectively. A two-way ANOVA was performed to compare the flight velocities under different temperature regimes according to the tested odor. Statistical results are depicted in Table I. Vertical bars represent S.E.M.

Ae. aegypti. Gopfert et al. (1999) and Arthur et al. (2014) recorded higher average frequencies, ranging from 445 to 510 Hz at medium temperature (23°C). More recently, Villarreal et al. (2017) showed an increase of 8–13 Hz in wing beat frequency for every 1°C gain in *Ae. aegypti*. Temperature also affected mosquito flight velocity in the olfactometer, which increased in response to some odors but not others. In our olfactometer exper-

iments, in both group and individual assays, we found that more mosquitoes made a choice toward the arm delivering the CO₂ at 30 than 20°C. It is worth mentioning that more mosquitoes were active at higher temperatures but the nature of their choice (i.e., the preference for the test odor arm among the mosquitoes that chose one of the two arms) also changed at higher temperatures.




Fig. 5 Violin plots of the EAG responses ($-mV$) to the different chemicals tested under different thermal conditions (20, 25, and $30^{\circ}C$). The central black dot indicates the mean, and the vertical bar represents S.E.M. Above each violin plot is a representative EAG response. The gray bar indicates the odor pulse. Note the typical “V” shape of the EAG responses as well as the absence of antennal response to the solvent (no odor) control, mineral oil.

Temperature is known to affect neuronal activity in both vertebrates and invertebrates (Martin et al. 2011). In the present study, we assessed whether the peripheral detection of chemicals by the antennae was affected by T_a . We found that responses of the antennae varied with T_a but were also odor-dependent. Our data indicate an optimum at $25^{\circ}C$ for antennal sensitivity. Martin et al. (2011) found that exposure to a higher T_a led to an increase in the EAG response amplitude in *Drosophila*. Moreover, using single sensillum recordings (SSRs), they noted that the ORNs were directly impacted by T_a . It is worth mentioning that although sensitivity peaks at $25^{\circ}C$ in the EAG, some odors still show higher behavioral attraction levels at $30^{\circ}C$, which could be due to an effect of the temperature at the central level, e.g., at the level of the antennal lobes. Moreover, a higher antennal sensitivity did not necessarily correlate with a

higher behavioral response, as expected for odors presented alone. This further supports the hypothesis of an additional layer of modulation of T_a , at the level of central olfactory processing. Future experiments including antennal lobe recordings will reveal whether T_a has an impact on signal processing in the brain. Moreover, testing the impact on temperature on ORNs sensitivity in mosquitoes could also help determine why T_a affects behavioral responses to certain odors but not others.

Aedes aegypti is a major disease vector and an invasive species, which global distribution is expected to shift because of global warming (Reinhold et al. 2018). Increased average daily temperatures will impact the mosquito's general activity, including flight, as well host-seeking and biting behavior, which could have major consequences for public health due to increased pathogen transmissions. It is thus critical to understand

how T_a affects mosquito behavior and in particular olfactory behaviors, which are central to host seeking. Finally, as we are facing challenges in mosquito control due, in part, to insecticide resistance, a better understanding of how environmental factors affect mosquito behaviors can provide important knowledge to trap and bait design. The present data provide critical insights on how T_a affects both the mosquito olfactory system and its flight kinematics, which can be integrated in traps relying on olfactory baits as well as acoustic traps (Andrés et al. 2020).

Funding

This work was supported by *The Journal of Experimental Biology Travelling Fellowship* awarded to C.L. J.L. and K.T. were supported by *The Mary Gates Fellowship*. We acknowledge the support of the Air Force Office of Scientific Research [grants FA9550-16-1-0167 (JAR), and FA9550-21-1-0101 (JAR)]; the National Science Foundation [grant 2121935 (JAR)]; and the National Institutes of Health [grant R01AI148300 (JAR)].

Acknowledgments

We thank Binh Nguyen and Mikayla Kiyokawa for taking care of the mosquito colony. We would like to thank Raymond Huey for helpful discussions.

Conflicts of Interest

The authors have no conflict of interest to declare.

Data availability

Data is available upon request.

References

- Andrés M, Su MP, Albert J, Cator LJ. 2020. Buzzkill: targeting the mosquito auditory system. *Current Opin Insect Sci* 40:11–7.
- Arthur BJ, Emr KS, Wyttenbach RA, Hoy RR. 2014. Mosquito (*Aedes aegypti*) flight tones: frequency, harmonicity, spherical spreading, and phase relationships. *J Acoust Soc Am* 135:933–41.
- Barrozo RB, Lazzari CR. 2004. The response of the blood-sucking bug *Triatoma infestans* to carbon dioxide and other host odours. *Chem Senses* 29:319–29.
- Beyenbach K, Masia R. 2002. Membrane conductances of principal cells in Malpighian tubules of *Aedes aegypti*. *J Insect Physiol* 48:375–86.
- Christophers S. 1960. *Aedes aegypti* (L.) the yellow fever mosquito: its life history, bionomics and structure. New York: Cambridge University Press. p. 750
- Cork A, Park KC. 1996. Identification of electrophysiologically active compounds for the malaria mosquito, *Anopheles gambiae*, in human sweat extracts. *Med Vet Entomol* 10:269–76.
- Clements AN. 1992. The biology of mosquitoes. Vol. I. Development, nutrition and reproduction. London: Chapman & Hall. p. 509.
- Eiras AE, Jepson PC. 1991. Host location by *Aedes aegypti* (Diptera: culicidae): a wind tunnel study of chemical cues. *Bull Entomol Res* 81:151–60.
- Eldridge BF. 1968. The effect of temperature and photoperiod on blood-feeding and ovarian development in mosquitoes of the *Culex pipiens* complex. *Am J Trop Med Hyg* 17:133–40.
- Geier M, Sass H, Boeckh J. 1996. A search for components in human body odour that attract females of *Aedes aegypti*. *Ciba Found Symp* 200:132–44.
- Gopfert MC, Briegel H, Robert D. 1999. Mosquito hearing: sound-induced antennal vibrations in male and female *Aedes aegypti*. *J Exp Biol* 202:2727–38.
- Götz KG. 1987. Course-control, metabolism and wing interference during ultralong tethered flight in *Drosophila melanogaster*. *J Exp Biol* 128:35–46.
- Heinrich B. 1976. Heat exchange in relation to blood flow between thorax and abdomen in bumblebees. *J Exp Biol* 64:561–85.
- Heinrich B. 1979. Keeping a cool head: honeybee thermoregulation. *Science* 205:1269–71.
- Heinrich B. 1993. The hot-blooded insects: mechanisms and evolution of thermoregulation. Berlin, Heidelberg: Springer.
- Herrera CM. 1997. Thermal biology and foraging responses of insect pollinators to the forest floor irradiance mosaic. *Oikos* 78:601–11.
- Huey RB, Stevenson RD. 1979. Integrating thermal physiology and ecology of ectotherms: a discussion of approaches. *Am Zool* 19:357–66.
- Jafari M, Ansari-Pour N. 2019. Why, when and how to adjust your P values? *Cell Journal (Yakhteh)* 20:604.
- Kirby MJ, Lindsay SW. 2004. Responses of adult mosquitoes of two sibling species, *Anopheles arabiensis* and *A. gambiae* ss (Diptera: culicidae), to high temperatures. *Bull Entomol Res* 94:441–8.
- Lahondère C. 2021. A step-by-step guide to mosquito electroantennography. *J Viz Exp* 169:e62042.
- Lahondère C, Bonizzoni M. 2022. Thermal biology of invasive *Aedes* mosquitoes in the context of climate change. *Curr Opin Insect Sci* 51:100920.
- Lahondère C, Lazzari CR. 2012. Mosquitoes cool down during blood feeding to avoid overheating. *Curr Biol* 22:40–5.
- Lahondère C, Insausti TC, Paim RM, Luan X, Belev G, Pereira MH, Ianowski JP, Lazzari CR. 2017. Countercurrent heat exchange and thermoregulation during blood-feeding in kissing bugs. *Elife* 6:e26107.
- Lazzari CR. 2009. Orientation towards hosts in haematophagous insects: an integrative perspective. *Adv in Insect Phys* 37: 1–58.
- Lazzari CR, Fauquet A, Lahondère C, Araújo RN, Pereira MH. 2021. Soft ticks perform evaporative cooling during blood-feeding. *J Insect Physiol* 130:104197.
- Lanciani CA, Le TM. 1995. Effect of temperature on the wing length-body weight relationship in *Anopheles quadrimaculatus*. *J Am Mosq Control Assoc* 11:241–3.
- Le CT. 2003. *Introductory biostatistics*. Hoboken, NJ: John Wiley and Sons Ltd.
- Lehmann FO, Dickinson MH. 1997. The changes in power requirements and muscle efficiency during elevated force

- production in the fruit fly *Drosophila melanogaster*. *J Exp Biol* 200:1133–43.
- Lyimo EO, Takken W, Koella JC. 1992. Effect of rearing temperature and larval density on larval survival, age at pupation and adult size of *Anopheles gambiae*. *Entomol Exp Appl* 63:265–71.
- Martin F, Riveron J, Alcorta E. 2011. Environmental temperature modulates olfactory reception in *Drosophila melanogaster*. *J Insect Physiol* 57:1631–42.
- May ML. 1976. Thermoregulation and adaptation to temperature in dragonflies (Odonata: anisoptera). *Ecol Monogr* 46:1–32.
- Paaijmans KP, Blanford S, Chan BH, Thomas MB. 2011. Warmer temperatures reduce the vectorial capacity of malaria mosquitoes. *Biol Lett* 8:rsbl20111075.
- R Development Core Team. 2022. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, <http://www.R-project.org>.
- Reinhold JM, Lazzari CR, Lahondère C. 2018. Effects of the environmental temperature on *Aedes aegypti* and *Aedes albopictus* mosquitoes: a review. *Insects* 9:158.
- Reinhold JM, Shaw R, Lahondère C. 2021. Beat the heat: *Culex quinquefasciatus* regulates its body temperature during blood feeding. *J Therm Biol* 96:102826.
- Reiser MB, Dickinson MH. 2008. A modular display system for insect behavioral neuroscience. *J Neurosci Methods* 167:127–39.
- Riveron J, Boto T, Alcorta E. 2009. The effect of environmental temperature on olfactory perception in *Drosophila melanogaster*. *J Insect Physiol* 55:943–51.
- Rowley WA, Graham CL. 1968. The effect of temperature and relative humidity on the flight performance of female *Aedes aegypti*. *J Insect Physiol* 14:1251–7.
- Rueda LM, Patel KJ, Axtell RC, Stinner RE. 1990. Temperature-dependent development and survival rates of *Culex quinquefasciatus* and *Aedes aegypti* (Diptera: culicidae). *J Med Entomol* 27:892–8.
- Schwaerzel M, Monastirioti M, Scholz H, Friggi-Grelin F, Birman S, Heisenberg M. 2003. Dopamine and octopamine differentiate between aversive and appetitive olfactory memories in *Drosophila*. *J Neurosci* 23:10495–502.
- Thomson RM. 1938. The reactions of mosquitoes to temperature and humidity. *Bull Entomol Res* 29:125–40.
- Trpis M, McClelland GAH, Gillett JD, Teesdale C, Rao TR. 1973. Diel periodicity in the landing of *Aedes aegypti* on man. *Bull World Health Organ* 48:623.
- Villarreal SM, Winokur O, Harrington L. 2017. The impact of temperature and body size on fundamental flight tone variation in the mosquito vector *Aedes aegypti* (Diptera: culicidae): implications for acoustic lures. *J Med Entomol* 54:1116–21.
- Vinauger C, Lutz EK, Riffell JA. 2014. Olfactory learning and memory in the disease vector mosquito *Aedes aegypti*. *J Exp Biol* 217:2321–30.
- Vinauger C, Lahondère C, Wolff GH, Locke LT, Liaw JE, Parrish JZ, Akbari OS, Dickinson MH, Riffell JA. 2018. Modulation of host learning in *Aedes aegypti* mosquitoes. *Curr Biol* 28:333–44.e8.
- Vinauger C, Van Breugel F, Locke LT, Tobin KK, Dickinson MH, Fairhall AL, Akbari OS, Riffell JA. 2019. Visual-olfactory integration in the human disease vector mosquito *Aedes aegypti*. *Curr Biol* 29:2509–16.e5.
- WHO. 2019. World health statistics 2019: monitoring health for the SDGs, sustainable development goals. Geneva: World Health Organization.