



No to *Neocosmospora*: Phylogenomic and Practical Reasons for Continued Inclusion of the *Fusarium solani* Species Complex in the Genus *Fusarium*

 Kerry O'Donnell,^a Abdullah M. S. Al-Hatmi,^b Takayuki Aoki,^c Balázs Brankovics,^d  José F. Cano-Lira,^{e,f}  Jeffrey J. Coleman,^g G. Sybren de Hoog,^{h,i} Antonio Di Pietro,^j Rasmus J. N. Frandsen,^k David M. Geiser,^l Connie F. C. Gibas,^m Josep Guarro,ⁿ Hye-Seon Kim,^a H. Corby Kistler,^o Imane Laraba,^a  John F. Leslie,^p Manuel S. López-Berges,^j Erik Lysøe,^q  Jacques F. Meis,^{h,i} Michel Monod,^r Robert H. Proctor,^a Martijn Rep,^s Carmen Ruiz-Roldán,^j Adnan Šišić,^t  Jason E. Stajich,^u Emma T. Steenkamp,^v Brett A. Summerell,^w Theo A. J. van der Lee,^d Anne D. van Diepeningen,^d Paul E. Verweij,^{h,i} Cees Waalwijk,^d Todd J. Ward,^a Brian L. Wickes,^x  Nathan P. Wiederhold,^m Michael J. Wingfield,^y Ning Zhang,^z  Sean X. Zhang^z

^aMycotoxin Prevention and Applied Microbiology (MPM) Research Unit, ARS-USDA, Peoria, Illinois, USA

^bMinistry of Health, Directorate General of Health Services, Ibri, Oman

^cNational Agriculture and Food Research Organization, Genetic Resources Center, Tsukuba, Ibaraki, Japan

^dWageningen Plant Research, Wageningen University & Research, Wageningen, The Netherlands

^eUniversitat Rovira i Virgili Medical School, Mycology Unit, Reus, Spain

^fISPV, Reus, Spain

^gDepartment of Entomology and Plant Pathology, Auburn University, Auburn, Alabama, USA

^hDepartment of Medical Microbiology, Radboud University Medical Centre, Nijmegen, The Netherlands

ⁱCenter of Expertise in Mycology Radboudumc/CWZ, Radboud University Medical Centre, Nijmegen, The Netherlands

^jDepartamento de Genética, Universidad de Córdoba, Córdoba, Spain

^kDepartment of Biotechnology and Biomedicine, Technical University of Denmark, Kongens Lyngby, Denmark

^lDepartment of Plant Pathology and Environmental Microbiology, Pennsylvania State University, University Park, Pennsylvania, USA

^mDepartment of Pathology, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

ⁿUnitat de Microbiologia, Departament de Ciències Mèdiques Bàsiques, Universitat Rovira i Virgili, Reus, Spain

^oUSDA ARS Cereal Disease Laboratory, University of Minnesota, St. Paul, Minnesota, USA

^pDepartment of Plant Pathology, Kansas State University, Manhattan, Kansas, USA

^qNorwegian Institute of Bioeconomy Research, Division of Biotechnology and Plant Health, Ås, Norway

^rService de Dermatologie, Laboratoire de Mycologie, Centre Hospitalier Universitaire Vaudois (CHUV), Lausanne, Switzerland

^sSwammerdam Institute for Life Science, University of Amsterdam, Amsterdam, The Netherlands

^tDepartment of Ecological Plant Protection, University of Kassel, Witzenhausen, Germany

^uDepartment of Microbiology and Plant Pathology, University of California–Riverside, Riverside, California, USA

^vDepartment of Biochemistry, Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa

^wAustralian Institute of Botanical Science, Royal Botanic Garden and Domain Trust, Sydney, Australia

^xDepartment of Microbiology, Immunology & Molecular Genetics, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

^yDepartment of Plant Biology, Rutgers University, New Brunswick, New Jersey, USA

^zDepartment of Pathology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

ABSTRACT This article is to alert medical mycologists and infectious disease specialists of recent name changes of medically important species of the filamentous mold *Fusarium*. *Fusarium* species can cause localized and life-threatening infections in humans. Of the 70 *Fusarium* species that have been reported to cause infections, close to one-third are members of the *Fusarium solani* species complex (FSSC), and they collectively account for approximately two-thirds of all reported *Fusarium* infections. Many of these species were recently given scientific names for the first time

Citation O'Donnell K, Al-Hatmi AMS, Aoki T, Brankovics B, Cano-Lira JF, Coleman JJ, de Hoog GS, Di Pietro A, Frandsen RJN, Geiser DM, Gibas CFC, Guarro J, Kim H-S, Kistler HC, Laraba I, Leslie JF, López-Berges MS, Lysøe E, Meis JF, Monod M, Proctor RH, Rep M, Ruiz-Roldán C, Šišić A, Stajich JE, Steenkamp ET, Summerell BA, van der Lee TAJ, van Diepeningen AD, Verweij PE, Waalwijk C, Ward TJ, Wickes BL, Wiederhold NP, Wingfield MJ, Zhang N, Zhang SX. 2020. No to *Neocosmospora*: phylogenomic and practical reasons for continued inclusion of the *Fusarium solani* species complex in the genus *Fusarium*. mSphere 5:e00810-20. <https://doi.org/10.1128/mSphere.00810-20>.

Editor Aaron P. Mitchell, University of Georgia
This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.
Address correspondence to Kerry O'Donnell, kerry.odonnell@usda.gov.

This is contribution no. 20-355-J from the Kansas Agricultural Experiment Station.

Published 16 September 2020

by a research group in the Netherlands, but they were misplaced in the genus *Neocosmospora*. In this paper, we present genetic arguments that strongly support inclusion of the FSSC in *Fusarium*. There are potentially serious consequences associated with using the name *Neocosmospora* for *Fusarium* species because clinicians need to be aware that fusaria are broadly resistant to the spectrum of antifungals that are currently available.

KEYWORDS clinical mycology, evolution, fungi, phylogenetics, taxonomy

Robust taxonomy facilitates communication and should be cogent and consider the scientific and practical ramifications. As presently circumscribed, the monophyletic genus *Fusarium* comprises over 300 phylogenetically distinct species (i.e., phylopecies) distributed among 23 evolutionary lineages referred to as species complexes (1; D. M. Geiser, A. M. S. Al-Hatmi, T. Aoki, T. Arie, et al., submitted for publication). Over 70 *Fusarium* species distributed among 12 of these complexes have been implicated in mycotic infections of humans and other animals (Fig. 1A) (2, 3). Of these, members of the *Fusarium solani* species complex (FSSC) account for approximately two-thirds of all reported fusarioses and at least 23 of the clinically relevant phylospecies (Fig. 1B). There has been a consensus for over a century that the FSSC is part of *Fusarium*, which was affirmed by molecular phylogenetic analyses (4) and codified in a proposal to recognize *Fusarium* as a monophyletic group that includes the FSSC (5). However, there has been a recent push to divide *Fusarium* into seven genera and specifically to move the FSSC, the most important group of fusarial human pathogens, into the genus *Neocosmospora* (6–8).

In this paper, we demonstrate that these nomenclatural changes were based on the incorrect interpretation of the phylogenetic data regarding the Geiser et al. (5) circumscription of *Fusarium* that includes the FSSC. In addition, these papers published an additional 31 FSSC species and 18 new combinations under the genus *Neocosmospora* (7, 8). A new 19-locus phylogeny, however, removes all doubt about inclusion of the FSSC in a taxonomically sound and historically consistent concept of *Fusarium* (Geiser et al., submitted). Thus, Aoki et al. (9) recombined these species in *Fusarium*, as the more robust phylogeny reaffirmed the Geiser et al. (5) circumscription of *Fusarium*. Here, we correct the record and argue that support for inclusion of the FSSC in *Fusarium* is more definitive than ever. We also cite concerns of medical mycologists if human-pathogenic fusaria are unnecessarily fragmented into multiple genera and urge continued use of the genus name *Fusarium* for human pathogens in the FSSC, along with the *F. dimerum* and *F. ventricosum* species complexes (10).

Fusarial infections most frequently encountered in healthy individuals include onychomycoses and keratitis; the latter often occurs after a traumatic introduction to the eye or is associated with soft contact lens wear (11). *Fusarium* keratitis outbreaks that occurred in 2005 and 2006 in the United States, Singapore, and Hong Kong were associated with the use of Bausch & Lomb's ReNu with MoistureLoc contact lens solution (12, 13). The finding that most of these infections were caused by FSSC species/multilocus sequence types (MLSTs) that are common in and apparently adapted to plumbing systems (14) suggests that patients inadvertently contaminated their contact lenses through poor hygiene. Though relatively rare compared to invasive candidiasis and aspergillosis (15), fusaria can cause life-threatening disseminated opportunistic infections in persistently neutropenic immunocompromised and immunosuppressed patients (16). Mortality among this patient population is high because fusaria are resistant to the broad spectrum of antifungals currently available and dissemination frequently occurs (17). Successful treatment of these patients often requires antifungal drug combinations in addition to the patient recovering from neutropenia (18).

Prior to the introduction of MLST-based diagnoses and recognition of phylospecies based on genealogical exclusivity (19), mycoses caused by members of the FSSC were typically reported as *F. solani* (20), due to a lack of diagnostic micromorphological




FIG 1 (A) Partitioned maximum likelihood bootstrapped (ML-BS) phylogeny of *Fusarium* inferred from exonic sequences of 19 housekeeping genes totaling 54.99 kb; 20.3 kb of characters were parsimony informative characters (PIC). The phylogram was rooted on sequences of two outgroup species, *Beauveria bassiana* and *Trichoderma brevicompactum*. Evolutionary relationships among the 12 species complexes that contain clinically relevant fusaria were completely resolved based on ML-BS values between 91 and 100%. The node identified by F1, which represents the generic limits of *Fusarium*, received 100% ML-BS support based on 5,000 pseudoreplicates of the data conducted with IQ-TREE (27). (B) ML-BS phylogeny of clade 3 of the *Fusarium solani* species complex (FSSC) inferred from portions of three loci (*TEF1*, *RPB2*, and the internal transcribed spacer [ITS] ribosomal DNA [rDNA]). All 23 species implicated in fusarioses of humans and other animals are nested in FSSC clade 3. Numbers on nodes represent ML-BS support based on 5,000 pseudoreplicates of the data. FSSC numbers in brackets represent the *ad hoc* nomenclature previously used to distinguish species (22). T, ex-type strains; IT, ex-isotype strain; NT, ex-neotype strain.

characters among FSSC species. The first robust MLST-based species-level studies on the FSSC, however, revealed two important facts: (i) the FSSC comprises at least 45 phylopecies distributed among three clades and (ii) all of the clinically relevant species were nested within clade 3, one of three major evolutionary subgroups that comprise the FSSC (21, 22). Because Latin binomials were not available for most of the species, Chang et al. (12) adopted an *ad hoc* species/multilocus haplotype nomenclature for the FSSC, in which species and multilocus haplotypes were distinguished, respectively, by Arabic numerals and a lowercase Roman letter. This informal nomenclature was subsequently extended to 35 species in FSSC clade 3 (e.g., FSSC 2-d denotes haplotype d of *F. solani* species complex undescribed species 2) to facilitate accurate communication within the clinical microbiological and phytopathological scientific communities (22).

Following the work of Short et al. (23), where names were assigned to two of the most important human-pathogenic FSSC species, *F. petrophilum* (FSSC 1) and *F. keratoplasticum* (FSSC 2), Sandoval-Denis and Crous (7) and Sandoval-Denis et al. (8) provided names and types for almost all of the remaining phylogenetically diagnosable species. However, these species were described under the name *Neocosmospora*. These authors went to great lengths to promote the use of this genus name but with arguments that either fail to address or inaccurately represent the work of O'Donnell et al. (4) and Geiser et al. (5). In addition, they recombined species originally described in *Fusarium* in this teleomorphic genus, thus promoting a *de facto* dual nomenclature that obscures the fact that *Neocosmospora* is nested phylogenetically within a monophyletic *Fusarium* (5, 24; Geiser et al., submitted). There are potentially serious consequences associated with this because clinicians need to be aware that these fungi are broadly resistant to the spectrum of antifungals that are currently available (25). An identification as *Neocosmospora* would thus fail to connect a case to this crucial knowledge, thereby disregarding the valuable clinical information that distinguishes it from an identification as *Fusarium*. Therefore, we are strongly opposed to this taxonomy because it will confuse stakeholders and may have negative ramifications in clinical microbiology and medicine.

The present article and a companion submission to *Phytopathology* (Geiser et al., submitted) are nearly unanimously supported by the global *Fusarium* research community. Here, we set the record straight by pointing out that the justification of Sandoval-Denis and Crous (7) for describing members of the FSSC as *Neocosmospora* was based on four errant conclusions (i to iv below).

Errant conclusion i. The molecular phylogenetic circumscription of *Fusarium* in the work of Geiser et al. (5) and, by extension, of O'Donnell et al. (4) is polyphyletic.

Response. The claim that the Geiser et al. (5) delimitation of *Fusarium* is polyphyletic is not supported by published data. Indeed, we know of no published data that even suggests this conclusion. The monophyletic circumscription of *Fusarium* presented by Geiser et al. (5) was based on a two-locus phylogeny published by O'Donnell et al. (4); similar clades have been resolved in many other publications (e.g., the "Terminal *Fusarium* Clade" [26]), including that of Lombard et al. (6). In fact, Geiser et al. (5) conservatively and critically evaluated the phylogeny presented in the work of O'Donnell et al. (4), which was based on sequences of the largest (*RPB1*) and second-largest (*RPB2*) RNA polymerase II subunit genes, to ensure that the delimitation of *Fusarium* was robustly connected to a monophyletic group. Two nodes in the *RPB1*/*RPB2* tree (named F1 and F2) were identified as potential taxonomic hypotheses for *Fusarium*, with F1 better representing longstanding use but receiving inferior statistical support compared to that of F2. It is important to note that both F1 and F2 included the FSSC, but F2 excluded the two most basal lineages (i.e., the *F. dimerum* and *F. ventricosum* species complexes). The Geiser et al. (5) hypothesis provisionally retained these species complexes in *Fusarium* (hypothesis F1 in Fig. 1A) until their inclusion or exclusion could be assessed by a more robust comparative phylogenomic data set.

The 19-gene phylogeny recently presented in the paper of Geiser et al. (submitted) and presented here in modified form (Fig. 1A) is such a data set. Both the F1 and F2 taxonomic hypotheses from Geiser et al. (5) received strong statistical support (i.e., 100% maximum likelihood [ML] bootstrap) in the 19-gene analysis. Details of the partitioned ML phylogenetic analysis of the 55.1-kb 19-gene data set conducted with IQ-TREE (27) are presented in the paper by Geiser et al. (submitted) and illustrate the power of comparative phylogenomics for discovering novel phylogenetically informative marker loci. In summary, the phylogenetic uncertainty openly considered by Geiser et al. (5) has been resolved in support of a taxonomic concept for *Fusarium* that includes the FSSC, consistent with longstanding use by medical mycologists and phytopathologists.

Errant conclusion ii. Segregation of the FSSC as *Neocosmospora* represents a "more natural classification."

Response. The term “natural classification” means that all members of a taxon share an ancestor; in other words, they should be monophyletic. Based on our response to errant claim i, *Fusarium*, inclusive of *Neocosmospora*, is monophyletic. The claim that separation of *Neocosmospora* from *Fusarium* represents a more natural classification is without merit because the concept of *Fusarium* that includes the FSSC is defined based on its monophyly, and monophyly is the principal criterion for taxon recognition in modern taxonomy. Even prior to the demise of the dual nomenclature system on 1 January 2013, when teleomorph genera had taxonomic priority, the FSSC had a longstanding taxonomic place within *Fusarium*, with virtually no controversy among stakeholders, especially among medical mycologists and plant pathologists. Certainly, there is great value in continuing to recognize the >90 species in this clade as the “*Fusarium solani* species complex,” representing its status as a distinct clade within a monophyletic *Fusarium*, and maintaining continuity of use. In contrast, use of “*Neocosmospora*” creates an unnecessary communication barrier between the FSSC and over a century of *Fusarium* research. Also note that because *Fusarium* was typified close to a century before *Neocosmospora* (1809 versus 1899), the name *Fusarium* has nomenclatural priority (https://www.iapt-taxon.org/nomen/pages/main/art_11.html).

Fusarium is by far the most widely used name applied to the FSSC, as reflected by a simple Google search. “*Fusarium solani*” received 100 times more hits than “*Neocosmospora*” (~3,190,000 to ~31,600; search conducted 27 May 2020). The longstanding status of the FSSC in *Fusarium* has also been reinforced by annual *Fusarium* educational workshops offered around the world since the 1970s, in particular, the *Fusarium* Laboratory Workshop (28) and the Tropical *Fusarium* Workshops. Highlighting the natural connection between the FSSC and *Fusarium*, in both of these workshops, the FSSC and *F. oxysporum* species complex (FOSC, which Lombard et al. [6] retains in *Fusarium*) are presented in tandem due to the ecological, genomic, morphological, and clinical connections that exist between these two groups.

Errant conclusion iii. Substantial molecular evidence indicates *Fusarium* and *Neocosmospora* are not congeneric.

Response. Absolutely nothing has been published to date to suggest that “*Neocosmospora*” is not nested phylogenetically within *Fusarium* as we define it (4) (Fig. 1). This unassailable fact reflects the central logic of the Geiser et al. (5) taxonomic hypothesis. Lombard et al. (6) hypothesize that *Fusarium* and *Neocosmospora* are two equally ranked clades within a broader clade consisting of at least seven genera, all of which have an established taxonomic history as *Fusarium* species. These competing hypotheses are supportable by the same phylogenetic evidence but are readily distinguished based on their relative practical merits. In fact, the phylogeny can be used as the basis for any number of taxonomic hypotheses within node F1 (Fig. 1A), either as a single genus as it has been treated for over a century, as seven genera, or as potentially dozens of genera.

Errant conclusion iv. Support of the *Fusarium* research community for the Geiser et al. (5) conservation of *Fusarium* in a way that includes the FSSC, with its adherents referred to as “some researchers,” has been understated.

Response. To the contrary, 66 scientists from 16 different countries spanning 6 continents, representing virtually all of the world leaders in *Fusarium* research and the disciplines of medical mycology, plant pathology, molecular biology and genomics, mycotoxicology, and other fields, coauthored Geiser’s proposal to circumscribe a monophyletic *Fusarium* such that it preserves longstanding use (5) (Fig. 1). Now that there is overwhelming statistical support for hypothesis F1 (Fig. 1A), the Geiser et al. (5) circumscription of *Fusarium* will gain even more support from end users.

Just prior to the demise of dual nomenclature, two scholarly studies opened the door to dividing *Fusarium* along teleomorph lines (26, 29), using competing genus names that would require breaking *Fusarium* up into at least seven genera. However, both studies also referred to the clade associated with the F1 node as “The Terminal *Fusarium* Clade” (italics theirs), raising the question, “Why did they use that name if that

clade represented something other than the ideal circumscription of *Fusarium*?" These papers also presented phylogenetics that provided a framework for a monophyletic *Fusarium* that includes the FSSC, as presented here. The FSSC is an integral part of FUSARIUM-ID (30; <http://isolate.fusariumdb.org/blast.php>) and *Fusarium* MLST (31; <http://fusarium.mycobank.org/>), two websites dedicated to identification of fusaria via the Internet. We urge the people who maintain these sites to respect the will of the *Fusarium* community and retain the FSSC in *Fusarium* and their databases. Formal recombinations of the newly proposed *Neocosmospora* names in *Fusarium* were registered in *Index Fungorum*, where they are now validly published (9; Geiser et al., submitted). In addition, we updated our nucleotide accessions in GenBank to reflect the current taxonomy.

End users will ultimately decide what names to use for these fungi (32), and we urge the clinical microbiological community to continue to refer to these etiological agents by their *Fusarium* binomials (see clade 3 in Fig. 1B for binomials of the 23 FSSC members of clinical relevance) or as a member of the FSSC when an identification is made only to the species complex level (33). Using the name "*Neocosmospora*" obscures rather than facilitates communication among basic and applied scientists and regulatory agencies in medicine and agriculture by promoting a taxonomy governed by *de facto* dual nomenclature. When infectious disease specialists need to access clinically relevant literature to manage fusarial infections, they will need to use "*Fusarium*" or "fusariosis," not "*Neocosmospora*" or "neocosmosporosis," as the search terms. With 191,000 Google hits (search conducted on 27 May 2020), fusariosis is well embedded in the medical/clinical literature, whereas "neocosmosporosis" failed to recover even a single clinically relevant record. Applying the work of Hawksworth (34) to this case, the name "*Fusarium*" is crucial for accessing the accumulated knowledge in clinical, medical, and other scientific literature.

ACKNOWLEDGMENTS

This paper has been communicated to C. Schoch, A. N. Miller, and C. Aime of the International Commission on the Taxonomy of Fungi (ICTF; <https://www.fungaltaxonomy.org/ictf-members/>) and to S. A. Redhead (Chair), P. M. Kirk, and Z. Wilhelm de Beer of the Nomenclature Committee for Fungi (NCF; <https://www.ima-mycology.org>).

Special thanks are due to Amy McGovern and Crystal Probyn for expert technical assistance in generating and mining the whole-genome sequence data in the MPM Research Unit, Peoria, IL.

The USDA-ARS National Program for Food Safety, grant NSF DEB-1655980, and Pennsylvania State Agricultural Experiment Station Project 4655 provided funding for this study.

Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

REFERENCES

1. Aoki T, O'Donnell K, Geiser DM. 2014. Systematics of key phytopathogenic *Fusarium* species: current status and future challenges. *J Gen Plant Pathol* 80:189–201. <https://doi.org/10.1007/s10327-014-0509-3>.
2. Al-Hatmi AMS, Meis JF, de Hoog GS. 2016. *Fusarium*: molecular diversity and intrinsic drug resistance. *PLoS Pathog* 12:e1005464. <https://doi.org/10.1371/journal.ppat.1005464>.
3. O'Donnell K, Sutton DA, Wiederhold N, Robert VARG, Crous PW, Geiser DM. 2016. Veterinary fusarioses within the United States. *J Clin Microbiol* 54:2813–2819. <https://doi.org/10.1128/JCM.01607-16>.
4. O'Donnell K, Rooney AP, Proctor RH, Brown DW, McCormick SP, Ward TJ, Frandsen RJN, Lysøe E, Rehner SA, Aoki T, Robert VARG, Crous PW, Groenewald JZ, Kang S, Geiser DM. 2013. *RPB1* and *RPB2* phylogeny support an early Cretaceous origin and a strongly supported clade comprising all agriculturally and medically important fusaria. *Fungal Genet Biol* 52:20–31. <https://doi.org/10.1016/j.fgb.2012.12.004>.
5. Geiser DM, Aoki T, Bacon CW, Baker SE, Bhattacharyya MK, Brandt ME, Brown DW, Burgess LW, Chulze S, Coleman JJ, Correll JC, Covert SF, Crous PW, Cuomo CA, De Hoog GS, Di Pietro A, Elmer WH, Epstein L, Frandsen RJN, Freeman S, Gagkaeva T, Glenn AE, Gordon TR, Gregory NF, Hammond-Kosack KE, Hanson LE, Jiménez-Gasco MDM, Kang S, Kistler HC, Kuldau GA, Leslie JF, Logrieco A, Lu G, Lysøe E, Ma L-J, McCormick SP, Migheli Q, Moretti A, Munaut F, O'Donnell K, Pfenning L, Ploetz RC, Proctor RH, Rehner SA, Robert VARG, Rooney AP, Bin Salleh B, Scandiani MM, Scauflaire J, Short DPG, et al. 2013. One fungus, one name: defining the genus *Fusarium* in a scientifically robust way that preserves long-standing use. *Phytopathology* 103:400–408. <https://doi.org/10.1094/PHYTO-07-12-0150-LE>.
6. Lombard L, van der Merwe NA, Groenewald JZ, Crous PW. 2015. Generic concepts in Nectriaceae. *Stud Mycol* 80:189–245. <https://doi.org/10.1016/j.simyco.2014.12.002>.
7. Sandoval-Denis M, Crous PW. 2018. Removing chaos from confusion: assigning names to common human and animal pathogens in *Neocos-*

- mospora*. Persoonia 41:109–129. <https://doi.org/10.3767/persoonia.2018.41.06>.
8. Sandoval-Denis M, Lombard L, Crous PW. 2019. Back to the roots: a reappraisal of *Neocosmospora*. Persoonia 43:90–185. <https://doi.org/10.3767/persoonia.2019.43.04>.
 9. Aoki T, Geiser DM, Kasson MT, O'Donnell K. 2020. Nomenclatural novelities. Index Fungorum 440:1–5.
 10. Schroers H-J, O'Donnell K, Lamprecht SC, Kammeyer PL, Johnson S, Sutton DA, Rinaldi MG, Geiser DM, Summerbell RC. 2009. Taxonomy and phylogeny of the *Fusarium dimerum* species group. Mycologia 101: 44–70. <https://doi.org/10.3852/08-002>.
 11. van Diepeningen AD, Al-Hatmi AMS, Brankovics B, de Hoog GS. 2014. Taxonomy and clinical spectra of *Fusarium* species: where do we stand in 2014? Curr Clin Micro Rep 1:10–18. <https://doi.org/10.1007/s40588-014-0003-x>.
 12. Chang DC, Grant GB, O'Donnell K, Wannemuehler KA, Noble-Wang J, Rao CY, Jacobson LM, Crowell CS, Sneed RS, Lewis FMT, Schaffzin JK, Kainer MA, Genese CA, Alfonso EC, Jones DB, Srinivasan A, Fridkin SK, Park BJ, Fusarium Keratitis Investigation Team. 2006. Multistate outbreak of *Fusarium* keratitis associated with use of a contact lens solution. JAMA 296:953–963. <https://doi.org/10.1001/jama.296.8.953>.
 13. Khor W-B, Aung T, Saw S-M, Wong T-Y, Tambyah PA, Tan A-L, Beuerman R, Lim LL, Chan W-K, Heng W-J, Lim J, Loh RSK, Lee S-B, Tan DTH. 2006. An outbreak of *Fusarium* keratitis associated with contact lens wear in Singapore. JAMA 295:2867–2873. <https://doi.org/10.1001/jama.295.24.2867>.
 14. Short DPG, O'Donnell K, Zhang N, Juba JH, Geiser DM. 2011. Widespread occurrence of diverse human pathogenic types of the fungus *Fusarium* detected in plumbing drains. J Clin Microbiol 49:4264–4272. <https://doi.org/10.1128/JCM.05468-11>.
 15. Zhang SX, O'Donnell K, Sutton DA. 2019. *Fusarium* and other opportunistic hyaline fungi, p 2132–2162. In Carroll KC, Pfaller MA, Landry ML, McAdam AJ, Patel R, Richter S, Warnock DW (ed), Manual of clinical microbiology, 12th ed ASM Press, Washington, DC.
 16. Nucci M, Anaissie E. 2007. *Fusarium* infections in immunocompromised patients. Clin Microbiol Rev 20:695–704. <https://doi.org/10.1128/CMR.00014-07>.
 17. Al-Hatmi AMS, Bonifaz A, Ranque S, Sybren de Hoog G, Verweij PE, Meis JF. 2018. Current antifungal treatment of fusariosis. Int J Antimicrob Agents 51:326–332. <https://doi.org/10.1016/j.ijantimicag.2017.06.017>.
 18. Guarro J. 2013. Fusariosis, a complex infection caused by a high diversity of fungal species refractory to treatment. Eur J Clin Microbiol Infect Dis 32:1491–1500. <https://doi.org/10.1007/s10096-013-1924-7>.
 19. Taylor JW, Jacobson DJ, Kroken S, Kasuga T, Geiser DM, Hibbett DS, Fisher MC. 2000. Phylogenetic species recognition and species concepts in fungi. Fungal Genet Biol 31:21–32. <https://doi.org/10.1006/fgbi.2000.1228>.
 20. Leslie JF, Summerell BA. 2006. The *Fusarium* laboratory manual. Blackwell Publishing, Ames, IA.
 21. Zhang N, O'Donnell K, Sutton DA, Nalim FA, Summerbell RC, Padhye AA, Geiser DM. 2006. Members of the *Fusarium solani* species complex that cause infections in both humans and plants are common in the environment. J Clin Microbiol 44:2186–2190. <https://doi.org/10.1128/JCM.00120-06>.
 22. O'Donnell K, Sutton DA, Fothergill A, McCarthy D, Rinaldi MG, Brandt ME, Zhang N, Geiser DM. 2008. Molecular phylogenetic diversity, multilocus haplotype nomenclature, and in vitro antifungal resistance within the *Fusarium solani* species complex. J Clin Microbiol 46:2477–2490. <https://doi.org/10.1128/JCM.02371-07>.
 23. Short DPG, O'Donnell K, Thrane U, Nielsen KF, Zhang N, Juba JH, Geiser DM. 2013. Phylogenetic relationships among members of the *Fusarium solani* species complex in human infections and the descriptions of *F. keratoplasticum* sp. nov. and *F. petrophilum* stat. nov. Fungal Genet Biol 53:59–70. <https://doi.org/10.1016/j.fgb.2013.01.004>.
 24. O'Donnell K. 2000. Molecular phylogeny of the *Nectria haematococca-Fusarium solani* species complex. Mycologia 92:919–938. <https://doi.org/10.1080/00275514.2000.12061237>.
 25. Al-Hatmi AM, de Hoog GS, Meis JF. 2019. Multiresistant *Fusarium* pathogens on plants and humans: solutions in (from) the antifungal pipeline? Infect Drug Resist 12:3727–3737. <https://doi.org/10.2147/IDR.S180912>.
 26. Gräfenhan T, Schroers H-J, Nirenberg HI, Seifert KA. 2011. An overview of the taxonomy, phylogeny and typification of nectriaceous fungi in *Cosmospora*, *Acremonium*, *Fusarium*, *Stilbella* and *Volutella*. Stud Mycol 68:79–113. <https://doi.org/10.3114/sim.2011.68.04>.
 27. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum likelihood phylogenies. Mol Biol Evol 32:268–274. <https://doi.org/10.1093/molbev/msu300>.
 28. Leslie JF, Summerell BA. 2006. *Fusarium* laboratory workshops—a recent history. Mycotoxin Res 22:73–74. <https://doi.org/10.1007/BF02956766>.
 29. Schroers H-J, Gräfenhan T, Nirenberg HI, Seifert KA. 2011. A revision of *Cyanonectria* and *Geejayessia* gen nov, and related species with *Fusarium*-like anamorphs. Stud Mycol 68:115–138. <https://doi.org/10.3114/sim.2011.68.05>.
 30. Geiser DM, del Mar Jiménez-Gasco M, Kang S, Makalowska I, Veeraraghavan N, Ward TJ, Zhang N, Kuldau GA, O'Donnell K. 2004. FUSARIUM-ID v 10: a DNA sequence database for identifying *Fusarium*. Eur J Plant Pathol 110:473–479. <https://doi.org/10.1023/B:EJPP.0000032386.75915.a0>.
 31. O'Donnell K, Ward TJ, Robert VARG, Crous PW, Geiser DM, Kang S. 2015. DNA sequence-based identification of *Fusarium*: current status and future directions. Phytoparasitica 43:583–595. <https://doi.org/10.1007/s12600-015-0484-z>.
 32. de Hoog GS, Chaturvedi V, Denning DW, Dyer PS, Frisvad JC, Geiser D, Gräser Y, Guarro J, Haase G, Kwon-Chung K-J, Meis JF, Meyer W, Pitt JI, Samson RA, Taylor JW, Tintelnot K, Vitale RG, Walsh TJ, Lackner M, the ISHAM Working Group on Nomenclature of Medical Fungi. 2015. Name changes in medically important fungi and their implications for clinical practice. J Clin Microbiol 53:1056–1062. <https://doi.org/10.1128/JCM.02016-14>.
 33. Balajee SA, Borman AM, Brandt ME, Cano J, Cuenca-Estrella M, Dannaoui E, Guarro J, Haase G, Kibbler CC, Meyer W, O'Donnell K, Petti CA, Rodriguez-Tudela JL, Sutton D, Velegriaki A, Wickes BL. 2009. Sequence-based identification of *Aspergillus*, *Fusarium*, and *Mucorales* species in the clinical mycology laboratory: where are we and where should we go from here? J Clin Microbiol 47:877–884. <https://doi.org/10.1128/JCM.01685-08>.
 34. Hawksworth DL. 2011. Naming *Aspergillus* species: progress towards one name for each species. Med Mycol 49:570–576. <https://doi.org/10.3109/13693786.2010.504753>.