

Novel Alphaproteobacteria transcribe genes for nitric oxide transformation at high levels in a marine oxygen-deficient zone

Claire E. Elbon,¹ Frank J. Stewart,² Jennifer B. Glass¹

AUTHOR AFFILIATIONS See affiliation list on p. 10.

ABSTRACT Marine oxygen-deficient zones (ODZs) are portions of the ocean where intense nitrogen loss occurs primarily via denitrification and anammox. Despite many decades of study, the identity of the microbes that catalyze nitrogen loss in ODZs is still being elucidated. Intriguingly, high transcription of genes in the same family as the nitric oxide dismutase (*nod*) gene from *Methylospirillum* has been reported in the anoxic core of ODZs. Here, we show that the most abundantly transcribed *nod* genes in the Eastern Tropical North Pacific ODZ belong to a new order (UBA11136) of Alphaproteobacteria, rather than *Methylospirillum* as previously assumed. Gammaproteobacteria and Planctomycetia also transcribe *nod*, but at lower relative abundance than UBA11136 in the upper ODZ. The *nod*-transcribing Alphaproteobacteria likely use formaldehyde and formate as a source of electrons for aerobic respiration, with additional electrons possibly from sulfide oxidation. They also transcribe multiheme cytochrome (here named *ptd*) genes for a putative porin-cytochrome protein complex of unknown function, potentially involved in extracellular electron transfer. Molecular oxygen for aerobic respiration may originate from nitric oxide dismutation via cryptic oxygen cycling. Our results implicate Alphaproteobacteria order UBA11136 as a significant player in marine nitrogen loss and highlight their potential in one-carbon, nitrogen, and sulfur metabolism in ODZs.

IMPORTANCE In marine oxygen-deficient zones (ODZs), microbes transform bioavailable nitrogen to gaseous nitrogen, with nitric oxide as a key intermediate. The Eastern Tropical North Pacific contains the world's largest ODZ, but the identity of the microbes transforming nitric oxide remains unknown. Here, we show that highly transcribed nitric oxide dismutase (*nod*) genes belong to Alphaproteobacteria of the novel order UBA11136, which lacks cultivated isolates. These Alphaproteobacteria show evidence for aerobic respiration, using oxygen potentially sourced from nitric oxide dismutase, and possess a novel porin-cytochrome protein complex with unknown function. Gammaproteobacteria and Planctomycetia transcribe *nod* at lower levels. Our results pinpoint the microbes mediating a key step in marine nitrogen loss and reveal an unexpected predicted metabolism for marine Alphaproteobacteria.

KEYWORDS

nitric oxide, Alphaproteobacteria, marine, oxygen-deficient zone, nitrogen, oxygen, denitrification

Marine oxygen-deficient zones (ODZs) contribute up to half of the ocean's nitrogen loss (1) and are a major source of marine emissions of the potent greenhouse gas nitrous oxide (N₂O) (2). The primary source of the N₂O at the oxic–anoxic interface and in anoxic waters in ODZs is denitrification (3, 4). The microbial enzyme responsible for N₂O production during denitrification is nitric oxide reductase (Nor), which uses electrons from cytochrome *c* (cNor) or quinol (qNor), to reduce nitric oxide (NO) to N₂O (5–7). In the qNor family, there are *bona fide* qNor enzymes and NO dismutase (Nod). Nod proteins lack the quinol-binding site, seemingly preventing the enzyme from taking up external electrons; instead, Nod is theorized to disproportionate NO into dinitrogen and O₂ in methane-oxidizing Methyloirabiolota bacteria (8, 9) and in the alkane-oxidizing Gammaproteobacterium HdN1 (10).

The Eastern Tropical North and South Pacific (ETNP and ETSP) ODZs are the world's largest and second largest ODZs, and the subjects of extensive microbial ecology studies. Abundant NO reductase-like genes and transcripts in the ETNP and ETSP ODZ cluster in the same enzyme subfamily as Nod (11–14). Due to the similarity of ODZ Nod proteins to those of Methyloirabiolota (NC10), it was initially presumed that ODZ bacteria also used Nod proteins to disproportionate NO into N₂ and O₂ for use in intra-aerobic methane oxidation (11, 13, 15). However, Fuchsman et al. (12) found that the peak of *nod* gene abundance in the ETNP ODZ correlates with a peak of modeled N₂O production (4) and does not correlate with abundance of methane monooxygenase genes, suggesting that Nod proteins in the ETNP ODZ are potentially an important source of N₂O and are unlikely to be involved in methane oxidation. The plausibility that Nod proteins can reduce NO to N₂O is supported by a study of a novel eukaryotic denitrification pathway in foraminifera (*Globobulimina* spp.) that produce N₂O while expressing Nod (16). Yet, the phylogenetic identity and metabolic context of marine Nod proteins, which are a key biological source of either N₂O or O₂+N₂ in marine ODZs, remain unresolved.

In this study, we sought to determine the identity, predicted metabolism, and environmental niche of the ODZ organism responsible for the highly transcribed *nod* genes first discovered by Padilla et al. (11). We found that the most abundantly transcribed *nod* genes in the ETNP ODZ belong to Alphaproteobacteria in the novel order UBA11136. Significant transcription of *nod* genes was limited to waters with <1 μM O₂. These *nod*-transcribing Alphaproteobacteria also transcribe genes involved in aerobic respiration, which was unexpected given that they inhabit anoxic waters, as well as genes involved in oxidation of formaldehyde, likely indicating methylotrophy. Genes encoding multiheme cytochrome proteins potentially implicated in nitrogen or iron cycling were also transcribed.

RESULTS

Transcribed *nod* sequences in the ETNP ODZ belong to Alphaproteobacteria, Gammaproteobacteria, and Planctomycetia

Our reanalysis of highly transcribed *nod* genes in the ETNP ODZ (11) shows that these genes belong to Alphaproteobacteria rather than a member of Methyloirabiolota as previously assumed. Querying the Nod amino acid sequences from Padilla et al. (11) against ETNP ODZ metagenomes in the JGI IMG/MER database returned multiple 100% identity matches, including a *nod* gene (Ga0066848_100037855) on a scaffold with hypothetical genes with 100% identity to Alphaproteobacteria metagenome-assembled genomes (MAGs) from the ETNP ODZ (17) (Table S1). We binned previously sequenced ETNP ODZ metagenomes Ga0066848 (ETNP201310SV72) and Ga0066829 (ETNP201306SV43) (18) into MAGs. Contigs with the most highly transcribed *nod* genes were present in two Alphaproteobacteria MAGs (GTDB taxonomy: UBA11136 sp002686135; species representative: Rhodospirillaceae bacterium isolate ARS27) with 97% average nucleotide identity. Querying the Nod amino acid sequences from Padilla et al. (11) against NCBI's nonredundant protein database returned matches to other MAGs assigned to Alphaproteobacteria order UBA11136 from low-oxygen

Editor Jennifer F. Biddle, University of Delaware, Newark, Delaware, USA

Address correspondence to Jennifer B. Glass, jennifer.glass@eas.gatech.edu.

The authors declare no conflict of interest.

See the funding table on p. 11.

Received 20 November 2023

Accepted 9 February 2024

Published 6 March 2024

Copyright © 2024 American Society for Microbiology. All Rights Reserved.

marine settings (ETNP, Saanich Inlet, and the Black Sea; 78%–80% identity), the marine magnetotactic alphaproteobacterium *Magnetovibrio blakemorei* MV-1 (75% identity), Gammaproteobacterium HdN1 (66% identity), and *Methyloirabilis* spp. (66% identity; Table S2).

To glean additional insights into evolutionary relationships, we updated a previous Nod phylogeny (19) with additional amino acid sequences from marine MAGs (20–22) and ETNP ODZ metagenomes (18), subdivided into cells that are free-living (FL; 0.2–1.6 μm) and from the particle fraction (PF; >1.6 μm ; Fig. 1A; Table S3). The Nod topology was generally consistent with a previous phylogeny from Fuchsman et al. (12), with




FIG 1 Marine Nod clades, gene neighborhoods, and depth profiles of transcription. (A) A maximum likelihood phylogeny of nitric oxide dismutase (Nod) amino acid sequences in marine (blue) and select terrestrial (brown) taxa, primarily from marine MAGs (20–22) and ETNP ODZ metagenomes (18). Branch support was evaluated using 1,000 rapid bootstrap replicates, with bootstrap values shown for deep branches. The tree is drawn to scale, with branch lengths in number of (Continued on next page)

FIG 1 (Continued) substitutions per site. Bold sequences represent those present in multiple ETNP ODZ metagenomes (see Table S3 for duplicate accession numbers). “PF” indicates genes from the particle fraction (>1.6 μ m fraction) of filters. “FL” indicates genes from the free-living fraction (0.2–1.6 μ m) collected on Sterivex filters.

The most highly transcribed ETNP ODZ sequence is indicated with an asterisk. The qNor sequence *Geobacillus stearothermophilus* was used as the outgroup. (B) Gene neighborhoods surrounding *nod* genes in select taxa. GenBank contigs: *Cecembia calidifontis* SGXG01000001, *Scalindua japonica* BAOS01000045, Gammaproteobacteria NP964 PBRC01000062, Gammaproteobacterium HdN1 FP929140, Deltaproteobacteria NZCL01000067, *Candidatus Methyloirabilis oxyfera* FP565575, and Rhodospirillaceae NP1106 PCBZ01000014. Unlabeled gray genes are hypothetical. (C, D) Oxygen concentrations (gray lines), nitrite concentrations (black circles), and *nod* transcripts (squares, as reads per kilobase per million mapped reads [RPKM]) with depth in ETNP ODZ P1 (onshore) and P2 (offshore) sites (25).

additional taxonomic data from MAGs in the *Tara Oceans* data set further constraining Nod placement (22). As expected based on the binning and BLAST results, the Nod sequence from Padilla et al. (11) (Ga0066848_100037855) clustered phylogenetically with marine Alphaproteobacteria (OTU III in Fuchsman et al. [12], hereafter “Alpha-type Nod”); this clade contained three unique sequences, all of which were present in multiple metagenomes and all from the free-living fraction, and one of which was identical to that of Rhodospirillaceae NP1106 (GenBank: MBV28360). Four unique ODZ Nod sequences clustered with marine Gammaproteobacteria (OTU II in Fuchsman et al. [12], hereafter “Gamma-type Nod”); these sequences were monophyletic with a cluster of Gammaproteobacteria Nod cluster sequences from sewage sludge, including Gammaproteobacterium HdN1 (23) and other wastewater Gammaproteobacteria. Multiple ETNP ODZ metagenomes contained Gamma-type Nod sequences identical to those of Gammaproteobacteria NP964 (GenBank: MBP20251). Gamma-type Nod had ~70% identity to Alpha-type Nod. Several ODZ Nod sequences, all from the particle fraction, clustered with marine Deltaproteobacteria in a clade of monophyletic *nod* genes from groundwater Methyloirabilota, Deltaproteobacteria, and Acidobacteria MAGs (~65% identity to Alpha-type Nod). Six unique Nod ODZ protein sequences (two of which were present in multiple metagenomes) clustered with Planctomycetia (OTU I in Fuchsman et al. [12], hereafter “Planctomycetia-type Nod”), were primarily found in free-living cells, and had ~40% identity to Alpha-type Nod. Intriguingly, two ODZ sequences clustered in the eukaryotic *Globobulimina* clade (~50% identity to Alpha-type Nod). Viral Nod sequences from Saanich Inlet (~55% identity to Alpha-type Nod) clustered with the viral Nod sequence previously reported by Gazitúa et al. (24) from the ETSP ODZ (St16 OMZ 317E-viral).

We investigated gene neighborhoods surrounding ODZ *nod* genes in the three main phylogenetic clusters of ODZ sequences: Planctomycetia-type Nod, Gamma-type Nod, and Alpha-type Nod. Although “unknown Nor-related” marine Bacteroidota sequences were located on an operon with other *nor* genes, there was no consistent gene neighborhood for *nod* sequences (Fig. 1B). Planctomycetia-type *nod* genes were not located in the vicinity of any genes with recognizable related function. Gamma-type *nod* gene neighborhoods contained ferredoxins and cytochrome *b561* genes for electron transport. Upstream of the Alpha-type *nod* in Rhodospirillaceae NP1106 is a cluster of formylmethanofuran dehydrogenase genes (*fmd/fwd*) used in C1 metabolism via tetrahydromethanopterin/methanofuran-linked reactions. Immediately upstream or downstream of *nod* genes, helix–turn–helix transcriptional regulators were common (Fig. 1B). Neighboring Gamma-type and *Methyloirabilis* *nod* genes, LuxR-type regulators were common; these regulators have diverse functions and their potential connection to Nod remains unclear. Neighboring Alpha-type and Bacteroidota (e.g., *Cecembia calidifontis*) *nod* genes, Rrf2-type regulators were present. The protein NsrR in the Rrf2 family regulates global cellular response to NO toxification by directly sensing NO with an iron-sulfur cluster (26, 27). The presence of this NsrR-like regulator suggests that Nod in marine Alphaproteobacteria and Bacteroidota may be involved in nitrosative stress response and NO detoxification.

Alphaproteobacterial *nod* is highly transcribed in anoxic waters

We assessed transcription of Alpha-, Gamma-, and Planctomycetia-type *nod* genes from the oxycline to upper ODZ (secondary nitrite maximum) using ETNP ODZ metatranscriptomes from an onshore station with a shallower oxycline (P1; Fig. 1C) and an offshore station with a deeper oxycline (P2; Fig. 1D) (25). In both oxyclines, transcription was low (4–10 reads per kilobase per million mapped reads [RPKM], $n = 8$) for all three *nod* types (Fig. 1C and D). Below the oxyclines, *nod* transcripts began to rise and were highest at the secondary nitrite

maxima, with Alpha-type (184–274 RPKM, $n = 4$) > Gamma-type (55–95 RPKM, $n = 4$) > Planctomycetia-type (13–19 RPKM, $n = 4$; Table S4).

MAGs with highly transcribed *nod* gene represent a new order of Alphaproteobacteria

In order to assess the phylogeny of the *nod*-containing Alphaproteobacteria MAGs, we constructed an alphaproteobacterial phylogeny using the conserved protein NADH ubiquinone oxidoreductase subunit L (NuoL) as in Cevallos and Degli Esposti (28), with additional representation of order UBA11136 including our MAG ETNP2013_S10_300m_22 (Fig. 2). MAG ETNP2013_S06_300m_15 was not included in the phylogeny because its *nucL* gene was truncated. The phylogeny confirmed that *nod*-containing Alphaproteobacteria belong to the order UBA11136 and showed that UBA11136 is situated near other Alphaproteobacteria orders found in ODZs.




FIG 2 Alphaproteobacteria phylogeny with order UBA11136 expanded and *nod*-containing MAGs bolded. The phylogeny was constructed using the alphaproteobacterial phylogenetic marker NADH ubiquinone oxidoreductase subunit L as in Cevallos and Degli Esposti (28). Taxonomic names are from Cevallos and Degli Esposti (28) and GTDB Release 08-RS214. The scale bar represents amino acid substitutions per site. The full phylogeny is shown in Fig. S1.

Alphaproteobacteria transcribe genes for formate metabolism, aerobic respiration, and a multiheme cytochrome complex

To glean insight into potential roles for Nod in a cellular context, we sought to reconstruct the electron transport chain of the Alphaproteobacteria with the most highly transcribed *nod* genes (Alphaproteobacterium MAG ETNP2013_S10_300m_22 and Alphaproteobacterium MAG ETNP2013_S06_300m_15, 73% and 69% estimated completeness, respectively) at the secondary nitrite maximum. Of total metagenomic reads, 0.38% map to ETNP2013_S10_300m_22 and 0.39% map to ETNP2013_S06_300m_15. In both MAGs, *nod* was in the top three most transcribed genes in the ETNP ODZ (~44,000 FPKM; Table S5), after a bacterial nucleoid DNA-binding protein and a potassium-gated

channel protein. In addition to *nod*, we found that genes for formaldehyde oxidation via tetrahydromethanopterin/methanofuran-linked reactions, including formylmethanofuran dehydrogenase (*fwd/fmd*) and formylmethanofuran–tetrahydromethanopterin N-formyltransferase (*ftr*), were transcribed in both MAGs (Table S5). Both MAGs also transcribed NAD-dependent formate dehydrogenase (Table S5). Thus, the alphaproteobacterium appears to be capable of conversion of formaldehyde to formate and use of formate as a source of electrons for NADH:ubiquinone oxidoreductase (Complex I; Fig. 3). The source of formaldehyde is likely methanol oxidation, as pyrroloquinoline quinone (PQQ)-dependent ethanol/methanol dehydrogenases were found in Alphaproteobacteria MAGs from low-oxygen marine settings (Table S6). Methane monooxygenase genes were not found in the partial Alphaproteobacteria MAGs, precluding our ability to rule out the possibility of these genes in the missing portions of the genomes. The Alphaproteobacteria PQQ-dependent dehydrogenase genes contained the motif DYDG (Table S6), which is characteristic of the lanthanide-containing form of the enzymes rather than the calcium form (29).

A full aerobic electron transport chain (Complex I, II, III, and IV) and F₀F₁-type ATP synthase were transcribed in both bins (Fig. 3; Table S5). Complex IV (cytochrome c oxidase) was type A1 according to the Sousa et al. (30) classification, and the *cox* operon in the GTDB species representative Rhodospirillaceae ARS27 was subtype b (COX2-COX1-CtaB-CtaG_Cox11-COX3-DUF983-SURF1-CtaA1-M32-Tsy-M16B) according to the Geiger et al. (31) classification. Sulfur oxidation genes, including flavocytochrome c sulfide dehydrogenase (FccAB), sulfane hydrogenase (SoxCD), and carrier protein SoxYZ, were also transcribed, as were numerous transposes (Fig. 3; Table S5).

Genes for a multiheme cytochrome complex were transcribed in both bins. To our knowledge, this putative operon has not been previously described. Hereafter, we designate it the *ptdABCDEFG* operon for its sequence of penta/tetra/deca-heme proteins, interspersed with other conserved proteins. *ptdAB* genes are highly transcribed in our Alphaproteobacteria MAGs, but it is unclear if the rest of the operon is also highly transcribed, because it was truncated in our MAGs' scaffolds. The *ptd* gene cluster consists of a penta-heme protein with a C-terminal beta-sandwich (PtdA), a porin (PtdB), a FAD/NAD(P)-binding oxidoreductase (PtdC), a periplasmic tetra-heme protein (PtdD), a cyclic nucleotide-binding domain protein with two 4Fe–4S clusters (PtdE), a cytoplasmic transmembrane ferric reductase-like protein (PtdF), and a periplasmic deca-heme protein (PtdG; Fig. 3; Table S7 and S8). The function of this complex is unknown, but the presence of genes encoding a porin and multiple multiheme proteins resembles porin-cytochrome protein complexes involved in extracellular reduction electron transfer during Fe(III) and Mn(IV) reduction (32, 33). PtdA has a homolog to a penta-heme cytochrome c552 protein of unknown function in a thermophilic purple sulfur gammaproteobacterium (34) and is in the same COG family (COG3303) as formate-dependent nitrite reductase, NrfA. *ptdABCDEFG* genes were prevalent in Alphaproteobacteria, Gammaproteobacteria, Nitrospirales, and Planctomycetes MAGs from marine or high salinity environments (Fig. 4; Table S7).




FIG 3 Schematic of the electron transport chain in *nod*-containing Alphaproteobacteria. Increasing transcriptional activity is indicated from lighter to darker blue (Table S5). Red circles with black lines indicate hemes. Hypothetical Ptd proteins are labeled A, B, C, D, E, F, and G (Table S8). Proposed electron transfer from formate to Complex I is shown. Highly transcribed Nod protein and predicted O₂ generation is shown as feeding into A1-type CCO Complex IV. Additional electrons for CytC and the electron transport chain are proposed to come from sulfur oxidation carried out by the flavocytochrome *c* sulfide dehydrogenase (FccAB, FCC), and sulfane-sulfur dehydrogenase (SoxCD) with the multi-enzyme carrier complex (SoxYZ).

DISCUSSION

This study predicts the previously ambiguous identity of the microorganisms that make the dominant nitric oxide-transforming protein (Nod) in the world's largest ODZ, the Eastern Tropical North Pacific. Extensive horizontal gene transfer of *nod* genes between microbial genomes is evident from the lack of conservation of gene neighborhood and patchy phylogeny (12), which may be mediated by viral infection (24). We found that the most transcriptionally active *nod* genes in the ETNP upper ODZ belong to



the novel Alphaproteobacteria order UBA11136. Alpha-type *nod* transcript abundances (~200 RPKM) are similar to those of dissimilatory nitrate reductase (*narG*) in the ODZ (35). The *nod*-transcribing Alphaproteobacteria are also transcribing genes for formaldehyde oxidation, likely as a source of electrons to the respiratory chain via NAD reduction by formate dehydrogenase. Sulfide may be used as a supplemental electron donor and/or may be concomitantly oxidized for detoxification (36, 37).

On the other end of the electron transport chain, high transcription of a heme/copper terminal oxidase suggests that O₂ is being used as the terminal electron acceptor in *nod*-transcribing Alphaproteobacteria MAGs. The transcribed heme/copper oxidase is A1-type (low O₂ affinity), also present in mitochondria, and adapted for high O₂ concentrations. Low O₂ affinity A1-type heme/copper oxidases are transcribed in other anoxic environments (41). Because ODZs have extremely low concentrations of O₂ below the oxycline, O₂ for aerobic respiration may be generated *in situ* and rapidly consumed. Given that the function of Nod is proposed to be dismutation of two NO molecules into N₂ and O₂ (8), it is possible that the O₂ source for aerobic respiration in the UBA11136 MAGs is NO dismutation, although other sources of O₂ (e.g., *in situ* photosynthesis, mixing) in anoxic waters are also conceivable (42). The physiological uses of Gamma-type and Planctomycetia-type Nod may be different from Alpha-type Nod, although this remains to be investigated.

Myxococcota MAG contained *nirS*, and one *Scalindua* MAG contained *nirS*). It is also possible that another uncharacterized enzyme produces NO.

This study suggests that marine Alphaproteobacteria from order UBA11136 are actively reducing NO under anoxia, as implied by their abundant transcription of *nod* genes. Although there is strong evidence that the substrate for Nod in ODZs is NO based on its abundance, the products of this enzyme (N₂O vs N₂+O₂) remain uncertain. Nod is theorized to disproportionate NO into N₂ and O₂ in methane-oxidizing *Methylomirabilota* bacteria (8, 9), but no biochemical characterizations of Nod have been published to date, and foraminifera expressing Nod produce N₂O (16). The apparent lack of other denitrification genes in *nod*-transcribing Alphaproteobacteria is consistent with the observation that denitrification in ODZs is largely divided into distinct microbial taxa (12, 13, 45). For example, although nitrate reductase (*narG*) genes are widely distributed amongst ODZ microbes (45), SAR11 bacteria appear to dominate in *narG* transcriptional activity (35). Our finding that the transcription of *nod* is catalyzed primarily by marine Alphaproteobacteria implies that this taxon contributes significantly to marine nitrogen loss.

MATERIALS AND METHODS

Nod phylogeny and gene neighborhood

Amino acid sequences of highly transcribed *nod* genes “ETNP 2014 Stn10 150m” and “ETNP 2013 Stn6 300m” were acquired from the authors of Padilla et al. (11) (see Table S2 for sequences). These sequences were used for BLASTP searches of ODZ metagenomes in the JGI IMG/MER database and the NCBI nonredundant protein (nr) database. Sequences ($n = 53$, 731 gap-free sites) were aligned using the MAFFT online server with the L-INS-i method (46). A phylogeny was generated with 1,000 bootstraps using model LG+I+G4 with W-IQ-TREE (47). The phylogeny was visualized using FigTree v.1.4.4, and the fasta file (Nod_alignment) is available as a supplemental data set. Gene neighborhoods were generated using the EFI Gene Neighborhood Tool (48) with single sequence BLAST of the UniProt database using the amino acid sequence Ga0066848_100037855 (JGI IMG/ MER) as the Nod query with an e-value cutoff of 10^{-5} and with 10 genes upstream and downstream the gene of interest.

Transcription of *nod* genes in ETNP ODZ depth profiles

Magic Basic Local Alignment Search Tool (49) was used to search ETNP ODZ metatranscriptomes (PRJNA727903; Mattes et al. [25]) using representative nucleotide sequences for Planctomycetia-like (Ga0066826_100064333 [JGI IMG/MER]), Gamma-like (PBRC01000062.1:19833–22205 [NCBI]), and Alpha-like (Ga0066848_100037855 [JGI IMG/MER]) *nod* genes. Default parameters were used except for the score threshold (18). Read hits were normalized to reads per kilobase million (RPKM).

Metagenomic binning

Binning of metagenome-assembled genomes (MAGs) was performed using the KBase platform (50). ETNP ODZ metagenomes were collected in 2013 and sequenced by Joint Genome Institute (JGI) using an Illumina HiSeq 2500 as described in Ruiz-Perez et al. (18). Assemblies for the ETNP ODZ metagenomes (18) containing

Alpha-type *nod* genes (ETNP201310SV72 [GOLD Analysis Project ID Ga0066848; stn10 300m] and ETNP201306SV43 [GOLD Analysis Project ID Ga0066829; stn6 300m]) were imported from JGI IMG/MER into KBase. Metagenomic assemblies were binned into MAGs using MaxBin2 v2.2.4 (51). The two MAGs containing *nod* genes (MAG ETNP2013_S10_300m_22 from ETNP201310SV72, and ETNP2013_S06_300m_15 from ETNP201306SV43) were selected for further analysis. Average nucleotide identity was calculated using FastANI (52). MAG taxonomy and genome quality were evaluated by GTDB-Tk v2.3.2 (53). MAGs were annotated with RASTtk v1.073 (54). Metagenomic reads were mapped to MAGs using Bowtie2 (55).

Alphaproteobacterial NuoL phylogeny

Alphaproteobacterial NADH ubiquinone oxidoreductase subunit L (NuoL) and mitochondrial ND5 marker proteins ($n = 320$) were aligned as in Cevallos and Degli Esposti (28), with additional representation of order UBA11136. A maximum likelihood phylogeny with 1000 bootstraps was constructed in IQ-tree (56) using the LG+F model with ultrafast bootstrap (57). Taxonomic names and clades are from Cevallos and Degli Esposti (28) and GTDB Release 08-RS214. The fasta file (NuoL_alignment) is available as a supplemental data set. Alphaproteobacteria MAGs containing *nod* genes (Table S2) were classified as belonging to order UBA11136 using GTDB-Tk v2.3.2 (53).

Mapping transcripts to metagenomic bins

Metatranscriptomic mapping to MAGs was performed using the KBase platform (50). RNA-seq data (25) were imported from the depth with the highest *nod* transcription, the secondary nitrite maximum (126 m, NCBI run SRR14460584), and aligned to MAGs using the Bowtie2 (55) app in KBase. The Cufflinks v2.2.1 (58) app in KBase was then used to assemble the aligned RNA-seq data into a set of transcripts and to calculate the relative abundances of the transcripts expressed in fragments per kilobase per million fragments mapped (FPKM).

Cellular localization and heme numbers

Cellular locations of Ptd proteins were predicted using PSORTb v3.0.3 analysis (59). Numbers of heme-binding motifs per protein were identified by counting CXXCH sequences. Ptd gene neighborhoods were generated using the EFI Gene Neighborhood Tool (48) with single sequence BLAST of the UniProt database using the amino acid sequence Ga0066848_100031354 (JGI IMG/MER) as the PtdA query with an e-value cutoff of 10^{-5} and with 10 genes upstream and downstream the gene of interest.

ACKNOWLEDGMENTS

We thank Laura Bristow for helpful discussions. We thank Mauro Degli Esposti for assistance with the NADH ubiquinone oxidoreductase subunit L phylogeny. We thank Cory Padilla, Anthony Bertagnolli, and Neha Sarode for sharing previous data.

We acknowledge funding from an NSF Graduate Research Fellowship to C.E.E., the Simons Foundation, and NSF Awards 2022991 and 2054927 to F.J.S.

AUTHOR AFFILIATIONS

¹School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA

²Department of Microbiology & Cell Biology, Montana State University, Bozeman, Montana, USA

AUTHOR ORCIDS

Claire E. Elbon  <http://orcid.org/0000-0002-2796-2475>

Frank J.  Stewart <http://orcid.org/0000-0002-8139-5176>

Jennifer B.  Glass <http://orcid.org/0000-0003-0775-2486>

FUNDING

Funder	Grant(s)	Author(s)
National Science Foundation (NSF)	2022991	Frank J. Stewart
National Science Foundation (NSF)	2054927	Frank J. Stewart
National Science Foundation (NSF)	Graduate Research Fellowship	Claire E. Elbon
Simons Foundation (SF)		Frank J. Stewart

AUTHOR CONTRIBUTIONS

Claire E. Elbon, Formal analysis, Investigation, Methodology, Visualization | Frank J. Stewart, Funding acquisition, Writing – review and editing | Jennifer B. Glass, Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Validation, Visualization, Writing – original draft, Writing – review and editing

DATA AVAILABILITY

The KBase bioinformatic pipeline and MAGs are at <https://narrative.kbase.us/narrative/106999>. Original metagenomic reads are available at BioProject [PRJNA375524](#) (ETNP201306SV43, [SAMN06344130](#)) and BioProject [PRJNA375542](#) (ETNP201310SV72, [SAMN06344148](#)). MAG Alphaproteobacteria bacterium ETNP2013_S06_300m_15 was deposited into BioProject [PRJNA375524](#) (BioSample [SAMN38229257](#), WGS Accession [JAZDBU000000000](#)) and Alphaproteobacteria bacterium ETNP2013_S10_300m_22 was deposited into BioProject [PRJNA375542](#) (BioSample [SAMN38228782](#), WGS Accession [JAZDCE000000000](#)). All ETNP ODZ data sets used in this manuscript are listed in Table S9.

ADDITIONAL FILES

The following material is available [online](#).

Supplemental Material

Supplemental figure (AEM02099-23-s0001.docx). Figure S1.

Supplemental FASTA file, NuoL (AEM02099-23-s0002.pdf). NuoL amino acid alignment FASTA file.

Supplemental FASTA file, Nod (AEM02099-23-s0003.pdf). Nod amino acid alignment FASTA file.

Supplemental tables (AEM02099-23-s0004.xlsx). Table S1–S9.

REFERENCES

- DeVries T, Deutsch C, Rafter PA, Primeau F. 2013. Marine denitrification rates determined from a global 3-D inverse model. *Biogeosciences* 10:2481–2496. <https://doi.org/10.5194/bg-10-2481-2013>
- Yang S, Chang BX, Warner MJ, Weber TS, Bourbonnais AM, Santoro AE, Kock A, Sonnerup RE, Bullister JL, Wilson ST, Bianchi D. 2020. Global reconstruction reduces the uncertainty of oceanic nitrous oxide emissions and reveals a vigorous seasonal cycle. *Proc Natl Acad Sci U S A* 117:11954–11960. <https://doi.org/10.1073/pnas.1921914117>
- Frey C, Bange HW, Achterberg EP, Jayakumar A, Löscher CR, Arévalo-Martínez DL, León-Palmero E, Sun M, Sun X, Xie RC, Oleynik S, Ward BB. 2020. Regulation of nitrous oxide production in low-oxygen waters off the coast of Peru. *Biogeosciences* 17:2263–2287. <https://doi.org/10.5194/bg-17-2263-2020>
- Babbin AR, Bianchi D, Jayakumar A, Ward BB. 2015. Rapid nitrous oxide cycling in the suboxic ocean. *Science* 348:1127–1129. <https://doi.org/10.1126/science.aaa8380>
- Wasser IM, de Vries S, Moëne-Loccoz P, Schröder I, Karlin KD. 2002. Nitric oxide in biological denitrification: Fe/Cu metalloenzyme and metal complex NO_x redox chemistry. *Chem Rev* 102:1201–1234. <https://doi.org/10.1021/cr0006627>
- Kraft B, Strous M, Tegetmeyer HE. 2011. Microbial nitrate respiration—genes, enzymes and environmental distribution. *J Biotechnol* 155:104–117. <https://doi.org/10.1016/j.jbiotec.2010.12.025>
- Zumft WG. 2005. Nitric oxide reductases of prokaryotes with emphasis on the respiratory, heme-copper oxidase type. *J Inorg Biochem* 99:194–215. <https://doi.org/10.1016/j.jinorgbio.2004.09.024>
- Ettwig KF, Butler MK, Le Paslier D, Pelletier E, Mangenot S, Kuypers MMM, Schreiber F, Dutilh BE, Zedelius J, de Beer D, Gloerich J, Wessels HJCT, van Alen T, Luesken F, Wu ML, van de Pas-Schoonen KT, Op den Camp HJM, Janssen-Megens EM, Francois K-J, Stunnenberg H, Weissenbach J, Jetten MSM, Strous M. 2010. Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* 464:543–548. <https://doi.org/10.1038/nature08883>
- Ettwig KF, Speth DR, Reimann J, Wu ML, Jetten MSM, Keltjens JT. 2012. Bacterial oxygen production in the dark. *Front Microbiol* 3:273. <https://doi.org/10.3389/fmicb.2012.00273>
- Zedelius J, Rabus R, Grundmann O, Werner I, Brodtkorb D, Schreiber F,

- Ehrenreich P, Behrends A, Wilkes H, Kube M, Reinhardt R, Widdel F. 2011. Alkane degradation under anoxic conditions by a nitrate-reducing bacterium with possible involvement of the electron acceptor in substrate activation. *Environ Microbiol Rep* 3:125–135. <https://doi.org/10.1111/j.1758-2229.2010.00198.x>
11. Padilla CC, Bristow LA, Sarode N, Garcia-Robledo E, Gómez Ramírez E, Benson CR, Bourbonnais A, Altabet MA, Girguis PR, Thamdrup B, Stewart FJ. 2016. NC10 bacteria in marine oxygen minimum zones. *ISME J* 10:2067–2071. <https://doi.org/10.1038/ismej.2015.262>
 12. Fuchsman CA, Devol AH, Saunders JK, McKay C, Rocap G. 2017. Niche partitioning of the N cycling microbial community of an offshore oxygen deficient zone. *Front Microbiol* 8:2384. <https://doi.org/10.3389/fmicb.2017.02384>
 13. Dalsgaard T, Stewart FJ, Thamdrup B, De Brabandere L, Revsbech NP, Ulloa O, Canfield DE, DeLong EF. 2014. Oxygen at nanomolar levels reversibly suppresses process rates and gene expression in anammox and denitrification in the oxygen minimum zone off northern Chile. *mBio* 5:e01966. <https://doi.org/10.1128/mBio.01966-14>
 14. Ganesh S, Parris DJ, DeLong EF, Stewart FJ. 2014. Metagenomic analysis of size-fractionated picoplankton in a marine oxygen minimum zone. *ISME J* 8:187–211. <https://doi.org/10.1038/ismej.2013.144>
 15. Thamdrup B, Steinsdóttir HGR, Bertagnolli AD, Padilla CC, Patin NV, Garcia-Robledo E, Bristow LA, Stewart FJ. 2019. Anaerobic methane oxidation is an important sink for methane in the ocean's largest oxygen minimum zone. *Limn & Ocean* 64:2569–2585. <https://doi.org/10.1002/lno.11235>
 16. Woehle C, Roy A-S, Glock N, Wein T, Weissenbach J, Rosenstiel P, Hiebenthal C, Michels J, Schönfeld J, Dagan T. 2018. A novel eukaryotic denitrification pathway in foraminifera. *Curr Biol* 28:2536–2543. <https://doi.org/10.1016/j.cub.2018.06.027>
 17. Uzun M, Alekseeva L, Krutkina M, Koziaeva V, Grouzdev D. 2020. Unravelling the diversity of magnetotactic bacteria through analysis of open genomic databases. *Sci Data* 7:252. <https://doi.org/10.1038/s41597-020-00593-0>
 18. Ruiz-Perez CA, Bertagnolli AD, Tsementzi D, Woyke T, Stewart FJ, Konstantinidis KT. 2021. Description of *Candidatus* mesopelagibacter carboxydoydys and *Candidatus* anoxipelagibacter denitrificans: nitrate-reducing SAR11 genera that dominate mesopelagic and anoxic marine zones. *Syst Appl Microbiol* 44:126185. <https://doi.org/10.1016/j.syapm.2021.126185>
 19. Hu Q-Q, Zhou Z-C, Liu Y-F, Zhou L, Mbadinga SM, Liu J-F, Yang S-Z, Gu JD, Mu B-Z. 2019. High microbial diversity of the nitric oxide dismutation reaction revealed by PCR amplification and analysis of the *nod* gene. *Inter Biod & Biod* 143:104708. <https://doi.org/10.1016/j.ibiod.2019.05.025>
 20. Lin H, Ascher DB, Myung Y, Lamborg CH, Hallam SJ, Gionfriddo CM, Holt KE, Moreau JW. 2021. Mercury methylation by metabolically versatile and cosmopolitan marine bacteria. *ISME J* 15:1810–1825. <https://doi.org/10.1038/s41396-020-00889-4>
 21. Cabello-Yeves PJ, Callieri C, Picazo A, Mehrshad M, Haro-Moreno JM, Roda-Garcia JJ, Dzhenbekova N, Slabakova V, Slabakova N, Moncheva S, Rodriguez-Valera F. 2021. The microbiome of the black sea water column analyzed by shotgun and genome centric metagenomics. *Environ Microbiome* 16:5. <https://doi.org/10.1186/s40793-021-00374-1>
 22. Tully BJ, Graham ED, Heidelberg JF. 2018. The reconstruction of 2,631 draft metagenome-assembled genomes from the global oceans. *Sci Data* 5:170203. <https://doi.org/10.1038/sdata.2017.203>
 23. Ehrenreich P, Behrends A, Harder J, Widdel F. 2000. Anaerobic oxidation of alkanes by newly isolated denitrifying bacteria. *Arch Microbiol* 173:58–64. <https://doi.org/10.1007/s002030050008>
 24. Gazitúa MC, Vik DR, Roux S, Gregory AC, Bolduc B, Widner B, Mulholland MR, Hallam SJ, Ulloa O, Sullivan MB. 2021. Potential virus-mediated nitrogen cycling in oxygen-depleted oceanic waters. *ISME J* 15:981–998. <https://doi.org/10.1038/s41396-020-00825-6>
 25. Mattes TE, Burke S, Rocap G, Morris RM. 2022. Two metatranscriptomic profiles through low-dissolved-oxygen waters (DO, 0 to 33 μ M) in the eastern tropical North Pacific ocean. *Microbiol Resour Announc* 11:e0120121. <https://doi.org/10.1128/mra.01201-21>
 26. Tucker NP, Le Brun NE, Dixon R, Hutchings MI. 2010. There's NO stopping NsrR, a global regulator of the bacterial NO stress response. *Trends Microbiol* 18:149–156. <https://doi.org/10.1016/j.tim.2009.12.009>
 27. Bodenmiller DM, Spiro S. 2006. The *yjeB* (*nsrR*) gene of *Escherichia coli* encodes a nitric oxide-sensitive transcriptional regulator. *J Bacteriol* 188:874–881. <https://doi.org/10.1128/JB.188.3.874-881.2006>
 28. Cevallos MA, Degli Esposti M. 2022. New Alphaproteobacteria thrive in the depths of the ocean with oxygen gradient. *Microorganisms* 10:455. <https://doi.org/10.3390/microorganisms10020455>
 29. Keltjens JT, Pol A, Reimann J, Op den Camp HJM. 2014. PQQ-dependent methanol dehydrogenases: rare-earth elements make a difference. *Appl Microbiol Biotechnol* 98:6163–6183. <https://doi.org/10.1007/s00253-014-5766-8>
 30. Sousa FL, Alves RJ, Ribeiro MA, Pereira-Leal JB, Teixeira M, Pereira MM. 2012. The superfamily of heme-copper oxygen reductases: types and evolutionary considerations. *Bio et Biophysica Acta (BBA) - Bio* 1817:629–637. <https://doi.org/10.1016/j.bbabi.2011.09.020>
 31. Geiger O, Sanchez-Flores A, Padilla-Gomez J, Degli Esposti M. 2023. Multiple approaches of cellular metabolism define the bacterial ancestry of mitochondria. *Sci Adv* 9:eadh0066. <https://doi.org/10.1126/sciadv.adh0066>
 32. Richardson DJ, Butt JN, Fredrickson JK, Zachara JM, Shi L, Edwards MJ, White G, Baiden N, Gates AJ, Marritt SJ, Clarke TA. 2012. The 'porincytochrome' model for microbe-to-mineral electron transfer. *Mol Microbiol* 85:201–212. <https://doi.org/10.1111/j.1365-2958.2012.08088.x>
 33. Shi L, Fredrickson JK, Zachara JM. 2014. Genomic analyses of bacterial porin-cytochrome gene clusters. *Front Microbiol* 5:657. <https://doi.org/10.3389/fmicb.2014.00657>
 34. Chen JH, Yu LJ, Boussac A, Wang-Otomo ZY, Kuang T, Shen JR. 2019. Properties and structure of a low-potential, penta-heme cytochrome C(552) from a thermophilic purple sulfur photosynthetic bacterium *Thermochromatium tepidum*. *Photosynth Res* 139:281–293. <https://doi.org/10.1007/s11120-018-0507-y>
 35. Tsementzi D, Wu J, Deutsch S, Nath S, Rodriguez-R LM, Burns AS, Ranjan P, Sarode N, Malmstrom RR, Padilla CC, Stone BK, Bristow LA, Larsen M, Glass JB, Thamdrup B, Woyke T, Konstantinidis KT, Stewart FJ. 2016. SAR11 bacteria linked to ocean anoxia and nitrogen loss. *Nature* 536:179–183. <https://doi.org/10.1038/nature19068>
 36. Callbeck CM, Canfield DE, Kuypers MMM, Yilmaz P, Lavik G, Thamdrup B, Schubert CJ, Bristow LA. 2021. Sulfur cycling in oceanic oxygen minimum zones. *Limn & Ocean* 66:2360–2392. <https://doi.org/10.1002/lno.11759>
 37. Schmitz RA, Peeters SH, Mohammadi SS, Berben T, van Erven T, Iosif CA, van Alen T, Versantvoort W, Jetten MSM, Op den Camp HJM, Pol A. 2023. Simultaneous sulfide and methane oxidation by an extremophile. *Nat Commun* 14:2974. <https://doi.org/10.1038/s41467-023-38699-9>
 38. Moffett JW, Goepfert TJ, Naqvi SWA. 2007. Reduced iron associated with secondary nitrite maxima in the Arabian sea. *Deep Sea Res Part I Oceanogr Res Pap* 54:1341–1349. <https://doi.org/10.1016/j.dsr.2007.04.004>
 39. Glass JB, Kretz CB, Ganesh S, Ranjan P, Seston SL, Buck KN, Landing WM, Morton PL, Moffett JW, Giovannoni SJ, Vergin KL, Stewart FJ. 2015. Metaomic signatures of microbial metal and nitrogen cycling in marine oxygen minimum zones. *Front Microbiol* 6:998. <https://doi.org/10.3389/fmicb.2015.00998>
 40. Simon J, Gross R, Einsle O, Kroneck PM, Kröger A, Klimmek O. 2000. A NapC/NirT-type cytochrome c (NrfH) is the mediator between the quinone pool and the cytochrome c nitrite reductase of *Wolinella succinogenes*. *Mol Microbiol* 35:686–696. <https://doi.org/10.1046/j.1365-2958.2000.01742.x>
 41. Berg JS, Ahmerkamp S, Pjevac P, Hausmann B, Milucka J, Kuypers MMM. 2022. How low can they go? Aerobic respiration by microorganisms under apparent anoxia. *FEMS Microbiol Rev* 46:1–14. <https://doi.org/10.1093/femsre/fuac006>
 42. Garcia-Robledo E, Padilla CC, Aldunate M, Stewart FJ, Ulloa O, Paulmier A, Gregori G, Revsbech NP. 2017. Cryptic oxygen cycling in anoxic marine zones. *Proc Natl Acad Sci U S A* 114:8319–8324. <https://doi.org/10.1073/pnas.1619844114>
 43. Lutterbeck HE, Arévalo-Martínez DL, Löscher CR, Bange HW. 2018. Nitric oxide (NO) in the oxygen minimum zone off Peru. *Deep Sea Res Part II Top Stud Oceanogr* 156:148–154. <https://doi.org/10.1016/j.dsr2.2017.12.023>
 44. Ward BB, Zafiriou OC. 1988. Nitrification and nitric oxide in the oxygen minimum of the eastern tropical North Pacific. *Deep Sea Res Part A*

- Oceanogr Res Pap 35:1127–1142. [https://doi.org/10.1016/0198-0149\(88\)90005-2](https://doi.org/10.1016/0198-0149(88)90005-2)
45. Zhang IH, Sun X, Jayakumar A, Fortin SG, Ward BB, Babbitt AR. 2023. Partitioning of the denitrification pathway and other nitrite metabolisms within global oxygen deficient zones. *ISME Commun* 3:76. <https://doi.org/10.1038/s43705-023-00284-y>
 46. Katoh K, Rozewicki J, Yamada KD. 2019. MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Brief Bioinform* 20:1160–1166. <https://doi.org/10.1093/bib/bbx108>
 47. Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ. 2016. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res* 44:W232–W235. <https://doi.org/10.1093/nar/gkw256>
 48. Zallot R, Oberg N, Gerlt JA. 2019. The EFI web resource for genomic enzymology tools: leveraging protein, genome, and metagenome databases to discover novel enzymes and metabolic pathways. *Biochemistry* 58:4169–4182. <https://doi.org/10.1021/acs.biochem.9b00735>
 49. Boratyn GM, Thierry-Mieg J, Thierry-Mieg D, Busby B, Madden TL. 2019. Magic-BLAST, an accurate RNA-seq aligner for long and short reads. *BMC Bioinformatics* 20:405. <https://doi.org/10.1186/s12859-019-2996-x>
 50. Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, Dehal P, Ware D, Perez F, Canon S, et al. 2018. KBase: the United States department of energy systems biology knowledgebase. *Nat Biotechnol* 36:566–569. <https://doi.org/10.1038/nbt.4163>
 51. Wu YW, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* 32:605–607. <https://doi.org/10.1093/bioinformatics/btv638>
 52. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 9:5114. <https://doi.org/10.1038/s41467-018-07641-9>
 53. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2022. GTDB-Tk V2: memory friendly classification with the genome taxonomy database. *Bioinformatics* 38:5315–5316. <https://doi.org/10.1093/bioinformatics/btac672>
 54. Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA III, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>
 55. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>
 56. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32:268–274. <https://doi.org/10.1093/molbev/msu300>
 57. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: improving the ultrafast bootstrap approximation. *Mol Biol Evol* 35:518–522. <https://doi.org/10.1093/molbev/msx281>
 58. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* 7:562–578. <https://doi.org/10.1038/nprot.2012.016>
 59. Yu NY, Wagner JR, Laird MR, Melli G, Rey S, Lo R, Dao P, Sahinalp SC, Ester M, Foster LJ, Brinkman FSL. 2010. PSORTb 3.0: improved protein subcellular localization prediction with refined localization subcategories and predictive capabilities for all prokaryotes. *Bioinformatics* 26:1608–1615. <https://doi.org/10.1093/bioinformatics/btq249>