

Gröbner bases, symmetric matrices, and type C Kazhdan–Lusztig varieties

Laura Escobar¹ | Alex Fink² | Jenna Rajchgot³ |
Alexander Woo⁴

¹Department of Mathematics,
Washington University in St. Louis, St.
Louis, Missouri, USA

²School of Mathematical Sciences, Queen
Mary University of London, London, UK

³Department of Mathematics and
Statistics, McMaster University,
Hamilton, ON, Canada

⁴Department of Mathematics and
Statistical Science, University of Idaho,
Moscow, Idaho, USA

Correspondence

Jenna Rajchgot, Department of
Mathematics and Statistics, McMaster
University, Hamilton, ON, Canada.
Email: rajchgot@mcmaster.ca

Funding information

NSF, Grant/Award Numbers: DMS
1855598, DMS 2142656; NSERC,
Grant/Award Number:
RGPIN-2017-05732; Simons Collaboration,
Grant/Award Number: 359792; European
Union's Horizon 2020 research and
innovation program, Grant/Award
Number: No 792432

Abstract

We study a class of combinatorially defined polynomial ideals that are generated by minors of a generic symmetric matrix. Included within this class are the symmetric determinantal ideals, the symmetric ladder determinantal ideals, and the symmetric Schubert determinantal ideals of A. Fink, J. Rajchgot, and S. Sullivant. Each ideal in our class is a type C analog of a Kazhdan–Lusztig ideal of A. Woo and A. Yong; that is, it is the scheme-theoretic defining ideal of the intersection of a type C Schubert variety with a type C opposite Schubert cell, appropriately coordinatized. The Kazhdan–Lusztig ideals that arise are exactly those where the opposite cell is 123-avoiding. Our main results include Gröbner bases for these ideals, prime decompositions of their initial ideals (which are Stanley–Reisner ideals of subword complexes), and combinatorial formulas for their multigraded Hilbert series in terms of pipe dreams.

MSC 2020

14M15, 14M12, 14N15 (primary)

Contents

1. INTRODUCTION	2
2. BACKGROUND	5
2.1. Gröbner bases and initial ideals	5

2.2. Torus actions, multigradings, and K -polynomials	5
2.3. Simplicial complexes and Stanley–Reisner ideals	6
3. KAZHDAN–LUSZTIG VARIETIES	6
3.1. Schubert cells and varieties	6
3.2. Permutations and left–right weak order	8
3.3. Opposite Schubert cells as spaces of matrices	10
3.4. Rank conditions on type C Kazhdan–Lusztig varieties	13
4. SMALL PATCHES	15
4.1. Small patches and symmetric matrices	15
4.2. Equations for type C Kazhdan–Lusztig varieties	21
4.3. Torus action of type C Kazhdan–Lusztig varieties and the weights for $v \geq v_{\square}$	23
5. TYPE C SUBWORD COMPLEXES AND VERTEX DECOMPOSITION	25
5.1. Subword complexes	25
5.2. Labeling the vertices with variables	26
6. PROOF OF THEOREM 4.15	28
6.1. The Stanley–Reisner ideal is contained in the initial ideal	29
6.2. Equality of K -polynomials	34
6.3. Proof of Theorem 4.15	36
7. K -POLYNOMIAL FORMULAS VIA PIPE DREAMS	37
7.1. Type C pipe dreams on small patches	37
7.2. Multidegrees and K -polynomials	40
7.3. Combinatorics of type C pipe dreams	43
8. BEYOND SMALL PATCHES	44
ACKNOWLEDGMENTS	47
JOURNAL INFORMATION	47
REFERENCES	48

1 | INTRODUCTION

Let $\mathbb{K}$ be a field of characteristic zero. In this paper, we study a class of generalized symmetric determinantal ideals. Each ideal in our class is defined by imposing certain *combinatorial* southwest rank conditions on an $n \times n$ symmetric matrix M whose i, j entry is either zero or an indeterminate $z_{ij} = z_{ji}$ and whose nonzero entries lie in a skew partition, in English conventions. Among the ideals in our class are the symmetric determinantal ideals, the symmetric ladder determinantal ideals [15, 16], and the symmetric Schubert determinantal ideals of [9]. We plan to describe in detail the connection with symmetric ladder determinantal ideals in a separate paper.

Let $R = \mathbb{K}[z_{ij}]$ be the polynomial ring in the variables appearing in a matrix M as above. We interpret this ring in terms of a type C opposite Schubert cell. Let G be the symplectic group $Sp_{2n}(\mathbb{K})$, represented as the group of $2n \times 2n$ matrices preserving the form $e_1 \wedge e_{2n} + \cdots + e_n \wedge e_{n+1}$. We will work with the following Borel subgroups of G :

$$B_G^+ := \{\text{upper triangular matrices in } G\} \quad \text{and} \quad B_G^- := \{\text{lower triangular matrices in } G\}.$$

The type C flag variety is G/B_G^+ , and an opposite Schubert cell is a B_G^- -orbit in G/B_G^+ . These cells are indexed by elements of the Weyl group C_n of G , which can be identified with the set of permutations

$$C_n = \{v_1 \dots v_{2n} \in S_{2n} : v_{2n+1-i} = 2n+1-v_i \text{ for } i = 1, \dots, n\}.$$

Then, the ring R is the coordinate ring of a type C opposite Schubert cell associated to some 123-avoiding permutation $v \in C_n$, with an appropriate choice of coordinates (see Proposition 4.7). From the point of view of Schubert cells in G/B_G^+ , our choices of symplectic form and coordinates have a long history. These coordinates were used extensively to study Schubert cells by W. Fulton and P. Pragacz [13], who were likely aware of the connection to symmetric matrices at least in some special cases.

This choice of coordinates allows us to study a large class of generalized symmetric determinantal ideals from the point of view of Kazhdan–Lusztig varieties in G/B_G^+ . Each ideal we encounter is obtained by imposing southwest rank conditions on M , using combinatorial rules encoded by some $w \in C_n$. Given $v, w \in C_n$, we denote by $\mathcal{N}_{v,w}$ the affine variety associated to one of our ideals; these varieties form a subclass of type C Kazhdan–Lusztig varieties. A Schubert variety is a B_G^+ -orbit closure in G/B_G^+ , and a Kazhdan–Lusztig variety is the intersection of a Schubert variety with an opposite Schubert cell.

In general, Kazhdan–Lusztig varieties provide affine neighborhoods of the T -fixed points of Schubert varieties, and they have been used to study singularities of Schubert varieties. One such instance is [42], in which A. Woo and A. Yong introduced Kazhdan–Lusztig ideals of type A for this purpose. Each Kazhdan–Lusztig ideal is the prime defining ideal of a type A Kazhdan–Lusztig variety, appropriately coordinatized. In [43], Woo and Yong showed that Kazhdan–Lusztig ideals of type A possess nice Gröbner bases for which the corresponding initial ideals are Stanley–Reisner ideals of vertex decomposable balls or spheres. They furthermore proved multiple combinatorial formulas for their multigraded Hilbert series.

Similarly, a type C Kazhdan–Lusztig ideal is the defining ideal of a type C Kazhdan–Lusztig variety. In our main theorem, we use the interpretation of $\mathcal{N}_{v,w}$ as a type C Kazhdan–Lusztig variety to give a Gröbner basis with squarefree initial terms for the ideals we encounter. We give R a term order which is *diagonal*. Roughly this means that the leading term of any minor is the product of the diagonal entries of the submatrix it arises from. One example of a diagonal term order is the lexicographic term order where $z_{ij} > z_{i'j'}$ if and only if either $i > i'$, or $i = i'$ and $j > j'$. Our main result is as follows. It is stated precisely as Theorem 4.15 in the main body of the paper.

Theorem. *Let $v, w \in C_n$ and v be 123-avoiding. The determinants defining the ideal of $\mathcal{N}_{v,w}$ form a Gröbner basis with respect to any diagonal term order.*

This result is proved in Section 6 using K -polynomials and the subword complexes of A. Knutson and E. Miller [29, 30]. Note that our conventions are upside-down from those of Knutson and Miller, so our diagonal term orders are indeed analogs of their antidiagonal term orders.

In [28], Knutson showed that the defining ideal of any Kazhdan–Lusztig variety has a Gröbner basis whose leading terms are squarefree, and, in [27], he determined that the resulting initial ideal is the Stanley–Reisner ideal of a certain subword complex. However, he did not provide a Gröbner basis. Up to sign, our coordinates agree with the Bott–Samelson coordinates of [28]. Hence, our results make [28, Theorem 7] more explicit by describing the coordinates and stating which minors in the Gröbner basis arise from each element of the essential set (which corresponds to Knutson’s “basic elements”).

In Section 7, we define type C pipe dreams and use them to give consequences to Theorem 4.15. Namely, in Corollary 7.2, we give prime decompositions of the initial ideals, in Proposition 7.3, we give combinatorial formulas for their multigraded Hilbert series, and in Proposition 7.4, we give combinatorial formulas for their K -polynomials. Up to a change in convention, these formulas

give, in the case where v is 123-avoiding, combinatorial models of S. Billey's formula [6] and its extension to K -theory by W. Graham [17] and M. Willems [40] for a particular choice of reduced word. (We note that Billey's formula was first stated by H. Andersen, J. Jantzen, and W. Soergel [1] and independently rediscovered in different but related context by Billey [6]; see J. Tymoczko's survey paper [39] for details and more recent developments.) We note that in recent work, E. Smirnov and A. Tutubalina [37] have studied pipe dreams in all classical groups; these differ from ours even in the special case we describe.

The polynomials given in these formulas also have an interpretation as a particular specialization of type C double Schubert and double Grothendieck polynomials, which are *stable* equivariant Chow [21, Theorem 10.8] and K -theory [26, Theorem 2] classes of type C Schubert varieties. Here, being stable classes means they are lifts of these classes, independent of the rank of the ambient flag variety, which satisfy certain recurrences and boundary conditions parallel to those holding in the type A case. From these polynomials, the multidegrees and K -polynomials of Kazhdan–Lusztig varieties are obtained in two equivalent ways, either geometrically by restricting to affine patches, which is equivalent to localization at torus fixed points, or algebraically according to particular restriction maps. T. Ikeda, L. Mihalcea, and H. Naruse [21] were the first to define type C double Schubert polynomials, and they gave several formulas including two using divided difference operators as well as an algebraic restriction map for recovering local classes. Type C double Grothendieck polynomials were explicitly defined by A. Kirillov in [25]. The connection to geometry was made by Kirillov and Naruse in [26]. Under this interpretation of our formulas as specializations of type C double Schubert and double Grothendieck polynomials, one can consider these formulas as type C analogs of the type A specialization formulas of A. Buch and R. Rimanyi [8].

We note that the *symmetric matrix Schubert varieties* defined and studied by Z. Hamaker, E. Marberg, and B. Pawłowski in [19, 34] are not special cases of the varieties that we consider in the present paper. The varieties studied by Hamaker, Marberg, and Pawłowski are defined by imposing northwest rank conditions on symmetric matrices, while we impose southwest (and northeast) conditions. Correspondingly, the pipe dreams they introduce are symmetric across an axis perpendicular to our axis of symmetry. The varieties in [19, 34] are related to Borel group orbit closures in G/K where $G = GL_n$ is a general linear group and $K = O_n$ is an orthogonal subgroup of G .

Outline of this paper

In Section 2, we give the commutative algebra background for the paper. In Section 3, we establish the notation and setup for type C Kazhdan–Lusztig varieties. In Section 4, we introduce the coordinates of Proposition 4.7 which we use for opposite Schubert cells associated to 123-avoiding permutations. Then, in Proposition 4.12, we use these coordinates to describe the defining ideal of $\mathcal{N}_{v,w}$ as a generalized symmetric determinantal ideal. In preparation for computing the K -polynomials of our ideals, we describe in Section 4.3 the weights of the coordinates with respect to the action of the torus of diagonal matrices. In Section 5, we give background on subword complexes, and, for the complexes related to our ideals, we describe how to label their vertices using our coordinates. We also state the vertex decomposition of subword complexes, which we will use to compare the K -polynomials of our ideals with the K -polynomials of the Stanley–Reisner ideals for subword complexes. We then proceed to prove Theorem 4.15 in Section 6. In Section 7, we introduce type C pipe dreams for small patches and give various consequences to Theorem 4.15.

Lastly, in Section 8, we show that our Gröbner basis result (Theorem 4.15) does not naturally extend beyond the small patch setting to general type C Kazhdan–Lusztig ideals.

2 | BACKGROUND

2.1 | Gröbner bases and initial ideals

A *term order* $<$ is a total order on the monomials in a polynomial ring R with respect to which 1 is minimal and such that if m, m', m'' are monomials such that $m' < m''$, then $mm' < mm''$. One class of term orders we will use are *lexicographic* term orders. Given a total ordering $x_1 > \dots > x_k$ on the variables of R , an exponent vector $(a_1, \dots, a_k) \in \mathbb{N}^k$ can be assigned to any monomial $m = x_1^{a_1} \dots x_k^{a_k}$; then two monomials compare in the lexicographic term order just as their exponent vectors compare in the lexicographic order on $\mathbb{N}^k$. More precisely, $x_1^{a_1} \dots x_k^{a_k} < x_1^{b_1} \dots x_k^{b_k}$ if and only if there is some $1 \leq i \leq k$ so that $a_1 = b_1, \dots, a_{i-1} = b_{i-1}$, and $a_i < b_i$. In particular, the variables themselves still compare as $x_1 > \dots > x_k$ in the lexicographic term order.

The *initial term* of a polynomial in R , with respect to a fixed term order $<$, is the maximum of the monomials in its support. If I is an ideal of R , then its *initial ideal*, denoted as $\text{in}_< I$, is the ideal generated by all initial terms of elements of I . A *Gröbner basis* for I is a generating set for I whose initial terms generate $\text{in}_< I$.

2.2 | Torus actions, multigradings, and K -polynomials

One reference for the material in this section is [35, Chapter 8].

Suppose that a torus $T = (\mathbb{K}^*)^n$ acts on affine space $\mathbb{A}^k = \text{Spec } \mathbb{K}[z_1, \dots, z_k]$ with weights $-a_1, \dots, -a_k \in \mathbb{Z}^n$. This means that, given $x = (x_1, \dots, x_n) \in T$ and $p \in \mathbb{A}^k$ with coordinates $p = \sum_{i=1}^k z_i(p) f_i$ (where $\{f_i\}$ denotes the dual basis to $\{z_i\}$),

$$x \cdot p = \sum_{i=1}^k x^{-a_i} z_i(p) f_i,$$

where

$$x^{-a_i} = \prod_{j=1}^n x_j^{-a_{ij}}.$$

Then T acts on the coordinate functions $z_1, \dots, z_k$ with weights $a_1, \dots, a_k$ respectively. This action induces a $\mathbb{Z}^n$ -grading on the ring $R = \mathbb{K}[z_1, \dots, z_k]$ given by setting the degree of z_i as a_i , so that $\deg(z_1^{b_1} \dots z_k^{b_k}) = \sum_{i=1}^k b_i a_i$.

Given $a \in \mathbb{Z}^n$ and a graded R -module M , let M_a denote the a th-graded piece of M . Suppose that $\dim_{\mathbb{K}}(M_a)$ is finite for all a , which will be the case if $a_1, \dots, a_k$ generate a pointed cone in $\mathbb{Z}^n \otimes \mathbb{R}$ and M is finitely generated. Then define the *Hilbert series* of M to be

$$\mathcal{H}(M; \mathbf{t}) = \sum_{a \in \mathbb{Z}^n} \dim_{\mathbb{K}}(M_a) \mathbf{t}^a.$$

Furthermore, define the K -polynomial of M as

$$\mathcal{K}(M; \mathbf{t}) = \mathcal{H}(M; \mathbf{t}) \prod_{i=1}^k (1 - \mathbf{t}^{\deg(z_i)}).$$

This is a Laurent polynomial in the variables t_i . Finally, the *multidegree* of M with its given multigrading, denoted as $C(M; \mathbf{t})$, is the sum of all lowest degree terms in $\mathcal{K}(M; 1 - t_1, \dots, 1 - t_k)$.

Note that an ideal and its initial ideal have equal K -polynomials and equal multidegrees. (This is called the *degenerative* property in [35].) Furthermore, if $N \subseteq M$ and $\mathcal{K}(N; \mathbf{t}) = \mathcal{K}(M; \mathbf{t})$, then $N = M$. Also, multidegrees are *additive* in the sense that the multidegree $C(R/I; \mathbf{t})$ is the sum $\sum_J C(R/J; \mathbf{t})$ where the sum is over those J in a primary decomposition of I such that $\sqrt{J}$ is a minimal prime of I that has the same height as I (see [35, §8.5]).

2.3 | Simplicial complexes and Stanley–Reisner ideals

A *simplicial complex* Δ on the vertex set V is a set of subsets of V , called *faces*, such that if $F \in \Delta$, then all subsets of F are in Δ . A *facet* of Δ is a maximal face under containment. If Δ is a simplicial complex on V , and $z \notin V$, then the *cone* $\text{cone}_z \Delta$ is the simplicial complex

$$\{F \subseteq V \cup \{z\} : F \cap V \in \Delta\}$$

on vertex set $V \cup \{z\}$.

The *Stanley–Reisner ideal* of Δ is the ideal I_Δ of the polynomial ring $R = \mathbb{K}[V]$ generated by products of variables that index nonfaces of Δ , that is,

$$I_\Delta := \left\langle \prod_{z \in Z} z : Z \subseteq V, Z \notin \Delta \right\rangle.$$

3 | KAZHDAN–LUSZTIG VARIETIES

In this section, we recall background on Schubert varieties in flag varieties of types A and C . In particular, we discuss Kazhdan–Lusztig varieties, which we define (following [42]) to be the intersection of a Schubert variety with an opposite Schubert cell. These are affine varieties.

3.1 | Schubert cells and varieties

Fix an integer $n \geq 1$, and let E be the $2n \times 2n$ matrix

$$E := \begin{bmatrix} 0 & J_n \\ -J_n & 0 \end{bmatrix},$$

where J_n is the $n \times n$ antidiagonal matrix with antidiagonal entries 1. The matrix E determines a nondegenerate, skew-symmetric bilinear form on $\mathbb{K}^{2n}$. The symplectic group $Sp_{2n}(\mathbb{K})$ is

$$Sp_{2n}(\mathbb{K}) := \{M \in GL_{2n}(\mathbb{K}) : E(M^t)^{-1}E^{-1} = M\},$$

or, equivalently, it is the fixed point set of the involution

$$\sigma : GL_{2n}(\mathbb{K}) \rightarrow GL_{2n}(\mathbb{K}), \quad \sigma(M) = E(M^t)^{-1}E^{-1}.$$

Following [33, Chapter 6], we let $H := GL_{2n}(\mathbb{K})$ and $G := Sp_{2n}(\mathbb{K})$. We will work with the following Borel subgroups of H :

$$B_H^+ := \{\text{upper triangular matrices in } H\} \quad \text{and} \quad B_H^- := \{\text{lower triangular matrices in } H\}.$$

These give rise to the following Borel subgroups of G :

$$B_G^+ = (B_H^+)^{\sigma} \quad \text{and} \quad B_G^- = (B_H^-)^{\sigma}.$$

Consider the type A flag variety H/B_H^+ . A *Schubert cell* in this flag variety is a B_H^+ -orbit for the left action of B_H^+ on H/B_H^+ by multiplication, and a *Schubert variety* is its closure. An *opposite Schubert cell* is a B_H^- -orbit in H/B_H^+ and an *opposite Schubert variety* is its closure. In the type C flag variety G/B_G^+ , Schubert cells and varieties are defined analogously by replacing appearances of H and $B_H^{\pm}$ in the above definitions by G and $B_G^{\pm}$, respectively.

Denote by S_{2n} the Weyl group of H . Given $w \in S_{2n}$, we denote by $P(w)$ the permutation matrix having its nonzero entries in positions $(w(j), j)$ for $j = 1, \dots, n$. We use this convention to be consistent with [43]. Each Schubert cell in the type A flag variety H/B_H^+ is equal to some orbit $B_H^+ \cdot P(w)B_H^+/B_H^+$ where $w \in S_{2n}$. The analog is true for opposite Schubert cells: every opposite Schubert cell in H/B_H^+ equals

$$\Omega_v^{A\circ} := B_H^- \cdot P(v)B_H^+/B_H^+$$

for some $v \in S_{2n}$. We remark that here, and throughout the remainder of the paper, we use the letter “ w ” for permutations indexing Schubert cells or varieties, and we use the letter “ v ” for permutations indexing opposite Schubert cells.

The Weyl group C_n of G can be identified with the set of permutations

$$C_n = \{v_1 \dots v_{2n} \in S_{2n} : v_i = 2n + 1 - v_{2n+1-i} \text{ for } i = 1, \dots, n\}. \quad (1)$$

Equivalently, C_n consists of the $v \in S_{2n}$ such that $w_0 v w_0 = v$, where w_0 is the longest element of S_{2n} . In the type C flag variety G/B_G^+ , Schubert and opposite Schubert cells and varieties are indexed by elements of C_n . Concretely, given $w \in C_n$, the permutation matrix $P(w)$ is an element of G , so $B_G^+ \cdot P(w)B_G^+/B_G^+$ is a Schubert cell, and every Schubert cell is of this form. The analogous statements hold for Schubert varieties and opposite Schubert cells and varieties. We denote the type C opposite Schubert cells by

$$\Omega_v^{\circ} := B_G^- \cdot P(v)B_G^+/B_G^+$$

for $v \in C_n$.

It is useful to note that type C Schubert cells and varieties are the σ -fixed point sets of type A Schubert cells and varieties. See also the treatment in [9, Theorem 2.5].

Theorem 3.1. [33, Proposition 6.1.1.1] *The involution σ induces a natural involution $\sigma : H/B_H^+ \rightarrow H/B_H^+$.[†] For $v \in C_n$, the opposite Schubert cell Ω_v° is stable under σ and*

$$\Omega_v^\circ = (\Omega_v^{A^\circ})^\sigma.$$

In other words, Ω_v° consists of the σ -fixed points of the type A opposite Schubert cell $\Omega_v^{A^\circ}$.

Let X_w^A denote the type A Schubert variety $\overline{B_H^+ \cdot P(w)B_H^+ / B_H^+}$ and let X_w denote the type C Schubert variety $\overline{B_G^+ \cdot P(w)B_G^+ / B_G^+}$. Following [42] (see also [43]), we refer to the intersection of a Schubert variety with an opposite Schubert cell as a *Kazhdan–Lusztig variety*. We denote the type A Kazhdan–Lusztig variety as

$$\mathcal{N}_{v,w}^A = X_w^A \cap \Omega_v^{A^\circ},$$

and the type C Kazhdan–Lusztig variety as

$$\mathcal{N}_{v,w} = X_w^A \cap \Omega_v^\circ. \quad (2)$$

Despite the appearances of $H = GL_{2n}(\mathbb{K})$ in the latter intersection above, $\mathcal{N}_{v,w}$ is indeed equal to the intersection of a type C Schubert variety with a type C opposite Schubert cell. This follows immediately from [33, Proposition 6.1.1.2], which says that

$$X_w = X_w^A \cap G/B_G^+,$$

as schemes, under the natural inclusion $G/B_G^+ \hookrightarrow H/B_H^+$.

We remark that Kazhdan–Lusztig varieties are useful for studying singularities of Schubert varieties using computational algebraic methods. This is because a neighborhood of a torus fixed point in a Schubert variety is isomorphic, up to a factor of an affine space, to a Kazhdan–Lusztig variety, which is an affine variety. This isomorphism is due to D. Kazhdan and G. Lusztig [24, Lemma A.4], and explained in [42, Section 3]. We will describe the prime defining ideals of Kazhdan–Lusztig varieties in Section 3.4.

3.2 | Permutations and left-right weak order

The *simple reflections* in S_m are the permutations $s_1, \dots, s_{m-1}$, where s_i transposes i and $i + 1$. In C_n , define the simple reflections to be $c_0, c_1, \dots, c_{n-1}$, where $c_0 \in C_n$ is the permutation that transposes n and $n + 1$, and for $i = 1, \dots, n - 1$, $c_i \in C_n$ is the permutation that transposes $n + i$ with $n + i + 1$ (so, it must also transpose $n - i$ and $n - i + 1$). We warn the reader that these indexing conventions for S_m and C_n are different: under the defining embedding $C_n \subseteq S_{2n}$, the simple reflection c_i is

[†] We abuse notation and use σ for both maps.

identified with s_n if $i = 0$ or $s_{n-i}s_{n+i}$ otherwise, not with something built from s_i . Both S_m and C_n are generated by their sets of simple reflections.

If $W = S_m$ (resp. C_n), a *reduced word* for $v \in W$ is a sequence $Q = (\alpha_1, \dots, \alpha_\ell)$ such that $v = s_{\alpha_1} \cdots s_{\alpha_\ell}$ (resp. $v = c_{\alpha_1} \cdots c_{\alpha_\ell}$) and ℓ is minimized. We denote by $\ell_W(v)$ the length of any reduced word for $v \in W$. When there is no chance for confusion, we omit the subscript W from our notation for length.

Throughout the paper, we let $<_R$ denote the *right weak order* on S_m ; namely, $u \leq_R v$ if some initial substring of some reduced word for v is a reduced word for u . Similarly, $<_L$ denotes the *left weak order* on S_m , which is defined by declaring that $u \leq_L v$ if some terminal substring of some reduced word for v is a reduced word for u . The *left-right weak order* on S_m is denoted throughout the paper by $<$ and defined by $u \leq v$ if $v = s_{\alpha_1} \cdots s_{\alpha_a} u s_{\beta_1} \cdots s_{\beta_b}$ and $\ell(v) = \ell(u) + a + b$. We write $u < v$ if v covers u in left-right weak order. Both weak orders as well as the two sided weak order on C_n are induced by that on S_{2n} . Thus, we use the same notation for them.

We let $<_{Br}$ denote the Bruhat order; namely, $v \geq_{Br} w$ if some reduced word Q for v has as a subword a reduced word for w . If this holds for one reduced word Q then it holds for all such reduced words.

A simple reflection c_k is a (right) *ascent* of $v \in C_n$ if $vc_k >_{Br} v$ and a (right) *descent* of v otherwise, namely, if $vc_k <_{Br} v$. The *last ascent* of v is the ascent c_k where k is maximized. Note that vc_k and v compare the same way in the Bruhat, right weak, and left-right weak orders: vc_k is either greater than v in all three or less than v in all three.

Our convention for the (Rothe) *diagram* of a permutation $w \in S_m$ is the set


$$D(w) = \{(w(j), i) : i < j \text{ and } w(i) < w(j)\}.$$

It is drawn by placing boxes in an $n \times n$ matrix in the positions given by elements of $D(w)$. There is a familiar pictorial procedure to obtain $D(w)$ from $P(w)$: one replaces each 1 by a $\bullet$, deletes all 0s, and draws at each $\bullet$ the “hook” that extends to the east and north of the $\bullet$. The entries of the matrix that no hook passes through are the elements of $D(w)$.

Example 3.2. The diagram of $w = 365124$ is

$$D(w) = \{(4, 1), (5, 1), (6, 1), (2, 4), (4, 4), (4, 5)\}$$

and it is drawn



Rothe diagrams are important to us for providing coordinates for opposite Schubert cells and Kazhdan–Lusztig varieties, as we see in the next subsection.

3.3 | Opposite Schubert cells as spaces of matrices

Let $H = GL_m(\mathbb{K})$ and $B_H^+ \subseteq H$ be the Borel subgroup of upper triangular matrices. For $v \in S_m$, let $\Sigma_v^A \subseteq H$ be the set of matrices M such that if $P(v)_{ij} = 1$, then $M_{ij} = 1$, and, otherwise, if $(i, j) \notin D(v)$, then $M_{ij} = 0$.

Proposition 3.3. [12, Section 10.2] *The map $\pi_H : H \rightarrow H/B_H^+$ sending a matrix M to its coset MB_H^+/B_H^+ induces a (scheme-theoretic) isomorphism from the space of matrices Σ_v^A to the opposite Schubert cell $\Omega_v^{A^\circ}$.*

We can similarly identify each type C opposite Schubert cell with a space of matrices using the map π_H . We now do this explicitly, to prepare for the explicit coordinate-dependent presentation needed in our main theorem. The material discussed in this section follows from general theory on algebraic groups and flag varieties, for example, [23, Chapter 13], and this particular presentation features in [5].

Let $v \in C_n$. Identifying Ω_v° as a closed subvariety of $\Omega_v^{A^\circ}$ by Theorem 3.1, define the space of matrices

$$\Sigma_v := \pi_H^{-1}(\Omega_v^\circ), \quad (3)$$

and note that Σ_v , which is a closed subvariety of Σ_v^A , is isomorphic to Ω_v° .

Furthermore, we identify Kazhdan–Lusztig varieties with spaces of matrices by letting

$$\Sigma_{v,w}^A := \pi_H^{-1}(\mathcal{N}_{v,w}^A)$$

and

$$\Sigma_{v,w} := \pi_H^{-1}(\mathcal{N}_{v,w}).$$

We now wish to describe Σ_v as the set of σ -fixed points of Σ_v^A . This description will follow from the containment $\sigma(\Sigma_v^A) \subseteq \Sigma_v^A$ for $v \in C_n$. In order to prove this containment, the following factorization of the matrices in Σ_v^A is useful. Let U_i^- be the unipotent subgroup of H consisting of matrices with 1s along the diagonal, 0s in all off-diagonal positions except for $(i+1, i)$, and an arbitrary element of $\mathbb{K}$ in position $(i+1, i)$.

Proposition 3.4. *Define $\tilde{\Sigma}_\alpha := P(s_\alpha)U_\alpha^-$. Given $v \in S_m$ and $(\alpha_1, \dots, \alpha_\ell)$ a reduced word for w_0v , the map*

$$\mathcal{M} : \tilde{\Sigma}_{\alpha_1} \times \dots \times \tilde{\Sigma}_{\alpha_\ell} \rightarrow \Sigma_v^A, \quad (a_1, \dots, a_\ell) \mapsto P(w_0)a_1 \dots a_\ell$$

is an isomorphism.

Proof. We proceed by induction on $\ell(w_0v)$. The base case is when $v = w_0$, and it is clear that the result holds in this case. For the inductive case, let $vs_i \succ v$ and write a reduced expression

$w_0 v s_i = s_{\alpha_1} \cdots s_{\alpha_\ell}$. By induction,

$$\mathcal{M} : \tilde{\Sigma}_{\alpha_1} \times \cdots \times \tilde{\Sigma}_{\alpha_\ell} \rightarrow \Sigma_{v s_i}^A, \quad (a_1, \dots, a_\ell) \mapsto P(w_0) a_1 \cdots a_\ell$$

is an isomorphism. So, it suffices to show that the image of the multiplication map

$$m : \Sigma_{v s_i}^A \times \tilde{\Sigma}_i \rightarrow H, \quad (a, b) \mapsto ab \quad (4)$$

is Σ_v^A and that it is an isomorphism upon restricting the codomain to Σ_v^A . To see this, let $a \in \Sigma_{v s_i}^A$ and $b \in \tilde{\Sigma}_i$. Assume that the (i, i) -entry of the matrix b is equal to $t \in \mathbb{K}$. Observe that ab is obtained from a by performing two elementary column operations: first swap columns i and $i + 1$, and then replace column i by column i plus t times column $i + 1$. Let $a_{[i, i+1]}$ and $(ab)_{[i, i+1]}$ be the submatrices of a and ab , respectively, consisting of columns i and $i + 1$. Because $v s_i \succ v$, after removing all rows of $a_{[i, i+1]}$ and $(ab)_{[i, i+1]}$ that do not have pivots, we are left with:

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \text{ (from } a) \quad \text{and} \quad \begin{bmatrix} 1 & 0 \\ t & 1 \end{bmatrix} \text{ (from } ab).$$

Thus, every matrix in the image of the map m of (4) can be factored uniquely as ab , and so, m is an isomorphism onto its image. Finally, a straightforward check shows that the locations of diagram boxes in rows without pivots of $a_{[i, i+1]}$ and $(ab)_{[i, i+1]}$ coincide. (Alternatively, see [43, Lemma 6.5].) Hence, the image of m is contained in Σ_v^A . As m is an isomorphism onto its image, and Σ_v^A and the domain of m are affine spaces of the same dimension, the proposition is proved. $\square$

Corollary 3.5. *Let $H = GL_{2n}(\mathbb{K})$. The map $\sigma : H \rightarrow H$ restricts to an isomorphism*

$$\sigma : \Sigma_v^A \rightarrow \Sigma_{w_0 v w_0}^A.$$

In particular, if $v \in C_n \subseteq S_{2n}$, then σ maps Σ_v^A isomorphically onto itself.

Proof. We first observe, by a straightforward direct check, that σ maps $\tilde{\Sigma}_\alpha$ isomorphically onto $\tilde{\Sigma}_{2n-\alpha}$. Let $v \in S_{2n}$ and let $(\alpha_1, \dots, \alpha_\ell)$ be a reduced word for $w_0 v$. By our observation, we have an isomorphism,

$$\tilde{\Sigma}_{\alpha_1}^A \times \cdots \times \tilde{\Sigma}_{\alpha_\ell}^A \rightarrow \tilde{\Sigma}_{2n-\alpha_1}^A \times \cdots \times \tilde{\Sigma}_{2n-\alpha_\ell}^A. \quad (5)$$

Noting that $(2n - \alpha_1, \dots, 2n - \alpha_\ell)$ is a reduced word for $w_0(w_0 v)w_0 = v w_0$, the first statement of the corollary follows by applying Proposition 3.4, which states that the domain of (5) is isomorphic to Σ_v^A and the codomain is isomorphic to $\Sigma_{w_0 v w_0}^A$.

The second statement follows immediately since $w_0 v w_0 = v$ for any $v \in C_n \subseteq S_{2n}$. $\square$

Corollary 3.6. *For $v \in C_n$, $\Sigma_v = (\Sigma_v^A)^\sigma$.*

Proof. Let $M \in \Sigma_v$. By (3) and Theorem 3.1, $MB_H^+/B_H^+ \in \Omega_v^\circ = (\Omega_v^{A^\circ})^\sigma$. This implies that $\sigma(M)B_H^+/B_H^+ = MB_H^+/B_H^+$. Since $M \in \Sigma_v^A$, by Corollary 3.5, $\sigma(M) \in \Sigma_v^A$. We can then apply Proposition 3.3 to deduce that $\sigma(M) = M$ and conclude that $M \in (\Sigma_v^A)^\sigma$. Conversely, let $M \in (\Sigma_v^A)^\sigma$.

Then, $\sigma(MB_H^+/B_H^+) = \sigma(M)B_H^+/B_H^+ = MB_H^+/B_H^+$. Furthermore, by Proposition 3.3, we have that $MB_H^+/B_H^+ \in (\Omega_v^{A^\circ})^\sigma = \Omega_v^\circ$. We conclude that $M \in \Sigma_v = \pi_H^{-1}(\Omega_v^\circ)$. $\square$

We end with two examples of computing Σ_v . The first shows that, in general, the space of matrices Σ_v can be complicated. The second shows that for particular choices of v , Σ_v is easy to describe. From Section 4, we will generally restrict to only these nice Σ_v .

Example 3.7. Given $v = 231\,645$, we have that

$$\Sigma_v^A = \pi_H^{-1}(\Omega_v^{A^\circ}) = \left\{ \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ a & 1 & 0 & 0 & 0 & 0 \\ b & c & d & 0 & 1 & 0 \\ e & f & g & 0 & h & 1 \\ i & j & k & 1 & 0 & 0 \end{bmatrix} : a, b, \dots, k \in \mathbb{K} \right\}.$$

Since

$$\sigma \left(\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ a & 1 & 0 & 0 & 0 & 0 \\ b & c & d & 0 & 1 & 0 \\ e & f & g & 0 & h & 1 \\ i & j & k & 1 & 0 & 0 \end{bmatrix} \right) = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ -h & 1 & 0 & 0 & 0 & 0 \\ -ch + f & c & j & 0 & 1 & 0 \\ -af + (ac - b)h + e & -ac + b & -aj + i & 0 & -a & 1 \\ -dh + g & d & k & 1 & 0 & 0 \end{bmatrix},$$

we can equate the entries of $\sigma(M)$ with the entries of $M \in \Sigma_v^A$ to obtain the conditions defining Σ_v . It is straightforward to verify that

$$\Sigma_v = \left\{ \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ a & 1 & 0 & 0 & 0 & 0 \\ b & c & d & 0 & 1 & 0 \\ e & -ac + b & -ad + i & 0 & -a & 1 \\ i & d & k & 1 & 0 & 0 \end{bmatrix} : a, b, c, d, e, i, k \in \mathbb{K} \right\}.$$

Example 3.8. By a similar computation to the one in the previous example, one can check that the space of matrices $\Sigma_{321\,654}$ is naturally identified with the space of 3×3 symmetric matrices. That is,

$$\Sigma_{321\,654} = \left\{ \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{11} & z_{12} & z_{13} & 0 & 0 & 1 \\ z_{12} & z_{22} & z_{23} & 0 & 1 & 0 \\ z_{13} & z_{23} & z_{33} & 1 & 0 & 0 \end{bmatrix} : z_{ij} \in \mathbb{K} \right\}.$$

3.4 | Rank conditions on type C Kazhdan–Lusztig varieties

Given $w \in S_{2n}$, let $r_w : \{1, \dots, 2n\} \times \{1, \dots, 2n\} \rightarrow \{1, \dots, 2n\}$ be the *rank function* of w , defined by

$$r_w(p, q) = |\{i \leq q : w(i) \geq p\}|,$$

so that $r_w(p, q)$ is the number of entries of w weakly southwest of (p, q) .

Given a matrix M , let $\tau_{p,q}(M)$ denote the submatrix of entries of M weakly southwest of position (p, q) . A matrix $M \in \Sigma_v^A$ is in $\Sigma_{v,w}^A$ if and only if, for all $p, q \in [2n]$, $\tau_{p,q}(M)$ has rank at most $r_w(p, q)$. Not all of these rank conditions are necessary to determine $\Sigma_{v,w}^A$. In type A, Fulton [11] defined the *essential set*, which gives a smaller set of sufficient conditions, as the set of boxes on the northeast [†] corners of the connected components of $D(w)$. To be precise, let


$$E^A(w) := \{(p, q) \in D(w) : (p-1, q), (p, q+1) \notin D(w)\}.$$

Equivalently, one can also define

$$E^A(w) = \{(p, q) : w(q) < p \leq w(q+1), w^{-1}(p-1) \leq q < w^{-1}(p)\}.$$

Then $M = \Sigma_{v,w}^A$ if and only if the size $r_w(p, q) + 1$ minors of $\tau_{p,q}(M)$ vanish for all $(p, q) \in E^A(w)$, and, in fact, these equations define $\Sigma_{v,w}^A$ as a subvariety of Σ_v^A scheme theoretically [42, Proposition 3.1].

Example 3.9. Let $w = 465213$. We have



so, the (type A) essential set of w is $E^A(w) = \{(5, 1), (3, 5)\}$. Furthermore, $r_w(5, 1) = 0$ and $r_w(3, 5) = 3$.

Suppose that $v = 231645$, as featured in Example 3.7. Then $M \in \Sigma_{v,w}^A$ if and only if $M \in \Sigma_v^A$ and the size $r_w(p, q) + 1$ minors of $\tau_{p,q}(M)$ vanish for all $(p, q) \in E^A(w)$. In particular, $e = i = 0$, and we have five additional equations coming from the 4×4 minors of

$$\begin{bmatrix} a & 1 & 0 & 0 & 0 \\ b & c & d & 0 & 1 \\ e & f & g & 0 & h \\ i & j & k & 1 & 0 \end{bmatrix}.$$

[†] Note that Fulton uses different conventions to ours. His hooks emanate east and south rather than east and north, and he works in $B^- \setminus G$ rather than G/B^- , so his permutation matrices are the transpose of ours.

Recall from (2) that $\mathcal{N}_{v,w}$ is the intersection of Ω_v with a type A Schubert variety. Hence, the rank conditions defining $\Sigma_{v,w}$ are the same as those defining $\Sigma_{v,w}^A$, but now applied to Σ_v instead of Σ_v^A . In type C, Anderson [2] showed that a smaller set suffices. (Some details were made more explicit in [41, Section 4].) First, for a permutation $w \in C_n$, boxes of $E^A(w)$ always come in pairs. If $(p, q) \in E^A(w)$, then $(2n + 2 - p, 2n - q) \in E^A(w)$, and, furthermore,[†]

$$r_w(2n + 2 - p, 2n - q) = p - q - 1 + r_w(p, q).$$

We will choose one box out of each pair by requiring that $p \geq n + 1$, and, if $p = n + 1$, $q \leq n$. Furthermore, if (p, q) and $(p, 2n - q)$ are both in $E^A(w)$ with $p > n + 1$ and $q < n$, and $r_w(p, q) = r_w(p, 2n - q) - (n - q)$, then $(p, 2n - q)$ is redundant.

Definition 3.10. Define $E(w)$ as the subset of $E^A(w)$ such that $(p, q) \in E^A(w)$ is in $E(w)$ if and only if both of the following conditions are satisfied:[‡]

- $p \geq n + 1$,
- If $q \geq n + 1$ and $(p, 2n - q)$ is also in $E^A(w)$, then $r_w(p, 2n - q) > r_w(p, q) + n - q$.

The second condition subsumes the redundancy condition for $p = n + 1$; we always will get equality instead of the desired inequality in that case.

Example 3.11. Let $w = 465\,213$ as in Example 3.9. The (type C) essential set of w is $E(w) = \{(5, 1)\}$. Suppose that $v = 321\,654$ as in Example 3.8. Then $M \in \Sigma_{v,w}$ if and only if $M \in \Sigma_v$ and the size 1 minors of $\tau_{5,1}(M)$ vanish. Thus,

$$\Sigma_{v,w} = \left\{ \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{11} & 0 & 0 & 0 & 0 & 1 \\ 0 & z_{22} & z_{23} & 0 & 1 & 0 \\ 0 & z_{23} & z_{33} & 1 & 0 & 0 \end{bmatrix} : z_{ij} \in \mathbb{K} \right\}.$$

Example 3.12. Let $w = 426\,153$. The type A essential set of w is $E^A(w) = \{(3, 2), (3, 4), (5, 2), (5, 4)\}$. The first condition that $p \geq n + 1$ eliminates $(3, 2)$ and $(3, 4)$ (whose conditions are equivalent to those given by (5, 4) and, respectively, (5, 2)). Note that $(p, q) = (5, 4)$ does not satisfy the second condition, since $q = 4 \geq n + 1$, $(p, 2n - q) = (5, 2) \in E^A(w)$, and $r_w(5, 2) = 0 = r_w(5, 4) + n - q = 1 + 3 - 4$. Hence, $E(w) = \{(5, 2)\}$.

If we let $v = 321\,654$ as in Example 3.8, we see that the condition $r_w(5, 2) = 0$ forces $z_{12} = z_{22} = z_{13} = z_{23} = 0$, and this automatically forces $r_w(5, 4) = 1$ (noting that z_{23} appears in two places in

[†] AW regrets his earlier failure in [41] in the perpetual quest to make an even number of sign errors.

[‡] Anderson in [2, Definition 1.2] and AW in [41, Section 4] choose the leftmost box in each pair, rather than the lower box as we do.

the matrix), indicating that the condition from (5,4) is redundant. In particular,

$$\Sigma_{v,w} = \left\{ \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{11} & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & z_{33} & 1 & 0 & 0 \end{bmatrix} : z_{ij} \in \mathbb{K} \right\}.$$

4 | SMALL PATCHES

Let $v_{\square} \in C_n$ denote the *square word* permutation, whose permutation matrix is

$$P(v_{\square}) = \begin{bmatrix} J_n & 0 \\ 0 & J_n \end{bmatrix}.$$

In this section, we discuss various properties of type C opposite Schubert cells Ω_v° where $v \geq v_{\square}$ in left-right weak order. We refer to such opposite Schubert cells as *small patches*.

The purpose of this section is for us to fix explicit coordinates and conventions. In addition to being crucial in our main theorem on Gröbner bases, our choice of coordinates yields a natural identification between small patches and symmetric ladders from the commutative algebra literature [15, 16].

4.1 | Small patches and symmetric matrices

To choose coordinates on type A opposite Schubert cells, it is enough to take a distinct indeterminate for each element of $D(v)$ (see [43, Section 2.2]). In this section, we put specific coordinates on type C opposite Schubert cells Ω_v° when $v \geq v_{\square}$.

By Theorem 3.1, the type C opposite Schubert cell Ω_v° of $v \in C_n$ consists of the σ -fixed points of the type A cell $\Omega_v^{A^{\circ}}$. For $v_{\square}$, the type A cell $\Omega_{v_{\square}}^{A^{\circ}}$ is isomorphic to the set of matrices

$$\Sigma_{v_{\square}}^A = \left\{ \begin{bmatrix} J_n & 0 \\ M & J_n \end{bmatrix} : M \text{ is any } n \times n \text{ matrix} \right\}.$$

Applying Corollary 3.6, we can directly compute the σ -fixed points of $\Sigma_{v_{\square}}^A$ to show the following result. (See Example 3.8 for the $n = 3$ case).

Proposition 4.1. $\Sigma_{v_{\square}} = \left\{ \begin{bmatrix} J_n & 0 \\ Z & J_n \end{bmatrix} : Z \text{ is a symmetric } n \times n \text{ matrix} \right\}.$

Our next goal is to give explicit coordinates for Σ_v whenever $v \geq v_{\square}$. This will show that, by deleting certain rows and columns with no variables, matrices in Σ_v can be identified with symmetric matrices with zeroes in prescribed entries, and our coordinates will be entries of these

matrices. For the rest of this paper, we restrict to such v . A permutation $v \in S_m$ is *123-avoiding* if there do not exist $i < j < k$ such that $v(i) < v(j) < v(k)$. A *left-to-right minimum* of a permutation is an index a such that $v(i) > v(a)$ for all $i < a$; a *right-to-left maximum* is an index b such that $v(k) < v(b)$ for all $k > b$.

Lemma 4.2. *If $v \in C_n$, then v is 123-avoiding if and only if there exist left-to-right minima $a_1 < \dots < a_n$ such that $a_i \neq 2n + 1 - a_j$ for any i and j . If we let $b_i = 2n + 1 - a_{n+1-i}$ for all i , then $b_1 < \dots < b_n$ are right-to-left maxima.*

Note that, by definition of left-to-right minima and right-to-left maxima, $v(a_1) > \dots > v(a_n)$ and $v(b_1) > \dots > v(b_n)$.

Proof. It is well known that every index in a 123-avoiding permutation $v \in S_m$ is a left-to-right minimum or a right-to-left maximum. Indeed, if j is neither, then there exists $i < j$ with $v(i) < v(j)$ and there exists $k > j$ with $v(k) > v(j)$, so v is not 123-avoiding. For $v \in C_n$, whenever a is a right-to-left maximum, $2n + 1 - a$ is a left-to-right minimum by definition. Hence, for all j with $1 \leq j \leq n$, if j is a left-to-right minimum, we let j be one of the a_i . Otherwise, $2n + 1 - j$ is a left-to-right minimum and we let $2n + 1 - j$ be one of the a_i . $\square$

Proposition 4.3. *If $v \in C_n$, then $v \geq v_{\square}$ if and only if v is 123-avoiding. Moreover, if $v = u_l v_{\square} u_r$, $\ell(v) = \ell(u_l) + \ell(v_{\square}) + \ell(u_r)$, and we set $a_i = u_r^{-1}(i)$ and $b_i = u_r^{-1}(n + i)$, then the a 's and b 's are as in Lemma 4.2.*

Proof. Supposing that $v \geq v_{\square}$, we will prove that v is 123-avoiding by induction on $\ell(v) - \ell(v_{\square})$. Notice that the statement is true for $v_{\square}$. For the inductive step, suppose $v = u_l v_{\square} u_r$ is 123-avoiding and $w \geq v \geq v_{\square}$. Then, there exists c_d such that $w = v c_d$ or $w = c_d v$. First, suppose that $w = v c_d$. Since $v c_d > v$, we must have picked d such that $v(n - d) < v(n - d + 1)$. Note $n - d$ cannot be a right-to-left maximum and $n - d + 1$ cannot be a left-to-right minimum, so $n - d = a_j$ and $n - d + 1 = b_{n+1-k}$ for some j, k (which implies that $n + d = a_k$ and $n + d + 1 = b_{n+1-j}$). Notice that we obtain the desired sequences $a'_1 < \dots < a'_n$ and $b'_1 < \dots < b'_n$ for $w = v c_d$ by taking $a'_i = a_i$ and $b'_i = b_i$ for all i , except that $a'_j = n - d + 1$, $b'_{n+1-k} = n - d$, $a'_k = n + d + 1$, and $b'_{n+1-j} = n + d$. We conclude that $w = v c_d$ is 123-avoiding. Furthermore, $w = u_l v_{\square} u'_r$ where $u'_r = u_r c_d$, and we have $a'_i = a_i = u_r^{-1}(i) = (u'_r)^{-1}(i)$ for $i \neq j, k$, while $a'_j = n - d + 1 = c_d(a_j) = c_d(u_r^{-1}(j)) = (u'_r)^{-1}(j)$, and similarly, $a'_k = (u'_r)^{-1}(k)$.

Continuing the inductive step, suppose that $w = c_d v$. Since $c_d v > v$, we must have picked d such that $v^{-1}(n - d) < v^{-1}(n - d + 1)$. Since $v(v^{-1}(n - d)) < v(v^{-1}(n - d + 1))$, then $v^{-1}(n - d)$ is not a right-to-left maximum and $v^{-1}(n - d + 1)$ is not a left-to-right minimum, so $v^{-1}(n - d) = a_j$ and $v^{-1}(n - d + 1) = b_{n+1-k}$ for some j, k (which implies that $v(a_k) = n + d$ and $v(b_{n+1-j}) = n + d + 1$). Observe then that for all i , $c_d v(a_i) > c_d v(a_{i+1})$ and $c_d v(b_i) > c_d v(b_{i+1})$. In this case, we obtain the desired sequences for $w = c_d v$ by keeping the a 's and b 's for v . Note that u_r is unchanged. This proves the second statement and the forward direction of the first statement.

Now suppose that $v \in C_n$ is 123-avoiding. Choose a 's and b 's for v , as in Lemma 4.2. We will provide an algorithm that produces $\alpha_1, \dots, \alpha_s$ and $\beta_1, \dots, \beta_t$ such that $v_{\square} = c_{\alpha_s} \dots c_{\alpha_1} v c_{\beta_1} \dots c_{\beta_t}$ and $\ell(v) = \ell(v_{\square}) + s + t$, thus proving that $v \geq v_{\square}$.




FIGURE 1 On the left, we have the diagram of $v = 462513 > v_{\square}$ and on the right its associated skew partition.

We start by finding the α 's. If $v(a_1) = n$, we will have no c_{α} 's. Otherwise, since $v(a_1) > \dots > v(a_n) \geq 1$, we must have $v(a_1) > n$. Let $j = \max\{k \in [n] : v(a_k) > n + 1 - k\}$, and let $c \in C_n$ be the generator that transposes $v(a_j) - 1$ with $v(a_j)$ (so, it transposes $v(2n + 1 - a_j)$ with $v(2n + 1 - a_j) + 1$). Since a_j is a left-to-right minimum, $v^{-1}(v(a_j) + 1) > a_j$, so $cv < v$. By the construction of j , we have $v(a_j) - 1 \neq v(a_{j+1})$, and hence, $v^{-1}(v(a_j) - 1)$ is not a left-to-right minimum for v . Hence, cv is also 123-avoiding as witnessed by the same indices a_i and b_i . Set α_1 so that $c = c_{\alpha_1}$. Iterate this process, without changing the a_i and b_i , until $v(a_k) = n + 1 - k$ for all $k \in [n]$.

Starting with the output v of the previous paragraph, we now find the β 's. Note that this output has the property that $v(a_i) = n + 1 - i$ (and $v(b_i) = 2n + 1 - i$) for all $i \in [n]$, and this property will be maintained throughout the process. Let $j = \min\{k \in [n] : a_k > k\}$, and let $c \in C_n$ be the generator that transposes a_j with $a_j - 1$ (so, it must also transpose b_{n+1-j} with $b_{n+1-j} + 1$). First, note that since a_j is a left-to-right minimum, $v(a_j - 1) > v(a_j)$, so $vc < v$. Second, note that $a_j - 1$ is not a left-to-right minimum by definition of j , so $a_j - 1 = b_k$ for some $k \in [n]$ and $v(a_j - 1) > n$. Take $a'_i = a_i$ and $b'_i = b_i$ for all i , except that $a'_j = b_k$, $b'_k = a_j$, $a'_{n+1-k} = b_{n+1-j}$, and $b'_{n+1-j} = a_{n+1-k}$. Then, vc is 123-avoiding and $vc(a'_j) = n + 1 - j$ (and $vc(b'_{n+1-j}) = 2n + 1 - j$). Set β_1 to be the index such that $c = c_{\beta_1}$. Iterate the process starting with $vc_{\beta_1} \dots c_{\beta_1}$ until no longer possible, so until $a_j = j$ for all $j \in [n]$. Note then that the algorithm terminates with $v_{\square}$ and the proposition follows. $\square$

One can consider Proposition 4.3 as a type C analog of [7, Theorem 2.1], which characterizes 321-avoiding permutations as those smaller (in left-right weak order) than the maximal grassmannian permutation for some descent. Since 321-avoiding type C permutations are 321-avoiding type A permutations, [7, Theorem 2.1] directly implies the “only if” direction of the first sentence of Proposition 4.3, but in type A, the choice of maximal grassmannian permutation can depend on v , whereas in type C, $v_{\square}$ is the only choice. Also unlike type A, the conditions of Proposition 4.3 are not equivalent to $w_0 v$ being fully commutative: $v = 1324 \in C_2$ does not satisfy the proposition although $w_0 v = 4231 = c_1 c_0 c_1$ is fully commutative.

The following corollary describes the diagram for $v \geq v_{\square}$; see Figure 1 for an example.

Corollary 4.4. *If $v \in C_n$, then $v \geq v_{\square}$ if and only if $D(v)$ becomes a skew partition after deleting all the rows and columns that do not contain boxes of $D(v)$.*

Proof. This is a direct consequence of the previous proposition together with the note following [7, Theorem 2.1]. We remark that we obtain 123-avoiding permutations instead of 321-avoiding ones due to the difference in our conventions for $D(v)$. $\square$

Throughout the paper, we denote by $\bar{v}$ a factorization $v = u_l v_{\square} u_r$ such that $\ell(v) = \ell(u_l) + \ell(v_{\square}) + \ell(u_r)$. We let

$$R_{\bar{v}} = \mathbb{K}[z_{ij} : i \leq j, u_r^{-1}(i) < u_r^{-1}(2n+1-j), u_l(n+1-i) < u_l(n+j)].$$

Furthermore, let $M_{\bar{v}}$ be the matrix with z_{ij} as the entries at $(u_l(n+j), u_r^{-1}(i))$ and $(u_l(n+i), u_r^{-1}(j))$ whenever $u_r^{-1}(i) < u_r^{-1}(2n+1-j)$ and $u_l(n+1-i) < u_l(n+j)$, 1s at $(v(i), i)$ for all i , and 0s at all other positions. Note that the z_{ij} only appear within $D(v)$. For some examples, the general element of Σ_{321654} in Example 3.8 and the matrix on the left-hand side of the equality in Example 4.8 are both of the form $M_{\bar{v}}$.

Note that given $v \geq v_{\square}$, a choice of $a_1, \dots, a_n$ and $b_1, \dots, b_n$ as in Lemma 4.2 is equivalent to choosing a factorization $\bar{v}$. In this language

$$R_{\bar{v}} = \mathbb{K}[z_{ij} : i \leq j, a_i < b_{n+1-j}, v(a_i) < v(b_{n+1-j})],$$

and $M_{\bar{v}}$ is the matrix with z_{ij} as the entries at $(v(b_{n+1-j}), a_i)$ and $(v(b_{n+1-i}), a_j)$ whenever $a_i < b_{n+1-j}$ and $v(a_i) < v(b_{n+1-j})$, 1s at $(v(i), i)$ for all i , and 0s at all other positions.

Example 4.5. This example shows how the labeling of the coordinates in $R_{\bar{v}}$ depends on the choice of our factorization $\bar{v}$. The factorization $\bar{v}^{(1)}$ corresponding to $v = 642\,531 > v_{\square}$, $(a_{\bullet}) = (1, 2, 3)$, and $(b_{\bullet}) = (4, 5, 6)$ is $\bar{v}^{(1)} = u_l^{(1)} v_{\square} u_r^{(1)}$ where $u_l^{(1)} = 246\,135$ and $u_r^{(1)} = 123\,456$. Then,

$$M_{\bar{v}^{(1)}} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & z_{23} & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & z_{23} & z_{33} & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

By comparison, for $(a_{\bullet}) = (2, 3, 6)$ and $(b_{\bullet}) = (1, 4, 5)$, we have that $\bar{v}^{(2)} = u_l^{(2)} v_{\square} u_r^{(2)}$ where $u_l^{(2)} = 124\,356$ and $u_r^{(2)} = 412\,563$. Then

$$M_{\bar{v}^{(2)}} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & z_{12} & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & z_{12} & z_{22} & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

One can relate the variables to the self-conjugate skew partition associated to v by Corollary 4.4 as follows. If the skew partition has n rows (or equivalently n columns), then there is only one choice for $\bar{v}$, and z_{ij} is a variable if and only if (i, j) (equivalently (j, i)) is a box of the skew partition. If the skew partition has fewer rows, then different choices of $\bar{v}$ will give rise to different (but always self-conjugate) placements of the self-conjugate skew partition in an $n \times n$ box corresponding to different coordinates.

The matrix $M_{\bar{v}}$ satisfies the following property.

Proposition 4.6. *If $1 \leq \delta \leq 2n$ is a left-to-right minimum of v , then the only nonzero entry of $M_{\bar{v}}$ in row $v(\delta)$ is the 1 at position $(v(\delta), \delta)$. Similarly, if $1 \leq \delta \leq 2n$ is a right-to-left maximum of v , then the only nonzero entry of $M_{\bar{v}}$ in column δ is the 1 at position $(v(\delta), \delta)$.*

Proof. Suppose that $1 \leq \delta \leq 2n$ is a left-to-right minimum of v . Entries to the right of this position lie on its hook, whereas an entry (ϵ, δ) to its left lies on the hook extending from $(\epsilon, v^{-1}(\epsilon))$ by the left-to-right minimum condition. An analogous argument proves the second part of the lemma. $\square$

We now give concrete coordinates for the coordinate ring of Σ_v .

Proposition 4.7. *If $v \geq v_{\square}$, then $R_{\bar{v}}$ is a coordinate ring of Σ_v and $M_{\bar{v}}$ is the generic matrix in Σ_v . (In other words, a matrix is in Σ_v if and only if it can be obtained by setting each variable in $M_{\bar{v}}$ to some element of $\mathbb{K}$.) Furthermore, if $v = u_l v_{\square} u_r$ and $\ell(v) = \ell(u_l) + \ell(v_{\square}) + \ell(u_r)$, then the rule $M \mapsto P(u_l^{-1})MP(u_r^{-1})$ induces the injective map from Σ_v to $\Sigma_{v_{\square}}$ which identifies the entry named z_{ij} in $M_{\bar{v}}$ with the entry named z_{ij} in $M_{\bar{v}_{\square}}$.*

Notice that $v \not\geq v_{\square}$ in Example 3.7, and indeed, the entries of the general matrix in Σ_v in that example cannot all be made to be variables. An interesting question is to describe the entries of Σ_v for general $v \in C_n$ and give a Gröbner basis for Kazhdan–Lusztig varieties arising from these cells.

Before proving this result, let us give an example and some necessary lemmas.

Example 4.8. Let $v = 462513$. Then $c_0 v_{\square} c_0 c_1 = v$. The following equality illustrates the “furthermore” part of Proposition 4.7:

$$P(c_0) \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & z_{12} & 0 & z_{13} & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{12} & 0 & z_{22} & 1 & 0 & 0 \\ z_{13} & 1 & 0 & 0 & 0 & 0 \end{bmatrix} P(c_1 c_0) = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & z_{12} & z_{13} & 0 & 0 & 1 \\ z_{12} & z_{22} & 0 & 0 & 1 & 0 \\ z_{13} & 0 & 0 & 1 & 0 & 0 \end{bmatrix}.$$

In general, the effect of right multiplication by $P(u_r^{-1})$ is to collect at the left side all columns containing any variable z_{ij} , and similarly, left multiplication by $P(u_l^{-1})$ collects rows with variables at the bottom.

The following lemmas will be used to prove Lemma 6.2. They are adaptations of [43, Lemma 6.5] to type C.

Lemma 4.9. *Let $v \in C_n$ and k be such that $vc_k \succ v$ in right weak order. The diagram $D(vc_k)$ is obtained from $D(v)$ as follows: $D(vc_k)$ agrees with $D(v)$ except in columns $n \pm k$ and $n \pm k + 1$. To obtain columns $n - k$ and $n - k + 1$ of $D(vc_k)$, move all the boxes of $D(v)$ in column $n - k$ strictly above row $v(n - k + 1)$ one unit to the right and delete the box in position $(v(n - k + 1), n - k)$. Repeat the analogous process in columns $n + k$ and $n + k + 1$ (if $k \neq 0$).*

We need the analogous lemma for left weak order as well:

Lemma 4.10. *Let $v \in C_n$ and k be such that $c_k v \succ v$ in left weak order. The diagram $D(c_k v)$ is obtained from $D(v)$ as follows: $D(c_k v)$ agrees with $D(v)$ except in rows $n \pm k$ and $n \pm k + 1$. To obtain rows $n - k$ and $n - k + 1$ of $D(c_k v)$, move all the boxes of $D(v)$ in row $n + 1 - k$ strictly right of column $v^{-1}(n - k)$ one unit up and delete the box in position $(n + 1 - k, v^{-1}(n - k))$. Repeat the analogous process in rows $n + k$ and $n + k + 1$ (if $k \neq 0$).*

Proof of Proposition 4.7. To prove that $M_{\bar{v}}$ is the generic matrix in Σ_v , it suffices to show that $\sigma(M_{\bar{v}}) = M_{\bar{v}}$ and the entries of $M_{\bar{v}}$ are as in the beginning of Section 3.3, implying that, regardless how we evaluate the z_{ij} in $M_{\bar{v}}$, we get a matrix in Σ_v^A . The second statement follows by definition, since $(M_{\bar{v}})_{ij} = 1$ whenever $P(v)_{ij} = 1$ and $(M_{\bar{v}})_{ij} = 0$ whenever $(i, j) \notin D(v)$. For the first statement, we proceed by induction on $\ell(v)$. The base case, when $v = v_{\square}$, is trivial. For the inductive case, consider some $v > v_{\square}$, and let c_k be a simple reflection such that either $vc_k \succ v$ or $c_k v \succ v$. By the inductive hypothesis, $M_{\bar{v}}$ is the generic matrix in Σ_v . Throughout the proof, we will fix a factorization $\bar{v} = u_l v_{\square} u_r$ with corresponding sequences $(a_{\cdot})$ and $(b_{\cdot})$.

First suppose that $vc_k \succ v$ so that $v(n - k) < v(n - k + 1)$ and $v(n + k) < v(n + k + 1)$. Let $\overline{vc_k}$ be the factorization $(u_l)v_{\square}(u_r c_k)$. Since v is 123-avoiding, $n \pm k$ are left-to-right minima, and $n \pm k + 1$ are right-to-left maxima. Therefore, $(v(n - k + 1), n - k) = (v(b_{n+1-j}), a_i)$ and $(v(n + k + 1), n + k) = (v(b_{n+1-i}), a_j)$ for some i, j , and the corresponding entries in $(M_{\bar{v}})$ are the variable z_{ij} or z_{ji} . Without loss of generality, we assume that $i \leq j$. Our goal is to show that $M_{\overline{vc_k}}P(c_k)$ is obtained from $M_{\bar{v}}$ by setting $z_{ij} = 0$. Right multiplication by $P(c_k)$ swaps column $n + k$ with $n + k + 1$ and column $n - k$ with $n - k + 1$. Since $P(vc_k)P(c_k) = P(v)$, then the positions of the 1s in $M_{\overline{vc_k}}P(c_k)$ and $M_{\bar{v}}$ agree. By Lemma 4.9,

$$D(vc_k) \setminus [n] \times \{n \pm k, n \pm k + 1\} = D(v) \setminus [n] \times \{n \pm k, n \pm k + 1\},$$

and therefore, the positions of 0 entries in $M_{\overline{vc_k}}P(c_k)$ and $M_{\bar{v}}$ agree on all columns, except possibly columns $n \pm k, n \pm k + 1$. This lemma also implies that, for $\delta \in \{n \pm k, n \pm k + 1\}$, if $(M_{\bar{v}})_{\epsilon\delta} = 0$, then $(M_{\overline{vc_k}}P(c_k))_{\epsilon\delta} = 0$.

It remains to analyze the variable entries of $M_{\bar{v}}$. Since $n \pm k + 1$ are right-to-left maxima for v , columns $n \pm k + 1$ of $M_{\bar{v}}$ do not contain any variables. Similarly, since $n \pm k$ are right-to-left maxima for vc_k , columns $n \pm k + 1$ of $M_{\bar{v}}P(c_k)$ do not contain any variables and are therefore equal to columns $n \pm k + 1$ of $M_{\bar{v}}$. Note that the sequences $(a'_{\cdot})$ and $(b'_{\cdot})$ from Proposition 4.3 for $\overline{vc_k}$ agree with the sequences $(a_{\cdot})$ and $(b_{\cdot})$ everywhere except

$$\begin{aligned} a'_i &= b_{n+1-j} = n - k + 1, & b'_{n+1-j} &= a_i = n - k, & a'_j &= b_{n+1-i} = n + k + 1, \text{ and} \\ b'_{n+1-i} &= a_j = n + k. \end{aligned}$$

If $\delta \neq n \pm k, n + 1 \pm k$ and $(M_{\bar{v}})_{\epsilon\delta} = z_{i'j'}$, then $(\epsilon, \delta) \in D(v)$ and $(\epsilon, \delta) \in \{(v(b_{n+1-j'}), a_{i'}), (v(b_{n+1-i'}), a_{j'})\}$. Combining $(\epsilon, \delta) \in D(vc_k)$, which follows from Lemma 4.9, with $vc_k(b'_m) = v(b_m)$ for all m , we have that $(\epsilon, \delta) \in \{(vc_k(b'_{n+1-j'}), a'_{i'}), (vc_k(b'_{n+1-i'}), a'_{j'})\}$. It follows that $(M_{\overline{vc_k}})_{\epsilon\delta} = z_{i'j'}$. Finally, assume $\delta = n \pm k$ and $(M_{\bar{v}})_{\epsilon\delta} = z_{i'j'}$ so that $(\epsilon, \delta) \in D(v)$ and by Lemma 4.9 $(\epsilon, \delta + 1) \in D(vc_k)$, except if $\epsilon = v(\delta)$. If $\epsilon = v(\delta)$, then $z_{i'j'} = z_{ij}$ and we obtain $(M_{\overline{vc_k}}P(c_k))_{\epsilon\delta}$ by setting $z_{ij} = 0$, as desired. If $\epsilon \neq v(\delta)$, then $\epsilon \in \{v(b_{n+1-j'}), v(b_{n+1-i'})\} = \{vc_k(b'_{n+1-j'}), vc_k(b'_{n+1-i'})\}$, and therefore, $(M_{\overline{vc_k}})_{\epsilon\delta} = z_{i'j'}$. We conclude that $M_{\overline{vc_k}}P(c_k)$ is obtained from $M_{\bar{v}}$ by setting $z_{ij} = 0$ if it lies in positions $(v(n \pm k + 1), n \pm k)$.

We are left with proving that $\sigma(M_{\overline{v c_k}}) = M_{\overline{v c_k}}$. However, by induction, $\sigma(M_{\overline{v}}) = M_{\overline{v}}$, and therefore, $\sigma(M_{\overline{v c_k}} P(c_k)) = M_{\overline{v c_k}} P(c_k)$ by the argument above. Hence, since σ is a group homomorphism and $\sigma(P(c_k)) = P(c_k)$, we conclude $\sigma(M_{\overline{v c_k}}) = M_{\overline{v c_k}}$. Finally, returning to the case $c_k v \succ v$, using Lemma 4.10 and other similar arguments, one can show that in this case, $P(c_k) M_{\overline{c_k v}}$ is obtained from $M_{\overline{v}}$ by setting $z_{ij} = 0$ if it lies in positions $(n \pm k + 1, v^{-1}(n \pm k))$, and $\sigma(M_{\overline{c_k v}}) = M_{\overline{c_k v}}$. $\square$

Given $v \in C_n$, $v \geq v_{\square}$, let $V_{\overline{v}}$ be the set of variables of $R_{\overline{v}}$, that is, the set of variables that appear as entries of $M_{\overline{v}}$.

Corollary 4.11. *Fix a factorization $\overline{v} = u_l v_{\square} u_r$. If c_k is an ascent of v and we set $\overline{v c_k} = (u_l) v_{\square} (u_r c_k)$, then $V_{\overline{v c_k}} \subseteq V_{\overline{v}}$ and $V_{\overline{v}} \setminus V_{\overline{v c_k}} = \{z_{ij}\}$, where z_{ij} is the entry of $M_{\overline{v}}$ in positions $(v(n \pm k + 1), n \pm k)$.*

Proof. This follows from the inductive step in Proposition 4.7. $\square$

4.2 | Equations for type C Kazhdan–Lusztig varieties

Let $R_{\overline{v}}$ and R_v^A denote, respectively, the coordinate rings $\mathbb{k}[\Sigma_v]$ and $\mathbb{k}[\Sigma_v^A]$. Similarly, let $M_{\overline{v}}$ and M_v^A denote, respectively, the generic matrices in $\mathbb{k}[\Sigma_v]$ and $\mathbb{k}[\Sigma_v^A]$. Let $I_{\overline{v}, w}$ be the ideal of $R_{\overline{v}}$ generated by the size $r_w(p, q) + 1$ minors of $\tau_{p, q}(M_{\overline{v}})$ over all (p, q) in $E(w)$. We call $I_{\overline{v}, w}$ a *Kazhdan–Lusztig ideal*.

Proposition 4.12. *We have $\Sigma_{v, w} = \text{Spec}(R_{\overline{v}}/I_{\overline{v}, w})$.*

Theorem 4.15 will give an independent proof of this proposition in the case $v \geq v_{\square}$.

Proof. The discussion of rank conditions in Section 3.4 proves equality as sets. Equality as schemes follows from [42, Proposition 3.1], which gives the analogous scheme-theoretic equality in type A, and Theorem 3.1 along with Equation (2) (which follows from [33, Proposition 6.1.1.2]). $\square$

Note that, for type B, the analogs of Proposition 4.12 and the statements from [33] it depends on are true only set-theoretically, not necessarily scheme-theoretically, so our work cannot be extended to that case. See Example 8.3 for an illustration of the potential difficulties in type B.

We now define the term orders we use in this paper. A *diagonal* term order on $R_{\overline{v}}$ is one where, given any minor in $M_{\overline{v}}$ where the diagonal term is nonzero, the diagonal term is the leading term. In notation, this means that, if $\epsilon_1 < \dots < \epsilon_r$, $\delta_1 < \dots < \delta_r$, D is the minor of $M_{\overline{v}}$ using rows $\{\epsilon_1, \dots, \epsilon_r\}$ and columns $\{\delta_1, \dots, \delta_r\}$, and $\prod_{i=1}^r (M_{\overline{v}})_{\epsilon_i \delta_i}$ is nonzero, then this product is the leading term of D . Note that there can be multiple distinct diagonal term orders. However, even if the diagonal term is zero for a given minor, there are restrictions on which term can be its leading term under a diagonal term order, since the condition applies to every subminor of the minor in question. As we are taking southwest minors rather than northwest minors in defining $I_{\overline{v}, w}$, our diagonal term orders are equivalent to the antidiagonal term orders of [30].

We note that our diagonal term orders on different sets of variables are compatible with each other.

Proposition 4.13. *Let c_k be an ascent of v , let $<$ be a diagonal term order on $R_{\bar{v}}$, and let $<' = <|_{R_{\bar{v}c_k}}$ be the restriction of $<$ to $R_{\bar{v}c_k}$. Then, $<'$ is a diagonal term order on $R_{\bar{v}c_k}$.*

Proof. Suppose that D' is a minor of $M_{\bar{v}c_k}$ with nonzero diagonal term using rows $\epsilon_1 < \dots < \epsilon_s$ and columns $\delta_1 < \dots < \delta_s$. Let D be the minor of $M_{\bar{v}}$ using rows $\epsilon_1 < \dots < \epsilon_s$ and columns $c_k(\delta_1) < \dots < c_k(\delta_s)$. If D' involves only one of the columns $n - k$ and $n - k + 1$, and D' involves only one of the columns $n + k$ and $n + k + 1$, then the diagonal term of D in $M_{\bar{v}}$ is the same as the diagonal term of D' in $M_{\bar{v}c_k}$, as the rows and columns are ordered in the same way. Note that the variable z_{ij} in $V_{\bar{v}} \setminus V_{\bar{v}c_k}$ cannot appear in the diagonal term of D , as that would imply the diagonal term of D is zero. Hence, the leading term of D under $<'$ must be the diagonal term, as the leading term of D' under $<$ is the diagonal term.

Otherwise, if D' involves both columns $n - k$ and $n - k + 1$ in $M_{\bar{v}}$, then, since c_k is an ascent of v , there is a right-to-left maximum in column $n - k + 1$ of $M_{\bar{v}}$ and column $n - k$ of $M_{\bar{v}c_k}$. So, by Proposition 4.6, $D' = \pm \widetilde{D'}$ where $\widetilde{D'}$ is the minor formed by removing column $n - k$ and row $vc_k(n - k)$ from D' . Similarly, $D = \pm \widetilde{D}$ where $\widetilde{D}$ is the minor formed by removing column $n - k + 1$ and row $v(n - k + 1)$ from D . Now the argument in the previous paragraph applies to $\widetilde{D}$ and $\widetilde{D'}$. A similar argument applies if D' involves both columns $n + k$ and $n + k + 1$. $\square$

We show that there is at least one diagonal term order, namely, the lexicographic term order $<_{\text{lex}}$ where $z_{ij} > z_{i'j'}$ if and only if either $i > i'$, or $i = i'$ and $j > j'$. One can see from the next section that $<_{\text{lex}}$ is the term order used in [27], made explicit for this case.

Proposition 4.14. *The term order $<_{\text{lex}}$ is a diagonal term order.*

Proof. We prove this by downward induction in length. The base case is where $v = w_0$, where $R_{\bar{v}}$ has no variables, and hence, the statement is vacuously true. Let c_k be the last ascent of v . By Corollary 4.11, $V_{\bar{v}} \setminus V_{\bar{v}c_k} = \{z_{ij}\}$, where z_{ij} is the entry of $M_{\bar{v}}$ in positions $(v(n \pm k + 1), n \pm k)$. Moreover, z_{ij} must appear as the south-most nonzero entry in its column, and the 1 appearing immediately to its right is a right-to-left maximum. Hence, there are no variables southeast of (either appearance, if there are two, of) z_{ij} in $M_{\bar{v}}$. It follows that z_{ij} is the largest variable in $R_{\bar{v}}$ under $<_{\text{lex}}$.

By induction, $<_{\text{lex}}$ restricted to $R_{\bar{v}c_k}$ is a diagonal term order. By Proposition 4.13, it suffices to show that if z_{ij} appears in a minor of $M_{\bar{v}}$ with a nonzero diagonal term, then it must appear in the diagonal term. Since there are no variables southeast of z_{ij} in $M_{\bar{v}}$, any minor of $M_{\bar{v}}$ with nonzero diagonal term and such that z_{ij} does not appear on the diagonal term must have as its southeast entry the 1 directly to the right of z_{ij} . By Proposition 4.6, the only nonzero entry of the rightmost column of the minor is this 1. It follows that z_{ij} does not appear in any term of the minor. $\square$

We now state our main theorem.

Theorem 4.15. *Given $v \geq v_{\square}$, the size $r_w(p, q) + 1$ minors of $\tau_{p,q}(M_{\bar{v}})$ over all (p, q) in $E(w)$ form a Gröbner basis for $I_{\bar{v},w}$ with respect to any diagonal term order.*

The proof appears in Section 6. The main technique is to show that K -polynomials of subword complexes, suitably weighted, satisfy the Kostant–Kumar recursion. (This technique follows [27].)

Examples 8.1 and 8.2 show that Theorem 4.15 can fail when the $v \geq v_{\square}$ hypothesis is dropped.

4.3 | Torus action of type C Kazhdan–Lusztig varieties and the weights for $v \geq v_{\square}$

Let T be the torus consisting of the diagonal matrices in $Sp_{2n}(\mathbb{K})$. Since any $(t_{ij}) \in T$ is fixed under σ , it satisfies $t_{ii} = t_{2n+1-i, 2n+1-i}^{-1}$ for all i . The torus T acts on Ω_v° by left multiplication, that is, given matrices $M \in Sp_{2n}(\mathbb{K})$ and $N \in T$,

$$N \cdot (MB_G^+/B_G^+) := (NM)B_G^+/B_G^+.$$

This action induces the following torus action on Σ_v : for $M \in \Sigma_v$ and $N \in T$, $N \cdot_v M$ is the matrix in Σ_v representing $(NM)B_G^+/B_G^+$. Let us describe the action more concretely. Notice that in general $NM \notin \Sigma_v$ because the entry of NM in position $(\pi(j), j)$ need not equal 1. Thus, to obtain an element of Σ_v , we need to multiply on the right by the appropriate element of T to make these entries 1.

It will be most convenient for us to denote by $(x_1, \dots, x_n)$ the element of T where

$$(x_1, \dots, x_n) := \text{diag}(x_n, \dots, x_1, x_1^{-1}, \dots, x_n^{-1}), \quad (6)$$

the diagonal matrix with diagonal entries $x_n, \dots, x_1, x_1^{-1}, \dots, x_n^{-1}$ from northeast to southwest.

Example 4.16. We describe the action of T on $\Sigma_{v_{\square}}$ for $v_{\square} = 321\,654$:

$$\begin{aligned} (x_1, x_2, x_3) \cdot_v \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{11} & z_{12} & z_{13} & 0 & 0 & 1 \\ z_{12} & z_{22} & z_{23} & 0 & 1 & 0 \\ z_{13} & z_{23} & z_{33} & 1 & 0 & 0 \end{bmatrix} &= (x_1, x_2, x_3) \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{11} & z_{12} & z_{13} & 0 & 0 & 1 \\ z_{12} & z_{22} & z_{23} & 0 & 1 & 0 \\ z_{13} & z_{23} & z_{33} & 1 & 0 & 0 \end{bmatrix} (x_3^{-1}, x_2^{-1}, x_1^{-1}) \\ &= \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ x_1^{-2}z_{11} & x_1^{-1}x_2^{-1}z_{12} & x_1^{-1}x_3^{-1}z_{13} & 0 & 0 & 1 \\ x_1^{-1}x_2^{-1}z_{12} & x_2^{-2}z_{22} & x_2^{-1}x_3^{-1}z_{23} & 0 & 1 & 0 \\ x_1^{-1}x_3^{-1}z_{13} & x_2^{-1}x_3^{-1}z_{23} & x_3^{-2}z_{33} & 1 & 0 & 0 \end{bmatrix}. \end{aligned}$$

Since T acts by left multiplication on each Schubert variety and on each opposite Schubert cell, the torus T also acts by left multiplication on each Kazhdan–Lusztig variety $\mathcal{N}_{v,w}$.

Let us now restrict to $v \geq v_{\square}$ and explicitly compute the weights on the coordinates z_{ij} of the action. We adopt the convention that the weight e_i denotes the homomorphism in $\text{Hom}(T, \mathbb{K})$ that sends the element $(x_1, \dots, x_n)$ to x_i . We will write weights additively. In addition, we will let $t_i = \exp(e_i)$ denote the formal exponential of the weight e_i , so that the group operation on the t_i (and monomials therein) is multiplication.

Lemma 4.17. *The coordinate function z_{ij} on $M_{\bar{v}_{\square}}$ has weight $e_i + e_j$.*

Proof. As one can see from Example 4.16, acting on $M \in \Sigma_{v_{\square}}$ by $(x_1, \dots, x_n)$ multiplies the entry $z_{ij}(M)$ by $x_i^{-1}x_j^{-1}$. Hence, the weight of the action on z_{ij} , the coordinate function on this entry, is $e_i + e_j$. $\square$

We next see that the analog of Lemma 4.17 holds for any $v \geq v_{\square}$. To do so, we need to consider the action of C_n on the weights induced from permuting the diagonal entries of $(x_1, \dots, x_n)$. Notice that given $u \in C_n$, the induced action is so that for $i \leq n$,

$$u \cdot e_i = \begin{cases} -e_{n+1-u(n+i)} & \text{if } u(n+i) \leq n, \\ e_{u(n+i)-n} & \text{if } u(n+i) \geq n+1. \end{cases}$$

Proposition 4.18. *Let $v \geq v_{\square}$ with $\bar{v} = u_l v_{\square} u_r$. The weight of a coordinate function of $M_{\bar{v}}$ depends only on its position (and not on $\bar{v}$). Furthermore, the coordinate function z_{ij} on $M_{\bar{v}}$ has weight $u_l \cdot e_i + u_l \cdot e_j$.*

Note that for $\bar{v} = u_l v_{\square} u_r$ and $i \leq n$, in the notation of Lemma 4.2, we have $u_l(i) = v(a_{n+1-i})$ and $u_l(n+i) = v(b_{n+1-i})$.

Proof. Define $y_{n+i} := x_i^{-1}$ and $y_{n+1-i} := x_i$, $1 \leq i \leq n$, so that $\text{diag}(y_1, \dots, y_{2n}) = (x_1, \dots, x_n)$. Then, the action of T on Σ_v , $v \in C_n$, is given by

$$\mathbf{y} \cdot_v M = \text{diag}(y_1, \dots, y_{2n}) M \text{diag}(y_{v(1)}^{-1}, \dots, y_{v(2n)}^{-1}), \quad \mathbf{y} \in T, M \in \Sigma_v.$$

Therefore, the weight for the coordinate function in position (ϵ, δ) is the weight corresponding to $y_{\epsilon} y_{\delta}^{-1}$, which depends only on (ϵ, δ) and not on $\bar{v}$.

Suppose that $v \geq v_{\square}$ and $\bar{v} = u_l v_{\square} u_r$. Let $M \in \Sigma_v$ and $\mathbf{y} \in T$. Given $i \leq j$, the variable z_{ij} appears as entries $(v(b_{n+1-j}), a_i)$ and $(v(b_{n+1-i}), a_j)$ of M . The entries in positions $(v(b_{n+1-j}), a_i)$ and $(v(b_{n+1-i}), a_j)$ of $\mathbf{y} \cdot_v M$ are

$$y_{v(b_{n+1-j})} y_{v(a_i)}^{-1} z_{ij} = y_{u_l(n+j)} y_{u_l(n+1-i)}^{-1} z_{ij} \quad \text{and} \quad y_{v(b_{n+1-i})} y_{v(a_j)}^{-1} z_{ij} = y_{u_l(n+i)} y_{u_l(n+1-j)}^{-1} z_{ij},$$

respectively. By definition of $\mathbf{y}$ and the fact $u_l \in C_n$,

$$y_{u_l(n+j)} y_{u_l(n+1-i)}^{-1} = y_{u_l(n+j)} y_{u_l(n+i)} = y_{u_l(n+i)} y_{u_l(n+1-j)}^{-1}.$$

Again by definition of $\mathbf{y}$, we conclude that the weight of z_{ij} is $u_l \cdot e_i + u_l \cdot e_j$. $\square$

Example 4.19. Let $v = 642531$ and $\bar{v}$ be the factorization associated to $(a_{\bullet}) = (1, 2, 3)$, and $(b_{\bullet}) = (4, 5, 6)$. We describe the weights of the coordinate functions of $M_{\bar{v}}$ via an explicit computation:

$$\begin{aligned}
(x_1, x_2, x_3) \cdot \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & z_{23} & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & z_{23} & z_{33} & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} &= (x_1, x_2, x_3) \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & z_{23} & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & z_{23} & z_{33} & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} (x_2^{-1}, x_1, x_3) \\
&= \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & x_3 \\ 0 & 0 & x_2 & 0 & 0 & 0 \\ 0 & 0 & x_1 z_{23} & 0 & x_1 & 0 \\ 0 & x_1^{-1} & 0 & 0 & 0 & 0 \\ 0 & x_2^{-1} z_{23} & x_2^{-1} z_{33} & x_2^{-1} & 0 & 0 \\ x_3^{-1} & 0 & 0 & 0 & 0 & 0 \end{bmatrix} (x_2^{-1}, x_1, x_3) \\
&= \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & x_2^{-1} x_1 z_{23} & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & x_1 x_2^{-1} z_{23} & x_2^{-1} x_2^{-1} z_{33} & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.
\end{aligned}$$

Thus, the weight of z_{23} is $-e_1 + e_2$ and the weight of z_{33} is $-e_2 - e_2$. We verify that for z_{23} , this agrees with Proposition 4.18. Since $v(b_{n+1-2}) \leq n$ and $v(b_{n+1-3}) \geq n+1$, we have $u_l \cdot e_2 = -e_1$ and $u_l \cdot e_3 = e_2$. One can verify that if we now take $\bar{v}$ to be the factorization associated to $(a_*) = (2, 3, 6)$ and $(b_*) = (1, 4, 5)$ (as in the second part of Example 4.5), the weight of z_{12} is $-e_1 + e_2$ and the weight of z_{22} is $-e_2 - e_2$.

We end by noting that this multigrading is positive, so that the only elements in $R_{\bar{v}}$ which have degree $\mathbf{0}$ are the constants.

Corollary 4.20. *Let $v \geq v_{\square}$ with $\bar{v} = u_l v_{\square} u_r$. The multigrading on $R_{\bar{v}}$ that assigns degree $u_l \cdot e_i + u_l \cdot e_j$ to coordinate function z_{ij} is a positive multigrading.*

Proof. The set of all vectors $e_i + e_j$ generates a pointed cone, so the images of these vectors under the action of a fixed u_l do also. $\square$

5 | TYPE C SUBWORD COMPLEXES AND VERTEX DECOMPOSITION

5.1 | Subword complexes

In [29, 30], A. Knutson and E. Miller defined a family of simplicial complexes, called subword complexes, for arbitrary Coxeter groups. Let $Q = (\alpha_1, \dots, \alpha_\ell)$ be a reduced word for $v \in C_n$, as defined in Section 3.2. The *subword complex* $S(Q, w)$ associated to Q and $w \in C_n$ is the

simplicial complex on the vertex set $[\ell] = \{1, \dots, \ell\}$ whose maximal faces are the sets $[\ell] \setminus \{i_1, \dots, i_k\}$ such that the subword $(\alpha_{i_1}, \dots, \alpha_{i_k})$ of Q is a reduced word for w . If $v \not\geq_{\text{Br}} w$, then $S(Q, w) = \emptyset$ is the simplicial complex with no faces. This must be distinguished from the complex $\{\emptyset\}$, which is $S(Q, v)$ whenever Q is a reduced word for v .

The next theorem is the key fact we use about subword complexes, describing their vertex decomposition, first proved as [30, Theorem E] (for every Coxeter group).

Theorem 5.1. *Let $v, w \in C_n$, and let $Q = (\alpha_1, \dots, \alpha_\ell)$ be a reduced word for v . Assume that $v \geq_{\text{Br}} w$. If $v = 1$, then $w = 1$, $Q = ()$, and $S(Q, w) = \{\emptyset\}$. Otherwise $\ell > 0$. Let $Q' = (\alpha_1, \dots, \alpha_{\ell-1})$ and $i = \alpha_\ell$. Then,*

$$S(Q, w) = \text{cone}_\ell S(Q', w) \cup S(Q', wc_i).$$

5.2 | Labeling the vertices with variables

Recall that $V_{\bar{v}}$ denotes the set of variables of $R_{\bar{v}}$, that is, the set of variables that appear as entries of $M_{\bar{v}}$.

Lemma 5.2. *Let c_k be an ascent of v , $\bar{v} = u_l v_{\square} u_r$, $\overline{vc_k} = (u_l) v_{\square} (u_r c_k)$, and $V_{\bar{v}} \setminus V_{\overline{vc_k}} = \{z_{ij}\}$. Then, $k = j - i$.*

Proof. By Corollary 4.11, z_{ij} is in positions $(v(n \pm k + 1), n \pm k)$ of $M_{\bar{v}}$. We also have that z_{ij} is in position $(v(b_{n+1-i}), a_j)$ and therefore $a_j = n + k$, $b_{n+1-i} = n + k + 1$. Since the number of columns to the right of column $n + k$ is counted by both $|\{a_{i+1}, \dots, a_n\}| + |\{b_{n+1-j}, \dots, b_n\}|$ and $2n - (n + k) = n - k$, we have

$$n - k = |\{a_{i+1}, \dots, a_n\}| + |\{b_{n+1-j}, \dots, b_n\}| = n - i + j.$$

We conclude that $k = j - i$. □

Proposition 5.3. *Let Q be the word $(j_1 - i_1, \dots, j_\ell - i_\ell)$, where $z_{i_1 j_1} <_{\text{lex}} \dots <_{\text{lex}} z_{i_\ell j_\ell}$ are the variables in $V_{\bar{v}}$. The word Q is a reduced word for $w_0 v$.*

Proof. Let $\bar{v} = u_l v_{\square} u_r$. We proceed by induction on $\ell(w_0 v)$. The base case $v = w_0$ is trivial. For the inductive case, let Q' equal Q without the last letter, which we denote by α_ℓ . By construction $\alpha_\ell = j - i$ where z_{ij} is the last variable in $V_{\bar{v}}$ under $<_{\text{lex}}$. Let the lowest box of $D(v)$ containing z_{ij} be in the $(n + k)$ th column. Since z_{ij} is the last variable in $V_{\bar{v}}$, there are no boxes of $D(v)$ weakly southeast of this box. This implies that c_k is the last ascent of v and, by Corollary 4.11, that $V_{\bar{v}} \setminus V_{\overline{vc_k}} = \{z_{ij}\}$, where $k = j - i$ by Lemma 5.2. (As in the proof of Proposition 4.7, $\overline{vc_k}$ denotes the factorization $(u_\ell) v_{\square} (u_r c_k)$, where $u_\ell v_{\square} u_r$ is the factorization denoted by $\bar{v}$.) Therefore, Q' is the word constructed from the variables in $V_{\overline{vc_k}}$. By the induction hypothesis, Q' is a reduced word for $w_0 vc_k$. Because c_{α_ℓ} is an ascent of vc_{α_ℓ} , that is, $\ell((w_0 vc_{\alpha_\ell}) s_{j-i}) = \ell(w_0 vc_{\alpha_\ell}) + 1$, we can append $\alpha_\ell = j - i$ to a reduced word for $w_0 vc_{\alpha_\ell}$ to obtain a reduced word for $w_0 vc_{\alpha_\ell} s_{j-i} = w_0 v$. It follows that Q is a reduced word for $w_0 v$. □




FIGURE 2 The simplicial complex $\Delta_{\overline{v}c_0, w}$ for $v = 321\ 654$ and $w = 635\ 241$.




FIGURE 3 The simplicial complex $\Delta_{\overline{v}, w} = \text{cone}_{z_{33}} \Delta_{\overline{v}c_0, w} \cup \Delta_{\overline{v}c_0, wc_0}$ for $v = 321\ 654$ and $w = 632\ 541$.

Define $\zeta : [\ell] \rightarrow V_{\overline{v}}$ to be the map that associates to k the k th smallest variable in $V_{\overline{v}}$ under $<_{\text{lex}}$. Let

$$\Delta_{\overline{v}, w} = \{\zeta(F) : F \in S(Q, w_0 w)\},$$

where Q is the reduced word for $w_0 v$ defined in Proposition 5.3. This is a simplicial complex isomorphic to $S(Q, w_0 w)$ but relabeled, so its vertex set is $V_{\overline{v}}$.

Proposition 5.4 translates Theorem 5.1 to the notation $\Delta_{\overline{v}, w}$, and breaks it into the cases that will appear in our proofs (which are also parallel to the cases in the statement of Theorem 6.8).

Proposition 5.4. *Let $v, w \in C_n$ and $\overline{v} = u_l v_{\square} u_r$.*

- If $v \not\leq_{\text{Br}} w$, then $\Delta_{\overline{v}, w} = \emptyset$.
- If $v = w_0$, then $w = w_0$ (or we are in the previous case), and $\Delta_{\overline{v}, w} = \{\emptyset\}$.
- Otherwise, let k be the last ascent of v , so $vc_k >_{\text{Br}} v$, and let $\overline{vc_k} = u_l v_{\square} (u_r c_k)$.
 - (1) If k is a descent of w , so $wc_k <_{\text{Br}} w$, then

$$\Delta_{\overline{v}, w} = \text{cone}_{z_{ij}} \Delta_{\overline{vc_k}, w},$$

where z_{ij} is the largest variable with respect to $<_{\text{lex}}$ on $R_{\overline{v}}$.

- (2) If k is an ascent of w , so $wc_k >_{\text{Br}} w$, then

$$\Delta_{\overline{v}, w} = \text{cone}_{z_{ij}} (\Delta_{\overline{vc_k}, w}) \cup \Delta_{\overline{vc_k}, wc_k}.$$

Example 5.5. For $v = v_{\square} = 321\ 654$ and $w = 635\ 241$, we have that $Q = (0, 1, 2, 0, 1, 0)$ and $w_0 w = c_0 c_1$. The last ascent of v is c_0 and this is a descent of w . In this case, $\Delta_{\overline{v}, w} = \text{cone}_{z_{33}} \Delta_{\overline{v}c_0, w}$, and $\Delta_{\overline{v}c_0, w}$ is pictured in Figure 2.

Example 5.6. For $v = v_{\square} = 321\ 654$ and $w = 632\ 541$, we have that $Q = (0, 1, 2, 0, 1, 0)$ and $w_0 w = c_0 c_1 c_0$. The last ascent of v is c_0 and this is an ascent of w . In this case, $\Delta_{\overline{v}, w} = \text{cone}_{z_{33}} (\Delta_{\overline{v}c_0, w}) \cup \Delta_{\overline{v}c_0, wc_k}$ as one can see in Figure 3.

Let $K_{\bar{v},w}$ be the Stanley–Reisner ideal of $\Delta_{\bar{v},w}$. This is the ideal generated by the monomials that are the nonfaces of $\Delta_{\bar{v},w}$, so

$$K_{\bar{v},w} = \left\langle \prod_{z \in Z} z : Z \subseteq V_{\bar{v}}, Z \notin \Delta_{\bar{v},w} \right\rangle.$$

Translating Proposition 5.4 to be in terms of $K_{\bar{v},w}$ gives the following.

Proposition 5.7. *Let $v, w \in C_n$ and $\bar{v} = u_l v \square u_r$.*

- *If $v \not\leq_{\text{Br}} w$, then $K_{\bar{v},w} = \langle 1 \rangle$.*
- *If $v = w_0$, then $w = w_0$ (or we are in the previous case), and $K_{\bar{v},w} = \langle 0 \rangle$.*
- *Otherwise, let k be the last ascent of v , so $vc_k >_{\text{Br}} v$, and let $\overline{vc_k} = u_l v \square (u_r c_k)$. Let z_{ij} be the largest variable with respect to $<_{\text{lex}}$ on $R_{\bar{v}}$.*
 - (1) *If k is a descent of w , so $wc_k <_{\text{Br}} w$, then*

$$K_{\bar{v},w} = K_{\overline{vc_k},w} R_{\bar{v}}.$$

- (2) *If k is an ascent of w , so $wc_k >_{\text{Br}} w$, then*

$$K_{\bar{v},w} = \langle z_{ij} m : m \in K_{\overline{vc_k},w} \rangle + K_{\overline{vc_k},wc_k} R_{\bar{v}}.$$

Example 5.8. In this example, we verify (1) and (2) in the proposition above. First, let v and w be as in Example 5.5. We compute that both ideals are generated by $z_{11}z_{22}, z_{11}z_{23}, z_{12}z_{23}$, although for $K_{\bar{v},w}$, these generators are interpreted in the ring $\mathbb{K}[z_{11}, z_{12}, z_{13}, z_{22}, z_{23}, z_{33}]$, and for $K_{\overline{vc_k},w}$ in the ring $\mathbb{K}[z_{11}, z_{12}, z_{13}, z_{22}, z_{23}]$.

Now, let v and w be as in Example 5.6. Direct computation shows that

$$K_{\bar{v},w} = \langle z_{11}z_{33}, z_{12}z_{33}, z_{22}z_{33}, z_{11}z_{22}, z_{11}z_{23}, z_{12}z_{23} \rangle,$$

and $K_{\overline{vc_0},w} = \langle z_{11}, z_{12}, z_{22} \rangle$, and $K_{\overline{vc_0},wc_0} = \langle z_{11}z_{22}, z_{11}z_{23}, z_{12}z_{23} \rangle$. Then, $K_{\bar{v},w} = z_{33}K_{\overline{vc_0},w} + K_{\overline{vc_0},wc_0}$.

6 | PROOF OF THEOREM 4.15

In this section, we prove Theorem 4.15. We explain the overall structure of the proof now, and dedicate subsections to the details.

Given $v, w \in C_n$ with $v \geq v \square$ in left–right weak order, a factorization $\bar{v}$ for v , and a diagonal term order $<$ on $R_{\bar{v}}$, let $\mathcal{G}_{\bar{v},w,<}$ be the set of initial monomials of the generators we used to define $I_{\bar{v},w}$. We recall that these generators are the size $r_w(p, q) + 1$ minors of the truncated matrix $\tau_{p,q}(M_{\bar{v}})$, running over all (p, q) in $E(w)$. Let $J_{\bar{v},w,<}$ be the ideal generated by $\mathcal{G}_{\bar{v},w,<}$. We will show that

$$K_{\bar{v},w} \subseteq J_{\bar{v},w,<} \subseteq \text{in}_{<} I_{\bar{v},w},$$

the former containment being Proposition 6.7 below and the latter clear from the definition of $J_{\bar{v},w}$. In Proposition 6.9, we will prove that the K -polynomials $\mathcal{K}(R_{\bar{v}}/K_{\bar{v},w}; \mathbf{t})$ and $\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \mathcal{K}(R_{\bar{v}}/\text{in}_{<} I_{\bar{v},w}; \mathbf{t})$ are equal. The containments above then imply

$$K_{\bar{v},w} = J_{\bar{v},w,<} = \text{in}_{<} I_{\bar{v},w},$$

and the latter equality is the statement of Theorem 4.15. Note that this shows that $J_{\bar{v},w,<}$ is independent of the choice of diagonal term order $<$.

6.1 | The Stanley–Reisner ideal is contained in the initial ideal

This subsection proves the containment $K_{\bar{v},w} \subseteq J_{\bar{v},w,<}$, which is Proposition 6.7. The proof will be by induction on the length $\ell(w_0 v)$. A factorization $v = u_l v_{\square} u_r$, where $\ell(v) = \ell(u_l) + \ell(v_{\square}) + \ell(u_r)$, can be extended to a factorization $w_0 = u_l v_{\square} u_r (w_0 v)^{-1}$, and if the induction were unrolled, it would descend to v from its base case $v = w_0$ in right weak order by acting by simple reflections at the right of this factorization. Thus, we can use right weak order to induct down from w_0 to any 123-avoiding permutation where every permutation along the way is 123-avoiding.

Throughout this section, we let $\bar{v}$ be the factorization $v = u_l v_{\square} u_r$, and, for c_k an ascent of v , we let $\overline{v c_k}$ be the factorization $v = u_l v_{\square} (u_r c_k)$.

Every term of the Leibniz formula for a minor of $M_{\bar{v}}$ is zero or a signed monomial in the variables z_{ij} . Our proofs in this section will rely on the fact that there are no cancelations among these terms. This is essentially the fact known in spectral graph theory as the Harary–Sachs theorem [20, 36].

Lemma 6.1. *Every coefficient of any minor of $M_{\bar{v}}$ is a signed power of 2. If $\prod_{j=1}^r (M_{\bar{v}})_{p_j q_j}$ is nonzero, then it is a monomial contained in the support of the $(\{p_1, \dots, p_r\}, \{q_1, \dots, q_r\})$ minor of $M_{\bar{v}}$.*

Proof. If a square submatrix N of $M_{\bar{v}}$ contains an entry 1, then by Proposition 4.6, expansion along either its row or column shows that every nonzero term in $\det N$ involves that entry 1 and N has the same determinant as a smaller submatrix, up to sign. So, we may assume that N contains no 1s.

Give the rows of N the names $\epsilon_1, \dots, \epsilon_r$ and its columns the names $\delta_1, \dots, \delta_r$, in such a way that whenever N contains a row and column that contribute the same weight to the action of T on Σ_v (see Proposition 4.18), then this row and column are ϵ_i and δ_i for some i . This ensures that, if a variable is repeated in N , the two positions at which it appears are (ϵ_i, δ_j) and (ϵ_j, δ_i) for some i and j .

We have

$$\pm \det(N) = \sum_{u \in S_r} \text{sgn}(u) \prod_j N_{\epsilon_j \delta_{u(j)}}.$$

Suppose $u, u' \in S_r$ index terms of this sum which are both equal, up to sign, to a monomial m . Because each nonzero entry of N is a variable, the multisets of variables entering the product for u and that for u' must be equal. Thus, for every $i, j \in [r]$, if $u(i) = j$, then $u'(i) = j$ or $u'(j) = i$ (and if $u(i) = j$ and $u(j) = i$, then $u'(i) = j$ and $u'(j) = i$). It follows that u' is obtained from u by inverting some of the cycles in its disjoint cycle decomposition. A cycle and its inverse have the same sign, so $\text{sgn}(u)$ is constant over all terms with $\prod_j N_{\epsilon_j \delta_{u(j)}} = m$. The coefficient of m in $\det N$

is $\text{sgn}(u)$ times the number of such terms. Because any set of disjoint cycles can be inverted independently, this number is 2^k , where k is the number of cycles C with $\prod_{j \in C} N_{\epsilon_j \delta_{c(j)}} = \prod_{j \in C} N_{\epsilon_{c(j)} \delta_j}$. This is nonzero in $\mathbb{K}$ because $\text{char } \mathbb{K} \neq 2$. $\square$

We now prove the main technical lemma of this section.

Lemma 6.2. Fix a diagonal term order $<$ on $R_{\bar{v}}$. Let c_k be the last ascent of v and fix $<'$, the restriction of $<$, as our diagonal term order on $R_{\bar{v}c_k}$. Suppose that m is the leading term (with respect to $<'$) of the minor of $M_{\bar{v}c_k}$ on row set $A = \{\epsilon_1, \dots, \epsilon_r\}$ and column set $B = \{\delta_1, \dots, \delta_r\}$, labeled so that the entries giving the leading term are the (ϵ_j, δ_j) entries, that is, $m = \prod_{j=1}^r (M_{\bar{v}c_k})_{\epsilon_j \delta_j}$. Let p, q be such that $A \subset [2n] \setminus [p-1]$ and $B \subset [q]$, and assume that row ϵ_j of $\tau_{p,q}(M_{\bar{v}c_k})$ contains an entry 1 exactly when $j = s+1, \dots, r$. Define

$$B' = \{c_k(\delta_1), \dots, c_k(\delta_s), v^{-1}(\epsilon_{s+1}), \dots, v^{-1}(\epsilon_r)\}.$$

- (1) The leading term m' (with respect to $<$) of the minor of $M_{\bar{v}}$ on row set A and column set B' divides m .
- (2) If c_k is a descent of w , $m \in \mathcal{G}_{\bar{v}c_k, w, <'}$, $(p, q) \in E(w)$, and $r = r_w(p, q) + 1$ then $m' \in \mathcal{G}_{\bar{v}, w, <}$.
- (3) If c_k is an ascent of w , $m \in \mathcal{G}_{\bar{v}c_k, w, <'}$, $(p, q) \in E(w)$, $r = r_w(p, q) + 1$, and $V_{\bar{v}} \setminus V_{\bar{v}c_k} = \{z_{ij}\}$, then there exists $m'' \in \mathcal{G}_{\bar{v}, w, <}$ with $m'' \mid z_{ij}m'$.

Proof. Let $1 \leq \delta \leq 2n$ be a column index. By Proposition 4.6, in any minor of $M_{\bar{v}}$ using row $v(\delta)$ and column δ , all nonvanishing terms use the $(v(\delta), \delta)$ entry. Applied to the (A, B') minor of $M_{\bar{v}}$, we conclude that this minor equals, up to sign, the

$$(\{\epsilon_1, \dots, \epsilon_s\}, \{c_k(\delta_1), \dots, c_k(\delta_s)\})$$

minor of $M_{\bar{v}}$.

Let us first show that the $(\{\epsilon_1, \dots, \epsilon_s\}, \{c_k(\delta_1), \dots, c_k(\delta_s)\})$ minor of $M_{\bar{v}}$ equals, up to sign, the $(\{\epsilon_1, \dots, \epsilon_s\}, \{\delta_1, \dots, \delta_s\})$ minor of $M_{\bar{v}c_k}$. By Corollary 4.11, $M_{\bar{v}c_k}$ is obtained from $M_{\bar{v}}P(c_k)$ by setting a single variable z_{ij} to 0, which appears in columns $n \pm k$ of $M_{\bar{v}}$. Proposition 4.7 shows that none of the other variables change names. Therefore, the claim is straightforward except in the case where z_{ij} shows up in the $(\{\epsilon_1, \dots, \epsilon_s\}, \{c_k(\delta_1), \dots, c_k(\delta_s)\})$ minor of $M_{\bar{v}}$. This occurs if $n \pm k + 1 \in \{\delta_1, \dots, \delta_s\}$. Without loss of generality, let $\delta_1 = n \pm k + 1$; we must then have $v c_k(\delta_1) = v(\delta_1 - 1) \in \{\epsilon_1, \dots, \epsilon_s\}$. Let $\epsilon_t = v(\delta_1 - 1)$. Since $\delta_1 \leq q$ and $p \leq \epsilon_t$, $\tau_{p,q}(M_{\bar{v}})$ contains a 1 in row ϵ_t , namely, at (ϵ_t, δ_1) . This contradicts the definition of s and we can thus conclude that $n \pm k + 1 \notin \{\delta_1, \dots, \delta_s\}$.

Let $\mu = \prod_{l=1}^s (M_{\bar{v}c_k})_{\epsilon_l \delta_l}$ and note $m/\mu = \prod_{l=s+1}^r (M_{\bar{v}c_k})_{\epsilon_l \delta_l}$. We now show the leading term m' of the minor in the previous paragraph must be μ . This is because any term m'' of this minor can be extended to a term $m''m/\mu$ of the (A, B) minor of $M_{\bar{v}c_k}$. Therefore, $m = m''m/\mu$ for some term m'' of the smaller minor, that is, $m'' = \mu$. The fact that monomial orders respect multiplication implies that $m' = \mu$. We conclude that $m' \mid m$ and (1) follows.

We now show (2). Suppose that c_k is a descent of w and $m \in \mathcal{G}_{\bar{v}c_k, w, <'}$. Since $(p, q) \in E(w)$, we cannot have $q = n \pm k$ and it follows that $B' \subset [q]$. We therefore have that m' is the leading term of a minor of $\tau_{p,q}(M_{\bar{v}})$ of size $r_w(p, q) + 1$ and (2) follows.

$$\begin{array}{ccccc}
 & & b & & q \\
 & a & z_{\diamondsuit} & \dots & z_{\diamondsuit j} \\
 \text{(i)} & & \vdots & & \vdots \\
 & \epsilon_{s+1} & z_{i\diamondsuit} & \dots & z_{ij}
 \end{array}
 \qquad
 \begin{array}{ccccc}
 & & b & & q \\
 & a & 0 & \dots & (\neq 0) \\
 & & \vdots & & \vdots \\
 \text{(ii)} & v(b) & 1 & \dots & 0 \\
 & & \vdots & & \vdots \\
 & \epsilon_{s+1} & z_{i\diamondsuit} & \dots & z_{ij}
 \end{array}$$

FIGURE 4 The entries of $M_{\bar{v}}$ in relevant rows and columns.

To show (3), first note that, if $B' \subseteq [q]$, then, as in (2), m' is the leading term of a minor of size $r_w(p, q) + 1$ lying inside $\tau_{p,q}(M_{\bar{v}})$, so we can take $m'' = m'$, and $m'' \mid z_{ij}m'$.

If $B' \not\subseteq [q]$, then we must have $q = n \pm k$. Since c_k is an ascent of v , we have $v_q < v_{q+1}$, so $q + 1$ is not a left-to-right minimum of v . Hence, $q + 1$ is a right-to-left maximum of v by Lemma 4.2, and, by Proposition 4.6, the only nonzero entry in column $q + 1$ of $M_{\bar{v}}$ is a 1 in position $(v(q + 1), q + 1)$. This implies that $v^{-1}(q + 1) = \epsilon_j$ for some $j > s$. Without loss of generality, suppose $v(\epsilon_{s+1}) = q + 1$.

Let $B'' = B' \setminus \{q + 1\} \cup \{q\}$. Note that $\#B'' = r_w(p, q) + 1$ since $q \notin B'$ as $c_k(q) = q + 1 > q$. Let $\delta_{s+1} = q$, so $c_k(\delta_{s+1}) = q + 1$. We have $(M_{\bar{v}})_{\epsilon_{s+1}c_k(\delta_{s+1})} = z_{ij}$.

We now construct a row set A' so that the (A', B'') -minor of $\tau_{p,q}(M_{\bar{v}})$ has leading term m'' such that $m'' \mid z_{ij}m'$. Without loss of generality, assume $p \leq v_{c_k}(\delta_j) < \epsilon_{s+1}$ when $j \leq t$, and $p > v_{c_k}(\delta_j)$ or $v_{c_k}(\delta_j) \geq \epsilon_{s+1}$ when $t < j \leq s$. Now let $A' = \{v_{c_k}(\delta_1), \dots, v_{c_k}(\delta_t), \epsilon_{t+1}, \dots, \epsilon_r\}$. We show that the leading term of the (A', B'') -minor of $M_{\bar{v}}$ is $m'' = \prod_{\ell=t+1}^{s+1} (M_{\bar{v}})_{\epsilon_{\ell}c_k(\delta_{\ell})}$ and that $m'' \mid z_{ij}m'$.

First, we show by contradiction that $(M_{\bar{v}})_{\epsilon_{s+1}c_k(\delta_{s+1})} = z_{ij}$ must be part of the leading term. Since c_k is the last ascent, every entry in column $c_k(\delta_{s+1}) = q$ below row ϵ_{s+1} must be 0. Hence, if the leading term of the (A', B'') -minor of $M_{\bar{v}}$ does not contain z_{ij} , then there must exist some row $a \in A'$ with $a < \epsilon_{s+1}$ and some column $b \in B''$ with $b < q$ such that the (a, q) and (ϵ_{s+1}, b) entries are in the leading term of the minor. Now note that $(M_{\bar{v}})_{ab} = 0$, as, otherwise, by the diagonalness of the term order, this entry and z_{ij} would give a larger term, as seen in Figure 4(i). Since $(M_{\bar{v}})_{ab} = 0$ but $(M_{\bar{v}})_{aq} \neq 0$ and $(M_{\bar{v}})_{\epsilon_{s+1}b} \neq 0$, we must have that $a < v(b) < \epsilon_{s+1}$, as seen in Figure 4(ii). By our labeling, we must have $b = c_k(\delta_j)$ for some $j \leq t$, and $v(b) \in A'$. Now note that b must be a left-to-right minimum, so the only nonzero entry in row $v(b)$ is the 1 at $(v(b), b)$. This contradicts our assumption that the (ϵ_{s+1}, b) entry of $M_{\bar{v}}$ is in the leading term.

Next, note that $\tilde{m} = \prod_{\ell=t+1}^s (M_{\bar{v}})_{\epsilon_{\ell}c_k(\delta_{\ell})}$ must be the leading term of the $(\{\epsilon_{t+1}, \dots, \epsilon_s\}, \{c_k(\delta_{t+1}), \dots, c_k(\delta_s)\})$ minor of $M_{\bar{v}}$, because any other term $\tilde{m}'$ can be extended to a term $\tilde{m}'m'/\tilde{m}$ of the (A, B') minor of $M_{\bar{v}}$ by multiplying by $m'/\tilde{m} = \prod_{\ell=1}^t (M_{\bar{v}})_{\epsilon_{\ell}c_k(\delta_{\ell})}$, but $m' \geq \tilde{m}'m'/\tilde{m}$ by (1), and the fact that monomial orders respect multiplication implies that $\tilde{m} \geq \tilde{m}'$.

We have left-to-right minima at $c_k(\delta_1), \dots, c_k(\delta_t)$. Hence, by Proposition 4.6, every term in the (A', B'') minor of $M_{\bar{v}}$ must use the $(v_{c_k}(\delta_{\ell}), c_k(\delta_{\ell}))$ th entries, which are all 1s. Therefore, the leading term is $z_{ij}\tilde{m}$. By definition, $m'' = z_{ij}\tilde{m}$. By construction, we have $m'' \mid z_{ij}m'$. $\square$

Example 6.3. Let $v = 326154$ and $w = 465213$ in C_3 . Then, the last ascent of v is c_1 : that is, in our one-line notation for v , position $4 = 3 + 1$ is the rightmost position of a digit that is followed by a larger digit. Since w has a descent at c_1 , we are in part (2) of Lemma 6.2. We have

$$M_{\bar{v}} = \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{11} & z_{12} & 0 & z_{13} & 0 & 1 \\ z_{12} & z_{22} & 0 & z_{23} & 1 & 0 \\ z_{13} & z_{23} & 1 & 0 & 0 & 0 \end{bmatrix}, \quad M_{\overline{vc_1}} = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ z_{11} & 0 & z_{12} & 0 & z_{13} & 1 \\ z_{12} & 0 & z_{22} & 1 & 0 & 0 \\ z_{13} & 1 & 0 & 0 & 0 & 0 \end{bmatrix},$$

So, the type C essential set of w is $E(w) = \{(5, 1)\}$. Thus, $\mathcal{G}_{\bar{v},w} = \{z_{13}, z_{12}\} = \mathcal{G}_{\overline{vc_1},w}$ in this case, and so, each element of $\mathcal{G}_{\overline{vc_1},w}$ is divisible by an element of $\mathcal{G}_{\bar{v},w}$.

Example 6.4. Let $v = v_{\square}c_0 \in C_5$ and let $w = a937\,654\,821$, where $a = 10$. Then, $E(w) = \{(8, 7)\}$, $r_w(8, 7) = 2$, the last ascent of v is c_1 , and this is a descent of w . We have

$$\tau_{8,7}(M_{\bar{v}}) = \begin{bmatrix} z_{13} & z_{23} & z_{33} & z_{34} & 0 & z_{35} & 0 \\ z_{14} & z_{24} & z_{34} & z_{44} & 0 & z_{45} & 1 \\ z_{15} & z_{25} & z_{35} & z_{45} & 1 & 0 & 0 \end{bmatrix}$$

and

$$\tau_{8,7}(M_{\overline{vc_1}}) = \begin{bmatrix} z_{13} & z_{23} & z_{33} & 0 & z_{34} & 0 & z_{35} \\ z_{14} & z_{24} & z_{34} & 0 & z_{44} & 1 & 0 \\ z_{15} & z_{25} & z_{35} & 1 & 0 & 0 & 0 \end{bmatrix}.$$

Observe that the 3×3 minor of $\tau_{8,7}(M_{\overline{vc_1}})$ coming from columns 3,5,7 is $z_{35}z_{44}z_{35}$, so that this term is in $\mathcal{G}_{\overline{vc_1},w,<}$ (no matter what $<$ is). This term is not in $\mathcal{G}_{\bar{v},w,<}$. In the proof of Lemma 6.2(1), we take the minor from the last three columns of $\tau_{8,7}(M_{\bar{v}})$, which constitute B' . This minor is z_{35} , which divides $z_{35}z_{44}z_{35}$.

Example 6.5. Let $v = c_1c_0v_{\square} = 6\,4218\,753 \in C_4$ and $w = 87436\,521$. Then, $E(w) = \{(5, 4)\}$, $r_w(5, 4) = 2$, the last ascent of v is c_0 , and this is an ascent of w . We have:

$$\tau_{5,4}(M_{\bar{v}}) = \begin{bmatrix} 0 & z_{22} & z_{23} & z_{24} \\ 1 & 0 & 0 & 0 \\ z_{13} & z_{23} & z_{33} & z_{34} \\ z_{14} & z_{24} & z_{34} & z_{44} \end{bmatrix}$$

and

$$\tau_{5,4}(M_{\overline{v_{c_0}}}) = \begin{bmatrix} 0 & z_{22} & z_{23} & 0 \\ 1 & 0 & 0 & 0 \\ z_{13} & z_{23} & z_{33} & 0 \\ z_{14} & z_{24} & z_{34} & 1 \end{bmatrix}.$$

Observe that the 3×3 minor of $\tau_{5,4}(M_{\overline{v_{c_0}}})$ using columns 1, 3, and 4 and rows 5, 7, and 8 is $z_{13}z_{23}$. The leading term of the corresponding minor in $M_{\overline{v}}$ uses column 5, which is outside $\tau_{5,4}(M_{\overline{v}})$. In the proof of Lemma 6.2(3), we first replace column 5 with column 4, then row 7 with row 6. The leading term of this minor of $\tau_{5,4}(M_{\overline{v}})$ is $z_{23}z_{44}$, which divides $z_{13}z_{23}z_{44}$.

Lemma 6.6. *Let $w \in C_n$ and c_k be an ascent of w . Let $(p, q) \in E(wc_k)$.*

- (i) *If $q \neq n + k + 1$ and $q \neq n + 1 - k$, then $(p, q) \in E(w)$ and $r_w(p, q) = r_{wc_k}(p, q)$.*
- (ii) *If $q = n + k + 1$ or $q = n + 1 - k$ and $(p, q - 1) \in D(wc_k)$, then $(p, q) \in E(w)$ and $r_w(p, q) = r_{wc_k}(p, q)$.*
- (iii) *If $q = n + k + 1$ or $q = n + 1 - k$ and $(p, q - 1) \notin D(wc_k)$, then $(p, q - 1) \in E(w)$ and $r_w(p, q - 1) = r_{wc_k}(p, q) - 1$.*

Proof. Note that we cannot have $q = n + k$ or $q = n - k$ since c_k is a descent of wc_k . The first statement then follows from Lemma 4.9.

Suppose $q = n + k + 1$ or $q = n + 1 - k$ and $(p, q - 1) \in D(wc_k)$. Since $(p, q) \in E^A(wc_k)$, $wc_k(q) < p \leq wc_k(q + 1)$. Since $(p, q - 1) \in D(wc_k)$, $wc_k(q - 1) < p$. Hence, $w(q) = wc_k(q - 1) < p \leq w(q + 1) = wc_k(q + 1)$, and $(p, q) \in E^A(w)$. Also, $r_w(p, q) = r_{wc_k}(p, q)$. Since $(p, q) \in E(wc_k)$, $p \geq n + 1$. Furthermore, if $q = n + k + 1$ and $(p, 2n - q) = (p, n - k - 1) \in E^A(w)$, then $(p, n - k - 1) \in E^A(wc_k)$. Applying the second condition of Definition 3.10 to $(p, q) \in E(wc_k)$, we have that $r_{wc_k}(p, n - k - 1) > r_{wc_k}(p, n + k + 1) - k - 1$. Since $r_w(p, n - k - 1) = r_{wc_k}(p, n - k - 1)$, we also have $r_w(p, 2n - q) > r_w(p, q) + n - q$. Hence, $(p, q) \in E(w)$.

Now suppose $q = n + k + 1$ or $q = n + 1 - k$ and $(p, q - 1) \notin D(wc_k)$. Then, $w(q) = wc_k(q - 1) > p$. Hence, $w(q - 1) < p < w(q)$, and $(p, q - 1) \in E^A(w)$. Also, $r_w(p, q - 1) = r_{wc_k}(p, q) - 1$. If $q = n + k + 1$ and $(p, 2n - (q - 1)) = (p, n - k) \in E^A(w)$, then $w(n + 1 - k) \geq p$, so $wc_k(n - k) \geq p$. We first wish to show that $(p, n - 1 - k) \in E^A(wc_k)$. Since $(p, n + k + 1) \in E(wc_k)$, we have $wc_k(n + k + 2) \geq p > n$. Using the symmetry of type C permutations, as noted in (1), $wc_k(n - 1 - k) \leq n < p$. This implies $(p, n - 1 - k) \in D(wc_k)$ and $(p, n - k) \notin D(wc_k)$. To conclude that $(p, n - 1 - k) \in E^A(wc_k)$, we must show that $(p - 1, n - 1 - k) \notin D(wc_k)$. Note that $(p - 1, n - 1 - k) \notin D(wc_k)$ if and only if $(p - 1, n - 1 - k) \notin D(w)$. We are assuming that $(p, n - k) \in E^A(w)$. Therefore, $w(j) \neq p - 1$ for $j > n - k$, and thus, we can only have $(p - 1, n - 1 - k) \in D(w)$ if $w(n - k) = p - 1$. However, since $w(q) > p > n$ and $w(n - k) = w(2n + 1 - q)$, (1) implies that $w(n - k) < n \leq p - 1$. It follows that $(p, n - 1 - k) \in E^A(wc_k)$.

Applying the second condition of Definition 3.10 to $(p, q) \in E(wc_k)$, we have that $r_{wc_k}(p, n - 1 - k) > r_{wc_k}(p, n + k + 1) - k - 1$. Since $r_w(p, n - k) = r_w(p, n - k - 1) = r_{wc_k}(p, n - k - 1)$ and $r_w(p, n + k) = r_{wc_k}(p, n + k + 1) - 1$, we see that $r_w(p, n - k) > r_w(p, n + k) - k$, and $(p, q - 1) \in E(w)$. $\square$

Proposition 6.7. *Let $v \geq v_{\square}$ and $w \in C_n$. Then, $K_{\overline{v}, w} \subseteq J_{\overline{v}, w, <}$ for any diagonal term order $<$.*

Proof. We induct on $\ell(w_0v)$. In the base case, $v = w_0$, these are both ideals of the polynomial ring $R_{w_0} = \mathbb{K}$ in zero variables, namely, the zero ideal if $w = w_0$ and the unit ideal otherwise.

If $v \neq w_0$, then it may occur that $v \not\leq w$ so that $K_{\bar{v},w}$ is the unit ideal. In this case, $r_v(p, q) > r_w(p, q)$ for some $(p, q) \in E(w)$. Hence, there will be at least $r_w(p, q) + 1$ entries equal to 1 in $\tau_{p,q}(M_{\bar{v}})$. Taking a size $r_w(p, q) + 1$ minor of $\tau_{p,q}(M_{\bar{v}})$ that contains a 1 in each column, we see that the minor evaluates to 1 (every entry above and to the right of a 1 being 0). Thus, $1 \in J_{\bar{v},w}$ as desired.

Henceforth, we assume $v \leq w$. Let c_k be the last ascent of v , let $\{z_{ij}\} = V_{\bar{v}} \setminus V_{\bar{v}c_k}$, and let $\prec'$ be the restriction of $\prec$ to $R_{\bar{v}c_k}$. We consider two cases according to whether c_k is an ascent or a descent of w . If a descent, Proposition 5.7 says

$$K_{\bar{v},w} = K_{\bar{v}c_k,w} R_{\bar{v}}.$$

By induction, $K_{\bar{v}c_k,w} \subseteq J_{\bar{v}c_k,w,\prec'}$, since $\prec'$ is a diagonal term order by Proposition 4.13. So, it suffices to show that $J_{\bar{v}c_k,w,\prec'} R_{\bar{v}} \subseteq J_{\bar{v},w,\prec}$; but this is part (2) of Lemma 6.2.

If c_k is an ascent of w , then Proposition 5.7 says

$$K_{\bar{v},w} = \langle z_{ij}m : m \in K_{\bar{v}c_k,w} \rangle + K_{\bar{v}c_k,wc_k} R_{\bar{v}}. \quad (7)$$

Let $\hat{m}$ be a monomial generator of $K_{\bar{v},w}$. We split into two cases again according to whether $\hat{m} \in K_{\bar{v}c_k,wc_k}$.

If $\hat{m} \in K_{\bar{v}c_k,wc_k}$, then by induction, $\hat{m} \in J_{\bar{v}c_k,wc_k,\prec'}$. Hence, there exists $m \in \mathcal{G}_{\bar{v}c_k,wc_k,\prec'}$ dividing $\hat{m}$. Since c_k is an ascent of w , it is a descent of wc_k , and therefore, by Lemma 6.2, there exists $m' \in \mathcal{G}_{\bar{v},wc_k,\prec}$ dividing m and hence $\hat{m}$. Since $m' \in \mathcal{G}_{\bar{v},wc_k,\prec}$, it is the leading term of some minor of $\tau_{p,q}(M_{\bar{v}})$, of size $r_{wc_k}(p, q) + 1$, for some $(p, q) \in E(wc_k)$. Call this minor D . We now apply Lemma 6.6, breaking the argument into its three cases.

If $q \neq n \pm k + 1$, then $(p, q) \in E(w)$ and $r_w(p, q) = r_{wc_k}(p, q)$, so $D \in I_{\bar{v},w}$, $m' \in \mathcal{G}_{\bar{v},w,\prec}$, and therefore, $m \in J_{\bar{v},w,\prec}$. Similarly, if $q = n \pm k + 1$ and $(p, q - 1) \in D(wc_k)$, then $m \in J_{\bar{v},w,\prec}$. Lastly, if $q = n \pm k + 1$ and $(p, q - 1) \notin D(wc_k)$, the lemma implies that $(p, q - 1) \in E(w)$ and $r_w(p, q - 1) = r_{wc_k}(p, q) - 1$. If column q is not used in D , then D is a minor of $\tau_{p,q-1}(M_{\bar{v}})$ of size $r_w(p, q - 1) + 2$. By Laplace expansion, m' , being the leading term of D , is divisible by the leading term m'' of some minor of size $r_w(p, q - 1) + 1$. Hence, $m'' \in \mathcal{G}_{\bar{v},w,\prec}$ and $\hat{m} \in J_{\bar{v},w,\prec}$. Otherwise, if column q is used in D , let (a, q) be the position in column q appearing in the leading term. Then $m'' := m' / (M_{\bar{v}})_{a,q}$ is the leading term of the minor of size $r_{wc_k}(p, q) - 1$ obtained by omitting row a and column q . Therefore, $m'' \in \mathcal{G}_{\bar{v},w,\prec}$ and $\hat{m} \in J_{\bar{v},w,\prec}$.

If instead $\hat{m} \notin K_{\bar{v}c_k,wc_k}$, then $\hat{m}/z_{ij} \in K_{\bar{v}c_k,w}$ and by induction $\hat{m}/z_{ij} \in J_{\bar{v}c_k,w,\prec'}$. Let $m \in \mathcal{G}_{\bar{v}c_k,w}$ be a generator dividing $\hat{m}/z_{ij}$, arising from a minor of $\tau_{p,q}(M_{\bar{v}c_k})$. Part (1) of Lemma 6.2 (which does not literally apply because c_k is not a descent of w) produces a minor D of $M_{\bar{v}}$ of size $r_w(p, q) + 1$ with a leading term m' that divides m , and part (3) produces $m'' \in \mathcal{G}_{\bar{v},w,\prec}$ dividing $z_{ij}m'$. Hence, we have $m'' \mid z_{ij}m' \mid z_{ij}m \mid \hat{m}$, and $\hat{m} \in J_{\bar{v},w,\prec}$. $\square$

6.2 | Equality of K -polynomials

We wish to show that, with respect to the weighting of the variables on $R_{\bar{v}}$ introduced in Section 4.3, the K -polynomial of the $R_{\bar{v}}$ -module $R_{\bar{v}}/I_{\bar{v},w}$ is equal to the K -polynomial of $R_{\bar{v}}/K_{\bar{v},w}$. We do this by showing that the K -polynomial of $R_{\bar{v}}/K_{\bar{v},w}$ satisfies the recursion of the next theorem, due to B. Kostant and S. Kumar [31, Proposition 2.4]. Kumar showed that the local K -classes of X_w ,

and hence, the K -polynomials of $R_{\bar{v}}/I_{\bar{v},w}$, follow this recursion. The version below is due to Knutson (see [27, Theorem 1]) and has been translated to our specific setting. See also [43, Theorem 6.12] for the type A analog of the statement below.

Throughout this subsection, we assume that $\mathbb{K} = \mathbb{C}$ as the recursion on K -polynomials that we reference is only proved in that setting. In the next subsection, we show that, despite this assumption, Theorem 4.15 holds over arbitrary $\mathbb{K}$ of characteristic zero.

Theorem 6.8. *Let $v, w \in C_n$.*

- *If $v \not\leq w$, then*

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = 0.$$

- *If $v = w_0$, then $w = w_0$ (or we are in the previous case). Then,*

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = 1.$$

- *Otherwise, let k be the last right ascent of v , so $vs_k > v$.*

- (1) *If k is a descent of w , so $cs_k < w$, then*

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \mathcal{K}(R_{\overline{vc_k}}/I_{\overline{vc_k},w}; \mathbf{t}).$$

- (2) *If k is an ascent of w , so $cs_k > w$, then*

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \mathcal{K}(R_{\overline{vc_k}}/I_{\overline{vc_k},w}; \mathbf{t}) + (1 - t_i t_j) \mathcal{K}(R_{\overline{vc_k}}/I_{\overline{vc_k},wc_k}; \mathbf{t}) - (1 - t_i t_j) \mathcal{K}(R_{\overline{vc_k}}/I_{\overline{vc_k},w}; \mathbf{t}),$$

where (i, j) is such that z_{ij} is the smallest variable with respect to $<_{\text{lex}}$ on $R_{\bar{v}}$.

Proof. This is [27, Theorem 1] in the type C setting, but we should explain two changes that arise in translating the statement to our setting.

First, the original statement is about the pullback of the class $[\mathcal{O}_{X_w}] \in K_T(G/B_G^+)$ to $K_T(vB_G^+/B_G^+)$, where vB_G^+/B_G^+ is the point in the stratification of G/B_G^+ . In commutative algebra terms, this is

$$\sum_i (-1)^i [\text{Tor}_i(\mathcal{O}_{X_w}, \mathcal{O}_{vB_G^+/B_G^+})] \in K_T(vB_G^+/B_G^+).$$

A class in $K_T(\text{pt})$ can be identified with its formal character. On the other hand, since vB_G^+/B_G^+ is the point in $M_{\bar{v}}$ with all coordinates set to 0, taking Tor with $\mathcal{O}_{vB_G^+/B_G^+}$ is the same as taking the K -polynomial.

Second, we need to match $e^{v(\alpha)}$ with $t_i t_j$. As shown in the proof of Proposition 4.7, we have $n + i = v(n + k + 1)$ and $n + 1 - j = v(n + k)$, so $v(e_k) = -e_j$ and $v(e_{k+1}) = e_i$. Since $\alpha = e_{k+1} - e_k$, $v(\alpha) = e_i + e_j$, so $e^{v(\alpha)} = t_i t_j$. (To be precise, there are two canceling sign differences from the original statement, one from our definition of X_w as B_G^+ -orbit closures rather than B_G^- -orbit closures, and the second from our use of $w_0 v$ instead of v .) $\square$

Proposition 6.9. *Given $v, w \in C_n$,*

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \mathcal{K}(R_{\bar{v}}/K_{\bar{v},w}; \mathbf{t}).$$

Proof. [35, Theorem 1.13] implies that given a simplicial complex Δ on the vertex set $V \cup \{z\}$ with its vertex decomposition at z , say $\Delta = \text{cone}_z \Lambda \cup \Pi$ where $\Lambda \subseteq \Pi$ are simplicial complexes on V (respectively, the link and deletion of z in Δ), we have

$$\mathcal{K}(R[V \cup \{z\}]/I_{\Delta}; \mathbf{t}) = \mathcal{K}(R[V]/I_{\Lambda}; \mathbf{t}) + (1 - \mathbf{t}^{\deg(z)})\mathcal{K}(R[V]/I_{\Pi}; \mathbf{t}) - (1 - \mathbf{t}^{\deg(z)})\mathcal{K}(R[V]/I_{\Lambda}; \mathbf{t}).$$

Applied to Proposition 5.4, this produces a recursive formula for $\mathcal{K}(R/K_{\bar{v},w}; \mathbf{t})$, which comes out identical to Theorem 6.8 with every appearance of I replaced by K . Thus, $\mathcal{K}(R/I_{\bar{v},w}; \mathbf{t})$ and $\mathcal{K}(R/K_{\bar{v},w}; \mathbf{t})$ satisfy the same recursion, and so are the same. $\square$

6.3 | Proof of Theorem 4.15

Proof. We first assume $\mathbb{K} = \mathbb{C}$, and return to an arbitrary field of characteristic zero in the last paragraph. Fix a diagonal term order $<$. By Proposition 6.7, $K_{\bar{v},w} \subseteq J_{\bar{v},w,<}$. By definition, $J_{\bar{v},w,<} \subseteq \text{in}_{<} I_{\bar{v},w}$. Hence, we have surjections

$$R/K_{\bar{v},w} \twoheadrightarrow R/J_{\bar{v},w,<} \twoheadrightarrow R/\text{in}_{<} I_{\bar{v},w}.$$

Now Proposition 6.9 states that

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \mathcal{K}(R_{\bar{v}}/K_{\bar{v},w}; \mathbf{t}).$$

Since

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \mathcal{K}(R_{\bar{v}}/\text{in}_{<} I_{\bar{v},w}; \mathbf{t}),$$

the above containments are actually equalities, and

$$J_{\bar{v},w,<} = \text{in}_{<} I_{\bar{v},w},$$

as desired.

To complete the proof, we note that since the essential minors in $I_{\bar{v},w}$ are polynomials with $\mathbb{Z}$ coefficients, the essential minors are a Gröbner basis over any field $\mathbb{K}$ of characteristic zero. Indeed, if f and g are essential minors, then the S -polynomial $S(f, g)$ reduces to 0 under division by the essential minors when working over $\mathbb{Q}$; hence, it does over any field of characteristic zero. $\square$

The above proof also gives the following corollary.

Corollary 6.10. *Under any diagonal term order, the initial ideal of $I_{\bar{v},w}$ is $K_{\bar{v},w}$.*

By Lemma 6.1, all coefficients of essential minors are powers of 2.




FIGURE 5 The pipe dream on the left is reduced and contains the pipe dream on the right, which is not reduced. These are both pipe dreams for 1432.

Conjecture 6.11. *Theorem 4.15 holds over an arbitrary field $\mathbb{K}$ of characteristic not 2.*

7 | K-POLYNOMIAL FORMULAS VIA PIPE DREAMS

7.1 | Type C pipe dreams on small patches

In this section, we recall the notion of pipe dreams associated to pairs of type A permutations v, w . We then define a closely related notion of type C pipe dreams, specifically for pairs of permutations $v, w \in C_n$ with $v \geq v_{\square}$. These are simply type A pipe dreams with symmetry imposed about the diagonal.

We begin with pipe dream complexes for pairs of permutations in S_m . Pipe dreams were invented by S. Fomin and A. Kirillov in [10] and further studied by N. Bergeron and S. Billey in [4]. Knutson and Miller [29, 30] endowed them with the structure of a simplicial complex, namely, a subword complex. A (type A) *pipe dream* is a tiling of the entries in the southwest triangle of an $m \times m$ matrix with the tiles *cross* $\begin{smallmatrix} + \\ + \end{smallmatrix}$, *elbow* $\begin{smallmatrix} \nearrow \\ \searrow \end{smallmatrix}$, and *half elbow* $\begin{smallmatrix} \nearrow \\ \nearrow \end{smallmatrix}$ such that:

- (1) the diagonal is tiled with $\begin{smallmatrix} \nearrow \\ \searrow \end{smallmatrix}$, and
- (2) the weak southwest triangle only uses $\begin{smallmatrix} + \\ + \end{smallmatrix}$ and $\begin{smallmatrix} \nearrow \\ \searrow \end{smallmatrix}$.

A pipe dream ρ induces an arrangement of m pseudolines and ρ is *reduced* if no two pseudolines cross twice. A pipe dream ρ is *contained* in another pipe dream ρ' if the set of positions of the elbows of ρ is contained in the set of positions of the elbows of ρ' : see Figure 5 for an example.

Label the west ends of the pseudolines $1, \dots, m$ bottom to top in the order of their incidence with the west boundary, as in Figure 5. Also label the south ends of the pseudolines with $1, \dots, m$ by transporting the west labels along the pseudolines, except ignoring all crossings subsequent to the first between each pair of pseudolines, that is, moving the labels as if such crosses were elbows instead. Then, a pipe dream for $w \in S_m$ is a pipe dream whose labels along the south boundary read w . See Figure 5 for two examples of pipe dreams for 1432.

Let $v \in S_m$ and consider the pictorial description for $D(v)$ described in Section 3.2. Let us denote by $D_{\sqcup}(v)$ the diagram obtained from $D(v)$ after left-aligning. The (type A) *pipe dream complex* $PD_{v,w}^A$ for $v, w \in S_m$ is the simplicial complex with vertices given by the boxes of $D_{\sqcup}(v)$, and one facet for each reduced pipe dream ρ for w whose crosses are contained in $D_{\sqcup}(v)$, the set of vertices in the facet being the set of positions of elbows in ρ . We will abuse notation and reuse the name $\rho \in PD_{v,w}^A$ for the facet. In Figure 6, we give an example of $D_{\sqcup}(v)$, and in Figure 7 an example of $PD_{v,w}^A$.




FIGURE 6 The diagram $D(v)$, left-aligned diagram $D_L(v)$, and corresponding reduced pipe dream ρ for $v = 426\ 153$. The word $Q_A(\rho)$ corresponding to ρ is $(2,4,5,1,3,4,2)$ that is a reduced word for w_0v .




FIGURE 7 The above simplicial complex is the pipe dream complex $PD_{v,w}^A$ for $v = 426\ 153$ and $w = 456\ 231$. The two shaded triangles are the maximal faces, and each is labeled with an appropriate reduced pipe dream.

The type A *Demazure product* $\mathbf{d}(Q)$ of a word $Q = (\alpha_1, \dots, \alpha_k)$ in the symbols $1, \dots, 2n - 1$ is an element of S_{2n} defined recursively by

$$\mathbf{d}(Q) = \begin{cases} \mathbf{d}(Q')s_{\alpha_k} & \text{if } \ell(\mathbf{d}(Q')s_{\alpha_k}) > \ell(\mathbf{d}(Q')) \\ \mathbf{d}(Q') & \text{if } \ell(\mathbf{d}(Q')s_{\alpha_k}) < \ell(\mathbf{d}(Q')) \end{cases}$$

where $Q' = (\alpha_1, \dots, \alpha_{k-1})$, with the base case that the Demazure product of the empty word is the identity permutation. Compare [29, Definition 3.1]. The Demazure product models ignoring crossings subsequent to the first, as in a pipe dream; in [43], the nilHecke algebra is used for the same purpose.

Associate to a pipe dream ρ a word $Q_A(\rho)$ in the symbols $1, \dots, 2n - 1$ as follows. To a cross tile in position (i, j) , we associate the symbol $i - j$. (This has the effect of making all instances of the symbol k associated to cross tiles on the k th diagonal below the main diagonal.) Then, we obtain the word $Q_A(\rho)$ by reading the symbols associated to cross tiles in the leftmost column from top to bottom, then the next column from top to bottom, and so on. See Figure 6 for an example of $Q_A(\rho)$. We remark that if ρ is a pipe dream for $w \in S_m$, then $\mathbf{d}(Q_A(\rho)) = w_0w$, and that the pipe dream with crosses exactly in $D_L(v)$ is a reduced pipe dream for v .

We now describe type C pipe dream complexes for permutations $v \geq v_{\square}$ and w . A *type C pipe dream* is a (type A) pipe dream whose crosses lie within $D_L(v_{\square}) = D(v_{\square})$ and which is symmetric



FIGURE 8 The diagram on the left is a type C pipe dream for 635 241; however, the diagram on the right is not.




FIGURE 9 On the left is the assignment of generators to the weak lower triangular part of $D(v_{\square})$. On the right is a reduced type C pipe dream ρ for $v = 53\,281\,764$. The word corresponding to ρ is $c_1c_2c_3c_0c_1c_2c_0c_1$ that is a reduced expression for w_0v .

about the diagonal of $D(v_{\square})$. Since all the tiles outside of $D(v_{\square})$ are elbows, we only draw the tiles inside this region, which is an $n \times n$ square (Figure 8). We give the positions in this region coordinates from $(1,1)$ to (n,n) , rather than using the coordinates $(n+1,1)$ to $(2n,n)$ they would inherit from their inclusion in the diagrams for type A pipe dreams.

The type C Demazure product of a word Q in the symbols $0, \dots, n-1$ is the type A Demazure product of the word obtained from Q by replacing each 0 with n and each other symbol i with the two consecutive entries $n-i, n+i$. This is an element of C_n . There being no possibility of confusion, we will also denote this by $\mathbf{d}(Q)$.

Just as in the type A case, we associate to a type C pipe dream ρ a word $Q_C(\rho)$ in the symbols $0, \dots, n-1$. We start by associating the symbol $j-i$ to the positions (j,i) and (i,j) in $D(v_{\square})$, these being the positions where the variable z_{ij} appears in the southwest quarter of $M_{v_{\square}}$. Next, given a type C pipe dream ρ , assign to each cross the symbol corresponding to the position of the cross. Last, for each cross in the weak lower triangular part of $D(v_{\square})$, read the generators in the leftmost column from top to bottom, then the next column from top to bottom, and so on. We define $Q_C(\rho)$ to be the resulting word. We say that ρ is a type C pipe dream for $w \in C_n$ if $\mathbf{d}(Q_C(\rho)) = w_0w$. A type C pipe dream is *reduced* if the word $Q_C(\rho)$ is reduced. We will prove in Section 7.3 that, just as for type A, if we transport labels $1, \dots, 2n$ from the north and west sides of the picture along the pseudolines, ignoring all crossings subsequent to the first between each pair of pseudolines, the resulting labels on the south side read w .

Let $v \in C_n$ with $v \geq v_{\square}$. Note that there is at least one type C pipe dream for v . For example, a factorization $\bar{v}$ of v gives a type C pipe dream, which we will call $\rho(\bar{v})$, consisting of crosses in the positions of z_{ij} in the southwest quarter of $M_{v_{\square}}$ for each $z_{ij} \in V_{\bar{v}}$.

Given a type C pipe dream ρ , we denote by ρ_L the set of positions of its crosses in the weak lower triangular part of $D(v_{\square})$. Given a reduced pipe dream ρ for v and some $w \in C_n$, the *type C pipe dream complex* $PD_{\rho,w}^C$ is the simplicial complex whose vertices are the boxes in ρ_L (Figure 10).




FIGURE 10 The type C pipe dream complex $PD_{\rho,w}^C$ for $w = 58\,372\,615$ and ρ the pipe dream in Figure 9 is obtained from the simplicial complex above by coning the vertex corresponding to the elbow in position (4,2).

The vertices of $PD_{\bar{v},w}^C$ are thus a subset of the entries in the weak lower triangular part of $D(v_{\square})$. The facets of $PD_{\rho,w}^C$ correspond to the reduced type C pipe dreams σ for w such that $\rho \subset \sigma$; the vertices in such a facet are the positions of the elbows of σ that lie in ρ_L . If $\rho = \rho(\bar{v})$, we abbreviate $PD_{\rho(\bar{v}),w}^C$ to $PD_{\bar{v},w}^C$. From the definition, we see that type C pipe dreams that contain a type C pipe dream for a reduced type C word for w are in bijection with faces of $\Delta_{\bar{v},w}$.

The following lemma follows from the definition of $PD_{\rho,w}^C$.

Lemma 7.1. *The simplicial complex $PD_{\bar{v},w}^C$ is the image of $\Delta_{\bar{v},w}$ under the isomorphism acting on vertices as $z_{ij} \mapsto (j, i)$. In particular, the facet F of $\Delta_{\bar{v},w}$ corresponds to the facet $\{(p, q) \in \rho(\bar{v})_L : z_{qp} \in F\}$ of $PD_{\bar{v},w}^C$.*

7.2 | Multidegrees and K-polynomials

In analogy with the type A setting [43, Theorem 3.2], we have that the prime components of the diagonal initial ideal of each $I_{\bar{v},w}$, with $v \geq v_{\square}$ in left–right weak order, are indexed by type C pipe dreams.

Corollary 7.2. *For any diagonal term order $\prec$, the initial ideal $\text{in}_{\prec} I_{\bar{v},w}$ has the prime decomposition*

$$\text{in}_{\prec} I_{\bar{v},w} = \bigcap_{\rho} \langle z_{qp} : (p, q) \in \rho_L \rangle,$$

where ρ ranges over all reduced elements of $PD_{\bar{v},w}^C$.

Proof. By Corollary 6.10, we have $\text{in}_{\prec} I_{\bar{v},w} = K_{\bar{v},w}$, the latter of which is the Stanley–Reisner ideal of the (type C) subword complex $\Delta_{\bar{v},w}$. Consequently,

$$\text{in}_{\prec} I_{\bar{v},w} = \bigcap_{F \in \text{Facets}(\Delta_{\bar{v},w})} \langle \zeta([\ell] \setminus F) \rangle,$$

where $\zeta : [\ell] \rightarrow V_{\bar{v}}$ is as defined immediately after the proof of Proposition 5.3. The result now follows by Lemma 7.1. $\square$

Our next goal is to provide multidegree and K -polynomial formulas for our type C Kazhdan–Lusztig varieties $X_{\bar{v},w}$, $v \geq v_{\square}$ in terms of type C pipe dreams. Our formulas are in the variables $t_1, \dots, t_n$ discussed in Section 4.3. We recall the action of C_n on this variable set described there:

given $u \in C_n$, the action is so that for $i \leq n$,

$$u \odot t_i^{-1} = \begin{cases} t_{n+1-u(n+i)} & \text{if } u(i+n) \leq n, \\ t_{u(n+i)-n}^{-1} & \text{if } u(i+n) \geq n+1. \end{cases}$$

We will also want an additive version of this action:

$$u \oplus -t_i = \begin{cases} t_{n+1-u(n+i)} & \text{if } u(i+n) \leq n, \\ -t_{u(n+i)-n} & \text{if } u(i+n) \geq n+1. \end{cases}$$

Corollary 7.2 immediately implies a positive multidegree formula for our type C Kazhdan-Lusztig varieties $X_{\bar{v},w}$, $v \geq v_{\square}$ in terms of type C pipe dreams. Note that this formula can also be recovered from the K -polynomial formula given below in Proposition 7.4.

Proposition 7.3. *The multidegree of $R_{\bar{v}}/I_{\bar{v},w}$ is*

$$C(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \sum_{\rho} \prod_{\substack{(i,j) \text{ is a} \\ \text{cross in } \rho_L}} ((u_i \oplus -t_i) + (u_j \oplus -t_j)),$$

where ρ ranges over all reduced elements of $PD_{\bar{v},w}^C$.

As noted in the introduction, Ikeda, Mihalcea, and Naruse also give formulas for these multidegrees [21, Theorem 1.1, Definition 6.1]. To aid the reader in aligning conventions, we observe that the multidegree of our $I_{\bar{v},w}$ is what is called in [21] the localization of $\mathfrak{S}_{w_0w}$ at w_0v , but our work and theirs agree on the meaning of t_i .

Remark. Our pipe dream formula specializes to a formula for multiplicities: since each $v \geq v_{\square}$ is 123-avoiding, each ideal $I_{\bar{v},w}$ is homogeneous with respect to the standard grading. Consequently, the multiplicity $\text{mult}_{P(v)}(X_w)$ of the Schubert variety X_w at the point $P(v)B_G^+/B_G^+$ is equal to the number of reduced type C pipe dreams in $PD_{\bar{v},w}^C$. See [43, Fact 5.1] for the analog of this observation in the type A setting.

We note that other combinatorial formulas for multiplicities in type C in special cases can be found in the works [3, 14, 22] and [32]. For example, in the recent paper [3], Anderson, Ikeda, Jeon, and Kawago provide a combinatorial formula for the multiplicity of a singularity of a covexillary Schubert variety (in a classical-type flag variety) in terms of excited Young diagrams.

Our next goal is to give a formula for the K -polynomial of $R_{\bar{v}}/I_{\bar{v},w}$ in terms of type C pipe dreams.

Proposition 7.4. *Let ρ be the reduced pipe dream for v associated to the factorization $\bar{v} = u_l v_{\square} u_r$. The K -polynomial of $R_{\bar{v}}/I_{\bar{v},w}$ is*

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \sum_{\sigma \in PD_{\bar{v},w}^C} (-1)^{\text{cr}(\sigma_L) - \ell(w_0w)} \prod_{\substack{(i,j) \text{ is a} \\ \text{cross in } \sigma_L}} (1 - (u_i \odot t_i^{-1})(u_j \odot t_j^{-1})),$$

where $\text{cr}(\sigma_L)$ is the number of crosses in σ_L , and the sum is over the nonboundary faces of $PD_{\bar{v},w}^C$.

Note that all factorizations of v yield the same K -polynomial.

Proof. By [29, Theorem 4.1], the K -polynomial of the Stanley–Reisner ideal I of the subword complex $S(Q, w_0w)$ is

$$\mathcal{K}(R/I; \mathbf{t}) = \sum_F (-1)^{|Q \setminus F| - \ell(w_0w)} \prod_{i \notin F} (1 - \mathbf{t}^{\deg(i)})$$

where the sum is over the faces F of $S(Q, w)$ such that $\mathbf{d}(Q \setminus F) = w_0w$. This is the same index set as in the proposition by [29, Theorem 3.7]. By Lemma 7.1, $PD_{\bar{v},w}^C$ is isomorphic to $\Delta_{\bar{v},w}$, which, in turn, is a relabeling of a subword complex $S(Q, w_0w)$, as in Section 5.2. We obtain the expression in the proposition by translating the equation above into the language of type C pipe dream complexes. Explicitly, for σ the pipe dream in $PD_{\bar{v},w}^C$ associated to the face F in $S(Q, w_0w)$:

- σ is a pipe dream for w if and only if $\mathbf{d}(Q \setminus F) = w_0w$,
- $|Q \setminus F| = \text{cr}(\sigma_L)$, and
- by Proposition 4.18, if (i, j) is a cross in σ_L , then $\mathbf{t}^{\deg(i,j)} = (u_i \odot t_i^{-1})(u_j \odot t_j^{-1})$.

□

We can also give a formula without the signs of Proposition 7.4. As noted in the previous proof, $PD_{\bar{v},w}^C$ is isomorphic to the subword complex $S(Q, w_0w)$, which is a shellable simplicial complex [29, Theorem 2.5]. That is, there exists a *shelling order* $F_1, \dots, F_m$ of the facets of $PD_{\bar{v},w}^C$, namely, an order such that, for each $i = 2, \dots, m$, the intersection of F_i with $\bigcap_{j < i} F_j$ is pure of dimension $\dim(PD_{\bar{v},w}^C) - 1$.

Proposition 7.5. *With notation as in Proposition 7.4, let $F_1, \dots, F_m$ be a shelling order for $PD_{\bar{v},w}^C$. Then, the K -polynomial of $R_{\bar{v}}/I_{\bar{v},w}$ is*

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \sum_{k=1}^m \prod_{\substack{(i,j) \text{ is a} \\ \text{cross in } (F_k)_L}} (1 - (u_i \odot t_i^{-1})(u_j \odot t_j^{-1})) \prod_{(i,j) \in \text{Abs}(F_k)} ((u_i \odot t_i^{-1})(u_j \odot t_j^{-1})),$$

where $\text{Abs}(F_k)$ is the set of elbows in $(i, j) \in F_k$ such that $F_k \setminus (i, j) \subset F_\ell$ for some $\ell < k$.

The corresponding result in [29] is Theorem 4.4, which uses the language of subword complexes and describes $\text{Abs}(F_k)$ differently: see [29, Remark 4.5].

Proof. This follows from Proposition 7.4 by collecting, for each $k = 1, \dots, m$, the summands indexed by faces σ of the form $\sigma = \bigcap_{\ell \in L} F_\ell$ where $k = \max L$, as described in the ungraded case in [38, Proposition 2.3]. □

In the case in which $v \geq_R v_\square$, we can always take a factorization $\bar{v}$ with $u_i = \text{id}$, which simplifies the appearance of the product in the formula in Proposition 7.4. A similar simplification can be written down for Proposition 7.5.

Corollary 7.6. For $v \geq_R v_{\square}$ the K -polynomial of $R_{\bar{v}}/I_{\bar{v},w}$ is

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = \sum_{\sigma \in PD_{\bar{v},w}^C} (-1)^{|\sigma_L| - \ell(w_0 w)} \prod_{(i,j) \in \sigma_L} (1 - t_i^{-1} t_j^{-1}).$$

The following example shows that this type C K -polynomial is not simply obtained from the type A K -polynomial by substituting the weights as suggested by the embedding of maximal tori from $Sp_{2n}(\mathbb{K})$ to $GL_{2n}(\mathbb{K})$.

Example 7.7. Let $v = v_{\square} = 2143, w = 3412$ in C_2 . Notice that the only pipe dream for w in $PD_{\bar{v},w}^C$ is



and therefore,

$$\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t}) = 1 - t_1^{-1} t_2^{-1}.$$

We now compare this to the type A K -polynomial using the embedding of tori above. To avoid confusion, let us denote the variables of the K -polynomials of type A Kazhdan–Lusztig varieties by $t_1^A, t_2^A, t_3^A, t_4^A$, where t_i^A corresponds to the i -th entry of the torus T consisting of diagonal matrices in $GL_{2n}(\mathbb{K})$. The pipe dream pictured above is also the only pipe dream for w in $PD_{v,w}^A$. By the type A version of [29, Theorem 4.1], we have that the K -polynomial of the type A Kazhdan–Lusztig variety associated to v, w is $(1 - t_3^A (t_1^A)^{-1})(1 - t_4^A (t_2^A)^{-1})$. The embedding of the torus in $Sp_4(\mathbb{K})$ is given by the substitution

$$t_1^A = t_2, \quad t_2^A = t_1, \quad t_3^A = t_1^{-1}, \quad t_4^A = t_2^{-1}.$$

Therefore, $\mathcal{K}(R_{\bar{v}}/I_{\bar{v},w}; \mathbf{t})$ is not the substitution of the type A K -polynomial.

7.3 | Combinatorics of type C pipe dreams

In this section, we discuss the combinatorics of type C pipe dreams. We begin by showing that we can recognize type C pipe dreams for w by following pseudolines, mirroring type A pipe dreams.

Proposition 7.8. A type C pipe dream is a pipe dream for w if and only if when we transport labels $1, \dots, 2n$ from the north and west sides of the picture along the pseudolines, ignoring all crossings subsequent to the first between each pair of pseudolines, the resulting labels on the south side read w .

Proof. Let ρ be a type C pipe dream for w , meaning that its associated word $Q_C(\rho) = (\alpha_1, \dots, \alpha_\ell)$ is a word with type C Demazure product $w_0 w$. Let Q^A be the word obtained by replacing each α_k with either one or two entries as follows:

$$\alpha_i \mapsto \begin{cases} n & \text{if } \alpha_k = 0, \\ n - \alpha_k, n + \alpha_k & \text{if } \alpha_k \neq 0. \end{cases} \quad (8)$$




FIGURE 11 Partway through applying commutation relations, in the proof of Proposition 7.8.

Note that if α_k corresponds to a cross of ρ in position (p, q) of $D(v_\square)$, then $\alpha_k = p - q$, and as a type A pipe dream, ρ has crosses at positions $(n + p, q)$ and $(n + q, p)$.

If $Q' = (\alpha_1, \dots, \alpha_j, \alpha_{j+1}, \dots, \alpha_k)$, $Q'' = (\alpha_1, \dots, \alpha_{j+1}, \alpha_j, \dots, \alpha_k)$, and $|\alpha_{j+1} - \alpha_j| \geq 2$, then $\mathbf{d}(Q') = \mathbf{d}(Q'')$; this is known as the *commutation relation* for the Demazure product.

Our goal is to use commutation relations to transform Q^A into the word $Q_A(\rho)$. Note then that the “only if” part of the proposition will follow from the combinatorics of type A pipe dreams.

The first entry of $Q_A(\rho)$ is $n + q - p = n - \alpha_1$, which is also the first entry of Q^A . Now consider $n - \alpha_k$ and suppose that for $i = 1, \dots, k - 1$, we have used the commutation relation to move $n - \alpha_i$ in Q^A to the correct position in $Q_A(\rho)$. We wish to move $n - \alpha_k = n + q - p$ to the position in $Q_A(\rho)$ associated to the cross in position $(n + q, p)$. If $\alpha_k = 0$, this is already the case, so let us suppose that $\alpha_k \neq 0$. We are allowed to use the commutation relation as long as we do not encounter $n + q - p + 1$ or $n + q - p - 1$. In Figure 11, the blue boxes represent the positions of the crosses that contribute $n + q - p \pm 1$ to $Q_A(\rho)$. For $i = 1, \dots, k - 1$ an $n - \alpha_i$ corresponding to a cross in the gray-shaded region has been moved to the correct position in $Q_A(\rho)$. Note that all the blue boxes outside the gray region correspond to entries in Q^A that appear after $n \pm \alpha_k$. Therefore, we can use commutation relations to move $n - \alpha_k$ to the left until it reaches the correct position in $Q_A(\rho)$. Continuing with this process, we transform Q^A into $Q_A(\rho)$ using only commutation relations.

Now let ρ be a type C pipe dream such that following the pseudolines yields w . Let $Q_A(\rho)$ be the word associated to ρ considered as a type A pipe dream. By undoing the process described above, we can transform $Q_A(\rho)$ into Q^A and lastly to $Q_C(\rho)$ by undoing the substitution (8). We leave the details to the reader. $\square$

8 | BEYOND SMALL PATCHES

In [28], Knutson showed that the defining ideal of any Kazhdan–Lusztig variety has a Gröbner basis whose leading terms are squarefree, and, in [27], he determined that the resulting initial ideal is the Stanley–Reisner ideal of the subword complex $S(Q, w_0 w)$, where Q is a reduced word for $w_0 v$. For small patches, our coordinates agree up to sign with the *Bott–Samelson coordinates* in [28], and our monomial order $<_{\text{lex}}$ agrees with the monomial order in [28]. Thus, Theorem 4.15 shows that the type C essential minors are a Gröbner basis in Knutson’s setup. (Knutson does not

provide a Gröbner basis in [28].) In this short section, we show that things are more mysterious beyond the case of small patches as the essential minors are *not* typically a Gröbner basis.

Example 8.1. Let $v = 231\,645$ as in Example 3.7. Observe that $Q = (0, 1, 0, 2, 1, 0, 2)$ is a reduced word for w_0v . Let

$$C_0^{(i)} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_i & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad C_1^{(i)} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & b_i & -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & b_i & 1 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad C_2^{(i)} = \begin{bmatrix} c_i & -1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & c_i & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 \end{bmatrix},$$

so that in *Bott–Samelson* coordinates, the opposite cell associated to v is identified with the space of matrices

$$w_0 C_0^{(1)} C_1^{(1)} C_0^{(2)} C_2^{(1)} C_1^{(2)} C_0^{(3)} C_2^{(2)} = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ c_2 & -1 & 0 & 0 & 0 & 0 \\ c_2 a_1 - b_1 & -a_1 & c_1 & 0 & 1 & 0 \\ c_2 b_1 - a_2 & -b_1 & b_2 & 0 & c_2 & 1 \\ c_2 c_1 - b_2 & -c_1 & a_3 & -1 & 0 & 0 \end{bmatrix},$$

where each a_i, b_i, c_i can take arbitrary values in $\mathbb{K}$. Let $w = 462\,513$. Then, now treating a_i, b_i, c_i as indeterminates, we see that the type C essential set (see Definition 3.10) is $\{(5, 1), (5, 3)\}$, and so, the type C essential minors are:

$$\mathcal{E} = \left\{ c_2 c_1 - b_2, \quad c_2 b_1 - a_2, \quad 2 \times 2 \text{ minors of } \begin{bmatrix} c_2 b_1 - a_2 & -b_1 & b_2 \\ c_2 c_1 - b_2 & -c_1 & a_3 \end{bmatrix} \right\}.$$

Using the lexicographic monomial order $c_2 > a_3 > b_2 > c_1 > a_2 > b_1 > a_1$, which is compatible with the vertex decomposition of the subword complex $S(Q, w_0w)$ described in Section 5.1, we see that

- the initial ideal of the Kazhdan–Lusztig ideal $\langle \mathcal{E} \rangle$ is the Stanley–Reisner ideal of $S(Q, w_0w)$ as expected, yet
- $\mathcal{E}$ is *not* a Gröbner basis.

Nevertheless, the set of type A essential minors is a Gröbner basis. There also exists a Gröbner basis consisting of type C essential minors that differ from the conventions introduced in Section 5, namely, the minors given by choosing the essential boxes $\{(5, 1), (3, 3)\}$.

In the next example, we see that unlike in the previous example, the type A essential minors need not be a Gröbner basis either.

Example 8.2. Consider $v = 213\,465$ so that $Q = (0, 1, 0, 2, 1, 0, 2, 1)$ is a reduced word for w_0v . Let $C_0^{(i)}, C_1^{(i)}, C_2^{(i)}$ as in the previous example. Then, the opposite Schubert cell associated to v is

identified with the space of matrices

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ c_2 & -b_3 & 1 & 0 & 0 & 0 \\ c_2 a_1 - b_1 & -b_3 a_1 + c_1 & a_1 & -1 & 0 & 0 \\ c_2 b_1 - a_2 & -b_3 b_1 + b_2 & b_1 & -c_2 & 0 & 1 \\ c_2 c_1 - b_2 & -b_3 c_1 + a_3 & c_1 & -b_3 & -1 & 0 \end{bmatrix}, \quad (9)$$

where $a_i, b_i, c_i \in \mathbb{K}$. Let $w = 632\,541$. There is a unique (type A or type C) essential box $\{(4, 3)\}$. If we treat a_i, b_i, c_i as indeterminates, the essential minors are then the 2×2 minors of the south-west 3×3 submatrix of (9). For the lexicographic monomial order with $b_3 > c_2 > a_3 > b_2 > c_1 > a_2 > b_1 > a_1$, we see that the ideal generated by the essential minors has initial ideal equal to the Stanley–Reisner ideal of $S(Q, w_0 w)$, yet the set of essential minors is *not* a Gröbner basis.

Consequently, it is still an open problem to find combinatorially defined Gröbner basis for type C Kazhdan–Lusztig ideals $\mathcal{N}_{v,w}$ when $v \not\leq v_\square$ in left–right weak order.

In type B, our methods fail because the analog of [33, Proposition 6.1.1.2] does not hold scheme-theoretically. In some cases, imposing the determinantal equations on the patches yields nonreduced schemes. If we apply our methods to type B small patches, we see that we end up taking the determinants of some skew-symmetric matrices, and the obvious solution to this problem is to take the pfaffians of those skew-symmetric matrices instead. We expect that results similar to ours can be proven with this modification in that case.

Beyond small patches in type B, some unpublished preliminary work of the fourth author and Alexander Yong suggests that the appropriate equations would still form a Gröbner basis under a diagonal term order, but we need to impose rank conditions on some submatrices that are non-trivially similar to a skew-symmetric one. Note that all formulas for the pfaffian require knowing the basis with respect to which a matrix is skew-symmetric, and we were not able to systematically determine the change of basis that turned these “secretly skew-symmetric” matrices into actually skew-symmetric matrices. The following example illustrates some of the difficulties.

Example 8.3. In this example, we work with the type B Weyl group. We embed B_4 into the symmetric group S_9 via $b_0 = s_4 s_5 s_4$, $b_1 = s_3 s_6$, $b_2 = s_2 s_7$, and $b_3 = s_1 s_8$. Consider $v = 13\,2456\,879 \in S_9$ and observe that $Q = (0, 1, 2, 3, 0, 1, 2, 3, 0, 1, 2, 0, 1, 2, 0, 1, 2, 3)$ is a reduced word for $w_0 v$. Let

$$B_0^{(i)} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{2}a_i^2 & a_i & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_i & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad B_1^{(i)} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & b_i & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & b_i & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$B_2^{(i)} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & c_i & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & c_i & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad B_3^{(i)} = \begin{bmatrix} d_i & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & d_i & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{bmatrix}.$$

Then, using these Bott–Samelson coordinates, the opposite cell associated to v is identified with the space of matrices $M_Q = w_0 B_0^{(1)} B_1^{(1)} B_2^{(1)} B_3^{(1)} B_0^{(2)} B_1^{(2)} B_2^{(2)} B_3^{(2)} B_0^{(3)} B_1^{(3)} B_2^{(3)} B_0^{(4)} B_1^{(4)} B_2^{(4)} B_3^{(4)}$.

Let $w = 381\,654\,927 \in S_9$. The type B essential set for w (which can be calculated in a similar way to the type C essential set; see also [2]) consists of three boxes in locations (7, 6), (7, 8), and (9, 6). Taking the associated essential minors of M_Q , where a_i, b_i, c_i, d_i appearing in M_Q are considered as indeterminates, we obtain an ideal generated by 107 minors. Using Macaulay2 [18], we see that this ideal can be presented as:

$$I = \langle d_1, c_2, b_4, b_3, a_4, c_3 c_4 - d_2 d_4, c_1 c_4 - b_1 d_4, a_3 c_4 - a_2 d_4, b_1 c_3 - c_1 d_2, a_2 c_3 - a_3 d_2, \\ a_3 b_1 - a_2 c_1, a_2 a_3 d_4 + 2 b_2 d_4, a_2^2 d_4 + 2 b_2 c_4, a_3^2 d_2 + 2 b_2 c_3, a_2 a_3 d_2 + 2 b_2 d_2, a_2 a_3 c_1 + 2 b_2 c_1, \\ a_2^2 c_1 + 2 b_1 b_2, a_2 a_3 b_2 + 2 b_2^2, a_2 a_3^2 + 2 a_3 b_2, a_2^2 a_3 + 2 a_2 b_2 \rangle.$$

This ideal is not radical, and hence, does not scheme-theoretically define a Kazhdan–Lusztig variety. So, in particular, the type B analog of Proposition 4.12 is false. We note that radical of the ideal I above is:

$$\langle d_1, c_2, b_4, b_3, a_4, c_3 c_4 - d_2 d_4, c_1 c_4 - b_1 d_4, a_3 c_4 - a_2 d_4, b_1 c_3 - c_1 d_2, a_2 c_3 - a_3 d_2, a_3 b_1 - a_2 c_1, \\ a_2 a_3 + 2 b_2 \rangle.$$

ACKNOWLEDGMENTS

We thank Patricia Klein, Allen Knutson, and Alex Yong for helpful discussions, and the Fields Institute for hosting us at the Thematic Program on Combinatorial Algebraic Geometry in Fall 2016, where this project was conceived. We also thank anonymous referees and Emmanuel Neye for helpful feedback on earlier versions of this paper. LE was partially supported by NSF Grants DMS 1855598 and DMS 2142656. JR was partially supported by NSERC Grant RGPIN-2017-05732. AW was partially supported by Simons Collaboration Grant 359792. AF received funding from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 792432.

JOURNAL INFORMATION

The *Journal of the London Mathematical Society* is wholly owned and managed by the London Mathematical Society, a not-for-profit Charity registered with the UK Charity Commission. All surplus income from its publishing programme is used to support mathematicians and mathematics research in the form of research grants, conference grants, prizes, initiatives for early career researchers and the promotion of mathematics.

ORCID

Alex Fink  <https://orcid.org/0000-0003-1562-4695>

REFERENCES

1. H. H. Andersen, J. C. Jantzen, and W. Soergel, *Representations of quantum groups at a p th root of unity and of semisimple groups in characteristic p : independence of p* , *Astérisque* **220** (1994), 321.
2. D. Anderson, *Diagrams and essential sets for signed permutations*, *Electron. J. Combin.* **25** (2018), no. 3, Paper 3.46, 23.
3. D. Anderson, T. Ikeda, M. Jeon, and R. Kawago, *The multiplicity of a singularity in a vexillary Schubert variety*, Preprint, arXiv:2112.07375, 2021.
4. N. Bergeron and S. Billey, *RC-graphs and Schubert polynomials*, *Experiment. Math.* **2** (1993), no. 4, 257–269.
5. S. Billey and M. Haiman, *Schubert polynomials for the classical groups*, *J. Amer. Math. Soc.* **8** (1995), no. 2, 443–482.
6. S. C. Billey, *Kostant polynomials and the cohomology ring for G/B* , *Duke Math. J.* **96** (1999), no. 1, 205–224.
7. S. C. Billey, W. Jockusch, and R. P. Stanley, *Some combinatorial properties of Schubert polynomials*, *J. Algebraic Combin.* **2** (1993), no. 4, 345–374.
8. A. S. Buch and R. Rimányi, *Specializations of Grothendieck polynomials*, *C. R. Math. Acad. Sci. Paris* **339** (2004), no. 1, 1–4.
9. A. Fink, J. Rajchgot, and S. Sullivant, *Matrix Schubert varieties and Gaussian conditional independence models*, *J. Algebraic Combin.* **44** (2016), no. 4, 1009–1046.
10. S. Fomin and A. N. Kirillov, *The Yang-Baxter equation, symmetric functions, and Schubert polynomials*, *Proceedings of the 5th Conference on Formal Power Series and Algebraic Combinatorics (Florence, 1993)*, vol. 153, 1996, pp. 123–143.
11. W. Fulton, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, *Duke Math. J.* **65** (1992), no. 3, 381–420.
12. W. Fulton, *Young tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, Cambridge, 1997. With applications to representation theory and geometry.
13. W. Fulton and P. Pragacz, *Schubert varieties and degeneracy loci*, *Lecture Notes in Mathematics*, vol. 1689, Springer, Berlin, 1998. Appendix J by the authors in collaboration with I. Ciocan-Fontanine.
14. S. R. Ghorpade and K. N. Raghavan, *Hilbert functions of points on Schubert varieties in the symplectic Grassmannian*, *Trans. Amer. Math. Soc.* **358** (2006), no. 12, 5401–5423.
15. E. Gorla, J. C. Migliore, and U. Nagel, *Gröbner bases via linkage*, *J. Algebra* **384** (2013), 110–134.
16. E. Gorla, *Symmetric ladders and G -biliaison*, *Liaison, Schottky problem and invariant theory*, *Progr. Math.*, vol. 280, Birkhäuser Verlag, Basel, 2010, pp. 49–62.
17. W. Graham, *Equivariant K -theory and Schubert varieties*, 2002. Unpublished.
18. D. R. Grayson and M. E. Stillman, *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2/>.
19. Z. Hamaker, E. Marberg, and B. Pawlowski, *Involution pipe dreams*, *Sém. Lothar. Combin.* **82B** (2020), Art. 63, 12.
20. F. Harary, *The determinant of the adjacency matrix of a graph*, *SIAM Rev.* **4** (1962), 202–210.
21. T. Ikeda, L. C. Mihalea, and H. Naruse, *Double Schubert polynomials for the classical groups*, *Adv. Math.* **226** (2011), no. 1, 840–886.
22. T. Ikeda and H. Naruse, *Excited Young diagrams and equivariant Schubert calculus*, *Trans. Amer. Math. Soc.* **361** (2009), no. 10, 5193–5221.
23. J. C. Jantzen, *Representations of algebraic groups*, *Mathematical Surveys and Monographs*, 2nd ed., vol. 107, American Mathematical Society, Providence, RI, 2003.
24. D. Kazhdan and G. Lusztig, *Representations of Coxeter groups and Hecke algebras*, *Invent. Math.* **53** (1979), no. 2, 165–184.
25. A. N. Kirillov, *On double Schubert and Grothendieck polynomials for classical groups*, Preprint, arXiv:1504.01469, 2015.
26. A. N. Kirillov and H. Naruse, *Construction of double Grothendieck polynomials of classical types using id Coxeter algebras*, *Tokyo J. Math.* **39** (2017), no. 3, 695–728.

27. A. Knutson, *Schubert patches degenerate to subword complexes*, Transform. Groups **13** (2008), no. 3–4, 715–726.
28. A. Knutson, *Frobenius splitting, point-counting, and degeneration*, Preprint, arXiv:0911.4941, 2009.
29. A. Knutson and E. Miller, *Subword complexes in Coxeter groups*, Adv. Math. **184** (2004), no. 1, 161–176.
30. A. Knutson and E. Miller, *Gröbner geometry of Schubert polynomials*, Ann. of Math. (2) **161** (2005), no. 3, 1245–1318.
31. B. Kostant and S. Kumar, *T-equivariant K-theory of generalized flag varieties*, J. Differential Geom. **32** (1990), no. 2, 549–603.
32. V. Kreiman, *Schubert classes in the equivariant K-theory and equivariant cohomology of the Lagrangian Grassmannian*, Preprint, arXiv:0602245, 2006.
33. V. Lakshmibai and K. N. Raghavan, *Standard monomial theory*, Encyclopaedia of Mathematical Sciences, vol. 137, Springer, Berlin, 2008. Invariant theoretic approach, Invariant Theory and Algebraic Transformation Groups, 8.
34. E. Marberg and B. Pawłowski, *K-theory formulas for orthogonal and symplectic orbit closures*, Adv. Math. **372** (2020), 107299, 43.
35. E. Miller and B. Sturmfels, *Combinatorial commutative algebra*, Graduate Texts in Mathematics, vol. 227, Springer, New York, 2005.
36. H. Sachs, *Über Teiler, Faktoren und charakteristische Polynome von Graphen. I*, Wiss. Z. Tech. Hochsch. Ilmenau **12** (1966), 7–12.
37. E. Smirnov and A. Tutubalina, *Pipe dreams for Schubert polynomials of the classical groups*, Preprint, arXiv:2009.14120, 2020.
38. R. P. Stanley, *Combinatorics and commutative algebra*, 2nd ed., Progress in Mathematics, vol. 41, Birkhäuser Boston, Inc., Boston, MA, 1996.
39. J. Tymoczko, *Billey's formula in combinatorics, geometry, and topology*, Schubert calculus—Osaka 2012, Adv. Stud. Pure Math., vol. 71, Math. Soc., Japan, Tokyo, 2016, pp. 499–518.
40. M. Willems, *Cohomologie et K-théorie équivariantes des variétés de Bott-Samelson et des variétés de drapeaux*, Bull. Soc. Math. France **132** (2004), no. 4, 569–589.
41. A. Woo, *Hultman elements for the hyperoctahedral groups*, Electron. J. Combin. **25** (2018), no. 2, Paper 2.41, 25.
42. A. Woo and A. Yong, *Governing singularities of Schubert varieties*, J. Algebra **320** (2008), no. 2, 495–520.
43. A. Woo and A. Yong, *A Gröbner basis for Kazhdan-Lusztig ideals*, Amer. J. Math. **134** (2012), no. 4, 1089–1137.