

**Modulation of Kinase Activities *In Vitro* by Hepatitis C Virus Protease NS3/NS4A
Mediated-Cleavage of Key Immune Modulator Kinases**

Mohd Amir F. Abdullah¹, Sarah M. McWhirter^{2,&}, and Zucai Suo^{1,3,*}

¹Department of Biochemistry, The Ohio State University, Columbus, OH 43210, U.S.A;

²Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA 02138,
U.S.A.

³Department of Biomedical Sciences, Florida State University College of Medicine, Tallahassee,
FL 32306, U.S.A

Running Title: Regulation of host kinase activities by hepatitis C protease complex

[&]Present address: Vice President, Head of Biology, Lycia Therapeutics, 400 East Jamie Court,
Suite 200, South San Francisco, CA 94080

*To whom correspondence should be addressed: Dr. Zucai Suo, Department of Biomedical
Sciences, Florida State University College of Medicine, Tallahassee, FL 32306, USA; Tel.: (850)
645-2501; E-mail: zucai.suo@med.fsu.edu

Key words: IKK α , IKK β , IKK ϵ , TBK1, HCV protease, kinase regulatory domain.

Abbreviations: HCV, hepatitis C virus; IKK α , Inhibitor of κ B kinase α ; I κ B α , Inhibitor of NF- κ B α ; IRF, interferon regulatory factor; MALDI-TOF, Matrix-Assisted Laser Desorption/Ionization Time-Of-Flight mass spectrometry; Nano-LC MS/MS, capillary-liquid chromatography tandem mass spectrometry; NF- κ B, nuclear factor- κ B; NS, nonstructural proteins; TBK1, TANK-binding kinase 1.

ABSTRACT

Hepatitis C Virus NS3/NS4A, a serine protease complex, has been found to interact with many host proteins and cause various adverse effects on cellular function and immune response. For example, the cleavage of important immune factors by NS3/NS4A has been suggested as a mechanism for hepatitis C virus to evade innate immunity. The spectrum of susceptible substrates for NS3/NS4A cleavage certainly includes important immune modulator kinases like IKK α , IKK β , IKK ϵ , and TBK1, as demonstrated in this paper. We show that the kinase activities of these four host kinases were transformed in unexpected ways by NS3/NS4A. Treatment with NS3/NS4A caused significant reduction of the kinase activities of both IKK α and IKK β , suggesting that HCV might use its NS3/NS4A protease activity to deactivate the NF- κ B-associated innate immune responses. In contrast, the kinase activities of both IKK ϵ and TBK1 were enhanced after NS3/NS4A treatment, and more strikingly, the enhancement was more than 10-fold within 20 min of treatment. Our mass spectroscopic results suggested that the cleavage after Cys89 in the kinase domain of IKK ϵ by NS3/NS4A led to their higher kinase activities, and three potential mechanisms were discussed. The observed kinase activity enhancement might facilitate the activation of both IKK ϵ - and TBK1-dependent cellular antiviral pathways, likely contributing to spontaneous clearance of the virus and observed acute HCV infection. After longer than 20 min cleavage, both IKK ϵ - and TBK1 gradually lost their kinase activities and the relevant antiviral pathways were expected to be inactivated, facilitating the establishment of chronic HCV infection.

INTRODUCTION

Liver diseases caused by hepatitis C virus (HCV) are a serious health problem. About 30% of HCV infection is acute infection since the infected people can clear the virus within 6 months of infection without any treatment. The remaining 70% of HCV infection is chronic infection that may persist for decades. It is estimated that 10% to 20% of chronically infected people will develop liver cirrhosis, while 1% to 5% will develop hepatocellular carcinoma [1]. Currently, an estimated 58 million people worldwide have chronic HCV infection, with about 1.5 million new infections occurring per year. Due to availability of cost-effective and efficacious generic direct-acting antiviral agents (DAAs), the World Health Organization (WHO) has set up an ambitious target of HCV elimination by 2030 and many countries are working towards the target [2-4].

The HCV genome is a plus-stranded RNA of about 10 kb in length, and its organization is similar to that of members of the family Flaviviridae [5-7]. Based on sequence analysis, seven major HCV genotypes have been identified and each of them has sub-genotypes [8]. The genome encodes a single precursor protein that includes structural and nonstructural proteins [6,9]. The precursor protein is proteolytically processed by both host signal peptidases and viral proteases to produce at least 10 viral proteins: Core, E1, E2, p7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B [9-11]. For example, the NS4A/NS4B, NS4B/NS5A, and NS5A/NS5B junctions are cleaved *in trans* by a serine protease complex, NS3/NS4A [12] while the NS3/NS4A junction is cleaved *in cis* by NS3 [13]. The NS3 protein is a multidomain protein of about 70 kDa. The amino-terminal third of NS3 forms a chymotrypsin-like serine protease domain [12,14,15] while the carboxyl-terminal two-third possesses both ATPase and RNA helicase activities [16,17]. Based on the analysis of 1,568 HCV NS3 sequences derived from patients infected with HCV (genotypes 1 to 6), 85 of 181 amino acid residues of the NS3 protease domain possess less than 1% variability [18]. The consensus sequence alignment demonstrates that the catalytic triad residues His57, Asp81, and Ser139 for the NS3 protease [19] is conserved across all HCV genotypes. NS4A, a 7-kDa and 54 amino acid residue protein, acts as a cofactor of the NS3 protease. The N-terminal 20 residues of NS4A are predicted to form a hydrophobic trans-membrane α -helix [8, 14], which presumably inserts into the ER membrane to anchor the HCV replicase [8]. The NS4A residues 22-31 (consensus sequence: SVVIVGRIIL) constitute the hydrophobic core, which interacts with two β -strands of the NS3 protease domain [7-13][19]. In the absence of NS4A, the NS3 protease activity decreases significantly, *e.g.* a ~950-fold decrease

at the cleavage of the NS4A/NS4B junction [11]. After several decades of development of direct acting anti-HCV drugs, quite a few small molecule inhibitors, targeting NS3/NS4A protease, NS5B RNA polymerase, and NS5A, have been approved by the Food and Drug Administration since 2011 and their various combinations have been widely used as therapies to treat HCV infection with good tolerability and greater than 90% sustained virological response [20,21].

Intracellularly, NS3 has been shown to bind to the catalytic subunit of protein kinase A (PKA) which prevents the nuclear translocation of this subunit and affects many PKA functions [22]. The activation of oncogene products via cleavage by the NS3/NS4A protease has been suggested as a possible role in the development of hepatocellular carcinoma by HCV [23]. NS3 is also shown to have oncogenic activity in NIH 3T3 mouse fibroblast cells [23], rat fibroblast cells [24], and human hepatocyte line [25]. Additionally, NS3 and HCV core have been found to activate inflammatory pathways in monocytes via toll-like receptors [26].

Interestingly, the protease activity of NS3/NS4A has been implicated in the inhibition of cellular antiviral pathways, leading to persistent HCV infections. Foy *et al.* first demonstrated that the NS3/4A protease blocks the phosphorylation and effector action of interferon regulatory factor-3 (IRF-3) [27]. They subsequently found that NS3/NS4A depressed retinoic acid-inducible gene I (RIG-I) signaling [28]. Meanwhile, Li *et al.* discovered that NS3/NS4A cleaves toll-like receptor 3 (TLR3) adaptor protein, TRIF [29]. Moreover, two other groups discovered that NS3/4A cleaves an adaptor protein in the RIG-I antiviral pathway, resulting in the shutting down of RIG-I-dependent antiviral responses [30,31].

In addition to these signaling molecules, we hypothesized that NS3/NS4A may cleave other proteins involved in innate immunity and antiviral pathways, *e.g.* cellular kinases such as Inhibitor of κB kinase α (IKK α), IKK β , IKK ϵ , and TBK1. These kinases are known to phosphorylate I κ B proteins to activate transcription factors nuclear factor-κB (NF-κB), IRF-3, and IRF-7 [32,33], which are central to cellular immune and inflammatory responses and cell survival in higher eukaryotes [34]. Specifically, the canonical NF-κB signaling pathway is mediated by IKK β while the non-canonical pathway depends on IKK α [35]. IKK β is primarily responsible for the phosphorylation of specific residues of prototypical I κ B proteins (I κ B α , I κ B β , and I κ B ϵ) bound to NF-κB as well as an atypical I κ B protein, p105, which is the precursor of the NF-κB subunit p50 [36]. IKK α phosphorylates specific residues of p100 to convert it to the NF-κB subunit p52 [35]. IKK ϵ and TBK1 directly phosphorylate IRF3 and IRF7 and such phosphorylation promotes the dimerization and nuclear translocation of these transcription

factors that stimulate production of type I interferons [37]. IKK ϵ and TBK1 have also been found to play a role in oncogenic transformation and thereby become novel drug targets with applications in the treatment of cancer and a variety of inflammatory diseases including rheumatoid arthritis and obesity-related metabolic disorders [38]. To confirm our hypothesis, we performed *in vitro* cleavage of these kinases by recombinant NS3/NS4A. Our results demonstrate that NS3/NS4A can cleave and regulate the kinase activities of IKK α , IKK β , IKK ϵ , and TBK1.

EXPERIMENTAL PROCEDURES

Peptide Synthesis. All peptides utilized in this study were chemically synthesized and HPLC purified by Peptides International Inc. (KY, USA). All synthetic peptides were confirmed to be the desired ones by analyses of amino acid composition and electron spray mass spectrometry. Their purities exceeded 95%, and they were stored frozen at -80 °C as 10 mM stock solutions in deionized water or 100% dimethyl sulfoxide.

Protein Expression and Purification of HCV NS3

The N-terminal domain (residues 1-182) of the NS3 protein is as efficient as the full-length NS3 protein in proteolytic activity [39]. We have cloned the N-terminal protease domain and expressed the protein fragment in *Escherichia coli* BL21 (DE3) Gold (Novagen). The truncated NS3 protein also contains a C-terminus His₆ tag to facilitate the purification process. The cells were induced with 0.4 mM IPTG at room temperature to increase the solubility of NS3 in the cytoplasmic extract [40]. The NS3 was purified through a Ni-NTA affinity column, a heparin sepharose column, and finally a Mono S 10/10 cation exchange column [40-43]. After concentration, the protein was flash-frozen with liquid nitrogen before storage at - 80°C. SDS-PAGE of the purified NS3 showed a single protein band at 20.1 kDa (Fig. 1) indicating that the protein was highly purified. Concentration of the purified NS3 protease domain was measured spectrophotometrically at 280 nm using the calculated molar extinction coefficient of 17,990 M⁻¹cm⁻¹.

Purification of Recombinant Human IKK α , IKK β , IKK ϵ , and TBK 1

IKK α , IKK β , and IKK ϵ were purified as described in [44,45]. Flag-TBK1 was cloned into pFastBac (Life Technologies). Recombinant bacmids and baculovirus were prepared according to the manufacturer's instructions. Flag-TBK1 was isolated from baculovirus-infected Sf9 cells and purified using M2-agarose Affinity Gel (Sigma) equilibrated with 50 mM Tris, pH 8.0, 250 mM NaCl, 0.5 mM NaF, 0.5 mM DTT, 0.1% Tween-20, and 10% Glycerol, and eluting it with 0.34 mg/ml flag peptide (Sigma) in the same buffer. The concentrations of the kinases were determined using the Bradford assay (Bio-Rad) using BSA as a standard.

Kinase Digestion Condition for Western Blot Analysis

Purified NS3 (6.4 μ M) was preincubated with 94 μ M of the NS4A synthetic peptide (KKKGSVVIVGRIILS) (Peptides International Inc., KY, USA) on ice for 40 min in the 1X digestion buffer (50 mM HEPES, pH 7.8, 0.5 mM EDTA, 25% glycerol, 10 mM DTT, 0.25 M KCl, 0.05 M NaCl, 0.2% CHAPS) to allow maximum interaction between NS3 and its cofactor NS4A peptide. The affinity of NS4A peptide to NS3 is lower in 20% glycerol-containing buffers than in solution containing 50% glycerol [46]. Water and kinases (1.2 μ g) were added to a final reaction volume of 25 μ l. The reaction was incubated at 23 °C for 29 h. The reactions were stopped by adding SDS loading dye and heated at 95 °C for 3 min and electrophoresed on an 8% SDS-PAGE gel. The separated proteins were transferred to a PVDF membrane, and immunoblotted with the following antibodies: anti-IKK α (sc7606, Santa Cruz Biotechnology), anti-IKK β (IMG-159, Imgenex Corp.), anti-TBK1 (IMG-139, Imgenex Corp.), and anti-IKK ϵ [45].

MALDI-TOF and Nano-LC MS/MS Analysis

Purified NS3 (6.4 μ M) was preincubated with 134 μ M NS4A synthetic peptide on ice for 40 min in the 1X digestion buffer to allow maximum interaction between NS3 and its cofactor NS4A peptide. Water and IKK ϵ (8 μ g) were added to a final reaction volume of 25 μ l. The reaction was incubated at 23 °C for 30 min. The reactions were stopped by adding SDS loading dye and heated at 95 °C for 3 min and electrophoresed on an 8% SDS-PAGE gel. The ~70-kDa cleavage product of IKK ϵ separated on the SDS-PAGE gel was *in gel* digested with trypsin and analyzed by Matrix-Assisted Laser Desorption/Ionization Time-Of-Flight mass spectrometry (MALDI-TOF) and Capillary-liquid chromatography tandem mass spectrometry (Nano-LC MS/MS) at the CCIC Mass Spectrometry and Proteomics Facility of The Ohio State University.

Kinase Digestion Conditions for Pre-steady-state Kinetic Assays

To achieve maximal interaction between NS3 and the NS4A peptide, high concentrations of NS4A peptide was used. Previously, the binding affinity of NS3 to NS4A was estimated at about 20 μ M [47]. Based on this result, we estimated that about 92 % of NS3 will form a protease complex with NS4A if we use 5 μ M NS3 and 244 μ M NS4A. We performed the single turnover kinetic assay using the same buffer condition as in the digestion above. Digestions were performed on IKK ϵ at 0.1 and 0.5 μ M concentrations, which are 50- and 10-fold less than the

NS3/NS4A concentration. At different digestion durations, a small aliquot was removed from the reaction and added to an excess amount of gel loading buffer and frozen on dry ice/methanol bath to stop the reaction. The reaction was electrophoresed on a 5% SDS-PAGE gel, transferred onto a PVDF membrane, and the remaining IKK ϵ substrate was detected using anti-IKK ϵ antibodies. The remaining IKK ϵ concentration was plotted versus time to obtain an exponential decay curve. The curve was fitted to a single exponential equation:

$$[\text{kinase}] = c + a * \exp(-k_{\text{obs}}t) \quad \text{Eq. 1}$$

The observed cleavage rate constant (k_{obs}) and a constant c were yielded. The equilibrium dissociation constant (K_d) of the binding between a kinase and NS3/NS4A was estimated based on the following equation:

$$K_d = c * (E_t - c) / (S_0 - c) \quad \text{Eq. 2}$$

where S_0 is initial kinase concentration, E_t is total concentration of the NS3/NS4A complex, and c is assumed to be the concentration of unbound kinase.

Kinase Digestion Condition for Kinase Activity Assay

Purified NS3 (1.25 μM) was preincubated with NS4A peptide (325 μM) on ice for 15 min followed by 15 min at 23 $^{\circ}\text{C}$ in the 5X concentrated digestion buffer. A kinase (0.5 μg) was added to a final reaction volume of 50 μl and incubated at 23 $^{\circ}\text{C}$. Aliquots (5 μl each) were removed from the reaction at 1, 4, 8, 16, 32, 60, 300, 900, 1200 min and flash-frozen in a dry ice/methanol bath and kept in storage at - 80 $^{\circ}\text{C}$ until use. A control reaction without the NS3/NS4A was also performed to test the stability of the kinases in the digestion buffer for the duration of the digestion.

Protease Inhibition Assay

Pefabloc SC (final concentration 2 mg/ml, Roche), was added at the beginning of the 15 min preincubation of NS3 and NS4A at 23 $^{\circ}\text{C}$ (before addition of IKK ϵ) or after the reaction was completed (prior to the flash-freezing step). Four reaction conditions were set up: i) IKK ϵ without NS3/NS4A; ii) IKK ϵ with NS3/NS4A; iii) IKK ϵ with NS3/NS4A pre-treated with Pefabloc; and iv) IKK ϵ with NS3/NS4A (Pefabloc was added after 16 min of digestion). The reactions were performed at 23 $^{\circ}\text{C}$ for 16 min, and the reaction was stopped by flash-freezing in

dry ice/methanol bath. Kinase activity of the treated IKK ϵ were measured using GST-I κ B α (residues 5-55) as substrate.

Kinase Activity Assay

Kinase activity assays were performed by incubating 1.5 μ l of the digested kinases with 10 μ l cocktail (20 mM HEPES, pH 7.6, 50 mM NaCl, 20 mM beta-glycerol phosphate, 1 mM sodium vanadate, 10 mM MgCl₂, 1 mM DTT, 0.1 mg/ml substrate (GST-I κ B α (residues 5-55), GST-I κ B α mutant S32A/S36A), 0.1 mM ATP, 0.015 μ Ci/ μ l [γ -³²P]ATP) at 23 °C for 30 min. The reactions were stopped by adding SDS loading dye. The samples were separated by SDS-PAGE, the gel was stained with Coomassie blue staining solution. Band intensities were determined to ensure equal sample loading. The gel was dried and then quantitated with a PhosphorImager 445 SI (Molecular Dynamics). Radioactive intensities were determined from the bands (normalized to the amount of substrate detected in the gel from the Coomassie stain) to measure kinase activity. Relative kinase activity was determined by calculating the ratio of the kinase activity of NS3/NS4A-treated kinase over the control kinase (in digestion buffer without NS3/NS4A) from the same digestion duration.

RESULTS

IKK α , IKK β , IKK ϵ , and TBK1 are substrates for NS3/NS4A protease in vitro

The consensus sequence of (D/E)XXXXC(A/S) (X, any amino acid residue; the scissile bond, between Cys and Ala or Ser) has been determined for all *trans* cleavage sites (Table 1) in viral processing of HCV polyprotein precursor by NS3/NS4A [9,48]. Interestingly, protein sequence alignments in Table 1 suggest that IKK α , IKK β , IKK ϵ , and TBK1 all contain potential NS3/NS4A cleavage sites which are similar in sequence to those junctions in the HCV polyprotein precursor. To examine whether NS3/NS4A can cleave these kinases, a recombinant NS3 protease domain (residues 1-182) containing a C-terminal hexahistidine tag was chromatographically purified from *E. coli* with at least 95% purity (Fig. 1). Since the NS4A residues 22-31 (SVVIVGRIIL) constitute the hydrophobic core, which binds to and activates NS3 [7-13], a synthetic peptide with the sequence of KKGSVVIVGRIILSGK (the underlined residues are from the consensus HCV NS4A sequence while the three Lys residues were added to increase the aqueous solubility of the peptide) was used as the NS4A cofactor in all *in vitro* digestion assays in this paper. To examine the protease activity of the NS3/NS4A complex, an internally quenched fluorogenic substrate (sequence: Ac-Asp-Glu-Asp(Edans)-Glu-Glu-Abu ψ [COO]-Ala-Ser-Lys(Dabcyl)-NH₂; Abu, 2-aminobutyric acid; Ac, acetylation; Dabcyl, 4-[[4'-(dimethylamino)phenyl]azo]benzoic acid; Edans, 5-[(2'-aminoethyl)amino]naphthalenesulfonic acid; ψ [COO], the ester bond between Abu and *L*-(+)-lactic acid) [49] was used as the substrate. After cleavage of the peptide bond Abu ψ [COO]-Ala by NS3/NS4A, the fluorescence of the donor (Edans) increased (data not shown). The steady-state protease activity ($355,000\text{ M}^{-1}\text{s}^{-1}$) of the NS3/NS4A complex was close to $345,000\text{ M}^{-1}\text{s}^{-1}$ measured by Taliani *et al.* [49], indicating that our assembled NS3/NS4A complex was active. To investigate if the steady-state protease activity was affected by the molar ratio of NS3/NS4A, 2 μM of the internally quenched fluorogenic peptide substrate was added to a preincubated solution of 20 nM NS3 and the NS4A peptide with a concentration varied from 800 nM to 6 μM for 0 to 3 min at 23 °C. The steady-state protease activity measured based on the time-dependent donor fluorescence change was comparable (data not shown), suggesting that the substrate cleavage kinetics was not significantly influenced by the change of the molar ratio of NS3/NS4A from 40- to 300-fold.

The four human kinases were expressed and purified from baculovirus infected insect cells (Experimental Procedures). Fig. 2 shows that IKK α , IKK β , IKK ϵ , and TBK1 proteins were

susceptible to degradation catalyzed by NS3/NS4A. To our knowledge, this is a first report of NS3/NS4A having direct proteolytic effect on these human kinases. Notably, IKK α and IKK β were cleaved into several small fragments after 29 hours at 23 °C while one large cleavage product was dominant in the degradation reactions of IKK ϵ and TBK1, e.g. the ~70-kDa cleavage product of IKK ϵ (the band immediately below the intact IKK ϵ protein band). As a negative control reaction, Dpo4, a 40-kDa Y-Family DNA polymerase from *Sulfolobus solfataricus* [50-52] that does not contain the NS3/NS4A cleavage sites (Table 1) was not cleaved by NS3/NS4A (data not shown). This indicates that NS3/NS4A did not randomly cleave proteins in our reaction conditions. When the ~70-kDa cleavage product of IKK ϵ was sequenced through *in gel* trypsin digestion and then mass spectrometric analysis (MALDI-TOF and Nano-LC MS/MS, see Experimental Procedures), we discovered that peptide coverage were from residue 118 (cleavage after Arg117 by trypsin) to the C-terminus of IKK ϵ (data not shown). This suggested that the two predicted NS3/NS4A cleavage sites (after Cys509, and Cys626) in IKK ϵ (Table 1) were inaccessible to the protease during degradation. Cleavage after Cys89 (the third predicted NS3/NS4A cleavage site in IKK ϵ) would produce a large ~70-kDa peptide, corresponding to the cleavage product observed in Fig. 2, and a small ~15-kDa peptide that was not observed in Fig. 2. The reason why residues 90-117 were not identified by our mass spectrometric analysis was not clear. Similarly, cleavage of TBK1 at the predicted NS3/NS4A cleavage site (after Cys267, Table 1) would produce a large ~50-kDa peptide (corresponding to the cleavage product observed in Fig. 2) and a smaller ~30-kDa peptide (not observed in Fig. 2). The generation of the ~50-kDa peptide was confirmed through MALDI-TOF and Nano-LC MS/MS analysis of the large TBK1 cleavage product, which covered residue 268 to the C-terminus of TBK1 (data not shown).

Estimation of the substrate specificity of IKK ϵ by pre-steady-state kinetic methods

To examine if NS3/NS4A cleaves these four kinases with reasonable efficiency and to avoid the complication of product inhibition, we measured the cleavage efficiency of IKK ϵ at 23 °C under single-turnover conditions. At least 10-fold excess of NS3/NS4A over IKK ϵ was used in the kinetic assay. As a kinetic strategy, the single-turnover kinetic assay was employed to estimate the cleavage rate of a human kinase by NS3/NS4A. It does not mean that the single-turnover reaction conditions would be certainly encountered *in vivo*. The single-turnover assay was

performed by using 5.0 μM NS3/NS4A with one of two different IKK ϵ concentrations and the data were fit into a single exponential equation, Eq. 1 (see Experimental Procedures). The observed cleavage rate constants (k_{obs}) at IKK ϵ concentrations of 0.1 and 0.5 μM were determined to be 0.0075 and 0.0145 s^{-1} , respectively (Fig. 3). Due to lack of enough protein substrate IKK ϵ , we did not determine the substrate concentration dependence of k_{obs} which can yield the maximum k_p and the equilibrium dissociation constant K_d for the binding of IKK ϵ to NS3/NS4A. However, we can estimate the K_d from these time courses shown in Fig. 3. By assuming the constant in Eq. 1 being the concentration of unbound IKK ϵ in a cleavage reaction, we estimated the K_d to be 4.57 and 1.68 μM in the presence of 0.1 and 0.5 μM IKK ϵ , respectively (see Experimental Procedures). These low micromolar K_d values indicate the binding between the protease complex NS3/NS4A and IKK ϵ is moderately strong. The substrate specificity k_{obs}/K_d for the degradation of 0.1 and 0.5 μM IKK ϵ was estimated to be 1,641 and 8,631 $\text{M}^{-1}\text{s}^{-1}$, respectively. Previously, steady-state kinetic analysis has been used to estimate the cleavage efficiency (k_{cat}/K_m) of peptide substrates representing the viral NS4A/NS4B (1,600 $\text{M}^{-1}\text{s}^{-1}$) and NS4B/NS5A (110 $\text{M}^{-1}\text{s}^{-1}$) cleavage sites catalyzed by NS3/NS4A at 30 $^{\circ}\text{C}$ [53]. These k_{obs}/K_d and k_{cat}/K_m values suggest that IKK ϵ is as good a substrate, if not better, to NS3/NS4A as those HCV junction-mimicking substrates. Thus, HCV NS3/NS4A protease is capable of cleaving IKK ϵ , and possibly IKK α , IKK β , and TBK1, *in vivo*.

NS3/NS4A modulates kinase activity

We expected that proteolytic digestion of these four kinases by NS3/NS4A would reduce their respective kinase activities. This assay is more sensitive than the detection of cleavage products since changes in the kinase activity will reflect the physiological changes on the kinases. Since we are only measuring the kinase activity and not the cleavage products, much lower concentrations of NS3 (1.25 μM) and the NS4A peptide (325 μM) were used. Each of those kinases was reacted with NS3/NS4A at 23 $^{\circ}\text{C}$ for 35 min and the reaction was stopped by flash-freezing the reaction in dry ice/methanol bath. The treated kinases (with or without NS3/NS4A) were then tested for their kinase activities to phosphorylate GST-I κ B α (residues 5-55). Fig. 4 shows that the kinase activities of IKK α and IKK β were reduced after treatment with NS3/NS4A, but surprisingly, the kinase activities of IKK ϵ and TBK1 increased after treatment with NS3/NS4A. Moreover, the NS3/NS4A treatment also increased TBK1 phosphorylation of

mutant GST-I κ B α (residues 5-55) containing Ser32Ala/Ser36Ala, a substrate that was normally not phosphorylated by TBK1, IKK α , IKK β , and IKK ϵ . This suggests the substrate specificity of TBK1 was altered by NS3/NS4A.

To test the stability of the kinases in the NS3/NS4A digestion buffer condition, a time-course digestion assay was set up at 23 °C. In the control reactions (without NS3/NS4A), the duration of incubation from 1 min to 20 hours significantly reduced the kinase activity of IKK β (Fig. 5B), suggesting that IKK β was not stable. In contrast, the kinase activities of IKK ϵ (Fig. 6B) and TBK1 (Fig. 7B) were stable over the 20-hour incubation, indicating that IKK ϵ and TBK1 were more stable than IKK β . In Fig. 5C, treatment with NS3/NS4A further reduced the kinase activity of IKK β compared to undigested IKK β . Amazingly, treatment of IKK ϵ with NS3/NS4A increased kinase activity up to 11-fold within the first 16 min, and then gradually returned to the level of control IKK ϵ over a period of 1,200 min (Fig. 6C). Similarly, treatment of TBK1 with NS3/NS4A increased kinase activity up to 14-fold within the first 8 min, and then gradually returned to the level of control TBK1 over a period of 1,200 min (Fig. 7C). The cause of the initial increase in kinase activities of both IKK ϵ and TBK1 by NS3/NS4A may be due to the degradation of a kinase regulatory domain (see Discussion). As these two kinases were further degraded by NS3/NS4A, their overall kinase activities dropped significantly with time. After five hours of cleavage by NS3/NS4A, the remaining kinase activities of digested IKK ϵ and TBK1 were still slightly higher than those of undigested kinases.

The NS3/NS4A proteolytic activity is essential for modulation of the kinase activities of both IKK ϵ and TBK1

Enhancement of the kinase activities of IKK ϵ and TBK1 during cleavage could be due to either NS3/NS4A mediated cleavage or protein/protein interactions such as the binding of IKK ϵ to NS3/NS4A, or both. To examine these possibilities, we used Pefabloc SC, a broad-spectrum serine protease inhibitor, to covalently modify NS3 and permanently inhibit its serine protease activity. Fig. 8 shows that the IKK ϵ kinase activity was not affected by NS3/NS4A which was pretreated with Pefabloc SC, but untreated NS3/NS4A was able to enhance the kinase activity of IKK ϵ . Similar results were also observed with TBK1 (data not shown), suggesting the cleavage of these two kinases by NS3/NS4A was likely the reason for their kinase activity enhancement.

DISCUSSION

There are many reports about the interactions of HCV proteins with host cell factors involved in cell signaling, apoptosis, transcriptional regulation, transformation, membrane rearrangements, and immuno-modulation, which lead to dysregulation of cellular functions [54]. Among the HCV proteins, NS3 actively interacts with cellular proteins through either direct binding interactions, or its protease activity, to influence cellular processes such as triggering inflammation [26], affecting PKA activity [22], inhibiting cellular antiviral responses [29,55], and cell transformation [23-25]. It is likely that more host proteins will be found to interact with NS3 through new assays or methodologies.

In this paper, we used pre-steady-state kinetic methods to examine the cleavage of NS3/NS4A under single turnover reactions, in which molar excess of the protease complex was reacted with each of the four human kinases (IKK α , IKK β , IKK ϵ , and TBK1). The single turnover conditions ensured that most of the examined kinase were bound and cleaved by NS3/NS4A during the first turnover. In contrast, many reports used steady-state kinetic assays in which a small amount of NS3/NS4A was used to digest molar excess of a protein substrate (kinase) for long times [29,41,47,56]. The steady-state kinetic parameters including k_{cat} and K_M from these assays are difficult to be kinetically interpreted because they are complex functions of all reactions occurring at the enzyme active site over multiple turnovers [57-61]. To make it worse, the fact that NS3/NS4A has a tendency to be bound and inhibited by one of its proteolytic products [62] makes the interpretation of the steady-state kinetic parameters more challenging. Under these steady-state reaction conditions, the cleavage of a protein substrate by a protease like NS3/NS4A may appear to be slow because of the product inhibition. Furthermore, the detection of interactions between the NS3 protease domain with cellular protein targets will be difficult if their interactions are transient and weak [41,47,56]. With these factors in mind, we used 10-fold more NS3/NS4A (6.4 μ M) over the kinase substrate (0.6 μ M) to enhance the binding interaction between the protease and its protein substrate. We were able to demonstrate that the NS3/NS4A indeed cleaved IKK α , IKK β , IKK ϵ , and TBK1 which were expressed and purified from baculovirus-infected insect cells (Fig. 2). To further support the results from the cleavage assays, the kinases were first treated with a small amount of NS3/NS4A for various times and then assayed for their remaining kinase activities (Figs. 4-7). Surprisingly, the kinase activities of IKK ϵ and TBK1 were significantly enhanced by appropriate levels of NS3/NS4A

cleavage while the opposite results were observed with IKK α and IKK β (Figs. 4-7). In contrast to our observation, the *in vitro* cleavage of IKK ϵ and TBK1 by NS3/NS4A was not observed previously by Li *et al.* [29]. In their digestion assays, the kinases were synthesized from *in vitro* translation reactions, rather than purified from baculovirus-infected insect cells, the cleavage reaction buffer was different from ours and not optimized, and the molar ratio between NS3/NS4A and each kinase was unknown. In addition to these factors, the components in Flexi-Rabbit reticulocyte lysate may inhibit the cleavage of IKK ϵ and TBK1 by NS3/NS4A in their *in vitro* reactions [29].

Notably, contradicting results for the cleavage of IKK ϵ and TBK1 by NS3/NS4A in cell-based assays can be found in the literature. In one publication, neither IKK ϵ nor TBK1 is cleaved by NS3/NS4A when each of these kinases is individually co-transfected with NS3/NS4A into an osteosarcoma cell line UNS3-4A-24 [29]. Similarly, Breiman *et al.* [63] who also perform co-transfection of NS3/NS4A with each of these kinases but in a different cell line (HEK 293T cells) conclude that NS3/NS4A does not cleave IKK ϵ and TBK1. Notably, in their Fig. 2B [63], it is shown that in cells transfected with 5 μ g IKK ϵ DNA, the IRF-3P level (phosphorylation of IRF-3) is considerably higher in the presence than in the absence of NS3/NS4A. However, the cells that are transfected with less IKK ϵ DNA (0.2, 1, and 2 μ g) do not show any difference in the level of IRF-3 phosphorylation in the presence or absence of NS3/NS4A. To reanalyze their results based on our current knowledge, we would suggest that a higher level of IKK ϵ expression, resulted from 5 μ g IKK ϵ DNA transfection, increases the binding interaction between IKK ϵ and NS3/NS4A and subsequent cleavage of IKK ϵ by the protease, leading to the enhanced phosphorylation of IRF-3 in the cells. This scenario is likely considering that the binding affinity between IKK ϵ and the protease domain of NS3/NS4A was moderate with a K_d in low micromolar range (Fig. 3).

More interestingly, our data (Figs. 4-7) indicate that NS3/NS4A mediated-cleavage of cellular kinases significantly affected their kinase activities in a time-dependent manner. For example, the kinase activities of IKK α and IKK β were reduced by NS3/NS4A treatment (Figs. 4 and 5) and cannot phosphorylate inhibitor I κ B α , leading to a direct inhibition to NF- κ B activation *in vivo* [32,33]. This represents another mechanism of viral immune evasion. In stark contrast, cleavage of IKK ϵ and TBK1 by NS3/NS4A within the first few minutes, the kinase activities of IKK ϵ (Fig. 6) and TBK1 (Fig. 7) to phosphorylate I κ B α were enhanced more than

10-fold. We further showed that the enhancement in kinase activity in IKK ϵ by NS3/NS4A was dependent upon its proteolytic activity because pretreatment of NS3/NS4A with Pefabloc SC eliminated any effect of the protease on the kinase activity of IKK ϵ (Fig. 8). Pefabloc SC is a serine protease inhibitor that acylated highly conserved Ser139, one of the NS3 protease catalytic triad residues [19], leading to the inactivation of this protease activity. If occurring *in vivo*, the initial cleavage-caused kinase activity enhancement will facilitate the activation of both IKK ϵ - and TBK1-dependent cellular antiviral pathways, leading to enhanced immune responses to the early stage of HCV infection. This could be one of the reasons for many patients whose acute HCV-infection is controlled and rapidly cleared from the blood [64]. After NS3/NS4A cleavage for extended times, the kinase activities of IKK ϵ (Fig. 6) and TBK1 (Fig. 7), just like those of IKK α (Fig. 4) and IKK β (Fig. 5), were eventually eliminated. Thus, NS3/NS4A eventually allows HCV to evade cellular immune responses and to establish chronic infection.

At present, we can only speculate on how the cleavage of four canonical and noncanonical IKK kinases (IKK α , IKK β , IKK ϵ , and TBK1) by NS3/NS4A enhanced or abolished their kinase activities (Figs. 4-7). Human IKK kinases share 28% sequence identity and a similar domain structural organization with an N-terminal kinase domain (KD) followed by a ubiquitin-like domain (ULD), a helical scaffold dimerization domain (SDD), and a C-terminal domain (CTD) as illustrated by the solved structures of C-terminal truncated IKK α , IKK β , and TBK1 (Fig. 9) [38]. Notably, the structure of IKK ϵ has not been available yet. The kinase activities of IKK α and IKK β are regulated by phosphorylation on two serine residues within their classical kinase activation loops [37]. In comparison, the phosphorylation of a single serine residue in the activation loops of IKK ϵ and TBK1 activates the kinase activities of IKK ϵ and TBK1 [37]. Several predicted HCV NS3/NS4A cleavage sites (Table 1) were mapped onto the structures of truncated IKK α , IKK β , and TBK1 with activated KDs (Fig. 9). Based on the KDs of IKK ϵ and TBK1 [37,38,45,65] (Figs. 9C and 10), the sizes of the main cleavage products (Fig. 2), and our MALDI-TOF and Nano-LC MS/MS analysis of the protein sequence of the 70 kDa cleavage product of IKK ϵ (Fig. 2), cleavage after Cys89 in IKK ϵ and after Cys267 in TBK1 occurred within their respective KDs (see Results). The first possible scenario is that the kinase activation loop in the cleaved KD of TBK1 (or IKK ϵ) can be more efficiently trans-autophosphorylated for its kinase activation than the same kinase activation loop in intact TBK1 (or IKK ϵ), which is believed to be trans-autophosphorylated through higher-order oligomerization of TBK1 (or

IKK ϵ) dimers [38]. The second possible scenario is that their KDs were cleaved but were reassembled due to protein-protein interaction between different kinase cleavage products that remained in the same reaction solution. Cleavage of the KDs might cause these kinases to adopt conformations that were more catalytically efficient than the intact IKK ϵ and TBK1. The third possible scenario is that both IKK ϵ and TBK1 possess an autoinhibitory domain. The NS3/NS4A cleavage of these autoinhibitory domains led to up-regulation of the kinase activities of IKK ϵ (Fig. 6) and TBK1 (Fig. 7). Interestingly, both N- and C-terminal autoinhibitory domains have been identified in other kinase families, and these autoinhibitory domains usually lie outside the kinase catalytic domain [66,67]. Because the likely cleavage site of TBK1 (after Cys267) by NS3/NS4A is near the C-terminus of the KD (Figs. 9C and 10), we speculate that the autoinhibitory domain is within the C-terminus of the KD in TBK1. In comparison, such an autoinhibitory domain was expected to lie within the N-terminal 89 amino acid residues of IKK ϵ since this region was cleaved off IKK ϵ by NS3/NS4A and not observed in Fig. 2 (see above discussion). Future studies are needed to further investigate these three mechanistic scenarios regarding the correlation between the cleavage of IKK ϵ and TBK1 by HCV NS3/NS4A and the kinase activities of the two noncanonical IKK kinases.

Human IKK α and IKK β are highly homologous canonical IKK kinases [35]. Notably, HCV NS3/NS4A cleaved both IKK α and IKK β into several fragments (Fig. 2). One of the predicted cleavage sites in both IKK α and IKK β is in their respective kinase activation loops containing the mitogen-activated protein (MAP)-kinase kinase consensus motif SXXXXS, where X is any amino acid residue [68-71] (S176 and S180 in IKK α , S177 and S181 in IKK β , Table 1). The X-ray crystal and cryo-EM structures of human IKK α [35], and the X-ray crystal structures of human IKK β [72,73] show that their classical kinase activation loops protrude outside of their KDs before the phosphorylation of two serine residues and subsequent kinase activation (data not shown), and are well exposed to solvent even after serine phosphorylation (Figs. 9A and 9B). In other words, the kinase activation loops in IKK α and IKK β , regardless of their serine phosphorylation status, will be accessible by HCV NS3/NS4A for cleavage. Consistently, the kinase activation loops are predicted to be cleaved by NS3/NS4A right after C178 in IKK α and C179 in IKK β (Table 1) and these cystine residues are on the surface of activated IKK α and IKK β (Figs. 9A and 9B). Thus, we hypothesize that the observed time-dependent decreases in

the kinase activities of IKK α and IKK β (Figs. 4 and 5) were mainly due to the gradual cleavage of the kinase activation loops in these kinases by HCV NS3/NS4A. It is possible that the kinase activity decreases were also contributed through the NS3/NS4A cleavage at other predicated sites of the KDs of IKK α and IKK β (Table 1), *e.g.* the cleavage of the ULD of IKK α (Fig. 9A). The exact NS3/NS4A cleavage sites in IKK α and IKK β will be unambiguously identified through future sequencing of the isolated peptide fragments by mass spectrometry analysis.

Conclusions

In this paper, we discovered that the HCV protease complex of NS3/NS4A was able to cleave human IKK α , IKK β , IKK ϵ , and TBK1. The cleavage of IKK α and IKK β decreased their kinase activities in a time-dependent manner. Surprisingly, the short time cleavage of IKK ϵ and TBK1 enhanced their phosphorylation of the protein substrate I κ B α and the scenarios for such increase of the kinase activities were discussed. After long time cleavage by NS3/NS4A, IKK ϵ and TBK1 were cleaved into small peptides and their kinase activities were gradually decreased and eventually lost.

Acknowledgements

We thank Sha-Mei Liao from Millennium Pharmaceuticals, Inc. for providing TBK1 protein. S. M. M. thanks Tom Maniatis for financial support. This work was supported by National Institutes of Health Grant (R01GM122093) and National Science Foundation Grant (MCB-1716168) to Z.S.

Conflict of interest

The authors declare that they have no conflicts of interest with the contents of this article.

REFERENCES

- [1] J. Cohen, The Scientific Challenge of Hepatitis C, *Science* 285 (1999) 26.
- [2] H.C.V.C. Polaris Observatory, Global change in hepatitis C virus prevalence and cascade of care between 2015 and 2020: a modelling study, *Lancet Gastroenterol Hepatol* 7 (2022) 396-415. 10.1016/S2468-1253(21)00472-6.
- [3] Y. Waheed, Progress on global hepatitis elimination targets, *World J Gastroenterol* 27 (2021) 8199-8200. 10.3748/wjg.v27.i47.8199.
- [4] R.K. Dhiman, M. Premkumar, Hepatitis C Virus Elimination by 2030: Conquering Mount Improbable, *Clin Liver Dis (Hoboken)* 16 (2020) 254-261. 10.1002/cld.978.
- [5] A. Ishihama, K. Nagata, Viral RNA polymerase, *Crit. Rev. Biochem.* 23 (1988).
- [6] C.M. Rice, E.M. Lenes, S.R. Eddy, S.J. Shin, R.L. Sheets, J.H. Strauss, Nucleotide sequence of yellow fever virus: implication for flavivirus gene expression and evolution, *Science* 229 (1985) 726-735.
- [7] A. Takamizawa, C. Mori, L. Fuke, S. Manabe, S. Murakami, J. Fujita, E. Onishi, T. Andoh, I. Yoshida, H. Okayama, Structure and organization of the hepatitis C virus genome isolated from human carriers, *J. Virol.* 65 (1991) 1105-1113.
- [8] T. Nakano, G.M. Lau, G.M. Lau, M. Sugiyama, M. Mizokami, An updated analysis of hepatitis C virus genotypes and subtypes based on the complete coding region, *Liver Int* 32 (2012) 339-345. 10.1111/j.1478-3231.2011.02684.x.
- [9] A. Grakoui, D.W. McCourt, C. Wychowski, S.M. Feinstone, C.M. Rice, Characterization of the hepatitis C virus-encoded serine proteinase: determination of proteinase-dependent polyprotein cleavage sites, *J. Virol.* 67 (1993) 2832-2843.
- [10] C. Lin, B.D. Lindenbach, B.M. Prágai, D.W. McCourt, C.M. Rice, Processing in the hepatitis C virus E2-NS2 region: identification of p7 and two distinct E2-specific products with different termini, *J. Virol.* 68 (1994) 5063-5073.
- [11] H. Mizushima, M. Hijikata, S. Asabe, M. Hirota, K. Kimura, K. Shimotohno, Two hepatitis C virus glycoprotein E2 products with different C termini, *J. Virol.* 68 (1994) 6215-6222.
- [12] C. Failla, L. Tomei, R. De Francesco, Both NS3 and NS4A are required for proteolytic processing of hepatitis C virus nonstructural proteins, *J. Virol.* 68 (1994) 3753-3760.
- [13] L. Tomei, C. Failla, E. Santolini, R.D. Francesco, N.L. Monica, NS3 is a serine protease required for processing of hepatitis C virus polyprotein, *J. Virol.* 67 (1993) 4017-4026.
- [14] R. Bartenschlager, V. Lohmann, T. Wilkinson, J.O. Koch, Complex formation between the NS3 serine-type proteinase of the hepatitis C virus and NS4A and its importance for polyprotein maturation, *J. Virol.* 69 (1995) 7519-7528.
- [15] B. Hahm, D.S. Han, S.H. Back, O.-K. Song, M.-J. Cho, C.-J. Kim, K. Shimotohno, S.K. Jang, NS3-4A of hepatitis C virus is a chymotrypsin-like protease, *J. Virol.* 69 (1995) 2534-2539.
- [16] L. Jin, D.L. Peterson, Expression, isolation and characterization of the hepatitis C virus ATPase/RNA helicase, *Arch. Biochem. Biophys.* 323 (1995) 47-53.
- [17] D.W. Kim, Y. Gwack, J.H. Han, J. Choe, C-terminal domain of the hepatitis C virus NS3 protein contains an RNA helicase activity, *Biochem. Biophys. Res. Commun.* 215 (1995) 160-166.
- [18] V. Cento, C. Mirabelli, R. Salpini, S. Dimonte, A. Artese, G. Costa, F. Mercurio, V. Slicher, L. Parrotta, A. Bertoli, M. Ciotti, D. Di Paolo, C. Sarrecchia, M. Andreoni, S. Alcaro, M. Angelico, C.F. Perno, F. Ceccherini-Silberstein, HCV genotypes are differently prone to the

development of resistance to linear and macrocyclic protease inhibitors, PLoS One 7 (2012) e39652. 10.1371/journal.pone.0039652.

[19] N. Yao, P. Reichert, S.S. Taremi, W.W. Prosise, P.C. Weber, Molecular views of viral polyprotein processing revealed by the crystal structure of the hepatitis C virus bifunctional protease-helicase, *Structure* 7 (1999) 1353-1363. 10.1016/s0969-2126(00)80025-8.

[20] N.M. Ganta, G. Gedda, B. Rathnakar, M. Satyanarayana, B. Yamajala, M.J. Ahsan, S.S. Jadav, T. Balaraju, A review on HCV inhibitors: Significance of non-structural polyproteins, *Eur J Med Chem* 164 (2019) 576-601. 10.1016/j.ejmech.2018.12.045.

[21] T. Kish, A. Aziz, M. Sorio, Hepatitis C in a New Era: A Review of Current Therapies, *P T* 42 (2017) 316-329.

[22] P. Borowski, K. Oehlmann, M. Heiland, R. Laufs, Nonstructural Protein 3 of hepatitis C virus blocks the distribution of the free catalytic subunit of cyclic AMP-dependent protein kinase, *J. Virol.* 71 (1997) 2838-2843.

[23] D. Sakamuro, T. Furukawa, T. Takegami, Hepatitis C Virus Nonstructural Protein NS3 Transforms NIH 3T3 Cells, *J. Virol.* 69 (1995) 3893-3896.

[24] R. Zemel, S. Gerechet, H. Greif, L. Bachmatove, Y. Birk, A. Golan-Goldhirsh, M. Kunin, Y. Berdichevsky, I. Benhar, R. Tur-Kaspa, Cell transformation induced by hepatitis C virus NS3 serine protease, *J. Viral Hepatitis* 8 (2001) 96-102.

[25] D.-Y. Feng, Y. Sun, R.-X. Cheng, X.-M. Ouyang, H. Zheng, Effect of hepatitis C virus nonstructural protein NS3 on proliferation and MAPK phosphorylation of normal hepatocyte line, *World J. Gastroenterol.* 11 (2005) 2157-2161.

[26] A. Dolganiuc, S. Oak, K. Kodys, D.T. Golenbock, R.W. Finberg, E. Kurt-Jones, G. Szabo, Hepatitis C core and nonstructural 3 proteins trigger toll-like receptor 2-mediated pathways and inflammation activation, *Gastroenterobiology* 127 (2004) 1513-1524.

[27] E. Foy, K. Li, C. Wang, R.J. Sumpter, M. Ikeda, S.M. Lemon, M.J. Gale, Regulation of interferon regulatory factor-3 by the hepatitis C virus serine protease, *Science* 300 (2003) 1145-1148.

[28] E. Foy, K. Li, R. Sumpter, Jr., Y.M. Loo, C.L. Johnson, C. Wang, P.M. Fish, M. Yoneyama, T. Fujita, S.M. Lemon, M. Gale, Jr., Control of antiviral defenses through hepatitis C virus disruption of retinoic acid-inducible gene-I signaling, *Proc. Natl. Acad. Sci. U.S.A.* 102 (2005) 2986-2991.

[29] K. Li, E. Foy, J.C. Ferreon, M. Nakamura, A.C.M. Ferreon, M. Ikeda, S.C. Ray, M.J. Gale, S.M. Lemon, Immune evasion by hepatitis C virus NS3/4A protease-mediated cleavage of the Toll-like receptor 3 adaptor protein TRIF, *Proc. Natl. Acad. Sci. U.S.A.* 102 (2005) 2992-2997.

[30] X.-D. Li, L. Sun, R.B. Seth, G. Pineda, Z.J. Chen, Hepatitis C virus protease NS3/4A cleaves mitochondrial antiviral signaling protein off the mitochondria to evade innate immunity, *Proc. Natl. Acad. Sci. U.S.A.* 102 (2005) 17717-17722.

[31] E. Meylan, J. Curran, K. Hofmann, D. Moradpur, M. Binder, R. Bartenschlager, J. Tschopp, Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by hepatitis C virus, *Nature* 437 (2005) 1167-1172.

[32] A.A. Stephenson, D.J. Taggart, G. Xu, J.D. Fowler, H. Wu, Z. Suo, The inhibitor of kappaB kinase beta (IKKbeta) phosphorylates IkappaBalphalpha twice in a single binding event through a sequential mechanism, *J Biol Chem* (2022) 102796. 10.1016/j.jbc.2022.102796.

[33] G. Xu, Y.C. Lo, Q. Li, G. Napolitano, X. Wu, X. Jiang, M. Dreano, M. Karin, H. Wu, Crystal structure of inhibitor of kappaB kinase beta, *Nature* 472 (2011) 325-330. 10.1038/nature09853.

[34] C. Xiao, S. Ghosh, NF-kappaB, an evolutionarily conserved mediator of immune and inflammatory responses, *Advances in experimental medicine and biology* 560 (2005) 41-45. 10.1007/0-387-24180-9_5.

[35] S. Polley, D.O. Passos, D.B. Huang, M.C. Mulero, A. Mazumder, T. Biswas, I.M. Verma, D. Lyumkis, G. Ghosh, Structural Basis for the Activation of IKK1/alpha, *Cell Rep* 17 (2016) 1907-1914. 10.1016/j.celrep.2016.10.067.

[36] M. Karin, Y. Ben-Neriah, Phosphorylation meets ubiquitination: the control of NF-[kappa]B activity, *Annu Rev Immunol* 18 (2000) 621-663. 10.1146/annurev.immunol.18.1.621.

[37] X. Ma, E. Helgason, Q.T. Phung, C.L. Quan, R.S. Iyer, M.W. Lee, K.K. Bowman, M.A. Starovasnik, E.C. Dueber, Molecular basis of Tank-binding kinase 1 activation by transautophosphorylation, *Proc Natl Acad Sci U S A* 109 (2012) 9378-9383. 10.1073/pnas.1121552109.

[38] A. Larabi, J.M. Devos, S.L. Ng, M.H. Nanao, A. Round, T. Maniatis, D. Panne, Crystal structure and mechanism of activation of TANK-binding kinase 1, *Cell Rep* 3 (2013) 734-746. 10.1016/j.celrep.2013.01.034.

[39] C. Failla, L. Tomei, R. De Francesco, An amino-terminal domain of the hepatitis C virus NS3 protease is essential for interaction with NS4A, *J. Virol.* 69 (1995) 1769-1777.

[40] L. Zhang, J.A. Brown, S.A. Newmister, Z. Suo, Polymerization fidelity of a replicative DNA polymerase from the hyperthermophilic archaeon *Sulfolobus solfataricus* P2, *Biochemistry* 48 (2009) 7492-7501. 10.1021/bi900532w.

[41] C. Steinkuhler, A. Urbani, L. Tomei, G. Biasiol, M. Sardana, E. Bianchi, A. Pessi, R. De Francesco, Activity of purified hepatitis C virus protease NS3 on peptide substrates, *J. Virol.* 70 (1996) 6694-6700.

[42] J.H. Wong, J.A. Brown, Z. Suo, P. Blum, T. Nohmi, H. Ling, Structural insight into dynamic bypass of the major cisplatin-DNA adduct by Y-family polymerase Dpo4, *EMBO J* 29 (2010) 2059-2069. emboj2010101 [pii] 10.1038/emboj.2010.101.

[43] J.A. Brown, S.A. Newmister, K.A. Fiala, Z. Suo, Mechanism of double-base lesion bypass catalyzed by a Y-family DNA polymerase, *Nucleic Acids Res* 36 (2008) 3867-3878.

[44] F.S. Lee, R.T. Peters, L.C. Dang, T. Maniatis, MEKK1 activates both IκB kinase α , IκB kinase β , *Proc. Natl. Acad. Sci. U.S.A.* 95 (1998) 9319-9324.

[45] R.T. Peters, S.-M. Liao, T. Maniatis, IKK ϵ is part of a novel PMA-inducible IκB kinase complex, *Molecular Cell* 5 (2000) 513-522.

[46] E. Bianchi, A. Urbani, G. Biasiol, M. Brunetti, A. Pessi, R. De Francesco, C. Steinkuhler, Complex formation between the hepatitis C virus serine protease and a synthetic NS4A cofactor peptide, *Biochemistry* 36 (1997) 7890-7897.

[47] V.V. Sardana, J.T. Blue, J. Zugay-Murphy, M.K. Sardana, L.C. Kuo, An uniquely purified HCV NS3 protease and NS4A(21-34) peptide form a highly active serine protease complex in peptide hydrolysis, *Protein Expr. Purif.* 16 (1999) 440-447.

[48] E. Pizzi, A. Tramontano, L. Tomei, N. La Monica, C. Failla, M. Sardana, T. Wood, R. De Francesco, Molecular model of the specificity pocket of the hepatitis C virus protease: implications for substrate recognition, *Proc. Natl. Acad. Sci. U.S.A.* 91 (1994) 888-892.

[49] M. Taliani, E. Bianchi, F. Narjes, M. Fossatelli, A. Urbani, C. Steinkuhler, R. De Francesco, A. Pessi, A continuous assay of hepatitis C virus protease based on resonance energy transfer depsipeptide substrates, *Anal. Biochem.* 240 (1996) 60-67.

[50] K.A. Fiala, Z. Suo, Pre-Steady-State Kinetic Studies of the Fidelity of *Sulfolobus solfataricus* P2 DNA Polymerase IV, *Biochemistry* 43 (2004) 2106-2115.

[51] S.M. Sherrer, D.C. Beyer, C.X. Xia, J.D. Fowler, Z. Suo, Kinetic Basis of Sugar Selection by a Y-Family DNA Polymerase from *Sulfolobus solfataricus* P2, *Biochemistry* 49 (2010) 10179-10186. 10.1021/bi101465n.

[52] B.A. Maxwell, Z. Suo, Kinetic Basis for the Differing Response to an Oxidative Lesion by a Replicative and a Lesion Bypass DNA Polymerase from *Sulfolobus solfataricus*, *Biochemistry* 51 (2012) 3485-3496. 10.1021/bi300246r.

[53] J.A. Landro, S.A. Raybuck, Y.P. Luong, E.T. O'Malley, S.L. Harbeson, K.A. Morgenstern, G. Rao, D.J. Livingston, Mechanistic role of an NS4A peptide cofactor with the truncated NS3 protease of hepatitis C virus: elucidation of the NS4A stimulatory effect via kinetic analysis and inhibitor mapping, *Biochemistry* 36 (1997) 9340-9348.

[54] T.L. Tellinghuisen, C.M. Rice, Interaction between hepatitis C virus proteins and host cell factors, *Curr. Opin. Microbiol.* 5 (2002) 419-427.

[55] M. Otsuka, N. Kato, M. Moriyama, H. Taniguchi, Y. Wang, N. Dharel, T. Kawabe, M. Omata, Interaction Between the HCV NS3 Protein and the Host TBK1 Protein Leads to Inhibition of Cellular Antiviral Responses, *Hepatology* 41 (2005) 1004-1012.

[56] A. Urbani, E. Bianchi, F. Narjes, A. Tramontano, R. De Francesco, C. Steinkuhler, A. Pessi, Substrate specificity of the hepatitis C virus serine protease NS3, *J. Biol. Chem.* 272 (1997) 9204-9209.

[57] K.A. Johnson, Transient-state kinetic analysis of enzyme reaction pathways, in: D.S. Sigman (Ed.) *The Enzymes*, Academic Press, Inc. 1992, pp. 1-61.

[58] W.J. Zahurancik, S.J. Klein, Z. Suo, Kinetic Mechanism of DNA Polymerization Catalyzed by Human DNA Polymerase epsilon, *Biochemistry* 52 (2013) 7041-7049. 10.1021/bi400803v.

[59] K.A. Fiala, Z. Suo, Mechanism of DNA Polymerization Catalyzed by *Sulfolobus solfataricus* P2 DNA Polymerase IV, *Biochemistry* 43 (2004) 2116-2125.

[60] J.A. Brown, Z. Suo, Elucidating the kinetic mechanism of DNA polymerization catalyzed by *Sulfolobus solfataricus* P2 DNA polymerase B1, *Biochemistry* 48 (2009) 7502-7511. 10.1021/bi9005336.

[61] A. Kumar, A.J. Reed, W.J. Zahurancik, S.M. Daskalova, S.M. Hecht, Z. Suo, Interlocking activities of DNA polymerase beta in the base excision repair pathway, *Proc Natl Acad Sci U S A* 119 (2022) e2118940119. 10.1073/pnas.2118940119.

[62] C. Steinkuhler, G. Biasiol, M. Brunetti, A. Urbani, U. Koch, R. Cortese, A. Pessi, R. De Francesco, Product inhibition of the hepatitis C virus NS3 protease, *Biochemistry* 37 (1998) 8899-8905.

[63] A. Breiman, N. Grandvaux, R. Lin, C. Ottone, S. Akira, M. Yoneyama, T. Fujita, J. Hiscott, E.F. Meurs, Inhibition of RIG-I-dependent signaling to the interferon pathway during hepatitis C virus expression and restoration of signaling by IKKepsilon, *J. Virol.* 79 (2005) 3969-3978.

[64] J.H. Hoofnagle, Management of hepatitis C: current and future perspective, *J. Hepatol.* 31 (1999) 264-268.

[65] T. Shimada, T. Kawai, K. Takeda, M. Matsumoto, J. Inoue, Y. Tatsumi, A. Kanamaru, S. Akira, IKK-i, a novel lipopolysaccharide-inducible kinase that is related to IkappaB kinases, *Int. Immunol.* 11 (1999) 1357-1362.

[66] L.A. Cohen, J.-L. Guan, Residues within the first subdomain of the FERM-like domain in Focal Adhesion Kinase are important in its regulation, *J. Biol. Chem.* 280 (2005) 8197-8207.

[67] Y.L. Zu, Y. Ai, C.K. Huang, Characterization of an autoinhibitory domain in human mitogen-activated protein kinase-activated protein kinase 2, *J. Biol. Chem.* 270 (1995) 202-206.

[68] J.A. DiDonato, M. Hayakawa, D.M. Rothwarf, E. Zandi, M. Karin, A cytokine-responsive IkappaB kinase that activates the transcription factor NF-kappaB, *Nature* 388 (1997) 548-554. 10.1038/41493.

[69] F. Mercurio, H. Zhu, B.W. Murray, A. Shevchenko, B.L. Bennett, J. Li, D.B. Young, M. Barbosa, M. Mann, A. Manning, A. Rao, IKK-1 and IKK-2: cytokine-activated IkappaB kinases essential for NF-kappaB activation, *Science* 278 (1997) 860-866. 10.1126/science.278.5339.860.

[70] J.D. Woronicz, X. Gao, Z. Cao, M. Rothe, D.V. Goeddel, IkappaB kinase-beta: NF-kappaB activation and complex formation with IkappaB kinase-alpha and NIK, *Science* 278 (1997) 866-869. 10.1126/science.278.5339.866.

[71] E. Zandi, D.M. Rothwarf, M. Delhase, M. Hayakawa, M. Karin, The IkappaB kinase complex (IKK) contains two kinase subunits, IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappaB activation, *Cell* 91 (1997) 243-252. 10.1016/s0092-8674(00)80406-7.

[72] S. Liu, Y.R. Misquitta, A. Olland, M.A. Johnson, K.S. Kelleher, R. Kriz, L.L. Lin, M. Stahl, L. Mosyak, Crystal structure of a human IkappaB kinase beta asymmetric dimer, *J Biol Chem* 288 (2013) 22758-22767. 10.1074/jbc.M113.482596.

[73] S. Polley, D.B. Huang, A.V. Hauenstein, A.J. Fusco, X. Zhong, D. Vu, B. Schrofelbauer, Y. Kim, A. Hoffmann, I.M. Verma, G. Ghosh, T. Huxford, A structural basis for IkappaB kinase 2 activation via oligomerization-dependent trans auto-phosphorylation, *PLoS Biol* 11 (2013) e1001581. 10.1371/journal.pbio.1001581.

[74] W.L. DeLano, The PyMOL Molecular Graphics System. DeLano Scientific, San Carlos, CA, USA., PYMOL; DeLano Scientific: San Carlos, CA, (2002).

Table 1. Sequence alignment of known NS3/NS4A cleavage sites on HCV polyprotein precursor and putative NS3/NS4A cleavage sites on IKK α , IKK β , IKK ϵ , and TBK1. The conserved residues are in bold-face and the arrows depict the cleavage sites.

Proteins	Residue number of P6	P6 P5 P4 P3 P2 P1 ↓ P1' P2' P3' P4'
NS4A/NS4B		D E M E E C ↓ A S H L
NS4B/NS5A		D C S T P C ↓ S G S W
NS5A/NS5B		E D V V C C ↓ S M S Y
IKK α	173	D Q G S L C ↓ T S F V
	374	D G V R G C ↓ D S Y M
	691	D H S L S C ↓ V V T P
	711	E E N L N C ↓ L G H L
IKK β	174	D Q G S L C ↓ T S F V
	711	E A H N L C ↓ T L L E
IKK ϵ	84	L V M E Y C ↓ S S G S
	504	E V L S R C ↓ S Q N I
	621	L R L V G C ↓ S V A A
TBK1	262	D M P V S C ↓ S L S R

Figure Legends

Figure 1. SDS-PAGE of purified truncated HCV NS3 protease domain expressed in *E. coli*. The NS3 was purified through a Ni-NTA affinity column, followed by a heparin sepharose column, and finally a MonoS 10/10 cation exchange column. The left lane are protein size markers, and the right lane is purified NS3 (20 kDa).

Figure 2. Western blot analysis of *in vitro* digestion of recombinant human IKK α , IKK β , IKK ϵ , and TBK1 with NS3/NS4A. The kinases (1.2 μ g) were digested with NS3/NS4A (6.4 μ M NS3 + 94 μ M NS4A) at 23 °C for 29 hours. Arrows indicate bands of intact kinases.

Figure 3. Pre-steady-state kinetic analysis of the cleavage of IKK ϵ by NS3/NS4A (5 μ M NS3 + 244 μ M NS4A) at 23°C. The IKK ϵ concentration was 0.1 μ M (A) and 0.5 μ M (B). In each time course, the concentration of remaining IKK ϵ was plotted against time. The data were fit to Eq. 1 (see Experimental Procedures) to yield 0.0075 and 0.0145 s^{-1} for the observed cleavage rate constants in (A) and (B), respectively.

Figure 4. Effect of the proteolytic activity of NS3/NS4A on IKK α , IKK β , IKK ϵ and TBK1 kinase activities. The kinases (0.5 μ g) were treated with NS3/NS4A (1.25 μ M NS3 + 325 μ M NS4A) at 23 °C for 35 min. The kinase activities were tested using GST-I κ B α (residues 5-55) and [γ -³²P]ATP as substrates. Phosphorylated GST-I κ B α bands were detected by exposing to X-ray film. wt, wild type GST-I κ B α (residues 5-55); m, GST-I κ B α S32A/S36A mutant; no enz, reaction without kinase.

Figure 5. Effect of the proteolytic activity of NS3/NS4A on IKK β kinase activity in a time-dependent manner. The kinases (0.5 μ g) were treated with NS3/NS4A (1.25 μ M NS3 + 325 μ M NS4A). (A) IKK β was treated with or without NS3/4A for up to 20 h. The kinase activity was tested using GST-I κ B α (residues 5-55) and [γ -³²P]ATP as substrates. Phosphorylated GST-I κ B α bands were detected by a phosphoimager. (B) The band intensities detected by the phosphoimager were plotted versus time. (C) The relative kinase activities (IKK β treated with NS3/NS4A versus untreated control) were plotted versus time.

Figure 6. Effect of NS3/NS4A cleavage of IKK ϵ in a time-dependent manner. The kinases (0.5 μ g) were treated with NS3/NS4A (1.25 μ M NS3 + 325 μ M NS4A). (A) IKK ϵ was treated with or without NS3/4A up to 20 h. The kinase activity was tested using GST-I κ B α (residues 5-55) and [γ -³²P]ATP as substrates. Phosphorylated GST-I κ B α bands were detected by a phosphoimager. (B) The band intensities detected by the phosphoimager were plotted versus time. (C) The relative kinase activities (IKK ϵ treated with NS3/4A versus untreated control) were plotted versus time.

Figure 7. Effect of NS3/NS4A digestion on TBK1 kinase activity in a time-dependent manner. The kinases (0.5 μ g) were treated with NS3/NS4A (1.25 μ M NS3 + 325 μ M NS4A). (A) TBK1 was treated with or without NS3/4A up to 20 h. The kinase activity was tested using GST-I κ B α (residues 5-55) and [γ -³²P]ATP as substrates. Phosphorylated GST-I κ B α bands were detected by a phosphoimager. (B) The band intensities detected by the phosphoimager were plotted versus time. (C) The relative kinase activities (TBK1 treated with NS3/NS4A versus untreated control) were plotted versus time.

Figure 8. Effect of Pefabloc SC on the capability of NS3/NS4A to enhance the kinase activity of IKK ϵ . Pefabloc SC was added at the beginning of the 15 min preincubation of NS3 and NS4A (1.25 μ M NS3 + 325 μ M NS4A) at 23 °C (before addition of IKK ϵ) or after the reaction was completed (prior to the flash-freezing step). Four reaction conditions were set up: i) IKK ϵ not treated with NS3/NS4A; ii) IKK ϵ treated with NS3/NS4A; iii) IKK ϵ treated with NS3/NS4A (pre-treated with Pefabloc); and iv) IKK ϵ treated with NS3/NS4A (Pefabloc added after completion of digestion). The digestions were performed at 23 °C for 16 min, and the reaction was stopped by flash-freezing in dry ice/methanol bath. The kinase activities were measured using GST-I κ B α as substrate.

Figure 9. Ribbon diagram representation of the X-ray crystal structures of three truncated human IKK kinases generated by using PYMOL[74]. The N-terminal KD (salmon) including its activation loops (cyan), ULD (blue), and SDD (green) are shown while the CTD is missing. (A) IKK α (residues 6-660) (PDB: 5EBZ) [35]. The phosphoacceptor sites S176 and S180 of the

kinase activation loop (cyan) were mutated to glutamate to mimic their phosphorylation status. HCV NS3/NS4A was predicted to cleave after C178 (magenta, spheres) within the kinase action loop and C379 (magenta, spheres) in the ULD (Table 1). (B) IKK β (residues 1-671) complexed with a staurosporine analog inhibitor K252a (black) (PDB: 4KIK) [72]. The kinase activation loop (cyan) has three disordered residues (residues 174-176) and its phosphoacceptor sites (S177 and S181) were phosphorylated. NS3/NS4A was predicted to cleave after C179 (magenta, spheres) within the kinase action loop (Table 1). (C) TBK1 (residues 1-657) complexed with a specific inhibitor BX7 (black) (PDB: 4IW0) [38]. The kinase activation loop (cyan) was well ordered and its sole phosphoacceptor site S172 was phosphorylated. NS3/NS4A was predicted to cleave after C267 (magenta, spheres) within the KD (Table 1).

Figure 10. Alignment of protein sequences of human IKK ϵ and TBK1. The solid backgrounds indicate identical amino acid residues. Vertical brackets define the KD, while the horizontal bracket denotes the kinase activation loop. The heptad repeats of hydrophobic residues within the putative leucine zipper domain are indicated with “+” symbols. Two helices of the potential helix-loop-helix domain are overlined. These domains were predicted previously by Shimada *et al.* [65] and Peters *et al.* [45] and mostly validated by the crystal structures of human TBK1 [37,38]. The predicated NS3/NS4A cleavage sites on IKK ϵ and TBK1 (Table 1) are represented by arrow heads on top (IKK ϵ) or bottom (TBK1) of their protein sequences.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7





Figure 8




* NS3/4A was pretreated with Pefabloc.


** Pefabloc was added after digestion was completed.

Figure 9

(A)



(B)



(C)




Figure 10

