

MITOGENOME REPORT



The mitochondrial genome of the deep-sea pyramid urchin *Echinocrepis rostrata* (Echinoidea: Holasteroidea: Pourtalesiidae)

Matthew Stephenson, Avery S. Hiley , Greg W. Rouse  and Nicolás Mongiardino Koch 

Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA

ABSTRACT

We present the mitochondrial genome of the deep-sea, epibenthic, irregular echinoid *Echinocrepis rostrata*, representing the first sequenced mitogenome of the order Holasteroidea. The length of the complete *E. rostrata* mitochondrial genome is 15,716 base pairs, and its GC content is 34.87%. It contains 13 protein-coding genes, two rRNA genes, and 22 tRNA genes, whose order is identical to that of all other available echinoid mitogenomes. Phylogenetic analysis of available mitochondrial genomes, based on all coding loci, places *E. rostrata* as the sister group to spatangoids (heart urchins).

ARTICLE HISTORY

Received 23 November 2023
Accepted 15 March 2024

KEYWORDS

Echinoid; irregular urchins;
Irregularia; mitogenome;
phylogeny

Introduction

The genus *Echinocrepis* A. Agassiz, 1881 contains two species of irregular echinoids characterized by a uniquely high and pyramidal test (i.e. the main structure of the echinoid skeleton, which is generally either spherical or flattened in shape). The genus is part of the family Pourtalesiidae A. Agassiz, 1881 (Kroh and Mooi 2023), which includes exclusively deep sea (bathyal to abyssal) taxa that feed by plowing through the upper layers of sediment and ingesting detritus (Schultz 2017). *Echinocrepis cuneata* A. Agassiz 1878 is known only from two specimens collected in Antarctic waters by the H.M.S *Challenger* expeditions (Agassiz 1878). The remaining species, *Echinocrepis rostrata* Mironov 1973 (Figure 1A), is much better known, and is found across the North Pacific, from the Kamchatka Peninsula to the northwest of the Channel Islands in California, where it inhabits waters 3,315–5,020 m deep (Mironov 1973; Schultz 2017). *E. rostrata* is considered a deep-sea ‘bulldozer’ species, whose bioturbating activity plays an important role in determining the fate of the particulate organic carbon (POC) that reaches the ocean bottom. Its abundance and feeding ecology fluctuate with POC influx (Vardaro et al. 2009; Miguez-Salas et al. 2022) potentially affecting sediment geochemistry, nutrient availability in benthic ecosystems, and rates of carbon sequestration. Distinct color morphotypes of the species are known, yet molecular barcoding suggests these belong to a single species (Vardaro 2010).

Materials & methods

A specimen of *E. rostrata* was collected from the North Pacific Ocean (40.771504° N, 127.548792° W) during research expedition TN-403 by remotely operated vehicle (ROV) *Jason* aboard R/V *Thomas G. Thompson* (<https://www.rvdata.us/search/cruise/TN403>). No ethical approval was required for sampling. The specimen was collected on June 7th, 2022 from a depth of 3,232 m, slightly shallower than the reported depth range for the species. Upon arrival to the vessel, gonadal material was dissected, placed in 95% ethanol, and kept refrigerated. Ethanol-preserved test fragments and internal tissues are deposited in the Scripps Institution of Oceanography Benthic Invertebrate Collection (SIO-BIC; <https://scripps.ucsd.edu/benthic-invertebrate-collection>, contact: Greg Rouse, grouse@ucsd.edu) under voucher SIO-BIC E11460.

DNA was extracted using the Zymo Research DNA-Tissue Miniprep kit (Zymo Research, Irvine, CA) and sequenced on the Illumina Novaseq 6000 platform (Illumina, San Diego, CA) following library preparation by Novogene (en.novogene.com). Species identity was confirmed by comparing the COX1 sequence (extracted from the assembled mitogenome) with publicly available data.

Genomic raw reads were trimmed or excluded using quality scores with Trimmomatic v0.39 (Bolger et al. 2014) under default settings. Mitogenome assembly was performed with MitoFinder v1.4 (Allio et al. 2020). The Echinoderm and Flatworm Mitochondrial Code (NCBI; transl_table = 9) was specified for the translation of protein-coding genes (PCGs).

CONTACT Nicolás Mongiardino Koch  nmongiardinokoch@ucsd.edu  Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA

 Supplemental data for this article can be accessed online at <https://doi.org/10.1080/23802359.2024.2333572>.

© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.




Figure 1. A. Aboral (top) and lateral (bottom) views of *Echinocrepis rostrata* (SIO-BIC E4015). Anterior margin toward the left. Photo by Greg Rouse. B. Phylogenetic position of *E. rostrata* (highlighted; accession number OR494080) among major echinoid clades using partitioned maximum likelihood inference with an amino acid dataset including all 13 mitochondrial protein-coding loci. Node numbers correspond to ultrafast bootstrap support values. The following sequences were used: *Eucidaris tribuloides* NC_050252 (unpublished), *Echinothrix calamaris* NC_050274 (Wakayama et al. 2019), *Strongylocentrotus droebachiensis* NC_009940 (unpublished), *Astriclypeus mannii* NC_061414 (Shin et al. 2022), *Sinaechinocyamus mai* QFS00168 – GQF00180 (Lin et al. 2020), *Echinocardium cordatum* NC_013881 (Perseke et al. 2010), and *Nacospatangus alta* NC_023255 (unpublished).

Complete records for all echinoid mitogenomes publicly available on NCBI's RefSeq (downloaded May 18th, 2023) were provided as a reference file during assembly (listed in Table S1). The assembled mitogenome was annotated using the integrated MitoFinder pipeline with MEGAHIT v1.2.9 (Li et al. 2016) and ARWEN v1.2 (Laslett and Canbäck 2008), and a second annotation was obtained using the MITOS Web server (Bernt et al. 2013). Coverage was estimated by mapping trimmed reads to the assembled contig using BBDMap (part of BBTools v35.34, Bushnell 2014). Assemblies and annotations were imported into Geneious v11.1.5 (Kearse et al. 2012), which was used to circularize the mitogenome and manually edit annotations by comparing against the aforementioned echinoid mitogenomes.

Amino acid sequences for the 13 mitochondrial PCGs of *E. rostrata* were extracted and compiled with those of all four available irregular echinoid mitogenomes, as well as three regular echinoid outgroups. Individual genes were aligned using MAFFT v7.505 (Katoh and Standley 2013) and concatenated. A maximum likelihood tree was constructed with IQ-TREE v2.0.3 (Minh et al. 2020) using the best-fit partitioned model of evolution (Chernomor et al. 2016; Kalyaanamoorthy et al. 2017), which merged loci into two partitions. One thousand replicates of ultrafast bootstrap (Hoang et al. 2018) were used to determine node support values.

Results

Percent identities of the obtained COX1 sequence to those published by Vardaro (2010) ranged between 99.5% and 100% (GenBank accession numbers GU228453–GU228465). This material was collected from Station M, West Pacific (35.16° N, 123.02° W), approximately 780 km south of the

sampling locality of the specimen employed for mitogenome sequencing, and only 85 km away from the type locality of *E. rostrata* (Kroh and Mooi 2023), confirming the identity of the sequenced specimen.

A total of 13,376,922 paired-end reads of 150 base pairs (bp) were sequenced, of which 13,085,912 (97.8%) passed quality controls and were used for assembly. A single circular contig with an average coverage of 27.03x was obtained (Figure S1; 100 bp sliding window coverage range = 13.04–48.9). The mitochondrial genome of *E. rostrata* contains 15,716 bp and includes 13 PCGs, two rRNA genes, and 22 tRNA genes. The GC content is 34.87% and the control region (located between tRNA-Thr and tRNA-Pro) is 135 bp long. As in other echinoids, the ND4L coding sequence is likely initiated with an alternative codon (Quek et al. 2021). The mitochondrial gene order of *E. rostrata* (shown in Figure S2) is identical to that of all other known echinoid mitogenomes, which represents the ancestral gene order for Echinodermata (Perseke et al. 2010).

The result of the phylogenetic inference agrees with recent phylotranscriptomic studies (Mongiardino Koch et al. 2018, 2022), placing holasteroids as the sister-group to spatangoids (Figure 1B). Atelostomata, the clade defined by these two orders, is included within a monophyletic Irregularia, whose most closely-related lineage of regular echinoids is Echinacea (represented here by *Strongylocentrotus droebachiensis*).

Discussion

While a large number of echinoid mitogenomes are available, sequencing effort has mostly focused on regular echinoids that are used as model organisms in developmental and

genomic research. Data for irregular echinoids (sand dollars, heart urchins, and related groups) is much scarcer. The data presented here represent the first available mitogenome for a deep-sea irregular taxon, as well as the first for the order Holasteroidea, within which pourtalesiid echinoids are placed. The structure of the mitogenome reinforces the known pattern of gene order conservation within Echinoidea. Phylogenetic results agree with the topologies of previous phylogenomic efforts, yet support values for deep nodes are low, including both Irregularia and the node directly subtending it, Carinacea (i.e. Irregularia + Echinacea). The fossil record of irregular echinoids stretches back to the Early Jurassic (Smith et al. 2006), while molecular estimates of divergence times place the origin of both aforementioned clades within the Triassic (Mongiardino Koch et al. 2022). Such ancient divergences might be beyond the reach of mitogenome datasets, suggesting that caution should be exerted when interpreting novel topologies obtained through such efforts (e.g. Shin et al. 2022; Sun et al. 2022).

Acknowledgments

We would like to thank the captain and crew of the R/V *Thomas G. Thompson*, as well as chief scientist Amy Gartman and scientists Amanda Demopoulos and Kira Mizell. The manuscript has been improved by the comments of two anonymous reviewers.

Author contributions

GWR and NMK conceived and designed the study. The mitochondrial genome data were sequenced, assembled, and annotated by ASH, MS, and NMK. The phylogenetic analysis was performed by NMK and MS. NMK and MS drafted the manuscript with revisions from ASH and GWR. All authors read and approved the final manuscript and agree to be accountable for all aspects of the work.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was supported by National Science Foundation grant DEB-2036186.

ORCID

Avery S. Hiley  <http://orcid.org/0000-0001-8956-5080>
 Greg W. Rouse  <http://orcid.org/0000-0001-9036-9263>
 Nicolás Mongiardino Koch  <http://orcid.org/0000-0001-6317-5869>

Data availability statement

All molecular data reported here is available from NCBI (<https://www.ncbi.nlm.nih.gov>). The assembled mitogenome is available under accession number OR494080. Genomic raw reads are deposited under SRA number SRR25731399, part of BioProject PRJNA1008243. Associated BioSample number is SAMN37113738.

References

- Agassiz A. 1878. Preliminary report on the Echini of the exploring expedition of H.M.S. "Challenger". Sir C. Wyville Thomson Chief of Civilian Staff. Proceedings of the American Academy of Arts and Sciences. New series 14:190–212. doi:10.2307/25138537.
- Allio R, Schomaker-Bastos A, Romiguier J, Prosdocimi F, Nabholz B, Delsuc F. 2020. MitoFinder: efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics. *Mol Ecol Resour.* 20(4):892–905. doi:10.1111/1755-0998.13160.
- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J, Middendorf M, Stadler PF. 2013. MITOS: improved de novo metazoan mitochondrial genome annotation. *Mol Phylogenet Evol.* 69(2):313–319. doi:10.1016/j.ympev.2012.08.023.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics.* 30(15):2114–2120. doi:10.1093/bioinformatics/btu170.
- Bushnell B. 2014. BBMap: a fast, accurate, splice-aware aligner. Berkeley (CA): Lawrence Berkeley National Lab. (LBNL).
- Chernomor O, von Haeseler A, Minh BQ. 2016. Terrace aware data structure for phylogenomic inference from supermatrices. *Syst Biol.* 65(6):997–1008. doi:10.1093/sysbio/syw037.
- Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: improving the ultrafast bootstrap approximation. *Mol Biol Evol.* 35(2):518–522. doi:10.1093/molbev/msx281.
- Kalyaanamoorthy A, Minh BQ, Wong TKF, von Haeseler A, Jermini LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods.* 14(6):587–589. doi:10.1038/nmeth.4285.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30(4):772–780. doi:10.1093/molbev/mst010.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics.* 28(12):1647–1649. doi:10.1093/bioinformatics/bts199.
- Kroh A, Mooi R. 2023. World Echinoidea Database. [accessed 2023 Jun 15]. <https://www.marinespecies.org/echinoidea>. doi:10.14284/355.
- Laslett D, Canbäck B. 2008. ARWEN: a program to detect tRNA genes in metazoan mitochondrial nucleotide sequences. *Bioinformatics.* 24(2):172–175. doi:10.1093/bioinformatics/btm573.
- Li D, Luo R, Liu CM, Leung CM, Ting HF, Sadakane K, Yamashita H, Lam TW. 2016. MEGAHIT v1.0: a fast and scalable metagenome assembler driven by advanced methodologies and community practices. *Methods.* 102:3–11. doi:10.1016/j.ymeth.2016.02.020.
- Lin J-P, Tsai M-H, Kroh A, Trautman A, Machado DJ, Chang L-Y, Reid R, Lin KT, Bronstein O, Lee SJ, et al. 2020. The first complete mitochondrial genome of the sand dollar *Sinacchinocyamus mai* (Echinoidea: clypeasteroidea). *Genomics.* 112(2):1686–1693. doi:10.1016/j.ygeno.2019.10.007.
- Miguez-Salas O, Vardaro MF, Rodríguez-Tovar FJ, Pérez-Claros JA, Huffard CL. 2022. Deep-sea echinoid trails and seafloor nutrient distribution: present and past implications. *Front Mar Sci.* 9:903864. doi:10.3389/fmars.2022.903864.
- Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol Biol Evol.* 37(5):1530–1534. doi:10.1093/molbev/msaa015.
- Mironov AN. 1973. New deep-sea species of sea urchins of the genus *Echinocrepis* and the distribution of the family Pourtalesiidae (Echinoidea, Meridosterniana). *Trudy Instituta Okeanologii Akademii Nauk SSSR.* 91:240–247.
- Mongiardino Koch N, Coppard SE, Lessios HA, Briggs DEG, Mooi R, Rouse GW. 2018. A phylogenomic resolution of the sea urchin tree of life. *BMC Evol Biol.* 18(1):189. doi:10.1186/s12862-018-1300-4.
- Mongiardino Koch N, Thompson JR, Hiley AS, McCowin MF, Armstrong AF, Coppard SE, Aguilera F, Bronstein O, Kroh A, Mooi R, et al. 2022. Phylogenomic analyses of echinoid diversification prompt a re-evaluation of their fossil record. *Elife.* 11:e72460. doi:10.7554/eLife.72460.

- Perseke M, Bernhard D, Fritzsche G, Brümmer F, Stadler PF, Schlegel M. 2010. Mitochondrial genome evolution in Ophiuroidea, Echinoidea, and Holothuroidea: insights in phylogenetic relationships of Echinodermata. *Mol Phylogenet Evol.* 56(1):201–211. doi:[10.1016/j.ympev.2010.01.035](https://doi.org/10.1016/j.ympev.2010.01.035).
- Quek ZBR, Chang JJM, Ip YCA, Chan YKS, Huang D. 2021. Mitogenomes reveal alternative initiation codons and lineage-specific gene order conservation in echinoderms. *Mol Biol Evol.* 38(3):981–985. doi:[10.1093/molbev/msaa262](https://doi.org/10.1093/molbev/msaa262).
- Schultz HA. 2017. Echinoidea: with bilateral symmetry. Irregularia: Walter de Gruyter GmbH & Co KG.
- Shin JS, Song CU, Choi H, Kwon KK, Kang DH, Eyun SI, Choi KS. 2022. The complete mitochondrial genome of the sand dollar *Astriclypeus manni* (Verrill, 1867) (Echinoidea: Astriclypeidae) in the subtidal sand flat in Jeju Island off the south coast of Korea. *Mitochondrial DNA B Resour.* 7(9):1602–1603. doi:[10.1080/23802359.2022.2116946](https://doi.org/10.1080/23802359.2022.2116946).
- Smith AB, Pisani D, Mackenzie-Dodds JA, Stockley B, Webster BL, Littlewood DTJ. 2006. Testing the molecular clock: molecular and paleontological estimates of divergence times in the Echinoidea (Echinodermata). *Mol Biol Evol.* 23(10):1832–1851. doi:[10.1093/molbev/msl039](https://doi.org/10.1093/molbev/msl039).
- Sun S, Xiao N, Sha Z. 2022. Complete mitochondrial genomes of four deep-sea echinoids: conserved mitogenome organization and new insights into the phylogeny and evolution of Echinoidea. *PeerJ.* 10: e13730. doi:[10.7717/peerj.13730](https://doi.org/10.7717/peerj.13730).
- Vardaro MF. 2010. Genetic and anatomic relationships among three morphotypes of the echinoid *Echinocrepis rostrata*. *Invertebr Biol.* 129(4): 368–375. doi:[10.1111/j.1744-7410.2010.00205.x](https://doi.org/10.1111/j.1744-7410.2010.00205.x).
- Vardaro MF, Ruhl HA, Smith JKL. 2009. Climate variation, carbon flux, and bioturbation in the abyssal North Pacific. *Limnol Oceanogra.* 54(6): 2081–2088. doi:[10.4319/lo.2009.54.6.2081](https://doi.org/10.4319/lo.2009.54.6.2081).
- Wakayama N, Kiyono Y, Matsumoto N, Saitoh M, Kanazawa K. 2019. Effective DNA extraction methods for mitochondrial phylogenomics of the sea urchins. *Zoosymposia.* 15(1):192–202. doi:[10.11646/zoosymposia.15.1.20](https://doi.org/10.11646/zoosymposia.15.1.20).