



On the Stability of Shear Flows in Bounded Channels, II: Non-monotonic Shear Flows

Alexandru D. Ionescu¹ · Sameer Iyer² · Hao Jia³

Received: 29 December 2022 / Accepted: 25 July 2023

© Vietnam Academy of Science and Technology (VAST) and Springer Nature Singapore Pte Ltd. 2023

Abstract

We give a proof of linear inviscid damping and vorticity depletion for non-monotonic shear flows with one critical point in a bounded periodic channel. In particular, we obtain quantitative depletion rates for the vorticity function without any symmetry assumptions.

Keywords Inviscid damping · Vorticity depletion · Non-monotonic shear flows

Mathematics Subject Classification (2010) 35B40 · 35Q31 · 35P25

1 Introduction

The study of stability problems in mathematical analysis of fluid dynamics has a long and distinguished history, dating back to the work of Kelvin [18], Orr [25] and Rayleigh [26] among many others, and continuing to the present day. Hydrodynamical stability problems can be considered in both two and three dimensions. In this paper we work with two dimensional inviscid flows.

For the Euler equations, there is significant recent progress on the asymptotic stability of monotonic shear flows and vortices, assuming spectral stability, see for example [3, 9, 14, 15, 17, 22, 28, 30, 34, 35] for linear results. The main mechanism of stabilization is the so called “inviscid damping”, which refers to the transfer of energy of vorticity to higher and higher frequencies leading to decay of the stream and velocity functions, as $t \rightarrow \infty$. Extending

Dedicated to Carlos Kenig, on the occasion of his 70th birthday.

✉ Hao Jia
jia@umn.edu

Alexandru D. Ionescu
aionescu@math.princeton.edu

Sameer Iyer
sameer@math.ucdavis.edu

¹ Department of Mathematics, Princeton University, Princeton, NJ, USA

² Department of Mathematics, University of California, Davis, CA, USA

³ School of Mathematics, University of Minnesota, Minneapolis, MN, USA

the linearized stability analysis for inviscid fluid equations to the full nonlinear setting is a challenging problem, and the only available results are on spectrally stable monotonic shear flows [2, 10, 12, 23], and on point vortices [11]. We refer also to the recent review article [13] for a more in-depth discussion of recent developments of both linear and nonlinear inviscid damping.

Many physically important shear flows are not monotonic, such as the Poiseuille flow and the Kolmogorov flows. For such flows on the linear inviscid level, there is an additional significant physical phenomenon called “vorticity depletion” which refers to the asymptotic vanishing of vorticity as $t \rightarrow \infty$ near the critical point where the derivative of the shear flow is zero, first predicted in Bouchet and Morita [5], and proved rigorously in Wei–Zhang–Zhao [31]. A similar phenomenon was proved in Bedrossian–Coti Zelati–Vicol [3] for the case of vortices. See also [14] by the first and third author for a refined description of the dynamics in Gevrey spaces as a step towards proving nonlinear vortex symmetrization.

In [31] by Wei–Zhang–Zhao, sharp linear inviscid damping estimates and quantitative depletion estimates were obtained for an important class of “symmetric shear flows” in a channel (see also [32] by Wei–Zhang–Zhao for a similar result for Kolmogorov flow). When no symmetry is assumed, only qualitative bounds are available. Heuristically the general case should be similar to the symmetric one, since the main vorticity depletion mechanism is completely local and asymptotically all shear flows approach symmetric ones at the (non-degenerate) critical points. However there are significant difficulties in using the approach of [31] to extend the quantitative depletion bounds of [31] to the general case, as the argument in [31] relies heavily on decomposition of functions into odd and even parts, which are specific to symmetric shear flows.

In this paper, we prove linear inviscid damping estimates and quantitative vorticity depletion estimates for a class of stable non-monotonic shear flows with one non-degenerate critical point. The main new features of our results are that we do not need symmetry condition on the background shear flow, and that our formulation on quantitative depletion for vorticity function seems to be new even for general symmetric shear flows (see however Wei–Zhang–Zhao [32] which contains a sharp depletion rate at the critical points for Kolmogorov flow), see Theorem 1.2 below for the precise statements. We begin with the description of our main equations and theorem.

1.1 Main Equations

Consider the two dimensional Euler equation linearized around a shear flow $(b(y), 0)$, in the periodic channel $(x, y, t) \in \mathbb{T} \times [0, 1] \times [0, \infty)$:

$$\begin{aligned} \partial_t \omega + b(y) \partial_x \omega - b''(y) u^y &= 0, \\ \operatorname{div} u &= 0 \quad \text{and} \quad \omega = -\partial_y u^x + \partial_x u^y, \end{aligned}$$

with the natural non-penetration boundary condition $u^y|_{y=0,1} = 0$.

For the linearized flow, $\int_{\mathbb{T} \times [0,1]} u^x(x, y, t) dx dy$ and $\int_{\mathbb{T} \times [0,1]} \omega(x, y, t) dx dy$ are conserved quantities. In this paper, we will assume that

$$\int_{\mathbb{T} \times [0,1]} u_0^x(x, y) dx dy = \int_{\mathbb{T} \times [0,1]} \omega_0 dx dy = 0.$$

These assumptions can be dropped by adjusting $b(y)$ with a linear shear flow $C_0y + C_1$. Then one can see from the divergence free condition on u that there exists a stream function $\psi(t, x, y)$ with $\psi(t, x, 0) = \psi(t, x, 1) \equiv 0$, such that

$$u^x = -\partial_y \psi, \quad u^y = \partial_x \psi.$$

The stream function ψ can be solved through

$$\Delta \psi = \omega, \quad \psi|_{y=0,1} = 0.$$

We summarize our equations as follows

$$\begin{cases} \partial_t \omega + b(y) \partial_x \omega - b''(y) \partial_x \psi = 0, \\ \Delta \psi(t, x, y) = \omega(t, x, y), \quad \psi(t, x, 0) = \psi(t, x, 1) = 0, \\ (u^x, u^y) = (-\partial_y \psi, \partial_x \psi), \end{cases} \quad (1.1)$$

for $t \geq 0$, $(x, y) \in \mathbb{T} \times [0, 1]$.

Our goal is to understand the long time behavior of $\omega(t)$ as $t \rightarrow \infty$, with Sobolev regular initial vorticity ω_0 .

1.2 The Main Results

We describe more precisely the main assumptions and our main conclusion. The main conditions we shall assume on the shear flow $b(y) \in C^4([0, 1])$ are as follows.

Assumption 1.1 *We assume that the background flow $b(y) \in C^4([0, 1])$ satisfies the following conditions.*

(1)

$$S := \{y \in [0, 1] : b'(y) = 0\} = \{y_*\} \subset (0, 1).$$

In addition, $b''(y_) \neq 0$. For the sake of concreteness, we assume that $b''(y_*) > 0$ in this paper.*

(2) *For $k \in \mathbb{Z} \setminus \{0\}$, the linearized operator $L_k : L^2(0, 1) \rightarrow L^2(0, 1)$ defined as*

$$L_k g(y) := b(y)g(y) + b''(y) \int_0^1 G_k(y, z)g(z) dz \quad (1.2)$$

has no discrete eigenvalues nor generalized embedded eigenvalues. In the above G_k is the Green's function for $k^2 - \frac{d^2}{dy^2}$ on the interval $(0, 1)$ with zero Dirichlet boundary condition, as defined in (3.1) below.

We refer to Section 2 below for the definition and more discussion about generalized embedded eigenvalues.

Our main result is the following theorem.

Theorem 1.2 *Assume that $\omega(t, \cdot) \in C([0, \infty), H^4(\mathbb{T} \times [0, 1]))$ with the associated stream function $\psi(t, \cdot)$ is the unique solution to (1.1), with initial data $\omega_0 \in H^4(\mathbb{T} \times [0, 1])$ satisfying for all $y \in [0, 1]$,*

$$\int_{\mathbb{T}} \omega_0(x, y) dx = 0.$$

Then we have the following bounds.

(i) *Inviscid damping estimates:*

$$\|\psi(t, \cdot)\|_{L^2(\mathbb{T} \times [0,1])} \lesssim \frac{1}{\langle t \rangle^2} \|\omega_0\|_{H^4(\mathbb{T} \times [0,1])}, \quad (1.3)$$

$$\|u^x(t, \cdot)\|_{L^2(\mathbb{T} \times [0,1])} \lesssim \frac{1}{\langle t \rangle} \|\omega_0\|_{H^4(\mathbb{T} \times [0,1])}, \quad \|u^y(t, \cdot)\|_{L^2(\mathbb{T} \times [0,1])} \lesssim \frac{1}{\langle t \rangle^2} \|\omega_0\|_{H^4(\mathbb{T} \times [0,1])}. \quad (1.4)$$

(ii) *Vorticity depletion estimates: there exists a decomposition*

$$\omega(t, x, y) := \omega_{\text{loc}}(t, x, y) + \omega_{\text{nloc}}(t, x, y),$$

where for $(x, y, t) \in \mathbb{T} \times [0, 1] \times [0, \infty)$,

$$|\omega_{\text{loc}}(t, x, y)| \lesssim |y - y_*|^{7/4} \|\omega_0\|_{H^4(\mathbb{T} \times [0,1])}, \quad |\omega_{\text{nloc}}(t, x, y)| \lesssim \frac{1}{\langle t \rangle^{7/8}} \|\omega_0\|_{H^4(\mathbb{T} \times [0,1])}. \quad (1.5)$$

1.3 Remarks and Main Ideas of Proof

We have the following remarks on Theorem 1.2. *Firstly*, in the above theorem we have not tracked the minimal regularity required for the bounds (1.3), (1.4) and (1.5) to hold, and a more careful argument can probably significantly reduce the number of derivatives needed on the initial data ω_0 . *Secondly*, we note also that the argument here can be applied to non-monotonic shear flows with multiple non-degenerate points, although the presentation will be more complicated. *Thirdly*, a more sophisticated analysis may yield a sharper rate of vorticity depletion with rate

$$|\omega_{\text{loc}}(t, x, y)| \lesssim |y - y_*|^{2-}, \quad |\omega_{\text{nloc}}(t, x, y)| \lesssim \langle t \rangle^{-1+}.$$

It is not clear to us though if one can reach the optimal rates of $|y - y_*|^2$ and $\langle t \rangle^{-1}$.

We briefly explain the main ideas of the proof.

By a standard spectral representation formula, see (2.4), it suffices to study the spectral density functions and the associated Rayleigh equation (2.5). There are two main cases to consider. When the spectral parameter λ is not close to the critical value $b(y_*)$, the situation is similar to monotonic shear flows and can be treated as in [15]. The main new case is when the spectral parameter λ is close to the critical value $b(y_*)$. In this case, the Rayleigh equation (2.5) is very singular, and the potential term $\frac{b''(y)}{b(y) - \lambda + i\varepsilon}$ has a quadratic singularity roughly of the form $\frac{2}{(y - y_*)^2 + (\lambda - b(y_*)) + i\varepsilon}$ for y close to y_* .

The key observation here, as in [14], is that the potential term $\frac{b''(y)}{b(y) - \lambda + i\varepsilon}$ is *critically singular* and has real part with a *favorable* sign for $1 \gg |y - y_*| \gg |\lambda - b(y_*)|^{1/2}$, which needs to be incorporated as part of the main term. We therefore define a modified Green's function for the main term, see (3.6)–(3.7), which has strong vanishing conditions near $y = y_*$, leading ultimately to vorticity depletion. After extracting the main terms in the Rayleigh equation (2.5), the rest of the terms can be treated as compact perturbations, and can be bounded using a limiting absorption principle, see Lemma 4.4, thanks to the spectral assumption 1.1.

The limiting absorption principle provides preliminary bounds on the spectral density functions $\psi_{k,\varepsilon}^t(y, \lambda)$ with $t \in \{\pm\}$. To obtain the desired quantitative decay rates, we take up to two derivatives in λ of the spectral density functions, and again use the limiting absorption

principle to estimate the resulting derivatives, after extracting the main singular terms. The procedure is more or less straightforward but the calculations are quite lengthy. We refer to [15] also for similar calculations in a simpler setting. Lastly, we note that there are important cancellations between $\psi_{k,\varepsilon}^+(y, \lambda)$ and $\psi_{k,\varepsilon}^-(y, \lambda)$ in the limit $\varepsilon \rightarrow 0+$, which is the reason why we need two versions of the limiting absorption principle, see Lemma 4.4, with different weighted spaces.

1.4 Notations

We summarize here some notations that are specific for this paper for the reader's conveniences. For positive numbers α, β , we set $\alpha \wedge \beta := \min\{\alpha, \beta\}$. We denote for $d > 0$, $\Sigma_d := \{b(y) : y \in [y_* - d, y_* + d]\}$, $S_d := [y_* - d, y_* + d]$. We also denote $\Sigma := \{b(y) : y \in [0, 1]\}$ and $I := [0, 1]$. For $k \in \mathbb{Z} \setminus \{0\}$, we define for $f \in H^1(I)$ the norm $\|f\|_{H_k^1(I)} := \|f\|_{L^2(I)} + |k|^{-1}\|f'\|_{L^2(I)}$.

2 Spectral Property and Representation Formula

Taking Fourier transform in x in the equation (1.1) for ω , we obtain that

$$\partial_t \omega_k + ikb(y)\omega_k - ikb''(y)\psi_k = 0, \quad (2.1)$$

for $k \in \mathbb{Z}$, $t \geq 0$, $y \in [0, 1]$. In the above, ω_k and ψ_k are the k -th Fourier coefficients of ω , ψ in x respectively. For each $k \in \mathbb{Z} \setminus \{0\}$, recall from (1.2) that for any $g \in L^2(0, 1)$,

$$L_k g(y) = b(y)g(y) + b''(y) \int_0^1 G_k(y, z)g(z)dz,$$

where G_k is the Green's function for the operator $k^2 - \frac{d^2}{dy^2}$ on $(0, 1)$ with zero Dirichlet boundary condition. Then (2.1) can be reformulated abstractly as

$$\partial_t \omega_k + ikL_k \omega_k = 0. \quad (2.2)$$

In contrast to the spectral property of the linearized operator around monotonic shear flows, the spectral property of L_k is less understood, especially on the generation of discrete eigenvalues and embedded eigenvalues. From general spectral theory, we know that the spectrum of L_k consists of the continuous spectrum

$$\Sigma := \{b(y) : y \in [0, 1]\},$$

together with some discrete eigenvalues with nonzero imaginary part which can only accumulate at the set of continuous spectrum Σ . Unlike the case of monotonic shear flows where the discrete eigenvalues can accumulate only at inflection points of the background shear flow, there appears no simple characterization of the possible accumulation points for non-monotonic shear flows.

Recall that $\lambda \in \Sigma$ is called an embedded eigenvalue if there exists a nontrivial $g \in L^2(0, 1)$, such that

$$L_k g = \lambda g.$$

For non-monotonic shear flows, this definition is too restrictive, as accumulation points of discrete eigenvalues may no longer be embedded eigenvalues. To capture the discrete

eigenvalues, we recall the following definition of “generalized embedded eigenvalues”, which can be found already in [31], adapted to our setting.

Definition 2.1 We call $\lambda \in \Sigma$ a generalized embedded eigenvalue, if one of the following conditions is satisfied.

- λ is an embedded eigenvalue.
- $\lambda \neq b(y_*)$ and there exists a nontrivial $\psi \in H_0^1(0, 1) : (0, 1) \rightarrow \mathbb{C}$ such that in the sense of distributions on $(0, 1)$,

$$(k^2 - \partial_y^2)\psi(y) + \text{P.V.} \frac{b''(y)\psi(y)}{b(y) - \lambda} + i\pi \sum_{z \in [0, 1], b(z) = \lambda} \frac{b''(z)\psi(z)}{|b'(z)|} \delta(y - z) = 0. \quad (2.3)$$

We remark that our assumption that the critical point y_* of $b(y)$ being non-degenerate implies that the sum in (2.3) is finite, and that the spectral assumption 1.1 is satisfied if $b'' > 0$ on $[0, 1]$.

Proposition 2.2 Suppose that $k \in \mathbb{Z} \setminus \{0\}$ and $\omega_0^k \in L^2([0, 1])$. Then the stream function $\psi_k(t, y)$ for $k \in \mathbb{Z} \setminus \{0\}$, $y \in [0, 1]$, $t \geq 0$ has the representation

$$\psi_k(t, y) = -\frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\Sigma} e^{-ik\lambda t} \left[\psi_{k,\varepsilon}^-(y, \lambda) - \psi_{k,\varepsilon}^+(y, \lambda) \right] d\lambda, \quad (2.4)$$

where $\psi_{k,\varepsilon}^{\iota}(y, \lambda)$ for $\iota \in \{+, -\}$, $y \in [0, 1]$, $\lambda \in \Sigma$, $k \in \mathbb{Z} \setminus \{0\}$, and sufficiently small $\varepsilon \in [-1/4, 1/4] \setminus \{0\}$, are the solutions to

$$-k^2 \psi_{k,\varepsilon}^{\iota}(y, \lambda) + \frac{d^2}{dy^2} \psi_{k,\varepsilon}^{\iota}(y, \lambda) - \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \psi_{k,\varepsilon}^{\iota}(y, \lambda) = \frac{-\omega_0^k(y)}{b(y) - \lambda + i\varepsilon}, \quad (2.5)$$

with zero Dirichlet boundary condition.

Proof By standard theory of spectral projection, from (2.2), we obtain that for $y \in [0, 1]$,

$$\omega_k(t, y) = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\Sigma} e^{i\lambda t} \left\{ [(\lambda + kL_k - i\varepsilon)^{-1} - (\lambda + kL_k + i\varepsilon)^{-1}] \omega_0^k \right\} (y) d\lambda.$$

We then obtain for $y \in [0, 1]$,

$$\begin{aligned} \psi_k(t, y) &= -\frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\Sigma} e^{-ik\lambda t} \int_0^1 G_k(y, z) \\ &\quad \times \left\{ [(-\lambda + L_k - i\varepsilon)^{-1} - (-\lambda + L_k + i\varepsilon)^{-1}] \omega_0^k \right\} (z) dz d\lambda \\ &= -\frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\Sigma} e^{-ik\lambda t} \left[\psi_{k,\varepsilon}^-(y, \lambda) - \psi_{k,\varepsilon}^+(y, \lambda) \right] d\lambda. \end{aligned}$$

In the above, for $y \in [0, 1]$ and $\lambda \in \Sigma$,

$$\begin{aligned} \psi_{k,\varepsilon}^+(y, \lambda) &:= \int_0^1 G_k(y, z) [(-\lambda + L_k + i\varepsilon)^{-1} \omega_0^k] (z) dz, \\ \psi_{k,\varepsilon}^-(y, \lambda) &:= \int_0^1 G_k(y, z) [(-\lambda + L_k - i\varepsilon)^{-1} \omega_0^k] (z) dz. \end{aligned}$$

Therefore for $\iota \in \{+, -\}$, $y \in [0, 1]$, $\lambda \in \Sigma$,

$$\left(k^2 - \frac{d^2}{dy^2} \right) \psi_{k,\varepsilon}^{\iota}(y, y_0) = (-\lambda + L_k + i\varepsilon)^{-1} \omega_0^k(y),$$

which implies

$$\begin{aligned}\omega_0^k(y) &= (-\lambda + L_k + i\epsilon) \left(k^2 - \frac{d^2}{dy^2} \right) \psi_{k,\epsilon}^t(y, \lambda) \\ &= (b(y) - \lambda + i\epsilon) \left(k^2 - \frac{d^2}{dy^2} \right) \psi_{k,\epsilon}^t(y, \lambda) + b''(y) \psi_{k,\epsilon}^t(y, \lambda).\end{aligned}\quad (2.6)$$

It follows from (2.6) that $\psi_{k,\epsilon}^+(y, \lambda)$, $\psi_{k,\epsilon}^-(y, \lambda)$ satisfy (2.5). The proposition is now proved. $\square$

Remark 2.3 The existence of $\psi_{k,\epsilon}^t$ for sufficiently small $\epsilon \neq 0$ follows from our spectral assumptions, which imply the solvability of (2.5) for sufficiently small $\epsilon \neq 0$, see also (4.4).

3 Bounds on the Green's Function and Modified Green's Function

3.1 Elementary Properties of the Standard Green's Function

For integers $k \in \mathbb{Z} \setminus \{0\}$, recall that the Green's function $G_k(y, z)$ solves

$$-\frac{d^2}{dy^2} G_k(y, z) + k^2 G_k(y, z) = \delta(y - z),$$

with Dirichlet boundary conditions $G_k(0, z) = G_k(1, z) = 0$, $z \in (0, 1)$. G_k has the explicit formula

$$G_k(y, z) = \frac{1}{k \sinh k} \begin{cases} \sinh(k(1-z)) \sinh(ky) & \text{if } y \leq z, \\ \sinh(kz) \sinh(k(1-y)) & \text{if } y \geq z, \end{cases} \quad (3.1)$$

and the symmetry

$$G_k(y, z) = G_k(z, y) \quad \text{for } k \in \mathbb{Z} \setminus \{0\}, y, z \in [0, 1].$$

We note the following bounds for G_k

$$\begin{aligned}& \sup_{y \in [0, 1], |A| \leq 10} \left[|k|^2 \|G_k(y, z)(\log |z - A|)^m\|_{L^1(z \in [0, 1])} + |k| \|\partial_{y,z} G_k(y, z)(\log |z - A|)^m\|_{L^1(z \in [0, 1])} \right] \\ & + \sup_{y \in [0, 1], \alpha \in \{0, 1\}} \left[|k|^{3/2-\alpha} \|\partial_{y,z}^\alpha G_k(y, z)\|_{L^2(z \in [0, 1])} \right] \lesssim |\log(k)|^m \quad \text{for } m \in \{0, 1, 2, 3\}.\end{aligned}\quad (3.2)$$

Define

$$F_k(y, z) = \frac{1}{\sinh k} \begin{cases} -k \cosh(k(1-z)) \cosh(ky), & 0 \leq y \leq z \leq 1, \\ -k \cosh(kz) \cosh(k(1-y)), & 1 \geq y > z \geq 0. \end{cases}$$

We note that

$$\partial_y \partial_z G_k(y, z) = \partial_z \partial_y G_k(y, z) = \delta(y - z) + F_k(y, z) \quad \text{for } y, z \in [0, 1]. \quad (3.3)$$

By direct computation, we see F_k satisfies the bounds

$$\begin{aligned}& \sup_{y \in [0, 1], |A| \leq 10} \left[\|F_k(y, z)(\log |z - A|)^m\|_{L^1(z \in [0, 1])} + |k|^{-1} \|\partial_{y,z} F_k(y, z)(\log |z - A|)^m\|_{L^1(z \in [0, 1])} \right] \\ & + \sup_{y \in [0, 1], \alpha \in \{0, 1\}} \left[|k|^{-1/2-\alpha} \|\partial_{y,z}^\alpha F_k(y, z)\|_{L^2(z \in [0, 1])} \right] \lesssim |\log(k)|^m \quad \text{for } m \in \{0, 1, 2, 3\}.\end{aligned}\quad (3.4)$$

The bounds (3.2) and (3.4) can be proved by explicit calculations and are useful in the proof of Lemma 4.1 below.

3.2 Bounds on the Modified Green's Function

It follows from Assumption 1.1 that there exists a $\delta_0 \in (0, 1/8)$ such that

$$\inf\{|y_*|, |y_* - 1|\} > 10\delta_0 \quad \text{and} \quad \sup_{y \in (y_* - 4\delta_0, y_* + 4\delta_0)} |b'''(y)|\delta_0 < |b''(y_*)|/10.$$

Define the set

$$\Sigma_{\delta_0} := \{b(y) : y \in [y_* - \delta_0, y_* + \delta_0]\},$$

and fix a standard smooth cutoff function $\varphi \in C_c^\infty(-2, 2)$ satisfying $\varphi \equiv 1$ on $[-3/2, 3/2]$. For simplicity of notations, we denote

$$I := (0, 1).$$

To simplify notations we define also for $d \in (0, 1/10)$,

$$S_d := [y_* - d, y_* + d]. \quad (3.5)$$

For applications below, we also need to study the “modified Green’s function” $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$ for $y, z \in [0, 1]$, $\lambda \in \Sigma_{\delta_0}$ and $\varepsilon \in [-1/8, 1/8] \setminus \{0\}$, which satisfies for $y, z \in (0, 1)$,

$$(k^2 - \partial_y^2) \mathcal{G}_k(y, z; \lambda + i\varepsilon) + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \left[\varphi\left(\frac{y - y_*}{\delta_0}\right) - \varphi\left(\frac{y - y_*}{\delta(\lambda)}\right) \right] \mathcal{G}_k(y, z; \lambda + i\varepsilon) = \delta(y - z), \quad (3.6)$$

with the boundary condition

$$\mathcal{G}_k(y, z; \lambda + i\varepsilon)|_{y \in \{0, 1\}} = 0. \quad (3.7)$$

In the above, we have used the notation that

$$\delta(\lambda) := 8\sqrt{|\lambda - b(y_*)|/b''(y_*)}. \quad (3.8)$$

Define the weight $\varrho(y; \lambda + i\varepsilon)$ for $y, z \in [0, 1]$, $\lambda \in \Sigma_{\delta_0}$ and $\varepsilon \in [-1/8, 1/8] \setminus \{0\}$ as

$$\varrho(y; \lambda + i\varepsilon) := |\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2} + |y - y_*|.$$

The crucial bounds we need for the modified Green’s function $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$ is the following.

Lemma 3.1 *Let $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$ for $y, z \in [0, 1]$, $\lambda \in \Sigma_{\delta_0}$ and $\varepsilon \in [-1/8, 1/8] \setminus \{0\}$ be defined as in (3.6). Then we have the identity for $y, z \in [0, 1]$,*

$$\mathcal{G}_k(y, z; \lambda + i\varepsilon) = \mathcal{G}_k(z, y; \lambda + i\varepsilon), \quad (3.9)$$

and the following statements hold.

(i) *We have the bounds*

$$\begin{aligned} \sup_{y \in [0, 1], |y - z| \leq \min\{\varrho(z; \lambda + i\varepsilon), 1/|k|\}} |\mathcal{G}_k(y, z; \lambda + i\varepsilon)| &\lesssim \min\{\varrho(z; \lambda + i\varepsilon), 1/|k|\}, \\ \sup_{y \in [0, 1], |y - z| \leq \min\{\varrho(z; \lambda + i\varepsilon), 1/|k|\}} |\partial_y \mathcal{G}_k(y, z; \lambda + i\varepsilon)| &\lesssim 1; \end{aligned} \quad (3.10)$$

- (ii) For $y_1, y_2 \in [0, 1]$ with $y_2 \in [\min\{y_1, z\}, \max\{y_1, z\}]$ and $\varrho(y_2; \lambda + i\varepsilon) \gtrsim 1/|k|$, we have the bounds with $\alpha \in \{0, 1\}$

$$\begin{aligned} & |\partial_y^\alpha \mathcal{G}_k(y_1, z; \lambda + i\varepsilon)| \\ & \lesssim \left[|k| + \varrho^{-1}(y_1; \lambda + i\varepsilon) \right]^\alpha e^{-|k||y_1 - y_2|} \left[|k| \int_{[y_2 - 1/|k|, y_2 + 1/|k|] \cap I} |\mathcal{G}_k(y, z; \lambda + i\varepsilon)|^2 dy \right]^{1/2}. \end{aligned} \quad (3.11)$$

- (iii) For $y_1, y_2 \in [0, 1]$ with $y_2 \in [\min\{y_1, z\}, \max\{y_1, z\}]$ and $\varrho(y_2; \lambda + i\varepsilon) \ll 1/|k|$, we have the bounds with $\alpha \in \{0, 1\}$

$$\begin{aligned} & |\partial_y^\alpha \mathcal{G}_k(y_1, z; \lambda + i\varepsilon)| \\ & \lesssim \left[|k| + \varrho^{-1}(y_1; \lambda + i\varepsilon) \right]^\alpha \min \left\{ \frac{\varrho^2(y_1; \lambda + i\varepsilon)}{\varrho^2(y_2; \lambda + i\varepsilon)}, \frac{\varrho(y_2; \lambda + i\varepsilon)}{\varrho(y_1; \lambda + i\varepsilon)} \right\} M, \end{aligned} \quad (3.12)$$

where

$$M := \left[\frac{1}{\varrho(y_2; \lambda + i\varepsilon)} \int_{[y_2 - \varrho(y_2; \lambda + i\varepsilon), y_2 + \varrho(y_2; \lambda + i\varepsilon)] \cap I} |\mathcal{G}_k(y, z; \lambda + i\varepsilon)|^2 dy \right]^{1/2}.$$

Proof The proof is based on energy estimates and “entanglement inequalities”, as in [16]. See also the earlier work [33] where this type of inequality was used. We divide the proof into several steps.

Step 1: the proof of (3.10). We first establish the bounds (3.10). For simplicity of notation, we suppress the dependence on $z, \lambda + i\varepsilon$ and set for $y \in [0, 1]$,

$$h(y) := \mathcal{G}_k(y, z; \lambda + i\varepsilon), \quad V(y) := \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \left[\varphi \left(\frac{y - y_*}{\delta_0} \right) - \varphi \left(\frac{y - y_*}{\delta} \right) \right]. \quad (3.13)$$

Multiplying $\bar{h}$ to (3.6) and integrating over $[0, 1]$, we obtain that

$$\int_0^1 |\partial_y h(y)|^2 + |k|^2 |h(y)|^2 dy + \int_0^1 V(y) |h(y)|^2 dy = \bar{h}(z). \quad (3.14)$$

Note that for $y \in [0, 1]$, $\Re V(y) \geq 0$, and in addition, for $y \in S_{\delta_0}$ and

$$|y - y_*| > C_0 (|\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2})$$

with sufficiently large $C_0 \gg 1$,

$$1 + \Re V(y) \gtrsim \frac{1}{\varrho^2(y; \lambda + i\varepsilon)}.$$

It follows from (3.14) that

$$\begin{aligned} & \int_0^1 |\partial_y h(y)|^2 + |k|^2 |h(y)|^2 dy + \int_{y \in S_{\delta_0}, |y - y_*| > C_0(\delta + |\varepsilon|^{1/2})} \frac{1}{[\varrho(y; \lambda + i\varepsilon)]^2} |h(y)|^2 dy \\ & \lesssim |h(z)|. \end{aligned} \quad (3.15)$$

Using the Sobolev type inequality

$$\|h\|_{L^\infty(J)} \lesssim \|h\|_{L^2(J_*)} |J|^{-1/2} + \|\partial_y h\|_{L^2(J)} |J|^{1/2}, \quad (3.16)$$

for any intervals J, J_* with $J_* \subseteq J$ and $|J_*| \gtrsim |J|$, and choosing the interval $J \subset I$ as an interval containing z with length of the size $C_1 \min\{1/|k|, \varrho(z; \lambda + i\varepsilon)\}$, we obtain from (3.15) that

$$\int_0^1 |\partial_y h(y)|^2 + |k|^2 |h(y)|^2 dy + \int_{y \in S_{\delta_0}, |y-y_*| > C_0(\delta+|\varepsilon|^{1/2})} \frac{1}{[\varrho(y; \lambda + i\varepsilon)]^2} |h(y)|^2 dy \lesssim \min\{1/|k|, \varrho(z; \lambda + i\varepsilon)\}. \quad (3.17)$$

The first inequality of (3.10) then follows from (3.17) and (3.16). To obtain the second inequality in (3.10), we notice that the (3.6) and the first inequality in (3.10) imply pointwise bounds on $\partial_y^2 \mathcal{G}_k$ for $y \neq z$. The desired bound in the second inequality of (3.10) then follows from interpolation between the pointwise bounds on $\mathcal{G}_k$ and $\partial_y^2 \mathcal{G}_k$.

Step 2: the proof of (3.11). Denote

$$M_1 := \left[|k| \int_{[y_2-1/|k|, y_2+1/|k|] \cap I} |\mathcal{G}_k(y, z; \lambda + i\varepsilon)|^2 dy \right]^{1/2}.$$

For the sake of concreteness, we assume that $y_1 > z$ (so $y_2 \in [z, y_1]$). We shall also assume that $y_1 - y_2 \gg 1/|k|$ as the other case is analogous but easier. For $\varphi \in C_p^1([y_2, 1])$, the space of piecewise C^1 functions, with $\varphi(y_2) = 0$, we multiply $\varphi^2 \bar{h}$ to (3.6) and integrate over $[y_2, 1]$ to obtain that

$$\int_{y_2}^1 |\partial_y h(y)|^2 \varphi^2(y) + 2\partial_y h(y) \overline{h(y)} \varphi(y) \partial_y \varphi(y) + |k|^2 \varphi^2(y) |h(y)|^2 + V(y) |h(y)|^2 \varphi^2(y) dy = 0. \quad (3.18)$$

Taking the real part of (3.18) and using Cauchy–Schwarz inequality, we get that

$$\int_{y_2}^1 [|\partial_y \varphi(y)|^2 - |k|^2 |\varphi(y)|^2] |h(y)|^2 dy \geq 0. \quad (3.19)$$

We now choose φ more specifically as follows. We require that

$$\begin{aligned} \varphi(y_2) = 0, \quad \varphi''(y) = 0 \quad \text{for } y \in [y_2, y_2 + 1/|k|], \quad \varphi(y_2 + 1/|k|) = 1, \\ \varphi'(y) = |k| \varphi(y) \quad \text{for } y \in [y_2 + 1/|k|, y_1 - 1/|k|], \quad \varphi'(y) = 0 \quad \text{for } y \in [y_1 - 1/|k|, 1]. \end{aligned} \quad (3.20)$$

It follows from (3.19)–(3.20) that

$$\int_{y_1-1/|k|}^1 |k|^2 \varphi^2(y) |h(y)|^2 dy \lesssim |k| M_1^2, \quad \varphi(y) \approx e^{|k||y_1-y_2|} \quad \text{for } y \in [y_1 - 1/|k|, y_1 + 1/|k|] \cap I. \quad (3.21)$$

The desired bounds (3.11) follow from (3.21) and (3.6).

Step 3: the the proof of (3.12). For the sake of concreteness, we assume that $y_1 > z$ (and so $y_2 \in [z, y_1]$). We shall also assume that $y_1 - y_2 \gg \varrho(y_2; \lambda + i\varepsilon)$ and that $y_2 > y_* + \delta + |\varepsilon|^{1/2}$ as the other cases are analogous.

For $\varphi \in C_p^1([y_2, 1])$ with $\varphi(y_2) = 0$, we multiply $\varphi^2 \bar{h}$ to (3.6) and integrate over $[y_2, 1]$ to obtain that

$$\int_{y_2}^1 |\partial_y h(y)|^2 \varphi^2(y) + 2\partial_y h(y) \overline{h(y)} \varphi(y) \partial_y \varphi(y) + |k|^2 \varphi^2(y) |h(y)|^2 + V(y) |h(y)|^2 \varphi^2(y) dy = 0.$$

Write for $y \in [y_2, 1]$

$$h(y) = (y - y_*)^{1/2} h^*(y). \quad (3.22)$$

Simple calculations show that

$$\int_{y_2}^1 (y - y_*) |\partial_y h^*(y)|^2 \varphi^2(y) + 2(y - y_*) \partial_y \varphi(y) \varphi(y) \partial_y h^*(y) \overline{h^*(y)} \\ + \frac{1}{4(y - y_*)} |h^*(y)|^2 \varphi^2(y) + |k|^2 |h(y)|^2 \varphi^2(y) + (y - y_*) V(y) \varphi^2(y) |h^*(y)|^2 dy = 0.$$

Therefore

$$\int_{y_2}^1 \left[\frac{1}{4(y - y_*)} + (y - y_*) \Re V(y) \right] \varphi^2(y) |h^*(y)|^2 dy \leq \int_{y_2}^1 (y - y_*) (\partial_y \varphi)^2(y) |h^*(y)|^2 dy,$$

which implies that

$$\int_{y_2}^1 \frac{1}{y - y_*} \left[((y - y_*) \partial_y \varphi)^2(y) - (1/4 + (y - y_*)^2 \Re V(y)) \varphi^2(y) \right] |h^*(y)|^2 dy \geq 0. \quad (3.23)$$

We notice the pointwise bounds for $y \in [y_2, 1]$,

$$1/4 + (y - y_*)^2 \Re V(y) \geq \max \left\{ 0, 9/4 - C_2 \frac{\varrho^2(y_2; \lambda + i\varepsilon)}{(y - y_*)^2} - C_2 |y - y_*| \right\}. \quad (3.24)$$

Now we choose $\varphi \in C_p^1([y_2, 1])$ more precisely as follows. We require that

$$\begin{aligned} \varphi(y_2) = 0, \quad \varphi''(y) = 0 \quad \text{for } y \in [y_2, y_2 + \varrho(y_2; \lambda + i\varepsilon)], \quad \varphi(y_2 + \varrho(y_2; \lambda + i\varepsilon)) = 1, \\ (y - y_*) \varphi'(y) = [1/4 + (y - y_*)^2 \Re V(y)]^{1/2} \varphi(y) \\ \text{for } y \in [y_2 + \varrho(y_2; \lambda + i\varepsilon), y_1 - \varrho(y_1; \lambda + i\varepsilon)], \\ \text{and } \varphi'(y) = 0 \quad \text{for } y \in [y_1 - \varrho(y_1; \lambda + i\varepsilon), 1]. \end{aligned} \quad (3.25)$$

It follows from (3.23)–(3.25) that

$$\begin{aligned} \int_{y_1 - \varrho(y_1; \lambda + i\varepsilon)}^{y_1} \frac{1}{\varrho(y_1; \lambda + i\varepsilon)} \varphi^2(y) |h^*(y)|^2 dy \lesssim M^2 / \varrho(y_2; \lambda + i\varepsilon), \\ \varphi(y) \approx \frac{(y_1 - y_*)^{3/2}}{\varrho^{3/2}(y_2; \lambda + i\varepsilon)} \quad \text{for } y \in [y_1 - \varrho(y_1; \lambda + i\varepsilon), y_1]. \end{aligned} \quad (3.26)$$

The desired bounds (3.12) follow from the change of variable (3.22), the bound (3.23), (3.26) and (3.6).

Lastly we indicate how to prove the identity (3.9). For any $y, z \in (0, 1)$, using the notation in (3.13) for V , we have by integration by parts

$$\begin{aligned} \mathcal{G}_k(y, z; \lambda + i\varepsilon) &= \int_{[0, 1]} (k^2 - \partial_\varrho^2 + V(\varrho)) \mathcal{G}_k(\varrho, y; \lambda + i\varepsilon) \mathcal{G}_k(\varrho, z; \lambda + i\varepsilon) d\varrho \\ &= \int_{[0, 1]} (k^2 - \partial_\varrho^2 + V(\varrho)) \mathcal{G}_k(\varrho, z; \lambda + i\varepsilon) \mathcal{G}_k(\varrho, y; \lambda + i\varepsilon) d\varrho \\ &= \mathcal{G}_k(z, y; \lambda + i\varepsilon), \end{aligned}$$

which completes the proof of (3.9). $\square$

As a corollary of Lemma 3.1, we have the following additional bounds on the modified Green's function.

Lemma 3.2 Let $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$ for $y, z \in [0, 1]$, $\lambda \in \Sigma_{\delta_0}$, $k \in \mathbb{Z} \setminus \{0\}$ and $\varepsilon \in [-1/8, 1/8] \setminus \{0\}$ be defined as in (3.6). Recall the definition (3.8) for $\delta = \delta(\lambda) > 0$. Define

$$h := 10(\delta + |\varepsilon|^{1/2}),$$

and also for $y, z \in [0, 1]$,

$$\mathcal{H}_k(y, z; \lambda + i\varepsilon) := \left[\partial_z + \varphi \left(\frac{y - y_*}{h} \right) \partial_y \right] \mathcal{G}_k(y, z; \lambda + i\varepsilon).$$

Then the following statements hold for $z \in S_{4\delta}$.

(i) We have the bounds

$$\begin{aligned} \sup_{y \in [0, 1], |y-z| \leq \min\{\varrho(z; \lambda + i\varepsilon), 1/|k|\}} |\mathcal{H}_k(y, z; \lambda + i\varepsilon)| &\lesssim 1, \\ \sup_{y \in [0, 1], |y-z| \leq \min\{\varrho(z; \lambda + i\varepsilon), 1/|k|\}} |\partial_y \mathcal{H}_k(y, z; \lambda + i\varepsilon)| &\lesssim 1/\min\{\varrho(z; \lambda + i\varepsilon), 1/|k|\}; \end{aligned}$$

(ii) For $y_1, y_2 \in [0, 1]$ with $y_2 \in [\min\{y_1, z\}, \max\{y_1, z\}]$ and $\varrho(y_2; \lambda + i\varepsilon) \gtrsim 1/|k|$, we have the bounds with $\alpha \in \{0, 1\}$

$$\begin{aligned} &[\min\{\varrho(y_1; \lambda + i\varepsilon), 1/|k|\}]^\alpha |\partial_y^\alpha \mathcal{H}_k(y_1, z; \lambda + i\varepsilon)| \\ &\lesssim \frac{e^{-|k||y_1 - y_2|}}{\min\{\varrho(z; \lambda + i\varepsilon), 1/|k|\}} \left[|k| \int_{[y_2 - 1/|k|, y_2 + 1/|k|] \cap I} |\mathcal{G}_k(y, z; \lambda + i\varepsilon)|^2 dy \right]^{1/2}. \end{aligned}$$

(iii) For $y_1, y_2 \in [0, 1]$ with $y_2 \in [\min\{y_1, z\}, \max\{y_1, z\}]$ and $\varrho(y_2; \lambda + i\varepsilon) \ll 1/|k|$, we have the bounds with $\alpha \in \{0, 1\}$

$$\begin{aligned} &[\min\{\varrho(y_1; \lambda + i\varepsilon), 1/|k|\}]^\alpha |\partial_y^\alpha \mathcal{H}_k(y_1, z; \lambda + i\varepsilon)| \\ &\lesssim \frac{1}{\min\{\varrho(z; \lambda + i\varepsilon), 1/|k|\}} \min \left\{ \frac{\varrho^2(y_1; \lambda + i\varepsilon)}{\varrho^2(y_2; \lambda + i\varepsilon)}, \frac{\varrho(y_2; \lambda + i\varepsilon)}{\varrho(y_1; \lambda + i\varepsilon)} \right\} M, \end{aligned}$$

where

$$M := \left[\frac{1}{\varrho(y_2; \lambda + i\varepsilon)} \int_{[y_2 - \varrho(y_2; \lambda + i\varepsilon), y_2 + \varrho(y_2; \lambda + i\varepsilon)] \cap I} |\mathcal{G}_k(y, z; \lambda + i\varepsilon)|^2 dy \right]^{1/2}.$$

Proof Denote with a slight abuse of notation for $y \in [0, 1]$,

$$\varphi^\dagger(y) := \varphi \left(\frac{y - y_*}{h} \right), \quad V(y) := \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \left[\varphi \left(\frac{y - y_*}{\delta_0} \right) - \varphi \left(\frac{y - y_*}{\delta(\lambda)} \right) \right].$$

Then $\mathcal{H}_{k,j}(y, z; \lambda + i\varepsilon)$ satisfies for $y \in [0, 1]$, $z \in S_{4\delta}$,

$$\begin{aligned} &(k^2 - \partial_y^2) \mathcal{H}_k(y, z; \lambda + i\varepsilon) + V(y) \mathcal{H}_k(y, z; \lambda + i\varepsilon) \\ &= -\partial_y^2 \varphi^\dagger(y) \partial_y \mathcal{G}_k(y, z; \lambda + i\varepsilon) - \partial_y V(y) \varphi^\dagger(y) \mathcal{G}_k(y, z; \lambda + i\varepsilon) \\ &\quad - 2\partial_y \varphi^\dagger(y) \partial_y^2 \mathcal{G}_k(y, z; \lambda + i\varepsilon). \end{aligned} \tag{3.27}$$

The desired bounds then follow from (3.27), Lemma 3.1 and standard elliptic regularity theory. $\square$

The bounds in Lemmas 3.1 and 3.2 are quite sharp, since we can exploit the decay coming from both k^2 and $\frac{b''(y)}{b(y) - \lambda + i\varepsilon} \left[\varphi \left(\frac{y - y_*}{\delta_0} \right) - \varphi \left(\frac{y - y_*}{\delta(\lambda)} \right) \right]$. It is however somewhat complicated to formulate a concrete bound that is easy to use. Instead, the following simple bounds are more often used.

Corollary 3.3 *Let $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$ for $y, z \in [0, 1]$, $\lambda \in \Sigma_{\delta_0}$ and $\varepsilon \in [-1/8, 1/8] \setminus \{0\}$ be defined as in (3.6). Then we have the following bounds.*

(i) *For $y, z \in [0, 1]$, we have the bounds with $\alpha \in \{0, 1\}$*

$$\begin{aligned} & [|k| + \varrho^{-1}(y; \lambda + i\varepsilon)]^{-\alpha} |\partial_y^\alpha \mathcal{G}_k(y, z; \lambda + i\varepsilon)| \\ & \lesssim \frac{1}{|k| + \varrho^{-1}(z; \lambda + i\varepsilon)} \min \left\{ e^{-|k||y-z|}, \frac{\varrho^2(y; \lambda + i\varepsilon)}{\varrho^2(z; \lambda + i\varepsilon)}, \frac{\varrho(z; \lambda + i\varepsilon)}{\varrho(y; \lambda + i\varepsilon)} \right\}. \end{aligned} \quad (3.28)$$

(iii) *For $y \in [0, 1]$, $z \in S_{4\delta}$, we have the bounds with $\alpha \in \{0, 1, 2\}$*

$$\begin{aligned} & [|k| + \varrho^{-1}(y; \lambda + i\varepsilon)]^{-\alpha} |\partial_y^\alpha \mathcal{H}_k(y, z; \lambda + i\varepsilon)| \\ & \lesssim \min \left\{ e^{-|k||y-z|}, \frac{\varrho^2(y; \lambda + i\varepsilon)}{\varrho^2(z; \lambda + i\varepsilon)}, \frac{\varrho(z; \lambda + i\varepsilon)}{\varrho(y; \lambda + i\varepsilon)} \right\}. \end{aligned} \quad (3.29)$$

Proof The desired bounds (3.28)–(3.29) follow directly from Lemma 3.1 and Lemma 3.2, by choosing, if necessary, another point y' between y and z such that $\varrho(y'; \lambda + i\varepsilon) \approx 1/|k|$, and applying (3.28)–(3.29) on intervals $[\min\{z, y'\}, \max\{z, y'\}]$ and $[\min\{y', y\}, \max\{y', y\}]$ successively. $\square$

4 The Limiting Absorption Principle

In this section, we study the solvability of the main Rayleigh equations (2.5). It turns out that the situation is very different for the spectral range $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$ (the non-degenerate case) and $\lambda \in \Sigma_{\delta_0}$ (the degenerate case). We first consider the non-degenerate case.

4.1 The Non-degenerate Case

Fix $\varepsilon \in [-1/4, 1/4] \setminus \{0\}$, $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$, $k \in \mathbb{Z} \setminus \{0\}$. Define for each $g \in L^2(0, 1)$ the operator

$$T_{k,\lambda,\varepsilon} g(y) := \int_0^1 G_k(y, z) \frac{b''(z)g(z)}{b(z) - \lambda + i\varepsilon} dz. \quad (4.1)$$

For applications below, we fix a smooth cutoff function $\Phi \in C_0^\infty(y_* - \delta_0/3, y_* + \delta_0/3)$ with $\Phi \equiv 1$ on $[y_* - \delta_0/4, y_* + \delta_0/4]$. To obtain the optimal dependence on the frequency variable k , we define

$$\|g\|_{H_k^1(I)} := \|g\|_{L^2(I)} + |k|^{-1} \|g'\|_{L^2(I)}.$$

Lemma 4.1 *For $\varepsilon \in [-1/4, 1/4] \setminus \{0\}$, $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$, $k \in \mathbb{Z} \setminus \{0\}$, the operator $T_{k,\lambda,\varepsilon}$ satisfies the bound*

$$\|T_{k,\lambda,\varepsilon} g\|_{H_k^1(I)} \lesssim |k|^{-1/3} \|g\|_{H_k^1(I)} \quad \text{for all } g \in H_k^1(I). \quad (4.2)$$

In addition, we have the more precise regularity structure

$$\begin{aligned} & \left\| \partial_y T_{k,\lambda,\varepsilon} g(y) + \frac{b''(y)(1 - \Phi(y))g(y)}{b'(y)} \log(b(y) - \lambda + i\varepsilon) \right\|_{W^{1,1}(\mathbb{R})} \\ & \lesssim \langle k \rangle^{4/3} \|g\|_{H_k^1(I)}. \end{aligned} \quad (4.3)$$

Proof We can decompose for $y \in [0, 1]$,

$$T_{k,\lambda,\varepsilon}g(y) := T_{k,\lambda,\varepsilon}^1g(y) + T_{k,\lambda,\varepsilon}^2g(y),$$

where

$$\begin{aligned} T_{k,\lambda,\varepsilon}^1g(y) &:= \int_0^1 G_k(y, z) \frac{\Phi(z)b''(z)g(z)}{b(z) - \lambda - i\varepsilon} dz, \\ T_{k,\lambda,\varepsilon}^2g(y) &:= \int_0^1 G_k(y, z) \frac{(1 - \Phi(z))b''(z)g(z)}{b(z) - \lambda + i\varepsilon} dz. \end{aligned}$$

It follows from the definition of Φ that $T_{k,\lambda,\varepsilon}^1g(y)$ satisfies the bound

$$\|T_{k,\lambda,\varepsilon}^1g(y)\|_{H_k^1(I)} \lesssim |k|^{-1/3} \|g\|_{H_k^1(I)}, \quad \|\partial_y T_{k,\lambda,\varepsilon}^1g(y)\|_{W^{1,1}(I)} \lesssim \langle k \rangle^{4/3} \|g\|_{H_k^1(I)}.$$

To bound $T_{k,\lambda,\varepsilon}^2g(y)$, we follow the approach in [15]. Using integration by parts, we obtain that

$$\begin{aligned} T_{k,\lambda,\varepsilon}^2g(y) &= \int_0^1 G_k(y, z) \frac{(1 - \Phi(z))b''(z)g(z)}{b'(z)} \partial_z \log(b(z) - \lambda + i\varepsilon) dz \\ &= - \int_0^1 \partial_z G_k(y, z) \frac{(1 - \Phi(z))b''(z)g(z)}{b'(z)} \log(b(z) - \lambda + i\varepsilon) dz \\ &\quad - \int_0^1 G_k(y, z) \partial_z \left[\frac{(1 - \Phi(z))b''(z)g(z)}{b'(z)} \right] \log(b(z) - \lambda + i\varepsilon) dz. \end{aligned}$$

The desired bounds follow from the bound (3.2), the formula (3.3) and (3.4). $\square$

We now prove the limiting absorption principle, using the assumption that there is no discrete or generalized embedded eigenvalues.

Lemma 4.2 *There exist $\varepsilon_0, \kappa > 0$, such that the following statement holds. For all $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$, $k \in \mathbb{Z} \setminus \{0\}$, $0 < |\varepsilon| < \varepsilon_0$ and any $g \in H_k^1(I)$, we have the bound*

$$\|g + T_{k,\lambda,\varepsilon}g\|_{H_k^1(I)} \geq \kappa \|g\|_{H_k^1(I)}. \quad (4.4)$$

Proof We prove (4.4) by contradiction. Assume that there exist for $j \geq 1$, a sequence of numbers $k_j \in \mathbb{Z} \setminus \{0\}$, $\lambda_j \in \Sigma \setminus \Sigma_{\delta_0/2}$, $\varepsilon_j \in \mathbb{R} \setminus \{0\} \rightarrow 0$ and functions $g_j \in H_{k_j}^1(I)$ with $\|g_j\|_{H_{k_j}^1(I)} = 1$, satisfying $k_j \rightarrow k_* \in (\mathbb{Z} \setminus \{0\}) \cup \{\pm\infty\}$, $\lambda_j \rightarrow \lambda_* \in \overline{\Sigma \setminus \Sigma_{\delta_0}}$ as $j \rightarrow \infty$, such that

$$\|g_j + T_{k_j,\lambda_j,\varepsilon_j}g_j\|_{H_{k_j}^1(I)} \rightarrow 0, \quad \text{as } j \rightarrow \infty. \quad (4.5)$$

The bounds (4.2) and (4.5) imply that $|k_j| \lesssim 1$. Thus $k_* \in \mathbb{Z} \setminus \{0\}$. Using $\|g_j\|_{H_{k_j}^1(I)} = 1$, the bounds (4.3) and the compact embedding $W^{1,1}(I) \rightarrow L^2(I)$, we conclude that by passing to a subsequence, $T_{k_j,\lambda_j,\varepsilon_j}g_j$ converges in $H^1(I)$. In view of (4.5) we can assume that $g_j \rightarrow g$ in $H^1(I)$, where $\|g\|_{H_{k_*}^1} = 1$.

Using formula (4.1), we obtain from (4.5) that for $y \in I$,

$$g(y) + \lim_{j \rightarrow \infty} \int_0^1 G_{k_*}(y, z) \frac{b''(z)g(z)}{b(z) - \lambda + i\varepsilon_j} dz = 0. \quad (4.6)$$

Applying $k_*^2 - \frac{d^2}{dy^2}$ to (4.6), we get that for $y \in I$,

$$k_*^2 g(y) - g''(y) + \lim_{j \rightarrow \infty} \frac{(b(y) - \lambda_*) b''(y) g(y)}{(b(y) - \lambda_*)^2 + \varepsilon_j^2} + i\pi \sum_{z \in [0, 1], b(z) = \lambda} \frac{b''(z) g(z)}{|b'(z)|} \delta(y - z) = 0,$$

in the sense of distributions for $y \in (0, 1)$, which contradicts our spectral assumption that λ_* is not a generalized embedded eigenvalue for L_k . The lemma is then proved. $\square$

4.2 The Degenerate Case $\lambda \in \Sigma_{\delta_0}$

Recall the definition (3.8) for $\delta = \delta(\lambda)$. For $\lambda \in \Sigma_{\delta_0}$, $k \in \mathbb{Z} \setminus \{0\}$, $y \in I$ and $\varepsilon \in [-1/8, 1/8] \setminus \{0\}$, we denote

$$d_k(\lambda, \varepsilon) := [\lambda - b(y_*)]^{1/2} + |\varepsilon|^{1/2} \wedge \frac{1}{|k|}, \quad \varrho_k(y; \lambda + i\varepsilon) := \varrho(y; \lambda + i\varepsilon) \wedge \frac{1}{|k|}. \quad (4.7)$$

Define the weight and the associated weighted Sobolev spaces X_{N, ϱ_k} and X_{L, ϱ_k} as

$$\begin{aligned} \|g\|_{X_{N, \varrho_k}(I)} &:= \sum_{\alpha \in \{0, 1\}} (\delta + |\varepsilon|^{1/2})^{-1/2} \left\| [d_k(\lambda, \varepsilon)]^{(-7/4+\alpha)} \partial_y^\alpha g \right\|_{L^2(S_{3(\delta+|\varepsilon|^{1/2})})} \\ &\quad + \sum_{\alpha \in \{0, 1\}} \left\| \varrho_k^{-7/4+\alpha}(\cdot; \lambda + i\varepsilon) \partial_y^\alpha g \right\|_{L^\infty(I \setminus S_{3(\delta+|\varepsilon|^{1/2})})}, \end{aligned} \quad (4.8)$$

and

$$\begin{aligned} \|g\|_{X_{L, \varrho_k}(I)} &:= \sum_{\alpha \in \{0, 1\}} (\delta + |\varepsilon|^{1/2})^{-1/2} \left\| d_k^\alpha(\lambda, \varepsilon) \partial_y^\alpha g \right\|_{L^2(S_{3(\delta+|\varepsilon|^{1/2})})} \\ &\quad + \sum_{\alpha \in \{0, 1\}} \left\| d_k(\lambda, \varepsilon)^{-1} \varrho_k^{\alpha+1}(\cdot; \lambda + i\varepsilon) \partial_y^\alpha g \right\|_{L^\infty(I \setminus S_{3(\delta+|\varepsilon|^{1/2})})}. \end{aligned} \quad (4.9)$$

We remark that the choice of the weights in (4.8)–(4.9) is closely related to the behavior of the modified Green's function $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$. In (4.8), we consider the case that the “source” z is of unit distance away from the critical point y_* , where the expected decay of $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$ towards the $y = y_*$ is given roughly by $\varrho_k^2(y; \lambda + i\varepsilon)$; similar considerations apply in (4.9) if one considers the case that the source z is near y_* and study the behavior of $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$ away from y_* . The choice of exponent as $7/4$ is somewhat arbitrary, as long as it is less than 2. The endpoint case of exponent being 2 though results in a subtle logarithmic divergence that seems more technical to handle.

Fix $\varepsilon \in [-1/4, 1/4] \setminus \{0\}$, $\lambda \in \Sigma_{\delta_0}$, $k \in \mathbb{Z} \setminus \{0\}$. Recall the definition (3.8) for $\delta = \delta(\lambda) > 0$. Define for each $g \in L^2(0, 1)$ the operator

$$T_k^*(\lambda + i\varepsilon)g(y) := \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \left[1 - \varphi\left(\frac{y - y_*}{\delta_0}\right) + \varphi\left(\frac{y - y_*}{\delta}\right) \right] \frac{b''(z)g(z)}{b(z) - \lambda + i\varepsilon} dz. \quad (4.10)$$

Then we have the following bounds for $T_k^*(\lambda + i\varepsilon)$.

Lemma 4.3 *For $\varepsilon \in [-1/4, 1/4] \setminus \{0\}$, $\lambda \in \Sigma_{\delta_0}$, $k \in \mathbb{Z} \setminus \{0\}$, the operator $T_k^*(\lambda + i\varepsilon)$ satisfies the bound for $X \in \{X_{N, \varrho_k}(I), X_{L, \varrho_k}(I)\}$*

$$\|T_k^*(\lambda + i\varepsilon)g\|_X \lesssim (1 + |k|(|\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}))^{-1/4} \|g\|_X \quad \text{for all } g \in H_k^1(I). \quad (4.11)$$

Proof We provide the detailed proof only for the case $X = X_{N, \varrho_k}(I)$ as the other case is analogous. Since k, λ, ε are fixed, for simplicity of notations, we suppress the dependence on k, λ, ε to write T^* as $T_k^*(\lambda + i\varepsilon)$, and decompose for $y \in I$,

$$T^*g(y) := T_1^*g(y) + T_2^*g(y),$$

where

$$\begin{aligned} T_1^*g(y) &:= \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \left[1 - \varphi\left(\frac{z - y_*}{\delta_0}\right) \right] \frac{b''(z)g(z)}{b(z) - \lambda + i\varepsilon} dz, \\ T_2^*g(y) &:= \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi\left(\frac{z - y_*}{\delta}\right) \frac{b''(z)g(z)}{b(z) - \lambda + i\varepsilon} dz. \end{aligned}$$

It follows from the bounds on modified Green's function $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$, see Lemma 3.1, that

$$\|T_1^*g\|_{X_{N, \varrho_k}(I)} \lesssim |k|^{-1/2} \|g\|_{X_{N, \varrho_k}(I)}. \quad (4.12)$$

To prove (4.11), it suffices to prove

$$\|T_2^*g\|_{X_{N, \varrho_k}(I)} \lesssim (1 + |k|(\delta + |\varepsilon|^{1/2}))^{-1/4} \|g\|_{X_{N, \varrho_k}(I)}. \quad (4.13)$$

We assume momentarily that $|\varepsilon| \lesssim |\lambda - b(y_*)|$ and explain how to remove this assumption at the end of the proof. We decompose further for $y \in I$,

$$\begin{aligned} T_2^*g(y) &= \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi\left(\frac{z - y_*}{\delta'}\right) \varphi\left(\frac{z - y_*}{\delta}\right) \frac{b''(z)g(z)}{b(z) - \lambda + i\varepsilon} dz \\ &\quad + \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \left[1 - \varphi\left(\frac{z - y_*}{\delta'}\right) \right] \varphi\left(\frac{z - y_*}{\delta}\right) \frac{b''(z)g(z)}{b(z) - \lambda + i\varepsilon} dz \\ &:= T_{2,R}^*g(y) + T_{2,S}^*g(y), \end{aligned}$$

where we have chosen $\delta' = \delta/C_3$ with a large constant C_3 so that $|b(y) - \lambda| \approx |\lambda - b(y_*)|$ for $|y - y_*| < \delta'$.

It suffices to prove for $\diamond \in \{R, S\}$

$$\|T_{2,\diamond}^*g\|_{X_{N, \varrho_k}(I)} \lesssim (1 + |k|(|\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}))^{-1/4} \|g\|_{X_{N, \varrho_k}(I)}. \quad (4.14)$$

Step 1. We first prove (4.14) with $\diamond = R$.

Case I: $1/|k| > |\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}$. In this case for $|z - y_*| \lesssim \delta$ and $|y - y_*| \lesssim 1$ we have the bound

$$|\mathcal{G}_k(y, z; \lambda + i\varepsilon)| \lesssim \frac{\delta^2 + |\varepsilon|}{|y - y_*| + \delta + |\varepsilon|^{1/2}}, \quad |\partial_y \mathcal{G}_k(y, z; \lambda + i\varepsilon)| \lesssim \frac{\delta^2 + |\varepsilon|}{(|y - y_*| + \delta + |\varepsilon|^{1/2})^2}. \quad (4.15)$$

It follows from the bound (4.15) that

$$\|T_{2,R}^*g\|_{X_{N, \varrho_k}(I)} \lesssim (1 + |k|(|\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}))^{-1/4} \|g\|_{X_{N, \varrho_k}(I)}.$$

Case II: $1/|k| \ll |\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}$. In this case, we have for $|z - y_*| \lesssim \delta$ and $|y - y_*| \lesssim 1$ that

$$|\mathcal{G}_k(y, z; \lambda + i\varepsilon)| + |k|^{-1} |\partial_y \mathcal{G}_k(y, z; \lambda + i\varepsilon)| \lesssim |k|^{-1} e^{-|k||y - z|}. \quad (4.16)$$

The desired bound

$$\|T_{2,R}^*g\|_{X_{N, \varrho_k}(I)} \lesssim (1 + |k|(|\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}))^{-1/4} \|g\|_{X_{N, \varrho_k}(I)}$$

follows from (4.16).

Step 2. We now turn to the proof of (4.14) with $\diamond = S$ and still consider two cases.

Case I: $1/|k| > |\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}$. Denoting for $y \in I$,

$$\varphi^* \left(\frac{y - y_*}{\delta} \right) := \left[1 - \varphi \left(\frac{z - y_*}{\delta'} \right) \right] \varphi \left(\frac{z - y_*}{\delta} \right),$$

we can rewrite

$$\begin{aligned} T_{2,S}^* g(y) &= \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi^* \left(\frac{z - y_*}{\delta} \right) \frac{b''(z)g(z)}{b'(z)} \partial_z \log \frac{b(z) - \lambda + i\varepsilon}{\delta^2} \\ &= - \int_0^1 \partial_z \left[\mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi^* \left(\frac{z - y_*}{\delta} \right) \frac{b''(z)g(z)}{b'(z)} \right] \log \frac{b(z) - \lambda + i\varepsilon}{\delta^2} dz. \end{aligned} \quad (4.17)$$

As a consequence of (4.17), we also have

$$\begin{aligned} \partial_y [T_{2,S}^* g(y)] &= \partial_y \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi^* \left(\frac{z - y_*}{\delta} \right) \frac{b''(z)g(z)}{b'(z)} \partial_z \log \frac{b(z) - \lambda + i\varepsilon}{\delta^2} dz \\ &= - \int_0^1 \left[\partial_y (\partial_z + \partial_y) \mathcal{G}_k(y, z; \lambda, \varepsilon) \varphi^* \left(\frac{z - y_*}{\delta} \right) \frac{b''(z)g(z)}{b'(z)} \right] \log \frac{b(z) - \lambda + i\varepsilon}{\delta^2} dz \\ &\quad + \int_0^1 \left[\partial_y^2 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi^* \left(\frac{z - y_*}{\delta} \right) \frac{b''(z)g(z)}{b'(z)} \right] \log \frac{b(z) - \lambda + i\varepsilon}{\delta^2} dz \\ &\quad - \int_0^1 \partial_y \mathcal{G}_k(y, z; \lambda + i\varepsilon) \partial_z \left[\varphi^* \left(\frac{z - y_*}{\delta} \right) \frac{b''(z)g(z)}{b'(z)} \right] \log \frac{b(z) - \lambda + i\varepsilon}{\delta^2} dz. \end{aligned} \quad (4.18)$$

Note that on the support of $\varphi^* \left(\frac{z - y_*}{\delta} \right)$, we have

$$|b'(z)| \approx \delta, \quad \varrho(z; \lambda + i\varepsilon) \approx \delta.$$

The desired bound (4.14) for $\diamond = S$ follows from (4.17)–(4.18) and Corollary 3.3, and we have, in addition,

$$\begin{aligned} &(\delta + |\varepsilon|^{1/2})^{-1/2} \left\| \partial_y \left\{ \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi^* \left(\frac{z - y_*}{\delta} \right) \frac{b''(z)g(z)}{b'(z)} \log \frac{b(z) - \lambda + i\varepsilon}{\delta^2} \right\} \right\|_{L^2(S_{3(\delta + |\varepsilon|^{1/2})})} \\ &\lesssim \delta^{-1/4} \left[1 + |k| \left(|\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2} \right) \right]^{-1/4} \|g\|_{X_{N, \varrho_k}(I)}. \end{aligned} \quad (4.19)$$

Case II: $1/|k| \ll |\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}$. This case is analogous to *Case I*, using Lemma 3.1 and Lemma 3.2.

Finally we turn to the assumption that $|\varepsilon|^{1/2} \lesssim \delta$. Suppose $|\varepsilon|^{1/2} \gg \delta$, then the factor $\frac{1}{b(z) - \lambda + i\varepsilon}$ is not truly singular, and the desired bounds (4.13) follow directly from the bounds on the modified Green's function $\mathcal{G}_k(y, z; \lambda + i\varepsilon)$ from Lemma 3.1 and Lemma 3.2. Indeed, we have the stronger bound

$$\|T_2^* g\|_{X_{N, \varrho_k}(I)} \lesssim \frac{\delta}{\sqrt{|\varepsilon|}} \|g\|_{X_{N, \varrho_k}(I)}, \quad (4.20)$$

which will be useful below. $\square$

The following limiting absorption principle plays an essential role in establishing the vorticity depletion phenomenon.

Lemma 4.4 *There exist positive numbers ε_0, κ such that the following statement holds.*

For $\varepsilon \in [-\varepsilon_0, \varepsilon_0] \setminus \{0\}$, $\lambda \in \Sigma_{\delta_0}$, $k \in \mathbb{Z} \setminus \{0\}$, and $X \in \{X_{N, \varrho_k}(I), X_{L, \varrho_k}(I)\}$,

$$\|(I + T_k^*(\lambda + i\varepsilon))g\|_X \geq \kappa \|g\|_X \quad \text{for all } g \in H_k^1(I). \quad (4.21)$$

Proof We only consider the case $X = X_{N, \partial k}(I)$ as the other case is analogous. We prove (4.21) by a contradiction argument. Assume (4.21) does not hold for any $\varepsilon_0 > 0$. Then there exist for $\ell \in \mathbb{Z} \cap [1, \infty)$,

$$\lambda_\ell \rightarrow \lambda_* \in \Sigma_{\delta_0}, \quad \varepsilon_\ell \neq 0 \quad \text{with} \quad \varepsilon_\ell \rightarrow 0, \quad k_\ell \rightarrow k_* \in (\mathbb{Z} \setminus \{0\}) \cup \{\pm\infty\},$$

and functions g_ℓ satisfying

$$\|g_\ell\|_{X_{N, \partial k_\ell}(I)} = 1 \quad (4.22)$$

such that

$$\|(I + T_{k_\ell}^*(\lambda_\ell + i\varepsilon_\ell))g_\ell\|_{X_{N, \partial k_\ell}(I)} \rightarrow 0. \quad (4.23)$$

We can assume that $\lambda_* = b(y_*)$, otherwise the proof follows from the argument in the non-degenerate case. We consider several cases.

Case I: $\limsup_{\ell \rightarrow \infty} \|g_\ell\|_{H^1(I \setminus S_{\delta_0})} > 0$. By the bound (4.12), we can assume that $k_* \in \mathbb{Z} \setminus \{0\}$. By the bounds (4.22) and (4.23), we can assume (passing to a subsequence if necessary) that

$$g_\ell \rightarrow g \quad \text{in} \quad H_{\text{loc}}^1(I \setminus \{y_*\}) \quad \text{as} \quad \ell \rightarrow \infty, \quad g(0) = g(1) = 0.$$

Then it follows from (4.22) and (4.23) that g satisfies

$$|g(y)| \lesssim |y - y_*|^{7/4},$$

and for $y \in (0, 1)$,

$$(k_*^2 - \partial_y^2)g(y) + \frac{b''(y)}{b(y) - b(y_*)}g(y) = 0,$$

which imply that $b(y_*)$ is an embedded eigenvalue for L_k , a contradiction to the spectral assumption.

Case II: $\limsup_{\ell \rightarrow \infty} \|g_\ell\|_{H^1(I \setminus S_{\delta_0})} = 0$. By the bound (4.11) we can assume that $|k_\ell|(\delta_\ell + |\varepsilon_\ell|^{1/2}) \lesssim 1$. In this case, using (4.23), we obtain that (passing to a subsequence if necessary)

$$\begin{aligned} & \|(|\lambda_\ell - b(y_*)| + |\varepsilon|)^{-9/8} g_\ell\|_{L^2([y_* - \delta_\ell - |\varepsilon|^{1/2}, y_* + \delta_\ell + |\varepsilon|^{1/2}])} \\ & + \left\| (|\lambda_\ell - b(y_*)| + |\varepsilon|)^{-5/8} \partial_y g_\ell \right\|_{L^2([y_* - \delta_\ell - |\varepsilon|^{1/2}, y_* + \delta_\ell + |\varepsilon|^{1/2}])} \geq \sigma > 0, \end{aligned}$$

where we recall from (3.8) that

$$\delta_\ell \approx |\lambda_\ell - b(y_*)|^{1/2}.$$

We divide into several subcases.

Subcase II.1: $|\varepsilon_\ell|^{1/2} \approx \delta_\ell$ for a subsequence.

Define the change of variables for $\ell \geq 1$, $y \in I$,

$$y - y_* = \delta_\ell Y, \quad g_\ell(y) := (|\lambda_\ell - b(y_*)| + |\varepsilon_\ell|)^{7/8} H_\ell(Y). \quad (4.24)$$

It follows from (4.19) that we can extract a nontrivial limit $H \in H^1(\mathbb{R})$ of H_ℓ satisfying for $Y \in \mathbb{R}$,

$$(\beta^2 - \partial_Y^2)H(Y) + \frac{b''(y_*)}{b''(y_*)Y^2/2 + \gamma + i\alpha}H(Y) = 0,$$

where $\beta \in \mathbb{R}$, $\alpha, \gamma \in \mathbb{R} \setminus \{0\}$. This is impossible since the shear flow $(b''(y_*)Y^2/2, 0)$, $Y \in \mathbb{R}$ is spectrally stable, thanks to Rayleigh's inflection point criteria.

Subcase II.2: $|\varepsilon_\ell|^{1/2} = o(\delta_\ell)$ for a subsequence. Passing to a subsequence and using rescaling as in (4.24) we can extract a nontrivial limit $H \in H^1(\mathbb{R})$, such that

$$(\beta^2 - \partial_Y^2)H(Y) + \lim_{\varepsilon \rightarrow 0} \frac{b''(y_*)}{b''(y_*)Y^2/2 + \gamma + i\varepsilon} H(Y) = 0.$$

This is again impossible since the shear flow $(b''(y_*)Y^2/2, 0)$, $Y \in \mathbb{R}$ is spectrally stable.

Subcase II.3: $\delta_\ell = o(|\varepsilon_\ell|^{1/2})$ for a subsequence. This case is not possible thanks to the bound (4.20). The lemma is now proved. $\square$

5 Bounds on $\psi_{k,\varepsilon}^t$: The Non-degenerate Case

In this section we obtain bounds on $\psi_{k,\varepsilon}^t(y, \lambda)$ in the non-degenerate case, i.e. when $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$. Since the arguments are analogous to those in [15], we will be brief in the proofs, and provide only comments on the main ideas involved.

We begin with the following preliminary bounds.

Lemma 5.1 For $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$, $k \in \mathbb{Z} \setminus \{0\}$, $\iota \in \{\pm\}$ and $0 < \varepsilon < \varepsilon_0$, we have the bounds

$$\|\psi_{k,\varepsilon}^t(\cdot, \lambda)\|_{H_k^1(I)} \lesssim |k|^{-1/2} \|\omega_{0k}\|_{H_k^1(I)}. \quad (5.1)$$

Proof The desired bounds (5.1) follow directly from the Rayleigh equation (2.5) and Lemma 4.2, once we use the Green's function G_k to invert $k^2 - \partial_y^2$ and formulate (2.5) as an integral equation. $\square$

To obtain control on $\partial_\lambda \psi_{k,\varepsilon}^t(\cdot, \lambda)$ for $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$, we take derivative in (2.5), and obtain that

$$(k^2 - \partial_y^2) \partial_\lambda \psi_{k,\varepsilon}^t(y, \lambda) + \frac{b''(y) \partial_\lambda \psi_{k,\varepsilon}^t(y, \lambda)}{b(y) - \lambda + i\varepsilon} = \frac{\omega_0^k(y)}{(b(y) - \lambda + i\varepsilon)^2} - \frac{b''(y) \psi_{k,\varepsilon}^t(z, \lambda)}{(b(y) - \lambda + i\varepsilon)^2} \quad (5.2)$$

for $y \in I$ with zero boundary value at $y \in \{0, 1\}$. Reformulating (5.2) as an integral equation, we obtain that

$$\begin{aligned} \partial_\lambda \psi_{k,\varepsilon}^t(y, \lambda) + \int_0^1 G_k(y, z) \frac{b''(z) \partial_\lambda \psi_{k,\varepsilon}^t(z, \lambda)}{b(z) - \lambda + i\varepsilon} dz \\ = \int_0^1 G_k(y, z) \frac{\omega_0^k(z)}{(b(z) - \lambda + i\varepsilon)^2} dz - \int_0^1 G_k(y, z) \frac{b''(z) \psi_{k,\varepsilon}^t(z, \lambda)}{(b(z) - \lambda + i\varepsilon)^2} dz. \end{aligned} \quad (5.3)$$

Recall the definition of the smooth cutoff function Φ below (4.1). We have the following bounds for $\partial_\lambda \psi_{k,\varepsilon}^t(y, \lambda)$ when $\lambda \in \Sigma \setminus \Sigma_{\delta_0}$.

Lemma 5.2 For $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$, $k \in \mathbb{Z} \setminus \{0\}$, $\iota \in \{\pm\}$ and $0 < \varepsilon < \varepsilon_0$, $\partial_\lambda \psi_{k,\varepsilon}^t(y, \lambda)$ satisfies the following decomposition

$$\begin{aligned} \partial_\lambda \psi_{k,\varepsilon}^t(y, \lambda) = & \left[\frac{b'(y_0) \omega_0^k(y)}{|b'(y)|^2} - \frac{b''(y) \psi_{k,\varepsilon}^t(y, \lambda)}{|b'(y)|^2} \right] (1 - \Phi(y)) \log(b(y) - \lambda + i\varepsilon) \\ & + \sum_{\sigma=0,1} \omega_0^k(\sigma) \Psi_{k,\sigma,\varepsilon}^t(y, \lambda) \log(b(\sigma) - \lambda + i\varepsilon) + \mathcal{R}_{\sigma,k,y_0,\varepsilon}^t(y). \end{aligned}$$

In the above for $\sigma \in \{0, 1\}$, $\iota \in \{\pm\}$, $0 < \varepsilon < \varepsilon_0$, and $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$,

$$\|\mathcal{R}_{\sigma,k,y_0,\varepsilon}^t\|_{H_k^1(I)} \lesssim |k|^{1/2} \|\omega_{0k}\|_{H_k^2(I)}, \quad \|\Psi_{k,\sigma,\varepsilon}^t(\cdot, \lambda)\|_{H_k^1(I)} \lesssim |k|^{-1/2}.$$

Proof The basic idea is to expand the right hand side of (5.3) using integration by parts, and apply Lemma 4.2 after removing the most singular parts. Indeed, denoting schematically,

$$\mathcal{U} := \int_0^1 G_k(y, z) \frac{\omega_0^k(z)}{(b(z) - \lambda + i\epsilon)^2} dz - \int_0^1 G_k(y, z) \frac{b''(z)\psi_{k,\epsilon}^t(z, \lambda)}{(b(z) - \lambda + i\epsilon)^2} dz,$$

we note that $\partial_\lambda \psi_{k,\epsilon}^t(y, \lambda) - \mathcal{U}$ satisfies the equation (recalling (4.1) for the definition of $T_{k,\lambda,\epsilon}$),

$$(I + T_{k,\lambda,\epsilon}) [\partial_\lambda \psi_{k,\epsilon}^t(y, \lambda) - \mathcal{U}] = -T_{k,\lambda,\epsilon} \mathcal{U}. \quad (5.4)$$

The term $T_{k,\lambda,\epsilon} \mathcal{U} \in H_k^1(I)$ (noting however that for the boundary terms we need to track the singular coefficient $\log(b(\sigma) - \lambda + i\epsilon)$, $\sigma \in \{0, 1\}$), and we can apply Lemma 4.2 to (5.4) in order to obtain the desired conclusions. We refer to [15] for the detailed proof. $\square$

To obtain bounds on $\partial_\lambda^2 \psi_{k,\epsilon}^t(y, \lambda)$ for $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$, we take two derivatives in (2.5) and obtain that

$$\begin{aligned} & (k^2 - \partial_y^2) \partial_\lambda^2 \psi_{k,\epsilon}^t(y, \lambda) + \frac{b''(y) \partial_\lambda^2 \psi_{k,\epsilon}^t(y, \lambda)}{b(y) - \lambda + i\epsilon} \\ &= 2 \frac{\omega_0^k(y)}{(b(y) - \lambda + i\epsilon)^3} - 2 \frac{b''(y) \psi_{k,\epsilon}^t(z, \lambda)}{(b(y) - \lambda + i\epsilon)^3} + \frac{b''(y) \partial_\lambda \psi_{k,\epsilon}^t(z, \lambda)}{(b(y) - \lambda + i\epsilon)^2} \end{aligned} \quad (5.5)$$

for $y \in I$ with zero boundary value at $y \in \{0, 1\}$. We can reformulate (5.5) in the integral form for $y \in I$, as

$$\begin{aligned} & \partial_\lambda^2 \psi_{k,\epsilon}^t(y, \lambda) + \int_0^1 G_k(y, z) \frac{b''(z) \partial_\lambda^2 \psi_{k,\epsilon}^t(z, \lambda)}{b(z) - \lambda + i\epsilon} dz \\ &= \int_0^1 G_k(y, z) \left[2 \frac{\omega_0^k(z)}{(b(z) - \lambda + i\epsilon)^3} - 2 \frac{b''(z) \psi_{k,\epsilon}^t(z, \lambda)}{(b(z) - \lambda + i\epsilon)^3} + \frac{b''(z) \partial_\lambda \psi_{k,\epsilon}^t(z, \lambda)}{(b(z) - \lambda + i\epsilon)^2} \right] dz. \end{aligned} \quad (5.6)$$

We have the following bounds on $\partial_\lambda^2 \psi_{k,\epsilon}^t(y, \lambda)$ for $\lambda \in \Sigma \setminus \Sigma_{\delta_0/2}$.

Lemma 5.3 For $k \in \mathbb{Z} \setminus \{0\}$, $\epsilon \in \{\pm\}$ and $0 < \epsilon < \epsilon_0$, we have the following bound

$$\begin{aligned} & \left\| \partial_\lambda^2 \psi_{k,\epsilon}^t(y, \lambda) - \frac{\omega_0^k(1) \Phi_{k,\epsilon}^{1\epsilon}(y, \lambda)}{b(1) - \lambda + i\epsilon} - \frac{\omega_0^k(0) \Phi_{k,\epsilon}^{0\epsilon}(y, \lambda)}{b(0) - \lambda + i\epsilon} \right. \\ & \quad \left. - \frac{b''(y) \psi_{k,\epsilon}^t(y, \lambda) - \omega_0^k(y)}{|b'(y)|^2 (b(y) - \lambda + i\epsilon)} \right\|_{L^2(y \in I, \lambda \in \Sigma \setminus \Sigma_{\delta_0/2})} \\ & \lesssim |k|^{3/2} \|\omega_{0k}\|_{H_k^3(I)}. \end{aligned}$$

In the above the functions $\Phi_{k,\epsilon}^{\sigma\epsilon}$, $\sigma \in \{0, 1\}$ satisfy the equation for $y \in I$

$$\begin{aligned} (I + T_{k,\lambda,\epsilon}) \Phi_{k,\epsilon}^{1\epsilon} &= \frac{\sinh(ky)}{|b'(1)|^2 \sinh k}, \\ (I + T_{k,\lambda,\epsilon}) \Phi_{k,\epsilon}^{0\epsilon} &= \frac{\sinh(k(1-y))}{|b'(0)|^2 \sinh k}. \end{aligned}$$

Proof The main idea of the proof is to expand the right-hand side of (5.6) and apply Lemma 4.2 after removing the most singular terms. Indeed, denoting schematically,

$$\mathcal{U}^* = \int_0^1 G_k(y, z) \left[2 \frac{\omega_0^k(z)}{(b(z) - \lambda + i\epsilon)^3} - 2 \frac{b''(z) \psi_{k,\epsilon}^t(z, \lambda)}{(b(z) - \lambda + i\epsilon)^3} + \frac{b''(z) \partial_\lambda \psi_{k,\epsilon}^t(z, \lambda)}{(b(z) - \lambda + i\epsilon)^2} \right] dz,$$

we have

$$(I + T_{k,\lambda,\iota\varepsilon}) [\partial_\lambda^2 \psi_{k,\varepsilon}^\iota(y, \lambda) - \mathcal{U}^* + T_{k,\lambda,\iota\varepsilon} \mathcal{U}^*] = [T_{k,\lambda,\iota\varepsilon}]^2 \mathcal{U}^*.$$

We note that $\partial_\lambda^2 \psi_{k,\varepsilon}^\iota(y, \lambda) - \mathcal{U}^* + T_{k,\lambda,\iota\varepsilon} \mathcal{U}^* \in H_k^1(I)$ (however we again need to track the singularities in λ in the boundary terms, involving $\log(b(\sigma) - \lambda + i\varepsilon)$ and $1/(b(\sigma) - \lambda + i\varepsilon)$ for $\sigma \in \{0, 1\}$), and we can apply Lemma 4.2 in order to obtain the desired conclusions. We refer to [15] for the detailed proof. $\square$

6 Bounds on $\psi_{k,\varepsilon}^\iota$: The Degenerate Case

In this section, we use the limiting absorption principle to study the Rayleigh equation (2.5) for $\lambda \in \Sigma_{\delta_0}$. More precisely, write for $k \in \mathbb{Z} \setminus \{0\}$, $\iota \in \{\pm\}$, $\lambda \in \Sigma_{\delta_0}$, $0 < \varepsilon < \varepsilon_0$ (recall the definition of ε_0 from Lemma 4.4),

$$\psi_{k,\varepsilon}^\iota(y, \lambda) = \phi_{k,\varepsilon}^\iota(y, \lambda) + \Psi(y) \frac{1}{b''(y)} \omega_{0k}(y), \quad (6.1)$$

where $\Psi \in C_c^\infty(S_{3\delta_0})$ and $\Psi \equiv 1$ on $S_{2\delta_0}$. Recall that $S_d = [y_* - d, y_* + d]$ for $d > 0$ from (3.5). Then $\phi_{k,\varepsilon}^\iota(y, \lambda)$ satisfies for $y \in I$,

$$(k^2 - \partial_y^2) \phi_{k,\varepsilon}^\iota(y, \lambda) + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \phi_{k,\varepsilon}^\iota(y, \lambda) = g_{k,\varepsilon}^\iota(y, \lambda), \quad (6.2)$$

where for $k \in \mathbb{Z} \setminus \{0\}$, $\iota \in \{\pm\}$, $\lambda \in \Sigma_{\delta_0}$, $0 < \varepsilon < \varepsilon_0$

$$g_{k,\varepsilon}^\iota(y, \lambda) := \frac{1 - \Psi(y)}{b(y) - \lambda + i\varepsilon} \omega_{0k}(y) - (k^2 - \partial_y^2) \left[\frac{\Psi(y)}{b''(y)} \omega_{0k}(y) \right]. \quad (6.3)$$

Our main results are bounds for the functions $\phi_{k,\varepsilon}^\iota(y, \lambda)$. We begin with the following preliminary bounds.

Lemma 6.1 *Assume that $k \in \mathbb{Z} \setminus \{0\}$, $\lambda \in \Sigma_{\delta_0}$ and let $\phi_{k,\varepsilon}^\iota(y, \lambda)$ with $\iota \in \{\pm\}$, $\varepsilon \in (0, \varepsilon_0)$ be as defined in (6.1)–(6.2). Recall from (3.8) and (4.7) that*

$$\delta := \delta(\lambda) = 8\sqrt{|\lambda - b(y_*)|/|b''(y_*)|}, \quad d_k = d_k(\lambda, \varepsilon) := [|\lambda - b(y_*)|^{1/2} + |\varepsilon|^{1/2}] \wedge \frac{1}{|k|}.$$

We have the bounds for $k \in \mathbb{Z} \setminus \{0\}$, $\varepsilon \in (0, \varepsilon_0)$, $\iota \in \{\pm\}$, $\lambda \in \Sigma_{\delta_0}$,

$$\begin{aligned} & \sum_{\alpha \in \{0,1\}} \left\| d_k^{-7/4+\alpha} \partial_y^\alpha \phi_{k,\varepsilon}^\iota(y, \lambda) \right\|_{L^2([y_*-3(\delta+|\varepsilon|^{1/2}), y_*+3(\delta+|\varepsilon|^{1/2})])} (\delta + |\varepsilon|^{1/2})^{-1/2} \\ & + \sum_{\alpha \in \{0,1\}} \left\| (|y - y_*| \wedge d_k)^{-7/4+\alpha} \partial_y^\alpha \phi_{k,\varepsilon}^\iota(y, \lambda) \right\|_{L^\infty([0,1] \setminus [y_*-3(\delta+|\varepsilon|^{1/2}), y_*+3(\delta+|\varepsilon|^{1/2})])} \\ & \lesssim |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \end{aligned} \quad (6.4)$$

Define for $y \in [0, 1]$, $k \in \mathbb{Z} \setminus \{0\}$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$,

$$\psi_k(y, \lambda) := \lim_{\varepsilon \rightarrow 0^+} [\psi_{k,\varepsilon}^+(y, \lambda) - \psi_{k,\varepsilon}^-(y, \lambda)] = \lim_{\varepsilon \rightarrow 0^+} [\phi_{k,\varepsilon}^+(y, \lambda) - \phi_{k,\varepsilon}^-(y, \lambda)].$$

Then we have the bounds for $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$,

$$\begin{aligned} & \sum_{\alpha \in \{0,1\}} \left\| (\delta \wedge |k|^{-1})^{-7/4+\alpha} \partial_y^\alpha \psi_k(y, \lambda) \right\|_{L^2([y_*-3\delta, y_*+3\delta])} \delta^{-1/2} \\ & + \sum_{\alpha \in \{0,1\}} \left\| (\delta \wedge |k|^{-1})^{-11/4} \left(|y - y_*| \wedge \frac{1}{|k|} \right)^{1+\alpha} \partial_y^\alpha \psi_k(y, \lambda) \right\|_{L^\infty([0,1] \setminus [y_*-3\delta, y_*+3\delta])} \\ & \lesssim |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \end{aligned} \quad (6.5)$$

Proof It follows from (6.3) and our assumptions on the initial data ω_{0k} that we have the bound for $k \in \mathbb{Z} \setminus \{0\}$, $\iota \in \{\pm 1\}$, $0 < \varepsilon < \varepsilon_0$ and $\lambda \in \Sigma_{\delta_0}$,

$$\|g_{k,\varepsilon}^\iota(y, \lambda)\|_{C(I)} \lesssim |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}.$$

We can reformulate (6.2) in the integral form as (recall the definition of $T^*(\lambda + i\varepsilon)$ from (4.10))

$$\phi_{k,\varepsilon}^\iota(y, \lambda) + T_k^*(\lambda + i\varepsilon)\phi_{k,\varepsilon}^\iota(y, \lambda) = \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) g_{k,\varepsilon}^\iota(z, \lambda) dz$$

for $y \in I$. By Lemma 4.4, we obtain the bound

$$\|\phi_{k,\varepsilon}^\iota(\cdot, \lambda)\|_{X_{N, \varrho_k}(I)} \lesssim \left\| \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) g_{k,\varepsilon}^\iota(z, \lambda) dz \right\|_{X_{N, \varrho_k}} \lesssim |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)},$$

which, by the definition of the space X_{N, ϱ_k} , see (4.8), implies the desired bounds (6.4).

For applications below on isolating the singularity at $\lambda = b(y)$, we fix $\varphi_\delta(y) \in C_c^\infty(S_{2\delta})$ as

$$\varphi_\delta(y) := \varphi\left(\frac{y}{\delta}\right) \left[1 - \varphi\left(\frac{y}{\delta'}\right)\right] \quad (6.6)$$

for $y \in I$, with $\delta' := \delta/M$ and an $M \gg 1$ sufficiently large such that $|b(y) - \lambda| \approx |\lambda - b(y_*)|$ for $|y - y_*| < \delta/M$.

To prove (6.5), we note from (6.2) that $\phi_{k,\varepsilon}^+(y, \lambda) - \phi_{k,\varepsilon}^-(y, \lambda)$ satisfies the equation for $y \in I$.

$$\begin{aligned} & (k^2 - \partial_y^2) [\phi_{k,\varepsilon}^+(y, \lambda) - \phi_{k,\varepsilon}^-(y, \lambda)] + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} [\phi_{k,\varepsilon}^+(y, \lambda) - \phi_{k,\varepsilon}^-(y, \lambda)] \\ & = g_{k,\varepsilon}^+(y, \lambda) - g_{k,\varepsilon}^-(y, \lambda) - \left[\frac{b''(y)}{b(y) - \lambda + i\varepsilon} - \frac{b''(y)}{b(y) - \lambda - i\varepsilon} \right] \phi_{k,\varepsilon}^-(y, \lambda). \end{aligned}$$

Denote for $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $\varepsilon \in (0, \varepsilon_0)$ and $y \in I$ the function $h_{k,\varepsilon}(y, \lambda)$ as the solution to

$$\begin{aligned} & (k^2 - \partial_y^2) h_{k,\varepsilon}(y, \lambda) + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} h_{k,\varepsilon}(y, \lambda) \\ & = \varphi_\delta(y) \left[\frac{b''(y)}{b(y) - \lambda - i\varepsilon} - \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \right] \phi_{k,\varepsilon}^-(y, \lambda), \end{aligned} \quad (6.7)$$

with zero Dirichlet boundary condition. Then it is clear that for $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $y \in I$,

$$\psi_k(y, \lambda) = \lim_{\varepsilon \rightarrow 0^+} h_{k,\varepsilon}(y, \lambda).$$

We can reformulate (6.7) as the following integral equation for $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $y \in I$,

$$\begin{aligned} & h_{k,\varepsilon}(y, \lambda) + T_k^*(\lambda + i\varepsilon)h_{k,\varepsilon}(y, \lambda) \\ &= - \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon)\varphi_\delta(z) \left[\frac{b''(z)}{b(z) - \lambda + i\varepsilon} - \frac{b''(z)}{b(z) - \lambda - i\varepsilon} \right] \phi_{k,\varepsilon}^-(z, \lambda) dz. \end{aligned}$$

It follows from the bound (6.4) that for $|\varepsilon| \lesssim (\delta \wedge \frac{1}{|k|})^4$,

$$\begin{aligned} & \left\| \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon)\varphi_\delta(z) \left[\frac{b''(z)}{b(z) - \lambda + i\varepsilon} - \frac{b''(z)}{b(z) - \lambda - i\varepsilon} \right] \phi_{k,\varepsilon}^-(z, \lambda) dz \right\|_{X_{L,\varrho_k}} \\ & \lesssim \left(\delta \wedge \frac{1}{|k|} \right)^{7/4}. \end{aligned}$$

The desired bound (6.5) then follows from Lemma 4.4 with $X = X_{L,\varrho_k}$. $\square$

To obtain higher order regularity bounds (in λ) of $\phi_{k,\varepsilon}^t(\cdot, \lambda)$, we take the derivative ∂_λ in (6.2). It follows that $\partial_\lambda \phi_{k,\varepsilon}^t(y, \lambda)$ satisfies for $y \in I$,

$$\left[k^2 - \partial_y^2 + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \right] \partial_\lambda \phi_{k,\varepsilon}^t(y, \lambda) = - \frac{b''(y)}{(b(y) - \lambda + i\varepsilon)^2} \phi_{k,\varepsilon}^t(y, \lambda) + \partial_\lambda g_{k,\varepsilon}^t(y, \lambda), \quad (6.8)$$

with zero Dirichlet boundary condition.

Recall the definition of φ_δ from (6.6). We have the following bounds on $\partial_\lambda \phi_{k,\varepsilon}^t(y, \lambda)$.

Lemma 6.2 Assume that $k \in \mathbb{Z} \setminus \{0\}$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$. Let $\psi_{k,\varepsilon}^t(y, \lambda)$ and $\phi_{k,\varepsilon}^t(y, \lambda)$ with $\iota \in \{\pm\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$ be as defined in (2.5) and (6.1) respectively. Recall from (3.8) that

$$\delta := \delta(\lambda) = 8\sqrt{|\lambda - b(y_*)|/b''(y_*)}.$$

Denote for $y \in [0, 1]$, $\iota \in \{\pm\}$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$,

$$\begin{aligned} \Lambda_{1,\varepsilon}^t(y, \lambda) &:= \phi_{k,\varepsilon}^t(y, \lambda)\varphi_\delta(y) \frac{b''(y)}{(b'(y))^2} \log \frac{b(y) - \lambda + i\varepsilon}{\delta^2}, \\ \Lambda_1(y, \lambda) &:= \psi_{k,\varepsilon}(y, \lambda)\varphi_\delta(y) \frac{b''(y)}{(b'(y))^2} \log \frac{b(y) - \lambda}{\delta^2}. \end{aligned}$$

We have the bounds for $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$, $\iota \in \{\pm\}$, and $\lambda \in \Sigma_{\delta_0}$ that

$$\begin{aligned} & \sum_{\alpha \in \{0,1\}} \left\| (\delta \wedge |k|^{-1})^{1/4+\alpha} \partial_y^\alpha [\partial_\lambda \phi_{k,\varepsilon}^t(y, \lambda) - \Lambda_{1,\varepsilon}^t(y, \lambda)] \right\|_{L^2([y_*-3\delta, y_*+3\delta])} \delta^{-1/2} \\ & + \sum_{\alpha \in \{0,1\}} \left\| (\delta \wedge |k|^{-1})^2 \left(|y - y_*| \wedge \frac{1}{|k|} \right)^{-7/4+\alpha} \partial_y^\alpha \partial_\lambda \phi_{k,\varepsilon}^t(y, \lambda) \right\|_{L^\infty([0,1] \setminus [y_*-3\delta, y_*+3\delta])} \\ & \lesssim |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \end{aligned} \quad (6.9)$$

In addition, we have the bounds for $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$ and $k \in \mathbb{Z} \setminus \{0\}$,

$$\begin{aligned} & \sum_{\alpha \in \{0,1\}} \left\| (\delta \wedge |k|^{-1})^{1/4+\alpha} \partial_y^\alpha [\partial_\lambda \psi_k(y, \lambda) - \Lambda_1(y, \lambda)] \right\|_{L^2([y_*-3\delta, y_*+3\delta])} \delta^{-1/2} \\ & + \sum_{\alpha \in \{0,1\}} \left\| (\delta \wedge |k|^{-1})^{-3/4} \left(|y - y_*| \wedge \frac{1}{|k|} \right)^{1+\alpha} \partial_y^\alpha \partial_\lambda \psi_k(y, \lambda) \right\|_{L^\infty([0,1] \setminus [y_*-3\delta, y_*+3\delta])} \\ & \lesssim |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \end{aligned} \quad (6.10)$$

Proof Define for $k \in \mathbb{Z} \setminus \{0\}$, $\iota \in \{\pm\}$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$, $y \in I$,

$$\begin{aligned} \partial_\lambda \phi_{k,\varepsilon}^t(y, \lambda) &:= \phi_{k,\varepsilon}^t(y, \lambda; 1) \\ &+ \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \left[\frac{-b''(z)}{(b(z) - \lambda + i\varepsilon)^2} \phi_{k,\varepsilon}^t(z, \lambda) + \partial_\lambda g_{k,\varepsilon}^t(z, \lambda) \right] dz. \end{aligned} \quad (6.11)$$

It follows from (6.8) that $\phi_{k,\varepsilon}^t(y, \lambda; 1)$ satisfies for $y \in I$,

$$\begin{aligned} & \phi_{k,\varepsilon}^t(y, \lambda; 1) + T_k^*(\lambda + i\varepsilon) \phi_{k,\varepsilon}^t(y, \lambda; 1) \\ &= -T_k^*(\lambda + i\varepsilon) \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \left[-\frac{b''(z)}{(b(z) - \lambda + i\varepsilon)^2} \phi_{k,\varepsilon}^t(z, \lambda) + \partial_\lambda g_{k,\varepsilon}^t(z, \lambda) \right] dz. \end{aligned}$$

Denote for $k \in \mathbb{Z} \setminus \{0\}$, $\iota \in \{\pm\}$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$, $z \in I$,

$$\begin{aligned} h_{k,\varepsilon}^t(z, \lambda; 1) &:= \frac{b''(z)}{(b(z) - \lambda + i\varepsilon)^2} \varphi_\delta(z) \phi_{k,\varepsilon}^t(z, \lambda), \\ h_{k,\varepsilon}^t(z, \lambda; 2) &:= \frac{b''(z)}{(b(z) - \lambda + i\varepsilon)^2} (1 - \varphi_\delta(z)) \phi_{k,\varepsilon}^t(z, \lambda), \quad h_{k,\varepsilon}^t(z, \lambda; 3) := \partial_\lambda g_{k,\varepsilon}^t(z, \lambda). \end{aligned}$$

It follows from the bound (6.4) and Lemma 3.1 that for $j \in \{2, 3\}$

$$\left\| T_k^*(\lambda + i\varepsilon) \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) h_{k,\varepsilon}^t(z, \lambda; j) dz \right\|_{X_{N, \varrho_k}} \lesssim (\delta \wedge |k|^{-1})^{-2} |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \quad (6.12)$$

Using integration by parts argument similar to (4.17)–(4.18), we have also

$$\left\| T_k^*(\lambda + i\varepsilon) \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) h_{k,\varepsilon}^t(z, \lambda; 1) dz \right\|_{X_{N, \varrho_k}} \lesssim (\delta \wedge |k|^{-1})^{-2} |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \quad (6.13)$$

It follows from (6.12)–(6.13) and Lemma 4.4 that for $\lambda \setminus \{b(y_*)\}$,

$$\left\| \phi_{k,\varepsilon}^t(y, \lambda; 1) \right\|_{X_{N, \varrho_k}} \lesssim (\delta \wedge |k|^{-1})^{-2} |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \quad (6.14)$$

The desired bound (6.9) follows, as a consequence of (6.14) and (6.11).

Using (6.8), we get that for $y \in I$,

$$\begin{aligned} & \left[k^2 - \partial_y^2 + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \right] \left[\partial_\lambda \phi_{k,\varepsilon}^+(y, \lambda) - \partial_\lambda \phi_{k,\varepsilon}^-(y, \lambda) \right] \\ &= - \left[\frac{b''(y)}{(b(y) - \lambda + i\varepsilon)^2} \phi_{k,\varepsilon}^+(y, \lambda) - \frac{b''(y)}{(b(y) - \lambda - i\varepsilon)^2} \phi_{k,\varepsilon}^-(y, \lambda) \right] \\ &+ \left[\partial_\lambda g_{k,\varepsilon}^+(y, \lambda) - \partial_\lambda g_{k,\varepsilon}^-(y, \lambda) \right] - \left[\frac{b''(y)}{b(y) - \lambda + i\varepsilon} - \frac{b''(y)}{b(y) - \lambda - i\varepsilon} \right] \partial_\lambda \phi_{k,\varepsilon}^-(y, \lambda), \end{aligned}$$

with zero Dirichlet boundary condition.

Denoting for $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$ and $y \in I$, $D\phi_{k,\varepsilon}(y, \lambda)$ as the solution to

$$\begin{aligned} & \left[k^2 - \partial_y^2 + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \right] D\phi_{k,\varepsilon}(y, \lambda) \\ &= -\varphi_\delta(y) \left[\frac{b''(y)}{(b(y) - \lambda + i\varepsilon)^2} \phi_{k,\varepsilon}^+(y, \lambda) - \frac{b''(y)}{(b(y) - \lambda - i\varepsilon)^2} \phi_{k,\varepsilon}^-(y, \lambda) \right] \\ & \quad - \varphi_\delta(y) \left[\frac{b''(y)}{b(y) - \lambda + i\varepsilon} - \frac{b''(y)}{b(y) - \lambda - i\varepsilon} \right] \partial_\lambda \phi_{k,\varepsilon}^-(y, \lambda), \end{aligned} \quad (6.15)$$

for $y \in I$ with zero Dirichlet boundary condition.

We notice the identity that for $y \in I$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$,

$$\partial_\lambda \psi_k(y, \lambda) = \lim_{\varepsilon \rightarrow 0^+} D\phi_{k,\varepsilon}(y, \lambda).$$

We can reformulate (6.15) as the integral equation for $y \in I$,

$$\begin{aligned} & D\phi_{k,\varepsilon}(y, \lambda) + T_k^*(\lambda + i\varepsilon) D\phi_{k,\varepsilon}(y, \lambda) \\ &= - \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi_\delta(z) \left[\frac{b''(z)}{(b(z) - \lambda + i\varepsilon)^2} \phi_{k,\varepsilon}^+(z, \lambda) - \frac{b''(z)}{(b(z) - \lambda - i\varepsilon)^2} \phi_{k,\varepsilon}^-(z, \lambda) \right] dz \\ & \quad - \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi_\delta(z) \left[\frac{b''(z)}{b(z) - \lambda + i\varepsilon} - \frac{b''(z)}{b(z) - \lambda - i\varepsilon} \right] \partial_\lambda \phi_{k,\varepsilon}^-(z, \lambda) dz \\ &:= R_{k,\varepsilon}(y, \lambda). \end{aligned} \quad (6.16)$$

We can write for $y \in I$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$,

$$D\phi_{k,\varepsilon}(y, \lambda) := R_{k,\varepsilon}(y, \lambda) + D\phi_{k,\varepsilon}(y, \lambda; 1). \quad (6.17)$$

Then $D\phi_{k,\varepsilon}(y, \lambda; 1)$ satisfies for $y \in I$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$, the equation

$$D\phi_{k,\varepsilon}(y, \lambda; 1) + T_k^*(\lambda + i\varepsilon) D\phi_{k,\varepsilon}(y, \lambda; 1) = -T_k^*(\lambda + i\varepsilon) R_{k,\varepsilon}(y, \lambda). \quad (6.18)$$

The desired bounds (6.21) follow from (6.16)–(6.18), and Lemma 3.2 with $X = X_{L, \mathcal{Q}k}$. $\square$

Lastly we turn to the highest order derivative $\partial_\lambda^2 \psi_{k,\varepsilon}^t(y, \lambda)$ that we need to control. To study $\partial_\lambda^2 \psi_{k,\varepsilon}^t(y, \lambda)$, we take the derivative ∂_λ in (6.8) and obtain that

$$\begin{aligned} & \left[k^2 - \partial_y^2 + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \right] \partial_\lambda^2 \phi_{k,\varepsilon}^t(\cdot, \lambda) = - \frac{2b''(y)}{(b(y) - \lambda + i\varepsilon)^2} \partial_\lambda \phi_{k,\varepsilon}^t(\cdot, \lambda) \\ & \quad - \frac{2b''(y)}{(b(y) - \lambda + i\varepsilon)^3} \phi_{k,\varepsilon}^t(y, \lambda) \\ & \quad + \partial_\lambda^2 g_{k,\varepsilon}^t(y, \lambda). \end{aligned} \quad (6.19)$$

Lemma 6.3 Assume that $k \in \mathbb{Z} \setminus \{0\}$, $\lambda \in \Lambda_{\delta_0} \setminus \{b(y_*)\}$ and let $\phi_{k,\varepsilon}^t(y, \lambda)$ with $\iota \in \{\pm\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$ be as defined in (6.2). Recall that

$$\delta := \delta(\lambda) = 8\sqrt{|\lambda - b(y_*)|/b''(y_*)}. \quad (6.20)$$

Denoting for $y \in [0, 1]$, $\lambda \in \Lambda_{\delta_0} \setminus \{b(y_*)\}$,

$$\begin{aligned} \Lambda_2(y, \lambda) &:= -\psi_k(y, \lambda) \varphi_\delta(y) \frac{b''(y)}{(b'(y))^2} \lim_{\varepsilon \rightarrow 0^+} \frac{1}{b(y) - \lambda + i\varepsilon} \\ &\quad - \varphi_\delta(y) \frac{b''(y)}{(b'(y))^2} \lim_{\varepsilon \rightarrow 0^+} \left[\frac{1}{b(y) - \lambda + i\varepsilon} - \frac{1}{b(y) - \lambda - i\varepsilon} \right] \phi_{k,\varepsilon}^-(y, \lambda), \end{aligned}$$

then we have the bounds for $\lambda \in \Lambda_{\delta_0} \setminus \{b(y_*)\}$,

$$\begin{aligned} &\sum_{\alpha \in \{0,1\}} \|(\delta \wedge |k|^{-1})^{9/4} [\partial_\lambda^2 \psi_k(y, \lambda) - \Lambda_2(y, \lambda)]\|_{L^2([y_*-3\delta, y_*+3\delta])} \delta^{-1/2} \\ &\quad + \sum_{\alpha \in \{0,1\}} \left\| (\delta \wedge |k|^{-1})^{5/4} \left(|y - y_*| \wedge \frac{1}{|k|} \right) \partial_\lambda^2 \psi_k(y, \lambda) \right\|_{L^\infty([0,1] \setminus [y_*-3\delta, y_*+3\delta])} \\ &\lesssim |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \end{aligned} \quad (6.21)$$

Proof Denote for $k \in \mathbb{Z} \setminus \{0\}$, $\lambda \in \Lambda_{\delta_0} \setminus \{b(y_*)\}$, $\iota \in \{\pm\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$ and $y \in I$,

$$\begin{aligned} h_{k,\varepsilon}^\iota(z, \lambda; 4) &:= -\frac{2b''(z)}{(b(z) - \lambda - i\iota\varepsilon)^2} \varphi_\delta(z) \partial_\lambda \phi_{k,\varepsilon}^\iota(z, \lambda), \\ h_{k,\varepsilon}^\iota(z, \lambda; 5) &= -\frac{2b''(z)}{(b(z) - \lambda - i\iota\varepsilon)^3} \varphi_\delta(z) \phi_{k,\varepsilon}^\iota(z, \lambda), \\ h_{k,\varepsilon}^\iota(z, \lambda; 6) &:= -\frac{b''(z)}{(b(z) - \lambda - i\iota\varepsilon)^2} (1 - \varphi_\delta(z)) \partial_\lambda \phi_{k,\varepsilon}^\iota(z, \lambda), \\ h_{k,\varepsilon}^\iota(z, \lambda; 7) &= -\frac{2b''(z)}{(b(z) - \lambda - i\iota\varepsilon)^3} (1 - \varphi_\delta(z)) \phi_{k,\varepsilon}^\iota(z, \lambda), \\ h_{k,\varepsilon}^\iota(z, \lambda; 8) &:= \partial_\lambda^2 g_{k,\varepsilon}^\iota(z, \lambda). \end{aligned} \quad (6.22)$$

Define for $k \in \mathbb{Z} \setminus \{0\}$, $\lambda \in \Lambda_{\delta_0} \setminus \{b(y_*)\}$, $\iota \in \{\pm\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$ and $z \in I$,

$$\begin{aligned} \partial_\lambda^2 \phi_{k,\varepsilon}^\iota(y, \lambda) &:= \phi_{k,\varepsilon}^\iota(y, \lambda; 2) + \sum_{j=4}^8 \int_0^1 \mathcal{G}_k(y, z; \lambda + i\iota\varepsilon) h_{k,\varepsilon}^\iota(z, \lambda; j) dz \\ &\quad - \sum_{j=4}^8 T_k^*(\lambda + i\iota\varepsilon) \int_0^1 \mathcal{G}_k(y, z; \lambda + i\iota\varepsilon) h_{k,\varepsilon}^\iota(z, \lambda; j) dz. \end{aligned} \quad (6.23)$$

It follows from (6.19) that $\phi_{k,\varepsilon}^\iota(y, \lambda; 2)$ satisfies for $y \in I$,

$$\begin{aligned} &\phi_{k,\varepsilon}^\iota(y, \lambda; 2) + T_k^*(\lambda + i\iota\varepsilon) \phi_{k,\varepsilon}^\iota(y, \lambda; 2) \\ &= \sum_{j=4}^8 [T_k^*(\lambda + i\iota\varepsilon)]^2 \int_0^1 \mathcal{G}_k(y, z; \lambda + i\iota\varepsilon) h_{k,\varepsilon}^\iota(z, \lambda; j) dz. \end{aligned} \quad (6.24)$$

It follows from Lemma 6.2 and Lemma 3.1 that for $j \in \{6, 7, 8\}$

$$\left\| [T_k^*(\lambda + i\varepsilon)]^2 \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) h_{k,\varepsilon}^\iota(z, \lambda; j) dz \right\|_{X_{N,\varrho_k}} \lesssim (\delta \wedge |k|^{-1})^{-4} |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \quad (6.25)$$

Using integration by parts argument similar to (4.17)–(4.18), we have also for $j \in \{4, 5\}$,

$$\left\| [T_k^*(\lambda + i\varepsilon)]^2 \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) h_{k,\varepsilon}^t(z, \lambda; j) dz \right\|_{X_{N,\ell k}} \lesssim (\delta \wedge |k|^{-1})^{-4} |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}. \quad (6.26)$$

It follows from (6.22)–(6.26) and Lemma 4.4 that for $\lambda \in \Lambda_{\delta_0} \setminus \{b(y_*)\}$, $\iota \in \{\pm\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$,

$$\|\phi_{k,\varepsilon}^\iota(y, \lambda; 2)\|_{X_{N,\ell k}} \lesssim (\delta \wedge |k|^{-1})^{-4} |k|^{5/2} \|\omega_{0k}\|_{H_k^1(I)}.$$

Using (6.19), we get that for $y \in I$,

$$\begin{aligned} & \left[k^2 - \partial_y^2 + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \right] \left(\partial_\lambda^2 \phi_{k,\varepsilon}^+(y, \lambda) - \partial_\lambda^2 \phi_{k,\varepsilon}^-(y, \lambda) \right) \\ &= \sum_{j=4}^8 \left[h_{k,\varepsilon}^+(y, \lambda; j) - h_{k,\varepsilon}^-(y, \lambda; j) \right]. \end{aligned}$$

Denoting $D^2 \phi_{k,\varepsilon}(y, \lambda)$, $h \in I$, $\lambda \in \Lambda_{\delta_0} \setminus \{b(y_*)\}$, as the solution to

$$\left[k^2 - \partial_y^2 + \frac{b''(y)}{b(y) - \lambda + i\varepsilon} \right] D^2 \phi_{k,\varepsilon}(y, \lambda) = \sum_{j=4}^5 \left[h_{k,\varepsilon}^+(y, \lambda; j) - h_{k,\varepsilon}^-(y, \lambda; j) \right] \quad (6.27)$$

for $y \in I$ with zero Dirichlet boundary condition.

We note the identity that for $y \in I$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$,

$$\partial_\lambda^2 \psi_k(y, \lambda) = \lim_{\varepsilon \rightarrow 0+} D^2 \phi_{k,\varepsilon}(y, \lambda).$$

We can reformulate (6.27) as the integral equation for $y \in I$

$$\begin{aligned} & D^2 \phi_{k,\varepsilon}(y, \lambda) + T_k^*(\lambda + i\varepsilon) D^2 \phi_{k,\varepsilon}(y, \lambda) \\ &= \int_0^1 \mathcal{G}_k(y, z; \lambda + i\varepsilon) \varphi_\delta(z) \sum_{j=4}^5 \left[h_{k,\varepsilon}^+(z, \lambda; j) - h_{k,\varepsilon}^-(z, \lambda; j) \right] dz := R_{k,\varepsilon}^*(y, \lambda). \end{aligned} \quad (6.28)$$

We can write for $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$, $0 < \varepsilon < \min\{|\lambda - b(y_*)|, \varepsilon_0\}$, $y \in I$,

$$D^2 \phi_{k,\varepsilon}(y, \lambda) := D^2 \phi_{k,\varepsilon}(y, \lambda; 2) + R_{k,\varepsilon}^*(y, \lambda) - T_k^*(\lambda + i\varepsilon) R_{k,\varepsilon}^*(y, \lambda). \quad (6.29)$$

Then $D^2 \phi_{k,\varepsilon}(y, \lambda; 2)$ satisfies for $y \in I$, $\lambda \in \Sigma_{\delta_0} \setminus \{b(y_*)\}$,

$$D^2 \phi_{k,\varepsilon}(y, \lambda; 2) + T_k^*(\lambda + i\varepsilon) D^2 \phi_{k,\varepsilon}(y, \lambda; 2) = [T_k^*(\lambda + i\varepsilon)]^2 R_{k,\varepsilon}^*(y, \lambda). \quad (6.30)$$

The desired bounds (6.21) follow from (6.28)–(6.30), and Lemma 3.2 with $X = X_{L,\ell k}$, using also the bound

$$\left\| [T_k^*(\lambda + i\varepsilon)]^2 R_{k,\varepsilon}^*(\cdot, \lambda) \right\|_{X_{L,\ell k}} \lesssim \left(\delta \wedge \frac{1}{|k|} \right)^{-4} |k|^{5/2} \|\omega_{0k}\|_{H_k^3(I)}.$$

□

7 Proof of Theorem 1.2

In this section, we prove Theorem 1.2. We can assume that $t \geq 1$. We first give the proof of (1.3)–(1.4). Using the representation formula (2.4), we have

$$\begin{aligned}\psi_k(t, y) &= \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0^+} \int_{\Sigma} e^{-ik\lambda t} \left[\psi_{k,\varepsilon}^+(y, \lambda) - \psi_{k,\varepsilon}^-(y, \lambda) \right] d\lambda \\ &= -\frac{1}{2\pi i k^2 t^2} \lim_{\varepsilon \rightarrow 0^+} \int_{\Sigma} e^{-ik\lambda t} \left[\partial_{\lambda}^2 \psi_{k,\varepsilon}^+(y, \lambda) - \partial_{\lambda}^2 \psi_{k,\varepsilon}^-(y, \lambda) \right] d\lambda.\end{aligned}$$

Fix $\Phi^* \in C_0^\infty(\Sigma_{\delta_0})$ with $\Phi^* \equiv 1$ on $\Sigma_{2\delta_0/3}$. We can decompose for $t \geq 1$, $y \in [0, 1]$,

$$\psi_k(t, y) := \psi_k^1(t, y) + \psi_k^2(t, y),$$

where

$$\begin{aligned}\psi_k^1(t, y) &:= -\frac{1}{2\pi i k^2 t^2} \lim_{\varepsilon \rightarrow 0^+} \int_{\Sigma} e^{-ik\lambda t} (1 - \Phi^*(\lambda)) \left[\partial_{\lambda}^2 \psi_{k,\varepsilon}^+(y, \lambda) - \partial_{\lambda}^2 \psi_{k,\varepsilon}^-(y, \lambda) \right] d\lambda, \\ \psi_k^2(t, y) &:= -\frac{1}{2\pi i k^2 t^2} \lim_{\varepsilon \rightarrow 0^+} \int_{\Sigma} e^{-ik\lambda t} \Phi^*(\lambda) \left[\partial_{\lambda}^2 \psi_{k,\varepsilon}^+(y, \lambda) - \partial_{\lambda}^2 \psi_{k,\varepsilon}^-(y, \lambda) \right] d\lambda.\end{aligned}$$

For (1.3), it suffices to prove that for $\sigma \in \{1, 2\}$, $k \in \mathbb{Z} \setminus \{0\}$ and $t \geq 1$,

$$\|\psi_k^\sigma(t, \cdot)\|_{L^2([0,1])} \lesssim \frac{|k|^3}{t^2} \|\omega_{0k}\|_{H_k^3([0,1])}. \quad (7.1)$$

The case $\sigma = 1$ in (7.1) corresponding to the non-degenerate case is analogous to the case of monotonic shear flows, see [15], and follow from Lemmas 5.1–5.3. We focus on the main new case $\sigma = 2$ in (7.1). Denote for $k \in \mathbb{Z} \setminus \{0\}$,

$$M_k := |k|^{5/2} \|\omega_{0k}\|_{H_k^3([0,1])}.$$

Our main tools are Lemmas 6.1, 6.2, and 6.3, which imply the following bounds for $y \in [0, 1]$, $\lambda \in \Sigma_{\delta_0}$.

– If $|\lambda - b(y_*)|^{1/2} < |y - y_*|/20$, then

$$\begin{aligned}|\psi_k(y, \lambda)| &\lesssim (\min\{|\lambda - b(y_*)|^{1/2}, |k|^{-1}\})^{11/4} (|y - y_*|^{-1} + |k|) M_k, \\ |\partial_{\lambda}^2 \psi_k(y, \lambda)| &\lesssim (\min\{|\lambda - b(y_*)|^{1/2}, |k|^{-1}\})^{-5/4} (|y - y_*|^{-1} + |k|) M_k;\end{aligned} \quad (7.2)$$

– If $|y - y_*|/20 < |\lambda - b(y_*)|^{1/2} < 20|y - y_*|$, then

$$\begin{aligned}|\psi_k(y, \lambda)| &\lesssim (\min\{|\lambda - b(y_*)|^{1/2}, |k|^{-1}\})^{5/4} |\lambda - b(y_*)|^{1/4} M_k, \\ |\psi_k(y, \lambda) - \psi_k(y, b(y))| &\lesssim |\lambda - b(y)|^{1/2} |\lambda - b(y_*)|^{3/8} M_k, \\ \|\partial_{\lambda}^2 \psi_k(\cdot, \lambda) - \Lambda_2(\cdot, \lambda)\|_{L^2(|y - y_*| \approx |\lambda - b(y_*)|^{1/2})} &\lesssim (|\lambda - b(y_*)|^{-1/2} + |k|)^{9/4} |\lambda - b(y_*)|^{1/4} M_k;\end{aligned} \quad (7.3)$$

– If $|\lambda - b(y_*)|^{1/2} > 20|y - y_*|$, then

$$\begin{aligned}|\psi_k(y, \lambda)| &\lesssim |\lambda - b(y_*)|^{1/4} (\min\{|\lambda - b(y_*)|^{1/2}, |k|^{-1}\})^{5/4} M_k, \\ \|\partial_{\lambda}^2 \psi_k(\cdot, \lambda) - \Lambda_2(\cdot, \lambda)\|_{L^2(|y - y_*| < |\lambda - b(y_*)|^{1/2}/20)} &\lesssim (|\lambda - b(y_*)|^{-1/2} + |k|)^{9/4} |\lambda - b(y_*)|^{1/4} M_k.\end{aligned} \quad (7.4)$$

It follows from (7.2)–(7.4) that for $y \in [0, 1]$, $t \geq 1$,

$$\left| \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(\lambda) \Lambda_2(y, \lambda) d\lambda \right| \lesssim |y - y_*|^{-1/4} \max \{1, |k|^{1/2} |y - y_*|^{1/2}\} M_k, \quad (7.5)$$

and, by considering the cases $|\lambda - b(y_*)| \ll |y - y_*|^2$, $|\lambda - b(y_*)| \approx |y - y_*|^2$ and $|\lambda - b(y_*)| \gg |y - y_*|^2$, also that for $y \in [0, 1]$, $t \geq 1$,

$$\left\| \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(\lambda) [\partial_\lambda^2 \psi_k(y, \lambda) - \Lambda_2(y, \lambda)] d\lambda \right\|_{L^2([0, 1])} \lesssim |k|^{9/4} M_k. \quad (7.6)$$

The desired bound (7.1) for $\sigma = 2$ follows from (7.5)–(7.6).

The proof of (1.4) is similar to the proof of (1.3), using Lemma 6.1 and Lemma 6.2.

We now turn to the proof of the depletion bounds (1.5). Assume that $k \in \mathbb{Z} \setminus \{0\}$. Applying $-k^2 + \partial_y^2$ to $\psi_k(t, y)$ in (2.4), and using (2.5), we get that for $y \in [0, 1]$, $t \geq 1$,

$$\omega_k(t, y) = \omega_k^*(t, y) + \omega_k^{**}(t, y),$$

where

$$\begin{aligned} \omega_k^*(t, y) &:= \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0^+} \int_{\Sigma} e^{-ik\lambda t} (1 - \Phi^*(y)) \\ &\quad \times \left[\frac{b''(y) \psi_{k,\varepsilon}^+(y, \lambda) - \omega_{0k}(y)}{b(y) - \lambda + i\varepsilon} - \frac{b''(y) \psi_{k,\varepsilon}^-(y, \lambda) - \omega_{0k}(y)}{b(y) - \lambda - i\varepsilon} \right] d\lambda, \\ \omega_k^{**}(t, y) &:= \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0^+} \int_{\Sigma} e^{-ik\lambda t} \Phi^*(y) \\ &\quad \times \left[\frac{b''(y) \psi_{k,\varepsilon}^+(y, \lambda) - \omega_{0k}(y)}{b(y) - \lambda + i\varepsilon} - \frac{b''(y) \psi_{k,\varepsilon}^-(y, \lambda) - \omega_{0k}(y)}{b(y) - \lambda - i\varepsilon} \right] d\lambda. \end{aligned}$$

We have the bound for $t \geq 1$,

$$\|\omega_k^*(t, y)\|_{L^\infty([0, 1])} \lesssim |k|^2 M_k.$$

For $|y - y_*| < \delta_0/10$, $t \geq 1$, since $(b(y) - \lambda + i\varepsilon)$ with $\varepsilon \in \{\pm\}$ is not singular in this case, we have in addition by integration by parts that

$$|\omega_k^*(t, y)| \lesssim |k|^2 \frac{1}{t} M_k.$$

We now turn to $\omega_k^{**}(t, y)$. Using (6.1), we can write for $y \in [0, 1]$, $t \geq 1$,

$$\begin{aligned} 2\pi i \omega_k^{**}(t, y) &= \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(\lambda) \\ &\quad \times \left[\frac{\phi_{k,\varepsilon}^+(y, \lambda) - (1 - \Psi(y)) \omega_{0k}(y)}{b(y) - \lambda + i\varepsilon} - \frac{\phi_{k,\varepsilon}^-(y, \lambda) - (1 - \Psi(y)) \omega_{0k}(y)}{b(y) - \lambda - i\varepsilon} \right] d\lambda \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(\lambda) \left[\frac{\phi_{k,\varepsilon}^+(y, \lambda)}{b(y) - \lambda + i\varepsilon} - \frac{\phi_{k,\varepsilon}^-(y, \lambda)}{b(y) - \lambda - i\varepsilon} \right] d\lambda + W_k(t, y), \end{aligned}$$

where $W_k(t, y)$ satisfies the bound for $t \geq 1$,

$$\|W_k(t, \cdot)\|_{L^\infty([0, 1])} \lesssim t^{-1} M_k, \quad (7.7)$$

which follows from simple integration by parts argument. We decompose for $y \in [0, 1] \setminus \{y_*\}$,

$$\begin{aligned} \omega_k^{**}(t, y) &= \frac{W_k(t, y)}{2\pi i} \\ &= \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(\lambda) \left[\frac{\psi_k(y, \lambda)}{b(y) - \lambda + i\varepsilon} \right] d\lambda \\ &\quad + \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(\lambda) \phi_{k,\varepsilon}^-(y, \lambda) \left[\frac{1}{b(y) - \lambda + i\varepsilon} - \frac{1}{b(y) - \lambda - i\varepsilon} \right] d\lambda. \end{aligned}$$

It follows from (7.2)–(7.4) that

$$\left| \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(\lambda) \phi_{k,\varepsilon}^-(y, \lambda) \left[\frac{1}{b(y) - \lambda + i\varepsilon} - \frac{1}{b(y) - \lambda - i\varepsilon} \right] d\lambda \right| \lesssim |y - y_*|^{7/4} M_k. \quad (7.8)$$

For $\gamma \in (1, \infty)$ to be fixed below, by considering the three ranges (I) $|\lambda - b(y_*)| \lesssim |y - y_*|^2$, (II) $|\lambda - b(y_*)| \geq \gamma |y - y_*|^2$, and (III) $|y - y_*|^2 \ll |\lambda - b(y_*)| < \gamma |y - y_*|^2$, and using Lemma 6.2 and Lemma 6.3, we get that

$$\begin{aligned} &\left| \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(y) \left[\frac{\psi_k(y, \lambda)}{b(y) - \lambda + i\varepsilon} \right] d\lambda \right| \\ &\lesssim |y - y_*|^{7/4} (1 + |k|^{1/2} |y - y_*|^{1/2}) \\ &\quad + \frac{1}{|k|t} (|k|^{1/2} + \gamma^{-1/8} |y - y_*|^{-1/4}) + \gamma^{7/8} |y - y_*|^{7/4} \Big] M_k. \end{aligned}$$

In the above, we used integration by part to get decay in t in range (II). Optimizing in γ , we get that for $t \geq 1$,

(i) if $t|y - y_*|^2 \lesssim 1$,

$$\left| \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(y) \left[\frac{\psi_k(y, \lambda)}{b(y) - \lambda + i\varepsilon} \right] d\lambda \right| \lesssim [t^{-1} + |k|^{1/2} |y - y_*|^{7/4} + t^{-7/8}]. \quad (7.9)$$

(ii) if $t|y - y_*|^2 \gg 1$,

$$\begin{aligned} &\left| \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0+} \int_{\mathbb{R}} e^{-ik\lambda t} \Phi^*(y) \left[\frac{\psi_k(y, \lambda)}{b(y) - \lambda + i\varepsilon} \right] d\lambda \right| \\ &\lesssim |y - y_*|^{7/4} (1 + |k|^{1/2} |y - y_*|^{1/2}) + \frac{1}{|k|^{1/2} t^{7/8}} + |y - y_*|^{7/4} \Big] M_k. \quad (7.10) \end{aligned}$$

The desired bounds (7.7), (7.8), (7.9)–(7.10). Theorem 1.2 is now proved.

Acknowledgements The first author is supported in part by NSF grant DMS-2007008. The second author is supported in part by NSF grant DMS-2306528. The third author is supported in part by NSF grant DMS-1945179.

References

1. Arnold, V., Khesin, B.: Topological Methods in Hydrodynamics. Springer-Verlag, New York (1998)
2. Bedrossian, J., Masmoudi, N.: Inviscid damping and the asymptotic stability of planar shear flows in the 2D Euler equations. Publ. Math. IHES **122**, 195–300 (2015)
3. Bedrossian, J., Coti Zelati, M., Vicol, V.: Vortex axisymmetrization, inviscid damping, and vorticity depletion in the linearized 2D Euler equations. Ann. PDE **5**, 4 (2019)

4. Bedrossian, J.: Nonlinear echoes and Landau damping with insufficient regularity. *Tunis. J. Math.* **3**, 121–205 (2021)
5. Bouchet, F., Morita, H.: Large time behavior and asymptotic stability of the 2D Euler and linearized Euler equations. *Phys. D* **239**, 948–966 (2010)
6. Case, K.: Stability of inviscid plane Couette flow. *Phys. Fluids* **3**, 143–148 (1960)
7. Deng, Y., Masmoudi, N.: Long-time instability of the Couette flow in low Gevrey spaces. *Commun. Pure Appl. Math.* **76**, 2804–2887 (2023)
8. Faddeev, L.: On the theory of the stability of plane-parallel flows of an ideal fluid. *Zap. Nauchn. Sem. LOMI* **21**, 164–172 (1971)
9. Grenier, E., Nguyen, T.T., Rousset, F., Soffer, A.: Linear inviscid damping and enhanced viscous dissipation of shear flows by using the conjugate operator method. *J. Funct. Anal.* **278**, 108339 (2020)
10. Ionescu, A., Jia, H.: Inviscid damping near the Couette flow in a channel. *Commun. Math. Phys.* **374**, 2015–2096 (2020)
11. Ionescu, A., Jia, H.: Axi-symmetrization near point vortex solutions for the 2D Euler equation. *Commun. Pure Appl. Math.* **75**, 818–891 (2022)
12. Ionescu, A., Jia, H.: Nonlinear inviscid damping near monotonic shear flows. *Acta Math.* **230**, 321–399 (2023)
13. Ionescu, A., Jia, H.: On the nonlinear stability of shear flows and vortices. *Proc. ICM 2022* (to appear) (2023)
14. Ionescu, A., Jia, H.: Linear vortex symmetrization: the spectral density function. *Arch. Rational Mech. Anal.* **246**, 61–137 (2022)
15. Jia, H.: Linear inviscid damping near monotone shear flows. *SIAM J. Math. Anal.* **52**, 623–652 (2020)
16. Jia, H.: Uniform linear inviscid damping and enhanced dissipation near monotonic shear flows in high Reynolds number regime (I): the whole space case. *J. Math. Fluid Mech.* **25**, 42 (2023)
17. Jia, H.: Linear inviscid damping in Gevrey spaces. *Arch. Rational Mech. Anal.* **235**, 1327–1355 (2020)
18. Kelvin, L.: Stability of fluid motion-rectilinear motion of viscous fluid between two plates. *Phil. Mag.* **24**, 155 (1887)
19. Kirchhoff, G.: *Vorlesungen über Mathematische Physik*. Teubner, Leipzig (1876)
20. Lin, Z., Zeng, C.: Inviscid dynamical structures near Couette flow. *Arch. Rational Mech. Anal.* **200**, 1075–1097 (2011)
21. Lin, Z.: Instability of some ideal plane flows. *SIAM J. Math. Anal.* **35**, 318–356 (2003)
22. Liu, X., Zeng, C.: Capillary gravity water waves linearized at monotone shear flows: eigenvalues and inviscid damping. [arXiv:2110.12604](https://arxiv.org/abs/2110.12604) (2021)
23. Masmoudi, N., Zhao, W.: Nonlinear inviscid damping for a class of monotone shear flows in finite channel. [arXiv:2001.08564](https://arxiv.org/abs/2001.08564) (2020)
24. Mouhot, C., Villani, C.: On Landau damping. *Acta Math.* **207**, 29–201 (2011)
25. Orr, W.: The stability or instability of steady motions of a perfect liquid and of a viscous liquid, Part I: a perfect liquid. *Proc. R. Ir. Acad. Sect. A Math. Phys. Sci.* **27**, 9–68 (1907)
26. Rayleigh, L.: On the stability or instability of certain fluid motions. *Proc. Lond. Math. Soc.* **s1-11**, 57–72 (1879)
27. Rosencrans, S., Sattinger, D.: On the spectrum of an operator occurring in the theory of Hydrodynamics stability. *J. Math. Phys.* **45**, 289–300 (1966)
28. Stepin, S.: Nonself-adjoint Friedrichs model in hydrodynamic stability. *Funct. Anal. Appl.* **29**, 91–101 (1995). Translated from *Funktsional. Anal. i Ego Prilozhen.* **29**, 22–35 (1995)
29. Yamanaka, T.: A new higher order chain rule and Gevrey class. *Ann. Glob. Anal. Geom.* **7**, 179–203 (1989)
30. Wei, D., Zhang, Z., Zhao, W.: Linear inviscid damping for a class of monotone shear flow in Sobolev spaces. *Commun. Pure Appl. Math.* **71**, 617–687 (2018)
31. Wei, D., Zhang, Z., Zhao, W.: Linear inviscid damping and vorticity depletion for shear flows. *Ann. PDE* **5**, 3 (2019)
32. Wei, D., Zhang, Z., Zhao, W.: Linear inviscid damping and enhanced dissipation for the Kolmogorov flow. *Adv. Math.* **362**, 106963 (2020)
33. Wei, D.: Diffusion and mixing in fluid flow via the resolvent estimate. *Sci. China Math.* **64**, 507–518 (2021)
34. Zillinger, C.: Linear inviscid damping for monotone shear flows. *Trans. Amer. Math. Soc.* **369**, 8799–8855 (2017)
35. Zillinger, C.: Linear inviscid damping for monotone shear flows in a finite periodic channel, boundary effects, blow-up and critical Sobolev regularity. *Arch. Rational Mech. Anal.* **221**, 1449–1509 (2016)
36. Coti Zelati, M., Zillinger, C.: On degenerate circular and shear flows: the point vortex and power law circular flows. *Commun. Partial Differ. Equ.* **44**, 110–155 (2019)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.