

Opinion: The importance of historical and paleoclimate aerosol radiative effects

Natalie Mahowald¹, Longlei Li¹, Samuel Albani², Douglas Hamilton³, Jasper F. Kok⁴

¹ Department of Earth and Atmospheric Sciences, Cornell University, Ithaca, NY, USA

5 ² Department of Environmental and Earth Sciences, University of Milano-Bicocca, Milan, Italy

³ Department of Marine, Earth and Atmospheric Sciences, North Carolina State, Raleigh, NC, USA.

⁴ Department of Atmospheric Sciences, University of California, Los Angeles, CA, USA

Correspondence to: Natalie Mahowald (mahowald@cornell.edu).

10 **Abstract.** Estimating the past aerosol radiative effects and their uncertainties is an important topic in climate science. Aerosol radiative effects propagate into large uncertainties in estimates of how present and future climate evolves with changing greenhouse gas emissions. A deeper understanding of how aerosols affected the atmospheric energy budget under past climates is hindered in part by a lack of relevant paleo observations and in part because less attention has been paid to the problem. Because of the lack of information we do not seek here to determine the change in the radiative forcing due to aerosol changes, but rather to estimate the uncertainties in those changes. Here we argue that current uncertainties from emission uncertainties (90% confidence interval range spanning 2.8 W/m²) are just as large as model spread uncertainties (2.8 W/m²) in calculating preindustrial to current day aerosol radiative effects. There are no estimates for radiative forcing for important aerosols such as wildfire and dust aerosols in most paleoclimate time periods. However, qualitative analysis of paleoclimate proxies suggests that changes in aerosols between different past climates are similar in magnitude to changes in aerosols between preindustrial and current day, plus there is the added uncertainty from the variability in aerosols and fires in the preindustrial. From the limited literature we crudely estimate a paleoclimate aerosol uncertainty for the last glacial maximum relative to preindustrial of 4.8 W/m² and we estimate the uncertainty in the aerosol feedback in the natural Earth system over the paleoclimate (last glacial maximum to preindustrial) to be about 3.2 W/m²/°K. In order to more accurately assess the uncertainty in historical aerosol radiative effects, we propose a new model intercomparison project, which would include multiple plausible emission scenarios tested across a range of state-of-the-art climate models over the historical period. These emission scenarios would then be compared to the available aerosol observations to constrain which are most probable. In addition, future efforts should work to characterize and constrain paleo-aerosol forcings and uncertainties. Careful propagation of aerosol uncertainties in the literature is required to ensure an accurate quantification of uncertainties in projections of future climate changes.

15
20
25
30

1 Introduction

While CO₂ radiative forcing has been the most important driver of the observed climate warming, aerosol interactions with radiation and cloud properties represent the largest source of uncertainty in the Intergovernmental Panel on Climate Change (IPCC) estimates of present day to preindustrial radiative forcing (e.g. IPCC, 2021). This large uncertainty is driven by the heterogeneity of aerosols in space, time, composition, size and shape in the current climate, complexity in aerosol impacts on radiation and clouds, poor knowledge of aerosols in historical and past climate conditions, and how aerosol processes have changed over time (Albani et al., 2018; Bellouin et al., 2020; IPCC, 2021; Carslaw et al., 2017; Gulev et al., 2021; Szopa et al., 2021). Substantial research emphasis has been placed on understanding the direct changes in emissions from human activities (e.g. fossil fuel combustion); these emissions are estimated based on many processes, including emission factors for different industries, estimates of the changes in location and intensity of different industrial facilities, as well as time dependent changes in technologies (e.g. (Bond et al., 2007; Klimont et al., 2017)). On the other hand, the uncertainty in aerosol emissions that are usually considered natural, such as from dust storms or wildfires, are likely larger, and contribute to larger uncertainties in aerosol radiative forcing (Carslaw et al., 2010; Mahowald et al., 2011a; Regayre et al., 2018). Because of limited paleo-observations, we currently rely on emission models that were calibrated using current observations and apply them to past climates, not only for industrial sources but also for wildfire and dust aerosols (Van Marle et al., 2017; Turnock et al., 2020; Zhao et al., 2022).

However, the dominant mechanisms for natural emission processes are unlikely to have remained constant over time, and thus using present day observations to constrain past model predictions is biased towards the present day and thus anthropogenic influenced aerosol behavior (e.g. Hamilton et al., 2018). We therefore advocate that past model simulations should, as much as possible, be constrained using paleoenvironmental archives of past aerosol. Note that for IPCC estimates, most models use the same emission dataset(s) to drive emissions meaning that the aerosol radiative forcing uncertainty estimates based on Climate Model Intercomparison Project (CMIP6) estimates tend to accurately represent the uncertainty due to model spread using the same emission dataset, but do not necessarily aim to include the uncertainty due to emission uncertainties (Hoesly et al., 2018; Bellouin et al., 2020; Thornhill et al., 2020; Pincus et al., 2016), which are especially large in preindustrial or paleoclimate climates (Li et al., 2019; Kok et al., 2023).

60

Here we argue that an improved characterization of the evolution of ‘natural’ aerosols, especially dust and wildfire aerosols, is needed to improve our understanding of aerosol radiative effects over the historical and paleoclimate

time periods. Before we can seek to constrain the uncertainties (e.g. Bellouin et al., 2020), we need to first characterize the different sources of uncertainties and their magnitude (Carslaw et al., 2017), especially emissions of natural aerosols (e.g. Hamilton et al., 2018; Kok et al., 2023), and propagate these uncertainties into the climate simulations. Additionally, we argue that radiative perturbations due to changes in natural aerosols that are affected by human actions, such as dust and wildfire aerosols, should be treated as a radiative forcing. Note that we use the IPCC glossary definition of radiative forcing, as the change in the top of atmosphere radiative balance due to the addition of a species (IPCC, 2021; Gulev et al., 2021; Szopa et al., 2021), and use the term radiative forcing instead of effective radiative forcing, for which the same arguments apply. In our discussion here, we include both direct effects and aerosol-cloud interactions, realizing that the largest uncertainties are often associated with aerosol-cloud interactions. We discuss 1) the limited available data constraining the changes in dust and wildfire aerosol emissions over the historical time period and how those compare to the CMIP6 emission datasets; 2) consider whether aerosols can be considered feedbacks or forcings; 3) propagating the uncertainty in emissions of natural aerosols to create more realistic estimates of aerosol radiative effect uncertainties, 4) estimates for the uncertainties in aerosol radiative effects for paleoclimate time periods, 5) other sources of aerosol uncertainties and 6) the importance of accurate aerosol radiative effect uncertainties for climate science. Finally, 7) we propose a path forward to improve the characterization of the uncertainties in aerosol radiative effects, which would then allow us to start constraining the uncertainties with observations.

80 **2 Observations of natural aerosol changes since 1850**

Natural aerosols include dust, wildfire emissions, sea salts and biogenic organic aerosols, among others (e.g. dimethyl sulfide, lightning NO_x). Aerosols such as dust or carbonaceous and sulfur species emitted by wildfires can be generated under natural conditions. As such, these aerosols can produce important feedbacks in a changing climate (Allen et al., 2016; Kok et al., 2018; Thornhill et al., 2021). However, some of these aerosols, notably dust and wildfire aerosols, are also directly or indirectly affected by human actions. For instance, dust emissions can be increased substantially both by human disturbance of the natural landscape and by anthropogenic diversions of surface water flows (Lee et al., 2012; Ginoux et al., 2012; Xi and Sokolik, 2016). Similarly, wildfire emissions can be increased by open fires set by humans as part of deforestation or agricultural practices (van der Werf et al., 2010). Additionally, natural wildfire emissions can be suppressed by human activity, for example active firefighting or removal of forests and grasslands due to agricultural and urban development (Knorr et al., 2014; Bistinas et al., 2013). As such, the radiative perturbation due to historical changes in these natural aerosols can be partially due to both human land use changes (a forcing) and natural and anthropogenic climate changes (a feedback).

2.1 Historical changes in desert and agricultural dust

95 While the global concentration of long lived and well-mixed species such as CO₂ can be retrieved from ice cores, aerosols in the troposphere have lifetimes of only a few days to two weeks (Textor and others, 2006), meaning that aerosol observations need to be available across the globe in dozens or hundreds of locations at a high temporal frequency before we can begin to build a reasonable understanding of their distribution and behavior. Unfortunately, for most aerosol species we do not have that kind of coverage for most time periods, including the present day (e.g. see spatial coverage in Naik et al., 2021). Dust is an exception in some respects, in that dust is preserved to some extent in several natural archives, and thus there are compilations of dust changes over different time periods. For paleoclimate conditions (e.g. the last glacial cycle, the Holocene, etc.) the main natural archives for dust include ice cores and marine and terrestrial (loess/paleosol) sediment records, for which paleodust compilations exist (e.g. Albani et al., 2015). Most of those natural archives do not have the potential to cover the last few centuries, including the late Holocene to preindustrial to modern transitions, due to issues with dating or disturbance of surface sediments (e.g. for many marine sediment records the core top is lost during retrieval and thus the last 100 or so years are not easily obtainable; loess provides the substrate for very fertile soils exploited for agriculture in the last millennia) (Maher et al., 2010). Other archives, such as in particular firn cores from the polar areas, ice cores from mountain glaciers, and ombrotrophic peat bogs, have the potential to preserve dust deposition records over the last decades/centuries, although they are still affected by major uncertainties when it comes to retrieving accurate deposition records (Albani et al., 2015). With these caveats in mind, there is still paleo data that can be compiled to infer the evolution of desert dust in different regions across the globe since the pre-industrial, albeit with large uncertainties (Kok et al., 2023).

115 Desert dust is generated in dry, unvegetated regions with strong winds, and the generation of dust may be enhanced in regions with human land use (Ginoux et al., 2012). Between the 1960s and 1980s dust was observed to have changed by a factor of 4 over the North Atlantic region, perhaps due to expansion of land use, due to precipitation changes during the Sahel drought, or due to changes in winds (Mahowald et al., 2002; Prospero and Lamb, 2003; Evan et al., 2016). Paleoclimate evidence also suggests that dust is very sensitive to both climate change as well as land use (Lambert et al., 2008; Neff et al., 2008; Mulitza et al., 2010).

A recent synthesis of dust deposition observations suggests a $55 \pm 30\%$ (90% confidence interval) increase in dust globally since preindustrial times (Kok et al., 2023). While there is not sufficient data for each source to have complete confidence in such estimates, they still represent our best knowledge of the state of changes in dust since

125 1850s. Unfortunately, the model simulations from the last Climate Model Intercomparison Project (CMIP6) that include prognostic dust do not match these changes, either because of the lack of correct inclusion of the impact of land use onto dust, or incorrect simulation of the feedbacks of natural and anthropogenic climate changes onto dust (Fig. 1). Indeed, the CMIP6 models show relatively constant dust amounts in contrast to the large change that is observed (Kok et al., 2023).

130

2.2 Historical changes in wildfire and open fires

Paleoenvironmental archives for wildfires and other open fires include charcoal records near the source regions (Marlon et al., 2008), ice cores (McConnell et al., 2007), tree ring scars, and speleothems. While wildfire is a natural ecosystem process, humans can modify the frequency and intensity of fires through many mechanisms (Bowman et al., 2009, Pechony and Shindell 2010). Humans are responsible for much of the ignition of fires today, but also convert land from natural to managed land, reducing the area available for wildfires. In addition, humans may suppress fires once they start (Bowman et al., 2009; Kloster et al., 2010). Observations over the last 20 years from satellites show strong interannual variability with 50% changes in emissions observed correlated with climate signals, although human contributions could have played a vital role (van der Werf et al., 2006). It has long been the assumption in the atmospheric chemistry and aerosol community that open fires have increased due to humans (Knorr et al., 2014; Bistinas et al., 2013). The open fire emission data used for the CMIP6 uses data from current satellites to predict emissions from 1997 onwards, however, prior to the Satellite fire Era the emissions are generated from fire models with limited additional proxy evidence. These models include a range of representations of how fires have evolved with increasing human population density, and not all account for active or passive fire suppression. Overall, this results in CMIP6 fire emissions increasing since 1850s (Van Marle et al., 2017). However, there is ample evidence that the relationship may be much more nuanced. Paleoclimate data from charcoal records suggests a maximum in open fires in the 1850s, and a decrease since then (Marlon et al., 2008). Satellite data shows a global decrease in burned area over the last decades, driven primarily from the conversion of natural lands to agricultural and pastoral lands (Andela et al., 2017, Jones et al. 2022). Indeed, those fire models which include a more realistic representation of how fires and human population density are related simulate a much higher amount of fires during 1850 than the CMIP6 (or AEROCOM) emission dataset suggests (Hamilton et al., 2018; Hoesly et al., 2018; Dentener et al., 2006). Including these fire emission simulations in aerosol models improves the match to the available ice core data of the deposition ratio between present day to preindustrial of black carbon (yellow and red symbols) than using the default CMIP6 datasets (blue symbols) (Hamilton et al., 2018; Liu et al., 2021) (Fig. 2). This suggests that it is probable that there were more open fire emissions during

the PI than accounted for in the emission inventories used for CMIP6 simulations (Hoesly et al., 2018; van Marle et al., 2017).

Is it plausible that emissions from wildfires and other open fires, such as agricultural fires, could be so much larger in preindustrial times than currently being accounted for in climate model emission datasets? It is difficult to know for certain of course, but a few examples show the possibility, using different mechanisms. One study suggests that higher fire amounts in 1850 relative to today could be due to more land use change today than in 1850. In other words, today there is less natural area available for fires than 1850 (van der Werf et al., 2013) which is consistent with the decreased fire burn area observed over the satellite era (Andela et al., 2017). Other studies have suggested that wildire suppression has been important for reducing fires in North America for example over the last 50 years (Marlon et al., 2012) . In addition, agricultural open burning could be important; today, northern India represents the region with some of the highest aerosol optical depths and worst air quality (Li et al., 2022; Burnett et al., 2018). Despite the area having large population centers and industrial emissions, one of the largest sources in that region is agricultural burning (Cusworth et al., 2018). Similarly, the ban on straw burning, a primary source of pollutants in central and eastern China (Wu et al., 2018), has become a national policy for air pollution control, as in many other countries. This suggests that indeed, the high emission factors of open burning make it a very effective source of aerosols.

In addition, wildfires and open fires represent some of the most important aerosols for direct and aerosol-cloud radiative effects, with a total radiative effect in the current climate of -2 W/m^2 (IPCC, 2021; Gulev et al., 2021; Szopa et al., 2021; Penner et al., 1992). Changes in wildfires and open fires represent about -1 W/m^2 or 50% of the anthropogenic aerosol radiative forcing since 1850, and all of these estimates have very large uncertainties (Unger et al., 2010; IPCC, 2019, 2021; Gulev et al., 2021; Szopa et al., 2021). The overall negative forcing results from the effect of an increasing aerosol burden increasing cloud albedo, and from the prevailing effects of preferentially scattering sulfates and particulate organic matter (secondary aerosol from fire emission of precursors such as biogenic volatile organic compounds), over the preferentially absorbing black carbon emissions from fires (Hamilton et al., 2018; Carslaw et al., 2017; Penner et al., 1992; IPCC, 2021; Gulev et al., 2021; Szopa et al., 2021).

In summary, trade-offs between the effects of climate change and land use make it difficult to estimate past changes in the loading of dust, and smoke from open fires. Changes in other aerosols (fossil fuels, biofuels, biogenic aerosols, sea spray), are difficult to estimate as well (e.g. Figure 2.9a and b in Gulev et al., 2021). Without

observations of these aerosols in past climates, it is difficult to have confidence in our existing emission models and their past and future emissions projections.

3 Are dust and other natural aerosols forcers or feedbacks?

The nomenclature that dust and wildfire aerosols are natural aerosols is perhaps misleading and might have caused the important changes that have occurred in these aerosols to receive insufficient consideration by climate scientists. For instance, in the Sixth Assessment Report, the radiative perturbation due to the ~50% increase in dust over the historical record (Hooper and Marx, 2018; Mahowald et al., 2010; Kok et al., 2023) is not explicitly accounted for as a radiative forcing of the climate system, although the dust-climate feedback was quantified (Forster et al., 2021; Naik et al., 2021) and the report does highlight that there is substantial uncertainty in this feedback (Szopa et al., 2021). There are several reasons why not explicitly accounting for the historical dust increase as a radiative forcing could be problematic. First, although the exact proportion of modern dust that can be considered anthropogenic is uncertain (Tegen et al., 2004; Mahowald, 2007; Ginoux et al., 2012; Stanelle et al., 2014), a large body of work indicates that human land use changes in semi-arid and arid lands can produce a large increase in dust aerosol emissions (Neff et al., 2008; Webb and Pierre, 2018). Such land use changes have been widespread since the Industrial Revolution (Klein-Goldewijk, 2001) making it likely that a substantial part of the historical dust increase – perhaps even most of it – was driven by human land use changes (Ginoux et al., 2012; Hooper and Marx, 2018; Kok et al., 2023), which thus constitutes a radiative forcing.

A second reason is that not accounting for the historical dust increase as a radiative forcing implicitly assumes that the historical dust increase has been due to a climate feedback. However, the dust change per degree global surface temperature warming is inconsistent between time periods. Indeed, the dust increase during the planetary warming of the past century is opposite to what is seen in the paleo-record, for which cold periods like the Last Glacial Maximum coincide with high dust loadings (Albani et al., 2014). Moreover, there is no model consensus on whether dust will increase or decrease under future climate warming, in part because of large uncertainties in how precipitation in arid regions will change (IPCC, 2019). This inconsistency in the dust change per unit global surface temperature warming could be due to a number of factors: (i) the historical dust increase was primarily driven by human land use changes, not climate changes; (ii) the dust feedback is highly dependent on the climate state; or (iii) the dust feedback occurs over much longer timescales than the observed ~century-scale dust increase. Whatever the reason, the fact that the dust change per unit surface temperature change is not consistent between

time periods undermines both the plausibility and the usefulness of classifying historical dust changes as a feedback in the context of future climate predictions.

220

The final, and most important, reason why not explicitly accounting for the historical dust increase as a radiative forcing might be problematic is that the dust increase indicated by dust deposition records (McConnell et al., 2007a; Mulitza et al., 2010; Mahowald et al., 2010; Hooper and Marx, 2018) is not captured by climate models (Kok et al., 2023) (Fig. 1). These models therefore also predict a dust-climate feedback that is indistinguishable from zero (Thornhill et al., 2021; Kok et al., 2023). As such, not explicitly accounting for dust changes as a radiative forcing has the net effect of omitting this potentially important perturbation to Earth's energy balance. This can bias climate sensitivity constraints and projections of future climate changes (Kok et al. 2023).

Considering these important issues, we argue that model simulations should include historical dust changes as an external forcing for two reasons: 1) because it is unlikely that the anthropogenic forcing and the climate feedback components of the historical dust loading change can be reliably separated in the near future and 2) because climate models currently cannot reproduce the historical dust changes (Kok et al., 2023; Mahowald et al., 2010). Since many models include prognostic schemes for dust, we propose that models add a temporally varying emissions factor obtained from constraints on the historical evolution of atmospheric dust deposition (e.g., Mahowald et al., 2010). This would enable models to both reproduce the historical change in dust, yet to also predict future changes in dust forced by climate and land use changes. A similar approach could be used for other natural aerosol emissions, such as from wildfires.

However, when models do treat changes in aerosols as feedbacks, the full uncertainty in the feedback should be included. For example, from this analysis, the feedback uncertainty should be $\pm 1.6 \text{ W/m}^2$ uncertainty from preindustrial to current (see Section 4), over which time temperatures increased by about 1°K , so that means an uncertainty in the feedback of $\pm 1.6 \text{ W/m}^2/^\circ\text{K}$ (90% confidence).

4 Characterizing preindustrial to current aerosol forcing uncertainty

245

Since CMIP6 aerosol simulations are not consistent with available observations for dust and open fires, it is clear that additional uncertainty needs to be added to the aerosol radiative forcing estimates for the preindustrial to present day in order to make sure that the uncertainty ranges includes available observations. This is a substantial

undertaking, but here we show schematically a back-of-the-envelope calculation of how including the observations
 250 would affect estimates of aerosol radiative forcing uncertainty (Fig. 3).

We use here a slightly different nomenclature than (Sherwood et al., 2020) for example, to emphasize the
 uncertainties in radiative forcing, without introducing too much nomenclature, and thus define ΔF as the change in
 radiative forcing between two different times, and Σ is the uncertainty in that estimate using the 90% confidence
 255 intervals.

If we sum the sources of uncertainty currently available in the literature ($\Sigma_{Fires}^{PD-PI} = 2.8 \text{ W/m}^2$ from fires,
 $\Sigma_{Dust}^{PD-PI} = 0.4 \text{ W/m}^2$ from dust (Hamilton et al., 2018; Wan et al., 2021; Kok et al., 2023) and here we assume a
 10% error for industrial emissions for $\Sigma_{Industry}^{PD-PI} = 0.2 \text{ W/m}^2$ using Eq. (1), we obtain 2.8 W/m^2 , clearly dominated
 260 by fires (see Table 1 for terms). This is consistent also with attribution literature, which shows that much of the
 anthropogenic radiative forcing is from the biomass burning source (Figure 1a in Unger et al., 2010).

$$\Sigma_{Emis}^{PD-PI} = [(\Sigma_{Fires}^{PD-PI})^2 + (\Sigma_{Dust}^{PD-PI})^2 + (\Sigma_{Industry}^{PD-PI})^2]^{1/2}, \quad (1)$$

265 The uncertainty in radiative forcing from uncertainty in preindustrial emissions ($\Sigma_{Emis}^{PD-PI} = 2.8 \text{ W/m}^2$) is similar in
 magnitude to the uncertainty from using one emission scenario for the historical time period (2.8 W/m^2 which is
 the unconstrained model uncertainty using 90% confidence intervals) (Bellouin et al., 2020; Sherwood et al., 2020),
 which we refer to here as the unconstrained aerosol process uncertainty for present day to preindustrial ($\Sigma_{Process}^{PD-PI}$).
 The model spread in radiative forcing with the same emission scenario is due to differences in model simulations
 270 of concentration, radiation and cloud interactions using the same emission change, which are large, because these
 processes are poorly understood (Li et al., 2022; Pincus et al., 2016). Note that here we use the same term ‘scenario’
 for what happened in the past as what we use for choices in the future, to emphasize that we do not know the
 emissions. The models make different assumptions about aerosol lifetime, size distribution, aerosol microphysics,
 which results in different radiative forcings, so we assume this spread in models is the uncertainty in processes
 275 ($\Sigma_{Process}^{PD-PI}$), which is 2.8 W/m^2 . One could also think of this uncertainty in the process as coming from two sources:
 variability in present day processes in simulating aerosols (e.g. that radiative forcing is sensitive not just to the total
 emissions, but also to where, what kind and what else is in the region (Li et al., 2022; Bellouin et al., 2020) and

one part that is proportional to the strength of the change in aerosols, which could be proportional to the change in radiative forcing (ΔF) times some factor γ (Eq. 2).

$$\Sigma_{Process}^{PD-PI} = [(\Sigma_{Process}^{PD})^2 + (\Delta F * \gamma)^2]^{1/2}, \quad (2)$$

We propose that future studies should identify the strength of the base uncertainty ($\Sigma_{Process}^{PD}$) and the portion of this process uncertainty that is proportional to the strength of the change in the radiative forcing ($(\Delta F * \gamma)^2$).

The total unconstrained uncertainty due to aerosol changes could be estimated as being 4 W/m² using equation 3, assuming the uncertainties are orthogonal (Eq. 3).

$$\Sigma_{TotalUNC}^{PD-PI} = [(\Sigma_{Process}^{PD-PI})^2 + (\Sigma_{Emis}^{PD-PI})^2]^{1/2}, \quad (3)$$

Emissions from industry (which are likely better known) have been increasing since 1850 in the CMIP6 simulations from which we estimated the aerosol radiative forcings. But aerosol from wildfires have also been increasing during this historical period in these simulations (van Marle et al., 2017). As discussed in Section 1, the paleoclimate data (and fire models which explicitly account for passive fire suppression effects of land use change) suggest that open fires have been decreasing since 1850, potentially offsetting the increase in industrial emissions (Fig. 4b). In contrast, the CMIP6 wildfire emissions assume large increases since 1850 in wildfires and open fires (Van Marle et al., 2017). This produces a large uncertainty in the time series of aerosol forcing over the historical period. Notice that since wildfire emissions can vary strongly over a couple of years or decades (van der Werf et al., 2004), it should not be assumed (without observations) that the radiative forcing from wildfires follows either the top or the bottom of the error bar, but rather could vary from one year to the other over the whole range.

Some of the difficulty of looking at preindustrial to present day aerosol changes is simply understanding what ‘natural’ aerosols would look like without humans. Unfortunately, there are strong fluctuations across the time period just before the industrial era (e.g. 1500-1850) in fires (Fig. 5), some potentially associated with humans (e.g. perhaps the increase in 1850 and decrease after this time period), but a large change during the little ice age suggests

that climate change can radically change the fires (van der Werf et al., 2013). While the IPCC has used 1850 to 1900 as the preindustrial period for climate simulations (e.g. Allen et al., 2018), for aerosols this is not an ideal
310 time period, as it is likely that aerosols are already elevated due to anthropogenic activities during this time period, while 1750 could be better, although still part of the little ice age. The issue of what is the right baseline for preindustrial aerosols is important also for considering paleoclimates (Section 5). As shown in Fig. 3 and 5, the uncertainty in preindustrial to present day emission changes in aerosols is driven by preindustrial emission uncertainties, which is partly associated with variability across the preindustrial time period.

315

There are, of course, constraints on present day radiative effects from aerosols from satellites and other tools, which can constrain the last 30-40 years (e.g. Bellouin et al., 2020). And there are energy constraints on the present day to preindustrial change in aerosol radiative forcing using energy balance constraints (Smith et al., 2021; Sherwood et al., 2020) which result in an 57% reduction in the uncertainty using fixed emissions ($\Sigma_{Const_Process}^{PD-PI}=1.6 \text{ W/m}^2$).
320 Unconstrained emission uncertainties and unconstrained process uncertainties have yet to be combined in a rigorous method like Bellouin et al. (2020) did for process uncertainties (in that study they assume that emissions are well known), but this should be done in the future. Adding in the uncertainties in emissions, especially from wildfires, would mean that while directly emitted anthropogenic aerosols are going up (as estimated in CMIP6), wildfire emissions may be going down. The wildfire aerosols resulting from these emissions would thus partially offset the
325 radiative cooling from the increase in anthropogenic aerosols. Thus, if we take the case of high wildfires in the preindustrial (-2 W/m^2 in Fig. 4), this could imply that estimates of aerosol radiative forcing from purely anthropogenic sources today which are large (-2 W/m^2) would be more likely; the 1850 to 2000 aerosol radiative forcing would be the same as assumed now, but the 1850 aerosol radiative forcing would already be large. This would have important implications for climate warming over the next few decades, as anthropogenic emissions of
330 aerosols are likely to decrease, leading to more warming than projected without including preindustrial aerosol emission uncertainties.

5 Characterizing paleoclimate aerosol forcing uncertainty

335 Unfortunately, except for dust or wildfires in certain time periods (Albani et al., 2015; Power et al., 2007; Zennaro et al., 2014; Marlon et al., 2008), there is very little information about the distribution or amount of aerosols in different climate regimes, and therefore we do not know the emissions well, nor then the impact of those emissions onto climate. We can envision these uncertainties are mostly unknown unknowns. We have some information that

they are likely to be large (since aerosol uncertainties today are so large, and we know less about paleoclimate aerosols), but we cannot yet directly constrain these. We do know that there were large fluctuations: for example, dust was likely 2-4x higher in the last glacial maximum than today (Lambert et al., 2015; Mahowald et al., 1999; Albani et al., 2014; Albani and Mahowald, 2019), while between preindustrial and present day, the change is smaller at only approximately 2x (Kok et al., 2023) (Fig. 6). For dust, we have estimates at the last glacial maximum (LGM) and 6000 bpa (Albani and Mahowald, 2019; Albani et al., 2014), which suggest that the changes in radiative forcing could be on the order of 0 to -2 W/m² (Albani et al., 2018), although studies using carefully compared dust optics show smaller radiative forcings, because dust both absorbs and reflects both short and long wave radiation (Albani and Mahowald, 2019; Braconnot et al., 2021).

But changes between preindustrial and present day aerosol radiative forcings are dominated by changes in fires (Section 1 and 2): are these changes are large in the paleodata as seen in the last 150 years? The limited paleodata suggests large changes in fires during different time periods in the past (Fischer et al., 2015; van der Werf et al., 2013; Zennaro et al., 2014; Arienzo et al., 2017). For example, in considering cold periods like the last glacial maximum, there is likely a large reduction in fires in high latitudes, due to the presence of the Laurentide and Fenno-Scandinavian icesheets, which is consistent with fire proxies in Greenland ice cores (e.g. the ammonium record for the North Greenland Ice core Project (Fischer et al., 2015) . Generally, the charcoal record suggests lower fire frequency in the last glacial maximum than preindustrial (Marlon et al., 2016), although ice sheets could have removed sediment records of wildfires (Fig. 6). For climate impacts, the low and mid latitude fires tend to be more important today (Hamilton et al., 2018), so more information on the frequency and extent of wildfires in those regions are the most important, and difficult to retrieve from ice cores. The changes seen in wildfires between preindustrial and for example, last glacial maximum are as large if not larger than those seen between preindustrial and present day (Fig. 6). These studies suggest qualitatively that the changes we have seen in fires over the preindustrial to present day are not unprecedented in size, but rather are similar to paleoclimate changes.

In addition, paleoclimate data such as temperature changes or aerosol changes are done relative to preindustrial changes, and as discussed in Section 2, and shown in Fig. 5, there is substantial variability in preindustrial fires. It is unclear what value to use for preindustrial aerosol emissions to compare to paleodata values: do we use 1850 values or values from the little ice age? Or some average? One can think of the variability in changes in emissions between some time (T) and PI as being shown in Equation 4. The uncertainties in emissions in PI are driving the uncertainties in PD-PI describe above and are about 2.8W/m² (Σ_{Emis}^{PI}): since these uncertainties are due to

variability not a lack of information, increased knowledge is unlikely to reduce these uncertainties from variability (only if the new studies show less variability). In addition, the changes in emissions between PI and any other time period are likely to be similarly large, but may not be orthogonal (Σ_{Emis}^T). Adding these together (using Eq. 4), we obtain $\Sigma_{Emis}^{T-PI}=4.0 \text{ W/m}^2$. Note that the uncertainties in emissions proposed here for different time periods could be constrained to some extent (Σ_{Emis}^T), but uncertainties due to the variability in PI emissions (Fig. 5) would be difficult to constrain, and there may be substantial variability as well as uncertainty in emissions during different time periods, so the values proposed here may actually underestimate the uncertainty.

$$\Sigma_{Emis}^{T-PI} = [(\Sigma_{Emis}^{PI})^2 + (\Sigma_{Emis}^T)^2]^{1/2}, \quad (4)$$

Once we have paleoproxies to provide data about changes in fire emissions, especially, we can constrain the emission uncertainties for paleo time periods relative to present day, hopefully. Unfortunately knowing the emissions does not translate into knowing the radiative forcing in past times, as we known from our experience simulating preindustrial to present day emission changes in existing models using the same emissions (Bellouin et al., 2020). There are uncertainties of translating these changes in emissions into changes in direct radiative and aerosol-cloud interactions or process uncertainties, which we assume here, since we do not have better information, that these are a similar size to present day to preindustrial uncertainties ($\Sigma_{Process}^{PD-PI}=\Sigma_{Process}^{T-PI}= 2.8 \text{ W/m}^2$). These uncertainties are due to differences in the modeling of aerosols, and assumptions about size and how aerosols interact with clouds which can be different depending on where the aerosols are emitting: this uncertainty will remain in paleoclimates, and might even become larger, since the aerosol size, composition and mixing state could be quite different and the very important impact of aerosols onto clouds is sensitive to the background conditions (Carslaw et al., 2017). Fires from different ecosystems, or even different types of fires in the same ecosystems, have very different emissions of black carbon, organic carbon and sulfate, and thus different effects, but we do not know how these will change in different time periods. Natural aerosols are the source of much the uncertainty in today's climate compared to anthropogenic aerosols, because of the difficulty of the estimating the exact timing and distribution of emissions, as well as the sources are more complicated in composition and location (e.g. Mahowald et al., 2011b; Carslaw et al., 2017; Rathod et al., 2020). Similar to the present day relative to preindustrial, we can estimate the paleotime to preindustrial radiative forcing uncertainty using equation 3 and obtain 4.8 W/m^2 as the range of uncertainty for paleotime periods. (Notice that if we can constrain the change in

400 radiative forcing from changing emissions of aerosols to be smaller than that between preindustrial and present day, using observations and equation 2, it is possible we could proportionately reduce the uncertainty in radiative forcing from aerosols from process uncertainties, see equation 2.) Converting this radiative forcing uncertainty into a feedback uncertainty requires knowing the temperature change, which is also uncertain, but if we use 3C as a reasonable value, the aerosol feedback uncertainty derived from the last glacial maximum to PI is $\pm 1.6 \text{ W/m}^2/\text{K}$,
405 similar to the value derived from the PD-PI time period. If we add in the uncertainty in temperature change between preindustrial and last glacial maximum, this estimate would be even larger, of course.

It seems likely that aerosol emissions from fires during the last glacial maximum are much smaller than preindustrial or present day values, while estimates suggest dust is $\sim 3\times$ larger in the last glacial maximum than the
410 present day. Will these changes in aerosols balance out? That is unlikely but vital to consider. Dust is by mass the most important aerosol in the atmosphere, and contributes substantial to direct forcing and ice nucleation processes, but fire emissions are important for liquid aerosol-cloud processes (Mahowald et al., 2011a; Carslaw et al., 2010). Because of the non-linearity in aerosol-cloud interactions, small changes in fire emissions in pristine environments, like the last glacial maximum, might be even more important than estimated here (Carslaw et al., 2017).
415 Understanding the aerosol interactions with clouds especially for the last glacial maximum is both important and intriguing.

Since today the largest uncertainties in the radiative forcing come from aerosol uncertainties, estimates in past climates should ensure that aerosol uncertainties remain one of the largest uncertainties in those times as well: how
420 could we know the change of aerosols from some paleoclimate to preindustrial better than we know the change in aerosol forcing between preindustrial to present day? If ice sheets or insolation or continental distributions are different and causing large changes in top of atmosphere fluxes and thus climate regime, most likely aerosol changes are equally large, but we do not know in what direction. More analysis might result in even larger changes in radiative forcing and its uncertainty in some time periods, since here we are assuming, without prior information,
425 that the radiative forcing of any paleotime period relative to preindustrial is around 0.0 W/m^2 . For the LGM, for example, if we only include dust, a more negative value should be chosen as the mean, since we have evidence of increased dust in the paleorecord (Albani et al., 2018). On the other hand, the limited data suggests that fires have substantially decreased relative to preindustrial, which would warm the climate. It is beyond the scope of this opinion piece to characterize the central estimate, but rather here we just point out the many uncertainties in these
430 estimates.

6 Aerosol processes and other sources of uncertainty

So far here we have focused on the more frequently studied processes of aerosol direct radiative effects, and aerosol-
435 cloud interactions with an emphasis on cloud condensation nuclei. However there remain substantial uncertainties
in these aerosol radiative effects even in the current climate (Bellouin et al., 2020; Li et al., 2022). Aerosols are
spatially and temporally heterogeneous in composition, size and amount, leading to vastly different physical and
chemical properties. They are in general poorly observed compared to meteorological phenomenon (e.g. Naik et
al., 2021). Not only the bulk composition matters but the details of the mixing state and size are vital for radiative
440 and cloud interactions (Matsui et al., 2018; Bond et al., 2013; Li et al., 2022, 2021). In addition, in preindustrial
times, the impact of aerosols, for example on cloud properties, can be different than present day because of a lower
background aerosol amount (Carslaw et al., 2017; Hamilton et al., 2014).

The impacts of large changes in important ice nuclei such as dust or primary biogenic particles is likely to be large
445 but has yet to be fully assessed (Burrows et al., 2013; Murray et al., 2021; Storelvmo, 2017). Another important
feedback that is relatively well known but not included in most climate models is due to nitrogen aerosols such as
ammonium or nitrate (e.g. Bauer et al., 2007; Paulot et al., 2016). Future concentrations of aerosols deriving from
land use practices such as ammonia or nitrate are not likely to decrease as quickly as from fossil fuels (Gidden et
al., 2019). Indeed, as sulfate is phased out, more nitric acid will form nitrate aerosols (due to higher pH), partially
450 buffering decreases in aerosol AOD (Paulot et al., 2016; Pye et al., 2009). Including better parameterizations of
ice nucleating particles and nitrogen aerosols is key to improving future aerosol projections.

In addition, aerosols can provide nutrients and pollutants to different ecosystems (Mahowald et al., 2017; Hamilton
et al., 2021), linking aerosol changes to changes in biogeochemistry and the carbon cycle. These effects could
455 potentially be quite large ($0.5 \text{ W/m}^2 + 0.4 \text{ W/m}^2$) (Mahowald, 2011), but are poorly constrained, and do not
explicitly appear in the standard radiative forcing diagram, since they reflect CO_2 that is not in the atmosphere, but
could have been (Mahowald, 2011).

Another natural emission to which estimates of radiative forcing are sensitive to is biogenic volatile organic
460 compound emissions (BVOCs). BVOCs are a major source of new aerosol particles in the atmosphere. (Guenther
et al., 2006; Arneth et al., 2010). Furthermore, biogenic particle formation processes contributed more to the aerosol

burden in the PI than the PD (Gordon et al., 2016). Estimates of the radiative forcing of BVOC are sensitive to how well characterized new particle formation processes are in a model. The recent addition of an organic particle formation pathway, which occurs in the absence of SO₂ (such the PI), results in an increased aerosol burden in the past than the present. Once more increasing the PI aerosol burden reduces the estimate of the aerosol forcing over the historical period, this time by reducing the cloud forcing by $\sim 0.2 \text{ W/m}^2$ (Zhu et al., 2019).

7 Implications of including uncertainty in emissions in radiative forcing estimates

Aerosol radiative forcing and its uncertainty is used extensively in climate change science, including to constrain climate sensitivity (Sherwood et al., 2020) and thereby future climate changes (IPCC, 2021; Gulev et al., 2021; Szopa et al., 2021). Because the published aerosol uncertainties tend not to include poorly constrained uncertainties such as discussed here, this information is not effectively passed to physical climate scientists who use these estimates. For example, a recent review of climate sensitivity (Sherwood et al., 2020) focused on using independent methods to reduce uncertainty in climate sensitivity. In that paper, aerosol radiative forcing uncertainties for different time periods are mentioned in several different places. They use the unconstrained model range of the aerosol radiative forcing obtained by (Bellouin et al., 2020), which as discussed above, does not account for emission uncertainties. Paleoclimate constraints are often used for constraining climate sensitivity, as discussed in (Sherwood et al., 2020). Currently there exist estimates for dust aerosol radiative forcing changes between last glacial maximum and current, which is included in (Sherwood et al., 2020) as $-1.0 \pm 1.6 \text{ W/m}^2$ (90% confidence intervals: they report 1 sigma values of $\pm 1 \text{ W/m}^2$ in Section 5.2.2 which are converted to 90% confidence here by multiplying by 1.6 as a first estimate) but no mention is made of the potential for changes in the more important wildfires. As discussed in Section 4, estimates for radiative forcing of aerosols for paleotime periods especially wildfires are missing, but should be estimated to be $0.0 \pm 2.4 \text{ W/m}^2$ (90% confidence interval). In addition, the aerosol feedback within the system is assumed in Sherwood et al., 2020 to have an uncertainty of $\pm 0.22 \text{ Wm}^2$ (they report 1 sigma values of $\pm 0.15 \text{ W/m}^2$ in section 3.2 which here we convert to 90% confidence intervals) whereas here we estimated the aerosol feedback uncertainty to be $\pm 1.6 \text{ W/m}^2$ (90% confidence), substantially larger. Including more realistic aerosol uncertainties into estimates of climate sensitivity should be done to ensure adequate propagation of errors, although they are unlikely to change the central estimates (Sherwood et al., 2020).

In addition, some authors argue that there were not significant changes in aerosol radiative forcing during the 1970s and 1980s, using standard CMIP6-type estimates, and try to estimate climate sensitivity in this time period

(Jiménez-de-la-Cuesta and Mauritsen, 2019). As noted above, however, the 1970s is a time period of the Sahel drought, and dust radiative forcing between 1960s and 1980s changed by perhaps $-0.57 \pm 0.46 \text{ W/m}^2$ (Mahowald et al., 2010), suggesting that is not an ideal time period to target. One should add onto this estimate the possibly important changes in wildfires which could have occurred over this time period but for which we do not have data.

In addition to climate sensitivity, some studies use the CMIP6 simulations to constrain past aerosol radiative forcing changes (e.g. Smith et al., 2021). Since the simulations do not include different spatial and temporal uncertainties in emissions, they are not including the real uncertainty in the aerosol forcing. Other studies use CMIP6 or similar simulations to attribute the change in temperatures or precipitation to different forcings (e.g. Biasutti and Giannini, 2006; Undorf et al., 2018; Hegerl et al., 2019), and these attempts to attribute changes could be based on a poor representation of the real uncertainty in the preindustrial of the aerosol forcing. Aerosol radiative forcing uncertainty cannot be constrained easily by temperature time series, since other uncertainties can be difficult to pull apart from aerosol uncertainties (Kiehl, 2007; Lee et al., 2016). Attributing climate at a regional scale is the next frontier of detection and attribution, but this cannot be done without accurate aerosol histories (Lehner and Coats, 2021).

In summary, we argue that it is critical that the full uncertainty deduced in the aerosol literature, including due to changes in natural aerosols, be passed to physical climate scientists so that they can accurately account for these in constraints on climate sensitivity and in projections of future climate changes.

8 Conclusions: Pathway to improve historical and paleoclimate characterization of uncertainties

How can we address the systematic underestimate in the uncertainty of changes in aerosol radiative effects between different time periods? Here we propose some steps towards first characterizing the true uncertainties, including emission uncertainties, and then using observations to constrain these aerosol pathways to constrain the radiative forcing and uncertainties.

a. Characterize historical uncertainties in aerosol and aerosol precursor emissions, using available knowledge of emissions, and how they might have changed. These estimates should include some versions which are consistent with available paleodata (e.g., for dust Kok et al., 2023; and for wildfires Hamilton et al., 2018 and Liu et al., 2021). The uncertainties from emissions should be combined with uncertainties in aerosol

- processes to create a more robust uncertainty bound for different time periods. Note that many ‘natural’ aerosols are prognostic in the models (e.g. dust), and therefore to in order to match available scenarios, the prognostic aerosol schemes may require using a temporally varying emission factor to enable them to simulate the correct temporal trends (e.g. Mahowald et al., 2010).
- b. Characterize paleoclimate emissions of aerosols and the resulting radiative forcing at important past climates, such as last glacial maximum and last interglacial. These estimates should be based as much as possible on observations, and possible ranges.
- c. We propose a new intercomparison project (AEROHISTMIP) which would include multiple emission pathways in the historical model simulations conducted for CMIP exercises. Several Earth system models should not only conduct ensemble members of one aerosol emission scenario, but also use multiple aerosol emission scenarios to better understand the uncertainty in aerosol radiative forcing and climate response due to the uncertainties in aerosol emissions in different time periods. The evolution of several related past model intercomparison projects under one umbrella (e.g. Composition, Air quality, Climate inTeractions Initiative: CACTI) provides the ideal opportunity now to include such simulations.
- d. Constrain preindustrial to present day aerosol radiative effects. From (c), combined with observations (a), the most likely past emissions scenarios can be identified, and we can make the first steps towards constraining uncertainty, similar to the efforts underway to characterize which of the climate models are most reliable (e.g. IPCC, 2021). Any inability of existing models to simulate observations, as well as other remaining uncertainties, should be carefully assessed: it may not be possible for the models to simulate the observed changes.
- e. Obtain more paleoclimate proxies for aerosol concentrations. Here we have focused mostly on wildfires and dust, since there is enough paleoclimate data to show that CMIP6 does not represent historical changes in these aerosols well, but indeed it is not possible currently to validate the changes in emissions for other natural (e.g., sea spray) and anthropogenic (e.g., sulfate) aerosols as well. We need the development of more proxies for historical and paleoclimate changes in aerosols to increase confidence in our estimates of aerosol radiative forcing.
- f. Continue to improve aerosol measurement databases, including more in situ observations of the aerosol composition in more locations (Snider et al., 2016), as well as continued use of satellite observations to constrain the magnitude of aerosol radiative forcing (e.g. Smith et al., 2021). We encourage more observations for variables directly related to the radiative forcing (e.g., aerosol optical depth) and those that could help narrow uncertainties in crucial parameters that describe related physiochemical processes. Some of these data

555 can also be used to assess the model performance, narrowing the model spread, and validate satellite retrievals. As aerosol number concentration, the determinant of the change in cloud properties to emission changes, cannot be retrieved from paleo proxies (d) there needs to be a simultaneous effort in understanding natural aerosol processes and impacts on clouds under pristine “PI-like” present day conditions (Hamilton et al., 2014; McCoy et al., 2020)

560 g. Characterize current model direct aerosol radiative effects and aerosol cloud interactions using new tools. Currently meteorological models are tightly connected to the aerosol models they host, making it difficult to independently evaluate the structural differences between aerosol models. Recent efforts to develop generalized chemical and aerosol interfaces would allow more effective evaluation of chemical and aerosol schemes separate from their host models (Hodzic et al., submitted). A generalized framework could also
565 allow artificial intelligence methods to be integrated into multiple models and model-data comparisons or assimilations to be used across models.

In summary: while there has been substantial progress in aerosol-climate science over the past 20 years, aerosols remain one of the most important uncertainties in climate change science, and are likely to continue to be important
570 to study for at least the next 20 years.

Data Availability

N/A

575 Author contribution

NMM conceived of the idea and created the first draft, with substantial improvements from LL, SA and JFK.

Competing interests

580 None

Acknowledgements

NMM and LL would like to acknowledge support from DOE DE-SC0021302. JFK acknowledges support from
585 NSF grants 1856389 and 2151093. We acknowledge helpful discussions with Flavio Lehner and Steve Sherwood.

References

- Albani, S. and Mahowald, N. M.: Paleodust insights onto dust impacts on climate, *J Clim*, 7897–7913, <https://doi.org/10.1175/jcli-d-18-0742.1>, 2019.
- Albani, S., Mahowald, N. M., Perry, A. T., Scanza, R. A., Zender, C. S., Heavens, N. G., Maggi, V., Kok, J. F., and Otto-
590 Bliesner, B. L.: Improved dust representation in the Community Atmosphere Model, *J Adv Model Earth Syst*, 6, 541–570, <https://doi.org/10.1002/2013MS000279>, 2014.
- Albani, S., Mahowald, N. M., Winckler, G., Anderson, R. F., Bradtmiller, L. I., Delmonte, B., Francois, R., Goman, M., Heavens, N. G., Hesse, P. P., Hovan, S. A., Kang, S. G., Kohfeld, K. E., Lu, H., Maggi, V., Mason, J. A., Mayeswki, P., McGee, D., Miao, X., Otto-Bliesner, B. L., Perry, A. T. A., Pourmand, A., Roberts, H. M., Rosenbloom, N.,
595 Stevens, T., Sun, J., François, R., Goman, M., Heavens, N. G., Hesse, P. P., Hovan, S. A., Kang, S. G., Kohfeld, K. E., Lu, H., Maggi, V., Mason, J. A., Mayewski, P. A., McGee, D., Miao, X., Otto-Bliesner, B. L., Perry, A. T. A., Pourmand, A., Roberts, H. M., Rosenbloom, N., Stevens, T., and Sun, J.: Twelve thousand years of dust: the Holocene global dust cycle constrained by natural archives, *Climate of the Past*, 11, 869–2015, <https://doi.org/10.5194/cp-11-869-2015>, 2015.
- 600 Albani, S., Balkanski, Y., Mahowald, N., Winckler, G., Maggi, V., and Delmonte, B.: Aerosol-Climate Interactions During the Last Glacial Maximum, *Curr Clim Change Rep*, 4, 99–114, <https://doi.org/10.1007/s40641-018-0100-7>, 2018.
- Allen, M. R., Dube, O. P., Solecki, W., Aragón-Durand, F., Cramer, W., Humphreys, S., Kainuma, M., Kala, J., Mahowald, N., Mulugetta, Y., Perez, R., Wairiu, M., and Zickfeld, K.: Framing and Context, in: *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global*
605 *greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, edited by: [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P. R. S., A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. L., and T. Maycock, M. Tignor, and T. W., World Meteorological Organization (WMO), Geneva, Switzerland, 45–91, 2018.
- Andela, N., Morton, D. C., Giglio, L., Chen, Y., Van Der Werf, G. R., Kasibhatla, P. S., DeFries, R. S., Collatz, G. J., Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Yue, C., and Randerson, J. T.: A human-driven decline in global burned area, *Science* (1979), 356, 1356–1362, <https://doi.org/10.1126/science.aal4108>, 2017.
- 610 Arienzo, M. M., McConnell, J. R., Murphy, L. N., Chellman, N., Das, S., Kipfstuhl, S., and Mulvaney, R.: Holocene black carbon in Antarctica paralleled Southern Hemisphere climate, *J Geophys Res*, 122, 6713–6728, <https://doi.org/10.1002/2017JD026599>, 2017.
- 615 Arneeth, A., Harrison, S., Zaehle, S., Tsigaridis, K., Menon, S., Bartlein, P., Feichter, J., Korhola, A., Kulmala, M., O'Donnell, D., Schurgers, G., Sorvari, S., and Vesala, T.: Terrestrial biogeochemical feedbacks in the climate system, *Nature-geoscience*, 3, 525–532; DOI: 10.1038/NGEO905, 2010.

- Bauer, S. E., Koch, D., Unger, N., Metzger, S. M., Shindell, D. T., and Streets, D. G.: Atmospheric Chemistry and Physics
 620 Nitrate aerosols today and in 2030: a global simulation including aerosols and tropospheric ozone, *Atmos. Chem. Phys.*, 5043–5059 pp., 2007.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M.,
 Daniaou, A. L., Dufresne, J. L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U.,
 625 Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M.,
 Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding
 Global Aerosol Radiative Forcing of Climate Change, <https://doi.org/10.1029/2019RG000660>, 1 March 2020.
- Biasutti, M. and Giannini, A.: Robust Sahel drying in response to late 20th century forcings, *Geophys Res Lett*, 33,
 doi:10.1029/2006GL026067, 2006.
- Bistasas, I., Oom, D., Sá, A. C. L., Harrison, S. P., Prentice, I. C., and Pereira, J. M. C.: Relationships between human
 630 population density and burned area at continental and global scales, *PLoS One*, 8, 1–12,
<https://doi.org/10.1371/journal.pone.0081188>, 2013.
- Bond, T., Bhardwaj, E., Dong, R., Jogani, R., Jung, S., Roden, C., Streets, D., and Trautman, N.: Historical emissions of black
 and organic carbon aerosol from energy-related combustion, 1850-2000, *Global Biogeochem Cycles*, 21,
 doi:10.1029/2006GB002840, 2007.
- 635 Bond, T., Doherty, S. J., Fahey, D., Forster, P., Bernsten, T., DeAngelo, B., Flanner, M., Ghan, S., Karcher, B., Koch, D.,
 Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M., Schultz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N.,
 Guttikunda, S., Hopke, P., Jacobson, M., Kaiser, J. W., Klimont, Z., Lohman, U., Schwartz, J., Shindel, D., Storelvmo,
 T., Warren, S., and Zender, C.: Bounding the role of black carbon in the climate system: A scientific assessment, *J*
Geophys Res, D118, 5380–5552; doi:10.1002/jgrd_50171, 2013.
- 640 Bowman, D. M. J. S., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. a, D-Antoio, C., DeFries, R. S., Doyle, J. C.,
 Harrison, S. P., Johnson, F., Keeley, J. E., Krawchuk, M. a, Kull, C. a, MArtson, J., Mortiz, M., Prentice, I. C., Roos,
 C. I., Scott, A. C., Swetnam, T. W., Werf, G. van der, Pyne, S. J., Blach, J., Bowman, D. M. J. S., Balch, J. K., Artaxo,
 P., Bond, W. J., Carlson, J. M., Cochrane, M. a, D’Antonio, C. M., DeFries, R. S., Doyle, J. C., Harrison, S. P.,
 Johnston, F. H., Keeley, J. E., Krawchuk, M. a, Kull, C. a, Marstone, J., Moritz, M. a, Prentice, C., Roos, C. I., Scott,
 645 A. C., Swetnam, T. W., G.R., van der W., Pyne, S. J., Marston, J. B., Moritz, M. a, Prentice, I. C., Roos, C. I., Scott,
 A. C., Swetnam, T. W., van der Werf, G. R., and Pyne, S. J.: Fire in the Earth System, *Science* (1979), 324,
 10.1126/science.1163886, <https://doi.org/10.1126/science.1163886>, 2009.
- Braconnot, P., Albani, S., Balkanski, Y., Cozic, A., Kageyama, M., Sima, A., Marti, O., and Peterschmitt, J. Y.: Impact of dust
 in PMIP-CMIP6 mid-Holocene simulations with the IPSL model, *Climate of the Past*, 17, 1091–1117,
 650 <https://doi.org/10.5194/cp-17-1091-2021>, 2021.
- Burnett, R., Chen, H., Szyszkowicz, M., Fann, N., Hubbell, B., Pope, C. A., Apte, J. S., Brauer, M., Cohen, A., Weichenthal,
 S., Coggins, J., Di, Q., Brunekreef, B., Frostad, J., Lim, S. S., Kan, H., Walker, K. D., Thurston, G. D., Hayes, R. B.,

- Lim, C. C., Turner, M. C., Jerrett, M., Krewski, D., Gapstur, S. M., Diver, W. R., Ostro, B., Goldberg, D., Crouse, D. L., Martin, R. V., Peters, P., Pinault, L., Tjepkema, M., Van Donkelaar, A., Villeneuve, P. J., Miller, A. B., Yin, P., Zhou, M., Wang, L., Janssen, N. A. H., Marra, M., Atkinson, R. W., Tsang, H., Thach, T. Q., Cannon, J. B., Allen, R. T., Hart, J. E., Laden, F., Cesaroni, G., Forastiere, F., Weinmayr, G., Jaensch, A., Nagel, G., Concin, H., and Spadaro, J. V.: Global estimates of mortality associated with longterm exposure to outdoor fine particulate matter, *Proc Natl Acad Sci U S A*, 115, 9592–9597, <https://doi.org/10.1073/pnas.1803222115>, 2018.
- Burrows, S. M., Hoose, C., Pöschl, U., and Lawrence, M. G.: Ice nuclei in marine air: Biogenic particles or dust?, *Atmos Chem Phys*, 13, 245–267, <https://doi.org/10.5194/acp-13-245-2013>, 2013.
- Carslaw, K. S., Boucher, O., Spracklen, D., Mann, G., Rae, J. G., Woodward, S., and Kumala, M.: A review of natural aerosol interactions and feedbacks within the Earth System, *Atmos Chem Phys*, 10, 1701–1737, <https://doi.org/10.5194/acp-10-1701-2010>, 2010.
- Carslaw, K. S., Gordon, H., Hamilton, D. S., Johnson, J. S., Regayre, L. A., Yoshioka, M., and Pringle, K. J.: Aerosols in the Pre-industrial Atmosphere, *Curr Clim Change Rep*, 3, 1–15, <https://doi.org/10.1007/s40641-017-0061-2>, 2017.
- Cusworth, D. H., Mickley, L. J., Sulprizio, M. P., Liu, T., Marlier, M. E., Defries, R. S., Guttikunda, S. K., and Gupta, P.: Quantifying the influence of agricultural fires in northwest India on urban air pollution in Delhi, India, *Environmental Research Letters*, 13, <https://doi.org/10.1088/1748-9326/aab303>, 2018.
- Dentener F, Kinne S, Bond T, Boucher O, Cofala J, Generoso S, Ginoux P, Gong S, Hoelzemann J J, Ito A, Marelli L, Penner J E, Putaud J-P, Textor C, Schulz M, Van Der Werf G R and Wilson J 2006 *Atmospheric Chemistry and Physics Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom* vol 6 Online: www.atmos-chem-phys.net/6/4321/2006/
- Ferretti, D. F., Miller, J. B., White, J. W. C., Etheridge, D. M., Lassey, K. R., Lowe, D. C., Meure, C. M. M., Dreier, M. F., Trudinger, C. M., Van Ommen, T. D., and Langenfelds, R. L.: Unexpected Changes to the Global Methane Budget over the Past 2000 Years, *Science* (1979), 309, 1714–1717, <https://doi.org/doi:10.1126/science.1115193>, 2005.
- Fischer, H., Schüpbach, S., Gfeller, G., Bigler, M., Röthlisberger, R., Erhardt, T., Stocker, T. F., Mulvaney, R., and Wolff, E. W.: Millennial changes in North American wildfire and soil activity over the last glacial cycle, *Nat Geosci*, 8, 723–727, <https://doi.org/10.1038/ngeo2495>, 2015.
- Forster, P., Alterskjaer, K., Smith, C., Colman, R., Damon Matthews, H., Ramaswamy, V., Storelvmo, T., Armour, K., Collins, W., Dufresne, J., Frame, D., Lunt, D., Mauritsen, T., Watanabe, M., Wild, M., Zhang, H., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.: Chapter 7: The Earth’s Energy Budget, Climate Feedbacks and Climate Sensitivity, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Masson-Delmotte, V. , P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and Zhou, B.,

Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 923–1054,
<https://doi.org/10.1017/9781009157896.009>, 2021.

- 690 Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., Van Vuuren, D. P., Van Den Berg, M., Feng, L.,
Klein, D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O., Harmsen, M., Hasegawa, T., Havlik, P., Hilaire, J.,
Hoesly, R., Horing, J., Popp, A., Stehfest, E., and Takahashi, K.: Global emissions pathways under different
socioeconomic scenarios for use in CMIP6: A dataset of harmonized emissions trajectories through the end of the
century, *Geosci Model Dev*, 12, 1443–1475, <https://doi.org/10.5194/gmd-12-1443-2019>, 2019.
- 695 Ginoux, P., Prospero, J., Gill, T. E., Hsu, N. C., and Zhao, M.: Global scale attribution of anthropogenic and natural dust
sources and their emission rates based on MODIS deep blue aerosol products, *Reviews of Geophysics*, 50,
DOI:10.1029/2012RG000388, 2012.
- 700 Gordon, H., Sengupta, K., Rap, A., Duplissy, J., Frege, C., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J.,
Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Dunne, E. M., Adamov, A., Amorim, A., Bernhammer, A.-K.,
Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J. S., Dias, A., Ehrhart, S., Fischer, L., Flagan, R. C.,
Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J.,
Kirkby, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Monks, S.
A., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K. J., Richards, N. A. D., Rissanen, M. P.,
Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sharma, S., Sipilä, M., Steiner, G., Stozhkov,
Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer,
705 D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U.,
Kulmala, M., Curtius, J., and Carslaw, K. S.: Reduced anthropogenic aerosol radiative forcing caused by biogenic
new particle formation, *Proceedings of the National Academy of Sciences*, 113, 12053–12058,
<https://doi.org/10.1073/pnas.1602360113>, 2016.
- 710 Gulev, S.K., P.W. Thorne, J. Ahn, F.J. Dentener, C.M. Domingues, S. Gerland, D. Gong, D.S. Kaufman, H.C. Nnamchi,
J. Quaas, J.A. Rivera, S. Sathyendranath, S.L. Smith, B. Trewin, K. von Schuckmann, and R.S. Vose, 2021: Changing
State of the Climate System. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group
I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai,
A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E.
Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge
715 University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 287–422,
doi:10.1017/9781009157896.004.
- Hamilton, D. S., Lee, L. A., Pringle, K. J., Reddington, C. L., Spracklen, D. V., and Carslaw, K. S.: Occurrence of pristine
aerosol environments on a polluted planet., *Proc Natl Acad Sci U S A*, 111, 18466–71,
720 <https://doi.org/10.1073/pnas.1415440111>, 2014.

- Hamilton, D. S., Hantson, S., Scott, C. E., Kaplan, J. O., Pringle, K. J., Nieradzik, L. P., Rap, A., Folberth, G. A., Spracklen, D. V., and Carslaw, K. S.: Reassessment of pre-industrial fire emissions strongly affects anthropogenic aerosol forcing, *Nat Commun*, 9, <https://doi.org/10.1038/s41467-018-05592-9>, 2018.
- 725 Hamilton, D. S., Perron, M. M. G., Bond, T. C., Bowie, A. R., Buchholz, R. R., Guieu, C., Ito, A., Maenhaut, W., Myriokefalitakis, S., Olgun, N., Rathod, S. D., Schepanski, K., Tagliabue, A., Wagner, R., and Mahowald, N. M.: Earth, Wind, Fire, and Pollution: Aerosol Nutrient Sources and Impacts on Ocean Biogeochemistry, <https://doi.org/10.1146/annurev-marine-031921>, 2021.
- Hegerl, G. C., Brönnimann, S., Cowan, T., Friedman, A. R., Hawkins, E., Iles, C., Müller, W., Schurer, A., and Undorf, S.: Causes of climate change over the historical record, <https://doi.org/10.1088/1748-9326/ab4557>, 9 December 2019.
- 730 Hodzic, A., Mahowald, N., Dawson, M., Johnson, J., Bernardet, L., Bosler, P. A., Clune, T., Fast, J. D., Fierce, L., Liu, X., Ma, P.-L., Murphy, B., Riemer, N., and Schulz, M.: Generalized Aerosol/chemistry iNterface (GIANT): a community effort to advance collaborative science across weather and climate models, submitted, *Bulletin of the American Meteorological Society*.
- 735 Hoesly R M, Smith S J, Feng L, Klimont Z, Janssens-Maenhout G, Pitkanen T, Seibert J J, Vu L, Andres R J, Bolt R M, Bond T C, Dawidowski L, Kholod N, Kurokawa J I, Li M, Liu L, Lu Z, Moura M C P, O'Rourke P R and Zhang Q 2018 Historical (1750-2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS) *Geosci Model Dev* **11** 369–408
- Hooper, J. and Marx, S.: A global doubling of dust emissions during the Anthropocene?, *Glob Planet Change*, 169, 70–91, <https://doi.org/10.1016/j.gloplacha.2018.07.003>, 2018.
- 740 IPCC: IPCC, 2019: Summary for Policymakers., in: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, edited by: Pörtner, H.-O., Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N. M., 2019.
- 745 IPCC: Summary for Policymakers, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou, Cambridge University Press, Cambridge, UK, 3–32, <https://doi.org/10.1017/9781009157896.001>, 2021.
- Jiménez-de-la-Cuesta, D. and Mauritsen, T.: Emergent constraints on Earth's transient and equilibrium response to doubled CO₂ from post-1970s global warming, *Nat Geosci*, 12, 902–905, <https://doi.org/10.1038/s41561-019-0463-y>, 2019.
- 750 Kiehl, J. T.: Twentieth century climate model response and climate sensitivity, *Geophys Res Lett*, 34, <https://doi.org/10.1029/2007GL031383>, 2007.
- Klein-Goldewijk, K.: Estimating global land use change over the past 300 years: the HYDE database, *Global Biogeochem Cycles*, 15, 417–434, 2001.

- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions of particulate matter including black carbon, *Atmos Chem Phys*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, 2017.
- Kloster, S., Mahowald, N. M., Randerson, J. T., Thornton, P. E., Hoffman, F. M., Levis, S., Lawrence, P. J., Feddema, J. J., Oleson, K. W., and Lawrence, D. M.: Fire dynamics during the 20th century simulated by the Community Land Model, *Biogeosciences*, 7, <https://doi.org/10.5194/bg-7-1877-2010>, 2010.
- Knorr, W., Kaminski, T., Arneth, A., and Weber, U.: Impact of human population density on fire frequency at the global scale, *Biogeosciences*, 11, 1085–1102, <https://doi.org/10.5194/bg-11-1085-2014>, 2014.
- Kok, J.F., Storelvmo, T., Karydis, V.A. *et al.* Mineral dust aerosol impacts on global climate and climate change. *Nat Rev Earth Environ* 4, 71–86 (2023). <https://doi.org/10.1038/s43017-022-00379-5>
- Lambert, F., Delmonte, B., Petit, J. R., Bigler, M., Kaufmann, P. R., Hutterli, A. M. A., Stocker, T. F., Ruth, U., Steffensen, J. P., and Maggi, V.: Dust-climate couplings over the past 800,000 years from the EPICA Dome C ice core, *Nature*, 452, 616–619, 2008.
- Lambert, F., Tagliabue, A., Shaffer, G., Lamy, F., Winckler, G., Farias, L., Gallardo, L., and De Pol-Holz, R.: Dust fluxes and iron fertilization in Holocene and Last Glacial Maximum climates, *Geophys Res Lett*, 42, 6014–6023, <https://doi.org/10.1002/2015GL064250>, 2015.
- Lee, L. A., Reddington, C. L., and Carslaw, K. S.: On the relationship between aerosol model uncertainty and radiative forcing uncertainty, *Proc Natl Acad Sci U S A*, 113, 5820–5827, <https://doi.org/10.1073/pnas.1507050113>, 2016.
- Lehner, F. and Coats, S.: Does Regional Hydroclimate Change Scale Linearly With Global Warming?, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2021GL095127>, 2021.
- Li, F., Val Martin, M., Andreae, M. O., Arneth, A., Hantson, S., Kaiser, J. W., Lasslop, G., Yue, C., Bachelet, D., Forrest, M., Kluzek, E., Liu, X., Mangeon, S., Melton, J. R., Ward, D. S., Darmanov, A., Hickler, T., Ichoku, C., Magi, B. I., Sitch, S., Van Der Werf, G. R., Wiedinmyer, C., and Rabin, S. S.: Historical (1700–2012) global multi-model estimates of the fire emissions from the Fire Modeling Intercomparison Project (FireMIP), *Atmos Chem Phys*, 19, 12545–12567, <https://doi.org/10.5194/acp-19-12545-2019>, 2019.
- Li, J., Carlson, B. E., Yung, Y. L., Lv, D., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, V., Kahn, R. A., Zhang, P., Dubovik, O., Ding, A., Lacis, A. A., Zhang, L., and Dong, Y.: Scattering and absorbing aerosols in the climate system, <https://doi.org/10.1038/s43017-022-00296-7>, 1 June 2022.
- Li, L., Mahowald, N. M., Miller, R. L., Pérez García-Pando, C., Klose, M., Hamilton, D. S., Gonçalves Ageitos, M., Ginoux, P., Balkanski, Y., Green, R. O., Kalashnikova, O., Kok, J. F., Obiso, V., Paynter, D., and Thompson, D. R.: Quantifying the range of the dust direct radiative effect due to source mineralogy uncertainty, *Atmos Chem Phys*, 21, 3973–4005, <https://doi.org/10.5194/acp-21-3973-2021>, 2021.

- Liu, P., Kaplan, J. O., Mickley, L. J., Li, Y., Chellman, N. J., Arienzo, M. M., Kodros, J. K., Pierce, J. R., Sigl, M., Freitag, J., Mulvaney, R., Curran, M. A. J., and McConnell, J. R.: Improved estimates of preindustrial biomass burning reduce the magnitude of aerosol climate forcing in the Southern Hemisphere, *Sci. Adv.*, 2021.
- 790 Maher, B., Prospero, J., Mackie, D., Gaiero, D., Hesse, P., and Balkanski, Y.: Global connections between aeolian dust, climate and ocean biogeochemistry at the present day and at the last glacial maximum, *Earth Sci Rev.*, 99, 61–97, 2010.
- Mahowald, N.: Anthropocene changes in desert area: sensitivity to climate model predictions, *Geophys Res Lett.*, 34, doi:10.1029/2007GL030472, <https://doi.org/10.1029/2007GL030472>, 2007.
- Mahowald, N.: Aerosol indirect effect on biogeochemical cycles and climate, *Science* (1979), 334, <https://doi.org/10.1126/science.1207374>, 2011.
- 795 Mahowald, N., Kohfeld, K., Hansson, M., Balkanski, Y., Harrison, S. P., Prentice, I. C., Schulz, M., and Rodhe, H.: Dust sources and deposition during the last glacial maximum and current climate: A comparison of model results with paleodata from ice cores and marine sediments, *Journal of Geophysical Research Atmospheres*, 104, <https://doi.org/https://doi.org/10.1029/1999JD900084>, 1999.
- 800 Mahowald, N., Ward, D. S., Kloster, S., Flanner, M. G., Heald, C. L., Heavens, N. G., Hess, P. G., Lamarque, J.-F., and Chuang, P. Y.: Aerosol impacts on climate and biogeochemistry, *Annu Rev Environ Resour.*, 36, <https://doi.org/10.1146/annurev-environ-042009-094507>, 2011a.
- Mahowald, N., Lindsay, K., Rothenberg, D., Doney, S. C., Moore, J. K., Thornton, P., Randerson, J. T., and Jones, C. D.: Desert dust and anthropogenic aerosol interactions in the Community Climate System Model coupled-carbon-climate model, *Biogeosciences*, 8, <https://doi.org/10.5194/bg-8-387-2011>, 2011b.
- 805 Mahowald, N. M., Zender, C. S., Luo, C., Savoie, D., Torres, O., Corral, J. D., and Del Corral, J.: Understanding the 30-year Barbados desert dust record, *Journal of Geophysical Research Atmospheres*, 107, <https://doi.org/10.1029/2002JD002097>, 2002.
- 810 Mahowald, N. M., Kloster, S., Engelstaedter, S., Moore, J. K., Mukhopadhyay, S., McConnell, J. R. R., Albani, S., Doney, S. C., Bhattacharya, A., Curran, M. A. J. A. J., Flanner, M. G. G., Hoffman, F. M. M., Lawrence, D. M. M., Lindsay, K., Mayewski, P. A. A., Neff, J., Rothenberg, D., Thomas, E., Thornton, P. E. E., and Zender, C. S. S.: Observed 20th century desert dust variability: impact on climate and biogeochemistry, *Atmos Chem Phys*, 10, 10875–10893, <https://doi.org/10.5194/acp-10-10875-2010>, 2010.
- Mahowald, N. M., Scanza, R., Brahney, J., Goodale, C. L., Hess, P. G., Moore, J. K., and Neff, J.: Aerosol Deposition Impacts on Land and Ocean Carbon Cycles, *Curr Clim Change Rep.*, 3, 16–31, <https://doi.org/10.1007/s40641-017-0056-z>, 2017.
- 815 Van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A. L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and Van Der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with

proxies and fire models (1750–2015), *Geosci Model Dev*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.

Marlon, J., Bartlein, P., Carcaillet, C., Gavin, D., Harrison, S., Higuera, P., Joos, F., Power, M., and Prentice, I. C.: Climate and human influences on global biomass burning over the past two millennia, *Nature-Geosciences*, 1, 697–701, <https://doi.org/https://doi.org/10.1029/PA005i001p00001>, 2008.

Marlon, J. R., Bartlein, P. J., Gavin, D. G., Long, C. J., Anderson, R. S., Briles, C. E., Brown, K. J., Colombaroli, D., Hallett, D. J., Power, M. J., Scharf, E. A., and Walsh, M. K.: Long-term perspective on wildfires in the western USA, *Proceedings of the National Academy of Sciences*, 109, E535–E543, <https://doi.org/10.1073/pnas.1112839109>, 2012.

Marlon, J. R., Kelly, R., Daniau, A. L., Vanni $\grave{e}$ re, B., Power, M. J., Bartlein, P., Higuera, P., Blarquez, O., Brewer, S., Br $\acute{e}$ cher, T., Feurdean, A., Romera, G. G., Iglesias, V., Yoshi Maezumi, S., Magi, B., Mustaphi, C. J. C., and Zhihai, T.: Reconstructions of biomass burning from sediment-charcoal records to improve data-model comparisons, *Biogeosciences*, 13, 3225–3244, <https://doi.org/10.5194/bg-13-3225-2016>, 2016.

Matsui, H., Hamilton, D. S., and Mahowald, N. M.: Black carbon radiative effects highly sensitive to emitted particle size when resolving mixing-state diversity, *Nat Commun*, 9, 3446, <https://doi.org/10.1038/s41467-018-05635-1>, 2018.

McConnell, J. R., Aristarain, A., Banta, J., Edwards, P., and Simoes, J.: 20th Century doubling in dust archived in an Antarctic Peninsula ice core parallels climate change and desertification in South America, *PNAS*, 104, 5743–5748, <https://doi.org/https://doi.org/10.1073/pnas.0607657104>, 2007a.

McConnell, J. R., Edwards, R., Kok, G., Flanner, M., Zender, C. S., Saltzman, E., Banta, J. R., Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th-century industrial black carbon emissions altered Arctic climate forcing, *Science* (1979), 317, 1381–1384, 2007b.

McCoy, S. L., McCoy, D. T., Wood, R., Regayre, L., Watson-Parris, D., Grosvenor, D., Mulcahy, J., Hu, Y., Bender, F., Field, P., Carslaw, K., and Gordon, H.: The hemispheric contrast in cloud microphysical properties constrains aerosol forcing., *Proceedings of the National Academy of Science*, 117, <https://doi.org/10.1073/pnas.1922502117/-/DCSupplemental>, 2020.

Mulitza, S., Heslop, D., Pittauerova, D., Fischer, H., Meyer, I., Stuut, J.-B., Zabel, M., Mollenhauer, G., Collins, J., Kuhnert, H., and Schulz, M.: Increase in African dust flux at the onset of commercial agriculture in the Sahel region., *Nature*, 466, doi:10.1038/nature09213, 226–228, 2010.

Murray, B. J., Carslaw, K. S., and Field, P. R.: Opinion: Cloud-phase climate feedback and the importance of ice-nucleating particles, *Atmos Chem Phys*, 21, 665–679, <https://doi.org/10.5194/acp-21-665-2021>, 2021.

Naik, V. , S., Szopa, S., Adhikary, B., P. Artaxo, T. Berntsen, W. D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler Scharr, Z. Klimont, H. Liao, N. Unger, and P. Zanis: Chapter 6: Short-Lived Climate Forcers, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 10–171, 2021.

- Neff, J. C., Ballantyne, A. P., Farmer, G. L., Mahowald, N. M., Conroy, J. L., Landry, C. C., Overpeck, J. T., Painter, T. H., Lawrence, C. R., and Reynolds, R. L.: Increasing eolian dust deposition in the western United States linked to human activity, *Nat Geosci*, 1, <https://doi.org/10.1038/ngeo133>, 2008.
- 855 Paulot, F., Ginoux, P., Cooke, W. F., Donner, L. J., Fan, S., Lin, M. Y., Mao, J., Naik, V., and Horowitz, L. W.: Sensitivity of nitrate aerosols to ammonia emissions and to nitrate chemistry: Implications for present and future nitrate optical depth, *Atmos Chem Phys*, 16, 1459–1477, <https://doi.org/10.5194/acp-16-1459-2016>, 2016.
- Penner, J., Dickinson, R., and O'Neill, C. A.: Penneretal1992.science.256.5062.1432, *Science* (1979), 256, 1432–1434, 1992.
- 860 Pincus, R., Forster, P. M., and Stevens, B.: The Radiative Forcing Model Intercomparison Project (RFMIP): experimental protocol for CMIP6, *Geosci Model Dev*, 9, 3447–3460, <https://doi.org/10.5194/gmd-9-3447-2016>, 2016.
- Power, M., Marlon, J., Ortiz, N., Bartlein, P., Harrison, S., Mayle, F., A. Ballouche, R. H. W. Bradshaw C. Carcaillet S. Mooney, P. I. Moreno, I. C. Prentice, C. C., K. Thonicke W. Tinner Y. Zhang, Y. Zhao, A. A. Ali, R. S. Anderson, R. Beer, C. W., H. Behling C. Briles A. Brunelle, M. Bush, P. Camill, G. Q. Chu, J. Clark, K. J. B., D. Colombaroli S. Connor M. Daniels, J. Dodson, E. Doughty, M. E. Edwards, A.-L. D., W. Finsinger D.
- 865 Foster M.-J. Gaillard, D. G. Gavin, E. Gobet, S. Haberle, J. F., D. J. Hallett P. Higuera S. Horn, J. Inoue, P. Kaltenrieder, L. Kennedy, Z. C. Kong, G. H., C. Larsen C. J. Long E. A. Lynch, M. McGlone, S. Meeks, S. Mensing, G. Meyer, J. L., T. Minckley J. Mohr J. New, R. Newnham, R. Noti, W. Oswald, J. Pierce, D. M. N., P. J. H. Richard C. Rowe B. N. Shuman, H. Takahara, J. Toney, M. F. S. G. i, C. Turney D. H. Urrego-Sanchez M. Vandergoes, B. Vanniere, E. Vescovi, C. U., and M. Walsh X. Wang J. Wilmshurst, J. H. Zhang, N. W.: Changes
- 870 in fire regimes since the Last Glacial Maximum: an assessment based on a global synthesis and analysis of charcoal data, *Clim Dyn*, DOI 10.1007/s00382-007-0334-x, 2007.
- Prospero, J. and Lamb, P.: African droughts and dust transport to the Caribbean: climate change implications, *Science* (1979), 302, 1024–1027, <https://doi.org/10.1126/science.1089915>, 2003.
- Pye, H., Liao, H., Wu, S., Mickley, L., Jacob, D., Henze, D., and Seinfeld, J.: Effect of changes in climate and emissions on
- 875 future sulfate-nitrate-ammonium aerosol levels in the United States, *J Geophys Res*, 114, doi:10.1029/2008JD010701, 2009.
- Rathod, S., Hamilton, D. S. D. S., N. Mahowald, Klimont, Z., Corbett, J. J., and Tami Bond: A mineralogy-based anthropogenic combustion-iron emission inventory, *Journal of Geophysical Research Atmospheres*, 125, <https://doi.org/10.1029/2019JD032114>, 2020.
- 880 Regayre, L. A., Johnson, J. S., Yoshioka, M., Pringle, K. J., Sexton, D. M. H., Booth, B. B. B., Lee, L. A., Bellouin, N., and Carslaw, K. S.: Aerosol and physical atmosphere model parameters are both important sources of uncertainty in aerosol ERF, *Atmos Chem Phys*, 18, 9975–10006, <https://doi.org/10.5194/acp-18-9975-2018>, 2018.
- Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., Hegerl, G., Klein, S. A., Marvel, K. D., Rohling, E. J., Watanabe, M., Andrews, T., Braconnot, P., Bretherton, C. S., Foster, G. L., Hausfather, Z., von
- 885 der Heydt, A. S., Knutti, R., Mauritsen, T., Norris, J. R., Proistosescu, C., Rugenstein, M., Schmidt, G. A., Tokarska,

- K. B., and Zelinka, M. D.: An Assessment of Earth's Climate Sensitivity Using Multiple Lines of Evidence, *Reviews of Geophysics*, 58, 1–92, <https://doi.org/10.1029/2019RG000678>, 2020.
- Smith, C. J., Harris, G. R., Palmer, M. D., Bellouin, N., Collins, W., Myhre, G., Schulz, M., Golaz, J. C., Ringer, M., Storelvmo, T., and Forster, P. M.: Energy Budget Constraints on the Time History of Aerosol Forcing and Climate Sensitivity, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2020JD033622>, 2021.
- Snider, G., Weagle, C. L., Murdymootoo, K. K., Ring, A., Ritchie, Y., Stone, E., Walsh, A., Akoshile, C., Anh, N. X., Balasubramanian, R., Brook, J., Qonitan, F. D., Dong, J., Griffith, D., He, K., Holben, B. N., Kahn, R., Lagrosas, N., Lestari, P., Ma, Z., Misra, A., Norford, L. K., Quel, E. J., Salam, A., Schichtel, B., Segev, L., Tripathi, S., Wang, C., Yu, C., Zhang, Q., Zhang, Y., Brauer, M., Cohen, A., Gibson, M. D., Liu, Y., Martins, J. V., Rudich, Y., and Martin, R. V.: Variation in global chemical composition of PM_{2.5}: emerging results from SPARTAN, *Atmos Chem Phys*, 16, 9629–9653, <https://doi.org/10.5194/acp-16-9629-2016>, 2016.
- Stanelle, T., Bey, I., Raddatz, T., Reick, C., and Tegen, I.: Anthropogenically induced changes in twentieth century mineral dust burden and the associated impact on radiative forcing, *J Geophys Res*, 119, 13,526–13,546, <https://doi.org/10.1002/2014JD022062>, 2014.
- Storelvmo, T.: Aerosol Effects on Climate via Mixed-Phase and Ice Clouds, *The Annual Review of Earth and Planetary Sciences* is online at earth.annualreviews.org, 45, 199–222, <https://doi.org/10.1146/annurev-earth-060115>, 2017.
- Szopa, S., V. Naik, B. Adhikary, P. Artaxo, T. Berntsen, W.D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler-Scharr, Z. Klimont, H. Liao, N. Unger, and P. Zanis, 2021: Short-Lived Climate Forcers. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 817–922, doi:10.1017/9781009157896.008.
- Tegen, I., Werner, M., Harrison, S. P., Kohfeld, K. E., Mahowald, N. M., Rivera Rivera, G. D., and Luo, C.: Relative importance of climate and land use in determining present and future global soil dust emission, *Geophys Res Lett*, 31, L05105, doi:10.1029/2003GL019216, 2004, <https://doi.org/10.1029/2004GL021272>, 2004.
- Textor, C. and others, and: Analysis and quantification of the diversities of aerosol life cycles within AeroCOM, *Atmos Chem Phys*, 6, 1777–1813, 2006.
- Thornhill, G., Collins, W., Olivié, D., Archibald, A., Bauer, S., Checa-Garcia, R., Fiedler, S., Folberth, G., Gjermundsen, A., Horowitz, L., Lamarque, J.-F., Michou, M., Mulcahy, J., Nabat, P., Naik, V., O'Connor, F., Paulot, F., Schulz, M., Scott, C., Seferian, R., Smith, C., Takemura, T., Tilmes, S., and Weber, J.: Climate-driven chemistry and aerosol feedbacks in CMIP6 Earth system models, *Atmos Chem Phys*, 1–36, <https://doi.org/10.5194/acp-2019-1207>, 2020.

- Thornhill, G., Collins, W., Olivié, D., B. Skeie, R., Archibald, A., Bauer, S., Checa-Garcia, R., Fiedler, S., Folberth, G.,
 920 Gjermundsen, A., Horowitz, L., Lamarque, J. F., Michou, M., Mulcahy, J., Nabat, P., Naik, V., M. O'Connor, F.,
 Paulot, F., Schulz, M., E. Scott, C., Séférian, R., Smith, C., Takemura, T., Tilmes, S., Tsigaridis, K., and Weber, J.:
 Climate-driven chemistry and aerosol feedbacks in CMIP6 Earth system models, *Atmos Chem Phys*, 21, 1105–1126,
<https://doi.org/10.5194/acp-21-1105-2021>, 2021.
- Turnock, S. T., Allen, R. J., Andrews, M., Bauer, S. E., Deushi, M., Emmons, L., Good, P., Horowitz, L., John, J. G., Michou,
 925 M., Nabat, P., Naik, V., Neubauer, D., O'Connor, F. M., Olivié, D., Oshima, N., Schulz, M., Sellar, A., Shim, S.,
 Takemura, T., Tilmes, S., Tsigaridis, K., Wu, T., and Zhang, J.: Historical and future changes in air pollutants from
 CMIP6 models, *Atmos Chem Phys*, 20, 14547–14579, <https://doi.org/10.5194/acp-20-14547-2020>, 2020.
- Undorf, S., Polson, D., Bollasina, M. A., Ming, Y., Schurer, A., and Hegerl, G. C.: Detectable Impact of Local and Remote
 Anthropogenic Aerosols on the 20th Century Changes of West African and South Asian Monsoon Precipitation,
 930 *Journal of Geophysical Research: Atmospheres*, 123, 4871–4889, <https://doi.org/10.1029/2017JD027711>, 2018.
- Unger, N., Bond, T., Wang, J. S., Koch, D. M., Menon, S., Shindell, D., and Bauer, S.: Attribution of climate forcing to
 economic sectors, *Proceedings of the National Academy of Science USA*, 107, 3382–3387, 2010.
- Wan, J. S., Hamilton, D. S., and Mahowald, N. M.: Importance of Uncertainties in the Spatial Distribution of Preindustrial
 Wildfires for Estimating Aerosol Radiative Forcing, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2020GL089758>,
 935 2021.
- Wang, Z., Chappellaz, J., Park, K., and Mak, J. E.: Large Variations in Southern, *Science* (1979), 205, 1663–1666,
<https://doi.org/10.1126/science.1197257>, 2010.
- Webb, N. P. and Pierre, C.: Quantifying Anthropogenic Dust Emissions, *Earths Future*, 6, 286–295,
<https://doi.org/10.1002/2017EF000766>, 2018.
- 940 van der Werf, G. R., Randerson, J., Collatz, G. J., Giglio, L., Kasibhatla, P., Arrellano, A., Olson, S., and Kasischke, E.:
 Continental-scale partitioning of fire emissions during the 1997 to 2001 El Nino/La Nina Period, *Science* (1979), 303,
 73–76, 2004.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F. : Interannual variability
 in global biomass burning emissions from 1997-2004, *Atmos Chem Phys*, 6, 3423–3441, doi: 10.5194/acp-6-3423–
 945 2006, 2006.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin,
 Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural
 and peat fires (1997-2009), *Atmos Chem Phys*, 10, 11707–11735, doi:10.5194/acp-10-11707–2010, 2010.
- van der Werf, G. R., Peters, W., van Leeuwen, T. T., and Giglio, L.: What could have caused pre-industrial biomass burning
 950 emissions to exceed current rates?, *Climate of the Past*, 9, 289–306, <https://doi.org/10.5194/cp-9-289-2013>, 2013.
- Van Marle M J E, Kloster S, Magi B I, Marlon J R, Daniau A L, Field R D, Arneth A, Forrest M, Hantson S, Kehrwald N M,
 Knorr W, Lasslop G, Li F, Mangeon S, Yue C, Kaiser J W and Van Der Werf G R 2017 Historic global biomass

burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015) *Geosci Model Dev* **10** 3329–57


955 Wu, J., Kong, S., Wu, F., Cheng, Y., Zheng, S., Yan, Q., Zheng, H., Yang, G., Zheng, M., Liu, D., Zhao, D., and Qi, S.: Estimating the open biomass burning emissions in central and eastern China from 2003 to 2015 based on satellite observation, *Atmos Chem Phys*, 18, 11623–11646, <https://doi.org/10.5194/acp-18-11623-2018>, 2018.

Zennaro, P., Kehrwald, N., McConnell, J. R., Schüpbach, S., Maselli, O. J., Marlon, J., Vallelonga, P., Leuenberger, D., Zangrando, R., Spolaor, A., Borrotti, M., Barbaro, E., Gambaro, A., and Barbante, C.: Fire in ice: Two millennia of
960 boreal forest fire history from the Greenland NEEM ice core, *Climate of the Past*, 10, 1905–1924, <https://doi.org/10.5194/cp-10-1905-2014>, 2014.

Zhao, A., Ryder, C. L., and Wilcox, L. J.: How well do the CMIP6 models simulate dust aerosols?, *Atmos Chem Phys*, 22, 2095–2119, <https://doi.org/10.5194/acp-22-2095-2022>, 2022.

Zhu, J., Penner, J. E., Yu, F., Sillman, S., Andreae, M. O., and Coe, H.: Decrease in radiative forcing by organic aerosol
965 nucleation, climate, and land use change, *Nat Commun*, 10, <https://doi.org/10.1038/s41467-019-08407-7>, 2019.

Figure captions



970 Figure 1. Annual global mean dust aerosol loading from the observationally based reconstruction (Kok et al., 2023) in black, (grey shading represents the 90% confidence interval), compared against 10-year running means from historical runs of Climate Model Intercomparison Project (CMIP6) ensemble members. Reproduced with permission from (Kok et al., 2023).





Figure 2: Observed and modeled present day to preindustrial ratios (PD/PI) for black carbon in ice cores using different open fire emissions. Ice cores sites are from Greenland (open square and plus signs), Wyoming (diamond) and France (x's) for 4 different model simulations: AEROCOM (purple), CMIP6 (blue), SIMFIRE-BLAZE (yellow) and LMfire (red) are taken from (Hamilton et al., 2018). Ice core sites from Bolivia (solid circle) and Antarctica (solid square) using CMIP6 (blue) and LMfire (red) are taken from (Liu et al., 2021). The solid black line shows the 1:1 line.

Full uncertainty of aerosol changes requires multiple emission pathways and comparisons to observations



980 Figure 3: Schematic of the sources of uncertainties in aerosol radiative effects, from emission models to modeled concentration changes to modeled direct and aerosol-cloud radiative effects. The CMIP6 unconstrained uncertainties using a single emission scenario have a 90% confidence interval range of 2.8 W/m^2 (Bellouin et al., 2020; Sherwood et al., 2020). For the uncertainty summed over different aerosol species (Σ) using different emission scenarios for the past climate, the 90% confidence interval ranges from wildfires of 2.8 W/m^2 (Hamilton et al., 2018; Wan et al., 2021) is added in quadrature to uncertainties from dust of 0.4 W/m^2 (Kok et al., 2023) and an estimate of industrial emission uncertainties (assuming 10% error) of 0.2 W/m^2 , obtaining a 2.8 W/m^2 uncertainty in emissions.




Figure 4: Estimates of the unconstrained aerosol radiative forcing and its uncertainties for (a) different paleoclimate time periods relative to preindustrial (PI; 1850) and (b) present day (PD) relative to 1850 based on CMIP6 model spread (schematic based on (Smith et al., 2021) in blue) and including the emission uncertainties (90% confidence intervals) from wildfires, dust and anthropogenic aerosols as described in Fig. 3 (green), using the time series for wildfires from (Marlon et al., 2008). The left vertical axis represents the present day minus preindustrial radiative forcing following (Smith et al., 2021; Sherwood et al., 2020), and the right axis adds in the emission uncertainties for the preindustrial (from Fig. 3; Hamilton et al., 2018), shifting the preindustrial baseline of aerosol radiative effects (black arrow). Notice that the size of the black arrow and shift in the preindustrial state is not known, and this is a schematic to illustrate how the uncertainties in emissions in the preindustrial impact understanding of the radiative forcing.




Figure. 5. Variability of biomass burning rates over the last centuries based on a worldwide compilation of charcoal records (Marlon et al., 2008), CO mixing ratios from fires using CO concentration measurements at the South Pole (SPO), its isotopic signature, and a mass balance model (Wang et al., 2010) and a similar approach but based on CH₄ (Ferretti et al., 2005). The CO ice core data ended in 1897 but were extended (dashed line) by Wang et al. (2010) to present-day using firn samples (1968 and 1986) as well as modelling (year 2000). Shaded areas indicate reported uncertainty. Note that the datasets have different footprints and that absolute values cannot be compared directly. Reproduced with permission from (van der Werf et al., 2013) under CCC3.0.




Figure 6: Relative size of paleoclimate and historical changes in aerosols. A) Based on z scores from charcoal records, the variability across preindustrial time periods (green), present day (blue) and last glacial maximum is shown in global, northern extratropics, tropics, and southern extratropics based on data from (Marlon et al., 2008, 2016). Charcoal reconstructions use z-scores, which are normalized by the mean value at a site, divided by the variability, and thus a -2 z-score for LGM suggests significantly lower charcoal amounts. B) global dust changes ratio of deposition between present day and preindustrial (blue oval; Mahowald et al., 2010; Kok et al., 2023), and for the last glacial maximum relative to preindustrial (gold oval) (Mahowald et al., 1999; Albani et al., 2014, 2018; Lambert et al., 2015).

Table 1. Estimates of aerosol radiative forcing uncertainty, symbols, and citations. These represent the 90% confidence intervals.

Aerosol radiative forcing uncertainty term	Symbol	Estimate (range of 90% confidence interval)	Source
Unconstrained process uncertainty: Present day to preindustrial unconstrained model spread with fixed emissions from CMIP6	$\Sigma_{Process}^{PD-PI}$	2.8 W/m ²	(Bellouin et al., 2020; Sherwood et al., 2020)
Unconstrained emission uncertainty: Present day to preindustrial uncertainty in emission changes unconstrained	Σ_{Emis}^{PD-PI}	2.8 W/m ²	Equation 1 and. (Hamilton et al., 2018; Wan et al., 2021; Kok et al., 2023)
Total unconstrained uncertainty: Present day to preindustrial	Σ_{Total}^{PD-PI}	4.0 W/m ²	Equation 2
Constrained process uncertainty: Present day to preindustrial constrained with	$\Sigma_{Const_Process}^{PD-PI}$	1.6 W/m ²	(Bellouin et al., 2020)

observations using CMIP6 emissions			
Unconstrained emission uncertainty: preindustrial time period	Σ_{Emis}^{PI}	2.8 W/m ²	PI uncertainties in emission drive uncertainties in PD-PI
Unconstrained process uncertainty: Paleotime T to preindustrial	$\Sigma_{Process}^{T-PI}$	2.8 W/m ²	Assume same as PD to PI
Unconstrained emission uncertainty: Paleotime T to preindustrial uncertainty	Σ_{Emis}^{T-PI}	4.0 W/m ²	Equation 4
Unconstrained total uncertainty: Paleotime T to preindustrial uncertainty	Σ_{Total}^{T-PI}	4.8 W/m ²	Equation 3, using Σ_{Emis}^{T-PI} and $\Sigma_{Process}^{T-PI}$