

# Evaluating manual versus automated benthic foraminiferal $\delta^{18}\text{O}$ alignment techniques for developing chronostratigraphies in marine sediment records

5 Jennifer L. Middleton<sup>1</sup>, Julia Gottschalk<sup>2</sup>, Gisela Winckler<sup>1,3</sup>, Jean Hanley<sup>1</sup>, Carol Knudson<sup>1</sup>, Jesse R. Farmer<sup>4</sup>, Frank Lamy<sup>5</sup>, Lorraine E. Lisiecki<sup>6</sup>, and Expedition 383 Scientists<sup>7\*</sup>

<sup>1</sup>Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA

<sup>2</sup>Institute of Geosciences, Kiel University, Kiel, Germany

<sup>3</sup>Department of Earth and Environmental Sciences, Columbia University, New York, NY, USA

10 <sup>4</sup>School for the Environment, University of Massachusetts Boston, Boston, MA, USA

<sup>5</sup>Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research, Bremerhaven, Germany

<sup>6</sup>Department of Earth Science, University of California, Santa Barbara, CA, USA

<sup>7</sup>International Ocean Discovery Program, Texas A&M University, College Station, TX, USA

\*A full list of authors appears at the end of the paper.

15 *Correspondence to:* Jennifer L. Middleton ([jennym@ldeo.columbia.edu](mailto:jennym@ldeo.columbia.edu))

25 **Abstract.** Paleoceanographic interpretations of Plio-Pleistocene climate variability over the past 5 million years rely on the evaluation of event timing of proxy changes in sparse records across multiple ocean basins. In turn, orbital-scale chronostratigraphic controls for these records are often built from stratigraphic alignment of benthic foraminiferal stable oxygen isotope ( $\delta^{18}\text{O}$ ) records to a preferred dated target stack or composite. This chronostratigraphic age model approach yields age model uncertainties  
30 associated with alignment method, target selection, the assumption that the undated record and target experienced synchronous changes in benthic foraminiferal  $\delta^{18}\text{O}$  values, and the assumption that any possible stratigraphic discontinuities within the undated record have been appropriately identified. However, these age model uncertainties and their impact on paleoceanographic interpretations are seldom reported or discussed. Here, we investigate and discuss these uncertainties for conventional manual and  
35 automated tuning techniques based on benthic foraminiferal  $\delta^{18}\text{O}$  records and evaluate their impact on sedimentary age models over the past 3.5 Myr using three sedimentary benthic foraminiferal  $\delta^{18}\text{O}$  records as case studies. In one case study, we present a new benthic foraminiferal  $\delta^{18}\text{O}$  record for International  
40 Ocean Discovery Program (IODP) Site U1541 (54°13'S, 125°25'W), recently recovered from the South Pacific on IODP Expedition 383. The other two case studies examine published benthic foraminiferal  $\delta^{18}\text{O}$  records of Ocean Drilling Program (ODP) Site 1090 and the ODP Site 980/981 composite. Our analysis suggests average age uncertainties of 3 to 5 kyr associated with manually-derived versus automated alignment, 1 to 3 kyr associated with automated probabilistic alignment itself, and 2 to 6 kyr associated with the choice of tuning target. Age uncertainties are higher near stratigraphic segment ends and where local benthic foraminiferal  $\delta^{18}\text{O}$  stratigraphy differs from the tuning target. We conclude with  
45 recommendations for community best practices for the development and characterization of age uncertainty of sediment core chronostratigraphies based on benthic foraminiferal  $\delta^{18}\text{O}$  records.

## 1 Introduction

50 Stratigraphic alignment, or tuning, of benthic foraminiferal stable oxygen isotope ( $\delta^{18}\text{O}$ ) data to an independently dated target record, such as the LR04 benthic stack (Lisiecki and Raymo, 2005) or the Cenozoic Global Reference benthic foraminiferal carbon and oxygen Isotope Dataset (CENOGGRID; Westerhold et al., 2020), has been used to develop orbital-scale age models for carbonate-bearing marine sediment cores for decades. This practice arose from early work to cross-link marine sedimentary records  
55 with sparse absolute age control using stratigraphic similarities among records that are likely mechanistically linked (e.g., Imbrie et al., 1984; Martinson et al., 1987; Shackleton et al., 1990). Absolute ages for these early targets were mostly inferred from well constrained orbital variations in Earth's radiative forcing (orbital tuning), while later targets have also been dated using astronomically forced global ice volume models (Imbrie et al., 1984; Lisiecki and Raymo, 2005; Ahn et al., 2017; Westerhold  
60 et al., 2020). When available, paleomagnetic reversals and radiometric dates have been used to provide additional absolute age control on these target records (e.g., Mankinen and Dalrymple, 1979; Stein et al., 1993; Thompson and Goldstein, 2006). Benthic foraminiferal  $\delta^{18}\text{O}$ -derived age models of marine sediment cores are widely relied upon in the interpretation of Plio-Pleistocene changes in climate, ocean circulation, and marine biogeochemistry (e.g., Shackleton and Hall, 1984; Ravelo et al., 2004; Elderfield  
65 et al., 2012; Hodell et al., 2023). However, benthic foraminiferal  $\delta^{18}\text{O}$ -derived age models are often presented without discussion of the impact of the alignment approach on assigned ages and without estimates of age model uncertainty arising from alignment method, uncertainties within the tuning target, or appropriateness of the chosen target for the specific study region.

70 Benthic foraminiferal  $\delta^{18}\text{O}$  stratigraphy is based on the assumption that benthic foraminifera record global variations in bottom water  $\delta^{18}\text{O}$  caused mainly by climatic changes in temperature, salinity, and global ice volume over orbital timescales (Pisias et al., 1984; Lisiecki and Raymo, 2005; Skinner and Shackleton, 2004). Benthic foraminifera of the genus *Uvigerina* are considered an ideal benthic foraminiferal species for the generation of  $\delta^{18}\text{O}$  stratigraphies because they are believed to calcify in equilibrium with seawater

75  $\delta^{18}\text{O}$ , although it occupies a shallow infaunal habitat (Shackleton, 1974). Other widely used species for these efforts include epibenthic foraminifera of the genus *Cibicidoides* or *Cibicides*, whose stable oxygen isotope composition is generally corrected by +0.64 ‰ to match presumable equilibrium seawater  $\delta^{18}\text{O}$  values (Shackleton and Opdyke, 1973). More recent studies have found disequilibrium effects between the  $\delta^{18}\text{O}$  values of *Uvigerina* and *Cibicidoides* species that range between 0.47 ‰ (Marchitto et al., 2014) 80 and 0.73 ‰ (Jöhnck et al., 2012), depending on local bottom water and pore water pH conditions. In addition, regional and temporal variations in bottom water temperature can introduce further discrepancies between the  $\delta^{18}\text{O}$  values of benthic foraminifera and the bottom water (e.g., Marchitto et al., 2014; Elderfield et al., 2012). However, the premise of benthic  $\delta^{18}\text{O}$  stratigraphy hinges on the representation of bottom water  $\delta^{18}\text{O}$  by the  $\delta^{18}\text{O}$  of benthic foraminiferal species of the genus *Uvigerina* 85 (such as *U. peregrina* and *U. hispida*) and *Cibicidoides* (such as *C. wuellerstorfi* and *C. kullenbergi*), with a constant correction factor between the two (Lisiecki and Raymo, 2005). In other words, benthic  $\delta^{18}\text{O}$  stratigraphy assumes the effect of bottom water and temperature variations in space and time to be minimal (Lisiecki and Raymo, 2005). Once a suitable benthic foraminiferal  $\delta^{18}\text{O}$  record has been generated, the undated record can be aligned to a dated benthic foraminiferal  $\delta^{18}\text{O}$  tuning target under the 90 assumptions that the new record is stratigraphically continuous and that deep ocean  $\delta^{18}\text{O}$  changes occur synchronously over glacial-interglacial timescales.

Each tuning target carries its own sampling biases and independent age uncertainties that impact the ages assigned to new records using benthic foraminiferal  $\delta^{18}\text{O}$ -based alignments. Community-preferred targets 95 for stratigraphic benthic foraminiferal  $\delta^{18}\text{O}$  alignment have continuously evolved over the past decades with the increasing number and spatial coverage of available benthic foraminiferal  $\delta^{18}\text{O}$  records (e.g., Imbrie et al., 1984; Shackleton et al., 1995; Huybers and Wunsch, 2004; Lisiecki and Raymo, 2005; Ahn et al., 2017; Westerhold et al., 2020). Early targets, such as SPECMAP, resulting from the SPECtral Mapping Project, were built by stacking the planktonic foraminiferal  $\delta^{18}\text{O}$  records of five or fewer 100 sediment cores and were typically limited to within the last 800 kyr of the late Pleistocene (Imbrie et al., 1984; Pisias et al., 1984; Martinson et al., 1987). These early targets were further restricted in their ability to reflect global changes in deep seawater  $\delta^{18}\text{O}$  because they may have been biased by surface water  $\delta^{18}\text{O}$

variations due to the inclusion of planktonic foraminiferal  $\delta^{18}\text{O}$  records. Later targets, such as LR04 (Lisiecki and Raymo, 2005) and Prob-stack (Ahn et al., 2017), improved upon these approaches by 105 stacking benthic foraminiferal  $\delta^{18}\text{O}$  records from 57 and 180 sediment cores around the world, respectively, and extending the interval covered by the benthic foraminiferal stack to  $\sim$ 5 Ma. Composite splice targets built by combining discrete high resolution benthic foraminiferal  $\delta^{18}\text{O}$  records in series, such as the S95 composite (Shackleton, 1995), the benthic  $\delta^{18}\text{O}$  megasplice (De Vleeschouwer et al., 2017), and the CENOGRID splice (Westerhold et al., 2020), have also been generated and provide a 110 means to examine the high-frequency variability within benthic foraminiferal  $\delta^{18}\text{O}$  data that may be smoothed over or lost in the generation of a global stack. On the other hand, such composite targets are heavily influenced by regional bottom water  $\delta^{18}\text{O}$  and other variability at the limited number of sites included in the record.

115 Increasing data coverage and the recovery of more core sites across the globe have revealed regional differences in the timing of benthic foraminiferal  $\delta^{18}\text{O}$  changes across different ocean basins and water masses (Lisiecki and Raymo, 2009; Skinner and Shackleton, 2005; Stern and Lisiecki, 2014; Waelbroeck et al., 2011; Labeyrie et al., 2005), which challenge one of the basic assumptions of benthic foraminiferal  $\delta^{18}\text{O}$  stratigraphies. Early work, based on two independently dated sediment cores from the Pacific and 120 Atlantic Oceans, identified a  $\sim$ 4 kyr lag in the earliest signal of glacial termination between the deep North Atlantic and the deep equatorial Pacific (Skinner and Shackleton, 2005). Subsequent development of Pacific- and Atlantic Ocean-specific benthic foraminiferal  $\delta^{18}\text{O}$  stacks (i.e., the LR09 stacks) confirmed these results and suggested an average lag of 1.6 kyr in deglacial benthic seawater  $\delta^{18}\text{O}$  changes in the Pacific relative to the Atlantic (Lisiecki and Raymo, 2009). More recently, the generation of seven 125 regional stacks over the last glacial cycle from 252 sediment cores has further refined our understanding of regional asynchronicity in the timing of the benthic foraminiferal  $\delta^{18}\text{O}$  signal (Stern and Lisiecki, 2014). Stern and Lisiecki (2014) find a maximum lag of 4 kyr between the earliest signals of glacial termination detected in the intermediate South Atlantic and the latest signals detected in the deep Indian Ocean and shorter lags occurring between other water masses and regions. If left unconsidered, such 130 regional differences in the timing of benthic foraminiferal  $\delta^{18}\text{O}$  changes relative to a global tuning target

can introduce systematic age uncertainties into the resulting benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphy.

Benthic foraminiferal  $\delta^{18}\text{O}$  records are traditionally aligned manually to the chosen target via user-defined identification and assignment of characteristic  $\delta^{18}\text{O}$  signals at specific tie points, with the assumption of a constant sedimentation rate between selected tie points. Tie points, or age markers, are typically assigned at mid-points of sharp transitions within the benthic foraminiferal  $\delta^{18}\text{O}$  record because the mid-point depth is easy to identify. However, tie points are also frequently assigned to local maxima and minima within the benthic foraminiferal  $\delta^{18}\text{O}$  record, where depth uncertainties may be higher due to limitations of sampling resolution, and no community-accepted guidelines exist for the establishment of benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies by manual tuning. Hence, the number of tie points identified and their exact location within each benthic foraminiferal  $\delta^{18}\text{O}$  record can vary among users, which may significantly impact the final age model of the sediment core. This user-induced variance is particularly significant in core intervals where the alignment of the benthic foraminiferal  $\delta^{18}\text{O}$  record to the tuning target is not straightforward. Further, conventional manual stratigraphic alignment efforts based on benthic foraminiferal  $\delta^{18}\text{O}$  records generally lack age uncertainty estimates, although in rare cases ad-hoc uncertainties are assigned to individual tie points. Consequently, benthic foraminiferal  $\delta^{18}\text{O}$ -derived age models are a useful, but subjective, dating tool for establishing age models of marine sedimentary records.


150

While alignment software such as AnalySeries (Paillard et al., 1996) compute quantitative correlation coefficients for user-defined alignments, such tools do not eliminate the subjectivity and related uncertainties inherent in the user-defined visual alignment approach and do not provide estimates of chronostratigraphic age uncertainties. In contrast, a probabilistic alignment algorithm like the Matlab-based Hidden Markov Model (HMM)-Match (Lin et al., 2014) largely eliminates the subjectivity resulting from reliance on multiple user-defined tie points. In fact, such algorithms can generate user-independent age estimates for every data point within an input benthic foraminiferal  $\delta^{18}\text{O}$  record without presuming constant sedimentation rates across periods of major climatic change or other predefined intervals. HMM-

Match additionally provides statistically informed estimates of alignment-derived age uncertainty based  
160 on reported benthic foraminiferal  $\delta^{18}\text{O}$  variance of the tuning target and based on differences in the benthic  
foraminiferal  $\delta^{18}\text{O}$  tuning target and the input  $\delta^{18}\text{O}$  record (Lin et al., 2014; Ahn et al., 2017).  
Consequently, HMM-Match has become an increasingly popular tool for the development of Plio-  
165 Pleistocene chronostratigraphies in marine sediment cores based on benthic foraminiferal  $\delta^{18}\text{O}$  (e.g.,  
Hodell et al., 2023; Ford and Raymo, 2020). However, as with the manual alignment approaches,  
170 alignment algorithms like HMM-Match (Lin et al., 2014) do not currently account for the age model  
uncertainties associated with regional offsets between the timing of benthic foraminiferal  $\delta^{18}\text{O}$  changes  
at a given location compared to globally averaged data set.

Here, we assess the uncertainties associated with each step of the stratigraphic alignment of benthic  
170 foraminiferal  $\delta^{18}\text{O}$  records and evaluate their impact on sediment core age models over the past 3.5 Myr  
using three example sites from the International Ocean Discovery Program (IODP) and its predecessor  
Ocean Drilling Program (ODP; Fig. 1). As part of this analysis, we present a new benthic foraminiferal  
 $\delta^{18}\text{O}$  record from IODP Site U1541 in the central South Pacific Ocean, recovered on IODP Expedition  
383, that is based on  $\sim$ 1600 benthic foraminiferal  $\delta^{18}\text{O}$  measurements over the last 3.5 Myr. We  
175 additionally examine ODP Site 1090 in the South Atlantic and ODP Site 980/981 composite in the North  
Atlantic using published benthic foraminiferal  $\delta^{18}\text{O}$  records (Venz and Hodell, 2002; Hodell et al., 2000;  
Oppo et al., 1998; Mcmanus et al., 1999; Flower et al., 2000; Raymo et al., 2004). We assess the impact  
of stratigraphic alignment approach (manual versus automated probabilistic methods) on age model  
output by independently aligning the benthic foraminiferal  $\delta^{18}\text{O}$  records of each example site to the LR04  
180 stack using first the conventional manual alignment method and then the open-source HMM-Match  
probabilistic alignment algorithm of Lin et al. (2014). We evaluate the impact of stratigraphic target  
choice on age model output by aligning the benthic foraminiferal  $\delta^{18}\text{O}$  records of each example site to a  
suite of reference targets (LR04, LR09, Prob-stack, and the CENOGRID splice) using the HMM-Match  
alignment algorithm (Lin et al., 2014). We additionally assess the sensitivity of such age models to the  
185 assumptions of global synchronicity in seawater  $\delta^{18}\text{O}$  changes and stratigraphic continuity within the

undated record. Finally, we present suggested best practices for the development and characterization of age uncertainty of sediment core age models based on benthic foraminiferal  $\delta^{18}\text{O}$  records.



**Figure 1: Location map for sediment cores included in the LR04 (smaller black circles; Lisiecki and Raymo, 2005) and Prob-stack global benthic foraminiferal stable oxygen isotope ( $\delta^{18}\text{O}$ ) tuning target stacks (larger open circles; Ahn et al., 2017). International Ocean Discovery Program (IODP) Site U1541, Ocean Drilling Program (ODP) Site 1090, and ODP Site 980/981 are denoted by stars. Color bar indicates the maximum age of the benthic foraminiferal  $\delta^{18}\text{O}$  records from each site included in the LR04 and Prob-stack. Four benthic foraminiferal  $\delta^{18}\text{O}$  records have significant data gaps and are numbered as follows: 1) MV0502-4JC data cover the intervals 2975-1575 ka, 679-441 ka and 210-23 ka; 2) ODP Site 1014 data cover the last 187 ka and 916-386 ka; 3) ODP Site 1241 data cover the intervals 5691-2443 ka and 2129-0 ka; 4) IODP Site U1313 data covers the intervals 3331-2414 ka, 913-788 ka, 700-600 ka and 356-549 ka were considered.**

## 2 Sampling and Measurements

### 190 2.1 Sediment Cores

190 IODP Site U1541 ( $54^{\circ}13' S$ ,  $125^{\circ}25' W$ , 3604 m water depth) was recovered on the western flank of the East Pacific Rise, ~300 km from the ridge axis, in the Subantarctic Pacific Ocean on IODP Expedition 383 (Fig. 1; Winckler et al., 2021). IODP Site U1541 is currently bathed in Lower Circumpolar Deep Water (LCDW) and represents one of the first continuous sedimentary sections from the Subantarctic 195 Pacific that spans the Plio-Pleistocene with orbital-scale resolution (Lamy et al., 2019; Winckler et al., 2021). As such, sediments from IODP Site U1541 can be used to investigate the dynamic atmospheric, oceanographic, and Antarctic processes impacting the large Pacific sector of the Southern Ocean across the wide range of climatic variability of the past 5 Myr and more. We present new benthic foraminiferal  $\delta^{18}O$  data from IODP Site U1541 for the past 3.5 Myr and the age model results of our stratigraphic 200 alignment analyses represent the first orbital-scale U1541 age models based on benthic foraminiferal  $\delta^{18}O$  tuning available for this site. Due to its location in the sparsely studied but climatically important South Pacific (Fig. 1), we use IODP Site U1541 as an end-member case study to investigate the age uncertainties associated with benthic foraminiferal  $\delta^{18}O$  stratigraphic alignment of a record from an under-sampled study region (i.e., the Pacific Ocean) to common alignment targets.

205

ODP Site 1090 ( $42^{\circ}55' S$ ,  $8^{\circ}54'E$ , 3702 m water depth) was recovered from the southern flank of the Agulhas Ridge in the Subantarctic Atlantic Ocean on ODP Leg 177 (Fig. 1; Gersonde et al., 1999). ODP Site 1090 is also bathed by LCDW, though it is positioned close to the lower boundary of North Atlantic Deep Water (NADW). ~~Among the many insights derived from ODP Site 1090, paleoceanographic reconstructions at this site provide critical constraints on changes in deep water circulation, the migration of the polar frontal system, and dust-borne iron delivery to the Subantarctic Atlantic across the Plio-Pleistocene (Venz and Hodell, 2002; Diekmann and Kuhn, 2002; Martínez-García et al., 2011).~~ The original stratigraphic alignments of the benthic foraminiferal  $\delta^{18}O$  records for ODP Site 1090 and its pre-site survey core, TTN057-6, were manually determined using the benthic foraminiferal  $\delta^{18}O$  record of 210 Deep Sea Drilling Project (DSDP) Site 607 as an alignment target (Venz and Hodell, 2002; Hodell et al., 2000). Sediment chronology of the DSDP Site 607 record itself was determined via orbital tuning of its

benthic foraminiferal  $\delta^{18}\text{O}$  record between paleomagnetic tie points (Raymo et al., 1990; Ruddiman et al., 1989; Raymo et al., 1989). We use ODP Site 1090 as a case study to investigate the age uncertainties inherent to the choice of alignment technique for a site in the Atlantic Ocean that is well represented by 220 common tuning targets and to address the impact of stratigraphic discontinuity on the resulting benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphy.

ODP Site 980 (55°29' N, 14°42' W, 2169 m water depth) and ODP Site 981 (55°29' N, 14°39' W, 2173 m water depth) were recovered from the Feni Drift in the Northeast Atlantic on ODP Leg 162 (Fig. 1; 225 Jansen et al., 1996). ODP Sites 980 and 981 are currently bathed in lower NADW. ~~The ODP Site 980/981 records have been valuable for investigating orbital and millennial scale climate and ocean circulation changes in the North Atlantic and their relationship with northern hemisphere ice sheets, as well as changes in the geomagnetic field, over the mid to late Pleistocene (Raymo et al., 2004; McManus et al., 1999; Channell and Raymo, 2003; Marino et al., 2011).~~ The original stratigraphic alignments for the 230 combined ODP Site 980/981 benthic foraminiferal  $\delta^{18}\text{O}$  records, spliced together at ~860 ka (Raymo et al., 2004), were manually determined using the Shackleton et al. (1990) target chronology for ODP Site 677 (Oppo et al., 1998; McManus et al., 1999; Flower et al., 2000; Raymo et al., 2004). The sediment 235 chronology of ODP Site 677 was determined via orbital tuning of its benthic foraminiferal  $\delta^{18}\text{O}$  record following the calculations of Berger (1989) and Berger and Loutre (1988) under the assumption of a constant phase relationship between astronomical radiative forcing and climatic response (Shackleton et al., 1990). Due to the high-resolution benthic foraminiferal  $\delta^{18}\text{O}$  record available for this site and its 240 location in the densely sampled North Atlantic, we use ODP Site 980/981 as a case study with maximum influence of NADW. This site also enables investigation into the age uncertainties associated with benthic foraminiferal  $\delta^{18}\text{O}$  stratigraphies under favorable alignment conditions, where regional seawater temperature and  $\delta^{18}\text{O}$  differences and associated temporal offsets in benthic foraminiferal  $\delta^{18}\text{O}$  changes between the undated record and the global LR04 and Prob-stack targets, which are heavily weighted by North Atlantic sampling, are expected to be minimal.

## 2.2 Benthic foraminiferal stable oxygen isotope analyses

The benthic foraminiferal  $\delta^{18}\text{O}$  data from IODP Site U1541 were measured in the upper 85 m core composite depth below seafloor (CCSF) of the U1541 shipboard splice (with identification code: CCSF\_383\_U1541\_ABC\_20190624; Winckler et al., 2021) at a 5 cm sampling resolution. This sampling resolution yields an average temporal resolution of  $\sim$ 2 kyr per sample over the 3.5 Myr record examined. These data were combined with benthic foraminiferal  $\delta^{18}\text{O}$  data spanning the last four glacial cycles from PS75/059-2 (Ullermann et al., 2016), which is the pre-site survey cored of IODP Site U1541. To combine the benthic foraminiferal  $\delta^{18}\text{O}$  records from both cores, the PS75/059-2 data were mapped onto a common U1541 depth scale via a manual stratigraphic alignment of high-resolution X-ray fluorescence (XRF) Fe intensity variations in PS75/059-2 (Lamy et al., 2014) and U1541 that result in 22 tie points (Supplementary Fig. S1 and Supplementary Table S1).

Bulk sediment samples from IODP Site U1541 were freeze-dried, sieved with deionized water over a 150  $\mu\text{m}$  sieve, and dried at  $\sim$ 45°C to prepared for benthic foraminiferal  $\delta^{18}\text{O}$  measurements. The  $>150\ \mu\text{m}$  sample fraction was then examined for benthic foraminifera under a stereomicroscope. One to five tests of benthic foraminifera of the genera *Cibicidoides*, *Cibicides*, and *Uvigerina*, following the morphotype description of Gottschalk et al. (2016) for *Cibicidoides* and *Cibicides*, were selected for each  $\delta^{18}\text{O}$  measurement with a wet brush. The samples were measured on a Thermo Scientific Delta V+ isotope ratio mass spectrometer with an automated Kiel IV Carbonate Device at Lamont-Doherty Earth Observatory (LDEO), Columbia University, USA. The data were corrected relative to the NBS (National Bureau of Standards)-19 calcite standard, which was analyzed every 10 samples, and are reported in  $\delta$ -notation with respect to the Vienna PeeDeeBee (VPDB) international standard. The long-term standard deviation for  $\delta^{18}\text{O}$  of the NBS-19 standard on this instrument is 0.06‰ VPDB. The mean offset between *Uvigerina* spp. and *Cib.* spp. at IODP Site U1541 is  $0.49 \pm 0.19\text{‰}$  VPDB ( $n=87$ ). We therefore corrected our *Cibicidoides/Cibicides* spp./sp.  $\delta^{18}\text{O}$  record for  $\delta^{18}\text{O}$ -disequilibrium effects according to the measured mean offset. Intra-species and intra-morphotype *Cibicidoides/Cibicides* spp./sp.  $\delta^{18}\text{O}$  variability at IODP Site U1541 is similar to the analytical uncertainty of the  $\delta^{18}\text{O}$  analyses ( $\sigma < 0.06\text{‰}$  VPDB).


We use previously published benthic foraminiferal  $\delta^{18}\text{O}$  data for ODP Site 1090 (Venz and Hodell, 2002; Hodell et al., 2000). The data from ODP Site 1090 were measured on one to three specimens per sample of the species *Cibicidoides wuellerstorfi* using a Finnigan MAT 252 mass spectrometer with an estimated  $1\sigma$  analytical uncertainty of  $\pm 0.06\text{\textperthousand}$  VPDB based on the reproducibility of repeat measurements of the 275 NBS-19 internal standard ( $n = 321$ ; Venz and Hodell, 2002). The benthic foraminiferal  $\delta^{18}\text{O}$  measurements of the ODP Site 1090 survey core, TTN057-6-PC4, were made on *Cibicidoides wuellerstorfi* and *C. kullenbergi* and carry an estimated  $1\sigma$  analytical uncertainty of  $< 0.10\text{\textperthousand}$  VPDB (Hodell et al., 2000). The average sampling resolution for the combined benthic foraminiferal  $\delta^{18}\text{O}$  record of ODP Site 1090 and TTN057-6-PC4, spliced together at  $\sim 400$  ka, is  $\sim 3$  kyr over the 2.9 Myr interval 280 examined (Venz and Hodell, 2002; Hodell et al., 2000). Where repeat analyses or measurements from both species were available, we employ the average foraminiferal  $\delta^{18}\text{O}$  value at each depth for our alignment analyses.

We use previously published benthic foraminiferal  $\delta^{18}\text{O}$  data for ODP Site 980/981. The data from ODO 285 Site 980 were measured on one to three specimens per sample of *C. wuellerstorfi* or *C. kullenbergi* on a Finnigan MAT 252 or Precision Isotope Ratio Mass Spectrometer (PRISM) with a  $1\sigma$  analytical uncertainty of  $\pm 0.08\text{\textperthousand}$  VPDB (McManus et al., 1999; Flower et al., 2000; Oppo et al., 1998). The ODP Site 981 data were measured on benthic foraminifera of the genus *Cibicidoides* (Raymo et al., 2004). ~~The original stratigraphic alignments for the combined benthic foraminiferal  $\delta^{18}\text{O}$  records of ODP Site 290 980/981, spliced together at  $\sim 860$  ka (Raymo et al., 2004), were manually determined using the Shackleton et al. (1990) target chronology for DSDP Site 677 (Oppo et al., 1998; McManus et al., 1999; Flower et al., 2000; Raymo et al., 2004).~~ The average sampling resolution for the combined ODP Site 980/981 record is  $\sim 1$  kyr over the past 1.8 Myr.

### 3 Stratigraphic Alignment

#### 295 3.1 Manual alignment

The manual alignment of the benthic foraminiferal  $\delta^{18}\text{O}$  record from IODP Site U1541 was generated by visual alignment of characteristic peaks and troughs in the record to the benthic foraminiferal LR04 stack using the publicly available QAnalySeries software (Kotov and Pälike, 2018), which is based on the original AnalySeries program of Paillard et al. (1996). The manual alignments of ODP Site 1090 and 300 ODP Site 980/981 were generated by realigning the original age model tie points for each site (e.g., Venz and Hodell, 2002; Raymo et al., 2004) to the LR04 stack using the QAnalySeries software. These manual alignment approaches yielded an average temporal spacing of  $\sim$ 30 kyr between tie points for IODP Site U1541 (116 tie points over the last 3.5 Myr; Fig. 2; Supplemental Table S2) and ODP Site 980/981 (66 tie points over the last 1.8 Myr; Fig. 2; Supplemental Table S4) and  $\sim$ 45 kyr between tie points for ODP 305 Site 1090 (65 tie points over the last 2.9 Myr; Fig. 2, Supplemental Tables S3). Due to the subjectivity inherent in user-defined alignments, the manual alignments presented in Supplemental Tables S2-S4 represent just one of many possible manual alignment outcomes for each record.



**Figure 2: Benthic foraminiferal  $\delta^{18}\text{O}$  records based on *Cibicidoides* and *Cibicides* sp. and spp. from IODP Site U1541 (purple, squares), ODP Site 1090 (dark blue, filled circles), and ODP Sites 980/981 (green, open circles). Data plotted using age models derived from manual alignment of each record to the LR04 global stack (Lisiecki and Raymo, 2005) for (a) 0-1200 ka; (b) 1200-2400 ka; and (c) 2400-3600 ka. The benthic foraminiferal  $\delta^{18}\text{O}$  records for ODP Site 1090 and ODP Site 980/981 are plotted with a -0.5‰ shift and a -1.0‰ shift, respectively, for clarity. Stratigraphic tie points for each manual alignment are denoted on the x-axis (Supplementary Table S1).**

### 3.2 Automated probabilistic alignment

310 HMM-Match iteratively aligns every data point of a benthic foraminiferal  $\delta^{18}\text{O}$  record along a depth scale to the benthic foraminiferal  $\delta^{18}\text{O}$  target record along an age scale using a probabilistic assessment of the

implied sedimentation rates associated with each alignment. The HMM-Match algorithm accounts for natural variance in a benthic foraminiferal  $\delta^{18}\text{O}$  data set (e.g., due to bioturbation, spatial variability, and measurement uncertainty) by assuming that the residual  $\delta^{18}\text{O}$  value between each input record and the target will fall along a normal probability distribution (Lin et al., 2014). The algorithm checks the implied relative sedimentation rate changes associated with each alignment fit against their natural likelihood based on the distribution of relative sedimentation rate changes observed in an independent compilation of 37 radiocarbon-derived sediment core chronologies over the last 40 kyr (Lin et al., 2014). The automated HMM-Match algorithm additionally generates Bayesian confidence bands for the resulting age model that provide age model uncertainties associated with the stratigraphic benthic foraminiferal  $\delta^{18}\text{O}$  alignment (Lin et al., 2014), in contrast to conventional manual stratigraphic alignment techniques.

While HMM-Match can generate objective and reproducible alignments of benthic foraminiferal  $\delta^{18}\text{O}$  records, the algorithm requires user-derived depth-age tie point estimates for the start and end points for the entire record or for each alignment segment. Alignment segments for each site were chosen to maximize relative stratigraphic continuity between gaps in sediment recovery (such as coring gaps) or data availability within each record. For IODP Site U1541, six input tie points were used to align the  $\sim 3.5$  Myr record that spans two coring gaps between 31.78-32.75 and 75.67-77.12 m CCSF-A (Winckler et al., 2021; Table 1). The specific depth-age values for these tie points were initially determined visually based on peaks and troughs in the benthic foraminiferal  $\delta^{18}\text{O}$  data near the top and bottom of each segment. Subsequently, where the visual alignment of segment ends was deemed ambiguous, the visually determined segment end point ages were varied by  $\pm 40$  kyr and rerun through the HMM-Match algorithm. The segment tie points of IODP Site U1541 were finalized based on the input tie points that yielded the lowest segment average age uncertainties in HMM-Match (Table 1).

335

**Table 1: Depth-age tie points for HMM-Match-based alignments.**

Core (Tie Point Source)	Depth (m CCSF-A or mcd)	Age (Ma)
IODP Site U1541 (this study)	0.00	0.000
	31.35	1.126

	32.90	1.198
	74.54	3.035
	77.32	3.135
	84.95	3.480
ODPT Site 980 (Raymo et al., 2004)	0.01	0.000
	93.81	0.860
ODP Site 981 (Raymo et al., 2004)	60.16	0.861
	109.90	1.812
TTNO57-6-PC4 (Venz and Hodell, 2002)	0.03	0.000
	9.00	0.339
ODP Site 1090 (Venz and Hodell, 2002)	12.76	0.416
	40.37	1.458
	40.42	1.476
	44.37	1.835
	44.42	1.866
	50.74	2.242
	50.94*	2.295
	51.84*	2.361
	51.89*	2.391
	53.68*	2.544
	53.73*	2.577
	56.08	2.720
	56.18	2.832
	57.33	2.903


\*Tie points were excluded from the HMM-Match to Prob-stack Alignment B for ODP Site 1090 to examine sensitivity to undetected hiatuses within the record.

For ODP Sites 1090 and 980/981, the depth-age inputs for HMM-Match were selected from the first and last tie points listed for each continuous section of published chronology (Table 1; Venz and Hodell, 2002; Raymo et al., 2004). At ODP Site 1090, where previous work has identified six mid-core hiatuses (Venz and Hodell, 2002), the HMM-Match alignment was constructed from seven segments bounded by these hiatuses. At ODP Site 980/981, where no such hiatuses have been reported, the HMM-Match alignment was constructed from only two segments- one segment each for the ODP Site 980 and ODP Site 981 records, respectively (Table 1; Raymo et al., 2004). We note that the automated HMM-Match algorithm typically assigns zero age uncertainty at the depth of each tie point used to bound the segments, thus the age uncertainty estimates reported at the start and end of each segment should be treated with caution.

Automated alignment algorithms, such as HMM-Match (Lin et al., 2014), introduce a risk of disconnecting users from nuances in the sedimentation history of a given site and possible stratigraphic discontinuities within a given benthic foraminiferal  $\delta^{18}\text{O}$  record. In order to test the HMM-Match 350 algorithm for sensitivities to hiatuses within the input data, we use the ODP Site 1090 benthic foraminiferal  $\delta^{18}\text{O}$  data to generate an additional alignment to the Prob-stack target (ODP Site 1090 Alignment B). The ODP Site 1090 Alignment B was based on the deliberate omission of some of the identified hiatuses in the benthic foraminiferal  $\delta^{18}\text{O}$  record of ODP Site 1090, using fewer tie point constraints and resulting in only four segments (Table 1).

### 355 3.3 Tuning Targets

LR04 – Since 2005, the LR04 stack (Fig. 3; Lisiecki and Raymo, 2005) has been the most commonly employed reference stack for global ocean  $\delta^{18}\text{O}$  variations on glacial-interglacial timescales over the past 5.3 Myr. While the LR04 benthic foraminiferal  $\delta^{18}\text{O}$  stack consists of 57 stacked records available at the 360 time of its development, the stack is heavily weighted towards Atlantic sediment core data due to limited core availability from the Pacific and Indian Oceans (Fig. 1). Due to decreasing record availability with increasing distance into the past, the resolution and number of records included in LR04 decrease moving further back in time (Lisiecki and Raymo, 2005). The late Pleistocene interval of the LR04 benthic foraminiferal  $\delta^{18}\text{O}$  stack is based on the average of 30 individual sediment core records yielding a 1 kyr stack resolution, while the Pliocene interval spanning 3 to 5.3 Ma is built on the average of 15 or fewer 365 individual records and has a 5 kyr stack resolution. Absolute ages for LR04 were determined from orbital tuning of a simple non-linear ice volume model to the 21 June isolation curve for 65°N based on the orbital solution of Laskar et al., 1993).



**Figure 3: Selected benthic foraminiferal  $\delta^{18}\text{O}$  alignment target records over the past 5 Ma.** Plotted targets include Prob-stack (black; Ahn et al., 2017), LR04 (blue; Lisiecki and Raymo, 2005), the CENOGRID splice (gray; Westerhold et al., 2020), and the LR09 Atlantic (yellow) and Pacific stacks (orange; Lisiecki and Raymo, 2009) for (a) 0-1000 ka; (b) 1000-2000 ka; (c) 2000-3000 ka; (d) 3000-4000 ka; and (e) 4000-5000 ka.

LR09 - The LR09 Pacific stack is composed of 14 benthic foraminiferal  $\delta^{18}\text{O}$  records, primarily from the equatorial and North Pacific Ocean, while the LR09 Atlantic stack is composed of 20 benthic foraminiferal  $\delta^{18}\text{O}$  records from the North and South Atlantic Ocean (Lisiecki and Raymo, 2009). These stacks were generated by averaging the available Pacific and Atlantic records from LR04 and three additional sites over the past 800 kyr, with a  $\pm 1$  kyr smoothing window for data points between 0 and 500 ka and a  $\pm 2$  kyr smoothing window between 500 and 800 ka (Fig. 3; Lisiecki and Raymo, 2009). The LR09 regional stacks were generated to examine regional differences in the timing of bottom water  $\delta^{18}\text{O}$  variability and only cover the last 800 kyr (Lisiecki and Raymo, 2009), but are included in our analysis to examine the impact of regional-specific targets on the HMM-Match-derived ages for our three example sites. The absolute age constraints on the LR09 Pacific and Atlantic stacks are the same as in the LR04 stack (Lisiecki and Raymo, 2009).

Prob-stack - The Prob-stack builds upon the LR04 global benthic foraminiferal  $\delta^{18}\text{O}$  stack covering the last 5.0 Myr with the inclusion of 123 additional records, including more data coverage from the Pacific Ocean, and a realignment of each record using the HMM-Match algorithm (Fig. 1 and 3; Ahn et al., 2017). As with the LR04 benthic foraminiferal  $\delta^{18}\text{O}$  stack, the resolution and number of records included in Prob-stack decreases further back in time, from 1 kyr resolution and more than 120 records in the Late Pleistocene to 5 kyr resolution and fewer than 20 records by 5 Ma (Fig. 1; Ahn et al., 2017). Although the exact structure of Prob-stack varies slightly from LR04 (Fig. 3), the Prob-stack target is based on the same absolute age constraints as the LR04 age model (Ahn et al., 2017).

CENOGRID - In contrast to stacked targets, such as LR04 (Lisiecki and Raymo, 2005), LR09 (Lisiecki and Raymo, 2009), and Prob-stack (Ahn et al., 2017), the CENOGRID composite represents a continuous splice of the benthic foraminiferal  $\delta^{18}\text{O}$  records with the highest resolution to span the entire Cenozoic (i.e., the past 66 Myr; Westerhold et al., 2020). CENOGRID covers the past 30 Myr with at least 2 kyr resolution (Westerhold et al., 2020). The CENOGRID splice has the advantage that it resolves high frequency variability in benthic foraminiferal  $\delta^{18}\text{O}$  and that it can be used for alignment of sedimentary

records that extend beyond the Plio-Pleistocene. The drawback of using CENORGRID as an alignment target, however, is that the composite splice consists entirely of low-latitude records and its 0-5 Ma  
400 interval is heavily biased towards benthic foraminiferal  $\delta^{18}\text{O}$  records from the Ceara Rise in the equatorial Atlantic (Fig. 3; Westerhold et al., 2020). Consequently, any regional variability in benthic foraminiferal  $\delta^{18}\text{O}$  observed in an undated record from a region outside the equatorial Atlantic would complicate millennial-scale probabilistic alignment to the CENOGGRID splice. The absolute age constraints for the CENOGGRID composite from 0 to 5 Ma are based on astronomical tuning of sediment images data,  
405 physical property data, and benthic foraminifera  $\delta^{18}\text{O}$  data from the Ceara Rise sites (i.e., ODP Sites 925, 926, 927, 928, and 929) to the Laskar et al. (2004) orbital solution (Wilkins et al., 2017; De Vleeschouwer et al., 2017; Westerhold et al., 2020). While there is general agreement between the LR04 and CENOGGRID ages over the past 3.5 Myr examined in this study, there is notable disagreement between them from 1.9 to 1.8 Ma (Wilkins et al., 2017).

410

Target Uncertainties - The alignment uncertainties computed by HMM-Match are based on the degree of misfit between the input benthic foraminiferal  $\delta^{18}\text{O}$  record and the target, which is a function of sampling resolution and signal to noise ratio of the input record as well as any uncertainties or estimated variance in the benthic foraminiferal  $\delta^{18}\text{O}$  value of the target record for each point in time (Lin et al., 2014). For  
415 alignments to Prob-stack and LR04, the published uncertainties in  $\delta^{18}\text{O}$  values for each reference target were used (Ahn et al., 2017; Lisiecki and Raymo, 2005). The  $1\sigma$  uncertainty of LR04 represents the mean standard error for each point within the stack and has an average value of 0.05 to 0.06 ‰ over the past 5.3 Myr (Lisiecki and Raymo, 2005). In contrast, the reported  $1\sigma$  uncertainty of Prob-stack represents the observed variance in the global records at each point within the stack, not the mean standard error, and  
420 has an average value of 0.18 ‰ over the past 5.0 Myr (Ahn et al., 2017).

No corresponding uncertainties were listed for the LR09 Pacific and Atlantic stacks (Lisiecki and Raymo, 2009) or for the CENOGGRID splice (Westerhold et al., 2020). Instead, we assign the  $1\sigma$   $\delta^{18}\text{O}$  uncertainties reported for 0-800 ka in LR04 to the LR09 Pacific and Atlantic stacks. We note that these values likely  
425 represent the upper end of the possible mean standard error of each regional stack because the LR09

Pacific stack ( $n = 14$ ) and the LR09 Atlantic stack ( $n = 20$ ) are both comprised of fewer and more regionally clustered records than the global LR04 stack ( $n = 57$ ). Similarly, we assign to the CENOGRID splice a  $1\sigma$   $\delta^{18}\text{O}$  uncertainty of 0.10‰ throughout the record, which is only slightly higher than analytical uncertainties of 0.06 to 0.08‰ reported for the Ceara Rise data sets used to generate the Plio-Pleistocene portion of the CENOGRID splice (ODP Sites 925, 926, 927, 928, and 929; Westerhold et al., 2020; Bickert et al., 1997; deMenocal et al., 1997; Tiedemann and Franz, 1997; Billups et al., 1998). As each of the stacked target records were developed with the assumption of synchronous changes in benthic foraminiferal  $\delta^{18}\text{O}$  values among the individual records that form the basis of each stack, none of the uncertainties in  $\delta^{18}\text{O}$  values reported or assigned for these stacks accounts for the regional variability in the timing of benthic foraminiferal  $\delta^{18}\text{O}$  changes in the global ocean. Nor do these uncertainties reflect potential offsets in the timing of benthic foraminiferal  $\delta^{18}\text{O}$  changes in an input record relative to the tuning target.

In the probabilistic HMM-Match alignment algorithm (Lin et al., 2014), tuning targets with higher designated benthic foraminiferal  $\delta^{18}\text{O}$  variance will yield stratigraphic alignment outputs with higher age model uncertainties. Given that the benthic foraminiferal  $\delta^{18}\text{O}$  uncertainties assigned to each target record are based on different assumptions and measures of variance (e.g., the  $1\sigma$   $\delta^{18}\text{O}$  values for Prob-stack reflect variance in the global data set while the  $1\sigma$   $\delta^{18}\text{O}$  values for CENOGRID reflect analytical uncertainty for measurements at a single site), we do not provide an in-depth comparison of the differences in HMM-Match-generated uncertainties for alignment of a single site to different tuning targets. Rather, we evaluate the impact of tuning target on resulting benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies by comparing the age offset between benthic foraminiferal  $\delta^{18}\text{O}$  alignments to Prob-stack with those to LR04, LR09 regional stacks, and CENOGRID, respectively. We employ Prob-stack as our benchmark tuning target because it is the most globally representative target, assuming globally synchronous changes in benthic  $\delta^{18}\text{O}$  (Ahn et al., 2017).

## 4 Results

For each example site (IODP Site U1541, ODP Site 1090, and ODP Site 980/981), we present the age models and age uncertainties (Supplementary Tables S5-7) generated by stratigraphic alignment of the benthic foraminiferal  $\delta^{18}\text{O}$  record via manual and automated probabilistic tuning (using the HMM-Match algorithm; Lin et al., 2014) to the LR04 stack and HMM-Match-based alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record to a selected suite of reference targets (LR04, LR09, Prob-stack, and the CENOGGRID splice). At ODP Site 1090, we additionally present the age models and age uncertainties associated with the Alignment B sensitivity test for stratigraphic discontinuity.


### 460 4.1 Age models and alignment outputs for IODP Site U1541

The manual and probabilistic HMM-Match-based alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record of IODP Site U1541 to the LR04 stack yield sedimentary age models that are largely consistent on the orbital scale, except for two apparent glacial cycle offsets in alignment, generating age model offsets up to 40 kyr and 47 kyr, at  $\sim 3.0$  Ma and  $\sim 3.4$  Ma, respectively (Fig. 4). The direction of age model offset (e.g., 465 whether HMM-Match-based ages are younger or older than the manually-derived ages) varies throughout the record. There is no systematic lead or lag associated with the age model of the HMM-Match-derived alignment relative to the age model of the manually-derived alignment. Including (excluding) the two intervals of complete cycle offsets, the mean value of the absolute difference between the manually aligned and HMM-Match aligned LR04 ages for IODP Site U1541 averages 4.6 kyr (3.2 kyr) across the 470 entire record. Periods with a high age model offset between the manually derived and HMM-Match-derived age models at IODP Site U1541 are indicative of intervals within the benthic foraminiferal  $\delta^{18}\text{O}$  record where stratigraphic similarities to the LR04 target stack are ambiguous due to regional differences in the shape of the benthic foraminiferal  $\delta^{18}\text{O}$  record, differences in temporal resolution between the input record and the target, and/or stratigraphic discontinuities (Fig. 4).

475

The stratigraphic alignment of the benthic foraminiferal  $\delta^{18}\text{O}$  record of IODP Site U1541 to the Prob-stack target using HMM-Match provides a sub orbital-scale chronology spanning the past 3.5 Myr in the upper 85 m (CCSF-A) of this site (Fig. 4). The HMM-Match generated  $1\sigma$  age uncertainties associated

with this alignment are on average 2 kyr across the record, although they are typically lower in the most 480 recent 1 Myr, where benthic foraminiferal  $\delta^{18}\text{O}$  variations occur at lower frequencies and with larger amplitude (Fig. 4). The highest age uncertainties in the record, reaching a maximum unidirectional  $1\sigma$  age uncertainty of 20 kyr, occur between 2190-2110 ka and likely result from a poorly defined Marine Isotope Stage (MIS) 82 in the benthic foraminiferal  $\delta^{18}\text{O}$  at IODP U1541 data relative to Prob-stack (Fig. 4).



**Figure 4: Stratigraphic benthic foraminiferal  $\delta^{18}\text{O}$  alignments for IODP Site U1541.** a) Prob-stack benthic foraminiferal  $\delta^{18}\text{O}$  target (Ahn et al., 2017) for comparison; b) Manual (light blue) and automated HMM-Match-based (dark blue; Lin et al., 2014) alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record from IODP Site U1541 to the LR04 benthic foraminiferal  $\delta^{18}\text{O}$  stack (Lisiecki and Raymo, 2005); c) Age model offset between the age models for IODP Site U1541 plotted in (b); d) Automated alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record from IODP Site U1541 to Prob-stack (black), CENOGGRID (gray; Westerhold et al., 2020), and the LR09 Pacific stack (brown; Lisiecki and Raymo, 2009) using HMM-Match; e)  $1\sigma$  age model uncertainties reported by HMM-Match for the benthic foraminiferal  $\delta^{18}\text{O}$  alignment of IODP Site U1541 to Prob-stack; f) Age offsets between the age models at IODP Site 1541 generated by the automated HMM-Match-based alignments to Prob-stack and LR04 (blue), Prob-stack and CENOGGRID (gray), and Prob-stack and LR09 Pacific stack (brown). The zero line is plotted in (c), (e), and (f) for reference.

485


Stratigraphic alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  data from IODP Site U1541 to the LR04, LR09, and CENOGGRID targets yield sedimentary chronostratigraphies that are similar on the orbital scale to the Prob-stack-derived U1541 age model (Fig. 4). However, there is a clear difference between alignments between 2220-2110 ka, overlapping with the period of high age model uncertainty for the benthic 490 foraminiferal  $\delta^{18}\text{O}$  alignment to Prob-stack reported by HMM-Match, when the magnitude of absolute age model offsets between the Prob-stack-based alignment and the IODP Site U1541 alignments to LR04 and to CENOGGRID reaches up to 60 kyr (Fig. 4). For the entire 3.5 Myr record, the average magnitude of absolute age offsets between the Prob-stack-based age model and those based on alignments to LR04 and to CENOGGRID are 5.2 and 5.7 kyr, respectively (Fig. 4, Supplemental Table S5). Over the last 495 800 kyr, the period for which the LR09 benthic  $\delta^{18}\text{O}$  stack is available for the Pacific Ocean, absolute age offsets between target alignment outputs are lower. Specifically, the average absolute age offsets between Prob-stack and LR04, LR09, and CENOGGRID-based alignment ages are 2.5, 2.3, and 4.7 kyr, respectively, across this interval (Supplemental Table S5). The largest offsets in this interval occur between 480-440 ka (i.e., MIS 12 and 13), when age offsets between the manually derived and HMM- 500 Match-derived alignments at IODP Site U1541 are also above average (Fig. 4).

The absolute values of the age offsets between the benthic foraminiferal  $\delta^{18}\text{O}$  alignment of IODP Site U1541 to Prob-stack and to the LR09 Pacific stack vary from near zero up to 16 kyr, with an average of 2.3 kyr (Fig. 4). The 16 kyr age offset occurs at 478 ka and coincides with relatively high age offsets 505 between the benthic foraminiferal  $\delta^{18}\text{O}$  alignment of IODP Site U1541 to Prob-stack and the LR04 (16 kyr) and CENOGGRID targets (14 kyr), respectively. This suggests that the age offset is unlikely to be caused by regional asynchronicity between Prob-stack and the LR09 Pacific stack over the past 800 kyr. The average age offset of 2.3 kyr between the benthic foraminiferal  $\delta^{18}\text{O}$  alignment of IODP Site U1541 510 to Prob-stack and to the LR09 Pacific stack is consistent with temporal differences of up to 4 kyr in benthic foraminiferal  $\delta^{18}\text{O}$  records from the Pacific and the North Atlantic (Lisiecki and Raymo, 2005; Lisiecki and Raymo, 2009; Skinner and Shackleton, 2005). While tuning to the closest regional stack may mitigate age uncertainties associated with spatially heterogenous and temporally diachronous  $\delta^{18}\text{O}$  records, the closest regional stack may still not be representative of a unique local record. Limited data availability has so far inhibited the development of a regional stack suitable for LCDW in the South 515 Pacific (e.g., Stern and Lisiecki, 2014). Further, high resolution benthic foraminiferal  $\delta^{18}\text{O}$  records, such as that of IODP Site U1541, may include sub-orbital-scale isotopic features that are not resolved in currently available Pleistocene stacks due to the relatively lower temporal resolution of most of the records used to construct the stacks.

## 4.2 Age models and alignment outputs for ODP Site 1090

520 Over the last 1.4 Myr, the manual and automated HMM-Match-based alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record from ODP Site 1090 to the LR04 stack yield similar chronostratigraphies with an average of the absolute values of age offsets equal to 3.7 kyr between them (Fig. 5). Our manual alignment of the benthic foraminiferal  $\delta^{18}\text{O}$  record at ODP Site 1090 resulted in different user-defined tie points and estimated stratigraphic hiatus durations compared to the original tie points defined by Venz 525 and Hodell (2002) and used for segmentation of the input record for automated HMM-Match-based alignment (Table 1). As a consequence, the age offsets between the manual and automated alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  at ODP Site 1090 are larger in the older part of the record (2.9 to 1.4 Ma), where multiple stratigraphic hiatuses are thought to occur (Venz and Hodell, 2002; Fig. 5). The maximum

age difference between the manual versus HMM-Match-based alignment methods of benthic foraminiferal  $\delta^{18}\text{O}$  for ODP Site 1090 across this interval is 110 kyr.



**Figure 5: Stratigraphic benthic foraminiferal  $\delta^{18}\text{O}$  alignments for ODP Site 1090.** a) Prob-stack benthic foraminiferal  $\delta^{18}\text{O}$  target (Ahn et al., 2017) for comparison; b) Manual (light blue) and automated HMM-Match-based (dark blue; Lin et al., 2014) alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record from ODP Site 1090 to the LR04 benthic foraminiferal  $\delta^{18}\text{O}$  stack (Lisiecki and Raymo, 2005); c) Age model offset between the age models for ODP Site 1090 plotted in (b); d) Automated alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record from ODP Site 1090 to Prob-stack (black), CENOGGRID (gray; Westerhold et al., 2020), and the LR09 Atlantic stack (brown; Lisiecki and Raymo, 2009) using HMM-Match

for benthic foraminiferal  $\delta^{18}\text{O}$  alignment of ODP Site 1090 to Prob-stack; f) Age offsets between the age models at ODP Site 1090 generated by the automated HMM-Match-based alignments to Prob-stack and LR04 (blue), Prob-stack and CENOGRID (gray), and Prob-stack and LR09 Atlantic stack (brown). The zero line is plotted in (c), (e), and (f) for reference.

The HMM-Match alignment of the benthic foraminiferal  $\delta^{18}\text{O}$  record at ODP Site 1090 to the Prob-stack target generates an orbital-scale age model over the past 2.9 Myr with an average reported  $1\sigma$  age uncertainty of  $\sim 2$  kyr and a maximum reported  $1\sigma$  age uncertainty of 11 kyr at 2870 ka (Fig. 5). These HMM-Match estimates of alignment uncertainties at ODP Site 1090 likely underestimate the true alignment uncertainty in the bottom portion of the record (2.9 to 1.4 Ma), where the HMM-Match algorithm assigns zero age uncertainty by default to the user-defined tie points at the start and stop of each hiatus, i.e., the end points of each segment input into the alignment algorithm.


540

Over the past 800 kyr, the average magnitude of absolute age offsets between the HMM-Match alignment of the benthic foraminiferal  $\delta^{18}\text{O}$  record at ODP Site 1090 to Prob-stack and similar alignments to the LR04, LR09 Atlantic, and CENOGRID tuning targets are 3.8 kyr, 3.5 kyr, and 5.7 kyr, respectively. The largest target-based age offset in the HMM-Match alignments of ODP Site 1090 is 26 kyr and occurs 545 between the alignment to Prob-stack and the alignment to LR04 at 2870 ka (Fig. 5). This large age offset coincides with the period of highest  $1\sigma$  age uncertainty in the HMM-Match-based alignment to Prob-stack for ODP Site 1090.

The HMM-Match-based Alignment B to Prob-stack, using four segments rather than seven to deliberately 550 skip known hiatuses in the sedimentary record, provides a broader perspective on the true uncertainties associated with the benthic foraminiferal  $\delta^{18}\text{O}$ -derived age models at ODP Site 1090 (Fig. 6). Specifically, Alignment B was generated without user-defined constraints on 3 mid-section hiatuses between 50.74 and 56.08 mcd (Table 1). In the absence of user-defined flags of these hiatuses into HMM-Match, Alignment B yields large age model uncertainties for ODP Site 1090 between 2.7 and 1.9 Ma, reaching 555 up to 60 kyr between 2.4 and 2.3 Ma (Fig. 6). While the reported age model uncertainties for the standard HMM-Match alignment to Prob-stack (based on 7 segments) at ODP Site 1090 are typically less than 10

kyr across this interval (Fig. 5), the age model offset between the standard alignment and Alignment B exceeds 160 kyr at 2.2 Ma (Fig. 6). This age model offset, a result of undiagnosed stratigraphic discontinuity, is the largest age model offset observed across all alignments investigated in this study.

560



**Figure 6: Comparison of age models at ODP Site 1090 generated by automated HMM-Match-based benthic foraminiferal  $\delta^{18}\text{O}$  alignments to Prob-stack using different segmentation.** a) HMM-Match-based benthic foraminiferal  $\delta^{18}\text{O}$  alignments to Prob-stack based on seven user-defined segments (blue; i.e., the standard 1090 HMM-Match alignment as plotted in Figure 5) versus four user-defined stratigraphic segments (orange; ODP Site 1090 Alignment B) compared to the Prob-stack target (black); b)  $1\sigma$  age model uncertainties reported by the automated HMM-Match alignment algorithm for the standard alignment and Alignment B of ODP Site 1090 to Prob-stack; c) Age model offset between the standard alignment and Alignment B of ODP Site 1090 to Prob-stack.


#### 4.3 Age models and alignment outputs for ODP Site 980/981

As observed at IODP Site U1541 and ODP Site 1090, the manual and automated HMM-Match-based benthic foraminiferal  $\delta^{18}\text{O}$  alignments of ODP Site 980/981 to the LR04 stack yield broadly similar 565 chronostratigraphies on the orbital scale (Fig. 7). The absolute difference in age models between these two alignment approaches reaches a maximum value of 24 kyr at 650 ka (leading into MIS 16) and has an average value of 2.8 kyr across the 1.8 Myr record at this site. The maximum age model offset at MIS 16 coincides with an interval of high temporal resolution in the benthic foraminiferal  $\delta^{18}\text{O}$  record at ODP 570 Site 980/981, with one sample every 0.6 kyr. Such an age offset may thus result from the inability of conventional manual alignment to resolve rapid changes in sedimentation rate across intervals with limited (i.e., smoothed) benthic foraminiferal  $\delta^{18}\text{O}$  variability in the tuning target. This observation demonstrates that age uncertainties associated with manual tuning cannot always be reduced by increasing the temporal resolution of the input data and suggests that a smoothing of a high-resolution input record prior to alignment may be beneficial for tuning.

575

The HMM-Match-based benthic foraminiferal  $\delta^{18}\text{O}$  alignment to the Prob-stack target at ODP Site 980/981 generates an orbital-scale age model over the past 1.8 Myr with an average and maximum reported  $1\sigma$  age uncertainty of 0.8 kyr and 5.2 kyr, respectively (Fig. 7). In comparison, the magnitude of absolute age offsets between the age models generated by automated HMM-Match-based benthic 580 foraminiferal  $\delta^{18}\text{O}$  alignments to the LR04, LR09 Atlantic, and CENOGGRID targets at ODP Site 980/981 are larger, with a maximum value of 18 kyr between the alignments to Prob-stack and to LR04 at 1.42 Ma (Fig. 7). Over the last 800 kyr, the average absolute age offsets between automated HMM-Match-based benthic foraminiferal  $\delta^{18}\text{O}$  alignments to Prob-stack and to LR04, LR09 Atlantic, and CENOGGRID at ODP Site 980/981 are 2.0 kyr, 1.4 kyr, and 4.7 kyr, respectively (Fig. 7).

585



**Figure 7: Stratigraphic benthic foraminiferal  $\delta^{18}\text{O}$  alignments for ODP Site 980/981.** a) Prob-stack benthic foraminiferal  $\delta^{18}\text{O}$  target (Ahn et al., 2017) for comparison; b) Manual (light blue) and automated HMM-Match-based (dark blue; Lin et al., 2014) alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record from ODP Site 980/981 to the LR04 benthic foraminiferal  $\delta^{18}\text{O}$  stack (Lisiecki and Raymo, 2005); c) Age model offset between the age models for ODP Site 980/981 plotted in (b); d) Automated alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record from ODP Site 980/981 to Prob-stack (black), CENOGGRID (gray; Westerhold et al., 2020), and the LR09 Atlantic stack (brown; Lisiecki and Raymo, 2009) using HMM-Match; e)  $1\sigma$  age model uncertainties reported by HMM-Match for the benthic foraminiferal  $\delta^{18}\text{O}$  alignment of ODP Site 980/981 to Prob-stack; f) Age offsets between the age models at ODP Site 980/981 generated by the automated HMM-Match-based alignments to Prob-stack and LR04 (blue), Prob-stack and CENOGGRID (gray), and Prob-stack and LR09 Atlantic stack (brown). The zero line is plotted in (c), (e), and (f) for reference.

## 5 Discussion

Based on the benthic foraminiferal  $\delta^{18}\text{O}$  alignment results presented above for IODP Site U1541, ODP Site 1090, and ODP Site 980/981, we discuss and synthesize the largest sources of uncertainty in benthic  
590 foraminiferal  $\delta^{18}\text{O}$ -based chronostratigraphies and suggest best community practices for the application of this approach.


### 5.1 Manual vs. probabilistic alignment

The probabilistic stratigraphic alignment algorithm HMM-Match (Lin et al., 2014) provides three advantages over manual alignment approaches for the development of benthic foraminiferal  $\delta^{18}\text{O}$   
595 (chrono)stratigraphies: 1) it reduces the amount of user subjectivity in assigning tie points; 2) it generates estimates of age uncertainty associated with the alignment method itself; and 3) by aligning every data point within an input benthic foraminiferal  $\delta^{18}\text{O}$  record, HMM-Match estimates high-resolution sedimentation rate changes throughout an entire record (Fig. 8). These sedimentation rate changes are more realistic than the limited average sedimentation rate estimates between user-defined tie points that  
600 result from a manual alignment of benthic foraminiferal  $\delta^{18}\text{O}$  to a tuning target. In addition, linear sedimentation rate records with high temporal resolution can provide useful information on the relationship between environmental and climatic variability on sediment accumulation, dissolution, and redistribution processes on the seafloor (e.g., Suman and Bacon, 1989; Costa et al., 2020; Francois et al., 1990).

605

The age offsets generated between manual and automated probabilistic (HMM-Match-based) alignments of the benthic foraminiferal  $\delta^{18}\text{O}$  record to the LR04 benthic stack at IODP Site U1541 (Fig. 4), ODP Site 1090 (Fig. 5), and ODP Site 980/981 (Fig. 7) provide a means to start to quantitatively assess the magnitude of alignment uncertainties incurred from manual alignments with user-defined tie points.  
610 These uncertainties typically average between 3 to 5 kyr across the three sites, but can reach values of 40 kyr or higher near segment ends (e.g., at 2.7 Ma at ODP Site 1090) and intervals where local benthic

foraminiferal  $\delta^{18}\text{O}$  changes differ from the reference target (e.g., at 3.0 Ma and 3.4 Ma at IODP Site U1541). This range of uncertainty is comparable to the 2.5 kyr tuning uncertainty estimated by Martinson et al. (1987) based on the manual alignment of five paleoclimatological indicators from a single site (planktonic and benthic foraminiferal  $\delta^{18}\text{O}$ , radiolarian abundance, sea surface temperature, stable carbon isotopes, and carbonate content) to a stacked  $\delta^{18}\text{O}$  target. As observed at ODP Site 980/981, high temporal resolution of the benthic foraminiferal  $\delta^{18}\text{O}$  record does not guarantee a reduction in this form of alignment uncertainty (Section 4.3).



**Figure 8: Linear sedimentation rate (LSR) estimates for IODP Site U1541, ODP Site 1090, and ODP Site 980/981.** Values are based on manual alignment (blue) and automated HMM-Match alignment (gray) of benthic foraminiferal  $\delta^{18}\text{O}$  records to the LR04 benthic stack (Lin et al., 2014; Lisiecki and Raymo, 2004).

The age model uncertainties generated by HMM-Match for the benthic foraminiferal  $\delta^{18}\text{O}$  alignments at IODP Site U1541 (Fig. 4), ODP Site 1090 (Fig. 5), and ODP Site 980/981 (Fig. 7) include uncertainties associated with the reported benthic foraminiferal  $\delta^{18}\text{O}$  variance in the target reference record and the stratigraphic alignment method itself (i.e., the process of ‘wiggle matching;’ Lin et al., 2014). Uncertainties in the latter are calculated under the assumptions of reasonable stratigraphic continuity and that user-defined ages for each segment start and end are certain (Lin et al., 2014). These values average ~1 to 3 kyr across the three example sites and typically decrease with increased temporal resolution of the undated benthic foraminiferal  $\delta^{18}\text{O}$  record and of the tuning target (which are generally better constrained in the late Pleistocene than in the Pliocene).

The point-by-point alignment uncertainties estimated by HMM-Match are valuable for assessing changes in the confidence level of benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies across multi-million year records, where long-term changes in the seawater  $\delta^{18}\text{O}$  signal, or temporal resolution of the input record and tuning target may cause systematic changes in age uncertainty. Although HMM-Match assigns unrealistically low age uncertainties at user-defined tie points for the start and end of an alignment segment, this issue has been resolved in more recent evolutions of automated alignment algorithms (Lee et al., 2023). For example, the new automated alignment software package, BIGMACS, allows users to specify the uncertainties of added tie points to generate more realistic estimates of alignment uncertainty (Lee et al., 2023). However, neither HMM-Match-based nor BIGMACS-based alignment age uncertainties account for the uncertainty associated with the absolute chronology of the target reference or regional variability in bottom water temperature and/or the timing of local changes in seawater (and resulting benthic foraminiferal)  $\delta^{18}\text{O}$  between the input record and the selected tuning target.

While automated alignment algorithms like HMM-Match and BIGMACS provide many advantages in the generation of benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies, we note that depositional environments with highly variable sedimentation rate changes and benthic foraminiferal  $\delta^{18}\text{O}$  records with long data gaps or temporal variations in sampling resolution may not abide by the assumptions included in these automated algorithms. Specifically, the HMM-Match algorithm is designed to minimize large

650 sedimentation rate changes between data points based on the probability distribution calculated from a compilation of 37 radiocarbon-dated sediment cores (Lin et al., 2014). This guiding principle may hinder the success of HMM-Match-generated alignments for benthic foraminiferal  $\delta^{18}\text{O}$  records with irregular sampling frequencies or from regions like the Antarctic Southern Ocean where very large sedimentation rate changes are expected across a deglaciation (e.g., Hasenfratz et al., 2019). Consequently, a close  
655 visual evaluation of the automated alignment outputs against their designated targets should be completed manually (i.e., by the user) to quality check the resulting benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies.

## 5.2 Choice of tuning target and the assumption of global synchronicity

We encourage the utilization of Prob-stack as a global tuning target because Prob-stack includes the most  
660 holistic estimate to date of spatial variance in benthic foraminiferal  $\delta^{18}\text{O}$  values across the last 5 Myr, under the assumption of global synchronicity in benthic foraminiferal  $\delta^{18}\text{O}$  variability, and therefore is best suited for calculating more realistic estimates of chronostratigraphic alignment uncertainties (Ahn et al., 2017). Our analyses reveal an average age offset of 2-6 kyr between benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies generated for IODP Site U1541, ODP Site 1090, and ODP Site 980/981 alignments  
665 to Prob-stack, LR04, and CENOGGRID using HMM-Match (Fig. 4, 5, and 7, respectively). These values suggest that the choice of tuning target, or comparing records based on alignments to different tuning targets, can introduce a level of age uncertainty comparable to the uncertainty associated with the alignment itself (Section 5.1). The magnitude of these age offsets is similar to that of regional differences in the timing of benthic foraminiferal  $\delta^{18}\text{O}$  changes in the global ocean across glacial/interglacial cycles  
670 (i.e., up to 4 kyr; Lisiecki and Raymo, 2009; Skinner and Shackleton, 2005; Stern and Lisiecki, 2014; Waelbroeck et al., 2011; Layberie et al., 2005).

The age model uncertainties caused by regional asynchronicity in temperature and/or seawater (i.e., benthic foraminiferal)  $\delta^{18}\text{O}$  changes across glacial cycles must also be considered when comparing event  
675 timing among benthic foraminiferal  $\delta^{18}\text{O}$ -tuned sediment records from different basins and water masses. The age offsets observed between the benthic foraminiferal  $\delta^{18}\text{O}$  alignments at IODP Site U1541, ODP

Site 1090, and ODP Site 980/981 to Prob-stack and their respective alignments to the LR09 Pacific and Atlantic regional stacks provide a qualitative assessment of the impact of regional differences on age model uncertainties and average 1.4 to 3.5 kyr over the past 800 kyr.

680

Regional stacks of individual benthic foraminiferal  $\delta^{18}\text{O}$  records with high-resolution radiocarbon dates over the last glacial cycle suggest that the spatial variability in seawater  $\delta^{18}\text{O}$  and its response to changing climate may induce systematic age offsets up to 1.7 kyr between the deep South Pacific IODP Site U1541 record and ODP Sites 1090 and 980/981 in the deep South Atlantic and deep North Atlantic, respectively, 685 during glacial terminations (Stern and Lisiecki, 2014). Significant age offsets, up to 1.0 and 3.5 kyr, should be considered when comparing the new IODP Site U1541 data to records from the intermediate Pacific and the deep Indian basin, respectively (Stern and Lisiecki, 2014). The development and refinement of a deep Pacific regional stack extending beyond the last glacial cycle would help further define the uncertainties associated with asynchronous  $\delta^{18}\text{O}$  changes in the global ocean.

690

While age uncertainties associated with the absolute chronology of the alignment target are less of a concern for the comparison of relative event timing between sedimentary records with similarly stratigraphically determined age models, estimates of such age uncertainties are critical for comparison with terrestrial or independently dated marine records. The absolute age uncertainty associated with the 695 LR04 stack is estimated to yield age model uncertainties of 4 kyr for 0-1 Ma, 6 kyr for 1-3 Ma, 15 kyr from 3-4 Ma, and 30 kyr or higher for 4-5.3 Ma (Lisiecki and Raymo, 2005). The absolute age uncertainty associated with the LR09 stacks and Prob-stacks is expected to be similar to the values for LR04, as these stacks are based on the same time domain as LR04 (Lisiecki and Raymo, 2009; Ahn et al., 20017). Absolute age uncertainties are not specifically reported for the 0-5 Ma interval of the CENOGRID 700 composite based on astronomical tuning of a number of high-resolution benthic foraminiferal  $\delta^{18}\text{O}$  records from Ceara Rise (i.e., ODP Sites 925, 926, 927, 928 and 929), however a broader absolute age uncertainty estimate of 10 kyr is reported for the late Miocene to Pleistocene interval of CENOGRID (Wilkens et al., 2017; Westerhold et al., 2020). Including the absolute age uncertainties of the tuning

targets in stratigraphic benthic foraminiferal  $\delta^{18}\text{O}$  alignments would significantly increase the total  
705 reported age uncertainties of marine sedimentary records.

### 5.3 Stratigraphic discontinuity

The largest source of age uncertainties identified by our analysis arises from stratigraphic discontinuities  
within the input benthic foraminiferal  $\delta^{18}\text{O}$  record, as shown by ODP Site 1090 (Fig. 5 and 6). Coring  
710 gaps, extended sampling breaks, and sedimentary hiatuses within an undated record invalidate the  
assumption of stratigraphic continuity on which benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies are built  
and require a sound assessment of potential hiatuses in the sedimentary record by the user. The  
uncertainties associated with these stratigraphic discontinuities are higher during the 40 kyr-glacial cycles  
of the Pliocene and early Pleistocene, when there are lower amplitude and higher frequency benthic  
715 foraminiferal  $\delta^{18}\text{O}$  changes than during the late Pleistocene. Missing cycles are, accordingly, harder to  
identify within this interval (Ahn et al., 2017; Lisiecki and Raymo, 2005; Westerhold et al., 2020).

Although the HMM-Match algorithm did not independently identify the specific mid-section hiatuses  
reported by Venz and Hodell (2002) in the benthic foraminiferal  $\delta^{18}\text{O}$  record at ODP Site 1090 in our  
sensitivity test, the algorithm reported large uncertainty estimates (reaching up to  $\sim$ 60 kyr) across this  
720 interval of discontinuity (Fig. 6). The increase in HMM-Match generated alignment uncertainties thereby  
flags intervals of suspected stratigraphic discontinuity at ODP Site 1090 and demonstrates the utility of  
this algorithm to detect intervals of stratigraphic complication within a long sedimentary record (Fig. 6).  
Such hiatuses may otherwise be difficult to detect and could introduce tens of kyr of systematic error into  
benthic foraminiferal  $\delta^{18}\text{O}$ -derived sediment chronologies. We caution, however, that HMM-Match has  
725 been documented to generate alignment errors and underestimated age uncertainties in high resolution  
benthic foraminiferal  $\delta^{18}\text{O}$  records with millennial-scale sampling gaps along the Iberian Margin (Lisiecki  
et al., 2022). Interruptions in the stratigraphic integrity of the sediment record under investigation can be  
assessed via core descriptions, imagery, x-ray fluorescence (XRF) scanning data or other high resolution  
indicators of sediment stratigraphy in order to make an informed refinement of the stratigraphic alignment  
730 beyond age uncertainties reported by HMM-Match.

One may hypothesize, based on the relatively high age uncertainties for the automated HMM-Match-based benthic foraminiferal  $\delta^{18}\text{O}$  alignment to Prob-stack at IODP Site U1541 between 2190-2110 ka (reaching up to 20 kyr), that the benthic foraminiferal  $\delta^{18}\text{O}$  record of this MIS 82 interval may be disrupted  
735 by a mid-section hiatus (Fig. 4). However, no such stratigraphic discontinuities are observed in the high resolution (1-2 cm) shipboard density or color records at IODP Site U1541 across this interval (Winckler et al., 2021). Rather, this section may reflect an interval where local changes in benthic foraminiferal  $\delta^{18}\text{O}$  values deviate significantly from the global stack.

#### 5.4 Suggestions for best practice

740 Our HMM-Match analyses of benthic foraminiferal  $\delta^{18}\text{O}$  alignments at IODP Site U1541, ODP Site 1090, and ODP Site 980/981 to common tuning targets suggest combined age uncertainties for the HMM-Match-based alignments to Prob-stack to be  $\pm$  5-6 kyr for 0 to 1 Ma,  $\pm$  7-9 kyr for 1 to 3 Ma, and  $\pm$  18 kyr for 3 to 3.5 Ma. These values include age uncertainties associated with the alignment process, benthic foraminiferal  $\delta^{18}\text{O}$  variance in the tuning target, and absolute age constraints on the tuning target (e.g.,  
745 the absolute age uncertainties reported by Lisiecki and Raymo, 2005). The largest contribution to these uncertainty estimates is the uncertainty on absolute ages of the target record (Section 5.2, Lisiecki and Raymo, 2005). Continued community efforts to provide independent age control on the absolute ages of benthic foraminiferal  $\delta^{18}\text{O}$  tuning targets, especially for sediments older than 3 Ma, will be critical for the further reduction of total uncertainty associated with benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphy.

750

In light of the existing caveats of benthic foraminiferal  $\delta^{18}\text{O}$  alignments, we suggest the following best practices for stratigraphically aligning benthic foraminiferal  $\delta^{18}\text{O}$  records to common targets to minimize alignment uncertainty and ensure consistent characterization of age model uncertainty during the Plio-Pleistocene.

755 1) Investigate possible disruptions in stratigraphic continuity using sediment core images, XRF, or other high-resolution scan data.

2) Utilize an automated and probabilistic alignment algorithm, like HMM-Match (Lin et al., 2014), to reduce user subjectivity in record alignment and to generate point-by-point estimates of age uncertainties and sedimentation rate variations resulting from the alignment.

760 3) Visually assess the quality of alignment outputs generated by automated algorithms for alignment mismatches or other irregularities, especially across glacial terminations.

765 4) Align benthic foraminiferal  $\delta^{18}\text{O}$  records to stacked records that include characterization of global or regional heterogeneity in  $\delta^{18}\text{O}$  variance (e.g., Prob-stack; Ahn et al., 2017; or an appropriate regional stack, if available) to better constrain uncertainties associated with the alignment itself.

5) Acknowledge the amount by which benthic foraminiferal  $\delta^{18}\text{O}$  changes at the study site may lead or lag the tuning target of choice. The development of updated and long-term regional stacks will help reduce these associated age uncertainties.

770 6) When working with multiple sediment cores with age models based on stratigraphic benthic foraminiferal  $\delta^{18}\text{O}$  alignments, ensure the age models for each record are calculated using the same approaches and tuning targets.

7) Publish sample depth data along with sample ages and measurement values so that it is transparent how published records may be updated or realigned to newer age models.

775 On a broader note, given the financial burden of programming platforms like MATLAB, we encourage the development or translation of automated alignment algorithms into license-free coding languages such as Python or R so that they may become more accessible to the research community.

## 6 Conclusions

Stratigraphic alignments based on new and published benthic foraminiferal  $\delta^{18}\text{O}$  records of IODP Site 780 U1541, ODP Site 1090, and ODP Site 980/981 provide valuable insights into the types and range of sediment age model uncertainties associated with benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies. Our results suggest typical age uncertainties of 3 to 5 kyr for manual alignment efforts and 1 to 3 kyr for probabilistic HMM-Match-based alignments (Lin et al., 2014) over the past 3.5 Myr. We identify an

average age offset of 2-6 kyr between stratigraphic benthic foraminiferal  $\delta^{18}\text{O}$  alignments to Prob-stack  
785 (Ahn et al., 2017), LR04 (Lisiecki and Raymo, 2005); and the CENOGRID target (Westerhold et al.,  
2020). The 2-6 kyr age uncertainty associated with tuning target choice is comparable to the estimated 1-  
4 kyr range of lag time in benthic foraminiferal  $\delta^{18}\text{O}$  changes between different ocean basins and water  
masses (Lisiecki and Raymo, 2009; Skinner and Shackleton, 2005; Stern and Lisiecki, 2014; Waelbroeck  
et al., 2011; Layberie et al., 2005). Stratigraphic discontinuities within the benthic foraminiferal  $\delta^{18}\text{O}$   
790 input record can contribute to large age uncertainties (reaching 60 kyr and higher) in marine sediment  
core chronostratigraphies.

The uncertainties associated with benthic foraminiferal  $\delta^{18}\text{O}$  chronostratigraphies and their impact on  
paleoceanographic interpretations can be reduced using best practices for age model uncertainty  
795 characterization and alignment. These practices include: investigation into the stratigraphic continuity of  
an undated record; utilization of probabilistic alignment algorithms (such as HMM-Match or BIGMACS;  
Lin et al., 2014; Lee et al., 2023); and alignment targets that suitably characterize heterogeneity in  $\delta^{18}\text{O}$   
variance (such as Prob-stack globally; Ahn et al., 2017; or recent regional stacks like the North Atlantic  
stack of Hobart et al., 2023); and acknowledgement of the amount by which benthic foraminiferal  $\delta^{18}\text{O}$   
800 changes of the study region may lead or lag the alignment target. Continued investigation into regional  
asynchronicity in seawater and benthic foraminiferal  $\delta^{18}\text{O}$  changes across the Plio-Pleistocene and  
independent constraints on the absolute ages of tuning target intervals, particularly older than 3 Ma,  
should also be supported to further reduce age uncertainties based on stratigraphic benthic foraminiferal  
 $\delta^{18}\text{O}$  alignments of marine sedimentary records to a common tuning target.

## 805 Data Availability

All data in this paper are available in the references and supplemental tables and from the PANGAEA  
database (DOI/URL to be updated).

## Team List

Expedition 383 Scientists not separately listed as authors include: Carlos A. Alvarez Zarikian  
810 (International Ocean Discovery Program, Texas A&M University, USA), Helge W. Arz (Leibniz Institute  
for Baltic Sea Research Warnemünde, Germany), Chandranath Basak (Department of Earth Sciences,  
University of Delaware, USA), Anieke Brombacher (Department of Earth and Planetary Sciences, Yale  
University, USA), Oliver M. Esper (Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine  
Research, Germany), Lisa C. Herbert (Department of Earth, Ocean, and Atmospheric Science, Florida  
815 State University, USA), Shinya Iwasaki (Research and Development Center for Global Change,  
JAMSTEC, Japan), Vera J. Lawson (Department of Earth and Planetary Sciences, Rutgers, USA), Lester  
Lembke-Jene (Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research, Germany),  
Li Lo (Department of Geosciences, National Taiwan University, Taiwan), Alexandre L. de Souza  
(Department of Geology, Federal University of Rio de Janeiro, Brazil), Elisa Malinverno (Department of  
820 Earth and Environmental Sciences, University of Milano-Bicocca, Italy), Elisabeth Michel (Laboratoire  
des Sciences du Climat et de l'Environnement LSCE, Institute Pierre Simon Laplace IPSL, CRNS-CEA-  
UVSQ, France), Simone Moretti (Climate Geochemistry Department, Max Planck Institute for Chemistry  
MPIC, Germany), Christopher M. Moy (Geology Department, University of Otago, New Zealand), Ana  
Christina Ravelo (Ocean Sciences Department, University of California Santa Cruz, USA), Christina R.  
825 Riesselman (Geology Department, University of Otago, New Zealand), Mariem Saavedra-Pellitero  
(School of the Environment, Geography and Geosciences, University of Portsmouth, United Kingdom),  
Inah Seo (Global Ocean Research Center, Korea Institute of Ocean Science and Technology KIOST,  
Republic of Korea), Raj K. Singh (School of Earth, Ocean and Climate Sciences, Indian Institute of  
Technology Bhubaneshwar, India), Rebecca A. Smith (Department of Geosciences, University of  
830 Massachusetts-Amherst, USA), Joseph S. Stoner (College of Earth, Ocean and Atmospheric Sciences,  
Oregon State University, USA), Igor M. Venancio (Department of Geochemistry, Universidade Federal  
Fluminense, Brazil), Sui Wan (South China Sea Institute of Oceanology, Chinese Academy of Sciences,  
China), Xiangyu Zhao (National Institute of Polar Research, Geoscience Group, Japan).

## 835 Author Contributions

JLM, GW, and JG designed the study and led the research. JLM, JG, JH, CK, JRF prepared the IODP Site U1541 sediment samples for benthic foraminiferal oxygen isotope analysis. Expedition 383 Scientists (FL, GW, HWA, JRF, JG, LLJ, JLM, CAZ, CB, AB, OME, LCH, SI, VJL, LL, EM, EM, SM, CMM, ACR, CRR, MS-P, IS, RKS, RAS, ALS, JSS, IM, SW, XYZ) collected the stratigraphic, 840 sedimentological, and physical properties data from IODP Site U1541 and contributed to the interpretation of results. FL provided the XRF-based alignment of IODP Site U1541 and pre-site survey core PS75/059-2. JLM, JG, and GW did the alignment analysis and wrote the manuscript with discussion and contributions from LEL and Expedition 383 Scientists.

## Acknowledgements

845 We are grateful to the captain, crew, and scientific party of the R/V *JOIDES Resolution* for their support during International Ocean Discovery Program (IODP) Expedition 383 “Dynamics of the Pacific Antarctic Circumpolar Current (DYNAPACC).” We thank W. Huang and B. Linsley for technical assistance with oxygen isotope analyses and T. Lamones for insightful discussions. We acknowledge the IODP U.S. Science Support Program for supplemental funding of post-expedition activities via post- 850 expedition activity awards to GW, JLM, JG, and JRF. We additionally thank two anonymous reviewers for their constructive feedback. GW and JLM acknowledge support from National Science Foundation (NSF) grant 2305426. JG acknowledges support from the Deutsche Forschungsgemeinschaft through grant GO 2294/3-1. JRF acknowledges support from NSF grant 2305427.

## 855 Competing Interests

The authors declare that they have no conflict of interest.

## References

Ahn, S., Khider, D., Lisiecki, L. E., and Lawrence, C. E.: A probabilistic Pliocene–Pleistocene stack of benthic  
860  $\delta^{18}\text{O}$  using a profile hidden Markov model, *Dynamics and Statistics of the Climate System*, 2,  
<https://doi.org/10.1093/climsys/dzx002>, 2017.

Berger, A.: Third International Conference on Paleo-oceanography, 16 pp., 1989.

Berger, A. and Loutre, M. F.: New Insolation values for the climate of the last 10 million years, *Scientific Report*  
865 1988/13. Institute d’Astronomie et de Geophysique Georges Lemaitre. Universite Catholique de Louvain-la-  
Neuve, 1988.

Bickert, T., Curry, W., and Wefer, G.: Late Pliocene to Holocene (2.6-0 Ma) western equatorial Atlantic deep-  
water circulation: Inferences from benthic stable isotopes, *Proceedings of the Ocean Drilling Program, Scientific Results*, 154, 239–254, 1997.

Billups, K., Ravelo, A. C., and Zachos, J. C.: Early Pliocene deep water circulation in the western equatorial  
870 Atlantic: Implications for high-latitude climate change, *Paleoceanography*, 13, 84–95, 1998.

Channell, J. E. T. and Raymo, M. E.: Paleomagnetic record at ODP Site 980 (Feni Drift, Rockall) for the past 1.2 Myrs, *Geochemistry, Geophysics, Geosystems*, 4, <https://doi.org/10.1029/2002GC000440>, 2003.

Costa, K. M., Hayes, C. T., Anderson, R. F., Pavia, F. J., Bausch, A., Deng, F., Dutay, J. C., Geibert, W., Heinze,  
875 C., Henderson, G., Hillaire-Marcel, C., Hoffmann, S., Jaccard, S. L., Jacobel, A. W., Kienast, S. S., Kipp, L.,  
Lerner, P., Lippold, J., Lund, D., Marcantonio, F., McGee, D., McManus, J. F., Mekik, F., Middleton, J. L.,  
Missiaen, L., Not, C., Pichat, S., Robinson, L. F., Rowland, G. H., Roy-Barman, M., Tagliabue, A., Torfstein,  
A., Winckler, G., and Zhou, Y.: 230Th Normalization: New Insights on an Essential Tool for Quantifying  
Sedimentary Fluxes in the Modern and Quaternary Ocean, *Paleoceanogr Paleoclimatol*, 35,  
<https://doi.org/10.1029/2019PA003820>, 2020.

880 deMenocal, P., Archer, D., and Leth, P.: Pleistocene variations in deep Atlantic circulation and calcite burial  
between 1.2 and 0.6 Ma: a combined data-model approach, in: *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 154, edited by: Shackleton, N. J., Curry, W. B., Richter, C., and Bralower, T. J., Ocean Drilling Program, <https://doi.org/10.2973/odp.proc.sr.154.113.1997>, 1997.

Diekmann, B. and Kuhn, G.: Sedimentary record of the mid-Pleistocene climate transition in the southeastern South  
885 Atlantic (ODP Site 1090), *Palaeogeography, Paleaeoclimatology, Palaeoecology*, 182, 241–258, 2002.

Elderfield, H., Ferretti, P., Greaves, M., Crowhurst, S., Mccave, I. N., Hodell, D., and Piotrowski, A. M.: Evolution of Ocean Temperature and Ice Volume Through the Mid-Pleistocene Climate Transition, *Science*, 337, 704–709, <https://doi.org/10.1126/science.1223055>, 2012.

Flower, B. P., Oppo, D. W., McManus, J. F., Venz, K. A., Hodell, D. A., and Cullen, J. L.: North Atlantic intermediate to deep water circulation and chemical stratification during the past 1 Myr, *Paleoceanography*, 15, 388–403, <https://doi.org/10.1029/1999PA000430>, 2000.

Ford, H. L. and Raymo, M. E.: Regional and global signals in seawater  $\delta^{18}\text{O}$  records across the mid-Pleistocene transition, *Geology*, 48, 113–117, <https://doi.org/10.1130/G46546.1>, 2020.

Francois, R., Bacon, M. P., and Suman, D. O.: Thorium 230 profiling in deep-sea sediments: High-resolution records of flux and dissolution of carbonate in the Equatorial Atlantic during the last 24,000 years, *Paleoceanography*, 5, 761–787, 1990.

Gersonde, R., Hodell, D. A., Blum, P., and et al.: Leg 177 Summary: Southern Ocean Paleoceanography, *Proceedings of the Ocean Drilling Program, Initial Reports*, 177, 1999.

Hobart, B., Lisiecki, L. E., Rand, D., Lee, T., and Lawrence, C. E.: Late Pleistocene 100-kyr glacial cycles paced by precession forcing of summer insolation, *Nat Geosci*, 16, 717–722, <https://doi.org/10.1038/s41561-023-01235-x>, 2023.

Hodell, D. A., Charles, C. D., and Ninnemann, U. S.: Comparison of interglacial stages in the South Atlantic sector of the southern ocean for the past 450 kyr: implications for Marine Isotope Stage (MIS) 11, *Glob Planet Change*, 24, 7–26, 2000.

Hodell, D. A., Crowhurst, S. J., Lourens, L., Margari, V., Nicolson, J., Rolfe, J. E., Skinner, L. C., Thomas, N. C., Tzedakis, P. C., Mlneck-Vautravers, M. J., and Wolff, E. W.: A 1.5-million-year record of orbital and millennial climate variability in the North Atlantic, *Climate of the Past*, 19, 607–636, <https://doi.org/10.5194/cp-19-607-2023>, 2023.

Huybers, P. and Wunsch, C.: A depth-derived Pleistocene age model: Uncertainty estimates, sedimentation variability, and nonlinear climate change, *Paleoceanography*, 19, <https://doi.org/10.1029/2002PA000857>, 2004.

Imbrie, J., Hays, J. D., Martinson, D. G., McIntyre, A., Mix, A. C., Morley, J. J., Pisias, N. G., Prell, W. L., and Shackleton, N. J.: The orbital theory of Pleistocene climate: Support from a revised chronology of the marine  $\delta^{18}\text{O}$  record, in: *Milankovitch and Climate, Part 1*, edited by: Berger, A., Springer, New York, 269–305, 1984.

Jansen, E., Raymo, M. E., Blum, P., and et al.: *Proceedings of the Ocean Drilling Program, Initial Reports*, edited by: Jansen, E., Raymo, M. E., and Blum, P., Ocean Drilling Program, College Station, TX, <https://doi.org/10.2973/odp.proc.ir.162.1996>, 1996.

Jöhnck, J., Holbourn, A., Kuhnt, W., Andersen, N.: Oxygen isotope offsets in deep-water benthic foraminifera. *Journal of Foraminiferal Research*, 51, 225–244, <https://doi.org/10.2113/gsjfr.51.3.225>, 2021.

Kotov, S. and Pälike, H.: QAnalySeries- a cross-platform time series tuning and analysis tool, in: AGU Fall Meeting Abstracts, PP53D-1230, 2018.

920 Labeyrie, L., Waelbroeck, C., Cortijo, E., Michel, E., and Duplessy, J. C.: Changements de l'hydrologie profonde pendant la dernière déglaciation, *Comptes Rendus - Geoscience*, 337, 919–927, <https://doi.org/10.1016/j.crte.2005.05.010>, 2005.

Lamy, F., Gersonde, R., Winckler, G., Esper, O., Jaeschke, A., Kuhn, G., Ullermann, J., Martinez-Garcia, A., 925 Lambert, F., and Kilian, R.: Increased dust deposition in the Pacific Southern Ocean during glacial periods, *Science* (1979), 343, 403–407, <https://doi.org/10.1126/science.1245424>, 2014.

Lamy, F., Winckler, G., Alvarez Zarikian, C. A., and Expedition 383 Scientists: Expedition 383 Preliminary Report: Dynamics of the Pacific Antarctic Circumpolar Current (DYNAPACC), International Ocean Discovery Program, <https://doi.org/10.14379/iodp.pr.383.2019>, 2019.

930 Laskar, J., Joutel, F., and Boudin, F.: Orbital, precessional, and insolation quantities for the Earth from -20 Myr to +10 Myr, *Astron Astrophys*, 270, 522–533, 1993.

Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *Astron Astrophys*, 428, 261–285, <https://doi.org/10.1051/0004-6361:20041335>, 2004.

935 Lee, T., Rand, D., Lisiecki, L. E., Gebbie, G., and Lawrence, C.: Bayesian age models and stacks: combining age inferences from radiocarbon and benthic  $\delta$  18 O stratigraphic alignment , *Climate of the Past*, 19, 1993–2012, <https://doi.org/10.5194/cp-19-1993-2023>, 2023.

Lin, L., Khider, D., Lisiecki, L. E., and Lawrence, C. E.: Probabilistic sequence alignment of stratigraphic records, *Paleoceanography*, 29, 976–989, <https://doi.org/10.1002/2014PA002713>, 2014.

940 Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta$  18O records, *Paleoceanography*, 20, 1–17, <https://doi.org/10.1029/2004PA001071>, 2005.

Lisiecki, L. E. and Raymo, M. E.: Diachronous benthic  $\delta$ 18O responses during late Pleistocene terminations, *Paleoceanography*, 24, <https://doi.org/10.1029/2009PA001732>, 2009.

945 Lisiecki, L. E., Jones, A. M., Rand, D., Lee, T., and Lawrence, C. E.: Comparing age model techniques for the last glacial cycle: A case study of ten Iberian Margin sediment cores, *Quat Sci Rev*, 287, <https://doi.org/10.1016/j.quascirev.2022.107559>, 2022.

Marchitto, T., Curry, W., Lynch-Stieglitz, J., Bryan, S., Cobb, K., and Lund, D.: Improved oxygen isotope temperature calibrations for cosmopolitan benthic foraminifera. *Geochemica et Cosmochimica Acta*, 130, 1-11, 2014.

950 Mankinen, E. A. and Dalrymple, G. B.: Revised geomagnetic polarity time scale for the interval 0-5 M.y.B.P., *J Geophys Res*, 84 B2, 615–626, <https://doi.org/10.1029/jb084ib02p00615>, 1979.

Marino, M., Maiorano, P., and Flower, B. P.: Calcareous nannofossil changes during the Mid-Pleistocene Revolution: Paleoecologic and paleoceanographic evidence from North Atlantic Site 980/981, *Palaeogeogr Palaeoclimatol Palaeoecol*, 306, 58–69, <https://doi.org/10.1016/j.palaeo.2011.03.028>, 2011.

955 Martínez-Garcia, A., Rosell-Melé, A., Jaccard, S. L., Geibert, W., Sigman, D. M., and Haug, G. H.: Southern Ocean dust-climate coupling over the past four million years, *Nature*, 476, 312–315, <https://doi.org/10.1038/nature10310>, 2011.

Martinson, D. G., Pisias, N. G., Hays, J. D., Imbrie, J., Moore, T. C., and Shackleton, N. J.: Age Dating and the Orbital Theory of the Ice Ages: Development of a High-Resolution 0 to 300,000-Year Chronostratigraphy, *Quat Res*, 27, 1–29, 1987.

960 McManus, J. F., Oppo, D. W., and Cullen, J. L.: A 0.5-Million-Year Record of Millennial-Scale Climate Variability in the North Atlantic, *Science* (1979), 283, 971–975, 1999.

Oppo, D. W., McManus, J. F., and Cullen, J. L.: Abrupt climate events 500,000 to 340,000 years ago: Evidence from subpolar North Atlantic sediments, *Science* (1979), 278, 1335–1338, 1998.

965 Paillard, D., Labeyrie, L., and Yiou, P.: Analyseries 1.0: a Macintosh software for the analysis of geographical time-series, *Eos (Washington DC)*, 77, 379, 1996.

Pisias, N. G., Martinson, D. G., Moore, T. C., Shackleton, N. J., Prell, W., Hays, J., and Boden, G.: High resolution stratigraphic correlation of benthic oxygen isotopic records spanning the last 300,000 years, *Mar Geol*, 56, 119–136, 1984.

970 Ravelo, A. C., Andreasen, D. H., Lyle, M., Lyle, A. O., and Wara, M. W.: Regional climate shifts caused by gradual global cooling in the Pliocene epoch, 2004.

Raymo, M. E., Ruddiman, W. F., Backman, J., Clement, B. M., and Martinson, D. G.: Late Pliocene variation in Northern Hemisphere ice sheets and North Atlantic deep water circulation, *Paleoceanography*, 413–446 pp., 1989.

975 Raymo, M. E., Ruddiman, W. F., Shackleton, N. J., and Oppo, D. W.: Evolution of Atlantic-Pacific d13C gradients over the last 2.5 M.y., *Earth and Planetary Science Letters*, 353–368 pp., 1990.

Raymo, M. E., Oppo, D. W., Flower, B. P., Hodell, D. A., McManus, J. F., Venz, K. A., Kleiven, K. F., and McIntyre, K.: Stability of North Atlantic water masses in face of pronounced climate variability during the Pleistocene, *Paleoceanography*, 19, <https://doi.org/10.1029/2003PA000921>, 2004.

980 Ruddiman, W. F., Raymo, M. E., Martinson, D. G., Clement, B. M., and Backman, J.: Pleistocene evolution: Northern hemisphere ice sheets and North Atlantic Ocean, *Paleoceanography*, 4, 353–412, <https://doi.org/10.1029/PA004i004p00353>, 1989.

985 Shackleton, N. J.: Attainment of isotopic equilibrium between ocean water and benthonic foraminifera genus *Uvigerina*: isotopic changes in the ocean during the last glacial. *Les méthodes quantitatives d'étude des variations du climat au cours du Pleistocène*, Gif-sur-Yvette, Colloque international du CNRS, 219, 203–210, 1974.

990 Shackleton, N. J. and Hall, M. A.: Oxygen and carbon isotope stratigraphy of DSDP Hole 552A: Plio-Pleistocene glacial history, *Initial Reports of the Deep Sea Drilling Project*, 81, 599–609, 1984.

Shackleton, N. J. and Opdyke, N. D.: Oxygen isotope paleomagnetic stratigraphy of Equatorial Pacific core V28-238: Oxygen isotope temperatures and ice volumes on a  $10^5$  year and  $10^6$  year scale, *Quat Res*, 3, 39–55, 1973.

Shackleton, N. J., Berger, A., and Peltier, W. R.: An alternative astronomical calibration of the lower Pleistocene timescale based on ODP Site 677, *Trans R Soc Edinb Earth Sci*, 81, 251–261, <https://doi.org/10.1017/S0263593300020782>, 1990.

995 Shackleton, N. J., Crowhurst, S., Hagelberg, T., Pisias, N. G., and Schneider, D. A.: A new late Neogene time scale: Application to Leg 138 sites, *Proceedings of the Ocean Drilling Program Scientific Results*, 138, 73–101, 1995.

Skinner, L. C. and Shackleton, N. J.: Rapid transient changes in northeast Atlantic deep water ventilation age across Termination I, *Paleoceanography*, 19, <https://doi.org/10.1029/2003PA000983>, 2004.

1000 Skinner, L. C. and Shackleton, N. J.: An Atlantic lead over Pacific deep-water change across Termination I: Implications for the application of the marine isotope stage stratigraphy, *Quat Sci Rev*, 24, 571–580, <https://doi.org/10.1016/j.quascirev.2004.11.008>, 2005.

Stein, M., Wasserburg, G. J., Aharon, P., Chen, J. H., Zhu, Z. R., Bloom, A., and Chappell, J.: TIMS U-series dating and stable isotopes of the last interglacial event in Papua New Guinea, *Geochimica Cosmochimica Acta*, 1005 57, 2541–2554, 1993.

Stern, J. V. and Lisiecki, L. E.: Termination 1 timing in radiocarbon-dated regional benthic  $\delta^{18}\text{O}$  stacks, *Paleoceanography*, 29, 1127–1142, <https://doi.org/10.1002/2014PA002700>, 2014.

Suman, D. O. and Bacon, M. P.: Variations in Holocene sedimentation in the North American Basin determined from  $^{230}\text{Th}$  measurements, *Deep-Sea Research*, 36, 869–878, 1989.

1010 1010 Thompson, W. G. and Goldstein, S. L.: A radiometric calibration of the SPECMAP timescale, *Quat Sci Rev*, 25, 3207–3215, <https://doi.org/10.1016/j.quascirev.2006.02.007>, 2006.

Tiedemann, R. and Franz, S. O.: Deepwater circulation, chemistry, and terrigenous sediment supply in the equatorial Atlantic during the Pliocene, 3.3–2.6 Ma and 5–4.5 Ma, *Proceedings of the Ocean Drilling Program, Scientific Results*, 154, 299–318, 1997.

1015 1015 Ullermann, J., Lamy, F., Ninnemann, U., Lembke-Jene, L., Gersonde, R., and Tiedemann, R.: Pacific-Atlantic Circumpolar Deep Water coupling during the last 500 ka, *Paleoceanography*, 31, 639–650, <https://doi.org/10.1002/2016PA002932>, 2016.

Venz, K. A. and Hodell, D. A.: New evidence for changes in Plio-Pleistocene deep water circulation from Southern Ocean ODP Leg 177 Site 1090, *Paleogeography, Paleoclimatology, Paleoecology*, 182, 197–220, 2002.

1020 1020 De Vleeschouwer, D., Vahlenkamp, M., Crucifix, M., and Pälike, H.: Alternating Southern and Northern Hemisphere climate response to astronomical forcing during the past 35 m.y., *Geology*, 45, <https://doi.org/10.1594/PANGAEA>, 2017.

Waelbroeck, C., Skinner, L. C., Labeyrie, L., Duplessy, J. C., Michel, E., Vazquez Riveiros, N., Gherardi, J. M., and Dewilde, F.: The timing of deglacial circulation changes in the Atlantic, *Paleoceanography*, 26, <https://doi.org/10.1029/2010PA002007>, 2011.

1025 1025 Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnet, J. S. K., Bohaty, S. M., De Vleeschouwer, D., Florindo, F., Frederichs, T., Hodell, D. A., Holbourn, A. E., Kroon, D., Lauretano, V., Littler, K., Lourens, L. J., Lyle, M., Pälike, H., Röhl, U., Tian, J., Wilkens, R. H., Wilson, P. A., and Zachos, J. C.: An astronomically dated record of Earth's climate and its predictability over the last 66 million years, *Science* (1979), 369, 1383–1388, <https://doi.org/10.1126/SCIENCE.ABA6853>, 2020.

1030 1030 Wilkens, R. H., Westerhold, T., Drury, A. J., Lyle, M., Gorgas, T., and Tian, J.: Revisiting the Ceara Rise, equatorial Atlantic Ocean: Isotope stratigraphy of ODP Leg 154 from 0 to 5 Ma, *Climate of the Past*, 13, 779–793, <https://doi.org/10.5194/cp-13-779-2017>, 2017.

Winckler, G., Lamy, F., Alvarez Zarikian, C. A., Arz, H. W., Basak, C., Brombacher, A., Esper, O. M., Farmer, J. R., Gottschalk, J., Herbert, L. C., Iwasaki, S., Lawson, V. J., Lembke-Jene, L., Lo, L., Malinverno, E., Michel, E., Middleton, J. L., Moretti, S., Moy, C. M., Ravelo, A. C., Riesselman, C. R., Saavedra-Pellitero, M., Seo, I., Singh, R. K., Smith, R. A., Souza, A. L., Stoner, J. S., Venancio, I. M., Wan, S., Zhao, X., and Foucher McColl, N.: Site U1541 in Dynamics of the Pacific Antarctic Circumpolar Current, in: *Proceedings of the International*

1040 Ocean Discovery Program, vol. 383, edited by: Lamy, F., Winckler, G., Alvarez Zarikian, C. A., and Expedition  
383 Scientists, International Ocean Discovery Program, College Station, TX,  
<https://doi.org/10.14379/iodp.proc.383.105.2021>, 2021.