

California's 2023 snow deluge: contextualizing an extreme snow year against future climate change

Authors: Adrienne M. Marshall¹, John T. Abatzoglou², Stefan Rahimi^{3,4}, Dennis P. Lettenmaier⁵, Alex Hall⁴

1. Hydrologic Science and Engineering, Colorado School of Mines, Golden, CO 80401.
adriennemarshall@mines.edu
2. Management of Complex Systems, University of California Merced, Merced, CA 95343
3. Atmospheric Science, University of Wyoming, Laramie, WY, 82071
4. Center for Climate Science, University of California, Los Angeles, CA 90095
5. Department of Geography, University of California Los Angeles, Los Angeles, CA 90095

Corresponding author: Adrienne Marshall

Major Classification: Physical Sciences

Minor Classification: Earth, Atmospheric, and Planetary Sciences

Abstract: The increasing prevalence of low snow conditions in a warming climate has attracted substantial attention in recent years, but a focus exclusively on low snow leaves high snow years relatively underexplored. However, these large snow years are hydrologically and economically important in regions where snow is critical for water resources. Here we introduce the term “snow deluge” and use anomalously high snowpack in California's Sierra Nevada during the 2023 water year as a case study. Snow monitoring sites across the state had a median 41-year return interval for April 1 snow water equivalent (SWE). Similarly, a process-based snow model showed a 54-year return interval for statewide April 1 SWE (90% confidence interval: 38-109 years). While snow droughts can result from either warm or dry conditions, snow deluges require both cool and wet conditions. Relative to the last century, cool season temperature and precipitation associated with California's 2023 snow deluge were both moderately anomalous, while temperature was highly anomalous relative to recent climatology. Downscaled climate models in the SSP-370 scenario indicate that California snow deluges – which we define as the 20-year April 1 SWE event - are projected to decline with climate change (58% decline by late century), although less so than median snow years (73% decline by late century). This pattern occurs across the western U.S. Changes to snow deluge, and discrepancies in snow deluge and median snow year changes, could impact water resources and ecosystems. Understanding these changes is therefore critical to appropriate climate adaptation.

Significance Statement: Snow in mountainous regions is critical for water resources and is declining due to climate warming. While low snow years have been extensively studied, we know relatively little about large snow years and their potential changes. Here, we introduce the term “snow deluge” to describe extreme snow years and show that the 2023 California snow deluge was roughly a 1-in-54 year event, and was a 1-in-320 year event (or greater) at 5% of snow monitoring stations. Snow deluges are projected to decline across the western U.S. in future climates, although less so than median snow years. Snow deluges can be both destructive and beneficial. Better understanding the phenomenon and its potential changes could improve management of snow-dependent ecosystems and economies.

Keywords: climate change, snow hydrology, hydroclimate

Introduction

In recent years, a mounting body of evidence has highlighted the importance of snow drought (1) - a period of anomalous low snowpack - and its consequences for water resources and ecosystem function (2). Snow droughts are becoming more common and will likely continue to do so with continued warming (3–7). However, an exclusive focus on snow drought precludes understanding of the other end of the spectrum: what we term here as “snow deluge” - that is, years in which unusually large quantities of snow water equivalent (SWE) accumulate on the land surface.

The 2023 water year in California presents a promising case study for the concept of snow deluge. California snow surveyors described the year’s April 1 SWE as at least the largest since 1952, though they highlighted challenges of comparisons due to changing observational networks (8). The scientific community highlighted the role of atmospheric rivers and uncertainty of potential climate change attribution (9). Immediate impacts of the years’ snow deluge included widespread flooding, with subsequent downstream impacts to communities and agriculture; increases in State Water Project deliveries; and rollbacks of emergency drought provisions (8).

Previous work has shown that very large snow accumulations can occur in years with a high frequency of atmospheric rivers in the western U.S. (10, 11), but most work on extreme snowfall has focused on the event scale rather than seasonal accumulations. Individual extreme snowfall events account for the majority of interannual snowfall variability and have regionally varying correlations with ENSO (12). Both theory and climate models project that extreme snowfall events will decline less than smaller snowfall events, and could even increase in colder regions (13, 14). Other work on large annual precipitation accumulations (pluvials) has focused on total precipitation, rather than snowfall or snow accumulation (15–19). Probabilities of large annual snow accumulations are substantially different from those of precipitation, particularly in the snow-to-rain transition zone that encompasses a large fraction of western watersheds (20). The lack of existing literature on future changes in extreme snowfall years leaves open the question of the probabilities and impacts of extreme snow accumulations on an annual scale.

The term “snow drought” was apparently first used by Wiesnet in 1981 (21), who published a brief (two-paragraph) commentary noting that snow-covered area in January of that year was at an all-time low since the inception of the satellite record. Since then, definitions have proliferated, to the extent that Gottlieb & Mankin (2) noted that the term is “amorphous” and ill-defined (Text S1). Snow droughts are commonly defined as “warm” or “dry” snow droughts based on the temperature and precipitation anomalies (1). In defining snow deluge, we consider the extent to which wet or cool conditions produce snow deluges. We hypothesize that snow deluges likely require precipitation anomalies high enough and temperature anomalies cool enough (22) to produce anomalous snowfall or snowpack, while snow droughts may be created by either warm temperatures or low precipitation alone. Individual snow deluge years may nevertheless be primarily driven by relatively large seasonal cold or wet anomalies contingent on the climatological context of the region of interest. As with snow droughts, these climate drivers could affect the spatial distribution and heterogeneity of snow deluge conditions (6).

In this study, we introduce the concept of snow deluge and use the 2023 water year in California as a case study. We ask: (1) How unusual was the 2023 California snow deluge in the context of the modern observational record? (2) To what extent are snow deluges - including the 2023 water year - driven by relatively cold temperatures or high precipitation? (3) How will snow deluges change in projected future climates? We calculate spatially distributed return

intervals of April 1 SWE to ascertain the approximate likelihood of the conditions in the 2023 snow year, using both in situ snowpack observations (Figure S1) and estimates modeled with the Variable Infiltration Capacity (VIC) model (23), using a Generalized Extreme Value (GEV) theoretical approach. We evaluate the extent to which temperature and precipitation conditions in snowy regions of California were anomalous in 2023 using a long-term climate record (24); we specifically consider the importance of wet-day and dry-day temperature anomalies (25) in addition to average cool season temperatures. We also evaluate the extent to which the snow deluge of 2023 was rare relative to three previous decades versus the previous century. Finally, we use bias-corrected dynamically downscaled simulations from the Sixth Coupled Model Intercomparison Project (CMIP6) (26) to estimate potential changes in snow deluges relative to median years, and to evaluate the change in probability of the 2023 snow deluge within future climate contexts.


Results

Characterizing the 2023 California snow deluge

In situ snow courses and automated snow sensors showed high values of April 1 SWE throughout the mountains of California, with the largest values in the central and southern Sierra (Figure 1a). The state-wide median across operational snow monitoring sites was 1.5 m (interquartile range; IQR = 0.8 m), with a maximum value of 2.9 m. Snow conditions were most anomalous in the central and southern Sierra, with return intervals at many sites exceeding 100 years (Figure 1b; see Figure S2 for study area map). In contrast, snow in the northern part of the state was relatively unexceptional. The median estimated return interval across sites was 41 years (IQR = 50 years); 5% of sites had return intervals exceeding 320 years. While the snow deluge was widespread across the Sierra Nevada, it also extended to other parts of the western U.S., with exceptionally high April 1 SWE observed in parts of Utah, southeast Idaho, Nevada, and the western slope of the Colorado Rockies commensurate with the path of numerous atmospheric rivers (Figure S3-S6).

In situ data also indicate that the 2023 snow deluge was more spatially heterogeneous than previous snow deluges. In 2023, more sites in California (42%) recorded their largest-ever April 1 SWE over their full period of record than any other year (Table S1). The next largest year by this metric was 1983, with 27% of active sites recording their largest April 1 SWE. However, 1983 had a far greater percentage of sites (92%) ranked within the top five years on record (Table S2). By this measure, 2023 was only the third largest on record, with 81% of California sites within the top five.

VIC simulations indicated that approximately 56 km³ of SWE was on the ground in California on April 1, 2023. This is 257% of the 1921-2023 simulated average and the largest April 1 SWE value in the 1921-2023 VIC simulations. Sierra Nevada April 1 SWE represented about 93% of total reservoir capacity draining that region (27). VIC simulations approximately reproduced the magnitude and return frequency of in situ April 1 SWE, but spatial distributions were somewhat different. The largest SWE accumulations were simulated around the southern part of the Sacramento River basin and the Central Lahontan, draining to Nevada (Figure 1c). Similarly, the highest estimated return intervals based on the VIC simulations were distributed slightly north of those in the observed data, although with comparable return intervals (Figure 1d). Peak SWE simulations from VIC were not the highest on record, but were otherwise comparably anomalous, with a California-total peak SWE of 59 km³, which is about 215% of the long-term average (Figure S7).



Climate context for the 2023 snow deluge

Climate data from the NCLimGrid product (23) aggregated over the region of California with at least 50 mm mean peak SWE over the VIC historical record (see Methods) indicates that the 2023 snow deluge was driven by relatively cold temperatures - particularly relative to the last

several decades - and moderately, though not exceptionally high precipitation (Figure 2a). Six snow deluges with a California-total April 1 SWE return interval greater than 20 years occurred between 1921-2023 (similar results are seen using peak SWE; Figure S8-S9). Relative to the full period of record, almost all deluge events were within the top 10th percentile of cool season (Nov-Mar) precipitation, with the exception of 1922, which had just above median precipitation (Table S3). Cool season temperatures were within the coldest 10th percentile for four of six deluges, while 1938 and 1983 had temperatures in approximately the 40th percentile. The requirement for relatively cool and wet conditions is also evident in Figure 2b; while there is large interannual variability in cool-season precipitation in California, all snow deluges fall above the mean precipitation and below the mean cool season average temperature. Wet-day and dry-day temperature anomalies were consistently negative in snow deluge years since 1952 when these data became available. An exception is the 1983 snow deluge, when wet-day temperature anomalies were in the 71st percentile (Table S3). In 2023, wet-day temperature anomalies had a slightly smaller departure from their average values suggesting that unusually cold storms were not specifically responsible for the deluge (Figure 2a). When aggregated to the large spatial scale of California and temporal scale of a full water year, cool-season temperature and precipitation appear to have a relatively strong influence on April 1 SWE, with about 80% of April 1 SWE variability explained as a linear function of temperature and precipitation (Table S4). Years with precipitation deluge but lacking snow deluge tended to have warmer January-March temperatures than snow deluge years, but did not have major differences in precipitation seasonality (Figures S10-S11).




Figure 2. Historical California snow deluge and associated climate. (a) Time series of spatially-averaged April 1 SWE; cool-season precipitation; and cool-season average temperature and dry-day and wet-day anomalies. Values are spatially averaged over locations with > 50 mm mean April 1 SWE over 1920-2023. Colors indicate the extent to which highlighted snow deluges are wet or cool relative to other snow deluge years. Dashed lines indicate mean values over the full period of record. **(b)** April 1 SWE as a function of cool-season (Nov-March) average temperature and total precipitation. The dashed horizontal and vertical lines in (a) and (b) denote the Nov-March mean temperature and cumulative precipitation averaged during 1952-2023 (the period of record over which all variables were available), respectively.

The climate variables contributing to the 2023 snow deluge had spatially heterogeneous anomalies, with colder dry-day temperature anomalies throughout the Sierra Nevada, and colder wet-day temperature anomalies in the northern part of the state where snow accumulations were least anomalous (Figure S12). Precipitation and wet-day anomalies were both greatest at high elevations in the central and southern Sierra Nevada (Figure S13). Average and dry-day temperature anomalies were colder at high elevations, while wet-day temperature anomalies were colder at lower elevations in the northern parts of the state. While these anomalies showed elevation dependence, the sparseness of observing networks at high elevations and geographic differences in anomalies suggest caution in interpreting these findings.

Return intervals for climate and snow deluge summed across California illustrate the extent to which the 2023 snow deluge was driven by unusually cool temperatures and high precipitation, as well as the importance of sampling uncertainty and hydroclimatic trends (Figure 3; Figure S14). Return intervals for 2023 precipitation were similar regardless of the period of record: relative to the 103-year period of record, we estimated a 20-year return interval (90% CI: 16-30 years); relative to the 1981-2010 period (selected to align with historical WRF simulations), we estimated a 15-year return interval (CI: 10-29 years; see Table S5 for results relative to a 1991-2020 historical period). In contrast, estimates of the infrequency of the cool temperatures of 2023 depended strongly on the period of record: relative to the 103-year period of record, we estimated a 31-year return interval (CI: 21-62 years); relative to the 1980-2010 distribution, we estimated a 610-year return interval (CI: 220-5.0x10³ years). These findings reflect non-stationarity in the temperature record over the last century associated with well documented warming trends. Interestingly, April 1 SWE reflected much less of this discrepancy between the full period and recent decades: relative to the full period of record, we estimate that April 1 SWE in 2023 was a 54-year event (CI: 38-109 years); relative to the 1981-2010 period, April 1 SWE was a 59-year event (CI: 40-370 years). Peak SWE had similar probabilities to April 1 SWE but with more of a change based on period of record; relative to the 103-year period 2023 peak SWE was a 42-year event (CI: 31-77 years) and a 54-year event (CI: 35-302 years) based on the 1981-2010 period (Figure S15).




Figure 3. Precipitation, temperature, and April 1 SWE return intervals. Estimated return intervals for cool-season (a) precipitation and (b) temperature from NCLimGrid; and (c) April 1 SWE from VIC. Shading indicates 90% CI for 1000 bootstrapped samples using $\frac{2}{3}$ of the data in each case, and color indicates the period of record from which samples are drawn. Horizontal gray dashed line denotes values for 2023; vertical lines denote associated return intervals. Gray

points show observations with empirical return intervals for the 103-year period; large black point denotes 2023.

Projected changes in snow deluge probabilities

We use dynamically downscaled GCM outputs (Table S6) from the CMIP6 project for the Shared Socioeconomic Pathway (SSP) 3.70 scenario to evaluate potential changes in snow deluge. Note that these are not directly comparable to the VIC simulations due to the use of a different land surface model (Noah-MP) in the Weather Research and Forecasting (WRF) simulations used for downscaling (See Figures S16-18 for a comparison of the two datasets). We fit the GEV separately on the SWE output from each of the nine dynamically downscaled GCMs, which somewhat circumvents biases inherited from choices of land surface models and parent GCMs. Two-year return interval snow years (approximately equivalent to the median) are projected to decline more on a percentage basis than 20-year snow deluges across California, though there is substantial inter-model variability (Figure S19). The 2-year event is projected to decline by 36% by mid-century and 73% by late-century (9-GCM median). In contrast, a 20-year event is projected to decline by 22% by mid-century and 58% by late century. For a 60-year event approximately matching the rarity of the 2023 snow deluge, percentage declines are similar to the 20-year event: by mid-century, these events are projected to decline by 21%, with a 59% decline by late century. Results are comparable for peak SWE (Figure S20). While the multi-GCM median shows a larger percentage decline for median years than deluges, that pattern is not consistent across all individual GCMs (Figure S21).

The finding that snow deluges will likely decline, but less so than median years, is consistent across the western U.S. (Figure 4). The 9-GCM median 2-year peak SWE events have the largest percentage declines in the lower elevation Sierra Nevada, northern Cascades, and lower elevation edges of the Rocky Mountains, with the smallest declines in the higher elevation Rocky Mountains in Colorado and Wyoming (Figure 4a). Spatial patterns are similar for 20-year return interval events. The percent declines in snow deluge years are predominantly smaller than percentage declines for median years: by mid-century, 92% of pixels have a smaller percentage decline in their snow deluge years than median years; by late-century, this rises to 98% of pixels (Figure 4b). Overall, these results suggest that while both median and deluge years will see declines in peak SWE, percentage declines in deluge years will be considerably smaller.




Figure 4. Projected changes in snow deluge. (a) WRF-simulated change in 2-year and 20-year return interval peak SWE values from historical (1981-2010) to mid-century (2031-2060) (top panels) and historical to late century (2070-2099) (bottom panels); median across 9 GCMs; (b) difference between 20-year and 2-year return interval percentiles for the same periods as in (a).

Discussion

April 1 SWE resulting from California's 2023 snow deluge had roughly a 54-year return interval, with considerable spatial variability in the observational record. While unusual, this is substantially less rare than the 2015 snow drought (28). The anomalous snow conditions were partially explained by spatially averaged cool temperatures and high precipitation, though these climate anomalies might not be collocated with the regions of greatest snow accumulation (Figure S14). Temporal averaging could also mask important variability: while wet- and dry-day temperature anomalies were similar in 2023, wet and dry day temperatures have been warming at different rates (25). Frequencies of compound climate extremes have also been changing (29, 30). These could create a non-stationary relationship between temperature anomalies and April 1 SWE. Future work could also evaluate the extent to which snow deluges the frequency and temporal clustering of atmospheric rivers and their synoptic rain-snow elevation relative to the land surface (10). Deluge formation could be impacted by winter melt prior to April 1 (31), which is in turn affected by variability in incoming shortwave radiation (32), snowfall intensity (33), atmospheric humidity (34), rain-on-snow events (35), and changes in snow albedo following disturbance events (36). Despite this range of factors impacting snow accumulation and melt dynamics, spatially aggregated cool-season temperature and precipitation explain at least 80% of variability in statewide April 1 SWE.

The projected smaller percentage decline of snow deluges relative to median years aligns with previous findings that large snowfall events are anticipated to decline less than median events, or even to increase in relatively cold locations (13, 14). This is fundamentally due to competing effects of warming, which decreases the fraction of precipitation falling as snow (20), and Clausius-Clapeyron effects, which increase precipitation intensity (14). Previous studies of this phenomena have been event-based, rather than annual. As the largest events are major contributors to annual snow accumulation (13), the corresponding result at the annual scale aligns with and extends previous findings. Because these changes are calculated relative to historical means, the smaller declines in deluges may or may not lead to an increase in interannual variability: indeed, while the smaller declines in deluges are consistent across the western U.S., interannual variability of peak SWE is projected to increase only in the highest elevations of the maritime regions and in colder interior continental snowpacks (3).

While some likely impacts of snow deluges are evident from existing literature, others are relatively poorly understood. The snow deluge of 2023 resulted in substantial flooding, particularly in the Tulare Lake basin (37); future work could evaluate the conditions under which snow deluges result in large or disruptive flooding. Also important is the extent to which snow deluge years result in high runoff and reservoir storage increases, including potential nonstationarities between snowfall and runoff in a warming climate (38). Indeed, the relative importance of snow deluge versus large total precipitation accumulations for total runoff and associated impacts is not particularly well understood (39). Impacts on reservoir recharge are even more uncertain and could depend on operations and the success of forecast-informed reservoir operations (40).

Snow deluges could also impact snow-dependent terrestrial and aquatic ecosystems. As snow droughts become more frequent yet perhaps remain punctuated by occasional deluges, we might consider two scenarios: in one, a snow-adapted forest ecosystem or river experiences persistent low snow years. In another scenario, these low snow years are occasionally interrupted by a deluge. Differences in ecosystem dynamics - including phenology and mortality rates (41) - between these systems could result in substantially different rates and types of landscape change. Such differences could also depend on underlying geology and capacity for soil profiles to store water throughout the critical zone (42). The relative loss of snow deluge could also be important for wildlife: while declining snowpacks have long motivated concerns for snow-dependent wildlife (43), very large snow events can also increase mortality for some species (44).

Snow deluge impacts on human communities could also be important. For instance, ski area economics could be substantially affected by occasional snow deluge years (45), though recent experience indicates that deluges can enhance winter recreation by extending the season, or reduce it due to infrastructure impacts. More widespread impacts to infrastructure (e.g., buildings, transport, utilities) and associated snow hazards during snow deluge events and rapid melt of such events can impact communities (46, 47). Summer recreation is impacted by the magnitude and timing of snow accumulation and melt in a preceding snow year (48, 49), although differences in human and ecological responses can cause phenological mismatches (50). Perhaps even more importantly, historical analysis of Colorado River negotiations finds a risk that high runoff years could breed complacency and delay planned adaptation (51); the impacts of such delays in adaptation response relative to benefits accrued from snow deluge are currently unknown. Snow deluges could also impact hydropower production (52); while many analyses of hydroclimate risks to the power grid have commonly focused on drought impacts

(53, 54), potential benefits or unknown drawbacks of snow deluges are relatively under characterized (55).

Multiple lines of evidence indicate that 2023 was exceptional in California with respect to April 1 SWE. While it was not alone in the historical record, it had more sites with record-breaking April 1 SWE than any other water year and a 54-year return interval aggregated over the state. Relative to only the last few decades, cool temperatures in winter and spring 2023 appear to be a truly exceptional driver of anomalous snow accumulations in 2023. However, against the backdrop of the last 103 years, both temperature and precipitation were unusual but not extreme, highlighting the importance of spatiotemporal variability, additional energy balance components, and compound extremes for generating snow deluge (56). Climate models indicate that snow deluges comparable to the 2023 California event will become increasingly rare, though will have smaller relative declines than typical years; such changes align with projections of increasing wet snow drought (7). Much as aridification has been posed as a more useful framing than drought (57), increasingly common conditions of snow drought may require that we pay new attention to the other end of the snow spectrum, with future work further exploring snow deluge and its consequences.

Materials and Methods

In situ April 1 SWE data

April 1 SWE observations were obtained from snow pillows and courses operated by the California Department of Water Resources (CA DWR) and the USDA NRCS. Snow courses in other states operated by other agencies were also obtained from the NRCS website (Figure S1; S3). Daily SWE data were gap filled using linear interpolation when five or fewer days were missing. This resulted in 0.9% (20/2314) of CA DWR site-years and 1 NRCS site-year (out of 23,687) having filled April 1 SWE data. Across the western U.S., 840 snow course sites operated by NRCS, 58 cooperator sites (including CA DWR), and 556 NRCS snow pillow sites had an April 1 SWE value in 2023 with at least 30 years of preceding data and were retained for analysis. In California, 215 snow course sites, 30 CA DWR snow pillows, and 23 NRCS snow pillows were retained. Following (58), no adjustment was made for cases when snow courses were obtained slightly before or after April 1. After data cleaning, snow courses had an average of 67 $\pm$ 18 (SD) years of data. CA DWR snow pillows had 40 $\pm$ 7 years of April 1 SWE observations, and NRCS snow pillow sites had 43 $\pm$ 5 years of data.

Modeled April 1 SWE

To provide a spatially complete estimate of the 2023 snow deluge, the Variable Infiltration Capacity (VIC) model was run using a “drought monitor” instantiation (59) that prioritizes homogeneity of a long record, with limited addition and removal of individual stations. Daily output from VIC covers the period 1921-2023 and was forced by 1/16th degree meteorological forcings (60) that have been extended through 2023 (61). VIC is a physically based hydrologic model that has shown good comparison with in-situ observations as well as with trends (62) and has comparable spatial patterns and SWE totals to other well-validated data products, though it slightly underestimates April 1 SWE relative to a higher-resolution product derived from satellite remote-sensing observations (63, 64; Figures S22-24; Text S3). VIC pixels were subset to locations with at least 50 mm average peak SWE simulated over the period of record (1921-2023). April 1 SWE values were extracted for each pixel, and statewide April 1 SWE values were summed across this region for each year.

Historical climate data

Monthly precipitation and average temperatures from the NClimGrid dataset were used to evaluate the extent to which snow deluges were driven by relatively cool or wet conditions (24). NClimGrid extends from 1895 to the present across the contiguous U.S. at a 1/24th degree resolution. NClimGrid is derived from Global Historical Climate Network-Daily stations, using substantial quality correction procedures and climatologically aided interpolation to develop the gridded product. Errors exist due to interpolation and sampling uncertainty but are well quantified (24). For each water year from 1921-present, average monthly temperature over November to March and total monthly precipitation over the same period were calculated for the masked snowy area of California derived from the VIC product. Throughout the manuscript, references to “temperature” or “precipitation” refer to Nov-March average and total, respectively. We interpret these values as estimates appropriate for relative comparisons and historical contextualization, rather than perfectly accurate values.

The daily NClimGrid product, available from water years 1952-2023, was used to evaluate the contributions of wet- and dry-day temperature anomalies and total wet-days. Following (25), the daily average temperature anomalies on wet and dry days were calculated and averaged over November-March. Daily NClimGrid was also used to obtain the number of wet days for each year.

Projected changes in snow deluge

Dynamically downscaled general circulation model (GCM) outputs from the Sixth Coupled Model Intercomparison (CMIP6) project (65) were used to contextualize the 2023 snow deluge against simulated historical and future climates, and to evaluate the expected change in snow deluges relative to median snow years using the methods of (66-68). Nine GCMs (Table S6) were bias corrected to the ERA5 (69) historical period for 1980-2010 prior to downscaling following the methods of (70). These bias-corrected GCMs were then used as forcing to the Weather Research and Forecasting (WRF) model (71), run at a 9-km grid length across the western U.S. WRF was coupled with the Noah-MP land surface model. Simulations were run for the historical period (1980-2013) and the Shared Socioeconomic Pathway (SSP) 3.70 period (2015-2099). A single scenario was used based on initial analyses indicating that inter-model variation was greater than inter-scenario variability, indicating that a suite of GCMs for a single scenario could adequately capture an appropriate range of uncertainty (72). The mid-century period was defined as 2031-2060, with a late-century period defined as 2070-2099. To evaluate projected changes in snow deluge, April 1 SWE and peak SWE were calculated for each year and GCM in the WRF dataset. As with VIC, statewide April 1 SWE was calculated by summing SWE over the snowy area of California, defined as grid cells that had at least 50 mm average peak SWE across 9 GCMs in the WRF record during model years 1980-2010.

Statistical Analysis

Return intervals for in situ and VIC-modeled April 1 SWE for both individual pixels and statewide aggregated data were estimated using a Generalized Extreme Value (GEV) distribution fit in the R programming language (73) using the extRemes package (74). Sampling uncertainty in the VIC record was calculated by bootstrapping 1000 samples, each with $\frac{2}{3}$ of the relevant period of record, and identified snow deluges as years with at least a 20-year return interval relative to the full period of record (see Text S2 for full details). For each of the three defined periods in the WRF dataset (historical, mid-century, and late-century), California-total April 1

and peak SWE values were calculated for 2-year, 20-year, and 60-year return intervals for each GCM, and median percentage changes in these return values are reported and mapped.

References

1. A. Harpold, M. Dettinger, S. Rajagopal, Defining Snow Drought and Why It Matters. *Eos* (2017) <https://doi.org/10.1029/2017EO068775> (June 28, 2023).
2. A. R. Gottlieb, J. S. Mankin, Observing, Measuring, and Assessing the Consequences of Snow Drought. *Bull. Am. Meteorol. Soc.* **103**, E1041–E1060 (2022).
3. A. M. Marshall, J. T. Abatzoglou, T. E. Link, C. J. Tennant, Projected Changes in Interannual Variability of Peak Snowpack Amount and Timing in the Western United States. *Geophys. Res. Lett.* **46**, 8882–8892 (2019).
4. L. S. Huning, A. AghaKouchak, Global snow drought hot spots and characteristics. *Proc. Natl. Acad. Sci.* **117**, 19753–19759 (2020).
5. R. R. Shrestha, B. R. Bonsal, J. M. Bonnyman, A. J. Cannon, M. R. Najafi, Heterogeneous snowpack response and snow drought occurrence across river basins of northwestern North America under 1.0°C to 4.0°C global warming. *Clim. Change* **164**, 40 (2021).
6. A. J. Heldmyer, N. R. Bjarke, B. Livneh, A 21st-Century perspective on snow drought in the Upper Colorado River Basin. *JAWRA J. Am. Water Resour. Assoc.* **59**, 396–415 (2023).
7. M. Cowherd, L. R. Leung, M. Girotto, Evolution of global snow droughts characteristics from 1850 to 2100. *Environ. Res. Lett.* (2023) <https://doi.org/10.1088/1748-9326/acd804> (June 5, 2023).
8. California Department of Water Resources, California’s Snowpack is Now One of the Largest Ever, Bringing Drought Relief, Flooding Concerns (2023) (July 12, 2023).
9. G. Conroy, What the science says about California’s record-setting snow. *Nature* (2023) <https://doi.org/10.1038/d41586-023-00937-x> (July 12, 2023).
10. B. Guan, N. P. Molotch, D. E. Waliser, E. J. Fetzer, P. J. Neiman, The 2010/2011 snow season in California’s Sierra Nevada: Role of atmospheric rivers and modes of large-scale variability. *Water Resour. Res.* **49**, 6731–6743 (2013).
11. B. Guan, N. P. Molotch, D. E. Waliser, E. J. Fetzer, P. J. Neiman, Extreme snowfall events linked to atmospheric rivers and surface air temperature via satellite measurements. *Geophys. Res. Lett.* **37** (2010).
12. A. C. Lute, J. T. Abatzoglou, Role of extreme snowfall events in interannual variability of snowfall accumulation in the western United States. *Water Resour. Res.* **50**, 2874–2888 (2014).
13. A. C. Lute, J. T. Abatzoglou, K. C. Hegewisch, Projected changes in snowfall extremes and interannual variability of snowfall in the western United States. *Water Resour. Res.* **51**, 960–972 (2015).
14. P. A. O’Gorman, Contrasting responses of mean and extreme snowfall to climate change. *Nature* **512**, 416–418 (2014).
15. D. Zamora-Reyes, *et al.*, The Unprecedented Character of California’s 20th Century Enhanced Hydroclimatic Variability in a 600-Year Context. *Geophys. Res. Lett.* **49**, e2022GL099582 (2022).
16. D. L. Swain, B. Langenbrunner, J. D. Neelin, A. Hall, Increasing precipitation volatility in twenty-first-century California. *Nat. Clim. Change* **8**, 427–433 (2018).
17. X. Huang, D. L. Swain, Climate change is increasing the risk of a California megaflood. *Sci. Adv.* **8**, eabq0995 (2022).

18. C. A. Woodhouse, K. E. Kunkel, D. R. Easterling, E. R. Cook, The twentieth-century pluvial in the western United States. *Geophys. Res. Lett.* **32** (2005).
19. R. S. Kangas, T. J. Brown, Characteristics of US drought and pluvials from a high-resolution spatial dataset. *Int. J. Climatol.* **27**, 1303–1325 (2007).
20. P. Z. Klos, T. E. Link, J. T. Abatzoglou, Extent of the rain-snow transition zone in the western U.S. under historic and projected climate. *Geophys. Res. Lett.* **41**, 4560–4568 (2014).
21. D. Wiesnet, Winter snow drought. *Eos Trans. Am. Geophys. Union* **62**, 137–137 (1981).
22. C. H. Luce, V. Lopez-Burgos, Z. Holden, Sensitivity of snowpack storage to precipitation and temperature using spatial and temporal analog models. *Water Resour. Res.* **50**, 9447–9462 (2014).
23. X. Liang, D. P. Lettenmaier, E. F. Wood, S. J. Burges, A simple hydrologically based model of land surface water and energy fluxes for general circulation models. *J. Geophys. Res.* **99**, 14415 (1994).
24. R. S. Vose, *et al.*, Improved Historical Temperature and Precipitation Time Series for U.S. Climate Divisions. *J. Appl. Meteorol. Climatol.* **53**, 1232–1251 (2014).
25. J. T. Abatzoglou, A. M. Marshall, A. C. Lute, M. Safeeq, Precipitation Dependence of Temperature Trends Across the Contiguous US. *Geophys. Res. Lett.* **49** (2022).
26. S. Rahimi, *et al.*, Evaluation of a Reanalysis-Driven Configuration of WRF4 Over the Western United States From 1980 to 2020. *J. Geophys. Res. Atmospheres* **127**, e2021JD035699 (2022).
27. State of California, CDEC Web Applications. *California Daily Statewide Reservoirs Report* (2023) (January 9, 2024).
28. S. Belmecheri, F. Babst, E. R. Wahl, D. W. Stahle, V. Trouet, Multi-century evaluation of Sierra Nevada snowpack. *Nat. Clim. Change* **6**, 2–3 (2016).
29. J. T. Abatzoglou, S. Z. Dobrowski, S. A. Parks, Multivariate climate departures have outpaced univariate changes across global lands. *Sci. Rep.* **10**, 3891 (2020).
30. S. Zhou, B. Yu, Y. Zhang, Global concurrent climate extremes exacerbated by anthropogenic climate change. *Sci. Adv.* **9**, eabo1638 (2023).
31. K. N. Musselman, N. Addor, J. A. Vano, N. P. Molotch, Winter melt trends portend widespread declines in snow water resources. *Nat. Clim. Change* **11**, 418–424 (2021).
32. K. N. Musselman, M. P. Clark, C. Liu, K. Ikeda, R. Rasmussen, Slower snowmelt in a warmer world. *Nat. Clim. Change* **7**, 214–219 (2017).
33. A. M. Marshall, T. E. Link, A. P. Robinson, J. T. Abatzoglou, Higher Snowfall Intensity is Associated with Reduced Impacts of Warming Upon Winter Snow Ablation. *Geophys. Res. Lett.* **47**, e2019GL086409 (2020).
34. A. A. Harpold, P. D. Brooks, Humidity determines snowpack ablation under a warming climate. *Proc. Natl. Acad. Sci.* **115**, 1215–1220 (2018).
35. K. N. Musselman, *et al.*, Projected increases and shifts in rain-on-snow flood risk over western North America. *Nat. Clim. Change* **8**, 808–812 (2018).
36. A. L. Koshkin, B. J. Hatchett, A. W. Nolin, Wildfire impacts on western United States snowpacks. *Front. Water* **4** (2022).
37. S. Hubler, M. Abramson, A Vast Lake Has Captivated California Where Farms Stood a Year Ago. *N. Y. Times* (2023) (August 24, 2023).
38. B. Livneh, A. M. Badger, Drought less predictable under declining future snowpack. *Nat. Clim. Change* **10**, 452–458 (2020).
39. B. L. Gordon, *et al.*, Why does snowmelt-driven streamflow response to warming vary? A data-driven review and predictive framework. *Environ. Res. Lett.* **17**, 053004 (2022).

40. C. J. Delaney, *et al.*, Forecast Informed Reservoir Operations Using Ensemble Streamflow Predictions for a Multipurpose Reservoir in Northern California. *Water Resour. Res.* **56**, e2019WR026604 (2020).
41. R. A. Slatyer, K. D. L. Umbers, P. A. Arnold, Ecological responses to variation in seasonal snow cover. *Conserv. Biol.* **36**, e13727 (2022).
42. M. L. Goulden, R. C. Bales, California forest die-off linked to multi-year deep soil drying in 2012–2015 drought. *Nat. Geosci.* **12**, 632–637 (2019).
43. J. J. Barsugli, *et al.*, Projections of Mountain Snowpack Loss for Wolverine Denning Elevations in the Rocky Mountains. *Earth's Future* **8**, e2020EF001537 (2020).
44. M. F. Rudolf, E. C. Wilson, J. N. Pauli, Anomalous snow events increase mortality for a winter-adapted species. *Can. J. Zool.* **100**, 574–582 (2022).
45. C. Wobus, *et al.*, Projected climate change impacts on skiing and snowmobiling: A case study of the United States. *Glob. Environ. Change* **45**, 1–14 (2017).
46. E. Cho, J. M. Jacobs, Extreme Value Snow Water Equivalent and Snowmelt for Infrastructure Design Over the Contiguous United States. *Water Resour. Res.* **56**, e2020WR028126 (2020).
47. A. Vajda, *et al.*, Severe weather affecting European transport systems: the identification, classification and frequencies of events. *Nat. Hazards* **72**, 169–188 (2014).
48. J. S. Jenkins, J. T. Abatzoglou, B. A. Peterson, Spring Snowpack Influences on the Volume and Timing of Spring and Peak-Season Overnight Visitation to Yosemite Wilderness. *Weather Clim. Soc.* **15**, 365–379 (2023).
49. A. M. Marshall, V. Butsic, J. Harte, The Phenology of Wilderness Use: Backcountry Recreation in a Changing Climate. *Weather Clim. Soc.* **10**, 209–223 (2018).
50. I. K. Breckheimer, *et al.*, Crowd-sourced data reveal social–ecological mismatches in phenology driven by climate. *Front. Ecol. Environ.* **18**, 76–82 (2020).
51. A. Sullivan, D. D. White, M. Hanemann, Designing collaborative governance: Insights from the drought contingency planning process for the lower Colorado River basin. *Environ. Sci. Policy* **91**, 39–49 (2019).
52. A. M. Marshall, J. M. Chen, Impacts of changing snowfall on seasonal complementarity of hydroelectric and solar power. *Environ. Res. Infrastruct. Sustain.* **2**, 021001 (2022).
53. S. Turner, N. Voisin, K. D. Nelson, V. C. Tidwell, “Drought impacts on hydroelectric power generation in the Western United States” (Pacific Northwest National Lab. (PNNL), Richland, WA (United States), 2022) <https://doi.org/10.2172/1887470> (October 11, 2022).
54. J. D. Kern, Y. Su, J. Hill, A retrospective study of the 2012–2016 California drought and its impacts on the power sector. *Environ. Res. Lett.* **15**, 094008 (2020).
55. E. Grubert, A. M. Marshall, Water for energy: Characterizing co-evolving energy and water systems under twin climate and energy system nonstationarities. *WIREs Water* **9**, e1576 (2022).
56. A. AghaKouchak, *et al.*, Climate Extremes and Compound Hazards in a Warming World. *Annu. Rev. Earth Planet. Sci.* **48**, 519–548 (2020).
57. J. T. Overpeck, B. Udall, Climate change and the aridification of North America. *Proc. Natl. Acad. Sci.* **117**, 11856–11858 (2020).
58. P. W. Mote, S. Li, D. P. Lettenmaier, M. Xiao, R. Engel, Dramatic declines in snowpack in the western US. *Npj Clim. Atmospheric Sci.* **1**, 1–6 (2018).
59. A. W. Wood, D. P. Lettenmaier, A Test Bed for New Seasonal Hydrologic Forecasting Approaches in the Western United States. *Bulletin of the American Meteorological Society* **87**, 1699–1712 (2006).

60. B. Livneh, *et al.*, A Long-Term Hydrologically Based Dataset of Land Surface Fluxes and States for the Conterminous United States: Update and Extensions. *J. Clim.* **26**, 9384–9392 (2013).
61. L. Su, *et al.*, Drought Variability over the Conterminous United States for the Past Century. *J. Hydrometeorol.* **22**, 1153–1168 (2021).
62. P. W. Mote, A. F. Hamlet, M. P. Clark, D. P. Lettenmaier, Declining mountain snowpack in western North America. *Bull. Am. Meteorol. Soc.* **86**, 39–49 (2005).
63. D. Li, D. P. Lettenmaier, S. A. Margulis, K. Andreadis, The Role of Rain-on-Snow in Flooding Over the Conterminous United States. *Water Resources Research* **55**, 8492–8513 (2019).
64. K. Yang, *et al.*, Intercomparison of snow water equivalent products in the Sierra Nevada California using airborne snow observatory data and ground observations. *Frontiers in Earth Science* **11** (2023).
65. V. Eyring, *et al.*, Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.* **9**, 1937–1958 (2016).
66. S. Rahimi, *et al.*, “An Overview of the Western United States Dynamically Downscaled Dataset (WUS-D3)” *Geosci. Model Dev. Preprint* <https://doi.org/10.5194/gmd-2023-162> (2024).
67. S. Rahimi, *et al.*, Dynamical downscaling should use physically explainable bias correction. *Geophys. Res. Lett.* (2023).
68. M. Risser, *et al.*, Is bias correction in dynamical downscaling defensible? *Geophys. Res. Lett.* (2023).
69. H. Hersbach, *et al.*, The ERA5 global reanalysis. *Q. J. R. Meteorol. Soc.* **146**, 1999–2049 (2020).
70. C. L. Bruyère, J. M. Done, G. J. Holland, S. Fredrick, Bias corrections of global models for regional climate simulations of high-impact weather. *Clim. Dyn.* **43**, 1847–1856 (2014).
71. C. Skamarock, *et al.*, A Description of the Advanced Research WRF Model Version 4.3 (2021) <https://doi.org/10.5065/1dfh-6p97> (September 15, 2023).
72. W. Krantz, D. Pierce, N. Goldenson, D. Cayan, Memorandum on Evaluating Global Climate Models for Studying Regional Climate Change in California (2021).
73. R Core Team, R: A Language and Environment for Statistical Computing (2021).
74. E. Gilleland, R. W. Katz, **extRemes 2.0**: An Extreme Value Analysis Package in R. *J. Stat. Softw.* **72** (2016).

Acknowledgements

AMM was partially supported by NSF Award #2241892 and the Alfred P. Sloan Foundation Award #2023-19665. JTA was partially supported by USDA NIFA Awards #2021-69012-35916 and 2021-67021-35344. SR was partially supported by the Department of Energy’s HyperFACETS (A23-1053-S003) project, the Strategic Environmental Research and Development Program (SERDP) Project, the California Energy Commission (EPC-20-006), and the University of California’s Climate Ecosystems Future (LRF-18-542511) project. We also thank the computational support through the NCAR-Wyoming Supercomputing Center (NWSC) and the Computational and Information Systems Laboratory (CISL). We thank Noah Molotch and one anonymous reviewer for constructive feedback that substantially improved this manuscript.

Data and materials availability: In situ April 1 SWE data are available from the Natural Resources Conservation Service (NRCS): <https://wcc.sc.egov.usda.gov/reportGenerator/>. VIC-

modeled April 1 and peak SWE are available at <https://zenodo.org/records/10602557>.
NClimGrid data is available from the National Oceanic and Atmospheric Administration (NOAA): <https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ncdc:C00332>. WRF simulations are available at: <https://aws.amazon.com/marketplace/pp/prodview-g4wqgpy2pa5dk#resources>.