

On a Nonlocal Integral Operator Commuting with the Laplacian and the Sturm-Liouville Problem: Low Rank Perturbations of the Operator

Lotfi Hermi^{1, a)} and Naoki Saito^{2, b)}

¹⁾Department of Mathematics and Statistics,
Florida International University,
Miami, FL, USA

²⁾Department of Mathematics,
University of California, One Shields Avenue,
Davis, CA 95616 USA

(*Electronic mail: saito@math.ucdavis.edu.)

(*Electronic mail: lhermi@fiu.edu.)

(Dated: 18 March 2024)

We reformulate all general real coupled self-adjoint boundary value problems as integral operators and show that they are all finite rank perturbations of the free space Green's function on the real line. This free space Green's function corresponds to the nonlocal boundary value problem proposed earlier by Saito [N. Saito, Appl. Comput. Harmonic Anal., **25**, 68–97 (2008)]. We prove these perturbations to be polynomials of rank up to 4. They encapsulate in a fundamental way the corresponding boundary conditions.

I. INTRODUCTION

In this article, we continue the program initiated in Ref. 1, and revisited in Ref. 2. In Ref. 1, Saito proposed the use the eigenfunctions of the integral operator commuting with the Laplace operator for analyzing functions and data defined on a general shape domain $\Omega \subset \mathbb{R}^d$. Computing eigenfunctions of the integral operator is simpler, more stable, and much faster with modern fast algorithms such as the Fast Multipole Method^{3,4} than directly solving the corresponding Laplacian eigenvalue problem (via the Helmholtz equation). This formulation imposes an interesting *nonlocal boundary condition in the Helmholtz equation formulation*, and we have been interested in investigating the nature of this boundary condition. Restricting the domain Ω to intervals in $\mathbb{R}$, but extending the kernel function from that of Saito's original proposal, we continued our analysis of the integral operators and their spectral properties².

This article addresses an old question of Gel'fand & Levitan⁵, namely the nature of the spectrum under perturbation. We treat this question in an integral operator setting. This formulation is expressed in terms of the Green's function, or fundamental solution, that corresponds to an elliptic operator. Green's functions are an important tool of Quantum Field Theory, Electrodynamics, Seismology, and Partial Differential Equations which found new applications in Machine Learning. In Quantum Field Theory, the Green's function is called the *propagator*⁶ or *two-point correlation function*⁷. It encodes the probability of measuring a field at a given point given its source at another. In Seismology, the Green's function plays a fundamental role in the solution of elastodynamic systems⁸. Integral operators are prominent in Machine Learning^{9–15}. They are ubiquitous in spectral clustering algorithms, kernel methods, and many manifold learning algorithms. Understanding their spectra is fundamental for various applications^{16–18}. Perturbation techniques are traditionally used to obtain closed forms of such kernels¹⁹. In this article we show that they can also reveal information about the boundary conditions more explicitly.

We focus on the one-dimensional setting of a *nonlocal boundary value problem defined by a symmetric kernel on a finite interval*. The problems we treat have roots in the work of Marcel Riesz^{20,21}, and in the work of Schoenberg^{22–24} in the discrete setting. A general formulation is given in Hellwig²⁵ (Chap. 4). We offer a unified alternative following Kato¹⁹ and more recent works by Gesztesy and his collaborators^{26–30}. The spectral framework proposed in this article provides an efficient and systematic way of obtaining the Green's function, especially in the case where $\lambda = 0$ is one of the eigenvalues (compare, e.g., with the Neumann boundary condition calculations in Porter & Stirling³¹), or the more recent Fucci *et al.*^{32,33} for a more general setting of the Kreĭn-von-Neumann eigenvalue problem. Our work aligns with Gesztesy & Kirsten^{29,30}. Our framework explains the complications in the expression of the Green's function in terms of Riesz projections.

In its simplest form, our operator of interest is a *nonlocal* operator defined by

$$\mathcal{H}f := -\frac{1}{2} \int_0^1 |x-y| f(y) dy. \quad (1)$$

^{a)}<https://faculty.fiu.edu/~lhermi>.

^{b)}<https://www.math.ucdavis.edu/~saito/>.

We are interested in the spectral properties of this operator. The eigenvalue problem associated with eq. (1), i.e., $\mathcal{K}u = (1/\lambda)u$, corresponds to the following *Sturm-Liouville eigenvalue problem*:

$$\begin{aligned} u'' + \lambda u &= 0 \quad \text{in } (0, 1), \\ u'(0) &= -u(0) - u(1), \\ u'(1) &= u(0) + u(1). \end{aligned} \tag{2}$$

We are also interested in the spectral properties of the general problem

$$\begin{aligned} u'' + \lambda u &= 0 \quad \text{in } (0, 1), \\ u'(0) &= \alpha u(0) + \beta u(1), \\ u'(1) &= \gamma u(0) + \delta u(1), \end{aligned} \tag{3}$$

where $\gamma = -\beta$ ($\neq 0$) to guarantee the self-adjointness of the operator (see Ref. 34, especially Sec. 3.5, 3.6). We let

$$C := \begin{pmatrix} \alpha & \beta \\ -\beta & \delta \end{pmatrix} \in \mathbb{R}^{2 \times 2}.$$

We observe that this boundary condition (BC) corresponds to the *General Self-Adjoint Real-Coupled* (“GSARC” for short) BC introduced in Refs. 35–37 (see also Refs. 38 and 39). For a given 2×2 matrix $C \in SL(2, \mathbb{R})$, i.e., the Special Linear Group acting on $\mathbb{R}$, the GSARC problem is also given by

$$\begin{pmatrix} u(1) \\ u'(1) \end{pmatrix} = B \begin{pmatrix} u(0) \\ u'(0) \end{pmatrix} \tag{4}$$

Note that the restriction $\det B = 1$ is imposed in the literature to simplify the problem. To see that eq. (4) follows from eq. (3), we calculate the matrix B , explicitly to obtain

$$B = \begin{pmatrix} -\frac{\alpha}{\beta} & \frac{1}{\beta} \\ -\frac{\beta^2 + \alpha\delta}{\beta} & \frac{\delta}{\beta} \end{pmatrix}$$

Remark I.1. The matrix B is sometimes written in terms of the sines and cosines to reflect the self-adjointness; see, e.g., the works of F. Gesztesy and his collaborators^{26,32,33}. To highlight this angular dependence, we note that one can indeed recast eq. (2) in the form eq. (3), using the KAN decomposition of B , viz.,

$$B = KAN \tag{5}$$

with $K = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$, $A = \begin{pmatrix} r & 0 \\ 0 & 1/r \end{pmatrix}$, and $N = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$, where $r > 0$, and $\theta, n \in \mathbb{R}$. (This is a unique decomposition called “the Iwasawa decomposition” in the Lie theory literature; it is a special case of QR factorization; see⁴⁰). The correspondence between eq. (5) and eq. (3) is explicitly given by

$$\alpha = \frac{\cos \theta + nr^2 \sin \theta}{nr^2 \cos \theta - \sin \theta}, \quad \beta = \frac{-r}{nr^2 \cos \theta - \sin \theta}, \quad \text{and} \quad \delta = \frac{r^2 \cos \theta}{nr^2 \cos \theta - \sin \theta}.$$

□

Remark I.2. When $\beta = -\gamma = 0$ in eq. (3) the problem corresponds to what can be termed as the *General Self-Adjoint Real-Separated* (“GSARS” for short) BC, detailed in the work of Folland³⁴ (Section 3.5, pp. 86-95). In the series of articles of Zettl and his co-authors^{36,37,41} on the geometric structures of spaces of boundary conditions, it sometimes takes the form

$$\begin{aligned} u'(0) \sin \theta_0 + u(0) \cos \theta_0 &= 0, \\ u'(1) \sin \theta_1 + u(1) \cos \theta_1 &= 0, \end{aligned} \tag{6}$$

where $\theta_0, \theta_1 \in [0, \pi)$. These correspond to $\alpha = -\cot \theta_0$ and $\delta = -\cot \theta_1$ in our notation. We shall discuss these particular GSARS cases in this article. A general framework for spaces of boundary conditions for ‘separated’ and ‘coupled’ self-adjoint BCs for higher order Sturm-Liouville problems can be found in the literature.^{42–44}

Recent activity^{45–47}, following⁴⁸, has focused on minimal extension of operators and their properties. We adopt a more classical point of view,^{49,50} focusing on *lower rank perturbation* of eq. (1). We will show that in the most general setting, all known examples of the Sturm-Liouville problem are polynomial perturbations of eq. (1) by operators of rank at most 4. We believe the result to be new. This article proposes:

- A unified view of all known boundary value problems (BVPs) recasting them in terms of the integral operator formulation (e.g., Green's functions);
- All of these BVPs are equivalent — up to the finite dimensional perturbation — to the base nonlocal integral operator eq. (1). The BCs for all these problems are indeed encoded in the perturbation in the following sense: The nonlocal operator eq. (1) corresponds to the Green's function on the whole real line; introducing BCs on $[0, 1]$ amounts to introducing a perturbation on this free-space Green's function.
- A unified approach to obtain the Green's functions from the expansion of the resolvents. This is an expansion on old ideas due to Kato¹⁹.

The spectral questions explored here connect in a fundamental way with the works^{27,28,32,33}.

Remark I.3. *We are interested in trace or sum rules for spectral functions associated with $\mathcal{K}$ in eq. (1), emanating from our recent work². We plan to treat the closed forms for the iterated kernel and spectral functions corresponding to various Sturm-Liouville problems in our upcoming series of articles.* $\square$

Remark I.4. *Perturbation questions of the types treated here have been visited in the literature in applications to the Laplacian of a graph (in particular, a tree) and its connection with the distance matrix; see^{51,52}.* $\square$

The integral operator framework works very well for the nonlocal operator eq. (1). The literature has focused much on Dirichlet and Neumann BVPs; see, e.g.,³¹. However, for the other BVPs we deal with in this paper, it is more difficult to derive the corresponding Green's functions. We propose a unified approach to obtain the Green's functions from the expansion of the resolvents. This has been explored by Kato¹⁹, but not fully exploited yet as far as we know.

Remark I.5. *Table I summarizes the values of α , β , and δ for each BC we deal with in our article, including the periodic and anti-periodic cases. They are coupled BCs and part of a different setting in the literature.*^{36,37,41}

TABLE I. Discriminant and rank of perturbation for values of α, β, δ for various BCs; KvN=Krein-von-Neumann; GSARC=General Self-Adjoint Real-Coupled; GSARS=General Self-Adjoint Real-Separated; $\Delta := \delta - 2\beta - \alpha + \beta^2 + \alpha\delta$

Problem	Nonlocal	KvN	Dirichlet	Neumann	Robin	Radoux	GSARC	GSARC	GSARS	GSARS	Periodic	Anti-Periodic
Conditions	$\alpha = -1$ $\beta = -1$ $\delta = 1$	$\alpha = -1$ $\beta = 1$ $\delta = 1$	$\alpha \rightarrow \infty$ $\beta = 0$ $\delta \rightarrow \infty$	$\alpha = 0$ $\beta = 0$ $\delta = 0$	$\alpha \rightarrow \infty$ $\beta = 0$ $\delta = 0$	$\alpha \rightarrow \infty$ $\beta = 0$ $\delta = 1$	$\Delta \neq 0$	$\Delta = 0$	$\beta = 0$ $\Delta \neq 0$	$\beta = 0$ $\Delta = 0$	-	-
Value of Δ	4	0	∞	0	$-\infty$	1	$\neq 0$	0	$\neq 0$	0	-	-
Perturbation Rank	-	4	2	2	2	2	2	2, 3, or 4	2	2	3	1

The organization of this article is as follows. In section II we show the equivalence of the problems eq. (1) and eq. (2). In section III we show how to obtain the integral operator from the resolvent. In section IV we show that all the problems of the form eq. (3) are lower rank perturbations of eq. (1) of up to rank 4.

II. FROM THE INTEGRAL OPERATOR TO THE DIFFERENTIAL EQUATION

To the best of our knowledge the properties of the operator eq. (1) were first explored in a very general setting in the works of Marcel Riesz^{20,21}. For simplicity, let

$$\kappa(x, y) := -\frac{1}{2} |x - y|. \quad (7)$$

To see that eq. (1) and eq. (2) are equivalent, observe, as in M. Riesz (see formulas of²⁰ (p. 41) and²¹ (p. 73)), that the function

$$u(x) = \int_0^1 \kappa(x, y) f(y) dy \quad (8)$$

is a solution of the Poisson equation

$$-u''(x) = f(x). \quad (9)$$

Note that

$$-\frac{\partial^2 \kappa(x, y)}{\partial y^2} = \delta(x - y), \quad (10)$$

where δ denotes the Dirac delta function. By eq. (9)

$$\begin{aligned} u(x) &= \int_0^1 \kappa(x, y) f(y) dy \\ &= - \int_0^1 \kappa(x, y) u''(y) dy \\ &= - \int_0^1 \frac{\partial^2 \kappa(x, y)}{\partial y^2} u(y) dy + \left[-u'(y) \kappa(x, y) + u(y) \frac{\partial \kappa}{\partial y} \right]_0^1 \\ &= u(x) + \left[-u'(y) \kappa(x, y) + u(y) \frac{\partial \kappa}{\partial y} \right]_0^1. \end{aligned} \quad (11)$$

Therefore, it must satisfy the BC

$$\left[-u'(y) \kappa(x, y) + u(y) \frac{\partial \kappa}{\partial y} \right]_0^1 = 0.$$

This means

$$-u'(1) \kappa(x, 1) + u(1) \frac{\partial \kappa}{\partial y} \Big|_{y=1} = -u'(0) \kappa(x, 0) + u(0) \frac{\partial \kappa}{\partial y} \Big|_{y=0}$$

and reduces to

$$-\frac{1}{2} u'(1) (x - 1) + \frac{1}{2} u(1) = \frac{1}{2} u'(0) x - \frac{1}{2} u(0)$$

for any $x \in [0, 1]$. Equating the coefficients leads the BCs in eq. (2).

III. FROM THE RESOLVENT TO THE INTEGRAL OPERATOR

The standard procedure to obtain the integral operator corresponding to a differential operator is to find the fundamental solution, or Green's function, first. Then the integral operator is just the formal solution of the problem. In our setting above, the fundamental solution of eq. (2) is $\kappa(x, y)$ (viz. eq. (9)) satisfying its BCs. Reversing the steps of eq. (11) proves this point.

Rather than following this procedure, we propose a unified framework for GSARC problems. They cover $SL(2, \mathbb{R})$ cases, and the prominent periodic and anti-periodic cases, which do not follow the GSARC setting. The various BCs are summarized in table I.

We propose a framework where the calculations proceed via the *resolvent*. The Green's function is then obtained from the expansion of this resolvent. We note four well-known examples appearing in the literature: Stakgold and Holst⁵³ (Example 1, pp. 416–420, eigenvalue problem (7.1.28)); Kato¹⁹ (Example 6.21, p. 183; Example 4.14, p. 293; and Example 1.4, p. 367). In these examples, the resolvent is calculated explicitly, but the full extent of the method we are proposing in this article is not exploited.

The proposed approach exploits in a fundamental way the Taylor and/or Laurent expansion of the resolvent function (or Neumann series) corresponding to linear operators. This approach aligns with the recent work of Gesztesy and Kirsten^{26,29,30}. Our procedure explicitly calculates expressions of these operators for all GSARC problems. This method turns out to be an efficient tool for obtaining closed forms of the zeta spectral function in terms of the iterated Brownian bridge⁵⁴.

A. Review of resolvents and Green's functions

For relevant materials, we cite the standard literature on the properties of the resolvent, and provide the necessary background we will need in this article; see Yosida⁵⁵ (Chap. 8); Kato¹⁹ (Chap. 1; Chap. 10); Dunford & Schwartz⁵⁶ (Chap. 7); and Taylor & Lay⁵⁷ (Chap. 5). These expansions have roots in Nagumo's 1916 article⁵⁸; see also⁵⁹. Let A be a densely defined and closed linear operator of trace class on a Hilbert space $\mathcal{H}$ with domain $\mathcal{D}(A)$ and range $\mathcal{R}(A)$ in $\mathcal{H}$. For the applications we envision in this paper, the *spectrum* consists of a discrete set of eigenvalues, $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \leq \dots$ accumulating to ∞ , i.e., $\sigma(A) = \{\lambda_k\}_{k=1}^\infty$. This is certainly the case of all the problems defined by the eigenvalue problem eq. (3). Finally, let $\rho(A) = \mathbb{C} \setminus \sigma(A)$ be the *resolvent set* of A , and $T = A^{-1}$ its inverse.

On $\rho(A)$, the *resolvent* $R(z) := (A - zI)^{-1}$ is a holomorphic function of z . Moreover, it admits the expansion (see, e.g.,⁵⁹ (p. 23; p. 193)):

$$\begin{aligned} R(z) &= (A(I - zA^{-1}))^{-1} \\ &= (I - zA^{-1})^{-1} A^{-1} \\ &= A^{-1} + A^{-2}z + A^{-3}z^2 + \dots \\ &= T + T^2z + T^3z^2 + \dots \end{aligned} \quad (12)$$

for $|z| \|A^{-1}\| = |z| \|T\| < 1$. Here I is the identity operator on $\mathcal{H}$. In the applications we propose T will correspond to the integral operator associated with A , while T^{k+1} is the associated iterated form of order k . The singularities of $R(z)$ are exactly the eigenvalues of A . If $z = 0$ is an eigenvalue, $R(z)$ admits the *Laurent series*

$$R(z) = \sum_{n=-\infty}^{\infty} z^n A_n. \quad (13)$$

The coefficients A_n are given by

$$A_n = \frac{1}{2\pi i} \oint_{C(0,\varepsilon)} z^{-n-1} R(z) dz. \quad (14)$$

where $C(0,\varepsilon)$ is a positively-oriented small circle of radius ε centered at $z = 0$, excluding all other eigenvalues of A . We note that the *Riesz projection* $P := -A_{-1}$. In more explicit terms, following Kato¹⁹ (Sec. III.6.5), if $z = 0$ is an isolated singularity, then

$$R(z) = -\frac{P}{z} - \sum_{n=1}^{\infty} \frac{D^n}{z^{n+1}} + \sum_{n=0}^{\infty} z^n S^{n+1}, \quad (15)$$

with

$$P := -\frac{1}{2\pi i} \oint_{C(0,\varepsilon)} R(z) dz, \quad D := AP = -\frac{1}{2\pi i} \oint_{C(0,\varepsilon)} z R(z) dz, \quad (16)$$

D is quasi-nilpotent with

$$D = DP = PD, \quad (17)$$

and

$$S := \frac{1}{2\pi i} \oint_{C(0,\varepsilon)} \frac{R(z)}{z} dz, \quad (18)$$

satisfying

$$AS = I - P, \quad SP = PS = 0. \quad (19)$$

The dimension of the range corresponding to P is the algebraic multiplicity, $m(T;0)$ of $z = 0$ ^{26,29,30}:

$$m(T;0) = \dim \mathcal{R}(P) = \text{tr}_{\mathcal{H}}(P). \quad (20)$$

When $z = 0$ is not an isolated singularity, $P \equiv 0$, $S = A^{-1} = T$ and we recover eq. (12).

B. Our one-dimensional setting

For the BVPs we deal with, we set $A = -\frac{d^2}{dx^2} + \text{BCs on } [0, 1]$. We present a unified approach to all GSARC problems. We also investigate a couple of non-GSARC problems that appear in the literature, namely the periodic and anti-periodic cases treated in³². Our results in this article focus on perturbation questions *not* pursued in^{26,29,30,32}.

For $z \in \rho(A)$, the resolvent $R(z)$ applied to a function $f(x)$ takes the form

$$R(z)f(x) = \left((A - zI)^{-1} f \right) (x) = \int_0^1 G(z, x, y) f(y) dy, \quad (21)$$

where $G(z, x, y)$ is the associated kernel. Explicit resolvent kernels for the various BCs considered in this paper are listed in table II. We will also denote by $G_0(x, y)$ the Green's function corresponding to the expansion eq. (15)

$$Sf(x) = (A^{-1} (I - P) f) (x) = \int_0^1 G_0(x, y) f(y) dy. \quad (22)$$

It is explicitly given in table III for the various BCs considered in this paper.

Remark III.1. *Though its study is relegated to our subsequent article, the iterated Green's function corresponding to S^{n+1} in eq. (15) is denoted by $G_n(x, y)$, viz.,*

$$S^{n+1}f(x) = \int_0^1 G_n(x, y) f(y) dy. \quad (23)$$

□

We end this discussion by deriving the condition under which eq. (3) admits $\lambda = 0$ as an eigenvalue (and hence leads to an isolated singularity leading to the Laurent series eq. (15) for the resolvent.) Clearly, for $\lambda = 0$, we are after the condition that all lines of the form $y = mx + b$ are non-trivial solutions of the BVP eq. (3) (with $\gamma = -\beta$). This means

$$\begin{aligned} m &= \alpha b + \beta(m + b) \\ m &= -\beta b + \delta(m + b) \end{aligned}$$

or

$$\begin{aligned} (1 - \beta)m - (\alpha + \beta)b &= 0 \\ (1 - \delta)m + (\beta - \delta)b &= 0. \end{aligned}$$

Hence the following secular equation must hold:

$$\begin{vmatrix} 1 - \beta & -(\alpha + \beta) \\ 1 - \delta & \beta - \delta \end{vmatrix} = 0. \quad (24)$$

Simplifying, we are led to the condition $\Delta := \delta - \alpha - 2\beta + \beta^2 + \alpha\delta = 0$. We call Δ the *discriminant* of the corresponding Sturm-Liouville problem.

Note that when $\beta = 1$, eq. (24) leads to $\alpha = -1$ or $\delta = 1$. This is a generalized instance of the Kreĭn-von Neumann problem where $\alpha = -1$, $\beta = 1$, and $\delta = 1$. In fact, for the BVPs we deal with, only the Kreĭn-von Neumann BVP belongs to the GSARC class with $\Delta = 0$ while the other BVPs belong to the GSARC class with $\Delta \neq 0$. The discriminant condition and various BCs, including the “separated” case ($\beta = 0$) are plotted in Fig. 1.

C. The Nonlocal Boundary Conditions eq. (2)

This is the eigenvalue problem eq. (2), which corresponds to the BVP eq. (3) with

$$C = \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix}$$




FIG. 1. Discriminant condition $\delta - \alpha - 2\beta + \beta^2 + \alpha\delta = 0$ corresponding to $\lambda = 0$ being an eigenvalue, with various BCs treated, including the discriminant condition for the “separated” BCs case ($\beta = 0$) indicated by the curve on the two sheets of the surface.

with discriminant $\Delta = 4$. Since $\lambda = 0$ is not an eigenvalue, its Riesz projection $P \equiv 0$. To obtain the expression of the resolvent kernel in table II, we first fix $0 \leq y \leq 1$, and let

$$u(x, y) := \begin{cases} u_1(x, y) & 0 \leq x \leq y \leq 1 \\ u_2(x, y) & 0 \leq y \leq x \leq 1 \end{cases} \quad (25)$$

be a solution of the Helmholtz equation

$$u'' + zu = -\delta(x - y) \quad (26)$$

with $'$ denote the derivative $\frac{d}{dx}$. Here, for simplicity, $G(z, x, y) = u(x, y)$, suppressing the dependence on z in the expressions of u_1 and u_2 . We now impose the BCs

$$\begin{aligned} u_1'(0, y) &= -u_1(0, y) - u_2(1, y) \\ u_2'(1, y) &= u_1(0, y) + u_2(1, y) \\ u_1(y^-, y) &= u_2(y^+, y) \\ u_2'(y^+, y) - u_1'(y^-, y) &= -1 \end{aligned} \quad (27)$$

TABLE II. Resolvent kernel $G(z, x, y)$ for various BCs; the formulas for GSARC with $\Delta = 0$, GSARS with $\Delta = 0$, and GSARS with $\Delta \neq 0$ are too long to list in this table; see appendix A, appendix C, and section IV J, respectively

Problem	Resolvent
Nonlocal (Saito)	$\frac{1}{\sqrt{z}(2+2\cos\sqrt{z}+\sqrt{z}\sin\sqrt{z})} \begin{cases} -\sqrt{z}\cos(\sqrt{z}x)\cos(\sqrt{z}(1-y))+2\cos\frac{\sqrt{z}}{2}\sin\left(\frac{\sqrt{z}}{2}(1+2x-2y)\right), & 0 \leq x \leq y \leq 1 \\ -\sqrt{z}\cos(\sqrt{z}y)\cos(\sqrt{z}(1-x))+2\cos\frac{\sqrt{z}}{2}\sin\left(\frac{\sqrt{z}}{2}(1+2y-2x)\right), & 0 \leq y \leq x \leq 1 \end{cases}$
Kreĭn-von Neumann	$\frac{1}{\sqrt{z}(-2+2\cos\sqrt{z}+\sqrt{z}\sin\sqrt{z})} \begin{cases} -\sqrt{z}\cos(\sqrt{z}x)\cos(\sqrt{z}(1-y))+2\sin\frac{\sqrt{z}}{2}\cos\left(\frac{\sqrt{z}}{2}(1+2x-2y)\right), & 0 \leq x \leq y \leq 1 \\ -\sqrt{z}\cos(\sqrt{z}y)\cos(\sqrt{z}(1-x))+2\sin\frac{\sqrt{z}}{2}\cos\left(\frac{\sqrt{z}}{2}(1+2y-2x)\right), & 0 \leq y \leq x \leq 1 \end{cases}$
Dirichlet ²⁸	$\frac{1}{\sqrt{z}\sin\sqrt{z}} \begin{cases} \sin(\sqrt{z}x)\sin(\sqrt{z}(1-y)), & 0 \leq x \leq y \leq 1 \\ \sin(\sqrt{z}y)\sin(\sqrt{z}(1-x)), & 0 \leq y \leq x \leq 1 \end{cases}$
Neumann	$-\frac{1}{\sqrt{z}\sin\sqrt{z}} \begin{cases} \cos(\sqrt{z}x)\cos(\sqrt{z}(1-y)), & 0 \leq x \leq y \leq 1 \\ \cos(\sqrt{z}y)\cos(\sqrt{z}(1-x)), & 0 \leq y \leq x \leq 1 \end{cases}$
Robin	$\frac{1}{(\alpha-\delta)z\cos\sqrt{z}-\sqrt{z}(z+\alpha\delta)\sin\sqrt{z}} \begin{cases} (\sqrt{z}\cos(\sqrt{z}x)+\alpha\sin(\sqrt{z}x))(\sqrt{z}\cos(\sqrt{z}(1-y))-\delta\sin(\sqrt{z}(1-y))), & 0 \leq x \leq y \leq 1 \\ (\sqrt{z}\cos(\sqrt{z}y)+\alpha\sin(\sqrt{z}y))(\sqrt{z}\cos(\sqrt{z}(1-x))-\delta\sin(\sqrt{z}(1-x))), & 0 \leq y \leq x \leq 1 \end{cases}$
Periodic	$-\frac{1}{2\sqrt{z}\sin\frac{\sqrt{z}}{2}} \begin{cases} \cos\left(\frac{\sqrt{z}}{2}(1+2x-2y)\right), & 0 \leq x \leq y \leq 1 \\ \cos\left(\frac{\sqrt{z}}{2}(1+2y-2x)\right), & 0 \leq y \leq x \leq 1 \end{cases}$
Anti-Periodic	$\frac{1}{2\sqrt{z}\cos\frac{\sqrt{z}}{2}} \begin{cases} \sin\left(\frac{\sqrt{z}}{2}(1+2x-2y)\right), & 0 \leq x \leq y \leq 1 \\ \sin\left(\frac{\sqrt{z}}{2}(1+2y-2x)\right), & 0 \leq y \leq x \leq 1 \end{cases}$
Radoux	$-\frac{1}{\sqrt{z}(\sqrt{z}\cos\sqrt{z}-\sin\sqrt{z})} \begin{cases} \sqrt{z}\sin(\sqrt{z}x)\cos(\sqrt{z}(1-y))-\sin(\sqrt{z}(1-y)), & 0 \leq x \leq y \leq 1 \\ \sqrt{z}\sin(\sqrt{z}y)\cos(\sqrt{z}(1-x))-\sin(\sqrt{z}(1-x)), & 0 \leq y \leq x \leq 1 \end{cases}$
GSARC, $\Delta \neq 0$	$\frac{1}{(-4\beta z+2z(-\alpha+\delta)\cos\sqrt{z}+2\sqrt{z}(z+\beta^2+\alpha\delta)\sin\sqrt{z})} \begin{cases} -(z+\beta^2+\alpha\delta)\cos(\sqrt{z}(1+x-y))+(-z+\beta^2+\alpha\delta)\cos(\sqrt{z}(1-x-y)) \\ +\sqrt{z}(-2\beta\sin(\sqrt{z}(x-y))+(-\alpha+\delta)\sin(\sqrt{z}(1+x-y))) \\ +(\alpha+\delta)\sin(\sqrt{z}(1-x-y))), & 0 \leq x \leq y \leq 1 \\ -(z+\beta^2+\alpha\delta)\cos(\sqrt{z}(1+y-x))+(-z+\beta^2+\alpha\delta)\cos(\sqrt{z}(1-y-x)) \\ +\sqrt{z}(-2\beta\sin(\sqrt{z}(y-x))+(-\alpha+\delta)\sin(\sqrt{z}(1+y-x))) \\ +(\alpha+\delta)\sin(\sqrt{z}(1-y-x))), & 0 \leq y \leq x \leq 1 \end{cases}$

TABLE III. Green's function $G(x, y)$ for various BCs; see appendix A and appendix C for the formulas of GSARC and GSARS with $\Delta = 0$

Problem	Green's function
Nonlocal (Saito)	$-\frac{1}{2} x-y $
Kreĭn-von Neumann	$-\frac{1}{2} x-y +\frac{2}{15}-\frac{3}{5}(x+y)+2(x^2+y^2)+\frac{6}{5}xy-3xy(x+y)-(x^3+y^3)+2xy(x^2+y^2)$
Dirichlet ²⁸	$-\frac{1}{2} x-y +\frac{1}{2}(x+y)-xy$
Neumann	$-\frac{1}{2} x-y +\frac{1}{3}-\frac{1}{2}(x+y)+\frac{1}{2}(x^2+y^2)$
Robin	$-\frac{1}{2} x-y -\frac{1}{2(\delta-\alpha+\alpha\delta)}(2-2\delta+(\alpha+\delta-\alpha\delta)(x+y)+2\alpha\delta xy)$
Periodic	$-\frac{1}{2} x-y +\frac{1}{12}+\frac{1}{2}(x-y)^2$
Anti-Periodic	$-\frac{1}{2} x-y +\frac{1}{4}$
Radoux	$-\frac{1}{2} x-y +\frac{1}{2}(x+y)-\frac{9}{5}xy+\frac{1}{5}xy(x^2+y^2)$
GSARC, $\Delta \neq 0$	$-\frac{1}{2} x-y +\frac{1}{\beta^2+\alpha\delta-\alpha+2\beta\delta}((1+\delta)+\frac{1}{2}(\beta^2+\alpha\delta-\alpha-\delta)(x+y)-(\beta^2+\alpha\delta)xy)$
GSARS, $\Delta \neq 0$	$-\frac{1}{2} x-y -\frac{(\cos\theta_0\cos\theta_1+\sin(\theta_0+\theta_1))(x+y)-2\gamma\cos\theta_0\cos\theta_1}{\cos(\theta_0-\theta_1)+\cos(\theta_0+\theta_1)-2\sin(\theta_0-\theta_1)}$

The last two equations of eq. (27) reflect, respectively, the continuity and jump condition at y (obtained by integrating eq. (26)). It is a common strategy⁶⁰ to set

$$\begin{aligned} u_1(x, y) &= a_1 \cos(\sqrt{z}x) + a_2 \sin(\sqrt{z}x), \quad 0 \leq x \leq y \leq 1; \\ u_2(x, y) &= b_1 \cos(\sqrt{z}x) + b_2 \sin(\sqrt{z}x), \quad 0 \leq y \leq x \leq 1, \end{aligned} \quad (28)$$

and then plug eq. (28) into eq. (27) to determine a_1, a_2, b_1 , and b_2 . These then lead to a solution $G(z, x, y)$, which depends on the spectral parameter z , as well as x and y :

$$G(z, x, y) = \frac{-\sqrt{z}\cos(\sqrt{z}x)\cos(\sqrt{z}(1-y))+2\cos\frac{\sqrt{z}}{2}\sin\left(\frac{\sqrt{z}}{2}(1+2x-2y)\right)}{\sqrt{z}(2+2\cos\sqrt{z}+\sqrt{z}\sin\sqrt{z})}, \quad (29)$$

for $0 \leq x \leq y \leq 1$, and by symmetry, the expression $G(z, y, x)$ when $0 \leq y \leq x \leq 1$. The poles of $G(z, x, y)$ as a function of z are exactly the eigenvalues of our differential operator. Note that $G_0(x, y) = \lim_{z \rightarrow 0^+} G(z, x, y) = \frac{1}{2}(x-y)$, for $0 \leq x \leq y \leq 1$,

thus recovering the expression of the integral operator eq. (1). Here, we use the convention of Kato¹⁹, for $0 \leq y \leq x \leq 1$, i.e., $G_0(x, y) = \frac{1}{2}(y - x)$ by symmetry. The full expressions for all BVPs considered in this paper are listed in tables II and III. Explicit expressions for the kernel of the Riesz projection, $p(x, y)$, are displayed in table IV for various conditions.

TABLE IV. Riesz Projection kernel for various BCs; the case of GSARC, $\Delta = 0$, $\alpha = -1$ reduces to a generalized Kreĭn-von Neumann problem (with δ as a free parameter); see remark III.8 for further details

Problem	Riesz Projection Kernel
Nonlocal (Saito)	0 (None)
Kreĭn-von Neumann	$-4 + 6x + 6y - 12xy$
Dirichlet ²⁸	0 (None)
Neumann	-1
Robin	0 (None)
Periodic	-1
Anti-Periodic	0 (None)
Radoux	$-3xy$
GSARC, $\Delta \neq 0$	0 (None)
GSARC, $\Delta = 0$ and $\alpha \neq -1$	$\frac{-3(1-\beta+(\alpha+\beta)x)(1-\beta+(\alpha+\beta)y)}{3+\alpha^2+\beta^2+3\alpha-3\beta-\alpha\beta}$
GSARS, $\Delta \neq 0$	0 (None)
GSARS, $\Delta = 0$	$\frac{6(x\cos\theta_0 - \sin\theta_0)(y\cos\theta_0 - \sin\theta_0)}{-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0}$

D. Kreĭn-von Neumann Boundary Conditions

This problem received special attention in the literature, both in one dimension, and in higher dimensions,^{27,28,45–48}, partially in connection with work on self-adjoint extensions of the operator $\mathcal{T}_\infty = -\frac{d^2}{dx^2}$. Note that the Kreĭn-von Neumann extension is the smallest (or soft) positive self-adjoint extension while the Friedrichs extension is the largest (or hard) positive self-adjoint extension. For the definitions and relevant characterizations we refer the reader to the cited literature as well as⁶¹.

In one dimension, it corresponds to the problem eq. (3) with

$$C = \begin{pmatrix} -1 & 1 \\ -1 & 1 \end{pmatrix}$$

with discriminant $\Delta = 0$. Strauss⁶⁰ (pg. 101, Exercise 12; pg. 145, Exercise 4) calls this problem an “unusual” eigenvalue problem. We owe this remark to M. Ashbaugh who brought it to our attention. Solving eq. (26) for u given by eq. (25) (and eq. (28)), and imposing the boundary, continuity, and jump conditions

$$\begin{aligned} u_1'(0, y) &= -u_1(0, y) + u_2(1, y) \\ u_2'(1, y) &= -u_1(0, y) + u_2(1, y) \\ u_1(y^-, y) &= u_2(y^+, y) \\ u_2'(y^+, y) - u_1'(y^-, y) &= -1 \end{aligned} \tag{30}$$

gives

$$G(z, x, y) = \frac{-\sqrt{z}\cos(\sqrt{z}x)\cos(\sqrt{z}(1-y)) + 2\sin\frac{\sqrt{z}}{2}\cos\left(\frac{\sqrt{z}}{2}(1+2x-2y)\right)}{\sqrt{z}(-2+2\cos\sqrt{z}+\sqrt{z}\sin\sqrt{z})}$$

for $0 \leq x \leq y \leq 1$, and by symmetry, the expression $G(z, y, x)$ when $0 \leq y \leq x \leq 1$. The kernel of the Riesz projection is given by

$$p(x, y) := \lim_{z \rightarrow 0^+} zG(z, x, y) = -4 + 6x + 6y - 12xy. \tag{31}$$

The quasi-nilpotent operator D defined in eq. (16) is null in this case because

$$\lim_{z \rightarrow 0^+} z^2 G(z, x, y) = 0.$$

The Green's function corresponding to this problem is obtained by finding the limit

$$\begin{aligned} G_0(x, y) &= \lim_{z \rightarrow 0^+} G(z, x, y) - \frac{p(x, y)}{z} \\ &= \frac{1}{30} (4 - 3x - 33y + 60x^2 + 60y^2 + 36xy - 30x^3 - 90x^2y - 90xy^2 - 30y^3 \\ &\quad + 60x^3y + 60xy^3) \end{aligned}$$

for $0 \leq x \leq y \leq 1$. A unified expression, valid for all $x, y \in [0, 1]$, and relating to the nonlocal operator appears in table III.

E. Dirichlet Boundary Conditions

This example is found in many sources, e.g.,^{1,19,28,31,50,62,63}. It is the easiest as it captures the essence of many features of GSARC problems. As a GSARC problem, this is the limiting case when $\alpha \rightarrow \infty$, $\beta = -\gamma = 0$, and $\delta \rightarrow \infty$ (e.g., $\alpha = \delta \rightarrow \infty$, thus $\Delta \rightarrow \infty$).

Solving eq. (26) for u given by eq. (25) (and eq. (28)), and imposing the boundary, continuity, and jump conditions

$$\begin{aligned} u_1(0, y) &= 0 \\ u_2(1, y) &= 0 \\ u_1(y^-, y) &= u_2(y^+, y) \\ u_2'(y^+, y) - u_1'(y^-, y) &= -1 \end{aligned} \tag{32}$$

yields

$$G(z, x, y) = \frac{\sin(\sqrt{z}x) \sin(\sqrt{z}(1-y))}{\sqrt{z} \sin \sqrt{z}}.$$

Note that $\lim_{z \rightarrow 0^+} z G(z, x, y) = 0$. Thus the Riesz projection $P \equiv 0$ and $G(z, x, y)$ admits a pure Taylor series at $z = 0$, from which we get

$$G_0(x, y) = \lim_{z \rightarrow 0^+} G(z, x, y) = x(1-y),$$

for $0 \leq x \leq y \leq 1$, and the symmetric expression for $0 \leq y \leq x \leq 1$. This fundamental solution assumes the unified form

$$G_0(x, y) = -\frac{1}{2}|x-y| + \frac{1}{2}(x+y) - xy,$$

valid for all $x, y \in [0, 1]$. The resolvent and Green's function are tabulated in tables II and III (see also^{19,25}).

F. Neumann Boundary Conditions

The expression for the Green's function $G_0(x, y)$ corresponding to this eigenvalue problem appear, for example, in^{1,31}. To obtain the resolvent kernel and Green's function expressions in tables II and III, we proceed as before, with the associated GSARC matrix $C \equiv 0$, and the discriminant $\Delta = 0$. For $0 \leq x \leq y \leq 1$, the resolvent kernel is explicitly given by

$$G(z, x, y) = -\frac{\cos(\sqrt{z}x) \cos(\sqrt{z}(1-y))}{\sqrt{z} \sin \sqrt{z}}.$$

The kernel of Riesz projection P is given by $p(x, y) = \lim_{z \rightarrow 0^+} z G(z, x, y) = -1$, while the nilpotent operator $D \equiv 0$ (since $\lim_{z \rightarrow 0^+} z^2 G(z, x, y) = 0$). The singularity $z = 0$ corresponds to a simple eigenvalue of the problem eq. (3). The Green's function corresponding to this problem is then

$$\begin{aligned} G_0(x, y) &= \lim_{z \rightarrow 0^+} G(z, x, y) - \frac{p(x, y)}{z} \\ &= \frac{1}{6} (2 - 6y + 3x^2 + 3y^2), \end{aligned}$$

for $0 \leq x \leq y \leq 1$. This agrees with the expressions in^{1,31}.

G. Robin Boundary Conditions

In the Robin boundary problem (named after the French mathematician Victor Gustave Robin; see, e.g.,^{64,65}), we still deal with the eigenvalue problem eq. (3) but now subject to (the separated) conditions where the flux at the boundary is proportional to its value, i.e., $u'(0) = \alpha u(0)$ and $u'(1) = \delta u(1)$. In the recent literature the problem is sometimes called the “Generalized Robin Problem”⁶⁶. We will first focus on the general case, then treat particular cases which received special attention in the literature^{19,53}. The discriminant is given by $\Delta = \delta - \alpha + \alpha\delta$. We will treat the case when $\Delta \neq 0$, relegating the null case to the most general setting. We solve for the resolvent kernel as before, to get, for $0 \leq x \leq y \leq 1$

$$G(z, x, y) = \frac{(\sqrt{z} \cos(\sqrt{z}x) + \alpha \sin(\sqrt{z}x)) (\sqrt{z} \cos(\sqrt{z}(1-y)) - \delta \sin(\sqrt{z}(1-y)))}{(\alpha - \delta) z \cos \sqrt{z} - \sqrt{z}(z + \alpha\delta) \sin \sqrt{z}},$$

and the symmetric expression $G(z, y, x)$ for $0 \leq y \leq x \leq 1$.

Since $\lim_{z \rightarrow 0^+} zG(z, x, y) = 0$, the Riesz projection P and nilpotent operator D are identically equal to zero. For $0 \leq x \leq y \leq 1$, the Green’s function corresponding to this problem is given by

$$\begin{aligned} G_0(x, y) &= \lim_{z \rightarrow 0^+} G(z, x, y) \\ &= -\frac{(1 + \alpha x)(1 - \delta(1 - y))}{\delta - \alpha + \alpha\delta}, \end{aligned}$$

and the symmetric expression $G_0(y, x)$ for $0 \leq y \leq x \leq 1$.

Example III.2. One of the simplest cases of the Robin BCs is: $u(0) = u'(1) = 0$, i.e., $\alpha \rightarrow \infty$ and $\delta \rightarrow 0$, $\Delta \rightarrow -\infty$, corresponding to Dirichlet BC at one end, and free Neumann BC at the other end. For $0 \leq x \leq y \leq 1$, the resolvent kernel is explicitly given by

$$G(z, x, y) = \frac{\sin(\sqrt{z}x) \cos(\sqrt{z}(1-y))}{\sqrt{z} \cos \sqrt{z}}.$$

The Green’s function corresponding to this problem is then

$$G_0(x, y) = \lim_{z \rightarrow 0^+} G(z, x, y) = x \quad \text{for } 0 \leq x \leq y \leq 1,$$

and the symmetric expression $G_0(y, x)$ for $0 \leq y \leq x \leq 1$. □

Example III.3. Another example appears in Kato’s work as Example 4.14¹⁹ (p. 293) where the BCs are given by $u(0) = 0$ and $\tau u'(1) - u(1) = 0$, and $0 < \tau < 1$ is a fixed parameter (our τ is the κ in Kato’s book). Note that $\alpha \rightarrow \infty$ and $\delta = 1/\tau$ in this example.

Hence,

$$G(z, x, y) = \frac{\sin(\sqrt{z}x)}{\tau z \cos \sqrt{z} - \sqrt{z} \sin \sqrt{z}} (\tau \sqrt{z} \cos(\sqrt{z}(1-y)) - \sin(\sqrt{z}(1-y))).$$

This expression agrees with the result in Kato’s book. For $0 \leq x \leq y \leq 1$, the Green’s function corresponding to this problem

$$G_0(x, y) = \lim_{z \rightarrow 0^+} G(z, x, y) = x - \frac{1}{1 - \tau} xy$$

and the symmetric expression for $0 \leq y \leq x \leq 1$. We note that this problem admits a unique negative eigenvalue; see¹⁹. □

Example III.4. Another variation of this Robin problem appears as Example 1.4¹⁹ (p. 367) where the BCs are replaced with $u(0) = 0$ and $u'(1) + \tau u(1) = 0$, i.e., $\alpha \rightarrow \infty$ and $\delta = -1/\tau$ in the general setting. Hence, we have

$$G(z, x, y) = \frac{\sin(\sqrt{z}x) (\sqrt{z} \cos(\sqrt{z}(1-y)) + \tau \sin(\sqrt{z}(1-y)))}{z \cos \sqrt{z} + \tau \sqrt{z} \sin \sqrt{z}}.$$

for $0 \leq x \leq y \leq 1$. This expression is also equivalent to Kato¹⁹. The Green’s function corresponding to this problem is given by

$$G_0(x, y) = \lim_{z \rightarrow 0^+} G(z, x, y) = x - \frac{\tau}{1 + \tau} xy \quad \text{for } 0 \leq x \leq y \leq 1. \quad (33)$$

Unlike example III.3, this problem does not exhibit negative eigenvalues. □

Example III.5. Yet another variation of this Robin eigenvalue problem also appears in the work of Stakgold and Holst⁵³ (Example 1, pp.416–420), with BCs of the form: $u(0) = 0$, and $u'(1) \sin \theta + u(1) \cos \theta = 0$. This is equivalent to example III.4 with $\tau = \cot \theta$ (or $\alpha \rightarrow \infty$ and $\delta = -\tan \theta$).

For $0 \leq x \leq y \leq 1$, and in terms of the angle θ , the resolvent kernel takes the form

$$G(z, x, y) = \frac{\sin(\sqrt{z}x) (\sqrt{z} \sin \theta \cos(\sqrt{z}(1-y)) + \cos \theta \sin(\sqrt{z}(1-y)))}{z \sin \theta \cos \sqrt{z} + \sqrt{z} \cos \theta \sin \sqrt{z}}$$

while, the Green's function corresponding to the integral operator is then given by

$$G_0(x, y) := \lim_{z \rightarrow 0^+} G(z, x, y) = x - \frac{1}{1 + \tan \theta} xy. \quad (34)$$

These expressions are equivalent to those appearing in⁵³. □

Remark III.6. For $\delta - \alpha + \alpha\delta = 0$, i.e., $\delta = \frac{\alpha}{\alpha+1}$ or $\alpha = \frac{\delta}{1-\delta}$, the Riesz projection P has a kernel given by

$$p(x, y) = -\frac{3(1+\alpha x)(1+\alpha y)}{3+3\alpha+\alpha^2}$$

while the nilpotent operator $D \equiv 0$. The Green's function corresponding to this problem is then

$$\begin{aligned} G_0(x, y) &= \lim_{z \rightarrow 0^+} G(z, x, y) - \frac{p(x, y)}{z} \\ &= c_{00} + c_{10}x + c_{01}y + c_{20}x^2 + c_{11}xy + c_{02}y^2 + c_{30}x^3 + c_{21}x^2y \\ &\quad + c_{12}xy^2 + c_{03}y^3 + c_{31}x^3y + c_{13}xy^3 \end{aligned}$$

for $0 \leq x \leq y \leq 1$, where the coefficients are explicitly given in appendix A, with $\beta = 0$. The expression $3 + 3\alpha + \alpha^2 > 0$ (its discriminant is negative). It appears in the expression of the Riesz projection kernel, and in all the expressions of the c_{ij} coefficients. The Green's function $G_0(x, y)$ is a fourth-order polynomial in x and y .

H. Periodic Boundary Conditions

Historically, this is a well-studied eigenvalue problem, due to its tight connection to Fourier series⁶⁷. Although it is not of GSARC type, we include it for comparison sake with the work of³². In lieu of the BCs in eq. (3), we treat the eigenvalue problem with $u(0) = u(1)$ and $u'(0) = u'(1)$.

Solving the Helmholtz equation eq. (26) for u with these BCs, and imposing the continuity and jump conditions as before, leads, for $0 \leq x \leq y \leq 1$, to

$$G(z, x, y) = -\frac{\cos\left(\frac{\sqrt{z}}{2}(1+2x-2y)\right)}{2\sqrt{z} \sin \frac{\sqrt{z}}{2}}.$$

The kernel of Riesz projection P is given by $p(x, y) = \lim_{z \rightarrow 0^+} z G(z, x, y) = -1$. The nilpotent operator $D \equiv 0$ since $\lim_{z \rightarrow 0^+} z^2 G(z, x, y) = 0$.

The Green's function corresponding to this periodic case is then

$$\begin{aligned} G_0(x, y) &= \lim_{z \rightarrow 0^+} G(z, x, y) - \frac{p(x, y)}{z} \\ &= \frac{1}{12} (1 + 6x - 6y + 6x^2 - 12xy + 6y^2), \end{aligned}$$

for $0 \leq x \leq y \leq 1$, and the corresponding expression by symmetry for $0 \leq y \leq x \leq 1$.

I. Anti-Periodic Boundary Conditions

This is one of the eigenvalue problems revisited in the recent article of Fucci, *et al.*³², though not an GSARC problem. The BCs are given by $u(0) = -u(1)$ and $u'(0) = -u'(1)$. The resolvent kernel ensuing from the proposed technique herein is given,

for $0 \leq x \leq y \leq 1$, by

$$G(z, x, y) = \frac{\sin\left(\frac{\sqrt{z}}{2}(1+2x-2y)\right)}{2\sqrt{z}\cos\frac{\sqrt{z}}{2}}$$

The kernel of Riesz projection $P \equiv 0$ since $p(x, y) = \lim_{z \rightarrow 0^+} z G(z, x, y) = 0$, and thus the nilpotent operator $D \equiv 0$.

The Green's function corresponding to the anti-periodic case is then

$$G_0(x, y) = \lim_{z \rightarrow 0^+} G(z, x, y) = \frac{1}{4}(1+2x-2y)$$

for $0 \leq x \leq y \leq 1$, and the corresponding expression by symmetry for $0 \leq y \leq x \leq 1$.

J. Radoux Boundary Conditions

The Radoux eigenvalue problem⁶⁸ appears in the literature in the context of obtaining sum rules for the roots for the transcendental equation $\tan x = x$. Sum rules are closed form expressions for the zeta function of these roots with a history that is traceable back to Rayleigh and others; see². Radoux used spectral techniques for solving eq. (3) with the BCs $u(0) = 0$, $u'(1) = u(1)$ to find these sum rules. This eigenvalue problem is a limiting case of section III G when $\alpha \rightarrow \infty$, and $\delta = 1$, (or when $\tau \rightarrow 1$ in example III.3). The discriminant $\Delta \rightarrow \infty$ and the formulas of section III G are no longer valid.

While the resolvent kernel is recoverable from this limit, the Green's function is not. Proceeding as before, we obtain

$$G(z, x, y) = \frac{\sin(\sqrt{z}x)(\sqrt{z}\cos(\sqrt{z}(1-y)) - \sin(\sqrt{z}(1-y)))}{z\cos\sqrt{z} - \sqrt{z}\sin\sqrt{z}}.$$

The kernel of Riesz projection P is given by $p(x, y) = \lim_{z \rightarrow 0^+} z G(z, x, y) = -3xy$, and the nilpotent operator $D \equiv 0$ since $\lim_{z \rightarrow 0^+} z^2 G(z, x, y) = 0$. For $0 \leq x \leq y \leq 1$, the Green's function is then

$$G_0(x, y) = \lim_{z \rightarrow 0^+} G(z, x, y) - \frac{p(x, y)}{z} = \frac{1}{10}(-10x + 18xy - 5x^3y - 5xy^3). \quad (35)$$

K. GSARC Boundary Conditions with Non-Zero Discriminant

We now focus on the general problem eq. (3), with $\Delta = \delta - \alpha - 2\beta + \beta^2 + \alpha\delta \neq 0$. The technique described yields a resolvent kernel where

The expression for

$$(-4\beta z + 2z(-\alpha + \delta)\cos\sqrt{z} + 2\sqrt{z}(z + \beta^2 + \alpha\delta)\sin\sqrt{z}) G(z, x, y)$$

is given by

$$\begin{aligned} & - (z + \beta^2 + \alpha\delta)\cos(\sqrt{z}(1+x-y)) + (-z + \beta^2 + \alpha\delta)\cos(\sqrt{z}(1-x-y)) \\ & + \sqrt{z}(-2\beta\sin(\sqrt{z}(x-y)) + (-\alpha + \delta)\sin(\sqrt{z}(1+x-y)) \\ & + (\alpha + \delta)\sin(\sqrt{z}(1-x-y))) \end{aligned}$$

for $0 \leq x \leq y \leq 1$, and thus by symmetry for full expression in table II. For $0 \leq x \leq y \leq 1$, the kernel of Riesz projection $P \equiv 0 \equiv D$ since $\lim_{z \rightarrow 0^+} z^n G(z, x, y) = 0$, for $n = 1, 2$. The Green's function is then

$$\begin{aligned} G_0(x, y) &= \lim_{z \rightarrow 0^+} G(z, x, y) \\ &= \frac{-1 + \delta - (\alpha + \beta)x + (\beta - \delta)y + (\beta^2 + \alpha\delta)x(1-y)}{\delta - \alpha - 2\beta + \beta^2 + \alpha\delta} \end{aligned}$$

for $0 \leq x \leq y \leq 1$, and the symmetric expression $G_0(y, x)$ for $0 \leq y \leq x \leq 1$.

L. GSARC Boundary Conditions with Zero Discriminant

We now discuss the case when the discriminant $\Delta = \delta - \alpha - 2\beta + \beta^2 + \alpha\delta = 0$. We first discuss the generic case, with $\delta = \frac{\alpha+2\beta-\beta^2}{\alpha+1}$ (equivalently $\alpha = \frac{\delta-2\beta+\beta^2}{1-\delta}$), then discuss the three limiting cases: (i) $\alpha \rightarrow -1$ (so that $\beta = 1$, and δ is arbitrary); (ii) $\delta \rightarrow 1$ (so that $\beta = 1$, and α is arbitrary); and (iii) the special case satisfying both (i) and (ii), i.e., $\alpha = -1$, $\beta = 1$, $\delta = 1$.

The expression of the resolvent kernel following from the same method is such that

$$2z(-2(1+\alpha)\beta - (\alpha^2 + \beta^2 - 2\beta) \cos \sqrt{z} + ((1+\alpha)z + (\alpha+\beta)^2) \sin \sqrt{z}) G(z, x, y)$$

is equal to

$$\begin{aligned} & -((1+\alpha)z + (\alpha+\beta)^2) \cos(\sqrt{z}(1+x-y)) + (-(1+\alpha)z + (\alpha+\beta)^2) \cos(\sqrt{z}(1-x-y)) \\ & + \sqrt{z} \left\{ -2(1+\alpha)\beta \sin(\sqrt{z}(x-y)) - (\alpha^2 + \beta^2 - 2\beta) \sin(\sqrt{z}(1+x-y)) \right. \\ & \left. + (2+\alpha-\beta)(\alpha+\beta) \sin(\sqrt{z}(1-x-y)) \right\} \end{aligned}$$

Upon calculating $\lim_{z \rightarrow 0^+} z^n G(z, x, y)$ for $n = 1, 2, \dots$, we realize that the kernel of the Riesz projection is given by

$$p(x, y) = -\frac{3(1-\beta + (\alpha+\beta)x)(1-\beta + (\alpha+\beta)y)}{3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta}$$

while the nilpotent operator $D \equiv 0$.

The Green's function corresponding to this problem is then

$$\begin{aligned} G_0(x, y) &= \lim_{z \rightarrow 0^+} G(z, x, y) - \frac{p(x, y)}{z} \\ &= c_{00} + c_{10}x + c_{01}y + c_{20}x^2 + c_{11}xy + c_{02}y^2 + c_{30}x^3 + c_{21}x^2y \\ &\quad + c_{12}xy^2 + c_{03}y^3 + c_{31}x^3y + c_{13}xy^3 \end{aligned}$$

for $0 \leq x \leq y \leq 1$. The coefficients depend on the parameters and are explicitly given in appendix A.

We opted not to include the expressions for $G(z, x, y)$ and $G_0(x, y)$ in table II and table III since the expressions are too long.

Remark III.7. Note that $c_{10} - c_{01} = 1$, $c_{20} = c_{02}$, $c_{21} = c_{12}$, $c_{30} = c_{03}$, and $c_{31} = c_{13}$. These identities will prove useful in the calculation of the perturbation in section IV H. $\square$

Remark III.8. We now treat the limiting cases (i), (ii), and (iii). Again, we list the results for $0 \leq x \leq y \leq 1$ and complete the expression for $0 \leq y \leq x \leq 1$ by symmetry.

(i) $\alpha \rightarrow -1$ (so that $\beta = 1$, and δ is arbitrary)

The expression of the resolvent kernel following from the same method is such that

$$(\sqrt{z}(-2 + (1+\delta) \cos \sqrt{z}) + (1-\delta+z) \sin \sqrt{z}) G(z, x, y)$$

is equal to

$$\begin{aligned} & (-\sqrt{z} \cos(\sqrt{z}(1-y)) + \delta \sin(\sqrt{z}(1-y)) + \sin(\sqrt{z}y)) \cos(\sqrt{z}x) \\ & + (\sqrt{z} \cos(\sqrt{z}(1-y)) - \cos(\sqrt{z}y) + (1-\delta) \sin(\sqrt{z}(1-y))) \frac{\sin(\sqrt{z}x)}{\sqrt{z}}. \end{aligned}$$

The Riesz projection has a kernel

$$p(x, y) = -3(1-x)(1-y)$$

and the corresponding Green's function is

$$\begin{aligned} G_0(x, y) &= -\frac{1}{20(1-\delta)} ((-9+4\delta) + (19-4\delta)x - 30(1-\delta)x^2 + 10(1-\delta)x^3 \\ &\quad + (39-24\delta)y + (-69+24\delta)y^2 + 10(1-\delta)y^3 \\ &\quad + 3(-23+8\delta)xy + 30(1-\delta)xy^2 + 30(1-\delta)x^2y \\ &\quad - 10(1-\delta)x^3y - 10(1-\delta)xy^3). \end{aligned}$$

(ii) $\delta \rightarrow 1$ (so that $\beta = 1$, and α is arbitrary). The expression of the resolvent kernel following from the same method is such that

$$(\sqrt{z}(-2 + (1 - \alpha)\cos\sqrt{z}) + (1 + \alpha + z)\sin\sqrt{z}) G(z, x, y)$$

is equal to

$$\begin{aligned} & (-\sqrt{z}\cos(\sqrt{z}(1-y)) + \sin(\sqrt{z}(1-y)) + \sin(\sqrt{z}y)) \cos(\sqrt{z}x) \\ & - \sqrt{z}(\alpha\cos(\sqrt{z}(1-y)) + \cos(\sqrt{z}y)) - (1 + \alpha)\sin(\sqrt{z}(1-y)) \frac{\sin(\sqrt{z}x)}{\sqrt{z}}. \end{aligned}$$

The Riesz projection has a kernel

$$p(x, y) = -3xy$$

and the corresponding Green's function is

$$\begin{aligned} G_0(x, y) = \frac{1}{20(1 + \alpha)} & (20 - 30x + 10(-1 + 2\alpha)y \\ & + (9 - 36\alpha)xy + 10(1 + \alpha)xy^3 + 10(1 + \alpha)x^3y). \end{aligned}$$

(iii) $\alpha = -1$, $\beta = 1$, and $\delta = 1$

This is the most particular of all these cases. The expression of the resolvent kernel following from the same method is such that

$$(2\sqrt{z}(-1 + \cos\sqrt{z} + z\sin\sqrt{z}) G(z, x, y)$$

is equal to

$$\begin{aligned} & (-\sqrt{z}\cos(\sqrt{z}(1-y)) + \sin(\sqrt{z}(1-y)) + \sin(\sqrt{z}y)) \cos(\sqrt{z}x) \\ & - \sin\left(\sqrt{z}\left(\frac{1}{2} - y\right)\right) \frac{\sin(\sqrt{z}x)}{2\sin\frac{\sqrt{z}}{2}}. \end{aligned}$$

The Riesz projection kernel is given by $p(x, y) = -4 + 6x + 6y - 12xy$. The corresponding Green's function is

$$\begin{aligned} G_0(x, y) = \frac{1}{30} & (4 - 3x + 60x^2 - 30x^3 - 33y + 36xy - 90x^2y \\ & + 60x^3y + 60y^2 - 90xy^2 - 30y^3 + 60xy^3). \end{aligned}$$

In all cases, the nilpotent operator $D \equiv 0$. □

IV. PERTUBATIONS OF THE NONLOCAL OPERATOR EQ. (I.1)

With $\mathcal{K}$ as defined in eq. (1), we are interested in understanding the nature of the integral operator $\mathcal{T} := \mathcal{K} - \mathcal{K}g$, where $\mathcal{K}g f(x) := \int_0^1 G_0(x, y)f(y) dy$, $G_0(x, y)$ is one of the Green's functions developed in section III for various GSARC problems. The perturbation $\mathcal{T}$ we aim to study is then given by

$$\mathcal{T}f(x) := \int_0^1 (\kappa(x, y) - G_0(x, y)) f(y) dy, \quad (36)$$

where $\kappa(x, y)$ was defined by eq. (7). The method consists of finding the eigenvalues and corresponding eigenfunctions of this operator using Linear Algebra. In each of the BVPs in this article, we proceed as follows:

Step 1: Determine the form of the kernel $T(x, y) = \kappa(x, y) - G_0(x, y)$ for $x \leq y$;

Step 2: Set the kernel $T(x, y)$, for $x \geq y$, by symmetry;

Step 3: Simplify the expression of

$$\int_0^1 T(x, y) f(y) dy = \int_0^x T(x, y) f(y) dy + \int_x^1 T(y, x) f(y) dy; \quad (37)$$

(This leads to the appropriate form of the function $f(x)$. It turns out to be a polynomial with a degree off by 1 from the size of the matrix corresponding to the operator $\mathcal{T}$.)

Step 4: Find the spectral resolution of the operator $\mathcal{T}$ by working out the spectral resolution of the corresponding matrix problem.

The range of the operator $\mathcal{T}$ is a space of polynomials. If n is the rank of the matrix $M_{\mathcal{T}}$ corresponding to the operator $\mathcal{T}$. The spectral resolution of this operator is given by

$$\mathcal{T}f(x) = \sum_{k=1}^n \langle f, u_k \rangle u_k(x) \quad (38)$$

where $\langle f, g \rangle := \int_0^1 f(x)g(x) dx$ is the usual dot product, and $u_1, \dots, u_n$ are the eigenvectors of the matrix $M_{\mathcal{T}}$ which we determine for the various eigenvalue problems discussed earlier.

In all cases, we will show that $T(x, y) = T(y, x)$, thus the split of the integral in eq. (37) is not needed. In fact the resolved kernel associated with $\mathcal{T}$ is symmetric as well in x and y .

Remark IV.1. *Perturbation is treated in a multiplicative framework in Section 2.8 of⁶⁹ where the finiteness of the rank of the perturbation implies the existence of boundary triplets (see Theorem 2.8.1). Kreĭn-type resolvent formulas (and thus Green functions) appear in⁷⁰ (Sec.4.3, 7.5, 11.3, 13.10, 14.13, Appendix D.6).*

A. Dirichlet Boundary Conditions

As noted in¹ and², the Green's function corresponding to the Dirichlet problem (consult table III) is given by

$$G_0(x, y) = \frac{1}{2}(x+y) - xy - \frac{1}{2}|x-y| = -\left(x - \frac{1}{2}\right)\left(y - \frac{1}{2}\right) + \frac{1}{4} - \frac{1}{2}|x-y|. \quad (39)$$

The latter form appears in the 1959 paper of Lidskiĭ⁴⁹. From eq. (39), we easily get

$$T(x, y) = xy - \frac{1}{2}(x+y).$$

Since this expression is symmetric, viz. $T(x, y) = T(y, x)$, the perturbation operator takes the simple expression

$$\begin{aligned} \mathcal{T}f(x) &= \int_0^1 \left(xy - \frac{1}{2}(x+y)\right) f(y) dy \\ &= -\left(\int_0^1 \frac{1}{2} y f(y) dy\right) - \left(\int_0^1 \left(\frac{1}{2} - y\right) f(y) dy\right) x. \end{aligned} \quad (40)$$

By virtue of eq. (40), the eigenvalue problem $\mathcal{T}f(x) = \lambda f(x)$ leads to a linear eigenfunctions of the form $f(x) = a_0 + a_1 x$ and the matrix eigenvalue problem

$$\begin{pmatrix} -\frac{1}{4} & -\frac{1}{6} \\ 0 & \frac{1}{12} \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \end{pmatrix} = \lambda \begin{pmatrix} a_0 \\ a_1 \end{pmatrix}.$$

Thus $\lambda_1 = -\frac{1}{4}$, $\lambda_2 = \frac{1}{12}$, and the corresponding eigenvectors are

$$u_1(x) = 1, \quad u_2(x) = x - \frac{1}{2}. \quad (41)$$

This agrees with Lidskiĭ⁴⁹ and our earlier work². We have the immediate theorem:

Theorem IV.2. *The Dirichlet operator $\mathcal{K}_{\mathcal{G}}$ is a rank 2 perturbation of $\mathcal{K}$. Moreover,*

$$\mathcal{T}f(x) = \langle f, u_1 \rangle u_1(x) + \langle f, u_2 \rangle u_2(x),$$

where $u_1(x), u_2(x)$ are defined in eq. (41).

B. Neumann Boundary Conditions

The calculation of the perturbation $\mathcal{T}$ appears as Remark 2.9 in², where we used the method of f^{71,72}. We offer here a succinct scheme. From the expression of Green's function,

$$G_0(x, y) = \frac{1}{6} (2 + 3x^2 - 6y + 3y^2),$$

the kernel of the perturbation is

$$T(x, y) = -\frac{1}{6} (2 - 3x + 3x^2 - 3y + 3y^2)$$

for $x \leq y$. Since this expression is symmetric, viz. $T(x, y) = T(y, x)$, the operator $\mathcal{T}$ acts as

$$\begin{aligned} \mathcal{T}f(x) &= \int_0^1 T(x, y)f(y) dy \\ &= \left(\int_0^1 -\frac{1}{6} (2 - 3y + 3y^2) f(y) dy \right) + \left(\int_0^1 \frac{1}{2} f(y) dy \right) (x - x^2). \end{aligned} \quad (42)$$

Hence, the eigenvalue problem corresponding to this operator, $\mathcal{T}f(x) = \lambda f(x)$, leads to a quadratic form of the eigenvector $f(x) = a_0 + a_1(x - x^2)$, and the matrix eigenvalue problem

$$\begin{pmatrix} -\frac{1}{4} & -\frac{7}{180} \\ \frac{1}{2} & \frac{1}{12} \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \end{pmatrix} = \lambda \begin{pmatrix} a_0 \\ a_1 \end{pmatrix}.$$

Thus $\lambda_1 = \frac{1}{60}(-5 - \sqrt{30})$, and $\lambda_2 = \frac{1}{60}(-5 + \sqrt{30})$. The corresponding eigenvectors are:

$$u_1(x) = -\frac{1}{30} (10 + \sqrt{30}) + x - x^2, \quad u_2(x) = \frac{1}{30} (-10 + \sqrt{30}) + x - x^2. \quad (43)$$

Hence we have:

Theorem IV.3. *The Neumann operator $\mathcal{H}_g$ is a rank 2 perturbation of $\mathcal{H}$. Moreover,*

$$\mathcal{T}f(x) = \langle f, u_1 \rangle u_1(x) + \langle f, u_2 \rangle u_2(x),$$

where $u_1(x)$ and $u_2(x)$ are defined in eq. (43).

Remark IV.4. *The expression of the perturbation operator $\mathcal{T}f(x)$ in eq. (42) is quadratic in x , and thus one can simply write $f(x) = a_0 + a_1x + a_2x^2$. This leads to an eigenvalue problem with a matrix $M_{\mathcal{T}}$ given by*

$$M_{\mathcal{T}} = \begin{pmatrix} -\frac{1}{4} & -\frac{1}{8} & -\frac{31}{361} \\ \frac{1}{2} & \frac{1}{4} & \frac{1}{6} \\ -\frac{1}{2} & -\frac{1}{4} & -\frac{1}{6} \end{pmatrix}.$$

Notice that the first column is twice the second, hence the column vectors are linearly dependent and the rank of this 3×3 matrix is 2. Indeed the eigenvalues of this matrix are λ_1 and λ_2 above, with the eigenfunctions given in eq. (43), and $\lambda_0 = 0$, with corresponding eigenfunction $u_0(x) = -\frac{1}{2} + x$. Alternatively one can perform a Singular Value Decomposition (SVD) of the matrix $M_{\mathcal{T}}$ which immediately gives the number of non-zero singular values in this case as 2. This is also the rank since the number of non-zero singular values is also the rank (see Theorem 5.1 of Ref. 73).

C. Krein-von Neumann Boundary Conditions

In section IIID, we derived the Green's function $G_0(x, y)$ corresponding to this problem with BCs $u'(0) = u'(1) = -u(0) + u(1)$. For $0 \leq x \leq y \leq 1$, it is explicitly given by

$$G_0(x, y) = \frac{1}{30} (4 - 3x - 33y + 60x^2 + 60y^2 + 36xy)$$

$$-30x^3 - 90x^2y - 90xy^2 - 30y^3 + 60x^3y + 60xy^3).$$

The expression of $T(x, y)$ is given by

$$T(x, y) = \frac{1}{15} (-2 + 9x - 30x^2 + 15x^3 + 45x^2y - 30x^3y - 18xy - 30x^3y + 45xy^2 + 15y^3 - 30y^2 + 9y).$$

This is again symmetric, viz. $T(x, y) = T(y, x)$. This is a cubic polynomial in x , and thus the eigenvector of the matrix problem is of the form $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3$. The corresponding matrix has the explicit form

$$M_{\mathcal{T}} = \begin{pmatrix} -\frac{1}{4} & -\frac{1}{6} & -\frac{23}{180} & -\frac{109}{1050} \\ \frac{1}{2} & \frac{1}{4} & \frac{1}{6} & \frac{87}{700} \\ -\frac{1}{2} & 0 & \frac{1}{12} & \frac{1}{10} \\ 0 & -\frac{1}{6} & -\frac{1}{6} & -\frac{3}{20} \end{pmatrix}. \quad (44)$$

Its eigenvalues are: $\lambda_1 = \frac{1}{60}(-5 - \sqrt{30})$, $\lambda_2 = \frac{1}{60}(-5 + \sqrt{30})$, $\lambda_3 = \frac{1}{420}(21 - \sqrt{462})$, and $\lambda_4 = \frac{1}{420}(21 + \sqrt{462})$. The corresponding eigenvectors are

$$\begin{aligned} u_1(x) &= \frac{1}{30} (10 + \sqrt{30}) - x + x^2, \\ u_2(x) &= \frac{1}{30} (10 - \sqrt{30}) - x + x^2, \\ u_3(x) &= \frac{1}{140} (14 - \sqrt{462}) - \frac{1}{70} (21 + \sqrt{462})x - \frac{3}{2}x^2 + x^3, \\ u_4(x) &= \frac{1}{140} (14 + \sqrt{462}) - \frac{1}{70} (21 - \sqrt{462})x - \frac{3}{2}x^2 + x^3. \end{aligned} \quad (45)$$

We now have:

Theorem IV.5. *The Kreĭn-von Neumann operator $\mathcal{K}_{\mathcal{G}}$ is a rank 4 perturbation of $\mathcal{K}$. Moreover,*

$$\mathcal{T}f(x) = \sum_{k=1}^4 \langle f, u_k \rangle u_k(x),$$

where $u_k(x)$, $k = 1, 2, 3, 4$, are defined in eq. (45).

D. Robin Boundary Conditions

Using the expression of the Green's function developed in section III G, the expression of the perturbation is given by

$$T(x, y) = \frac{2 - 2\delta + (\alpha + \delta - \alpha\delta)x + (\alpha + \delta - \alpha\delta)y + 2\alpha\delta xy}{2(\delta - \alpha + \alpha\delta)}$$

for $0 \leq x \leq y \leq 1$, with $T(x, y) = T(y, x)$ for all $x, y \in [0, 1]$. The eigenvectors are then of the form $f(x) = a_0 + a_1x$. The matrix is given by,

$$M_{\mathcal{T}} = \begin{pmatrix} \frac{4+\alpha-3\delta-\alpha\delta}{4(\delta-\alpha+\alpha\delta)} & \frac{3+\alpha-2\delta-\alpha\delta}{6(\delta-\alpha+\alpha\delta)} \\ \frac{\alpha+\delta}{2(\delta-\alpha+\alpha\delta)} & \frac{3\alpha+3\delta+\alpha\delta}{12(\delta-\alpha+\alpha\delta)} \end{pmatrix}.$$

with eigenvalues

$$\lambda_1 = \frac{6+3\alpha-3\delta-\alpha\delta-2\sqrt{(3+3\alpha+\alpha^2)(3-3\delta+\delta^2)}}{12(\delta-\alpha+\alpha\delta)}, \lambda_2 = \frac{6+3\alpha-3\delta-\alpha\delta+2\sqrt{(3+3\alpha+\alpha^2)(3-3\delta+\delta^2)}}{12(\delta-\alpha+\alpha\delta)},$$

and corresponding eigenvectors

$$u_1(x) = \frac{3 - 3\delta - \alpha\delta - \sqrt{(3+3\alpha+\alpha^2)(3-3\delta+\delta^2)}}{3(\alpha+\delta)} + x$$

and

$$u_2(x) = \frac{3 - 3\delta - \alpha\delta + \sqrt{(3 + 3\alpha + \alpha^2)(3 - 3\delta + \delta^2)}}{3(\alpha + \delta)} + x.$$

Example IV.6. In example III.2 the limiting BC case $u(0) = u'(1) = 0$ example of the Robin problem was discussed. The perturbation kernel takes the form

$$T(x, y) = -\frac{1}{2}(x + y)$$

for $0 \leq x \leq y \leq 1$, with $T(x, y) = T(y, x)$ for all $x, y \in [0, 1]$. The eigenvectors are then of the form $f(x) = a_0 + a_1x$. The matrix is given by,

$$M_{\mathcal{T}} = \begin{pmatrix} -\frac{1}{4} & -\frac{1}{6} \\ -\frac{1}{2} & -\frac{1}{4} \end{pmatrix}.$$

with eigenvalues $\lambda_1 = -\frac{3+2\sqrt{3}}{12}$ and $\lambda_2 = -\frac{3-2\sqrt{3}}{12}$, and corresponding eigenvectors

$$u_1(x) = \frac{1}{\sqrt{3}} + x$$

and

$$u_2(x) = -\frac{1}{\sqrt{3}} + x.$$

Example IV.7. In Kato's first variation on the Robin example discussed in example III.3, the perturbation kernel takes the form

$$T(x, y) = -\frac{1}{2}(x + y) + \frac{1}{1-\tau}xy$$

for $0 \leq x \leq y \leq 1$. Clearly, $T(x, y) = T(y, x)$ for all $x, y \in [0, 1]$. The eigenvectors are then of the form $f(x) = a_0 + a_1x$. The matrix of the finite rank operator is 2×2 ,

$$M_{\mathcal{T}} = \begin{pmatrix} -\frac{1}{4} & -\frac{1}{6} \\ \frac{\tau}{2(1-\tau)} & \frac{1+3\tau}{12(1-\tau)} \end{pmatrix}.$$

with eigenvalues $\lambda_1 = -\frac{1-3\tau-2\sqrt{1-3\tau+3\tau^2}}{12(1-\tau)}$ and $\lambda_2 = -\frac{1-3\tau+2\sqrt{1-3\tau+3\tau^2}}{12(1-\tau)}$, and corresponding eigenvectors

$$u_1(x) = -\frac{1 - \sqrt{1 - 3\tau + 3\tau^2}}{3\tau} + x$$

and

$$u_2(x) = -\frac{1 + \sqrt{1 - 3\tau + 3\tau^2}}{3\tau} + x.$$

Example IV.8. In Kato's second variation on the Robin BC of example III.4, the perturbation kernel takes the form

$$T(x, y) = -\frac{1}{2}(x + y) + \frac{\tau}{1+\tau}xy$$

for $0 \leq x \leq y \leq 1$. We also have $T(x, y) = T(y, x)$ for all $x, y \in [0, 1]$. The eigenvectors are then of the form $f(x) = a_0 + a_1x$. The associated matrix is also 2×2 ,

$$M_{\mathcal{T}} = \begin{pmatrix} -\frac{1}{4} & -\frac{1}{6} \\ -\frac{1}{2(1+\tau)} & -\frac{3-\tau}{12(1+\tau)} \end{pmatrix}.$$

with eigenvalues $\lambda_1 = -\frac{3+\tau+2\sqrt{3+3\tau+\tau^2}}{12(1+\tau)}$ and $\lambda_2 = -\frac{3+\tau-2\sqrt{3+3\tau+\tau^2}}{12(1+\tau)}$, and corresponding eigenvectors

$$u_1(x) = \frac{\tau + \sqrt{3+3\tau+\tau^2}}{3} + x$$

and

$$u_2(x) = \frac{\tau - \sqrt{3+3\tau+\tau^2}}{3} + x.$$

Example IV.9. In example III.5 of Stakgold and Holst, the perturbation kernel takes the form

$$T(x, y) = -\frac{1}{2}(x+y) + \frac{\cot \theta}{1 + \cot \theta} xy$$

for $0 \leq x \leq y \leq 1$, and again $T(x, y) = T(y, x)$ for all $x, y \in [0, 1]$. The eigenvectors are then of the form $f(x) = a_0 + a_1 x$. The matrix of the finite rank operator is 2×2 ,

$$M_{\mathcal{T}} = \begin{pmatrix} -\frac{1}{4} & -\frac{1}{6} \\ -\frac{1}{2(1+\cot \theta)} & -\frac{3-\cot \theta}{12(1+\cot \theta)} \end{pmatrix}.$$

with eigenvalues $\lambda_1 = -\frac{3+\cot \theta+2\sqrt{3+\cot \theta+\cot^2 \theta}}{12(1+\cot \theta)}$ and $\lambda_2 = -\frac{3+\cot \theta-2\sqrt{3+\cot \theta+\cot^2 \theta}}{12(1+\cot \theta)}$, and corresponding eigenvectors

$$u_1(x) = \frac{\cot \theta + \sqrt{3+3\cot \theta+\cot^2 \theta}}{3} + x$$

and

$$u_2(x) = \frac{\cot \theta - \sqrt{3+3\cot \theta+\cot^2 \theta}}{3} + x.$$

Remark IV.10. When $\delta - \alpha + \alpha\delta = 0$, and by the virtue of remark III.6, the perturbation kernel takes the form

$$\begin{aligned} T(x, y) &= -c_{00} + \left(\frac{1}{2} - c_{10}\right)x - \left(c_{01} + \frac{1}{2}\right)y - c_{20}x^2 - c_{11}xy - c_{02}y^2 - c_{30}x^3 \\ &\quad - c_{21}x^2y - c_{12}xy^2 - c_{03}y^3 - c_{31}x^3y - c_{13}xy^3 \\ &:= d_{00} + d_{10}x + d_{10}y + d_{20}x^2 + d_{11}xy + d_{20}y^2 + d_{30}x^3 + d_{21}x^2y \\ &\quad + d_{21}xy^2 + d_{30}y^3 + d_{31}x^3y + d_{31}xy^3. \end{aligned} \tag{46}$$

where the c_{ij} and d_{ij} coefficients are again explicitly given in appendix A, with $\beta = 0$. The eigenfunctions have the form $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3$, and the corresponding matrix $M_{\mathcal{T}} = (m_{ij})$, is 4×4 with elements given by appendix B with $\beta = 0$. This is a rank 2 matrix with $\lambda = 0$ as a double eigenvalue; see also section IV J.

E. Periodic Boundary Conditions

The Green's function in this case is given in section III H, for $0 \leq x \leq y \leq 1$, by

$$G_0(x, y) = \frac{1}{12} (1 + 6x - 6y + 6x^2 - 12xy + 6y^2).$$

The expression of $T(x, y)$ is given by

$$T(x, y) = \frac{1}{12} (-1 - 6x^2 + 12xy - 6y^2).$$

It is symmetric, viz. $T(x, y) = T(y, x)$. Since it is a quadratic polynomial in x , the eigenvector of the matrix problem is of the form $f(x) = a_0 + a_1x + a_2x^2$. The corresponding matrix is

$$M_{\mathcal{T}} = \begin{pmatrix} -\frac{1}{4} & -\frac{1}{6} & -\frac{23}{180} \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{4} \\ -\frac{1}{2} & -\frac{1}{4} & -\frac{1}{6} \end{pmatrix}.$$

The eigenvalues are: $\lambda_1 = \frac{1}{12}$, $\lambda_2 = \frac{1}{60}(-5 - \sqrt{30})$, $\lambda_3 = \frac{1}{60}(-5 + \sqrt{30})$. The corresponding eigenvectors are

$$u_1(x) = -\frac{1}{2} + x, \quad u_2(x) = \frac{1}{30}(10 + \sqrt{30}) - x + x^2, \quad u_3(x) = \frac{1}{30}(10 - \sqrt{30}) - x + x^2. \quad (47)$$

We have:

Theorem IV.11. *The periodic operator $\mathcal{K}_{\mathcal{G}}$ is a rank 3 perturbation of $\mathcal{K}$. Moreover,*

$$\mathcal{T}f(x) = \sum_{k=1}^3 \langle f, u_k \rangle u_k(x),$$

where $u_k(x)$, $k = 1, 2, 3$, are defined in eq. (47).

F. Anti-Periodic Boundary Conditions

The anti-periodic integral operator turns out to be the “closest” to the nonlocal integral operator. To see this, note that for $0 \leq x \leq y \leq 1$, the derived Green’s function takes the form

$$G_0(x, y) = \frac{1}{4}(1 + 2x - 2y) \quad (48)$$

and $T(x, y) = -\frac{1}{4}$. Thus $\mathcal{T}$ has eigenvalue $\lambda = -\frac{1}{4}$ and eigenvector $u_1(x) \equiv 1$. Hence, we have:

Theorem IV.12. *The anti-periodic operator $\mathcal{K}_{\mathcal{G}}$ is a rank 1 perturbation of $\mathcal{K}$. Moreover,*

$$\mathcal{T}f(x) = \langle f, u_1 \rangle u_1(x) = \int_0^1 f(y) dy,$$

which is simply a constant often called the “DC” component of $f(x)$.

G. Radoux Boundary Conditions

The Green’s function $G_0(x, y)$ corresponding to this problem is explicitly given in section III J as

$$G_0(x, y) = \frac{1}{10}(10x - 18xy + 5xy^3 + 5x^3y),$$

for $0 \leq x \leq y \leq 1$. That of $T(x, y)$ is given by

$$T(x, y) = \frac{1}{10}(-5x - 5y + 18xy - 5xy^3 - 5x^3y).$$

This is again symmetric, viz. $T(x, y) = T(y, x)$, and we only need a single integral to calculate the corresponding matrix operator.

$$\begin{aligned} \mathcal{T}f(x) &= \int_0^1 \frac{1}{10}(-5x - 5y + 18xy - 5xy^3 - 5x^3y) f(y) dy \\ &= \frac{x}{10} \int_0^1 (-5 + 18y - 5y^3) f(y) dy - \frac{1+x^3}{2} \int_0^1 y f(y) dy. \end{aligned}$$

Hence, the eigenvalue problem corresponding to this operator, $\mathcal{T}f(x) = \lambda f(x)$, leads to a cubic form of the eigenvector $f(x) = a_0x + a_1(1+x^3)$, and the matrix eigenvalue problem

$$\begin{pmatrix} \frac{307}{700} & \frac{1}{4} \\ -\frac{7}{20} & -\frac{1}{6} \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \end{pmatrix} = \lambda \begin{pmatrix} a_0 \\ a_1 \end{pmatrix}.$$

Its eigenvalues are $\lambda_1 = \frac{1}{420}(-21 - \sqrt{2982})$, and $\lambda_2 = \frac{1}{420}(-21 + \sqrt{2982})$. The corresponding eigenvectors are

$$\begin{aligned} u_1(x) &= x - \frac{5}{921}(126 + \sqrt{2982})(1+x^3), \\ u_2(x) &= x + \frac{5}{921}(-126 + \sqrt{2982})(1+x^3). \end{aligned} \quad (49)$$

In conclusion, we have the following theorem.

Theorem IV.13. *The Radoux operator $\mathcal{K}_g$ is a rank 2 perturbation of $\mathcal{K}$. Moreover,*

$$\mathcal{T}f(x) = \langle f, u_1 \rangle u_1(x) + \langle f, u_2 \rangle u_2(x),$$

where $u_1(x)$ and $u_2(x)$ are defined in eq. (49).

H. GSARC Boundary Conditions with Non-Zero Discriminant

For $\Delta = \delta - \alpha - 2\beta + \beta^2 + \alpha\delta \neq 0$ the Green's function developed in section III K is a quadratic polynomial given, for $0 \leq x \leq y \leq 1$ by

$$G_0(x, y) = \frac{-1 + \delta - (\alpha + \beta)x + (\beta - \delta)y + (\beta^2 + \alpha\delta)x(1-y)}{\delta - \alpha - 2\beta + \beta^2 + \alpha\delta}.$$

The kernel of the perturbation is then

$$T(x, y) = c_{00} + c_{10}x + c_{01}y + c_{11}xy$$

where

$$c_{00} = \frac{1 - \delta}{\Delta},$$

$$c_{10} = c_{01} = \frac{\alpha + \delta - \beta^2 - \alpha\delta}{2\Delta},$$

and

$$c_{11} = \frac{\beta^2 + \alpha\delta}{2\Delta}.$$

The eigenvectors corresponding to this finite rank operator are of the form $f(x) = a_0 + a_1x$, and the 2×2 matrix of the operator $M_{\mathcal{T}}$ has the entry $m_{11} = \frac{-4 - \alpha + 3\delta + \beta^2 + \alpha\delta}{4\Delta}$, $m_{12} = \frac{-3 - \alpha + 2\delta + \beta^2 + \alpha\delta}{6\Delta}$, $m_{21} = \frac{\alpha + \delta}{2\Delta}$, and $m_{22} = \frac{3\alpha + 3\delta + \beta^2 + \alpha\delta}{12\Delta}$. Its eigenvalues are $\lambda_1 = \frac{-6 - 3\alpha + 3\delta + \beta^2 + \alpha\delta - 2\sqrt{\tilde{\Delta}}}{12\Delta}$, $\lambda_2 = \frac{-6 - 3\alpha + 3\delta + \beta^2 + \alpha\delta + 2\sqrt{\tilde{\Delta}}}{12\Delta}$, where

$$\tilde{\Delta} := 9 + 9\alpha + 3\alpha^2 - 6\beta^2 - 3\alpha\beta^2 + \beta^4 - 9\delta - 9\alpha\delta + 3\alpha^2\delta + 3\beta^2\delta + 2\alpha\beta^2\delta + 3\delta^2 + 3\alpha\delta^2 + \alpha^2\delta^2$$

and the corresponding eigenvectors are

$$u_1(x) = \frac{2(3 - \beta^2 - 3\delta - \alpha\delta - \sqrt{\tilde{\Delta}})}{3(\alpha + \delta)} + x$$

and

$$u_2(x) = \frac{2(3 - \beta^2 - 3\delta - \alpha\delta + \sqrt{\tilde{\Delta}})}{3(\alpha + \delta)} + x.$$

I. GSARC Boundary Conditions with Zero Discriminant

The Green's function $G_0(x, y)$ is detailed in section III L. It assumes the form $G_0(x, y) = -\frac{1}{2}|x - y| - T(x, y)$, where $T(x, y)$ is the kernel of the perturbation given by the expression

$$\begin{aligned} T(x, y) &= -c_{00} + \left(\frac{1}{2} - c_{10}\right)x - \left(c_{01} + \frac{1}{2}\right)y - c_{20}x^2 - c_{11}xy - c_{02}y^2 - c_{30}x^3 \\ &\quad - c_{21}x^2y - c_{12}xy^2 - c_{03}y^3 - c_{31}x^3y - c_{13}xy^3 \\ &:= d_{00} + d_{10}x + d_{10}y + d_{20}x^2 + d_{11}xy + d_{20}y^2 + d_{30}x^3 + d_{21}x^2y \\ &\quad + d_{21}xy^2 + d_{30}y^3 + d_{31}x^3y + d_{31}xy^3. \end{aligned} \quad (50)$$

By virtue of remark III.7, $T(x, y) = T(y, x)$ for all $x, y \in [0, 1]$. This is also a cubic polynomial in x . The eigenvector of the matrix problem is of the form $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3$. The corresponding matrix is 4×4

$$M_{\mathcal{T}} = (m_{ij}).$$

where explicit forms of these entries, obtained using Wolfram Mathematica[®] are given in appendix B.

Its four eigenvalues λ_k , and four eigenvectors, $u_k(x)$, $k = 1, 2, 3, 4$ are too complicated to list.

We now have:

Theorem IV.14. *The GSARC operator $\mathcal{K}_{\mathcal{G}}$ is an up-to-rank 4 perturbation of $\mathcal{K}$. Moreover,*

$$\mathcal{T}f(x) = \sum_{k=1}^{\text{rank}(\mathcal{K}_{\mathcal{G}})} \langle f, u_k \rangle u_k(x).$$

Remark IV.15. *The case $\beta = -\alpha$ merits special attention since $T(x, y)$ in eq. (50) reduces to a quadratic polynomial; see also the expressions of c_{ij} 's in appendix A. Since $\Delta = 0$, we also have $\delta = -\alpha$. For $\alpha \neq 0, -1$, the Green's function reduces to*

$$G_0(x, y) = \frac{4 + \alpha}{12(1 + \alpha)} + \frac{\alpha}{2(1 + \alpha)}x - \frac{2 + \alpha}{2(1 + \alpha)}y - \frac{\alpha}{1 + \alpha} + \frac{1}{2}x^2 + \frac{1}{2}y^2. \quad (51)$$

The kernel of the Riesz projection is given by $p(x, y) = -1$, and therefore

$$T(x, y) = -\frac{4 + \alpha}{12(1 + \alpha)} + \frac{1}{2(1 + \alpha)}x + \frac{1}{2(1 + \alpha)}y + \frac{\alpha}{1 + \alpha}xy + \frac{1}{2}x^2 + \frac{1}{2}y^2. \quad (52)$$

Proceeding as before, we obtain a rank 3 perturbation with eigenvalues $\lambda_1 = \frac{\alpha}{12(1 + \alpha)}$,

$\lambda_2 = \frac{-5 - \sqrt{30}}{60}$, $\lambda_3 = \frac{-5 + \sqrt{30}}{60}$ and corresponding eigenvectors

$$\begin{aligned} u_1(x) &= -\frac{1}{2} + x \\ u_2(x) &= \frac{10 - \sqrt{30}}{30} - x + x^2 \\ u_3(x) &= \frac{10 + \sqrt{30}}{30} - x + x^2. \end{aligned}$$

The cases $\alpha = -1$ and $\alpha = 0$ have already been treated, i.e., the Kreĭn-von Neumann case (rank 4) and the Neumann case (rank 2), respectively. In light of these discussions, we note that rank 1 perturbation in the GSARC problem with $\Delta = 0$ cannot occur.

J. GSARS Boundary Conditions

We briefly discuss the GSARS BC case, in the form of eq. (6). It corresponds in our notation to $\beta = -\gamma = 0$ (see Remark I.2). It is indeed equivalent to the generalized Robin problem treated in section III G and section IV D. We focus on the dependence on the angles θ_0 and θ_1 . The discriminant condition for not having a zero eigenvalue is $\Delta = \delta - \alpha + \alpha\delta \neq 0$, which is equivalent to $\tilde{\Delta} = \cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 2\sin(\theta_0 - \theta_1) \neq 0$. In this case, the Riez projection $P \equiv 0$. For $0 \leq x \leq y \leq 1$, with

$$c = (-1 + z)\sin(\sqrt{z} - \theta_0 - \theta_1) + (-1 + 2\sqrt{z} - z)\sin(\sqrt{z} + \theta_0 - \theta_1)$$

$$+ (-1 - 2\sqrt{z} - z) \sin(\sqrt{z} - \theta_0 + \theta_1) + (-1 + z) \sin(\sqrt{z} + \theta_0 + \theta_1)$$

the resolvent kernel is such that

$$c \sqrt{z} G(z, x, y)$$

is equal to

$$((-1 - \sqrt{z}) \sin(\sqrt{z}x - \theta_0) + (-1 + \sqrt{z}) \sin(\sqrt{z}x + \theta_0)) ((-1 - \sqrt{z}) \sin(\sqrt{z}(-1 + y) - \theta_1) + (-1 + \sqrt{z}) \sin(\sqrt{z}(-1 + y) + \theta_1)).$$

The Green's function corresponding to the GSARS BC with non-zero discriminant is then

$$G_0(x, y) = \frac{2(x \cos \theta_0 - \sin \theta_0) ((1 - y) \cos \theta_1 + \sin \theta_1)}{\cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 2 \sin(\theta_0 - \theta_1)}.$$

The kernel of the perturbation is given by

$$T(x, y) = \frac{(\cos \theta_0 \cos \theta_1 + \sin(\theta_0 + \theta_1))(x + y) - 2xy \cos \theta_0 \cos \theta_1}{\cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 2 \sin(\theta_0 - \theta_1)}.$$

The matrix of the associated finite rank operator is 2×2 , and is given by

$$M_{\mathcal{T}} = \begin{pmatrix} \frac{3 \cos(\theta_0 - \theta_1) - 5 \cos(\theta_0 + \theta_1) + 4 \sin(\theta_0 - \theta_1) + 2 \sin(\theta_0 + \theta_1)}{4(\cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 2 \sin(\theta_0 - \theta_1))} & \frac{2 \cos(\theta_0 - \theta_1) - 4 \cos(\theta_0 + \theta_1) + 3 \sin(\theta_0 - \theta_1) + \sin(\theta_0 + \theta_1)}{6(\cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 2 \sin(\theta_0 - \theta_1))} \\ \frac{-\sin(\theta_0 + \theta_1)}{\cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 2 \sin(\theta_0 - \theta_1)} & -\frac{\cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 6 \sin(\theta_0 + \theta_1)}{12(\cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 2 \sin(\theta_0 - \theta_1))} \end{pmatrix}.$$

The eigenvalues of the rank 2 perturbation in this case, are solutions of the quadratic equation

$$A\lambda^2 + 2B\lambda + C = 0$$

where $A = 48\tilde{\Delta}$, $B = -20 \cos(\theta_0 - \theta_1) + 29 \cos(\theta_0 + \theta_1) - 24 \sin(\theta_0 - \theta_1)$, and $C = -\tilde{\Delta}$.

The various GSARS cases are displayed in fig. 2 for $(\theta_0, \theta_1) \in [0, \pi) \times [0, \pi)$: Dirichlet $(0, 0)$, Neumann $(\pi/2, \pi/2)$, Robin $(0, \pi/2)$, and Radoux $(0, 3\pi/4)$. The curve represents the level sets of the discriminant condition $\tilde{\Delta} = 0$.

The case $(\theta_0, \theta_1) = (\pi/4, 0)$ represents an interesting case not treated in the literature, with $\tilde{\Delta} = 0$, corresponding to the boundary conditions $u'(0) + u(0) = u(1) = 0$. Its resolvent has the kernel

$$G(z, x, y) = \frac{(\sqrt{z} \cos(\sqrt{z}x) - \sin(\sqrt{z}x)) (\sin(\sqrt{z}(1 - y)))}{z \cos \sqrt{z} - \sqrt{z} \sin \sqrt{z}}.$$

Its Riesz projection has the kernel $p(x, y) = -3(1 - x)(1 - y)$, and the nilpotent operator $D \equiv 0$. The Green's function is given by

$$G_0(x, y) = \frac{1}{10} (2 - 2x + 15x^2 - 5x^3 - 12y + 12xy - 15x^2y + 5x^3y + 15y^2 - 15xy^2 - 5y^3 + 5xy^3).$$

Working as before, we obtain the kernel of the perturbation

$$T(x, y) = \frac{1}{10} (-2 + 7x - 15x^2 + 5x^3 + 7y - 12xy + 15x^2y - 5x^3y - 15y^2 + 15xy^2 + 5y^3 - 5xy^3). \quad (53)$$

Note that $T(x, y) = T(y, x)$. The operator $\mathcal{T}$ then acts as

$$\begin{aligned} \mathcal{T}f(x) &= \int_0^1 T(x, y) f(y) dy \\ &= \left(\int_0^1 \frac{1}{10} (-2 + 7y - 15y^2 + 5y^3) f(y) dy \right) (1 - x) \\ &\quad + \left(\int_0^1 \frac{1}{10} (1 - y) f(y) dy \right) (5x - 15x^2 + 5x^3). \end{aligned}$$

The eigenvalue problem $\mathcal{T}f(x) = \lambda f(x)$ leads to a linear eigenfunctions of the form $f(x) = a_0(-1-x) + a_1(5x - 15x^2 + 5x^3)$ and the matrix eigenvalue problem

$$\begin{pmatrix} -\frac{1}{12} & \frac{199}{420} \\ \frac{1}{30} & -\frac{1}{70} \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \end{pmatrix} = \lambda \begin{pmatrix} a_0 \\ a_1 \end{pmatrix}.$$

Thus $\lambda_1 = \frac{-21-\sqrt{2982}}{420}$, $\lambda_2 = \frac{-21+\sqrt{2982}}{420}$, and the corresponding eigenvectors are

$$u_1(x) = \frac{-14-\sqrt{2982}}{14}(1-x) + (5x - 15x^2 + 5x^3), \quad u_2(x) = \frac{-14+\sqrt{2982}}{14}(1-x) + (5x - 15x^2 + 5x^3). \quad (54)$$

Remark IV.16. The expression of the perturbation in eq. (53) is cubic in x . An alternative analysis with an eigenfunction of the form $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3$, with a corresponding matrix

$$M_{\mathcal{T}} = \begin{pmatrix} -\frac{9}{40} & -\frac{17}{120} & -\frac{13}{120} & -\frac{31}{350} \\ \frac{19}{40} & \frac{9}{40} & \frac{3}{20} & \frac{159}{1400} \\ -\frac{3}{4} & -\frac{1}{4} & -\frac{1}{8} & -\frac{3}{40} \\ \frac{1}{4} & \frac{1}{12} & \frac{1}{24} & \frac{1}{40} \end{pmatrix}. \quad (55)$$

This leads to the same eigenvalues and eigenfunctions as above, and $\lambda_0 = 0$, a double eigenvalue. Its SVD gives two non-zero singular values, hence a rank of 2. Notice that the determinant of $M_{\mathcal{T}}$ is zero, and so is the case of the 3×3 subdeterminant extracted from the first three columns (but not the 2×2 subdeterminant of the first two columns), another way to conclude that the rank is indeed 2. This is a general fact for every point on the level sets in Fig. 2.

Remark IV.17. The discriminant condition $\tilde{\Delta} = 0$ corresponds to the level sets on Fig. 2. It is equivalent to $\tan \theta_1 = -1 + \tan \theta_0$, leading to two possibilities: $\theta_1 = \pi + \arctan(-1 + \tan \theta_0)$ for $\theta_0 \in [0, \pi/4) \cup (\pi/2, \pi)$, and $\theta_1 = \arctan(-1 + \tan \theta_0)$ for $\theta_0 \in [\pi/4, \pi/2)$. The case $(\theta_0, \theta_1) = (\pi/2, \pi/2)$ corresponds to Neumann BCs. Curious enough, along these level sets the expressions of resolvent, Riesz projection, Green's function, and perturbation kernel are different from the calculations in this subsection (since the discriminant is zero). The problem leads to a 4×4 matrix $M_{\mathcal{T}}$, but the rank of the perturbation is 2. The calculations are no different from the case $(\theta_0, \theta_1) = (\pi/4, 0)$. The details of these expressions are in appendix C.

K. Square of the Volterra Operator

The earliest treatment of the eigenvalues of the operator eq. (1) appears in the seminal paper of Lidskii⁴⁹, in the context of the Volterra operator—specifically the context of nonselfadjoint operators not satisfying the completeness property for Hilbert-Schmidt operators. We find it also in the book of Gohberg and Kreĭn⁵⁰ (pp. 208–210), where it is referred to as a “Volterra operator with a two dimensional imaginary component”.

With $\mathcal{V}f(x) := \int_x^1 f(y) dy$, one immediately obtains $\mathcal{V}^2 f(x) = \int_x^1 (x-y) f(y) dy$. This operator is then decomposed into its Hermitian components^{49,50,74}:

$$\mathcal{V}^2 = (\mathcal{V}^2)_R + i(\mathcal{V}^2)_I \quad (56)$$

where

$$\begin{aligned} (\mathcal{V}^2)_R f &:= \frac{1}{2} (\mathcal{V}^2 + \mathcal{V}^{2*}) f = -\frac{1}{2} \int_0^1 |x-y| f(y) dy, \\ (\mathcal{V}^2)_I f &:= \frac{1}{2i} (\mathcal{V}^2 - \mathcal{V}^{2*}) f = \frac{1}{2i} \int_0^1 (x-y) f(y) dy. \end{aligned} \quad (57)$$

The real part $(\mathcal{V}^2)_R$ is clearly our operator $\mathcal{K}$ in eq. (1). In this light, $\mathcal{K} = \mathcal{V}^2 + \mathcal{T}_{\mathcal{V}}$ where $\mathcal{T}_{\mathcal{V}} f = -i(\mathcal{V}^2)_I f = -\frac{1}{2} \int_0^1 (x-y) f(y) dy$.




FIG. 2. Examples of GSARS cases treated with respect to the discriminant condition $\cos(\theta_0 - \theta_1) + \cos(\theta_0 + \theta_1) - 2\sin(\theta_0 - \theta_1) = 0$, for $\theta_0, \theta_1 \in [0, \pi]$.

Theorem IV.18. *The operator $\mathcal{T}_\psi$ is a rank 2 perturbation of $\mathcal{K}$.*

Proof. To see this, we calculate the eigenvalues and eigenvectors of this perturbation, i.e., solve the eigenvalue problem $\mathcal{T}_\psi f = \lambda f$. This statement is equivalent to the problem

$$-\frac{1}{2} \int_0^1 (x-y) f(y) dy = \lambda f(x).$$

Note that $\lambda \neq 0$ otherwise $f(x) \equiv 0$. This immediately implies that $f(x) = mx + b$, where $m = -\frac{1}{2\lambda} \int_0^1 f(y) dy$ and $b = -\frac{1}{2\lambda} \int_0^1 y f(y) dy$. The problem then reduces to the matrix eigenvalue problem

$$\begin{pmatrix} -\frac{1}{4} & -\frac{1}{2} \\ \frac{1}{6} & \frac{1}{4} \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \lambda \begin{pmatrix} m \\ b \end{pmatrix}.$$

Thus $\lambda_1 = \frac{i}{2\sqrt{12}}$, $\lambda_2 = -\frac{i}{2\sqrt{12}}$, and the corresponding complex-valued eigenvectors $u_1(x) = \frac{1}{2}(-3 + i\sqrt{3})x + 1$, $u_2(x) = \frac{1}{2}(-3 - i\sqrt{3})x + 1$. This agrees with Lidskii⁴⁹. $\square$

V. CONCLUSION

In this article we offered a unified framework of looking at general real coupled self-adjoint BVPs, where the use of the resolvent kernel was essential to the analysis, and where the Green's formula corresponding to the integral operator formulation of Sturm-Liouville problems was obtained using an abstract and very classical formulation in¹⁹. While apparently similar in formulation to the nonlocal integral operator commuting with the Laplacian corresponding to the free space Green's function, the Krein-von Neumann problem was shown to be the “farthest”, while the anti-periodic problem proved to be the “closest” when seen as finite-rank perturbations. In⁵⁴, we will develop the spectral theory of the associate iterated *Brownian bridge kernels* corresponding to this GSARC framework, and show how to recover the values of the spectral zeta function for the nonlocal operator eq. (1) from the power series of the resolvent in a unified way; see also^{27,28,63,75}.

Appendix A: Details of the Green's Function for GSARC BCs with $\Delta=0$

The coefficients of $G_0(x,y)$ were generated using Wolfram Mathematica[®]. Noting also remark III.7, they are given by the following equations.

$$\begin{aligned} 20(3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta)^2 c_{00} &= 4(15 + 30\alpha + 21\alpha^2 + 5\alpha^3) \\ &\quad - (75 + 87\alpha + 28\alpha^2) \beta \\ &\quad + 2(27 + 17\alpha + 2\alpha^2) \beta^2 - (23 + 7\alpha) \beta^3 + 4\beta^4 \end{aligned}$$

$$\begin{aligned} 20(3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta)^2 c_{10} &= 4\alpha(15 + 30\alpha + 21\alpha^2 + 5\alpha^3) \\ &\quad - (30 + 135\alpha + 123\alpha^2 + 34\alpha^3) \beta \\ &\quad + 3(5 + 24\alpha + 11\alpha^2) \beta^2 + (9 - 7\alpha) \beta^3 - 4\beta^4 \end{aligned}$$

$$\begin{aligned} \frac{20}{3}(3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta)^2 c_{01} &= -4(15 + 25\alpha + 15\alpha^2 + 3\alpha^3) \\ &\quad + (110 + 115\alpha + 39\alpha^2 + 2\alpha^3) \beta \\ &\quad - (95 + 56\alpha + 9\alpha^2) \beta^2 + (43 + 11\alpha) \beta^3 - 8\beta^4 \end{aligned}$$

$$\begin{aligned} \frac{20}{3}(3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta)^2 c_{11} &= -4\alpha(15 + 25\alpha + 15\alpha^2 + 3\alpha^3) \\ &\quad + \alpha(100 + 95\alpha + 27\alpha^2) \beta \\ &\quad + 2(10 - 25\alpha - 11\alpha^2) \beta^2 - (25 - 7\alpha) \beta^3 + 8\beta^4 \end{aligned}$$

$$c_{20} = c_{02} = \frac{3(1 - \beta)^2}{2(3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta)}$$

$$c_{30} = c_{03} = \frac{(1 - \beta)(\alpha + \beta)}{2(3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta)}$$

$$c_{21} = c_{12} = \frac{3(1 - \beta)(\alpha + \beta)}{2(3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta)}$$

$$c_{31} = c_{13} = \frac{(\alpha + \beta)^2}{2(3 + \alpha^2 + \beta^2 - \alpha\beta + 3\alpha - 3\beta)}$$

Appendix B: Details of the perturbation kernel for GSARC BCs with $\Delta=0$

The following is the list of entries of the matrix associated with the GSARC case with zero discriminant in section IV I. They were calculated using Wolfram Mathematica[®]. The d_{ij} coefficients are defined by eq. (50). They are explicitly given by

$$\begin{aligned} d_{00} &= -c_{00} \\ d_{10} &= \frac{1}{2} - c_{10} = -c_{01} - \frac{1}{2} \\ d_{20} &= -c_{20} = -c_{02} \end{aligned}$$

$$\begin{aligned}
d_{11} &= -c_{11} \\
d_{30} &= -c_{30} = -c_{03} \\
d_{21} &= -c_{21} = -c_{12} \\
d_{31} &= -c_{31} = -c_{13}
\end{aligned} \tag{B1}$$

$$m_{11} = d_{00} + \frac{d_{10}}{2} + \frac{d_{20}}{3} + \frac{d_{30}}{4}$$

$$m_{12} = \frac{d_{00}}{2} + \frac{d_{10}}{3} + \frac{d_{20}}{4} + \frac{d_{30}}{5}$$

$$m_{13} = \frac{d_{00}}{3} + \frac{d_{10}}{4} + \frac{d_{20}}{5} + \frac{d_{30}}{6}$$

$$m_{14} = \frac{d_{00}}{4} + \frac{d_{10}}{5} + \frac{d_{20}}{6} + \frac{d_{30}}{7}$$

$$m_{21} = d_{10} + \frac{d_{11}}{2} + \frac{d_{31}}{4}$$

$$m_{22} = \frac{d_{10}}{2} + \frac{d_{11}}{3} + \frac{d_{31}}{5}$$

$$m_{23} = \frac{d_{10}}{3} + \frac{d_{11}}{4} + \frac{d_{31}}{6}$$

$$m_{24} = \frac{d_{10}}{4} + \frac{d_{11}}{5} + \frac{d_{31}}{7}$$

$$m_{31} = d_{20}$$

$$m_{32} = \frac{d_{20}}{2}$$

$$m_{33} = \frac{d_{20}}{3}$$

$$m_{34} = \frac{d_{20}}{4}$$

$$m_{41} = d_{30} + \frac{d_{31}}{2}$$

$$m_{42} = \frac{d_{30}}{2} + \frac{d_{31}}{3}$$

$$m_{43} = \frac{d_{30}}{3} + \frac{d_{31}}{4}$$

$$m_{44} = \frac{d_{30}}{4} + \frac{d_{31}}{5}.$$

In terms of the GSARC parameters, α, β (when the discriminant is equal to zero), after a lot of simplifications using Wolfram Mathematica[®], the matrix elements reduce to the following:

$$\begin{aligned} 40(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)^2 m_{11} = & -(90+195\alpha+173\alpha^2+69\alpha^3+10\alpha^4) \\ & + (165+224\alpha+114\alpha^2+19\alpha^3)\beta \\ & - (143+120\alpha+31\alpha^2)\beta^2+3(19+7\alpha)\beta^3-9\beta^4 \end{aligned}$$

$$\begin{aligned} 120(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)^2 m_{12} = & -(135+291\alpha+273\alpha^2+120\alpha^3+20\alpha^4) \\ & + 2(147+201\alpha+108\alpha^2+20\alpha^3)\beta \\ & - 3(90+77\alpha+21\alpha^2)\beta^2+4(27+10\alpha)\beta^3-17\beta^4 \end{aligned}$$

$$\begin{aligned} 240(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)^2 m_{13} = & -2(93+198\alpha+189\alpha^2+86\alpha^3+15\alpha^4) \\ & + (438+597\alpha+325\alpha^2+62\alpha^3)\beta \\ & - (411+352\alpha+97\alpha^2)\beta^2+(165+61\alpha)\beta^3-26\beta^4 \end{aligned}$$

$$\begin{aligned} 2800(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)^2 m_{14} = & -4(420+885\alpha+850\alpha^2+393\alpha^3+70\alpha^4) \\ & + (4125+5605\alpha+3064\alpha^2+592\alpha^3)\beta \\ & - 2(1955+1673\alpha+462\alpha^2)\beta^2+(1573+581\alpha)\beta^3-248\beta^4 \end{aligned}$$

$$\begin{aligned} 40(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)^2 m_{21} = & (180+420\alpha+345\alpha^2+117\alpha^3+11\alpha^4) \\ & - 2(150+270\alpha+147\alpha^2+29\alpha^3)\beta \\ & + 3(65+87\alpha+20\alpha^2)\beta^2-4(12+13\alpha)\beta^3-\beta^4 \end{aligned}$$

$$\begin{aligned} 40(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)^2 m_{22} = & 2(45+120\alpha+109\alpha^2+42\alpha^3+5\alpha^4) \\ & - (150+329\alpha+199\alpha^2+44\alpha^3)\beta \\ & + (83+160\alpha+41\alpha^2)\beta^2-(7+31\alpha)\beta^3-6\beta^4 \end{aligned}$$

$$\begin{aligned} 80(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)^2 m_{23} = & 4(30+85\alpha+80\alpha^2+32\alpha^3+4\alpha^4) \\ & - (200+480\alpha+301\alpha^2+69\alpha^3)\beta \\ & + 2(50+117\alpha+31\alpha^2)\beta^2+3(1-15\alpha)\beta^3-12\beta^4 \end{aligned}$$

$$2800(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)^2 m_{24} = 2(1575+4620\alpha+4425\alpha^2+1800\alpha^3+229\alpha^4)$$

$$-(5250 + 13275\alpha + 8475\alpha^2 + 1978\alpha^3)\beta + 3(815 + 2160\alpha + 581\alpha^2)\beta^2 + (285 - 1243\alpha)\beta^3 - 382\beta^4$$

$$m_{31} = -\frac{3(1-\beta)^2}{2(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)}$$

$$m_{32} = -\frac{3(1-\beta)^2}{4(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)}$$

$$m_{33} = -\frac{(1-\beta)^2}{2(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)}$$

$$m_{34} = -\frac{3(1-\beta)^2}{8(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)}$$

$$m_{41} = -\frac{(2+\alpha-\beta)(\alpha+\beta)}{4(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)}$$

$$m_{42} = -\frac{(3+2\alpha-\beta)(\alpha+\beta)}{12(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)}$$

$$m_{43} = -\frac{(4+3\alpha-\beta)(\alpha+\beta)}{24(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)}$$

$$m_{44} = -\frac{(5+4\alpha-\beta)(\alpha+\beta)}{40(3+3\alpha-3\beta-\alpha\beta+\alpha^2+\beta^2)}.$$

Appendix C: Details of the perturbation kernel for GSARS BCs with $\tilde{\Delta}=0$

In either case $\theta_1 = \pi + \arctan(-1 + \tan \theta_0)$ for $\theta_0 \in [0, \pi/4) \cup (\pi/2, \pi)$, or $\theta_1 = \arctan(-1 + \tan \theta_0)$ for $\theta_0 \in [\pi/4, \pi/2)$, for $0 \leq x \leq y \leq 1$, the resolvent is such that

$$(2\sqrt{z}\cos\sqrt{z}\cos^2\theta_0 + \sin\sqrt{z}(-1-z-(1-z)\cos 2\theta_0 + z\sin 2\theta_0))G(z, x, y)$$

is equal to

$$\begin{aligned} & 2\sin\theta_0(\cos\theta_0\sin(\sqrt{z}(1-y)) + \sqrt{z}\cos(\sqrt{z}(1-y))(\sin\theta_0 - \cos\theta_0))\cos(\sqrt{z}x) \\ & + 2\cos\theta_0(-\cos\theta_0\sin(\sqrt{z}(1-y)) + \sqrt{z}\cos(\sqrt{z}(1-y))(\cos\theta_0 - \sin\theta_0))\frac{\sin(\sqrt{z}x)}{\sqrt{z}}. \end{aligned}$$

Calculating $\lim_{z \rightarrow 0^+} z^n G(z, x, y)$ for $n = 1, 2, \dots$ give a Riesz projection with a kernel

$$p(x, y) = \frac{6(x\cos\theta_0 - \sin\theta_0)(y\cos\theta_0 - \sin\theta_0)}{-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0}$$

and a nilpotent operator $D \equiv 0$. Note by simple calculus that the expression $-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0$ is never zero, since it has a maximum value of $-4 + \sqrt{13} \approx -0.394$ at $\theta_0 = \frac{1}{2}\arctan \frac{3}{2}$ and minimum value of $-4 + \sqrt{13} \approx -7.606$ at $\theta_0 = \frac{\pi}{2} + \frac{1}{2}\arctan \frac{3}{2}$. The Green's function is such that

$$-20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 G_0(x, y)$$

is equal to

$$a_0(x, y) + a_2(x, y) \cos 2\theta_0 + a_4(x, y) \cos 4\theta_0 + b_2(x, y) \sin 2\theta_0 + b_4(x, y) \sin 4\theta_0,$$

where

$$\begin{aligned} a_0(x, y) &= -132 - 90x + 360y - 150x^2 + 204xy - 150y^2 - 15x^3 - 45x^2y - 45xy^2 - 15y^3 - 30x^3y - 30xy^3 \\ a_2(x, y) &= 120 - 40x - 360y + 180x^2 + 72xy + 180y^2 - 20x^3y - 20xy^3 \\ a_4(x, y) &= 12 + 50x - 30x^2 - 132xy - 30y^2 + 15x^3 + 45x^2y + 45xy^2 + 15y^3 + 10x^3y + 10xy^3 \\ b_2(x, y) &= 140 + 144x - 336y + 90x^2 - 360xy + 90y^2 + 40x^3 + 120x^2y + 120xy^2 + 40y^3 + 30x^3y + 30xy^3 \\ b_4(x, y) &= -50 + 12x + 132y - 45x^2 - 45y^2 - 10x^3 - 30x^2y - 30xy^2 - 10y^3 + 15x^3y + 15xy^3 \end{aligned}$$

With $T(x, y) = \kappa(x, y) - G_0(x, y)$, the expression of the integral operator $\mathcal{T}f(x)$ is a cubic polynomial, hence the eigenvector takes the form $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3$ and the corresponding 4×4 matrix $M_{\mathcal{T}} = (m_{ij})$ is such that

$$\begin{aligned} -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{11} &= \frac{473}{4} - 80\cos 2\theta_0 - \frac{73}{4}\cos 4\theta_0 - 132\sin 2\theta_0 + \frac{63}{2}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{12} &= \frac{123}{4} - \frac{115}{3}\cos 2\theta_0 - \frac{59}{6}\cos 4\theta_0 - \frac{137}{2}\sin 2\theta_0 + \frac{57}{4}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{13} &= \frac{171}{4} - 26\cos 2\theta_0 - \frac{27}{4}\cos 4\theta_0 - \frac{142}{3}\sin 2\theta_0 + \frac{28}{3}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{14} &= \frac{232}{7} - 20\cos 2\theta_0 - \frac{36}{7}\cos 4\theta_0 - \frac{1278}{35}\sin 2\theta_0 + \frac{246}{35}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{21} &= -\frac{429}{2} + 169\cos 2\theta_0 + \frac{47}{2}\cos 4\theta_0 + \frac{457}{2}\sin 2\theta_0 - \frac{263}{4}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{22} &= -\frac{473}{4} + 80\cos 2\theta_0 + \frac{73}{4}\cos 4\theta_0 + 132\sin 2\theta_0 - \frac{63}{2}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{23} &= -82 + 52\cos 2\theta_0 + 14\cos 4\theta_0 + 93\sin 2\theta_0 - \frac{41}{2}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{24} &= -\frac{8787}{140} + \frac{1346}{35}\cos 2\theta_0 + \frac{1571}{140}\cos 4\theta_0 + \frac{502}{7}\sin 2\theta_0 - \frac{106}{7}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{31} &= \frac{345}{2} - 180\cos 2\theta_0 + \frac{15}{2}\cos 4\theta_0 - 150\sin 2\theta_0 + 60\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{32} &= 90 - 90\cos 2\theta_0 - 85\sin 2\theta_0 + \frac{65}{2}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{33} &= \frac{245}{4} - 60\cos 2\theta_0 - \frac{5}{4}\cos 4\theta_0 - 60\sin 2\theta_0 + \frac{45}{2}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{34} &= \frac{93}{2} - 45\cos 2\theta_0 - \frac{3}{2}\cos 4\theta_0 - \frac{93}{2}\sin 2\theta_0 + \frac{69}{4}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{41} &= 30 + 10\cos 2\theta_0 - 20\cos 4\theta_0 - 55\sin 2\theta_0 + \frac{5}{2}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{42} &= \frac{35}{2} + \frac{20}{3}\cos 2\theta_0 - \frac{65}{6}\cos 4\theta_0 - 30\sin 2\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{43} &= \frac{25}{2} + 5\cos 2\theta_0 - \frac{15}{2}\cos 4\theta_0 - \frac{125}{6}\sin 2\theta_0 + \frac{5}{12}\sin 4\theta_0 \\ -20(-4 + 2\cos 2\theta_0 + 3\sin 2\theta_0)^2 m_{44} &= \frac{39}{4} + 4\cos 2\theta_0 - \frac{23}{4}\cos 4\theta_0 - 16\sin 2\theta_0 - \frac{1}{2}\sin 4\theta_0. \end{aligned}$$

The determinant of $M_{\mathcal{T}}$ is zero and so is the case of the 3×3 subdeterminant extracted from the first three columns (but not the 2×2 subdeterminant of the first two columns). Hence the rank of the perturbation is 2. A calculation on Wolfram Mathematica[®] confirms this, and shows that $\lambda = 0$ is a double eigenvalue.

Acknowledgments

We would like to thank Fritz Gesztesy for introducing us to some aspects of this problem and for sharing an advanced copy of⁷⁰; Jussi Behrndt for reference⁶⁹; and Mark Ashbaugh for useful discussions. We acknowledge email correspondence with Walter A. Strauss regarding the “unusual” eigenvalue problem discussed in Sec. III D and thank him for his feedback. The authors thank the anonymous referee whose comments helped improve the article significantly.

L.H. thanks and acknowledges N.S.’s NSF grant DMS-1912747 during his visits to the UC Davis campus in 2019 and 2022 where some of this work was done.

N.S. was partially supported by his NSF grants DMS-1912747, CCF-1934568, and his ONR grant N00014-20-1-2381.

REFERENCES

- ¹N. Saito, “Data analysis and representation on a general domain via eigenfunctions of Laplacian,” *Appl. Comput. Harmon. Anal.* **25**, 68–97 (2008).
- ²L. Hermi and N. Saito, “On Rayleigh-type formulas for a non-local boundary value problem associated with an integral operator commuting with the Laplacian,” *Appl. Comput. Harmon. Anal.* **45**, 59–83 (2018).
- ³L. Greengard and V. Rokhlin, “A fast algorithm for particle simulation,” *J. Comput. Phys.* **73**, 325–348 (1987).
- ⁴X. Xue, *On a Fast Algorithm for Computing the Laplacian Eigenpairs via Commuting Integral Operators*, Ph.D. thesis, Dept. Math., UC Davis (2007), 96 pp.
- ⁵I. Gel’fand and B. Levitan, “On a simple identity for the characteristic values of a differential operator of the second order,” *Doklady Akad. Nauk SSSR. (N.S.)* **88**, 593–596 (1953), (Russian).
- ⁶M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (CRC Press, 1995).
- ⁷D. J. Griffiths and D. F. Schroeter, *Introduction to Quantum Mechanics* (Cambridge University Press, 2018).
- ⁸K. Aki and P. G. Richards, *Quantitative Seismology* (University Science Books, Sausalito, CA, 2002).
- ⁹M. A. Carreira-Perpiñán, “A review of dimension reduction techniques,” *Tech. Rep. CS-96-09* (Dept. of Computer Science, University of Sheffield, UK, 1996) 69 pp.
- ¹⁰C. K. Chui and J. Z. Wang, “Dimensionality reduction of hyperspectral imagery data for feature classification,” in *Handbook of Geomathematics*, Vol. 2, edited by W. Freeden, M. Z. Nashed, and T. Sonar (2010) pp. 1005–1049.
- ¹¹L. Rosasco, M. Belkin, and E. D. Vito, “On learning with integral operators,” *J. of Machine Learning Research* **11**, 905–934 (2010).
- ¹²U. von Luxburg, “A tutorial on spectral clustering,” *Stat. Comput.* **17**, 395–416 (2007).
- ¹³U. von Luxburg, M. Belkin, and O. Bousquet, “Consistency of spectral clustering,” *Ann. Statist.* **36**, 555–586 (2008).
- ¹⁴U. von Luxburg, O. Bousquet, and M. Belkin, “On the convergence of spectral clustering on random samples: the normalized case,” in *Proceedings of the 17th Annual Conference on Learning Theory (COLT 2004)* (Springer, 2004) pp. 457–471.
- ¹⁵Y. Yao, L. Rosasco, and A. Caponnetto, “On early stopping in gradient descent learning,” *Constr. Approx.* **26**, 289–315 (2007).
- ¹⁶J. A. Lee and M. Verleysen, *Nonlinear Dimensionality Reduction* (Springer, 2007).
- ¹⁷L. Lieu and N. Saito, “Signal ensemble classification using low-dimensional embeddings and Earth Mover’s Distance,” in *Wavelets and Multiscale Analysis: Theory and Applications*, Applied and Numerical Harmonic Analysis, edited by J. Cohen and A. I. Zayed (Birkhäuser, Boston, MA, 2011) Chap. 11, pp. 227–256.
- ¹⁸L. J. P. van der Maaten, E. O. Postma, and H. J. van den Herik, “Dimensionality reduction: A comparative review,” *Technical Report TiCC-TR 2009-005* (Tilburg Centre for Creative Computing, Tilburg Univ., 2009).
- ¹⁹T. Kato, *Perturbation Theory for Linear Operators*, Classics in Mathematics (Springer-Verlag, Berlin, Germany, 1995).
- ²⁰M. Riesz, “L’intégrale de Riemann-Liouville et potentiels,” *Acta Sci. Math. (Szeged)*, 1–42 (1938–40), (French).
- ²¹M. Riesz, “L’intégrale de Riemann-Liouville et le problème de Cauchy,” *Acta Math.* **81**, 1–223 (1949), (French).
- ²²I. J. Schoenberg, “On certain metric spaces arising from Euclidean spaces by a change of metric and their imbedding in Hilbert space,” *Ann. of Math.* **38**, 787–793 (1937).
- ²³I. J. Schoenberg, “Metric spaces and completely monotone functions,” *Ann. of Math.* **39**, 811–841 (1938).
- ²⁴I. J. Schoenberg, “Metric spaces and positive definite functions,” *Trans. Amer. Math. Soc.* **44**, 522–536 (1938).
- ²⁵G. Hellwig, *Differential Operators of Mathematical Physics* (Addison-Wesley, Reading, Massachusetts, 1964).
- ²⁶F. Gesztesy and K. Kirsten, “Effective computation of traces, determinants, and ζ -functions for Sturm-Liouville operators,” *J. Funct. Anal.* **276**, 520–562 (2019).
- ²⁷M. S. Ashbaugh, F. Gesztesy, L. Hermi, K. Kirsten, L. Littlejohn, and H. Tossounian, “Trace formulas applied to the Riemann ζ -function,” in *Integrability, Supersymmetry and Coherent States*, CRM Ser. Math. Phys. (Springer, Cham, 2019) pp. 231–253.
- ²⁸M. S. Ashbaugh, F. Gesztesy, L. Hermi, K. Kirsten, L. Littlejohn, and H. Tossounian, “Green’s functions and Euler’s formula for $\zeta(2n)$,” in *Schrödinger Operators, Spectral Analysis and Number Theory: In Memory of Erik Balslev*, edited by S. Albeverio and R. Weder (Springer, Cham, 2021) pp. 27–45.
- ²⁹F. Gesztesy and K. Kirsten, “Computing traces, determinants, and ζ -functions for Sturm-Liouville operators: a survey,” in *Analysis and Operator Theory*, Springer Optim. Appl., Vol. 146 (Springer, Cham, 2019) pp. 93–148.
- ³⁰F. Gesztesy and K. Kirsten, “On traces and modified Fredholm determinants for half-line Schrödinger operators with purely discrete spectra,” *Quart. Appl. Math.* **77**, 615–630 (2019).
- ³¹D. Porter and D. S. G. Stirling, *Integral Equations: A Practical Treatment from Spectral Theory to Applications* (Cambridge University Press, Cambridge, UK, 1990).
- ³²G. Fucci, F. Gesztesy, K. Kirsten, L. L. Littlejohn, R. Nichols, and J. Stanfill, “The Krein–von Neumann extension revisited,” *Appl. Anal.* **101**, 1593–1616 (2022).
- ³³G. Fucci, F. Gesztesy, K. Kirsten, and J. Stanfill, “Spectral ζ -functions and ζ -regularized functional determinants for regular Sturm-Liouville operators,” *Res. Math. Sci.* **8**, Paper No. 61, 46 (2021).
- ³⁴G. B. Folland, *Fourier Analysis and Its Applications* (Amer. Math. Soc., Providence, RI, 1992) republished by AMS, 2009.
- ³⁵P. B. Bailey, W. N. Everitt, and A. Zettl, “Regular and singular Sturm-Liouville problems with coupled boundary conditions,” *Proc. Roy. Soc. Edinburgh Sect. A* **126**, 505–514 (1996).
- ³⁶Q. Kong and A. Zettl, “Dependence of eigenvalues of Sturm-Liouville problems on the boundary,” *J. Differential Equations* **126**, 389–407 (1996).

- ³⁷Q. Kong and A. Zettl, “Eigenvalues of regular Sturm-Liouville problems,” *J. Differential Equations* **131**, 1–19 (1996).
- ³⁸M. S. P. Eastham, Q. Kong, H. Wu, and A. Zettl, “Inequalities among eigenvalues of Sturm-Liouville problems,” *J. Inequal. Appl.* **3**, 25–43 (1999).
- ³⁹A. Zettl, *Sturm-Liouville theory* (American Mathematical Society, Providence, RI, 2005).
- ⁴⁰R. J. de la Cruz and E. Reyes, “On the Iwasawa decomposition of a perplectic matrix,” *Comm. Algebra* **49**, 932–947 (2021).
- ⁴¹Q. Kong, H. Wu, and A. Zettl, “Geometric aspects of Sturm-Liouville problems. I. Structures on spaces of boundary conditions,” *Proc. Roy. Soc. Edinburgh Sect. A* **130**, 561–589 (2000).
- ⁴²F. V. Atkinson, *Discrete and Continuous Boundary Problems*, Mathematics in Science and Engineering, Vol. 8 (Academic Press, New York, 1964).
- ⁴³A. Wang, J. Sun, and A. Zettl, “The classification of self-adjoint boundary conditions: separated, coupled, and mixed,” *J. Funct. Anal.* **255**, 1554–1573 (2008).
- ⁴⁴A. Wang, J. Sun, and A. Zettl, “The classification of self-adjoint boundary conditions of differential operators with two singular endpoints,” *J. Math. Anal. Appl.* **378** (2011), 493–506 (2011).
- ⁴⁵M. S. Ashbaugh, F. Gesztesy, A. Laptev, M. Mitrea, and S. Sukhtaiev, “A bound for the eigenvalue counting function for Krein-von Neumann and Friedrichs extensions,” *Adv. Math.* **304**, 1108–1155 (2017).
- ⁴⁶M. S. Ashbaugh, F. Gesztesy, M. Mitrea, R. Shterenberg, and G. Teschl, “A survey on the Krein–von Neumann extension, the corresponding abstract buckling problem, and Weyl-type spectral asymptotics for perturbed Krein Laplacians in nonsmooth domains,” in *Mathematical Physics, Spectral Theory and Stochastic Analysis*, Oper. Theory Adv. Appl., Adv. Partial Differ. Equ. (Basel), Vol. 232 (Birkhäuser/Springer Basel AG, Basel, 2013) pp. 1–106.
- ⁴⁷M. S. Ashbaugh, F. Gesztesy, M. Mitrea, R. Shterenberg, and G. Teschl, “The Krein-von Neumann extension and its connection to an abstract buckling problem,” *Math. Nachr.* **283**, 165–179 (2010).
- ⁴⁸A. Alonso and B. Simon, “The Birman-Kreĭn-Vishik theory of selfadjoint extensions of semibounded operators,” *J. Operator Theory* **4**, 251–270 (1980).
- ⁴⁹V. B. Lidskii, “Conditions for completeness of a system of root subspaces for non-selfadjoint operators with discrete spectra,” *Tr. Mosk. Mat. Obs.* **8**, 83–120 (1959), english transl.in: Amer. Math. Soc. Translations: Series 2, Vol. 34: Eleven Papers on Analysis. Amer. Math. Soc., Providence, RI, 1963, pp. 241–281.
- ⁵⁰I. C. Gohberg and M. G. Kreĭn, *Introduction to the Theory of Linear Nonselfadjoint Operators in Hilbert Space*, Translations of Mathematical Monographs, Vol. 18 (Amer. Math. Soc., Providence, R.I., 1969) translated from the Russian by A. Feinstein.
- ⁵¹R. Bapat, S. J. Kirland, and M. Neumann, “On distance matrices and Laplacians,” *Linear Algebra Appl.* **401**, 193–209 (2005).
- ⁵²W. So, “Rank one perturbation and its application to the Laplacian spectrum of a graph,” *Linear and Multilinear Algebra* **46**, 193–198 (1999).
- ⁵³I. Stakgold and M. Holst, *Green’s functions and boundary value problems*, Pure and Applied Mathematics (John Wiley & Sons, Inc., Hoboken, NJ, 2011).
- ⁵⁴L. Hermi and N. Saito, “On a nonlocal integral operator commuting with the Laplacian and the Sturm-Liouville problem II: Iterated Brownian bridge kernels,” Unpublished (2024).
- ⁵⁵K. Yosida, *Functional Analysis*, 6th ed. (Springer-Verlag, Berlin, 1980).
- ⁵⁶N. Dunford and J. T. Schwartz, *Linear Operators - Part I: General Theory* (Interscience Publishers, Inc., New York, 1963).
- ⁵⁷A. E. Taylor and D. C. Lay, *Introduction to Functional Analysis*, 2nd ed. (Robert E. Krieger Publishing Company, Malabar, FL, 1980).
- ⁵⁸M. Nagumo, “Einige analytische Untersuchungen in linearen metrischen Ringen,” *Japan J. Math.* **13**, 61–80 (1936).
- ⁵⁹R. Kress, *Linear Integral Equations*, 3rd ed., Applied Mathematical Sciences, Vol. 82 (Springer-Verlag, New York, 2014).
- ⁶⁰W. A. Strauss, *Partial Differential Equations: An Introduction*, 2nd ed. (John Wiley & Sons, Inc., New York, 2008).
- ⁶¹K. Schmüdgen, *Unbounded Self-adjoint Operators on Hilbert Space*, Graduate Texts in Mathematics, Vol. 265 (Springer, New York, 2012).
- ⁶²L. A. Dikii, “Trace formulas for Sturm-Liouville differential operators,” *Amer. Math. Soc. Transl. (2)* **18**, 81—115 (1961), translated by J. B. Diaz.
- ⁶³R. Cavoretto, G. E. Fasshauer, and M. McCourt, “An introduction to the Hilbert-Schmidt SVD using iterated Brownian bridge kernels,” *Numer. Algor.* **68**, 393–422 (2015).
- ⁶⁴K. Gustafson and T. Abe, “The third boundary condition — was it Robin’s?” *Math. Intell.* **20**, 63–71 (1998).
- ⁶⁵K. Gustafson and T. Abe, “(Victor) Gustave Robin: 1855–1897,” *Math. Intell.* **20**, 47–53 (1998).
- ⁶⁶F. Gesztesy and M. Mitrea, “Generalized Robin boundary conditions, Robin-to-Dirichlet maps, and Krein-type resolvent formulas for Schrödinger operators on bounded Lipschitz domains,” in *Perspectives in partial differential equations, harmonic analysis and applications*, Proc. Sympos. Pure Math., Vol. 79 (Amer. Math. Soc., Providence, RI, 2008) pp. 105–173.
- ⁶⁷W. Freeden, *Metaharmonic Lattice Point Theory*, Pure and Applied Mathematics, Vol. 299 (CRC Press, Boca Raton, FL, 2011).
- ⁶⁸C. Radoux, “Seies liees aux racines de l’equation $\operatorname{tg} x = x$,” *Bull. Soc. Math. Belg., ser. B* **40**, 199–205 (1988).
- ⁶⁹J. Behrndt, S. Hassi, and H. de Snoo, *Boundary value problems, Weyl functions, and differential operators*, Monographs in Mathematics, Vol. 108 (Birkhäuser/Springer, Cham, 2020) pp. vii+772.
- ⁷⁰F. Gesztesy, R. Nichols, and M. Zinchenko, *Sturm-Liouville Operators, Their Spectral Theory, and Some Applications* (American Mathematical Society, 2023).
- ⁷¹Y. A. Abramovich and C. D. Aliprantis, *An Invitation to Operator Theory*, Graduate Studies in Mathematics (Amer. Math. Soc., Providence, RI, 2002).
- ⁷²Y. A. Abramovich and C. D. Aliprantis, *Problems in Operator Theory*, Graduate Studies in Mathematics (Amer. Math. Soc., Providence, RI, 2002).
- ⁷³L. N. Trefethen and D. Bau, III, *Numerical Linear Algebra* (SIAM, Philadelphia, PA, 1997).
- ⁷⁴Y. Arlinskii and E. Tsekanovskii, “Non-self-adjoint Jacobi matrices with a rank-one imaginary part,” *J. Funct. Anal.* **241**, 383–438 (2006).
- ⁷⁵G. Fasshauer and M. McCourt, *Kernel-based Approximation Methods using MATLAB*, Interdisciplinary Mathematical Sciences, Vol. 19 (World Sci. Publ., Singapore, 2015).