

ARTICLE

Facile Synthesis of Diiodoheteroindenes and Understanding Their Sonogashira Cross-Coupling Selectivity for the Construction of Unsymmetrical Enediynes†

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Alexander V. Ponomarev,^a Natalia A. Danilkina,^a Julia S. Okuneva,^a Aleksandra A. Vidyakina,^a Ekaterina A. Khmelevskaya,^a Alexander S. Bunev,^b Andrey M. Rumyantsev,^c Anastasia I. Govdi,^a Thomas Suarez,^d Igor V. Alabugin,^{*d} Irina A. Balova^{*a}

Electrophile-promoted cyclizations of functionalized alkynes offer a useful tool for constructing halogen-substituted heterocycles primed for further derivatization. Preinstallation of an iodo-substituent at the alkyne prior to the iodocyclization opens access to *ortho* di-iodinated heterocyclic precursors for the preparation of unsymmetrical heterocycle-fused enediynes. This general approach was used to prepare 2,3-diiodobenzothiophene, 2,3-diidoindole, and 2,3-diiodobenzofuran, a useful family of substrates for systematic studies for the role of heteroatom on the regioselectivity of cross-coupling reactions. Diiodobenzothiophene showed much higher regioselectivity for Sonogashira cross-coupling at C2 than diidoindole and diiodobenzofuran. As a result, the benzothiophene can be conveniently involved in a one-pot sequential coupling with two different alkynes, yielding unsymmetrical benzothiophen-fused enediynes. On other hand, the Sonogashira reaction of diidoindole and diiodobenzofuran formed considerable amounts of di-substituted enediyne in addition to the monoalkyne product at coupling at C2. Interestingly, no C3-monocoupling products was observed for all of the diiodides, suggesting that incorporation of the 1st alkyne at C2 activates the C3 position for the 2nd coupling. Additional factors affecting regioselectivity were detected, discussed and connected, through computational analysis, to transmetalation being the rate-determining step for the Sonogashira reaction. Several enediynes synthesized showed cytotoxic activity, which is not associated with DNA strand breaks typical of natural enediyne antibiotics.

Introduction

Electrophile-promoted cyclization of functionalized alkynes is a valuable tool for the synthesis of halogen-substituted heterocycles.^{1–7} The reaction is suitable for different nucleophilic functional groups and alkynes. The most recent examples of electrophile-promoted cyclization for «halogen» electrophiles open access to oxazinoindolones,⁸ 3-sulfonamido-4-haloisocoumarins,⁹ sulfonamido-substituted polycyclic aromatic compounds,¹⁰ medium-sized oxacycles,¹¹ iodofuranones,¹² iodothiazines,¹³ highly distorted fused helicenes^{14,15} and optically active tricyclic piperazine scaffolds through the dearomatic *ipso*-cyclization.¹⁶ Electrophile-promoted cyclization of various functionalized alkynes has been applied to construct polyheterocycles targeting nucleic acids.¹⁷

Earlier we have shown that iodocyclization of *ortho*-functionalized arylidacetylenes followed by the Sonogashira reaction is a convenient approach for the construction of unsymmetrically substituted enediyne systems fused to heteroindenes with the highest yields and functional group tolerance in the case of benzothiophene and indole (Fig. 1A).¹⁸ This «diacetylenic approach» was used in the synthesis of 10-membered heterocyclic analogues of enediyne antibiotics^{19–22}. However, this method has some limitations associated with the possibility of iodination of the second triple bond in the case of benzofuran and isocoumarin derivatives.^{18,23,24} An alternative way to obtain unsymmetrically substituted enediynes fused to heterocycles could be the regioselective stepwise substitution of different halogens, which has been applied for pyrimidine-fused enediynes (Fig. 1B).²⁵ Here we investigate the applicability of this approach under one-pot conditions for 2,3-diiodoheteroindenes (Fig. 1C).

The 2,3-diiodoheteroindenes can be obtained by several ways (Figure 2). The first method is the direct electrophilic iodination which has been applied to indoles^{26,27}, benzothiophene²⁸, and 2-iodoindole.²⁹ The second approach is based on the sequences of metalation / iodine exchange steps which has been reported for diidoindoles.³⁰ All three compounds can also be obtained by iodinative decarboxylation techniques, either as the desired products (reported for indoles,^{31,32} benzothiophene³³) or as by-products (reported for benzo-

^a Institute of Chemistry, Saint Petersburg State University (SPbU), Saint Petersburg, 199034, Russia. E-mail: i.balova@spbu.ru

^b Medicinal Chemistry Center, Tolyatti State University, 445020 Tolyatti, Russia

^c Department of Genetics and Biotechnology, Saint Petersburg State University (SPbU), Saint Petersburg, 199034, Russia

^d Department of Chemistry and Biochemistry, Florida State University, Tallahassee, FL 32306, USA. E-mail: alabugin@chem.fsu.edu

† Dedicated to the 300th anniversary of the Saint Petersburg University foundation
Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x




Fig. 1 Known examples of the synthesis of unsymmetrical acyclic and cyclic enediyne systems through the "diacylenic" (A), "stepwise" (B) approaches and proposed «iodoalkyne» approaches (C).




Fig. 2 Literature approaches to the synthesis of 2,3-diiodoheteroindenes.

thiophene and benzofuran,³⁴ albeit without the support of spectral data for 2,3-diiodobenzofuran). The highest yield for diiodobenzothiophene is reachable by the synthesis of 3-iodo-2-trimethylsilyl derivative followed by the Si-iodine exchange using ICl.^{35,36}

In principle, iodocyclization of the *ortho*-functionalized iodoethynylarenes can be considered as a convenient alternative synthetic approach to diiodoheteroindenes (Figure

1). However, to the best of our knowledge, the possibility of using electrophile-promoted cyclizations as a route to diiodoheteroindenes, has not been studied, despite the fact that electrophile-promoted cyclization have been used as a synthetic tool for benzo[b]thiophenes with different halogen substituents,³⁷ as well as dihalothieno[2,3-*b*]quinoline and dihaloselenopheno[2,3-*b*]quinoline.³⁸ The regioselectivity in CC cross-coupling reaction is an important problem that is being studied both theoretically and experimentally.³⁹ It is known that the halogen atoms in 2,3-diiodo- and 2,3-dibromothiophenes as well as in 2,3-dibromoindoles have different reactivity in cross-coupling reactions.⁴⁰ Because the C2 position is more reactive than the C3 position, regioselective Sonogashira reactions at the C2 position have been reported for 2,3-dibromothiophene derivatives^{41–43}, 2,3-dibromofuran,⁴³ 2,3-diiodothiophenes,⁴⁴ diiodothienopyridine,⁴⁵ 2,3-dibromobenz[*b*]thiophene (Figure 3).⁴⁶ Even tetrabromothiophene can undergo regioselective C2,C5 Sonogashira coupling.⁴³ Not surprisingly, regioselective C2-I Sonogashira coupling has also been shown for C2-iodo/C3-bromo derivatives: 3-bromo-2-iodobenz[*b*]thiophene⁴³ and 3-bromo-2-iodoindole⁴⁷ where the more reactive halogen was placed at an intrinsically more reactive position. In the case of C2-bromo/C3-iodo derivatives, where the two above factors are in conflict, the C2/C3 regioselectivity is reversed. Thus, C3-iodo regioselectivity has been observed for 2-bromo-3-iodothiophenes,^{48–51} 2-bromo-3-iodofuran and -benzofuran.⁵² However, the selectivity of cross-coupling in 2,3-diiodoheteroindenes (benzo[*b*]thiophene, indole, and benzo[*b*]furan) has not been previously studied. 2,3-Diiodoindoles were only used in the synthesis of symmetrical diethynylindoles.⁵³

In this work, we report on the study of electrophile-promoted cyclization of functionalized iodoethynylarenes and the investigation of the regioselectivity of the Sonogashira reaction for 2,3-diiodoheteroindenes for the development of a new «iodoalkyne» approach to unsymmetrically substituted enediynes fused to heteroindenes (Figure 1).




Fig. 3 Known regioselectivity for the Sonogashira reaction of dihalo thiophenes, furans and heteroindenes.

Results and Discussion


Study on Electrophile-Promoted Cyclization of Iodoalkynes

The starting 2-(iodoethynyl) derivatives of thioanisole **3a**, *N,N*-dimethylaniline **3b** and anisole **3c** were obtained from the corresponding TMS-alkynes **1a–c** (see the Supporting Information) by a stepwise desilylation and iodination of terminal alkynes **2a–c** (Scheme 1).

Iodoalkynes **3a–c** turned out to be quite sensitive substances. They require mild reaction conditions and should be handled carefully during work up and isolation in order to avoid protodeiodination (see the SI). Thus, the iodoalkynes should not be stored in solution for prolonged time and the water bath temperature for a rotary evaporator should not exceed 35 °C. With these rules, the key iodoalkynes **3a–c** could be obtained in high yields (Scheme 1) and stored neat in a freezer for several months without any degradation.

Then, we turned to the study of iodocyclization of iodoalkynes **3a–c** using the conditions which gave the highest yields in the case of alkyl- /aryl-substituted *ortho*-functionalized alkynes.^{54–56} (Iodoethynyl)thioanisole **3a** underwent the iodocyclization smoothly with iodine as an electrophile: 2,3-diiodobenzothiophene **4** was isolated in almost quantitative yield (Table 1, entry 1). The cyclization of aniline **3b** using iodine gave a complex mixture of products. Switching to *N*-iodosuccinimide, gave indole **5** in 86% yield (Table 1, entry 2). The reaction of anisole **3c** with either iodine (under a variety of conditions) or ICl ended with electrophilic addition to the triple bond to form the dihalogenated alkenes **6** and **7**, respectively (Table 1, entries 3,4). When NIS was used, no conversion of the starting (iodoethynyl)anisole was observed even after 6 hours (Table 1, entry 5). However, we could overcome this obstacle by using 2-ethynylated phenols instead of anisoles and by switching to NIS/PPh₃ as the electrophilic system as described previously by Li et al.⁵⁷


Interestingly, when optimizing synthesis of (iodoethynyl)phenol **3d**, we have inadvertently found that desilylation/iodination of TMS-substituted ethynylphenol **1d** under AgF/NIS conditions⁵⁸



Conditions: **A**: K₂CO₃ (2 equiv), MeOH or EtOH, rt, 3–18 h; **B**: N-iodomorhpoline hydroiodide (1.25–3 equiv), CuI (15 mol%), THF, 5 °C, 2–3 h (for **2a,b**); **C**: DBU (1.1 equiv), NIS (1.1 equiv) MeCN, rt, 15 min (for **2c**)

Scheme 1 Synthesis of *o*-functionalized (iodoethynyl)arenes **3a–c**.

Table 1 The study of iodine-promoted cyclization of iodoalkynes **3a–c**.



Entry	Iodoalkyne	X	R	Conditions	Yield, %	Product
1	3a	S	H	A	97	4
2	3b	NMe	COOEt	B	86	5
3	3c	O	H	C	76	6 (Y = I)
				D	99	
				E	77	
4	3c	O	H	F	82	7 (Y = Cl)
5	3c	O	H	B	–	–


Conditions: A – I₂ (1.05 equiv), DCM, rt; B – NIS (2 eq.), DCM, rt; C – I₂ (1.05 equiv), DCM, rt; D – I₂ (3.00 equiv), DCM, 40 °C; E – I₂ (3.00 equiv), DCE, 80 °C; F – ICl (1.05 equiv), DCE, 80 °C.

gave the desired 2,3-diiodobenzofuran **8**, albeit in a low yield (13%) and along with a mixture of unidentified products. Iodocyclization of (iodoethynyl)phenol **3d** under NIS/PPh₃ conditions gave 2,3-diiodobenzofuran **8** in 37% isolated yield (Scheme 2).

Thus, the preinstallation of iodine at the triple bond of *ortho*-functionalized ethynylarenes enables direct synthesis of 2,3-dioiodobenzothiophene and 2,3-diiodoindole. Although iodination of 2-(iodoethynyl)anisole leads instead to the 1,2-diiodination of the triple bond, the target 2,3-diiodobenzofuran can be synthesized using 2-(iodoethynyl)phenol as a starting material and NIS/PPh₃ system as the electrophilic reagent.

Regioselectivity of the Sonogashira Coupling in 2,3-Diiodoheteroindenes

Next, we explored the regioselectivity of Sonogashira coupling^{59,60} for the diiodides **4**, **5**, **8**. The idea of selective substitution of one iodine atom in vicinal diiodoheteroindenes is based on the known regioselective Sonogashira reaction for 2,3-diiodothiophenes⁴⁴ and 2,3-dibromobenzothiophene.⁴⁶ The substitution is known to occur first at the C2, and then at the C3 position.



Scheme 2 Synthesis of 2,3-diiodobenzofuran **8**.

From the theoretical point of view, many factors influence the regioselectivity of oxidative addition step, which is known to be the limiting stage of the many Pd cross-coupling reactions.³⁹ An interesting selectivity feature is that the reaction does not prefer the *weaker* of the two C-X bonds as indicated by the data shown in Figure 4 for C-Br and C-I bonds in the two dihalides. Interestingly, the shorter and stronger C2-Hal bonds react first. Such observations are consistent with the earlier suggestion that, the rate-limiting step for the Sonogashira cross-coupling of aryl iodides is likely to be not the C-X bond activation via oxidative addition to the catalyst but trans-metallation of Cu-acetylides formed in the initial reaction stage.⁶¹

Therefore, we were aimed to compare experimentally the regioselectivity of the cross-coupling for all three diiodoheteroindenes **4**, **5**, **8** with the goal of evaluating how general the above trend is. From a practical point of view, this information is useful for deciding which heterocycles are appropriate for the synthesis of unsymmetrical enediyne systems under one-pot conditions.

To study the regioselectivity in the Sonogashira reaction of diiodobenzothiophene **4**, we chose 3-methoxypropyne **9a** as a partner because its regioselective cross-coupling would lead to the corresponding C2-(methoxymethyl)ethynyl derivative, which is an important starting material for the synthesis of cyclic alkynes through the Ni-catalyzed reaction (Table 2).¹⁹

All reactions were carried out using $\text{Pd}(\text{PPh}_3)_4/\text{CuI}$ catalytic system, while the nature of the base, solvent, temperature and

reagents ratio were varied. The yields of all products and the conversion of diiodide **4** were estimated using ^1H NMR spectroscopy of the reaction mixture with 1,3,5-trimethoxybenzene as the internal standard. The Sonogashira coupling of diiodide **4** with alkyne **9a** using potassium carbonate as a base at 45 °C gave C2-monosubstituted product **10a** in 76% yield. However, a non-desired disubstituted product **11a** was also formed in 11% yield (Table 3, Entry 1). Raising the temperature from 45 °C to 60 °C decreased the yield of monoethynylated benzothiophene **10a** (Table 3, Entry 2). Among the various bases tested (Table 3, Entries 3–5), the best result was achieved with K_3PO_4 (80% yield of **10a**, entry 3), albeit with the concomitant formation of 9% of enediyne **11a**. In order to diminish the formation of undesired enediyne **11a**, we reduced the acetylene/diiodide ratio from 1.2 to 1.0 equiv. (Table 3, Entry 6) but found these conditions led to a decrease in the yield of product **10a**. Thus, the conditions from entry 3 were chosen for the regioselective synthesis of 2-ethynyl-3-iodobenzothiophene **10a**: the reaction should be carried out in THF at 45 °C using K_3PO_4 as a base and maintaining the starting diiodide **4** to alkyne **9a** ratio of 1:1.25.

Importantly, the product of monosubstitution at C3 has never been observed. Thus, the ethynyl substituent in the C2-position of benzothiophene has an activating effect on the rate of cross-coupling at the C3 position. The activation can be associated either with a decrease in electron density at the C3 position for 2-ethynyl-3-iodobenzothiophenes compared to 2,3-diodobenzothiophene because of the electron withdrawing properties of Csp atoms, or/and with other factors such as coordination of palladium at the triple bond at the oxidative addition step.

Studies on regioselectivity of indole 5 and benzofuran 8 derivatives were carried out under optimized conditions for diiodobenzothiophene 4. We found that 2,3-diiodoindole 5 cannot be converted to the corresponding C2-ethynylindoles with the same selectivity as it has been observed for 2,3-diiodobenzothiophene 4 (Scheme 3). Although the desired C2-ethynylated indole 13a was the main reaction product (52 % yield) of the Sonogashira reaction with alkyne 9a (Scheme 11), the enediyne 14a (18%) along with the remaining starting diiodide 5 (conversion 87%) was also isolated. Raising the temperature by 10 °C led to the increased formation of symmetrically substituted enediyne 14a, with starting diiodide 5 still present in the reaction mixture.

The Sonogashira coupling of benzofuran diiodide **8** with methyl propargyl ether **9a** proceeded mostly through the C2-position with similar regioselectivity as for diiodindole **5**.






Fig. 4 Energetic and electronic parameters for the two C-Hal bonds in 2,3-dibromo- and 2,3-diiodobenzo[*b*]thiophenes

Table 2 Conditions optimization for the Sonogashira reaction between diiodobenzothiophene **4** and alkyne **9a**.^a



Entry	Base	T, °C	Conversion of 4 % ^b	Yield, %		
				10a	11a	12
1	K ₂ CO ₃	45	94	76	11	<1
2	K ₂ CO ₃	60	70	64	8	–
3	K ₃ PO ₄	45	>95	80	9	1.5
4	KF	45	39	45	2	4
5 ^c	KF	45	55	55	2	4
6 ^d	K ₃ PO ₄	45	73	66	7	1

^aReaction conditions: **4** (0.25 mmol, 1 equiv), **9a** (0.30 mmol, 1.2 equiv), Pd(PPh₃)₄ (5 mol.%), CuI (10 mol.%), base (1.00 mmol, 4.00 equiv), THF (2.50 mL, *c* = 0.1 M), 24 h; ^bConversion and yield were determined by ¹H NMR spectroscopy using 1,3,5-trimethoxybenzene as an internal standard; ^c DMF was used as a solvent; ^dthe alkyne **9a** was used in amount of 0.25 mmol, 1.00 equiv.



Scheme 3 Study of the Sonogashira reaction regioselectivity for 2,3-diodoindole 5 and 2,3-diodobenzofuran 8.

The Sonogashira coupling of benzofuran diiodide **8** with methyl propargyl ether **9a** proceeded mostly through the C2-position with similar regioselectivity as for diiodindole **5** to be 3 / 1, with the isolated yields of 59% for 2-ethynylbenzofuran **15a** and 27% for enediyne **16**.


Moderate yields and less selectivity for monoethynylation in the case of indole **5** and benzofuran **8** compared to benzothiophene **4** against the formation of diethynylated byproduct can be explained by higher sensitivity of the C3-I position towards additional activation by the initially introduced C2-alkyne moiety for these heterocycles.

Finally, we decided to check the possibility to reverse the regioselectivity for the benzothiophene series from C2-I to C3-I by replacing active halogen (I) with less reactive halogen (Br) in more activated position (C2). Thus corresponding C3-I regioselective Sonogashira reaction has been reported for 2-bromo-3-iodothiophenes (Fig. 3).⁴⁸⁻⁵¹ 2-Bromo-3-iodobenz[*b*]thiophene **17** was obtained similarly to its diiodide analogue **4** starting from bromoethynylthioanisole S1 (See SI). The Sonogashira reaction of 2-bromo-3-iodobenz[*b*]thiophene **17** with 3-methoxypropyne **9a** under optimized condition for the diiodobenzothiophene **4** (Table 3, Entry 3) proceeded at C3-I bond instead of C2-Br bond. However, the selectivity for the mono- vs. bis-substitution was significantly lower than C2-selectivity observed for substitution to diiodide **4**. Along with the iodine substitution at the C3 position (product **18a**), a significant amount of the disubstituted benzothiophene **11a** was formed: the ratio of alkyne **18a** to enediyne **11a** was found to be 2:1 (Scheme 4). It should be noted, that C2-Br substitution in the starting material was not observed, so the difference in intrinsic reactivity of C-Br and C-I bonds was more important than the inherent C2-preference. It also indicates once again, that, even for the C2-Br benzothiophene derivative, a triple bond has an activation effect at the adjacent position.


Additional Selectivity Studies of Dihalobenzothiophenes

To explore the activating influence of the firstly introduced triple bond on the remaining C-Hal in the Sonogashira coupling, we carried out additional experiments. In particular, we studied interactions of the two dihaloheteroindenes with the worst mono- vs. bis-ethynylation selectivity (i.e., **8**, **17**) with hept-1-yne **9b**, an alkyne without any additional functionalities (Scheme 5).

In the case of di-I-benzofuran **8**, the mono- to bis-ethynylation ratio increased from 3/1 (for methyl propargyl ether **9a**) to 5/1 (for heptyne **9b**).



Scheme 4 Study of the Sonogashira reaction regioselectivity for 2-bromo-3-iodobenz[*b*]thiophenes **17**.




Scheme 5 Study of the influence of alkyne nature on mono- vs. bis-ethynylation selectivity.

The analogous trend was observed for C2-bromo benzothiophene **17**: mono- **18b** to bis-ethynylation **11b** ratio also increased from 2/1 (for methyl propargyl ether **9a**) to 3/1 (for heptyne **9b**) (Scheme 5). The results are intriguing because the first alkyne is introduced at the different carbons (C2 for **8**, and C3 for **17**) in these Sonogashira reactions.

Exploratory Computational Analysis

Several interesting observations can be made from the experimental results discussed in the previous sections. First, introduction of the first alkyne group accelerates the 2nd ethynylation step. Second, the activation effect of the firstly incorporated triple bond is dependent on the alkyne substituent, i.e. the acceleration is less pronounced for heptyne **9b** in comparison to methyl propargyl ether **9a**.

In order to understand the electronic consequences of alkyne introduction, we have used DFT calculations to evaluate charge distribution in 2,3-diodobenzothiophene **4** and the two products of Sonogashira mono-substitution at C2 – 2-ethynyl-3-iodobenzothiophens **10a** and **10b'** (Scheme 6). At this stage, we have limited our computational analysis only to the benzothiophene systems where the experimental data are the most complete. Note that the C2-carbon in the starting diiodide **4** has much more electron density than C3 (natural charges of -0.48 and -0.30 e). Upon the introduction of a C2-alkyne substituents, the electron density at both C2 and C3 is significantly lowered. This interesting finding illustrates the uniqueness of sp-hybridized carbon and consistent with the high acceptor ability of alkynes.^{62,63}



Scheme 6 The activating effect of alkyne moiety and $\pi_{C=C} \rightarrow \sigma_{C=C}$ hyperconjugation on the C3 carbon in benzothiophenes.


The introduction of an oxygen atom at the propargylic position further decreases electron density at the C3 atom. This effect stems from a combination of the inductive effect and $\pi_{C=C} \rightarrow \sigma^*_{CO}$ hyperconjugation, which both accentuate the π accepting ability of the sp carbon. If this remote stereoelectronic effect can facilitate a reaction at the remaining C-Hal bond, it can explain the lower mono- vs. bis-ethynylation selectivity for the propargylic ether.

As transmetalation is known to be the rate-limiting step in Sonogashira reactions of aryl iodides,^{64,65} we evaluated substituent effects on this step with the hope that they will help to explain the observed reactivity trends. Scheme 7 illustrates differences in the calculated exergonicities for transmetalation. In this context, it is important to remember that, as long as the same elementary processes are compared, exergonicities are known to correlate with activation barriers. Such connection between reaction thermodynamics and kinetics readily follows from the Marcus theory or Bell-Evans-Polany postulate.^{66,67} For more recent examples of such connections between kinetics and thermodynamics, see the references.^{68,69}



The results of these calculations provided three conclusions, all in a good agreement with the experimental results. In particular, comparison of Eqs. 1 and 2 explains the preference for reaction at C2 as this process is more favorable. The more exergonic transmetalation at C2 (-19.4 kcal/mol) is expected to be faster than the less exergonic reaction at C3 (-17.8 kcal/mol). On the other hand, comparison of Eqs. 2 and 3 rationalizes the accelerating effect of alkyne moiety which complicates reaction designs that rely on the selective formation of monoalkynylated Sonogashira products from the diiodides. Finally, comparison of Eqs. 3 and 4 shows how the propargylic acceptor group at C2 further promotes transmetalation at C3, rendering selective monoalkynylation even more difficult.

Preparation of Non-Symmetric Enediynes

Next, we applied the optimized conditions for the Sonogashira cross-coupling of diiodide **4** with different alkynes **9a-f** (Scheme 8). In all cases, the reaction proceeded with the full conversion of diiodide **4** and gave 2-ethynyl-3-iodobenzothiophenes **10a-f** in good isolated yields.



Scheme 7 Thermodynamics of transmetalations in relevant benzothiophenes.




Scheme 8 Synthesis of 2-ethynylbenzo[b]thiophenes **10a-f**.

The enediyne by-products were formed in insignificant amounts, therefore they were not isolated chromatographically. Consequently, 2,3-diiodobenzothiophene shows minimal dependence of regioselectivity on the nature of the alkyne.

In order to demonstrate the utility of the regioselective Sonogashira reactions, we have carried out a two-step synthesis of unsymmetrically substituted enediyne systems from 2,3-diiodobenzothiophene **4** via two successive cross-coupling reactions with different alkynes under one-pot conditions, i.e., without isolating the product of the first Sonogashira reaction on the C2 atom. It turned out that this approach allowed quick and efficient preparation of the unsymmetrically substituted enediynes **19a-e**. In this sequence, the first substitution at C2 was carried out under the optimized conditions (Table 1, Entry 3), while the subsequent C3-substitution of iodine was accomplished by adding the second alkyne to the reaction mixture and running the reaction at 60 °C until the complete conversion. In this way, it was possible to obtain a series of unsymmetrically substituted enediynes, including isomeric enediynes **19a / 19b** and **19c / 19d** (Scheme 9). Because the resulting enediynes have substituents with different polarity at the triple bonds, chromatographic separation of the target unsymmetrical enediynes **19** from symmetrically substituted by-products **11a,b** is straightforward. The isolation of the trace amounts of symmetrical enediyne minor by-products in the syntheses of compounds **19c-e** was not carried out.

Thus, unsymmetrically substituted enediynes fused to a benzothiophene **19** can be synthesized from the easily synthetically accessible 2,3-diiodobenzothiophene **4** within a

ARTICLE




Scheme 9 Synthesis of unsymmetrical enediynes 19 using a one-pot approach

single synthetic stage using a one-pot procedure involving two subsequent Sonogashira reactions. The advantage of the developed synthetic route compared to the previously described approach based on the iodocyclization of *o*-(buta-1,3-diylyn)thioanisole followed by the Pd-catalyzed cross-coupling with alkynes is its higher divergence without the need to start from different diacetylenes in the first synthetic steps.

We have compared the overall yield of enediyne 19a synthesized by two different routes – through the «diacetylenic» approach^{19,24} and via the «iodoalkyne»

approach (Scheme 10). The new method gives a somewhat higher overall yield; however, it still requires separating traces of symmetrical endiynes in the last step. Therefore, we would recommend using the developed «iodoalkyne» approach for the synthesis of libraries of unsymmetrical benzothiophene-fused enediynes, while the «diacetylenic» route seems to be more suitable for the gram-scale synthesis of enediynes with an identically substituted triple bond at the C2 position.²²



Scheme 10 Comparison the efficiency of the «diacetylenic» and the «iodoalkyne» routes towards acyclic enediynes fused to a benzothiophene. The two routes are defined in Scheme 1.


ARTICLE

Preparation of Symmetric Enediynes

The symmetrically substituted enediynes can be synthesized directly from diiodide **4,5,8** carrying out the Sonogashira coupling at 60 °C. Following this way, several enediynes fused to benzothiophene **11a–d**, indole **14a**, and benzofuran **16a** were synthesized mostly in good yields (Scheme 11).

Biological Activity of Acyclic Enediynes

It has recently been shown that several acyclic $C_6F_5S(O)$ -substituted enediynes display cytotoxic activity unrelated to the Bergman cyclization. It is assumed that cytotoxicity is associated



Scheme 11 Synthesis of symmetrically substituted enediynes fused to a benzothiophene **11a–d**, indole **14a** and benzofuran **16a**

with disruption of the microtubule network.⁷⁰ Although the presence of pentafluoroaryl moiety was thought to be responsible for the binding of enediynes to tubulin through nucleophilic aromatic substitution, an enediyne molecule lacking this group was also active against human pancreatic tumor cells MIAPaCa-2 ($IC_{50} = 15.9\text{ }\mu M$). Furthermore, we have independently demonstrated that dialkylaminomethyl-substituted benzothiophene-fused acyclic enediynes show cytotoxic activity against NCI-H460 lung carcinoma.⁷¹

Therefore, we decided to test cytotoxicity for the series of acyclic enediynes fused to benzothiophene and indole. The list of compounds included enediynes **11a–d**, **19a–e** obtained in this work, as well as enediynes **19f–h**²², **19i,j** (see the Supporting Information), and **20**²⁴ synthesized in our group earlier using the «diacetylenic» route. Cytotoxicity of these compounds was tested against NCI-H460 lung carcinoma and WI-26 VA4 lung epithelial-like cell lines using the MTT colorimetric test^{72,73}. All the enediynes were studied at a concentration of $75\text{ }\mu M$. Most of unsymmetrically substituted enediynes fused to a benzothiophene **19** and an indole derivative **20** showed cytotoxic activity against NCI-H460 lung carcinoma cells. Furthermore, enediynes **19b,c,g, h**, **20** were more cytotoxic towards the NCI-H460 cells than to WI-26 VA4 lung epithelial-like fibroblasts (Figure 5).

Earlier, the Hu group showed that acyclic enediynes having a double bond as a part of a maleimide ring and with a σ -acceptor atom at the propargylic position are able to undergo MARACA transformation (a cascade acetylene-allene isomerization) under polar conditions. This reaction cascade leads to enyne allenes, prone to the formation of reactive σ,π -diradicals through the Mayers-Saito cyclization that induce DNA breaks.^{74,75} Taking into account, that most of the enediynes tested here have a propargylic σ -acceptor, we decided to check the ability of the acyclic enediynes to induce DNA strand breaks. Enediynes **19a–c,f,i,j, 20** were chosen and investigated in the assay with pBR322 DNA plasmid under the same conditions as for cyclic analogs of benzothiophene-fused enediynes.²⁰ The obtained data revealed that only enediyne **19i** possessed a weak DNA damaging activity: a trace amount of open circular DNA was detected in the electrophorogram (Figure 6). Other enediynes did not display DNA damaging activity. This finding confirms that cytotoxicity of acyclic enediynes studied here is associated neither with the Bergman cyclization nor with the Mayers-Saito cyclization. It may be worth noting that enediynes can also form polar intermediates that can also damage biomolecules.⁷⁶ Therefore, revealing the mechanism of cytotoxicity of enediynes requires further studies.

ARTICLE




Fig. 5 Screening for the cytotoxicity of the enediynes fused to benzothiophene 11, 19 and indole 20.




Fig. 6 Electrophorogram of pBR322 plasmid after exposure to 250 μ M concentrations of enediynes 19a-c,f,i,j, 20. CC – covalently closed circular plasmid; OC – open circular.

Conclusions

In summary, the iodocyclization of *ortho*-functionalized (iodoethynyl)arenes along with the regioselective Sonogashira coupling opens a convenient synthetic route towards heteroindene-fused unsymmetrical enediynes. 2,3-Diiodobenzothiophene and 2,3-diiodoindole were easily accessible via the iodocyclization. Although

(iodoethynyl)anisole underwent iodination of the triple bond instead of cyclization, 2,3-diiodobenzofuran could be prepared, instead, from (iodoethynyl)phenol and NIS/PPh₃.

The regioselective outcome of the Sonogashira reaction for 2,3-diiodoheteroindenes depends on the nature of the heterocycle, as well as on the nature of alkynes. The Sonogashira coupling for all derivatives follows the C2-regioselectivity rule, without producing a C3 mono-substitution byproduct. However, the regioselectivity suffers from the activating influence of the initially introduced alkyne on the adjacent C-Hal position resulting in the formation of disubstituted byproduct. This effect is stronger for diiodoindole and diiodobenzofuran than for diiodobenzothiophene. The ratio of mono- and disubstituted products also depends on the nature of the alkyne. Alkynes with propargylic acceptors at C2 have a greater activating effect at the C3 position towards the subsequent Sonogashira reaction.

As the result, diethynylation does not preclude synthesis of unsymmetrical benzothiophene-fused enediynes based on a sequential one-pot C2/C3 ethynylation, but limits the use of the one-pot approach for the indole and benzofuran derivatives. In

this case, either the “diacetylene” route or the stepwise two-pot substitution of both iodine atoms by the Sonogashira cross-coupling are recommended.

In addition, these experimental selectivity trends provided a tool to probe mechanistic features of Sonogashira cross-coupling. A diverse set of DFT data, including C-X bond strength, C-X bond polarizations, electron density distributions, and reaction exergonicities converge on transmetalation, rather than oxidative addition, as the be rate-limiting step of this reaction cascade. Computational analysis using various levels of DFT revealed a few significant points about the selectivity and provided insight into the mechanistic basis of the observed selectivities in the Sonogashira coupling.

Importantly, the alkyne moiety is an activating substituent (more than the heavier halogens). Furthermore, a remote (propargylic) acceptor can make this activating effect more pronounced. This observation illustrates once again unique properties of the alkyne functionality – it is both a significantly electronegative substituent and a perfect conduit for stereoelectronic hyperconjugative effects.⁷⁷

The acyclic unsymmetrical enediynes showed cytotoxic activity against NCI-H460 lung carcinoma cell lines along with the lower cytotoxicity toward WI-26 VA4 lung epithelial-like cell lines. Interestingly, the mechanism of the cytotoxicity is not associated with the induction of DNA strands breaks.

Author Contributions

AVP – Investigation, Writing – original draft; NAD - Data curation, Methodology, Project administration, Supervision, Visualization, Writing – original draft, Writing – review & editing; JSO – Investigation; AAV – Investigation, Validation; EAK – Investigation, Resources; ASB – Investigation, Visualization; AMR - Investigation, Visualization; AIG - Investigation, Funding acquisition; TS – Investigation, Writing – original draft; IVA – Data curation, Methodology, Supervision, Visualization, Writing – original draft, Writing – review & editing; IAB – Conceptualization, Methodology, Funding acquisition, Project administration, Supervision, Visualization, Writing – original draft, Writing – review & editing

Conflicts of interest

There are no conflicts to declare

Acknowledgements

This work represents an integration of two diverse projects funded by the Russian Science Foundation: IAB grant 21-13-00218 diiodides synthesis and reactivity; AIG grant 19-73-10077-Π biological applications. The study was carried out using equipment from the SPbU Research Park: the Centre for Magnetic Resonance and the Centre for Chemical Analysis and Materials Research. The authors would like to thank Alexander Ivanov (SPbU) and Sergey Smirnov (SPbU) for measuring the NMR and Oussama Abdelhamid Mammeri (SPbU) for measuring

HR MS. Authors gratefully acknowledge the Spectral-Analytical Center of FRC Kazan Scientific Center of RAS for providing necessary facilities to carry out GC-MS and HRMS-ESI measurements. Computational studies at FSU were supported by the National Science Foundation (NSF) (CHE-2102579).

Notes and references

- 1 T. Aggarwal, S. Kumar and A. K. Verma, *Org. Biomol. Chem.*, 2016, **14**, 7639–7653.
- 2 B. Godoi, R. F. Schumacher and G. Zeni, *Chem. Rev.*, 2011, **111**, 2937–2980.
- 3 Elsevier, *Compr. Org. Synth. II*, 2014, **4**, 412–494.
- 4 S. Singh and S. Chimni, *Synthesis (Stuttg.)*, 2015, **47**, 1961–1989.
- 5 K. Gilmore, R. K. Mohamed and I. V. Alabugin, *Wiley Interdiscip. Rev. Comput. Mol. Sci.*, 2016, **6**, 487–514.
- 6 T. Khan and S. Yaragorla, *Eur. J. Org. Chem.*, 2019, **2019**, 3989–4012.
- 7 J. Tummatorn, S. Ruchirawat and C. Thongsornkleeb, *Synthesis (Stuttg.)*, 2023, **55**, 3073–3089.
- 8 G. Leonel, I. Klann, D. F. Back, B. A. Iglesias, C. W. Nogueira and G. Zeni, *Chem. Eur. J.*, 2023, **29**, e202202847.
- 9 L. Habert, I. Diachenko, P. Retailleau and I. Gillaizeau, *Org. Biomol. Chem.*, 2021, **19**, 6623–6627.
- 10 T. Okitsu, M. Itoh, A. Inui, E. Muta, R. Nakamoto, Y. Adachi, A. Wada and M. Hatano, *Asian J. Org. Chem.*, 2022, **11**, e202200275.
- 11 T. Okitsu, A. Namura, S. Kondo, S. Tada, M. Yanagida and A. Wada, *Org. Chem. Front.*, 2020, **7**, 879–884.
- 12 D. Bag, H. Kour, N. Saha, Kamal, H. Holla, P. V. Bharatam and S. D. Sawant, *J. Org. Chem.*, 2023, **88**, 2377–2384.
- 13 I. Misiūnaitė, R. Bukšnaitienė, J. Pošiūnas, A. Brukštus and I. Žutautė, *J. Heterocycl. Chem.*, 2022, **59**, 1712–1722.
- 14 R. K. Mohamed, S. Mondal, J. V. Guerrera, T. M. Eaton, T. E. Albrecht-Schmitt, M. Shatruk and I. V. Alabugin, *Angew. Chem. Int. Ed.*, 2016, **55**, 12054–12058.
- 15 H.-C. Huang, Y.-C. Hsieh, P.-L. Lee, C.-C. Lin, Y.-S. Ho, W.-K. Shao, C.-T. Hsieh, M.-J. Cheng and Y.-T. Wu, *J. Am. Chem. Soc.*, 2023, **145**, 10304–10313.
- 16 T. Okitsu, A. Horike, N. Shimazawa and A. Wada, *Org. Biomol. Chem.*, 2020, **18**, 3501–3511.
- 17 S. Chen, D. L. Priebbenow, J. Somkhit, C. V. Scullino, K. Agama, Y. Pommier and B. L. Flynn, *Chem. Eur. J.*, 2022, **28**, e202201925.
- 18 N. A. Danilkina, A. E. Kulyashova, A. F. Khlebnikov, S. Bräse and I. A. Balova, *J. Org. Chem.*, 2014, **79**, 9018–9045.
- 19 A. G. Lyapunova, N. A. Danilkina, A. F. Khlebnikov, B. Köberle, S. Bräse and I. A. Balova, *Eur. J. Org. Chem.*, 2016, **2016**, 4842–4851.
- 20 N. Danilkina, A. Rumyantsev, A. Lyapunova, A. D'yachenko, A. Khlebnikov and I. Balova, *Synlett*, 2019, **30**, 161–166.
- 21 A. G. Lyapunova, N. A. Danilkina, A. M. Rumyantsev, A. F. Khlebnikov, M. V. Chislov, G. L. Starova, E. V. Sambuk, A. I. Govdi, S. Bräse and I. A. Balova, *J. Org. Chem.*, 2018, **83**, 2788–2801.
- 22 N. A. Danilkina, E. A. Khmelevskaya, A. G. Lyapunova, A. S. D'yachenko, A. S. Bunev, R. E. Gasanov, M. A. Gureev and I. A. Balova, *Molecules*, 2022, **27**, 6071.
- 23 N. A. Danilkina, L. Y. Gurskaya, A. V. Vasilyev and I. A. Balova, *Eur. J. Org. Chem.*, 2016, **2016**, 739–747.
- 24 N. A. Danilkina, A. S. D'yachenko, A. I. Govdi, A. F. Khlebnikov, I. V. Kornyakov, S. Bräse and I. A. Balova, *J. Org. Chem.*, 2020, **85**, 9001–9014.
- 25 N. Choy, B. Blanco, J. Wen, A. Krishan and K. C. Russell, *Org. Lett.*, 2000, **2**, 3761–3764.

26 S. Hamri, J. Rodríguez, J. Basset, G. Guillaumet and M. Dolors Pujol, *Tetrahedron*, 2012, **68**, 6269–6275.

27 J. Kour, P. Khajuria, A. Sharma and S. D. Sawant, *Chem. Asian J.*, 2022, **17**, e202200778.

28 J. Zhang, S. Li, G.-J. Deng and H. Gong, *ChemCatChem*, 2018, **10**, 376–380.

29 J. Bergman and L. Venemalm, *J. Org. Chem.*, 1992, **57**, 2495–2497.

30 Y. Liu and G. W. Gribble, *Tetrahedron Lett.*, 2000, **41**, 8717–8721.

31 A. Putey, F. Popowycz and B. Joseph, *Synlett*, 2007, **2007**, 419–422.

32 H. Hamamoto, Y. Miki, H. Umemoto, M. Umemoto, C. Ohta, E. Fujita, A. Nakamura and T. Maegawa, *Heterocycles*, 2015, **91**, 561.

33 Y. Yang, L. Zhang, G.-J. Deng and H. Gong, *Sci. Rep.*, 2017, **7**, 40430.

34 G. J. P. Perry, J. M. Quibell, A. Panigrahi and I. Larrosa, *J. Am. Chem. Soc.*, 2017, **139**, 11527–11536.

35 S. Mehta and R. C. Larock, *J. Org. Chem.*, 2010, **75**, 1652–1658.

36 T. R. Puleo and J. S. Bandar, in *Encyclopedia of Reagents for Organic Synthesis*, Wiley, 2022, pp. 1–2.

37 T. Kesharwani, C. Kornman, A. Tonnaer, A. Hayes, S. Kim, N. Dahal, R. Romero and A. Royappa, *Tetrahedron*, 2018, **74**, 2973–2984.

38 A. D. Sonawane, D. R. Garud, T. Udagawa and M. Koketsu, *Org. Biomol. Chem.*, 2018, **16**, 245–255.

39 J. Lu, S. Donnecke, I. Paci and D. C. Leitch, *Chem. Sci.*, 2022, **13**, 3477–3488.

40 M. F. Ibad, D. S. Zinad, M. Hussain, A. Ali, A. Villinger and P. Langer, *Tetrahedron*, 2013, **69**, 7492–7504.

41 S. Tšupova, M. M. Hansmann, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Chem. - A Eur. J.*, 2016, **22**, 16286–16291.

42 M. M. Hansmann, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Angew. Chem. Int. Ed.*, 2013, **52**, 2593–2598.

43 S. Tšupova, M. M. Hansmann, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Chem. Eur. J.*, 2016, **22**, 16286–16291.

44 N. B. Vinh, S. M. Devine, L. Munoz, R. M. Ryan, B. H. Wang, H. Krum, D. K. Chalmers, J. S. Simpson and P. J. Scammells, *ChemistryOpen*, 2015, **4**, 56–64.

45 Z. Szlavik, M. Csekei, A. Paczal, Z. B. Szabo, S. Sipos, G. Radics, A. Proszenyak, B. Balint, J. Murray, J. Davidson, I. Chen, P. Dokurno, A. E. Surgenor, Z. M. Daniels, R. E. Hubbard, G. Le Toumelin-Braizat, A. Claperon, G. Lysiak-Auvity, A.-M. Girard, A. Bruno, M. Chanrion, F. Colland, A.-L. Maragno, D. Demarles, O. Geneste and A. Kotschy, *J. Med. Chem.*, 2020, **63**, 13762–13795.

46 G. A. Salman, N. N. Pham, T. N. Ngo, P. Ehlers, A. Villinger and P. Langer, *Adv. Synth. Catal.*, 2017, **359**, 1402–1406.

47 X. Yin, Y. Li, Y. Zhu, Y. Kan, Y. Li and D. Zhu, *Org. Lett.*, 2011, **13**, 1520–1523.

48 Z. U. Levi and T. D. Tilley, *J. Am. Chem. Soc.*, 2009, **131**, 2796–2797.

49 Y. Morisaki, M. Tominaga and Y. Chujo, *Chem. Eur. J.*, 2012, **18**, 11251–11257.

50 A. Fukazawa, D. Kishi, Y. Tanaka, S. Seki and S. Yamaguchi, *Angew. Chem. Int. Ed.*, 2013, **52**, 12091–12095.

51 F. Guo, X. Yin, F. Pammer, F. Cheng, D. Fernandez, R. A. Lalancette and F. Jäkle, *Macromolecules*, 2014, **47**, 7831–7841.

52 DE102013214399, 2014.

53 C. Shu, C.-Y. Shi, Q. Sun, B. Zhou, T.-Y. Li, Q. He, X. Lu, R.-S. Liu and L.-W. Ye, *E*, 2019, **9**, 1019–1025.

54 D. Yue and R. C. Larock, *J. Org. Chem.*, 2002, **67**, 1905–1909.

55 D. Yue and R. C. Larock, *Org. Lett.*, 2004, **6**, 1037–1040.

56 D. Yue, T. Yao and R. C. Larock, *J. Org. Chem.*, 2005, **70**, 10292–10296.

57 Y.-L. Li, J. Li, S.-N. Yu, J.-B. Wang, Y.-M. Yu and J. Deng, *Tetrahedron*, 2015, **71**, 8271–8277.

58 E. J. M. Maduli, S. J. Edeson, S. Swanson, P. A. Procopiou and J. P. A. Harrity, *Org. Lett.*, 2015, **17**, 390–392.

59 K. Sonogashira, Y. Tohda and N. Hagihara, *Tetrahedron Lett.*, 1975, **16**, 4467–4470.

60 I. Kanwal, A. Mujahid, N. Rasool, K. Rizwan, A. Malik, G. Ahmad, S. A. A. Shah, U. Rashid and N. M. Nasir, *Catalysts*, 2020, **10**, 443.

61 R. Chinchilla and C. Nájera, *Chem. Rev.*, 2007, **107**, 874–922.

62 C. Hu, J. Mena and I. V. Alabugin, *Nat. Rev. Chem.*, 2023, **7**, 405–423.

63 I. Alabugin, E. Gonzalez-Rodriguez, R. Kawade, A. Stepanov and S. Vasilevsky, *Molecules*, 2019, **24**, 1036.

64 X. Wang, Y. Song, J. Qu and Y. Luo, *Organometallics*, 2017, **36**, 1042–1048.

65 C. He, J. Ke, H. Xu and A. Lei, *Angew. Chemie*, 2013, **125**, 1567–1570.

66 R. P. Bell, *Proc. R. Soc. Lond. A. Math. Phys. Sci.*, 1936, **154**, 414–429.

67 M. G. Evans and M. Polanyi, *Trans. Faraday Soc.*, 1938, **34**, 11–24.

68 I. V. Alabugin and M. Manoharan, *J. Am. Chem. Soc.*, 2005, **127**, 12583–12594.

69 G. dos P. Gomes and I. Alabugin, in *Applied Theoretical Organic Chemistry*, WORLD SCIENTIFIC (EUROPE), 2018, pp. 451–502.

70 C. Borie, S. Mondal, T. Arif, M. Briand, H. Lingua, F. Dumur, D. Gigmes, P. Stocker, B. Barbarat, V. Robert, C. Nicoletti, D. Olive, M. Maresca and M. Néchab, *Eur. J. Med. Chem.*, 2018, **148**, 306–313.

71 A. I. Govdi, S. O. Anisimov, N. A. Danilkina, A. S. Bunev and I. A. Balova, *Mendeleev Commun.*, 2023, **33**, 328–330.

72 T. Mosmann, *J. Immunol. Methods*, 1983, **65**, 55–63.

73 A. Kazantsev, O. Bakulina, D. Dar'in, G. Kantin, A. Bunev and M. Krasavin, *Org. Lett.*, 2022, **24**, 4762–4765.

74 M. Zhang, B. Li, H. Chen, H. Lu, H. Ma, X. Cheng, W. Wang, Y. Wang, Y. Ding and A. Hu, *J. Org. Chem.*, 2020, **85**, 9808–9819.

75 M. Zhang, H. Lu, B. Li, H. Ma, W. Wang, X. Cheng, Y. Ding and A. Hu, *J. Org. Chem.*, 2021, **86**, 1549–1559.

76 P. W. Peterson, R. K. Mohamed and I. V. Alabugin, *Eur. J. Org. Chem.*, 2013, **2013**, 2505–2527.

77 I. V. Alabugin, G. dos Passos Gomes and M. A. Abdo, *WIREs Comput. Mol. Sci.*, 2019, **9**, 1–66.