

1 **Detrending tree-ring widths in closed-canopy forests for climate and disturbance history**

2 **reconstructions**

3 Daniel L. Druckenbrod ^{a,*}, Edward R. Cook ^b, Neil Pederson ^c, Dario Martin-Benito ^d

4 ^a Rider University, Lawrenceville, NJ, USA

5 ^b Lamont-Doherty Earth Observatory of Columbia University, NY, USA

6 ^c Harvard Forest, Harvard University, Petersham, MA, USA

7 ^d Institute of Forest Sciences (ICIFOR) INIA-CSIC, Madrid, Spain

8 * Corresponding author, Email address: ddruckenbrod@rider.edu (D.L. Druckenbrod).

10 **Abstract**

11 Canopy disturbance events in forests often increase light availability and growth rates for
12 surviving trees. Using ring widths, release-detection methods identify the initiation of rapid
13 growth associated with these events enabling reconstructions of forest disturbance history.
14 Conversely, dendroclimate reconstructions minimize these rapid growth responses by detrending
15 entire ring-width time series to resolve underlying climate signals. Incorporating advantages
16 from both of these approaches, we present a canopy disturbance detrending method that
17 quantifies the subsequent, additional growth from release events as a discrete time series. Our
18 method uses radial-growth averaging to detect release events and then power transformation and
19 age-dependent smoothing splines to detrend individual release events, separating canopy
20 disturbance responses from climatic effects. We test our canopy disturbance detrending method
21 on both a coniferous and a broadleaf ring-width chronology from two contrasting temperate
22 forests in eastern North America with known disturbance histories. The resulting disturbance
23 growth time series agrees with the documented forest histories for each forest. Removing the

effects of canopy disturbance from the ring-width series in each chronology improves climate correlations with monthly values of temperature and precipitation and may recover more low-frequency variation with past climate compared to other common detrending methods. Our method provides an alternative approach for detrending disturbance events in closed-canopy forests that should be accessible and useful for both ecological and climatological reconstructions.

Keywords

dendroclimatology, dendroecology, age-dependent spline, radial-growth averaging, power transformation, canopy disturbance

1. Introduction

Ring widths often record growth releases in trees receiving increased light availability after surviving forest canopy disturbance events. Ranging from asynchronous endogenous tree mortality such as gap-phase dynamics, to more synchronous exogenous stand-wide mortality (Shugart, 1984; White, 1979), canopy disturbance events are important in the long-term development of forests from tropical to boreal latitudes (McCarthy, 2001; Seidl and Turner, 2022). Growth releases can be detected in tree-ring widths and used to reconstruct canopy disturbance events and their long-term legacies in forest composition and structure (Henry and Swan, 1974; Lorimer and Frelich, 1989; Marshall, 1927; Nowacki and Abrams, 1997; Oliver and Stephens, 1977; Stephens, 1956). Because growth patterns created by these release events add non-climatic information to ring-width time series, these events also degrade reconstructions of

past climate, presenting a significant challenge for paleoclimate studies in forested regions (Cook and Kairiukstis, 1990).

Climate and disturbance are recognized as two important drivers of tree growth in a conceptual model of tree growth (Cook, 1987) representing ring width (R_t) as a summation of the effects of a biological age trend (A_t), climate (C_t), endogenous and exogenous disturbance events (D_t), and other factors (E_t) on radial growth. Radial-growth averaging methods identify the frequency and relative magnitude of past canopy disturbance events, but these methods do not isolate a tree's growth response as a discrete time series such as represented by D_t . Conversely, dendroclimatic methods minimize A_t (Fritts, 1976) typically using negative exponential functions and also D_t using smoothing splines (Cook and Peters, 1981) and data-adaptive power transformations (Cook and Peters, 1997) on ring-width time series, but not in a manner that explicitly identifies canopy disturbance events. Warren (1980) introduced an additive detrending approach for removing canopy disturbance events using a general exponential curve; however, more recently age-dependent splines were developed to capture the rapid increase in a tree's earliest ring-widths followed by decreased growth as tree age increases as a time-varying response (Melvin et al., 2007). Melvin et al. (2007) applied age-dependent splines to reconstruct climate using regional curve standardization (Briffa et al., 1992; Helama et al., 2017); but they also suggested that time-varying splines may have other applications in dendrochronology. Here, we recognize that while canopy disturbance events may result in various ring-width patterns (*sensu* Lorimer and Frelich, 1989), these patterns also often feature a rapid increase in ring widths followed by a gradual decrease. Lastly, we also note that time-series statistical methods using autoregressive outliers and intervention detection (Druckenbrod, 2005; Druckenbrod et al., 2013) have previously been the only method capable of both identifying release events and detrending disturbance-growth

time series; however, a recent experimental test suggests that these methods may not be more accurate than radial-growth averaging in detecting the timing of these events (Trotsiuk et al., 2018).

Building on these previous methods, we develop a new approach that incorporates radial-growth averaging to identify release events and age-dependent splines with data-adaptive power transformation to detrend a ring-width series into its discrete underlying components, including the responses of trees to canopy disturbance and climatic variation. Using two example chronologies from eastern North America, an coniferous species from a diverse, mesic forest (Palmaghatt Ravine, New York) and a broadleaf species from a drier forest (Democrat Ridge, Missouri), we (1) present our canopy disturbance detrending method with examples using individual tree-ring width series, (2) compare reconstructions of canopy disturbance events with documented forest histories, and (3) evaluate monthly climate correlations with tree-ring widths and how those correlations vary with increasing time period using wavelets.

These two forest case studies exemplify many of the typical drivers of release events in closed-canopy forests and their ensuing disturbance regimes. Temperate forests in eastern North America are often dominated by gap dynamics at the scales of individual trees, but at larger spatial and temporal scales may be affected by more extensive disturbance agents including external agents such as pests and pathogens as well as physical agents including fire, windstorms, and drought (Delcourt et al. 1982, Druckenbrod et al. 2019). Natural disturbance regimes in most temperate forests typically have a return interval of 50 to 200 years; however, trees in these forests often require multiple release events to attain canopy status, enabling some trees to reach older ages (Runkle 1985). Human land use and other anthropogenic activities are

now also a major driver of disturbance regimes, whether directly or indirectly, in temperate forests (Gilliam 2016, Sommerfeld et al 2018).

2. Material and methods

2.1. Forest descriptions and documented canopy disturbance histories

Forests in the Palmaghatt Ravine (41.71N, 74.25W) were broadly classified as Hemlock-White Pine-Northern Hardwoods by Braun (1950) and then re-classified as Northern Hardwoods-Hemlock by Dyer (2006). Palmaghatt Ravine is located in the Shawangunk Mountains of New York State near where the first quantitative study of past hydroclimate from closed-canopy tree species in temperate mesic regions (Cook and Jacoby, 1977). Forests at Democrat Ridge (37.41N 92.00W), located in the Ozark Plateau, were classified as Oak-Hickory forest by both Braun (1950) and Dyer (2006).

Numerous extensive historical human impacts have been documented in the Palmaghatt Ravine forest and the larger Shawangunk Mountain region involving both harvesting and nonnative, invasive species. Selective logging of large hemlock trees (*Tsuga canadensis* (L.) Carr.) for the tanning industry began within the Shawangunk Mountains in 1849 (Thompson and Huth, 2011), leaving mostly smaller hemlocks in Palmaghatt Ravine by the end of that century (Pike, 1892). While logging was ongoing, the Palmaghatt Ravine was purchased privately for conservation in the late nineteenth century (ca. 1897 in Knickerbocker (1937)) and is now part of Minnewaska State Park. Across the Shawangunk Mountains, most of the merchantable timber was logged by 1892 and the arrival of chestnut blight (*Cryphonectria parasitica* (Murrill) Barr) in 1912 led to severe mortality of American chestnut trees and salvage logging into the 1920s (Thompson and Huth 2011; Huth and Smiley, 1985). After large-scale logging ceased in the Shawangunk

Mountains in the 1930s, defoliation by another introduced organism, spongy moth (*Lymantria dispar* L.) led to widespread documented oak mortality in 1986 and 1987 (Thompson and Huth 2011) and an earlier outbreak in 1981 that was so severe that moths in southern New York defoliated other species like hemlock in addition to hardwood species (Souto and Shields, 2000; D'Arrigo et al., 2001). Similarly, Hemlock Woolly Adelgid (*Adelges tsugae* Annand) was first documented regionally in 1991 (Morin et al., 2009) and led to widespread hemlock mortality by the end of the century (Orwig et al., 2002). Ice storms were reported to have impacted trees within Palmaghatt Ravine in 1942 and November 2002 (Yarrow, 2003).

Extensive historical logging occurred in the Ozark Plateau in the early twentieth century as European settlement reached this region more recently (Cleaveland and Stahle 1989; Guldin, 2008); however, remnant old post oak (*Quercus stellata* Wangenh.) forests remain in the Ozarks (Stahle, 1996). The chestnut blight also spread to this region from eastern North America by 1940, decreasing the abundance of Ozark chinkapin (*Castanea ozarkensis* Ashe) (Chapman et al., 2006).

2.2. Forest sampling design

These two sites also differed in sampling designs. Democrat Ridge followed a classic dendroclimatology-oriented design sampling trees that appeared old while Palmaghatt Ravine used an ecological plot design where all trees were sampled above diameter thresholds. At Democrat Ridge, post oaks were selectively sampled for an annual streamflow reconstruction (Cleaveland and Stahle, 1989). At Palmaghatt Ravine, five circular plots were surveyed along a 1-km transect extending along a moisture availability gradient (*sensu* Whittaker, 1956). Sampling at Palmaghatt Ravine surveyed a closed-canopy, temperate deciduous forest with ecological communities that have a stronger resemblance to the Mesophytic and Appalachian

oak section of Dyer (2006). Tree species in the wetter end of our transect are dominated by eastern hemlock and yellow birch (*Betula alleghaniensis* Britton). The dominant species in the plot at the drier end were chestnut oak (*Quercus montana* Willd.) and northern red oak (*Q. rubra* L.). In the intermediate plots, additional species included red maple (*Acer rubrum* L.), sweet birch (*B. lenta* L.), white ash (*Fraxinus americana* L.), and tulip poplar (*Liriodendron tulipifera* L.). Here, we use *T. canadensis* samples from the two most mesic plots along this transect for analysis.

In the plots at Palmaghatt, live and dead trees were sampled based on diameter at breast height (DBH) within nested circular plots to sample trees across size classes: 8 m radius for trees ≥ 13.5 cm DBH, 20 m for trees ≥ 25.0 cm, and 30 m for trees ≥ 40 cm following a design similar to Martin-Benito et al (2020). This approach samples trees proportionally across canopy strata to capture both the fading record of forest canopy disturbances present only in older canopy trees as well as the fading forward record typically present in younger, canopy subordinate trees (Druckenbrod et al., 2019). For both sites, two cores were typically collected per tree, visually crossdated, and statistically checked using Cofecha (Holmes, 1983).

2.3. Example of canopy disturbance detrending method on an eastern hemlock ring-width series

Canopy disturbance detrending involves consecutive steps applied onto each ring-width series (Table 1). Figure 1 shows an example application of this method to a time series from a core extracted from an older eastern hemlock at Palmaghatt Ravine, which had an earliest ring of 1737 (see Appendix Figure A1 for an example post oak series). Release events are identified following the earlier canopy accession settings of 15-year running means and a 50% percentage growth increase threshold using radial-growth averaging (Lorimer and Frelich, 1989). When

more than one adjacent year was greater than this threshold, the maximum percentage growth value was used to determine the year of release in this example (Altman 2020).

The ring-width series is then power transformed to reduce its heteroscedasticity since the variance of ring-width series generally increase with their mean (Cook and Kairiukstis 1990). Quantifying a series' heteroscedasticity enables an adjustment of the variance of a series when its mean values are reduced through subtraction from detrending (Cook and Peters, 1997). This enables a re-scaling of the variance at a new mean level after canopy disturbance detrending.

Beginning with the most recent event, the ring-width series is detrended from the initial year of each release event to the last year of the series using an age-dependent spline with an initial stiffness set to 10 years to capture the often rapid growth response after a canopy disturbance event. The initial value of each detrended release is set to the transformed ring-width value of the first year of the release. This process iterates until all release events have been detrended (Figure 1B-D). The core in Figure 1 detected release events in 1786, 1893, and 1937. This series also shows how an age-dependent spline is capable of detrending both transient (e.g., 1786) and sustained (e.g., 1893 and 1937) canopy release events. For this method, sustained release events occur when disturbance detrended ring-width values remain less than initial ring-width values for the remainder of the series after the release event. Conversely, transient events occur when detrended values only decrease initial ring-width values for a certain sequence of years after the release event instead of the entire remaining series. Transient release events terminate when the detrended value would be greater than the initial ring-width value, indicating that a tree likely returned to suppressed growth conditions.

After removing endogenous and exogenous disturbance events (D_t) using radial-growth averaging and an age-dependent spline, one can employ common dendroclimatic methods to

detrend an age trend across the entire series. Here we use an age-dependent spline across the entire power-transformed series to remove A_t . The transformed, D_t detrended series along with D_t and A_t detrended series may also be re-expressed back in the original units of tree-ring width (Figure 1E), producing a disturbance and age standardized series resembling the underlying age trend common in many trees. The gray-shaded difference between the initial and disturbance-detrended series can be viewed as a disturbance-growth time series measured in units of ring-widths per year that may have ecological applications (*sensu* Druckenbrod et al., 2013). Dividing the disturbance-standardized series by A_t creates a climate index (Figure 1F), C_t , as traditionally used in dendroclimatology.

2.4. Climate data, comparison detrending methods, and tree-ring indices

We compared correlations of both the detrended Palmaghatt eastern hemlock and Democrat Ridge post oak tree-ring width chronologies with monthly climate variables between our canopy disturbance detrending method and two detrending methods commonly used in dendroclimatology (negative exponential curve and a two-thirds spline). At Palmaghatt Ravine, we used monthly climate data from (Cook et al., 2010), a homogenous record collected approximately 10 km NE within adjacent preserved forest land (Mohonk Lake USHCN substation 305426) while we used monthly climate data from PRISM (PRISM Climate Group, Oregon State University, <https://prism.oregonstate.edu>, data accessed 21 Dec 2022) for Democrat Ridge. R package dplR (Bunn 2008; Bunn et al., 2022) was used to generate three standard, ring-width index chronologies for climate comparisons for both sites: a negative exponential curve, a two-thirds spline, and the canopy disturbance detrending method presented in this paper. All sampled cores, including replicates from individual trees, were included in each chronology at both sites. The seacorr function (Meko et al., 2011) in R package treeclim (Zang

and Biondi, 2022) was used to investigate possible correlations with monthly precipitation and temperature as either primary or secondary climate variables. Partial correlations between secondary climate variables and tree ring chronologies suggest additional association between a chronology that is not explained by correlation between the two climate variables (Meko et al. 2011).

Bradshaw and Spies (1992) used wavelets to investigate spatial patterns of canopy gap structure, recognizing that these patterns are aperiodic. Similarly, we recognize that canopy disturbance events are also often aperiodic through time and use wavelets to express patterns over increasing time periods. Indeed, it is this lack of regularity that prevents the use of traditional spectral processing to isolate disturbance signals in tree-ring time series and provides the genesis for the climate reconstruction methods compared in this paper. We use wavelet coherence in R package biwavelet (Gouhier et al. 2021) between each chronology and significant climate variables correlated with these chronologies to investigate how these correlations vary by frequency (see Allen et al. (2017) for an example with tree-ring chronologies) and whether this method retains more decadal-scale agreement with climate.

3. Results

3.1. Canopy disturbance detrending examples with an eastern hemlock chronology and a post oak ring-width chronology

In the eastern hemlock ring-width chronology, 137 disturbance events were detected in 59 series out of 67 total series. Release events clustered in the following decades 1870s, 1920s and 1930s, and 1980s (Figure 2A). In the post oak chronology, 94 disturbance events were detected across 52 series out of 61 total series in the chronology with the largest peaks in the decades of 1800 and 1900 (Figure 3A.). Detrending those series created a mean disturbance-growth increment

time series for both chronologies (Figure 2B and 3B gray shaded areas). The mean of the disturbance-standardized series decreases during these decades with the detrending of release events using an age-dependent spline; however, a smaller growth increase still remains from the recruitment of young, rapidly-growing trees during these disturbance events (blue line). These trees are evident by the histogram of tree ages using earliest rings per tree for the plot-based hemlock chronology (Figure 2C). Subsequently dividing each series by its age-trend produces tree-ring width indices that may be averaged across series to isolate climate signals in both chronologies (Figure 2D and 3C).

Re-expressing the transformed, disturbance-detrended series back to ring widths led to a negative value in year 2003 for 21 of 67 eastern hemlock series during an episode of severe suppressed growth. Ring-width values for this year were unusually small initially (with seven series recording a missing ring at that year prior to analysis) and reduction of growth from detrending with an age-dependent spline decreased these to negative values. For this example, values for this year were set to 0.001 mm to allow for the smallest possible measured growth rate. With the modified negative exponential detrending method and two-thirds spline methods, one hemlock series each also produced negative values and were detrended instead by their mean within dplR. In contrast, none of the canopy disturbance detrended series in the post oak ring-width chronology had negative values when re-expressed from transformed units to ring widths nor did the modified negative exponential, or two-thirds spline detrending produce negative values.

3.2. Effect of detrending methods on climate-growth correlations

Both eastern hemlock and post oak ring-width chronologies using traditional negative exponential detrending, two-thirds spline, and our canopy disturbance detrending method showed significant correlations with monthly temperature or precipitation (Table 2. Current and

prior year correlations are presented in Appendix Table A2 and time series comparison in Appendix Figure A2). For the hemlock chronology, temperatures during March correlated significantly using all three detrending methods; however, higher magnitude correlations occurred with the disturbance detrended chronology whether considering months independently or averaged from January through March.

For the post oak chronology, the highest magnitude correlations occurred with May and June precipitation totals across all three detrending methods, but the canopy disturbance detrending method also correlated more strongly with these months than the other two methods. The canopy disturbance detrending method also had the strongest negative correlation with June temperatures.

Descriptive statistics of ring-width indices from both chronologies show that while the average interseries correlation ($\bar{r}$) and estimated population signal (EPS) across the entire chronology is slightly lower using the modified negative exponential detrending, mean first-order autocorrelation is lower in the canopy disturbance detrending method (Table 3). Considering just the common interval with the climate data, $\bar{r}$ and EPS were both slightly higher with the canopy disturbance detrending method.

Wavelet coherence of the eastern hemlock chronology with and January-March mean monthly temperature showed an increase in significant correlations at periods between 8-32 years for the canopy disturbance detrending method compared with the other two detrending methods (Figure 4). For the post oak chronology, wavelet coherence showed significant correlations at periods between 8-32 years across all three detrending methods (Figure 5) with only a slight increase using the canopy disturbance detrending method. At both study sites, this correlation increase

occurred more recently in the chronology, when more release events are also detrended using this method.

4. Discussion

4.1. Advantages of canopy disturbance detrending

Climate correlations indicate that our new approach isolates climate signals from both a coniferous and a deciduous ring-width chronology as well as if not slightly better than previous methods. Canopy disturbance detrending shows a lower mean first-order autocorrelation and higher common signal ($\bar{r}$ and EPS) among ring-width indices than the other detrending methods suggesting that it is removing persistence from release events that obscure the underlying climate signal. Wavelet analyses show that our approach still retains low frequency variation in growth that is significantly correlated with low frequency variation of past climate. Wavelet coherence of the eastern hemlock population at the Palmaghatt Ravine revealed greater information at periods between 8 and 32 years than prior methods. Retention of a greater range of past climatic variation over varying time-scales is the predominant motivation of detrending methods in tree-ring analysis (RCS, modified negative exponential, etc). Here, in perhaps one of the more challenging forest types, disturbance regimes, and climatologies for the extraction of past climatic variation, the diverse and dense mixed-mesophytic forest, we find reasonable improvement from prior approaches. Notably, the Palmaghatt Ravine experienced a wide range of canopy disturbances and disturbance intensities over the last 140 years, a period upon which the entire calibration of tree rings into climatic information are made. This suggests that for the near term, past few decades, and prior century, our method made substantial improvements. For the post oak population in the drier region containing Democratic Ridge, we see only a slight

improvement in the recovery of past growth variation as all three methods contain significant correlations with climate at periods between 8 and 32 years. The reasons for this are not clear, but could potentially be explained by environmental or species-specific differences. First, the more open-grown post oak canopy at Democratic Ridge might lessen the impact of changes in competition on climate signals (*sensu* Phipps 1982). Second, temperature data may have more red-noise information than precipitation (Pelletier and Turcotte 1997). Third, shade tolerant tree species, including eastern hemlock, have also been shown to respond more positively to canopy disturbance events (Jones et al., 2009) and canopy disturbance events were more frequently observed at Palmaghatt Ravine than at Democrat Ridge. We cannot be sure, but can envision experiments to determine the utility of our method in different forests with different densities, species compositions, and forest histories or testing using pseudoproxies with known disturbances. At this point with these data, we recommend our approach for the recovery of climatic signals and variations of growth patterns over a range of periodicities in closed-canopy forests.

4.2. Comparing documentary and tree-ring forest disturbance histories

The documented history of the Palmaghatt Ravine forests is broadly reflected in the eastern hemlock reconstruction of canopy disturbance history (Figure 2). Extraction of hemlock for tanning likely led to the pronounced release events and subsequent recruitment beginning in the 1870s and continuing into the 1880s (Knickerbocker 1937); chestnut blight and subsequent salvage logging to the releases in the 1920s and 1930s (Thompson and Huth 2011; Huth and Smiley, 1985). Spongy moth defoliation may have led to the releases in the 1980s for hemlocks (Thompson and Huth 2011), which may have been less impacted than oaks and thus able to increase their subsequent growth rate.

320 Conversely eastern hemlock trees as evergreens were likely severely impacted from the arrival of
321 the Hemlock Woolly Adelgid and the November 2002 ice storm. The pronounced growth
322 suppression during the subsequent growing season led to negative detrended growth values when
323 re-expressed as ring-widths. Retaining transformed ring-width residuals in climate analyses
324 would avoid this artifact in detrended series in addition to the advantages of residuals outlined in
325 Cook and Peters (1997). The complexities of sequential canopy release events and canopy
326 suppression within this forest make it challenging to detrend, but a useful example for evaluating
327 our canopy disturbance detrending method.

328 Other studies of eastern hemlock in northeastern North America have found similar responses in
329 release events and recruitment to canopy disturbance events (Abrams et al., 2000; Abrams and
330 Orwig, 1996; Black and Abrams, 2005), particularly considering that eastern hemlock often lags
331 in its growth response to canopy disturbance events. Carter et al. (2021) detected releases in 74 -
332 86% of hemlock trees within two years of a known harvest with a maximum growth rate
333 occurring 6.8 ± 0.3 years after harvest. In a separate experiment, hemlock responded with a
334 maximum growth rate within 5 years of harvest and most species responded with maximum
335 growth rates within 3 to 15 years post-harvest (Jones et al. 2009).

336 In the post oak chronology, the ca. 1900 canopy disturbance recorded also aligns with intensive
337 forest logging in the region (Cleaveland and Stahle 1989). The ca. 1800 event is not documented
338 and may have resulted from a natural disturbance event. These two canopy disturbances are the
339 most pronounced events over the duration of this chronology. While the earlier canopy
340 disturbance appears to have had more of a transient growth release on the chronology, the ca.
341 1900 event led to a sustained growth release in the chronology suggesting that these sampled
342 trees obtained a dominant canopy position after this logging event.

4.3. Comparing tree-ring climate reconstructions from three detrending methods

This canopy disturbance detrending method isolated climate signals from both a coniferous and a broadleaf tree-ring chronology removing the effect of disturbances on tree growth and increasing its correlations with monthly climate variables at least as well as previous methods such as negative exponential curves and smoothing splines (Table 2). The climate signal of eastern hemlock at Palmaghatt is similar to a previous analysis using this species within the Shawangunk Mountains with a positive correlation with March temperatures but unlike that previous sampling, this chronology did not express a correlation with precipitation across prior and current months (Blasing et al., 1984; Cook and Jacoby, 1977). This difference may arise from the use of partial correlations in this analysis with the secondary precipitation variable, which reduce the influence of collinearity across climate variables (Meko et al., 2011). The post oak climate signal from Democrat Ridge with its positive correlations to May-June precipitation and negative correlations to June-Sept. temperatures aligns with other climate reconstructions from this species in the Ozarks (Cleaveland and Stahle, 1989; LeBlanc and Berland, 2019; Prolic and Goldblum, 2016).

Conceptually, these results also align with previous research showing that tree-ring series with release events had stronger climate reconstructions if the series was split into pre and post release segments that were separately detrended with splines (Blasing et al., 1983). However, this method detects and detrends the response of tree growth to canopy disturbance events by explicitly targeting adaptive splines to account for these aperiodic events within a time series.

4.4. Application of method

Intervention detection methods have shown the ability to minimize canopy disturbance events in climate reconstructions based on tree-ring widths (Rydval et al., 2017, 2016). Adding to this, we

hope that this method will be more accessible to other existing tree-ring resources such as the dplR and TRADER packages in the R programming language (Altman et al., 2014; Bunn et al., 2022), which will broaden the applicability of this approach to explicitly quantifying disturbance and climate time series from closed-canopy forests.

Our method also has the flexibility to incorporate other canopy release detection methods and detrending methods for climate analysis. The Lorimer and Frelich (1989) settings could be adjusted or replaced with other release detection approaches as reviewed by Altman (2020). Similarly, canopy suppression events could also potentially be detected and detrended, assisting with estimating the impacts from ice storms as shown with the eastern hemlock chronology. Other detrending or growth increment functions could also replace an age-dependent spline for removing canopy disturbance or age trends. For example, regional curve standardization (Helama et al., 2017), could also be considered for removing age trends from disturbance-detrended series as they may more likely approximate growth patterns of open-canopy forests after removing the canopy disturbance signal.

Tree-ring widths respond to multiple external environmental forcing events including climate and canopy disturbance events. Quantifying the responses from these independent drivers of tree growth remains as important now (e.g., Anderson-Teixeira et al., 2022) as it did when Cook (1987) proposed the conceptual growth model. This method provides an integrative approach to partitioning tree growth that incorporates previously separate advances from dendroclimatology and forest ecology.

5. Conclusions

We present a canopy disturbance detrending method that combines radial-growth averaging with data-adaptive power transformation and age-dependent splines to detrend ring widths into explicit time series quantifying the effect of canopy disturbance and climate on tree growth. Building from previous research on intervention detection methods in tree-ring width series, this more straightforward and accessible approach disentangles canopy disturbance and climate signals from trees in closed canopy forests. Comparing with traditional dendroclimatic detrending methods of a negative exponential curve and a two-thirds spline, this method not only effectively resolves a past climate signal, but it also detects release events and generates time series of disturbance growth increments. Results from eastern hemlock and post oak chronologies indicate that this method should have potential applications in studies within closed canopy forests of either past climate or ecological interactions or studies that consider both of these factors that influence tree growth.

Acknowledgements

We thank Malcolm Cleaveland and Dave Stahle for sharing their Democrat Ridge samples (mo024) with the ITRDB. For sampling Palmaghatt forests, we thank Jaclyn Testani, Javier Martin Fernandez, Ana Camila Gonzalez, Kristina Seftigen, Jesper Björklund, Adam Wolff, and, for one plot, the 2013 North American Dendroecological Fieldweek (NADEF) ecological dynamics group while Jaclyn Testani, Javier Martin Fernandez, Ana Camila Gonzalez led core processing. Permission to sample was given by resource managers of Minnewaska State Park Preserve. We also thank Dave Meko for sharing his Tree-Ring Toolbox for Matlab and three reviewers for their comments. A collaborative NSF P2C2 Award supported Daniel Druckenbrod (#2102790) and Ed Cook (#2102759). Dario Martin-Benito was partly funded by projects

AGL2015-73190-JIN, PID2019-110273RB-I00 and contract RYC-2017-23389 by the Spanish Ministry of Science and Innovation MCIN/AEI.

References

Abrams, M.D., Van de Gevel, S., Dodson, R.C., Copenheaver, C.A., 2000. The dendroecology and climatic impacts for old-growth white pine and hemlock on the extreme slopes of the Berkshire Hills, Massachusetts, U.S.A. *Can. J. Bot.* 78, 851–861.

<https://doi.org/10.1139/b00-057>

Abrams, M.D., Orwig, D.A., 1996. A 300-Year History of Disturbance and Canopy Recruitment for Co-Occurring White Pine and Hemlock on the Allegheny Plateau, USA. *Journal of Ecology* 84, 353–363. <https://doi.org/10.2307/2261198>

Allen, K.J., Fenwick, P., Palmer, J.G., Nichols, S.C., Cook, E.R., Buckley, B.M., Baker, P.J., 2017. A 1700-year *Athrotaxis selaginoides* tree-ring width chronology from southeastern Australia. *Dendrochronologia* 45, 90–100. <https://doi.org/10.1016/j.dendro.2017.07.004>

Altman, J., 2020. Tree-ring-based disturbance reconstruction in interdisciplinary research: Current state and future directions. *Dendrochronologia* 63, 125733.

<https://doi.org/10.1016/j.dendro.2020.125733>

Altman, J., Fibich, P., Dolezal, J., Aakala, T., 2014. TRADER: A package for Tree Ring Analysis of Disturbance Events in R. *Dendrochronologia* 32, 107–112.

<https://doi.org/10.1016/j.dendro.2014.01.004>

Anderson-Teixeira, K.J., Herrmann, V., Rollinson, C.R., Gonzalez, B., Gonzalez-Akre, E.B., Pederson, N., Alexander, M.R., Allen, C.D., Alfaro-Sánchez, R., Awada, T., 2022. Joint

433 effects of climate, tree size, and year on annual tree growth derived from tree-ring records of
434 ten globally distributed forests. *Global Change Biology* 28, 245–266.

435 Black, B.A., Abrams, M.D., 2005. Disturbance history and climate response in an old-growth
436 hemlock-white pine forest, central Pennsylvania. *Journal of the Torrey Botanical Society*
437 132, 103–114.

438 Blasing, T.J., Duvick, D.N., Cook, E.R., 1983. Filtering the effects of competition from ring-
439 width series. *Tree-Ring Bulletin* 43, 19–31.

440 Blasing, T.J., Solomon, A.M., Duvick, D.N., 1984. Response functions revisited. *Tree-Ring*
441 *Bulletin* 44, 1–15.

442 Bradshaw, G.A., Spies, T.A., 1992. Characterizing canopy gap structure in forests using wavelet
443 analysis. *Journal of Ecology* 205–215.

444 Braun, E.L., 1950. *Deciduous Forests of Eastern North America*. Hafner, New York.

445 Briffa, K.R., Jones, P.D., Bartholin, T.S., Eckstein, D., Schweingruber, F.H., Karlén, W.,
446 Zetterberg, P., Eronen, M., 1992. Fennoscandian summers from AD 500: temperature
447 changes on short and long timescales. *Climate Dynamics* 7, 111–119.
448 <https://doi.org/10.1007/BF00211153>

449 Bunn, A.G., 2008. A dendrochronology program library in R (dplR). *Dendrochronologia* 26,
450 115–124. <https://doi.org/10.1016/j.dendro.2008.01.002>

451 Bunn, A., Korpela, M., Biondi, F., Campelo, F., Mérian, P., Qeadan, F., Zang, C., Buras, A.,
452 Cecile, A., Mudelsee, M., Schulz, M., Klesse, S., Frank, D., Visser, R., Cook, E.,
453 Anchukaitis, K., 2022. dplR: Dendrochronology Program Library in R.

454 Carter, D.R., Bialecki, M.B., Windmuller-Campione, M., Seymour, R.S., Weiskittel, A., Altman,
455 J., 2021. Detecting growth releases of mature retention trees in response to small-scale gap

disturbances of known dates in natural-disturbance-based silvicultural systems in Maine.
Forest Ecology and Management 502, 119721. <https://doi.org/10.1016/j.foreco.2021.119721>

Chapman, R.A., Heitzman, E., Shelton, M.G., 2006. Long-term changes in forest structure and
species composition of an upland oak forest in Arkansas. Forest Ecology and Management
236, 85–92.

Cleaveland, M.K., Stahle, D.W., 1989. Tree ring analysis of surplus and deficit runoff in the
White River, Arkansas. Water Resources Research 25, 1391–1401.
<https://doi.org/10.1029/WR025i006p01391>

Cook, B.I., Cook, E.R., Anchukaitis, K.J., Huth, P.C., Thompson, J.E., Smiley, S.F., 2010. A
Homogeneous Record (1896–2006) of Daily Weather and Climate at Mohonk Lake, New
York. Journal of Applied Meteorology and Climatology 49, 544–555.
<https://doi.org/10.1175/2009JAMC2221.1>

Cook, E.R., 1987. The decomposition of tree-ring series for environmental studies. Tree-Ring
Bulletin 47, 37–59.

Cook, E.R., Jacoby, G.C., 1977. Tree-ring drought relationships in the Hudson Valley, New
York. Science 198, 399–401.

Cook, E.R., Kairiukstis, L.A. (Eds.), 1990. Methods of Dendrochronology. International Institute
for Applied Systems Analysis, Netherlands.

Cook, E.R., Peters, K., 1997. Calculating unbiased tree-ring indices for the study of climatic and
environmental change. Holocene 7, 361–370.

Cook, E.R., Peters, K., 1981. The smoothing spline: a new approach to standardizing forest
interior tree-ring width series for dendroclimatic studies. Tree Ring Bulletin 41, 45–53.

478 D'Arrigo, R.D., Schuster, W.S.F., Lawrence, D.W., Cook, E.R., Wiljanen, M., Thetford, R.D.,
479 2001. Climate-growth relationships of eastern hemlock and chestnut oak from Black Rock
480 Forest in the highlands of southeastern New York. *Tree-Ring Research* 57, 183–190.

481 Delcourt, H.R., Delcourt, P.A., Webb, T., 1982. Dynamic Plant Ecology: the spectrum of
482 vegetational change in space and time. *Quaternary Science Reviews* 1, 153–175.

483 Druckenbrod, D.L., 2005. Dendroecological reconstructions of forest disturbance history using
484 time-series analysis with intervention detection. *Canadian Journal of Forest Research* 35,
485 868–876.

486 Druckenbrod, D.L., Martin-Benito, D., Orwig, D.A., Pederson, N., Poulter, B., Renwick, K.M.,
487 Shugart, H.H., 2019. Redefining temperate forest responses to climate and disturbance in the
488 eastern United States: New insights at the mesoscale. *Global Ecology and Biogeography* 28,
489 557–575. <https://doi.org/10.1111/geb.12876>

490 Druckenbrod, D.L., Pederson, N., Rentch, J., Cook, E.R., 2013. A comparison of times series
491 approaches for dendroecological reconstructions of past canopy disturbance events. *Forest*
492 *Ecology and Management* 302, 23–33. <https://doi.org/10.1016/j.foreco.2013.03.040>

493 Dyer, J.M., 2006. Revisiting the Deciduous Forests of Eastern North America. *Bioscience* 56,
494 341–352. [https://doi.org/10.1641/0006-3568\(2006\)56\[341:rtdfoe\]2.0.co;2](https://doi.org/10.1641/0006-3568(2006)56[341:rtdfoe]2.0.co;2)

495 Fritts, H.C., 1976. *Tree rings and climate*. Academic Press, New York.

496 Gilliam, F.S., 2016. Forest ecosystems of temperate climatic regions: from ancient use to climate
497 change. *New Phytologist* 212, 871–887. <https://doi.org/10.1111/nph.14255>

498 Gouhier, T.C., Grinsted, A., Simko, V. 2021. biwavelet: Conduct Univariate and Bivariate
499 Wavelet Analyses

500 Guldin, J.M., 2008. A history of forest management in the Ozark Mountains, in: Pioneer Forest:
501 A Half-Century of Sustainable Uneven-Aged Forest Management in the Missouri Ozarks.
502 Gen. Tech. Rep. SRS-108. USDA Forest Service, Southern Research Station, Asheville,
503 North Carolina. pp. 3–8.

504 Helama, S., Melvin, T.M., Briffa, K.R., 2017. Regional curve standardization: State of the art.
505 The Holocene 27, 172–177. <https://doi.org/10.1177/0959683616652709>

506 Henry, J.D., Swan, J.M.A., 1974. Reconstructing forest history from live and dead plant material
507 - an approach to the study of forest succession in southwest New Hampshire. Ecology 55,
508 772–783.

509 Holmes, R.L., 1983. Computer-assisted quality control in tree-ring dating and measurement.
510 Tree-Ring Bulletin 44, 69–75.

511 Huth, P.C., Smiley, D., 1985. Forest Vegetation Changes: A Plant Community Transition - per
512 stipes (Research Report). Mohonk Preserve, Inc.

513 Jones, T.A., Domke, G.M., Thomas, S.C., 2009. Canopy tree growth responses following
514 selection harvest in seven species varying in shade tolerance. Can. J. For. Res. 39, 430–440.
515 <https://doi.org/10.1139/X08-186>

516 Knickerbocker, N.L., 1937. Minnewaska, “Frozen Waters”: The Intimate Story of the Origin and
517 Growth of the Lake Minnewaska Mountain Houses. The author.

518 LeBlanc, D.C., Berland, A.M., 2019. Spatial variation in oak (*Quercus* spp.) radial growth
519 responses to drought stress in eastern North America. Canadian Journal of Forest Research
520 48, 986–993.

521 Lorimer, C.G., Frelich, L.E., 1989. A methodology for estimating canopy disturbance frequency
522 and intensity in dense temperate forests. Canadian Journal of Forest Research 19, 651–663.

523 Marshall, R., 1927. The Growth of Hemlock before and after Release from Suppression. Harvard
524 University, Cambridge, MA.

525 Martin-Benito, D., Pederson, N., Lanter, C., Köse, N., Doğan, M., Bugmann, H., Bigler, C.,
526 2020. Disturbances and climate drive structure, stability, and growth in mixed temperate old-
527 growth rainforests in the Caucasus. *Ecosystems* 23, 1170–1185.

528 McCarthy, J., 2001. Gap dynamics of forest trees: A review with particular attention to boreal
529 forests. *Environmental Reviews* 9, 1–59. <https://doi.org/10.1139/a00-012>

530 Meko, D.M., Touchan, R., Anchukaitis, K.J., 2011. Seascorr: A MATLAB program for
531 identifying the seasonal climate signal in an annual tree-ring time series. *Computers &*
532 *Geosciences* 37, 1234–1241. <https://doi.org/10.1016/j.cageo.2011.01.013>

533 Melvin, T.M., Briffa, K.R., Nicolussi, K., Grabner, M., 2007. Time-varying-response smoothing.
534 *Dendrochronologia* 25, 65–69. <https://doi.org/10.1016/j.dendro.2007.01.004>

535 Morin, R.S., Liebhold, A.M., Gottschalk, K.W., 2009. Anisotropic spread of hemlock woolly
536 adelgid in the eastern United States. *Biol Invasions* 11, 2341–2350.
537 <https://doi.org/10.1007/s10530-008-9420-1>

538 Nowacki, G.J., Abrams, M.D., 1997. Radial-growth averaging criteria for reconstructing
539 disturbance histories from presettlement-origin oaks. *Ecological Monographs* 67, 225–249.

540 Oliver, C.D., Stephens, E.P., 1977. Reconstruction of a mixed-species forest in central New
541 England. *Ecology* 58, 562–572.

542 Orwig, D.A., Foster, D.R., Mausel, D.L., 2002. Landscape Patterns of Hemlock Decline in New
543 England due to the Introduced Hemlock Woolly Adelgid. *Journal of Biogeography* 29, 1475–
544 1487.

545 Pelletier, J.D., Turcotte, D.L., 1997. Long-range persistence in climatological and hydrological
546 time series: analysis, modeling and application to drought hazard assessment. *Journal of*
547 *hydrology* 203, 198–208.

548 Phipps, R.L., 1982. Comments on interpretation of climatic information from tree rings, eastern
549 North America. *Tree-ring Bulletin* 42, 11–22.

550 Pike, M.H., 1892. Shongum - III. *Garden and Forest* V, 483–484.

551 Prolic, C., Goldblum, D., 2016. Spatial and temporal patterns of post oak's (*Quercus stellata*)
552 radial growth-climate relationship. *Southeastern Geographer* 56, 171–186.

553 Runkle, J., 1985. Disturbance Regimes in Temperate Forests, in: Pickett, S.T.A., White, P.S.
554 (Eds.), *The Ecology of Natural Disturbance and Patch Dynamics*. Academic Press, New
555 York, pp. 17–33.

556 Rydval, M., Druckenbrod, D., Anchukaitis, K.J., Wilson, R., 2016. Detection and removal of
557 disturbance trends in tree-ring series for dendroclimatology. *Canadian Journal of Forest*
558 *Research* 46, 387–401. <https://doi.org/10.1139/cjfr-2015-0366>

559 Rydval, M., Loader, N.J., Gunnarson, B.E., Druckenbrod, D.L., Linderholm, H.W., Moreton,
560 S.G., Wood, C.V., Wilson, R., 2017. Reconstructing 800 years of summer temperatures in
561 Scotland from tree rings. *Clim Dyn* 49, 2951–2974. [https://doi.org/10.1007/s00382-016-](https://doi.org/10.1007/s00382-016-3478-8)
562 [3478-8](https://doi.org/10.1007/s00382-016-3478-8)

563 Seidl, R., Turner, M.G., 2022. Post-disturbance reorganization of forest ecosystems in a
564 changing world. *Proceedings of the National Academy of Sciences* 119, e2202190119.

565 Shugart, H.H., 1984. *A theory of forest dynamics: The ecological implications of forest*
566 *succession models*. Springer-Verlag, New York, New York.

567 Sommerfeld, A., Senf, C., Buma, B., D'Amato, A.W., Després, T., Díaz-Hormazábal, I., Fraver,
568 S., Frelich, L.E., Gutiérrez, Á.G., Hart, S.J., 2018. Patterns and drivers of recent disturbances
569 across the temperate forest biome. *Nature communications* 9, 4355.

570 Souto, D.R., Shields, K.S., 2000. Overview of hemlock health. In: McManus, Katherine A.;
571 Shields, Kathleen S.; Souto, Dennis R., eds. *Proceedings: Symposium on sustainable*
572 *management of hemlock ecosystems in eastern North America*. Gen. Tech. Rep. NE-267.
573 Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest
574 Experiment Station. 76-80. 267.

575 Stahle, D.W., 1996. Tree rings and ancient forest relics. *Arnoldia* 56, 2–10.

576 Stephens, E.P., 1956. The uprooting of trees: a forest process. *Soil Science Society of America*
577 *Proceedings* 20, 113–116.

578 Thompson, J.E., Huth, P.C., 2011. *Vegetation dynamics in the northern Shawangunk Mountains.*
579 *Environmental History of the Hudson River: Human Uses that Changed the Ecology,*
580 *Ecology that Changed Human Uses*. SUNY Press, Albany, NY 135–151.

581 Trotsiuk, V., Pederson, N., Druckenbrod, D.L., Orwig, D.A., Bishop, D.A., Barker-Plotkin, A.,
582 Fraver, S., Martin-Benito, D., 2018. Testing the efficacy of tree-ring methods for detecting
583 past disturbances. *Forest Ecology and Management* 425, 59–67.
584 <https://doi.org/10.1016/j.foreco.2018.05.045>

585 Warren, W.G., 1980. On removing the growth trend from dendrochronological data. *Tree-Ring*
586 *Bulletin* 40, 35–44.

587 White, P.S., 1979. Pattern, Process, and Natural Disturbance in Vegetation. *Botanical Review*
588 45, 229–299.

Whittaker, R.H., 1956. Vegetation of the Great Smoky Mountains. Ecological Monographs 26,
1–80.

Yarrow, D., 2003. Palmaghatt survey report. Eastern Native Tree Society.

Zang, C., Biondi, F., 2022. treeclim: Numerical Calibration of Proxy-Climate Relationships.

Figures




Figure 1: Example of the canopy disturbance detrending method on ring widths from an eastern hemlock tree in Palmaghatt Ravine, NY, USA. Three canopy disturbance events (triangles) detected from radial-growth averaging (A) that are then sequentially detrended using an age-dependent spline (B through D) on power-transformed ring widths. Ring widths re-expressed in original units with initial ring widths shown by the upper line and the disturbance-detrended series represented by the lower line. Gray shading depicts the disturbance growth increment beginning after the earliest release in 1786 and fitting an age-dependent spline to the entire series (blue line) represents the age-trend (A_t) for this tree (E). Dividing the disturbance-detrended series by the age trend produces a series that is both age and disturbance standardized index values (F).




Figure 2: Example *Tsuga canadensis* chronology at Palmaghatt Ravine, New York, USA using the canopy disturbance detrending method. Decadal histograms of canopy release events and gray line showing series sample size (A). Disturbance history chronology showing the difference of the initial average series values and the average disturbance-detrended ring widths with the gray shading representing the average disturbance growth increment across all series and an average age trend (blue line) fit to the average disturbance growth series (B). Histogram of tree recruitment with hemlock trees shown in black and other species in white (C) and resulting climate index after removing age trends from the disturbance-detrended series for all eastern hemlock series (D).




Figure 3: Example *Quercus stellata* chronology from Democrat Ridge, Missouri, USA using the canopy disturbance detrending method with decadal histograms of canopy release events and gray line showing series sample size (A). Disturbance history chronology showing the difference of the initial average series values and the average disturbance-detrended ring widths with the gray shading representing the average disturbance growth increment across all series and an average age trend (blue line) fit to the average disturbance growth series (B). Resulting climate index after removing age trends from the disturbance-detrended series for all post oak series (C).




Figure 4: Wavelet coherence between the *Tsuga canadensis* ring-width chronology and January-March mean monthly temperature from 1896-2004 using three detrending methods: Modified Negative Exponential (A), two-thirds spline (B), canopy disturbance detrending (C). After 1940, the correlation increases with the disturbance and age standardized chronology over periods of 8 to 32 years.




Figure 5: Wavelet coherence between the *Quercus stellata* ring-width chronology and June-July total monthly precipitation from 1896-1992 using three detrending methods: Modified Negative Exponential (A), two-thirds spline (B), and canopy disturbance detrending (C). Correlations are similar between the three detrending methods with only a slight increase after 1940 over periods of 8 to 32 years with the disturbance and age standardized chronology towards the present.

Table 1: Description of general steps in canopy disturbance detrending method.

Step	Description
1	Sequentially select each ring-width series for processing (similar to other dendrochronology programs such as Cofecha, ARSTAN, etc.).
2	Detect initiation of release events using radial growth averaging
3	Power transform ring-width series to reduce heteroscedasticity.
4	Using age-dependent splines, iteratively detrend any canopy release events beginning with the most recent event, which quantifies the duration and magnitude of transient and sustained release events.
5	Re-express disturbance growth increment and D_t standardized series in original growth units
6	Detrend across the entire D_t standardized series to minimize A_t using an age-dependent spline, producing a tree-ring index that is age and disturbance standardized (A_t and D_t).

Table 2: Selected simple correlations between detrending methods and primary seasonal climate variables (sensu, Meko 2011). showing significant relationships with total precipitation for a *Q. stellata* (1896-1992) ring-width chronology and with mean temperature for a *T. canadensis* ring-width chronology (1896-2004). * $p \leq 0.05$

<i>Q. stellata</i> Chronology	Total Precipitation		
	May	Jun	May-Jun
Modified Negative Exponential	0.303*	0.383*	0.443*
Two-Thirds Spline	0.292*	0.412*	0.456*
Canopy Disturbance	0.310*	0.475*	0.510*

<i>T. canadensis</i> Chronology	Mean Temperature			
	Jan	Feb	Mar	Jan-Mar
Modified Negative Exponential	0.161	0.180	0.224*	0.295*
Two-Thirds Spline	0.165	0.095	0.226*	0.253
Canopy Disturbance	0.249*	0.201	0.334*	0.409*

Table 3. Descriptive statistics of detrended ring-width indices from two study sites compared by detrending methods showing mean correlation between all series ($\bar{r}$), estimated population signal (EPS), and mean first-order autocorrelation (AR1) for either the entire years of the chronology or for the interval in common with the climate data.

Species	Detrending Method	Entire Chronology			Climate Interval	
		mean AR1	$\bar{r}$	EPS	$\bar{r}$	EPS
<i>Tsuga canadensis</i>	Mod. Neg. Exp.	0.774	0.267	0.960	0.272	0.961
<i>Tsuga canadensis</i>	Two-Thirds Spline	0.650	0.286	0.963	0.290	0.964
<i>Tsuga canadensis</i>	Canopy Disturbance	0.518	0.286	0.963	0.298	0.966
<i>Quercus stellata</i>	Mod. Neg. Exp.	0.567	0.378	0.974	0.365	0.972
<i>Quercus stellata</i>	Two-Thirds Spline	0.425	0.430	0.979	0.453	0.980
<i>Quercus stellata</i>	Canopy Disturbance	0.241	0.432	0.979	0.457	0.981

Appendix




Figure A1: Canopy disturbance detrending of a post oak (*Q. stellata*) series from Democrat Ridge, Missouri, USA, with a release (triangle) detected in 1914 (A). Detrending using an age-dependent spline on power-transformed ring widths (B). Ring widths re-expressed in original units with initial ring widths shown by the upper line and the disturbance-detrended series represented by the lower line. Gray shading depicts the disturbance growth increment beginning after 1914 and fitting an age-dependent spline to the entire series (blue line) represents the age-trend (A_t) for this tree (C). Dividing the disturbance-detrended series by the age trend produces a series that is both age and disturbance standardized index values (D).

930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948

Table A1. Descriptive statistics of both ring-width chronologies.

Species	No. trees	No. series	Timespan (yrs)	Av. series length (yrs)
<i>Tsuga canadensis</i>	26	67	1683-2013	125
<i>Quercus stellata</i>	26	61	1620-1992	186

Table A2. Climate correlations extending from prior year through current year growing season for two example ring-width chronologies using the seacorr function in R package treeclim (Zang and Biondi, 2022) based on Meko et al. (2011). Significant months ($p \leq 0.05$) shown in bold.

Eastern hemlock chronology shown by detrending method with av. temperature as primary response variable using simple correlations. Partial correlations with precipitation totals were not significant for any month with any method at this site.

Method	Var.	Prior Year Months					Current Year Months								
		Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep
ModNegExp	TEMP	-0.042	-0.069	0.084	0.034	0.014	0.161	0.180	0.224	0.055	-0.068	0.124	0.102	0.025	0.035
2/3 Spline	TEMP	-0.150	-0.105	0.045	-0.019	-0.019	0.165	0.095	0.226	0.012	-0.125	0.048	0.036	-0.093	-0.009
Canopy Dist.	TEMP	-0.211	-0.151	0.058	0.091	0.061	0.249	0.201	0.334	0.054	-0.161	0.087	-0.003	-0.090	0.029

Post oak chronology shown by detrending method with precipitation totals as primary response variable using simple correlations and av. temperature as secondary partial correlations.

Method	Var.	Prior Year Months					Current Year Months								
		Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep
ModNegExp	PRCP	0.273	0.000	0.058	0.026	0.003	0.205	0.148	0.051	0.082	0.303	0.383	0.006	0.180	-0.016
Spline	PRCP	0.305	0.020	0.030	0.001	-0.024	0.257	0.148	0.087	0.112	0.292	0.412	0.014	0.209	0.007
Canopy Dist.	PRCP	0.287	-0.008	0.043	0.047	0.004	0.281	0.147	0.162	0.098	0.310	0.475	-0.005	0.179	-0.025
ModNegExp	TEMP	-0.188	-0.198	0.119	0.048	-0.015	-0.075	0.165	0.027	-0.119	-0.027	-0.334	-0.275	-0.260	-0.317
2/3 Spline	TEMP	-0.161	-0.171	0.127	0.119	-0.016	-0.039	0.115	0.084	-0.138	-0.022	-0.349	-0.283	-0.225	-0.266
Canopy Dist.	TEMP	-0.201	-0.182	0.178	0.065	-0.032	-0.063	0.103	0.043	-0.162	-0.002	-0.391	-0.274	-0.259	-0.279




Figure A2. Normalized time series of detrended ring-width chronologies and seasonal climate variables: eastern hemlock with January-March average temperature (A) and post oak with May-June total precipitation in gray (B).