

INTEGRATED LOCAL ENERGY DECAY FOR THE DAMPED WAVE
EQUATION ON STATIONARY SPACE-TIMES*

COLLIN KOFROTH†

Abstract. We prove integrated local energy decay for the damped wave equation on stationary, asymptotically flat space-times in $(1+3)$ dimensions. Local energy decay constitutes a powerful tool in the study of dispersive partial differential equations on such geometric backgrounds. By utilizing the geometric control condition to handle trapped trajectories, we are able to recover high frequency estimates without any loss. We may then apply known estimates from the work of Metcalfe, Sterbenz, and Tataru in the medium and low frequency regimes in order to establish local energy decay. This generalizes the integrated version of results established by Bouclet and Royer from the setting of asymptotically Euclidean manifolds to the full Lorentzian case.

Key words. local energy estimates, asymptotically flat, damped wave equation, trapping, geometric control

MSC codes. 35L05, 58J45, 35Q93

DOI. 10.1137/22M1490545

1. Introduction.

1.1. Background. The goal of this paper is to establish local energy decay for the damped wave equation on asymptotically flat space-times with time-independent metrics (in the sense that ∂_t is a Killing field) subject to the geometric control condition. The primary advance in this work is recovering the high frequency local energy estimate present in [MST20] for waves on nontrapping space-times. Since the aforementioned work only utilized the nontrapping assumption at high frequencies, this establishes the key step in extending time-integrated versions of previously known results for damped waves on product manifolds (see [BR14]) to the full Lorentzian case. From the proven high frequency estimate, we may apply known results in [MST20] to conclude local energy decay to complete this extension.

Local energy estimates are a collection of rich and well-studied quantities within the field of dispersive partial differential equations, originally introduced on Minkowski space in classical works such as [Mor66, Mor68, Mor75], [MRS77]. A particularly important class of local energy estimates is the *integrated local energy estimates*; if u solves the homogeneous flat wave equation

$$(\partial_t^2 - \Delta)u = 0, \quad \Delta = \sum_{j=1}^n \partial_{x_j}^2$$

in spatial dimension $n \geq 3$, then the integrated local energy estimate which we are interested in takes the form

*Received by the editors April 26, 2022; accepted for publication (in revised form) March 20, 2023; published electronically September 27, 2023.

<https://doi.org/10.1137/22M1490545>

Funding: The work of the author was partially supported by the F. Ivy Carroll Summer Research Fellowship awarded by the UNC-CH Graduate School and by National Science Foundation (NSF) grant DMS-2054910 (PI: Jason Metcalfe).

†Department of Mathematics, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599 USA (ckofroth@asu.edu).

$$(1.1) \quad \sup_{j \geq 0} \left(\left\| \langle x \rangle^{-1/2} \partial u \right\|_{L_t^2 L_x^2(\mathbb{R}_+ \times \{\langle x \rangle \approx 2^j\})} + \left\| \langle x \rangle^{-3/2} u \right\|_{L_t^2 L_x^2(\mathbb{R}_+ \times \{\langle x \rangle \approx 2^j\})} \right) \lesssim \|\partial u(0)\|_{L^2},$$

where $\partial = (\partial_t, \nabla)$ denotes the space-time gradient, and $\langle x \rangle = (1 + |x|^2)^{1/2}$ denotes the Japanese bracket of x . This estimate is known to hold in the flat setting through a positive commutator argument using the multiplier introduced in the appendix of [Ste05]. In such a case, we will say that *(integrated) local energy decay* holds. This is a quantitative statement on *dispersion*, and it heuristically expresses that the energy of the wave must decay quickly enough within compact spatial sets to be integrable in time. Estimates of this form have significant utility, as they have been used to prove other important measures of dispersion, such as Strichartz estimates (see [BT07], [BT08], [JSS90], [JSS91], [MMT08], [MMTT10], [MT09], [MT12], [Tat08], [Toh12], and the references therein) and pointwise decay estimates (see [Hin22], [Loo22], [MTT12], [Mor20], [MW21], [Tat13], and references in these works). Additionally, local energy estimates have applications to nonlinear wave equations where one can develop estimates on an appropriate linearization of the problem, viewing the nonlinearity as a perturbation. These techniques have been applied in many works; see, e.g., [BH10], [KSS02], [KSS04], [MS06], [MS07], [SW10], and the citations contained in them. We will be focused on establishing local energy decay rather than demonstrating its utility via applications.

In [MST20], the authors proved that local energy decay holds for a broad class of stationary wave operators if and only if

1. *the space-time is nontrapping*: there are no null bicharacteristic rays which stay within a compact set for all time;
2. *the operator satisfies certain spectral assumptions*: upon replacing time derivatives in the wave operator with a complex parameter, one requires that this family of operators have no eigenvalues in the lower half-plane nor real resonances/embedded eigenvalues (see [MST20] for more precise definitions); equivalently, one requires analytic continuation of the inverse (*resolvent*) of this family of operators to the entire lower half-plane and continuous extension to the real line.

They also established results for *almost* stationary operators, though that is not the context of the work presented here. While the authors employed a nontrapping hypothesis, their work did not require product structure on their space-times, which makes their work highly influential in our own.

Although the absence of trapping is known to be necessary for waves to experience local energy decay (see [Ral69], [Sbi15]), one can recover weak local energy decay estimates with a prescribed loss at high frequencies for certain types of trapping (see [Bur98], [Chr08], [Ika82], [Ika88], [MMTT10], [NZ09], [Toh12], [WZ11], and the contained references). When the trapping is sufficiently weak/unstable, then this loss is nominal (in fact, logarithmic); this is the case for both the *Schwarzschild* [MMTT10] and *Kerr* [Toh12] space-times. Both space-times possess nontrivial trapped sets, which constitute regions where light remains for all time. Although one can extract weak local energy decay estimates, the trapping still generates an immutable barrier to full local energy decay. We will not be working in a scenario that generates loss, although we would be remiss if we did not briefly mention weak local energy decay and essential space-times that enjoy it.

The study of damped waves also possesses a deep history, especially on compact manifolds. The seminal work [RT74] introduced the *geometric control condition*,

which required that all null bicharacteristic rays intersect the damping region, and the authors used it to show that the energy of solutions to damped hyperbolic equations on compact product manifolds enjoys exponential decay in time. The uniform exponential bound is equivalent to so-called strong stabilization, whereby one can bound the energy at an arbitrary time by the initial energy multiplied by a monotone-decreasing, nonnegative function tending to zero as $t \rightarrow \infty$. This established the sufficiency of geometric control for strong stabilization in such settings, while [Ral69] demonstrated necessity (also, see [Leb96]). The work [BLR92] showed sufficiency for observability and control on compact manifolds with boundary where the observability/control region is contained within the boundary. While there is notably less literature in the noncompact setting, it was proven in [BR14] that local energy decay holds for the damped wave equation on asymptotically *Euclidean* space-times with time-independent metrics under the assumption of geometric control on trapped geodesics. The authors proved dissipative Mourre estimates to obtain uniform resolvent bounds in different frequency regimes in order to apply a limiting absorption argument. This approach is highly dependent on the metric coefficients being independent of time and the product structure (asymptotically Euclidean metrics contain no metric cross terms). This result was improved in [Roy18] to estimates in the (weighted) energy space.

In this article, we combine the approaches of [BR14] and [MST20] to establish high frequency local energy estimates for damped waves on stationary, asymptotically flat space-times and explain how such a result can be readily combined with the existing work in [MST20] to prove local energy decay. We underscore that we are not requiring the product structure evident in [BR14] nor [Roy18] but, instead, allow for the full Lorentzian formulation. Nonproduct metrics possess nontrivial cross terms and are called *nonstatic*, of which the Kerr metric constitutes an important example. We most closely keep to the framework present in [MST20], which does not assume product structure and has results for even more general asymptotically flat nontrapping space-times (such as nonstationary ones). We again stress their use of a nontrapping hypothesis, which we replace by imposing geometric control. Trapping is an intrinsically high frequency phenomenon, so only their high frequency work is affected by the trapping. Hence, this is the portion of the argument that needs modification to ensure local energy decay, and this is where the influence of [BR14] comes into play. Since the medium and low frequency analyses (as well as the procedure of combining the different frequency regime estimates into the full local energy decay estimate) do not depend on the nontrapping hypothesis nor use the damping themselves, the corresponding results in [MST20] readily apply (i.e., our problem essentially becomes a special case here). We omit the details of such results in this work, but we will explain why they apply in our context.

1.2. Problem setup and main results. Let (\mathbb{R}^4, g) be a Lorentzian manifold with coordinates $(t, x) \in \mathbb{R} \times \mathbb{R}^3$, where g has signature $(- + + +)$. We will consider *damped wave operators* of the form

$$P = \square_g + iaD_t, \quad \square_g = D_\alpha g^{\alpha\beta} D_\beta,$$

where $a \in C_c^\infty(\mathbb{R}^3)$ is nonnegative and positive on an open set, and $D_\alpha = \frac{1}{i}\partial_\alpha$, $\alpha = 0, 1, 2, 3$. Greek indices will generally range over such values, whereas Latin indices will run over the integers 1, 2, and 3. Notice that we are using the standard Einstein summation convention, which we will do throughout this work. We will also subject g to an *asymptotic flatness* condition. More precisely, we first define the norm

$$\|h\|_{AF} = \sum_{|\alpha| \leq 2} \left\| \langle x \rangle^{|\alpha|} \partial^\alpha h \right\|_{\ell_j^1 L^\infty([0, T] \times A_j)},$$

where $A_j = \{\langle x \rangle \approx 2^j\}$ for $j \geq 0$ denote inhomogeneous dyadic regions, and ℓ_j^1 denotes the ℓ^1 norm over the j index. The notation $A \lesssim B$ means that $A \leq CB$ for some $C > 0$, and the notation $A \approx B$ means that $B \lesssim A \lesssim B$. In the definition of the A_j 's, we require that these implicit constants are compatible to cover \mathbb{R}^3 . This allows us to define the AF topology.

DEFINITION 1.1. *We say that P is asymptotically flat if $\|g - m\|_{AF} < \infty$, where m denotes the Minkowski metric, and*

$$\left\| \langle x \rangle^{|\alpha|} \partial^\alpha g \right\|_{\ell_j^1 L^\infty([0, T] \times A_j)} \lesssim_\alpha 1$$

for all $\alpha \in \mathbb{N}^3$ with $|\alpha| \geq 3$.

The latter condition will be necessary for certain functions appearing in this work to be symbolic in the Kohn–Nirenberg sense. We remark that the dyadic summability assumptions on our metric are weaker than the long-range perturbation condition present in [BR14] (which provides a symbolic-type decay estimate for derivatives of the metric in x in terms of $\langle x \rangle^{-\rho}$ with $\rho > 0$ fixed). In [BR14], the damping is not assumed to be compactly supported, but rather nonnegative everywhere and subject to a similar symbolic estimate to the metric (with an additional power of decay). Since the damping is a *helpful* term which will only be necessary within a compact spatial set (this is made explicit with the introduction of the parameter R_0 on the next page), it is unnecessarily beneficial for us to assume that it is nonnegative everywhere.

We will primarily be interested in when ∂_t is a Killing field for g , in which case we say that P is *stationary*.

DEFINITION 1.2. *We say that P is stationary if g is independent of t .*

Next, we introduce the following:

- The parameters R_0 and \mathbf{c} , which are such that

$$\|g - m\|_{AF_{>R_0}} \leq \mathbf{c} \ll 1,$$

where the subscript denotes the restriction of the norm to $\{|x| > R_0\}$. The parameter \mathbf{c} should be viewed as being fixed first, after which we find an R_0 for which the above holds. Without loss of generality, we will assume that $\text{supp } a \subset \{|x| \leq R_0\}$ (as it is unnecessarily beneficial outside of this set).

- The sequence $(c_j)_{j \geq \log_2 R_0}$ satisfying

$$\|g - m\|_{AF(A_j)} \lesssim c_j, \quad \sum_j c_j \lesssim \mathbf{c},$$

where $\|\cdot\|_{AF(A_j)}$ denotes the restriction of the AF norm to the dyadic region A_j . We may assume, without any loss of generality, that the sequence is slowly varying, i.e.,

$$c_j/c_k \leq 2^{\delta|k-j|}, \quad \delta \ll 1.$$

This sequence will be utilized when working in spatial weights within dyadic regions.

These parameters tell us that, outside of a large enough spatial ball, the operator P is a uniformly small perturbation of the flat wave operator $\square_m = \partial_t^2 - \Delta$ (which we simply denote as \square). The sequence (c_j) provides a quantitative measure on the size of the AF norm throughout each spatial dyadic region outside of this ball.

We will also assume throughout that the vector field ∂_t is uniformly time-like, which essentially constitutes a choice of coordinates. This condition, coupled with the signature of the metric, ensures that $D_i g^{ij} D_j$ is uniformly elliptic, i.e.,

$$(1.2) \quad g^{ij} \xi_i \xi_j \approx |\xi|^2, \quad \xi \neq 0,$$

where $|\cdot|$ denotes the standard Euclidean norm. This follows from the positive-definiteness of the momentum-energy tensor when applied to time-like vector fields.

Next, we define the local energy norms

$$\begin{aligned} \|u\|_{LE} &= \sup_{j \geq 0} \left\| \langle x \rangle^{-1/2} u \right\|_{L_t^2 L_x^2(\mathbb{R}_+ \times A_j)}, \\ \|u\|_{LE^1} &= \|\partial u\|_{LE} + \left\| \langle x \rangle^{-1} u \right\|_{LE}. \end{aligned}$$

A predual-type norm to the LE norm is the LE^* norm, which is defined as

$$\|f\|_{LE^*} = \sum_{j=0}^{\infty} \left\| \langle x \rangle^{1/2} f \right\|_{L_t^2 L_x^2(\mathbb{R}_+ \times A_j)}.$$

Here, $L_t^p L_x^q$ denotes the Bochner space $L^p(\mathbb{R}_+, L^q(\mathbb{R}^3))$. In the particular case of $p, q = 2$, then this is a Hilbert space; we will use $\langle \cdot, \cdot \rangle$ to denote the inner product on $L_t^2 L_x^2$. Last, we define the sum-space norm

$$\|f\|_{LE^* + L_t^1 L_x^2} = \inf_{f=f_1+f_2} \left(\|f_1\|_{LE^*} + \|f_2\|_{L_t^1 L_x^2} \right).$$

If we wish for the time interval to be, e.g., $[0, T]$ in the above norms, then we will use the notation $\|u\|_{LE[0, T]}$, $\|u\|_{LE^1[0, T]}$, $\|u\|_{LE^*[0, T]}$, $\|u\|_{LE^* + L_t^1 L_x^2[0, T]}$ (although we will write $LE^*[0, T] + L_t^1 L_x^2[0, T]$ when referring to this space outside of norm subscripts), etc. A subscript of c on any of these spaces denotes compact spatial support.

There are two additional function spaces that will be utilized extensively in this work. The first is the class of Schwartz functions $\mathcal{S}(\mathbb{R}^4)$, which will be useful for approximation arguments. The second is a particular collection of functions which is often the natural class to study wave equations.

DEFINITION 1.3. *Let $T > 0$. We define the class \mathcal{W}_T to be the space of all functions $u \in C^2([0, T] \times \mathbb{R}^3)$ for which there exists $R > 0$ so that $u(t, x) = 0$ for all $t \in [0, T]$ and $|x| > R$. That is,*

$$\mathcal{W}_T = \{u \in C^2([0, T] \times \mathbb{R}^3) : (\exists R > 0)(\forall |x| > R)(\forall t \in [0, T]) \quad u(t, x) = 0\}.$$

We are interested in Cauchy problems of the form

$$\begin{cases} Pu = f \in LE^*[0, T] + L_t^1 L_x^2[0, T], \\ u[0] = (u(0), \partial_t u(0)) \in \dot{H}^1 \oplus L^2. \end{cases}$$

Remark 1.4. The decay conditions on $u \in \mathcal{W}_T$ are not as restrictive as they might initially appear. If the Cauchy data is compactly supported, then the condition is free

by finite speed of propagation. If it is not, then one can approximate the data (which generically lives in the energy space) by compactly supported data. The regularity conditions on u are also not restrictive, as one can perform density arguments to reduce to the case of increased regularity.

Now, we state the pertinent local energy estimates for such problems.

DEFINITION 1.5. *We say that (integrated) local energy decay holds for an asymptotically flat wave operator if the following estimate holds for all $T > 0$:*

$$(1.3) \quad \|u\|_{LE^1[0,T]} + \|\partial u\|_{L_t^\infty L_x^2[0,T]} \lesssim \|\partial u(0)\|_{L^2} + \|Pu\|_{LE^* + L_t^1 L_x^2[0,T]}$$

for all $u \in \mathcal{W}_T$ such that $u[0] \in \dot{H}^1 \oplus L^2$, with the implicit constant being independent of T .

The notion of an asymptotically flat wave operator is more broad than an asymptotically flat damped wave operator. They need not feature a damping term, and they are allowed to possess general lower-order terms which are asymptotically flat in an appropriate sense (see Definition 1.1 in [MST20] for a precise definition).

Note that, due to global energy conservation for the flat wave problem, the general definition of local energy decay that we have given here is consistent with the integrated local energy estimate for the flat wave equation in (1.1) (in the inhomogeneous case, one applies Hölder's inequality to the forcing). This estimate is known to hold whenever P is a small asymptotically flat perturbation of \square (see [Ali06], [MS06], [MS07], [MT12]). In [MST20], the authors considered large *AF* perturbations and proved that, for stationary problems, the local energy decay estimate (1.3) is equivalent to assuming that the wave operator P is nontrapping and has no negative eigenfunctions (L^2 eigenfunctions with corresponding eigenvalues in the lower half-plane) nor real resonant states (outgoing non- L^2 eigenfunctions with real “eigenvalues,” which are called resonances); see Definitions 2.2, 2.4, and 2.8 in [MST20] for more precise statements. The nontrapping hypothesis only arose during their proof of a high frequency estimate (Theorem 2.11 in [MST20]), which took the form

$$(1.4) \quad \|u\|_{LE^1[0,T]} + \|\partial u\|_{L_t^\infty L_x^2[0,T]} \lesssim \|\partial u(0)\|_{L^2} + \left\| \langle x \rangle^{-2} u \right\|_{LE[0,T]} + \|Pu\|_{LE^* + L_t^1 L_x^2[0,T]}.$$

The implicit constant in the above estimate is crucially independent of T . This estimate does not require u to be truncated to large time frequencies, but this is the context in which it is used in proving local energy decay.

The added spatial weight in the error term does not play a particular role in making this high frequency. Rather, it is the weight that naturally arises when performing a bootstrapping argument in the proof of the estimate; it is largely unimportant since this estimate can be reduced to studying solutions with compact spatial support (see section 2.5).

Remark 1.6. To see this as an estimate on the high frequencies, let $u \in \mathcal{S}(\mathbb{R}^4)$ be frequency-supported in time for τ in the range $1 \ll \tau_1 \leq |\tau| < \infty$ (1.4) is only applied for such u in the proof of local energy decay). Then, we can use Plancherel's theorem in t to obtain that

$$(1.5) \quad \left\| \langle x \rangle^{-2} u \right\|_{LE_{t,x}} \approx \left\| \langle x \rangle^{-2} \hat{u}(\tau, x) \right\|_{LE_{\tau,x}} \lesssim \frac{1}{\tau_1} \left\| \langle x \rangle^{-2} \tau \hat{u}(\tau, x) \right\|_{LE_{\tau,x}} \lesssim \frac{1}{\tau_1} \|u\|_{LE_{t,x}^1}.$$

For large enough τ_1 , this term can be absorbed into the left-hand side of (1.4), providing local energy decay for solutions restricted to high frequencies.

In fact, we may apply the high frequency estimate (1.4) to $u(t-T/2)$ to get (after dropping the uniform energy piece) that

$$\|u\|_{LE^1[-T/2, T/2]} \lesssim \|\partial u(-T/2)\|_{L^2} + \left\| \langle x \rangle^{-2} u \right\|_{LE[-T/2, T/2]} + \|Pu\|_{LE^*[-T/2, T/2]}.$$

Since the implicit constant is independent of T , we may take the limit as $T \rightarrow \infty$ and apply the prior work in (1.5) to obtain that

$$\|u\|_{LE^1} \lesssim \|Pu\|_{LE^*}.$$

This is the context that we will apply the estimate to establish local energy decay.

Our first main theorem is the following, which states that we recover the high frequency estimate (1.4) of [MST20] when working with damped waves and replacing the nontrapping hypothesis with the geometric control condition.

THEOREM 1.7. *Let P be a stationary, asymptotically flat damped wave operator satisfying the geometric control condition, and suppose that ∂_t is uniformly time-like. Then, the high frequency local energy estimate (1.4) holds for all $u \in \mathcal{W}_T$ such that $u[0] \in \dot{H}^1 \oplus L^2$. The implicit constant is independent of T .*

The geometric control condition, initially introduced in [RT74] for dissipative hyperbolic equations on compact product manifolds, requires that every trapped null bicharacteristic ray intersect the damping region. We will make this more precise in section 2.2.

Remark 1.8. The implicit constant in the bound depends on R_0 . In fact, much of our work will implicitly depend on R_0 due to our applications of asymptotic flatness. It is essential to note that this parameter is fixed second (with \mathbf{c} being fixed first), after which our other parameters (such as the scaling parameter γ and the high frequency parameter λ , which will both be introduced in section 3) will be chosen (and hence depend on it). We will not track the dependence on R_0 within our implicit constants any longer. Our constants throughout will *not* depend on T , however.

Our second main theorem is local energy decay.

THEOREM 1.9. *Let P be a stationary, asymptotically flat damped wave operator satisfying the geometric control condition, and suppose that ∂_t is uniformly time-like. Then, local energy decay holds, with the implicit constant in (1.3) independent of T .*

Remark 1.10. Unlike [MST20], we do not require any spectral hypotheses. The damping eliminates the possibility of nonzero real resonances, and a zero resonance cannot occur since $P|_{D_t=0}$ is elliptic, which is independent of the damping. The latter is discussed further in section 3.2.

This follows rather directly from our high frequency estimate and the existing work in [MST20]. We will cite the necessary result from [MST20] and explain why they apply here.

The structure of the paper is as follows. In section 2.2, we will introduce the Hamiltonian formalism required to define trapping and geometric control, and then we will state a key lemma (Lemma 2.4) for the proof of Theorem 1.7; namely, we construct an appropriate escape function and lower-order correction to allow for a positive commutator argument proof of the theorem. In section 2.3, we establish various

results on the bicharacteristic flow that are vital for proving Lemma 2.4, which we establish in section 2.4. In section 2.5, we will establish and cite supplemental energy estimates and provide multiple case reductions to simplify the proof of Theorem 1.7. In section 2.6, we will prove Theorem 1.7. Finally, section 3 will present the applicable theorems from [MST20] which are required to establish local energy decay for our problem and discuss why they apply in our setting. Sections 3.1 and 3.2 introduce the relevant medium and low frequency estimates, respectively, and section 3.3 provides a discussion on the proof of Theorem 1.9.

1.3. Cutoff notation. For the remainder of paper, we will fix the cutoffs

$$\chi \in C_c^\infty \text{ nonincreasing, } \chi \equiv 1 \text{ for } |x| \leq 1, \chi \equiv 0 \text{ for } |x| > 2,$$

$$\chi_{$$

and

$$\chi_R \in C_c^\infty, \quad 0 \leq \chi_R \leq 1, \quad \text{supp } \chi_R \subset \{|x| \approx R\}.$$

We will choose χ such that its square root is smooth (otherwise, we replace χ with χ^2 ; we only use χ for notational convenience). When working in frequency variables, we will often add the variable into the subscript to make the dependence clear (e.g., $\chi_{|\xi|>\lambda}$). We will also occasionally write $r = |x|$.

2. High frequency analysis.

2.1. Introduction. In this section, we will establish Theorem 1.7. The notions of trapping and geometric control are intrinsically dynamical, so we will provide a thorough discussion of the relevant theory. Namely, we must introduce the bicharacteristic flow generated by the principal symbol of the damped wave operator and the properties that it satisfies. From here, we will construct an escape function and correction term in order to apply a utilize commutator argument and prove the theorem. The constructed symbols will satisfy an appropriate positivity bound, which will allow us to apply the sharp Gårding inequality upon swapping to the framework of pseudodifferential operators. The symbols will also be supported in an unbounded range of frequencies $[\lambda, \infty)$ with $\lambda \gg 1$. This will be fundamentally important for bootstrapping error terms resulting from employment of pseudodifferential calculus.

2.2. Dynamical framework. In order to state the geometric control condition more precisely, we must first outline our dynamical framework, which is rooted in the Hamiltonian dynamics pertaining to the principal symbol of the operator P . Since we assumed that ∂_t is uniformly time-like, the signature of the metric and the cofactor expansion for the inverse metric tell us that $g^{00} \lesssim -1$, as well. This allows us to divide through by $-g^{00}$ and preserve the assumptions on the operator coefficients (see [MT12]). Hence, we may assume (without loss of generality) that $g^{00} = -1$.

After these modifications, the principal symbol of P is

$$p(\tau, x, \xi) = -(\tau^2 - 2\tau g^{0j}(x)\xi_j - g^{ij}(x)\xi_i\xi_j).$$

This is considered as a smooth function on $T^*\mathbb{R}^4 \setminus o$, where o denotes the zero section. This symbol generates a bicharacteristic/Hamiltonian flow on $\mathbb{R} \times T^*\mathbb{R}^4$ given by $\varphi_s(w) = (t_s(w), \tau_s(w), x_s(w), \xi_s(w))$ which solves

$$\begin{cases} \dot{t}_s = \partial_\tau p(\varphi_s(w)), & \dot{\tau}_s = -\partial_t p(\varphi_s(w)), \\ \dot{x}_s = \nabla_\xi p(\varphi_s(w)), & \dot{\xi}_s = -\nabla_x p(\varphi_s(w)), \end{cases}$$

with initial data $w \in T^*\mathbb{R}^4$. Explicitly, one can write the system as

$$\begin{cases} \dot{t}_s = -2\tau_s + 2g^{0j}(x_s)[\xi_s]_j, & \dot{\tau}_s = 0, \\ (\dot{x}_s)_k = 2\tau_s g^{0k}(x_s) + 2g^{kj}(x_s)[\xi_s]_j, & (\dot{\xi}_s)_k = -2\tau_s \partial_{x_k} g^{0j}(x_s)[\xi_s]_j - \partial_{x_k} g^{ij}(x_s)[\xi_s]_i [\xi_s]_j. \end{cases}$$

Since g is smooth and asymptotically flat, and ∂_t is uniformly time-like, we have a unique, smooth, globally defined flow with smooth dependence on the data. We will have particular interest in *null* bicharacteristics, i.e., those with initial data lying in the zero set of p (also called the *characteristic set* of P and denoted $\text{Char}(P)$). Using the flow φ_s , we define the *forward* and *backward trapped* and *nontrapped* sets with respect to φ_s , respectively, as

$$\begin{aligned} \Gamma_{tr} &= \left\{ w \in T^*\mathbb{R}^4 \setminus o : \sup_{s \geq 0} |x_s(w)| < \infty \right\} \cap \text{Char}(P), \\ \Lambda_{tr} &= \left\{ w \in T^*\mathbb{R}^4 \setminus o : \sup_{s \geq 0} |x_{-s}(w)| < \infty \right\} \cap \text{Char}(P), \\ \Gamma_\infty &= \left\{ w \in T^*\mathbb{R}^4 \setminus o : |x_s(w)| \rightarrow \infty \text{ as } s \rightarrow \infty \right\} \cap \text{Char}(P), \\ \Lambda_\infty &= \left\{ w \in T^*\mathbb{R}^4 \setminus o : |x_{-s}(w)| \rightarrow \infty \text{ as } s \rightarrow \infty \right\} \cap \text{Char}(P). \end{aligned}$$

The *trapped* and *nontrapped* sets are defined as

$$\Omega_{tr}^p = \Gamma_{tr} \cap \Lambda_{tr} \quad \text{and} \quad \Omega_\infty^p = \Gamma_\infty \cap \Lambda_\infty,$$

respectively.

DEFINITION 2.1. *The flow is said to be nontrapping if $\Omega_{tr}^p = \emptyset$.*

Now, we may state the geometric control condition precisely. Recall that our damping function was denoted a .

DEFINITION 2.2. *We say that geometric control holds if*

$$(2.1) \quad (\forall w \in \Omega_{tr}^p) (\exists s \in \mathbb{R}) \quad a(x_s(w)) > 0.$$

In contrast to the definition given in Assumption (A) in [RT74], we apply this condition specifically to the trapped null bicharacteristic rays (since *all* null bicharacteristic rays are trapped when the manifold is compact, such a specification was unnecessary in [RT74]). We will assume that (2.1) holds. Note that if (2.1) holds and $a \equiv 0$, then Ω_{tr}^p must be empty, meaning that the flow is nontrapping. In this case, we are back in the setting of [MST20]. For this reason, we will assume that $a > 0$ on an open set.

It will be beneficial to utilize a scaling property of P . Given a solution u to $Pu = f$, consider

$$\tilde{v}(t, x) := \gamma^{-2} u(\gamma t, \gamma x), \quad \gamma > 0.$$

If we call

$$\tilde{P} = D_\alpha \tilde{g}^{\alpha\beta} D_\beta + i\gamma \tilde{a} D_t, \quad \tilde{g}^{\alpha\beta}(x) = g^{\alpha\beta}(\gamma x), \quad \tilde{a}(x) = a(\gamma x),$$

then \tilde{v} solves

$$\tilde{P}\tilde{v} = \tilde{f}, \quad \tilde{f}(t, x) = f(\gamma t, \gamma x)$$

if and only if u solves $Pu = f$ (we can similarly undo the scaling to move between the frameworks). Notice that the scaled problem allows for an arbitrarily large constant γ in front of the damping.

Analogous Hamiltonian systems and trapped sets exist for the principal symbol \tilde{p} of \tilde{P} , and this amounts to simply replacing g by \tilde{g} . If we assume that geometric control holds for the flow generated by p , then we must check that it holds for the scaled problem.

PROPOSITION 2.3. *Assume that (2.1) holds. Then, for any $\gamma > 0$, (2.1) holds for the flow generated by \tilde{p} , with a replaced by \tilde{a} .*

Note that since $g^{00} \equiv -1$, it follows that $\tilde{g}^{00} \equiv -1$.

Proof. The flow generated by \tilde{p} solves the system

$$\left\{ \begin{array}{l} \frac{d}{ds} \tilde{t}_s = -2\tilde{\tau}_s + 2\tilde{g}^{0j}(\tilde{x}_s)[\tilde{\xi}_s]_j, \\ \frac{d}{ds} \tilde{\tau}_s = 0, \\ \frac{d}{ds} (\tilde{x}_s)_k = 2\tilde{\tau}_s \tilde{g}^{0k}(\tilde{x}_s) + 2\tilde{g}^{kj}(\tilde{x}_s)[\tilde{\xi}_s]_j, \\ \frac{d}{ds} (\tilde{\xi}_s)_k = -2\tilde{\tau}_s \partial_{x_k} \tilde{g}^{0j}(\tilde{x}_s)[\tilde{\xi}_s]_j - \partial_{x_k} \tilde{g}^{ij}(\tilde{x}_s)[\tilde{\xi}_s]_i [\tilde{\xi}_s]_j, \\ (\tilde{t}_s, \tilde{\tau}_s, \tilde{x}_s, \tilde{\xi}_s) \big|_{s=0} = (t, \tau, x, \xi). \end{array} \right.$$

Applying the chain rule and multiplying through by γ provides us with the system

$$\left\{ \begin{array}{l} \frac{d}{ds} (\gamma \tilde{t}_s) = -2(\gamma \tilde{\tau}_s) + 2g^{0j}(\gamma \tilde{x}_s)[\gamma \tilde{\xi}_s]_j, \\ \frac{d}{ds} (\gamma \tilde{\tau}_s) = 0, \\ \frac{d}{ds} (\gamma \tilde{x}_s)_k = 2(\gamma \tilde{\tau}_s) g^{0k}(\gamma \tilde{x}_s) + 2g^{kj}(\gamma \tilde{x}_s)[\gamma \tilde{\xi}_s]_j, \\ \frac{d}{ds} (\gamma \tilde{\xi}_s)_k = -2(\gamma \tilde{\tau}_s) [(\partial_{x_k} g^{0j})(\gamma \tilde{x}_s)] [\gamma \tilde{\xi}_s]_j - [(\partial_{x_k} g^{ij})(\gamma \tilde{x}_s)] [\gamma \tilde{\xi}_s]_i [\gamma \tilde{\xi}_s]_j, \\ ((\gamma \tilde{t}_s, (\gamma \tilde{\tau}_s, (\gamma \tilde{x}_s, (\gamma \tilde{\xi}_s) \big|_{s=0} = (\gamma t, \gamma \tau, \gamma x, \gamma \xi). \end{array} \right.$$

This is the same system that is solved by the Hamiltonian flow generated by p with initial data $(\gamma t, \gamma \tau, \gamma x, \gamma \xi)$. By uniqueness, we can conclude that

$$\left\{ \begin{array}{ll} \gamma \tilde{t}_s(t, \tau, x, \xi) = t_s(\gamma t, \gamma \tau, \gamma x, \gamma \xi), & \gamma \tilde{\tau}_s(t, \tau, x, \xi) = \tau_s(\gamma t, \gamma \tau, \gamma x, \gamma \xi), \\ \gamma \tilde{x}_s(t, \tau, x, \xi) = x_s(\gamma t, \gamma \tau, \gamma x, \gamma \xi), & \gamma \tilde{\xi}_s(t, \tau, x, \xi) = \xi_s(\gamma t, \gamma \tau, \gamma x, \gamma \xi). \end{array} \right.$$

Now, let $w = \Omega_{tr}^{\tilde{p}}$. From the above, we have that

$$\tilde{x}_s(w) = \gamma^{-1} x_s(\tilde{w}), \quad \tilde{w} = \gamma w.$$

Since

$$\sup_{s \in \mathbb{R}} |x_s(\tilde{w})| = \gamma \sup_{s \in \mathbb{R}} |\tilde{x}_s(w)| < \infty,$$

it follows that $\tilde{w} \in \Omega_{tr}^p$. By (2.1), there exists $s' \in \mathbb{R}$ so that $a(x_{s'}(\tilde{w})) > 0$, and so

$$\tilde{a}(\tilde{x}_{s'}(w)) = a(\gamma \tilde{x}_{s'}(w)) = a(x_{s'}(\tilde{w})) > 0,$$

which completes the proof. \square

Now that we have shown that geometric control is invariant under scaling, we will fix a large $\gamma > 0$ and study the problem from the scaled perspective (where our damping is now multiplied by γ) while reverting back to our original notation (x and ξ , no tildes, etc.). More precise conditions on the size of γ will come in section 2.4. It is readily seen that it is equivalent to prove Theorem 1.7 for the scaled problem, where we now have a large constant in front of the damping term.

Our proof of Theorem 1.7 is a positive commutator argument. At the symbolic level, this requires the construction of an escape function (as well as a lower-order correction). We must consider the skew-adjoint contribution of P , which will be a purely beneficial term due to the presence of the damping. Let p and s_{skew} represent the principal symbols of the self and skew-adjoint parts of P , respectively. Namely,

$$\begin{aligned} p(\tau, x, \xi) &= -(\tau^2 - 2\tau g^{0j}(x)\xi_j - g^{ij}(x)\xi_i\xi_j), \\ s_{skew}(\tau, x, \xi) &= i\gamma\tau a(x). \end{aligned}$$

The multiplication by γ in s_{skew} will prove advantageous for a bootstrapping argument, which is precisely why we implement the γ -scaling. Now, we are ready to state our escape function result, which we will prove in section 2.4.

LEMMA 2.4. *For all $\lambda > 1$, there exist symbols $q_j \in S^j(T^*\mathbb{R}^3)$ and $m \in S^0(T^*\mathbb{R}^3)$, all supported in $|\xi| \geq \lambda$, so that*

$$H_p q - 2is_{skew}q + pm \gtrsim \mathbb{1}_{|\xi| \geq \lambda} \langle x \rangle^{-2} (\tau^2 + |\xi|^2),$$

where $q = \tau q_0 + q_1$.

Here, $S^m(T^*\mathbb{R}^n)$ denotes the m th-order Kohn–Nirenberg symbol class. To each $q \in S^m(T^*\mathbb{R}^n)$, we will associate the pseudodifferential operator $q^w(x, D) : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$, namely the *Weyl quantization* of p , which is defined via the action

$$q^w(x, D)u(x) = (2\pi)^{-n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i(x-y)\cdot\xi} q\left(\frac{x+y}{2}, \xi\right) u(y) dy d\xi.$$

In the proof of Lemma 2.4, it will be useful to work with the half-wave decomposition, which allows us to avoid the cross terms in the principal symbol. To that end, we factor p as

$$p(\tau, x, \xi) = -(\tau - b^+(x, \xi))(\tau - b^-(x, \xi)),$$

where

$$b^\pm(x, \xi) = g^{0j}\xi_j \pm \sqrt{(g^{0j}\xi_j)^2 + g^{ij}\xi_i\xi_j}.$$

Observe that b^\pm are both homogeneous of degree 1 in ξ . Additionally, they are both signed.

PROPOSITION 2.5. *For any $(x, \xi) \in T^*\mathbb{R}^3 \setminus o$, we have that $b^+(x, \xi) > 0 > b^-(x, \xi)$.*

Proof. Let $\xi \neq 0$. First, we show that $b^+ > b^-$. Indeed, observe that

$$b^+ - b^- = 2\sqrt{(g^{0j}\xi_j)^2 + g^{ij}\xi_i\xi_j} > 0$$

using the ellipticity (see (1.2)). Using ellipticity again, we have that

$$\sqrt{(g^{0j}\xi_j)^2 + g^{ij}\xi_i\xi_j} > |g^{0j}\xi_j|.$$

Thus,

$$b^+ > g^{0j} \xi_j + |g^{0j} \xi_j| \geq 0, \quad b^- < g^{0j} \xi_j - |g^{0j} \xi_j| \leq 0. \quad \square$$

We will call $p^\pm = \tau - b^\pm$, so that $p = -p^+ p^-$. In particular, $p = 0$ if and only if $p^+ = 0$ or $p^- = 0$; due to Proposition 2.5, it cannot be the case that $p^+(w) = p^-(w) = 0$ for any $w \in T^* \mathbb{R}^4 \setminus o$. The Hamiltonians p^\pm also generate flows $\varphi_s^\pm(w) = (t_s^\pm(w), \tau_s^\pm(w), x_s^\pm(w), \xi_s^\pm(w))$ on $\mathbb{R} \times T^* \mathbb{R}^4$ which solve the Hamiltonian systems

$$\begin{cases} \dot{t}_s^\pm = \partial_\tau p^\pm(\varphi_s^\pm(w)), & \dot{\tau}_s^\pm = -\partial_t p^\pm(\varphi_s^\pm(w)), \\ \dot{x}_s^\pm = \nabla_\xi p^\pm(\varphi_s^\pm(w)), & \dot{\xi}_s^\pm = -\nabla_x p^\pm(\varphi_s^\pm(w)), \end{cases}$$

with initial data $w \in T^* \mathbb{R}^4$. Note that

$$\begin{cases} \dot{t}_s^\pm = 1, & \dot{\tau}_s^\pm = 0, \\ \dot{x}_s^\pm = -\nabla_\xi b^\pm(\varphi_s^\pm(w)), & \dot{\xi}_s^\pm = \nabla_x b^\pm(\varphi_s^\pm(w)). \end{cases}$$

There is a direct correspondence between null bicharacteristics for φ_s and null bicharacteristics for φ_s^\pm .

PROPOSITION 2.6. *Every null bicharacteristic for the flow generated by p is a null bicharacteristic for the flow generated by either p^+ or p^- . The converse is also true.*

Proof. Recall that for any $(t', \tau', x', \xi') =: w \in T^* \mathbb{R}^4 \setminus o$, we have that $p(w) = 0$ if and only if either $p^+(w) = 0$ or $p^-(w) = 0$. Without loss of generality, suppose that $p^+(w) = 0$. The Hamiltonians p and p^+ generate the systems

$$(2.2) \quad \begin{cases} \dot{t}_s = -p^+(\varphi_s) - p^-(\varphi_s), & \dot{\tau}_s = 0, \\ (\dot{x}_s)_k = -p^+(\varphi_s)p_{\xi_k}^-(\varphi_s) - p^-(\varphi_s)p_{\xi_k}^+(\varphi_s), & (\dot{\xi}_s)_k = p^+(\varphi_s)p_{x_k}^-(\varphi_s) + p^-(\varphi_s)p_{x_k}^+(\varphi_s), \end{cases}$$

and

$$(2.3) \quad \begin{cases} \dot{t}_s^+ = 1, & \dot{\tau}_s^+ = 0, \\ (\dot{x}_s^+)_k = p_{\xi_k}^+(\varphi_s^+), & (\dot{\xi}_s^+)_k = -p_{x_k}^+(\varphi_s^+), \end{cases}$$

respectively. We will take both systems to have initial data w .

We claim that since $p^+(w) = 0$, we must have that $p^+(\varphi_s(w)) = 0$ for all s . If not, then there would exist s' so that $p^-(\varphi_{s'}(w)) = 0$, i.e., $\tau_{s'}^-(w) = b^-(x_{s'}(w), \xi_{s'}(w)) < 0$. However, τ^- is constant and $p^+(w) = 0$, which implies that $\tau_s^-(w) = \tau' > 0$ for all s .

Thus, we can rewrite (2.2) as

$$(2.4) \quad \begin{cases} \dot{t}_s = -p^-(\varphi_s), & \dot{\tau}_s = 0, \\ (\dot{x}_s)_k = -p^-(\varphi_s)p_{\xi_k}^+(\varphi_s), & (\dot{\xi}_s)_k = p^-(\varphi_s)p_{x_k}^+(\varphi_s), \end{cases}$$

with initial data $(t_s, \tau_s, x_s, \xi_s)|_{s=0} = w$. Notice that $t^+ = t' + s$, and so we may reparameterize (2.3) in terms of t^+ :

$$\begin{cases} \frac{d}{dt^+} t_{t^+ - t'}^+ = 1, & \frac{d}{dt^+} \tau_{t^+ - t'}^+ = 0, \\ \left(\frac{d}{dt^+} x_{t^+ - t'}^+ \right)_k = p_{\xi_k}^+(\varphi_{t^+ - t'}^+), & \left(\frac{d}{dt^+} \xi_{t^+ - t'}^+ \right)_k = -p_{x_k}^+(\varphi_{t^+ - t'}^+), \end{cases}$$

with initial data $(t_{t^+ - t'}^+, \tau_{t^+ - t'}^+, x_{t^+ - t'}^+, \xi_{t^+ - t'}^+)|_{t^+ = t'} = w$.

Next, we reparameterize (2.4) to change the flow variable from s to t (which can be done since t_s is strictly increasing and hence invertible), generating the system

$$\begin{cases} \frac{d}{dt} t_{s(t)} = 1, & \frac{d}{dt} \tau_{s(t)} = 0, \\ \left(\frac{d}{dt} x_{s(t)} \right)_k = p_{\xi_k}^+ (\varphi_{s(t)}), & \left(\frac{d}{dt} \xi_{s(t)} \right)_k = -p_{x_k}^+ (\varphi_{s(t)}), \end{cases}$$

which has initial data $(t_{s(t)}, \tau_{s(t)}, x_{s(t)}, \xi_{s(t)})|_{t=t'} = w$. An application of uniqueness theory yields that $\varphi_{s(t)}(w) = \varphi_{t+ -t'}^+(w)$. The converse is similar by reversing the above process. \square

When working with the factored flow, the decoupling of (t, τ) and (x, ξ) allows us to project onto the (x, ξ) components of the flow without worrying about loss of information. For this reason, we will write $\Pi_{x, \xi} \circ \varphi^\pm$ as simply φ^\pm , where $\Pi_{x, \xi}(t, \tau, x, \xi) = (x, \xi)$. Notice that when we project, we are no longer looking at null bicharacteristics but, rather, bicharacteristics with initial data having nonzero ξ component.

Now, we may define all of the corresponding forward and backward trapped and nontrapped sets for the half-wave flows as

$$\begin{aligned} \Gamma_{tr}^\pm &= \left\{ w \in T^* \mathbb{R}^3 \setminus o : \sup_{s \geq 0} |x_s^\pm(w)| < \infty \right\}, \\ \Lambda_{tr}^\pm &= \left\{ w \in T^* \mathbb{R}^3 \setminus o : \sup_{s \geq 0} |x_{-s}^\pm(w)| < \infty \right\}, \\ \Gamma_\infty^\pm &= \left\{ w \in T^* \mathbb{R}^3 \setminus o : |x_s^\pm(w)| \rightarrow \infty \text{ as } s \rightarrow \infty \right\}, \\ \Lambda_\infty^\pm &= \left\{ w \in T^* \mathbb{R}^3 \setminus o : |x_{-s}^\pm(w)| \rightarrow \infty \text{ as } s \rightarrow \infty \right\}. \end{aligned}$$

The trapped and nontrapped sets are

$$\begin{aligned} \Omega_{tr}^\pm &= \Gamma_{tr}^\pm \cap \Lambda_{tr}^\pm \quad \text{and} \quad \Omega_\infty^\pm = \Gamma_\infty^\pm \cap \Lambda_\infty^\pm, \\ \Omega_{tr} &= \Omega_{tr}^+ \cup \Omega_{tr}^- \quad \text{and} \quad \Omega_\infty = \Omega_\infty^+ \cup \Omega_\infty^-. \end{aligned}$$

Note that the identities

$$\Omega_{tr} = \Pi_{x, \xi}(\Omega_{tr}^p) \quad \text{and} \quad \Omega_\infty = \Pi_{x, \xi}(\Omega_\infty^p)$$

hold as an immediate consequence of the factoring. Additionally, the factoring allows us to restate the geometric control condition as

$$(w \in \Omega_{tr}^+ \implies (\exists s \in \mathbb{R}) \ (a(x_s^+(w)) > 0)) \quad \text{and} \quad (w \in \Omega_{tr}^- \implies (\exists s \in \mathbb{R}) \ (a(x_s^-(w)) > 0)).$$

If $w \in \Omega_{tr}$, then it is trapped with respect to the flow generated by either p^+ or p^- by Proposition 2.6. If it is trapped with respect to p^+ , then there is a time so that w is flowed along a p^+ -bicharacteristic ray to a place where the damping is positive, and similarly if it is trapped with respect to p^- .

2.3. Results on the flow. Here, we establish results regarding the trapped/nontrapped sets and scalings for the flows, culminating in an extension of geometric control to bicharacteristic rays bounded either forward or backward in time. These results largely follow the path outlined in [BR14], although we require certain scaling results in order to utilize homogeneity arguments in later proofs (which were

unnecessary in [BR14] due to their use of semiclassical rescaling). In particular, Lemma 2.10 and Propositions 2.11 and 2.12 are analogous to results in Chapter 8 of [BR14] (namely, Lemma 8.2 and Propositions 8.3 and 8.4, respectively).

We will start with a scaling result on the flow.

PROPOSITION 2.7. *The flows generated by p^\pm satisfy the scalings*

$$\begin{cases} x_s^\pm(x, \xi) = x_s^\pm(x, \lambda\xi), \\ \lambda\xi_s^\pm(x, \xi) = \xi_s^\pm(x, \lambda\xi) \end{cases}$$

for any $\lambda > 0$.

Proof. Label the functions on the right-hand side as $x_{s,\lambda}^\pm$ and $\xi_{s,\lambda}^\pm$, respectively. Using the homogeneity of b^\pm , the left-hand side $(x_s^\pm, \lambda\xi_s^\pm)$ solves the system

$$\begin{cases} \frac{d}{ds}x_s^\pm = \nabla_\xi p^\pm(x_s^\pm, \xi_s^\pm) = \nabla_\xi p^\pm(x_s^\pm, \lambda\xi_s^\pm), \\ \frac{d}{ds}(\lambda\xi_s^\pm) = -\lambda\nabla_x p^\pm(x_s^\pm, \xi_s^\pm) = -\nabla_x p^\pm(x_s^\pm, \lambda\xi_s^\pm), \\ (x_s^\pm, \lambda\xi_s^\pm)|_{s=0} = (x, \lambda\xi), \end{cases}$$

while the right-hand side solves

$$\begin{cases} \frac{d}{ds}x_{s,\lambda}^\pm = \nabla_\xi p^\pm(x_{s,\lambda}^\pm, \xi_{s,\lambda}^\pm), \\ \frac{d}{ds}(\xi_{s,\lambda}^\pm) = -\nabla_x p^\pm(x_{s,\lambda}^\pm, \xi_{s,\lambda}^\pm), \\ (x_{s,\lambda}^\pm, \xi_{s,\lambda}^\pm)|_{s=0} = (x, \lambda\xi). \end{cases}$$

Applying uniqueness theory completes the proof. \square

This scaling implies that the trapped/nontrapped sets, and hence geometric control, are entirely determined by unit speed null bicharacteristics, i.e., by what happens on the unit cosphere bundle $S^*\mathbb{R}^3 = \{(x, \xi) \in T^*\mathbb{R}^3 : |\xi| = 1\}$. Indeed, observe that

$$x_s^\pm(x, \xi) = x_s^\pm(x, \xi/|\xi|).$$

The forward/backward trapped sets are defined in terms of supremums of the above over s , while the forward/backward nontrapped sets are defined via limits in s , and the prior equation shows that all of these are unaffected by the scaling in the ξ component of the initial data. A more pertinent scaling is given by the function

$$\Phi^\pm(x, \xi) = \left(x, \frac{\xi}{|b^\pm(x, \xi)|} \right).$$

The utility of this scaling comes from the fact that b^\pm is a constant of motion under the corresponding projected Hamiltonian flows and that

$$\left| \frac{\xi}{b^\pm(x, \xi)} \right| \approx 1,$$

which we now prove.

PROPOSITION 2.8. *For any $(x, \xi) \in T^*\mathbb{R}^3 \setminus o$,*

$$\left| \frac{\xi}{b^\pm(x, \xi)} \right| \approx 1.$$

Proof. By homogeneity,

$$\left| \frac{\xi}{b^\pm(x, \xi)} \right| = \frac{1}{\left| b^\pm \left(x, \frac{\xi}{|\xi|} \right) \right|}.$$

Write

$$\left| b^\pm \left(x, \frac{\xi}{|\xi|} \right) \right| = \left| g^{0j} \frac{\xi_j}{|\xi|} \pm \sqrt{\left(g^{0j} \frac{\xi_j}{|\xi|} \right)^2 + (g^{ij} - m^{ij}) \frac{\xi_i}{|\xi|} \frac{\xi_j}{|\xi|} + m^{ij} \frac{\xi_i}{|\xi|} \frac{\xi_j}{|\xi|}} \right|.$$

Since $\|g - m\|_{AF(|x| > R_0)} \ll 1$, asymptotic flatness guarantees that g^{0j} and $g^{ij} - m^{ij}$ are small in the exterior region $\{|x| > R_0\}$. Hence,

$$\left| b^\pm \left(x, \frac{\xi}{|\xi|} \right) \right| \approx \sqrt{m^{ij} \frac{\xi_i}{|\xi|} \frac{\xi_j}{|\xi|}} = 1$$

when $|x| > R_0$.

In the interior region, we are considering b^\pm on the compact set $\{|x| \leq R_0\} \times \{|\xi| = 1\}$. Since we know that $|b^\pm| > 0$ for all $\xi \neq 0$ from Proposition 2.5, continuity guarantees the desired boundedness here. \square

In view of Proposition 2.8, it follows that

$$\frac{\xi_j}{b^\pm(x, \xi)} \in S_{\text{hom}}^0(T^*\mathbb{R}^3 \setminus o), \quad j = 1, 2, 3,$$

where $S_{\text{hom}}^0(T^*\mathbb{R}^3 \setminus o)$ denotes the 0th-order homogeneous symbol class.

Remark 2.9. As a consequence of the scaling, the sets

$$\dot{\Gamma}_{tr}^\pm = \Gamma_{tr}^\pm \cap \Phi^\pm(T^*\mathbb{R}^3 \setminus o) \quad \text{and} \quad \dot{\Lambda}_{tr}^\pm = \Lambda_{tr}^\pm \cap \Phi^\pm(T^*\mathbb{R}^3 \setminus o)$$

are invariant under the flow. Indeed, it is readily seen that the (semi) trapped nature is preserved. Further, since b^\pm is constant along the flow, it follows from Proposition 2.7 that

$$\xi_s^\pm \left(x, \frac{\xi}{|b^\pm(x, \xi)|} \right) = \frac{1}{|b^\pm(x, \xi)|} \xi_s^\pm(x, \xi) = \frac{1}{|b^\pm(x_s^\pm(x, \xi), \xi_s^\pm(x, \xi))|} \xi_s^\pm(x, \xi).$$

Now, we prove a key result on nontrapped trajectories.

LEMMA 2.10. *If $R \geq R_0$ and*

$$|x_{\pm s'}^\pm(x, \xi)| \geq \max\{2R, |x| + \delta\}$$

for some $(x, \xi) \in T^\mathbb{R}^3 \setminus o$, $\delta > 0$, and $s' > 0$, then it holds for all $s \geq s'$, and*

$$|x_{\pm s}^\pm(x, \xi)| \rightarrow \infty$$

as $s \rightarrow \infty$.

That is, if we can get sufficiently far away from the origin and move radially outward from the initial position, then the trajectories are necessarily nontrapped. This can be proven directly, but the computations are simpler if one uses the correspondence between null bicharacteristics for p and p^\pm .

Proof. Without loss of generality, we will work with the x^+ bicharacteristic ray. By Proposition 2.6, it suffices to prove the result for the null bicharacteristic ray $x_{\pm s}$ with initial data \tilde{w} , where \tilde{w} is the lift of w to $T^*\mathbb{R}^4 \setminus o$ which is consistent with the comment immediately following the aforementioned proposition (in particular, the τ component is strictly positive). For any $z \in T^*\mathbb{R}^4 \setminus o$, we explicitly calculate that

$$\frac{1}{2} \frac{\partial^2}{\partial s^2} |x_{\pm s}(z)|^2 = \left| \frac{\partial}{\partial s} x_{\pm s}(z) \right|^2 + x_{\pm s}(z) \cdot \frac{\partial^2}{\partial s^2} x_{\pm s}(z),$$

where

$$\begin{aligned} \left| \frac{\partial}{\partial s} x_{\pm s}(z) \right|^2 &= 4\tau_{\pm s}^2(z) \left(\sum_{k=1}^3 g^{0k}(x_{\pm s}(z)) \right)^2 \\ &\quad + 4 \sum_{k=1}^3 [(g^{ki}(x_{\pm s}(z))(\xi_{\pm s}(z))_i)][(g^{kj}(x_{\pm s}(z))(\xi_{\pm s}(z))_j)] \\ &\quad + 8 \sum_{k=1}^3 \tau_{\pm s}(z) g^{0k}(x_{\pm s}(z)) g^{kj}(x_{\pm s}(z)) (\xi_j(z))_{\pm s}, \end{aligned}$$

and

$$\begin{aligned} x_{\pm s}(z) \cdot \frac{\partial^2}{\partial s^2} x_{\pm s}(z) &= 4\tau_{\pm s}(z)(x_{\pm s}(z))_k [\partial_\ell g^{0k}(x_{\pm s}(z))] (\tau_{\pm s}(z)g^{0\ell}(x_{\pm s}(z)) + g^{\ell j}(x_{\pm s}(z))(\xi_{\pm s}(z))_j) \\ &\quad + 4(x_{\pm s}(z))_j [\partial_\ell g^{kj}(x_{\pm s}(z))] (\tau_{\pm s}(z)g^{0\ell}(x_{\pm s}(z)) + g^{\ell j}(x_{\pm s}(z))(\xi_{\pm s}(z))_j) (\xi_{\pm s}(z))_j \\ &\quad - 2(x_{\pm s}(z))_k g^{kj}(x_{\pm s}(z)) (2\tau_{\pm s}(z) \partial_j g^{0i}(x_{\pm s}(z))(\xi_{\pm s}(z))_i \\ &\quad + \partial_j g^{i\ell}(x_{\pm s}(z))(\xi_{\pm s}(z))_i (\xi_{\pm s}(z))_\ell). \end{aligned}$$

Since τ_s is constant for stationary P , it follows that

$$\tau_{\pm s}(z) = \tau_0 = b^+(\Pi_{x,\xi}(z)) = b^\pm(x_{\pm s}^+(\Pi_{x,\xi}(z)), \xi_{\pm s}^+(\Pi_{x,\xi}(z))) \approx |\xi_{\pm s}^+(\Pi_{x,\xi}(z))|,$$

and so

$$\frac{\partial^2}{\partial s^2} |x_{\pm s}(z)|^2 \gtrsim |\xi_{\pm s}^+(\Pi_{x,\xi}(z))|^2 \left(1 - \|g - m\|_{AF_{>R}} \right)$$

provided that $|x_{\pm s}(z)| > R$. In such a case, we have that $\|g - m\|_{AF_{>R}} \ll 1$, and thus

$$\frac{\partial^2}{\partial s^2} |x_{\pm s}(z)|^2 > 0.$$

Using the condition that $|x_{\pm s'}^+(x, \xi)| \geq \max\{2R, |x| + \delta\}$ in a typical mean value theorem argument (elementary and, thus, omitted) establishes the existence of an $s'' \in [0, s']$ such that

$$|x_{\pm s''}(\tilde{w})|^2 > R^2 \quad \text{and} \quad \left(\frac{\partial}{\partial s} |x_{\pm s}(\tilde{w})|^2 \right) \Big|_{s=s''} > 0.$$

All together, we have that $|x_{\pm s}(\tilde{w})|^2$ has positive derivative at $s = s''$, and its derivative is increasing for all $s \geq s''$. In particular, $|x_{\pm s}(\tilde{w})|^2$ is increasing for all $s \geq s''$, which implies the result. \square

As a consequence, we can use the trapped and nontrapped sets to partition phase space.

PROPOSITION 2.11.

(a) *We can partition $T^*\mathbb{R}^3 \setminus o$ as*

$$\begin{aligned} T^*\mathbb{R}^3 \setminus o &= \Gamma_{tr}^\pm \sqcup \Gamma_\infty^\pm = \Lambda_{tr}^\pm \sqcup \Lambda_\infty^\pm, \\ T^*\mathbb{R}^3 \setminus o &= \Gamma_{tr}^\pm \cup \Lambda_{tr}^\pm \cup \Omega_\infty^\pm. \end{aligned}$$

(b) $\Gamma_\infty^\pm, \Lambda_\infty^\pm, \Omega_\infty^\pm$ are open in $T^*\mathbb{R}^3 \setminus o$, and $\Gamma_{tr}^\pm, \Lambda_{tr}^\pm, \Omega_{tr}^\pm$ are closed.

(c) *If $K \subset \Omega_\infty^\pm$ is compact, then for every $R \geq R_0$, there exists $T' \geq 0$ so that*

$$|x_s^\pm(v)| > R$$

for every $|s| \geq T'$ and $v \in K$. Also,

$$\bigcup_{s \in \mathbb{R}} \varphi_s^\pm(K)$$

is closed in $T^*\mathbb{R}^3 \setminus o$.

We omit the proof of this result, as it follows directly from using Lemma 2.10 and continuity of the flow in both the evolution parameter and the data, as in [BR14] (see Proposition 8.3 in the aforementioned work).

Finally, we show that if one assumes geometric control for bounded bicharacteristic rays, then it holds for semibounded bicharacteristic rays (that is, those that are bounded forward or backward in time). Although the proof is similar to that given in [BR14], we include it here due to its importance in our work and the seemingly increased complexity of our sets (at least notationally).

PROPOSITION 2.12. *Assume that the geometric control condition (2.1) holds. If $w \in \dot{\Gamma}_{tr}^\pm$, then there exists $s_\pm \geq 0$ so that $a(x_{s_\pm}^\pm(w)) > 0$. The same is true for $w \in \dot{\Lambda}_{tr}^\pm$, but with $s_\pm \leq 0$.*

Proof. We will only demonstrate this for $\dot{\Gamma}_{tr}^+$, as the work to establish the remaining cases is similar. If $w \in \dot{\Gamma}_{tr}^+$, then

$$\alpha := \sup_{s \geq 0} |x_s^+(w)| < \infty.$$

According to Remark 2.9, $|\xi_s^+(w)| \approx 1$ for all $s \in \mathbb{R}$. Thus,

$$\sup_{s \geq 0} |\varphi_s^+(w)| < \infty.$$

Then, there exists a point $w' \in T^*\mathbb{R}^3$ and a sequence (s_n) of nonnegative real numbers such that $\varphi_{s_n}^+(w) \rightarrow w'$ as $s_n \rightarrow \infty$. For any $s \in \mathbb{R}$, the group law for the flow tells us that $\varphi_{s+s_n}^+(w) = \varphi_s^+(\varphi_{s_n}^+(w))$, and so

$$x_{s+s_n}^+(w) = \Pi_x \circ \varphi_s^+(\varphi_{s_n}^+(w)) \rightarrow x_s^+(w') \text{ as } s_n \rightarrow \infty.$$

Since $s + s_n \geq 0$ for large enough n , it follows that $|x_s^+(w')| \leq \alpha$ for all $s \in \mathbb{R}$. By (2.1), there exists $s' \in \mathbb{R}$ for which $a(x_{s'}^+(w')) > 0$. Recall that $x_{s'+s_n}^+(w) \rightarrow x_{s'}^+(w')$ as $n \rightarrow \infty$. Since a is continuous and $s' + s_n \geq 0$ for n large enough, we conclude that $a(x_{s'+s_N}^+(w)) > 0$ for some large N . \square

2.4. Escape function construction. We will construct our symbols in multiple steps:

1. **On the characteristic set.** Since we are utilizing the half-wave decomposition, working on the characteristic set amounts to working on each individual light cone, then combining together. There are three regions of interest, two subregions of the *interior region* $\{|x| \leq R\}$ and the *exterior region* $\{|x| > R\}$. Here, $R \geq R_0$.
 - (a) **Interior, semibounded null bicharacteristics.** As opposed to working with the trapped and nontrapped sets, we will first work with the semibounded null bicharacteristics with initial data living in the interior region $\{|x| \leq R\}$. Working with the trapped and nontrapped sets can be difficult, since one can have nontrapped trajectories which are bounded forward or backward in time (but not both). Heuristically, these trajectories constitute the boundary of the nontrapped set. Instead, we will explicitly work with trajectories which are bounded forward or backward in time. This is where geometric control is used. This step is inspired by the work in [BR14].
 - (b) **The remainder of the interior region.** Since there is no trapping here, we construct a symbol similar to the one constructed in [BR14], [Doi96], and [MST20]. We will need to make an appropriate modification to avoid trapped trajectories while working with the half-wave symbols.
 - (c) **The exterior region.** As a consequence of asymptotic flatness, there are no trapped trajectories here. Hence, this follows from a similar multiplier to that used to prove local energy decay for the flat wave equation, although the multiplier must be appropriately adapted to the geometry. Here, we are motivated by prior work in [MMT08] and [MST20].
2. **On the elliptic set.** Here, we construct a correction term. That is, we will construct a lower-order symbol which provides no contribution on the characteristic set and provides positivity off of it. This is based on the work in [MST20].

We will break this construction up into a sequence of lemmas, starting with 1(a). While our construction follows that of [BR14], we reason differently. Their argument utilizes semiclassical rescaling, which provides compactness for their interior, semi-trapped set. Since we are sticking with the microlocal framework, we instead utilize homogeneity arguments to obtain this compactness. This is one of the reasons to work with the half-wave decomposition (the other being related to step 1(b), which we will outline once we get there).

With this in mind, we will utilize the sets

$$\begin{aligned}\Omega_R^\pm &:= (\Gamma_{tr}^\pm \cup \Lambda_{tr}^\pm) \cap \{|x| \leq R\}, \\ \dot{\Omega}_R^\pm &:= \Omega_R^\pm \cap \Phi^\pm(T^*\mathbb{R}^3 \setminus o).\end{aligned}$$

As a consequence of Propositions 2.8 and 2.11(b), the latter set is compact.

LEMMA 2.13 (semibounded escape function construction). *There exist $q^\pm \in C^\infty(T^*\mathbb{R}^3 \setminus o)$, an open set $V_R^\pm \supset \Omega_R^\pm$, and $C^\pm \in \mathbb{R}_+$ so that*

$$H_{p^\pm} q^\pm + C^\pm a \gtrsim_R \mathbb{1}_{V_R^\pm}.$$

Further, $q^\pm = q_1^\pm \circ \Phi^\pm$, where $q_1^\pm \in C_c^\infty(T^*\mathbb{R}^3 \setminus o)$.

Here, $\Phi^\pm \in S_{\text{hom}}^0(T^*\mathbb{R}^3 \setminus o)$ is the scaling function introduced in section 2.3. The fact that we omit the zero section is unavoidable, but it is nonproblematic; we will introduce high frequency cutoffs to our symbols later on which allow for smooth extensions to all of phase space.

Proof. We will first construct a symbol q_1^\pm and an open set $\dot{V}_R^\pm \supset \dot{\Omega}_R^\pm$ such that

$$H_{p^\pm} q_1^\pm + C^\pm a \gtrsim_R \mathbb{1}_{\dot{V}_R^\pm}.$$

To that end, let $w^\pm \in \dot{\Omega}_R^\pm$. By Proposition 2.12, there exists $s_{w^\pm} \in \mathbb{R}$ for which $a(x_{s_{w^\pm}}^\pm(w^\pm)) > 0$. Say that $2\alpha_{w^\pm} := a(x_{s_{w^\pm}}^\pm(w^\pm))$. By the continuity of the flow in the initial data, there exists a neighborhood U_{w^\pm} of w^\pm so that $a(x_{s_{w^\pm}}^\pm(z)) > \alpha_{w^\pm}$ for all $z \in U_{w^\pm}$. Select a smooth cutoff $\chi_{w^\pm} \in C_c^\infty(T^*\mathbb{R}^3)$ so that $\text{supp } \chi_{w^\pm} \subset U_{w^\pm}$ and $\chi_{w^\pm} \equiv 1$ on a smaller neighborhood V_{w^\pm} of w^\pm . Now, we define a symbol on $T^*\mathbb{R}^3 \setminus o$ given by

$$q_{w^\pm}(x, \xi) = \int_0^{s_{w^\pm}} (\chi_{w^\pm} \circ \varphi_{-s}^\pm)(x, \xi) ds.$$

Such a symbol is readily seen to be well-defined, and it is smooth by the aforementioned smooth flow dependence on data. Next, we demonstrate its symbolic nature. By continuity of the flow, $\varphi_{[0, s_{w^\pm}]}^\pm(\overline{U_{w^\pm}}) := \varphi^\pm([0, s_{w^\pm}] \times \overline{U_{w^\pm}})$ is compact. If $(x, \xi) \notin \varphi_{[0, s_{w^\pm}]}^\pm(\overline{U_{w^\pm}})$, then $(x, \xi) \notin \varphi_s^\pm(\overline{U_{w^\pm}})$ for any $s \in [0, s_{w^\pm}]$. Then, $\varphi_{-s}^\pm(x, \xi) \notin \overline{U_{w^\pm}}$ for any $s \in [0, s_{w^\pm}]$, implying that $q_{w^\pm}(x, \xi) = 0$. Hence, $q_{w^\pm} \in C_c^\infty(T^*\mathbb{R}^3 \setminus o)$.

Applying the Hamiltonian vector field H_{p^\pm} gives us

$$H_{p^\pm} q_{w^\pm} = \int_0^{s_{w^\pm}} H_{p^\pm}(\chi_{w^\pm} \circ \varphi_{-s}^\pm) ds = - \int_0^{s_{w^\pm}} \partial_s (\chi_{w^\pm} \circ \varphi_{-s}^\pm) ds = \chi_{w^\pm} - \chi_{w^\pm} \circ \varphi_{-s_{w^\pm}}^\pm.$$

Notice that the term $-\chi_{w^\pm} \circ \varphi_{-s_{w^\pm}}^\pm$ is nonpositive and that

$$\text{supp}(\chi_{w^\pm} \circ \varphi_{-s_{w^\pm}}^\pm) \subset \{v : \varphi_{-s_{w^\pm}}^\pm(v) \in U_{w^\pm}\} = \{v : v \in \varphi_{s_{w^\pm}}^\pm(U_{w^\pm})\} \subset \{x : a(x) > \alpha_{w^\pm}\}.$$

Using this support property, we can use the damping to absorb the poorly signed term and obtain nonnegativity of $H_{p^\pm} q_{w^\pm}$. Indeed, if we call $C_{w^\pm} = 2(\alpha_{w^\pm})^{-1}$, then we have

$$\chi_{w^\pm} \circ \varphi_{-s_{w^\pm}}^\pm + C_{w^\pm} a(x) \geq 0.$$

Thus,

$$H_{p^\pm} q_{w^\pm} + C_{w^\pm} a \gtrsim \mathbb{1}_{V_{w^\pm}}.$$

Since $\dot{\Omega}_R^\pm$ is compact, we can reduce the open cover $\{V_{w^\pm}\}_{w^\pm \in \dot{\Omega}_R^\pm}$ to a finite subcover $\{V_{w_j^\pm}\}_{j=1}^m$ with each $w_j^\pm \in \dot{\Omega}_R^\pm$. Call

$$\dot{V}_R^\pm = \bigcup_{j=1}^m V_{w_j^\pm}, \quad q_1^\pm = \sum_{j=1}^m q_{w_j^\pm}, \quad \text{and} \quad C^\pm = \sum_{j=1}^m C_{w_j^\pm}.$$

This provides us with a symbol $q_1^\pm \in C_c^\infty(T^*\mathbb{R}^3 \setminus o)$ so that

$$H_{p^\pm} q_1^\pm + C^\pm a \gtrsim \mathbb{1}_{\dot{V}_R^\pm}, \quad \dot{V}_R^\pm \supset \dot{\Omega}_R^\pm.$$

Finally, we will extend the above estimate from an indicator on \dot{V}_R^\pm to an indicator on a neighborhood $V_R^\pm \supset \Omega_R^\pm$. Consider the function $q^\pm : T^*\mathbb{R}^3 \setminus o \rightarrow \mathbb{R}$ given by

$$q^\pm = q_1^\pm \circ \Phi^\pm.$$

Since geometric control is invariant under Φ^\pm , we can see that $q^\pm \neq 0$. By definition,

$$H_{p^\pm} q^\pm \Big|_{(x,\xi)} = \frac{d}{ds} (q^\pm(x_s^\pm, \xi_s^\pm)) \Big|_{s=0}.$$

Since b^\pm is a constant of motion for the Hamiltonian system generated by p^\pm , it follows that

$$(\nabla_x b^\pm)(x_s^\pm, \xi_s^\pm) \dot{x}_s^\pm + (\nabla_\xi b^\pm)(x_s^\pm, \xi_s^\pm) \dot{\xi}_s^\pm = 0$$

for all s . Using this, we calculate that

$$\begin{aligned} \frac{d}{ds} (q^\pm(x_s^\pm, \xi_s^\pm)) &= \frac{d}{ds} \left(q_1^\pm \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \right) \\ &= (\nabla_x q_1^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \cdot (\dot{x}_s^\pm) \\ &\quad + (\nabla_\xi q_1^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \\ &\quad \cdot \frac{(|b^\pm(x_s^\pm, \xi_s^\pm)| \dot{\xi}_s^\pm - \xi_s^\pm ((\nabla_x b^\pm)(x_s^\pm, \xi_s^\pm) \dot{x}_s^\pm + (\nabla_\xi b^\pm)(x_s^\pm, \xi_s^\pm) \dot{\xi}_s^\pm))}{|b^\pm(x_s^\pm, \xi_s^\pm)|^2} \\ &= (\nabla_x q_1^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \cdot (\nabla_\xi p^\pm)(x_s^\pm, \xi_s^\pm) \\ &\quad - \frac{1}{|b^\pm(x_s^\pm, \xi_s^\pm)|} (\nabla_\xi q_1^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \cdot (\nabla_x p^\pm)(x_s^\pm, \xi_s^\pm) \\ &= (\nabla_x q_1^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \cdot (\nabla_\xi p^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \\ &\quad - (\nabla_\xi q_1^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \cdot (\nabla_x p^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right) \\ &= H_{p^\pm} q_1^\pm \Big|_{\left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right)}, \end{aligned}$$

where we have used homogeneity to obtain that

$$(\nabla_\xi p^\pm)(x_s^\pm, \xi_s^\pm) = (\nabla_\xi p^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right)$$

and

$$\frac{1}{|b^\pm(x_s^\pm, \xi_s^\pm)|} (\nabla_x p^\pm)(x_s^\pm, \xi_s^\pm) = (\nabla_x p^\pm) \left(x_s^\pm, \frac{\xi_s^\pm}{|b^\pm(x_s^\pm, \xi_s^\pm)|} \right).$$

If we define $V_R^\pm = (\Phi^\pm)^{-1}(\dot{V}_R^\pm)$, then we have an open neighborhood of Ω_R^\pm such that

$$H_{p^\pm} q^\pm \Big|_{(x,\xi)} + C^\pm a(x) = H_{p^\pm} q_1^\pm \Big|_{\left(x, \frac{\xi}{|b^\pm(x, \xi)|} \right)} + C^\pm a(x) \gtrsim \left(\mathbb{1}_{\dot{V}_R^\pm} \circ \Phi^\pm \right) (x, \xi) \geq \mathbb{1}_{V_R^\pm},$$

since $\Phi^\pm(V_R^\pm) \subset \dot{V}_R^\pm$. \square

Now that we have completed step 1(a), we move on to parts 1(b) and 1(c). Step 1(b) pertains to nontrapped null bicharacteristics in the interior region. The symbol that we produce follows the construction appearing in [Doi96] and utilized in many other works, such as [BR14] and [MST20]. Like in [MST20], we perform a factoring argument. The reason for studying the half-wave decomposition is due to the presence of a cutoff needed to make our constructed “symbol” genuinely symbolic. In the unfactored setting, cross terms in the metric arise when differentiating the cutoff in the computation of the Poisson bracket, generating an error term that is difficult to control. In the factored setting, this error can be handled straightforwardly.

Step 1(c) takes place in the exterior region. This is of little concern, as we possess robust exterior estimates. We utilize this symbol as a means of bootstrapping the aforementioned error term, which will be compactly supported in the region where the exterior symbol has strictly positive Poisson bracket with p^\pm .

To these ends, we will analyze both half-waves simultaneously (as in Lemma 2.13). While this portion of the argument follows the one given in [MST20], it does require a modification; the escape function on interior, nontrapped null bicharacteristics needs an appropriate adjustment to ensure that it avoids trapped trajectories. We start with a proposition where we construct a function that will be used for the previously described error absorption. The construction of this function comes from, e.g., [MST20], [Tat08].

PROPOSITION 2.14. *Let $\sigma > 0$. Then, there exists $f \in C^\infty$ satisfying $f(r) \approx_\sigma 1$ when $r > R_0$ and $f'(r) \approx \sigma c_j 2^{-j} f(r)$ when $r \approx 2^j > R_0$.*

Here, (c_j) is the slow-varying sequence introduced in section 1.2.

Remark 2.15. Although the sequence (c_j) is not defined for all natural numbers, the indices where it is not defined index finitely many dyadic regions (in particular, they omit where the operator P need not be a small AF perturbation). Since this region is compact, we can extend the sequence to such indices in an arbitrary manner. The typical way that this sequence is extended is by choosing c_j so that $\|g - m\|_{AF(A_j)} \lesssim c_j$ for the previously undefined indices j .

Proof. As in [Tat08], we can construct a smooth function $c(s)$ from the sequence (c_j) such that $c(s) \in (c_j, 2c_j)$ for each $s \in (2^j, 2^{j+1})$ and $|c'(s)| \leq \delta s^{-1} c(s)$. Since (c_j) is a positive sequence which converges to zero, it has a positive maximum, say c_N . Then, we observe that

$$\mathbf{c} \lesssim c_N \leq c(2^N + 2^{N-1}) = \left| \int_{2^N + 2^{N-1}}^{\infty} c'(s) ds \right| \lesssim \int_1^{\infty} \frac{c(s)}{s} ds$$

and

$$\int_1^{\infty} \frac{c(s)}{s} ds \leq \sum_{j=0}^{\infty} \int_{2^j}^{2^{j+1}} \frac{2c_j}{2^j} ds = 2 \sum_{j=0}^{\infty} c_j \lesssim \mathbf{c}.$$

That is,

$$\int_1^{\infty} \frac{c(s)}{s} ds \approx \mathbf{c}.$$

Now, set

$$f(r) = \exp \left(\sigma \int_1^r \frac{c(s)}{s} ds \right).$$

From our prior estimate, it is immediate that

$$f(r) \approx e^{\sigma c} \approx_\sigma 1$$

for $r > R_0$ and

$$f'(r) = \sigma \frac{c(r)}{r} f(r) \approx \sigma c_j 2^{-j} f(r)$$

for $r \approx 2^j$. □

Now, we complete steps 1(b) and 1(c).

LEMMA 2.16 (nontrapped escape function construction). *Let $R \geq R_0$. Then, there exist $q^\pm \in C^\infty(T^*\mathbb{R}^3 \setminus o)$ and $W^\pm \subset \Omega_\infty^\pm$ so that $V_R^\pm \cup W^\pm = T^*\mathbb{R}^3 \setminus o$ and*

$$H_{p^\pm} q^\pm \gtrsim c_j 2^{-j} \mathbb{1}_{W^\pm}, \quad |x| \approx 2^j.$$

Further, $q^\pm = \varepsilon q_{in}^\pm + q_{out}^\pm$, where $q_{in}^\pm = \tilde{q}_{in}^\pm \circ \Phi^\pm$ with $\tilde{q}_{in}^\pm \in C^\infty(T^*\mathbb{R}^3 \setminus o)$ is supported in $\{|x| \leq 4R\}$, $q_{out}^\pm \in S_{\text{hom}}^0(T^*\mathbb{R}^3 \setminus o)$, and $\varepsilon > 0$ is sufficiently small.

The inclusion of the sequence (c_j) is necessitated by the prior proposition, which is used for bootstrapping purposes in the exterior region. Its slowly varying nature allows one to work in the weight $\langle x \rangle^{-2}$ from the powers $|x| \approx 2^{-j}$ which will arise in the exterior (there is no trouble working in the weight $\langle x \rangle^{-2}$ in the interior region by compactness).

Proof. Recall from Proposition 2.8 that $|\xi| \approx |b^\pm(x, \xi)|$ on $T^*\mathbb{R}^3 \setminus o$; let $c^\pm, C^\pm > 0$ denote the respective lower and upper bound implicit constants in the inequalities and take δ^\pm such that $c^\pm - \delta^\pm > 0$. Now, choose $\psi^\pm \in C_c^\infty(T^*\mathbb{R}^3 \setminus o)$ such that

$$\begin{aligned} \text{supp } \psi^\pm &\subset \Omega_\infty^\pm \cap \{|x| \leq R\} \cap \{c^\pm - \delta^\pm < |\xi| < C^\pm + 1\}, \\ \psi^\pm &\equiv 1 \text{ on } U_R^\pm := (\Omega_\infty^\pm \cap \{|x| \leq R\} \cap \Phi^\pm(T^*\mathbb{R}^3 \setminus o)) \setminus \dot{V}_R, \end{aligned}$$

where $R \geq R_0$. Now, we define the function

$$\tilde{q}_{in}^\pm(x, \xi) = -\chi_{<2R}(|x|) \int_0^\infty \psi^\pm \circ \varphi_s^\pm(x, \xi) ds, \quad (x, \xi) \in T^*\mathbb{R}^3 \setminus o.$$

Since nontrapped null bicharacteristic rays must exit any compact set after a finite amount of time, this integral is well-defined for each $(x, \xi) \in T^*\mathbb{R}^3 \setminus o$, which establishes \tilde{q}_{in}^\pm as a well-defined function. It takes more work to show that \tilde{q}_{in}^\pm is smooth. Similar to [BR14], we will begin by establishing a maximal amount of time that bicharacteristic rays can remain in the support of the integrand. We already know that $\text{supp } \psi^\pm$ is compact. Let V^\pm be an open neighborhood of $\text{supp } \psi^\pm$ such that $\overline{V^\pm} \subset \Omega_\infty^\pm$. Take $\overline{V^\pm} = K$ in Proposition 2.11(c), and let T' be as given in the proposition. We claim that every point $w^\pm \in T^*\mathbb{R}^3 \setminus o$ has a neighborhood U_{w^\pm} of w^\pm and a time $s_{w^\pm} \geq 0$ such that $(\psi^\pm \circ \varphi_s^\pm)(z) = 0$ for every $z \in U_{w^\pm}$ and $s \in \mathbb{R}_+ \setminus [s_{w^\pm}, s_{w^\pm} + T']$. That is, all bicharacteristics (with speed ≈ 1) can spend no more than time T' within

$\text{supp } \psi^\pm$. The time s_{w^\pm} bears no similarity to the variable of the same name in the proof of Proposition 2.13. As a direct consequence of Proposition 2.11, we may take $U_{w^\pm} = V^\pm$ and $s_{w^\pm} = 0$ whenever $w^\pm \in \text{supp } \psi^\pm \subset V^\pm$. If

$$w^\pm \notin \bigcup_{s \in \mathbb{R}} \varphi_{-s}^\pm(\text{supp } \psi^\pm) =: \mathcal{X}^\pm,$$

then the fact that \mathcal{X}^\pm is closed provides an open neighborhood U_{w^\pm} of w^\pm such that $\mathcal{X}^\pm \cap U_{w^\pm} = \emptyset$. For each $z \in U_{w^\pm}$, we have that $\varphi_s^\pm(z) \notin \text{supp } \psi^\pm$ for all $s \in \mathbb{R}$, i.e., $(\psi^\pm \circ \varphi_s^\pm)(z) = 0$ for $s \in \mathbb{R}$. Hence, this case holds with U_{w^\pm} as defined and $s_{w^\pm} = 0$. Finally, let $w^\pm \in \mathcal{X}^\pm \setminus \text{supp } \psi^\pm$. Then, $\varphi_{s'}^\pm(w^\pm) \in \text{supp } \psi^\pm$ for some $s' \in \mathbb{R} \setminus \{0\}$. If $s' > 0$, then we can combine this with the fact that $\varphi_0^\pm(w) \notin \text{supp } \psi^\pm$ and the continuity of the flow to obtain $s_{w^\pm} > 0$ such that $\varphi_{s_{w^\pm}}^\pm(w) \in V^\pm$ and $\varphi_s^\pm(w) \notin \text{supp } \psi^\pm$ for all $s \in [0, s_{w^\pm}]$. By continuity of the flow in the data, we can extend the above to a neighborhood U_{w^\pm} . That is, there exists a neighborhood U_{w^\pm} of w^\pm so that for all $z \in U_{w^\pm}$, we have that $\varphi_{s_{w^\pm}}^\pm(z) \in V^\pm$ and $(\psi^\pm \circ \varphi_s^\pm)(z) = 0$ for all $s \in [0, s_{w^\pm}]$. Applying Proposition 2.11 to $K = V^\pm$ implies that $(\psi^\pm \circ \varphi_s^\pm)(z) = 0$ for all $z \in U_{w^\pm}$ and $s \in [0, s_{w^\pm}] \cup [s_{w^\pm} + T', \infty)$. It remains to consider if we cannot assume that $s' > 0$. In this case,

$$w^\pm \notin \bigcup_{s \in \mathbb{R}_+} \varphi_{-s}^\pm(\text{supp } \psi^\pm) =: \mathcal{X}_-^\pm.$$

Note that \mathcal{X}_-^\pm is closed by the same logic which showed that \mathcal{X}^\pm is closed (see the proof in Proposition 2.11(c)). From here, one can simply proceed as in the case where $w^\pm \notin \mathcal{X}^\pm$.

Using this result, we know that the integral present in \tilde{q}_{in}^\pm is always over an interval of maximal length T' . Hence, differentiation under the integral sign is nonproblematic, and in view of the regularity of the flow map, we conclude that $\tilde{q}_{in}^\pm \in C^\infty(T^*\mathbb{R}^3 \setminus o)$. Additionally, it is supported in $\{|x| \leq 4R\}$. In particular, it is smooth and bounded in all derivatives on the compact set

$$\{|x| \leq 4R\} \cap \Phi^\pm(T^*\mathbb{R}^3 \setminus o).$$

Now, consider the smooth function

$$q_{in}^\pm = \tilde{q}_{in}^\pm \circ \Phi^\pm$$

defined on $T^*\mathbb{R}^3 \setminus o$. As in the proof of Lemma 2.13, we get that

$$H_{p^\pm} q_{in}^\pm|_{(x,\xi)} = H_{p^\pm} \tilde{q}_{in}^\pm|_{\Phi^\pm(x,\xi)}.$$

Now, we calculate that

$$\begin{aligned} H_{p^\pm} \tilde{q}_{in}^\pm|_{\Phi^\pm(x,\xi)} &= \chi_{<2R}(|x|) \psi^\pm \left(x, \frac{\xi}{|b^\pm(x, \xi)|} \right) \\ &\quad + \frac{1}{2R} b_{\xi_k}^\pm \left(x, \frac{\xi}{|b^\pm(x, \xi)|} \right) \frac{x_k}{|x|} \chi' \left(\frac{|x|}{2R} \right) \int_0^\infty \psi^\pm \circ \varphi_s^\pm \left(x, \frac{\xi}{|b^\pm(x, \xi)|} \right) ds. \end{aligned}$$

The first term is nonnegative, supported in $\Omega_\infty^\pm \cap \{|x| \leq R\}$, and equal to 1 on $U^\pm := \Phi^{-1}(\dot{U}_R^\pm)$. The second term is an error term which is supported in $\{2R \leq |x| \leq 4R\}$.

The primary purpose of the exterior multiplier is to absorb this error term. To that end, let

$$q_{out}^\pm = -\chi_{>R}(|x|)f(|x|)b_{\xi_k}^\pm \frac{x_k}{|x|},$$

where f is the function constructed in Proposition 2.14. It is easy to see that $q_{out}^\pm \in S_{hom}^0(T^*\mathbb{R}^3 \setminus o)$, as it is smooth, bounded in all x derivatives due to asymptotic flatness, and homogeneous of degree 0, and it satisfies the appropriate symbol estimate. One can readily compute that

$$\begin{aligned} H_{p^\pm} q_{out}^\pm &= b_{\xi_k}^\pm \frac{x_k}{|x|} \chi_{>R}(|x|) f'(|x|) b_{\xi_j}^\pm \frac{x_j}{|x|} \\ &\quad + b_{\xi_k}^\pm \left(\delta_{jk} - \frac{x_j x_k}{|x|^2} \right) \chi_{>R}(|x|) \frac{f(|x|)}{|x|} \left(\delta_{jl} - \frac{x_j x_l}{|x|^2} \right) b_{\xi_l}^\pm \\ &\quad + R^{-1} \chi' \left(\frac{|x|}{R} \right) b_{\xi_k}^\pm \frac{x_k}{|x|} f(|x|) b_{\xi_j}^\pm \frac{x_j}{|x|} + \mathcal{O}(\langle x \rangle |\partial g|) \chi_{>R}(|x|) |x|^{-1}. \end{aligned}$$

We remark that the last term is small for $|x| > R$ by asymptotic flatness (and it is localized to this region due to the cutoff), while the remaining terms are all nonnegative. The third term is nonnegative and supported in the annulus $\{R \leq |x| \leq 2R\}$ due to the support of χ' . Making σ large enough and using asymptotic flatness provides that, for any $|x| \approx 2^j$,

$$\begin{aligned} H_{p^\pm} q_{out}^\pm &> \frac{\sigma}{2} c_j 2^{-j} f(|x|) \chi_{>R}(|x|) \frac{|x \cdot \nabla_\xi b^\pm|^2}{|x|^2} + \chi_{>R}(|x|) \frac{f(|x|)}{|x|} \left(|\nabla_\xi b^\pm|^2 - \frac{|x \cdot \nabla_\xi b^\pm|^2}{|x|^2} \right) \\ &\gtrsim c_j 2^{-j} \chi_{>R}(|x|) |\nabla_\xi b^\pm|^2 \\ &\gtrsim c_j 2^{-j} \chi_{>R}(|x|). \end{aligned}$$

Thus, $H_{p^\pm} q_{out}^\pm$ is nonnegative, is strictly positive for $|x| > R$, and

$$H_{p^\pm} q_{out}^\pm \gtrsim c_j 2^{-j} \chi_{>R}, \quad |x| \approx 2^j.$$

Recall that the error term in $H_{p^\pm} q_{in}^\pm$ is bounded and supported in $\{2R \leq |x| \leq 4R\}$, and $H_{p^\pm} q_{out}^\pm$ is strictly positive on the support of this error (with a uniform bound from below on this set).

Define

$$q^\pm = \varepsilon q_{in}^\pm + q_{out}^\pm \in C^\infty(T^*\mathbb{R}^3 \setminus o),$$

where $0 < \varepsilon \ll 1$. By choosing ε sufficiently small, we may absorb the aforementioned error due to our prior discussion, obtaining that $H_{p^\pm} q^\pm$ is nonnegative everywhere and positive on

$$W^\pm := U^\pm \cup \{(x, \xi) \in T^*\mathbb{R}^3 \setminus o : |x| > R\}.$$

By Proposition 2.11(a),

$$\begin{aligned} V_R^\pm \cup U^\pm &= V_R^\pm \cup ((\Omega_\infty^\pm \cap \{|x| \leq R\}) \setminus V_R^\pm) \\ &\supset (\Omega_R^\pm \cup \Omega_\infty) \cap \{|x| \leq R\} \\ &= (T^*\mathbb{R}^3 \setminus o) \cap \{|x| \leq R\}, \end{aligned}$$

and so

$$\begin{aligned} V_R^\pm \cup U^\pm &= (T^*\mathbb{R}^3 \setminus o) \cap \{|x| \leq R\}, \\ V_R^\pm \cup W^\pm &= T^*\mathbb{R}^3 \setminus o. \end{aligned}$$

We have already shown that

$$H_{p^\pm} q^\pm \approx 1, \quad (x, \xi) \in U^\pm,$$

and

$$H_{p^\pm} q^\pm \gtrsim c_j 2^{-j} \chi_{>R}, \quad |x| \approx 2^j.$$

The latter estimate readily extends to

$$H_{p^\pm} q^\pm \gtrsim c_j 2^{-j} \mathbb{1}_{W^\pm}, \quad |x| \approx 2^j,$$

by the compactness of the interior region $\{|x| \leq R\}$. \square

Now, we combine on the light cones to get our desired symbol q , as well as obtain positivity on the elliptic set (step 2). This largely follows the steps present in [MST20], although we have additional technicalities resulting from the damping.

Proof of Lemma 2.4. Let q_1^\pm denote the symbol q^\pm constructed in Lemma 2.13 (not the symbol q_1^\pm from the same lemma) and q_2^\pm denote the symbol q^\pm constructed in Lemma 2.16. We remark that, as a consequence of the chain rule, both symbols satisfy the standard S^0 bounds for $|\xi| \geq 1$. First, we truncate to the high frequency regime via the symbols

$$q_{j,>\lambda}^\pm = e^{-\sigma q_j^\pm} \chi_{>\lambda}(|b^\pm|), \quad j = 1, 2,$$

where σ is the parameter in Proposition 2.14. We assume that $\lambda > 1$. The exponentiation is implemented for bootstrapping: taking derivatives of the exponential will provide multiplication by $\sigma \gg 1$. Since $|b^\pm| \approx |\xi|$, these cutoffs genuinely truncate to high frequencies when λ is large. Further, the truncation to $|\xi| \gtrsim 1$ eliminates the singularities of q_j^\pm , i.e., $q_j^\pm \chi_{>\lambda}(|b^\pm|)$ smoothly extends to an element of $S^0(T^*\mathbb{R}^3)$.

We claim that exponentiation preserves the symbol class, so that $q_{j,>\lambda}^\pm \in S^0(T^*\mathbb{R}^3)$. We can immediately see that $q_{j,>\lambda}^\pm$ is smooth. Note that for $|\xi| \geq \lambda$, the exponentials $e^{-\sigma q_j^\pm}$ are bounded since q_j^\pm are bounded, and for $|\xi| < \lambda$, we immediately have that $q_{j,>\lambda}^\pm \equiv 0$. When checking the symbolic nature of $q_{j,>\lambda}^\pm$, we only need to study the boundedness of the ξ derivatives since our symbols q_j^\pm are bounded in all derivatives in x . Taking a partial derivative in ξ provides that

$$\partial_{\xi_k} q_{j,>\lambda}^\pm = -\sigma(\partial_{\xi_k} q_j^\pm) q_{j,>\lambda}^\pm \mp \frac{e^{-\sigma q_j^\pm}}{\lambda} (\partial_{\xi_k} b^\pm) \chi' \left(\frac{|b^\pm|}{\lambda} \right).$$

The first term is $\mathcal{O}(\langle \xi \rangle^{-1})$, and the second term is compactly supported in ξ . Due to the aforementioned compact support, we only need consider further ξ differentiation of $\sigma(\partial_{\xi_k} q_j^\pm) q_{j,>\lambda}^\pm$. If the ξ derivative lands on the exponential, then the result is $\mathcal{O}(\langle \xi \rangle^{-2})$ by the prior argument. If the derivative lands on $\partial_{\xi_k} q_j^\pm$, then the same asymptotics hold since $\partial_{\xi_k} q_j^\pm \in S^{-1}(T^*\mathbb{R}^3)$. If the derivative lands on the cutoff, then the result is compactly supported in ξ . Inducting establishes that $q_{j,>\lambda}^\pm \in S^0(T^*\mathbb{R}^3)$.

Now, we combine the symbols constructed on each light cone together as

$$q(\tau, x, \xi) = (\tau - b^+)(q_{1,>\lambda}^- + q_{2,>\lambda}^-) + (\tau - b^-)(q_{1,>\lambda}^+ + q_{2,>\lambda}^+).$$

Calling

$$q_j = (\tau - b^+)(q_{j,>\lambda}^- + (\tau - b^-)q_{j,>\lambda}^+),$$

we can see that

$$\begin{aligned} (H_p q + 2\gamma \tau a q) \big|_{\tau=b^\pm} &= (H_p q_1 + 2\gamma \tau a q_1) \big|_{\tau=b^\pm} + (H_p q_2 + 2\gamma \tau a q_2) \big|_{\tau=b^\pm} \\ &= H_p q_1 \big|_{\tau=b^\pm} \pm 2\gamma b^\pm (b^+ - b^-) a q_{1,>\lambda}^\pm \\ &\quad + H_p q_2 \big|_{\tau=b^\pm} \pm 2\gamma b^\pm (b^+ - b^-) a q_{2,>\lambda}^\pm. \end{aligned}$$

We will work with each term in the last equality separately. First, we compute that

$$\begin{aligned} H_p q_j \big|_{\tau=b^\pm} &= -(b^+ - b^-)^2 H_{p^\pm} q_{j,>\lambda}^\pm \\ &\quad - (b^\pm - b^\mp) q_{j,>\lambda}^\pm (b_{\xi_j}^\pm b_{x_j}^\mp - b_{x_j}^\pm b_{\xi_j}^\mp) \\ &= \sigma (b^+ - b^-)^2 q_{j,>\lambda}^\pm H_{p^\pm} q_j^\pm \\ &\quad - (b^\pm - b^\mp) q_{j,>\lambda}^\pm (b_{\xi_j}^\pm b_{x_j}^\mp - b_{x_j}^\pm b_{\xi_j}^\mp). \end{aligned}$$

By making σ sufficiently large, we get that

$$H_p q_j \big|_{\tau=b^\pm} \geq \frac{1}{2} \sigma (b^+ - b^-)^2 q_{j,>\lambda}^\pm H_{p^\pm} q_j^\pm + E_j^\pm,$$

where E_j^\pm are error terms which are supported in a neighborhood of the region where $H_{p^\pm} q_j^\pm = 0$. These terms are nonproblematic, as they are readily absorbed into the above estimate with differing j when we combine the estimates together. Hence, we will drop the E_j^\pm 's for ease of notation.

Observe that

$$\frac{b^\pm}{b^\pm - b^\mp} \approx 1.$$

By choosing γ large enough, we may apply Lemma 2.13 to obtain that

$$\begin{aligned} (2.5) \quad (H_p q_1 + 2\gamma \tau a q_1) \big|_{\tau=b^\pm} &\geq \frac{1}{2} \sigma (b^+ - b^-)^2 q_{1,>\lambda}^\pm H_{p^\pm} q_1^\pm \pm 2\gamma b^\pm (b^+ - b^-) a q_{1,>\lambda}^\pm \\ &= \frac{1}{2} \sigma (b^+ - b^-)^2 q_{1,>\lambda}^\pm \left(H_{p^\pm} q_1^\pm + \left(\frac{4\gamma}{\sigma} \right) \frac{b^\pm}{b^\pm - b^\mp} a \right) \\ &\gtrsim |\xi|^2 q_{1,>\lambda}^\pm \left(H_{p^\pm} q_1^\pm + \frac{\gamma}{\sigma} a \right) \\ &\gtrsim \mathbb{1}_{|\xi| \geq \lambda} \mathbb{1}_{V_R^\pm} |\xi|^2. \end{aligned}$$

Notice that γ depends on \mathbf{c} (and σ). For the $j = 2$ term, we use the prior computation, the fact that the damping term has positive sign, and Lemma 2.16:

$$\begin{aligned} (2.6) \quad (H_p q_2 + 2\gamma \tau a q_2) \big|_{\tau=b^\pm} &\geq \frac{1}{2} \sigma (b^+ - b^-)^2 q_{2,>\lambda}^\pm H_{p^\pm} q_2^\pm \pm 2\gamma b^\pm (b^+ - b^-) a q_{2,>\lambda}^\pm \\ &\gtrsim \mathbb{1}_{|\xi| \geq \lambda} \mathbb{1}_{W^\pm} c_j 2^{-j} |\xi|^2, \quad |x| \approx 2^j. \end{aligned}$$

Recall that $V_R^\pm \cup W^\pm = T^*\mathbb{R}^3 \setminus o$. Combining (2.5) and (2.6) together, we conclude that

$$(H_p q + 2\gamma\tau a q)|_{\tau=b^\pm} \gtrsim \mathbb{1}_{|\xi| \geq \lambda} \langle x \rangle^{-2} |\xi|^2,$$

where we have used the slowly varying, summable nature of (c_j) .

This provides the desired bound over the characteristic set. To extend it to all of phase space, we must construct a lower-order correction term. Explicitly, we seek an $m \in S^0$ so that

$$H_p q + 2\gamma\tau a q + mp \gtrsim \mathbb{1}_{|\xi| \geq \lambda} \langle x \rangle^{-2} |\xi|^2.$$

If we write

$$H_p q + 2\gamma a \tau q = a_0 \tau^2 + a_1 \tau + a_2,$$

where $a_j \in S^j$, then we have already established that

$$(2.7) \quad a_0(x, \xi)(b^\pm(x, \xi))^2 + a_1(x, \xi)b^\pm(x, \xi) + a_2(x, \xi) \gtrsim \mathbb{1}_{|\xi| \geq \lambda} \langle x \rangle^{-2} |\xi|^2.$$

So, we must analyze the quantity

$$a_0 \tau^2 + a_1 \tau + a_2 + pm = (a_0 - m) \tau^2 + (a_1 + (b^+ + b^-)m) \tau + (a_2 - b^+ b^- m).$$

If we choose m so that

$$(2.8) \quad a_0 - m > 0, \quad |\xi| \geq \lambda$$

and

$$(2.9) \quad (a_1 + (b^+ + b^-)m)^2 - 4(a_0 - m)(a_2 - b^+ b^- m) < 0, \quad |\xi| \geq \lambda,$$

then we will have that $a_0 \tau^2 + a_1 \tau + a_2 + mp$ is positive for $|\xi| \geq \lambda$ (the first condition on m guarantees that this polynomial in τ is concave up, and the second guarantees that there are no real zeros).

Let us begin by focusing on (2.9). The function

$$\begin{aligned} P(m) &= (a_1 + (b^+ + b^-)m)^2 - 4(a_0 - m)(a_2 - b^+ b^- m) \\ &= (b^+ - b^-)^2 m^2 + (2a_1(b^+ + b^-) + 4a_0 b^+ b^- + 4a_2)m + (a_1^2 - 4a_0 a_2) \end{aligned}$$

is a quadratic polynomial in m with a positive coefficient on the quadratic term, so it will achieve a minimal value at

$$m = -\frac{a_1(b^+ + b^-) + 2(a_0 b^+ b^- + a_2)}{(b^+ - b^-)^2}.$$

It is readily seen that $m \in S^0$ and that m is supported where $|\xi| \geq \lambda$. This minimal value is

$$\begin{aligned} P(m) &= \left(a_1 - (b^+ - b^-) \frac{a_1(b^+ + b^-) + 2(a_0 b^+ b^- + a_2)}{(b^+ - b^-)^2} \right)^2 \\ &\quad - 4 \left(a_0 + \frac{a_1(b^+ + b^-) + 2(a_0 b^+ b^- + a_2)}{(b^+ - b^-)^2} \right) \\ &\quad \times \left(a_2 + b^+ b^- \frac{a_1(b^+ + b^-) + 2(a_0 b^+ b^- + a_2)}{(b^+ - b^-)^2} \right) \\ &= -4 \frac{(a_0(b^+)^2 + a_1 b^+ + a_2)(a_0(b^-)^2 + a_1 b^- + a_2)}{(b^+ - b^-)^2} \\ &= -4(b^+ - b^-)^{-2} \left((H_p q + 2\gamma\tau a q)|_{\tau=b^+} \right) \left((H_p q + 2\gamma\tau a q)|_{\tau=b^-} \right) \\ &< 0, \end{aligned}$$

where we have used (2.7). So, (2.9) is satisfied. To establish (2.8), one can readily check that

$$\begin{aligned} a_0 - m &= a_0 + \frac{a_1(b^+ + b^-) + 2(a_0b^+b^- + a_2)}{(b^+ - b^-)^2} \\ &= (b^+ - b^-)^{-2} ((H_pq + 2\gamma\tau aq)|_{\tau=b^+} + (H_pq + 2\gamma\tau aq)|_{\tau=b^-}) \\ &> 0 \end{aligned}$$

for $|\xi| \geq \lambda$.

This gives us that

$$H_pq + 2\gamma\tau aq + mp > 0$$

for $|\xi| \geq \lambda$. In fact, we can check that the minimal value of the above in τ for $|\xi| \geq \lambda$ is

$$\begin{aligned} &\frac{(a_0(b^+)^2 + a_1b^+ + a_2)(a_0(b^+)^2 + a_1b^+ + a_2)}{(a_0(b^+)^2 + a_1b^+ + a_2) + (a_0(b^-)^2 + a_1b^- + a_2)} \\ &= \frac{((H_pq + 2\gamma\tau aq)|_{\tau=b^+}) (H_pq + 2\gamma\tau aq)|_{\tau=b^-})}{(H_pq + 2\gamma\tau aq)|_{\tau=b^+} + (H_pq + 2\gamma\tau aq)|_{\tau=b^-}}. \end{aligned}$$

The numerator is bounded below by $\langle x \rangle^{-4} |\xi|^4$. In view of the support and symbolic properties of q , the denominator satisfies the bounds

$$(H_pq + 2\gamma\tau aq)|_{\tau=b^+} + (H_pq + 2\gamma\tau aq)|_{\tau=b^-} \approx \langle x \rangle^{-2} |\xi|^2.$$

Since $|b^\pm(x, \xi)| \approx |\xi|$ and $|\tau| = |b^\pm(x, \xi)|$ in the above, we conclude the desired result. \square

2.5. Starting energy estimates and case reductions. In this section, we will establish various useful energy estimates, then reduce the proof of Theorem 1.7 to a simpler problem. Our starting point is a standard uniform energy inequality.

PROPOSITION 2.17. *Let P be a stationary damped wave operator, ∂_t be uniformly time-like, and $T > 0$. Then, we have the estimate*

$$(2.10) \quad \|\partial u(t)\|_{L^2}^2 \lesssim \|\partial u(0)\|_{L^2}^2 + \int_0^T \int_{\mathbb{R}^3} |Pu \partial_t u| dx dt, \quad 0 \leq t \leq T,$$

for all $u \in \mathcal{W}_T$.

Proof. Call $Pu = f$, and define the energy functional

$$E[u](t) = \int_{\mathbb{R}^3} D_i g^{ij} D_j u \bar{u} - g^{00} |\partial_t u|^2 dx.$$

After integrating the first term by parts, this functional is readily seen to be coercive due to the uniformly time-like nature of ∂_t , i.e.,

$$E[u](t) \approx \|\partial u(t)\|_{L^2}^2.$$

Differentiating in t and integrating by parts, gives that

$$\begin{aligned} \frac{d}{dt} E[u](t) &= - \int_{\mathbb{R}^3} g^{00} (\partial_t^2 u \partial_t \bar{u} + \partial_t u \partial_t^2 \bar{u}) dx + \int_{\mathbb{R}^3} D_i g^{ij} D_j \partial_t u \bar{u} + D_i g^{ij} D_j u \partial_t \bar{u} dx \\ &= \int_{\mathbb{R}^3} (g^{00} D_t^2 + D_i g^{ij} D_j) u \partial_t \bar{u} + \partial_t u \overline{(g^{00} D_t^2 + D_i g^{ij} D_j) u} dx \\ &= \int_{\mathbb{R}^3} (-(g^{0j} D_j D_t + D_j g^{0j} D_t + i a D_t) u + f) \partial_t \bar{u} \\ &\quad + \partial_t u \overline{(-(g^{0j} D_j D_t + D_j g^{0j} D_t + i a D_t) u + f)} dx \\ &= 2 \operatorname{Re} \int_{\mathbb{R}^3} \bar{f} \partial_t u dx - 2 \int_{\mathbb{R}^3} a |\partial_t u|^2 dx. \end{aligned}$$

Dropping the damping term and integrating the resulting estimate in time yields the inequality

$$E[u](t) \lesssim E[u](0) + \int_0^T \int_{\mathbb{R}^3} |f \partial_t u| dx dt, \quad 0 \leq t \leq T.$$

Applying the coercivity allows us to conclude. \square

Remark 2.18. Note that when $Pu = 0$, we have the energy dissipation statement

$$\frac{d}{dt} E[u](t) = -2 \int_{\mathbb{R}^3} a |\partial_t u|^2 dx \leq 0.$$

Applying the Schwarz inequality to Proposition 2.17 provides us with estimates which will prove useful throughout this work.

COROLLARY 2.19. *Under the same assumptions as Proposition 2.17, the uniform energy estimates*

$$\begin{aligned} \|\partial u\|_{L_t^\infty L_x^2} &\lesssim \|\partial u(0)\|_{L^2} + \|Pu\|_{L_t^1 L_x^2}, \\ \|\partial u\|_{L_t^\infty L_x^2} &\lesssim \|\partial u(0)\|_{L^2} + \|Pu\|_{LE^*}^{1/2} \|u\|_{LE^1}^{1/2}, \end{aligned}$$

and

$$\|\partial u\|_{L_t^\infty L_x^2} \lesssim \|\partial u(0)\|_{L^2} + \varepsilon^{-1} \|Pu\|_{LE^* + L_t^1 L_x^2} + \varepsilon \|u\|_{LE^1} \quad \forall \varepsilon > 0$$

hold.

Proof. By Proposition 2.17, we have the estimate

$$\|\partial u(t)\|_{L^2}^2 \lesssim \|\partial u(0)\|_{L^2}^2 + \int_0^T \int_{\mathbb{R}^3} |Pu \partial_t u| dx dt, \quad 0 \leq t \leq T.$$

To obtain the first estimate that we claimed, one applies the Schwarz inequality, takes a supremum in time of $\partial_t u$, and uses Young's inequality for products. To obtain the second estimate, write

$$|Pu \partial_t u| = \left(\langle x \rangle^{1/2} |Pu| \right) \left(\langle x \rangle^{-1/2} |\partial_t u| \right)$$

in (2.10) and apply the Schwarz inequality and Hölder's inequality applied to ℓ^1 with conjugate exponents $(p, q) = (1, \infty)$. \square

Next, we cite an exterior estimate from [MST20] (see [MMT08] for a similar result for the Schrödinger equation). Outside of a large enough compact spatial set, the operator P is a small *AF* perturbation of \square , in which case we obtain good energy estimates with a necessary truncation error (here, we must measure the energy at time T , which is nonproblematic in view of Corollary 2.19).

PROPOSITION 2.20 (Proposition 3.2 in [MST20]). *If P is asymptotically flat and $R \geq R_0$, then*

$$(2.11) \quad \|u\|_{LE_{>R}^1} \lesssim \|\partial u(0)\|_{L_{>R}^2} + \|\partial u(T)\|_{L_{>R}^2} + R^{-1} \|u\|_{LE_R} + \|Pu\|_{LE_{>R}^*}.$$

Since the damping is identically zero in this region, the result holds without any modification to the proof given in [MST20]. For this reason, we will omit the details here, although we will provide the overarching idea: Their proof is a positive commutator argument using the multiplier $Q_1 + Q_2$, where

$$Q_1 = \chi_{>2R}(|x|) f(|x|) \frac{x_j}{|x|} g^{jk} D_k + D_k \chi_{>2R}(|x|) f(|x|) \frac{x_j}{|x|} g^{jk}$$

is the principal term, and

$$Q_2 = \chi_{>2R}(|x|) f'(|x|)$$

is the lower-order correction term. Here, $f(|x|) = \frac{|x|}{|x|+2^j}$, and j is chosen so that $2^j \geq R$.

Before moving on to proving Theorem 1.7, we will simplify its proof through case reductions. As in section 4 of [MST20], one can readily reduce to the case of u having zero Cauchy data at times 0 and T , as well as $f \in LE_c^*$. To do this, one constructs an approximate solution to a problem with the same data and forcing with a wave operator which is a small *AF* perturbation of \square and agrees with P for $|x| > R_0$, then considers approximations using a unit time interval partition of unity and matching the initial (respectively, final time) Cauchy data on the first (respectively, last) solution granted by the partition. In view of these reductions, it is enough to establish

$$(2.12) \quad \|u\|_{LE^1[0,T]} \lesssim \left\| \langle x \rangle^{-2} u \right\|_{LE[0,T]} + \|Pu\|_{LE_c^*[0,T]}$$

for all $u \in \mathcal{W}_T$ satisfying that $u[0] = u[T] = 0$ in order to prove Theorem 1.7. The implicit constant in (2.12) is still independent of T .

We will explicitly establish an additional reduction, namely to solutions with compact spatial support, using Proposition 2.20.

Claim. It suffices to prove (2.12), and hence Theorem 1.7, for u supported in $\{|x| \leq 2R_0\}$.

Proof. Write $u = \chi_{<R_0} u + \chi_{>R_0} u$. On the exterior piece $\chi_{>R_0} u$, we apply Proposition 2.20 and Corollary 2.19 to get that

$$\|\chi_{>R_0} u\|_{LE^1[0,T]} \lesssim R_0^{-1} \|\chi_{>R_0} u\|_{LE_{R_0}[0,T]} + \|P(\chi_{>R_0} u)\|_{LE_{>R_0}^*[0,T]}.$$

The first term on the right is directly bounded by $\|u\|_{LE_{R_0}^1[0,T]}$. For the second term, we write

$$P(\chi_{>R_0}u) = (\chi_{>R_0})Pu + [P, \chi_{>R_0}]u,$$

and one can calculate that

$$[P, \chi_{>R_0}]u(t, x) = \mathcal{O}(R_0^{-1})\chi' \left(\frac{|x|}{R_0} \right) \partial u(t, x) + \mathcal{O}(R_0^{-2})\chi'' \left(\frac{|x|}{R_0} \right) u(t, x).$$

In LE^* , this term bounded by $\|u\|_{LE_{R_0 \leq |\cdot| \leq 2R_0}^1[0,T]}$, and so we have

$$\|\chi_{>R_0}u\|_{LE^1[0,T]} \lesssim \|u\|_{LE_{R_0}^1[0,T]}.$$

Suppose that (2.12) holds for $\chi_{<R_0}u$. From this, we get the estimate

$$\|\chi_{<R_0}u\|_{LE^1[0,T]} \lesssim \left\| \langle x \rangle^{-2} \chi_{<R_0}u \right\|_{LE[0,T]} + \|Pu\|_{LE_c^*[0,T]} + \|[P, \chi_{<R_0}]u\|_{LE_c^*[0,T]},$$

and so

$$\begin{aligned} \|u\|_{LE^1[0,T]} &\leq \|\chi_{<R_0}u\|_{LE^1[0,T]} + \|\chi_{>R_0}u\|_{LE^1[0,T]} \\ &\lesssim \left\| \langle x \rangle^{-2} u \right\|_{LE[0,T]} + \|Pu\|_{LE_c^*[0,T]} + \|u\|_{LE_{R_0 \leq |\cdot| \leq 2R_0}^1[0,T]}. \end{aligned}$$

The last term is readily estimated via Proposition 2.20, which establishes (2.12). \square

We record the results of these case reductions in the following proposition.

PROPOSITION 2.21. *In order to establish Theorem 1.7, it is sufficient to prove the estimate*

$$(2.13) \quad \|v\|_{LE^1[0,T]} \lesssim \|v\|_{L_t^2 L_x^2[0,T]} + \|Pv\|_{LE^*[0,T]}$$

for v supported in $\{|x| \leq 2R_0\}$ with $v[0] = v[T] = 0$.

Again, this implicit constant is independent of T but will depend on R_0 . Using the compact support of v to transition between the weighted and unweighted spaces will inherently generate multiplication by powers of R_0 , but this does not matter since the constant in the above may depend on such a parameter.

2.6. Proof of the high frequency estimate. Armed with the established case reductions, we will proceed with a proof of Theorem 1.7. Recall that it is equivalent to analyze the scaled problem.

Proof of Theorem 1.7. We will break this proof into a sequence of steps.

Step 1: Setting up the positive commutator and the frequency decomposition.

First, we remark that in view of Proposition 2.21, it will suffice to prove (2.13) for v supported in $\{|x| < 2R_0\}$ with $v[0] = v[T] = 0$. We can extend v by zero to be defined for $t \in \mathbb{R}$ and vanish for $t \notin (0, T)$. Then,

$$\begin{aligned} (2.14) \quad &2\text{Im} \left\langle Pv, \left(q^w - \frac{i}{2}m^w \right) v \right\rangle + \frac{i\gamma}{2} \langle [aD_t, m^w]v, v \rangle = \langle i[\square_g, q^w]v, v \rangle \\ &+ \gamma \langle (q^w aD_t + aD_t q^w)v, v \rangle \\ &+ \frac{1}{2} \langle (\square_g m^w + m^w \square_g)v, v \rangle. \end{aligned}$$

The right-hand side of (2.14) can be written as

$$\begin{aligned} & \langle i[\square_g, q^w]v, v \rangle + \gamma \langle (q^w a D_t + a D_t q^w)v, v \rangle + \frac{1}{2} \langle (\square_g m^w + m^w \square_g)v, v \rangle \\ & = \langle (H_p q + 2\gamma\tau a q + mp)^w v, v \rangle + \langle A_0 v, v \rangle, \end{aligned}$$

where $A_0 \in \Psi^0$. Recall that $2\gamma\tau a = -2is_{skew}$. By choosing $\gamma > 0$ large enough, we can apply Lemma 2.4 to get

$$(2.15) \quad H_p q - 2is_{skew} q + mp - C\mathbb{1}_{|\xi| \geq \lambda} \langle x \rangle^{-2} (|\xi|^2 + \tau^2) \geq 0,$$

where $C > 0$ is the implicit constant in Lemma 2.4. We can readily replace $\mathbb{1}_{|\xi| \geq \lambda}$ with $\chi_{|\xi| > \lambda}$.

We will split v into frequency components via $v = v_{>>\lambda} + v_{<<\lambda}$, where

$$\begin{aligned} v_{>>\lambda} &= \chi_{|\xi| + |\tau| > \lambda}(\partial) v, \\ v_{<<\lambda} &= \chi_{|\xi| + |\tau| < \lambda}(\partial) v. \end{aligned}$$

Since the desired estimate is a high frequency estimate, we will first analyze the high frequency components of v .

Step 2a: High frequencies—applying the Gårding inequality. By (2.15), we may apply the sharp Gårding inequality to obtain that

$$\begin{aligned} \langle (H_p q - 2is_{skew} q + mp)^w v_{>>\lambda}, v_{>>\lambda} \rangle &\gtrsim \left\langle \left(\chi_{|\xi| > \lambda} \langle x \rangle^{-2} (|\xi|^2 + \tau^2) \right)^w v_{>>\lambda}, v_{>>\lambda} \right\rangle \\ &\quad - \|v_{>>\lambda}\|_{H_{t,x}^{1/2}}^2. \end{aligned}$$

We remark that the implicit constant may be chosen independently of λ since there is a $\chi_{|\xi| > \lambda}$ cutoff embedded into q and m , and hence differentiation occurring in asymptotic expansion calculations possesses coefficients which either are independent of λ or feature inverse powers of λ (one can also entirely ignore the potential λ dependence and argue via Cauchy–Schwarz and Young’s inequality for products, although this introduces more parameters to track).

Since $\chi_{|\xi| + |\tau| < \lambda} \in S^{-\infty}$, it follows that

$$\langle (H_p q - 2is_{skew} q + mp)^w v, v \rangle = \langle (H_p q - 2is_{skew} q + mp)^w v_{>>\lambda}, v_{>>\lambda} \rangle + \langle S_0 v, v \rangle,$$

where $S_0 \in \Psi^{-\infty}$. In particular,

$$\begin{aligned} (2.16) \quad \langle (H_p q - 2is_{skew} q + mp)^w v, v \rangle &\gtrsim \left\langle \left(\chi_{|\xi| > \lambda} \langle x \rangle^{-2} (|\xi|^2 + \tau^2) \right)^w v_{>>\lambda}, v_{>>\lambda} \right\rangle \\ &\quad - \|v_{>>\lambda}\|_{H_{t,x}^{1/2}}^2 + \langle S_0 v, v \rangle. \end{aligned}$$

Using the pseudodifferential composition formula, we compute that

$$(2.17) \quad \left(\chi_{|\xi| > \lambda} \langle x \rangle^{-2} (|\xi|^2 + \tau^2) \right)^w = \left(\chi_{|\xi| > \lambda} (D_x) \right)^{1/2} D_\alpha \langle x \rangle^{-2} D_\alpha \left(\chi_{|\xi| > \lambda} (D_x) \right)^{1/2} + A_1,$$

where $A_1 \in \Psi^1$ arises from nonprincipal terms in the asymptotic expansion of the Moyal product (and the expansion features terms that are either independent of λ or involve inverse powers of λ). Integrating by parts once gives that

$$\begin{aligned} (2.18) \quad \left\langle \left(\chi_{|\xi| > \lambda} \langle x \rangle^{-2} (|\xi|^2 + \tau^2) \right)^w v_{>>\lambda}, v_{>>\lambda} \right\rangle &= \left\| \langle x \rangle^{-1} \partial v_{>\lambda} \right\|_{L_t^2 L_x^2}^2 + \langle A_1 v_{>>\lambda}, v_{>>\lambda} \rangle \\ &\gtrsim \|\partial v_{>\lambda}\|_{LE_{<2R_0}}^2 + \langle A_1 v_{>>\lambda}, v_{>>\lambda} \rangle, \end{aligned}$$

where $v_{>\lambda} = \chi_{|\xi|>\lambda}(D_x)v$ and $A_1 \in \Psi^1$ is now a modification of the previous version of the same variable in (2.17) (to be explained momentarily). One might expect the term $\partial((\chi_{|\xi|>\lambda}(D_x))^{1/2}v_{>>\lambda})$ to appear instead of $\partial v_{>\lambda}$, but it is readily seen that

$$(\chi_{|\xi|>\lambda}(|\xi|))^{1/2} \chi_{|\xi|+|\tau|>\lambda}(|(\tau, \xi)|) \approx \chi_{|\xi|>\lambda}(|\xi|) \chi_{|\xi|+|\tau|>\lambda}(|(\tau, \xi)|) = \chi_{|\xi|>\lambda}(|\xi|).$$

In particular, the τ has no effect on the resulting cutoff, and $\chi, \chi^{1/2}$ are both smooth and nondecreasing and have the same support properties (and differ only on a compact set). The only effect in exchanging these terms is modifying A_1 in order to account for the error resulting from this switch (i.e. the difference between the cutoffs); hence, the A_1 in (2.18) is different than in (2.17). For this reason, none of our analysis changes by working with $v_{>\lambda}$, and we will stick with this for notational convenience.

After incorporating (2.18) into (2.16), we have that

$$(2.19) \quad \langle (H_p q - 2is_{skew} aq + mp)^w v, v \rangle + \langle A_0 v, v \rangle \gtrsim \|\partial v_{>\lambda}\|_{LE_{<2R_0}}^2 - \|v_{>>\lambda}\|_{H_{t,x}^{1/2}}^2 - |\langle A_1 v_{>>\lambda}, v_{>>\lambda} \rangle| - |\langle A_0 v, v \rangle| - |\langle S_0 v, v \rangle|.$$

Step 2b: High frequencies—handling the error terms in (2.19). We will first analyze the term $\langle A_1 v_{>>\lambda}, v_{>>\lambda} \rangle$. Since $A_1 \in \Psi^1$, it is bounded from $H_{t,x}^1$ to $L_t^2 L_x^2$ (and the operator norm will yield no positive-power λ contributions due to the previous comment on the asymptotic expansion of the symbol). By using the Schwarz inequality and this mapping property, we have that

$$(2.20) \quad |\langle A_1 v_{>>\lambda}, v_{>>\lambda} \rangle| \lesssim \|v_{>>\lambda}\|_{H_{t,x}^1} \|v_{>>\lambda}\|_{L_t^2 L_x^2}.$$

Using Plancherel's theorem in (t, x) , the frequency localization, and the compact support of v , we obtain the bounds

$$(2.21) \quad \|v_{>>\lambda}\|_{H_{t,x}^1} \lesssim \|\langle(\tau, \xi)\rangle \chi_{|\xi|+|\tau|>\lambda} \hat{v}\|_{L_\tau^2 L_\xi^2} \lesssim \|\langle(\tau, \xi)\rangle \hat{v}\|_{L_\tau^2 L_\xi^2} = \|v\|_{H_{t,x}^1} \lesssim \|v\|_{LE^1}$$

and

$$(2.22) \quad \begin{aligned} \|v_{>>\lambda}\|_{L_t^2 L_x^2} &\approx \|\chi_{|\xi|+|\tau|>\lambda} \hat{v}\|_{L_\tau^2 L_\xi^2} \lesssim \left\| \frac{|\tau| + |\xi|}{\lambda} \chi_{|\xi|+|\tau|>\lambda} \hat{v} \right\|_{L_\tau^2 L_\xi^2} \\ &\lesssim \lambda^{-1} \|\partial v\|_{L_t^2 L_x^2} \lesssim \lambda^{-1} \|v\|_{LE^1}. \end{aligned}$$

Applying (2.21) and (2.22) to (2.20) yields that

$$|\langle A_1 v_{>>\lambda}, v_{>>\lambda} \rangle| \lesssim \lambda^{-1} \|v\|_{LE^1}^2.$$

For the term $\|v_{>>\lambda}\|_{H_{t,x}^{1/2}}^2$, note that

$$\begin{aligned} \|v_{>>\lambda}\|_{H_{t,x}^{1/2}}^2 &\lesssim \left\| \langle(\tau, \xi) \rangle^{1/2} \chi_{|\xi|+|\tau|>\lambda} \hat{v} \right\|_{L_\tau^2 L_\xi^2}^2 = \left\| \langle(\tau, \xi) \rangle^{-1/2} \langle(\tau, \xi) \rangle \chi_{|\xi|+|\tau|>\lambda} \hat{v} \right\|_{L_\tau^2 L_\xi^2}^2 \\ &\lesssim \lambda^{-1} \|\langle(\tau, \xi) \rangle \chi_{|\xi|+|\tau|>\lambda} \hat{v}\|_{L_\tau^2 L_\xi^2}^2 \lesssim \lambda^{-1} \|v\|_{LE^1}^2. \end{aligned}$$

For the $\langle A_0 v, v \rangle$ term, we can use L^2 -boundedness and the compact support of v to get

$$(2.23) \quad |\langle A_0 v, v \rangle| \leq \|A_0 v\|_{L_t^2 L_x^2} \|v\|_{L_t^2 L_x^2} \lesssim C(\lambda) \|v\|_{L_t^2 L_x^2}^2.$$

While this bound is λ -dependent, such terms appear on the *upper bound side* of the desired inequality, and hence can depend on λ in an arbitrary manner (as opposed to the LE^1 terms, which need an inverse power of λ for bootstrapping). The meaning of $C(\lambda)$ will change fluidly, just as one continuously re-notates a potentially changing constant by C when calculating successive inequalities.

The smoothing term $\langle S_0 v, v \rangle$ can be bounded in the same way as $\langle A_0 v, v \rangle$ (in particular, $S_0 \in \Psi^0$). Thus, we have the lower bound

(2.24)

$$\langle (H_p q - 2i s_{skew} q + mp)^w v, v \rangle + \langle A_0 v, v \rangle \gtrsim \|\partial v_{>\lambda}\|_{LE_{<2R_0}}^2 - C(\lambda) \|v\|_{L_t^2 L_x^2}^2 - \lambda^{-1} \|v\|_{LE^1}^2.$$

Next, we look at the left-hand side of (2.14).

Step 3: Bounding the left-hand side of (2.14). Since $[aD_t, m^w] \in \Psi^0$, performing the same work as in (2.23) provides that

$$(2.25) \quad \left| \frac{i\gamma}{2} \langle [aD_t, m^w] v, v \rangle \right| \lesssim C(\lambda) \|v\|_{L_t^2 L_x^2}^2.$$

For the remaining term on the left-hand side of (2.14), we split v into high and low frequency components once again to get that

$$2\text{Im} \left\langle Pv, \left(q^w - \frac{i}{2} m^w \right) v \right\rangle = 2\text{Im} \left\langle Pv, \left(q^w - \frac{i}{2} m^w \right) v_{>>\lambda} \right\rangle + \langle S_1 v, v \rangle,$$

where $S_1 \in \Psi^{-\infty}$. We have already demonstrated how to bound smoothing operator terms. For the other (primary) piece, we apply the Schwarz inequality, use the Ψ DO mapping properties of $q^w \in \Psi^1$ and $m^w \in \Psi^0$, and leverage the compact support of v (just as performed previously) to get

(2.26)

$$\left| 2\text{Im} \left\langle Pv, \left(q^w - \frac{i}{2} m^w \right) v_{>>\lambda} \right\rangle \right| \lesssim C(\lambda) \|Pv\|_{L_t^2 L_x^2} \|v\|_{LE^1} \lesssim C(\lambda) \|Pv\|_{LE_c^*} \|v\|_{LE^1}.$$

Step 4: Combining the established bounds into a high frequency bound. Putting (2.14), (2.24), (2.25), and (2.26) together, we obtain that

$$\|\partial v_{>\lambda}\|_{LE_{<2R_0}} \lesssim C(\lambda) \left(\|Pv\|_{LE^*}^{1/2} \|v\|_{LE^1}^{1/2} + \|v\|_{L_t^2 L_x^2} \right) + \lambda^{-1/2} \|v\|_{LE^1}.$$

Completing the $LE_{<2R_0}^1$ norm on the left-hand side of the above,

$$\|v_{>\lambda}\|_{LE_{<2R_0}^1} \lesssim C(\lambda) \left(\|Pv\|_{LE^*}^{1/2} \|v\|_{LE^1}^{1/2} + \|v\|_{L_t^2 L_x^2} \right) + \lambda^{-1/2} \|v\|_{LE^1} + \left\| \langle x \rangle^{-1} v_{>\lambda} \right\|_{LE}.$$

We note that

$$\left\| \langle x \rangle^{-1} v_{>\lambda} \right\|_{LE} \lesssim \|v\|_{L_t^2 L_x^2},$$

once again using Plancherel's theorem. Thus,

$$(2.27) \quad \|v_{>\lambda}\|_{LE_{<2R_0}^1} \lesssim C(\lambda) \left(\|Pv\|_{LE^*}^{1/2} \|v\|_{LE^1}^{1/2} + \|v\|_{L_t^2 L_x^2} \right) + \lambda^{-1/2} \|v\|_{LE^1}.$$

This establishes an estimate on the high frequencies. We must add in the lower frequencies to the left-hand side. That is, we must add $\|v_{<\lambda}\|_{LE_{<2R_0}^1}$ to both sides of (2.27).

Step 5a: Lower frequencies—further frequency splitting and bounding the low-low term. First, we get the bound

$$\left\| \langle x \rangle^{-1} v_{<\lambda} \right\|_{LE} \lesssim \|v\|_{L_t^2 L_x^2}$$

via Plancherel's theorem. For the term $\|\partial v_{<\lambda}\|_{LE}$, we write

$$v_{<\lambda} = v_{<>\sigma\lambda} + v_{<<\sigma\lambda},$$

where

$$\begin{aligned} v_{<>\sigma\lambda} &= \chi_{|\xi|<\lambda}(D_x) \chi_{|\tau|>\sigma\lambda}(D_t) v, \\ v_{<<\sigma\lambda} &= \chi_{|\xi|<\lambda}(D_x) \chi_{|\tau|<\sigma\lambda}(D_t) v, \end{aligned}$$

and $\sigma \gg 1$ will be chosen later (and does not denote the same σ as used in the construction of the escape function). Applying Plancherel's theorem, frequency localization, and the compact support of v again yields

$$\|\partial v_{<<\sigma\lambda}\|_{LE} \lesssim \|(|\tau| + |\xi|) \chi_{|\xi|<\lambda} \chi_{|\tau|<\sigma\lambda} \hat{v}\|_{L_\tau^2 L_\xi^2} \lesssim \sigma\lambda \|v\|_{L_t^2 L_x^2}.$$

Step 5b: Lower frequencies—bounding the low-high term. For $v_{<>\sigma\lambda}$, we compute that

$$(2.28) \quad \|\partial v_{<>\sigma\lambda}\|_{LE} \lesssim \|(|\tau| + |\xi|) \chi_{|\xi|<\lambda} \chi_{|\tau|>\sigma\lambda} \hat{v}\|_{L_\tau^2 L_\xi^2} \lesssim \lambda \|v\|_{L_t^2 L_x^2} + (\sigma\lambda)^{-1} \|(\partial_t^2 v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2}.$$

For the last term on the right, we utilize the expression for Pv to write

$$(2.29) \quad \begin{aligned} \|(\partial_t^2 v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} &\lesssim \|(Pv)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} + \|((g^{0j} D_j + D_j g^{0j}) D_t v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \\ &\quad + \|(D_i g^{ij} D_j v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} + \|(a D_t v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2}. \end{aligned}$$

One can readily check that

$$(2.30) \quad \|(Pv)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \lesssim \|Pv\|_{LE^*}$$

and

$$(2.31) \quad \|(a D_t v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \lesssim \|\partial v\|_{LE}.$$

For the other terms, we note that as *functions*, one has that $g^{\alpha j}, D_j g^{\alpha j} \in S^0$ for all $\alpha \in \{0, 1, 2, 3\}$ and $j \in \{1, 2, 3\}$, and so

$$[\chi_{|\xi|<\lambda}(D_x) \chi_{|\tau|>\sigma\lambda}(D_t), g^{\alpha j}] \in \Psi^{-1}, \quad [\chi_{|\xi|<\lambda}(D_x) \chi_{|\tau|>\sigma\lambda}(D_t), D_j g^{\alpha j}] \in \Psi^{-1}.$$

In particular, the above two operators are bounded on $L_t^2 L_x^2$. Pairing this with the fact that Fourier multipliers commute, we have that

$$(2.32) \quad \begin{aligned} \|((g^{0j} D_j + D_j g^{0j}) D_t v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} &\lesssim \|(D_j g^{0j}) (D_t v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \\ &\quad + \|g^{0j} (D_j D_t v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \\ &\quad + \|[\chi_{|\xi|<\lambda}(D_x) \chi_{|\tau|>\sigma\lambda}(D_t), (D_j g^{0j})] D_t v\|_{L_t^2 L_x^2} \\ &\quad + \|[\chi_{|\xi|<\lambda}(D_x) \chi_{|\tau|>\sigma\lambda}(D_t), g^{0j}] D_j D_t v\|_{L_t^2 L_x^2} \\ &\lesssim \lambda \|\partial v\|_{LE} + C(\lambda) \|v\|_{L_t^2 L_x^2} \end{aligned}$$

and

$$\begin{aligned}
 (2.33) \quad & \|(D_i g^{ij} D_j v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \lesssim \|(D_i g^{ij})(D_j v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} + \|g^{ij}(D_i D_j v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \\
 & \quad + \|([\chi_{|\xi|<\lambda}(D_x)\chi_{|\tau|>\sigma\lambda}(D_t), (D_i g^{ij})](D_j v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \\
 & \quad + \|([\chi_{|\xi|<\lambda}(D_x)\chi_{|\tau|\geq\sigma\lambda}(D_t), g^{ij}](D_i D_j v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \\
 & \lesssim C(\lambda) \|v\|_{L_t^2 L_x^2}.
 \end{aligned}$$

Applying (2.30)–(2.33) to (2.29) gives that

$$\|(\partial_t^2 v)_{<>\sigma\lambda}\|_{L_t^2 L_x^2} \lesssim C(\lambda) \|v\|_{L_t^2 L_x^2} + \lambda \|\partial v\|_{LE} + \|Pv\|_{LE^*}.$$

Plugging the resulting estimate into (2.28) implies that

$$\|\partial v_{<>\sigma\lambda}\|_{LE} \lesssim C(\lambda) \|v\|_{L_t^2 L_x^2} + (\sigma\lambda)^{-1} \|Pv\|_{LE^*} + \sigma^{-1} \|\partial v\|_{LE}.$$

Step 5c: Lower-frequency—combining all lower frequency contributions. Thus, the full low frequency contribution yields

$$\begin{aligned}
 (2.34) \quad & \|\partial v_{<\lambda}\|_{LE} \lesssim \max\{C(\lambda), \sigma\lambda\} \|v\|_{L_t^2 L_x^2} + (\sigma\lambda)^{-1} \|Pv\|_{LE^*} + \sigma^{-1} \|\partial v\|_{LE} \\
 & \lesssim \max\{C(\lambda), \sigma\lambda\} \|v\|_{L_t^2 L_x^2} + (\sigma\lambda)^{-1} \|Pv\|_{LE^*} + \sigma^{-1} \|v\|_{LE^1}.
 \end{aligned}$$

Step 6: Combining the high and lower-frequency bounds. Now, we can combine the high frequency work (2.27) with the low frequency work (2.34) and apply Young's inequality for products with parameter $\delta > 0$ to obtain that

$$\begin{aligned}
 \|v\|_{LE^1_{<2R_0}} & \lesssim C(\lambda) \|Pv\|_{LE^*}^{1/2} \|v\|_{LE^1}^{1/2} + \max\{C(\lambda), \sigma\lambda\} \|v\|_{L_t^2 L_x^2} + (\sigma\lambda)^{-1} \|Pv\|_{LE^*} \\
 & \quad + \left(\sigma^{-1} + \lambda^{-1/2}\right) \|v\|_{LE^1} \\
 & \lesssim \max\{C(\lambda), \sigma\lambda\} \|v\|_{L_t^2 L_x^2} + \left([C(\lambda)]^2 \delta^{-1} + (\sigma\lambda)^{-1}\right) \|Pv\|_{LE^*} \\
 & \quad + \left(\delta + \sigma^{-1} + \lambda^{-1/2}\right) \|v\|_{LE^1}.
 \end{aligned}$$

Due to the support of v in x , we know that $\|v\|_{LE^1_{<2R_0}} = \|v\|_{LE^1}$. Picking δ sufficiently small and λ, σ sufficiently large (all of which will depend on R_0) allows us to absorb the $\|v\|_{LE^1}$ term on the right-hand side into the left-hand side, providing (2.13) and completing the proof. \square

3. Local energy decay. In this section, we explain how recovering the high frequency estimate in [MST20] allows us to establish local energy decay by appealing to existing estimates in their work (which makes it easier to perform time-frequency localization). By an extension argument outlined in section 3.3, it is sufficient to reduce to the case of Schwartz functions, which allows for the removal of data terms. The simplified version of local energy decay for Schwartz functions can be readily proven by combining the proven high frequency estimates with appropriate medium and low frequency estimates, then utilizing a time-frequency partition of unity. The medium and low frequency estimates that we require come from the work in [MST20] and do not depend on the trapping nor the damping. We remark that such analyses are also *independent* of the stationarity of P .

3.1. Medium frequencies. The goal of the medium frequency estimate is to establish a weighted estimate which implies local energy decay for solutions supported at any range of time frequencies bounded away from both zero and infinity. This is rooted in the notion of a *Carleman estimate*, which is weighted $L_t^2 L_x^2$ estimates where the weight is pseudoconvex. The constants in our inequalities will depend on the parameter \mathbf{c} introduced in section 1.2, but they will (and must) be independent of the parameters in φ ; the Carleman weights for our estimates are radial. The Carleman weights which we use here are constructed in, e.g., [Boo18], [KT01].

The main medium frequency estimate is the following, and the corresponding theorem in [MST20] is Theorem 5.4.

THEOREM 3.1. *Let P be an asymptotically flat damped wave operator, and suppose that ∂_t is uniformly time-like. Then, for any $\delta > 0$, there exists a bounded, nondecreasing radial weight $\varphi = \varphi(\ln(1+r))$ so that for all $u \in \mathcal{S}(\mathbb{R}^4)$, we have the bound*

$$(3.1) \quad \begin{aligned} & \left\| (1 + \varphi''_+)^{1/2} e^\varphi (\nabla u, \langle r \rangle^{-1} (1 + \varphi') u) \right\|_{LE} + \left\| (1 + \varphi')^{1/2} e^\varphi \partial_t u \right\|_{LE} \\ & \lesssim \|e^\varphi P u\|_{LE^*} + \delta \left(\left\| (1 + \varphi')^{1/2} e^\varphi u \right\|_{LE} + \left\| \langle r \rangle^{-1} (1 + \varphi''_+)^{1/2} (1 + \varphi') e^\varphi \partial_t u \right\|_{LE} \right). \end{aligned}$$

Remark 3.2. We will justify *why* this is an appropriate estimate on the medium frequencies. To that end, suppose that u is supported at time frequencies τ such that $0 < \tau_0 \leq |\tau| \leq \tau_1$, where $\tau_0 < \tau_1$. For compatibility with the other frequency regimes, we will want $\tau_0 \ll 1 \ll \tau_1$. Plancherel's theorem yields that

$$\delta \left\| (1 + \varphi')^{1/2} e^\varphi u \right\|_{LE} \lesssim \frac{\delta}{\tau_0} \left\| (1 + \varphi')^{1/2} e^\varphi \partial_t u \right\|_{LE},$$

while

$$\delta \left\| \langle r \rangle^{-1} (1 + \varphi''_+)^{1/2} (1 + \varphi') e^\varphi \partial_t u \right\|_{LE} \lesssim \delta \tau_1 \left\| \langle r \rangle^{-1} (1 + \varphi''_+)^{1/2} (1 + \varphi') e^\varphi u \right\|_{LE}.$$

By choosing δ sufficiently small, both terms absorb into the left-hand side of (3.1) in a direct fashion. We can translate our work immediately into a local energy decay estimate for u , with an implicit constant which depends on φ . Notice that since δ can be chosen arbitrarily, (3.1) allows for any interval of frequencies bounded away from both zero and infinity.

The proof of this theorem is broken up into two Carleman estimates, one that applies within a large compact set and one that applies outside of this compact set. Within the compact set, the damping term is well-signed and readily absorbable as a perturbation due to the conditions on the weight φ . Here, the weight will be convex. Outside of the compact set, the damping is zero, so the proof in [MST20] follows through without any modification. In this region, one desires to use Proposition 2.20, which requires a constant weight. To that end, one breaks up the exterior into three regions: one where the Carleman weight is convex, a transition region where the conditions break in order to bend the weight to be constant near infinity, and a region near infinity where the weight is constant.

In both regions (the compact set and its exterior), the work in [MST20] allows for more general (e.g., unsigned) lower-order terms than a damping term. The proofs of these estimates within the aforementioned regions are based on positive commutator arguments utilizing the self- and skew-adjoint parts of the conjugated operator $P_\varphi = e^\varphi P e^{-\varphi}$. The work is then combined using a cutoff argument.

3.2. Low frequencies.

We will set

$$P_0 := P|_{D_t=0} = D_i g^{ij} D_j.$$

This represents P at time frequency zero, and one can utilize it to obtain information in a neighborhood of this frequency. Since ∂_t is uniformly time-like, P_0 is uniformly elliptic. The operator P_0 is a special case of that found in [MST20], so all of the results in their work apply here with almost no modification. Notice that the damping does not arise in P_0 .

At low frequencies, the obstruction to local energy decay arises when P has a resonance at frequency zero.

DEFINITION 3.3. *A function u is called a zero resonant state for P if $u \in \mathcal{LE}_0$ is nonzero and $P_0 u = 0$. If, in addition, $u \in L^2$, then we call u a zero eigenfunction.*

Here, the space \mathcal{LE} is a variant of the LE space where there is no time dependence (i.e., the time is fixed, and there is no time derivative arising in the norm), and \mathcal{LE}_0 is the closure of C_c^∞ in the \mathcal{LE} norm.

For a general wave operator P , such resonant states are annihilated by P while having finite energy. However, they also possess an infinite LE^1 norm when integrating in t over $[0, \infty)$, which violates local energy decay. Such states are ruled out in the context of this paper due to the uniform ellipticity of P_0 . A quantitative condition on the existence of such resonant states is as follows.

DEFINITION 3.4. *P is said to satisfy a zero resolvent bound/zero nonresonance condition if there exists some K_0 , independent of t , such that*

$$(3.2) \quad \|u\|_{\dot{H}^1} \leq K_0 \|P_0 u\|_{\dot{H}^{-1}} \quad \forall u \in \dot{H}^1.$$

Proposition 2.10 of [MST20] demonstrates that a stationary wave operator P has no zero resonant states/zero eigenfunctions if and only if the zero nonresonance condition holds. In our problem, this condition is satisfied due to the uniform ellipticity of P_0 .

The relevant low frequency estimate is the following, and the corresponding theorem in [MST20] is Theorem 6.1.

THEOREM 3.5. *Let P be an asymptotically flat damped wave operator, and suppose that ∂_t is uniformly time-like. Then,*

$$(3.3) \quad \|u\|_{LE^1} \lesssim \|\partial_t u\|_{LE_c^1} + \|Pu\|_{LE^*}$$

for all $u \in \mathcal{S}(\mathbb{R}^4)$.

Remark 3.6. The error term $\|\partial_t u\|_{LE_c^1}$ has the unfortunate effect of requiring information on the size of first-order derivatives of $\partial_t u$. However, this estimate will only be used when the time frequency is close to zero, in which case this term will be absorbable into the left-hand side of the inequality. Indeed, if we consider $u \in \mathcal{S}(\mathbb{R}^4)$ with frequency support $0 \leq |\tau| \leq \tau_0 \ll 1$, then we may apply Plancherel's theorem to obtain that

$$\|\partial_t u\|_{LE_c^1} \lesssim \tau_0 \|u\|_{LE_c^1}.$$

If τ_0 is sufficiently small, then we may absorb this term into the lower-bound side of (3.3) to obtain local energy decay for such u .

The proof leverages weighted elliptic estimates for the flat Laplacian Δ in order to get similar estimates for AF perturbations. Once again, the damping does not play a harmful (or even meaningful) role. At frequency zero, it provides no contribution, and near frequency zero, it can be readily absorbed by the error term in (3.3).

3.3. Establishing local energy decay. Now, we discuss the second main theorem, local energy decay. First, the authors in [MST20] show that it is sufficient to remove the Cauchy data at times 0 and T . This makes it significantly easier to perform frequency localization. The corresponding theorem in [MST20] is Theorem 7.1.

THEOREM 3.7. *Let P be a stationary, asymptotically flat damped wave operator satisfying the geometric control condition (2.1), and suppose that ∂_t is uniformly timelike. Then, the estimate*

$$(3.4) \quad \|u\|_{LE^1} \lesssim \|Pu\|_{LE^*}$$

holds for all $u \in \mathcal{S}(\mathbb{R}^4)$.

In order to prove Theorem 3.7, one splits u into its low, medium, and high frequency parts using a time-frequency partition of unity. In each relevant frequency regime, one applies the corresponding frequency estimate (as in Remarks 1.6, 3.2, and 3.6), and then sums them together. For the medium frequency estimate to be compatible with the low and high frequency regimes, one needs that it applies to any range of time frequencies bounded away from both zero (compatibility with low) and infinity (compatibility with high), which is the utility of the δ parameter in Theorem 3.1. The commutators of the time-frequency cutoffs and P are zero since P is stationary.

As in section 7 of [MST20], one proves that Theorem 3.7 implies Theorem 1.9 by fixing u and constructing a function v which matches the Cauchy data of u at times 0 and T (and satisfies an appropriate bound) which allows one to apply (3.4) to $u - v$. This construction is performed using a partition of unity on the support of $u[0]$, $u[T]$, and Pu . In particular, one splits into an interior region $\{|x| < 4R_0\}$ and an exterior region $\{|x| > 2R_0\}$. The damping is nonproblematic in the interior (here, one uses the uniform energy bounds) and is zero in the exterior. It is important to highlight the latter fact since the authors use a time reversal symmetry argument in the exterior region, and time reversal turns the damping into a driving force (and hence a harmful term). However, our damping is zero in the exterior region, rendering such an argument nonproblematic by choosing an appropriate small AF perturbation of \square which matches P in the exterior (where the damping is zero). In the context of [MST20], this provides the “two point” local energy estimate

$$\|u\|_{LE^1[0,T]} + \|\partial u\|_{L_t^\infty L_x^2[0,T]} \lesssim \|\partial u(0)\|_{L^2} + \|\partial u(T)\|_{L^2} + \|Pu\|_{LE^* + L_t^1 L_x^2[0,T]}.$$

In view of Corollary 2.19, this implies local energy decay.

Acknowledgments. This work was performed as a portion of the Ph.D. dissertation of the author at the University of North Carolina at Chapel Hill (see [Kof22]). The author would like to thank his advisor, Jason Metcalfe, for suggesting this problem, engaging in countless helpful conversations, and carefully reading multiple drafts of this work. He would also like to thank Yaiza Canzani, Hans Christianson, Jeremy Marzuola, and both referees of this article for providing many beneficial comments and references.

REFERENCES

- [Ali06] S. ALINHAC, *On the Morawetz-Keel-Smith-Sogge inequality for the wave equation on a curved background*, Publ. Res. Inst. Math. Sci., 42 (2006), pp. 705–720.
- [BH10] J. F. BONY AND D. HÄFNER, *The semilinear wave equation on asymptotically Euclidean manifolds*, Comm. Partial Differential Equations, 35 (2010), pp. 23–67.
- [BLR92] C. BARDOS, G. LEBEAU, AND J. RAUCH, *Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary*, SIAM J. Control Optim., 30 (1992), pp. 1024–1065.
- [Boo18] R. BOOTH, *An Investigation of Non-Trapping, Asymptotically Euclidean Wave Equations*, Ph.D. dissertation, University of North Carolina at Chapel Hill, 2018.
- [BR14] J. M. BOUCLET AND J. ROYER, *Local energy decay for the damped wave equation*, J. Funct. Anal., 266 (2014), pp. 4538–4615.
- [BT07] J. M. BOUCLET AND N. TZVETKOV, *Strichartz estimates for long range perturbations*, Amer. J. Math., 129 (2007), pp. 1565–1609.
- [BT08] J. M. BOUCLET AND N. TZVETKOV, *On global Strichartz estimates for non trapping metrics*, J. Funct. Anal., 254 (2008), pp. 1661–1682.
- [Bur98] N. BURQ, *Décroissance de l'énergie locale de l'équation des ondes pour le problème extérieur et absence de résonance au voisinage du réel*, Acta Math., 180 (1998), pp. 1–29.
- [Chr08] H. CHRISTIANSON, *Dispersive estimates for manifolds with one trapped orbit*, Comm. Partial Differential Equations, 33 (2008), pp. 1147–1174.
- [Doi96] S. DOI, *Remarks on the Cauchy problem for Schrödinger-type equations*, Comm. Partial Differential Equations, 21 (1996), pp. 163–178.
- [Hin22] P. HINTZ, *A sharp version of Price's law for wave decay on asymptotically at spacetimes*, Comm. Math. Phys., 389 (2022), pp. 491–542.
- [Ika82] M. IKAWA, *Decay of solutions of the wave equation in the exterior of two convex obstacles*, Osaka Math. J., 19 (1982), pp. 459–509.
- [Ika88] M. IKAWA, *Decay of solutions of the wave equation in the exterior of several convex bodies*, Ann. Inst. Fourier (Grenoble), 38 (1988), pp. 113–146.
- [JSS90] J. L. JOURNÉ, A. SOFFER, AND C. G. SOGGE, *$L^p \rightarrow L^{p'}$ estimates for time dependent Schrödinger operators*, Bull. Amer. Math. Soc. (N.S.), 23 (1990), pp. 519–524.
- [JSS91] J. L. JOURNÉ, A. SOFFER, AND C. G. SOGGE, *Decay estimates for Schrödinger operators*, Comm. Pure Appl. Math., 44 (1991), pp. 573–604.
- [Kof22] C. KOFROTH, *Local Energy Decay for Damped Waves on Stationary, Asymptotically at Space-Times*, Ph.D. dissertation, University of North Carolina at Chapel Hill, 2022.
- [KSS02] M. KEEL, H. F. SMITH, AND C. D. SOGGE, *Almost global existence for some semilinear wave equations*, J. Anal. Math., 87 (2002), pp. 265–279.
- [KSS04] M. KEEL, H. F. SMITH, AND C. D. SOGGE, *Almost global existence for quasilinear wave equations in three space dimensions*, J. Amer. Math. Soc., 17 (2004), pp. 109–153.
- [KT01] H. KOCH AND D. TATARU, *Carleman estimates and unique continuation for second-order elliptic equations with nonsmooth coefficients*, Comm. Pure Appl. Math., 54 (2001), pp. 339–360.
- [Leb96] G. LEBEAU, *Équation des ondes amorties*, in Algebraic and Geometric Methods in Mathematical Physics (Kaciveli, 1993), 1996, pp. 73–109.
- [Loo22] S. Z. LOOI, *Pointwise decay for the wave equation on nonstationary spacetimes*, J. Math. Anal. Appl., (2022), 126939.
- [MMT08] J. MARZUOLA, J. METCALFE, AND D. TATARU, *Strichartz estimates and local smoothing estimates for asymptotically at Schrödinger equations*, J. Funct. Anal., 255 (2008), pp. 1497–1553.
- [MMTT10] J. MARZUOLA, J. METCALFE, D. TATARU, AND M. TOHANEANU, *Strichartz estimates on Schwarzschild black hole backgrounds*, Comm. Math. Phys., 293 (2010), pp. 37–83.
- [Mor20] K. MORGAN, *The Effect of Metric Behavior at Spatial Infinity on Pointwise Wave Decay in the Asymptotically at Stationary Setting*, preprint, <https://arxiv.org/abs/2006.11324>, 2020.
- [Mor66] C. S. MORAWETZ, *Exponential decay of solutions of the wave equation*, Comm. Pure Appl. Math., 19 (1966), pp. 439–444.
- [Mor68] C. S. MORAWETZ, *Time decay for the nonlinear Klein-Gordon equations*, Proc. A, 306 (1968), pp. 291–296.
- [Mor75] C. S. MORAWETZ, *Decay for solutions of the exterior problem for the wave equation*, Comm. Pure Appl. Math., 28 (1975), pp. 229–264.
- [MRS77] C. S. MORAWETZ, J. V. RALSTON, AND W. A. STRAUSS, *Decay of solutions of the wave equation outside nontrapping obstacles*, Comm. Pure Appl. Math., 30 (1977), pp. 447–508.

- [MS06] J. METCALFE AND C. D. SOGGE, *Long-time existence of quasilinear wave equations exterior to star-shaped obstacles via energy methods*, SIAM J. Math. Anal., 38 (2006), pp. 188–209.
- [MS07] J. METCALFE AND C. D. SOGGE, *Global existence of null-form wave equations in exterior domains*, Math. Z., 256 (2007), pp. 521–549.
- [MST20] J. METCALFE, J. STERBENZ, AND D. TATARU, *Local energy decay for scalar fields on time dependent non-trapping backgrounds*, Amer. J. Math., 142 (2020), pp. 821–883.
- [MT09] J. METCALFE AND D. TATARU, *Decay estimates for variable coefficient wave equations in exterior domains*, in *Advances in Phase Space Analysis of Partial Differential Equations*, Springer, New York, 2009, pp. 201–216.
- [MT12] J. METCALFE AND D. TATARU, *Global parametrices and dispersive estimates for variable coefficient wave equations*, Math. Ann., 353 (2012), pp. 1183–1237.
- [MTT12] J. METCALFE, D. TATARU, AND M. TOHANEANU, *Price’s law on nonstationary space-times*, Adv. Math., 230 (2012), pp. 995–1028.
- [MW21] K. MORGAN AND J. WUNSCH, *Generalized Price’s Law on Fractional-Order Asymptotically at Stationary Spacetimes*, preprint, <https://arxiv.org/abs/2105.02305>, 2021.
- [NZ09] S. NONNENMACHER AND M. ZWORSKI, *Quantum decay rates in chaotic scattering*, Acta Math., 203 (2009), pp. 149–233.
- [Ral69] J. V. RALSTON, *Solutions of the wave equation with localized energy*, Comm. Pure Appl. Math., 22 (1969), pp. 807–823.
- [Roy18] J. ROYER, *Local decay for the damped wave equation in the energy space*, J. Inst. Math. Jussieu, 17 (2018), pp. 509–540.
- [RT74] J. RAUCH AND M. E. TAYLOR, *Exponential decay of solutions to hyperbolic equations in bounded domains*, Indiana Univ. Math. J., 24 (1974), pp. 79–86.
- [Sbi15] J. SBIERSKI, *Characterisation of the energy of Gaussian beams on Lorentzian manifolds: With applications to black hole spacetimes*, Anal. PDE, 8 (2015), pp. 1379–1420.
- [Ste05] J. STERBENZ, *Angular regularity and Strichartz estimates for the wave equation*, Int. Math. Res. Not., 4 (2005), pp. 187–231, With an appendix by Igor Rodnianski.
- [SW10] C. D. SOGGE AND C. WANG, *Concerning the wave equation on asymptotically Euclidean manifolds*, J. Anal. Math., 112 (2010), pp. 1–32.
- [Tat08] D. TATARU, *Parametrices and dispersive estimates for Schrödinger operators with variable coefficients*, Amer. J. Math., 130 (2008), pp. 571–634.
- [Tat13] D. TATARU, *Local decay of waves on asymptotically at stationary space-times*, Amer. J. Math., 135 (2013), pp. 361–401.
- [Toh12] M. TOHANEANU, *Strichartz estimates on Kerr black hole backgrounds*, Trans. Amer. Math. Soc., 364 (2012), pp. 689–702.
- [WZ11] J. WUNSCH AND M. ZWORSKI, *Resolvent estimates for normally hyperbolic trapped sets*, Ann. Henri Poincaré, 12 (2011), pp. 1349–1385.