

Subsets of rectifiable curves in Banach spaces I: Sharp exponents in traveling salesman theorems

Matthew Badger and Sean McCurdy

.....

Abstract The analyst’s traveling salesman problem (TSP) is to find a characterization of subsets of rectifiable curves in a metric space. This problem was introduced and solved in the plane by Jones in 1990 and subsequently solved in higher-dimensional Euclidean spaces by Okikiolu in 1992 and in the infinite-dimensional Hilbert space ℓ_2 by Schul in 2007. In this paper, we establish sharp extensions of Schul’s necessary and sufficient conditions for a bounded set $E \subset \ell_p$ to be contained in a rectifiable curve from $p = 2$ to $1 < p < \infty$. While the necessary and sufficient conditions coincide when $p = 2$, we demonstrate that there is a strict gap between the necessary condition and sufficient condition when $p \neq 2$. We also identify and correct technical errors in the proof by Schul. This investigation is partly motivated by recent work of Edelen, Naber, and Valtorta on Reifenberg-type theorems in Banach spaces and complements work of Hahlomaa and recent work of David and Schul on the analyst’s TSP in general metric spaces.

Contents

1. Introduction	1
2. Modulus of smoothness and proof of the sufficient conditions	12
3. Modulus of convexity and proof of the necessary conditions	25
4. Analyst’s TSP in finite-dimensional Banach spaces	43
5. Sharpness of the exponents via examples	46
Appendix. Schul’s prefiltration lemma in a metric space	62
References	68

1. Introduction

Given a set in a path-connected metric space, we may ask whether or not the set is contained in a curve of finite length (also called a *rectifiable curve*), and if so, ask how to find a curve containing the set that is (essentially) as short as possible. This problem was introduced and solved in the Euclidean plane by Jones [48] and is now commonly known as the *analyst’s traveling salesman problem*. While it is immediate that a set contained in a rectifiable curve is necessarily bounded and has finite 1-dimensional Hausdorff measure $\mathcal{H}^1$, this pair of conditions is not sufficient. To decide when a set is

Illinois Journal of Mathematics

Advance publication—final volume, issue, and page numbers to be assigned.

DOI 10.1215/00192082-10592363, © 2023 by the University of Illinois Urbana-Champaign

Received February 28, 2020. Received in final form January 26, 2023.

2020 Mathematics Subject Classification: Primary 28A75; Secondary 26A16, 28A80, 46B20, 65D10.

M. Badger was partially supported by NSF DMS grants 1650546 and 2154047.

contained in a rectifiable curve requires additional information about the local geometry of the set within the space. Full solutions to the analyst's traveling salesman problem (TSP) are currently available in $\mathbb{R}^n$ [58], Carnot groups [53], infinite-dimensional Hilbert space ℓ_2 [65], graph inverse limit spaces [33], and for Radon measures in $\mathbb{R}^n$ [17] and Carnot groups [13]; partial solutions are available in general metrics spaces [32, 43] as well as for higher-dimensional curves [16, 20] and surfaces [7, 10, 36, 38, 41, 47, 67]. Beyond the intrinsic interest of the analyst's TSP in metric geometry, finding tests to determine when a set is contained in a rectifiable curve or "nice" surface has led to applications in complex analysis, dynamics and probability, geometric analysis, and harmonic analysis. For a sample of applications, see [6, 8, 9, 11, 15, 18, 19, 25, 26, 27, 42, 55, 56, 57, 66].

In this paper, we establish sharp extensions of Schul's necessary and sufficient conditions for a bounded set $E \subset \ell_p$ to be contained in a rectifiable curve from $p = 2$ to $1 < p < \infty$ (see Theorems 1.6 and 1.7). As usual, ℓ_p denotes the (real) Banach space of p -summable sequences,

$$x = (x_1, x_2, \dots) \in \ell_p \quad \text{if and only if} \\ x_1, x_2, \dots \in \mathbb{R} \text{ and } |x|_p = \left(\sum_1^\infty |x_i|^p \right)^{1/p} < \infty.$$

While the necessary and sufficient conditions in Schul's theorem coincide when $p = 2$, we demonstrate that there is a strict gap between the necessary condition and sufficient condition when $p \neq 2$. En route, we prove that the classes of rectifiable curves in the infinite-dimensional spaces ℓ_p and ℓ_q differ when $p \neq q$.

PROPOSITION 1.1

Let $1 < p < \infty$. Every rectifiable curve in ℓ_p is a rectifiable curve in ℓ_q for all $q \geq p$. However, there exists a curve Γ in ℓ_p such that Γ is rectifiable (that is, $\mathcal{H}^1(\Gamma) < \infty$) in ℓ_q for every $q > p$, but Γ is not rectifiable (that is, $\mathcal{H}^1(\Gamma) = \infty$) in ℓ_p .

Proposition 1.1, Theorem 1.6, and Theorem 1.7 capture a special infinite-dimensional phenomenon. In particular, they imply that a solution of the analyst's TSP in ℓ_p cannot be neatly derived from the solution in ℓ_2 . By contrast, bi-Lipschitz equivalence of finite-dimensional Banach spaces ensures that a set in $\mathbb{R}^n$ is (a subset of) a rectifiable curve independent from the choice of underlying norm, even though the actual length of the curve depends on the norm. This paper serves to clarify the difference between the finite and infinite-dimensional settings. An essential reason for us to study (subsets of) rectifiable curves in ℓ_p for $1 < p < \infty$ is that the spaces interpolate between ℓ_2 , where the analyst's TSP is solved, and ℓ_∞ , which contains an isometric copy of any separable metric space. Thus, a resolution of the analyst's TSP in ℓ_p may provide insight into the analyst's TSP in general metric spaces. For further discussion and description of related research, see Section 1.3.

1.1. Analyst's TSP in Euclidean space and ℓ_2

To solve the analyst's TSP in the plane, Jones introduced unilateral linear approximation numbers, now universally called *Jones' beta numbers*, which measure how “flat” a set is in a given window. The Jones' beta numbers are well-defined in any Banach space. Let $\mathbb{X}$ be a Banach space,¹ let $E \subset \mathbb{X}$ be a nonempty set, and let $Q \subset \mathbb{X}$ be a set of positive diameter. If $E \cap Q \neq \emptyset$, define

$$(1.1) \quad \beta_E(Q) = \inf_L \sup_{x \in E \cap Q} \frac{\text{dist}(x, L)}{\text{diam } Q} \in [0, 1],$$

where the infimum ranges over all 1-dimensional affine subspaces (lines) $L \subset \mathbb{X}$; and, if $E \cap Q = \emptyset$, define $\beta_E(Q) = 0$. At one extreme, if $\beta_E(Q) = 0$, then $E \cap Q$ is contained in a line. At the other extreme, if $\beta_E(Q) \gtrsim 1$, then the set $E \cap Q$ is uniformly far away from every line passing through Q . From the definition, it immediately follows that

$$(1.2) \quad \beta_E(R) \leq \frac{\text{diam } Q}{\text{diam } R} \beta_F(Q) \quad \text{for all } E \subset F \text{ and } R \subset Q.$$

In view of the fact that rectifiable curves (having parameterizations of bounded variation) admit tangent lines $\mathcal{H}^1$ -a.e., one may expect that sets contained in a finite length curve have “vanishing beta numbers” at typical points of those sets. The *analyst's traveling salesman theorem* makes this idea precise and provides a characterization of subsets of rectifiable curves with an estimate on the shortest length of a curve containing the set as follows.

THEOREM 1.2 (Jones [48] in $\mathbb{R}^2$; Okikiolu [58] in $\mathbb{R}^n$)

Let $n \geq 2$ and let $E \subset \mathbb{R}^n$. Then E is contained in a rectifiable curve if and only if

$$(1.3) \quad S_E(\mathbb{R}^n) := \text{diam } E + \sum_{Q \in \Delta(\mathbb{R}^n)} \beta_E(3Q)^2 \text{diam } Q < \infty,$$

where the sum ranges over all dyadic cubes Q in $\mathbb{R}^n$ and $3Q$ denotes the concentric dilate of the cube Q with scaling factor 3. More precisely, if $S_E(\mathbb{R}^n) < \infty$, then E is contained in a curve Γ in $\mathbb{R}^n$ with

$$(1.4) \quad \mathcal{H}^1(\Gamma) \lesssim_n S_E(\mathbb{R}^n).$$

If $\Sigma \subset \mathbb{R}^n$ is a connected set, then

$$(1.5) \quad S_\Sigma(\mathbb{R}^n) \lesssim_n \mathcal{H}^1(\Sigma).$$

The constant 3 in (1.3) can be replaced with any constant $A > 1$. Then (1.4) and (1.5) hold with implicit constants depending on n and A .

For refinements of (1.4) and (1.5) for rectifiable Jordan arcs in $\mathbb{R}^n$ and in Hilbert space, see [24] and [52]. Time complexity of the analyst's traveling salesman algorithm for constructing the rectifiable curve in (1.4) is investigated in [62].

1. All Banach spaces in this paper are real Banach spaces of dimension at least 2.

REMARK 1.3

If E is a subset of a rectifiable curve in $\mathbb{R}^n$, then the analyst's traveling salesman theorem ensures that

$$(1.6) \quad \sum_{\substack{Q \in \Delta(\mathbb{R}^n) \\ \beta_E(3Q) \geq \epsilon}} \text{diam } Q \leq \epsilon^{-2} S(E) < \infty \quad \text{for all } \epsilon > 0.$$

It follows that

$$(1.7) \quad \lim_{\substack{Q \in \Delta(\mathbb{R}^n) \\ Q \downarrow x}} \beta_E(3Q) = 0 \quad \text{at } \mathcal{H}^1\text{-a.e. } x \in E,$$

or equivalently by (1.2),

$$(1.8) \quad \lim_{r \downarrow 0} \beta_E(B(x, r)) = 0 \quad \text{at } \mathcal{H}^1\text{-a.e. } x \in E.$$

Thus, subsets of rectifiable curves in $\mathbb{R}^n$ have “vanishing beta numbers” at typical points in the sense of (1.8). It is possible to construct examples of generalized von Koch snowflake curves (with carefully chosen angles) to show that (1.8) is also satisfied by certain curves of infinite length (Proposition 5.1). By contrast, the analyst's traveling salesman theorem guarantees that every curve Γ in $\mathbb{R}^n$ of infinite length satisfies $S_\Gamma(\mathbb{R}^n) = \infty$. In other words, the finiteness of $S_\Gamma(\mathbb{R}^n)$ is a perfect test to determine rectifiability of a curve Γ in $\mathbb{R}^n$.

REMARK 1.4

Let V be a finite set of points in $\mathbb{R}^n$ (equipped with the Euclidean norm) that is *1-separated* in the sense $|v - w| \geq 1$ for all distinct $v, w \in V$. Assume that L is a line in $\mathbb{R}^n$ such that $\text{dist}(v, L) \leq \beta \ll 1$ for all $v \in V$. Then the set $V = \{v_1, \dots, v_k\}$ may be enumerated according to its orthogonal projection onto L . For simplicity, let us further assume that $v_1, v_k \in \ell$. By the triangle inequality and a simple computation with the Pythagorean theorem (Figure 1),

$$(1.9) \quad |v_1 - v_k| \leq |v_1 - v_2| + \dots + |v_{k-1} - v_k| \leq (1 + C_1 \beta^2) |v_1 - v_k|$$

for some universal constant C_1 . Conversely, if $\text{dist}(v_i, L) \geq \alpha$ for some $1 \leq i \leq k$, then the Pythagorean theorem yields

$$(1.10) \quad |v_1 - v_i| + |v_i - v_k| \geq (1 + C_2 \alpha^2) |v_1 - v_k|$$

for some universal constant C_2 . At a high level, the estimates (1.9) and (1.10) correspond to (1.4) and (1.5) in the analyst's traveling salesman theorem, respectively. Informally, we say that the Pythagorean theorem is responsible for the exponent 2 on $\beta_E(3Q)^2$ in (1.3).

The dependence on the ambient dimension in the implicit constants in (1.4) and (1.5) is ultimately a consequence of using dyadic cubes in the sum $S_E(\mathbb{R}^n)$ to index “all” locations and scales in $\mathbb{R}^n$. By bi-Lipschitz equivalence, Theorem 1.2 persists in every finite-dimensional Banach space with implicit constants that also depend on the norm as well as a choice of coordinates; see Section 4 for a detailed statement. To formulate a




Figure 1. To use the Pythagorean theorem to verify (1.9) and (1.10) (shown with $k = 4$), first draw right triangles formed from line segments $[v_i, v_{i+1}]$ between consecutive points and the line passing through v_1 and v_k .

dimension-independent version of the analyst's traveling salesman theorem in ℓ_2 , Schul [65] replaced $S_E(\mathbb{R}^n)$ with a sum $S_E(\mathcal{G})$ indexed over a multiresolution family $\mathcal{G}$ of "all" locations and scales in the set E .

Let $\mathbb{X}$ be a metric space and let $E \subset \mathbb{X}$ be a nonempty set. For any $\rho > 0$, a ρ -net X_ρ in $\mathbb{X}$ is a set such that $\text{dist}(y, z) \geq \rho$ for all distinct $y, z \in X_\rho$ and $\text{dist}(x, X_\rho) < \rho$ for all $x \in \mathbb{X}$. Following [65], we define a *multiresolution family* $\mathcal{G}$ for E with *inflation factor* $A_{\mathcal{G}} > 1$ to be a collection of closed balls of the form

$$(1.11) \quad \mathcal{G} = \{B(x, A_{\mathcal{G}} 2^{-k}) : x \in X_k, k \in \mathbb{Z}\},$$

where $(X_k)_{k \in \mathbb{Z}}$ is a nested family of 2^{-k} -nets for E . Analogously, if each set X_k has the property $\text{dist}(y, z) \geq 2^{-k}$ for all distinct $y, z \in X_k$, but one or more of the sets X_k are not 2^{-k} -nets, then we call $\mathcal{G}$ a *partial multiresolution family* for E .

For any nonempty set $E \subset \ell_2$ and (partial) multiresolution family $\mathcal{G}$ for E , define

$$(1.12) \quad S_E(\mathcal{G}) := \text{diam } E + \sum_{Q \in \mathcal{G}} \beta_E(Q)^2 \text{diam } Q.$$

THEOREM 1.5 (Schul [65])

If $E \subset \ell_2$ and $S_E(\mathcal{G}) < \infty$ for some multiresolution family $\mathcal{G}$ for E with inflation factor $A_{\mathcal{G}} > 200$, then E is contained in a rectifiable curve Γ in ℓ_2 with

$$(1.13) \quad \mathcal{H}^1(\Gamma) \lesssim_{A_{\mathcal{G}}} S_E(\mathcal{G}).$$

If $\Sigma \subset \ell_2$ is a connected set and $\mathcal{H}$ is a (partial) multiresolution family for Σ with inflation factor $A_{\mathcal{H}} > 1$, then

$$(1.14) \quad S_{\Sigma}(\mathcal{H}) \lesssim_{A_{\mathcal{H}}} \mathcal{H}^1(\Sigma).$$

Note that the implicit constants in (1.13) and (1.14) depend on the inflation factor of the multiresolution family but are otherwise dimension free. Once again, the Pythagorean theorem in ℓ_2 determines the exponent 2 on $\beta_E(Q)^2$ in Theorem 1.5 à la Remark 1.4. More generally, Theorem 1.5 holds in *any* Hilbert space (including nonseparable spaces).

1.2. Schul's theorem in ℓ_p

For any nonempty set $E \subset \ell_p$ and (partial) multiresolution family $\mathcal{G}$ for E , define

$$(1.15) \quad S_{E,r}(\mathcal{G}) := \text{diam } E + \sum_{Q \in \mathcal{G}} \beta_E(Q)^r \text{diam } Q \quad \text{for all } 0 < r < \infty.$$

Note that in the notation of the previous section, $S_{E,2}(\mathcal{G}) \equiv S_E(\mathcal{G})$. Moreover, by (1.2),

$$(1.16) \quad S_{E,r}(\mathcal{G}) \leq S_{F,s}(\widehat{\mathcal{G}}) \quad \text{for all } E \subset F \text{ and } s \leq r,$$

where $\widehat{\mathcal{G}}$ is a multiresolution family for F extending $\mathcal{G}$.

The following pair of theorems, extending Schul's theorem from ℓ_2 to ℓ_p with $1 < p < \infty$, constitutes our main result. We emphasize that when $p \neq 2$, there is a strict gap between the necessary and sufficient conditions for a set to be contained in a rectifiable curve.

THEOREM 1.6 (Sharp sufficient conditions in ℓ_p)

Let $1 < p < \infty$. If $E \subset \ell_p$ and $S_{E,\min(p,2)}(\mathcal{G}) < \infty$ for some multiresolution family $\mathcal{G}$ for E with inflation factor $A_{\mathcal{G}} \geq 240$, then E is contained in a curve Γ in ℓ_p with

$$(1.17) \quad \mathcal{H}^1(\Gamma) \lesssim_{p,A_{\mathcal{G}}} S_{E,\min(p,2)}(\mathcal{G}).$$

The exponent $\min(p, 2)$ on beta numbers in (1.17) is sharp.

THEOREM 1.7 (Sharp necessary conditions in ℓ_p)

Let $1 < p < \infty$. If $\Sigma \subset \ell_p$ is a connected set and $\mathcal{H}$ is a (partial) multiresolution family for Σ with inflation factor $A_{\mathcal{H}} > 1$, then

$$(1.18) \quad S_{\Sigma,\max(2,p)}(\mathcal{H}) \lesssim_{p,A_{\mathcal{H}}} \mathcal{H}^1(\Sigma).$$

The exponent $\max(2, p)$ on beta numbers in (1.18) is sharp.

REMARK 1.8

In this paper, we prove Theorem 1.6 in full and we reduce the proof of Theorem 1.7 to Theorem 3.30; the proof of the latter theorem is deferred to Part II of this paper [14]. More generally, we show that an analogue of Theorem 1.6 holds in uniformly smooth Banach spaces (see Theorem 2.26) and an analogue of Theorem 1.7 holds in uniformly convex spaces (see Theorem 3.22). For example, L^p spaces of p -power integrable functions over a measure space, which include ℓ_p , are uniformly smooth and uniformly convex for all $1 < p < \infty$. In addition, we prove that a universal sufficient condition with exponent 1 is valid in arbitrary Banach spaces (see Theorem 2.10). Each of these results applies to both separable and nonseparable Banach spaces.

An essential feature of ℓ_2 is that the unit ball and induced distance are *rotationally invariant*. In particular, to compute the distance of a point x to a line L in Hilbert space, one may first translate and rotate so that $L = \text{span}(e_1)$ and $x \in \text{span}(e_1, e_2)$ if convenient. In ℓ_p , when $p \neq 2$, rotational-invariance is no longer available, and computation of the distance of a point to a line is sensitive to the position of the line and




Figure 2. (Color online) Unit balls in $\ell_{5/4}^2$ (blue), $\mathbb{R}^2$ (orange), ℓ_5^2 and ℓ_5^3 (green), where ℓ_p^n denotes $(\mathbb{R}^n, |\cdot|_p)$.

geometry of the unit ball. We remark that the gain in complexity witnessed when moving from ℓ_p^2 to ℓ_p^3 (for example, consider the shape of slices of their unit balls; see Figure 2) continues to increase when moving from the finite-dimensional spaces ℓ_p^n to the infinite-dimensional space ℓ_p . For instance, although the norms in ℓ_2^n and ℓ_p^n are $C(n, p)$ -bi-Lipschitz equivalent for each pair n and p , the bi-Lipschitz constant $C(n, p)$ degenerates as $n \rightarrow \infty$ for each $p \neq 2$.

EXAMPLE 1.9

To illustrate the essential idea behind the exponents in Theorems 1.6 and 1.7, let's estimate the length gain in $\ell_5^2 = (\mathbb{R}^2, |\cdot|_5)$ of isosceles triangles

$$\begin{aligned}
 (1.19) \quad T_h : a_h &= (0, 0), & b_h &= (l/2, h), & c_h &= (l, 0), & \text{and} \\
 T_d : a_d &= (0, 0), & b_d &= (2^{-6/5}l - 2^{-1/5}h, 2^{-6/5}l + 2^{-1/5}h), \\
 c_d &= (2^{-1/5}l, 2^{-1/5}l)
 \end{aligned}$$

with horizontal and diagonal bases $\overline{ac}$ of length l and height $\text{dist}(b, \overline{ac}) = h$. On one hand,

$$\begin{aligned}
 (1.20) \quad |\overline{a_h b_h}|_5 + |\overline{b_h c_h}|_5 - |\overline{a_h c_h}|_5 &= 2((l/2)^5 + h^5)^{1/5} - l \\
 &= l((1 + 32(h/l)^5)^{1/5} - 1) \simeq l(h/l)^5
 \end{aligned}$$

for $h \ll l$ by Taylor's theorem for $x \mapsto x^{1/5}$ at $x = 1$. On the other hand,

$$\begin{aligned}
 (1.21) \quad |\overline{a_d b_d}|_5 + |\overline{b_d c_d}|_5 - |\overline{a_d c_d}|_5 &= 2((2^{-6/5}l - 2^{-1/5}h)^5 + (2^{-6/5}l + 2^{-1/5}h)^5)^{1/5} - l \\
 &= l((1 + 40(h/l)^2 + 80(h/l)^4)^{1/5} - 1) \simeq l(h/l)^2
 \end{aligned}$$

when $h \ll l$ by Taylor's theorem, again. This computation indicates that on the axial directions, the excess in the triangle inequality for "flat" triangles is comparable to $l(h/l)^5$, while on the diagonal directions, the excess is comparable to $l(h/l)^2$. In fact, as long as the base of the triangle is neither horizontal nor vertical, the excess is comparable to $l(h/l)^2$ provided that $h \ll l$ (depending on proximity of the base to an axial direction). In higher dimensions, the excess in the triangle inequality also depends on the direction of the altitude of a triangle in addition to the direction of its base.

The strict gap in the exponents witnessed in Example 1.9 is an essential feature of ℓ_p geometry when $p \neq 2$. To prove Theorems 1.6 and 1.7, we employ tools from functional analysis such as modulus of smoothness, modulus of convexity, and normalized duality mappings to carry out the estimates outlined in Remark 1.4 and Example 1.9 in the setting of ℓ_p in full generality. In fact, we work in arbitrary uniformly smooth and uniformly convex Banach spaces, which include the ℓ_p spaces when $1 < p < \infty$. To establish the sufficient condition (1.17), we modify the "parametric proof" of the sufficient condition in ℓ_2 recently developed by Badger, Naples, and Vellis [16]. This approach also yields new sufficient criteria for a set in a Banach space to be contained in a $(1/s)$ -Hölder curve with $s > 1$ (see Theorem 2.7). To verify the necessary condition (1.18), we follow the proof of the necessary condition in ℓ_2 originally introduced by Schul [65], indicating which parts of the proof are metric, which parts are Banach, and which parts rely on the uniform convexity of the norm. The adaptation of the Hilbert space proofs to ℓ_p and related Banach spaces is nontrivial; see Sections 2 (sufficient conditions) and 3 (necessary conditions) for details. While revising an earlier draft of this manuscript, we encountered some mistakes in the proof of the necessary conditions in [65]; see Remark 3.8 and [14, Appendix C]. In Section 3.3, we show how to correct the error in a minimal way, leaving the outline of virtually all of the original proofs intact; see Remark 3.24. We complete the proof of Theorem 1.7 in [14].

To show that the exponents in Theorems 1.6 and 1.7 are sharp and to prove Proposition 1.1, we construct Koch-snowflake-like curves in Section 5. While the examples of sharpness when $\min(p, 2) = 2$ or $\max(2, p) = 2$ can be built inside any 2-dimensional subspace of ℓ_p , the examples of sharpness when $\min(p, 2) = p \neq 2$ or $\max(2, p) = p \neq 2$ require curves that extend outside of every finite-dimensional subspace of ℓ_p . This is plausible because we know the critical exponent in any finite-dimensional Banach space is 2 by the analyst's traveling salesman theorem in $\mathbb{R}^n$ and bi-Lipschitz equivalence of norms in finite dimensions (see Section 4 for details). Thus, it is natural to expect examples showing sharpness of an exponent $p \neq 2$ to live in infinite dimensions.

Sections 4 (the analyst's TSP in finite-dimensional Banach spaces) and 5 (examples) may be read independently of Sections 2 and 3 (proofs of the main theorems).

1.3. Related work

The inception for this investigation is a recent paper of Edelen, Naber, and Valtorta [38] that extends *Reifenberg's topological disk theorem* [60] (also see [36]) from the Euclidean to infinite-dimensional Hilbert and Banach spaces. The original formulation

of the theorem says that if a closed set $\Sigma \subset \mathbb{R}^n$ is uniformly bilaterally $\delta(k, n)$ -close² to some k -dimensional affine plane at all locations in Σ and on all sufficiently small scales, then Σ is locally homoeomorphic to open subsets of $\mathbb{R}^k$ (that is, Σ is locally a topological disk). Reifenberg proved the topological disk theorem to establish existence and regularity of the plateau problem in arbitrary dimension and codimension. Underpinning the theorem is an algorithm that takes a collection of planes approximating the set Σ and patches them together using orthogonal projections and partitions of unity to construct a parameterization. Edelen, Naber, and Valtorta solve the problem of how to implement this algorithm in a Banach space with dimension independent estimates. The main application of the Reifenberg algorithm in [38] is a structure theorem for measures in Banach spaces, which we now briefly describe.

Following the convention used in [38], for every Borel regular measure μ on a Banach space $\mathbb{X}$, location $x \in \mathbb{X}$, and scale $r > 0$, define the k -dimensional L^2 Jones beta number $\beta_\mu^k(x, r)$ by

$$(1.22) \quad \beta_\mu^k(x, r)^2 := \inf_{p+V^k} r^{-k-2} \int_{B(x, r)} \text{dist}(z, p+V)^2 d\mu(z),$$

where the infimum ranges over all k -dimensional affine subspaces of $\mathbb{X}$. Beta numbers associated to measures were originally introduced by David and Semmes [34, 35] to build a bridge between singular integral operators and quantitative rectifiability of sets.

THEOREM 1.10 (Edelen, Naber, and Valtorta [38, Theorem 2.1])

Let $\mathbb{X}$ be a Banach space, let μ be a finite Borel measure supported in $B(0, 1)$, let $S \subset B(0, 1)$ be a set with $\mu(B(0, 1) \setminus S) = 0$, and let $r : S \rightarrow (0, 1)$. Assume that μ satisfies

$$(1.23) \quad \int_{r(x)}^2 \beta_\mu^k(x, s)^\alpha \frac{ds}{s} \leq M^{\alpha/2} \quad \text{for all } x \in S,$$

where the exponent α is determined as follows:

- if $\mathbb{X}$ is a generic Banach space, then $\alpha = 1$;
- if $\mathbb{X}$ is a Hilbert space, then $\alpha = 2$;
- if $\mathbb{X}$ is a uniformly smooth Banach space and $k = 1$, then α is the smoothness power of $\mathbb{X}$ (for example, the smoothness power of ℓ_p is $\min(p, 2)$ for $1 < p < \infty$).

Then there is a subset $S' \subset S$ so that we have the following packing/measure estimate:

$$(1.24) \quad \mu\left(B(0, 1) \setminus \bigcup_{x \in S'} B(x, r(x))\right) \lesssim_{k, \rho_{\mathbb{X}}} M \quad \text{and} \quad \sum_{x \in S'} r(x)^k \lesssim_{k, \rho_{\mathbb{X}}} 1,$$

where $\rho_{\mathbb{X}}$ denotes the modulus of smoothness of $\mathbb{X}$ (see Section 2 below).

2. For a comparison of unilateral (Jones) versus bilateral (Reifenberg) flatness of a set, see [12].

Roughly speaking, condition (1.23) allows one to control the tilt of the approximating planes in the Reifenberg algorithm and construct local bi-Lipschitz parameterizations. For our present discussion, the most interesting aspect of Theorem 1.10 is the dependence of the exponent α on the geometry of the Banach space $\mathbb{X}$ and the dimension k of the approximating planes. In any given space, one would like to identify the largest possible exponent such that the theorem holds. The exponent $\alpha = 1$ corresponds to the triangle inequality, which holds in any Banach space, and the exponent $\alpha = 2$ corresponds to the Pythagorean theorem, which holds in any Hilbert space. In an intermediate scenario, Edelen, Naber, and Valtorta prove that when $\mathbb{X}$ is a smooth Banach space *and* $k = 1$, the exponent α can be taken to be the smoothness power of the Banach space. For example, $\alpha = p$ when $\mathbb{X} = \ell_p$ and $1 < p < 2$. Furthermore, Edelen, Naber, and Valtorta prove that the restriction to $k = 1$ is necessary to obtain $\alpha > 1$ in non-Hilbert spaces. This is tied up with the existence of good projections onto lines when $k = 1$ and the absence of good projections onto subspaces when $k \geq 2$; see [38, Sections 3.6 and 5.3] for details.

A strength of the Reifenberg algorithm over the analyst's traveling salesman theorem is that it gives conditions to build parameterizations of *every* dimension k . An advantage of the analyst's traveling salesman theorem over the Reifenberg algorithm is that it provides *necessary and sufficient* conditions for parameterizations of dimension $k = 1$. Edelen, Naber, and Valtorta's successful implementation of the Reifenberg algorithm in smooth Banach spaces with $k = 1$ provided our initial motivation to look for an analyst's traveling salesman theorem in Banach spaces.

A separate vein of research by Hahlomaa [43] and David and Schul [32] (also see [44, 63]) focuses on the analyst's TSP in the setting of an arbitrary metric space. Because metric spaces are not necessarily path connected, it is natural to reformulate the analyst's TSP as stated above and instead ask which sets in a given metric space $\mathbb{X}$ are contained in *rectifiable curve fragments*—that is, images of Lipschitz maps $f : S \rightarrow \mathbb{X}$ from some set $S \subset [0, 1]$. Hahlomaa's original work in this direction established an analogue of the *sufficient half* of the analyst's traveling salesman theorem by redefining Jones' beta numbers using Menger curvature, or equivalently, using the excess in the triangle inequality. For different perspectives on rectifiability in measure metric spaces, see, for example, [4, 21, 22, 23, 50, 56, 59].

Following [32], for a given metric space E and ball $Q = B(p, r)$, define $\beta_\infty^E(Q)$ by

$$(1.25) \quad \beta_\infty^E(Q)^2 := r^{-1} \sup \{ \text{dist}(x, y) + \text{dist}(y, z) - \text{dist}(x, z) : \\ x, y, z \in E \cap Q \text{ and } \text{dist}(x, y) \leq \text{dist}(y, z) \leq \text{dist}(x, z) \}.$$

The *metric beta number* $\beta_\infty^E(Q)$ measures the normalized excess in the triangle inequality among triples of points in $E \cap Q$. The exponent 2 on the left-hand side of (1.25) is a convention that is imposed to make the statement of Theorem 1.11 look similar to Theorem 1.2 when $E \subset \mathbb{R}^N$ is endowed with the Euclidean metric.

THEOREM 1.11 (Hahlomaa [43, Theorem 5.3])

Let E be a metric space and let $\mathcal{G}$ be a multiresolution family for E with inflation factor $A_{\mathcal{G}} \simeq 1$. If

$$(1.26) \quad S_{\infty}^E(\mathcal{G}) := \text{diam } E + \sum_{Q \in \mathcal{G}} \beta_{\infty}^E(Q)^2 \text{diam } Q < \infty,$$

then there exists a set $A \subset [0, 1]$ and a surjective Lipschitz map $f : A \rightarrow E$ with Lipschitz constant $\text{Lip}(f) \lesssim S_{\infty}^E(\mathcal{G})$.

REMARK 1.12

It is known that the converse to Hahlomaa's theorem is (quantitatively) false for certain rectifiable curves in $\ell_1^2 = (\mathbb{R}^2, |\cdot|_1)$; see [64, Example 3.3.1]. A similar phenomenon occurs in graph inverse limit spaces; see [33, Section 7]. This issue is not fully understood and merits further investigation.

David and Schul recently announced a partial converse to Hahlomaa's theorem, which is the first nontrivial necessary condition for the analyst's TSP in a metric space. Together, Theorems 1.11 and 1.13 are quite striking and indicate the rough shape that a full solution to the analyst's TSP in a general metric space might take. Recall that a metric space is *doubling* if every ball of radius r can be covered by at most D balls of radius $r/2$.

THEOREM 1.13 (David and Schul [32, Theorem A])

Let Σ be a connected, doubling metric space with doubling constant D and let $\mathcal{H}$ be a multiresolution family for Σ with inflation factor $A_{\mathcal{H}} > 1$. For every $\epsilon > 0$,

$$(1.27) \quad S_{\infty}^{\Sigma, \epsilon}(\mathcal{H}) := \text{diam } Q + \sum_{Q \in \mathcal{H}} \beta_{\infty}^{\Sigma}(Q)^{2+\epsilon} \text{diam } Q \lesssim_{\epsilon, D, A_{\mathcal{H}}} \mathcal{H}^1(\Sigma).$$

REMARK 1.14

The doubling assumption in Theorem 1.13 allows the authors to simplify the overall proof of theorem. David and Schul conjecture (see [32, Remark 1.6]) that the doubling assumption can be dropped by implementing the techniques in [65].

David and Schul present several corollaries to Theorem 1.13 with alternative definitions of the metric beta numbers $\beta_{\infty}^{\Sigma}(Q)$ (see (1.25)). In particular, they obtain necessary conditions for the analyst's TSP in ℓ_p , with $1 < p < \infty$, using traditional Jones' beta numbers $\beta_{\Sigma}(Q)$ (see (1.1)). Also see [32, Corollary D] for a more general statement on uniformly convex Banach spaces.

COROLLARY 1.15 (David and Schul [32])

Let $1 < p < \infty$, let $\Sigma \subset \ell_p$ be a connected set with doubling constant D , and let $\mathcal{H}$ be

a multiresolution family for Σ with inflation factor $A_{\mathcal{H}} > 1$. For all $\epsilon > 0$,

$$(1.28) \quad \begin{aligned} S_{E, \max(2, p) + \epsilon}(\mathcal{H}) &= \text{diam } E + \sum_{Q \in \mathcal{H}} \beta_E(Q)^{\max(2, p) + \epsilon} \text{diam } Q \\ &\lesssim_{\epsilon, D, p, A_{\mathcal{H}}} \mathcal{H}^1(\Sigma). \end{aligned}$$

In Theorem 1.7, we remove the doubling assumption *and* the error ϵ from Corollary 1.15. This is accomplished by following the strategy in [65].

2. Modulus of smoothness and proof of the sufficient conditions

2.1. Ordering flat sets in Banach spaces

A simple but important ingredient in all proofs of the analyst's traveling salesman theorem is that "almost flat" sets of points can be linearly ordered. To implement a generic Banach space version of the sufficient half of the analyst's traveling salesman theorem with universal constants (in the spirit of Hahlmaa [43]), we first develop an instance of this principle. The following lemma is modeled after [17, Lemma 8.3].

LEMMA 2.1 (Flatness implies order)

Let $\mathbb{X}$ be a Banach space. Suppose that $V \subset \mathbb{X}$ is a δ -separated set with $\#V \geq 2$ and there exist lines L_1 and L_2 and a number $0 \leq \alpha < 1/6$ such that

$$(2.1) \quad \text{dist}(v, L_i) \leq \alpha \delta \quad \text{for all } v \in V \text{ and } i = 1, 2.$$

Let $\pi_i : \mathbb{X} \rightarrow L_i$ denote a metric projection onto L_i ; that is, any map satisfying

$$(2.2) \quad \text{dist}(x, L_i) = \text{dist}(x, \pi_i(x)) \quad \text{for all } x \in \mathbb{X}.$$

There exist compatible identifications of L_1 and L_2 with $\mathbb{R}$ such that $\pi_1(v') \leq \pi_1(v'')$ if and only if $\pi_2(v') \leq \pi_2(v'')$ for all $v', v'' \in V$. If $v_1, v_2 \in V$, then

$$(2.3) \quad (1 + 2\alpha)^{-1} |\pi_1(v_1) - \pi_1(v_2)| \leq |v_1 - v_2| \leq (1 + 3\alpha) |\pi_1(v_1) - \pi_1(v_2)|.$$

Proof

Without loss of generality, it suffices to assume $\delta = 1$. Let $V \subset \mathbb{X}$ be a 1-separated set with at least two points. Assume that there exist 1-dimensional affine subspaces L_1 and L_2 in $\mathbb{X}$ and a number $0 \leq \alpha < 1/6$ such that

$$\text{dist}(v, L_i) \leq \alpha \quad \text{for all } v \in V \text{ and } i = 1, 2.$$

Let π_i denote a metric projection onto L_i . For any distinct pair of points $v_1, v_2 \in V$,

$$1 \leq |v_1 - v_2| \leq |\pi_i(v_1) - \pi_i(v_2)| + 2\alpha$$

because V is 1-separated and the distance of points in V to L_i is bounded by α . Hence,

$$(2.4) \quad |\pi_i(v_1) - \pi_i(v_2)| \geq |v_1 - v_2| - 2\alpha \geq 1 - 2\alpha > 2/3.$$

In particular, $2\alpha \leq 3\alpha |\pi_i(v_1) - \pi_i(v_2)|$, and it follows that

$$|v_1 - v_2| \leq (1 + 3\alpha) |\pi_i(v_1) - \pi_i(v_2)|.$$

This establishes the right half of (2.3). Similarly,

$$|\pi_i(v_1) - \pi_i(v_2)| \leq |v_1 - v_2| + 2\alpha \leq (1 + 2\alpha)|v_1 - v_2|,$$

which yields the left half of (2.3). In particular, note that

$$(2.5) \quad |v_1 - v_2| \geq |\pi_i(v_1) - \pi_i(v_2)| - 2\alpha.$$

Suppose for contradiction that there are identifications of L_1 and L_2 with $\mathbb{R}$ and distinct points $v, v', v'' \in V$ such that $\pi_1(v) < \pi_1(v') < \pi_1(v'')$, but $\pi_2(v') < \pi_2(v) < \pi_2(v'')$. Set

$$x := |v - v'|, \quad y := |v'' - v'|, \quad z := |v'' - v|,$$

$$x_1 := |\pi_1(v) - \pi_1(v')|, \quad y_1 := |\pi_1(v'') - \pi_1(v')|, \quad z_1 := |\pi_1(v'') - \pi_1(v)|,$$

$$x_2 := |\pi_2(v) - \pi_2(v')|, \quad y_2 := |\pi_2(v'') - \pi_2(v')|, \quad z_2 := |\pi_2(v'') - \pi_2(v)|.$$

Heuristically, since $\pi_1(v) < \pi_1(v') < \pi_1(v'')$, we have $z \approx x + y$, and since $\pi_2(v') < \pi_2(v) < \pi_2(v'')$, we have $y \approx x + z$. Hence, $z \approx z + 2x$, which yields a contradiction if α is sufficiently small. More precisely, by repeated application of (2.4) and (2.5):

$$\begin{aligned} z &\geq z_1 - 2\alpha = x_1 + y_1 - 2\alpha \geq x_1 + y - 4\alpha \geq x_1 + y_2 - 6\alpha \\ &= x_1 + x_2 + z_2 - 6\alpha \geq x_1 + x_2 + z - 8\alpha. \end{aligned}$$

Rearranging, we obtain $4/3 < x_1 + x_2 \leq 8\alpha < 8/6$, which is absurd. Therefore, under any choice of identifications of L_1 and L_2 with $\mathbb{R}$, either $\pi_1(v) \leq \pi_1(v')$ if and only if $\pi_2(v) \leq \pi_2(v')$ for all $v, v' \in V$, or $\pi_1(v) \leq \pi_1(v')$ if and only if $\pi_2(v) \geq \pi_2(v')$ for all $v, v' \in V$. Thus, we can choose compatible identifications of L_1 and L_2 with $\mathbb{R}$ such that $\pi_1(v') \leq \pi_1(v'')$ if and only if $\pi_2(v') \leq \pi_2(v'')$ for all $v', v'' \in V$. $\square$

COROLLARY 2.2 (cf. [16, Lemma 2.2])

Let $\mathbb{X}$ be a Banach space. Suppose that $V \subset \mathbb{X}$ is a δ -separated set with $\#V \geq 2$ and there exists a line L and a number $0 \leq \alpha < 1/6$ such that

$$(2.6) \quad \text{dist}(v, L) \leq \alpha\delta \quad \text{for all } v \in V.$$

Enumerate $V = \{v_1, \dots, v_n\}$ so that v_{i+1} lies to the right of v_i for all $1 \leq i \leq n-1$. Then

$$(2.7) \quad \sum_{i=1}^{n-1} |v_{i+1} - v_i|^s \leq (1 + 3\alpha)^{2s} |v_1 - v_n|^s \quad \text{for all } s \in [1, \infty).$$

Proof

Let π denote a metric projection onto L . For all $1 \leq i \leq n$, set $x_i := \pi(v_i)$. Then

$$(1 + 3\alpha)^{-1} |x_{i+1} - x_i| \leq |v_{i+1} - v_i| \leq (1 + 3\alpha) |x_{i+1} - x_i| \quad \text{for all } 1 \leq i \leq n-1$$

by Lemma 2.1. Assume $s \in [1, \infty)$ and $\#V \geq 2$. Then

$$\begin{aligned} \sum_{i=1}^{n-1} \frac{|v_{i+1} - v_i|^s}{(1 + 3\alpha)^s} &\leq \sum_{i=1}^{n-1} |x_{i+1} - x_i|^s \\ &\leq \left(\sum_{i=1}^{n-1} |x_{i+1} - x_i| \right)^s = |x_1 - x_n|^s \leq (1 + 3\alpha)^s |v_1 - v_n|^s \end{aligned}$$

because $s \geq 1$ and $x_1, \dots, x_n$ appear in the given order on the line L by Lemma 2.1. This proves (2.7). $\square$

REMARK 2.3

In a general Banach space, the metric projection is not unique and may be norm increasing. For example, in $\ell_\infty^2 = (\mathbb{R}^2, |\cdot|_\infty)$, consider the horizontal line L through the origin (“the x -axis”) and the point $v = (1, \alpha)$ for some $0 < \alpha \leq 1$. Then $|v|_\infty = 1$, $\text{dist}(v, L) = \alpha$, and a metric projection from v to L can be any point on the line segment $[1 - \alpha, 1 + \alpha] \times \{0\}$. In particular, if $\pi_L(v) = (1 + \alpha, 0)$, then $|\pi_L(v)|_\infty = (1 + \alpha)|v|_\infty > |v|_\infty$. This shows that in Lemma 2.1 for an arbitrary Banach space, we cannot expect to replace the lower bound in (2.3) with a 1-Lipschitz bound.

2.2. Lipschitz and Hölder continuous traveling salesman parameterizations in Banach spaces

Throughout this section, let $\mathbb{X}$ denote an arbitrary Banach space.

DEFINITION 2.4 (Doubling scales)

Let $0 < \xi_1 \leq \xi_2 < 1$. A (ξ_1, ξ_2) -doubling sequence of scales is a sequence $(\rho_k)_{k=0}^\infty$ of positive numbers such that $\rho_0 = 1$ and for all $k \geq 0$, $\xi_1 \rho_k \leq \rho_{k+1} \leq \xi_2 \rho_k$. In the special case when $\xi_1 = \xi_2$, we may call $(\rho_k)_{k=0}^\infty$ a *geometric sequence of scales*.

Following [16], let $\mathcal{V} = (V_k, \rho_k)_{k=0}^\infty$ be a sequence consisting of nonempty finite sets V_k in $\mathbb{X}$ and positive numbers ρ_k . Assume that there exist $x_0 \in \mathbb{X}$, $r_0 > 0$, $C^* \geq 1$, and $0 < \xi_1 \leq \xi_2 < 1$ such that $\mathcal{V}$ has the following properties:

- (V0) The numbers $(\rho_k)_{k=0}^\infty$ are a (ξ_1, ξ_2) -doubling sequence of scales.
- (V1) When $k = 0$, we have $V_0 \subset B(x_0, C^* r_0)$.
- (V2) For all $k \geq 0$, we have $V_k \subset V_{k+1}$.
- (V3) For all $k \geq 0$ and all distinct $v, v' \in V_k$, we have $|v - v'| \geq \rho_k r_0$.
- (V4) For all $k \geq 0$ and all $v \in V_{k+1}$, there exists $v' \in V_k$ such that $|v - v'| < C^* \rho_{k+1} r_0$.

With C^* and ξ_2 given, define the associated constant

$$(2.8) \quad A^* := \frac{C^*}{1 - \xi_2}.$$

In addition to (V0)–(V4), assume that for each $k \geq 0$ and $v \in V_k$, we are given a number $\alpha_{k,v} \geq 0$ and a line $L_{k,v}$ in $\mathbb{X}$ such that

$$(V5) \quad \sup_{x \in V_{k+1} \cap B(v, 30A^* \rho_k r_0)} \text{dist}(x, L_{k,v}) \leq \alpha_{k,v} \rho_{k+1} r_0.$$

DEFINITION 2.5 (Flat pairs; see [16])

Fix a parameter $\alpha_0 \in (0, 1/6)$. For all $k \geq 0$, define $\text{Flat}(k)$ to be the set of pairs $(v, v') \in V_k \times V_k$ such that

- (F1) $\rho_k r_0 \leq |v - v'| < 14A^* \rho_k r_0$, and
- (F2) $\alpha_{k,v} < \alpha_0$ and v' is the first point in $V_k \cap B(v, 14A^* \rho_k r_0)$ to the left or to the right of v with respect to the ordering induced by $L_{k,v}$.

Given a pair $(v, v') \in \text{Flat}(k)$, let $V_{k+1}(v, v')$ denote the set of all points $x \in V_{k+1} \cap B(v, 14A^* \rho_k r_0)$ such that x lies between v and v' (inclusive) with respect to the ordering induced by $L_{k,v}$.

DEFINITION 2.6 (Variation excess; see [16])

For all $s \in [1, \infty)$, for all $k \geq 0$, and for all $(v, v') \in \text{Flat}(k)$, define the s -variation excess $\tau_s(k, v, v')$ by

$$(2.9) \quad \tau_s(k, v, v') |v - v'|^s = \max \left\{ \left(\sum_{i=1}^{n-1} |v_{i+1} - v_i|^s \right) - |v - v'|^s, 0 \right\},$$

where $V_{k+1}(v, v') = \{v_1, \dots, v_n\}$ is enumerated so that $v_1 = v$ and for all $1 \leq i \leq n-1$, $v_{i+1} \in V_{k+1}(v, v')$ is the first point after v_i in the direction from v to v' with respect to the ordering induced by $L_{k,v}$ (hence $v_n = v'$).

The following theorem extends Badger, Naples, and Vellis; see [16, Theorem 5.1]. In its original form, the theorem was stated for $\mathbb{X} = \ell_2$ with a weaker restriction on α_1 , achieved through targeted use of the Pythagorean theorem.

THEOREM 2.7 (Hölder traveling salesman parameterizations for nets in Banach spaces)

Let $\mathbb{X}$ be a Banach space. In addition to (V0)–(V5), assume that

$$(2.10) \quad \alpha_0 \leq \alpha_1 := \frac{\xi_1(1 - \xi_2)}{87C^*}.$$

If the sum

$$(2.11) \quad S_{\mathbb{V}}^s := \sum_{k=0}^{\infty} \sum_{(v,v') \in \text{Flat}(k)} \tau_s(k, v, v') \rho_k^s + \sum_{k=0}^{\infty} \sum_{\substack{v \in V_k \\ \alpha_{k,v} \geq \alpha_0}} \rho_k^s < \infty,$$

then there exists a $(1/s)$ -Hölder continuous map $f : [0, 1] \rightarrow \mathbb{X}$ such that $f([0, 1])$ contains $\bigcup_{k \geq 0} V_k$ and the $(1/s)$ -Hölder constant of f satisfies $H \lesssim_{s, C^*, \xi_1, \xi_2} r_0(1 + S_{\mathbb{V}}^s)$.

Proof

Repeat the proof of [16, Theorem 5.1] (see [16, Sections 2–5]) *mutatis mutandis*. Use Lemma 2.1 and Corollary 2.2 above in place of [16, Lemmas 2.1 and 2.2]. $\square$

REMARK 2.8 (Outline of key modifications)

We specify some details to aid the reader with the proof of Theorem 2.7. The reader is first urged to read through [16, Section 2], followed by item (C0) in [16, Section 3.10], which indicates how key parameters are chosen.

In Hilbert space, the initial upper bound $1/16$ on the size of α_0 is made so that

$$1 + 3\alpha_0^2 \leq 1 + 3(1/16)^2 < 1.1.$$

See [16, Lemma 2.3]. In generic Banach space, let us initially require $\alpha_0 \leq 1/31$ so that $1 + 3\alpha_0 \leq 1 + 3(1/31) < 1.1$. Then using Lemma 2.1 instead of [16, Lemma 2.1], the rest of the proof and conclusion of [16, Lemma 2.3] goes through as written.

The algorithm presented in [16, Section 3] requires no changes in the Banach setting.

The principal estimates in the proof of the theorem occur in [16, Section 4]. No changes are required until we reach the proof of [16, Lemma 4.6], where we need to use Corollary 2.2 instead of [16, Lemma 2.2]. This time, we require that $\alpha_0 \leq 1/62$ so that we can replace the original estimate $1 + 3\alpha_0^2 < 1.1$ with the estimate $(1 + 3\alpha_0)^2 < 1.1$. The next required change occurs at the end of the proof of [16, Lemma 4.9]. Using Corollary 2.2 once again, we see the original requirement $1 + 3\alpha_0^2 - \xi_1/14A^* \leq 1$ becomes $(1 + 3\alpha_0)^2 - \xi_1/14A^* \leq 1$, or equivalently $6\alpha_0 + 9\alpha_0^2 \leq \xi_1/14A^*$. With our *a priori* bound $\alpha_0 \leq 1/62$, this certainly holds provided $(6 + 9/62)\alpha_0 \leq \xi_1/14A^*$. Thus, after noting that $(6 + 9/62)14 = 86.03\dots$, it suffices to take

$$\alpha_1 \leq \frac{\xi_1}{87A^*} = \frac{\xi_1(1 - \xi_2)}{87C^*}.$$

Note that

$$\min\left\{\frac{1}{31}, \frac{1}{62}, \frac{\xi_1(1 - \xi_2)}{87C^*}\right\} = \frac{\xi_1(1 - \xi_2)}{87C^*}$$

because $0 < \xi_1 \leq \xi_2 < 1$ and $C^* \geq 1$. There are two final uses of Lemma 2.1 instead of [16, Lemma 2.1] to estimate the separation of points after projection onto an approximating line $\ell_{k,v}$, once in the proof of [16, Proposition 4.11] and once in the proof of [16, (4.3)]. This change affects the value of the implicit constant in [16, Proposition 4.11], but not dependencies of the constant.

To finish the proof of Theorem 2.7, repeat the argument in [16, Section 5.1] verbatim.

COROLLARY 2.9

Let $\mathbb{X}$ be a Banach space. Assume $\mathcal{V} = (V_k, \rho_k)_{k=0}^\infty$ satisfies (V0)–(V5) above. If the sum

$$(2.12) \quad S_{\mathcal{V}} := \sum_{k=0}^{\infty} \sum_{v \in V_k} \alpha_{k,v} \rho_k < \infty,$$

then there exists a rectifiable curve Γ containing $\bigcup_{k \geq 0} V_k$ such that

$$(2.13) \quad \mathcal{H}^1(\Gamma) \lesssim_{C^*, \xi_1, \xi_2} r_0(1 + S_{\mathcal{V}}).$$

Proof

Set $\alpha_0 = \alpha_1$, which depends only on ξ_1, ξ_2 , and C^* . By Corollary 2.2 with $s = 1$, we have $\tau_1(k, v, v') \leq 6\alpha_{k,v} + 9\alpha_{k,v}^2 \leq 7\alpha_{k,v}$ for every flat pair $(v, v') \in \text{Flat}(k)$. Thus,

$$\begin{aligned} S_{\mathcal{V}}^1 &= \sum_{k=0}^{\infty} \sum_{(v, v') \in \text{Flat}(k)} \tau_1(k, v, v') \rho_k + \sum_{k=0}^{\infty} \sum_{\substack{v \in V_k \\ \alpha_{k,v} \geq \alpha_1}} \rho_k \\ &\leq \sum_{k=0}^{\infty} \sum_{(v, v') \in \text{Flat}(k)} 7\alpha_{k,v} \rho_k + \alpha_1^{-1} \sum_{\substack{v \in V_k \\ \alpha_{k,v} \geq \alpha_1}} \alpha_{k,v} \rho_k \leq \alpha_1^{-1} S_{\mathcal{V}} < \infty. \end{aligned}$$

By Theorem 2.7, there exists a Lipschitz map $f : [0, 1] \rightarrow \mathbb{X}$ such that $\Gamma := f([0, 1])$ contains $\bigcup_{k=0}^{\infty} V_k$ and $\mathcal{H}^1(\Gamma) \leq \text{Lip}(f) \lesssim_{C^*, \xi_1, \xi_2} r_0(1 + S_{\mathcal{V}}^1) \lesssim_{C^*, \xi_1, \xi_2} r_0(1 + S_{\mathcal{V}})$. $\square$

For completeness, we show how to use Corollary 2.9 to derive a beta number criterion for a set in a Banach space to be contained in a rectifiable curve. The following theorem is best viewed as the Banach space analogue of Theorem 1.11, expressed with the geometric Jones' beta numbers (1.1) instead of metric beta numbers (1.25). To recall the definition of the sum $S_{E,1}(\mathcal{G})$ of beta numbers over a multiresolution family $\mathcal{G}$ for E , see (1.15).

THEOREM 2.10 (Sufficient half of Schul's theorem in arbitrary Banach spaces)

Let $\mathbb{X}$ be a Banach space. If $E \subset \mathbb{X}$ and $S_{E,1}(\mathcal{G}) = \text{diam } E + \sum_{Q \in \mathcal{G}} \beta_E(Q) \text{diam } Q < \infty$ for some multiresolution family $\mathcal{G}$ for E with inflation factor $A_{\mathcal{G}} \geq 240$, then E is contained in a rectifiable curve Γ in $\mathbb{X}$ with

$$(2.14) \quad \mathcal{H}^1(\Gamma) \lesssim_{A_{\mathcal{G}}} S_{E,1}(\mathcal{G}).$$

Proof

Let $\mathbb{X}$ be a Banach space, let $E \subset \mathbb{X}$, let $\mathcal{G}$ be a multiresolution family for E with inflation factor $A_{\mathcal{G}} \geq 240$, and assume that $S_{E,1}(\mathcal{G}) < \infty$. Then E is bounded and there exists a unique integer $k_0 \in \mathbb{Z}$ such that

$$(2.15) \quad 2^{-k_0} \leq \text{diam } E < 2 \cdot 2^{-k_0}.$$

For all $k \geq 0$, define $\rho_k = 2^{-k}$ and $V_k = X_{k_0+k}$, where $(X_j)_{j \in \mathbb{Z}}$ are the 2^{-j} -nets for E used to define $\mathcal{G}$. Set parameters $C^* = 2$, $\xi_1 = \xi_2 = \frac{1}{2}$, and $r_0 = 2^{-k_0}$, and choose any $x_0 \in V_0 = X_{k_0}$. Then the sequence $\mathcal{V} = (V_k, \rho_k)_{k=0}^{\infty}$ satisfies properties (V0)–(V4) above. Note that $A^* = 4C^* = 8$ and $30A^* = 240$ since $\xi_1 = \xi_2 = \frac{1}{2}$. For each $k \geq 0$ and $v \in V_k$, set $\alpha_{k,v} = 8A_{\mathcal{G}}\beta_E(B(v, A_{\mathcal{G}}2^{-(k_0+k)}))$ and choose $L_{k,v}$ to be any line

such that

$$(2.16) \quad \begin{aligned} & \sup_{x \in E \cap B(v, A_{\mathcal{G}} 2^{-(k_0+k)})} \text{dist}(x, L_{k,v}) \\ & \leq 2\beta_E(B(v, A_{\mathcal{G}} 2^{-(k_0+k)})) \text{diam } B(v, A_{\mathcal{G}} 2^{-(k_0+k)}). \end{aligned}$$

Because $A_{\mathcal{G}} \geq 240 = 30A^*$, it follows that for all $k \geq 0$ and $v \in V_k$,

$$(2.17) \quad \begin{aligned} & \sup_{x \in V_{k+1} \cap B(v, 30A^* \rho_k r_0)} \text{dist}(x, L_{k,v}) \\ & \leq 2\beta_E(B(v, A_{\mathcal{G}} 2^{-(k_0+k)})) \text{diam } B(v, A_{\mathcal{G}} 2^{-(k_0+k)}) \\ & \leq 8A_{\mathcal{G}} \beta_E(B(v, A_{\mathcal{G}} 2^{-(k_0+k)})) 2^{-(k_0+k+1)} \\ & = \alpha_{k,v} \rho_{k+1} r_0. \end{aligned}$$

Thus, property (V5) is satisfied, as well. To proceed, observe that

$$(2.18) \quad \begin{aligned} S_{\mathcal{V}} &= \sum_{k=0}^{\infty} \sum_{v \in V_k} \alpha_{k,v} \rho_k = \sum_{k=0}^{\infty} \sum_{x \in V_k} 8A_{\mathcal{G}} \beta_E(B(v, A_{\mathcal{G}} 2^{-(k_0+k)})) 2^{-k} \\ &\leq 8A_{\mathcal{G}} 2^{k_0} S_{E,1}(\mathcal{G}). \end{aligned}$$

Since $S_{\mathcal{V}} \leq 8A_{\mathcal{G}} S_{E,1}(\mathcal{G}) < \infty$, there is a rectifiable curve Γ containing $\bigcup_{k=0}^{\infty} V_k$ such that

$$(2.19) \quad \mathcal{H}^1(E) \lesssim_{C^*, \xi_1, \xi_2} r_0(1 + S_{\mathcal{V}}) \lesssim r_0(1 + 8A_{\mathcal{G}} r_0^{-1} S_{E,1}) \lesssim_{A_{\mathcal{G}}} S_{E,1}$$

by Corollary 2.9 and (2.15). Finally, note that since $(V_k)_{k=0}^{\infty}$ is a sequence of $2^{-(k_0+k)}$ -nets for E and Γ is closed, Γ contains the set $\overline{\bigcup_{k=0}^{\infty} V_k} \supset E$, as well. $\square$

REMARK 2.11

The constant 240 in Theorem 2.10 has not been optimized and can be at least somewhat reduced at the cost of growing the implicit constant in (2.14). In the future event that a smaller constant is needed, the reader should first consult [16, Section 3.10]

2.3. Triangle inequality excess in uniformly smooth Banach spaces

Our goal in this section is to prove that in a uniformly smooth Banach space of power type $p \in (1, 2]$, the exponent 1 in Corollary 2.9 and Theorem 2.10 may be replaced with the exponent p . In the process, we will verify (1.17) in Theorem 1.6. The essential step is to improve the exponent in the bound (2.3) in Lemma 2.1. To achieve this, we follow the strategy used by Edelen, Naber, and Valtorta [38] in their proof of 1-dimensional Reifenberg-type theorems in uniformly smooth Banach spaces. The approach utilizes a special projection operator, which is available in uniformly smooth Banach spaces.

DEFINITION 2.12

Let $\mathbb{X}$ be a Banach space. The *modulus of smoothness* $\rho_{\mathbb{X}}$ of $\mathbb{X}$ is the function $\rho_{\mathbb{X}} : [0, \infty) \rightarrow [0, \infty)$ defined by

$$(2.20) \quad \rho_{\mathbb{X}}(t) := \sup_{|x|=1, |y|=t} \frac{1}{2}(|x+y| + |x-y|) - 1 \quad \text{for all } t \in [0, \infty).$$

DEFINITION 2.13

We say that $\mathbb{X}$ is *uniformly smooth* if $\rho_{\mathbb{X}}(t)/t = o(t)$ as $t \rightarrow 0$. In this case, we say that $\mathbb{X}$ is *smoothness power type* $p \in (1, 2]$ if there exists $C > 0$ such that $\rho_{\mathbb{X}}(t) \leq C t^p$ for all $t > 0$.

REMARK 2.14 (Essential facts)

For general background on the modulus of smoothness and uniformly smooth spaces, including the following inequalities, see, for example, [37, Chapter Three] or [54, Chapter 1, Section e]. On any Banach space $\mathbb{X}$, the modulus of smoothness $\rho_{\mathbb{X}}$ is a nondecreasing convex function such that $\rho_{\mathbb{X}}(0) = 0$ and

$$(2.21) \quad \sqrt{1+t^2} - 1 = \rho_{\ell_2}(t) \leq \rho_{\mathbb{X}}(t) \leq t \quad \text{for all } t \geq 0.$$

Since $\rho_{\mathbb{X}}$ is convex and $\rho_{\mathbb{X}}(0) = 0$,

$$(2.22) \quad \frac{\rho_{\mathbb{X}}(t_1)}{t_1} \leq \frac{\rho_{\mathbb{X}}(t_2)}{t_2} \quad \text{for all } 0 < t_1 \leq t_2.$$

Furthermore, there exists a constant $1 < L_0 < 3.18$ (see [40, Proposition 10]) such that

$$(2.23) \quad \frac{\rho_{\mathbb{X}}(t_2)}{t_2^2} \leq L_0 \frac{\rho_{\mathbb{X}}(t_1)}{t_1^2} \quad \text{for all } 0 < t_1 \leq t_2.$$

The modulus of smoothness $\rho_{\mathbb{X}}$ and modulus of convexity $\delta_{\mathbb{X}}$ (see Section 3.2) are related by

$$(2.24) \quad \rho_{\mathbb{X}^*}(t) = \sup \left\{ \frac{1}{2} t \epsilon - \delta_{\mathbb{X}}(\epsilon) : 0 < \epsilon \leq 2 \right\} \quad \text{for all } t > 0.$$

Hence, the dual $\mathbb{X}^*$ of a uniformly convex Banach space $\mathbb{X}$ (see Section 3.2) is uniformly smooth. Finally, every uniformly smooth Banach space is reflexive.

EXAMPLE 2.15

By Hanner's inequalities [45], $\rho_{L^p}(t) = p^{-1} t^p + o(t^p)$ when $1 < p \leq 2$; and $\rho_{L^p}(t) = \frac{1}{2}(p-1)t^2 + o(t^2)$ when $2 \leq p < \infty$. In particular, the L^p spaces are uniformly smooth with power type $\min(p, 2)$ when $1 < p < \infty$.

We now present a class of 1-Lipschitz projections onto a line in a Banach space. Given a real Banach space $\mathbb{X}$, let $\mathbb{X}^*$ denote the dual of $\mathbb{X}$ and let $J : \mathbb{X} \rightarrow \mathbb{X}^*$ denote a *normalized duality mapping*—that is, a (nonlinear) map satisfying

$$(2.25) \quad |J(x)|_{\mathbb{X}^*} = |x| \quad \text{and} \quad \langle J(x), x \rangle = |x|^2 \quad \text{for all } x \in \mathbb{X},$$

where $\langle f, x \rangle \equiv f(x) \in \mathbb{R}$ denotes the natural pairing of $f \in \mathbb{X}^*$ and $x \in \mathbb{X}$. Alternatively, J is a subgradient of the convex function $x \in \mathbb{X} \mapsto \frac{1}{2}|x|^2$ (see [5, 49]). The norm on any (uniformly) smooth Banach space $\mathbb{X}$ is Gateaux (uniformly Fréchet) differentiable, and thus, J is uniquely determined (see, for example, [37, Chapter Two]) when $\mathbb{X}$ is smooth. For example, when $\mathbb{X} = \ell_p$ with $1 < p < \infty$,

$$J(x) = |x|_{\ell_p}^{2-p} y \in \ell_p^* = \ell_{p'},$$

where $y = (|x_1|^{p-2}x_1, |x_2|^{p-2}x_2, \dots)$ and p' is the conjugate exponent to p .

DEFINITION 2.16 ([38, Definition 3.31])

Let $\mathbb{X}$ be a Banach space and let L be a 1-dimensional linear subspace of $\mathbb{X}$. Define the *J-projection* Π_L onto L by

$$(2.26) \quad \Pi_L(x) := \langle J(v), x \rangle v \quad \text{for all } x \in \mathbb{X},$$

where J is a normalized dual mapping and v is a point in L with $|v| = 1$. When L is a 1-dimensional affine subspace of $\mathbb{X}$, define $\Pi_L \equiv p + \Pi_{L-p}(\cdot - p)$ for any choice of $p \in L$. For all lines L , we also define $\Pi_L^\perp \equiv \text{Id}_{\mathbb{X}} - \Pi_L$.

Let us record some elementary properties of J -projection.

LEMMA 2.17

Let $\mathbb{X}$ be a Banach space and let L be a 1-dimensional linear subspace of $\mathbb{X}$. The *J-projection* Π_L satisfies each of the following properties:

- (1) For all $x \in L$, we have $\Pi_L(x) = x$.
- (2) For all $x \in L^\perp \equiv \Pi_L^\perp(\mathbb{X})$, we have $\langle Jv, x \rangle = 0$ and $\Pi_L(x) = 0$.
- (3) For all $x \in \mathbb{X}$, we have $|\Pi_L(x)| \leq |x|$.
- (4) For all $x \in \mathbb{X}$, we have $\text{dist}(x, L) \leq |\Pi_L^\perp(x)| \leq 2 \text{dist}(x, L)$.

Proof

Let $v \in L$ be the unit vector in the definition of Π_L . If $x \in L$, say $x = cv$, then

$$\Pi_L(x) = \langle Jv, cv \rangle v = c \langle Jv, v \rangle v = c|v|^2 v = cv = x.$$

This gives the first point. To see the second point, for any $x \in L^\perp$, say $x = \Pi_L^\perp(y)$,

$$\begin{aligned} \langle Jv, x \rangle &= \langle Jv, y - \langle Jv, y \rangle v \rangle = \langle Jv, y \rangle - \langle Jv, y \rangle \langle Jv, v \rangle \\ &= \langle Jv, y \rangle - \langle Jv, y \rangle |v|^2 = \langle Jv, y \rangle - \langle Jv, y \rangle = 0. \end{aligned}$$

Hence, $\Pi_L(x) = \langle Jv, x \rangle v = 0v = 0$. For the third point, observe that for any $x \in \mathbb{X}$,

$$|\Pi_L(x)| = |\langle Jv, x \rangle v| \leq |Jv|_{\mathbb{X}^*} |x| |v| = |v|^2 |x| = |x|.$$

To see the last point, suppose that $x \in \mathbb{X}$. Clearly, $|\Pi_L^\perp(x)| = |\Pi_L(x) - x| \geq \text{dist}(x, L)$. Choose $y \in L$ such that $|x - y| = \text{dist}(x, L)$. Then

$$\begin{aligned} |\Pi_L(x) - x| &\leq |\Pi_L(x) - y| + |x - y| \\ &= |\Pi_L(x) - \Pi_L(y)| + |x - y| \leq 2|x - y| = 2 \text{dist}(x, L). \quad \square \end{aligned}$$

We now check that in any Banach space, the J -projection Π_L induces a well-defined order on a sufficiently flat, separated set of points, by checking compatibility with the order induced by a metric projection π_L . The importance of this fact for us is that in the definition of flat pairs in the traveling salesman algorithm (see Definition 2.5), it does not matter whether we order points by π_L or Π_L .

LEMMA 2.18 (Order compatibility for Π_L and π_L)

Let $\mathbb{X}$ be a Banach space. Suppose that V is a δ -separated set with $\#V \geq 2$ and there exists a line L and a number $0 \leq \alpha < 1/8$ such that $\text{dist}(x, L) \leq \alpha\delta$ for all $x \in V$. Then the J -projection Π_L induces an order on V compatible with the order induced by a metric projection π_L onto L .

Proof

Let $V \subset \mathbb{X}$ be a δ -separated set with $\#V \geq 2$, let L be a line in $\mathbb{X}$, let $0 \leq \alpha < 1/6$, and assume that $\text{dist}(x, L) \leq \alpha\delta$ for all $x \in V$. Without loss of generality, we may assume that L is a linear subspace of $\mathbb{X}$. Fix any metric projection π_L onto L . The restriction on α ensures that π_L induces a unique order on V by Lemma 2.1. If $x, y \in V$ are distinct, then

$$(2.27) \quad \begin{aligned} |x - y| &\geq |\Pi_L(x) - \Pi_L(y)| \geq |x - y| - |\Pi_L^\perp(x)| - |\Pi_L^\perp(y)| \\ &\geq (1 - 4\alpha)\delta > \frac{1}{3}\delta > 0 \end{aligned}$$

by the triangle inequality and Lemma 2.17. We may now check that the J -projection Π_L induces an order on V that is compatible with the order induced by π_L . To that end, suppose that $x, y, z \in V$ are distinct points, write $x' = \pi_L(x)$, $y' = \pi_L(y)$, $z' = \pi_L(z)$ and $x'' = \Pi_L(x)$, $y'' = \Pi_L(y)$, $z'' = \Pi_L(z)$, and suppose to get a contradiction that there exists identifications of L with $\mathbb{R}$ such that $x' < y' < z'$ and $y'' < x'' < z''$. Heuristically, because of the order of the triples on the line L ,

$$|x - z| \approx |x - y| + |y - z| \approx |x - y| + \underbrace{|x - y| + |x - z|},$$

which is impossible if α is sufficient small. More precisely, on one hand, by (2.4) and (2.5) from the proof of Lemma 2.1 (recalling that there we normalized $\delta = 1$),

since $x' < y' < z'$. On the other hand, by (2.27),

since $y'' < x'' < z''$. Combining the previous two displayed equations and recalling V is δ -separated, we obtain

$$|x - z| \geq 2|x - y| + |x - z| - 14\alpha\delta \geq |x - z| + 2\delta - 14\alpha\delta,$$

or equivalently $(2 - 14\alpha)\delta \leq 0$. This is a contradiction because $\alpha < 1/8$ implies that $2 - 14\alpha > 2 - 14/8 = 1/4 > 0$. It readily follows that there is an identification of L with $\mathbb{R}$ such that $\pi_L(x) \leq \pi_L(y)$ if and only if $\Pi_L(x) \leq \Pi_L(y)$ for all $x, y \in V$. $\square$

REMARK 2.19 (Geometric interpretation in smooth spaces)

Assume that $\mathbb{X}$ is smooth. For a line L , spanned by a unit vector $v \in \mathbb{X}$, the J -projection Π_L admits the following geometric description. Let $T_v \partial B(0, 1)$ denote the tangent hyperplane to $\partial B(0, 1)$ at the point v (which exists because $\mathbb{X}$ is smooth). By Lemma 2.17(2), a point $x \in \mathbb{X}$ satisfies $\Pi_L(x) = cv$ if and only if $x \in (cv + T_v \partial B(0, 1))$. See Figure 3.




Figure 3. (Color online) Fibers (cyan) of the J -projection Π_L of $\mathbb{X} = \ell_{5/4}^2$ onto the (red) line $L = \text{span}(v)$, where $v = (1/4, y_0)$, $(1/4)^{5/4} + y_0^{5/4} = 1$. The dot (black) is the metric projection $\pi_L(w)$ onto L of $w = (1/4, y_1)$ with $\Pi_L(w) = (0, 0)$.

LEMMA 2.20

If $\mathbb{X}$ is a smooth Banach space, then

$$\frac{d}{dt}|x + ty|^2 = 2\langle J(x + ty), y \rangle$$

for all $x, y \in \mathbb{X}$ and $t \in \mathbb{R}$ with $x + ty \neq 0$ and $y \neq 0$.

Proof

Since $2J$ is the subgradient of $x \in \mathbb{X} \mapsto |x|^2$, we have

$$|z|^2 - |x + ty|^2 \geq \langle 2J(x + ty), z - (x + ty) \rangle \quad \text{for all } z \in \mathbb{X}.$$

The claim follows by applying the inequality to the difference quotient

$$\frac{|x + (t + h)y|^2 - |x + ty|^2}{h}$$

along values $h \rightarrow 0^+$ and for $h \rightarrow 0^-$, where the limit exists by Gateaux differentiability of the norm. $\square$

The following estimate by Alber is crucial for our application below.

LEMMA 2.21 (see Alber [2, Remark 7.3] or [1, (2.13)])

Let $\mathbb{X}$ be a uniformly smooth Banach space and write $h_{\mathbb{X}}(t) \equiv \rho_{\mathbb{X}}(t)/t$. Then

$$(2.28) \quad |J(x) - J(y)|_{\mathbb{X}^*} \leq 4C_0 h_{\mathbb{X}}(8C_0 L_0 |x - y|) \quad \text{for all } x, y \in \mathbb{X},$$

where $C_0 = 2 \max(1, \sqrt{\frac{1}{2}(|x|^2 + |y|^2)})$ and L_0 is Figiel's constant (2.23).

The next estimate is similar to [38, Lemma 3.27]. Unfortunately, the proof of the lemma given in [38] is partially based on [38, Lemma 3.26], which appears to misstate Lemma 2.21. Thus, we supply a proof of the estimate.

LEMMA 2.22

Let $\mathbb{X}$ be a uniformly smooth Banach space and let L be a 1-dimensional linear subspace of $\mathbb{X}$. If $x \in \mathbb{X}$, $|x| \leq 1$, and $\text{dist}(x, L) \leq \alpha|x|$, then

$$(2.29) \quad |x| \leq |\Pi_L(x)| + \frac{8\alpha}{1-4\alpha} h_{\mathbb{X}}(51\alpha|x|),$$

where $h_{\mathbb{X}}(t) \equiv \rho_{\mathbb{X}}(t)/t$.

Proof

For brevity, write $y = \Pi_L(x)$ and $z = \Pi_L^\perp(x) = x - y$. If $\text{dist}(x, L) \leq \alpha|x|$, then $|y| \leq |x|$ and $|z| \leq 2\alpha|x|$ by Lemma 2.17. Thus, by Lemma 2.20 and the fundamental theorem of calculus,

$$\begin{aligned} |x| - |y| &= \int_0^1 \frac{d}{dt} |y + tz| dt = \int_0^1 \frac{1}{2} |y + tz|^{-1} \frac{d}{dt} |y + tz|^2 dt \\ &= \int_0^1 |y + tz|^{-1} \langle J(y + tz), z \rangle dt \leq \frac{1}{(1-4\alpha)|x|} \int_0^1 \langle J(y + tz), z \rangle dt, \end{aligned}$$

where we used the rough estimate $|y + tz| \geq |y| - |z| \geq |x| - 2|z| \geq (1-4\alpha)|x|$.

Now, by Lemma 2.17, $\langle J(y), z \rangle = 0$. Therefore, by Alber's inequality (2.28),

$$\begin{aligned} \int_0^1 \langle J(y + tz), z \rangle dt &= \int_0^1 \langle J(y + tz) - J(y), z \rangle dt \leq \int_0^1 |J(y + tz) - J(y)|_{\mathbb{X}^*} |z| dt \\ &\leq 4C_0 |z| \int_0^1 h_{\mathbb{X}}(8C_0 L_0 |tz|) dt \leq 4C_0 |z| h_{\mathbb{X}}(8C_0 L_0 |z|), \end{aligned}$$

where $C_0 = \sup_{0 \leq t \leq 1} \max\{1, \sqrt{\frac{1}{2}(|y|^2 + |y + tz|^2)}\} \leq 1$ (since $|x| \leq 1$) and $1 < L_0 < 3.18$. Recall that $h_{\mathbb{X}}(t)$ is nondecreasing (see Remark 2.14) and $|z| \leq 2\alpha|x|$. Thus,

$$\int_0^1 \langle J(y + tz), z \rangle dt \leq 8\alpha|x| h_{\mathbb{X}}(51\alpha|x|)$$

because $16L_0 < 50.88$. Combining the displayed estimates yields (2.29). $\square$

We may now improve Lemma 2.1 in uniformly smooth Banach spaces.

LEMMA 2.23 (cf. Lemma 2.1)

Let $\mathbb{X}$ be a uniformly smooth Banach space. Suppose that V is a δ -separated set with $\#V \geq 2$ and there exists a line L and a number $0 \leq \alpha \leq 43/1224 = 0.0351\dots$ such that $\text{dist}(x, L) \leq \alpha\delta$ for all $x \in V$. If $v_1, v_2 \in V$, then

$$(2.30) \quad |\Pi_L(v_1) - \Pi_L(v_2)| \leq |v_1 - v_2| \leq (1 + \rho_{\mathbb{X}}(102\alpha)) |\Pi_L(v_1) - \Pi_L(v_2)|.$$

Proof

Let $V \subset \mathbb{X}$ be a δ -separated set with $\#V \geq 2$, let L be a line in $\mathbb{X}$, let $\alpha \geq 0$, and assume that $\text{dist}(x, L) \leq \alpha\delta$ for all $x \in V$. Fix any pair of distinct points $v_1, v_2 \in V$. By applying two translations and invoking the triangle inequality, we may assume that L is a linear subspace of $\mathbb{X}$, $0 = v_1 \in L$, and $v := v_2$ satisfies $|v| \geq \delta$ and $\text{dist}(v, L) \leq 2\alpha\delta$. Applying a dilation, we may further assume that $|v| = 1$. Then $\text{dist}(v, L) \leq 2\alpha\delta \leq 2\alpha|v|$ and by Lemma 2.22,

$$|v| \leq |\Pi(v)| + \frac{16\alpha}{1-8\alpha} h_{\mathbb{X}}(102\alpha) = |\Pi(v)| + \frac{8}{51(1-8\alpha)} \rho_{\mathbb{X}}(102\alpha).$$

Recall that $\rho_{\mathbb{X}}(t) \leq t$ in any Banach space; see (2.21). Hence,

$$|\Pi(v)| \geq |v| - \frac{8}{51(1-8\alpha)} \rho_{\mathbb{X}}(102\alpha) \geq 1 - \frac{16\alpha}{1-8\alpha} = \frac{1-24\alpha}{1-8\alpha}.$$

We now require that

$$\frac{8}{51(1-8\alpha)} \leq \frac{1-24\alpha}{1-8\alpha},$$

or equivalently, $\alpha \leq 43/1224$. Then combining the displayed equations yields the right-hand side of (2.30). The left-hand side of (2.30) follows immediately from Lemma 2.17. $\square$

COROLLARY 2.24 (cf. Corollary 2.2)

Let $\mathbb{X}$ be a uniformly smooth Banach space. Suppose that $V \subset \mathbb{X}$ is a δ -separated set with $\#V \geq 2$ and there exists a line L and a number $0 \leq \alpha \leq 43/1224$ such that $\text{dist}(x, L) \leq \alpha\delta$ for all $x \in V$. Enumerate $V = \{v_1, \dots, v_n\}$ so that v_{i+1} lies to the right of v_i for all $1 \leq i \leq n-1$, relative to the ordering induced by the metric projection π_L or the J -projection Π_L (see Lemma 2.18). Then

$$(2.31) \quad \sum_{i=1}^{n-1} |v_{i+1} - v_i|^s \leq (1 + \rho_{\mathbb{X}}(102\alpha))^s |v_1 - v_n|^s \quad \text{for all } s \in [1, \infty).$$

Proof

Repeat the proof of Corollary 2.2 *mutatis mutandis*. Use Lemma 2.23 instead of Lemma 2.1. $\square$

THEOREM 2.25 (Analyst's traveling salesman parameterizations for nets in uniformly smooth Banach spaces; cf. Corollary 2.9)

Let $\mathbb{X}$ be a uniformly smooth Banach space and let $\rho_{\mathbb{X}}$ denote its modulus of smoothness (see Definition 2.12). Assume that $\mathcal{V} = (V_k, \lambda_k)_{k=0}^{\infty}$ satisfies (V0)–(V5) in Section 2.2. If the sum

$$(2.32) \quad S_{\rho_{\mathbb{X}}}(\mathcal{V}) := \sum_{k=0}^{\infty} \sum_{v \in V_k} \rho_{\mathbb{X}}(102\alpha_{k,v}) \lambda_k < \infty,$$

then there exists a rectifiable curve Γ containing $\bigcup_{k=0}^{\infty} V_k$ such that

$$(2.33) \quad \mathcal{H}^1(\Gamma) \lesssim_{C^*, \xi_1, \xi_2} r_0 (1 + S_{\rho_{\mathbb{X}}}(\mathcal{V})).$$

Proof

As in the proof of Corollary 2.9, set $\alpha_0 = \alpha_1$, which only depends on C^* , ξ_1 , and ξ_2 . Note that $\alpha_1 \leq 1/384 < 43/1224$. Thus, by Corollary 2.24 with $s = 1$ and the bound $\rho_{\mathbb{X}}(102\alpha_1) \geq \rho_{\ell_2}(102\alpha_1) \simeq \alpha_1^2 \simeq_{C^*, \xi_1, \xi_2} 1$ (see (2.21)),

$$\begin{aligned} S_{\mathcal{V}}^1 &= \sum_{k=0}^{\infty} \sum_{(v, v') \in \text{Flat}(k)} \tau_1(k, v, v') \lambda_k + \sum_{k=0}^{\infty} \sum_{\substack{v \in V_k \\ \alpha_k, v \geq \alpha_1}} \lambda_k \\ &\leq \sum_{k=0}^{\infty} \sum_{(v, v') \in \text{Flat}(k)} \rho_{\mathbb{X}}(102\alpha_{k,v}) \lambda_k + \rho_{\ell_2}(102\alpha_1)^{-1} \sum_{\substack{v \in V_k \\ \alpha_k, v \geq \alpha_1}} \rho_{\mathbb{X}}(102\alpha_{k,v}) \lambda_k \\ &\lesssim_{C^*, \xi_1, \xi_2} S_{\rho_{\mathbb{X}}}(\mathcal{V}) < \infty. \end{aligned}$$

By Theorem 2.7, there exists a Lipschitz map $f : [0, 1] \rightarrow \mathbb{X}$ such that $\Gamma := f([0, 1])$ contains $\bigcup_{k=0}^{\infty} V_k$ and $\mathcal{H}^1(\Gamma) \leq \text{Lip}(f) \lesssim_{C^*, \xi_1, \xi_2} r_0(1 + S_{\mathcal{V}}^1) \lesssim_{C^*, \xi_1, \xi_2} r_0(1 + S_{\rho_{\mathbb{X}}}(\mathcal{V}))$. $\square$

THEOREM 2.26 (Sufficient half of Schul's theorem in uniformly smooth Banach spaces)

Let $\mathbb{X}$ be a uniformly smooth Banach space of power type $p \in (1, 2]$. If $E \subset \mathbb{X}$ and $S_{E,p}(\mathcal{G}) = \text{diam } E + \sum_{Q \in \mathcal{G}} \beta_E(Q)^p \text{diam } Q < \infty$ for some multiresolution family $\mathcal{G}$ for E with inflation factor $A_{\mathcal{G}} \geq 240$, then E is contained in a rectifiable curve Γ in $\mathbb{X}$ with

$$(2.34) \quad \mathcal{H}^1(\Gamma) \lesssim_{\rho_{\mathbb{X}}, A_{\mathcal{G}}} S_{E,p}(\mathcal{G}).$$

Proof

Repeat the proof of Theorem 2.10 *mutatis mutandis*. Use Theorem 2.25 in lieu of Corollary 2.9. $\square$

Because the Banach space ℓ_p is uniformly smooth of power type $\min(p, 2)$ when $1 < p < \infty$, the sufficient condition (1.17) in Theorem 1.6 follows immediately from Theorem 2.26.

3. Modulus of convexity and proof of the necessary conditions

3.1. Canonical parameterization of finite length continua and beta numbers associated to a parameterization

At the heart of the proof of necessary conditions in analyst's traveling salesman theorems is the existence of parameterizations of finite-length continua by Lipschitz curves. An excellent source for the essential background is the recent paper [3] by Alberti and Ottolini.

Given a continuous map $f : [0, 1] \rightarrow \mathbb{X}$ into a metric space and a closed, nondegenerate interval $I \subset [0, 1]$, the *variation* $\text{var}(f, I)$ of f over I is defined by

$$(3.1) \quad \text{var}(f, I) := \sup_{a_1 \leq \dots \leq a_{n+1}} \sum_{i=1}^n |f(a_{i+1}) - f(a_i)| \in [0, \infty],$$

where the supremum ranges over all finite increasing sequences in I . Associated to f , define the *multiplicity function* $m(f, \cdot) : X \rightarrow [0, \infty]$, $m(f, x) = \#f^{-1}(x)$ for all $x \in \mathbb{X}$. The following proposition records the well-known connection between the variation of f (intrinsic length) and the Hausdorff measure of the image of f (extrinsic length).

PROPOSITION 3.1 ([3, Proposition 3.5])

Let $f : [0, 1] \rightarrow \mathbb{X}$ be a continuous map into a metric space and let $I \subset [0, 1]$ be a closed, nondegenerate interval. If $\text{var}(f, I) < \infty$, then $m(f|_I, \cdot)$ is a Borel function and

$$(3.2) \quad \text{var}(f, I) = \int_{\mathbb{X}} m(f|_I, x) d\mathcal{H}^1(x).$$

A theorem of Ważewski [68] asserts that every connected, compact metric space Σ with finite 1-dimensional Hausdorff measure $\mathcal{H}^1$ admits a Lipschitz parameterization by the interval $[0, 1]$ with Lipschitz constant $\text{Lip}(f) = \sup_{x \neq y} |f(x) - f(y)|/|x - y|$ at most $2\mathcal{H}^1(\Sigma)$.³ Alberti and Ottolini have recently proved the following refinement of Ważewski's theorem (in particular, that f has degree zero). Property (2) says that the parameterization f of Σ is *essentially 2-to-1*.

THEOREM 3.2 ([3, Theorem 4.4])

Let Σ be a connected, compact metric space with $\mathcal{H}^1(\Sigma) < \infty$. Then there exists a continuous function $f : [0, 1] \rightarrow \Sigma$ such that

- (1) f is closed, Lipschitz, surjective, and has degree zero (see [3]);
- (2) $m(f, x) = 2$ for $\mathcal{H}^1$ -a.e. $x \in \Sigma$, and $\text{var}(f, [0, 1]) = 2\mathcal{H}^1(\Sigma)$; and,
- (3) f has constant speed equal to $2\mathcal{H}^1(\Sigma)$.

In any Banach space $\mathbb{X}$, a connected set $\Sigma \subset \mathbb{X}$ has the property that $\mathcal{H}^1(\Sigma) = \mathcal{H}^1(\overline{\Sigma})$, where $\overline{\Sigma}$ denotes the closure of Σ in $\mathbb{X}$. Moreover, if $\Sigma \subset \mathbb{X}$ is closed, connected, and $\mathcal{H}^1(\Sigma) < \infty$, then Σ is compact. The proofs of these facts are simple exercises with the definitions, using convexity of $\mathbb{X}$; see, for example, [65, Section 5] (although stated there for $\mathbb{X} = \ell_2$, the proofs there hold in any Banach space).

COROLLARY 3.3

Let $\mathbb{X}$ be a Banach space. If $\Sigma \subset \mathbb{X}$ is connected and $\mathcal{H}^1(\Sigma) < \infty$, then there exists an essentially 2-to-1 Lipschitz surjection $f : [0, 1] \rightarrow \overline{\Sigma}$ with $\text{Lip}(f) = 2\mathcal{H}^1(\Sigma)$.

For the remainder of Section 3, we fix a connected set Σ in a Banach space $\mathbb{X}$ with $\mathcal{H}^1(\Sigma) < \infty$ and we fix a parameterization $f : [0, 1] \rightarrow \overline{\Sigma}$ given by Corollary 3.3. Following [58] and [65], we refer to subcurves of f as “arcs”; note that we do not require arcs be 1-to-1.

3. This fails dramatically for higher-dimensional curves; see, for example, the “Cantor ladders” in [16, Section 9.2].

DEFINITION 3.4 (Arcs and associated quantities)

An arc, $\tau = f|_{[a,b]}$, of $\overline{\Sigma}$ is the restriction of f to some interval $[a, b] \subset [0, 1]$. Given an arc $\tau : [a, b] \rightarrow \overline{\Sigma}$, define

- $\text{Domain}(\tau) = [a, b]$, $\text{Image}(\tau) = \tau([a, b]) = f([a, b])$,
 $\text{Diam}(\tau) = \text{diam Image}(\tau)$;
- $\text{Start}(\tau) = \tau(a) = f(a)$, $\text{End}(\tau) = \tau(b) = f(b)$; and
- $\text{Edge}(\tau) = [f(a), f(b)]$, i.e. $\text{Edge}(\tau)$ is the line segment in $\mathbb{X}$ from $f(a)$ to $f(b)$.

REMARK 3.5

On any interval $[a, b] \subset [0, 1]$, the map $t \mapsto \text{Diam}(f|_{[a, a+t]})$ is a continuous function on $[0, b-a]$. Thus, by the intermediate value theorem, any arc τ with $\text{Diam}(\tau) > \alpha$ can be partitioned into a finite number of arcs σ , all of which save one have $\text{Diam}(\sigma) = \alpha$, and the final of which has $\text{Diam}(\sigma)$ between α and 2α .

DEFINITION 3.6 ([58, 65])

Given an arc τ of $\overline{\Sigma}$, we define the *arc beta number*

$$(3.3) \quad \tilde{\beta}(\tau) := \sup_{x \in \text{Image}(\tau)} \frac{\text{dist}(x, \text{Edge}(\tau))}{\text{Diam}(\tau)} \in [0, 1].$$

EXAMPLE 3.7

Unlike the Jones' beta numbers, which satisfy (1.2), the arc beta numbers are highly non-monotone. To see this, we construct a family of simple examples in the Euclidean plane $\mathbb{R}^2$. Consider the three collinear points

$$x = (0, 0), \quad y = (\epsilon, 0), \quad z = (1, 0) \quad \text{for some } \epsilon \in (0, 1).$$

Let τ' be an arc whose image traces a piecewise linear path

$$x \rightarrow y \rightarrow z \rightarrow y.$$

Let τ be an arc with $\text{Domain}(\tau') \subset \text{Domain}(\tau)$ such that the image of τ traces a piecewise linear path

$$z \rightarrow \underbrace{x \rightarrow y \rightarrow z}_{\text{Image}(\tau')} \rightarrow y.$$

We have $\text{Image}(\tau') = \text{Image}(\tau) = [0, 1] \times \{0\}$, whence $\text{Diam}(\tau') = \text{Diam}(\tau) = 1$. However, $\text{Edge}(\tau') = [x, y]$, while $\text{Edge}(\tau) = [y, z]$, so that

$$\tilde{\beta}(\tau') = \text{dist}(z, [x, y]) = 1 - \epsilon \quad \text{and} \quad \tilde{\beta}(\tau) = \text{dist}(x, [y, z]) = \epsilon.$$

Thus, $\tilde{\beta}(\tau')/\tilde{\beta}(\tau) = (1 - \epsilon)/\epsilon$ can be arbitrarily large or arbitrarily small by choosing the parameter ϵ sufficiently near 0 or 1, respectively. Similar examples can be made with 1-to-1 arcs by allowing the trace of τ' and τ to lie inside $[0, 1] \times [-h, h]$ with $h \ll \epsilon$.

REMARK 3.8

In [65] (at the very end of the statement of Lemma 3.13), the author mistakenly asserts that $\text{Domain}(\tau') \subset \text{Domain}(\tau)$ and $\text{Diam}(\tau) \lesssim \text{Diam}(\tau')$ imply that $\tilde{\beta}(\tau) \gtrsim \tilde{\beta}(\tau')$. This claim is then applied in the proofs of Lemmas 3.14 and 3.16 of that paper. By modifying the definition of almost flat arcs (see Remark 3.24 below), one can avoid this trap entirely.

Under special circumstances, the arc beta numbers are almost monotone. We do not use Lemma 3.9 or 3.10 below, but we include them for completeness.

LEMMA 3.9 (Almost monotonicity I)

If τ' and τ are arcs in $\overline{\Sigma}$ with $\text{Domain}(\tau') \subset \text{Domain}(\tau)$, then

$$(3.4) \quad \tilde{\beta}(\tau') \text{Diam}(\tau') \leq \tilde{\beta}(\tau) \text{Diam}(\tau) + \text{excess}(\text{Edge}(\tau'), \text{Edge}(\tau)),$$

where $\text{excess}(A, B) = \sup_{x \in A} \inf_{y \in B} |x - y|$ denotes the excess of A over B .

In particular, if $\epsilon \in (0, 1)$ and τ is obtained by concatenating an arc of (image) diameter no more than $\epsilon \tilde{\beta}(\tau') \text{Diam}(\tau')$ to each endpoint of τ' , then

$$(3.5) \quad \tilde{\beta}(\tau) \leq (1 - \epsilon)^{-1} \left(\frac{\text{Diam}(\tau)}{\text{Diam}(\tau')} \right) \tilde{\beta}(\tau').$$

Proof

Let $z \in \text{Image}(\tau')$ and $w \in \text{Edge}(\tau')$ be any pair of points such that $\tilde{\beta}(\tau') \text{Diam}(\tau') = \text{dist}(z, \text{Edge}(\tau')) = |z - w|$. By definition of the excess and compactness of line segments, there exists a point $v \in \text{Edge}(\tau)$ such that $|v - w| \leq \text{excess}(\text{Edge}(\tau'), \text{Edge}(\tau)) =: E$. Hence, $\tilde{\beta}(\tau) \text{Diam}(\tau) \geq |z - v| \geq |z - w| - |w - v| \geq \tilde{\beta}(\tau') \times \text{Diam}(\tau') - E$.

Let $\epsilon \in (0, 1)$ and suppose that τ is obtained from τ' by concatenating arcs of diameter no more than $\eta := \epsilon \tilde{\beta}(\tau') \text{Diam}(\tau')$ to the endpoints of τ' . Then $|\text{Start}(\tau) - \text{Start}(\tau')|$ and $|\text{End}(\tau) - \text{End}(\tau')|$ are bounded by η . Given $p \in \text{Edge}(\tau')$, there is $0 \leq t \leq 1$ such that $p = t \text{Start}(\tau') + (1 - t) \text{End}(\tau')$. Then $q = t \text{Start}(\tau) + (1 - t) \text{End}(\tau) \in \text{Edge}(\tau)$ and

$$\text{dist}(p, \text{Edge}(\tau)) \leq |p - q| \leq t |\text{Start}(\tau) - \text{Start}(\tau')| + (1 - t) |\text{End}(\tau) - \text{End}(\tau')| \leq \eta.$$

Because p was an arbitrary point in $\text{Edge}(\tau')$, it follows that $E \leq \eta$. Thus,

$$(1 - \epsilon) \tilde{\beta}(\tau') \text{Diam}(\tau') \leq \tilde{\beta}(\tau') \text{Diam}(\tau') - E \leq \tilde{\beta}(\tau) \text{Diam}(\tau). \quad \square$$

LEMMA 3.10 (Almost monotonicity II)

Let τ' and τ be arcs of $\overline{\Sigma}$ with $\text{Domain}(\tau') \subset \text{Domain}(\tau)$, let $x' = \text{Start}(\tau')$, let $y' = \text{End}(\tau')$, and let $z' \in \text{Image}(\tau')$ be a point such that

$$(3.6) \quad \tilde{\beta}(\tau') \text{Diam}(\tau') \leq \lambda \text{dist}(z', \text{Edge}(\tau')).$$

Let x, y, z denote (possibly nonunique) points in $\text{Edge}(\tau)$, which realize the distance of x', y', z' to $\text{Edge}(\tau)$, respectively. If z lies between x and y in $\text{Edge}(\tau)$, then

$$(3.7) \quad \tilde{\beta}(\tau') \leq 2\lambda \left(\frac{\text{Diam}(\tau)}{\text{Diam}(\tau')} \right) \tilde{\beta}(\tau).$$

Proof

Assign $\delta := \max_{w' \in \{x', y', z'\}} \text{dist}(w', \text{Edge}(\tau))$. If $z \in [x, y]$, then we can write $z = tx + (1-t)y$ for some $t \in [0, 1]$. Then

$$\text{dist}(z, \text{Edge}(\tau')) \leq |z - (tx' + (1-t)y')| \leq t|x - x'| + (1-t)|y - y'| \leq \delta.$$

Hence,

$$\begin{aligned} \lambda^{-1} \beta(\tau') \text{Diam}(\tau') &\leq \text{dist}(z', \text{Edge}(\tau')) \\ &\leq |z' - z| + \text{dist}(z, \text{Edge}(\tau')) \leq 2\delta \leq 2\tilde{\beta}(\tau) \text{Diam}(\tau), \end{aligned}$$

where the final inequality holds because $x', y', z' \in \text{Image}(\tau)$. $\square$

Let us now recall a key element in the proof of the necessary conditions in Theorems 1.2 and 1.5, first introduced by Okikiolu and later formalized by Schul.

DEFINITION 3.11 ([58, 65])

A *filtration* $\mathcal{F} = \bigcup_{n=n_0}^{\infty} \mathcal{F}_n$ is a family of arcs in $\overline{\Sigma}$ with the following properties.

- (1) *Tree structure*: If $\tau' \in \mathcal{F}_{n+1}$, then there exists a unique arc $\tau \in \mathcal{F}_n$ such that $\text{Domain}(\tau') \subset \text{Domain}(\tau)$.
- (2) *Geometric diameters*: For every $\tau \in \mathcal{F}_n$, $\underline{A}\rho^{-n} \leq \text{Diam}(\tau) \leq A\rho^{-n}$ for some constants $\rho > 1$ and $0 < \underline{A} < A < \infty$ independent of τ .
- (3) *Trivial overlaps*: For all $\tau, \tau' \in \mathcal{F}_n$, either $\tau = \tau'$, or $\text{Domain}(\tau)$ and $\text{Domain}(\tau')$ intersect in at most one point.
- (4) *Partitioning*: $\bigcup_{\tau \in \mathcal{F}_n} \text{Domain}(\tau) = \bigcup_{\tau \in \mathcal{F}_{n_0}} \text{Domain}(\tau)$ for every $n \geq n_0$.

LEMMA 3.12 ([58, 65])

Suppose that $\mathbb{X}$ is a Hilbert space. If $\mathcal{F} = \bigcup_{n=n_0}^{\infty} \mathcal{F}_n$ is a filtration, then

$$(3.8) \quad \sum_{\tau \in \mathcal{F}} \tilde{\beta}(\tau)^2 \text{Diam}(\tau) \lesssim_{(A/\underline{A}), \rho} \mathcal{H}^1\left(\bigcup_{\tau \in \mathcal{F}_{n_0}} \text{Image}(\tau)\right).$$

The exponent 2 in (3.8) is a consequence of the Pythagorean theorem or parallelogram law in Hilbert space. With Lemma 3.12 in hand, the remainder of the proof of necessary conditions in the analyst's traveling salesman theorem in $\mathbb{R}^n$ or ℓ_2 is essentially metric, with a strong harmonic analysis flavor. We outline these last steps in Section 3.3.

3.2. Okikiolu's filtration lemma in uniformly convex spaces

We now develop an analogue of Lemma 3.12 in uniformly convex spaces by following the proof in ℓ_2 from [65] (which is based on [58]) and replacing the parallelogram law in Hilbert space with a suitable inequality in uniformly convex spaces from [32].

DEFINITION 3.13

Let $\mathbb{X}$ be a Banach space. The *modulus of convexity* $\delta_{\mathbb{X}}$ of $\mathbb{X}$ is the function $\delta_{\mathbb{X}} : [0, 2] \rightarrow$

$[0, 1]$ defined by

$$(3.9) \quad \delta_{\mathbb{X}}(\epsilon) := \inf \left\{ 1 - \frac{|x + y|}{2} : |x| = |y| = 1 \text{ and } |x - y| \geq \epsilon \right\}.$$

DEFINITION 3.14

A Banach space is called *uniformly convex* if $\delta(\epsilon) > 0$ for all $\epsilon \in (0, 2]$. In this case, we say that $\mathbb{X}$ is *convexity power type* $p \in [2, \infty)$ if there exists $c > 0$ such that $\delta(\epsilon) \geq c\epsilon^p$ for all $\epsilon \in (0, 2]$.

REMARK 3.15 (Essential facts)

For general background on the modulus of convexity and uniformly convex spaces, we again refer the reader to [37, Chapter Three] or [54, Chapter 1, Section e]. On any Banach space $\mathbb{X}$, the modulus of convexity satisfies the inequality

$$(3.10) \quad \delta_{\mathbb{X}}(\epsilon) \leq \delta_{\ell_2}(\epsilon) = 1 - \sqrt{1 - \epsilon^2/4}.$$

Moreover, for all $\epsilon \in (0, 2]$,

$$(3.11) \quad \delta_{\mathbb{X}}(\epsilon) = \inf \left\{ 1 - \frac{|x + y|}{2} : |x| \leq 1, |y| \leq 1, \text{ and } |x - y| \geq \epsilon \right\};$$

see, for example, [54, p. 60]. In addition,

$$(3.12) \quad \frac{\delta_{\mathbb{X}}(\epsilon_1)}{\epsilon_1} \leq \frac{\delta_{\mathbb{X}}(\epsilon_2)}{\epsilon_2} \quad \text{for all } 0 < \epsilon_1 \leq \epsilon_2 \leq 2;$$

see, for example, [54, Lemma 1.e.8]. In contrast with the modulus of smoothness $\rho_{\mathbb{X}}$ (see Section 2.3), the modulus of convexity $\delta_{\mathbb{X}}$ is not necessarily a convex function. The dual $\mathbb{X}^*$ of a uniformly smooth Banach space $\mathbb{X}$ is uniformly convex and every uniformly convex Banach space is reflexive.

EXAMPLE 3.16

By Hanner's inequalities [45], $\delta_{L^p}(\epsilon) = \frac{1}{8}(p-1)\epsilon^2 + o(\epsilon^2)$ when $1 < p \leq 2$; and $\delta_{L^p}(\epsilon) = p^{-1}2^{-p}\epsilon^p + o(\epsilon^p)$ when $2 \leq p < \infty$. In particular, the L^p spaces are uniformly convex with power type $\max(2, p)$ when $1 < p < \infty$.

In [32], David and Schul observed that the modulus of convexity on a uniformly smooth space of power type p can be used to control the triangle inequality excess from below. Because of its centrality to the proof of (1.18), we include a proof of their estimate here for reference. Actually, we provide a slightly stronger statement. There is a large literature on related substitutes for the Pythagorean theorem and parallelogram law in Banach spaces; see, for example, [28, 29, 30, 31].

LEMMA 3.17 (see [32, Lemma 8.2])

Suppose $\mathbb{X}$ is a uniformly convex Banach space. If $x, y, z \in \mathbb{X}$, then

$$(3.13) \quad \begin{aligned} &|x - y| + |y - z| - |x - z| \\ &\geq 2r\delta_{\mathbb{X}}\left(\frac{\text{dist}(y, [x, z])}{r}\right) \quad \text{for all } r \geq \max\{|x - y|, |y - z|\}. \end{aligned}$$

Proof

By (3.10), $\delta_{\mathbb{X}}(1) \leq 1 - (\sqrt{3}/2) < 1$. Thus, in the degenerate case $x = z$, we have $|x - y| + |y - z| - |x - z| = 2|x - y| > 2\delta_{\mathbb{X}}(1) = 2\delta_{\mathbb{X}}(\text{dist}(y, [x, z]) / \max\{|x - y|, |y - z|\})$. This establishes (3.13) with $r = \max\{|x - y|, |y - z|\}$; the general case follows from (3.12). For the remainder of the proof, we assume that $x \neq z$.

If $y \in [x, z]$, then (3.13) holds trivially (both sides vanish). Thus, we may also assume $\text{dist}(y, [x, z]) > 0$.

Because $x \neq z$, the function $g(w) = |w - x|/|x - z|$ is continuous along $[x, z]$ with $g(x) = 0$ and $g(z) = 1$. Hence, there exists $y_0 \in [x, z]$ such that

$$(3.14) \quad \frac{|y_0 - x|}{|x - z|} = \frac{|y - x|}{|x - y| + |y - z|}.$$

Because $y_0 \in [x, z]$, we have $|x - y_0| + |y_0 - z| = |x - z|$, and it follows that

$$(3.15) \quad \frac{|y_0 - z|}{|x - z|} = \frac{|y - z|}{|x - y| + |y - z|}.$$

Rearranging (3.14), we see that

$$|y_0 - x| = |y - x| \frac{|x - z|}{|x - y| + |y - z|} \leq |y - x|$$

by the triangle inequality. By a parallel argument, starting from (3.15), $|y_0 - z| \leq |y - z|$. Therefore, $y, y_0 \in B(x, |y - x|) \cap B(z, |y - z|)$.

Now, let $y' = (y + y_0)/2$ and $h = |y - y_0| = 2|y - y'|$. Invoking (3.11) on a scaled and translated copy of $B(x, |y - x|)$ and similarly on $B(z, |y - z|)$, we obtain

$$\delta_{\mathbb{X}}\left(\frac{h}{|x - y|}\right) \leq 1 - \frac{|y' - x|}{|y - x|} \quad \text{and} \quad \delta_{\mathbb{X}}\left(\frac{h}{|z - y|}\right) \leq 1 - \frac{|y' - z|}{|y - z|}.$$

Therefore, if $|x - y| \leq r$ and $|y - z| \leq r$,

$$\begin{aligned} |x - y| + |y - z| - |x - z| &\geq |x - y| + |y - z| - |x - y'| - |z - y'| \\ &\geq |x - y| \delta_{\mathbb{X}}\left(\frac{h}{|x - y|}\right) + |z - y| \delta_{\mathbb{X}}\left(\frac{h}{|z - y|}\right) \\ &\geq 2r \delta_{\mathbb{X}}(h/r) \end{aligned}$$

by (3.12). Since $y_0 \in [x, z]$, $\text{dist}(y, [x, z]) \leq |y - y_0| = h$ and (3.13) follows. $\square$

LEMMA 3.18 (Filtration lemma, cf. Lemma 3.12)

Let $\mathbb{X}$ be a uniformly convex Banach space of power type $p \in [2, \infty)$, say $\delta_{\mathbb{X}}(\epsilon) \geq c\epsilon^p$ for all $\epsilon \in (0, 2]$. If $\mathcal{F} = \bigcup_{n=n_0}^{\infty} \mathcal{F}_n$ is a filtration in the sense of Definition 3.11, then

$$\begin{aligned} (3.16) \quad &\sum_{\tau \in \mathcal{F}} \tilde{\beta}(\tau)^p \text{Diam}(\tau) \\ &\lesssim_{c,p,(A/\underline{A}),\rho} \sum_{\tau \in \mathcal{F}_{n_0}} \text{var}(f, \text{Domain}(\tau)) - \sum_{\tau \in \mathcal{F}_{n_0}} \mathcal{H}^1(\text{Edge}(\tau)) \\ &\lesssim_{c,p,(A/\underline{A}),\rho} \mathcal{H}^1\left(\bigcup_{\tau \in \mathcal{F}_{n_0}} \text{Image}(\tau)\right). \end{aligned}$$

The implicit constant is of the form $c^{-1}(A/\underline{A})^{p-1}C(\rho, p)$ and blows up as either $c \downarrow 0$ or $p \uparrow \infty$; see (3.23).

Proof

We mimic the proof of Lemma 3.12 in [65], invoking Lemma 3.17 at a critical juncture to replace estimates depending on Hilbert space geometry. For every $\tau \in \mathcal{F}_n$ and $k \in \mathbb{N}$, we let $\mathcal{F}_{\tau,k}$ denote the k th generation descendants of τ ,

$$(3.17) \quad \mathcal{F}_{\tau,k} := \{\tau' \in \mathcal{F}_{n+k} : \text{Domain}(\tau') \subset \text{Domain}(\tau)\}.$$

Also, for every $\tau \in \mathcal{F}$, define

$$(3.18) \quad \Delta(\tau) := \left(\sum_{\tau' \in \mathcal{F}_{\tau,1}} |\text{Start}(\tau') - \text{End}(\tau')| \right) - |\text{Start}(\tau) - \text{End}(\tau)| \in [0, \infty).$$

We immediately see that for each $n_1 \geq n_0$,

$$\begin{aligned} \sum_{\tau \in \mathcal{F}_{n_0}} |\text{Start}(\tau) - \text{End}(\tau)| + \sum_{n=n_0}^{n_1} \sum_{\tau \in \mathcal{F}_n} \Delta(\tau) &= \sum_{\tau' \in \mathcal{F}_{n_1+1}} |\text{Start}(\tau') - \text{End}(\tau')| \\ &\leq \sum_{\tau \in \mathcal{F}_{n_0}} \text{var}(f, \text{Domain}(\tau)). \end{aligned}$$

Thus, because $|\text{Start}(\tau) - \text{End}(\tau)| = \mathcal{H}^1(\text{Edge}(\tau))$ for each arc and f is essentially 2-to-1, recalling (3.2), we obtain,

$$\begin{aligned} (3.19) \quad \sum_{\tau \in \mathcal{F}} \Delta(\tau) &\leq \sum_{\tau \in \mathcal{F}_{n_0}} \text{var}(f, \text{Domain}(\tau)) - \sum_{\tau \in \mathcal{F}_{n_0}} \mathcal{H}^1(\text{Edge}(\tau)) \\ &\leq \sum_{\tau \in \mathcal{F}_{n_0}} \int_{\mathbb{X}} m(\tau, x) d\mathcal{H}^1(x) \leq 2\mathcal{H}^1\left(\bigcup_{\tau \in \mathcal{F}_{n_0}} \text{Image}(\tau)\right). \end{aligned}$$

Next, for every $\tau \in \mathcal{F}$, we define the *discretized edge distance* d_τ by

$$(3.20) \quad d_\tau := \sup_{\tau' \in \mathcal{F}_{\tau,1}} \sup_{x \in \text{Edge}(\tau')} \text{dist}(x, \text{Edge}(\tau)).$$

In addition, for every arc τ and $k \in \mathbb{N}$, choose an arc $\tau_k \in \mathcal{F}_{\tau,k}$ such that d_{τ_k} is maximal among all arcs in $\mathcal{F}_{\tau,k}$. We also write $\tau_0 \equiv \tau$. We claim that

$$(3.21) \quad \tilde{\beta}(\tau) \text{Diam}(\tau) \leq \sum_{k=0}^{\infty} d_{\tau_k}.$$

To see this, choose an auxiliary sequence $\tau^k \in \mathcal{F}_{\tau,k}$ inductively so that $\tau^0 \equiv \tau$ and $\tilde{\beta}(\tau^{k+1}) \text{Diam}(\tau^{k+1})$ is maximal over all arcs in $\mathcal{F}_{\tau^k,1}$. Then

$$\tilde{\beta}(\tau) \text{Diam}(\tau) = \sum_{k=0}^{\infty} (\tilde{\beta}(\tau^k) \text{Diam}(\tau^k) - \tilde{\beta}(\tau^{k+1}) \text{Diam}(\tau^{k+1}))$$

because the series is telescoping and absolutely convergent by our assumption that the arcs have geometrically decaying diameters. Moreover,

$$\tilde{\beta}(\tau^k) \text{Diam}(\tau^k) - \tilde{\beta}(\tau^{k+1}) \text{Diam}(\tau^{k+1}) \leq d_{\tau^k}$$

by the triangle inequality and definition of the sequence τ^k . Hence,

$$\tilde{\beta}(\tau) \text{Diam}(\tau) \leq \sum_{k=0}^{\infty} d_{\tau^k} \leq \sum_{k=0}^{\infty} d_{\tau_k}$$

by maximality of the distance d_{τ_k} among all arcs in $\mathcal{F}_{\tau,k}$. This verifies (3.21).

The proof up until this point is valid in any Banach space. To continue, we now suppose that $\delta_{\mathbb{X}}$ is convexity power type $p \in [2, \infty)$, say $\delta_{\mathbb{X}}(\epsilon) \geq c\epsilon^p$ for all $\epsilon \in (0, 2]$. By Lemma 3.17 and the triangle inequality,

$$(3.22) \quad \begin{aligned} 2c \frac{d_{\tau}^p}{\text{Diam}(\tau)^{p-1}} &\leq \left(\sum_{\tau' \in \mathcal{F}_{\tau,1}} |\text{Start}(\tau') - \text{End}(\tau')| \right) - |\text{Start}(\tau) - \text{End}(\tau)| \\ &= \Delta(\tau) \end{aligned}$$

for any arc $\tau \in \mathcal{F}$. Now, by (3.21) and Minkowski's inequality for ℓ_p ,

$$\begin{aligned} \underbrace{\left(\sum_{\tau \in \mathcal{F}} \tilde{\beta}(\tau)^p \text{Diam}(\tau) \right)^{\frac{1}{p}}}_I &\leq \left(\sum_{\tau \in \mathcal{F}} \left(\sum_{k=0}^{\infty} d_{\tau_k} \right)^p \text{Diam}(\tau)^{1-p} \right)^{\frac{1}{p}} \\ &\leq \sum_{k=0}^{\infty} \left(\sum_{\tau \in \mathcal{F}} \frac{d_{\tau_k}^p}{\text{Diam}(\tau)^{p-1}} \right)^{1/p}. \end{aligned}$$

If $\tau \in \mathcal{F}$, say $\tau \in \mathcal{F}_n$, and $k \in \mathbb{N}$, then

$$\text{Diam}(\tau_k) \leq A\rho^{-(n+k)} = A\rho^{-k}\rho^{-n} \leq (A/\underline{A})\rho^{-k} \text{Diam}(\tau).$$

Hence,

$$I \leq \sum_{k=0}^{\infty} ((A/\underline{A})\rho^{-k})^{\frac{p-1}{p}} \left(\sum_{\tau \in \mathcal{F}} \frac{(d_{\tau_k})^p}{\text{Diam}(\tau_k)^{p-1}} \right)^{\frac{1}{p}}.$$

Thus, by (3.22),

$$(3.23) \quad \begin{aligned} I &\leq \sum_{k=0}^{\infty} (2c)^{-\frac{1}{p}} ((A/\underline{A})\rho^{-k})^{\frac{p-1}{p}} \left(\sum_{\tau \in \mathcal{F}} \Delta(\tau_k) \right)^{\frac{1}{p}} \\ &\leq \sum_{k=0}^{\infty} (2c)^{-\frac{1}{p}} ((A/\underline{A})\rho^{-k})^{\frac{p-1}{p}} \left(\sum_{\tau \in \mathcal{F}} \Delta(\tau) \right)^{\frac{1}{p}} \\ &= \frac{(2c)^{-\frac{1}{p}} (A/\underline{A})^{\frac{p-1}{p}}}{1 - \rho^{\frac{1-p}{p}}} \left(\sum_{\tau \in \mathcal{F}} \Delta(\tau) \right)^{\frac{1}{p}}. \end{aligned}$$

Therefore, combining (3.19) and (3.23), we obtain (3.16). $\square$

REMARK 3.19

The proof of Lemma 3.18 used the standing assumption that $f : [0, 1] \rightarrow \overline{\Sigma}$ is essentially at most 2-to-1. Instead, if one starts with a parameterization f that is essentially

at most m -to-1 for some $m \geq 1$, then (3.16) still holds, but with the implicit constant in the second inequality multiplied by a factor $m/2$.

3.3. Filtration design: Bounding sums of $\beta_\Sigma(Q)^p \text{diam } Q$ from above

The following decoration of a lemma by Schul is an essential tool for constructing filtrations from certain families of arcs, which we think of as *prefiltrations*. Although originally stated in Hilbert space, it is clear upon reading the proof that the lemma is valid in any metric space. Unfortunately, the statement in [65] contains a mistake, claiming erroneously that $\text{Diam}(\tau) \lesssim \text{Diam}(\tau')$ implies $\tilde{\beta}(\tau) \gtrsim \tilde{\beta}(\tau')$, which is false (even in $\mathbb{R}^2$) by Example 3.7. Also, Schul claims that one may transform a prefiltration into $2CJ$ or fewer filtrations with $J \gtrsim 1$ when $\underline{A} = 1$ and $\rho = 2$, but after writing down the details it seems to us that one must break apart the prefiltration into a larger number of families depending on A and take $J \gtrsim_A 1$ in order to verify (3.24) and (3.25). Thus, we supply a corrected statement and detailed proof, whose outline is due to Schul. We defer the proof to the appendix.

LEMMA 3.20

Prefiltration lemma, cf. [65, Lemma 3.13]] Let $\mathbb{X}$ be a metric space and let $f : [0, 1] \rightarrow \overline{\Sigma}$ be a continuous parameterization of a set $\overline{\Sigma} \subset \mathbb{X}$. Assume that $\rho > 1$, $0 < \underline{A} < A < \infty$, and $J \geq 1$ is any integer such that $\rho^J > 6A/\underline{A}$. Then for every family $\mathcal{F}^0 = \bigcup_{n=n_0}^\infty \mathcal{F}_n^0$ of arcs in $\overline{\Sigma}$ with $\mathcal{F}_{n_0}^0 \neq \emptyset$ satisfying

- (1) *bounded overlap: for every arc $\tau \in \mathcal{F}_n^0$, there exists no more than C arcs $\tau' \in \mathcal{F}_n^0$ such that $\text{Domain}(\tau) \cap \text{Domain}(\tau') \neq \emptyset$ for some constant C independent of τ ,*
- (2) *geometric diameters: for every arc $\tau \in \mathcal{F}_n^0$, we have $\underline{A}\rho^{-n} \leq \text{Diam}(\tau) \leq A\rho^{-n}$,*

we can construct $5(A/\underline{A})CJ$ or fewer filtrations $\mathcal{F}^1 = \bigcup_{n=n_1}^\infty \mathcal{F}_n^1$, $\mathcal{F}^2 = \bigcup_{n=n_2}^\infty \mathcal{F}_n^2$, $\dots$, with starting index $n_j \in \{n_0, n_0 + 1, \dots, n_0 + J - 1\}$ for all j and

$$(3.24) \quad \frac{1}{4}(\underline{A}\rho^{(J-1)n_j})\rho^{-Jn} \leq \text{Diam}(\tau) < 2(A\rho^{(J-1)n_j})\rho^{-Jn} \quad \text{for all } j, \tau \in \mathcal{F}_n^j, n \geq n_j,$$

such that for every index $n \geq n_0$ and arc $\tau' \in \mathcal{F}_n^0$, there exists $\mathcal{F}^j$ (in the list of filtrations), an index N with $n - n_j = J(N - n_j)$, and an arc $\tau \in \mathcal{F}_N^j$ such that

$$(3.25) \quad \text{Domain}(\tau') \subset \text{Domain}(\tau) \quad \text{and} \quad \text{Diam}(\tau) < 2 \text{Diam}(\tau').$$

The assignment $(n, \tau') \mapsto (\mathcal{F}^j, N, \tau)$ is injective.

REMARK 3.21

In Lemma 3.20, we may allow an arc to appear in some $\mathcal{F}_n^0$ several times, so long as the bounded overlap condition in the hypothesis is computed with multiplicity. During preprocessing (see the proof), each instance of a repeated arc $\tau' \in \mathcal{F}_n^0$ will be assigned

to a different intermediate family $\mathcal{D}_n^j$, and their extensions τ will land in different filtrations $\mathcal{F}_N^j$. We use this observation in the proofs of Lemmas 3.28 and 3.29 below.

Because the ℓ_p spaces are uniformly convex of power type $\max(2, p)$ when $1 < p < \infty$, the necessary condition (1.18) in Theorem 1.7 is an immediate consequence of the following theorem.

THEOREM 3.22 (Necessary half of Schul's theorem in uniformly convex Banach spaces)

Let $\mathbb{X}$ be a uniformly convex Banach space of power type $p \in [2, \infty)$. If $\Sigma \subset \mathbb{X}$ is connected and $\mathcal{H}$ is a (partial) multiresolution family for Σ with inflation factor $A_{\mathcal{H}} > 1$, then

$$(3.26) \quad S_{\Sigma, p}(\mathcal{H}) = \text{diam } \Sigma + \sum_{Q \in \mathcal{H}} \beta_{\Sigma}(Q)^p \text{diam } Q \lesssim_{p, \delta_{\mathbb{X}}, A_{\mathcal{H}}} \mathcal{H}^1(\Sigma).$$

In the remainder of this section, we outline the proof of Theorem 3.22 in detail. Because (3.26) is trivial when $\mathcal{H}^1(\Sigma) = \infty$, we may continue to assume that $\mathcal{H}^1(\Sigma) < \infty$ and work with the essentially 2-to-1 Lipschitz parameterization $f : [0, 1] \rightarrow \overline{\Sigma}$ fixed above in Section 3.1. Also, because Σ is connected, $\text{diam } \Sigma \leq \mathcal{H}^1(\Sigma)$. Thus, the essential task is to bound the beta number sum from above in terms of $\mathcal{H}^1(\Sigma)$. To carry this out, we modify the proof from [65, Section 3] with the correction noted in Remark 3.24 below. Most of the argument works in any Banach space, and we make sure to explicitly state wherever we need uniform convexity. We work with two classes of almost flat arcs and one class of non-almost-flat arcs, which we call dominant arcs. Flatness is measured with respect to a best approximating line for the image of the arc.

DEFINITION 3.23

For any ball $Q \in \mathcal{H}$ and scaling factor $\lambda \in \{1, 5, 7\}$, let

$$(3.27) \quad \Lambda(\lambda Q) := \{f|_{[a, b]} : [a, b] \text{ is a connected component of } f^{-1}(\overline{\Sigma} \cap 2\lambda Q) \\ \text{such that } \lambda Q \cap f([a, b]) \neq \emptyset\}.$$

The elements in $\Lambda(\lambda Q)$ are arcs in $2\lambda Q$ that touch λQ . Agree to write $\beta_{\Lambda(\lambda Q)}(2\lambda Q)$ as shorthand for $\beta_{\bigcup\{\text{Image}(\tau) : \tau \in \Lambda(\lambda Q)\}}(2\lambda Q)$.

An arc $\tau \in \Lambda(\lambda Q)$ is called **-almost flat* if

$$(3.28) \quad \beta(\tau) := \inf_L \sup_{z \in \text{Image}(\tau)} \frac{\text{dist}(z, L)}{\text{Diam}(\tau)} \leq 50\epsilon_2 \beta_{\Lambda(\lambda Q)}(2\lambda Q),$$

where L ranges over all lines in $\mathbb{X}$ and $0 < \epsilon_2 \ll 1$ is a parameter, ultimately chosen to depend on at most the inflation factor $A_{\mathcal{H}}$ of $\mathcal{H}$ (shortly after Lemma 3.30). Denote the set of *-almost flat arcs in $\Lambda(\lambda Q)$ by $S^*(\lambda Q)$.

An arc $\tau \in \Lambda(\lambda Q)$ is called *almost flat* if

$$(3.29) \quad \beta(\tau) \leq \epsilon_2 \beta_{\Sigma}(Q).$$

Denote the set of almost flat arcs in $\Lambda(\lambda Q)$ by $S(\lambda Q)$. An arc $\tau \in \Lambda(\lambda Q) \setminus S(\lambda Q)$ is called *dominant*.

REMARK 3.24

We have modified the definition of almost flat arcs from [58, 65], which instead required $\tilde{\beta}(\tau) \leq \epsilon_2 \beta_\Sigma(Q)$. Note that the quantity $\beta(\tau)$ is nothing other than the Jones' beta number $\beta_{\text{Image}(\tau)}(\text{Image}(\tau))$ of the image of τ in its own window. By (1.2), it follows that $\text{Domain}(\tau') \subset \text{Domain}(\tau)$ and $\text{Diam}(\tau) \lesssim \text{Diam}(\tau')$ imply $\beta(\tau) \gtrsim \beta(\tau')$. Further, it is apparent that $\beta(\tau) \leq \tilde{\beta}(\tau)$ for every arc. By using the Jones' beta number $\beta(\tau)$ instead of the arc beta number $\tilde{\beta}(\tau)$ wherever possible in the proof below, we avoid the issue in Remark 3.8. We work with $\tilde{\beta}(\tau)$ in only two spots, at the end of proofs of Lemmas 3.28 and 3.29, when we must invoke Lemma 3.18.

REMARK 3.25

For every $Q \in \mathcal{H}$ and $\lambda \in \{1, 5, 7\}$, $S(\lambda Q) \subset S^*(\lambda Q) \subset \Lambda(\lambda Q)$ by (1.2). Domains of distinct arcs in $\Lambda(\lambda Q)$ are disjoint, although the images of distinct arcs in $\Lambda(\lambda Q)$ may coincide because we do not require f be 1-to-1. If $\tau \in \Lambda(\lambda Q)$ is the “only arc” in $\Lambda(\lambda Q)$ in the sense that the image of each arc in $\Lambda(\lambda Q)$ is contained in $\text{Image}(\tau)$ and $\beta_\Sigma(Q) > 0$, then τ is dominant, since $\text{Diam}(\tau) \leq \text{diam } 14Q$ and $\epsilon_2 \ll 1$ imply

$$(3.30) \quad \beta_\Sigma(Q) = \beta_{\text{Image}(\tau)}(Q) \leq 14\beta(\tau) < (1/\epsilon_2)\beta(\tau).$$

One should think of the latter situation as being the infinitesimally generic case. Note that almost flat arcs are defined relative to $\beta_\Sigma(Q)$ and $*$ -almost flat arcs are defined relative to $\beta_{\Lambda(\lambda Q)}(2\lambda Q)$. This is intentional. The choice of the scaling factors $\lambda \in \{1, 5, 7\}$ and definition of $*$ -almost flat arcs are made to implement the proof of Lemma 3.29. Lastly, we have defined $\Lambda(\lambda Q)$ as arcs in $2\lambda Q$ touching λQ so that arcs $\tau \in \Lambda(\lambda Q)$ have uniformly large diameter: $\text{Diam}(\tau) \geq \lambda A_{\mathcal{H}} 2^{-k}$ whenever $Q = B(x, A_{\mathcal{H}} 2^{-k})$, $x \in X_k$, and $\Sigma \setminus 2\lambda Q \neq \emptyset$.

To proceed, we categorize the balls Q in the multiresolution family $\mathcal{H}$ according to the behavior of the associated arcs $\Lambda(Q) \cup \Lambda(5Q) \cup \Lambda(7Q)$. First, let $\mathcal{H}_0$ denote the collection of all balls $Q \in \mathcal{H}$ such that

$$(3.31) \quad \beta_\Sigma(Q) = 0 \quad \text{or} \quad \Sigma \subset 14Q,$$

where $\lambda Q = B(x, \lambda r)$ denotes the concentric dilate of the ball $Q = B(x, r)$ by $\lambda > 0$. Next, we group $\mathcal{H} \setminus \mathcal{H}_0$ into three (overlapping) subfamilies $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{C}$, as follows.

Let $(X_k)_{k \in \mathbb{Z}}$ denote the sequence of 2^{-k} -separated sets for Σ that is used to define the (partial) multiresolution family $\mathcal{H}$. For each $k \in \mathbb{Z}$, let $\mathcal{N}_k = \{B(x, (1/3)2^{-k}) : x \in X_k\}$ denote the collection of *net balls* of level k . By the triangle inequality,

$$(3.32) \quad \text{gap}(B_1, B_2) \geq \frac{1}{3} 2^{-k} \quad \text{for all } B_1 \neq B_2 \text{ in } \mathcal{N}_k,$$

where the reader may recall that $\text{gap}(S, T) = \inf\{\text{dist}(s, t) : s \in S, t \in T\}$. Suppose $Q = B(x, A_{\mathcal{H}} 2^{-k})$ for some $k \in \mathbb{Z}$ and $x \in X_k$. Let $\lambda \in \{1, 5, 7\}$. We say that $Q \in \mathcal{H}_1^\lambda$

if

(3.33) there exists $\tau \in \Lambda(\lambda Q) \setminus S(\lambda Q)$ such that $\text{Image}(\tau) \cap B(x, (1/3)2^{-k}) \neq \emptyset$;

that is, $Q \in \mathcal{H}_1^\lambda$ if there exists a dominant arc in $\Lambda(\lambda Q)$ that touches the net ball at the center of Q . Otherwise, $Q \notin \mathcal{H}_1^\lambda$ and every arc in $\Lambda(\lambda Q)$ that passes through the net ball at the center of Q is an almost flat arc in $\Lambda(\lambda Q)$; in this case, we assign Q to either $\mathcal{H}_2^\lambda$ or $\mathcal{H}_3^\lambda$, depending on global geometry of the $*$ -almost flat arcs in $\Lambda(\lambda Q)$. Following a convention from [65, p. 345], we write $\beta_{S^*(\lambda Q)}(2\lambda Q)$ as shorthand for $\beta_{\bigcup\{\text{Image}(\tau) : \tau \in S^*(\lambda Q)\}}(2\lambda Q)$. Fix a constant $0 < \epsilon_1 \ll 1$ depending on at most $A_{\mathcal{H}}$ to be specified below. We say that $Q \in \mathcal{H}_2^\lambda$ if

$$(3.34) \quad Q \notin \mathcal{H}_1^\lambda \quad \text{and} \quad \beta_{S^*(\lambda Q)}(2\lambda Q) > \epsilon_1 \beta_{\Lambda(\lambda Q)}(2\lambda Q).$$

We say that $Q \in \mathcal{H}_3^\lambda$ if

$$(3.35) \quad Q \notin \mathcal{H}_1^\lambda \quad \text{and} \quad \beta_{S^*(\lambda Q)}(2\lambda Q) \leq \epsilon_1 \beta_{\Lambda(\lambda Q)}(2\lambda Q).$$

When a ball $Q \in \mathcal{H}_2^\lambda$, $\beta_{S^*(\lambda Q)}(Q)$ dominates $\epsilon_1 \beta_{\Lambda(\lambda Q)}(2\lambda Q)$. By contrast, when a ball $Q \in \mathcal{H}_3^\lambda$, the $*$ -almost flat arcs in $\Lambda(\lambda Q)$ are collectively much flatter inside of the window $2\lambda Q$ than the union of all of the arcs in $\Lambda(\lambda Q)$.

We now define

$$(3.36) \quad \mathcal{A} := \{Q \in \mathcal{H} \setminus \mathcal{H}_0 : Q \in \mathcal{H}_1^\lambda \text{ for } \lambda = 1, \lambda = 5, \text{ or } \lambda = 7\},$$

$$(3.37) \quad \mathcal{B} := \{Q \in \mathcal{H} \setminus \mathcal{H}_0 : Q \in \mathcal{H}_2^\lambda \text{ for } \lambda = 1 \text{ or } \lambda = 5\},$$

$$(3.38) \quad \mathcal{C} := \{Q \in \mathcal{H} \setminus \mathcal{H}_0 : Q \in \mathcal{H}_3^\lambda \text{ for } \lambda = 1 \text{ and } \lambda = 5, \text{ and } Q \notin \mathcal{H}_1^7\}.$$

(We neither use $\mathcal{H}_2^7$ nor $\mathcal{H}_3^7$.) Note that $\mathcal{H} \setminus \mathcal{H}_0 \subset \mathcal{A} \cup \mathcal{B} \cup \mathcal{C}$. While the family $\mathcal{C}$ is disjoint from $\mathcal{A} \cup \mathcal{B}$, some balls in $\mathcal{H} \setminus \mathcal{H}_0$ could belong to both $\mathcal{A}$ and $\mathcal{B}$.

REMARK 3.26

Our classification of balls in the family $\mathcal{H}$ is slightly different than in [65], but roughly speaking our $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{C}$ balls correspond to Schul's $\mathcal{G}_1$, $\mathcal{G}_2$, and $\mathcal{G}_3$ balls. See [65, Figure 4, p. 346] for an illustration of the different families. More specifically, our class $\mathcal{B}$ corresponds to Schul's Δ_1 and $\Delta_{2,1}$ balls. Schul also defines $\Delta_{2,2}$ balls in $\mathcal{G}_2$, but our definition includes these in $\mathcal{A}$. We introduced the net balls to consolidate the estimates for $\mathcal{G}_1$ and $\Delta_{2,2}$ balls.

With the families $\mathcal{H}_0$, $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{C}$ now defined, the proof of Theorem 3.22 reduces to establishing an estimate like (3.26) for each category.

LEMMA 3.27 (Counting)

If $\mathbb{X}$ is an arbitrary Banach space, then for all $s > 1$,

$$(3.39) \quad \sum_{Q \in \mathcal{H}_0} \beta_\Sigma(Q)^s \text{diam}(Q) \lesssim_s A_{\mathcal{H}} \mathcal{H}^1(\Sigma),$$

where the implicit constant blows up as $s \downarrow 1$.

Proof

Modify the proof of [65, Lemma 3.9]. Set $D = \text{diam } \Sigma$. Suppose that $k \in \mathbb{Z}$, $x \in X_k$, and $Q = B(x, A_{\mathcal{H}} 2^{-k}) \in \mathcal{H}$. If $\text{diam } 14Q = 28A_{\mathcal{H}} 2^{-k} < D$, then $\Sigma \not\subset 14Q$. Hence, if $Q \in \mathcal{H}_0$ and $\beta_{\Sigma}(Q) > 0$, then $28A_{\mathcal{H}} 2^{-k} \geq D$. Thus, letting $\mathcal{H}(k)$ denote all $Q \in \mathcal{H}$ of radius $A_{\mathcal{H}} 2^{-k}$, we have

$$\sum_{Q \in \mathcal{H}_0} \beta_{\Sigma}(Q)^s \text{diam } Q \leq 2A_{\mathcal{H}} \sum_{k=-\infty}^{k_0} \sum_{Q \in \mathcal{H}(k)} \beta_{\Sigma}(Q)^s 2^{-k},$$

where k_0 is the unique integer such that

$$28A_{\mathcal{H}} 2^{-(k_0+1)} < D \leq 28A_{\mathcal{H}} 2^{-k_0}.$$

By considering any line passing through the ball's center, we obtain the trivial bound $\beta_{\Sigma}(Q) \leq (\text{diam } \Sigma)/(\text{diam } Q) = D/2A_{\mathcal{H}} 2^{-k}$ for all $Q \in \mathcal{H}(k)$. It follows that

$$\sum_{Q \in \mathcal{H}_0} \beta_{\Sigma}(Q)^s \text{diam } Q \leq (2A_{\mathcal{H}})^{1-s} D^s \sum_{k=-\infty}^{k_0} 2^{k(s-1)} \#\mathcal{H}(k).$$

Because $\#\mathcal{H}(k) = \#X_k \leq \#X_{k_0}$ for all $k \geq k_0$, it suffices to bound $\#X_{k_0}$. We picked k_0 so that $14A_{\mathcal{H}} 2^{-k_0} < D$, which more than certainly implies $(1/2)2^{-k_0} < D/2$. In particular, $\mathcal{H}^1(\Sigma \cap U(x, (1/2)2^{-k_0})) \geq (1/2)2^{-k_0}$ because $x \in \Sigma$, Σ is connected, and Σ is not trapped inside the ball. By definition of a net, the open balls $U(x, (1/2)2^{-k_0})$ centered on points $x \in X_{k_0}$ are pairwise disjoint. Ergo,

$$(1/2)2^{-k_0} \#X_{k_0} \leq \sum_{x \in X_{k_0}} \mathcal{H}^1(\Sigma \cap U(x, (1/2)2^{-k_0})) \leq \mathcal{H}^1(\Sigma).$$

Therefore, $\#X_{k_0} \leq 2^{k_0+1} \mathcal{H}^1(\Sigma) \leq 56A_{\mathcal{H}} \mathcal{H}^1(\Sigma)/D$. Assembling all of the pieces, we have

$$\begin{aligned} & \sum_{Q \in \mathcal{H}_0} \beta_{\Sigma}(Q)^s \text{diam } Q \\ & \leq 56A_{\mathcal{H}} \mathcal{H}^1(\Sigma) (D/2A_{\mathcal{H}})^{s-1} \sum_{k=-\infty}^{k_0} 2^{k(s-1)} \\ & = 56A_{\mathcal{H}} \mathcal{H}^1(\Sigma) (D/2A_{\mathcal{H}})^{s-1} \frac{2^{k_0(s-1)}}{1-2^{1-s}} \leq 56A_{\mathcal{H}} \mathcal{H}^1(\Sigma) \frac{14^{s-1}}{1-2^{1-s}}, \end{aligned}$$

where in the final inequality we used once again that $2^{k_0} \leq 28A_{\mathcal{H}}/D$. $\square$

LEMMA 3.28 (Filtrations I)

If $\mathbb{X}$ is uniformly convex of power type $p \in [2, \infty)$, then

$$(3.40) \quad \sum_{Q \in \mathcal{A}} \beta_{\Sigma}(Q)^p \text{diam } Q \lesssim_{p, \delta_{\mathbb{X}}, A_{\mathcal{H}}, \epsilon_2} \mathcal{H}^1(\Sigma).$$

Proof

Modify the proof of [65, Lemma 3.14]. We start with an auxiliary observation. Let $\mathbb{X}$ be

any Banach space. Suppose that ξ is an arc of $\overline{\Sigma}$ such that $\text{Image}(\xi)$ intersects N net balls in $\mathcal{N}_k$. We claim that if N is sufficiently large, then $\beta(\xi) \text{Diam}(\xi) \geq (1/18)2^{-k}$. To see this, choose points $v_1, \dots, v_N \in \text{Image}(\xi)$, one inside each net ball intersecting ξ . Then $v_1, \dots, v_N$ are $\delta = (1/3)2^{-k}$ separated by (3.32). Suppose that L is a line such that $\text{dist}(v_i, L) < (1/6)\delta$ for all i . By Corollary 2.2 with $s = 1$, after reordering $v_1, \dots, v_N$,

$$\frac{1}{3}(N-1)2^{-k} = (N-1)\delta \leq \sum_{i=1}^{N-1} |v_i - v_{i+1}| < (1 + 3/6)^2 |v_1 - v_N| \leq \frac{9}{4} \text{Diam}(\xi),$$

whence $N < 1 + (27/4) \text{Diam}(\xi)/2^{-k}$. Thus, if $N \geq 1 + (27/4) \text{Diam}(\xi)/2^{-k}$, then $\beta(\xi) \text{Diam}(\xi) \geq \inf_L \sup_i \text{dist}(v_i, L) \geq (1/6)\delta = (1/18)2^{-k}$. In particular, suppose that

$$(3.41) \quad \text{Diam}(\xi) \leq 28A_{\mathcal{H}}2^{-k} \quad \text{and} \quad N \geq 190A_{\mathcal{H}},$$

which guarantees $N \geq 1 + 189A_{\mathcal{H}} = 1 + (27/4)28A_{\mathcal{H}} \geq 1 + (27/4) \text{Diam}(\xi)/2^{-k}$. Then

$$(3.42) \quad \beta(\xi) \geq (1/18)2^{-k} / \text{Diam}(\xi) \geq 1/504A_{\mathcal{H}}.$$

Let $Q \in \mathcal{A}$. Then there exists $k \in \mathbb{Z}$, $x \in X_k$, and $\lambda \in \{1, 5, 7\}$ such that $\lambda Q = B(x, \lambda A_{\mathcal{H}}2^{-k}) \in \mathcal{H}_1^\lambda$. Thus, we may choose a dominant arc $\gamma_Q \in \Lambda(\lambda Q) \setminus S(\lambda Q)$ such that $\text{Image}(\gamma_Q)$ intersects the net ball $B(x, (1/3)2^{-k})$ and pick a point $y_Q \in \text{Domain}(\gamma_Q)$ such that $f(y_Q) \in B(x, (1/3)2^{-k})$. Let N_Q be the number of net balls $R \in \mathcal{N}_k$ such that R intersects $\text{Image}(\gamma_Q)$. We know that $N_Q \geq 1$, but if $\mathbb{X}$ is infinite-dimensional, then N_Q could be arbitrarily large. We now define an arc τ'_Q according to one of two alternatives. If $N_Q < \lceil 190A_{\mathcal{H}} \rceil$, then we set $\tau'_Q = \gamma_Q$ and have $\beta(\tau'_Q) > \epsilon_2 \beta_\Sigma(Q)$ since γ_Q is dominant. Otherwise, if $N_Q \geq \lceil 190A_{\mathcal{H}} \rceil$, then we choose τ'_Q to be any arc such that $y_Q \in \text{Domain}(\tau'_Q) \subset \text{Domain}(\gamma_Q)$ and $\text{Image}(\tau'_Q)$ touches precisely $\lceil 190A_{\mathcal{H}} \rceil$ of the net balls from $\mathcal{N}_k$ that intersect $\text{Image}(\gamma_Q)$; then $\beta(\tau'_Q) \geq 1/504A_{\mathcal{H}} \geq (1/504A_{\mathcal{H}})\beta_\Sigma(Q)$ by the auxiliary observation from above. In both cases, the arc τ'_Q satisfies $y_Q \in \text{Domain}(\tau'_Q)$, $\beta(\tau'_Q) \gtrsim_{A_{\mathcal{H}}, \epsilon_2} \beta_\Sigma(Q)$, and τ'_Q intersects no more than $191A_{\mathcal{H}}$ net balls in $\mathcal{N}_k$. Include an instance of the arc τ'_Q in the set $\mathcal{F}_k^0$. (In the unlikely event that $\tau'_R = \tau'_Q$ for some $R \neq Q$, we treat $\mathcal{F}_k^0$ as a multiset; see Remark 3.21.)

We claim that the family $\mathcal{F}^0 = \bigcup_{k=k_0}^\infty \mathcal{F}_k^0$ of arcs is a prefiltration (see Lemma 3.20), where k_0 is the smallest integer such that $\mathcal{A}$ contains a ball of radius $A_{\mathcal{H}}2^{-k_0}$. To see this, first note that if $\tau'_Q \in \mathcal{F}_k^0$, then

$$\frac{1}{3}2^{-k} \leq \text{Diam}(\tau'_Q) \leq 28A_{\mathcal{H}}2^{-k}.$$

The upper bound follows since τ'_Q is contained in $14Q$. The lower bound holds because either $\tau'_Q = \gamma_Q$ and $\text{Diam}(\gamma_Q) \geq \lambda A_{\mathcal{H}}2^{-k}$ by Remark 3.25, or τ'_Q touches at least 2 balls in $\mathcal{N}_k$ and (3.32) is in effect. Next, let's confirm that the arcs in $\mathcal{F}_k^0$ have uniformly bounded overlap for each k . Fix $k \geq k_0$ and $\tau'_Q \in \mathcal{F}_k^0$. Consider the set

$$Y = \{y_R : \tau'_R \in \mathcal{F}_k^0 \text{ and } \text{Domain}(\tau'_R) \cap \text{Domain}(\tau'_Q) \neq \emptyset\} \subset [0, 1].$$

Because each y_R belongs to the net ball at the center of R and (3.32) holds, there is a bijection between Y and arcs $\tau'_R \in \mathcal{F}_k^0$ such that τ'_R and τ'_Q intersect in their domains. Let $\tau'_S, \tau'_T \in \mathcal{F}_k^0$ be the unique arcs such that $y_S = \min Y$ and $y_T = \max Y$. Observe that

$$Y \subset \text{Domain}(\tau'_S) \cup \text{Domain}(\tau'_Q) \cup \text{Domain}(\tau'_T)$$

because the latter set is an interval containing y_S and y_T . Hence, $\#Y$ is bounded from above by the number of net balls in $\mathcal{N}_k$ intersected by τ'_S , τ'_Q , or τ'_T . Therefore, we know τ'_Q intersects at most $\#Y \leq 3 \cdot 191A_{\mathcal{H}} = 573A_{\mathcal{H}}$ arcs $\tau'_R \in \mathcal{F}_k^0$.

Invoking Lemma 3.20 with parameters $\rho = 2$, $A = 28A_{\mathcal{H}}$, $\underline{A} = 1/3$, $C = 573A_{\mathcal{H}}$, and $J = 1 + \lceil \log_2(504A_{\mathcal{H}}) \rceil$ (so that $2^J > 6A/\underline{A}$), we can find $O(A_{\mathcal{H}}^2 \log(A_{\mathcal{H}}))$ filtrations

$$\mathcal{F}^1 = \bigcup_{k=k_1}^{\infty} \mathcal{F}_k^1, \quad \mathcal{F}^2 = \bigcup_{k=k_2}^{\infty} \mathcal{F}_k^2, \quad \mathcal{F}^3 = \bigcup_{k=k_3}^{\infty} \mathcal{F}_k^2, \quad \dots$$

with starting indices $k_0 \leq k_j \leq k_0 + J - 1$ such that

$$\frac{1}{12} 2^{(J-1)k_j} 2^{-Jk} \leq \text{Diam}(\tau) < 56A_{\mathcal{H}} 2^{(J-1)k_j} 2^{-Jk} \quad \text{for all } \tau \in \mathcal{F}_k^j.$$

Note that the ratio of the upper and lower bounds for $\text{Diam}(\tau)$ depends only on $A_{\mathcal{H}}$. Moreover, for all $\tau'_Q \in \mathcal{F}_k^0$, there exists $\mathcal{F}^j$ (in the list of filtrations), an index K with $k - k_j = J(K - k_j)$ and an arc $\tau_Q \in \mathcal{F}_K^j$ such that

$$\text{Domain}(\tau'_Q) \subset \text{Domain}(\tau) \quad \text{and} \quad \text{Diam}(\tau_Q) < 2 \text{Diam}(\tau'_Q).$$

The assignment $(k, \tau'_Q) \mapsto (\mathcal{F}^j, K, \tau_Q)$ is injective. Now, for any $\tau'_Q \in \mathcal{F}_k^0$, we have

$$\beta_{\Sigma}(Q) \lesssim_{A_{\mathcal{H}}, \epsilon_2} \beta(\tau'_Q) \leq 2\beta(\tau_Q) \leq 2\tilde{\beta}(\tau_Q),$$

where the first inequality holds by our choice of τ'_Q , the second inequality holds by (1.2) because $\text{Diam}(\tau) \leq 2 \text{Diam}(\tau')$, and the third inequality holds generally. Thus,

$$\begin{aligned} \sum_{Q \in \mathcal{A}} \beta_{\Sigma}(Q)^p \text{diam } Q &\lesssim_{p, A_{\mathcal{H}}, \epsilon_2} \sum_{\tau' \in \mathcal{F}^0} \beta(\tau')^p \text{Diam}(\tau') \\ &\lesssim_p \sum_j \sum_{\tau \in \mathcal{F}^j} \beta(\tau)^p \text{Diam}(\tau) \leq \sum_j \sum_{\tau \in \mathcal{F}^j} \tilde{\beta}(\tau)^p \text{Diam}(\tau). \end{aligned}$$

Up until this point, the proof is valid in any Banach space and for any $1 \leq p < \infty$. Finally, if $\mathbb{X}$ is uniformly convex of power type $p \in [2, \infty)$, then by Lemma 3.18 and the bound on the total number of filtrations,

$$\sum_{Q \in \mathcal{A}} \beta_{\Sigma}(Q)^p \text{diam } Q \lesssim_{p, A_{\mathcal{H}}, \epsilon_2} \sum_j \sum_{\tau \in \mathcal{F}^j} \tilde{\beta}(\tau)^p \text{Diam}(\tau) \lesssim_{p, \delta_{\mathbb{X}}, A_{\mathcal{H}}, \epsilon_2} \mathcal{H}^1(\Sigma).$$

This verifies (3.40). □

LEMMA 3.29 (Filtrations II)

Assume that ϵ_1 is sufficiently small depending only on $A_{\mathcal{H}}$; $\epsilon_1 = 1/126A_{\mathcal{H}}$ will suffice.

If $\mathbb{X}$ is uniformly convex of power type $p \in [2, \infty)$, then

$$(3.43) \quad \sum_{Q \in \mathcal{C}} \beta_{\Sigma}(Q)^p \operatorname{diam} Q \lesssim_{p, \delta_{\mathbb{X}}, A_{\mathcal{H}}, \epsilon_2} \mathcal{H}^1(\Sigma).$$

Proof

Modify the proof of [65, Lemma 3.16]. We initially assume that $\mathbb{X}$ is any Banach space. Let $Q \in \mathcal{C}$, say $Q = B(x_Q, A_{\mathcal{H}} 2^{-k})$ for some $x_Q \in X_k$. Because

$$\beta_{S^*(Q)}(2Q) \leq \epsilon_1 \beta_{\Lambda(Q)}(2Q),$$

we have $\Lambda(Q) \setminus S^*(Q) \neq \emptyset$ provided that $\epsilon_1 < 1$. Choose any $\tau'_Q \in \Lambda(Q) \setminus S^*(Q)$. Then

$$A_{\mathcal{H}} 2^{-k} \leq \operatorname{Diam}(\tau'_Q) \leq 4A_{\mathcal{H}} 2^{-k} \quad \text{and} \quad \beta_{\Sigma}(Q) \leq 2\beta_{\Lambda(Q)}(2Q) < \frac{1}{25\epsilon_2} \beta(\tau'_Q)$$

by definition of $\Lambda(Q)$ and $S^*(Q)$, respectively. Include an instance of τ'_Q in the set $\mathcal{F}_k^0$.

We claim that the family $\mathcal{F}^0 = \bigcup_{k=k_0}^{\infty} \mathcal{F}_k^0$ of arcs is a prefiltration (see Lemma 3.20), where k_0 is the smallest integer such that $\mathcal{C}$ contains a ball of radius $A_{\mathcal{H}} 2^{-k_0}$. We already noted that arcs in $\mathcal{F}_k^0$ have good bounds on their diameters. To check the bounded overlap property, fix Q as above. Using $\beta_{S^*(5Q)}(10Q) \leq \epsilon_1 \beta_{\Lambda(5Q)}(10Q) \leq \epsilon_1$ for balls $Q \in \mathcal{C}$, we may choose a line L_Q in $\mathbb{X}$ such that

$$(3.44) \quad \begin{aligned} \operatorname{dist}(x, L_Q) &< (21/20)\epsilon_1 \operatorname{diam} 10Q \\ &\leq 21\epsilon_1 A_{\mathcal{H}} 2^{-k} \quad \text{for all } x \in \bigcup_{\tau \in S^*(5Q)} \operatorname{Image}(\tau). \end{aligned}$$

Suppose that $R \in \mathcal{C}$, $R = B(x_R, A_{\mathcal{H}} 2^{-k})$ for some $x_R \in X_k$, and τ'_R and τ'_Q intersect in their domains. Since τ'_Q is contained in $2Q$, τ'_R is contained in $2R$, and the arcs intersect, the triangle inequality yields $x_R \in 4Q$ and $x_Q \in 4R$. Hence, $R \subset 5Q$ and $Q \subset 5R$, as well. Our strategy is to show that x_R is close to L_Q relative to the scale 2^{-k} of separation between points in X_k . By (3.44), it suffices to exhibit an arc $\xi_R \in S^*(5Q)$ containing x_R and demand that ϵ_1 be sufficiently small relative to $A_{\mathcal{H}}$. To find ξ_R , first let $\gamma_R \in \Lambda(7R)$ be any arc containing x_R . Because x_R lies in the net ball $B(x_R, (1/3)2^{-k})$ and $R \notin (\mathcal{H}_0 \cup \mathcal{A})$, we know that $\gamma_R \in S(7R)$; that is, $\beta(\gamma_R) \leq \epsilon_2 \beta_{\Sigma}(R)$. Choose ξ_R to be a subarc of the arc γ_R such that $x_R \in \operatorname{Image}(\xi_R)$ and $\xi_R \in \Lambda(5Q)$, which exists because $x_R \in \operatorname{Image}(\gamma_R) \cap 4Q \subset 5Q$ and $10Q \subset 14R$. On the one hand, since $x_R \in \operatorname{Image}(\xi_R)$, $x_R \in 4Q$, and the endpoints of ξ_R are contained in the boundary of $10Q$, we have $\operatorname{Diam}(\xi_R) \geq 6A_{\mathcal{H}} 2^{-k}$. On the other hand, since $\operatorname{Image}(\gamma_R) \subset 14R$, $\operatorname{Diam}(\gamma_R) \leq 28A_{\mathcal{H}} 2^{-k}$. Thus, because $28/6 < 5$ and $\Sigma \cap R \subset \Sigma \cap 5Q \subset \bigcup_{\tau \in \Lambda(5Q)} \operatorname{Image}(\tau)$,

$$(3.45) \quad \beta(\xi_R) \leq 5\beta(\gamma_R) \leq 5\epsilon_2 \beta_{\Sigma}(R) \leq 25\epsilon_2 \beta_{\Sigma}(5Q) \leq 50\epsilon_2 \beta_{\Lambda(5Q)}(10Q)$$

by three applications of (1.2) and the observation that γ_R is almost flat. This computation confirms that the arc ξ_R belongs to $S^*(5Q)$. (In fact, this computation is the *raison d'être* for $*$ -almost flat arcs!) By (3.44), it follows that $\operatorname{dist}(x_R, L_Q) < (1/6)2^{-k}$

provided that $\epsilon_1 \leq 1/126A_{\mathcal{H}}$. For concreteness, we specify that $\epsilon_1 = 1/126A_{\mathcal{H}}$. Let $\{x_1, \dots, x_N\}$ be an enumeration of the centers of balls $R \in \mathcal{C}$ of the same generation as Q such that τ'_R intersects τ'_Q in the domain. Each $x_i \in 4Q$ and satisfies $\text{dist}(x_i, L_Q) < (1/6)2^{-k}$. By Corollary 2.2, after reordering $x_1, \dots, x_N$, we have

$$(N-1)2^{-k} \leq \sum_{i=1}^{N-1} |x_i - x_{i+1}| \leq (1+3/6)^2 |x_1 - x_N| \leq (9/4) \cdot 8A_{\mathcal{H}}2^{-k} = 18A_{\mathcal{H}}2^{-k}.$$

We conclude that $N \leq 1 + 18A_{\mathcal{H}} \leq 19A_{\mathcal{H}}$. Therefore, $\mathcal{F}^0$ is a prefiltration.

Invoking Lemma 3.20 with parameters $\rho = 2$, $A = 4A_{\mathcal{H}}$, $\underline{A} = A_{\mathcal{H}}$, $C = 19A_{\mathcal{H}}$, and $J = 5$ (so that $2^J > 6A/\underline{A}$), we can find $O(A_{\mathcal{H}})$ filtrations

$$\mathcal{F}^1 = \bigcup_{k=k_1}^{\infty} \mathcal{F}_k^1, \quad \mathcal{F}^2 = \bigcup_{k=k_2}^{\infty} \mathcal{F}_k^2, \quad \mathcal{F}^3 = \bigcup_{k=k_3}^{\infty} \mathcal{F}_k^2, \quad \dots$$

with starting indices $k_0 \leq k_j \leq k_0 + 4$ such that

$$\frac{1}{4}A_{\mathcal{H}}2^{(J-1)k_j}2^{-Jk} \leq \text{Diam}(\tau) < 8A_{\mathcal{H}}2^{(J-1)k_j}2^{-Jk} \quad \text{for all } \tau \in \mathcal{F}_k^j.$$

Note that the ratio of the upper and lower bounds for $\text{Diam}(\tau)$ is universal. Moreover, for all $\tau'_Q \in \mathcal{F}_k^0$, there exists $\mathcal{F}^j$ (in the list of filtrations), an index K with $k - k_j = J(K - k_j)$, and an arc $\tau_Q \in \mathcal{F}_K^j$ such that

$$\text{Domain}(\tau'_Q) \subset \text{Domain}(\tau) \quad \text{and} \quad \text{Diam}(\tau_Q) < 2\text{Diam}(\tau'_Q).$$

The assignment $(k, \tau'_Q) \mapsto (\mathcal{F}^j, K, \tau_Q)$ is injective. Now, for any $\tau'_Q \in \mathcal{F}_k^0$, we have that

$$\beta_{\Sigma}(Q) \lesssim_{\epsilon_2} \beta(\tau'_Q) \leq 2\beta(\tau_Q) \leq 2\tilde{\beta}(\tau_Q),$$

where the first inequality holds by our choice of τ'_Q , the second inequality holds by (1.2) because $\text{Diam}(\tau) \leq 2\text{Diam}(\tau')$, and the third inequality holds generally. Also, we have $\text{diam } Q \simeq \text{Diam}(\tau'_Q) \simeq \text{Diam}(\tau_Q)$. Thus,

$$\begin{aligned} \sum_{Q \in \mathcal{C}} \beta_{\Sigma}(Q)^p \text{diam } Q &\lesssim_{p, \epsilon_2} \sum_{\tau' \in \mathcal{F}^0} \beta(\tau')^p \text{Diam}(\tau') \\ &\lesssim_p \sum_j \sum_{\tau \in \mathcal{F}^j} \beta(\tau)^p \text{Diam}(\tau) \leq \sum_j \sum_{\tau \in \mathcal{F}^j} \tilde{\beta}(\tau^p) \text{Diam}(\tau). \end{aligned}$$

Up until this point, the proof is valid in any Banach space and for any $1 \leq p < \infty$. Finally, if $\mathbb{X}$ is uniformly convex of power type $p \in [2, \infty)$, then by Lemma 3.18 and the bound on the total number of filtrations,

$$\sum_{Q \in \mathcal{C}} \beta_{\Sigma}(Q)^p \text{diam } Q \lesssim_{p, \epsilon_2} \sum_j \sum_{\tau \in \mathcal{F}^j} \tilde{\beta}(\tau)^p \text{Diam}(\tau) \lesssim_{p, \delta_{\mathbb{X}}, A_{\mathcal{H}}, \epsilon_2} \mathcal{H}^1(\Sigma).$$

This verifies (3.43). □

THEOREM 3.30 (Geometric martingales)

Assume that ϵ_2 is sufficiently small depending only on $A_{\mathcal{H}}$ and ϵ_1 ; the value $\epsilon_2 =$

$2^{-55}\epsilon_1/A_{\mathcal{H}}$ will suffice. If $\mathbb{X}$ is any Banach space, then for all $q > 0$,

$$(3.46) \quad \sum_{Q \in \mathcal{B}} \beta_{\Sigma}(Q)^q \operatorname{diam} Q \lesssim_{q, A_{\mathcal{H}}, \epsilon_1} \mathcal{H}^1(\Sigma),$$

where the implicit constant blows up as $q \downarrow 0$.

Note the arbitrary power q in (3.46)! This indicates that $Q \in \mathcal{B}$ (imagine Q is centered at the intersection of two or more crossing line segments) occurs relatively infrequently. This result is universal insofar as it is valid in an *arbitrary* Banach space! We defer the (rather long) proof of Theorem 3.30 to the sequel to this paper [14].

Putting it all together, on a uniformly convex space of power type $p \in [2, \infty)$, choose parameters $0 < \epsilon_1 \ll 1$ and $0 < \epsilon_2 \ll 1$ with $\epsilon_1 \simeq A_{\mathcal{H}}^{-1}$ and $\epsilon_2 \simeq A_{\mathcal{H}}^{-2}$ so that Lemma 3.29 and Theorem 3.30 are valid. Then

$$\sum_{Q \in \mathcal{H}} \beta_{\Sigma}(Q)^p \operatorname{diam} Q \leq I + II + III + IV \lesssim_{p, \delta_{\mathbb{X}}, A_{\mathcal{H}}} \mathcal{H}^1(\Sigma),$$

where

$$I := \sum_{Q \in \mathcal{H}_0} \beta_{\Sigma}(Q)^p \operatorname{diam} Q \leq \sum_{Q \in \mathcal{H}_0} \beta_{\Sigma}(Q)^2 \operatorname{diam} Q$$

$$\lesssim_{A_{\mathcal{H}}} \mathcal{H}^1(\Sigma) \quad \text{by Lemma 3.27,}$$

$$II := \sum_{Q \in \mathcal{A}} \beta_{\Sigma}(Q)^p \operatorname{diam} Q \lesssim_{p, \delta_{\mathbb{X}}, A_{\mathcal{H}}, \epsilon_2} \mathcal{H}^1(\Sigma) \quad \text{by Lemma 3.28,}$$

$$III := \sum_{Q \in \mathcal{B}} \beta_{\Sigma}(Q)^p \operatorname{diam} Q \leq \sum_{Q \in \mathcal{B}} \beta_{\Sigma}(Q)^2 \operatorname{diam} Q$$

$$\lesssim_{A_{\mathcal{H}}, \epsilon_1} \mathcal{H}^1(\Sigma) \quad \text{by Theorem 3.30,} \quad \text{and}$$

$$IV := \sum_{Q \in \mathcal{C}} \beta_{\Sigma}(Q)^p \operatorname{diam} Q \lesssim_{p, \delta_{\mathbb{X}}, A_{\mathcal{H}}, \epsilon_2} \mathcal{H}^1(\Sigma) \quad \text{by Lemma 3.29.}$$

This completes the proof of Theorem 3.22.

4. Analyst's TSP in finite-dimensional Banach spaces

In this section, we record some additional observations about the analyst's TSP in arbitrary finite-dimensional Banach spaces. It is well known that diameter, distance of a point to a set, and Hausdorff measures are bi-Lipschitz metric invariants in the sense that if $f : \mathbb{X} \rightarrow \mathbb{Y}$ is a C -bi-Lipschitz map between metric spaces $\mathbb{X}$ and $\mathbb{Y}$, then each of the quantities in the source and target spaces are quantitatively equivalent with multiplicative constants determined by C . As a consequence, the Jones' beta numbers in a normed vector space are bi-Lipschitz invariant in the following sense.

LEMMA 4.1 (Bi-Lipschitz equivalence of Jones' beta numbers)

Let V be a vector space and let $|\cdot|_{\mathbb{X}}$ and $|\cdot|_{\mathbb{Y}}$ be equivalent norms on V , say that there exists a constant $1 \leq C < \infty$ such that $C^{-1}|x|_{\mathbb{X}} \leq |x|_{\mathbb{Y}} \leq C|x|_{\mathbb{X}}$ for all $x \in V$.

Let $\beta_{E,\mathbb{X}}(Q)$ and $\beta_{E,\mathbb{Y}}(Q)$ denote the Jones' beta numbers in $\mathbb{X} = (V, |\cdot|_{\mathbb{X}})$ and $\mathbb{Y} = (V, |\cdot|_{\mathbb{Y}})$, respectively—that is, for all nonempty sets $E \subset V$ and positive diameter sets $Q \subset V$ with $E \cap Q \neq \emptyset$,

$$\beta_{E,\mathbb{X}}(Q) = \inf_L \sup_{x \in E \cap Q} \frac{\text{dist}_{\mathbb{X}}(x, L)}{\text{diam}_{\mathbb{X}} Q} \quad \text{and} \quad \beta_{E,\mathbb{Y}}(Q) = \inf_L \sup_{x \in E \cap Q} \frac{\text{dist}_{\mathbb{Y}}(x, L)}{\text{diam}_{\mathbb{Y}} Q},$$

where L runs over all lines in V . Then $C^{-2}\beta_{E,\mathbb{X}}(Q) \leq \beta_{E,\mathbb{Y}}(Q) \leq C^2\beta_{E,\mathbb{X}}(Q)$ for all admissible E and Q .

Proof

Let $E \subset V$ be nonempty and let $Q \subset V$ have positive diameter. By convention, $\beta_{E,\mathbb{X}}(Q) = \beta_{E,\mathbb{Y}}(Q) = 0$ if $E \cap Q = \emptyset$; in this case, the conclusion is trivial. Thus, we may assume that $E \cap Q \neq \emptyset$. By equivalence of the norms, $\text{diam}_{\mathbb{X}} Q = \sup\{|x - y|_{\mathbb{X}} : x, y \in Q\}$ and $\text{diam}_{\mathbb{Y}} Q = \sup\{|x - y|_{\mathbb{Y}} : x, y \in Q\}$ are also equivalent:

$$(4.1) \quad C^{-1} \text{diam}_{\mathbb{X}} Q \leq \text{diam}_{\mathbb{Y}} Q \leq C \text{diam}_{\mathbb{X}} Q.$$

Similarly, for any nonempty set $S \subset V$ and $x \in V$, $\text{dist}_{\mathbb{X}}(x, S) = \inf\{|x - s|_{\mathbb{X}} : s \in S\}$ and $\text{dist}_{\mathbb{Y}}(x, S) = \inf\{|x - s|_{\mathbb{Y}} : s \in S\}$ are equivalent:

$$(4.2) \quad C^{-1} \text{dist}_{\mathbb{X}}(x, S) \leq \text{dist}_{\mathbb{Y}}(x, S) \leq C \text{dist}_{\mathbb{X}}(x, S).$$

The set of all lines (that is, 1-dimensional affine subspaces) in V is independent of a choice of norm. Fix a line L in V . By definition of $\beta_{E,\mathbb{X}}(Q)$, (4.1), and (4.2),

$$\beta_{E,\mathbb{X}}(Q) \leq \sup_{x \in E \cap Q} \frac{\text{dist}_{\mathbb{X}}(x, L)}{\text{diam}_{\mathbb{X}} Q} \leq C^2 \sup_{x \in E \cap Q} \frac{\text{dist}_{\mathbb{Y}}(x, L)}{\text{diam}_{\mathbb{Y}} Q}.$$

Taking the infimum over all possible lines, we obtain $\beta_{E,\mathbb{X}}(Q) \leq C^2\beta_{E,\mathbb{Y}}(Q)$. Interchanging the roles of $\mathbb{X}$ and $\mathbb{Y}$ yields $\beta_{E,\mathbb{Y}}(Q) \leq C^2\beta_{E,\mathbb{X}}(Q)$. $\square$

A basic fact in functional analysis is that every pair of finite-dimensional normed vector spaces of the same dimension has equivalent norms. Therefore, the original formulation of the analyst's traveling salesman theorem due to Jones [48] ($\dim V = 2$) and Okikiolu [58] ($\dim V \geq 3$) persists in any finite-dimensional vector space.

THEOREM 4.2 (Jones' and Okikiolu's theorems in finite-dimensional spaces)

Let $\mathbb{X}$ be a finite-dimensional Banach space with $\dim \mathbb{X} \geq 2$, let $\Delta(\mathbb{X})$ be a system of dyadic cubes with respect to some set of coordinates on $\mathbb{X}$, and let $E \subset \mathbb{X}$. Then E is contained in a rectifiable curve if and only if

$$(4.3) \quad S_E(\mathbb{X}) := \text{diam } E + \sum_{Q \in \Delta(\mathbb{X})} \beta_E(3Q)^2 \text{diam } Q < \infty,$$

where $3Q$ denotes the concentric dilate of the cube Q with scaling factor 3 and $\beta_E(3Q)$ denotes the Jones' beta number with respect to $\mathbb{X}$. More precisely, if $S_E(\mathbb{X}) < \infty$, then E is contained in a curve Γ in $\mathbb{X}$ with

$$(4.4) \quad \mathcal{H}^1(\Gamma) \lesssim_{\mathbb{X}, \Delta(\mathbb{X}), \dim \mathbb{X}} S_E(\mathbb{X}).$$

If $\Sigma \subset \mathbb{X}$ is a connected set, then

$$(4.5) \quad S_\Sigma(\mathbb{X}) \lesssim_{\mathbb{X}, \Delta(\mathbb{X}), \dim \mathbb{X}} \mathcal{H}^1(\Sigma).$$

The constant 3 in (4.3) can be replaced with any constant $A > 1$. Then (4.4) and (4.5) hold with implicit constants depending on the norm of $\mathbb{X}$, $\Delta(\mathbb{X})$ (that is, a choice of coordinates), $\dim \mathbb{X}$, and A .

Proof

Suppose that $\dim \mathbb{X} = n$. After fixing coordinates on $\mathbb{X}$, we may identify $\mathbb{X}$ with $\mathbb{R}^n$ and identify $\Delta(\mathbb{X})$ with $\Delta(\mathbb{R}^n)$. Then $(\mathbb{X}, |\cdot|) = (\mathbb{R}^n, |\cdot|)$ and $(\mathbb{R}^n, |\cdot|_2)$ are bi-Lipschitz equivalent, where $|\cdot|_2$ denotes the standard Euclidean norm. (The bi-Lipschitz constant depends on the norm $|\cdot|$ and on the choice of coordinates for $\mathbb{X}$.) Each of the quantities $\text{diam } E$, $\beta_E(3Q)$, and $\mathcal{H}^1(\Gamma)$ defined relative to $|\cdot|$ are bi-Lipschitz equivalent to the respective quantities defined relative to $|\cdot|_2$. Thus, the rectifiability of a curve Γ containing E is independent of the choice of norm (although the length of Γ depends on the norm), and the theorem follows immediately from Theorem 1.2. $\square$

EXAMPLE 4.3

Given linearly independent vectors $\vec{v}$ and $\vec{w}$ in $\mathbb{R}^2$, let $\Delta(\vec{v}, \vec{w})$ denote the system of dyadic parallelograms corresponding to the lattice generated by $\vec{v}$ and $\vec{w}$; for example, $P = \{s\vec{v} + t\vec{w} : 0 \leq s, t \leq 1\}$ and $P' = \{s\vec{v} + t\vec{w} : 0 \leq s, t \leq 1/2\}$ belong to $\Delta(\vec{v}, \vec{w})$, and P' is one of four children of P . A bounded set $E \subset \mathbb{R}^2$ is contained in a rectifiable curve if and only if

$$\sum_{P \in \Delta(\vec{v}, \vec{w})} \beta_E(3P)^2 \text{diam } P < \infty.$$

To see this, note that every $P \in \Delta(\vec{v}, \vec{w})$ is a dyadic square in $\mathbb{R}^2$ with respect to the coordinates induced by $\vec{v}$ and $\vec{w}$ and apply Theorem 4.2.

In a finite-dimensional Banach space, the analyst's traveling salesman theorem can be formulated using cubes, as in Theorems 1.2 and 4.2, and using multiresolution families, as in Theorems 1.5, 1.6, and 1.7. It is always possible to pass between one and the other, at the expense of growing implicit constants. For a more general formulation of this principle, see [24, Appendix B].

LEMMA 4.4

Let $\mathbb{X}$ be a finite-dimensional Banach space with $\dim \mathbb{X} \geq 2$, let $\Delta(\mathbb{X})$ be a system of dyadic cubes with respect to some set of coordinates on $\mathbb{X}$, let $a > 1$ be a scaling factor for cubes, let $E \subset \mathbb{X}$, and let $\mathcal{G}$ be a multiresolution family for E with inflation factor $A_{\mathcal{G}} > 1$. For all $0 < p < \infty$,

$$(4.6) \quad \sum_{Q \in \Delta(\mathbb{X})} \beta_E(aQ)^p \text{diam } Q \simeq_{p, a, A_{\mathcal{G}}, \dim \mathbb{X}} \sum_{B \in \mathcal{G}} \beta_E(B)^p \text{diam } B.$$

Proof

The key point is that any bounded set in $\mathbb{X}$ is contained in some dilate aQ of a cube $Q \in \Delta(\mathbb{X})$ with comparable diameters. As long as the bounded set intersects E , that set is also contained in some ball $B \in \mathcal{G}$ with comparable diameters. We will use this in conjunction with monotonicity of Jones' beta numbers and volume doubling in finite-dimensional spaces.

To normalize scales, let $D = \text{diam } Q_0$ for any choice of $Q_0 \in \Delta(\mathbb{X})$. Fix $Q \in \Delta(\mathbb{X})$, say with $\text{diam } Q = D2^{-j}$ for some $j \in \mathbb{Z}$. Fix $k \in \mathbb{Z}$ to be determined. If $aQ \cap E = \emptyset$, then $\beta_E(aQ) = 0$, so the cube Q is irrelevant to the sum on the left-hand side of (4.6). Suppose that $aQ \cap E \neq \emptyset$ and fix any $z \in aQ \cap E$. Choose $x \in X_k$ (the 2^{-k} -net appearing in the definition of $\mathcal{G}$) such that $|x - z| < 2^{-k}$, and set $B = B(x, A_{\mathcal{G}}2^{-k}) \in \mathcal{G}$. Then for any $y \in aQ$,

$$|y - x| \leq |y - z| + |z - x| < \text{diam } aQ + 2^{-k} = aD2^{-j} + 2^{-k} \leq A_{\mathcal{G}}2^{-k}$$

and $aQ \subset B$ provided that $aD2^{-j} \leq (A_{\mathcal{G}} - 1)2^{-k}$. We now specify that k is the unique integer such that

$$(4.7) \quad (A_{\mathcal{G}} - 1)2^{-(k+1)} < aD2^{-j} \leq (A_{\mathcal{G}} - 1)2^{-k}.$$

As noted, this ensures that $aQ \subset B$. Furthermore,

$$\text{diam } aQ \leq \text{diam } B \leq 2A_{\mathcal{G}}2^{-k} \leq \frac{4A_{\mathcal{G}}}{A_{\mathcal{G}} - 1} \text{diam } aQ.$$

Thus, $\beta_E(aQ) \lesssim_{A_{\mathcal{G}}} \beta_E(B)$ by (1.2). Now, by volume doubling, the sets of the form aR , where $R \in \Delta(\mathbb{X})$ is a dyadic cube of the same generation as Q , have bounded overlap determined by a and $\dim \mathbb{X}$. Also, by (4.7), each $k \in \mathbb{Z}$ is associated to a unique $j \in \mathbb{Z}$. (This is a simple expression of the fact that 2^{-k} -nets and dyadic cubes have the same scaling ratios.) All together, we conclude that

$$\sum_{Q \in \Delta(\mathbb{X})} \beta_E(aQ)^p \text{diam } Q \lesssim_{p,a,A_{\mathcal{G}},\dim \mathbb{X}} \sum_{B \in \mathcal{G}} \beta_E(B)^p \text{diam } B.$$

The reverse inequality holds by similar considerations. □

5. Sharpness of the exponents via examples

Our goal in this section is to verify the sharpness of the exponents on beta numbers in Theorems 1.6 and 1.7. To do so, we build Koch-snowflake-like curves Γ , for which we can estimate beta number sums over *arbitrary* multiresolution families. This type of construction is not new; see, for example, [16, 25, 38, 61] for motivating examples, but the details are subtle.

The organization is as follows. In Section 5.1, we verify sharpness of the exponent 2 in (1.17) when $2 \leq p < \infty$ and in (1.18) when $1 < p \leq 2$ by building curves in the Euclidean plane. In Section 5.2, we verify sharpness of the exponent p in (1.17) when $1 < p \leq 2$ and in (1.18) when $2 \leq p < \infty$ by building curves in infinite-dimensional Banach spaces. Finally, in Section 5.3, we carry out additional estimates to record a proof of Proposition 1.1.




Figure 4. (Color online) The snowflake map $\hat{\gamma}$, displayed with $|\vec{v}| = 1$, $\eta = 1/4$, and $s = 5/16$.

5.1. Examples with critical exponent 2

The main results of this subsection are the following two propositions. Recall that ℓ_p^n denotes $(\mathbb{R}^n, |\cdot|_p)$.

PROPOSITION 5.1

There exists a curve Γ in ℓ_2^2 such that $\mathcal{H}^1(\Gamma) = \infty$ and $S_{\Gamma, 2+\epsilon}(\mathcal{G}) < \infty$ for every multiresolution family $\mathcal{G}$ for Γ and every $\epsilon > 0$. In particular, the exponent 2 in (1.17) when $2 \leq p < \infty$ is sharp.

PROPOSITION 5.2

There exists a curve Γ in ℓ_2^2 such that $\mathcal{H}^1(\Gamma) < \infty$ and $S_{\Gamma, 2-\epsilon}(\mathcal{G}) = \infty$ for every multiresolution family $\mathcal{G}$ for Γ and every $\epsilon > 0$. In particular, the exponent 2 in (1.18) when $1 < p \leq 2$ is sharp.

Sharpness of the exponent 2 in (1.17) when $2 < p < \infty$ and in (1.18) when $1 < p < 2$ follows from the case $p = 2$ because ℓ_p contains a subspace isomorphic to ℓ_p^2 , which in turn is bi-Lipschitz equivalent to ℓ_2^2 . (Use Lemmas 4.1 and 4.4 to pass between multiscale sums of beta numbers in ℓ_p^2 and ℓ_2^2 . In particular, one may use a fixed system of dyadic squares as an intermediary between multiresolution families for Γ in ℓ_p^2 and in ℓ_2^2 .) Therefore, in this section, we focus on $\ell_2^2 \equiv \mathbb{R}^2$, the standard Euclidean plane. We wish to emphasize that in both statements the curve is independent of ϵ . We build the curves using the following procedure.

ALGORITHM 5.3 (Snowflake-like curves in $\mathbb{R}^2$)

Suppose that $\vec{p}, \vec{q} \in \mathbb{R}^2$ and $\gamma : [a, b] \rightarrow \mathbb{R}^2$ is a constant-speed parameterization of $I = [\vec{p}, \vec{q}]$ from $\gamma(a) = \vec{p}$ to $\gamma(b) = \vec{q}$; that is,

$$\gamma(t) = \vec{p} + \frac{t-a}{b-a}(\vec{q} - \vec{p}) \quad \text{for all } t \in [a, b].$$

Write $\vec{v} := \vec{q} - \vec{p}$ and let $\vec{y} \in \mathbb{R}^2$ denote the unique unit vector with $\vec{y} \perp \vec{v}$ such that $\vec{y}$ points to the left of the oriented line segment from $\vec{p}$ to $\vec{q}$. Given a *relative height* $0 \leq \eta \leq 1/\sqrt{12}$, we define a piecewise linear path $\hat{\gamma} : [a, b] \rightarrow \mathbb{R}^2$, as follows. Set $s := 1/4 + \eta^2 \in [1/4, 1/3]$. Divide $[a, b]$ into quarters. For all $a \leq t \leq (3/4)a + (1/4)b$,

$$\hat{\gamma}(t) = \vec{p} + 4\left(\frac{t-a}{b-a}\right)s\vec{v}$$

(see Figure 4). For all $(3/4)a + (1/4)b \leq t \leq (1/2)a + (1/2)b$,

$$\hat{\gamma}(t) = \vec{p} + s\vec{v} + 4\left(\frac{t - (3/4)a - (1/4)b}{b - a}\right)((1/2 - s)\vec{v} + \eta|\vec{v}|\vec{y}).$$

For all $(1/2)a + (1/2)b \leq t \leq (1/4)a + (3/4)b$,

$$\hat{\gamma}(t) = \vec{p} + (1/2)\vec{v} + \eta|\vec{v}|\vec{y} + 4\left(\frac{t - (1/2)a - (1/2)b}{b - a}\right)((1/2 - s)\vec{v} - \eta|\vec{v}|\vec{y}).$$

Finally, for all $(1/4)a + (3/4)b \leq t \leq b$,

$$\hat{\gamma}(t) = \vec{p} + (1 - s)\vec{v} + 4\left(\frac{t - (1/4)a - (3/4)b}{b - a}\right)s\vec{v}.$$

We say that $\hat{\gamma}$ is obtained from γ by *adding a bump of relative height η (on the left side)*. On each quarter of $[a, b]$, $\hat{\gamma}(t)$ traces a line segment of length $s|\vec{v}|$ at constant speed. Thus, $\hat{\gamma}$ has Lipschitz constant $4s|\vec{v}|/(b - a) = (1 + 4\eta^2)|\vec{v}|/(b - a)$. Additionally, we have $\|\gamma - \hat{\gamma}\|_\infty = |\gamma((a + b)/2) - \hat{\gamma}((a + b)/2)| = \eta|\vec{v}|$.

Starting from the parameterization $\gamma_0 : [0, 1] \rightarrow \mathbb{R}^2$ of the line segment $I_0 = [\vec{0}, \vec{e}_1]$ from $\gamma_0(0) = \vec{0}$ to $\gamma_0(1) = \vec{e}_1$, we now define a sequence of piecewise linear maps $\gamma_i : [0, 1] \rightarrow \mathbb{R}^2$ by iteratively adding bumps of relative height η_i . Suppose that γ_i has been defined for some $i \geq 0$ so that $\gamma_i|_{J_{i,k}}$ is a constant-speed parameterization of a line segment on each interval $J_{i,k} = [a_{i,k}, b_{i,k}]$ of the form

$$J_{i,k} = \left[a + \frac{(k-1)}{4^i}(b-a), a + \frac{k}{4^i}(b-a) \right] \quad (1 \leq k \leq 4^i, k \in \mathbb{Z}).$$

For each index $1 \leq k \leq 4^i$, define $\gamma_{i+1}|_{J_{i,k}} = \widehat{\gamma_i|_{J_{i,k}}}$ by adding a bump of relative height η_{i+1} . This defines a map γ_{i+1} . By induction, we obtain the full sequence $\gamma_0, \gamma_1, \gamma_2, \dots$; the image of each map γ_n is composed of 4^n line segments of length $r_n := (\text{Lip } \gamma_n)4^{-n}$. Evidently, for all $n = 0, 1, 2, \dots$

$$(5.1) \quad \text{Lip}(\gamma_n) = \prod_{i=1}^n (1 + 4\eta_i^2) \leq (4/3)^n,$$

$$(5.2) \quad r_{n+1} \leq (1/3)r_n,$$

$$(5.3) \quad \|\gamma_n - \gamma_{n+1}\|_\infty \leq \eta_{n+1}r_n \leq 3^{-n}\eta_{n+1} < 3^{-n}.$$

Therefore, $\gamma : [0, 1] \rightarrow \mathbb{R}^2$, which is defined pointwise by

$$(5.4) \quad \gamma(t) = \gamma_0(t) + \sum_{n=0}^{\infty} (\gamma_{n+1}(t) - \gamma_n(t)) \quad \text{for all } t \in [0, 1],$$

is continuous as the uniform limit of the maps γ_n by (5.3). We denote $\gamma([0, 1])$ by Γ , and for each n , we denote $\gamma_n([0, 1])$ by Γ_n . For all integers $n, k \geq 0$ with $0 \leq k \leq 4^n$, we call the point $\gamma_n(k/4^n)$ a *vertex* of Γ_n .

REMARK 5.4 (Modulus of continuity)

In fact, (5.1) and (5.3) imply that the maps γ_n and γ are uniformly $\log_4(3)$ -Hölder continuous (see, for example, [16, Appendix B]). This is the optimal modulus of continuity of the von Koch snowflake curve, which corresponds to the choice of relative

heights $\eta_n = 1/\sqrt{12}$ for all n . Furthermore, if $\sum_{n=1}^{\infty} \eta_n^2 < \infty$, then (5.1) implies that γ_n and γ are uniformly Lipschitz with Lipschitz constant at most $\prod_{n=1}^{\infty} (1 + 4\eta_n^2) \simeq \exp(\sum_{n=1}^{\infty} \eta_n^2)$.

REMARK 5.5 (Vertices)

For each integer $n \geq 0$, let V_n denote the set of vertices in Γ_n and let $\widetilde{V}_n := V_n \setminus V_{n-1}$ denote the set of “new vertices” in Γ_n with the convention that $\widetilde{V}_0 = V_0$. For later use, we observe that

$$(5.5) \quad V_0 \subset V_1 \subset V_2 \subset \cdots,$$

$$(5.6) \quad V_n = \widetilde{V}_0 \cup \widetilde{V}_1 \cup \cdots \cup \widetilde{V}_{n-1} \cup \widetilde{V}_n \quad (\widetilde{V}_i \cap \widetilde{V}_j = \emptyset \text{ when } i \neq j),$$

$$(5.7) \quad \#\widetilde{V}_0 = 2, \quad \#\widetilde{V}_n = 3 \cdot 4^{n-1} (n \geq 1), \quad \#V_n = 1 + 4^n.$$

REMARK 5.6 (Injectivity)

The restriction $\eta_i \leq 1/\sqrt{12}$ on the relative heights ensures that the parameterization γ of Γ constructed by Algorithm 5.3 is injective. This can be shown by a geometric argument similar to the proof that the standard von Koch snowflake curve is the attractor of an iterated function system that satisfies the open set condition (that is, draw equilateral triangles on the left side of each segment in Γ_n). We leave details to the dedicated reader. For other models of generalized von Koch curves, the question of injectivity of a parameterization is quite subtle; see, for example, [51].

LEMMA 5.7

If Γ is constructed by Algorithm 5.3, then

$$(5.8) \quad \exp\left(3.33 \sum_{n=1}^{\infty} \eta_n^2\right) \leq \mathcal{H}^1(\Gamma) = \prod_{n=1}^{\infty} (1 + 4\eta_n^2) \leq \exp\left(4 \sum_{n=1}^{\infty} \eta_n^2\right).$$

In particular, Γ is a rectifiable curve if and only if $\sum_{n=1}^{\infty} \eta_n^2 < \infty$. Moreover, in that case, Γ is Ahlfors regular with constants depending only on $\mathcal{H}^1(\Gamma)$.

Proof

The 1-dimensional Hausdorff measure $\mathcal{H}^1$ enjoys the bound $\mathcal{H}^1(K) \geq \text{diam } K$ for every connected set $K \subset \mathbb{R}^2$ (see, for example, [3, Lemma 2.11]). Thus, for each $n \geq 1$,

$$\begin{aligned} \mathcal{H}^1(\Gamma) &= \sum_{k=1}^{4^n} \mathcal{H}^1(\gamma([(k-1)4^{-n}, k4^{-n}])) \geq \sum_{k=1}^{4^n} \text{diam } \gamma([(k-1)4^{-n}, k4^{-n}])) \\ &\geq \sum_{k=1}^{4^n} \text{diam } \gamma_n([(k-1)4^{-n}, k4^{-n}])) = \sum_{k=1}^{4^n} \prod_{i=1}^n (1/4 + \eta_i^2) = \prod_{i=1}^n (1 + 4\eta_i^2), \end{aligned}$$

where the initial equality holds by Remark 5.6. Hence, $\mathcal{H}^1(\Gamma) \geq \prod_{n=1}^{\infty} (1 + 4\eta_n^2)$. Conversely, $\mathcal{H}^1(\Gamma) \leq \text{Lip } \gamma \leq \prod_{n=1}^{\infty} (1 + 4\eta_n^2)$. Therefore,

$$\mathcal{H}^1(\Gamma) = \prod_{n=1}^{\infty} (1 + 4\eta_n^2) = \exp\left(\sum_{n=1}^{\infty} \log(1 + 4\eta_n^2)\right).$$

Finally, since $0 \leq 4\eta_n^2 \leq 1/3$, we may use Taylor's theorem to bound $\log(1 + 4\eta_n^2) \leq 4\eta_n^2$ and $\log(1 + 4\eta_n^2) \geq 4\eta_n^2 - \frac{1}{2}(4\eta_n^2)^2 \geq (10/3)\eta_n^2$.

Suppose that $\mathcal{H}^1(\Gamma) < \infty$. By (5.2), (5.3), and the bound $\eta_n \leq 1/\sqrt{12}$ for all $n \geq 1$,

$$\|\gamma_n - \gamma\|_{\infty} \leq \sum_{j=n}^{\infty} \eta_{j+1} r_j \leq \frac{1}{\sqrt{12}} \sum_{k=0}^{\infty} r_{n+k} \leq \frac{1}{\sqrt{12}} r_n \sum_{k=0}^{\infty} 3^{-k} = \frac{\sqrt{3}}{4} r_n < 0.45 r_n.$$

Thus, given $x \in \Gamma$ and $n \geq 1$, we may pick $y \in \Gamma_n$ such that $|x - y| \leq (\sqrt{3}/4)r_n$. Next, let $v \in V_n$ be an endpoint of the segment in Γ_n containing y that is closest to y so that $|v - y| \leq (1/2)r_n$. Hence, $B(x, 0.05r_n) \subset B(v, r_n)$ and we may use (5.8) to estimate

$$\begin{aligned} \mathcal{H}^1(\Gamma \cap B(x, 0.05r_n)) &\leq \mathcal{H}^1(\Gamma \cap B(v, r_n)) \leq 2 \exp\left(\sum_{i=n}^{\infty} 4\eta_i^2\right) r_n \\ (5.9) \quad &\leq 2e^{6/5} \mathcal{H}^1(\Gamma) r_n \end{aligned}$$

because $\Gamma_n \cap B(v, r_n)$ consists of one or two line segments of length r_n . From (5.9), one easily deduces that $\mathcal{H}^1(\Gamma \cap B(x, r)) \simeq_{\mathcal{H}^1(\Gamma)} r$ for every $x \in \Gamma$ and every $0 < r \leq 1 = \text{diam } \Gamma$. Therefore, Γ is Ahlfors regular with constants depending only on $\mathcal{H}^1(\Gamma)$. $\square$

LEMMA 5.8

Assume that Γ is constructed by Algorithm 5.3. For all $i \geq 0$ and $j \geq i$,

$$(5.10) \quad \frac{\sqrt{3}}{4} \eta_i \leq \beta_{\Gamma_j}(B(v, r_j)) \leq 2\eta_i \quad \text{for all } v \in \widetilde{V}_i,$$

where $\widetilde{V}_i$ denote the “new vertices” in Γ_i (see Remark 5.5) and $\eta_0 = 0$. Furthermore,

$$(5.11) \quad \frac{\sqrt{3}}{4} \eta_i \leq \beta_{\Gamma}(\overline{B(v, r_i)}) \quad \text{for all } i \geq 0 \text{ and } v \in \widetilde{V}_i.$$

Proof

Note that when $v \in \widetilde{V}_0$ is an endpoint of Γ_j , the set $\Gamma_j \cap B(v, r_j)$ is a line segment. Thus, $\beta_{\Gamma_j}(B(v, r_j)) = 0$ for all $v \in \widetilde{V}_0$ and $j \geq 0$. Next, suppose that $i \geq 1$ and $v \in \widetilde{V}_i$. Using the line containing the segment in $\Gamma_{i-1} \cap B(v, r_i)$ to approximate the beta number, we obtain the estimate

$$\beta_{\Gamma_i}(B(v, r_i)) \leq \frac{\eta_i}{2(\frac{1}{4} + \eta_i^2)} \leq 2\eta_i.$$

When $j \geq i$, the set $\Gamma_j \cap B(v, r_j)$ agrees up to a dilation centered at v with $\Gamma_i \cap B(v, r_i)$. Thus, $\beta_{\Gamma_j}(B(v, r_j)) \leq 2\eta_i$ for all $v \in \widetilde{V}_i$ and $j \geq i$, as well.

To establish the lower bound, one can use symmetry to find the best-fitting line for each of the two triangles formed by consecutive segments of side length s in Figure 4. The best-fitting lines are parallel to the third side of each triangle, and the altitudes of the triangles are η and $\sqrt{s}\eta = (\frac{1}{4} + \eta^2)^{1/2}\eta$. It follows that for all $v \in \tilde{V}_i$ and $j \geq i$,

$$\beta_{\Gamma_j}(B(v, r_j)) = \beta_{\Gamma_i}(B(v, r_i)) \geq \frac{\frac{1}{2}(\frac{1}{4} + \eta_i^2)^{1/2}\eta_i}{2(\frac{1}{4} + \eta_i^2)} = \frac{\eta_i}{4(\frac{1}{4} + \eta_i^2)^{1/2}} \geq \frac{\sqrt{3}}{4}\eta_i$$

since $\eta_i \leq 1/\sqrt{12}$. Finally, because $\Gamma \cap \overline{B(v, r_i)}$ contains the vertices a, b, c of one of the two triangles, we find that $\beta_\Gamma(\overline{B(v, r_i)}) \geq \beta_{\{a,b,c\}}(\overline{B(v, r_i)}) \geq (\sqrt{3}/4)\eta_i$, as well. $\square$

5.1.1. Proof of Proposition 5.1. Build Γ using Algorithm 5.3 with relative heights $4\eta_i^2 := 1/(i+15)\log(i+15)$ for all $i \geq 1$. See Remark 5.9 below for an explanation of the logarithmic factor in the relative heights. Note that $\eta_{i+1} \leq \eta_i < 1/8$ for all i . Because $\sum_{i=1}^\infty \eta_i^2 = \infty$, we have $\mathcal{H}^1(\Gamma) = \infty$ by Lemma 5.7. To proceed, let $(X_k)_{k \in \mathbb{Z}}$ be any family of nested 2^{-k} -nets for Γ , and let $\mathcal{G} = \{B(x, A2^{-k}) : x \in X_k, k \in \mathbb{Z}\}$ be the associated multiresolution family with inflation factor $A > 1$. Since Γ is bounded, to prove that $S_{\Gamma, 2+\epsilon}(\mathcal{G}) < \infty$, it suffices to show that (cf. Lemma 3.27)

$$S_{\Gamma, 2+\epsilon}(\mathcal{G}') := \sum_{B \in \mathcal{G}'} \beta_\Gamma(B)^{2+\epsilon} \text{diam } B < \infty,$$

where $\mathcal{G}'$ is the subfamily of all balls starting from some initial generation k_0 .

Recall that each intermediate curve Γ_n consists of 4^n line segments of length

$$r_n = 4^{-n} \prod_{i=1}^n (1 + 4\eta_i^2) = 4^{-n} \exp\left(\sum_{i=1}^n \log(1 + 4\eta_i^2)\right).$$

By Taylor's theorem, we have

$$4\eta_i^2 - \frac{1}{2}(4\eta_i^2)^2 \leq \log(1 + 4\eta_i^2) \leq 4\eta_i^2 \quad \text{for all } i \geq 1.$$

Combined with the elementary bounds

$$\begin{aligned} \sum_{i=1}^n \frac{1}{(i+15)\log(i+15)} &\leq \frac{1}{16\log(16)} + \int_{16}^{n+15} \frac{1}{x\log(x)} dx \\ &< \frac{1}{16} + \log(\log(n+15)) \end{aligned}$$

and

$$\begin{aligned} \sum_{i=1}^n \frac{1}{(i+15)\log(i+15)} - \frac{1}{2(i+15)^2\log(i+15)^2} \\ \geq \int_{16}^{n+16} \left(\frac{1}{x\log(x)} - \frac{1}{2x^2\log(x)^2} \right) dx > \log(\log(n+16)) - 1.2, \end{aligned}$$

we obtain the rough estimate (for the record, $0.3 < e^{-1.2}$ and $e^{1/16} < 1.1$):

$$(5.12) \quad 0.3\log(n+16)4^{-n} < r_n < 1.1\log(n+15)4^{-n} \quad \text{for all } n \geq 1.$$

To continue, let $k \geq k_0$ (with k_0 sufficiently large depending on A to be specified below) and let $x \in X_k$. Our immediate goal is to estimate $\beta_\Gamma(B(x, A2^{-k}))$ from above in terms of $\beta_{\Gamma_m}(B(v, r_m))$ for some suitably chosen generation $m = m(A, k) \geq 1$ and vertex $v \in \Gamma_m$. Write $c := \log_4 A > 0$ so that $A2^{-k} = 4^{-(\frac{1}{2}k-c)}$. We now require $\frac{1}{2}k_0 - c \geq 3$, which ensures

$$A2^{-k_0} \leq 4^{-3} < 0.03 \log(16)4^{-1}.$$

Since $k \geq k_0$, there exists a unique integer $m \geq 1$ with

$$(5.13) \quad 0.03 \log(m+16)4^{-(m+1)} < A2^{-k} \leq 0.03 \log(m+15)4^{-m}.$$

By (5.12), it follows that

$$(5.14) \quad r_m \lesssim A2^{-k} < \frac{1}{10}r_m.$$

Next, by (5.2), (5.3), and fact that $\eta_{i+1} \leq \eta_i < 1/8$ for all i ,

$$\begin{aligned} \|\gamma_m - \gamma\|_\infty &\leq \sum_{j=m}^{\infty} \eta_{j+1}r_j \leq \eta_{m+1} \sum_{l=0}^{\infty} r_{m+l} \leq \eta_{m+1}r_m \sum_{k=0}^{\infty} 3^{-k} \\ &= (3/2)\eta_{m+1}r_m < (3/16)r_m. \end{aligned}$$

In particular, we can find $y \in \Gamma_m$ with $|x - y| < (3/16)r_m$ and then choose a vertex v in Γ_m such that $|y - v| \leq (1/2)r_m$ (that is, v is an endpoint of the segment containing y). Since

$$\frac{1}{10} + \frac{3}{16} + \frac{1}{2} < \frac{3}{4},$$

$B(x, A2^{-k}) \subset B(v, (3/4)r_m)$. Invoking the bound $\|\gamma - \gamma_m\|_\infty \leq (3/2)\eta_{m+1}r_m < (3/16)r_m$ again, we conclude that

$$E := \text{excess}(\Gamma \cap B(x, A2^{-k}), \Gamma_m \cap B(v, r_m)) \leq (3/2)\eta_{m+1}r_m,$$

where $\text{excess}(S, T) = \sup_{s \in S} \inf_{t \in T} |s - t|$ denotes the *excess of S over T* . Hence,

$$\begin{aligned} (5.15) \quad \beta_\Gamma(B(x, A2^{-k})) &\leq \frac{E}{2A2^{-k}} + \frac{r_m}{A2^{-k}} \cdot \beta_{\Gamma_m}(B(v, r_m)) \\ &\lesssim \eta_{m+1} + \beta_{\Gamma_m}(B(v, r_m)) \end{aligned}$$

by the triangle inequality and (5.14). Note that by Lemma 5.8,

$$(5.16) \quad \eta_{m+1} + \beta_{\Gamma_m}(B(v, r_m)) \leq \eta_{m+1} + 2\eta_i \quad \text{when } v \in \widetilde{V}_i \subset V_m,$$

where $\eta_0 = 0$. In particular, $\beta_\Gamma(B(x, A2^{-k})) \lesssim \eta_{m+1}$ when $v \in \widetilde{V}_0$ and $\beta_\Gamma(B(x, A2^{-k})) \lesssim \eta_i$ when $v \in \widetilde{V}_i$ for some $1 \leq i \leq m$.

Next, we bound the number of times a scale m and vertex $v \in V_m$ are associated to a point $x \in X_k$. On one hand, $\#X_k \cap B(v, r_m) \lesssim A^2$ for each vertex $v \in V_m$ by (5.14) since X_k is 2^{-k} -separated and our construction takes place in $\mathbb{R}^2$. (We could remove the dimension dependence by using Lemma 2.1, but we do not require a sharp upper bound.) On the other hand, the scale m associated to $k \geq k_0$ satisfies

$$0.03 \log(m+16)4^{-(m+1)} < 4^{-(\frac{1}{2}k-c)} \leq 0.03 \log(m+15)4^{-m},$$

where $c = \log_4 A$. Taking logarithms and rearranging, we have

$$2(m + c - \log_4(0.03 \log(m + 15))) \leq k < 2(m + 1 + c - \log_4(0.03 \log(m + 16))).$$

Thus, the number of integers k associated to a given integer m is at most

$$2(m + 1 + c - \log_4(0.03 \log(m + 16))) - 2(m + c - \log_4(0.03 \log(m + 15))) \leq 2.$$

To finish, fix a parameter $\epsilon > 0$. In view the previous paragraph, (5.13), and (5.15), we see that

$$\begin{aligned} S_{\Gamma, 2+\epsilon}(\mathcal{G}') &= \sum_{k=k_0}^{\infty} 2A2^{-k} \sum_{x \in X_k} \beta_{\Gamma}(B(x, A2^{-k}))^{2+\epsilon} \\ (5.17) \quad &\lesssim_{A, \epsilon} \sum_{m=1}^{\infty} \log(m + 15) 4^{-m} \sum_{v \in V_m} (\eta_{m+1} + \beta_{\Gamma_m}(B(v, r_m)))^{2+\epsilon}. \end{aligned}$$

Decomposing $V_m = \tilde{V}_0 \cup \dots \cup \tilde{V}_m$ (see Remark 5.5) and invoking (5.16),

$$\begin{aligned} (5.18) \quad \sum_{v \in V_m} (\eta_{m+1} + \beta_{\Gamma_m}(B(v, r_m)))^{2+\epsilon} &\lesssim_{\epsilon} \sum_{v \in \tilde{V}_0} \eta_{m+1}^{2+\epsilon} + \sum_{i=1}^m \sum_{v \in \tilde{V}_i} \eta_i^{2+\epsilon} \\ &\lesssim_{\epsilon} 1 + \sum_{i=1}^m 4^{i-1} \eta_i^{2+\epsilon}. \end{aligned}$$

Combining the previous two displayed equations, it follows that

$$(5.19) \quad S_{\Gamma, 2+\epsilon}(\mathcal{G}') \lesssim_{A, \epsilon} \underbrace{\sum_{m=1}^{\infty} \log(m + 15) 4^{-m}}_I + \underbrace{\sum_{m=1}^{\infty} \log(m + 15) 4^{-m} \sum_{i=1}^m 4^{i-1} \eta_i^{2+\epsilon}}_{II}.$$

It is apparent that $I \lesssim 1$. To bound II , exchange the order of summation:

$$\begin{aligned} (5.20) \quad II &= \sum_{i=1}^{\infty} 4^{i-1} \eta_i^{2+\epsilon} \sum_{m=i}^{\infty} \log(m + 15) 4^{-m} \lesssim \sum_{i=1}^{\infty} 4^{i-1} \eta_i^{2+\epsilon} \log(i + 15) 4^{-i} \\ &\lesssim \sum_{i=1}^{\infty} \log(i + 15) \eta_i^{2+\epsilon} = \sum_{i=1}^{\infty} \log(i + 15) \left(\frac{1}{4(i + 15) \log(i + 15)} \right)^{1+\frac{1}{2}\epsilon}. \end{aligned}$$

Hence $II \lesssim_{\epsilon} 1$ and $S_{\Gamma, 2+\epsilon}(\mathcal{G}') \lesssim_{A, \epsilon} 1$. Therefore, by our initial discussion, $S_{\Gamma, 2+\epsilon}(\mathcal{G}) < \infty$ for every $\epsilon > 0$ and every multiresolution family $\mathcal{G}$ for Γ . This completes the proof of Proposition 5.1.

REMARK 5.9 (Importance of the logarithmic factor)

Seeking out examples verifying the sharpness of (1.17), it is natural to first look at snowflake curves Γ built with relative heights $\eta_i^2 \simeq 1/i$. By carrying out the outline above with relative heights $4\eta_i^2 = \delta/(i + i_0)$ with parameters $\delta > 0$ and $i_0 \geq 1$, one obtains

$$II \lesssim \sum_{i=1}^{\infty} (i + i_0)^{\delta} \left(\frac{1}{4(i + i_0)} \right)^{1+\frac{1}{2}\epsilon},$$

where the latter expression is finite precisely when $\delta < (1/2)\epsilon$. Thus, for every $\epsilon > 0$, we could build a curve Γ_ϵ with $\mathcal{H}^1(\Gamma_\epsilon) = \infty$ and $S_{\Gamma_\epsilon, 2+\epsilon}(\mathcal{G}) < \infty$ by selecting $\delta = \delta(\epsilon)$ sufficiently small. The logarithmic correction used in the proof of Proposition 5.1 allows us to find a single curve Γ such that $\mathcal{H}^1(\Gamma) = \infty$ and $S_{\Gamma, 2+\epsilon}(\mathcal{G}) < \infty$ for all $\epsilon > 0$.

5.1.2. Proof of Proposition 5.2. Construct Γ using Algorithm 5.3 with relative heights $4\eta_i^2 = 1/(i+2)\log(i+2)^2$ for all $i \geq 1$. Then $\eta_{i+1} \leq \eta_i < 1/\sqrt{12}$ for all $i \geq 1$ and

$$(5.21) \quad \sum_{i=1}^{\infty} 4\eta_i^2 = \sum_{n=3}^{\infty} \frac{1}{n \log(n)^2} = 1.069\dots < \infty.$$

Hence, $\mathcal{H}^1(\Gamma) \leq \exp(\sum_{i=1}^{\infty} 4\eta_i^2) < \exp(1.07) < 3$ by Lemma 5.7. Similarly, we have that the intermediate curves Γ_n consist of 4^n segments of length r_n , where

$$(5.22) \quad 4^{-n} < r_n \leq \exp\left(\sum_{i=1}^n 4\eta_i^2\right) 4^{-n} < 3 \cdot 4^{-n} \quad \text{for all } n \geq 1.$$

Let $(X_k)_{k \in \mathbb{Z}}$ be an arbitrary family of nested 2^{-k} -nets for Γ , let $A > 1$, and let $\mathcal{G} = \{B(x, A2^{-k}) : x \in X_k, k \in \mathbb{Z}\}$ be the associated multiresolution family for Γ . We wish to show that $S_{\Gamma, 2-\epsilon}(\Gamma) = \infty$ for all $\epsilon > 0$. By Lemma 5.7, Γ is Ahlfors regular with constants determined by $\mathcal{H}^1(\Gamma)$. In particular, we know that

$$\#X_k \simeq 2^k \quad \text{for every } k \geq 0.$$

Thus, writing $\beta(k) := \inf_{x \in X_k} \beta_\Gamma(B(x, A2^{-k}))$, we have

$$(5.23) \quad S_{\Gamma, 2-\epsilon}(\mathcal{G}) \geq \sum_{k=k_1}^{\infty} \sum_{x \in X_k} \beta_\Gamma(B(x, A2^{-k}))^{2-\epsilon} A2^{-k} \gtrsim_A \sum_{k=k_1}^{\infty} \beta(k)^{2-\epsilon}.$$

To proceed, we will bound $\beta(k)$ from below in terms of η_m for sufficiently large k .

Choose k_0 sufficiently large such that $A2^{-k_0} < 6$. Suppose that $k \geq k_0$. Let $m(k) \geq 1$ be the unique integer such that $6 \cdot 4^{-m} \leq A2^{-k} < 6 \cdot 4^{-(m-1)}$. By (5.22), we have

$$(5.24) \quad 2r_m \leq A2^{-k} \lesssim r_m.$$

Given $x \in X_k$, choose $v \in V_m$ such that $|x - v| < r_m$ (cf. proof of Lemma 5.7). Then $B(v, r_m) \subset B(x, 2r_m) \subset B(x, A2^{-k})$. Hence,

$$\beta_\Gamma(B(x, A2^{-k})) \geq \frac{r_m}{A2^{-k}} \beta_\Gamma(B(v, r_m)) \gtrsim \eta_m$$

by Lemma 5.8. Because $x \in X_k$ was arbitrary, $\beta(k) \gtrsim \eta_m$. Now, $m \leq \frac{1}{2}k + \log_4(6/A) + 1$. Choose $k_1 \geq k_0$ sufficiently large such that $\log_4(6/A) + 3 \leq \frac{1}{2}k_1$. Then, for every $k \geq k_1$, we have $m + 2 \leq k$ and

$$\beta(k) \gtrsim \eta_m \gtrsim \frac{1}{(m+2)^{1/2} \log(m+2)} \gtrsim \frac{1}{k^{1/2} \log(k)}.$$

Therefore, for every $\epsilon > 0$,

$$S_{\Gamma, 2-\epsilon}(\mathcal{G}) \gtrsim_A \sum_{k=k_1}^{\infty} \beta(k)^{2-\epsilon} \gtrsim_A \sum_{k=k_1}^{\infty} k^{\frac{1}{2}\epsilon-1} \log(k)^{\epsilon-2} = \infty.$$

This completes the proof of Proposition 5.2.

5.2. Examples with critical exponent $p \neq 2$

To complete the proof of Theorems 1.6 and 1.7, we return to the infinite-dimensional setting. Our goal is to establish the following.

PROPOSITION 5.10

For all $1 < p < \infty$, there is a curve Γ in ℓ_p such that $\mathcal{H}^1(\Gamma) = \infty$ and $S_{\Gamma, p+\epsilon}(\mathcal{G}) < \infty$ for every multiresolution family $\mathcal{G}$ for Γ and every $\epsilon > 0$. In particular, the exponent p in (1.17) when $1 < p \leq 2$ is sharp.

PROPOSITION 5.11

For all $1 < p < \infty$, there is a curve Γ in ℓ_p such that $\mathcal{H}^1(\Gamma) < \infty$ and $S_{\Gamma, p-\epsilon}(\mathcal{G}) = \infty$ for every multiresolution family $\mathcal{G}$ for Γ and every $\epsilon > 0$. In particular, the exponent p in (1.18) when $2 \leq p < \infty$ is sharp.

We construct the curves in both propositions using the following algorithm, which is inspired by examples of Edelen, Naber, and Valtorta [38, Section 5.2] in $L^p([0, 1])$. The key point is that because we are working in an infinite-dimensional space, we may build each intermediate iteration of the snowflake by adding bumps in a new coordinate direction.

ALGORITHM 5.12 (Snowflake-like curves in ℓ_p with bumps along coordinate directions)

Let $\{e_i\}_{i=1}^{\infty}$ denote the standard basis in ℓ_p ; that is, $e_i(j) = \delta_{ij}$. Suppose $x, y \in \text{span}\{e_1, \dots, e_k\}$ and $\gamma : [a, b] \rightarrow \text{span}\{e_1, \dots, e_k\} \cong \ell_p^k$ is a constant-speed parameterization of $I = [x, y]$ from $\gamma(a) = x$ to $\gamma(b) = y$; that is,

$$\gamma(t) = x + \frac{t-a}{b-a}(y-x) \quad \text{for all } t \in [a, b].$$

Write $v := y - x$. Given a *relative height* $0 \leq \eta < 1/2$, we define a piecewise linear path $\hat{\gamma} : [a, b] \rightarrow \text{span}\{e_1, \dots, e_{k+1}\} \cong \ell_p^{k+1}$, as follows. Define $s \in [1/4, 1/2)$ to be the unique solution of $s^p = (\frac{1}{2} - s)^p + \eta^p$. Divide $[a, b]$ into quarters. For all $a \leq t \leq (3/4)a + (1/4)b$,

$$\hat{\gamma}(t) = x + 4\left(\frac{t-a}{b-a}\right)sv$$

(cf. Algorithm 5.3). For all $(3/4)a + (1/4)b \leq t \leq (1/2)a + (1/2)b$,

$$\hat{\gamma}(t) = x + sv + 4\left(\frac{t - (3/4)a - (1/4)b}{b-a}\right)((1/2 - s)v + \eta|v|_p e_{k+1}).$$

For all $(1/2)a + (1/2)b \leq t \leq (1/4)a + (3/4)b$,

$$\hat{\gamma}(t) = x + (1/2)v + \eta|v|e_{k+1} + 4\left(\frac{t - (1/2)a - (1/2)b}{b - a}\right)((1/2 - s)v - \eta|v|_p e_{k+1}).$$

Finally, for all $(1/4)a + (3/4)b \leq t \leq b$,

$$\hat{\gamma}(t) = x + (1 - s)v + 4\left(\frac{t - (1/4)a - (3/4)b}{b - a}\right)sv.$$

We say that $\hat{\gamma}$ is obtained from γ by *adding a bump of relative height η in the direction e_{k+1}* . On each quarter of $[a, b]$, $\hat{\gamma}(t)$ traces a line segment in ℓ_p of length $s|v|_p$ at constant speed. Thus, $\hat{\gamma}$ has Lipschitz constant $4s|v|_p/(b - a)$. Additionally, we have $\|\gamma - \hat{\gamma}\|_\infty = |\gamma((a + b)/2) - \hat{\gamma}((a + b)/2)|_p = \eta|v|_p$.

Starting from the arc-length parameterization $\gamma_0 : [0, 1] \rightarrow \text{span}\{e_1\}$ of the line segment $I_0 = [0, e_1]$ from $\gamma_0(0) = 0$ to $\gamma_0(1) = e_1$, we now define a sequence of piecewise linear maps $\gamma_i : [0, 1] \rightarrow \text{span}\{e_1, \dots, e_{i+1}\}$ by iteratively adding bumps of relative height η_i in the direction e_{i+1} . Suppose that γ_i has been defined for some $i \geq 0$ so that $\gamma_i|_{J_{i,k}}$ is a constant-speed parameterization of a line segment in $\text{span}\{e_1, \dots, e_{i+1}\}$ on each interval $J_{i,k} = [a_{i,k}, b_{i,k}]$ of the form

$$J_{i,k} = \left[a + \frac{(k-1)}{4^i}(b-a), a + \frac{k}{4^i}(b-a) \right] \quad (1 \leq k \leq 4^i, k \in \mathbb{Z}).$$

For each index $1 \leq k \leq 4^i$, define $\gamma_{i+1}|_{J_{i,k}} = \widehat{\gamma_i|_{J_{i,k}}}$ by adding a bump of relative height η_{i+1} in the direction e_{i+2} . This defines a map γ_{i+1} . By induction, we obtain the full sequence $\gamma_0, \gamma_1, \gamma_2, \dots$; the image of each map γ_n is composed of 4^n line segments in $\text{span}\{e_1, \dots, e_{n+1}\}$ of equal length $r_n = (\text{Lip } \gamma_n)4^{-n} = s_1 \cdots s_n$, where $s_i \in [1/4, 1/2)$ denotes the solution to $s_i^p = (\frac{1}{2} - s_i)^p + \eta_i^p$. Moreover,

$$(5.25) \quad \text{Lip}(\gamma_n) = \prod_{i=1}^n 4s_i < 2^n,$$

$$(5.26) \quad r_{n+1} \leq (1/2)r_n,$$

$$(5.27) \quad \|\gamma_n - \gamma_{n+1}\|_\infty \leq \eta_{n+1}r_n \leq 2^{-n}\eta_{n+1} < 2^{-n}.$$

Therefore, $\gamma : [0, 1] \rightarrow \ell_p$, which is defined pointwise by

$$(5.28) \quad \gamma(t) = \gamma_0(t) + \sum_{n=0}^{\infty} (\gamma_{n+1}(t) - \gamma_n(t)) \quad \text{for all } t \in [0, 1],$$

is continuous as the uniform limit of the maps γ_n by (5.27). We denote $\gamma([0, 1])$ by Γ , and for each n , we denote $\gamma_n([0, 1])$ by Γ_n . For all integers $n, k \geq 0$ with $0 \leq k \leq 4^n$, we call the point $\gamma_n(k/4^n)$ a *vertex* of Γ_n . Remark 5.5 holds in this setting, as well.

REMARK 5.13 (Modulus of continuity and injectivity)

In fact, (5.25) and (5.27) imply that the maps γ_n and γ are uniformly $(1/2)$ -Hölder continuous. The parameterization γ is injective, and this can be verified by showing that $\pi_1 \circ \gamma$ is injective, where $\pi_1 : \ell_p \rightarrow \mathbb{R}$ is projection onto the first coordinate. We leave details for the reader.

REMARK 5.14 (Improved bounds on r_n)

Let Γ in ℓ_p ($1 < p < \infty$) be constructed by Algorithm 5.12 with relative heights $0 \leq \eta_i \leq \bar{\eta}$ for all i , for some universal constant $\bar{\eta}$ to be specified below. Let $s_i \in [1/4, 1/2]$ be defined by $s_i^p = (\frac{1}{2} - s_i)^p + \eta_i^p$. Then

$$s_i \leq ((1/4)^p + \eta_i^p)^{1/p} = \frac{1}{4}(1 + (4\eta_i)^p)^{1/p}.$$

Since $p > 1$, Taylor's theorem with remainder gives

$$1 + \frac{1}{p}\delta - \frac{p-1}{2p^2}\delta^2 \leq (1 + \delta)^{1/p} \leq 1 + \frac{1}{p}\delta \quad \text{for all } 0 \leq \delta < 1.$$

On the one hand, assume that $\bar{\eta} < 1/4$. Then $(4\eta_i)^p < 4\bar{\eta} < 1$, and we obtain

$$s_i \leq \frac{1}{4}\left(1 + \frac{1}{p}(4\eta_i)^p\right) = \frac{1}{4} + \frac{1}{4p}(4\eta_i)^p.$$

On the other hand, assume that $\bar{\eta} < 1/8$. Then $s_i \leq 1/4 + (1/4p)2^{-p} < 3/8$. Hence, $1/8 < \frac{1}{2} - s_i \leq 1/4$, and using the Taylor bound we can write

$$\begin{aligned} s_i &= \left(\frac{1}{2} - s_i\right)\left(1 + \left(\frac{1}{2} - s_i\right)^{-p}\eta_i^p\right)^{1/p} \\ &\geq \frac{1}{2} - s_i + \frac{1}{p}\left(\frac{1}{2} - s_i\right)^{1-p}\eta_i^p - \frac{p-1}{2p^2}\left(\frac{1}{2} - s_i\right)^{1-2p}\eta_i^{2p} \\ &\geq \frac{1}{2} - s_i + \frac{1}{4p}(4\eta_i)^p - \frac{p-1}{16p^2}(8\eta_i)^{2p}. \end{aligned}$$

Rearranging the inequality, we obtain

$$s_i \geq \frac{1}{4} + \frac{1}{8p}(4\eta_i)^p - \frac{p-1}{32p^2}(8\eta_i)^{2p} = \frac{1}{4} + \left[\frac{1}{8p} - \frac{p-1}{32p^2}(16\eta_i)^p\right](4\eta_i)^p.$$

We now specify that $\bar{\eta} = 1/16$ so that

$$\frac{1}{8p} - \frac{p-1}{32p^2}(16\eta_i)^p \geq \frac{1}{8p} - \frac{p-1}{32p^2} \geq \frac{3}{32p}.$$

Therefore,

$$(5.29) \quad \frac{1}{4} + \frac{3}{32p}(4\eta_i)^p \leq s_i \leq \frac{1}{4} + \frac{1}{4p}(4\eta_i)^p \quad \text{whenever } \eta_i \leq \frac{1}{16}.$$

Thus, if $\eta_i \leq 1/16$ for all i , then the length $r_n = \prod_{i=1}^n s_i$ of each segment in Γ_n satisfies

$$(5.30) \quad 4^{-n} \prod_{i=1}^n \left(1 + \frac{3}{8p}(4\eta_i)^p\right) \leq r_n \leq 4^{-n} \prod_{i=1}^n \left(1 + \frac{1}{p}(4\eta_i)^p\right).$$

The reader may verify that when $p = 2$, the bound (5.30) is compatible with (5.1).

LEMMA 5.15

If Γ is constructed by Algorithm 5.12 with relative heights $\eta_i \leq 1/16$, then

$$(5.31) \quad \exp\left(\frac{1}{4p} \sum_{n=1}^{\infty} (4\eta_n)^p\right) \leq \mathcal{H}^1(\Gamma) \leq \exp\left(\frac{1}{p} \sum_{n=1}^{\infty} (4\eta_n)^p\right).$$

In particular, Γ is a rectifiable curve if and only if $\sum_{n=1}^{\infty} \eta_n^p < \infty$. Moreover, in that case, Γ is Ahlfors regular with constants depending only on $\mathcal{H}^1(\Gamma)$.

Proof

The outline is similar to the proof of Lemma 5.7. For each $n \geq 1$,

$$\begin{aligned} \mathcal{H}^1(\Gamma) &\stackrel{\text{Rk 5.13}}{=} \sum_{k=1}^{4^n} \mathcal{H}^1(\gamma([(k-1)4^{-n}, k4^{-n}])) \geq \sum_{k=1}^{4^n} \text{diam } \gamma([(k-1)4^{-n}, k4^{-n}])) \\ &\stackrel{(5.30)}{\geq} \sum_{k=1}^{4^n} 4^{-n} \prod_{i=1}^n \left(1 + \frac{3}{8} p^{-1} (4\eta_i)^p\right) = \prod_{i=1}^n \left(1 + \frac{3}{8} p^{-1} (4\eta_i)^p\right). \end{aligned}$$

We conclude that

$$\prod_{n=1}^{\infty} \left(1 + \frac{3}{8p} (4\eta_n)^p\right) \leq \mathcal{H}^1(\Gamma) \leq \text{Lip } \gamma \leq \liminf_{m \rightarrow \infty} \text{Lip } \gamma_m \stackrel{(5.30)}{\leq} \prod_{n=1}^{\infty} \left(1 + \frac{1}{p} (4\eta_n)^p\right).$$

To derive (5.31), rewrite each infinite product as the exponential of an infinite sum and use Taylor's theorem bounds for $\log(1+x)$ with $0 \leq x \leq 1/4$.

Suppose that $\mathcal{H}^1(\Gamma) < \infty$. By (5.26), (5.27), and assumption $\eta_n \leq 1/16$ for all $n \geq 1$,

$$\|\gamma_n - \gamma\|_{\infty} \leq \sum_{j=n}^{\infty} \eta_{j+1} r_j \leq \frac{1}{16} \sum_{k=0}^{\infty} r_{n+k} \leq \frac{1}{16} r_n \sum_{k=0}^{\infty} 2^{-k} = \frac{1}{8} r_n.$$

Thus, given $x \in \Gamma$ and $n \geq 1$, we may pick $y \in \Gamma_n$ such that $|x - y| \leq (1/8)r_n$. Next, let $v \in V_n$ be an endpoint of the segment in Γ_n containing y that is closest to y so that $|v - y| \leq (1/2)r_n$. Hence, $B(x, (3/8)r_n) \subset B(v, r_n)$ and we may use (5.8) to estimate

$$\begin{aligned} \mathcal{H}^1\left(\Gamma \cap B\left(x, \frac{3}{8}r_n\right)\right) &\leq \mathcal{H}^1(\Gamma \cap B(v, r_n)) \leq 2 \exp\left(\sum_{i=n}^{\infty} \frac{1}{p} (4\eta_i)^p\right) r_n \\ (5.32) \quad &\leq 2e^4 \mathcal{H}^1(\Gamma) r_n \end{aligned}$$

because $\Gamma_n \cap B(v, r_n)$ consists of one or two line segments of length r_n . It follows that Γ is Ahlfors regular with constants depending only on $\mathcal{H}^1(\Gamma)$. $\square$

LEMMA 5.16

Assume that Γ is constructed by Algorithm 5.12 with $\eta_i < 1/8$ for all i . For all $i \geq 0$ and $j \geq i$,

$$(5.33) \quad \frac{1}{4} \eta_i \leq \beta_{\Gamma_j}(B(v, r_j)) \leq 2\eta_i \quad \text{for all } v \in \widetilde{V}_i,$$

where $\widetilde{V}_i$ denote the “new vertices” in Γ_i (see Remark 5.5) and $\eta_0 = 0$. Furthermore,

$$(5.34) \quad \frac{1}{4} \eta_i \leq \beta_{\Gamma}(\overline{B(v, r_i)}) \quad \text{for all } i \geq 0 \text{ and } v \in \widetilde{V}_i.$$

Proof

The upper bound agrees with the case $p = 2$ above. Note that when $v \in \widetilde{V}_0$ is an

endpoint of Γ_j , the set $\Gamma_j \cap B(v, r_j)$ is a line segment. Thus, $\beta_{\Gamma_j}(B(v, r_j)) = 0$ for all $v \in \widetilde{V}_0$ and $j \geq 0$. Next, suppose that $i \geq 1$ and $v \in \widetilde{V}_i$. Using the line containing the segment in $\Gamma_{i-1} \cap B(v, r_i)$ to approximate the beta number, we obtain the estimate

$$\beta_{\Gamma_i}(B(v, r_i)) \leq \frac{\eta_i r_{i-1}}{2r_i} = \frac{\eta_i}{2s_i} \leq 2\eta_i,$$

where $\sup_{x \in \Gamma_i} \text{dist}(x, \ell) \leq \eta_i r_{i-1}$ because at stage i in the construction, we added the bump in the direction e_{i+1} and we recall that $s_i \geq \frac{1}{4}$. When $j \geq i$, the set $\Gamma_j \cap B(v, r_j)$ agrees up to a dilation centered at v with $\Gamma_i \cap B(v, r_i)$. Thus, $\beta_{\Gamma_j}(B(v, r_j)) \leq 2\eta_i$ for all $v \in \widetilde{V}_i$ and $j \geq i$, as well.

Before we determine the lower bounds in (5.33) and (5.34), we recall two basic properties of the beta numbers. First, for any set $E \subset B(v, r)$, we have $\beta_E(B(v, r)) = \beta_{\overline{E}}(\overline{B(v, r)})$. (This may fail if $E \not\subset B(v, r)$!) Second, $\beta_E(B(v, r))$ is increasing in E . Because $V_i \subset \Gamma_j$ for all $j \geq i$ and $V_i \subset \Gamma$, for any $v_k \in V_i$ the vertices v_{k-1} and v_{k+1} that are adjacent with respect to the global parametrization satisfy

$$\{v_{k-1}, v_k, v_{k+1}\} \subset \overline{\Gamma_j \cap B(v_k, r_i)} \quad \text{for any } j \geq i,$$

and the inclusion also holds with Γ in place of Γ_j . Therefore, up to a translation and dilation, there are two relevant configurations of vertices:

$$E_v(\eta) := \{0\} \cup \{sv\} \cup \left\{ \frac{1}{2}v + \eta e_{n+1} \right\},$$

$$F_v(\eta) := \{sv\} \cup \left\{ \frac{1}{2}v + \eta e_{n+1} \right\} \cup \{(1-s)v\},$$

where $v \in \text{span}\{e_1, \dots, e_n\}$ is an arbitrary vector with $|v|_p = 1$ and $s^p = (\frac{1}{2} - s)^p + \eta^p$ with $\eta < 1/8$. By Remark 5.14, $1/4 \leq s < 3/8$. The optimal lower bound in (5.33) and (5.34) is given by $\beta(\eta_i)$, where

$$\beta(\eta) := \min \left\{ \beta_{E_v(\eta)}(\overline{B(sv, s)}), \beta_{F_v(\eta)}\left(\overline{B\left(\frac{1}{2}v + \eta e_{n+1}, s\right)}\right) \right\}.$$

We work separately for each configuration, beginning with $E_v(\eta)$. Let L_1 be the line containing $[0, \frac{1}{2}v + \eta e_{n+1}]$. In the $v e_{n+1}$ -plane, the line L_1 is the locus of points (x, y) satisfying $y = 2\eta x$. When $x \geq \frac{1}{4} - \frac{1}{2}\eta$,

$$y \geq 2\eta \left(\frac{1}{4} - \frac{1}{2}\eta \right) \geq \frac{1}{2}\eta - \frac{1}{16}\eta = \frac{7}{16}\eta.$$

Since $s \geq \frac{1}{4}$, it follows that $B(sv, (7/16)\eta) \cap L_1 = \emptyset$. Hence,

$$(5.35) \quad \text{dist}(sv, L_1) \geq (7/16)\eta.$$

We now argue that $\beta_{E_v(\eta)}(\overline{B(sv, s)}) \geq (3/32s)\eta$ by way of contradiction. Assume that we can find a line L such that $E_v(\eta) \subset B_{(3/16)\eta}(L)$. By convexity, $L_1 \cap B(sv, s) \subset B_{(3/16)\eta}(L)$, as well. Let $x_1 \in L \cap B(0, (3/16)\eta)$ and $x_2 \in L \cap B(\frac{1}{2}v + \eta e_{n+1}, (3/16)\eta)$. Once again, by convexity, the segment of L that falls between x_1 and x_2 is contained in $B_{(3/16)\eta}(L_1)$. Since $\text{dist}(x, sv) \geq \frac{1}{2}s \geq (3/16)\eta$ for all $x \in L \setminus B_{(3/16)\eta}(L_1)$, the points $x \in L$ such that $\text{dist}(x, sv) \leq (3/16)\eta$ are contained

in $B_{(3/16)\eta}(L_1)$. Thus, by the triangle inequality,

$$(5.36) \quad E_v(\eta) \subset B_{(6/16)\eta}(L_1).$$

Since (5.35) and (5.36) are incompatible, we have reached a contradiction. Therefore,

$$\beta_{E_v(\eta)}(\overline{B(sv, s)}) \geq \frac{3}{32s}\eta \geq \frac{1}{4}\eta.$$

Using symmetry, one sees the best-fitting line for $F_v(\eta)$ is $\frac{1}{2}\eta e_{n+1} + \text{span}(v)$, whence

$$\beta_{F_v(\eta)}\left(\overline{B\left(\frac{1}{2}v + \eta e_{n+1}, s\right)}\right) \geq \frac{\eta}{4s} \geq \frac{2}{3}\eta.$$

Thus, the extremal configuration is given by $E_v(\eta)$ and $\beta(\eta) \geq (1/4)\eta$. $\square$

5.2.1. Proof of Proposition 5.10. Let $1 < p < \infty$. Let Γ be the curve constructed by Algorithm 5.12 with relative heights $\eta_i^p = \delta/(i + i_0) \log(i + i_0)$ for all $i \geq 1$, where $\delta > 0$ and i_0 are chosen so that $\eta_1 \leq 1/16$. Then $\mathcal{H}^1(\Gamma) = \infty$ in ℓ_p by Lemma 5.15. To prove that $S_{\Gamma, p+\epsilon}(\mathcal{G}) < \infty$ for every multiresolution family $\mathcal{G}$ for Γ and $\epsilon > 0$, repeat the proof of Proposition 5.1, *mutatis mutandis*, using Lemma 5.16 in lieu of Lemma 5.8.

5.2.2. Proof of Proposition 5.11. Let $1 < p < \infty$. Let Γ be the curve constructed by Algorithm 5.12 with relative heights $\eta_i^p = \delta/(i + i_0) \log(i + i_0)^2$ for all $i \geq 1$, where $\delta > 0$ and i_0 are chosen so that $\eta_1 \leq 1/16$. Then $\mathcal{H}^1(\Gamma) < \infty$ in ℓ_p and Γ is Ahlfors regular by Lemma 5.15. To prove that $S_{\Gamma, p-\epsilon}(\mathcal{G}) = \infty$ for every multiresolution family $\mathcal{G}$ for Γ and $\epsilon > 0$, repeat the proof of Proposition 5.2, *mutatis mutandis*, again substituting Lemma 5.16 for Lemma 5.8.

5.3. Proof of Proposition 1.1

Let $1 < p < q < \infty$. To begin, we verify that if $\Gamma \subset \ell_p$ is a rectifiable curve, then Γ is also rectifiable when viewed as a curve embedded in ℓ_q . As is well known, $\ell_p \subset \ell_q$ and $|v|_q \leq |v|_p$ for every $v \in \ell_p$. Hence, the diameter of a set in ℓ_p does not increase when embedded into ℓ_q . Thus, $\mathcal{H}_{\ell_q}^s(E) \leq \mathcal{H}_{\ell_p}^s(E)$ for every $s > 0$ and $E \subset \ell_p$, where $\mathcal{H}_{\ell_r}^s$ denotes the s -dimensional Hausdorff measure in ℓ_r . In particular, every rectifiable curve in ℓ_p is also a rectifiable curve in ℓ_q , possibly with shorter length.

We now construct a curve Γ such that $\mathcal{H}_{\ell_p}^1(\Gamma) = \infty$ and $\mathcal{H}_{\ell_q}^1(\Gamma) < \infty$ for every $q > p$. Build Γ in ℓ_p using Algorithm 5.12 with relative heights

$$\eta_i^p = \frac{\delta}{i \log(i + i_0)} \quad \text{for all } i \geq 1,$$

where $\delta > 0$ and $i_0 \in \mathbb{N}$ are chosen so that $0 < \eta_1 \leq 1/16$. Note that $\sum_{i=1}^{\infty} \eta_i^p = \infty$. Therefore, $\mathcal{H}_{\ell_p}^1(\Gamma) = \infty$ by Lemma 5.15.

Fix an exponent q with $p < q < \infty$. We break the proof that $\mathcal{H}_{\ell_q}^1(\Gamma) < \infty$ into a series of lemmata. First, we calculate the $\mathcal{H}_{\ell_q}^1$ -growth of a line segment under the snowflaking procedure $\gamma \mapsto \hat{\gamma}$ used in Algorithm 5.12. We emphasize that the estimate in the following lemma is independent of n .

LEMMA 5.17

Given $v \in \text{span}\{e_1, \dots, e_n\}$ with $|v|_p = 1$, let $\gamma : [0, 1] \rightarrow \text{span}\{e_1, \dots, e_n\}$ be the unit speed parameterization of the line segment $I = [0, v]$ in ℓ_p . Let $\hat{\gamma}$ be obtained from γ by adding a bump of relative height η in the direction e_{n+1} (see Algorithm 5.12). If $\eta \leq 1/16$, then $\mathcal{H}_{\ell_q}^1(I) = |v|_q$ and

$$(5.37) \quad \mathcal{H}_{\ell_q}^1(\hat{\gamma}([0, 1])) - \mathcal{H}_{\ell_q}^1(I) \lesssim_q \left(\eta/|v|_q\right)^q |v|_q.$$

Proof

The estimate is nearly identical to the calculation in Remark 5.14. Since $\eta \leq 1/16$, each of the four edges of $\hat{\gamma}(I)$ in ℓ_p have length $s \in [1/4, 3/8)$ given implicitly by the equation $s^p = (\frac{1}{2} - s)^p + \eta^p$. In ℓ_q , $\mathcal{H}_{\ell_q}^1(I) = |v|_q$ (since I is a segment) and the four edges of $\hat{\gamma}(I)$ have total length

$$\begin{aligned} \mathcal{H}_{\ell_q}^1(\hat{\gamma}([0, 1])) &= 2s|v|_q + 2\left(\left(\frac{1}{2} - s\right)^q |v|_q^q + \eta^q\right)^{1/q} \\ &= 2s|v|_q + (1 - 2s)|v|_q \left(1 + \left(\frac{1}{2} - s\right)^{-q} |v|_q^{-q} \eta^q\right)^{1/q} \\ &\leq |v|_q \left(1 + \left(\frac{1}{2} - s\right)^{-q} |v|_q^{-q} \eta^q\right)^{1/q}. \end{aligned}$$

Applying $(1 + \delta)^{1/q} \leq 1 + \frac{1}{q}\delta$ for all $\delta \geq 0$ with $\delta = (\frac{1}{2} - s)^{-q} |v|_q^{-q} \eta^q$, we conclude

$$\mathcal{H}_{\ell_q}^1(\hat{\gamma}([0, 1])) \leq |v|_q + \frac{1}{q} \left(\frac{\eta}{(\frac{1}{2} - s)|v|_q}\right)^q |v|_q \leq |v|_q + \frac{1}{q} \left(\frac{8\eta}{|v|_q}\right)^q |v|_q. \quad \square$$

The next estimate is elementary and left as an exercise for the reader.

LEMMA 5.18

For all $n \in \mathbb{N}$ and $v \in \ell_p^n$, $|v|_q \geq n^{\frac{1}{q} - \frac{1}{p}} |v|_p$.

We may now give a uniform estimate on the length of the intermediate curves Γ_n approximating Γ in ℓ_q .

LEMMA 5.19

With the exponents and the parameter $\eta_i^p = \delta/(i \log(i + i_0))$ fixed as above, the intermediate curves produced by Algorithm 5.12 satisfy

$$\mathcal{H}_{\ell_q}^1(\Gamma_n) \leq C(p, q) < \infty \quad \text{for all } n \geq 1,$$

where $C(p, q)$ is a constant depending on p and q with $C(p, q) \uparrow \infty$ as $q \downarrow p$.

Proof

The initial segment $\Gamma_0 = [0, e_1]$ satisfies $\mathcal{H}_{\ell_q}^1(\Gamma_0) = 1$. Let $n \geq 0$, let $1 \leq k \leq 4^n$, and let $J = \gamma_n([(k-1)4^{-n}, k4^{-n}])$ be an edge in Γ_n . Assign $J^+ := \gamma_{n+1}([(k-1)$

$1)4^{-n}, k4^{-n}]$). By Lemmas 5.17 and 5.18,

$$\mathcal{H}_{\ell_q}^1(J^+) - \mathcal{H}_{\ell_q}^1(J) \lesssim_q \left(\frac{\eta_{n+1} \mathcal{H}_{\ell_p}^1(J)}{\mathcal{H}_{\ell_q}^1(J)} \right)^q \mathcal{H}_{\ell_q}^1(J) \lesssim_q \left(\frac{\eta_{n+1}}{(n+1)^{\frac{1}{q}-\frac{1}{p}}} \right)^q \mathcal{H}_{\ell_q}^1(J).$$

Summing over the 4^n edges J in Γ_n , it follows that

$$\mathcal{H}_{\ell_q}^1(\Gamma_{n+1}) \leq \left(1 + C_q \left(\frac{\eta_{n+1}}{(n+1)^{\frac{1}{q}-\frac{1}{p}}} \right)^q \right) \mathcal{H}_{\ell_q}^1(\Gamma_n),$$

where C_q is a positive constant depending only on q . Therefore,

$$\mathcal{H}_{\ell_q}^1(\Gamma_n) \leq \prod_{i=1}^n \left(1 + C_q \left(\frac{\eta_i}{i^{\frac{1}{q}-\frac{1}{p}}} \right)^q \right) \leq \exp \left(C_q \sum_{i=1}^n \left(\frac{\eta_i}{i^{\frac{1}{q}-\frac{1}{p}}} \right)^q \right).$$

Finally, recalling our choice of η_i ,

$$\sum_{i=1}^n \left(\frac{\eta_i}{i^{\frac{1}{q}-\frac{1}{p}}} \right)^q \leq \sum_{i=1}^{\infty} \left(\frac{\delta^{1/p} i^{-1/p} \log(i + i_0)^{-1/p}}{i^{\frac{1}{q}-\frac{1}{p}}} \right)^q = \sum_{i=1}^{\infty} \frac{\delta^{q/p}}{i \log(i + i_0)^{q/p}} < \infty$$

because $q/p > 1$. All together, $\mathcal{H}_{\ell_q}^1(\Gamma_n) \leq C(p, q) < \infty$, where $C(p, q) \uparrow \infty$ when $q \downarrow p$. $\square$

By (5.27), γ_n converges uniformly to γ in ℓ_p , and thus in ℓ_q . Hence, Γ_n converges to Γ in the Hausdorff metric on compact subsets of ℓ_q . Therefore, by Gołab's semicontinuity theorem (see, for example, [3, Theorem 2.9]), $\mathcal{H}_{\ell_q}^1(\Gamma) \leq \liminf_{n \rightarrow \infty} \mathcal{H}_{\ell_q}^1(\Gamma_n) \leq C(p, q) < \infty$ for all $q > p$. This completes the proof of Proposition 1.1.

Appendix. Schul's prefiltration lemma in a metric space

Before we start the proof of Lemma 3.20, we record proofs of a couple of metric lemmas. Recall that a *pseudometric* on $\mathbb{X}$ is a function $d : \mathbb{X} \times \mathbb{X} \rightarrow [0, \infty)$ such that $d(x, x) = 0$, $d(x, y) = d(y, x)$, and $d(x, z) \leq d(x, y) + d(y, z)$ for all $x, y, z \in \mathbb{X}$.

LEMMA A.1

Let $\mathbb{X}$ be a metric space. If $f : [0, 1] \rightarrow \overline{\Sigma}$ is a continuous parameterization of a set $\overline{\Sigma} \subset \mathbb{X}$, then $d(a, b) = \text{diam } f([a, b])$ is a pseudometric on $[0, 1]$. Furthermore, with respect to this pseudometric, if $0 \leq a \leq b \leq x \leq y \leq 1$, then $\text{diam}_d([a, b]) = d(a, b)$ and $\text{gap}_d([a, b], [x, y]) = d(b, x)$.

Proof

To verify that d is a pseudometric on $[0, 1]$, only the triangle inequality requires some thought. Given $a, b, c \in [0, 1]$, set

$$A = f([a, b]), \quad B = f([b, c]), \quad C = f([a, c]),$$

so that $\text{diam } A = d(a, b)$, $\text{diam } B = d(b, c)$, and $\text{diam } C = d(a, c)$. Up to relabeling, there are two cases.

Case 1. If $a \leq c \leq b$, then $C \subset A$ and $d(a, c) \leq d(a, b) \leq d(a, b) + d(b, c)$.

Case 2. Suppose $a \leq b \leq c$. We now argue as in [3, Lemma 2.18], a step in a proof of Gołȃb's semicontinuity theorem. Let p and q be points in C such that $\text{dist}(p, q) = d(a, c)$. Consider the 1-Lipschitz function $F : C \rightarrow \mathbb{R}$ defined by $F(x) = \text{dist}(x, p)$. Then

$$d(a, b) + d(b, c) \geq \text{diam } F(A) + \text{diam } F(B) \geq \text{diam } F(C) \geq d(a, c),$$

where the first inequality holds since $\text{Lip}(F) = 1$, the second inequality holds since $F(A)$, $F(B)$, $F(C)$ are intervals in $\mathbb{R}$ with $F(C) \subset F(A) \cup F(B)$, and the third inequality holds since $F(p) = 0$ and $F(q) = \text{dist}(p, q) = d(a, c)$.

Suppose that $I = [a, b] \subset [0, 1]$. On the one hand, for any $a \leq a' \leq b' \leq b$, we know that the distance $d(a', b') = \text{diam } f([a', b']) \leq \text{diam } f([a, b]) = d(a, b)$ because $[a', b'] \subset [a, b]$. Hence, $\text{diam}_d I \leq d(a, b)$. On the other hand, $\text{diam}_d I \geq d(a, b)$ because $a, b \in I$. A similar argument yields the formula for the gap between intervals. $\square$

REMARK A.2

If the parameterization $f : [0, 1] \rightarrow \overline{\Sigma}$ has positive constant speed, then the pseudometric d on $[0, 1]$ is a metric. The pseudometric d on $[0, 1]$ bears some similarity to the intrinsic diameter distance on $\overline{\Sigma}$; see, for example, [39, 46].

EXAMPLE A.3

Let $\{e_i\}_{i=1}^\infty$ denote the standard basis in ℓ_2 . Fix a large integer $n > 1$ and let $f : [0, 1] \rightarrow \ell_2$ be a continuous map, which traces a constant-speed, piecewise linear path from e_1 to e_2 to e_3 and so on until reaching e_{n+1} . Decompose $[0, 1]$ into n intervals $I_i = f^{-1}([e_i, e_{i+1}])$. With respect to the metric d of Lemma A.1, we have $\text{diam}_d I_i = \text{diam } f(I_i) = \sqrt{2}$ for all i and $\text{diam}_d[0, 1] = \text{diam } f([0, 1]) = \sqrt{2}$. This shows that diam_d is not superadditive. Similarly, even though there are $\gtrsim n/2$ disjoint intervals of diameter $\sqrt{2}$ lying between I_1 and I_n , we only have $\text{gap}_d(I_1, I_n) = |e_2 - e_n|_{\ell_2} = \sqrt{2}$.

The second auxiliary lemma is related to [63, Lemma 2.16] and [14, Lemma A.1], but with an alternative gap condition that is motivated by the previous example.

LEMMA A.4

Let $\mathbb{X}$ be a metric space, let $f : [0, 1] \rightarrow \overline{\Sigma}$ be a continuous parameterization of a set $\overline{\Sigma} \subset \mathbb{X}$, let $\xi > 6$, and let $0 < r_- < r_+ < \infty$. Suppose $\{J_i\}_{i=1}^I$ is a finite ($I < \infty$) or infinite ($I = \infty$) sequence of closed intervals in $[0, 1]$ and $(k_i)_{i=1}^I$ is a sequence of integers bounded from below such that

- (1) *chain property*: for all $1 \leq i < I$, we have $J_i \cap J_{i+1} \neq \emptyset$;
- (2) *geometric decay*: for all $i \geq 1$, we have $\text{diam } f(J_i) \leq \xi^{-k_i} r_+$; and
- (3) *separation within levels*: for all $i, j \geq 1$ with $i \neq j$, if $k_i = k_j = k$, then there exists at least $\lceil 3r_+/r_- \rceil$ pairwise disjoint closed intervals K with $\text{diam } f(K) \geq \xi^{-k} r_-$ and such that K lies between J_i and J_j as subsets of $\mathbb{R}$.

If $\xi > r_+/r_-$, then there exists a unique $M \geq 1$ such that $k_M = \min_{i \geq 1} k_i$; moreover,

$$(A.1) \quad \sum_{i=1}^I \text{diam } f(J_i) \leq (1 + 3/\xi) \xi^{-k_M} r_+.$$

Proof

Without loss of generality, we may assume that $r_- = 1$ and $1 < r_+ < \infty$. Suppose that $\xi > 6$ is sufficiently large, ultimately depending only on r_+ , to be specified below. Fix sequences $\{J_i\}_{i=1}^I$ and $(k_i)_{i=1}^I$ as above. We claim that for every integer $1 \leq n \leq I$, there exists a unique $1 \leq m \leq n$ such that $k_m = \min_{i=1}^n k_i$ and

$$(A.2) \quad \sum_{i=1}^n \text{diam } f(J_i) \leq (1 + 2\xi^{-1} + 4\xi^{-2} + 8\xi^{-3} + \dots) \xi^{-k_m} r_+.$$

The base case $n = 1$ is trivial. Suppose for induction that the claim holds whenever $1 \leq n \leq N$. Put $n = N + 1$ and choose an index $1 \leq m \leq N + 1$ such that $k_m = \min_{i=1}^{N+1} k_i$. (We do not yet know that m is unique.) Separate $\{J_i : 1 \leq i \leq N + 1\} \setminus \{J_m\}$ into two chains $\{J_1, \dots, J_{m-1}\}$ and $\{J_{m+1}, \dots, J_{N+1}\}$, each of which contains no more than N intervals. Working with the first chain (unless it is empty), the induction hypothesis and fact $\xi > 6$ implies that there exists a unique $1 \leq j \leq m - 1$ such that $k_j = \min_{i=1}^{m-1} k_i$ and

$$\sum_{i=1}^{m-1} \text{diam } f(J_i) \leq (1 + 2\xi^{-1} + 4\xi^{-2} + 8\xi^{-3} + \dots) \xi^{-k_j} r_+ < (3/2) \xi^{-k_j} r_+.$$

We know that $k_j \geq k_m$ by our selection of m . Suppose for contradiction that $k_j = k_m$. By property (3), we can locate $P = \lceil 3r_+ \rceil$ pairwise disjoint closed intervals $K_1, K_2, \dots, K_P$ such that each K_p lies between J_j and J_m and $\text{diam } f(K_p) \geq \xi^{-k_j}$. This immediately implies that J_j and J_m do not intersect, so $j \leq m - 2$. For any $j + 1 \leq i \leq m - 1$, we have $k_i \geq k_j + 1$ by our specification of j ; hence, $\text{diam } f(J_i) \leq \xi^{-k_i} r_+ \leq \xi^{-1} \xi^{-k_j} r_+ < \xi^{-k_j}$ provided that $\xi > r_+$. This ensures that each interval J_i with $j + 1 \leq i \leq m - 1$ intersects at most two of the intervals K_p . (Otherwise, if some J_i intersected three or more intervals, then $J_i \supset K_p$ for some p and $\xi^{-k_j} > \text{diam } f(J_i) \geq \text{diam } f(K_p) \geq \xi^{-k_j}$. We remark that if we allowed $\text{diam } f(J_i) \geq \text{diam } f(K_p)$ for some p , then we could not control the overlap; see Example A.3.) Now, $\bigcup_{p=1}^P K_p \subset \bigcup_{i=j+1}^{m-1} J_i$ because the latter intervals connect J_j and J_m by (1). All together, these observations yield

$$P \xi^{-k_j} \leq \sum_{p=1}^P \text{diam } f(K_p) \leq 2 \sum_{i=j+1}^{m-1} \text{diam } f(J_i) < 3 \xi^{-k_j} r_+.$$

Thus, $P < 3r_+ \leq \lceil 3r_+ \rceil = P$, which is a contradiction. Therefore, we have $k_j \geq k_m + 1$. It follows that $k_i > k_m$ for all $1 \leq i \leq m - 1$ and

$$\sum_{i=1}^{m-1} \text{diam } f(J_i) \leq \xi^{-1} (1 + 2\xi^{-1} + 4\xi^{-2} + 8\xi^{-3} + \dots) \xi^{-k_m} r_+.$$

A similar conclusion holds for $m + 1 \leq i \leq N + 1$. Adding the estimates verifies the induction step.

To conclude, let $M \geq 1$ be an index such that $k_M = \min_{i=1}^I k_i$, which exists because the sequence $(k_i)_{i=1}^I$ is bounded from below. Because for all integers $1 \leq n \leq I$, there is a unique index $1 \leq m \leq n$ such that $k_m = \min_{i=1}^n k_i$, we discover that M is unique and $k_m = k_M$ for all $n \geq M$. Finally, (A.1) follows from (A.2) since $\xi > 6$. $\square$

Proof of Lemma 3.20

In the construction below, we view $\text{Diam}(\tau)$, the diameter of the image of an arc τ , as a weight assigned to $\text{Domain}(\tau)$, a closed interval lying in the domain of the map f . From this vantage point, a filtration is simply a nested family of finite partitions of some set $K \subset [0, 1]$ into intervals with geometrically decaying weights. This heuristic can be formalized using the pseudometric from Lemma A.1. As shorthand, we may speak of the intersection or union of arcs, but we always mean the arc formed by taking the intersection or union in the domains of the arcs.

Let $\rho > 1$, let $0 < \underline{A} < A < \infty$, and let $J \geq 1$ be an integer, with J sufficiently large depending on ρ and $A/\underline{A}$ to be specified below. Without loss of generality, we may assume that $\underline{A} = 1$ and $A > 1$. Let $\mathcal{F}^0$ be an admissible family of arcs in $\overline{\Sigma}$. Our plan is to first preprocess $\mathcal{F}^0$, partitioning it into a finite number of families $\mathcal{D}^1, \mathcal{D}^2, \dots$ of well-separated arcs. We then transform each family $\mathcal{D}^j$ into a nested family $\mathcal{E}^j$ of arcs satisfying (3.24) and (3.25). (This is where we use Lemma A.4.) Afterwards, we describe how to extend each family $\mathcal{E}^j$ to a filtration $\mathcal{F}^j$ with the same conclusions.

To begin, using the bounded overlap assumption, break $\mathcal{F}^0$ into C or fewer nonempty families such that within each family, arcs in the same level n are pairwise disjoint. Note that some levels of a family may be empty. By splitting each family into

$$\lceil 3A \rceil + 1 \quad (< 5A)$$

families, as necessary, we may assume that any pair of distinct arcs in level n of a family is separated in the domain by at least $\lceil 3A \rceil$ disjoint arcs σ with $\text{Diam}(\sigma) \geq \rho^{-n}$. Next, break apart each family of arcs into J (or fewer) nonempty families by jumping J generations at a time (that is, within each family), group together all levels n with $n - n_0 \equiv m \pmod{J}$, relabeling so that each original level $n = (n_0 + m) + kJ$ is assigned to level $N = (n_0 + m) + k$. Denote the resulting $5ACJ$ or fewer families by $\mathcal{D}^1 = \bigcup_{n=n_1}^{\infty} \mathcal{D}_n^1, \mathcal{D}^2 = \bigcup_{n=n_2}^{\infty} \mathcal{D}_n^2, \dots$, with starting index $n_j \in \{n_0, n_0 + 1, \dots, n_0 + J - 1\}$ for all j . Note that level n in $\mathcal{D}^j$ corresponds to level $n_j + J(n - n_j) = Jn - (J - 1)n_j$ in $\mathcal{F}^0$. By our assumption on the diameters of arcs in $\mathcal{F}^0$, if $n \geq n_j$ and $\tau \in \mathcal{D}_n^j$, then

$$(A.3) \quad \rho^{(J-1)n_j} \rho^{-Jn} \leq \text{Diam}(\tau) \leq (A\rho^{(J-1)n_j}) \rho^{-Jn}.$$

Further, any pair of distinct arcs in $\mathcal{D}_n^j$ is separated in the domain by at least $\lceil 3A \rceil$ disjoint intervals σ with $\text{Diam}(\sigma) \geq \rho^{(J-1)n_j} \rho^{-Jn}$.

Next, we transform $\mathcal{D}^1, \mathcal{D}^2, \dots$ into nested families $\mathcal{E}^1, \mathcal{E}^2, \dots$ of arcs. This will require us to join certain overlapping arcs, thereby increasing the diameter of arcs. By choosing J to be sufficiently large, we can control the growth of diameters.

Fix a family $\mathcal{D}^j$. For each $n \geq n_j$, let $\mathcal{D}_{n+}^j = \mathcal{D}_{n+1}^j \cup \mathcal{D}_{n+2}^j \cup \mathcal{D}_{n+3}^j \cup \dots$. Fix a level $n \geq n_j$ and an arc $\tau \in \mathcal{D}_n^j$. We inductively define a sequence of arcs by fusing the arc and all overlapping arcs from future generations relative to τ . Formally, we set

$$\begin{aligned} \tau_0 &= \tau, & \tau_{k+1} &= f \left| \bigcup_{\{\sigma \in \mathcal{D}_{n+}^j : \sigma \cap \tau_k \neq \emptyset\}} \text{Domain}(\tau) \right. \quad (k \geq 0), \\ \tau_\infty &= f \left| \overline{\bigcup_{k=0}^{\infty} \text{Domain}(\tau_k)} \right. \end{aligned}$$

where $\sigma \cap \tau_k \neq \emptyset$ means $\text{Domain}(\sigma) \cap \text{Domain}(\tau_k) \neq \emptyset$ and $f|I$ denotes the restriction of f to I . Let us upper bound $\text{Diam}(\tau_\infty)$ in terms of $\text{Diam}(\tau)$. Each arc τ_{k+1} can be obtained by taking the union of τ_k and up to two arcs σ_k^- and σ_k^+ that intersect, but are not contained in τ_k , one of each side. Thus, τ_∞ can be obtained by concatenating τ and two arcs, τ^- and τ^+ , with domains on either side of $\text{Domain}(\tau)$, where τ^- and τ^+ can each be expressed as the union of a finite or infinite chain of arcs in $\mathcal{D}_{n+}^j$. By Lemma A.1, $\text{Diam}(\tau^\pm) \leq \sum_k \text{Diam}(\sigma_k^\pm)$. To continue, we need J to be sufficiently large. Assume that $\rho^J > 6A \geq \max\{6, A\}$. By Lemma A.4, with $\xi = \rho^J$, $r_- = \rho^{(J-1)n_j}$, and $r_+ = A\rho^{(J-1)n_j}$,

$$\begin{aligned} \text{Diam}(\tau^\pm) &\leq (1 + 3/\rho^J)(A\rho^{(J-1)n_j})\rho^{-J(n+1)} \\ \text{(A.4)} \quad &< \frac{1}{4}\rho^{(J-1)n_j}\rho^{-Jn} \leq \frac{1}{4}\text{Diam}(\tau). \end{aligned}$$

Therefore,

$$\begin{aligned} \text{Diam}(\tau_\infty) &\leq \text{Diam}(\tau^-) + \text{Diam}(\tau) + \text{Diam}(\tau^+) \\ \text{(A.5)} \quad &< \frac{3}{2}\text{Diam}(\tau) \leq \frac{3}{2}(A\rho^{(J-1)n_j})\rho^{-Jn}. \end{aligned}$$

Recall that distinct arcs τ and $\tilde{\tau}$ in $\mathcal{D}_n^j$ are separated by an arc (in fact, several arcs) σ of $\overline{\Sigma}$ with $\text{Diam}(\sigma) \geq \rho^{(J-1)n_j}\rho^{-Jn}$. Thus, by (A.4) and the triangle inequality, for any distinct $\tau, \tilde{\tau} \in \mathcal{D}_n^j$, the domains of τ_∞ and $\tilde{\tau}_\infty$ are separated by some arc σ in $\overline{\Sigma}$ with

$$\text{(A.6)} \quad \text{Diam}(\sigma) > \frac{1}{2}\rho^{(J-1)n_j}\rho^{-Jn}.$$

For each j and $n \geq n_j$, define $\mathcal{E}_n^j$ by including the arc τ_∞ in $\mathcal{E}_n^j$ for each arc $\tau \in \mathcal{D}_n^j$. By (A.3) and (A.5), the families $\mathcal{E}^j = \bigcup_{n=n_j}^{\infty} \mathcal{E}_n^j$ satisfy—even better inequalities than—(3.24) and (3.25), where the roles of “ $\tau' \in \mathcal{F}_n^0$ ” and “ $\tau \in \mathcal{F}_N^j$ ” in the statement of the lemma are played by $\tau \in \mathcal{D}_n^j$ and $\tau_\infty \in \mathcal{E}_n^j$, respectively. Let us verify that $\mathcal{E}^j$ is a nested family of arcs in the sense that if (the domains of) τ_∞ and $\tilde{\tau}_\infty$ in $\mathcal{E}^j$ intersect in more than one point, then $\text{Domain}(\tau_\infty)$ is contained in $\text{Domain}(\tilde{\tau}_\infty)$ or vice versa. Suppose that $t \geq n \geq n_j$, $\tau \in \mathcal{D}_n^j$, $\tilde{\tau} \in \mathcal{D}_t^j$, and τ_∞ and $\tilde{\tau}_\infty$ intersect in at least two points. Then the interiors of $\text{Domain}(\tau_\infty)$ and $\text{Domain}(\tilde{\tau}_\infty)$ intersect. Hence, $\text{Domain}(\tau_k)$ and $\text{Domain}(\tilde{\tau}_l)$ intersect for some $k, l < \infty$, as well. Because $\tilde{\tau}_l$

is a finite union of overlapping arcs $\mathcal{D}_{(t-1)+}^j$ and $t \geq n$, it follows that $\tilde{\tau}_l \subset \tau_{k+m}$ for some sufficiently large positive integer m . Thus, $\tilde{\tau}_{l+i} \subset \tau_{k+m+i}$ for all $i \geq 0$ by the construction, and therefore, $\tilde{\tau}_\infty \subset \tau_\infty$. As an immediate corollary, distinct arcs in each level $\mathcal{E}_n^j$ intersect in at most one point. However, by (A.6), we know even more: for every j and $n \geq n_j$, the arcs in $\mathcal{E}_n^j$ are pairwise disjoint. Reviewing the construction so far, we see that the map that assigns each arc $\tau \in \mathcal{F}_n^0$ to an arc $\tau_\infty \in \mathcal{E}_{n_j+(n-n_j)/J}^j$ is injective.

To build filtrations $\mathcal{F}^j$ out of the families $\mathcal{E}^j$, it remains to add arcs to each level of a family, as necessary, to ensure that the domain of any arc is the union of the domains of its children. Fix a family $\mathcal{E}^j$. We will work top-down, starting with level $n = n_j$. Let $\tau_1, \dots, \tau_k$ denote the arcs in $\mathcal{E}_n^j$, ordered from left to right using the natural order of their domains as subsets of $\mathbb{R}$. It is possible that this list is empty. From left to right, expand

$$f = \sigma_0 \cup \tau_1 \cup \sigma_1 \cup \tau_2 \cup \dots \cup \tau_{k-1} \cup \sigma_{k-1} \cup \tau_k \cup \sigma_k$$

with domains of consecutive arcs intersecting in exactly one point; the initial and final arc σ_0 and σ_k may be empty or nonempty. If $\mathcal{E}_n^j$ is empty, write $f = \sigma_0$. Suppose that $\sigma = \sigma_i$ is nonempty. There are three alternatives.

Alternative 1. If it happens that $\text{Diam}(\sigma)$ is between $(1/4)\rho^{(J-1)n_j}\rho^{-n}$ and $\rho^{(J-1)n_j}\rho^{-n}$, we make no modification and include σ in $\mathcal{F}_n^j$.

Alternative 2. Another possibility is that $\text{Diam}(\sigma) < (1/4)\rho^{(J-1)n_j}\rho^{-n}$. If $\sigma = \sigma_i$ for some $0 \leq i \leq k-1$, set $\tau = \tau_{i+1}$, the first arc to the right; if $\sigma = \sigma_k$, set $\tau = \tau_k$, the first arc to the left. Replace τ by $\sigma \cup \tau$. By (A.5), the modified arc satisfies

$$\text{Diam}(\tau) < (3/2)A\rho^{(J-1)n_j}\rho^{-n} + (1/4)\rho^{(J-1)n_j}\rho^{-n} < (7/4)A\rho^{(J-1)n_j}\rho^{-n}.$$

(Exceptionally, if $f = \sigma_0 \cup \tau_1 \cup \sigma_1$ and both σ_0 and σ_1 have small diameters, replace τ by $\sigma_0 \cup \tau \cup \sigma_1$ instead; in this case

$$\text{Diam}(\tau) < (3/2)A\rho^{(J-1)n_j}\rho^{-n} + 2(1/4)\rho^{(J-1)n_j}\rho^{-n} < 2A\rho^{(J-1)n_j}\rho^{-n}.)$$

Similarly, if the original arc $\tau \in \mathcal{E}_n^j$ came from extending an arc $\tau' \in \mathcal{D}_n^j$, then the modified arc τ satisfies $\text{Diam}(\tau) < (7/4)\text{Diam}(\tau')$ (exceptionally, $\text{Diam}(\tau) < 2\text{Diam}(\tau')$). Include the modified arc τ in $\mathcal{F}_n^j$.

Alternative 3. Lastly, suppose that $\text{Diam}(\sigma) > \rho^{(J-1)n_j}\rho^{-n}$. We will partition σ into a finite number of shorter arcs ζ_i of diameter between $(1/4)\rho^{(J-1)n_j}\rho^{-n}$ and $\rho^{(J-1)n_j}\rho^{-n}$ (see Remark 3.5) but need to do this in an intelligent way in order to maintain nested levels. Let $\{\xi_1, \xi_2, \dots\}$ be an enumeration of the maximal arcs in $\mathcal{E}_{n+}^j = \mathcal{E}_{n+1}^j \cup \mathcal{E}_{n+2}^j \cup \dots$ whose domains are contained in $\text{Domain}(\sigma)$. (There may be none.) Earlier we declared that $\rho^J > 6A$. Hence, $\text{Diam}(\xi_l) < (3/2)A\rho^{(J-1)n}\rho^{-(n+1)} < (1/4)\rho^{(J-1)n_j}\rho^{-n}$. We need to make sure that the endpoints of arcs in our partitions of σ do not lie in the interior of $\text{Domain}(\xi_l)$ for any l . Proceed as follows: Let a be the left endpoint of $\text{Domain}(\sigma)$. Choose $t > 0$ as large as possible so that $\text{Diam}(f|_{[a, a+t]}) = (1/4)\rho^{(J-1)n_j}\rho^{-n}$, which we may do because the diameter varies continuously in t and $\text{Diam}(\sigma)$ is large. If $a + t$ does not lie in the interior of $\text{Domain}(\xi_l)$ for any l , then we set $\zeta_1 = f|_{[a, a+t]}$ and have $\text{Diam}(\zeta_1) = (1/4)\rho^{(J-1)n_j}\rho^{-n}$. Otherwise, if $a + t$ lies in

the interior ξ_l for some l , set $\zeta_1 = f|_{[a,a+t]} \cup \xi_l$, which satisfies $(1/4)\rho^{(J-1)n_j}\rho^{-n} \leq \text{Diam}(\zeta_1) < (1/2)\rho^{(J-1)n_j}\rho^{-n}$. Repeat a similar construction on $\sigma \setminus \zeta_1$, this time letting a be the left endpoint of $\text{Domain}(\sigma \setminus \zeta_1)$ and choosing $t > 0$ as large as possible so that $\text{Diam}(f|_{[a,a+t]}) \leq (1/4)\rho^{(J-1)n_j}\rho^{-n}$. If $\text{Diam}(f|_{[a,a+t]}) = (1/4)\rho^{(J-1)n_j}$, then continue as before and iterate. Otherwise, at some stage, $\sigma \setminus (\zeta_1 \cup \dots \cup \zeta_m) \neq \emptyset$, but $\text{Diam}(f|_{[a,a+t]}) < (1/4)\rho^{(J-1)n_j}\rho^{-n}$, where t is the right endpoint of the domain of σ . Replace ζ_m by $\zeta_m \cup f|_{[a,a+t]}$. This may increase the diameter of ζ_m , but in any event, the modified arc satisfies $(1/4)\rho^{(J-1)n_j}\rho^{-n} \leq \text{Diam}(\zeta_m) < (3/4)\rho^{(J-1)n_j}\rho^{-n}$. Include each of the intervals $\zeta_1, \dots, \zeta_m$ from the partition of σ in $\mathcal{F}_n^j$.

Carry out the indicated construction for each nonempty $\sigma = \sigma_i$. Also include any arc τ_i in $\mathcal{F}_n^j$ if it was not already included. This completes the definition of $\mathcal{F}_n^j$. To define the next level $\mathcal{F}_{n+1}^j$, repeat the construction from above on each arc τ in $\mathcal{F}_n^j$ independently, with f replaced by τ . By induction, we obtain a definition of $\mathcal{F}^j = \bigcup_{n=n_j}^{\infty} \mathcal{F}_n^j$ on each level. The family $\mathcal{F}^j$ is the desired filtration. $\square$

Acknowledgments. The authors thank an anonymous referee for helpful comments that led us to improve the exposition in Section 3 and include Section 4. The referee initially brought the gap in [65], which we report in Remark 3.8, to our attention; for the correction, see Remark 3.24. We thank Raanan Schul for encouraging us to write out the details in Section 3 and the appendix in full and hope this will assist others in learning the proof. Finally, the first author would like to thank Raanan for his past mentorship and for imparting some of his good mathematical taste.

References

- [1] Y. I. Alber, *A bound for the modulus of continuity for metric projections in a uniformly convex and uniformly smooth Banach space*, J. Approx. Theory **85** (1996), no. 3, 237–249. MR 1391134. DOI 10.1006/jath.1996.0040.
- [2] Y. I. Alber, *Metric and generalized projection operators in Banach spaces: properties and applications*, Theory and applications of nonlinear operators of accretive and monotone type, Lect. Notes Pure Appl. Math. **178**, Dekker, New York, 1996, pp. 15–50. MR 1386667.
- [3] G. Alberti and M. Ottolini, *On the structure of continua with finite length and Golab’s semicontinuity theorem*, Nonlinear Anal. **153** (2017), 35–55. MR 3614660. DOI 10.1016/j.na.2016.10.012.
- [4] L. Ambrosio and B. Kirchheim, *Rectifiable sets in metric and Banach spaces*, Math. Ann. **318** (2000), no. 3, 527–555. MR 1800768. DOI 10.1007/s002080000122.
- [5] E. Asplund, *Positivity of duality mappings*, Bull. Amer. Math. Soc. **73** (1967), 200–203. MR 0206663. DOI 10.1090/S0002-9904-1967-11678-1.
- [6] J. Azzam, *Harmonic measure and the analyst’s traveling salesman theorem*, preprint, arXiv:1905.09057v3 [math.CA].
- [7] J. Azzam and R. Schul, *An analyst’s traveling salesman theorem for sets of dimension larger than one*, Math. Ann. **370** (2018), nos. 3–4, 1389–1476. MR 3770170. DOI 10.1007/s00208-017-1609-0.

- [8] J. Azzam and R. Schul, *How to take shortcuts in Euclidean space: Making a given set into a short quasi-convex set*, Proc. Lond. Math. Soc. (3) **105** (2012), MR 2959930. DOI 10.1112/plms/pds005.
- [9] J. Azzam and X. Tolsa, *Characterization of n -rectifiability in terms of Jones' square function: Part II*, Geom. Funct. Anal. **25** (2015), no. 5, 1371–1412. MR 3426057. DOI 10.1007/s00039-015-0334-7.
- [10] J. Azzam and M. Villa, *Quantitative comparisons of multiscale geometric properties*, Anal. PDE **14** (2021), no. 6, 1873–1904. MR 4308668. DOI 10.2140/apde.2021.14.1873.
- [11] M. Badger, *Generalized rectifiability of measures and the identification problem*, Complex Anal. Synerg. **5** (2019), art. no. 2. MR 3941625. DOI 10.1007/s40627-019-0027-3.
- [12] M. Badger and S. Lewis, *Local set approximation: Mattila–Vuorinen type sets, Reifenberg type sets, and tangent sets*, Forum Math. Sigma **3** (2015), e24, 63. MR 3482273. DOI 10.1017/fms.2015.26.
- [13] M. Badger, S. Li, and S. Zimmerman, *Identifying 1-rectifiable measures in Carnot groups*, preprint, arXiv:2109.06753v3 [math.MG], 2023.
- [14] M. Badger and S. McCurdy, *Subsets of rectifiable curves in Banach spaces II: Universal estimates for almost flat arcs*, preprint, arXiv:2208.10288 [math.CA].
- [15] M. Badger and L. Naples, *Radon measures and Lipschitz graphs*, Bull. Lond. Math. Soc. **53** (2021), no. 3, 921–936. MR 4275101. DOI 10.1112/blms.12473.
- [16] M. Badger, L. Naples, and V. Vellis, *Hölder curves and parameterizations in the analyst's traveling salesman theorem*, Adv. Math. **349** (2019), 564–647. MR 3941391. DOI 10.1016/j.aim.2019.04.011.
- [17] M. Badger and R. Schul, *Multiscale analysis of 1-rectifiable measures II: Characterizations*, Anal. Geom. Metr. Spaces **5** (2017), 1–39. MR 3627148. DOI 10.1515/agms-2017-0001.
- [18] M. Badger and V. Vellis, *Geometry of measures in real dimensions via Hölder parameterizations*, J. Geom. Anal. **29** (2019), no. 2, 1153–1192. MR 3935254. DOI 10.1007/s12220-018-0034-2.
- [19] M. Badger and V. Vellis, *Hölder parameterization of iterated function systems and a self-affine phenomenon*, Anal. Geom. Metr. Spaces **9** (2021), no. 1, 90–119. MR 4289517. DOI 10.1515/agms-2020-0125.
- [20] Z. M. Balogh and R. Züst, *Box-counting by Hölder's traveling salesman*, Arch. Math. (Basel) **114** (2020), no. 5, 561–572. MR 4088553. DOI 10.1007/s00013-019-01415-5.
- [21] D. Bate, *Structure of measures in Lipschitz differentiability spaces*, J. Amer. Math. Soc. **28** (2015), no. 2, 421–482. MR 3300699. DOI 10.1090/S0894-0347-2014-00810-9.
- [22] D. Bate, M. Csörnyei, and B. Wilson, *The Besicovitch–Federer projection theorem is false in every infinite-dimensional Banach space*, Israel J. Math. **220** (2017), no. 1, 175–188. MR 3666823. DOI 10.1007/s11856-017-1514-y.
- [23] D. Bate and S. Li, *Characterizations of rectifiable metric measure spaces*, Ann. Sci. Éc. Norm. Supér. (4) **50** (2017), no. 1, 1–37. MR 3621425. DOI 10.24033/asens.2314.

- [24] C. J. Bishop, *The traveling salesman theorem for Jordan curves*, Adv. Math. **404** (2022), paper no. 108443. MR 4420442. DOI 10.1016/j.aim.2022.108443.
- [25] C. J. Bishop and P. W. Jones, *Harmonic measure, L^2 estimates and the Schwarzian derivative*, J. Anal. Math. **62** (1994), 77–113. MR 1269200. DOI 10.1007/BF02835949.
- [26] C. J. Bishop and P. W. Jones, *Wiggly sets and limit sets*, Ark. Mat. **35** (1997), no. 2, 201–224. MR 1478778. DOI 10.1007/BF02559967.
- [27] C. J. Bishop, P. W. Jones, R. Pemantle, and Y. Peres, *The dimension of the Brownian frontier is greater than 1*, J. Funct. Anal. **143** (1997), no. 2, 309–336. MR 1428819. DOI 10.1006/jfan.1996.2928.
- [28] W. L. Bynum, *Weak parallelogram laws for Banach spaces*, Canad. Math. Bull. **19** (1976), no. 3, 269–275. MR 0442647. DOI 10.4153/CMB-1976-042-4.
- [29] W. L. Bynum and J. H. Drew, *A weak parallelogram law for l_p* , Amer. Math. Monthly **79** (1972), 1012–1015. MR 0317008. DOI 10.2307/2318073.
- [30] R. Cheng, J. Mashreghi, and W. T. Ross, *Optimal weak parallelogram constants for L^p spaces*, Math. Inequal. Appl. **21** (2018), no. 4, 1047–1058. MR 3868101. DOI 10.7153/mia-2018-21-71.
- [31] R. Cheng and W. T. Ross, *Weak parallelogram laws on Banach spaces and applications to prediction*, Period. Math. Hungar. **71** (2015), no. 1, 45–58. MR 3374700. DOI 10.1007/s10998-014-0078-4.
- [32] G. C. David and R. Schul, *A sharp necessary condition for rectifiable curves in metric spaces*, Rev. Mat. Iberoam. **37** (2021), no. 3, 1007–1044. MR 4236801. DOI 10.4171/rmi/1216.
- [33] G. C. David and R. Schul, *The analyst’s traveling salesman theorem in graph inverse limits*, Ann. Acad. Sci. Fenn. Math. **42** (2017), no. 2, 649–692. MR 3701642. DOI 10.5186/aasfm.2017.4260.
- [34] G. David and S. Semmes, *Analysis of and on uniformly rectifiable sets*, Math. Surveys Monogr. **38**, Amer. Math. Soc., Providence, 1993. MR 1251061. DOI 10.1090/surv/038.
- [35] G. David and S. Semmes, *Singular integrals and rectifiable sets in $\mathbf{R}^n$: Beyond Lipschitz graphs*, Astérisque **193**, Soc. Math. France, Paris, 1991. MR 1113517.
- [36] G. David and T. Toro, *Reifenberg parameterizations for sets with holes*, Mem. Amer. Math. Soc. **215** (2012), no. 1012, vi+102. MR 2907827. DOI 10.1090/S0065-9266-2011-00629-5.
- [37] J. Diestel, *Geometry of Banach spaces—Selected topics*, Lect. Notes Math. **485**, Springer, Berlin, 1975. MR 0461094.
- [38] N. Edelen, A. Naber, and D. Valtorta, *Effective Reifenberg theorems in Hilbert and Banach spaces*, Math. Ann. **374** (2019), nos. 3–4, 1139–1218. MR 3985109. DOI 10.1007/s00208-018-1770-0.
- [39] D. M. Freeman and D. A. Herron, *Bilipschitz homogeneity and inner diameter distance*, J. Anal. Math. **111** (2010), 1–46. MR 2747060. DOI 10.1007/s11854-010-0011-6.
- [40] T. Figiel, *On the moduli of convexity and smoothness*, Studia Math. **56** (1976), no. 2, 121–155. MR 0425581. DOI 10.4064/sm-56-2-121-155.

- [41] S. Ghinassi, *Sufficient conditions for $C^{1,\alpha}$ parametrization and rectifiability*, Ann. Acad. Sci. Fenn. Math. **45** (2020), no. 2, 1065–1094. MR 4112275. DOI 10.5186/aasfm.2020.4557.
- [42] M. Goering and S. McCurdy, *Wild examples of countably rectifiable sets*, Ann. Fenn. Math. **46** (2021), no. 1, 553–570. MR 4277828. DOI 10.5186/aasfm.2021.4632.
- [43] I. Hahlomaa, *Menger curvature and Lipschitz parametrizations in metric spaces*, Fund. Math. **185** (2005), no. 2, 143–169. MR 2163108. DOI 10.4064/fm185-2-3.
- [44] I. Hahlomaa, *Menger curvature and rectifiability in metric spaces*, Adv. Math. **219** (2008), no. 6, 1894–1915. MR 2456269. DOI 10.1016/j.aim.2008.07.013.
- [45] O. Hanner, *On the uniform convexity of L^p and l^p* , Ark. Mat. **3** (1956), 239–244. MR 0077087. DOI 10.1007/BF02589410.
- [46] D. A. Herron, *Geometry and topology of intrinsic distances*, J. Anal. **18** (2010), 197–231. MR 2850242.
- [47] M. Hyde, *The restricted content and the d -dimensional analyst’s travelling salesman Theorem for general sets*, Adv. Math. **397** (2022), paper no. 108189. MR 4366226. DOI 10.1016/j.aim.2022.108189.
- [48] P. W. Jones, *Rectifiable sets and the traveling salesman problem*, Invent. Math. **102** (1990), no. 1, 1–15. MR 1069238. DOI 10.1007/BF01233418.
- [49] B. T. Kien, *The normalized duality mapping and two related characteristic properties of a uniformly convex Banach space*, Acta Math. Vietnam **27** (2002), no. 1, 53–67. MR 1914408.
- [50] B. Kirchheim, *Rectifiable metric spaces: Local structure and regularity of the Hausdorff measure*, Proc. Amer. Math. Soc. **121** (1994), no. 1, 113–123. MR 1189747. DOI 10.2307/2160371.
- [51] T. Keleti and E. Paquette, *The trouble with von Koch curves built from n -gons*, Amer. Math. Monthly **117** (2010), no. 2, 124–137. MR 2590192. DOI 10.4169/000298910X476040.
- [52] J. Krandel, *The traveling salesman theorem for Jordan curves in Hilbert spaces*, preprint, arXiv:2107.07017v3 [math.CA], 2022.
- [53] S. Li, *Stratified β -numbers and traveling salesman in Carnot groups*, J. London Math. Soc. **106** (2022), no. 2, 662–703. MR 4477201. DOI 10.1112/jlms.12582.
- [54] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces, II: Function Spaces*, Ergeb. Math. Grenzgeb. **97**, Springer, Berlin, 1979. MR 0540367.
- [55] T. H. McNicholl, *The power of backtracking and the confinement of length*, Proc. Amer. Math. Soc. **141** (2013), no. 3, 1041–1053. MR 3003695. DOI 10.1090/S0002-9939-2012-11385-1.
- [56] L. Naples, *Rectifiability of pointwise doubling measures in Hilbert space*, preprint, arXiv:2002.07570 [math.CA].
- [57] A. Naber and D. Valtorta, *Rectifiable-Reifenberg and the regularity of stationary and minimizing harmonic maps*, Ann. of Math. (2) **185** (2017), no. 1, 131–227. MR 3583353. DOI 10.4007/annals.2017.185.1.3.

- [58] K. Okikiolu, *Characterization of subsets of rectifiable curves in $\mathbf{R}^n$* , J. London Math. Soc. (2) **46** (1992), no. 2, 336–348. MR 1182488. DOI 10.1112/jlms/s2-46.2.336.
- [59] D. Preiss and J. Tišer, *On Besicovitch’s $\frac{1}{2}$ -problem*, J. London Math. Soc. (2) **45** (1992), no. 2, 279–287. MR 1171555. DOI 10.1112/jlms/s2-45.2.279.
- [60] E. R. Reifenberg, *Solution of the plateau problem for m -dimensional surfaces of varying topological type*, Acta Math. **104** (1960), 1–92. MR 0114145. DOI 10.1007/BF02547186.
- [61] S. Rohde, *Quasicircles modulo bilipschitz maps*, Rev. Mat. Iberoamericana **17** (2001), no. 3, 643–659. MR 1900898. DOI 10.4171/RMI/307.
- [62] A. Ramierez and V. Vellis, *Time complexity in the analyst’s traveling salesman algorithm*, preprint, arXiv:2202.10314, 2022.
- [63] R. Schul, *Ahlfors-regular curves in metric spaces*, Ann. Acad. Sci. Fenn. Math. **32** (2007), no. 2, 437–460. MR 2337487.
- [64] R. Schul, “Analyst’s traveling salesman theorems. A survey” in *In the Tradition of Ahlfors–Bers, IV*, Contemp. Math. **432**, Amer. Math. Soc., Providence, 2007, 209–220. MR 2342818. DOI 10.1090/conm/432/08310.
- [65] R. Schul, *Subsets of rectifiable curves in Hilbert space—The analyst’s TSP*, J. Anal. Math. **103** (2007), 331–375. MR 2373273. DOI 10.1007/s11854-008-0011-y.
- [66] X. Tolsa, *Painlevé’s problem and the semiadditivity of analytic capacity*, Acta Math. **190** (2003), no. 1, 105–149. MR 1982794. DOI 10.1007/BF02393237.
- [67] M. Villa, *Higher dimensional Jordan curves*, preprint, arXiv:1908.10289v2 [math.CA], 2020.
- [68] T. Ważewski, *Kontinua prostowalne w związku z funkcjami i odwzorowaniami absolutnie ciągłymi (Rectifiable continua in connection with absolutely continuous functions and mappings)*, Dodatek do Rocznika Polskiego Towarzystwa Matematycznego (Supplement to the Annals of the Polish Mathematical Society) (1927), 9–49.

Badger: Department of Mathematics, University of Connecticut, Storrs, Connecticut, USA;
matthew.badger@uconn.edu

McCurdy: Department of Mathematics, National Taiwan Normal University, Taipei City, Taiwan (R.O.C.); smccurdy@ntnu.edu.tw