

WIENER INDICES OF MINUSCULE LATTICES

COLIN DEFANT, VALENTIN FÉRAY, PHILIPPE NADEAU, AND NATHAN WILLIAMS

ABSTRACT. The *Wiener index* of a finite graph G is the sum over all pairs (p, q) of vertices of G of the distance between p and q . When P is a finite poset, we define its *Wiener index* as the Wiener index of the graph of its Hasse diagram. In this paper, we find exact expressions for the Wiener indices of the distributive lattices of order ideals in minuscule posets. For infinite families of such posets, we also provide results on the asymptotic distribution of the distance between two random order ideals.

1. INTRODUCTION

1.1. Background: the Wiener index of the noncrossing partition lattice. Meandric systems are collections of non-crossing loops (up to isotopy) intersecting the horizontal axis at $2n$ fixed points—they have connections to several areas of mathematics, physics, and biology. There is a simple bijection between meandric systems and pairs of noncrossing partitions of size n .

Let $\text{NC}(n)$ be the lattice of noncrossing partitions of n . In the paper [GNP20]—motivated by asymptotic problems about meanders and meandric systems—Goulden, Nica, and Puder raised the following question: what is the average distance between two (uniform) random partitions in $\text{NC}(n)$? The question was answered for large n by Thévenin and the second author in [FT22], where it was proved that this average distance behaves as κn for some constant κ .

It is natural to ask similar questions for other families of posets, looking either for an exact nice formula or for an asymptotic answer. When the number of elements is known (which is the case for $\text{NC}(n)$), one can equivalently ask for the sum of distances between all pairs of elements.

In general, let $G = (V, E)$ be a finite connected graph, and for $p, q \in V$, write $d(p, q)$ for the distance in G from p to q . The *Wiener index* of G is defined to be

$$(1.1) \quad d(G) := \sum_{(p,q) \in V \times V} d(p, q).$$

This definition has its origin as the *Wiener index* predicting the boiling point of certain organic compounds [Wie47, Wik22, Rou02], and it has also been called the *distance* of the graph G [EJS76]. When P is a poset, we define $d(P) := d(G(P))$ for the Wiener index of the Hasse diagram $G(P)$ of P (that is, the vertices of $G(P)$ are the elements of P , and there is an edge in $G(P)$ between p and q when there is a cover relation in P between p and q).

In the case of the noncrossing partition lattice, the results in [FT22] imply that

$$d(\text{NC}(n)) \sim |\text{NC}(n)|^2 \kappa n \sim \frac{\kappa 16^n}{\pi n^2},$$

but no exact enumeration appears possible. (There does not even seem to be a simple formula for κ ; see the discussion in [FT22] and the related open problem in [Wil22].)

1.2. Wiener indices of other lattices. Computer experiments suggest that there are few non-trivial families of graphs $\{G_n\}_{n \geq 1}$ of combinatorial objects for which it is possible to find *exact* formulas for the Wiener index. First, there are families with elementary exact solutions, for which $|G_n|$ is a relatively small polynomial in n (for example: path graphs, grid graphs, etc.), or in which each graph G_n has a transitive underlying symmetry group (for example: the weak order on a finite Coxeter group, a boolean lattice or hypercube, etc.). A short list of examples is given in [Wei21].

There is, however, one class of posets in algebraic combinatorics that demonstrates consistently exceptional enumerative behavior: the minuscule lattices [Pro84]. For example, both the number of elements and the number of maximal chains in a minuscule lattice have simple (uniformly stated and proven) product formulas, and the minuscule lattices are well understood from the perspective of dynamical algebraic combinatorics (promotion, rowmotion, etc.) [SW12, Hop20]. Because of

their rich algebraic, representation-theoretic, and combinatorial properties, such posets serve as useful test cases for results and methods that might apply to more complicated structures.

It turns out that the Wiener indices of minuscule lattices also admit simple exact formulas, and one of the goals of the present paper is to establish such formulas.

For completeness, we first recall the definition of minuscule lattices (note that we will only use here their classification, and not the algebraic definition). Let $\mathfrak{g}$ be a complex simple Lie algebra with Weyl group W . Fix a set Φ^+ of positive roots of $\mathfrak{g}$, and let Λ^+ be the set of dominant weights. The finite-dimensional irreducible complex representations V_λ of $\mathfrak{g}$ are indexed by dominant weights $\lambda \in \Lambda^+$; λ is called *minuscule* if the W -orbit of λ is the set of *all* weights in V_λ . The minuscule weights are exactly those fundamental weights whose corresponding simple roots appear exactly once in the simple root expansion of the highest root. For more information, we refer the reader to [Sta80, Pro84].

For λ minuscule, define a poset on the weights in V_λ by introducing a cover relation $\mu < \nu$ whenever $\mu + \alpha = \nu$ for some simple root $\alpha \in \Phi^+$. This poset on V_λ is a distributive lattice, which we call a *minuscule lattice* [Pro84].

There are three infinite families of minuscule lattices—the order ideals in:

- a rectangle (type A ; in type B , minuscule lattices are chains and thus particular cases of rectangles),
- a shifted staircase (types C and D), and
- a “double tailed diamond” (type D)

—as well as two exceptional minuscule lattices (of types E_6 and E_7).

In this paper, we show that the Wiener indices of the infinite families of minuscule lattices admit simple product formulas, although we regrettably have been unable to find a unifying expression for these formulas. We also provide information about the asymptotic distribution of the distance between random elements in these posets (in the rectangle and shifted staircase cases), and we give a method for computing the higher moments exactly.

Note added in revision: after submitting our paper to the arXiv, our exact enumerative results have been used by Bourn and Erickson to obtain a close formula for a generating series of pairs of compositions weighted by the so-called “Earth Moving Distance” (or mass transportation distance) between them [BE23].

1.3. Rectangles. Write $J(P)$ for the lattice of order ideals in a finite poset P , ordered by inclusion. By Birkhoff’s representation theorem, any finite distributive lattice is of this form. In Sections 2 to 4, we consider the Wiener index of $P_{m,k} = J([m] \times [k])$, the lattice of order ideals in an $m \times k$ rectangle. The elements of $P_{m,k}$ can be represented as lattice paths in an $m \times k$ grid, so $|P_{m,k}| = \binom{m+k}{k}$. In this case (and, more generally, for any distributive lattice), it is easy to see that $d(p, q) = |p \Delta q|$, where $p \Delta q = (p \setminus q) \cup (q \setminus p)$ is the symmetric difference of the order ideals p and q .


It will be convenient to draw the elements of $P_{m,k}$ as lattice paths from $(0, 0)$ to $(m+k, m-k)$ using steps of the form $U = (1, 1)$ and $D = (1, -1)$. Writing p_i, q_i for the heights of p and q after the ends of their i th steps, the number of squares in column i between the lattice paths p and q is given as $\left\lfloor \frac{q_i - p_i}{2} \right\rfloor$, so that

$$(1.2) \quad d(p, q) = \left\lfloor \frac{q_1 - p_1}{2} \right\rfloor + \left\lfloor \frac{q_2 - p_2}{2} \right\rfloor + \cdots + \left\lfloor \frac{q_{m+k} - p_{m+k}}{2} \right\rfloor.$$

Example 1.1. The graph $G(P_{2,2})$ is drawn in Figure 1. Its Wiener index is

$$\begin{aligned} 56 = \frac{4}{18} \binom{10}{5} = & (0+1+2+2+3+4) + (1+0+1+1+2+3) + (2+1+0+2+1+2) + \\ & (2+1+2+0+1+2) + (3+2+1+1+0+1) + (4+3+2+2+1+0). \end{aligned}$$

We have three results that completely describe the Wiener index of the lattice of order ideals of a rectangle.

FIGURE 1. The Hasse diagram $G(P_{2,2})$ of order ideals in a 2×2 square.

Theorem 1.2. *The generating function for the Wiener index of all posets $P_{m,k} = J([m] \times [k])$ is given by*

$$(1.3) \quad \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} d(P_{m,k}) x^m y^k = \frac{2xy}{(x^2 - 2xy + y^2 - 2x - 2y + 1)^2}.$$

This theorem is obtained via classical first return decomposition for lattice paths. The fact that this generating series is rational comes as a surprise, since several intermediate computation steps involve algebraic non-rational functions. Extracting the coefficient of $x^m y^k$ from Equation (1.3), we obtain a formula for $d(P_{m,k})$.

Corollary 1.3. *The Wiener index of $P_{m,k}$ is*

$$d(P_{m,k}) = \frac{mk}{4m + 4k + 2} \binom{2m + 2k + 2}{2k + 1}.$$

For fixed α , we obtain the asymptotic expected value of $d(p, q)$ in a $(\alpha n) \times n$ rectangle as $n \rightarrow \infty$. To keep notation simple, we assume throughout the paper that αn is an integer (otherwise, it suffices to replace αn by its integer value).

Corollary 1.4. *We have*

$$\frac{d(P_{\alpha n, n})}{|P_{\alpha n, n}|^2} \sim \frac{\sqrt{\pi \alpha (1 + \alpha)}}{4} n^{3/2} \text{ as } n \rightarrow \infty.$$

In Section 7, we also describe in this regime the asymptotic distribution of the distance $D_{\alpha, n}$ between two independent uniform random elements of $P_{\alpha n, n}$.

Proposition 1.5. *The random variable $n^{-3/2} D_{\alpha, n}$ converges in distribution and in moments to $\sqrt{2\alpha(1 + \alpha)} \cdot \int_0^1 |B_0(t)| dt$, where $B_0(t)$ is a Brownian bridge on $[0, 1]$.*

Informally, a Brownian bridge on $[0, 1]$ is a Brownian motion conditioned to have value 0 at time 1. Alternatively, if B is a Brownian motion, then $B_0(t) := B(t) - tB(1)$ is a Brownian bridge. Brownian bridges have been extensively studied in the probabilistic literature. In particular, much is known on the random variable $\int_0^1 |B_0(t)| dt$; see [Jan07, Section 20] for a survey of results including numerous references. In particular, a table of the first few moments can be found in [Jan07, Table 2]. We copy here the first three:

$$\mathbb{E} \left[\int_0^1 |B_0(t)| dt \right] = \frac{1}{4} \sqrt{\frac{\pi}{2}}, \quad \mathbb{E} \left[\left(\int_0^1 |B_0(t)| dt \right)^2 \right] = \frac{7}{60}, \quad \mathbb{E} \left[\left(\int_0^1 |B_0(t)| dt \right)^3 \right] = \frac{21}{512} \sqrt{\frac{\pi}{2}}.$$

Together with Proposition 1.5, this implies

$$\begin{aligned} \frac{1}{|P_{\alpha n, n}|^2} \sum_{p, q \in P_{\alpha n, n}} d(p, q) &= \mathbb{E}[D_{\alpha, n}] \sim \frac{\sqrt{\pi\alpha(1+\alpha)}}{4} n^{3/2}, \\ \frac{1}{|P_{\alpha n, n}|^2} \sum_{p, q \in P_{\alpha n, n}} d(p, q)^2 &= \mathbb{E}[D_{\alpha, n}^2] \sim \frac{7}{30} \alpha(1+\alpha) n^3, \\ \frac{1}{|P_{\alpha n, n}|^2} \sum_{p, q \in P_{\alpha n, n}} d(p, q)^3 &= \mathbb{E}[D_{\alpha, n}^3] \sim \frac{21}{256} \sqrt{\pi} \alpha^{3/2} (1+\alpha)^{3/2} n^{9/2}. \end{aligned}$$

Note that the first estimate is nothing but Corollary 1.4. This gives a second derivation of this asymptotic result, which does not go through the exact expression. Exact expressions for such higher moments can also be obtained through combinatorial means, see Section 8 for a derivation of the second moment.

1.4. Shifted staircases. In Sections 5 and 6, we consider the Wiener index of Q_n , the distributive lattice of order ideals in the n th *shifted staircase* poset. Explicitly, Q_n is the set of order ideals in the poset $\{(i, j) : 1 \leq i \leq j \leq n\}$ under componentwise ordering. The following results determine $d(Q_n)$ exactly.

The elements of Q_n can be represented as lattice paths starting at $(0, 0)$, ending somewhere on the line $x = n$, and using steps of the form $U = (1, 1)$ and $D = (1, -1)$. In particular, $|Q_n| = 2^n$. Similarly as for rectangles, if p and q are elements in Q_n , writing p_i, q_i for the heights of p and q after the ends of their i th steps, we have

$$(1.4) \quad d(p, q) = \sum_{i=1}^n \left| \frac{q_i - p_i}{2} \right|.$$

Example 1.6. The graph $G(Q_3)$ is plotted in Figure 2. Its Wiener index is

$$140 = \frac{6 \cdot 7}{3} \binom{5}{3} = 24 + 18 + 14 + 14 + 14 + 14 + 18 + 24.$$

Theorem 1.7. *The generating function for the Wiener index of all lattices Q_n is given by*

$$(1.5) \quad \sum_{n=0}^{\infty} d(Q_n) x^n = \frac{8x(1 + \sqrt{1-4x} - x(3 + \sqrt{1-4x}))}{(1-4x)(1-4x + \sqrt{1-4x})^3}.$$

Corollary 1.8. *The Wiener index of Q_n is*

$$d(Q_n) = \frac{2n(2n+1)}{3} \binom{2n-1}{n}.$$


Consequently, as n tends to $+\infty$, we have $d(Q_n) \sim \frac{2}{3\sqrt{\pi}} 4^n n^{3/2}$.

In Section 7, we turn to the asymptotic distribution of the distance E_n between two random order ideals of Q_n .

Proposition 1.9. *The random variable $n^{-3/2} E_n$ converges in distribution and in moments to $\frac{1}{\sqrt{2}} \cdot \int_0^1 |B(t)| dt$, where $B(t)$ is a Brownian motion on $[0, 1]$.*

Again, much is known about the random variable $\int_0^1 |B(t)| dt$, and a comprehensive literature review appears in [Jan07, Section 21]. In particular, the first few moments are given in [Jan07, Table 3]:

$$\mathbb{E} \left[\int_0^1 |B(t)| dt \right] = \frac{2}{3} \sqrt{\frac{2}{\pi}}, \quad \mathbb{E} \left[\left(\int_0^1 |B(t)| dt \right)^2 \right] = \frac{3}{8}, \quad \mathbb{E} \left[\left(\int_0^1 |B(t)| dt \right)^3 \right] = \frac{263}{630} \sqrt{\frac{2}{\pi}}.$$



Together with Proposition 1.9, this implies

Again, recalling that $|Q_n| = 2^n$, this allows us to recover the asymptotic behaviour of $d(Q_n)$ given in Corollary 1.8 without going through its exact expression.

Let R_n be the n th “double tailed diamond”—that is, the distributive lattice of order ideals in the minuscule poset of type D_n corresponding to the first fundamental weight.

$$d(R_n) = \frac{2}{3}(n+3)(4n^2+9n+8).$$

The proof in the case of R_n is elementary and left to the reader. The cases of E_6 and E_7 are treated by computer. The expressions for Wiener indices given in Corollary 1.3, Corollary 1.8, and Theorem 1.10 suggest that there may be a uniform formula for $d(P)$ for P a minuscule lattice—but we regrettably have been unable to find such an expression.

2. LATTICE PATH BIJECTIONS

We continue to use the steps $U = (1, 1)$ and $D = (1, -1)$, but we will also make use of two versions (or colors) of the step $(1, 0)$, denoted O_1 and O_2 . For $(p, q) \in P_{k, n-k} \times P_{k, n-k}$, define a lattice path $A(p, q)$ using the following dictionary between the i th pair of steps in (p, q) and the i th step in $A(p, q)$:

$$(2.1) \quad \begin{array}{c|c|c|c} (p, q) & A(p, q) & \frac{q_{i+1}-p_{i+1}}{2} - \frac{q_i-p_i}{2} & r_{i+1} - r_i \\ \hline (D, U) & U & +1 & +1 \\ (U, D) & D & -1 & -1 \\ (U, U) & O_1 & 0 & 0 \\ (D, D) & O_2 & 0 & 0 \end{array}.$$

Two examples of this bijection are illustrated in Figure 3. Given a lattice path r of length n with steps from the set $\{U, D, O_1, O_2\}$, write r_i for the height (i.e. the y -coordinate) of r at the end of its i th step. The unsigned area between r and the x -axis is $d(r) = |r_0| + |r_1| + |r_2| + \cdots + |r_{n-1}| + \frac{1}{2}|r_n|$. Then, comparing with Equation (1.2), it is clear that $d(p, q) = d(A(p, q))$: certainly $\frac{q_0-p_0}{2} = 0 = r_0$, so suppose that $\frac{q_i-p_i}{2} = r_i$; then the difference in height at the $(i+1)$ st step in $A(p, q)$ matches the difference in height at the $(i+1)$ st steps of p and q , as shown in the rightmost two columns of (2.1). Let us also write

$$(2.2) \quad \bar{d}(r) = |r_0| + |r_1| + |r_2| + \cdots + |r_n| = d(r) + \frac{1}{2}|r_n|.$$




FIGURE 3. Illustration of the bijection A from the table in (2.1). The horizontal steps O_1 and O_2 are indicated with lines of different thickness.

Write the set of all ordered pairs of paths in $P_{k, n-k}$ as

$$P_{k, n-k}^\times := P_{k, n-k} \times P_{k, n-k},$$

and denote the restriction of $P_{k, n-k}^\times$ to those pairs (p, q) with $p \leq q$ as

$$P_{k, n-k}^\leq := \{(p, q) \in P_{k, n-k}^\times : p \leq q\}.$$

We begin by converting the total area between pairs of paths in $P_{k, n-k}^\times$ and $P_{k, n-k}^\leq$ into the (unsigned) area under a single Motzkin path.

Definition 2.1. Write $\mathcal{W}$ for the set of *bilateral Motzkin paths*—that is, lattice paths from $(0, 0)$ to $(n, 0)$ for some $n \in \mathbb{Z}_{\geq 0}$ that use step set $\{U, D, O_1, O_2\}$. We write $\mathcal{W}_{n, k}$ for the set of bilateral Motzkin paths that end at $(n, 0)$ and use exactly k steps of the form U or O_1 . A *bicolored Motzkin path* is a bilateral Motzkin path that stays weakly above the x -axis. Write $\mathcal{M}$ (resp. $\mathcal{M}_{n, k}$) for the set of bicolored Motzkin paths in $\mathcal{W}$ (resp. $\mathcal{W}_{n, k}$).

Proposition 2.2. *The map $A: P_{n,n-k}^\times \rightarrow \mathcal{W}_{n,k}$ is a bijection satisfying $d(p, q) = d(A(p, q))$, and it restricts to a bijection from $P_{n,n-k}^\leq$ to $\mathcal{M}_{n,k}$.*

Proof. The dictionary in (2.1) shows that A is a bijection. When $p \leq q$, $A(p, q)$ never goes below the x -axis, so A maps $P_{n,n-k}^\leq$ onto $\mathcal{M}_{n,k}$. The claim about d was proven immediately after (2.1). $\square$

3. PROOF OF THEOREM 1.2

In this section, we use recurrences on Motzkin paths and their associated generating functions to deduce Theorem 1.2.

3.1. Bicolored Motzkin paths. Define the generating function for bicolored Motzkin paths to be

$$\mathcal{M}(x, u) = \sum_{n \geq 0} \sum_{k \geq 0} |\mathcal{M}_{n,k}| x^n u^k,$$

so that the coefficient of $x^n u^k$ counts bicolored Motzkin paths of total length n with exactly k steps of the form U and O_1 .

Proposition 3.1. *The generating function $\mathcal{M}(x, u)$ satisfies the functional equation*

$$\mathcal{M}(x, u) = 1 + x(1 + u)\mathcal{M}(x, u) + ux^2\mathcal{M}(x, u)^2,$$

with the explicit solution

$$\mathcal{M}(x, u) = \frac{1 - (u+1)x - \sqrt{(u^2 - 2u + 1)x^2 - 2(u+1)x + 1}}{2ux^2}.$$

Proof. The functional equation comes from decomposing a lattice path $r \in \mathcal{M}$ by first return to the x -axis: r is empty; or r starts with an O_1 step; or r starts with an O_2 step; or r starts with a U step. This is illustrated in Figure 4. The explicit solution is easily obtained by solving this quadratic equation in $\mathcal{M}(x, u)$. $\square$




FIGURE 4. The decompositions of lattice paths in $\mathcal{M}$, $\mathcal{M}$, $\mathcal{W}$, and $\mathcal{W}$ from Proposition 3.1, Proposition 3.2, Proposition 3.3, and Proposition 3.5. The multiplication by 2 comes from the symmetry of reflecting the first factor across the x -axis.

Define the generating function for the total area of paths in $\mathcal{M}$ by

$$\mathcal{M}(x, u) := \sum_{n \geq 0} \sum_{k \geq 0} x^n u^k \sum_{p \in \mathcal{M}_{n,k}} d(p).$$

Proposition 3.2. *The generating function $\mathcal{M}(x, u)$ satisfies the functional equation*

$$\mathcal{M}(x, u) = x(1 + u)\mathcal{M}(x, u) + ux^2(2\mathcal{M}(x, u)\mathcal{M}(x, u) + \mathcal{M}(x, u)\frac{d}{dx}(x\mathcal{M}(x, u))),$$

with the explicit solution

$$\mathcal{M}(x, u) = \frac{(u^2 + 1)x^2 - (u + 1)x + ((u + 1)x - 1)\left(\sqrt{(u - 1)^2 x^2 - 2(u + 1)x + 1} - 1\right)}{2ux^2((u - 1)^2 x^2 - 2(u + 1)x + 1)}.$$

Proof. As in Proposition 3.1, the functional equation comes from decomposing a lattice path r in $\mathcal{M}$ by first return to the x -axis: either r is empty (in which case it contributes zero area); or r starts with an O_1 step; or r starts with an O_2 step; or r starts with a U step. This is illustrated in Figure 4. The explicit solution is easily obtained by using the explicit form of $\mathcal{M}(x, u)$ from Proposition 3.1 and solving the linear equation in $\mathcal{M}(x, u)$. $\square$

3.2. Bilateral Motzkin paths. Define the generating function for bilateral Motzkin paths to be

$$\mathcal{W}(x, u) = \sum_{n \geq 0} \sum_{k \geq 0} |\mathcal{W}_{n,k}| x^n u^k,$$

so that the coefficient of $x^n u^k$ counts bilateral Motzkin paths of total length n with exactly k steps of the form U or O_1 .

Proposition 3.3. *The generating function $\mathcal{W}(x, u)$ satisfies the functional equation*

$$\mathcal{W}(x, u) = 1 + x(1 + u)\mathcal{W}(x, u) + 2ux^2\mathcal{M}(x, u)\mathcal{W}(x, u),$$

with the explicit solution

$$\mathcal{W}(x, u) = (u^2 x^2 - 2ux^2 - 2ux + x^2 - 2x + 1)^{-1/2}.$$

Proof. As in Proposition 3.1, the functional equation comes from decomposing a lattice path r in $\mathcal{W}$ by first return to the x -axis: r is empty; or r starts with an O_1 step; or r starts with an O_2 step; or r starts with a U or D step. This is illustrated in Figure 4. The explicit solution is easily obtained by using the explicit form of $\mathcal{M}(x, u)$ from Proposition 3.1 and solving the linear equation in $\mathcal{W}(x, u)$. $\square$

Remark 3.4. Because $\mathcal{W}_{n,k}$ encodes $P_{k,n-k}^\times$ by Proposition 2.2, we have

$$(3.1) \quad \mathcal{W}(x, u) = \sum_{n,k \geq 0} \binom{n}{k}^2 x^n u^k.$$

Define the generating function for the total area of paths in $\mathcal{W}$ to be

$$\mathcal{W}(x, u) := \sum_{n \geq 0} \sum_{k \geq 0} x^n u^k \sum_{p \in \mathcal{W}_{n,k}} d(p).$$

Proposition 3.5. *The generating function $\mathcal{W}(x, u)$ satisfies the functional equation*

$$\mathcal{W}(x, u) = x(1 + u)\mathcal{W}(x, u) + 2ux^2(\mathcal{M}(x, u)\mathcal{W}(x, u) + \mathcal{M}(x, u)\mathcal{W}(x, u) + \mathcal{W}(x, u)\frac{d}{dx}(x\mathcal{M}(x, u))),$$

with the explicit solution

$$(3.2) \quad \mathcal{W}(x, u) = \frac{2ux^2}{((u - 1)^2 x^2 - 2(u + 1)x + 1)^2}.$$

Proof. As in Proposition 3.2, the functional equation comes from decomposing a lattice path r in $\mathcal{W}$ by first return to the x -axis: if r is empty, then it counts for zero area; otherwise, r starts with an O_1 step; or r starts with an O_2 step; or r starts with a U or D step. This is illustrated in Figure 4. The explicit solution is easily obtained by using the explicit forms of $\mathcal{M}(x, u)$, $\mathcal{W}(x, u)$, and $\mathcal{M}(x, u)$ from Propositions 3.1 to 3.3 and solving the linear equation in $\mathcal{W}(x, u)$. $\square$

Substituting $u = y/x$ into Equation (3.2), we obtain Equation (1.3) and thus complete the proof of Theorem 1.2 for the generating function for the Wiener indices of the lattices $P_{m,k} = J([m] \times [k])$.

4. PROOFS OF COROLLARIES 1.3 AND 1.4

We note that

$$x^2 - 2xy + y^2 - 2x - 2y + 1 = (q - t - 1)(q - t + 1)(q + t - 1)(q + t + 1),$$

where $q^2 = x$ and $t^2 = y$. Then, by performing a partial fraction decomposition with a computer algebra system, we get

$$(4.1) \quad \frac{2xy}{(x^2 - 2xy + y^2 - 2x - 2y + 1)^2} = \frac{1}{32} \left(\frac{1}{(-1 - t + q)^2} + \frac{1}{(1 - t + q)^2} + \frac{1}{(-1 + t + q)^2} + \frac{1}{(1 + t + q)^2} \right) \\ + \frac{1}{32t(t+1)} \left(\frac{1+t+t^2}{(1+t-q)} + \frac{1+t+t^2}{(1+t+q)} \right) \\ + \frac{1}{32t(t-1)} \left(\frac{1-t+t^2}{(1-t+q)} + \frac{-1+t-t^2}{(-1+t+q)} \right).$$

Taking the n th coefficient in q from the right-hand side of Equation (4.1) gives

$$(4.2) \quad \frac{n+1}{32} \left((1+t)^{-2-n} + (-1+t)^{-2-n} + (1-t)^{-2-n} + (-1-t)^{-2-n} \right) \\ + \frac{1}{32t} \left((t^2+t+1)(t+1)^{-n-2} + (t^2+t+1)(-t-1)^{-n-2} \right) \\ + \frac{1}{32t} \left((-t^2+t-1)(1-t)^{-n-2} + (-t^2+t-1)(t-1)^{-n-2} \right).$$

Taking the j th coefficient in t from (4.2) and simplifying gives

$$(4.3) \quad \frac{((-1)^n + 1)((-1)^j + 1)}{32} \left(\frac{(n+j+1)!}{n!j!} - \frac{(n+j)!((n+1)^2 + j^2 + j(n+2))}{(n+1)!(j+1)!} \right).$$

Restricting (4.3) to $n = 2m$ and $j = 2k$ even, we obtain the desired expression in Corollary 1.3 for the coefficient of q^{nt^m} , giving the coefficient for $x^m y^k$:

$$\frac{1}{8} \left(\frac{(n+j+1)!}{n!j!} - \frac{(n+j)!((n+1)^2 + j^2 + j(n+2))}{(n+1)!(j+1)!} \right) \\ = \frac{(n+j)!}{8n!j!} \left(n+j+1 - \frac{j^2 + (n+1)^2 + j(n+2)}{(n+1)(j+1)} \right) \\ = \frac{nj}{8(n+j+1)} \binom{n+j+2}{j+1} \\ = \frac{mk}{4m+4k+2} \binom{2m+2k+2}{2k+1}.$$

Given the exact expression for $d(P_{m,k})$, Corollary 1.4 is routine using Stirling's asymptotic equivalent for factorials.

5. SHIFTED STAIRCASES

As for rectangles, we can view elements of Q_n as lattice paths of length n that start at $(0, 0)$ and use steps of the form $U = (1, 1)$ and $D = (1, -1)$. The main difference is that paths representing different order ideals can have different endpoints. Let

$$Q_n^\times := Q_n \times Q_n \quad \text{and} \quad Q_n^\leq := \{(p, q) \in Q_n^\times : p \leq q\}.$$

Definition 5.1. Define a *bilateral Motzkin prefix* to be a lattice path that starts at $(0, 0)$ and uses the steps of the form U, D, O_1, O_2 . Let $\mathcal{V}$ denote the set of bilateral Motzkin prefixes, and let $\mathcal{V}_n$ be the set of bilateral Motzkin prefixes that use exactly n steps. A *bicolored Motzkin prefix* is a bilateral Motzkin prefix that stays weakly above the x -axis. Write $\mathcal{N}$ for the set of bicolored Motzkin prefixes, and let $\mathcal{N}_n = \mathcal{N} \cap \mathcal{V}_n$.

Throughout this section, we write $d(p, q)$ for the length of a geodesic between bilateral Motzkin prefixes p and q in the Hasse diagram of Q_n .

By applying the same rules as in the table in (2.1), we can transform a pair $(p, q) \in Q_n \times Q_n$ into a path $A(p, q)$ that uses steps U, D, O_1, O_2 . The path $A(p, q)$ is similar to a bilateral Motzkin path, except it does not necessarily end on the x -axis. Recall the definition of $\bar{d}$ from (2.2).

Proposition 5.2. *The map $A: Q_n^\times \rightarrow \mathcal{V}_n$ is a bijection satisfying $d(p, q) = \bar{d}(A(p, q))$, and it restricts to a bijection from $Q_n^{\leq}$ to $\mathcal{N}_n$.*

Proof. The proof is essentially the same as that of Proposition 2.2. $\square$

6. PROOF OF THEOREM 1.7 AND COROLLARY 1.8

6.1. Bicolored Motzkin prefixes. Let

$$\mathcal{N}(x) := \sum_{n \geq 0} |\mathcal{N}_n| x^n$$

be the generating function for bicolored Motzkin prefixes.

Proposition 6.1. *The generating function $\mathcal{N}(x)$ satisfies the functional equation*

$$\mathcal{N}(x) = 1 + 3x\mathcal{N}(x) + x^2\mathcal{M}(x, 1)\mathcal{N}(x).$$

Thus,

$$\mathcal{N}(x) = \frac{2}{1 - 4x + \sqrt{1 - 4x}}.$$

Proof. The expression $1 + x\mathcal{N}(x)$ counts (possibly empty) bicolored Motzkin prefixes that only touch the x -axis at $(0, 0)$, while the expression $2x\mathcal{N}(x) + x^2\mathcal{M}(x, 1)\mathcal{N}(x)$ counts bicolored Motzkin prefixes that touch the x -axis at some point other than $(0, 0)$. This is illustrated on the first line of Figure 5. It is routine to derive the explicit solution from the functional equation and Proposition 3.1. $\square$




FIGURE 5. The decompositions of lattice paths in $\mathcal{N}$, $\bar{\mathcal{N}}$, $\mathcal{V}$, and $\bar{\mathcal{V}}$ from Proposition 6.1, Proposition 6.2, Proposition 6.3, and Proposition 6.4.

Define the generating function

$$\mathcal{N}(x) := \sum_{n \geq 0} x^n \sum_{p \in \mathcal{N}_n} \bar{d}(p).$$

Proposition 6.2. *The generating function $\mathcal{N}(x)$ satisfies the functional equation*

$$\begin{aligned} \mathcal{N}(x) &= 2x\mathcal{N}(x) + x^2\mathcal{M}(x, 1)\mathcal{N}(x) + x^2\mathcal{M}(x, 1)\mathcal{N}(x) + x^2\mathcal{N}(x)\frac{\partial}{\partial x}(x\mathcal{M}(x, 1)) \\ &\quad + x\mathcal{N}(x) + x\frac{\partial}{\partial x}(x\mathcal{N}(x)). \end{aligned}$$

Thus,

$$\mathcal{N}(x) = \frac{4x(1 + \sqrt{1-4x} - x(1 - \sqrt{1-4x}))}{\sqrt{1-4x}(1-4x + \sqrt{1-4x})^3}.$$

Proof. Following the same ideas used in the proof of Proposition 3.2, we find that

$$2x\mathcal{N}(x) + x^2\mathcal{M}(x, 1)\mathcal{N}(x) + x^2\mathcal{M}(x, 1)\mathcal{N}(x) + x^2\mathcal{N}(x)\frac{\partial}{\partial x}(x\mathcal{M}(x, 1))$$

counts bicolored Motzkin prefixes that touch the x -axis at some point other than $(0, 0)$, with each path p weighted by $\bar{d}(p)$. The generating function for bicolored Motzkin prefixes that only touch the x -axis at $(0, 0)$ (with each path p weighted by $\bar{d}(p)$) is

$$x\mathcal{N}(x) + x\frac{\partial}{\partial x}(x\mathcal{N}(x)).$$

This is illustrated on the second line of Figure 5. This yields the functional equation, from which the explicit solution is straightforward to obtain via Propositions 3.1 to 3.2. $\square$

Let

$$\mathcal{V}(x) := \sum_{n \geq 0} |\mathcal{V}_n| x^n.$$

Proposition 6.3. *The generating function $\mathcal{V}(x)$ satisfies the functional equation*

$$\mathcal{V}(x) = 1 + 2x\mathcal{N}(x) + 2x\mathcal{V}(x) + 2x^2\mathcal{M}(x, 1)\mathcal{V}(x).$$

Thus,

$$\mathcal{V}(x) = \frac{1 + \sqrt{1-4x}}{\sqrt{1-4x}(1-4x + \sqrt{1-4x})}.$$

Proof. The expression $1 + 2x\mathcal{N}(x)$ counts (possibly empty) bilateral Motzkin prefixes that only touch the x -axis at $(0, 0)$, while the expression $2x\mathcal{V}(x) + 2x^2\mathcal{M}(x, 1)\mathcal{V}(x)$ counts bilateral Motzkin prefixes that touch the x -axis at some point other than $(0, 0)$. This is illustrated on the third line of Figure 5. The explicit solution can be derived from the functional equation using Propositions 3.1 and 6.1. $\square$

Finally, consider the generating function

$$\mathcal{V}(x) := \sum_{n \geq 0} x^n \sum_{p \in \mathcal{V}_n} \bar{d}(p).$$

Proposition 6.4. *The generating function $\mathcal{V}(x)$ satisfies the functional equation*

$$\begin{aligned} \mathcal{V}(x) &= 2x\mathcal{V}(x) + 2x^2\mathcal{M}(x, 1)\mathcal{V}(x) + 2x^2\mathcal{M}(x, 1)\mathcal{V}(x) + 2x^2\mathcal{V}(x)\frac{\partial}{\partial x}(x\mathcal{M}(x, 1)) \\ &\quad + 2x\mathcal{N}(x) + 2x\frac{\partial}{\partial x}(x\mathcal{N}(x)). \end{aligned}$$

Thus,

$$(6.1) \quad \mathcal{V}(x) = \frac{8x(1 + \sqrt{1-4x} - x(3 + \sqrt{1-4x}))}{(1-4x)(1-4x + \sqrt{1-4x})^3}.$$

Proof. Following the same ideas used in the proof of Proposition 6.2, we find that

$$2x\mathcal{V}(x) + 2x^2\mathcal{M}(x, 1)\mathcal{V}(x) + 2x^2\mathcal{M}(x, 1)\mathcal{V}(x) + 2x^2\mathcal{V}(x)\frac{\partial}{\partial x}(x\mathcal{M}(x, 1))$$

counts bilateral Motzkin prefixes that touch the x -axis at some point other than $(0,0)$, with each path p weighted by $\bar{d}(p)$. Furthermore, the generating function for bilateral Motzkin prefixes that only touch the x -axis at $(0,0)$ (with each path p weighted by $\bar{d}(p)$) is

$$2x\mathcal{N}(x) + 2x\frac{\partial}{\partial x}(x\mathcal{N}(x)).$$

This is illustrated on the fourth line of Figure 5. This yields the functional equation, from which one can derive the explicit solution using Propositions 3.1, 6.1, 3.2 and 6.3. $\square$

We conclude Corollary 1.8 by expanding the explicit generating function for $\mathcal{V}(x)$ given in Equation (6.1) as

$$\frac{8x(1 + \sqrt{1-4x} - x(3 + \sqrt{1-4x}))}{(1-4x)(1-4x + \sqrt{1-4x})^3} = \sum_{n \geq 0} a_n x^n.$$

Using a computer algebra system, the coefficients a_n satisfy the difference equation

$$(2n+3)2^{-2n-1}a_n - (4n+5)2^{-2n-3}a_{n+1} + (2+2n)2^{-2n-5}a_{n+2} = 0$$

with initial conditions $a_0 = 0$ and $a_1 = 2$. It is easily checked that $\frac{2n(2n+1)}{3} \binom{2n-1}{n}$ satisfies this equation and initial conditions.

7. ASYMPTOTIC DISTRIBUTIONS

In this section, we prove Propositions 1.5 and 1.9, which describe the asymptotic distribution of the distance between 2 random points (also called 2-point distance) in $P_{\alpha n, n}$ and Q_n , respectively. We start with the case of shifted staircases, which is easier.

7.1. 2-point distance in Q_n . Recall that the elements in Q_n are exactly the lattice paths starting at $(0,0)$, ending somewhere on the line $x = n$, and using steps of the form $U = (1, 1)$ and $D = (1, -1)$. Let p^n and q^n be independent uniform random elements in Q_n . Seeing p^n and q^n as lattice paths, we write p_i^n and q_i^n for their heights after i steps. Clearly, for all $n \geq 1$ and $i \leq n$, one has $p_i^n = X_1 + \dots + X_i$ and $q_i^n = Y_1 + \dots + Y_i$, where $(X_j)_{j \geq 1}$ and $(Y_j)_{j \geq 1}$ are independent sequences of i.i.d. Rademacher random variables of parameter $1/2$. Using Equation (1.4), we write

$$d(p^n, q^n) = \frac{1}{2} \sum_{i=1}^n |p_i^n - q_i^n| = \frac{n}{2} \int_0^1 |p_{[nt]}^n - q_{[nt]}^n| dt.$$

By Donsker's theorem, the processes

$$\left(\frac{1}{\sqrt{n}} p_{[nt]}^n \right)_{t \leq 1} \quad \text{and} \quad \left(\frac{1}{\sqrt{n}} q_{[nt]}^n \right)_{t \leq 1}$$

converge in distribution to independent Brownian motions $(B_t)_{t \leq 1}$ and $(B'_t)_{t \leq 1}$ in Skorokhod space $D[0, 1]$ (see [Bil99, Chapter 3] for background on Skorokhod space). Since integration is a continuous functional on $D[0, 1]$, we have

$$n^{-3/2} d(p^n, q^n) = \frac{1}{2} \int_0^1 \left| n^{-1/2} p_{[nt]}^n - n^{-1/2} q_{[nt]}^n \right| dt \xrightarrow{d} \frac{1}{2} \int_0^1 |B_t - B'_t| dt,$$

where $\xrightarrow{d}$ means convergence in distribution. But $B_t - B'_t \stackrel{d}{=} \sqrt{2} B_t$, proving that $n^{-3/2} d(p^n, q^n)$ converges in distribution to $\frac{1}{\sqrt{2}} \int_0^1 |B_t| dt$, as claimed in Proposition 1.9.

It remains to prove moment convergence. By [Bil12, Corollary of Theorem 25.12], it suffices to show that for each $s > 1$, the sequence of s th moments of $n^{-3/2} d(p^n, q^n)$ is bounded as n tends to $+\infty$. We have

$$n^{-3/2} d(p^n, q^n) \leq n^{-1/2} \max_{i \leq n} |p_i^n| + n^{-1/2} \max_{i \leq n} |q_i^n|.$$

Both terms in the upper bound are identically distributed, so we only consider the first one. By Doob's maximal inequality, we have

$$\mathbb{E} \left[\left(\max_{i \leq n} |p_i^n| \right)^s \right] \leq \left(\frac{s}{s-1} \right)^s \mathbb{E} [|p_n^n|^s].$$

Since p_n^n is a sum of n i.i.d. *centered* random variables, we have the following classical bound on its moments (see, e.g., [Pet89]):

$$\mathbb{E}[|p_n^n|^s] \leq C(s) n^{s/2} \mathbb{E}[|X_1|^s],$$

where $C(s)$ is a constant depending only on s . In particular the s th moment of $n^{-1/2}p_n^n$ is bounded (as n tends to $+\infty$). Consequently, the s th moment of $n^{-1/2} \max_{i \leq n} p_i^n$ is bounded, and that of $n^{-3/2}d(p^n, q^n)$ is as well. This proves that the convergence of $n^{-3/2}d(p^n, q^n)$ to $\frac{1}{\sqrt{2}} \int_0^1 |B_t| dt$ holds also in moments, concluding the proof of Proposition 1.9. $\square$

7.2. 2-point distance in $P_{\alpha,n}$. We now turn to the case of rectangles. Let p^n and q^n be independent uniform random elements in $P_{\alpha,n}$, seen as lattice paths from $(0,0)$ to $((\alpha+1)n, (\alpha-1)n)$. These paths p^n and q^n can be constructed as partial sums of sequences of i.i.d. random variables *under some conditioning*. To this end, let $(X_j)_{j \geq 1}$ and $(Y_j)_{j \geq 1}$ be independent sequences of i.i.d. Rademacher random variables of parameter $\alpha/(\alpha+1)$. We also let $(\tilde{X}_j^n)_{j \geq 1}$ have the distribution of $(X_j)_{j \geq 1}$ conditioned to the event $\sum_{j \leq (\alpha+1)n} X_j = (\alpha-1)n$. Then one has the equality in distribution

$$(p_i^n)_{i \leq (\alpha+1)n} \stackrel{d}{=} \left(\sum_{j \leq i} \tilde{X}_j^n \right)_{i \leq (\alpha+1)n}.$$

Recall that we are interested in the quantity

$$(7.1) \quad n^{-3/2} D_{\alpha,n} = n^{-3/2} d(p^n, q^n) = \frac{1}{2n^{3/2}} \sum_{i=1}^{(\alpha+1)n} |p_i^n - q_i^n| = \frac{\alpha+1}{2\sqrt{n}} \int_0^1 |p_{\lceil(\alpha+1)nt\rceil}^n - q_{\lceil(\alpha+1)nt\rceil}^n| dt.$$

A version of Donsker's theorem for conditioned partial sums has been proved by Liggett [Lig68] (see in particular the corollary of Theorem 4 there). In our case, the centered process

$$\left(\frac{1}{\sigma \sqrt{(1+\alpha)n}} (p_{\lceil(\alpha+1)nt\rceil}^n - \lceil nt \rceil (\alpha-1)) \right)_{0 \leq t \leq 1}$$

converges in distribution to $B_0(t)$ in Skorokhod space $D[0,1]$, where $\sigma^2 = \text{Var}(X_1)$ and $B_0(t)$ is a Brownian bridge. A similar convergence result holds for q^n with an independent Brownian bridge $B'_0(t)$.

Using the continuity of taking integrals on $D[0,1]$, the quantity in (7.1) converges in distribution to

$$\frac{1}{2} (\alpha+1) \sigma \sqrt{\alpha+1} \int_0^1 |B_0(t) - B'_0(t)| dt.$$

An easy computation gives $\sigma = 2\sqrt{\alpha}/(\alpha+1)$, while $B_0(t) - B'_0(t) \stackrel{d}{=} \sqrt{2}B_0(t)$ in distribution. Consequently, $n^{-3/2}D_{\alpha,n}$ converges in distribution to $\sqrt{2\alpha(\alpha+1)} \int_0^1 |B_0(t)| dt$, as claimed in Proposition 1.5.

It remains to prove moment convergence. As above, we shall prove that for any $s > 1$, the random variable $n^{-3/2}D_{\alpha,n}$ has a bounded s th moment as n tends to $+\infty$. Using the convexity of the map $t \mapsto |t|^s$, we obtain

$$(7.2) \quad n^{-s} D_{\alpha,n}^s = 2^{-s} \left(\frac{1}{n} \sum_{i \leq (\alpha+1)n} |p_i^n - q_i^n| \right)^s \leq \frac{2^{-s}}{n} \sum_{i \leq (\alpha+1)n} |p_i^n - q_i^n|^s \leq \frac{1}{n} \sum_{i \leq (\alpha+1)n} \frac{|\bar{p}_i^n|^s + |\bar{q}_i^n|^s}{2},$$

where $\bar{p}_i^n = p_i^n - i \frac{\alpha-1}{\alpha+1}$ is the centered version of p_i^n (and idem for q). Writing $\bar{X}_i = X_i - \frac{\alpha-1}{\alpha+1}$, we have

$$\mathbb{E}[|\bar{p}_i^n|^s] = \mathbb{E} \left[\left| \sum_{j \leq i} \bar{X}_j \right|^s \middle| \sum_{j \leq (\alpha+1)n} \bar{X}_j = 0 \right] = \sum_k |k|^s \mathbb{P} \left[\sum_{j \leq i} \bar{X}_j = k \middle| \sum_{j \leq (\alpha+1)n} \bar{X}_j = 0 \right],$$

where the sum runs over possible values k for $\sum_{j \leq i} \bar{X}_j$. Using the independence of the $\bar{X}_j$, we have

$$\begin{aligned} \mathbb{P} \left[\sum_{j \leq i} \bar{X}_j = k \mid \sum_{j \leq (\alpha+1)n} \bar{X}_j = 0 \right] &= \frac{\mathbb{P} \left[\sum_{j \leq i} \bar{X}_j = k \wedge \sum_{j \leq (\alpha+1)n} \bar{X}_j = 0 \right]}{\mathbb{P} \left[\sum_{j \leq (\alpha+1)n} \bar{X}_j = 0 \right]} \\ &= \mathbb{P} \left[\sum_{j \leq i} \bar{X}_j = k \right] \cdot \frac{\mathbb{P} \left[\sum_{i < j \leq (\alpha+1)n} \bar{X}_j = -k \right]}{\mathbb{P} \left[\sum_{j \leq (\alpha+1)n} \bar{X}_j = 0 \right]}. \end{aligned}$$

Take $i \leq (\alpha+1)n/2$. The probabilities in the fraction can be evaluated asymptotically—uniformly in k —through the local limit theorem (see, e.g., [Dur19, Theorem 3.5.2]), which yields

$$\begin{aligned} \mathbb{P} \left[\sum_{j \leq (\alpha+1)n} \bar{X}_j = 0 \right] &\sim \frac{2}{\sigma \sqrt{2\pi(\alpha+1)n}}; \\ \mathbb{P} \left[\sum_{i < j \leq (\alpha+1)n} \bar{X}_j = -k \right] &= \frac{2e^{-k^2/2((\alpha+1)n-i)\sigma^2}}{\sigma \sqrt{2\pi((\alpha+1)n-i)}} + o(n^{-1/2}) \leq \frac{2+o(1)}{\sigma \sqrt{\pi(\alpha+1)n}}. \end{aligned}$$

In particular, the quotient is bounded by 2 for n large enough, uniformly in k . Bringing everything together, we obtain that for n large enough and $i \leq (\alpha+1)n/2$,

$$\mathbb{E}[\bar{p}_i^n |^s] \leq \sum_k |k|^s \cdot 2\mathbb{P} \left[\sum_{j \leq i} \bar{X}_j = k \right] = 2\mathbb{E} \left[\left| \sum_{j \leq i} \bar{X}_j \right|^s \right].$$

Since the $\bar{X}_j$ are i.i.d. *centered* random variables with finite moments, we have (see, e.g., [Pet89])

$$\mathbb{E} \left[\left| \sum_{j \leq i} \bar{X}_j \right|^s \right] \leq C(s) i^{s/2} \mathbb{E}[|X_1|^s],$$

where $C(s)$ is a constant depending only on s (and α in the sequel), which may change from line to line. Therefore, for n large enough and $i \leq (\alpha+1)n/2$, we have

$$\mathbb{E}[\bar{p}_i^n |^s] \leq C(s) n^{s/2}.$$

By symmetry, this holds also for $i \geq (\alpha+1)n/2$ (we have $p_i^n \stackrel{d}{=} p_{(\alpha+1)n-i}^n$ for all $i \leq (\alpha+1)n$). Going back to (7.2), we get

$$n^{-s} \mathbb{E}[D_{\alpha,n}^s] \leq (\alpha+1) C(s) n^{s/2}.$$

Thus $n^{-3/2} D_{\alpha,n}$ has bounded moments, and the convergence to $\sqrt{2\alpha(\alpha+1)} \int_0^1 |B_0(t)| dt$ holds also in moments. Proposition 1.9 is proved. $\square$

8. HIGHER MOMENTS

Given a finite graph $G = (V, E)$ and a positive integer r , let $d^r(G)$ denote the moment $d^r(G) = \sum_{(p,q) \in V \times V} d(p,q)^r$. The convergence of the distance between two random elements in distribution and in moments established in the previous section yields some asymptotic estimates for $d^r(P_{\alpha,n})$ and $d^r(Q_n)$. In this section, we give an exact expression of $d^2(P_{k,n-k})$. The same method can, in principle, be used to compute the moments $d^r(P_{k,n-k})$ one by one. Similarly, one could use a similar method, drawing from the ideas in Sections 5 and 6, to compute the moments $d^r(Q_n)$. For the sake of brevity, we merely state the explicit formula for $d^2(Q_n)$ and omit the computation.

Proposition 8.1. *We have*

$$d^2(P_{m,k}) = \frac{1}{30} \frac{m+k+1}{m+k} \binom{m+k}{m-1} \binom{m+k}{k-1} (7mk^2 + 7m^2k + 3m^2 + 10mk + 3k^2 + 3m + 3k + 4)$$

and

$$d^2(Q_n) = 2^{2n-4} n (20 + 15(n-2) + 3(n-2)^2).$$

Proof. As mentioned above, we will only prove the first formula. Given a bilateral Motzkin path p , let $\text{len}(p)$ denote the length of p , and let $\mathcal{U}(p)$ be the number of steps in p of the form U or O_1 . Recall that

$$\begin{aligned}\mathcal{W}(x, u) &= \sum_{p \in \mathcal{W}} x^{\text{len}(p)} u^{\mathcal{U}(p)}, & \mathbf{W}(x, u) &= \sum_{p \in \mathcal{W}} x^{\text{len}(p)} u^{\mathcal{U}(p)} d(p), \\ \mathcal{M}(x, u) &= \sum_{p \in \mathcal{M}} x^{\text{len}(p)} u^{\mathcal{U}(p)}, & \mathbf{M}(x, u) &= \sum_{p \in \mathcal{M}} x^{\text{len}(p)} u^{\mathcal{U}(p)} d(p).\end{aligned}$$

Let

$$\mathbb{W}(x, u) = \sum_{p \in \mathcal{W}} x^{\text{len}(p)} u^{\mathcal{U}(p)} d(p)^2 \quad \text{and} \quad \mathbb{M}(x, u) = \sum_{p \in \mathcal{M}} x^{\text{len}(p)} u^{\mathcal{U}(p)} d(p)^2.$$

It follows from Proposition 2.2 that $\mathbb{W}(x, u) = \sum_{n \geq 0} \sum_{k \geq 0} d^2(P_{k, n-k}) x^n u^k$.

The contribution to $\mathbb{M}(x, u)$ coming from paths that start with O_1 or O_2 is $x(u+1)\mathbb{M}(x, u)$. The other paths that contribute to $\mathbb{M}(x, u)$ begin with U and have the form $UpDq$ for some $p, q \in \mathcal{M}$. We find that

$$(8.1) \quad \mathbb{M}(x, u) = x(u+1)\mathbb{M}(x, u) + ux^2\Sigma,$$

where

$$\Sigma = \sum_{p, q \in \mathcal{M}} x^{\text{len}(p)+\text{len}(q)} u^{\mathcal{U}(p)+\mathcal{U}(q)} (d(p) + d(q) + \text{len}(p) + 1)^2.$$

We can write

$$\begin{aligned}(d(p) + d(q) + \text{len}(p) + 1)^2 &= (d(p)^2 + d(q)^2) + (\text{len}(p) + 1)^2 + 2d(p)(\text{len}(p) + 1) \\ &\quad + 2d(q)(\text{len}(p) + 1) + 2d(p)d(q)\end{aligned}$$

to find that

$$\begin{aligned}(8.2) \quad \Sigma &= 2\mathbb{M}(x, u)\mathcal{M}(x, u) + \mathcal{M}(x, u) \frac{\partial}{\partial x} \left(x \frac{\partial}{\partial x} (x\mathcal{M}(x, u)) \right) + 2\mathcal{M}(x, u) \frac{\partial}{\partial x} (x\mathbf{M}(x, u)) \\ &\quad + 2\mathbf{M}(x, u) \frac{\partial}{\partial x} (x\mathcal{M}(x, u)) + 2\mathbf{M}(x, u)^2.\end{aligned}$$

A similar argument yields the functional equation

$$(8.3) \quad \mathbb{W}(x, u) = x(u+1)\mathbb{W}(x, u) + 2ux^2\Sigma',$$

where

$$\begin{aligned}(8.4) \quad \Sigma' &= \sum_{\substack{p \in \mathcal{M} \\ q \in \mathcal{W}}} x^{\text{len}(p)+\text{len}(q)} u^{\mathcal{U}(p)+\mathcal{U}(q)} (d(p) + d(q) + \text{len}(p) + 1)^2 \\ &= \mathbb{M}(x, u)\mathcal{W}(x, u) + \mathbb{W}(x, u)\mathcal{M}(x, u) + \mathcal{W}(x, u) \frac{\partial}{\partial x} \left(x \frac{\partial}{\partial x} (x\mathcal{M}(x, u)) \right) \\ &\quad + 2\mathcal{W}(x, u) \frac{\partial}{\partial x} (x\mathbf{M}(x, u)) + 2\mathbf{W}(x, u) \frac{\partial}{\partial x} (x\mathcal{M}(x, u)) + 2\mathbf{M}(x, u)\mathbf{W}(x, u).\end{aligned}$$

We already computed explicit formulas for $\mathcal{M}(x, u)$, $\mathbf{M}(x, u)$, $\mathcal{W}(x, u)$, and $\mathbf{W}(x, u)$ in Propositions 3.1 to 3.3 and 3.5. Combining those formulas with Equations (8.1) to (8.4), we can derive the explicit formula

$$\mathbb{W}(x, u) = \frac{2ux^2 ((u-1)^2(u+1)x^3 - ((u-8)u+1)x^2 - (u+1)x + 1)}{((ux+x-1)^2 - 4ux^2)^{7/2}}.$$

Setting $u = y/x$ and extracting coefficients yields the desired explicit formula for $d^2(P_{m,k})$. $\square$

9. OPEN PROBLEMS

Comparing the results of Proposition 3.3 and 3.5, we get the intriguing equation

$$(9.1) \quad \mathcal{W} = 2ux^2 \mathcal{W}^4.$$

A direct proof of this would be interesting in itself and could lead to a bijective proof of Corollary 1.3 via the explicit formula (3.1). Recall that $\mathcal{W}$ counts pairs of paths where a cell in the symmetric difference is marked. The connected component where this cell occurs corresponds to a part where the two paths only meet at their beginning and end (this forms a *parallelogram polyomino*), and the generating function $\mathcal{W}^2$ naturally enumerates the rest of the paths. It follows that a bijective proof of (9.1) reduces to a bijective proof that $\mathcal{W}^2$ enumerates the total area of parallelogram polyominoes. The specialization $u = 1$ is known [DLNPR04].

As minuscule lattices arise as the weak order on certain maximal parabolic quotients of finite Coxeter groups, it would be interesting to extend our results to other parabolic quotients. Minuscule lattices also appear as certain crystal graphs; one could also ask about the Wiener indices of more general crystals.

ACKNOWLEDGEMENTS

V.F., P.N., and N.W. thank the organizers of the [2022 Oberwolfach conference in Enumerative Combinatorics](#) for inviting them to a wonderful conference. P.N. was partially supported by the project ANR19-CE48-011-01 (COMBINÉ). N.W. was partially supported by the National Science Foundation under Award No. 2246877. V.F. is partially supported by the Program “Future Leader” of the Initiative “Lorraine Université d’Excellence” (LUE), and by the ANR project CORTIPOM (ANR-21-CE40-0019). C.D. was supported by the National Science Foundation under Award No. 2201907 and by a Benjamin Peirce Fellowship at Harvard University.

Computer exploration to discover the simple exact formulas established in this paper have been carried out with the computer algebra software SageMath [The23], and we are grateful to its developers for their work.

REFERENCES

- [BE23] R. Bourn and W. Q. Erickson. Proof of a conjecture of bourn and willenbring concerning a family of palindromic polynomials, 2023.
- [Bil99] P. Billingsley. *Convergence of probability measures*. Wiley Ser. Probab. Stat. Chichester: Wiley, 2nd ed. edition, 1999.
- [Bil12] P. Billingsley. *Probability and measure. Anniversary edition*. Hoboken, NJ: John Wiley & Sons, 2012.
- [DLNPR04] A. Del Lungo, M. Nivat, R. Pinzani, and S. Rinaldi. A bijection for the total area of parallelogram polyominoes. *Discrete Appl. Math.*, 144(3):291–302, 2004.
- [Dur19] R. Durrett. *Probability. Theory and examples*, volume 49 of *Camb. Ser. Stat. Probab. Math.* Cambridge University Press, 5th edition, 2019.
- [EJS76] R. C. Entringer, D. E. Jackson, and D. Snyder. Distance in graphs. *Czechoslovak Math. J.*, 26(2):283–296, 1976.
- [FT22] V. Féray and P. Thévenin. Components in meandric systems and the infinite noodle. *Int. Math. Res. Not. IMRN*, 2022. doi:10.1093/imrn/rnac156.
- [GNP20] I. Goulden, A. Nica, and D. Puder. Asymptotics for a class of meandric systems, via the Hasse diagram of $\text{NC}(n)$. *Int. Math. Res. Not. IMRN*, 2020(4):983–1034, 2020.
- [Hop20] S. Hopkins. Order polynomial product formulas and poset dynamics. *arXiv preprint arXiv:2006.01568*, 2020.
- [Jan07] S. Janson. Brownian excursion area, wright’s constants in graph enumeration, and other Brownian areas. *Probab. Surv.*, 4:80–145, 2007.
- [Lig68] T. M. Liggett. An invariance principle for conditioned sums of independent random variables. *J. Math. Mech.*, 18:559–570, 1968.
- [Pet89] V. V. Petrov. Moments of sums of independent random variables. *J. Sov. Math.*, 61(1):1, 1989.
- [Pro84] R. A. Proctor. Bruhat lattices, plane partition generating functions, and minuscule representations. *European J. Combin.*, 5(4):331–350, 1984.

- [Rou02] D. H. Rouvray. The rich legacy of half a century of the Wiener index. In *Topology in Chemistry*, pages 16–37. Elsevier, 2002.
- [Sta80] R. P. Stanley. Weyl groups, the hard Lefschetz theorem, and the Sperner property. *SIAM J. Alg. Disc. Meth.*, 1(2):168–184, 1980.
- [SW12] J. Striker and N. Williams. Promotion and rowmotion. *European J. Combin.*, 33(8):1919–1942, 2012.
- [The23] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 9.8)*, 2023. <https://www.sagemath.org>.
- [Wei21] E. W. Weisstein. Wiener index. MathWorld—A Wolfram Web Resource, accessed 2021.
- [Wie47] H. Wiener. Structural determination of paraffin boiling points. *J. Am. Chem. Soc.*, 69(1):17–20, 1947.
- [Wik22] Wikipedia contributors. Wiener index — Wikipedia, the free encyclopedia, 2022. [Online; accessed 22-December-2022].
- [Wil22] N. Williams. Oberwolfach problem session: Enumerative combinatorics 2022. https://personal.utdallas.edu/~nxw170830/docs/Papers/Misc/Oberwolfach_EC_2022_problem_session.pdf, 2022. Accessed: December 20, 2022.

(C. Defant) DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02139, USA
Email address: `colindefant@gmail.com`

(V. Féray) UNIVERSITÉ DE LORRAINE, CNRS, IECL, F-54000, NANCY, FRANCE
Email address: `valentin.feray@univ-lorraine.fr`

(P. Nadeau) UNIV LYON, CNRS, UNIVERSITÉ CLAUDE BERNARD LYON 1, INSTITUT CAMILLE JORDAN, F-69622 VILLEURBANNE CEDEX, FRANCE
Email address: `nadeau@math.univ-lyon1.fr`

(N. Williams) UNIVERSITY OF TEXAS AT DALLAS
Email address: `nathan.williams1@utdallas.edu`