

Geometric Decomposition and Efficient Implementation of High Order Face and Edge Elements

Chunyu Chen¹, Long Chen², Xuehai Huang^{3,*} and Huayi Wei¹

¹ School of Mathematics and Computational Science, Xiangtan University; National Center of Applied Mathematics in Hunan, Hunan Key Laboratory for Computation and Simulation in Science and Engineering Xiangtan 411105, China.

² Department of Mathematics, University of California at Irvine, Irvine, CA 92697, USA.

³ School of Mathematics, Shanghai University of Finance and Economics, Shanghai 200433, China.

Received 24 September 2023; Accepted (in revised version) 22 February 2024

Abstract. This study investigates high-order face and edge elements in finite element methods, with a focus on their geometric attributes, indexing management, and practical application. The exposition begins by a geometric decomposition of Lagrange finite elements, setting the foundation for further analysis. The discussion then extends to $H(\text{div})$ -conforming and $H(\text{curl})$ -conforming finite element spaces, adopting variable frames across differing sub-simplices. The imposition of tangential or normal continuity is achieved through the strategic selection of corresponding bases. The paper concludes with a focus on efficient indexing management strategies for degrees of freedom, offering practical guidance to researchers and engineers. It serves as a comprehensive resource that bridges the gap between theory and practice.

AMS subject classifications: 65N30, 35Q60

Key words: Implementation of finite elements, nodal finite elements, $H(\text{curl})$ -conforming, $H(\text{div})$ -conforming.

1 Introduction

This paper introduces node-based basis functions for high-order finite elements, specifically focusing on Lagrange, BDM (Brezzi-Douglas-Marini) [8, 9, 18], and second-kind Nédélec elements [6, 18]. These elements are subsets of the spaces H^1 , $H(\text{div})$, and

*Corresponding author. Email addresses: cbtxs@smail.xtu.edu.cn (C. Chen), chenlong@math.uci.edu (L. Chen), huang.xuehai@sufe.edu.cn (X. Huang), weihuayi@xtu.edu.cn (H. Wei)

$H(\text{curl})$, with their shape functions being the full polynomial space $\mathbb{P}_k^n$, where k represents the polynomial degree, and n the geometric dimension. Notably, varying continuity across these elements gives rise to distinct characteristics. When $n=3$, for the lowest degree $k=1$, degrees of freedom (DoFs) of the $H(\text{div})$ -conforming finite element are posed on faces and DoFs of the $H(\text{curl})$ -conforming finite element are posed on edges. Therefore, conventionally an $H(\text{div})$ -conforming finite element is referred to as a face element, and an $H(\text{curl})$ -conforming element is an edge element. They are also known as the second family of face and edge element as the shape function space is the full polynomial space while the first family consists of incomplete polynomial spaces [6].

In the realm of Lagrange finite elements, nodal basis functions stand out for their simplicity and ease of computation. In contrary, constructing basis functions for face and edge elements is more intricate. Traditional approaches involve the Piola transformations, where basis functions are first devised on a reference element and subsequently mapped to the actual element using either covariant (to preserve tangential continuity, in the case of edge elements) or contravariant (to maintain normal continuity, for face elements) Piola transformations. Detailed explanations of this approach can be found in [17, 20], and implementation is in open-source software such as MFEM [5] and Fenics [4].

Arnold, Falk and Winther, in [7], introduced a geometric decomposition of polynomial differential forms. Basis functions based on Bernstein polynomials were proposed, paving the way for subsequent advancements. In [1, 3], basis functions founded on Bernstein polynomials were explored, accompanied by fast algorithms for the matrix assembly. Additionally, hierarchical basis functions for $H(\text{curl})$ -conforming finite elements were introduced in [2, 22-24].

While these methods offer valuable insights, they can be quite complex. Researchers have thus ventured into simpler approaches. In [14], a method multiplying scalar nodal finite element methods by vectors was introduced, resulting in $H(\text{div})$ and $H(\text{curl})$ conforming finite elements that exhibit continuity on both vertices and edges.

We propose a straightforward method to construct nodal bases for the second family face and edge elements. Initially, we clarify the basis of Lagrange elements by considering them dual to the degrees of freedom, which are determined by values at interpolation points. We then extend this principle to vector polynomial spaces, wherein each interpolation point establishes a frame that includes both tangential and normal (t-n) directions. We impose the continuity of either tangential or normal components by appropriate choice of the t-n decomposition. We explicitly derive the dual basis functions for these elements from the basis functions of Lagrange elements.

This idea has been previously explored in [10, 15, 16] for constructing a hierarchical basis of $H(\text{div})$ elements in two and three dimensions. Through a rotation process, it was also adapted for $H(\text{curl})$ elements in two dimensions, as detailed in [15, 16]. However, extending their methodology to higher dimensions presented significant challenges. Our work advances this field by developing a geometric decomposition of the second family face and edge elements in arbitrary dimensions and orders. Additionally, we introduce

degrees of freedom that are dual to the basis functions, aspects not covered in the aforementioned studies.

An essential aspect of practical finite element implementations is the management of local DoFs, ensuring proper mapping to global DoFs. This is critical for maintaining the correct continuity between elements and correctly integrating matrices and vectors across the system. Despite its importance, there is a scarcity of literature specifically addressing DoF management for high-order finite elements. In this paper, we explore the global indexing of interpolation points in Lagrange finite elements of arbitrary degree. Our discussion comprehensively covers DoF management for Lagrange, the second family face and edge elements. Our goal is to facilitate the implementation of high-order finite element methods by simplifying their DoF management.

The subsequent sections of this paper are organized as follows. Section 2 elucidates foundational concepts such as simplicial lattices, interpolation points, bubble polynomials, and triangulations. Sections 3, 4, and 5 delve into the construction of basis functions for Lagrange, the second family face and edge elements, respectively. Section 6 addresses the management of degrees of freedom, a critical aspect of effective matrix assembly. Finally, Section 7 presents two numerical examples, solving the mixed Poisson problem and Maxwell problem using the second family face and edge elements, respectively, to validate the correctness of the proposed basis function construction method.

2 Preliminaries

In this section, we provide essential foundations for our study. We introduce multi-indices, simplicial lattices, and interpolation points. We explain sub-simplices and their relations. Furthermore, we introduce the dictionary ordering of simplicial lattices and the concept of bubble polynomials.

2.1 Simplicial lattice

A multi-index of length $n+1$ is an array of non-negative integers:

$$\alpha = (\alpha_0, \alpha_1, \dots, \alpha_n), \quad \alpha_i \in \mathbb{N}, i = 0, \dots, n.$$

The degree or sum of the multi-index is $|\alpha| = \sum_{i=0}^n \alpha_i$ and factorial is $\alpha! = \prod_{i=0}^n (\alpha_i!)$. The set of all multi-indices of length $n+1$ and degree k will be called a simplicial lattice and denoted by $\mathbb{T}_k^n$, i.e.,

$$\mathbb{T}_k^n = \{\alpha \in \mathbb{N}^{n+1} : |\alpha| = k\}.$$

The elements in $\mathbb{T}_k^n$ can be linearly indexed by the dictionary ordering R_n :

$$R_n(\alpha) = \sum_{i=1}^n \binom{\alpha_i + \alpha_{i+1} + \dots + \alpha_n + n - i}{n+1-i}.$$

For example, for an element α in $\mathbb{T}_k^2$, the index is given by the mapping:

$$R_2(\alpha) = \frac{(\alpha_1 + \alpha_2)(\alpha_1 + \alpha_2 + 1)}{2} + \alpha_2.$$

Notice that α_0 is not used in the calculation of $R_n(\alpha)$.

2.2 Interpolation points

Let $x_0, x_1, \dots, x_n$ be $n+1$ points in $\mathbb{R}^n$ and

$$T = \text{Convex}(x_0, x_1, \dots, x_n) = \left\{ \sum_{i=0}^n \lambda_i x_i : 0 \leq \lambda_i \leq 1, \sum_{i=0}^n \lambda_i = 1 \right\},$$

be an n -simplex, where $\lambda = (\lambda_0, \dots, \lambda_n)$ is called the barycentric coordinate. We can have a geometric embedding of the algebraic set $\mathbb{T}_k^n$ as follows:

$$\mathcal{X}_T = \left\{ x_\alpha = \frac{1}{k} \sum_{i=0}^n \alpha_i x_i : \alpha \in \mathbb{T}_k^n \right\},$$

which is called the set of interpolation points with degree k on T ; see Fig. 1 for $k=4$ and $n=2$. In literature [19], it is called the k -th order principal lattice of n -simplex T . Given $\alpha \in \mathbb{T}_k^n$, the barycentric coordinate of α is given by

$$\lambda(\alpha) = (\alpha_0, \alpha_1, \dots, \alpha_n) / k.$$

The ordering of $\mathcal{X}_T$ is also given by $R_n(\alpha)$. Note that the indexing map $R_n(\alpha)$ is only a local ordering of the interpolation points on one n -simplex. In Section 6, we will discuss the global indexing of all interpolation points on the triangulation composed of simplexes.

By this geometric embedding, we can apply operators to the geometric simplex T . For example, $\mathcal{X}_{\hat{T}}$ denotes the set of interpolation points in the interior of T and $\mathcal{X}_{\partial T}$ is the set of interpolation points on the boundary of T .




Figure 1: The interpolation points of $\mathbb{T}_4^2$ and their linear indices and multi-indices.

2.3 Sub-simplices and sub-simplicial lattices

Following [7], let $\Delta(T)$ denote all the subsimplices of T , while $\Delta_\ell(T)$ denotes the set of subsimplices of dimension ℓ for $\ell=0, \dots, n$. The capital letter F is reserved for an $(n-1)$ -dimensional face of T and $F_i \in \Delta_{n-1}(T)$ denotes the face opposite to x_i for $i=0, 1, \dots, n$.

For a sub-simplex $f \in \Delta_\ell(T)$, following [13], we will overload the notation f for both the geometric simplex and the algebraic set of indices. Namely $f = \{f(0), \dots, f(\ell)\} \subseteq \{0, 1, \dots, n\}$ and

$$f = \text{Convex}(x_{f(0)}, \dots, x_{f(\ell)}) \in \Delta_\ell(T)$$

is the ℓ -dimensional simplex spanned by the vertices $x_{f(0)}, \dots, x_{f(\ell)}$.

If $f \in \Delta_\ell(T)$, then $f^* \in \Delta_{n-\ell-1}(T)$ denotes the sub-simplex of T opposite to f . When treating f as a subset of $\{0, 1, \dots, n\}$, $f^* \subseteq \{0, 1, \dots, n\}$ so that $f \cup f^* = \{0, 1, \dots, n\}$, i.e., f^* is the complement of set f . Geometrically,

$$f^* = \text{Convex}(x_{f^*(1)}, \dots, x_{f^*(n-\ell)}) \in \Delta_{n-\ell-1}(T)$$

is the $(n-\ell-1)$ -dimensional simplex spanned by vertices not contained in f .

Given a sub-simplex $f \in \Delta_\ell(T)$, through the geometric embedding $f \hookrightarrow T$, we define the prolongation/extension operator $E: \mathbb{T}_k^\ell \rightarrow \mathbb{T}_k^n$ as follows:

$$E(\alpha)_{f(i)} = \alpha_i, \quad i=0, \dots, \ell, \quad \text{and} \quad E(\alpha)_j = 0, \quad j \notin f. \quad (2.1)$$

Take $f = \{1, 3, 4\}$ for example, then the extension $E(\alpha) = (0, \alpha_0, 0, \alpha_1, \alpha_2, \dots, 0)$ for $\alpha = (\alpha_0, \alpha_1, \alpha_2) \in \mathbb{T}_k^\ell(f)$. With a slight abuse of notation, for a node $\alpha_f \in \mathbb{T}_k^\ell(f)$, we still use the same notation $\alpha_f \in \mathbb{T}_k^n(T)$ to denote such extension. The geometric embedding $x_{E(\alpha)} \in f$ which justifies the notation $\mathbb{T}_k^\ell(f)$ and its geometric embedding will be denoted by $\mathcal{X}_f$, which consists of interpolation points on f . $\mathbb{T}_k^\ell(f^\circ)$ is the set of lattice points whose geometric embedding is in the interior of f , i.e., $\mathcal{X}_f^\circ$.

The Bernstein representation of polynomial of degree k on a simplex T is

$$\mathbb{P}_k(T) := \text{span}\{\lambda^\alpha = \lambda_0^{\alpha_0} \lambda_1^{\alpha_1} \dots \lambda_n^{\alpha_n}, \alpha \in \mathbb{T}_k^n\}.$$

The bubble polynomial of f is a polynomial of degree $\ell+1$:

$$b_f := \lambda_f = \lambda_{f(0)} \lambda_{f(1)} \dots \lambda_{f(\ell)} \in \mathbb{P}_{\ell+1}(f).$$

2.4 Triangulation

Let Ω be a polyhedral domain in $\mathbb{R}^n$, $n \geq 1$. A geometric triangulation $\mathcal{T}_h$ of Ω is a set of n -simplices such that

$$\cup_{T \in \mathcal{T}_h} T = \Omega, \quad \hat{T}_i \cap \hat{T}_j = \emptyset, \quad \forall T_i, T_j \in \mathcal{T}_h, \quad T_i \neq T_j.$$

The subscript h denotes the diameter of each element and can be understood as a piecewise constant function on $\mathcal{T}_h$. A triangulation is conforming if the intersection of two

simplexes are common lower dimensional sub-simplex. We shall restrict to conforming triangulations in this paper.

The interpolation points on a conforming triangulation $\mathcal{T}_h$ is

$$\mathcal{X} = \bigcup_{T \in \mathcal{T}_h} \mathcal{X}_T. \quad (2.2)$$

Note that a lot of duplications exist in (2.2). A direct sum of the interpolation set is given by

$$\mathcal{X} = \Delta_0(\mathcal{T}_h) \oplus \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(\mathcal{T}_h)} \mathcal{X}_f, \quad (2.3)$$

where $\Delta_\ell(\mathcal{T}_h)$ denotes the set of ℓ -dimensional subsimplices of $\mathcal{T}_h$. In implementation, computation of local matrices on each simplex is based on (2.2) while to assemble a matrix equation, (2.3) should be used. As Lagrange element is globally continuous, the indexing of interpolation points on vertices, edges, faces should be unique and a mapping from the local index to the global index is needed. The index mapping from (2.2) to (2.3) will be discussed in Section 6.

3 Geometric decompositions of Lagrange elements

In this section, we present a geometric decomposition for Lagrange finite elements on n -dimensional simplices. We introduce the concept of Lagrange interpolation basis functions, where function values at interpolation points serve as degrees of freedom.

3.1 Geometric decomposition

For the polynomial space $\mathbb{P}_k(T)$ with $k \geq 1$ on an n -dimensional simplex T , we have the following geometric decomposition of Lagrange element [7, (2.6)] and a proof can be found in [12]. The integral at a vertex is understood as the function value at that vertex and $\mathbb{P}_k(\mathbf{x}) = \mathbb{R}$. When $k < 0$, $\mathbb{P}_k(T) = \emptyset$.

Theorem 3.1 (Geometric decomposition of Lagrange element). *For the polynomial space $\mathbb{P}_k(T)$ with $k \geq 1$ on an n -dimensional simplex T , we have the following decomposition*

$$\mathbb{P}_k(T) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} b_f \mathbb{P}_{k-(\ell+1)}(f). \quad (3.1)$$

The function $u \in \mathbb{P}_k(T)$ is uniquely determined by DoFs

$$\int_f u p \, ds, \quad p \in \mathbb{P}_{k-(\ell+1)}(f), \quad f \in \Delta_\ell(T), \quad \ell = 0, 1, \dots, n. \quad (3.2)$$

Introduce the bubble polynomial space of degree k on a sub-simplex f as

$$\mathbb{B}_k(f) := b_f \mathbb{P}_{k-(\ell+1)}(f), \quad f \in \Delta_\ell(T), \quad 1 \leq \ell \leq n.$$

When $\ell \geq k$, $\mathbb{B}_k(f) = \emptyset$. It is called a bubble space as

$$\text{tr}^{\text{grad}} u := u|_{\partial f} = 0, \quad u \in \mathbb{B}_k(f).$$

Then we can write (3.1) as

$$\mathbb{P}_k(T) = \mathbb{P}_1(T) \oplus \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_k(f). \quad (3.3)$$

That is a polynomial of degree k can be decomposed into a linear polynomial plus bubbles on edges, faces, and all sub-simplices.

Based on a conforming triangulation $\mathcal{T}_h$, the k -th order Lagrange finite element space $V_k^L(\mathcal{T}_h)$ is defined as

$$V_k^L(\mathcal{T}_h) = \{v \in C(\Omega) : v|_T \in \mathbb{P}_k(T), T \in \mathcal{T}_h, \text{ and DoFs (3.2) are single valued}\},$$

and will have a geometric decomposition

$$V_k^L(\mathcal{T}_h) = V_1^L(\mathcal{T}_h) \oplus \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(\mathcal{T}_h)} \mathbb{B}_k(f). \quad (3.4)$$

Here we extend the polynomial on f to each element T containing f by the Bernstein form and extension of multi-index; see $E(\alpha)$ defined in (2.1). Consequently the dimension of $V_k^L(\mathcal{T}_h)$ is

$$\dim V_k^L(\mathcal{T}_h) = \sum_{\ell=0}^n |\Delta_\ell(\mathcal{T}_h)| \binom{k-1}{\ell},$$

where $|\Delta_\ell(\mathcal{T}_h)|$ is the cardinality of number of $\Delta_\ell(\mathcal{T}_h)$, i.e., the number of ℓ -dimensional simplices in $\mathcal{T}_h$. We understand $\binom{k-1}{\ell} = 0$ if $\ell > k-1$. That is the degree of the polynomial dictates the dimension of the sub-simplex in the geometric decompositions (3.1) and (3.4).

The geometric decomposition (3.1) can be naturally extended to vector Lagrange elements. For $k \geq 1$, define

$$\mathbb{B}_k^n(f) := b_f \mathbb{P}_{k-(\ell+1)}(f) \otimes \mathbb{R}^n.$$

Clearly we have

$$\mathbb{P}_k^n(T) = \mathbb{P}_1^n(T) \oplus \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_k^n(f). \quad (3.5)$$

For an $f \in \Delta_\ell(T)$, we choose a $t-n$ coordinate $\{\mathbf{t}_i^f, \mathbf{n}_j^f, i=1, \dots, \ell, j=1, \dots, n-\ell\}$ so that

- $\mathcal{T}^f := \text{span}\{\mathbf{t}_1^f, \dots, \mathbf{t}_\ell^f\}$ is the tangential plane of f ;
- $\mathcal{N}^f := \text{span}\{\mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$ is the normal plane of f .

When $\ell=0$, i.e., for vertices, no tangential component, and for $\ell=n$, no normal component. We abbreviate $\mathbf{n}_1^F$ as $\mathbf{n}_F$ for $F \in \Delta_{n-1}(T)$, and $\mathbf{t}_1^e$ as $\mathbf{t}_e$ for $e \in \Delta_1(T)$. We have the trivial decompositions

$$\mathbb{R}^n = \mathcal{T}^f \oplus \mathcal{N}^f, \quad \mathbb{B}_k^n(f) = [\mathbb{B}_k(f) \otimes \mathcal{T}^f] \oplus [\mathbb{B}_k(f) \otimes \mathcal{N}^f]. \quad (3.6)$$

Restricted to an ℓ -dimensional sub-simplex $f \in \Delta_\ell(T)$, define

$$\mathbb{B}_k^\ell(f) := \mathbb{B}_k(f) \otimes \mathcal{T}^f,$$

which is a space of ℓ -dimensional vectors on the tangential space $\mathcal{T}^f$ with vanishing trace tr^{grad} on ∂f .

When move to a triangulation $\mathcal{T}_h$, we shall call a basis of $\mathcal{T}^f$ or $\mathcal{N}^f$ is global if it depends only on f not the element T containing f . Otherwise it is called local and may vary in different elements.

3.2 Lagrange interpolation basis functions

Previously DoFs (3.2) are given by moments on sub-simplexes. Now we present a set of DoFs as function values on the interpolation points and give its dual basis for the k -th order Lagrange element on an n -simplex.

Lemma 3.1 (Lagrange interpolation basis functions [19]). *A basis function of the k -th order Lagrange finite element space on T is:*

$$\phi_\alpha(x) = \frac{1}{\alpha!} \prod_{i=0}^n \prod_{j=0}^{\alpha_i-1} (k\lambda_i(x) - j), \quad \alpha \in \mathbb{T}_k^n,$$

with the DoFs defined as the function value at the interpolation points:

$$N_\alpha(u) = u(x_\alpha), \quad x_\alpha \in \mathcal{X}_T.$$

Proof. It is straightforward to verify the duality of the basis and DoFs

$$N_\beta(\phi_\alpha) = \phi_\alpha(x_\beta) = \delta_{\alpha,\beta} = \begin{cases} 1 & \text{if } \alpha = \beta, \\ 0 & \text{otherwise.} \end{cases}$$

As

$$|\mathbb{T}_k^n| = \binom{n+k}{k} = \dim \mathbb{P}_k(T),$$

$\{\phi_\alpha, \alpha \in \mathbb{T}_k^n\}$ is a basis of $\mathbb{P}_k(T)$ and $\{N_\alpha, \alpha \in \mathbb{T}_k^n\}$ is a basis of the dual space $\mathbb{P}_k^*(T)$. $\square$

Given a triangulation $\mathcal{T}_h$ and degree k , recall that

$$\mathcal{X}_{\mathcal{T}_h} = \bigcup_{T \in \mathcal{T}_h} \mathcal{X}_T = \Delta_0(\mathcal{T}_h) \oplus \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(\mathcal{T}_h)} \mathcal{X}_f^\ell.$$

Denote by

$$\mathbb{T}_k^n(\mathcal{T}_h) := \bigoplus_{x_i \in \Delta_0(\mathcal{T}_h)} \mathbb{T}_k^0(x_i) \oplus \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(\mathcal{T}_h)} \mathbb{T}_k^\ell(f).$$

For a lattice $\alpha \in \mathbb{T}_k^\ell(f)$, we use extension operator E defined in (2.1) to extend α to each simplex T containing f . We also extend the polynomial on f to T by the Bernstein form. Hereafter, for simplicity, we set $\mathcal{X}_{\hat{v}} := \mathcal{X}_v$ and $\mathbb{T}_k^0(\hat{v}) := \mathbb{T}_k^0(v)$ for vertex $v \in \Delta_0(\mathcal{T}_h)$.

Theorem 3.2 (DoFs of Lagrange finite element on $\mathcal{T}_h$). *A basis for the k -th Lagrange finite element space $V_k^L(\mathcal{T}_h)$ is*

$$\{\phi_\alpha, \alpha \in \mathbb{T}_k^n(\mathcal{T}_h)\}$$

with DoFs

$$N_\alpha(u) = u(\mathbf{x}_\alpha), \quad \mathbf{x}_\alpha \in \mathcal{X}_{\mathcal{T}_h}.$$

Proof. For $F \in \Delta_{n-1}(\mathcal{T}_h)$, thanks to Lemma 3.1, $\phi_\alpha|_F$ is uniquely determined by DoFs $\{u(\mathbf{x}_\alpha), \mathbf{x}_\alpha \in \mathcal{X}_F\}$ on face F , hence $\phi_\alpha \in V_k^L(\mathcal{T}_h)$. Clearly the cardinality of $\{\phi_\alpha, \alpha \in \mathbb{T}_k^n(\mathcal{T}_h)\}$ is same as the dimension of space $V_k^L(\mathcal{T}_h)$. Then we only need to show these functions are linearly independent, which follows from the fact $N_\beta(\phi_\alpha) = \phi_\alpha(\mathbf{x}_\beta) = \delta_{\alpha,\beta}$ for $\alpha, \beta \in \mathbb{T}_k^n(\mathcal{T}_h)$. $\square$

We now generalize the basis for a scalar Lagrange element to a vector Lagrange element. For an interpolation point $\mathbf{x} \in \mathcal{X}_T$, let $\{\mathbf{e}_i^x, i = 0, \dots, n-1\}$ be a basis of $\mathbb{R}^n$, and its dual basis is denoted by $\{\hat{\mathbf{e}}_i^x, i = 0, \dots, n-1\}$, i.e.,

$$(\hat{\mathbf{e}}_i^x, \mathbf{e}_j^x) = \delta_{i,j},$$

where $(\mathbf{a}, \mathbf{b}) = \mathbf{a} \cdot \mathbf{b}$ is the inner product of two vectors in $\mathbb{R}^n$. When $\{\mathbf{e}_i^x, i = 0, \dots, n-1\}$ is orthonormal, its dual basis is itself.

Corollary 3.1. *A polynomial function $\mathbf{u} \in \mathbb{P}_k^n(T)$ can be uniquely determined by the DoFs:*

$$N_\alpha^i(\mathbf{u}) := \mathbf{u}(\mathbf{x}_\alpha) \cdot \mathbf{e}_i^{x_\alpha}, \quad \mathbf{x}_\alpha \in \mathcal{X}_T, \quad i = 0, \dots, n-1.$$

The basis function on T dual to this set of DoFs can be explicitly written as:

$$\boldsymbol{\phi}_\alpha^i(\mathbf{x}) = \phi_\alpha(\mathbf{x}) \hat{\mathbf{e}}_i^{x_\alpha}, \quad \alpha \in \mathbb{T}_k^n, \quad i = 0, \dots, n-1.$$

Proof. It is straightforward to verify the duality

$$N_\beta^j(\boldsymbol{\phi}_\alpha^i) = \boldsymbol{\phi}_\alpha^i(\mathbf{x}_\beta) \cdot \mathbf{e}_j^{x_\beta} = \phi_\alpha(\mathbf{x}_\beta) \hat{\mathbf{e}}_i^{x_\alpha} \cdot \mathbf{e}_j^{x_\beta} = \delta_{i,j} \delta_{\alpha,\beta},$$

for $\alpha, \beta \in \mathbb{T}_k^n, i, j = 0, \dots, n-1$. $\square$

If the basis $\{\mathbf{e}_i^f, i = 0, \dots, n-1\}$ is global in the sense that it is independent of element T containing $\mathbf{x}$, we get the continuous vector Lagrange elements. Choose different basis will give different continuity.

4 Geometric decompositions of face elements

Define $H(\operatorname{div}, \Omega) := \{v \in L^2(\Omega; \mathbb{R}^n) : \operatorname{div} v \in L^2(\Omega)\}$. For a subdomain $K \subseteq \Omega$, the trace operator for the div operator is

$$\operatorname{tr}_K^{\operatorname{div}} v = \mathbf{n} \cdot v|_{\partial K} \quad \text{for } v \in H(\operatorname{div}, \Omega),$$

where $\mathbf{n}$ denotes the outwards unit normal vector of ∂K . Given a triangulation $\mathcal{T}_h$ and a piecewise smooth function $\mathbf{u}$, it is well known that $\mathbf{u} \in H(\operatorname{div}, \Omega)$ if and only if $\mathbf{n}_F \cdot \mathbf{u}$ is continuous across all faces $F \in \Delta_{n-1}(\mathcal{T}_h)$, which can be ensured by having DoFs on faces. An $H(\operatorname{div})$ -conforming finite element is thus also called a face element.

4.1 Geometric decomposition

Define the polynomial div bubble space

$$\mathbb{B}_k(\operatorname{div}, T) = \ker(\operatorname{tr}_T^{\operatorname{div}}) \cap \mathbb{P}_k^n(T).$$

Recall that $\mathbb{B}_k^\ell(f) = \mathbb{B}_k(f) \otimes \mathcal{T}^f$ consists of bubble polynomials on the tangential plane of f . For $\mathbf{u} \in \mathbb{B}_k^\ell(f)$, as $\mathbf{u}$ is on the tangential plane, $\mathbf{u} \cdot \mathbf{n}_F = 0$ for $f \subseteq F$. When $f \not\subseteq F$, $b_f|_F = 0$. So $\mathbb{B}_k^\ell(f) \subseteq \mathbb{B}_k(\operatorname{div}, T)$ for $k \geq 2$ and $\dim f \geq 1$. In [12], we have proved that the div-bubble polynomial space has the following decomposition.

Lemma 4.1. For $k \geq 2$,

$$\mathbb{B}_k(\operatorname{div}, T) = \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_k^\ell(f).$$

Notice that as no tangential plane on vertices, there is no div-bubble associated to vertices and consequently the degree of a div-bubble polynomial is greater than or equal to 2. Next we present two geometric decompositions of a div-element.

Theorem 4.1. For $k \geq 1$, we have

$$\mathbb{P}_k^n(T) = \mathbb{P}_1^n(T) \oplus \left(\bigoplus_{\ell=1}^{n-1} \bigoplus_{f \in \Delta_\ell(T)} (\mathbb{B}_k(f) \otimes \mathcal{N}^f) \right) \oplus \mathbb{B}_k(\operatorname{div}, T), \quad (4.1)$$

$$\mathbb{P}_k^n(T) = \left(\bigoplus_{F \in \Delta_{n-1}(T)} (\mathbb{P}_k(F) \mathbf{n}_F) \right) \oplus \mathbb{B}_k(\operatorname{div}, T). \quad (4.2)$$

Proof. The first decomposition (4.1) is a rearrangement of (3.5) by merging the tangential component $\mathbb{B}_k^\ell(f)$ into the bubble space $\mathbb{B}_k(\operatorname{div}, T)$.

Next we prove the decomposition (4.2). For an ℓ -dimensional, $0 \leq \ell \leq n-1$, sub-simplex $f \in \Delta_\ell(T)$, we choose the $n-\ell$ face normal vectors $\{\mathbf{n}_F : F \in \Delta_{n-1}(T), f \subseteq F\}$ as the basis of $\mathcal{N}^f$. Therefore we have

$$\mathbb{B}_k(f) \otimes \mathcal{N}^f = \bigoplus_{F \in \Delta_{n-1}(T), f \subseteq F} \mathbb{B}_k(f) \mathbf{n}_F.$$

Then (4.1) becomes

$$\mathbb{P}_k^n(T) = \mathbb{P}_1^n(T) \oplus \left(\bigoplus_{\ell=1}^{n-1} \bigoplus_{f \in \Delta_\ell(T)} \bigoplus_{F \in \Delta_{n-1}(T), f \subseteq F} \mathbb{B}_k(f) \mathbf{n}_F \right) \oplus \mathbb{B}_k(\operatorname{div}, T).$$

At a vertex v , we choose $\{\mathbf{n}_F : F \in \Delta_{n-1}(T), v \in F\}$ as the basis of $\mathbb{R}^n$ and write

$$\begin{aligned}\mathbb{P}_1^n(T) &= \bigoplus_{x \in \Delta_0(T)} \bigoplus_{F \in \Delta_{n-1}(T), x \in \Delta_0(F)} \text{span}\{\lambda_x \mathbf{n}_F\} \\ &= \bigoplus_{F \in \Delta_{n-1}(T)} \bigoplus_{x \in \Delta_0(F)} \text{span}\{\lambda_x \mathbf{n}_F\} \\ &= \bigoplus_{F \in \Delta_{n-1}(T)} \mathbb{P}_1(F) \mathbf{n}_F.\end{aligned}$$

Then by swapping the ordering of f and F in the direct sum, i.e.,

$$\bigoplus_{\ell=1}^{n-1} \bigoplus_{f \in \Delta_\ell(T)} \bigoplus_{F \in \Delta_{n-1}(T), f \subseteq F} \rightarrow \bigoplus_{F \in \Delta_{n-1}(T)} \bigoplus_{\ell=1}^{n-1} \bigoplus_{f \in \Delta_\ell(F)},$$

and using the decomposition (3.1) of Lagrange element, we obtain the decomposition (4.2). $\square$

In decomposition (4.1), we single out $\mathbb{P}_1^n(T)$ to emphasize an $H(\text{div})$ -conforming element can be obtained by adding div-bubble and normal component on sub-simplexes starting from edges. In (4.2), we group all normal components facewisely which leads to the classical BDM element.

As $H(\text{div}, \Omega)$ -conforming elements require the normal continuity across each F , the normal vector $\mathbf{n}_F$ is chosen globally. Namely $\mathbf{n}_F$ depends on F only. It may coincide with the outwards or inwards normal vector for an element T containing F . On the contrary, all tangential basis for $\mathcal{T}^f$ are local and thus the tangential component are multi-valued and merged to the element-wise div bubble function $\mathbb{B}_k(\text{div}, T)$.

4.2 A nodal basis for the BDM element

For the efficient implementation, we employ the DoFs of the function values at interpolation nodes and combine with $t-n$ decompositions to present a nodal basis for the BDM element.

Given an $f \in \Delta_\ell(T)$, we choose $\{\mathbf{n}_F, F \in \Delta_{n-1}(T), f \subseteq F\}$ as the basis for its normal plane $\mathcal{N}^f$ and an arbitrary basis $\{\mathbf{t}_i^f, i=1, \dots, \ell\}$ for the tangential plane. We shall choose $\{\hat{\mathbf{n}}_{F_i} \in \mathcal{N}^f, i \in f^*\}$ a basis of $\mathcal{N}^f$ dual to $\{\mathbf{n}_{F_i} \in \mathcal{N}^f, i \in f^*\}$, i.e.,

$$(\mathbf{n}_F, \hat{\mathbf{n}}_{F'}) = \delta_{F, F'}, \quad F, F' \in \Delta_{n-1}(T), \quad f \in F \cap F'.$$

Similarly choose a basis $\{\hat{\mathbf{t}}_i^f\}$ dual to $\{\mathbf{t}_i^f\}$.

We can always choose an orthonormal basis for the tangential plane $\mathcal{T}^f$ but for the normal plane $\mathcal{N}^f$ with basis $\{\mathbf{n}_{F_i} \in \mathcal{N}^f, i \in f^*\}$, we use Lemma 4.2 to find its dual basis.

For $f \in \Delta_\ell(T)$ and $e \in \partial f$, let $\mathbf{n}_f^e$ be a unit normal vector of e but tangential to f . When $\ell=1$, f is an edge and e is a vertex. Then $\mathbf{n}_f^e$ is the edge vector of f . Using these notations we can give an explicit expression of the dual basis $\{\hat{\mathbf{n}}_{F_i} \in \mathcal{N}^f, i \in f^*\}$.

Lemma 4.2. For $f \in \Delta_\ell(T)$,

$$\left\{ \hat{n}_{F_i} = \frac{1}{\mathbf{n}_{f+i}^f \cdot \mathbf{n}_{F_i}} \mathbf{n}_{f+i}^f, i \in f^* \right\}, \quad (4.3)$$

where $f+i$ denotes the $(\ell+1)$ -dimensional face in $\Delta_{\ell+1}(T)$ with vertices $f(0), \dots, f(\ell)$ and i for $i \in f^*$, is a basis of $\mathcal{N}^f$ dual to $\{\mathbf{n}_{F_i} \in \mathcal{N}^f, i \in f^*\}$.

Proof. Clearly $\mathbf{n}_{f+i}^f \in \mathcal{N}^f$ for $i \in f^*$. It suffices to prove

$$\mathbf{n}_{f+i}^f \cdot \mathbf{n}_{F_j} = 0 \quad \text{for } i, j \in f^*, i \neq j,$$

which follows from $\mathbf{n}_{f+i}^f \in \mathcal{T}^{f+i}$ and $f+i \subseteq F_j$. $\square$

By Corollary 3.1, we obtain a nodal basis for BDM face elements.

Theorem 4.2. For each $f \in \Delta_\ell(T)$, we choose $\{\mathbf{e}_i^f, i=0, \dots, n-1\} = \{\mathbf{t}_i^f, i=1, \dots, \ell, \mathbf{n}_{F_i}, i \in f^*\}$ and its dual basis $\{\hat{\mathbf{e}}_i^f, i=0, \dots, n-1\} = \{\hat{\mathbf{t}}_i^f, i=1, \dots, \ell, \hat{\mathbf{n}}_{F_i}, i \in f^*\}$. A basis function of the k -th order BDM element space on T is:

$$\{\phi_\alpha(\mathbf{x}) \hat{\mathbf{e}}_i^f, i=0, 1, \dots, n-1, \alpha \in \mathbb{T}_k^\ell(\hat{f})\}_{f \in \Delta_\ell(T), \ell=0, 1, \dots, n},$$

with the DoFs at the interpolation points defined as:

$$\{\mathbf{u}(\mathbf{x}_\alpha) \cdot \mathbf{e}_i^f, i=0, 1, \dots, n-1, \mathbf{x}_\alpha \in \mathcal{X}_f^f\}_{f \in \Delta_\ell(T), \ell=0, 1, \dots, n}. \quad (4.4)$$

By choosing a global normal basis in the sense that $\mathbf{n}_F$ depending only on F not the element containing F , we impose the continuity on the normal direction. We choose a local $\mathbf{t}^f$, i.e., for different element T containing f , $\phi_\alpha(\mathbf{x}) \mathbf{t}^f(T)$ is different, then no continuity is imposed for the tangential direction.

Define the global finite element space

$$V_{h,k}^{\text{div}} := \{\mathbf{u} \in L^2(\Omega; \mathbb{R}^n) : \mathbf{u}|_T \in \mathbb{P}_k(T; \mathbb{R}^n) \text{ for each } T \in \mathcal{T}_h, \\ \text{all the DoFs } \mathbf{u}(\mathbf{x}_\alpha) \cdot \mathbf{n}_F \text{ in (4.4) across } F \in \Delta_{n-1}(\mathcal{T}_h) \text{ are single-valued}\}.$$

By Theorem 4.1, $V_{h,k}^{\text{div}} \subset H(\text{div}, \Omega)$ and is equivalent to the BDM space.

The novelty is that we only need a basis of Lagrange element which is well documented; see Section 3.2. Coupling with different $t-n$ decomposition at different sub-simplex, we obtain the classical face elements. This concept has been explored in the works of [15, 16] and [10], focusing on the implementation of the $H(\text{div})$ element in two and three dimensions. Additionally, the adaptation of a 2D $H(\text{curl})$ element through rotation is discussed in [15, 16]. As we will demonstrate in the subsequent section, extending the edge element to higher dimensions presents significant challenges. This extension necessitates a comprehensive characterization of the curl operator and its associated polynomial bubble space.

5 Geometric decompositions of edge elements

In this section we present geometric decompositions of $H(\text{curl})$ -conforming finite element space on an n -dimensional simplex. We first generalize the curl differential operator to n dimensions and study its trace which motivates our decomposition. We then give an explicit basis dual to function values at interpolation points.

5.1 Differential operator and its trace

Denote by $\mathbb{S}$ and $\mathbb{K}$ the subspace of symmetric matrices and skew-symmetric matrices of $\mathbb{R}^{n \times n}$, respectively. For a smooth vector function $\mathbf{v}$, define

$$\text{curl} \mathbf{v} := 2\text{skw}(\text{grad} \mathbf{v}) = \text{grad} \mathbf{v} - (\text{grad} \mathbf{v})^\top,$$

which is a skew-symmetric matrix function. In two dimensions, for $\mathbf{v} = (v_1, v_2)^\top$,

$$\text{curl} \mathbf{v} = \begin{pmatrix} 0 & \partial_{x_2} v_1 - \partial_{x_1} v_2 \\ \partial_{x_1} v_2 - \partial_{x_2} v_1 & 0 \end{pmatrix} = \text{mskw}(\text{rot} \mathbf{v}),$$

where

$$\text{mskw} \mathbf{u} := \begin{pmatrix} 0 & -u \\ u & 0 \end{pmatrix}, \quad \text{rot} \mathbf{v} := \partial_{x_1} v_2 - \partial_{x_2} v_1.$$

In three dimensions, for $\mathbf{v} = (v_1, v_2, v_3)^\top$,

$$\text{curl} \mathbf{v} = \begin{pmatrix} 0 & \partial_{x_2} v_1 - \partial_{x_1} v_2 & \partial_{x_3} v_1 - \partial_{x_1} v_3 \\ \partial_{x_1} v_2 - \partial_{x_2} v_1 & 0 & \partial_{x_3} v_2 - \partial_{x_2} v_3 \\ \partial_{x_1} v_3 - \partial_{x_3} v_1 & \partial_{x_2} v_3 - \partial_{x_3} v_2 & 0 \end{pmatrix} = \text{mskw}(\nabla \times \mathbf{v}),$$

where

$$\text{mskw} \mathbf{u} := \begin{pmatrix} 0 & -u_3 & u_2 \\ u_3 & 0 & -u_1 \\ -u_2 & u_1 & 0 \end{pmatrix}$$

with $\mathbf{u} = (u_1, u_2, u_3)^\top$. Hence we can identify $\text{curl} \mathbf{v}$ as scalar $\text{rot} \mathbf{v}$ in two dimensions, and vector $\nabla \times \mathbf{v}$ in three dimensions. In general, $\text{curl} \mathbf{u}$ is understood as a skew-symmetric matrix.

Define Sobolev space

$$H(\text{curl}, \Omega) := \{\mathbf{v} \in L^2(\Omega; \mathbb{R}^n) : \text{curl} \mathbf{v} \in L^2(\Omega; \mathbb{K})\}.$$

Given a face $F \in \Delta_{n-1}(T)$, define the trace operator of curl as

$$\text{tr}_F^{\text{curl}} \mathbf{v} = 2\text{skw}(\mathbf{v} \mathbf{n}_F^\top)|_F = (\mathbf{v} \mathbf{n}_F^\top - \mathbf{n}_F \mathbf{v}^\top)|_F.$$

We define tr^{curl} as a piecewise defined operator as

$$(\text{tr}^{\text{curl}} \mathbf{v})|_F = \text{tr}_F^{\text{curl}} \mathbf{v}, \quad F \in \Delta_{n-1}(T).$$

For a vector $\mathbf{v} \in \mathbb{R}^n$ and an $(n-1)$ -dimensional face F , the tangential part of $\mathbf{v}$ on F is

$$\Pi_F \mathbf{v} := \mathbf{v}|_F - (\mathbf{v}|_F \cdot \mathbf{n}_F) \mathbf{n}_F = \sum_{i=1}^{n-1} (\mathbf{v}|_F \cdot \mathbf{t}_i^F) \mathbf{t}_i^F,$$

where $\{\mathbf{t}_i^F, i=1, \dots, n-1\}$ is an orthonormal basis of F . As we treat $\text{curl} \mathbf{v}$ as a matrix, so is the trace $\text{tr}_F^{\text{curl}} \mathbf{v}$, while the tangential component of $\mathbf{v}$ is a vector. Their relation is given in the following lemma.

Lemma 5.1. *For face $F \in \Delta_{n-1}(T)$, we have*

$$\text{tr}_F^{\text{curl}} \mathbf{v} = 2 \text{skw}((\Pi_F \mathbf{v}) \mathbf{n}_F^\top), \quad \Pi_F \mathbf{v} = (\text{tr}_F^{\text{curl}} \mathbf{v}) \mathbf{n}_F. \quad (5.1)$$

Proof. By the decomposition $\mathbf{v}|_F = \Pi_F \mathbf{v} + (\mathbf{v}|_F \cdot \mathbf{n}_F) \mathbf{n}_F$,

$$\text{tr}_F^{\text{curl}} \mathbf{v} = 2 \text{skw}((\Pi_F \mathbf{v}) \mathbf{n}_F^\top + (\mathbf{v}|_F \cdot \mathbf{n}_F) \mathbf{n}_F \mathbf{n}_F^\top) = 2 \text{skw}((\Pi_F \mathbf{v}) \mathbf{n}_F^\top),$$

which implies the first identity. Then by $\mathbf{n}_F^\top \mathbf{n}_F = 1$ and $(\Pi_F \mathbf{v})^\top \mathbf{n}_F = 0$,

$$(\text{tr}_F^{\text{curl}} \mathbf{v}) \mathbf{n}_F = ((\Pi_F \mathbf{v}) \mathbf{n}_F^\top - \mathbf{n}_F (\Pi_F \mathbf{v})^\top) \mathbf{n}_F = \Pi_F \mathbf{v},$$

i.e. the second identity holds. $\square$

Thanks to (5.1), the vanishing tangential part $\Pi_F \mathbf{v}$ and the vanishing tangential trace $(\text{tr}_F^{\text{curl}} \mathbf{v})$ are equivalent.

Lemma 5.2. *Let $\mathbf{v} \in L^2(\Omega; \mathbb{R}^n)$ and $\mathbf{v}|_T \in H^1(T; \mathbb{R}^n)$ for each $T \in \mathcal{T}_h$. Then $\mathbf{v} \in H(\text{curl}, \Omega)$ if and only if $\Pi_F \mathbf{v}|_{T_1} = \Pi_F \mathbf{v}|_{T_2}$ for all interior face $F \in \Delta_{n-1}(\mathcal{T}_h)$, where T_1 and T_2 are two elements sharing F .*

Proof. It is an immediate result of Lemma 5.1 in [6] and (5.1). $\square$

5.2 Polynomial bubble space

Define the polynomial bubble space of degree k for the curl operator as

$$\mathbb{B}_k(\text{curl}, T) = \ker(\text{tr}^{\text{curl}}) \cap \mathbb{P}_k^n(T).$$

For Lagrange bubble space $\mathbb{B}_k^n(T)$, all components of the vector function vanish on ∂T and consequently on all sub-simplex with dimension $\leq n-1$. For $\mathbf{u} \in \mathbb{B}_k(\text{curl}, T)$, only the tangential component vanishes, which will imply $\mathbf{u}$ vanishes on sub-simplex with dimension $\leq n-2$.

Lemma 5.3. *For $\mathbf{u} \in \mathbb{B}_k(\text{curl}, T)$, it holds $\mathbf{u}|_f = \mathbf{0}$ for all $f \in \Delta_\ell(T)$, $0 \leq \ell \leq n-2$. Consequently $\mathbb{B}_k(\text{curl}, T) \subset \mathbb{B}_k^n(T) \oplus \bigoplus_{F \in \Delta_{n-1}(T)} \mathbb{B}_k^n(F)$.*

Proof. It suffices to consider a sub-simplex $f \in \Delta_{n-2}(T)$. Let $F_1, F_2 \in \Delta_{n-1}(T)$ such that $f = F_1 \cap F_2$. By $\text{tr}_{F_i}^{\text{curl}} \mathbf{u} = \mathbf{0}$ for $i = 1, 2$, we have $\Pi_{F_i} \mathbf{u} = \mathbf{0}$ and consequently

$$(\mathbf{u} \cdot \mathbf{t}_i^f)|_f = 0, (\mathbf{u} \cdot \mathbf{n}_{F_1}^f)|_f = (\mathbf{u} \cdot \mathbf{n}_{F_2}^f)|_f = 0 \quad \text{for } i = 1, \dots, n-2,$$

where $\mathbf{n}_{F_i}^f$ is a normal vector f sitting on F_i . As $\text{span}\{\mathbf{t}_1^f, \dots, \mathbf{t}_{n-2}^f, \mathbf{n}_{F_1}^f, \mathbf{n}_{F_2}^f\} = \mathbb{R}^n$, we acquire $\mathbf{u}|_f = \mathbf{0}$. By the property of face bubbles, we conclude $\mathbf{u}$ is a linear combination of element bubble and $(n-1)$ -dimensional face bubbles. $\square$

Obviously $\mathbb{B}_k^n(T) \subset \mathbb{B}_k(\text{curl}, T)$. As tr^{curl} contains the tangential component only, the normal component $\mathbb{B}_k(F)\mathbf{n}_F$ is also a curl bubble. The following result says their sum is precisely all curl bubble polynomials.

Theorem 5.1. For $k \geq 1$, it holds that

$$\mathbb{B}_k(\text{curl}, T) = \mathbb{B}_k^n(T) \oplus \bigoplus_{F \in \Delta_{n-1}(T)} \mathbb{B}_k(F)\mathbf{n}_F, \quad (5.2)$$

and

$$\text{tr}^{\text{curl}} : \mathbb{P}_1^n(T) \oplus \bigoplus_{\ell=1}^{n-2} \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_k^n(f) \oplus \bigoplus_{F \in \Delta_{n-1}(T)} \mathbb{B}_k^{n-1}(F) \rightarrow \text{tr}^{\text{curl}} \mathbb{P}_k^n(T) \quad (5.3)$$

is a bijection.

Proof. It is obvious that

$$\mathbb{B}_k^n(T) \oplus \bigoplus_{F \in \Delta_{n-1}(T)} \mathbb{B}_k(F)\mathbf{n}_F \subseteq \mathbb{B}_k(\text{curl}, T).$$

Then apply the trace operator to the decomposition (3.5) to conclude that the map tr^{curl} in (5.3) is onto.

Now we prove it is also injective. Take a function $\mathbf{u} \in \mathbb{P}_1^n(T) \oplus \bigoplus_{\ell=1}^{n-2} \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_k^n(f) \oplus \bigoplus_{F \in \Delta_{n-1}(T)} \mathbb{B}_k^{n-1}(F)$ and $\text{tr}^{\text{curl}} \mathbf{u} = \mathbf{0}$. By Lemma 5.3, we can assume $\mathbf{u} = \sum_{F \in \Delta_{n-1}(T)} \mathbf{u}_k^F$ with $\mathbf{u}_k^F \in \mathbb{B}_k^{n-1}(F)$. Take $F \in \Delta_{n-1}(T)$. We have $\mathbf{u}|_F = \mathbf{u}_k^F|_F \in \mathbb{B}_k^{n-1}(F)$. Hence $(\mathbf{u}_k^F \cdot \mathbf{t})|_F = (\mathbf{u} \cdot \mathbf{t})|_F = 0$ for any $\mathbf{t} \in \mathcal{T}^F$, which results in $\mathbf{u}_k^F = \mathbf{0}$. Therefore $\mathbf{u} = \mathbf{0}$.

Once we have proved the map tr in (5.3) is bijection, we conclude (5.2) from the decomposition (3.5). $\square$

We will use curl_f to denote the curl operator restricted to a sub-simplex f with $\dim f \geq 1$. For $f \in \Delta_\ell(T)$, $\ell = 2, \dots, n-1$, by applying Theorem 5.1 to f , we have

$$\mathbb{B}_k(\text{curl}_f, f) = \mathbb{B}_k^\ell(f) \oplus \bigoplus_{e \in \partial f} \mathbb{B}_k(e)\mathbf{n}_f^e. \quad (5.4)$$

Notice that the curl_f -bubble function is defined for $\ell \geq 2$ not including edges. Indeed, for an edge e and a vertex x of e , $\mathbf{n}_e^x$ is $\mathbf{t}_e$ if x is the ending vertex of e and $-\mathbf{t}_e$ otherwise. Then for $\ell = 1$

$$\mathbb{B}_k(e)\mathbf{t}_e \oplus \bigoplus_{x \in \partial e} \text{span}\{\lambda_x \mathbf{n}_e^x\} = \mathbb{P}_k(e)\mathbf{t}_e, \quad (5.5)$$

which is the full polynomial not vanishing on ∂e .

5.3 Geometric decompositions

For $e \in \Delta_\ell(T)$, we choose the basis $\{\mathbf{n}_f^e : f = e + i, i \in e^*\}$ for $\mathcal{N}^e$ and a basis $\{\mathbf{t}_i^e, i = 1, \dots, \ell\}$ for $\mathcal{T}^e$. So we have the following geometric decompositions of $\mathbb{P}_k^n(T)$.

Theorem 5.2. For $k \geq 1$, we have

$$\begin{aligned} \mathbb{P}_k^n(T) = & \mathbb{P}_1^n(T) \oplus \bigoplus_{\ell=1}^n \bigoplus_{e \in \Delta_\ell(T)} (\mathbb{B}_k(e) \otimes \text{span}\{\mathbf{t}_i^e\}_{i=1}^\ell) \\ & \oplus (\mathbb{B}_k(e) \otimes \text{span}\{\mathbf{n}_{e+i}^e, i \in e^*\}), \end{aligned} \quad (5.6)$$

$$\mathbb{P}_k^n(T) = \bigoplus_{e \in \Delta_1(T)} \mathbb{P}_k(e) \mathbf{t}_e \oplus \bigoplus_{\ell=2}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_k(\text{curl}_f, f). \quad (5.7)$$

Proof. Decomposition (5.6) is the component form of decomposition (3.5). We can write $\mathbb{B}_k(e) \otimes \text{span}\{\mathbf{n}_{e+i}^e, i \in e^*\} = \bigoplus_{f \in \Delta_{\ell+1}(T), e \subseteq f} \mathbb{B}_k(e) \mathbf{n}_f^e$. Then in the summation (5.6), we shift the normal component one level up and switch the sum of e and f :

$$\begin{aligned} & \bigoplus_{\ell=1}^n \bigoplus_{e \in \Delta_\ell(T)} [\mathbb{B}_k^\ell(e) \oplus \bigoplus_{f \in \Delta_{\ell+1}(T), e \subseteq f} \mathbb{B}_k(e) \mathbf{n}_f^e] \\ & = \bigoplus_{e \in \Delta_1(T)} \mathbb{B}_k(e) \mathbf{t}_e \oplus \bigoplus_{\ell=2}^n \bigoplus_{f \in \Delta_\ell(T)} [\mathbb{B}_k^\ell(f) \oplus \bigoplus_{e \subseteq \partial f} \mathbb{B}_k(e) \mathbf{n}_f^e]. \end{aligned}$$

Then by the characterization (5.4) of $\mathbb{B}_k(\text{curl}_f, f)$ and (5.5), we get the decomposition (5.7). $\square$

Decomposition (5.7) is the counterpart of (3.3) for Lagrange element.

5.4 Tangential-normal decomposition of the second family of edge elements

Recall that for $e \in \Delta_{\ell-1}(T)$, the basis $\{\mathbf{n}_f^e : f \in \Delta_\ell(T), e \subseteq f\}$ of $\mathcal{N}^e$ is dual to the basis $\{\mathbf{n}_F : F \in \Delta_{n-1}(T), e \subseteq F\}$; see Lemma 4.2

Theorem 5.3. Take $\mathbb{P}_k^n(T)$ as the shape function space. Then it is determined by the following DoFs

$$\int_e \mathbf{u} \cdot \mathbf{t} \, p \, ds, \quad p \in \mathbb{P}_k(e), \quad e \in \Delta_1(T), \quad (5.8a)$$

$$\int_f \mathbf{u} \cdot \mathbf{p} \, ds, \quad \mathbf{p} \in \mathbb{B}_k(\text{curl}_f, f), \quad f \in \Delta_\ell(T), \quad \ell = 2, \dots, n. \quad (5.8b)$$

Proof. Based on the decomposition (5.6), the shape function $\mathbb{P}_k^n(T)$ is determined by the following DoFs

$$\int_e \mathbf{u} \cdot \mathbf{t} \, p \, ds, \quad p \in \mathbb{P}_k(e), \quad e \in \Delta_1(T), \quad (5.9a)$$

$$\int_f (\mathbf{u} \cdot \mathbf{t}_i^f) \, p \, ds, \quad p \in \mathbb{P}_{k-(\ell+1)}(f), \quad i = 1, \dots, \ell, \quad (5.9b)$$

$$\int_e (\mathbf{u} \cdot \mathbf{n}_f^e) \, p \, ds, \quad p \in \mathbb{P}_{k-\ell}(e), \quad e \in \partial f \quad (5.9c)$$

for $f \in \Delta_\ell(T)$, $\ell = 2, \dots, n$. By (5.2) and (5.4), DoFs (5.9) are equivalent to DoFs (5.8). $\square$

Remark 5.1. The DoFs of the second kind Nédélec edge element in [6,18] are

$$\begin{aligned} \int_e \mathbf{u} \cdot \mathbf{t} \, p \, ds, \quad p \in \mathbb{P}_k(e), \quad e \in \Delta_1(T), \\ \int_f \mathbf{u} \cdot \mathbf{p} \, ds, \quad \mathbf{p} \in \mathbb{P}_{k-\ell}^\ell(f) + (\Pi_f \mathbf{x}) \mathbb{P}_{k-\ell}(f), \quad f \in \Delta_\ell(T), \quad \ell = 2, \dots, n. \end{aligned}$$

There is an isomorphism between $\mathbb{P}_{k-\ell}^\ell(f) + (\Pi_f \mathbf{x}) \mathbb{P}_{k-\ell}(f)$ and $\mathbb{B}_k(\text{curl}_f, f)$ for $f \in \Delta_\ell(T)$ with $\ell = 2, \dots, n$, that is $\mathbb{B}_k(\text{curl}_f, f)$ is uniquely determined by DoF

$$\int_f \mathbf{u} \cdot \mathbf{p} \, ds, \quad \mathbf{p} \in \mathbb{P}_{k-\ell}^\ell(f) + (\Pi_f \mathbf{x}) \mathbb{P}_{k-\ell}(f),$$

whose proof can be found in Lemma 4.7 in [6].

Given an $e \in \Delta_{\ell-1}(T_h)$, we choose a global $\{\mathbf{n}_f^e, e \subseteq f \in \Delta_\ell(T_h)\}$ as the basis for the normal plane $\mathcal{N}^e$ and a global basis $\{\mathbf{t}_i^e\}$ of $\mathcal{T}^e$. Define the global finite element space

$$\begin{aligned} V_{h,k}^{\text{curl}} = \{ \mathbf{u} \in L^2(\Omega; \mathbb{R}^n) : \mathbf{u}|_T \in \mathbb{P}_k(T; \mathbb{R}^n) \text{ for each } T \in \mathcal{T}_h, \\ \text{all the DoFs (5.8) are single-valued} \}. \end{aligned}$$

Lemma 5.4. We have $V_h^{\text{curl}} \subset H(\text{curl}, \Omega)$.

Proof. For an $F \in \Delta_{n-1}(T)$, DoFs (5.9) related to $\Pi_F \mathbf{u}$ are

$$\begin{aligned} \int_e (\Pi_F \mathbf{u} \cdot \mathbf{t}_i^e) \, p \, ds, \quad i = 1, \dots, \ell-1, \quad p \in \mathbb{P}_{k-\ell}(e), \quad e \in \Delta_{\ell-1}(F), \quad \ell = 2, \dots, n, \\ \int_e (\Pi_F \mathbf{u} \cdot \mathbf{n}_f^e) \, p \, ds, \quad f \in \Delta_\ell(F), \quad e \subseteq f, \quad p \in \mathbb{P}_{k-\ell}(e), \quad e \in \Delta_{\ell-1}(F), \quad \ell = 1, \dots, n-1, \end{aligned}$$

thanks to DoFs (5.9), which uniquely determine $\Pi_F \mathbf{u}$. □

5.5 The second family of edge elements

By Corollary 3.1, we obtain a nodal basis for the second-kind Nédélec edge element.

Theorem 5.4. For each $e \in \Delta_\ell(T)$, $\ell = 0, 1, \dots, n$, we choose $\{\mathbf{e}_i^e, i = 0, \dots, n-1\} = \{\mathbf{t}_i^e, i = 1, \dots, \ell, \mathbf{n}_f^e, f = e+i, i \in e^*\}$ and its dual basis $\{\hat{\mathbf{e}}_i^e, i = 0, \dots, n-1\} = \{\hat{\mathbf{t}}_i^e, i = 1, \dots, \ell, \mathbf{n}_{F_i} / (\mathbf{n}_{F_i} \cdot \mathbf{n}_{e+i}^e), i \in e^*\}$. A basis function of the k -th order second kind Nédélec edge element space on T is:

$$\{\phi_\alpha(\mathbf{x}) \hat{\mathbf{e}}_i^e, i = 0, 1, \dots, n-1, \alpha \in \mathbb{T}_k^\ell(\hat{\mathbf{e}})\}_{e \in \Delta_\ell(T), \ell = 0, 1, \dots, n},$$

with the DoFs at the interpolation points defined as:

$$\{\mathbf{u}(\mathbf{x}_\alpha) \cdot \mathbf{e}_i^e, i = 0, 1, \dots, n-1, \mathbf{x}_\alpha \in \mathcal{X}_\ell\}_{e \in \Delta_\ell(T), \ell = 0, 1, \dots, n}. \quad (5.10)$$

Notice that the basis $\{t_i^e\}$ depends only on e and n_f^e depends on e and f but independent T . By asking the corresponding DoFs single valued, we obtain the tangential continuity. We can define the edge finite element space as

$$V_{h,k}^{\text{curl}} = \{u \in L^2(\Omega; \mathbb{R}^n) : u|_T \in \mathbb{P}_k(T; \mathbb{R}^n) \text{ for each } T \in \mathcal{T}_h, \text{ the DoFs } u(x_\alpha) \cdot t_i^e, u(x_\alpha) \cdot n_f^e \text{ in (5.10) are single-valued across all } e \text{ and } f \text{ with } \dim e, \dim f \leq n-1\}.$$

5.5.1 2D basis on triangular meshes

Let T be a triangle, for lattice point x located in different sub-simplices, we shall choose different frame $\{e_x^0, e_x^1\}$ at x and its dual frame $\{\hat{e}_x^0, \hat{e}_x^1\}$ as follows:

1. If $x \in \Delta_0(T)$, assume the two adjacent edges are e_0 and e_1 , then

$$e_x^0 = t_{e_0}, \quad e_x^1 = t_{e_1}, \quad \hat{e}_x^0 = \frac{n_{e_1}}{n_{e_1} \cdot t_{e_0}}, \quad \hat{e}_x^1 = \frac{n_{e_0}}{n_{e_0} \cdot t_{e_1}}.$$

2. If $x \in \mathcal{X}_{\hat{e}}, e \in \Delta_1(T)$, then

$$e_x^0 = t_e, \quad e_x^1 = n_e, \quad \hat{e}_x^0 = t_e, \quad \hat{e}_x^1 = n_e.$$

3. If $x \in \mathcal{X}_{\hat{T}}$, then

$$e_x^0 = (1, 0), \quad e_x^1 = (0, 1), \quad \hat{e}_x^0 = (1, 0), \quad \hat{e}_x^1 = (0, 1).$$




Figure 2: The left figure shows $\{e_0, e_1\}$ at each interpolation point, the right figure shows $\{\hat{e}_0, \hat{e}_1\}$ at each interpolation point.

5.5.2 3D basis on tetrahedron meshes

Let T be a tetrahedron, for any $x \in \mathcal{X}_T$, define a frame $\{e_x^0, e_x^1, e_x^2\}$ at x and its dual frame $\{\hat{e}_x^0, \hat{e}_x^1, \hat{e}_x^2\}$ as follows:




Figure 3: The left figure shows $\{e_0, e_1, e_2\}$ at each interpolation point, the right figure shows $\{\hat{e}_0, \hat{e}_1, \hat{e}_2\}$ at each interpolation point.

1. If $x \in \Delta_0(T)$ and adjacent edges of x are e_0, e_1, e_2 , adjacent faces of x are f_0, f_1, f_2 , then

$$\begin{aligned} e_x^0 &= t_{e_0}, & e_x^1 &= t_{e_1}, & e_x^2 &= t_{e_2}, \\ \hat{e}_x^0 &= \frac{n_{f_0}}{n_{f_0} \cdot t_{e_0}}, & \hat{e}_x^1 &= \frac{n_{f_1}}{n_{f_1} \cdot t_{e_1}}, & \hat{e}_x^2 &= \frac{n_{f_2}}{n_{f_2} \cdot t_{e_2}}. \end{aligned}$$

2. If $x \in \mathcal{X}_{\hat{e}}, e \in \Delta_1(T)$ and adjacent faces are f_0, f_1 then

$$\begin{aligned} e_x^0 &= t_e, & e_x^1 &= n_{f_0} \times t_e, & e_x^2 &= n_{f_1} \times t_e, \\ \hat{e}_x^0 &= t_e, & \hat{e}_x^1 &= \frac{n_{f_1}}{n_{f_1} \cdot (n_{f_0} \times t_e)}, & \hat{e}_x^2 &= \frac{n_{f_0}}{n_{f_0} \cdot (n_{f_1} \times t_e)}. \end{aligned}$$

3. If $x \in \mathcal{X}_{\hat{f}}, f \in \Delta_2(T)$, the first edge of f is e , then

$$\begin{aligned} e_x^0 &= t_e, & e_x^1 &= t_e \times n_f, & e_x^2 &= n_f, \\ \hat{e}_x^0 &= t_e, & \hat{e}_x^1 &= t_e \times n_f, & \hat{e}_x^2 &= n_f. \end{aligned}$$

4. If $x \in \mathcal{X}_{\hat{T}}$, then

$$\begin{aligned} e_x^0 &= (1, 0, 0), & e_x^1 &= (0, 1, 0), & e_x^2 &= (0, 0, 1), \\ \hat{e}_x^0 &= (1, 0, 0), & \hat{e}_x^1 &= (0, 1, 0), & \hat{e}_x^2 &= (0, 0, 1). \end{aligned}$$

6 Indexing management of degrees of freedom

In Section 2 we have already discussed the dictionary indexing rule for interpolation points in each element. In this section, we will address the global indexing rules for Lagrange interpolation points, ensuring that interpolation points on a sub-simplex, shared

by multiple elements, have a globally unique index. Given the one-to-one relationship between interpolation points and DoFs, this is equivalent to providing an indexing rule for global DoFs in the scalar Lagrange finite element space. Based on this, we will further discuss the indexing rules for DoFs in face and edge finite element spaces.

6.1 Lagrange finite element space

We begin by discussing the data structure of the tetrahedral mesh, denoted by $\mathcal{T}_h$. Let the numbers of nodes, edges, faces, and cells in $\mathcal{T}_h$ be represented as NN , NE , NF , and NC , respectively. We utilize two arrays to represent $\mathcal{T}_h$:

- `node` (shape: $(NN, 3)$): `node[i, j]` represents the j -th component of the Cartesian coordinate of the i -th vertex.
- `cell` (shape: $(NC, 4)$): `cell[i, j]` gives the global index of the j -th vertex of the i -th cell.

Given a tetrahedron denoted by $[0, 1, 2, 3]$, we define its local edges and faces as:

- `SEdge` = $[(0, 1), (0, 2), (0, 3), (1, 2), (1, 3), (2, 3)]$;
- `SFace` = $[(1, 2, 3), (0, 2, 3), (0, 1, 3), (0, 1, 2)]$;
- `OFace` = $[(1, 2, 3), (0, 3, 2), (0, 1, 3), (0, 2, 1)]$.

Here, we introduced two types of local faces. The prefix `S` implies sorting, and `O` indicates outer normal direction. Both `SFace[i, :]` and `OFace[i, :]` represent the face opposite to the i -th vertex but with varied ordering. The normal direction as determined by the ordering of the three vertices of `OFace` matches the outer normal direction of the tetrahedron. This ensures that the outer normal direction of a boundary face points outward from the mesh. Meanwhile, `SFace` aids in determining the global index of the interpolation points on the face. For an in-depth discourse on indexing, ordering, and orientation, we direct readers to `sc3` in *iFEM* [11].

Leveraging the unique algorithm for arrays, we can derive the following arrays from `cell`, `SEdge`, and `OFace`:

- `edge` (shape: $(NE, 2)$): `edge[i, j]` gives the global index of the j -th vertex of the i -th edge.
- `face` (shape: $(NF, 3)$): `face[i, j]` provides the global index of the j -th vertex of the i -th face.
- `cell2edge` (shape: $(NC, 6)$): `cell2edge[i, j]` indicates the global index of the j -th edge of the i -th cell.
- `cell2face` (shape: $(NC, 4)$): `cell2face[i, j]` signifies the global index of the j -th face of the i -th cell.




Figure 4: Local indexing (left) and global indexing (right) of DoFs for a face of tetrahedron, where the local vertex order is $[17, 0, 21]$ and global vertex order is $[0, 17, 21]$. Due to the different ordering of vertices in local and global representation of the face, the ordering of the local indexing and the global indexing is different.

Having constructed the edge and face arrays and linked cells to them, we next establish indexing rules for interpolation points on $\mathcal{T}_h$. Let k be the degree of the Lagrange finite element space. The number of interpolation points on each cell is

$$\text{ldof} = \dim \mathbb{P}_k(T) = \frac{(k+1)(k+2)(k+3)}{6},$$

and the total number of interpolation points on $\mathcal{T}_h$ is

$$\text{gdof} = \text{NN} + n_e^k \cdot \text{NE} + n_f^k \cdot \text{NF} + n_c^k \cdot \text{NC},$$

where

$$n_e^k = k-1, \quad n_f^k = \frac{(k-2)(k-1)}{2}, \quad n_c^k = \frac{(k-3)(k-2)(k-1)}{6},$$

are numbers of interpolation points inside edge, face, and cell, respectively. We need an index mapping from $[0:\text{ldof}-1]$ to $[0:\text{gdof}-1]$. See Fig. 4 for an illustration of the local index and the global index of interpolation points.

The tetrahedron's four vertices are ordered according to the right-hand rule, and the interpolation points adhere to the dictionary ordering map $R_3(\alpha)$. As Lagrange element is globally continuous, the indexing of interpolation points on the boundary ∂T should be global. Namely a unique index for points on vertices, edges, faces should be used and a mapping from the local index to the global index is needed.

We first set a global indexing rule for all interpolation points. We sort the index by vertices, edges, faces, and cells. For the interpolation points that coincide with the vertices, their global index are set as $0, 1, \dots, \text{NN}-1$. When $k > 1$, for the interpolation points

that inside edges, their global index are set as

$$\begin{matrix} 0 \\ 1 \\ \vdots \\ \text{NE}-1 \end{matrix} \begin{pmatrix} 0 \cdot n_e^k & \cdots & 1 \cdot n_e^k - 1 \\ 1 \cdot n_e^k & \cdots & 2 \cdot n_e^k - 1 \\ \vdots & & \vdots \\ (\text{NE}-1) \cdot n_e^k & \cdots & \text{NE} \cdot n_e^k - 1 \end{pmatrix} + \text{NN}.$$

Here recall that $n_e^k = k-1$ is the number of interior interpolation points on an edge. When $k > 2$, for the interpolation points that inside each face, their global index are set as

$$\begin{matrix} 0 \\ 1 \\ \vdots \\ \text{NF}-1 \end{matrix} \begin{pmatrix} 0 \cdot n_f^k & \cdots & 1 \cdot n_f^k - 1 \\ 1 \cdot n_f^k & \cdots & 2 \cdot n_f^k - 1 \\ \vdots & & \vdots \\ (\text{NF}-1) \cdot n_f^k & \cdots & \text{NF} \cdot n_f^k - 1 \end{pmatrix} + \text{NN} + \text{NE} \cdot n_e^k,$$

where $n_f^k = (k-2)(k-1)/2$ is the number of interior interpolation points on f . When $k > 3$, the global index of the interpolation points that inside each cell can be set in a similar way.

Then we use the two-dimensional array named `cell2ipoint` of shape (NC, ldof) for the index map. On the j -th interpolation point of the i -th cell, we aim to determine its unique global index and store it in `cell2ipoint[i, j]`.

For vertices and cell interpolation points, the mapping is straightforward by the global indexing rule. Indeed `cell` is the mapping of the local index of a vertex to its global index. However, complications arise when the interpolation point is located within an edge or face due to non-unique representations of an edge and a face.

We use the more complicated face interpolation points as a typical example to illustrate the situation. Consider, for instance, the scenario where the j -th interpolation point lies within the 0th local face F_0 of the i -th cell. Let $\alpha = \mathbf{m} = [\mathbf{m}_0, \mathbf{m}_1, \mathbf{m}_2, \mathbf{m}_3]$ be its lattice point. Given that F_0 is opposite to vertex 0, we deduce that $\lambda_0|_{F_0} = 0$, which implies $\mathbf{m}_0$ is 0. The remaining components of $\mathbf{m}$ are non-zero, ensuring that the point is interior to F_0 .

Two representations for the face with global index `cell2face[i, 0]` are subsequently acquired:

- `LFace = cell[i, SFace[0, :]]` (local representation)
- `GFace = face[cell2face[i, 0], :]` (global representation)

Although `LFace` and `GFace` comprise identical vertex numbers, their ordering differs. For example, `LFace = [5 6 10]` while `GFace = [10 6 5]`.

The array $\mathbf{m} = [\mathbf{m}_1, \mathbf{m}_2, \mathbf{m}_3]$ has a one-to-one correspondence with the vertices of `LFace`. To match this array with the vertices of `GFace`, a reordering based on argument sorting is performed:

```
1 i0 = argsort(argsort(GFace));
```

```

2 i1 = argsort(LFace);
3 i2 = i1[i0];
4 m = m[i2];

```

In the example of $\text{LFace} = [5 \ 6 \ 10]$ while $\text{GFace} = [10 \ 6 \ 5]$, the input $\mathbf{m} = [m_1, m_2, m_3]$ will be reordered to $\mathbf{m} = [m_3, m_2, m_1]$.

From the reordered $\mathbf{m} = [m_1, m_2, m_3]$, the local index ℓ of the j -th interpolation point on the global face $\mathbf{f} = \text{cell2face}[i, 0]$ can be deduced:

$$\begin{aligned}\ell &= \frac{(m_2 - 1 + m_3 - 1)(m_2 - 1 + m_3 - 1 + 1)}{2} + m_3 - 1 \\ &= \frac{(m_2 + m_3 - 2)(m_2 + m_3 - 1)}{2} + m_3 - 1.\end{aligned}$$

It's worth noting that the index of interpolation points solely within the face needs consideration. Finally, the global index for the j -th interpolation point within the 0th local face of the i -th cell is:

$$\text{cell2ipoint}[i \ j] = \text{NN} + n_e^k \cdot \text{NE} + n_f^k \cdot \mathbf{f} + \ell.$$

Here, we provide a specific example. Consider a 5th-degree Lagrange finite element space on the c -th tetrahedron in a mesh depicted in Fig. 4. The vertices of this tetrahedron are $[5, 17, 0, 21]$, and its 0th face is denoted as $\mathbf{f}$, with vertices $[0, 17, 21]$. Assuming that

$$\text{NN} + n_e^k \cdot \text{NE} + n_f^k \cdot \mathbf{f} = 1240.$$

For the 39th local DoF on tetrahedron, its corresponding multi-index is $[0, 3, 1, 1]$ on cell and $[3, 1, 1]$ on face. Since the local face is $[17, 0, 21]$ and the global face is $[0, 17, 21]$, then $\mathbf{m} = [1, 3, 1]$ and $\ell = 3$ thus

$$\text{cell2ipoint}[c, 39] = 1243.$$

Similar, for the 43th local DoF on tetrahedron, $\mathbf{m} = [1, 2, 2]$ and $\ell = 4$ thus

$$\text{cell2ipoint}[c, 43] = 1244.$$

Fig. 4 shows the correspondence between the local and the global indexing of DoFs for the cell.

In conclusion, we have elucidated the construction of global indexing for interpolation points inside cell faces. This method can be generalized for edges and, more broadly, for interior interpolation points of the low-dimensional sub-simplex of an n -dimensional simplex. Please note that for scalar Lagrange finite element spaces, the `cell2ipoint` array is the mapping array from local DoFs to global DoFs.

6.2 Face and edge finite element spaces

First, we want to emphasize that the management of DoFs is to manage the continuity of finite element space.

The face element space and edge element space are vector finite element spaces, which define DoFs of vector type by defining a vector frame on each interpolation point. At the same time, they define their vector basis functions by combining the Lagrange basis function and the dual frame of the DoFs.

Alternatively, we can say each DoF in the face element or edge element corresponds to a unique interpolation point p and a unique vector e , and each basis function also corresponds to a unique Lagrange basis function which is defined on p and a vector $\hat{e}$ which is the dual to e .

The management of DoFs is essentially a counting problem. First of all, we need to set global and local indexing rules for all DoFs. We can globally divide the DoFs into shared and unshared among simplexes. The DoFs shared among simplexes can be further divided into on-edge and on-face according to the dimension of the sub-simplex where the DoFs locate. First count the shared DoFs on each edge according to the order of the edges, then count the shared DoFs on each face according to the order of the faces, and finally count the unshared DoFs in the cell. On each edge or face, the DoFs' order can follow the order of the interpolation points. Note that, for the face element space and edge element space there are no DoFs shared on nodes. And for the 3D face element space there are no DoFs shared on edges. So the global numbering rule is similar with the Lagrange interpolation points.

According to the global indexing rule, we also can get a array named `dof2vector` with shape (gdof, GD) , where `gdof` is the number of global DoFs and `GD` represent geometry dimensions. And `dof2vector[i, :]` store the vector of the i -th DoF.

Next we need to set a local indexing rules and generate an array `cell2dof` with shape (NC, ldof) , where `ldof` is the number of local DoFs on each cell. Note that each DoF was determined by an interpolation point and a vector. And for each interpolation point, there is a frame (including `GD` vectors) on it. Given a DoF on the i -th cell, denote the local index of its interpolation point as p , and the local index of its vector in the frame denote as q , then one can set a unique local index number j by p and q , for example $j = n \cdot q + p$, where n is the number of interpolation points in the i -th cell. Furthermore, we can compute the `cell2dof[i, j]` by the global index `cell2ipoint[i, p]`, the sub-simplex that the interpolation point locate, and the global indexing rule.

Remark 6.1. Note that the local and global number rules mentioned above are not unique. Furthermore, with the array `cell2dof`, the implementation of these higher-order finite element methods mentioned in this paper is not fundamentally different from the conventional finite element in terms of matrix vector assembly and boundary condition handling.

7 Numerical examples

In this section, we numerically verify the 3-dimensional face element basis and edge element basis using two test problems over the domain $\Omega = (0,1)^3$ partitioned into a structured tetrahedron mesh $\mathcal{T}_h$. All the algorithms and numerical examples are implemented based on the FEALPy package [21].

7.1 High order elements for Poisson equation in the mixed formulation

Consider the Poisson problem:

$$\begin{cases} \mathbf{u} + \nabla p = 0 & \text{in } \Omega, \\ \operatorname{div} \mathbf{u} = f & \text{in } \Omega, \\ p = g & \text{on } \partial\Omega. \end{cases}$$

The variational problem is : find $\mathbf{u} \in H(\operatorname{div}, \Omega)$, $p \in L^2(\Omega)$ to satisfy:

$$\begin{aligned} \int_{\Omega} \mathbf{u} \cdot \mathbf{v} \, dx - \int_{\Omega} p \operatorname{div} \mathbf{v} \, dx &= - \int_{\partial\Omega} g(\mathbf{v} \cdot \mathbf{n}) \, ds, \quad \forall \mathbf{v} \in H(\operatorname{div}, \Omega), \\ - \int_{\Omega} (\operatorname{div} \mathbf{u}) q \, dx &= - \int_{\Omega} f q \, dx, \quad \forall q \in L^2(\Omega). \end{aligned}$$

Let $V_{h,k}^{\operatorname{div}}$ be the BDM space with degree k on $\mathcal{T}_h$ and piecewise polynomial space of degree $k-1$ on $\mathcal{T}_h$ by V_{k-1} . The corresponding finite element method is: find $\mathbf{u}_h \in V_{h,k}^{\operatorname{div}}$, $p_h \in V_{k-1}$, such that

$$\int_{\Omega} \mathbf{u}_h \cdot \mathbf{v}_h \, dx - \int_{\Omega} p \operatorname{div} \mathbf{v}_h \, dx = - \int_{\partial\Omega} g(\mathbf{v}_h \cdot \mathbf{n}) \, ds, \quad \forall \mathbf{v}_h \in V_{h,k}^{\operatorname{div}}, \quad (7.1)$$

$$- \int_{\Omega} (\operatorname{div} \mathbf{u}_h) q_h \, dx = - \int_{\Omega} f q_h \, dx, \quad \forall q_h \in V_{k-1}. \quad (7.2)$$

To test the convergence order of BDM space, we set

$$\begin{aligned} \mathbf{u} &= \begin{pmatrix} -\pi \sin(\pi x) \cos(\pi y) \cos(\pi z) \\ -\pi \cos(\pi x) \sin(\pi y) \cos(\pi z) \\ -\pi \cos(\pi x) \cos(\pi y) \sin(\pi z) \end{pmatrix}, \\ p &= \cos(\pi x) \cos(\pi y) \cos(\pi z), \quad f = 3\pi^2 \cos(\pi x) \cos(\pi y) \cos(\pi z). \end{aligned}$$

The numerical results are shown in Fig. 5 and it is clear that

$$\|\mathbf{u} - \mathbf{u}_h\|_0 \leq Ch^{k+1}, \quad \|p - p_h\|_0 \leq Ch^k.$$




Figure 5: Errors $\|u - u_h\|_0$ and $\|p - p_h\|_0$ of finite element method (7.1) and (7.2) on uniformly refined mesh with $k=2,3,4$.

7.2 High order elements for Maxwell equations

Consider the time harmonic Maxwell equation

$$\begin{cases} \nabla \times (\mu^{-1} \nabla \times E) - \omega^2 \tilde{\epsilon} E = J & \text{in } \Omega, \\ n \times E = 0 & \text{on } \partial\Omega, \end{cases}$$

where μ is the relative permeability, $\tilde{\epsilon}$ is the relative permittivity, and ω is the wavenumber. The variational problem is: find $E \in H_0(\text{curl}, \Omega)$ to satisfy

$$\int_{\Omega} \mu^{-1} (\nabla \times E) \cdot (\nabla \times v) \, dx - \int_{\Omega} \omega^2 \tilde{\epsilon} E \cdot v \, dx = \int_{\Omega} J \cdot v \, dx, \quad \forall v \in H_0(\text{curl}, \Omega).$$

Let $\mathring{V}_{h,k}^{\text{curl}} = V_{h,k}^{\text{curl}} \cap H_0(\text{curl}, \Omega)$, where $V_{h,k}^{\text{curl}}$ is the edge element space defined in Section 5.5. The corresponding finite element method is: find $E_h \in \mathring{V}_{h,k}^{\text{curl}}$ s.t.

$$\int_{\Omega} \mu^{-1} (\nabla \times E_h) \cdot (\nabla \times v_h) \, dx - \int_{\Omega} \omega^2 \tilde{\epsilon} E_h \cdot v_h \, dx = \int_{\Omega} J \cdot v_h \, dx, \quad \forall v_h \in \mathring{V}_{h,k}^{\text{curl}}. \quad (7.3)$$

To test the convergence rate of second-kind Nédélec space, we choose

$$\begin{aligned} \omega = \tilde{\epsilon} = \mu = 1, & \quad E = (f, \sin(x)f, \sin(y)f), \\ f = (x^2 - x)(y^2 - y)(z^2 - z), & \quad J = \nabla \times \nabla \times E - E. \end{aligned}$$

The numerical results are shown in Fig. 6 and it is clear that

$$\|E - E_h\|_0 \leq Ch^{k+1}, \quad \|\nabla \times (E - E_h)\|_0 \leq Ch^k.$$




Figure 6: Errors $\|E - E_h\|_0$ and $\|\nabla \times (E - E_h)\|_0$ of finite element method (7.3) on uniformly refined mesh with $k = 2, 3, 4$.

Acknowledgments

The first and fourth authors were supported by the National Natural Science Foundation of China (NSFC) (Grant Nos. 12371410, 12261131501), and the construction of innovative provinces in Hunan Province (Grant No. 2021GK1010). The second author was supported by NSF DMS-2012465 and DMS-2309785. The third author was supported by the National Natural Science Foundation of China (NSFC) (Grant No. 12171300), and the Natural Science Foundation of Shanghai (Grant No. 21ZR1480500).

References

- [1] M. Ainsworth, G. Andriamaro, and O. Davydov, A Bernstein–Bézier basis for arbitrary order Raviart–Thomas finite elements, *Constr. Approx.*, 41 (2015), 1-22.
- [2] M. Ainsworth and J. Coyle, Hierarchic finite element bases on unstructured tetrahedral meshes, *Internat. J. Numer. Methods Engrg.*, 58 (2003), 2103-2130.
- [3] M. Ainsworth and G. Fu, Bernstein–Bézier bases for tetrahedral finite elements, *Comput. Methods Appl. Mech. Engrg.*, 340 (2018), 178-201.
- [4] M. Alnæs, J. Blechta, J. Hake, A. Johansson, B. Kehlet, A. Logg, C. Richardson, J. Ring, M. E. Rognes, and G. N. Wells., The FEniCS project version 1.5, *Archive of Numerical Software*, 3 (2015), 9-23.
- [5] R. Anderson, J. Andrej, A. Barker, J. Bramwell, J.-S. Camier, J. Cervený, V. Dobrev, Y. Doudouit, A. Fisher, T. Kolev, et al, MFEM: A modular finite element methods library, *Comput. Math. Appl.*, 81 (2021), 42-74.
- [6] D. N. Arnold, R. S. Falk, and R. Winther, Finite element exterior calculus, homological techniques, and applications, *Acta Numer.*, 15 (2006), 1-155.

- [7] D. N. Arnold, R. S. Falk, and R. Winther, Geometric decompositions and local bases for spaces of finite element differential forms, *Comput. Methods Appl. Mech. Engrg.*, 198 (2009), 1660-1672.
- [8] F. Brezzi, J. Douglas, Jr., R. Durán, and M. Fortin, Mixed finite elements for second order elliptic problems in three variables, *Numer. Math.*, 51 (1987), 237-250.
- [9] F. Brezzi, J. Douglas, Jr., and L. D. Marini, Two families of mixed finite elements for second order elliptic problems, *Numer. Math.*, 47 (1985), 217-235.
- [10] D. A. Castro, P. R. B. Devloo, A. M. Farias, S. M. Gomes, D. de Siqueira, and O. Durán, Three dimensional hierarchical mixed finite element approximations with enhanced primal variable accuracy, *Comput. Methods Appl. Mech. Engrg.*, 306 (2016), 479-502.
- [11] L. Chen, *iFEM: An integrated finite element methods package in MATLAB*, Technical report, University of California at Irvine, <https://github.com/lyc102/ifem>, 2009.
- [12] L. Chen and X. Huang, *H(div)-conforming finite element tensors*, arXiv preprint arXiv:2112.14351v1, 2021.
- [13] L. Chen and X. Huang, Finite element de Rham and Stokes complexes in three dimensions, *Math. Comp.*, 93 (2024), 55-110.
- [14] S. H. Christiansen, J. Hu, and K. Hu, Nodal finite element de Rham complexes, *Numer. Math.*, 139 (2018), 411-446.
- [15] D. De Siqueira, P. R. B. Devloo, and S. M. Gomes, Hierarchical high order finite element approximation spaces for $H(\text{div})$ and $H(\text{curl})$, in G. Kreiss, P. Lötstedt, A. Målqvist, and M. Neytcheva, editors, *Numerical Mathematics and Advanced Applications 2009*, pages 269-276, Springer Berlin, Heidelberg, 2010.
- [16] D. De Siqueira, P. R. B. Devloo, and S. M. Gomes, A new procedure for the construction of hierarchical high order $H\text{div}$ and $H\text{curl}$ finite element spaces, *J. Comput. Appl. Math.*, 240 (2013), 204-214.
- [17] V. Ervin, Computational bases for RT_k and BDM_k on triangles, *Comput. Math. Appl.*, 64 (2012), 2765-2774.
- [18] J.-C. Nédélec, A new family of mixed finite elements in $\mathbf{R}^3$, *Numer. Math.*, 50 (1986), 57-81.
- [19] R. Nicolaides, On a class of finite elements generated by Lagrange interpolation, *SIAM J. Numer. Anal.*, 9 (1972), 435-445.
- [20] M. E. Rognes, R. C. Kirby, and A. Logg, Efficient assembly of $H(\text{div})$ and $H(\text{curl})$ conforming finite elements, *SIAM J. Sci. Comput.*, 31 (2010), 4130-4151.
- [21] H. Wei, C. Chen, and Y. Huang, *FEALPy: Finite Element Analysis Library in Python*, <https://github.com/weihuayi/fealpy>, Xiangtan University, 2017-2024.
- [22] J. Xin and W. Cai, A well-conditioned hierarchical basis for triangular $H(\text{curl})$ -conforming elements, *Commun. Comput. Phys.*, 9 (2011), 780-806.
- [23] J. Xin, W. Cai, and N. Guo, On the construction of well-conditioned hierarchical bases for (div) -conforming $\mathbb{R}^n$ simplicial elements, *Commun. Comput. Phys.*, 14 (2013), 621-638.
- [24] J. Xin, N. Guo, and W. Cai, On the construction of well-conditioned hierarchical bases for tetrahedral $H(\text{curl})$ -conforming Nédélec element, *J. Comput. Math.*, 29 (2011), 526-542.