



Stable phase retrieval in function spaces

D. Freeman¹ · T. Oikhberg² · B. Pineau³ · M. A. Taylor⁴

Received: 13 February 2023 / Revised: 15 October 2023 / Accepted: 18 October 2023

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023

Abstract

Let (Ω, Σ, μ) be a measure space, and $1 \leq p \leq \infty$. A subspace $E \subseteq L_p(\mu)$ is said to do *stable phase retrieval (SPR)* if there exists a constant $C \geq 1$ such that for any $f, g \in E$ we have

$$\inf_{|\lambda|=1} \|f - \lambda g\| \leq C \||f| - |g|\|. \quad (0.1)$$

In this case, if $|f|$ is known, then f is uniquely determined up to an unavoidable global phase factor λ ; moreover, the phase recovery map is C -Lipschitz. Phase retrieval appears in several applied circumstances, ranging from crystallography to quantum mechanics. In this article, we construct various subspaces doing stable phase retrieval, and make connections with $\Lambda(p)$ -set theory. Moreover, we set the foundations for an analysis of stable phase retrieval in general function spaces. This, in particular, allows us to show that Hölder stable phase retrieval implies stable phase retrieval, improving the stability bounds in a recent article of M. Christ and the third and fourth authors.

D. Freeman was supported by the Simons Foundation award 706481 and the NSF award 2154931.
T. Oikhberg was supported by the NSF award 1912897.

✉ T. Oikhberg
oikhberg@illinois.edu

D. Freeman
daniel.freeman@slu.edu

B. Pineau
bpineau@berkeley.edu

M. A. Taylor
mitchell.taylor@math.ethz.ch

¹ Department of Mathematics and Statistics, St Louis University, St Louis, MO, USA

² Department of Mathematics, University of Illinois, Urbana, IL 61801, USA

³ Department of Mathematics, University of California at Berkeley, Berkeley, USA

⁴ Department of Mathematics, ETH Zürich, Zürich, Switzerland

We also characterize those compact Hausdorff spaces K such that $C(K)$ contains an infinite dimensional SPR subspace.

Mathematics Subject Classification 46B42 · 43A46 · 42C15

Contents

1	Introduction	· · · · ·
1.1	Motivation and applications	· · · · ·
1.2	An overview of the results	· · · · ·
2	Preliminaries	· · · · ·
2.1	The Kadec–Pelczynski dichotomy	· · · · ·
2.2	Complex Banach lattices	· · · · ·
3	General theory	· · · · ·
3.1	Connections with almost disjoint pairs and sequences	· · · · ·
3.2	Hölder stable phase retrieval and witnessing failure of SPR on orthogonal vectors	· · · · ·
4	Examples	· · · · ·
4.1	Building SPR subspaces via isometric theory	· · · · ·
4.2	Existence of SPR embeddings into sequence spaces	· · · · ·
4.3	Explicit constructions of SPR subspaces using random variables	· · · · ·
4.4	Stability of SPR subspaces under ultraproducts and small perturbations	· · · · ·
5	SPR in L_p -spaces	· · · · ·
6	$C(K)$ -spaces with SPR subspaces	· · · · ·
7	Open problems	· · · · ·
	References	· · · · ·

1 Introduction

There are many situations in mathematics, science, and engineering where the goal is to recover some vector f from $|Tf|$, where T is a linear transformation into a function space. Note that if $|\lambda| = 1$ then it is impossible to distinguish f and λf in this way. The linear transformation T is said to do phase retrieval if this ambiguity is the only obstruction to recovering f . That is, given a vector space H and function space X , a linear operator $T : H \rightarrow X$ does *phase retrieval* if whenever $f, g \in H$ satisfy $|Tf| = |Tg|$ then $f = \lambda g$ for some scalar λ with $|\lambda| = 1$. Phase retrieval naturally arises in situations where one is only able to obtain the magnitude of linear measurements, and not the phase. Notable examples in physics and engineering which require phase retrieval include X-ray crystallography, electron microscopy, quantum state tomography, and cepstrum analysis in speech recognition. The study of phase retrieval in mathematical physics dates back to at least 1933 when in his seminal work *Die allgemeinen Prinzipien der Wellenmechanik* [65] W. Pauli asked whether a wave function is uniquely determined by the probability densities of position and momentum. In other words, Pauli asked whether $|f|$ and $|\widehat{f}|$ determine $f \in L_2(\mathbb{R})$ up to multiplication by a unimodular scalar. The mathematics of phase retrieval has since grown to be an important and well-studied topic in applied harmonic analysis.

As any application of phase retrieval would involve error, it is of fundamental importance that the recovery of f from $|Tf|$ not only be possible, but also be stable.

We say that T does *stable phase retrieval* if the recovery (up to a unimodular scalar) of f from $|Tf|$ is Lipschitz. If X is finite dimensional, then T does phase retrieval if and only if it does stable phase retrieval [12, 22]. However, if X is infinite dimensional and T is the analysis operator of a frame or a continuous frame, then T cannot do stable phase retrieval [2, 23]. Here, a collection of vectors $(\psi_t)_{t \in \Omega} \subseteq H$ is a *continuous frame* of a Hilbert space H over a measure space (Ω, Σ, μ) if the map $f \mapsto (\langle f, \psi_t \rangle)_{t \in \Omega}$ is an embedding of H into $L_2(\mu)$. One of the main goals of this paper is to use the theory of subspaces of Banach lattices to present a unifying framework for stable phase retrieval which encompasses the previously studied cases and allows for stable phase retrieval in infinite dimensions.

Let $X = L_p(\mu)$, or, more generally, a Banach lattice. Let $E \subseteq X$ be a subspace. We say that E does *phase retrieval* as a subspace of X if whenever $|f| = |g|$ for some $f, g \in E$ we have that $f = \lambda g$ for some scalar λ with $|\lambda| = 1$. Given a constant $C > 0$, we say that E does C -stable phase retrieval as a subspace of X if

$$\inf_{|\lambda|=1} \|f - \lambda g\| \leq C \| |f| - |g| \| \quad \text{for all } f, g \in E. \quad (1.1)$$

We may define an equivalence relation \sim on E by $f \sim g$ if $f = \lambda g$ for some scalar λ with $|\lambda| = 1$. Then, E does phase retrieval as a subspace of X if and only if the map $f \mapsto |f|$ from E/\sim to X is injective. Furthermore, E does C -stable phase retrieval as a subspace of X if and only if the map $f \mapsto |f|$ from E/\sim to X is injective and the inverse is C -Lipschitz. By introducing stable phase retrieval into the setting of Banach lattices, we are able to apply established methods from the subject to attack problems in phase retrieval, and conversely we hope that the ideas and questions in phase retrieval will inspire a new avenue of research in the theory of Banach lattices. Before starting the meat of the paper, we present some additional motivation, give an outline of our major results, and state some of the important ideas and theorems from Banach lattices which we will be applying. We conclude the paper by listing many open questions concerning stable phase retrieval in this new setting.

1.1 Motivation and applications

The inequality (1.1) arises in various circumstances. For instance, in crystallography and optics, one seeks to recover an unknown function $F \in L_2(\mathbb{R}^d)$ from the absolute value of its Fourier transform \widehat{F} . If one also seeks stability, this translates into an inequality of the form

$$\inf_{|\lambda|=1} \|F - \lambda G\|_{L_2} \leq C \| |\widehat{F}| - |\widehat{G}| \|_{L_2}, \quad (1.2)$$

which one would want to be valid for F, G in a subspace $E \subseteq L_2(\mathbb{R}^d)$ which incorporates the additional constraints F, G are known to satisfy. Using Plancherel's theorem to write $\|F - \lambda G\|_{L_2} = \|\widehat{F} - \lambda \widehat{G}\|_{L_2}$, one sees that the inequality (1.2) reduces to (1.1), up to passing to Fourier space and making the change of notation $f = \widehat{F}$ and $g = \widehat{G}$. We refer the reader to the surveys [42, 47] and references therein for a fur-

ther explanation of the importance of phase retrieval in optics, crystallography, and other areas. In particular, these articles explains why, in practice, physical experiments are often able to measure the magnitude of the Fourier transform, but are unable to measure the phase.

A second scenario where phase retrieval appears is quantum mechanics. In this case, one wants to identify situations where $|f|$ and $|\widehat{f}|$ determine $f \in L_2(\mathbb{R})$ uniquely. As already mentioned, Pauli asked whether this could true for all $f \in L_2(\mathbb{R})$. However, a counterexample to this conjecture was given in 1944: There exists $f, g \in L_2(\mathbb{R})$ such that $|f| = |g|$ and $|\widehat{f}| = |\widehat{g}|$ but f is not a multiple of g . This leads to the natural question of whether one can build “large” subspaces $G \subseteq L_2(\mathbb{R})$ for which $|f|$ and $|\widehat{f}|$ determine $f \in G \subseteq L_2(\mathbb{R})$ uniquely. By passing to the phase space $L_2(\mathbb{R}) \times L_2(\mathbb{R})$, we see that G has the above property if and only if $E := \{(f, \widehat{f}) : f \in G\}$ does phase retrieval as a subspace of $L_2(\mathbb{R}) \times L_2(\mathbb{R})$, i.e., knowing $h, k \in E$ and $|h| = |k|$ implies h is a unimodular multiple of k . This also naturally leads to the question of stability of Pauli phase retrieval, by requiring (1.1) hold on E . In this case, using Plancherel’s theorem to return to G , (1.1) on E translates into the inequality

$$\inf_{|\lambda|=1} \|f - \lambda g\|_{L_2} \leq C (\| |f| - |g| \|_{L_2} + \| |\widehat{f}| - |\widehat{g}| \|_{L_2}) \text{ for } f, g \in G. \quad (1.3)$$

For a non-exhaustive collection of results on Pauli phase retrieval and its generalizations, see [8, 42, 49, 50] and references therein. To our knowledge, the question of stability in the Pauli Problem is essentially unexplored. However, the results presented here in conjunction with [28] give a relatively large class of subspaces of $L_2(\mathbb{R}^d)$ satisfying (1.3).

Finally, we mention that phase retrieval has grown to become an exciting and important topic of research in frame theory [12–14, 18, 27, 34, 42]. A frame for a separable Hilbert space H is a sequence of vectors $(\phi_j)_{j \in J}$ in H such that there exists uniform bounds $A, B > 0$ so that

$$A \|f\|^2 \leq \sum_{j \in J} |\langle f, \phi_j \rangle|^2 \leq B \|f\|^2 \quad \text{for all } f \in H. \quad (1.4)$$

The *analysis operator* of a frame $(\phi_j)_{j \in J}$ of H is the map $\Theta : H \rightarrow \ell_2(J)$ given by $\Theta(f) = (\langle f, \phi_j \rangle)_{j \in J}$. Note that the uniform upper bound B in the frame inequality (1.4) guarantees that $\Theta : H \rightarrow \ell_2(J)$ is bounded, and the uniform lower bound A gives that Θ is an embedding of H into $\ell_2(J)$. Given a frame $(\phi_j)_{j \in J}$ of H , the *canonical dual frame* $(\widetilde{\phi}_j)_{j \in J}$ is defined by $\widetilde{\phi}_j = (\Theta^* \Theta)^{-1} \phi_j$ for all $j \in J$ and satisfies

$$f = \sum_{j \in J} \langle f, \widetilde{\phi}_j \rangle \phi_j = \sum_{j \in J} \langle f, \phi_j \rangle \widetilde{\phi}_j \quad \text{for all } f \in H. \quad (1.5)$$

Frames have many applications and play a fundamental role in signal processing and applied harmonic analysis. One important reason for this is that the analysis operator Θ is an embedding of H into $\ell_2(J)$, which allows for the application of filters, thresholding, and other signal processing techniques. Another reason is that

(1.5) gives a linear, stable, and unconditional reconstruction formula for a vector in terms of the frame coefficients.

A frame $(\phi_j)_{j \in J}$ is said to do *phase retrieval* if whenever $f, g \in H$ and $(|\langle f, \phi_j \rangle|)_{j \in J} = (|\langle g, \phi_j \rangle|)_{j \in J}$, there exists a unimodular scalar λ such that $f = \lambda g$. A frame is said to do *stable phase retrieval* if there exists a constant $C \geq 1$ such that for all $f, g \in H$,

$$\inf_{|\lambda|=1} \|f - \lambda g\|_H \leq C \|\Theta(f) - \Theta(g)\|_{\ell_2(J)}. \quad (1.6)$$

Using the fact that the analysis operator $\Theta : H \rightarrow \ell_2(J)$ is an embedding, we see that a frame does stable phase retrieval if and only if the subspace $\Theta(H) \subseteq \ell_2(J)$ does stable phase retrieval in the sense of (1.1). In finite dimensions, phase retrieval for frames is automatically stable. However, in infinite dimensions, it is necessarily unstable. As we will see, this is due to the fact that the ambient Hilbert lattice $\ell_2(J)$ is atomic, whereas the construction of SPR subspaces from [24] is done in the non-atomic lattice $L_2(\mathbb{R})$. For further investigations on the instability of phase retrieval for frames—including generalizations to continuous frames and frames in Banach spaces—see [2, 23].

As mentioned previously, phase retrieval problems arise in applications when considering an operator $T : H \rightarrow X$, which embeds a Hilbert space H into a function space X . In particular, the inequality (1.2) arises by taking T to be the Fourier transform, and (1.6) arises by taking T to be the analysis operator of a frame. A particularly important choice for T is the Gabor transform (see [3, 44] for recent advances in Gabor phase retrieval). As should now be evident, the question of stability for each of these phase retrieval problems can be translated into a special case of (1.1), by taking $E := T(H)$.

1.2 An overview of the results

The examples from Sect. 1.1 show that the inequality (1.1) unifies various phase retrieval problems. However, as mentioned previously, phase retrieval for frames is unstable in infinite dimensions, and it was only recently that the first examples of infinite dimensional SPR subspaces of real $L_2(\mu)$ spaces were constructed [24]. The purpose of this article is twofold. First, we construct numerous examples of subspaces of $L_p(\mu)$ doing stable phase retrieval. For this, we use various isometric Banach space techniques, modifications of the “almost disjointness” methods in classical Banach lattice theory, random constructions, and analogues of some constructions from harmonic analysis. Secondly, we prove several structural results about SPR subspaces of $L_p(\mu)$, and even general Banach lattices. Notably, both the characterization of real SPR in terms of almost disjoint pairs (Theorem 3.4), as well as the equivalence of SPR and its Hölder analogue (Corollary 3.12) hold for all Banach lattices. Our results also extend those in the recent article [28], which uses orthogonality and combinatorial arguments akin to Rudin’s work [68] on $\Lambda(p)$ -sets to produce examples of subspaces of (real or complex) $L_p(\mu)$ doing Hölder stable phase retrieval.

We now briefly overview the paper. In Sect. 2, we recall some basic terminology and results from Banach lattice theory in order to make the paper accessible to a

wider audience. Most notably, in Sect. 2.1 we collect basic facts related to the Kadec–Pelczynski dichotomy. Such results give structural information about closed subspaces of Banach lattices that are *dispersed*, i.e., that do not contain normalized almost disjoint sequences. As we will show in Theorem 3.4, a subspace of a (real) Banach lattice does stable phase retrieval if and only if it does not contain normalized almost disjoint *pairs*. In Theorem 2.1, we collect various facts about dispersed subspaces; finding SPR analogues of these results will occupy much of the paper. In particular, although SPR is much stronger than being dispersed, in Theorem 5.1 we will show that every closed infinite dimensional dispersed subspace of an order continuous Banach lattice contains a further closed infinite dimensional subspace doing SPR. The preliminary section finishes with Sect. 2.2, which recalls basic facts about complex Banach lattices.

Section 3 collects various results on stable phase retrieval that hold for general Banach lattices. In particular, in Sect. 3.1 we make the aforementioned connection between stable phase retrieval and almost disjoint pairs (see Theorem 3.4). In Sect. 3.2, we show that if the phase recovery map is Hölder continuous on the ball, then it is Lipschitz continuous on the whole space (Corollary 3.12). This follows from Theorem 3.10, which shows that failure of stable phase retrieval can be witnessed by “well-separated” vectors. The equivalence between stable phase retrieval and Hölder stable phase retrieval allows us to improve some results from [28], yielding the first examples of infinite dimensional closed subspaces of complex $L_2(\mu)$ doing stable phase retrieval.

In Sect. 4, we build infinite dimensional SPR subspaces using a variety of different techniques. In particular, we prove in Corollary 4.8 an analogue of statement (iii) of Theorem 2.1; namely, that for every dispersed subspace $E \subseteq L_p[0, 1]$ ($1 \leq p \leq \infty$), we can build a closed subspace $E' \subseteq L_p[0, 1]$ isomorphic to E , and doing stable phase retrieval. Moreover, for $p < 2$ and $q \in (p, 2]$, we will show that any closed subspace of $L_p(\mu)$ isometric to $L_q(\mu)$ does SPR in $L_p(\mu)$, see Proposition 4.1. Regarding sequence spaces, in Sect. 4.2 we show that ℓ_∞ embeds into itself in an SPR way, while no infinite dimensional subspace of ℓ_p does SPR when $1 \leq p < \infty$. Sect. 4.3 constructs SPR subspaces of rearrangement invariant spaces using random variables. This, in particular, tells us that subspaces spanned by iid Gaussian and q -stable random variables will do SPR in a variety of spaces, including all L_p -spaces in which they can be found. Here, we recall that “iid” stands for “independent and identically distributed”, and by a sequence of iid Gaussians we mean a sequence of independent random variables all of which have standard normal distribution (see [24] for more precise definitions as well as other applications of iid vectors to phase retrieval). Finally, Sect. 4.4 provides some basic stability properties of SPR subspaces.

Section 5 contains a study of the structure of SPR subspaces of $L_p(\mu)$, for a finite measure μ . We begin with the aforementioned Theorem 5.1, which is applicable for general order continuous Banach lattices, but for which much of the proof occurs in $L_1(\mu)$. Indeed, we will show that the generalization to order continuous Banach lattices follows from the result in $L_1(\mu)$ by arguing via the Kadec–Pelczynski dichotomy.

Note that from the classical results in Theorem 2.1 (a)-(d) it follows that if E is dispersed in $L_p(\mu)$ and $1 \leq q < p < \infty$, then E may be viewed as a closed subspace of $L_q(\mu)$, and it is dispersed in $L_q(\mu)$. In Theorem 5.3 we show that if $2 \leq p < \infty$, there are closed subspaces $E \subseteq L_p(\mu)$ which do SPR (and hence are dispersed in

$L_q(\mu)$ for all $1 \leq q \leq p$), but fail to do SPR when viewed as a closed subspace of $L_q(\mu)$ for all $1 \leq q < p$. However, by Theorem 5.6, if $p < 2$, then any SPR subspace $E \subseteq L_p(\mu)$ also does SPR when viewed as a closed subspace of $L_q(\mu)$ for any $1 \leq q \leq p$. Whether there is an SPR analogue of statement (v) of Theorem 2.1 remains an open problem.

Section 6 is devoted to the study of infinite dimensional SPR subspaces of $C(K)$. The main result is Theorem 6.1 which states that for a compact Hausdorff space K , the space $C(K)$ of continuous functions over K admits a (closed) infinite dimensional SPR subspace if and only if the Cantor–Bendixson derivative K' of K is infinite. The paper finishes with Sect. 7, which discusses various avenues for further research.

2 Preliminaries

As many of our results hold in the generality of Banach lattices, we briefly summarize some of the standard notations and conventions from this theory. For the most part, our conventions align with the Refs. [7, 61]. Moreover, the statements of our results require minimal knowledge of Banach lattices to understand; it is simply the proofs that use the technology and terminology from this theory. Unless otherwise mentioned, all L_p -spaces, $C(K)$ -spaces and Banach lattices are real. When a result is applicable for complex scalars, we will explicitly state this. The word “subspace” is to be interpreted in the vector space sense. If a result requires the subspace to be closed or (in)finite dimensional, we will state this.

Recall that a *vector lattice* is a vector space, equipped with a compatible lattice-ordering (see [7] for a precise definition). For a vector lattice X , the positive cone of X is denoted by $X_+ := \{f \in X : f \geq 0\}$. The infimum of $f, g \in X$ is denoted by $f \wedge g$, and the supremum is denoted by $f \vee g$. The modulus of f is defined as $|f| := f \vee (-f)$, and elements $f, g \in X$ are said to be *disjoint* if $|f| \wedge |g| = 0$. A *weak unit* is an element $e \in X_+$ for which $|f| \wedge e = 0$ implies $f = 0$. For a net (f_α) in X , the notation $f_\alpha \downarrow 0$ means that f_α is decreasing and has infimum 0. A subspace $E \subseteq X$ is a *sublattice* if it is closed under finite lattice operations; it is an *ideal* if $f \in E$ and $|g| \leq |f|$ implies $g \in E$.

A *Banach lattice* is a Banach space which is also a vector lattice, and for which one has the compatibility condition $\|f\| \leq \|g\|$ whenever $|f| \leq |g|$. Note that the SPR inequality (1.1) remains well-defined when $L_p(\mu)$ is replaced by an arbitrary Banach lattice. As we will see, several of our results on SPR are also valid in this level of generality. Common examples of Banach lattices include L_p -spaces, $C(K)$ -spaces, Orlicz spaces, and various sequence spaces. In this case, the ordering is pointwise, i.e., $f \leq g$ means $f(t) \leq g(t)$ for all (or almost all in the case of measurable functions) t in the domain of f and g .

A Banach lattice X is *order continuous* if for each net (f_α) satisfying $f_\alpha \downarrow 0$ we have $f_\alpha \xrightarrow{\|\cdot\|_X} 0$. L_p -spaces are order continuous for $1 \leq p < \infty$, but $C(K)$ -spaces are not (unless they are finite dimensional). To transfer results from $L_1(\mu)$ to more general Banach lattices, we will make use of the *AL-representation* procedure. For this, let X be an order continuous Banach lattice with a weak unit e . It is known that X

can be represented as an order and norm dense ideal in $L_1(\mu)$ for some finite measure μ . That is, there is a vector lattice isomorphism $T : X \rightarrow L_1(\mu)$ such that $\text{Range } T$ is an order and norm dense ideal in $L_1(\mu)$. Note that T need not be a norm isomorphism, though T may be chosen to be continuous with $T e = \mathbb{1}$. Moreover, $\text{Range } T$ contains $L_\infty(\mu)$ as a norm and order dense ideal. It is common to identify X with $\text{Range } T$ and view X as an ideal of $L_1(\mu)$. Such an inclusion of X into $L_1(\mu)$ is called an *AL-representation* of X . We refer to [61, Theorem 1.b.14] or [39, Section 4] for details on AL-representations.

2.1 The Kadec–Pelczynski dichotomy

Here, we briefly recap the literature on subspaces which do not contain almost disjoint normalized sequences. Recall that a sequence (x_n) in a Banach lattice X is said to be a *normalized almost disjoint sequence* if $\|x_n\|_X = 1$ for all n , and there exists a disjoint sequence (d_n) in X such that $\|x_n - d_n\|_X \rightarrow 0$. Following [15, 40, 41], a closed subspace of a Banach lattice that fails to contain normalized almost disjoint sequences will be called *dispersed*. The classical *Kadec–Pelczynski dichotomy* (c.f. [61, Proposition 1.c.8]) states that for a subspace E of an order continuous Banach lattice X with weak unit, either

- (i) E fails to be dispersed, i.e., E contains an almost disjoint normalized sequence, or,
- (ii) E is isomorphic to a closed subspace of $L_1(\Omega, \Sigma, \mu)$.

As we will see in Theorem 3.4, for real scalars, a subspace does stable phase retrieval if and only if it does not contain normalized almost disjoint pairs. Hence, the Kadec–Pelczynski dichotomy will provide a tool to analyze such subspaces.

In $L_p(\mu)$ for $1 \leq p < \infty$ and a probability measure μ , the Kadec–Pelczynski dichotomy can be improved. Indeed, we summarize the literature in the following theorem.

Theorem 2.1 *Let $1 \leq p < \infty$ and μ be a probability measure. For a closed subspace E of $L_p(\mu)$, the following are equivalent:*

- (a) E is dispersed, i.e., E contains no almost disjoint normalized sequences;
- (b) There exists $0 < q < p$ such that $\|\cdot\|_{L_p} \sim \|\cdot\|_{L_q}$ on E ;
- (c) For all $0 < q < p$, $\|\cdot\|_{L_p} \sim \|\cdot\|_{L_q}$ on E ;
- (d) E is strongly embedded in $L_p(\mu)$, i.e., convergence in measure coincides with norm convergence on E .

Moreover,

- (i) For $p \neq 2$, a closed subspace of $L_p[0, 1]$ is dispersed if and only if it contains no subspace isomorphic to ℓ_p .
- (ii) For $p > 2$, a closed subspace of $L_p[0, 1]$ is dispersed if and only if it is isomorphic to a Hilbert space.
- (iii) For $p < 2$ and any $q \in (p, 2]$, there is a closed subspace of $L_p[0, 1]$ which is both dispersed and isometric to $L_q[0, 1]$.

- (iv) For $p \neq 2$, $L_p[0, 1]$ cannot be written as the direct sum of two dispersed subspaces.
- (v) There exists an orthogonal decomposition $L_2[0, 1] = E \oplus E^\perp$ with both E and E^\perp dispersed in $L_2[0, 1]$.

Proof The equivalence of (b), (c) and (d) is [4, Proposition 6.4.5]. Other than the isometric portion of statement (iii), the rest of the statements are neatly summarized in [40, Propositions 3.4 and 3.5], with references to various textbooks for proofs. An isometric embedding of $L_q[0, 1]$ into $L_p[0, 1]$ for $q \in (p, 2)$ is given in [61, Corollary 2.f.5]. An isometric embedding of ℓ_2 into $L_p[0, 1]$ for $1 \leq p < \infty$ is given in [4, Proposition 6.4.12]. \square

Remark 2.2 One of the goals of this article is to find SPR analogues of the results in Theorem 2.1. However, we should mention that the connection between Theorem 2.1 and SPR has already been implicitly made in [28]. Recall that a subset $\Lambda \subseteq \mathbb{Z}$ is called a $\Lambda(p)$ -set if the closed subspace generated by the set of exponentials $\{e^{2\pi i n x} : n \in \Lambda\} \subseteq L_p(\mathbb{T})$ satisfies the equivalent conditions in Theorem 2.1. Such sets have been deeply studied [9, 20, 69], and have many interesting properties. For example, Rudin [68] showed that for all integers $n > 1$, there are $\Lambda(2n)$ -sets that are not $\Lambda(q)$ -sets for every $q > 2n$. Moreover, Bourgain [19] extended Rudin's theorem to all $p > 2$. On the other hand, when $p < 2$, and Λ is $\Lambda(p)$, then it is automatically $\Lambda(p + \varepsilon)$ for some $\varepsilon > 0$ [11, 45]. Since $|e^{2\pi i n x}| \equiv 1$, complex exponentials cannot do stable phase retrieval. However, by replacing $e^{2\pi i n x}$ by $\sin(2\pi n x)$ or other trigonometric polynomials with non-constant moduli, [28] is able to use combinatorial arguments in the spirit of Rudin to produce SPR subspaces of $L_p(\mu)$ when the dilation set Λ is sufficiently sparse.

2.2 Complex Banach lattices

Complex Banach lattices are defined as complexifications of real Banach lattices, and in the case of complex function spaces like $C(K)$ and $L_p(\mu)$, agree with the standard definition. More precisely, by a *complex Banach lattice* we mean the complexification $X_{\mathbb{C}} = X \oplus iX$ of a real Banach lattice, X , endowed with the norm $\|x + iy\|_{X_{\mathbb{C}}} = \|(x + iy)\|_X$, where the modulus $|\cdot| : X_{\mathbb{C}} \rightarrow X_+$ is the mapping given by

$$|x + iy| = \sup_{\theta \in [0, 2\pi]} \{x \cos \theta + y \sin \theta\}, \text{ for } x + iy \in X_{\mathbb{C}}. \quad (2.1)$$

We refer to [1, Section 3.2] and [70, Section 2.11] for a proof that the modulus function is well-defined, and behaves as expected.

With the above definition, one can define complex sublattices, complex ideals, etc. However, we will not need this. We do, however, note that if $T : X \rightarrow Y$ is a real linear operator between real Banach lattices, then we may define the *complexification* $T_{\mathbb{C}} : X_{\mathbb{C}} \rightarrow Y_{\mathbb{C}}$ of T via $T_{\mathbb{C}}(x + iy) = Tx + iTy$. The map $T_{\mathbb{C}}$ is \mathbb{C} -linear, bounded, and if T is a lattice homomorphism then $T_{\mathbb{C}}$ preserves moduli, i.e., $T|z| = |T_{\mathbb{C}}z|$ for $z \in X_{\mathbb{C}}$. When we work with complex Banach lattices $X_{\mathbb{C}}$, we will use these facts

to identify $X_{\mathbb{C}}$ as a space of measurable functions on some measure space, and then work pointwise. How to do this will be explained later in the paper.

3 General theory

In this section, we present several results on (stable) phase retrieval that are valid in general Banach lattices. We begin with the definitions:

Definition 3.1 Let E be a subspace of a vector lattice X . We say that E does *phase retrieval* if for each $f, g \in E$ with $|f| = |g|$ there is a scalar λ such that $f = \lambda g$.

Definition 3.2 Let E be a subspace of a real or complex Banach lattice X . We say that E does *C-stable phase retrieval* if for each $f, g \in E$ we have

$$\inf_{|\lambda|=1} \|f - \lambda g\| \leq C \||f| - |g|\|. \quad (3.1)$$

If E does *C-stable phase retrieval* for some C , we simply say that E does *stable phase retrieval* (*SPR* for short).

Note that if a subspace E of a real or complex Banach lattice X does *C-stable phase retrieval*, then so does its closure.

3.1 Connections with almost disjoint pairs and sequences

When considering whether a subspace $E \subseteq X$ does phase retrieval, there is one obvious obstruction. If $f, g \in E$ are non-zero disjoint vectors, then $|f - g| = |f + g| = |f| + |g|$, but $f - g$ cannot be a multiple of $f + g$. Hence, if E is to do phase retrieval, then it cannot contain disjoint pairs. Similarly, if E is to do stable phase retrieval, then it cannot contain “almost” disjoint pairs. As we will now see, in the real case, these are the *only* obstructions to (stable) phase retrieval.

Definition 3.3 Let E be a subspace of a real or complex Banach lattice X . We say that E contains ε -*almost disjoint pairs* if there are $f, g \in S_E$ (here and below, $S_E = \{e \in E : \|e\| = 1\}$) stands for the unit sphere of E) such that $\||f| \wedge |g|\| < \varepsilon$. If E contains ε -almost disjoint pairs for all $\varepsilon > 0$, we say that E contains *almost disjoint pairs*.

Theorem 3.4 Let E be a subspace of a Banach lattice X , $C \geq 1$ and $\varepsilon > 0$. Then,

- (i) If E does *C-stable phase retrieval*, then it contains no $\frac{1}{C}$ -*almost disjoint pairs*;
- (ii) If E contains no ε -*almost disjoint pairs*, then it does $\frac{2}{\varepsilon}$ -*stable phase retrieval*.

In particular, E does *stable phase retrieval* if and only if it does not contain *almost disjoint pairs*.

Proof (i): Suppose that E does *C-stable phase retrieval*, but there are $f, g \in E$ such that $\|f\| = \|g\| = 1$ but $\||f| \wedge |g|\| < \frac{1}{C}$. Define $h_1 = f + g$ and $h_2 = f - g$. Then since the identity

$$\||f + g| - |f - g|\| = 2(\||f| \wedge |g|\|)$$

holds in any vector lattice by [7, Theorem 1.7], we have

$$\| |h_1| - |h_2| \| = 2 \| |f| \wedge |g| \| < \frac{2}{C}.$$

On the other hand, $h_1 + h_2 = 2f$ has norm 2, and $h_1 - h_2 = 2g$ also has norm 2. This contradicts that E does C -stable phase retrieval.

(ii): A classical Banach lattice fact (see, e.g., [10, Remark after Lemma 3.3]) is that every Banach lattice embeds lattice isometrically into some space of the form

$$\left(\bigoplus_{i \in I} L_1(\Omega_i, \Sigma_i, \mu_i) \right)_\infty.$$

Since both stable phase retrieval and existence of almost disjoint pairs are invariant under passing to and from closed sublattices, we may assume without loss of generality that X is of this form.

Suppose E does not do $\frac{2}{\varepsilon}$ -stable phase retrieval. Find $f = (f_i), g = (g_i) \in E$ such that $\|f - g\|, \|f + g\| > \frac{2}{\varepsilon} \| |f| - |g| \|$. For each $i \in I$ let

$$I_i = \{t \in \Omega_i : \text{sign}(f_i(t)) = \text{sign}(g_i(t)), \text{ or one of } f_i(t), g_i(t) \text{ is zero}\}.$$

Then

$$I_i^c := \Omega_i \setminus I_i = \{t \in \Omega_i : \text{sign}(f_i(t)) = -\text{sign}(g_i(t))\}.$$

We compute that

$$|f| - |g| = (|f_i| - |g_i|)_{i \in I} = (|f_i|_{I_i} - |g_i|_{I_i})_{i \in I} + (|f_i|_{I_i^c} - |g_i|_{I_i^c})_{i \in I}.$$

So, since the modulus is additive on disjoint vectors,

$$|f| - |g| = (|f_i|_{I_i} - |g_i|_{I_i})_{i \in I} + (|f_i|_{I_i^c} - |g_i|_{I_i^c})_{i \in I}.$$

Now, by definition of I_i we have

$$(|f_i|_{I_i} - |g_i|_{I_i})_{i \in I} = (|f_i|_{I_i} - g_i|_{I_i})_{i \in I}$$

and

$$(|f_i|_{I_i^c} - |g_i|_{I_i^c})_{i \in I} = (|f_i|_{I_i^c} + g_i|_{I_i^c})_{i \in I}.$$

Notice next that $d_1 := (f_{i|I_i^c} - g_{i|I_i^c})_{i \in I}$ and $d_2 := (f_{i|I_i} + g_{i|I_i})_{i \in I}$ are disjoint. Moreover,

$$\begin{aligned}\|f - g - (f_{i|I_i^c} - g_{i|I_i^c})_{i \in I}\| &= \|(f_{i|I_i} - g_{i|I_i})_{i \in I}\| = \|(|f_{i|I_i}| - |g_{i|I_i}|)\|_{i \in I} \\ &\leq \||f| - |g|\| < \frac{\varepsilon}{2} \|f - g\|.\end{aligned}$$

Similarly,

$$\begin{aligned}\|f + g - (f_{i|I_i} + g_{i|I_i})_{i \in I}\| &= \|(f_{i|I_i^c} + g_{i|I_i^c})_{i \in I}\| = \|(|f_{i|I_i^c}| - |g_{i|I_i^c}|)\|_{i \in I} \\ &\leq \||f| - |g|\| < \frac{\varepsilon}{2} \|f + g\|.\end{aligned}$$

By assumption, we have that both $f + g$ and $f - g$ are non-zero. Hence, by [7, Lemma 1.4], and the fact that $|d_1| \wedge |d_2| = 0$ we have

$$\begin{aligned}\frac{|f - g|}{\|f - g\|} \wedge \frac{|f + g|}{\|f + g\|} &\leq \frac{|f - g - d_1|}{\|f - g\|} \wedge \frac{|f + g|}{\|f + g\|} + \frac{|d_1|}{\|f - g\|} \wedge \frac{|f + g|}{\|f + g\|} \\ &\leq \frac{|f - g - d_1|}{\|f - g\|} \wedge \frac{|f + g|}{\|f + g\|} + \frac{|d_1|}{\|f - g\|} \wedge \frac{|f + g - d_2|}{\|f + g\|}.\end{aligned}$$

It follows that

$$\left\| \frac{|f - g|}{\|f - g\|} \wedge \frac{|f + g|}{\|f + g\|} \right\| \leq \frac{\|f - g - d_1\|}{\|f - g\|} + \frac{\|f + g - d_2\|}{\|f + g\|} < \varepsilon.$$

Thus, we have constructed normalized ε -almost disjoint vectors $\frac{f+g}{\|f+g\|}$ and $\frac{f-g}{\|f-g\|}$ in E . \square

Remark 3.5 After this paper was complete, Bilokopytov [16] contacted us with a streamlined proof of Theorem 3.4, which also avoids the extra constant 2. We expect the proof to be published independently.

Remark 3.6 Implication (i) of Theorem 3.4 holds when the Banach lattice X is replaced by any vector lattice equipped with an absolute norm. Here, a norm on a vector lattice X is *absolute* if $\|f\| = \|f\|$ for all $f \in X$; see [17, 56, 66] for more information. The proof of Theorem 3.4 also shows that a subspace of a Banach lattice does phase retrieval if and only if it does not contain disjoint non-zero vectors. A compactness argument then yields that in finite dimensions, phase retrieval implies stable phase retrieval. Indeed, consider the map $S_E \times S_E \rightarrow \mathbb{R}$, $(f, g) \mapsto \||f| \wedge |g|\|$. Then this map is continuous, so its image is compact, which allows one to conclude that the existence of almost disjoint pairs implies the existence of a disjoint pair. In infinite dimensions, it is relatively easy to construct subspaces doing phase retrieval but failing stable phase retrieval.

Proposition 3.7 *Every infinite dimensional Banach lattice has a closed subspace which does phase retrieval but not stable phase retrieval.*

Proof By [6, p. 46, Exercise 13], any infinite dimensional Banach lattice X contains a normalized disjoint positive sequence, which we shall index as consisting of vectors $(u_i)_{i \in \mathbb{N}}$ and $(v_s)_{s \in \mathcal{S}}$; here, \mathcal{S} denotes the set of all two-element subsets of \mathbb{N} (the order is not important). We fix an injection $\phi : \mathbb{N}^2 \rightarrow \mathbb{N}$ and consider the vectors

$$f_i = u_i + \sum_{j \neq i} 2^{-4\phi(i,j)} v_{\{i,j\}}.$$

The sum above converges, and we have

$$\|u_i - f_i\| \leq \varepsilon_i, \text{ where } \varepsilon_i = \sum_j 2^{-4\phi(i,j)}.$$

Then $\sum_i \varepsilon_i = \sum_{i,j} 2^{-4\phi(i,j)} \leq \sum_m 2^{-4m} = 1/15$, hence, by [4, Theorem 1.3.9], (f_i) is a Schauder basic sequence. Also, $1 \leq \|f_i\| \leq 16/15$ for each i , so this basis is semi-normalized. We shall show that $E = \overline{\text{span}}[f_i : i \in \mathbb{N}]$ fails stable phase retrieval, but has phase retrieval.

To show the failure of SPR, let, for $i \neq j$, $\psi(i, j) = \max\{\phi(i, j), \phi(j, i)\}$. Clearly $\psi(i, j) = \psi(j, i)$, and $\lim_j \psi(i, j) = \infty$ for any i . Note that $f_i \wedge f_j = 2^{-4\psi(i,j)} v_{\{i,j\}}$, hence

$$\|f_i \wedge f_j\| = 2^{-4\psi(i,j)} \xrightarrow[i,j \rightarrow \infty]{} 0.$$

Next we show that E does phase retrieval. Pick non-zero $f, g \in E$, with $|f| = |g|$; we have to show that $f = \pm g$. To this end, write $f = \sum_i a_i f_i$ and $g = \sum_i b_i f_i$. We can expand

$$f = \sum_i a_i u_i + \sum_{\{i,j\} \in \mathcal{S}} (a_i 2^{-4\phi(i,j)} + a_j 2^{-4\phi(j,i)}) v_{\{i,j\}},$$

and likewise for g . Comparing the coefficients with u_i , we conclude that, for every i , $|a_i| = |b_i|$. By switching signs in front of f and g , and by re-indexing, we can assume that $a_1 = b_1 > 0$. We have to show that the equality $a_i = b_i$ holds for every $i > 1$.

The preceding reasoning shows that $a_i = 0$ iff $b_i = 0$. Suppose both a_i and b_i are different from 0. Comparing the coefficients with $v_{\{i,j\}}$, we see that

$$|2^{-4\phi(1,i)} a_1 + 2^{-4\phi(i,1)} a_i| = |2^{-4\phi(1,i)} b_1 + 2^{-4\phi(i,1)} b_i|,$$

which is only possible if $\text{sign } a_i = \text{sign } b_i$. □

Example 3.8 Theorem 3.4 fails for complex spaces. Indeed, define E as the complex span of $\{(1, 1, 1), (i, 1, -1)\} \subseteq \mathbb{C}^3$, where we equip \mathbb{C}^3 with the modulus $|(a, b, c)| := (|a|, |b|, |c|)$. Clearly, E contains vectors f, g with $|f| = |g|$ but such that $f - \lambda g$ is not zero for any $\lambda \in \mathbb{C}$. Hence, E fails phase retrieval. However, one can easily compute that E contains no disjoint vectors, which by compactness yields the non-existence

of almost disjoint vectors. Moreover, as observed in [28], a complex subspace that contains two linearly independent real vectors cannot do complex phase retrieval. In particular, if $E \subseteq X$ is subspace of a Banach lattice X with $\dim E \geq 2$, then the canonical subspace $E_{\mathbb{C}} \subseteq X_{\mathbb{C}}$ fails to do phase retrieval.

Remark 3.9 Theorem 3.4 shows that for real scalars, the study of subspaces doing stable phase retrieval is equivalent to the study of subspaces lacking almost disjoint pairs. As mentioned in Sect. 2.1, there is a vast literature on closed subspaces lacking almost disjoint normalized *sequences*. Clearly, if E contains an almost disjoint normalized sequence, then it fails to do stable phase retrieval. However, the converse is not true. For example, the standard Rademacher sequence (r_n) in $L_p[0, 1]$, $1 \leq p < \infty$, is dispersed by Khintchine's inequality, but $|r_n| \equiv 1$ for all n . Moreover, if one adds a single disjoint vector to a dispersed subspace, one produces a dispersed subspace failing phase retrieval. Nevertheless, as mentioned in Sect. 1.2, many of the results in Theorem 2.1 have SPR analogues.

3.2 Hölder stable phase retrieval and witnessing failure of SPR on orthogonal vectors

In [28], the following terminology was introduced in the setting of L_p -spaces: A subspace E of a real or complex Banach lattice X is said to do γ -Hölder stable phase retrieval with constant C if for all $f, g \in E$ we have

$$\inf_{|\lambda|=1} \|f - \lambda g\|_X \leq C \| |f| - |g| \|_X^\gamma (\|f\|_X + \|g\|_X)^{1-\gamma}. \quad (3.2)$$

The utility of this definition arose from a construction in [28] of SPR subspaces of $L_4(\mu)$ which are dispersed in $L_6(\mu)$. Applying certain Hölder inequality arguments, [28] was then able to deduce that such subspaces do $\frac{1}{4}$ -Hölder stable phase retrieval in $L_2(\mu)$. The idea in [28] is to begin with an orthonormal sequence (r_k) , and instead of comparing $|f|$ to $|g|$, one compares $|f|^2$ to $|g|^2$. Assuming the integrability condition $r_k \in L_4(\mu)$ with uniformly bounded norm, and various orthogonality and mean-zero conditions on the products $r_k \bar{r}_j$, the orthogonal expansion $f = \sum_k a_k r_k$ leads to an orthogonal expansion

$$|f|^2 = \sum_{k \neq j} a_k \bar{a}_j r_k \bar{r}_j + \sum_k |a_k|^2 s_k + \|f\|_{L_2}^2 \mathbb{1}, \quad s_k = |r_k|^2 - 1.$$

The products $r_k \bar{r}_j$ encode how the subspace “sits” in $L_4(\mu)$, i.e., they encode the lattice structure. However, analyzing $|f|^2$ rather than $|f|$ allows one to work algebraically. As was shown in [28], if one imposes appropriate orthogonality conditions, the subspace E spanned by r_k will do stable phase retrieval in $L_4(\mu)$. Reference [28] then gives examples of such r_k built from dilates of a single function P , with $|P|$ not identically constant. Verifying that such sequences (r_k) satisfy the required orthogonality conditions is then a combinatorial exercise, using sparseness of the dilates to get non-overlapping supports with respect to the basis expansion. This sparseness

naturally leads to E lying in higher L_p -spaces, so that by interpolating, one concludes that E does Hölder stable phase retrieval in $L_2(\mu)$ with $\gamma = \frac{1}{4}$ if $p = 6$, and $\gamma \rightarrow \frac{1}{2}$ as $p \rightarrow \infty$.

The purpose of this section is to show that—at the cost of dilating the constant—Hölder stable phase retrieval is equivalent to stable phase retrieval. For real scalars, this can already be deduced from the almost disjoint pair characterization in Theorem 3.4. However, the proof below works equally well for complex scalars. The following theorem was proven in [5] for phase retrieval using a continuous frame for a Hilbert space. We extend it here to subspaces of Banach lattices.

Theorem 3.10 *Let $(X, \|\cdot\|)$ be a Banach lattice, real or complex. Fix linearly independent $f, g \in X$, and suppose that $Y = \text{span}\{f, g\}$ is equipped with a Hilbert space norm $\|\cdot\|_H$, which is K -equivalent to $\|\cdot\|$. Then there exist $f', g' \in Y$ so that*

$$\min_{|\lambda|=1} \|f - \lambda g\| \leq K \min_{|\lambda|=1} \|f' - \lambda g'\|, \quad (3.3)$$

and

$$(\|f'\|^2 + \|g'\|^2)^{\frac{1}{2}} \leq K \min_{|\lambda|=1} \|f' - \lambda g'\|, \quad (3.4)$$

and

$$\||f| - |g|\| \geq \||f'| - |g'|\|. \quad (3.5)$$

Remark 3.11 Conditions (3.3) and (3.5) state that replacing (f, g) by (f', g') tightens the SPR inequality up to the universal factor K . The condition (3.4) states that f' and g' are “almost orthogonal” (in fact, the proof shows that they are orthogonal in the Hilbert space $H = (Y, \|\cdot\|_H)$); it also permits us to witness the failure of SPR on f', g' with controlled norm.

In general, by John’s Theorem, every 2-dimensional space is $\sqrt{2}$ -equivalent to a Hilbert space, but in certain cases a better estimate can be obtained. For instance, if $X = L_2(\mu)$, then for the inherited norm on Y we have $K = 1$ and f' orthogonal to g' . If X is a Banach lattice, which is r -convex and s -concave ($1 < r \leq 2 < s < \infty$) with constants $M^{(r)}(X)$ and $M_{(s)}(X)$ respectively, then, by [71, Theorem 28.6], there exists $\|\cdot\|_H$ for which

$$K \leq M^{(r)}(X)M_{(s)}(X)2^\alpha, \quad \text{where } \alpha = \max \left\{ \frac{1}{r} - \frac{1}{2}, \frac{1}{2} - \frac{1}{s} \right\}.$$

In particular, for $X = L_p$, there exists $\|\cdot\|_H$ for which $K \leq 2^{|1/p-1/2|}$. [71, Corollary 28.7] provides similar results for operator ideals (Schatten spaces).

In certain applications of Theorem 3.10 (such as Theorem 5.3), the norm $\|\cdot\|_H$ arises not from the Hilbert space with the minimal Banach–Mazur distance to Y , but from an equivalent Euclidean norm on some subspace E (with $Y \subseteq E \subseteq X$). In this setting K may exceed $\sqrt{2}$.

To prove Theorem 3.10, we need to represent elements of X as measurable functions. As mentioned in the proof of Theorem 3.4, every (real) Banach lattice X embeds lattice isometrically into a space of the form $(\bigoplus_{i \in I} L_1(\Omega_i, \Sigma_i, \mu_i))_\infty$. Hence, throughout the proof we can assume that elements of X are functions on a measure space. In the complex case, a similar reduction is possible. Indeed, let X be a complex Banach lattice. By the discussion in Sect. 2.2, we can assume that $X = Z_{\mathbb{C}}$ is the complexification of some (real) Banach lattice Z . We can then let $T : Z \rightarrow (\bigoplus_{i \in I} L_1(\Omega_i, \Sigma_i, \mu_i))_\infty$ be a lattice isometric embedding. The complexification $T_{\mathbb{C}}$ maps X into the complexification of $(\bigoplus_{i \in I} L_1(\Omega_i, \Sigma_i, \mu_i))_\infty$. The codomain of this map is still $(\bigoplus_{i \in I} L_1(\Omega_i, \Sigma_i, \mu_i))_\infty$, but now interpreted as a Banach lattice over the complex field (cf. [1, Exercises 3 and 5 on page 110]). Since T is one-to-one, the definition of $T_{\mathbb{C}}$ tells us that $T_{\mathbb{C}}$ is one-to-one. Moreover, as mentioned in Sect. 2.2, $T_{\mathbb{C}}$ preserves moduli. Finally, by [1, Lemma 3.18 or Corollary 3.23], $T_{\mathbb{C}}$ preserves norm. Thus, everything in the SPR inequality is preserved, so, analogously to the real case, we may assume throughout the proof that the complex Banach lattice X is a space of complex-valued functions.

Proof of Theorem 3.10 By scaling, we assume that, on Y , $\|\cdot\| \leq \|\cdot\|_H \leq K\|\cdot\|$. By replacing g by a unimodular scalar times g , we assume

$$\min_{|\lambda|=1} \|f - \lambda g\|_H = \|f - g\|_H.$$

This latter condition is equivalent to $\langle f, g \rangle \geq 0$. Indeed,

$$\|f - \lambda g\|_H^2 = \langle f, f \rangle + \langle g, g \rangle - 2\Re(\bar{\lambda}\langle f, g \rangle).$$

This is minimized when $\bar{\lambda}$ is the conjugate phase of $\langle f, g \rangle$. This is minimized when $\lambda = 1$ iff $\langle f, g \rangle \geq 0$.

Consider $f_r := f - r(f + g)$ and $g_r := g - r(f + g)$ for $r \in [0, 1/2]$. We let R be the first instance of $\langle f - r(f + g), g - r(f + g) \rangle = 0$. This is possible since when $r = 0$, the inner product is non-negative, and when $r = \frac{1}{2}$, it is negative. Note that

$$\|f_r - g_r\|_H = \|f - g\|_H.$$

Thus, since f_R and g_R are orthogonal,

$$\min_{|\lambda|=1} \|f_R - \lambda g_R\|_H = \min_{|\lambda|=1} \|f - \lambda g\|_H.$$

We will take $f' = f_R$ and $g' = g_R$. To see (3.3), we compute

$$\min_{|\lambda|=1} \|f - \lambda g\| \leq \min_{|\lambda|=1} \|f - \lambda g\|_H = \min_{|\lambda|=1} \|f' - \lambda g'\|_H \leq K \min_{|\lambda|=1} \|f' - \lambda g'\|. \quad (3.6)$$

Moreover, as f' and g' are orthogonal in H ,

$$K \min_{|\lambda|=1} \|f' - \lambda g'\| \geq \min_{|\lambda|=1} \|f' - \lambda g'\|_H = (\|f'\|_H^2 + \|g'\|_H^2)^{\frac{1}{2}} \geq (\|f'\|^2 + \|g'\|^2)^{\frac{1}{2}}. \quad (3.7)$$

This gives (3.4).

We now verify (3.5). To see this, we prove

$$||f_r| - |g_r|| \leq ||f| - |g|| \text{ for } r \in \left[0, \frac{1}{2}\right]. \quad (3.8)$$

We represent $X \subseteq L^0(\Omega)$ and let $t \in \Omega$. We will prove that

$$||f_r(t)| - |g_r(t)|| \leq ||f(t)| - |g(t)|| \text{ for } r \in \left[0, \frac{1}{2}\right]. \quad (3.9)$$

Note that (3.9) is simply a claim that an elementary inequality holds for complex numbers. Write $f(t) = a + ib$ and $g(t) = c + id$. Multiplying $f(t)$ and $g(t)$ by a unimodular scalar, we rotate so that $d = -b$. WLOG, $|a| \geq |c|$; then, multiplying by -1 if necessary, we also assume $a \geq 0$. We have

$$f_r(t) = a - r(a + c) + ib, \quad g_r(t) = c - r(a + c) - ib.$$

Now, we note that our assumptions give $||f_r(t)| - |g_r(t)|| = |f_r(t)| - |g_r(t)|$ for $0 \leq r \leq \frac{1}{2}$. Indeed, $\Im(f_r(t)) = -\Im(g_r(t))$ and $(\Re(f_r(t)))^2 \geq (\Re(g_r(t)))^2$ for $0 \leq r \leq \frac{1}{2}$ by elementary computations. Taking $r = 0$, $||f(t)| - |g(t)|| = |f(t)| - |g(t)|$. Hence, we must prove

$$|f_r(t)| - |g_r(t)| \leq |f(t)| - |g(t)| \text{ for } r \in \left[0, \frac{1}{2}\right].$$

This inequality is true for all $r \geq 0$. Indeed, recall first that $a \geq c$. By the Fundamental Theorem of Calculus, for any convex function ϕ and $w \geq 0$, we have $\phi(a - w) - \phi(c - w) \leq \phi(a) - \phi(c)$. In our case, the function $h(s) = \sqrt{s^2 + b^2}$ is convex and $r(a + c) \geq 0$; therefore,

$$\begin{aligned} |f_r(t)| - |g_r(t)| &= h(a - r(a + c)) - h(c - r(a + c)) \leq h(a) - h(c) \\ &= |f(t)| - |g(t)|. \end{aligned}$$

□

Corollary 3.12 *Let E be a subspace of a real or complex Banach lattice X , and $\gamma \in (0, 1]$. If E does γ -Hölder stable phase retrieval in X with constant $C > 0$ then E does stable phase retrieval in X with constant $\sqrt{2}(\sqrt{8}C)^{\frac{1}{\gamma}}$.*

Proof Let $f, g \in E$ with $\|f\| = 1$ and $\|g\| \leq 1$ such that

$$(\|f\|^2 + \|g\|^2)^{\frac{1}{2}} \leq \sqrt{2} \inf_{|\lambda|=1} \|f - \lambda g\|. \quad (3.10)$$

In particular,

$$2^{-1/2} \leq \inf_{|\lambda|=1} \|f - \lambda g\| \leq 2.$$

As E does C -stable γ -Hölder phase retrieval, we have that

$$2^{-1/2} \leq \inf_{|\lambda|=1} \|f - \lambda g\| \leq C \||f| - |g|\|^{\gamma} (\|f\| + \|g\|)^{1-\gamma} \leq 2^{1-\gamma} C \||f| - |g|\|^{\gamma}. \quad (3.11)$$

Thus, we have that $C^{1/\gamma} 2^{3/(2\gamma)-1} \||f| - |g|\| \geq 1$ and $\inf_{|\lambda|=1} \|f - \lambda g\| \leq 2$. It follows that

$$\inf_{|\lambda|=1} \|f - \lambda g\| \leq (2^{3/2} C)^{1/\gamma} \||f| - |g|\|. \quad (3.12)$$

To prove (3.12) we have assumed that $\|f\| = 1$ and $\|g\| \leq 1$. However, by scaling we have that any $f, g \in E$ which satisfy (3.10) also satisfy (3.12).

We now consider any pair of linearly independent vectors $x, y \in E$. By Theorem 3.10 there exists $f, g \in E$ which satisfy (3.10) such that

$$\min_{|\lambda|=1} \|x - \lambda y\| \leq \sqrt{2} \min_{|\lambda|=1} \|f - \lambda g\| \text{ and } \||x| - |y|\| \geq \||f| - |g|\|.$$

Thus, we have that

$$\min_{|\lambda|=1} \|x - \lambda y\| \leq 2^{1/2} (2^{3/2} C)^{1/\gamma} \||x| - |y|\|.$$

This proves that E does $2^{1/2} (2^{3/2} C)^{1/\gamma}$ -stable phase retrieval. \square

Remark 3.13 The constant $\sqrt{2}(\sqrt{8}C)^{\frac{1}{\gamma}}$ in Corollary 3.12 arises by using the worst case scenario $K = \sqrt{2}$ from Theorem 3.10. This constant can certainly be optimized; for example, if one also takes into account the distance from E to a Hilbert space.

To conclude this section we give a simple proof that in finite dimensions, phase retrieval is automatically stable.

Corollary 3.14 *Let X be a real or complex Banach lattice, and E a finite dimensional subspace of X . If E does phase retrieval, then E does stable phase retrieval.*

Proof The real case has already been dealt with in Remark 3.6, but the argument we provide below works for both real and complex scalars. Indeed, by Theorem 3.10, if

E fails to do stable phase retrieval then we can find, for each $N \in \mathbb{N}$, functions f_N, g_N with $\|f_N\| = 1, \|g_N\| \leq 1$,

$$(\|f_N\|^2 + \|g_N\|^2)^{\frac{1}{2}} \leq \sqrt{2} \min_{|\lambda|=1} \|f_N - \lambda g_N\|, \quad (3.13)$$

and

$$2 \geq \min_{|\lambda|=1} \|f_N - \lambda g_N\| > N\|f_N\| - \|g_N\|. \quad (3.14)$$

By compactness, after passing to subsequences, we may assume that $f_N \xrightarrow{\|\cdot\|} f$ and $g_N \xrightarrow{\|\cdot\|} g$, for some $f, g \in E$. Since $\|f_N\| = 1$ for all N , it follows that $\|f\| = 1$. Moreover, from (3.14) and continuity of lattice operations, we see that $\|f\| - \|g\| = 0$. Hence, $|f| = |g| \neq 0$. Fix a phase λ . By (3.13), we have

$$(\|f_N\|^2 + \|g_N\|^2)^{\frac{1}{2}} \leq \sqrt{2}\|f_N - \lambda g_N\|.$$

Passing to the limit, we see that

$$1 \leq (\|f\|^2 + \|g\|^2)^{\frac{1}{2}} \leq \sqrt{2}\|f - \lambda g\|.$$

Hence, $f \neq \lambda g$. It follows that E fails to do phase retrieval. \square

Remark 3.15 Note that the Banach lattice X in Corollary 3.14 is not assumed to be finite dimensional. This is of some note, as, unlike for closed spans, the closed sublattice generated by a finite set can be infinite dimensional.

4 Examples

4.1 Building SPR subspaces via isometric theory

As mentioned in Theorem 2.1, when $1 \leq p < 2$ and $q \in (p, 2]$, one can find isometric copies of $L_q[0, 1]$ in $L_p[0, 1]$. As we will now see, such subspaces must do SPR.

Proposition 4.1 Suppose $p, q \in [1, \infty)$, and either (1) $1 \leq p < q \leq 2$, or (2) $q = 2 < p < \infty$. There exists an $\varepsilon > 0$ such that if $E \subseteq L_p[0, 1]$ is $(1 + \varepsilon)$ -isomorphic to $F \subseteq L_q[0, 1]$, then E does SPR in $L_p[0, 1]$.

Proof We only handle case (1), as (2) is very similar. Suppose, for the sake of contradiction, that E fails SPR. Then by Theorem 3.4, E contains c -isomorphic copies of ℓ_p^2 , for any $c > 1$. Consequently, for any such c we can find norm one $f, g \in E$ so that $\|f + g\|_{L_p}, \|f - g\|_{L_p} \geq c^{-1}2^{1/p}$. However, by the Clarkson inequality in L_q ,

$$\|f + g\|_{L_q}^{q'} + \|f - g\|_{L_q}^{q'} \leq 2(\|f\|_{L_q}^q + \|g\|_{L_q}^q)^{q'-1} \leq (1 + \varepsilon)^{q'}2^{q'},$$

where $1/q + 1/q' = 1$. However, the left side is $\geq c^{-q'} 2^{1+q'/p}$, and it is easy to see that $1 + q'/p > q'$. Hence, we get a contradiction if $\varepsilon > 0$ is sufficiently small. \square

Corollary 4.2 *If either $1 \leq p < q \leq 2$, or $q = 2 < p < \infty$, then $L_p[0, 1]$ contains an SPR subspace isometric to $L_q[0, 1]$.*

Proof It is well known (see e.g. [51, Section 9]) that, under the above conditions, $L_p[0, 1]$ contains an isometric copy of $L_q[0, 1]$. By Proposition 4.1, that copy does SPR. \square

4.2 Existence of SPR embeddings into sequence spaces

Proposition 4.3 *If a Banach space E embeds into $\ell_\infty(\alpha)$ for some infinite cardinal α (which happens, in particular, when E itself has density character α), then there is an isomorphic SPR embedding of E inside of $\ell_\infty(\alpha)$.*

The fact that any Banach space E of density character α embeds isometrically into $\ell_\infty(\alpha)$ is standard. We recall the construction for the sake of completeness: Let $(x_i)_{i \in I}$ be a dense subset of E of cardinality α ; for each i find $x_i^* \in S_{E^*}$ so that $x_i^*(x_i) = \|x_i\|$. Then $E \rightarrow \ell_\infty(\alpha) : x \mapsto (x_i^*(x))_{i \in I}$ is the desired embedding. Similarly, one can show that if E is a dual space, with a predual of density character α , then E embeds isometrically into $\ell_\infty(\alpha)$.

To establish Proposition 4.3, it therefore suffices to prove:

Lemma 4.4 *For any infinite cardinal α , there exists an isometric SPR embedding of $\ell_\infty(\alpha)$ into itself.*

To prove Lemma 4.4, we rely on the following.

Lemma 4.5 *Suppose E is a (real or complex) Banach space, and $x, y \in E$ have norm 1. Then there exists a norm 1 functional $f \in E^*$ so that $|f(x)| \wedge |f(y)| \geq 1/5$.*

Proof Suppose first that $\text{dist}(y, \mathbb{F}x) \leq 2/5$ (here \mathbb{F} is either \mathbb{R} or \mathbb{C}). Find $t \in \mathbb{F}$ so that $\|y - tx\| \leq 2/5$. By the triangle inequality, $|t| \geq 3/5$. Find $f \in E^*$ so that $\|f\| = 1 = f(x)$. Then $|f(y)| \geq |t||f(x)| - \|y - tx\| \geq 1/5$. The case of $\text{dist}(x, \mathbb{F}y) \leq 2/5$ is handled similarly.

Now suppose $\text{dist}(x, \mathbb{F}y), \text{dist}(y, \mathbb{F}x) > 2/5$. By Hahn-Banach Theorem, there exist norm one $g, h \in E^*$ so that $g(x) \geq 2/5$, $g(y) = 0$, $h(y) \geq 2/5$, and $h(x) = 0$. Then $f := (g + h)/\|g + h\|$ has the desired properties. Indeed, $\|g + h\| \leq 2$, hence

$$|f(x)| \geq \frac{1}{2}(|g(x)| - |h(x)|) \geq \frac{1}{5},$$

and likewise, $|f(y)| \geq 1/5$. \square

Proof of Lemma 4.4 For the sake of brevity, we shall use the notation $E = \ell_\infty(\alpha)$, and $E_* = \ell_1(\alpha)$. Pick a dense set $(f_i)_{i \in I}$ in S_{E_*} , with $|I| = \alpha$. Define an isometric embedding $J : E \rightarrow \ell_\infty(I) : x \mapsto (f_i(x))_{i \in I}$. We shall show that, for every

$x, y \in S_E$ and $\varepsilon > 0$, there exists i so that $|f_i(x)| \wedge |f_i(y)| \geq 1/5 - \varepsilon$. Once this is done, we will conclude that $\|Jx \wedge Jy\| \geq 1/5$ for any $x, y \in S_E$, which by Theorem 3.4 tells us that J is indeed an SPR embedding.

By Lemma 4.5, there exists $f \in S_{E^*}$ so that $|f(x)| \wedge |f(y)| \geq 1/5$. By Goldstine's Theorem, there exists $f' \in S_{E^*}$ so that $|f'(x)| \wedge |f'(y)| \geq 1/5 - \varepsilon/2$. Find i so that $\|f' - f_i\| \leq \varepsilon/2$. Then

$$|f_i(x)| \wedge |f_i(y)| \geq |f'(x)| \wedge |f'(y)| - \|f' - f_i\| \geq \frac{1}{5} - \varepsilon,$$

which proves our claim. \square

Remark 4.6 We can define the canonical embedding of E into $C(B_{E^*})$ (with $B_{E^*} = \{e^* \in E^* : \|e^*\| \leq 1\}$ equipped with its weak* topology) by sending $e \in E$ to the function $e^* \mapsto e^*(e)$. The above reasoning shows that this embedding is SPR. For separable E , more can be said—see Proposition 6.2 below.

Remark 4.7 If an atomic lattice is order continuous (which ℓ_∞ of course is not), then the “gliding hump” argument shows the non-existence of infinite dimensional dispersed subspaces. The lattice c is not order continuous, but it has no infinite dimensional dispersed subspaces. This is because c contains c_0 as a subspace of finite codimension, hence any infinite dimensional subspace of c has an infinite dimensional intersection with c_0 .

Combining the results from this and the previous subsection, we see that, often, the collection of dispersed subspaces of a Banach lattice coincides with those that do SPR, up to isomorphism (cf. Question 7.1 below). Indeed, we have the following:

Corollary 4.8 *For every dispersed subspace $E \subseteq L_p[0, 1]$ ($1 \leq p \leq \infty$), there exists a closed subspace $E' \subseteq L_p[0, 1]$ isomorphic to E , and doing stable phase retrieval. The same result holds with $L_p[0, 1]$ replaced by $C[0, 1]$, $C(\Delta)$, c or any order continuous atomic Banach lattice.*

Proof By Theorem 2.1, for $1 \leq p < \infty$ and $p \neq 2$, a closed subspace of $L_p[0, 1]$ is dispersed if and only if it contains no subspace isomorphic to ℓ_p . A result of Rosenthal [67] states that for $1 \leq p < 2$, a subspace of $L_p[0, 1]$ that does not contain ℓ_p must be isomorphic to a subspace of L_r for some $r \in (p, 2]$. By Corollary 4.2, one can build an SPR copy of L_r in L_p .

In the case $2 \leq p < \infty$, Theorem 2.1 states that any dispersed subspace of $L_p[0, 1]$ must be isomorphic to a Hilbert space. By Corollary 4.2, $L_p[0, 1]$ contains an SPR copy of ℓ_2 . To deal with the case $p = \infty$, note that $L_\infty[0, 1]$ is isomorphic (as a Banach space) to ℓ_∞ , and use Lemma 4.4 together with the fact that ℓ_∞ lattice isometrically embeds in $L_\infty[0, 1]$.

For order continuous atomic lattices and c , there are no infinite dimensional dispersed subspaces by Remark 4.7. The claim for $C[0, 1]$ and $C(\Delta)$ will be proven in Proposition 6.2 below, when we analyze SPR subspaces of $C(K)$ -spaces. As we will see in the proof of Proposition 6.2, the fact that every separable Banach space embeds into $C[0, 1]$ and $C(\Delta)$ in an SPR fashion ultimately follows from Remark 4.6. \square

4.3 Explicit constructions of SPR subspaces using random variables

In this subsection, we construct SPR subspaces of a rather general class of function spaces by considering the closed span of certain independent random variables. The use of sub-Gaussian random vectors has been widely successful in building random frames for finite dimensional Hilbert spaces which do stable phase retrieval whose stability bound is independent of the dimension [25, 26, 35, 54, 55]. However, different distributions for random variables will allow for the construction of subspaces which do stable phase retrieval and are not isomorphic to Hilbert spaces. We begin by presenting a technical criterion for SPR.

Proposition 4.9 *Suppose X is a Banach lattice of measurable functions on a probability measure space (Ω, μ) which contains the indicator functions and has the property that for every $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon) > 0$ so that $\|\chi_S\| > \delta$ whenever $\mu(S) > \varepsilon$. Suppose, furthermore, that E is a subspace of X , which has the following property: There exist $\alpha > 1/2$ and $\beta > 0$ so that, for any norm one $f \in E$, we have*

$$\mu(\{\omega \in \Omega : |f(\omega)| \geq \beta\}) \geq \alpha. \quad (4.1)$$

Then E is an SPR-subspace.

Proof Suppose $f, g \in E$ have norm 1. By the Inclusion–Exclusion Principle,

$$\mu(\{\omega \in \Omega : |f(\omega)| \geq \beta, |g(\omega)| \geq \beta\}) \geq 2\alpha - 1.$$

Thus, $\||f| \wedge |g|\| \geq \beta\delta(2\alpha - 1)$. □

The above proposition is applicable, for instance, when X is a rearrangement invariant (r.i. for short; see [61] for an in-depth treatment) space on $(0, 1)$, equipped with the canonical Lebesgue measure λ . Examples include L_p spaces, and, more generally, Lorentz and Orlicz spaces (once again, described in great detail in [61]; for Lorentz spaces, see also [33]). Below we describe some SPR subspaces, spanned by independent identically distributed random variables.

Suppose f is a random variable, realized as a measurable function on $(0, 1)$ (with the usual Lebesgue measure λ). Then independent copies of f —denoted by f_1, f_2, \dots —can be realized on $((0, 1), \lambda)^{\aleph_0}$. By Caratheodory's Theorem (see e.g. [58, p. 121]), there exists a measure-preserving bijection between $((0, 1), \lambda)^{\aleph_0}$ and $((0, 1), \lambda)$. Therefore, we view f_1, f_2, \dots as functions on $(0, 1)$.

Suppose now that, in the above setting, the following statements hold:

- (i) f belongs to X , and has norm one in that space;
- (ii) There exists r so that, if f_1, \dots, f_n are independent copies of f , and $\sum_i |a_i|^r = 1$, then $\sum_i a_i f_i$ is equidistributed with f ;
- (iii) There exists $\beta > 0$ so that $\mathbb{P}(|f| > \beta) > 1/2$.

In this situation, if f_1, f_2, \dots are independent copies of f (viewed as elements of X , per the preceding paragraph), then $\overline{\text{span}}[f_i : i \in \mathbb{N}]$ is an SPR copy of ℓ_r in X .

We should mention two examples of random variables with the above properties: Gaussian ((ii) holds with $r = 2$) and q -stable ($q \in (1, 2)$; (ii) holds with $r = q$). The details can be found in [4, Section 6.4]. For the Gaussian variables, the probability density function is $d_f(x) = ce^{-x^2/2}$, with c depending on the normalization. For the q -stable variables with characteristic function $t \mapsto ce^{-|t|^q}$ (with c ensuring normalization), the Fourier inversion formula gives the density function

$$d_f(x) = \frac{c}{\pi} \int_0^\infty \cos(tx) e^{-t^q} dt.$$

In both cases, d_f is continuous (in the latter case, due to Dominated Convergence Theorem), hence there exists $\beta > 0$ so that

$$\mathbb{P}(|f| > \beta) = 1 - \int_{-\beta}^\beta d_f > \frac{3}{4}.$$

It is known that Gaussian random variables belong to L_p for $p \in [1, \infty)$, while the r -stable random variables ($1 < r < 2$) lie in L_p if and only $p \in [1, r)$. Moreover, the results from [61, p. 142–143] tell us that $L_s(0, 1) \subset L_{p,q}(0, 1)$ for $s > p$ (this is a continuous inclusion, not an isomorphic embedding). If $r > p$, then the r -stable variables belong to $L_{p,q}(0, 1)$ (indeed, take $s \in (p, r)$); then the r -stable variables live in $L_s(0, 1)$, which in turn sits inside of $L_{p,q}(0, 1)$. Likewise, one shows that any Lorentz space $L_{p,q}(0, 1)$ contains Gaussian random variables.

The above reasoning implies:

Proposition 4.10 *Suppose $1 \leq p < \infty$ and $1 \leq q \leq \infty$ (when $p = 1$, assume in addition $q < \infty$). Then $L_{p,q}(0, 1)$ contains a copy of ℓ_2 that does SPR. If, in addition, $1 \leq p < r < 2$, then $L_{p,q}(0, 1)$ contains a copy of ℓ_r that does SPR.*

4.4 Stability of SPR subspaces under ultraproducts and small perturbations

We show that SPR subspaces are stable under ultraproducts, and under small perturbations (in the sense of Hausdorff distance). These results hold for both real and complex spaces.

Proposition 4.11 *Suppose \mathfrak{U} is an ultrafilter on a set I , and, for each $i \in I$, E_i is a C -SPR subspace of a Banach lattice X_i . Then $\prod_{\mathfrak{U}} E_i$ is a C -SPR subspace of $\prod_{\mathfrak{U}} X_i$.*

We refer the reader to [46] or [32, Chapter 8] for information on ultraproducts of Banach spaces and Banach lattices.

Proof We have to show that, for any $x, y \in \prod_{\mathfrak{U}} E_i$, there exists a modulus one λ so that $\|x - \lambda y\| \leq C\|x\| - \|y\|$. To this end, find families (x_i) and (y_i) , representing x and y respectively. Then for each i there exists λ_i so that $|\lambda_i| = 1$ and $\|x_i - \lambda_i y_i\| \leq C\|x_i\| - \|y_i\|$. As ultraproducts preserve lattice operations, $|x|$ and $|y|$ are represented by $(|x_i|)$ and $(|y_i|)$, respectively, hence $\|x\| - \|y\| = \lim_{\mathfrak{U}} \|x_i\| - \|y_i\|$.

By the compactness of the unit torus, there exists $\lambda = \lim_{\mathfrak{U}} \lambda_i$, with $|\lambda| = 1$. Then $\|x - \lambda y\| = \lim_{\mathfrak{U}} \|x_i - \lambda_i y_i\|$, which leads to the desired inequality. \square

Remark 4.12 Proposition 4.11 can be used to give an alternative proof of Corollary 4.2. First find a family of finite dimensional subspaces $F_k \subseteq L_q(0, 1)$, ordered by inclusion, so that $\cup_k F_k$ is dense in $L_q(0, 1)$, and each F_k is isometric to $\ell_q^{n_k}$ for some n_k (one can, for instance, take subspaces spanned by certain step functions). A reasoning similar to that of [32, Theorem 8.8] permits us to find a free ultrafilter \mathfrak{U} so that $\prod_{\mathfrak{U}} F_k$ contains an isometric copy of $L_q(0, 1)$. A fortiori, $\prod_{\mathfrak{U}} \ell_q$ contains an isometric copy of $L_q(0, 1)$ (call it E).

Proposition 4.10 proves that $L_p(0, 1)$ contains a subspace, isometric to ℓ_q (spanned by Gaussian random variables for $q = 2$, q -stable random variables for $q < 2$) which does SPR. By Proposition 4.11, $\prod_{\mathfrak{U}} \ell_q$ embeds isometrically into $\prod_{\mathfrak{U}} L_p(0, 1)$, in an SPR fashion. By [46], $\prod_{\mathfrak{U}} L_p(0, 1)$ can be identified (as a Banach lattice) with $L_p(\Omega, \mu)$, for some measure space (Ω, μ) .

Let X be the (separable) sublattice of $L_p(\Omega, \mu)$ generated by E . By [61, Corollary 1.b.4], X is an L_p space. [58, Corollary, p. 128] gives a complete list of all separable L_p spaces; all of them lattice embed into $L_p(0, 1)$. Thus, we have established the existence of an SPR embedding of $E = L_q(0, 1)$ into $L_p(0, 1)$.

To examine stability of SPR under small perturbations, we introduce the notion of *one-sided Hausdorff distance* between subspaces of a given Banach space. If E, F are subspaces of X , define $d_{1H}(E, F)$ as the infimum of all $\delta > 0$ so that, for every $x \in F$ with $\|x\| \leq 1$ there exists $x' \in E$ with $\|x - x'\| < \delta$ (this “distance” is not reflexive, hence “one-sided”). Note also that, for x as above, there exists $x'' \in E$ with $\|x''\| = \|x\|$ and $\|x - x''\| < 2\delta$; indeed, one can take $x'' = \frac{\|x\|}{\|x'\|} x'$.

By “symmetrizing” d_{1H} , we obtain the classical *Hausdorff distance*: if E and F are subspaces of X , let $d_H(E, F) = \max\{d_{1H}(E, F), d_{1H}(F, E)\}$. For interesting properties of d_H , see [21], and references therein.

Proposition 4.13 *Suppose E is an SPR subspace of a Banach lattice X . Then there exists $\delta > 0$ so that any subspace F with $d_{1H}(E, F) < \delta$ is again SPR.*

From this we immediately obtain:

Corollary 4.14 *For any Banach lattice X , the set of its SPR subspaces is open in the topology determined by the Hausdorff distance.*

Remark 4.15 See [40, Proposition 3.10] for a similar stability result for dispersed subspaces of a Banach lattice.

Proof of Proposition 4.13 Suppose E does C-SPR. We shall show that, if $d_{1H}(E, F) < 1/(2\sqrt{2}(C + 1))$, then F does C' -SPR, with

$$\frac{1}{C'} = \frac{1}{C} \left(\frac{1}{\sqrt{2}} - 2d_{1H}(E, F) \right) - 2d_{1H}(E, F).$$

Suppose, for the sake of contradiction, that F fails to do C' -SPR. Find $f, g \in F$ so that $\min_{|\lambda|=1} \|f - \lambda g\| = 1$ and $\|f\| - \|g\| = c < 1/C'$. By Theorem 3.10, we

can find $f', g' \in F$ so that

$$\min_{|\lambda|=1} \|f' - \lambda g'\| \geq \frac{1}{\sqrt{2}}, \quad \||f'| - |g'|\| \leq c, \quad \text{and} \quad \|f'\| + \|g'\| \leq 2.$$

For any $\delta > d_{1H}(E, F)$, there exist $f'', g'' \in E$ so that $\|f'' - f'\| < \delta \|f'\|$ and $\|g'' - g'\| < \delta \|g'\|$. The triangle inequality implies:

$$\||f''| - |g''|\| \leq \||f'| - |g'|\| + \delta(\|f'\| + \|g'\|) \leq c + 2\delta;$$

$$\min_{|\lambda|=1} \|f'' - \lambda g''\| \geq \min_{|\lambda|=1} \|f' - \lambda g'\| - \delta(\|f'\| + \|g'\|) \geq \frac{1}{\sqrt{2}} - 2\delta.$$

As E does C -SPR, we conclude that

$$\frac{1}{\sqrt{2}} - 2\delta \leq C(c + 2\delta),$$

and consequently,

$$\frac{1}{\sqrt{2}} - 2d_{1H}(E, F) \leq C(c + 2d_{1H}(E, F)) < C \left(\frac{1}{C'} + 2d_{1H}(E, F) \right),$$

which contradicts our choice of C' . \square

Balan proved that frames which do stable phase retrieval for finite dimensional Hilbert spaces are stable under small perturbations [12]. The following extends this to infinite dimensional subspaces of Banach lattices.

Corollary 4.16 *Suppose (e_i) is a semi-normalized basic sequence in a Banach lattice X , so that $\overline{\text{span}}[e_i : i \in \mathbb{N}]$ does SPR in X . Then there exists $\varepsilon > 0$ so that if $(f_i) \subseteq X$ and $\sum_i \|e_i - f_i\| < \varepsilon$ then $\overline{\text{span}}[f_i : i \in \mathbb{N}]$ does SPR in X .*

Remark 4.17 In real L_2 , Corollary 4.16 can be strengthened. Suppose (e_i) is a sequence of normalized independent mean-zero random variables, spanning an SPR-subspace of L_2 . Then there exists an $\varepsilon > 0$ with the following property: if (f_i) is a collection of normalized independent mean-zero random variables so that (e_i, f_j) are independent whenever $i \neq j$, and $\sup_i \|e_i - f_i\| \leq \varepsilon$, then $\text{span}[f_i : i \in \mathbb{N}] \subseteq L_2$ does SPR as well. For the proof, recall that there exists $\gamma > 0$ so that the inequality $\|u \wedge v\| \geq \gamma$ holds for any norm one $u, v \in \text{span}[e_i : i \in \mathbb{N}]$. Let $\varepsilon = \gamma/4$. We will show that, for any norm one $x, y \in F = \text{span}[f_i : i \in \mathbb{N}]$, we have $\|x \wedge y\| \geq \gamma/2$.

Write $x = \sum_i \alpha_i f_i$ and $y = \sum_i \beta_i f_i$, and define $x' = \sum_i \alpha_i e_i$, $y' = \sum_i \beta_i e_i$. Then

$$\|x - x'\|^2 = \left\| \sum_i \alpha_i (f_i - e_i) \right\|^2 = \sum_i |\alpha_i|^2 \|f_i - e_i\|^2 \leq \varepsilon^2 \sum_i \alpha_i^2 = \varepsilon^2.$$

Similarly, $\|y - y'\| \leq \varepsilon$. Therefore, $\||x| - |x'|\|, \||y| - |y'|\| \leq \varepsilon$, hence $\||x| \wedge |y|\| \geq \||x'| \wedge |y'|\| - 2\varepsilon$. But $\||x'| \wedge |y'|\| \geq \gamma$, hence $\||x| \wedge |y|\| \geq \gamma/2$.

5 SPR in L_p -spaces

In this section, we investigate the relations between dispersed and SPR subspaces of L_p , as well as the relation between doing SPR in L_p versus doing SPR in L_q .

Theorem 5.1 *Every infinite dimensional dispersed subspace of an order continuous Banach lattice X contains a further closed infinite dimensional subspace that does SPR.*

Proof We first prove the claim for $L_1(\Omega, \mu)$, with μ a finite measure. Let E be a closed infinite dimensional subspace of $L_1(\Omega, \mu)$ containing no normalized almost disjoint sequence. By Theorem 2.1, E also does not contain ℓ_1 . By [57], every closed infinite dimensional subspace of $L_1(\Omega, \mu)$ almost isometrically contains ℓ_r for some $1 \leq r \leq 2$. Since E does not contain ℓ_1 , it follows that there exists $r > 1$ such that for all $\varepsilon > 0$, ℓ_r is $(1 + \varepsilon)$ -isomorphic to a subspace of E . Let $\alpha > 0$ be such that ℓ_1^2 is not $(1 + \alpha)$ -isomorphic to a subspace of ℓ_r . Such an α exists by the Clarkson argument in Proposition 4.1. We now claim that for $0 < \varepsilon < \alpha$, every subspace of L_1 that is $(1 + \varepsilon)$ -isomorphic to ℓ_r must do stable phase retrieval. Indeed, if E failed SPR, it would contain for all $\gamma > 0$ a $(1 + \gamma)$ -copy of ℓ_1^2 . Thus, for all $\gamma > 0$, we have that ℓ_1^2 is $(1 + \gamma)(1 + \varepsilon)$ -isomorphic to a subspace of ℓ_r . However, this gives a contradiction if $\gamma > 0$ is small enough such that $(1 + \gamma)(1 + \varepsilon) < 1 + \alpha$.

Now let E be a closed infinite dimensional dispersed subspace of an order continuous Banach lattice X . Replacing E be a separable subspace of E , we may assume that E is separable. Using that every closed sublattice of an order continuous Banach lattice is order continuous, replacing X by the closed sublattice generated by E in X , we may assume that X is separable. It follows in particular that X has a weak unit. By the AL-representation theory, there exists a finite measure space (Ω, μ) such that X can be represented as an ideal of $L_1(\Omega, \mu)$ satisfying

- (i) X is dense in $L_1(\Omega, \mu)$ and $L_\infty(\Omega, \mu)$ is dense in X ;
- (ii) $\|f\|_1 \leq \|f\|_X$ and $\|f\|_X \leq 2\|f\|_\infty$ for all $f \in X$.

Since E contains no almost disjoint normalized sequence, the Kadec–Pelczynski dichotomy [61, Proposition 1.c.8] guarantees that $\|\cdot\|_X \sim \|\cdot\|_{L_1}$ on E . In particular, we may view E as a closed infinite dimensional subspace of $L_1(\mu)$. We claim that E contains no almost disjoint sequence when viewed as a subspace of L_1 . Indeed, suppose there exists a sequence (x_n) in E with $\|x_n\|_{L_1} = 1$ for all n , and a disjoint sequence (d_n) in L_1 with $\|x_n - d_n\|_{L_1} \rightarrow 0$. Then in particular, x_n converges to 0 in measure. By [31, Theorem 4.6], $x_n \xrightarrow{un} 0$ in X . That is, for all $u \in X$, we have that $\|x_n \wedge u\|_X \rightarrow 0$. Thus, by [31, Theorem 3.2] there exists a subsequence (x_{n_k}) and a disjoint sequence (d_k) in X such that $\|x_{n_k} - d_k\|_X \rightarrow 0$. Since $\|x_n\|_{L_1} = 1$ and $\|\cdot\|_X \sim \|\cdot\|_{L_1}$ on E , this contradicts that E contains no normalized almost disjoint sequence.

By the beginning part of the proof, we may select an infinite dimensional closed subspace E' of E that does SPR in L_1 . In other words, there exists $\varepsilon > 0$ such that for all $f, g \in E'$ with $\|f\|_{L_1} = \|g\|_{L_1} = 1$ we have

$$\||f| \wedge |g|\|_{L_1} \geq \varepsilon.$$

Since $\|\cdot\|_X \sim \|\cdot\|_{L_1}$ on E , the same is true on E' , so we may view E' as a closed infinite dimensional subspace of X . We claim that it contains no normalized almost disjoint pairs. Indeed, if $f, g \in E'$ with $\|f\|_X = \|g\|_X = 1$, then $\|f\|_{L_1} \sim \|g\|_{L_1} \sim 1$. Now, using that E' does SPR in L_1 and property (ii) of the embedding, we have

$$\| |f| \wedge |g| \|_X \geq \| |f| \wedge |g| \|_{L_1} \gtrsim \varepsilon.$$

Thus, E' contains no normalized almost disjoint pairs when viewed as a subspace of X . It follows that E' does SPR in X . \square

Question 5.2 With Corollary 4.8 and Theorem 5.1 in mind, we ask the following: If a Banach lattice X contains an infinite dimensional dispersed subspace E , does it contain an infinite dimensional SPR subspace? If so, can we construct an infinite dimensional SPR subspace E' with $E' \subseteq E \subseteq X$?

Our next results are motivated by the equivalence between statements (a)-(d) in Theorem 2.1 and the discussion in Remark 2.2. Note that it follows from Theorem 2.1 (a)-(d) that if E is dispersed in $L_p(\mu)$ and $1 \leq q < p$, then E may be viewed as a closed subspace of $L_q(\mu)$, and it is dispersed in $L_q(\mu)$. It is then natural to ask the following question: Let μ be a finite measure and $1 \leq q < p$. Let E be a subspace of $L_p(\mu) \subseteq L_q(\mu)$. What is the relation between E doing SPR in $L_p(\mu)$ versus E doing SPR in $L_q(\mu)$? It is easy to see that if E does SPR in $L_q(\mu)$, then E does SPR in $L_p(\mu)$ if and only if $\|\cdot\|_{L_p} \sim \|\cdot\|_{L_q}$ on E . We will now show that E doing SPR in $L_p(\mu)$ does not imply E does SPR in $L_q(\mu)$, even though the property of being dispersed passes from $L_p(\mu)$ to $L_q(\mu)$.

Theorem 5.3 *For all $2 \leq p < \infty$ there exists a closed subspace $E \subseteq L_p[0, 1]$ such that E does stable phase retrieval in $L_p[0, 1]$ but E fails to do stable phase retrieval in $L_q[0, 1]$ for all $1 \leq q < p$.*

Proof Let $2 \leq p < \infty$. It will be convenient to build the subspace $E \subseteq L_p[0, 2]$ instead of $L_p[0, 1]$. Let $(r_j)_{j=1}^\infty$ be the Rademacher sequence of independent, mean-zero, ± 1 random variables on $[0, 1]$. For all $j \in \mathbb{N}$, we let $g_j = r_j + 2^{j/p} \mathbb{1}_{[1+2^{-j}, 1+2^{-j+1}]}$. Let $E = \overline{\text{span}}_{j \in \mathbb{N}} g_j$.

We first prove for all $1 \leq q < p$ that E fails to do stable phase retrieval in $L_q[0, 2]$. We have for all $j \neq i$ that $\|g_j - g_i\|_{L_q}^q = \|g_j + g_i\|_{L_q}^q \geq 2^{q-1}$. On the other hand, $|r_j| = |r_{j+1}|$ and $\lim \|2^{j/p} \mathbb{1}_{[1+2^{-j}, 1+2^{-j+1}]}\|_{L_q}^q = 0$. Thus, $\lim \| |g_j| - |g_{j+1}| \|_{L_q}^q = 0$. This shows that E fails to do stable phase retrieval in $L_q[0, 2]$.

We now prove that E does stable phase retrieval in $L_p[0, 2]$. Note that by Khintchine's Inequality there exists $B \geq 0$ so that $(\sum |a_j|^2)^{1/2} \leq \|\sum a_j r_j\|_{L_p} \leq B(\sum |a_j|^2)^{1/2}$ for all scalars $(a_j) \in \ell_2$. Thus, we have for all $f = \sum a_j r_j$ and

$x = f + \sum a_j 2^{j/p} \mathbb{1}_{[1+2^{-j}, 1+2^{-j+1})} \in E$ that

$$\begin{aligned} \|f\|_{L_2([0,1])}^p &\leq \|x\|_{L_p([0,2])}^p = \left\| \sum a_j r_j \right\|_{L_p([0,1])}^p + \sum |a_j|^p \\ &\leq B^p \left(\sum |a_j|^2 \right)^{p/2} + \left(\sum |a_j|^2 \right)^{p/2} \\ &= (B^p + 1) \|f\|_{L_2([0,1])}^p. \end{aligned}$$

This computation shows that the map $g_j \mapsto r_j$ extends linearly to a map $E \subseteq L_p[0, 2] \rightarrow L_2[0, 1]$, $x \mapsto f$, establishing an isomorphism between E and a Hilbert space. By Theorem 3.10 and Remark 3.11 it suffices to prove that there exists a constant $\delta > 0$ so that if $x, y \in E$ and $f, g \in L_2[0, 1]$ with $f = \mathbb{1}_{[0,1]}x$ and $g = \mathbb{1}_{[0,1]}y$ such that $\|f\|_{L_2} = 1$, $\|g\|_{L_2} \leq 1$, and $\langle f, g \rangle = 0$ then $\|x - y\|_{L_p} \geq \delta$.

We now claim that it suffices to prove that there exists $\varepsilon > 0$ such that

$$\text{if } \| |x \mathbb{1}_{(1,2)}| - |y \mathbb{1}_{(1,2)}| \|_{L_p} < \varepsilon \text{ then } \| |f|^2 - |g|^2 \|_{L_2}^2 \geq \delta. \quad (5.1)$$

Indeed, as all the L_q norms are equivalent on the span of the Rademacher sequence, there exists a uniform constant $K > 0$ so that the following holds:

$$\begin{aligned} \| |f|^2 - |g|^2 \|_{L_2}^2 &= \int (|f|^2 - |g|^2)^2 \\ &= \int (|f| - |g|)(|f| + |g|)(|f|^2 - |g|^2) \\ &\leq \| |f| - |g| \|_{L_2} \| (|f| + |g|)(|f|^2 - |g|^2) \|_{L_2} \\ &\leq K \| |f| - |g| \|_{L_2} \leq K \| |f| - |g| \|_{L_p}. \end{aligned}$$

Here, the constant K comes from bounding

$$\| (|f| + |g|)(|f|^2 - |g|^2) \|_{L_2} \leq K. \quad (5.2)$$

To get this upper estimate, note that, by Hölder's Inequality,

$$\begin{aligned} \| (|f| + |g|)(|f|^2 - |g|^2) \|_{L_2} &= \| (|f| + |g|)(|f| + |g|)(|f| - |g|) \|_{L_2} \\ &\leq \| |f| + |g| \|_{L_6}^2 \| |f| - |g| \|_{L_6}, \end{aligned}$$

hence, by Triangle Inequality,

$$\| (|f| + |g|)(|f|^2 - |g|^2) \|_{L_2} \leq (\|f\|_{L_6} + \|g\|_{L_6})^3. \quad (5.3)$$

Further, both f and g belong to the span of independent Rademachers, on which all the L_p norms are equivalent (for finite p). Since we know that $\|f\|_{L_2} = 1$ and $\|g\|_{L_2} \leq 1$, this gives a bound for the right-hand side of (5.3), which, in turn, implies (5.2).

To finish the proof of the claim, note that if $\| |x\mathbb{1}_{(1,2)}| - |y\mathbb{1}_{(1,2)}| \|_{L_p} \geq \varepsilon$ then $\| |x| - |y| \|_{L_p} \geq \varepsilon$ and if $\| |x\mathbb{1}_{(1,2)}| - |y\mathbb{1}_{(1,2)}| \|_{L_p} < \varepsilon$ then $\| |x| - |y| \|_{L_p} \geq \delta K^{-1}$.

We now establish (5.1) with $\varepsilon = 1/8$ and $\delta = 1$. Let $x = \sum a_j(r_j + 2^{j/p}\mathbb{1}_{[1+2^{-j}, 1+2^{-j+1}]})$ and $y = \sum b_j(r_j + 2^{j/p}\mathbb{1}_{[1+2^{-j}, 1+2^{-j+1}]})$. We let $f = \sum a_j r_j$ and $g = \sum b_j r_j$ and assume that $\|f\|_{L_2}^2 = \sum |a_j|^2 = 1$, $\|g\|_{L_2}^2 = \sum |b_j|^2 \leq 1$ and $\langle f, g \rangle = \sum a_j b_j = 0$. We may assume that

$$\left(\sum |a_j| - |b_j| \right)^{1/p} = \| |x\mathbb{1}_{(1,2)}| - |y\mathbb{1}_{(1,2)}| \|_{L_p} < \varepsilon = 1/8. \quad (5.4)$$

All that remains is to prove that $\| |f|^2 - |g|^2 \|_{L_2}^2 \geq \delta$. We have from (5.4) that $|a_j| - |b_j| \leq 1/8$ for all $j \in \mathbb{N}$. Hence, $|a_j|^2 - |b_j|^2 \leq 1/4$ for all $j \in \mathbb{N}$ as $|a_j| + |b_j| \leq 2$. As $r_j^2 = \mathbb{1}_{[0,1]}$ for all $j \in \mathbb{N}$, we have that

$$f^2 - g^2 = (f - g)(f + g) = 2 \sum_{j>i} (a_j a_i - b_j b_i) r_j r_i + \sum_{j>i} (a_j^2 - b_j^2) \mathbb{1}. \quad (5.5)$$

Note that (5.5) gives an expansion for $f^2 - g^2$ in terms of the ortho-normal collection of vectors $\{\mathbb{1}_{[0,1]}\} \cup \{r_j r_i\}_{j>i}$. Thus we have that

$$\begin{aligned} 2^{-1} \| |f|^2 - |g|^2 \|_{L_2}^2 &\geq 2 \sum_{j>i} |a_j a_i - b_j b_i|^2 \\ &= \sum_{j \in \mathbb{N}} \sum_{i \in \mathbb{N}} |a_j a_i - b_j b_i|^2 - \sum_{j \in \mathbb{N}} |a_j^2 - b_j^2|^2 \\ &= \sum_{j \in \mathbb{N}} \left(\left(\sum_{i \in \mathbb{N}} |a_j a_i|^2 + |b_j b_i|^2 \right) - \left(2 a_j b_j \sum_{i \in \mathbb{N}} a_i b_i \right) \right) - \sum_{j \in \mathbb{N}} |a_j^2 - b_j^2|^2 \\ &= \sum_{j \in \mathbb{N}} \left(\sum_{i \in \mathbb{N}} |a_j a_i|^2 + |b_j b_i|^2 \right) - \sum_{j \in \mathbb{N}} |a_j^2 - b_j^2|^2 \quad \text{as } \sum a_i b_i = 0 \\ &= \left(\|f\|_{L_2}^4 + \|g\|_{L_2}^4 \right) - \sum_{j \in \mathbb{N}} |a_j^2 - b_j^2|^2 \\ &\geq \left(\|f\|_{L_2}^4 + \|g\|_{L_2}^4 \right) - \frac{1}{4} \sum_{j \in \mathbb{N}} |a_j^2 - b_j^2|^2 \quad \text{as } |a_j^2 - b_j^2| \leq 1/4 \\ &\geq \left(\|f\|_{L_2}^4 + \|g\|_{L_2}^4 \right) - \frac{1}{4} (\|f\|_{L_2}^2 + \|g\|_{L_2}^2) \\ &= \frac{3}{4} + \|g\|_{L_2}^2 \left(\|g\|_{L_2}^2 - \frac{1}{4} \right) \quad \text{as } \|f\|_{L_2} = 1 \\ &\geq \frac{3}{4} - \frac{1}{8} \quad \text{as } \|g\|_{L_2} \leq 1. \end{aligned}$$

Hence, $\| |f|^2 - |g|^2 \|_{L_2}^2 \geq 3/2 - \frac{1}{4} > 1 = \delta$. \square

Example 5.4 In the special case $p = 2$, Theorem 5.3 could have been proven using a result in [24]. Indeed, as above, let (r_j) denote the Rademacher sequence, realized on the interval $[0, 1]$. Define $g_j = r_j + 2^{\frac{j}{2}} \mathbb{1}_{[1+2^{-j}, 1+2^{-j+1}]}$. We can think of the sequence (g_j) as being defined on a finite measure space. Note that $\|2^{\frac{j}{2}} \mathbb{1}_{[1+2^{-j}, 1+2^{-j+1}]}\|_{L_2} = 1$. Hence, for the same reason as in [24], $\overline{\text{span}}\{g_j\}$ does SPR in L_2 . However, recall that the Rademacher sequence does not do phase retrieval; we've also scaled the additional indicator functions to be perturbative in L_1 . Hence, for $i \neq j$ we have $\||g_i| - |g_j|\|_{L_1} = \frac{1}{2^i} + \frac{1}{2^j}$, whereas the other side of the SPR inequality is of order 1. This provides an example of a subspace $E \subseteq L_2(\mu) \subseteq L_1(\mu)$ that does SPR in $L_2(\mu)$ but not in $L_1(\mu)$.

The next result contrasts with Theorem 5.3 by showing, in particular, that if E does SPR in both L_p and L_q then we can both “interpolate” the SPR, and “extrapolate it downward”.

Theorem 5.5 *Suppose μ is a probability measure and $1 \leq q < p < \infty$. Let E be a closed subspace of L_p (real or complex). Assume that $\|\cdot\|_{L_p} \sim \|\cdot\|_{L_q}$ on E , and E does stable phase retrieval in L_q . Then for all $1 \leq r \leq p$, $\|\cdot\|_{L_r} \sim \|\cdot\|_{L_p}$ on E , and E does stable phase retrieval in L_r .*

Proof From the discussion on the Kadec–Pelczynski dichotomy (see Sect. 2.1), we know that, as $E \subseteq L_q(\mu)$ does SPR, then $\|\cdot\|_{L_p} \sim \|\cdot\|_{L_r}$ on E , whenever $r < p$.

Assume first that $q < r \leq p$. Let $C > 0$ so that the L_q and L_p norms are C -equivalent on E , and let $K > 0$ so that E does K -stable phase retrieval in L_q . As $q < r \leq p$ we have for all $f, g \in E$ that

$$\inf_{|\lambda|=1} \|f - \lambda g\|_{L_r} \leq C \inf_{|\lambda|=1} \|f - \lambda g\|_{L_q} \leq CK \||f| - |g|\|_{L_q} \leq CK \||f| - |g|\|_{L_r}. \quad (5.6)$$

Thus, E does stable phase retrieval in L_r .

We now turn to the case $1 \leq r < q$. By the previous argument, E does stable phase retrieval in L_p . Hence, the L_p norm is equivalent to the L_1 norm on E , and hence the L_p norm is equivalent to the L_r norm on E . Let $C > 0$ so that the L_p and L_r norms are C -equivalent on E , and let $K > 0$ so that E does K -stable phase retrieval in L_p . Let θ be the value so that $q^{-1} = \theta r^{-1} + (1 - \theta)p^{-1}$. By Hölder’s inequality, for any $f, g \in E$,

$$\begin{aligned} \||f| - |g|\|_{L_q} &\leq \||f| - |g|\|_{L_r}^\theta (\|f\|_{L_p} + \|g\|_{L_p})^{1-\theta} \\ &\leq C \||f| - |g|\|_{L_r}^\theta (\|f\|_{L_r} + \|g\|_{L_r})^{1-\theta}. \end{aligned} \quad (5.7)$$

Therefore, for any $f, g \in E$, we have

$$\begin{aligned} \inf_{|\lambda|=1} \|f - \lambda g\|_{L_r} &\leq \inf_{|\lambda|=1} \|f - \lambda g\|_{L_q} \leq K \||f| - |g|\|_{L_q} \\ &\leq CK \||f| - |g|\|_{L_r}^\theta (\|f\|_{L_r} + \|g\|_{L_r})^{1-\theta}. \end{aligned}$$

Thus, E does θ -Hölder stable phase retrieval in L_r . By Corollary 3.12, it follows that E does stable phase retrieval in L_r . \square

In Theorem 5.3, we showed that when $2 \leq p < \infty$ an SPR-subspace $E \subseteq L_p[0, 1]$ need not do SPR in $L_q[0, 1]$ for any $1 \leq q < p$. Our next result shows that the case $1 \leq p < 2$ is completely different.

Theorem 5.6 *Let (Ω, μ) be a probability space and let E be a closed infinite dimensional subspace of $L_p(\Omega, \mu)$. Consider the following statements:*

- (i) *E does stable phase retrieval in $L_p(\Omega, \mu)$.*
- (ii) *E does stable phase retrieval in $L_1(\Omega, \mu)$ and $\|\cdot\|_{L_p} \sim \|\cdot\|_{L_1}$ on E .*
- (iii) *There exists $\alpha > 0$ such that for all $x, y \in E$,*

$$\mu(\{t \in \Omega : |x(t)| \geq \alpha \|x\|_{L_p} \text{ and } |y(t)| \geq \alpha \|y\|_{L_p}\}) > \alpha. \quad (5.8)$$

Then for all $1 \leq p < \infty$, (iii) \Leftrightarrow (ii) \Rightarrow (i). Moreover, if $1 \leq p < 2$, all three statements are equivalent.

Proof (iii) \Rightarrow (ii): Note that condition (iii) implies that E contains no normalized $\alpha^{1+\frac{1}{p}}$ -disjoint pairs, when viewed in the L_p norm. Hence, E does SPR in L_p , which implies that $\|\cdot\|_{L_p} \sim \|\cdot\|_{L_1}$ on E . Using this in condition (iii), we conclude that E contains no normalized almost disjoint pairs, when viewed in the L_1 norm, hence does SPR in L_1 .

(ii) \Rightarrow (i): Let $C > 0$ so that $\|x\|_{L_p} \leq C \|x\|_{L_1}$ for all $x \in E$. Let $K > 0$ so that E does K -stable phase retrieval in L_1 . Thus, for all $x, y \in E$ we have that

$$\min_{|\lambda|=1} \|x - \lambda y\|_{L_p} \leq C \min_{|\lambda|=1} \|x - \lambda y\|_{L_1} \leq CK \| |x| - |y| \|_{L_1} \leq CK \| |x| - |y| \|_{L_p}.$$

Thus, E does CK -stable phase retrieval in $L_p(\Omega)$.

(i) \Rightarrow (iii): Let $1 \leq p < 2$ and assume that (i) is true but (iii) is false. We first note that condition (i) implies that $\|\cdot\|_{L_1} \sim \|\cdot\|_{L_p}$ on E . We may choose a sequence of pairs $(x_n, y_n)_{n=1}^\infty$ in E and $\alpha > 0$ such that $\|x_n\|_{L_p} = \|y_n\|_{L_p} = 1$, with

$$\mu(\{t \in \Omega : |x_n| \wedge |y_n| \geq n^{-1}\}) \rightarrow 0, \text{ but } \| |x_n| \wedge |y_n| \|_{L_p} \geq 2\alpha. \quad (5.9)$$

As $(|x_n| \wedge |y_n|)_{n=1}^\infty$ converges in measure to 0 and is uniformly bounded below in L_p norm, after passing to a subsequence we may find a sequence of disjoint subsets $(\Omega_n)_{n=1}^\infty \subseteq \Omega$ such that

$$\| (|x_n| \wedge |y_n|) \mathbb{1}_{\Omega_n^c} \|_{L_p} = \| |x_n| \wedge |y_n| - (|x_n| \wedge |y_n|) \mathbb{1}_{\Omega_n} \|_{L_p} \rightarrow 0. \quad (5.10)$$

Let $\varepsilon_n \searrow 0$ with $\varepsilon_1 < \alpha/2$. After passing to a subsequence, we may assume that $\|x_n \mathbb{1}_{\Omega_n}\|_{L_p} \geq \alpha$ for all $n \in \mathbb{N}$. As $(\Omega_n)_{n=1}^\infty$ is a sequence of disjoint subsets of the probability space (Ω, μ) , we have that $\mu(\Omega_n) \rightarrow 0$. Thus, after passing to a further subsequence we may assume that $\|x_j \mathbb{1}_{\Omega_n}\|_{L_p} < \varepsilon_n$ for all $j < n$. Again,

after passing to a further subsequence we may assume that there exists values $(\beta_n)_{n=1}^\infty$ such that $\lim_{j \rightarrow \infty} \|x_j \mathbb{1}_{\Omega_n}\|_{L_p} = \beta_n$ for all $n \in \mathbb{N}$. Furthermore, we may assume that $\|x_j \mathbb{1}_{\Omega_n}\|_{L_p} < \beta_n + \varepsilon_n/2$ for all $j > n$. As $(\Omega_j)_{j=1}^\infty$ is a sequence of disjoint sets, we have for all $N \in \mathbb{N}$ that

$$\lim_{j \rightarrow \infty} \|x_j\|_{L_p}^p \geq \lim_{j \rightarrow \infty} \sum_{n=1}^N \|x_j \mathbb{1}_{\Omega_n}\|_{L_p}^p = \sum_{n=1}^N \beta_n^p.$$

In particular, we have that $\beta_n \rightarrow 0$. Hence, after passing to a further subsequence of $(x_n)_{n=1}^\infty$ we may assume that $\beta_n < \varepsilon_n/2$ for all $n \in \mathbb{N}$. Thus, $\|x_j \mathbb{1}_{\Omega_n}\|_{L_p} < \varepsilon_n$ for all $j > n$. In summary, we have that for all $n \in \mathbb{N}$, $\|x_n \mathbb{1}_{\Omega_n}\|_{L_p} \geq \alpha$ and for all $j \neq n$, we have $\|x_j \mathbb{1}_{\Omega_n}\|_{L_p} < \varepsilon_n$.

As $\varepsilon_1 < \alpha/2$, we have in particular that $\|x_j - x_n\|_{L_p} \geq \alpha/2$ for all $j \neq n$. We have that $(x_n)_{n=1}^\infty$ is a semi-normalized sequence in a closed subspace of L_p which does not contain ℓ_p . Thus, by [67, Theorem 8], $(x_n)_{n=1}^\infty$ is equivalent to a semi-normalized sequence in $L_{p'}(\nu)$ for some $p < p' \leq 2$ and probability measure ν . We may assume after passing to a subsequence that $(x_n)_{n=1}^\infty$ is weakly convergent in $L_{p'}(\nu)$. Thus, the sequence $(x_{2n} - x_{2n-1})_{n=1}^\infty$ converges weakly to 0 in $L_{p'}(\nu)$. As $L_{p'}(\nu)$ has an unconditional basis, after passing to a further subsequence, we may assume that $(x_{2n} - x_{2n-1})_{n=1}^\infty$ is C -unconditional for some constant C .

As $L_{p'}(\nu)$ has type p' and $(x_{2n} - x_{2n-1})_{n=1}^\infty$ is unconditional, we have that $(x_{2n} - x_{2n-1})_{n=1}^\infty$ is dominated by the unit vector basis of $\ell_{p'}$. We will prove that there exists a constant K so that for all $N \in \mathbb{N}$ there exists $k \in \mathbb{N}$ such that the finite sequence $(x_{2n} - x_{2n-1})_{n=k+1}^{k+N}$ K -dominates the unit vector basis of ℓ_p^N . As $p < p'$, this would contradict that $(x_{2n} - x_{2n-1})_{n=1}^\infty$ is dominated by the unit vector basis of $\ell_{p'}$. Alternatively, one could use that L_p has type p , the uniform containment of ℓ_p^N , and [67, Theorem 13] to get that E contains a subspace isomorphic to ℓ_p , which, in view of Theorem 2.1, contradicts that E does stable phase retrieval in L_p .

Let $N \in \mathbb{N}$ and $\varepsilon > 0$. Let $k \in \mathbb{N}$ be large enough so that $2\varepsilon_k N < 2^{-1}\alpha$. Let $(a_j)_{j=k+1}^{k+N}$ be a sequence of scalars. We have that

$$\begin{aligned} & \left\| \sum_{j=k+1}^{k+N} a_j (x_{2j} - x_{2j-1}) \right\|_{L_p}^p \geq \sum_{n=k+1}^{k+N} \left\| \sum_{j=k+1}^{k+N} a_j (x_{2j} - x_{2j-1}) \right\|_{L_p(\Omega_{2n})}^p \\ & \geq \sum_{n=k+1}^{k+N} \left(2^{1-p} \|a_n (x_{2n} - x_{2n-1})\|_{L_p(\Omega_{2n})}^p - \left\| \sum_{j \neq n} a_j (x_{2j} - x_{2j-1}) \right\|_{L_p(\Omega_{2n})}^p \right) \\ & \geq \sum_{n=k+1}^{k+N} \left(2^{1-p} \alpha^p |a_n|^p - 2^p \varepsilon_k^p \left(\sum_{j \neq n} |a_j|^p \right) \right) \\ & \geq \sum_{n=k+1}^{k+N} \left(2^{1-p} \alpha^p |a_n|^p - 2^p \varepsilon_k^p N^{p-1} \sum_{j \neq n} |a_j|^p \right) \end{aligned}$$

$$\begin{aligned} &\geq \left(2^{1-p}\alpha^p - 2^p\varepsilon_k^p N^p\right) \sum_{n=k+1}^{k+N} |a_n|^p \\ &\geq 2^{-p}\alpha^p \sum_{n=k+1}^{k+N} |a_n|^p. \end{aligned}$$

Now that we have established that all three statements in Theorem 5.6 are equivalent for $1 \leq p < 2$, we can show the implication (ii) \Rightarrow (iii) for $1 \leq p < \infty$. Indeed, we assume (ii) holds. Since E does SPR in L_1 , by (ii) \Rightarrow (iii) for $p = 1$, we deduce that there exists $\alpha > 0$ such that for all $x, y \in E$,

$$\mu(\{t \in \Omega : |x(t)| \geq \alpha\|x\|_{L_1} \text{ and } |y(t)| \geq \alpha\|y\|_{L_1}\}) > \alpha. \quad (5.11)$$

Now we use the second assumption of (ii) to replace the L_1 norm with the L_p norm in (5.11). \square

6 $C(K)$ -spaces with SPR subspaces

Throughout this section, subspaces are assumed to be closed and infinite dimensional, unless otherwise mentioned. Recall that a non-empty compact Hausdorff space is called *perfect* if it has no isolated points, and *scattered* (or *dispersed*) if it contains no perfect subsets. For a compact Hausdorff space K , we define its *Cantor–Bendixson derivative* K' to be the set of all non-isolated points of K . Clearly K' is closed, and $K = K'$ iff K is perfect; otherwise, K' is a proper subset of K . Also, if K contains a perfect set S , then S lies inside of K' as well.

Theorem 6.1 *Suppose K is a compact Hausdorff space. Then $C(K)$ has an SPR subspace if and only if K' is infinite.*

The proof depends on an auxiliary result, strengthening Remark 4.6.

Proposition 6.2 *Every separable Banach space embeds isometrically into $C(\Delta)$, and into $C[0, 1]$, as a 10-SPR subspace (here Δ is the Cantor set).*

Proof Fix a separable Banach space E . Let K be the unit ball of E^* , with its weak* topology. By Lemma 4.5 and Remark 4.6, the natural isometric embedding $j : E \rightarrow C(K)$ (taking e into the function $K \rightarrow \mathbb{R} : e^* \mapsto e^*(e)$) is such that $\| |jx| \wedge |jy| \| \geq 1/5$ whenever $\|x\| = 1 = \|y\|$. As K is compact and metrizable, there exists a continuous surjection $\Delta \rightarrow K$ [53, Theorem 4.18]; this generates a lattice isometric embedding of $C(K)$ into $C(\Delta)$, hence one can find an isometric copy of $E \subseteq C(\Delta)$ so that $\| |x| \wedge |y| \| \geq 1/5$ whenever x, y are norm one elements of E .

View Δ as a subset of $[0, 1]$. Then there exists a positive unital isometric extension operator $T : C(\Delta) \rightarrow C[0, 1]$ —that is, for $f \in C(\Delta)$, $Tf|_{\Delta} = f$; $T1 = 1$; $\|T\| = 1$; and $Tf \geq 0$ whenever $f \geq 0$. The “standard” construction of T involves piecewise-affine extensions of functions from Δ to $[0, 1]$; for a more general approach, see the proof of [4, Theorem 4.4.4]. One observes that $\|Tx| \wedge |Ty\| \| \geq \| |x| \wedge |y| \|$,

hence, if $E \subseteq C(\Delta)$ has the property described in the preceding paragraph, then $\| |Tx| \wedge |Ty| \| \geq 1/5$ whenever $x, y \in E$ have norm 1.

By Theorem 3.4, the copies of E in $C(\Delta)$ and $C[0, 1]$ described above do 10-SPR. \square

The next result is standard topological fare (cf. [63, Theorem 29.2]).

Lemma 6.3 *Suppose K is a compact Hausdorff space, and $t \in U \subseteq K$, where U is an open set. Then there exists an open set V so that $t \in V \subseteq \overline{V} \subseteq U$.*

Proof of Theorem 6.1 Suppose first that K' is finite (in this case, K must be scattered). To show that any subspace $E \subseteq C(K)$ fails SPR, consider $C_0(K, K') = \{f \in C(K) : f|_{K'} = 0\}$. Then $\dim C(K)/C_0(K, K') = |K'| < \infty$, hence $E \cap C_0(K, K')$ is infinite dimensional as well. It suffices therefore to show that every infinite dimensional subspace of $C_0(K, K')$ fails SPR.

Note that, in the case of finite K' , $C_0(K, K')$ can be identified with $c_0(K \setminus K')$ as a Banach lattice. Indeed, any $f \in c_0(K \setminus K')$ is continuous on $K \setminus K'$, since this set consists of isolated points only. Extend f to a function $\tilde{f} : K \rightarrow \mathbb{R}$ with $\tilde{f}|_{K'} = 0$, $\tilde{f}|_{K \setminus K'} = f$. Note that for any $c > 0$, the set $\{t \in K \setminus K' : |f(t)| \geq c\} = \{t \in K : |f(t)| \geq c\}$ is finite, hence closed; consequently, $\{t \in K : |\tilde{f}(t)| < c\}$ is an open neighborhood of any element of K' . From this it follows that f is continuous.

On the other hand, pick $h \in C_0(K, K')$. We claim that $h|_{K \setminus K'} \in c_0(K \setminus K')$ —that is, $\{t \in K \setminus K' : |h(t)| > c\}$ is finite for any $c > 0$. Suppose, for the sake of contradiction, that this set is infinite for some c . By the compactness of K , this set must have an accumulation point, which must lie in K' . This, however, contradicts the continuity of h .

A “gliding hump” argument shows that no subspace of $c_0(K \setminus K')$ does SPR. From this we conclude that no subspace of $C(K)$ does SPR if K' is finite.

Now suppose K contains a perfect set. By [58, Theorem 2, p. 29], there exists a continuous surjection $\phi : K \rightarrow [0, 1]$. This map generates a lattice isometric embedding $T : C[0, 1] \rightarrow C(K) : f \mapsto f \circ \phi$. However, $C[0, 1]$ contains SPR subspaces, by Proposition 6.2.

It remains to prove that $C(K)$ contains an SPR copy of c_0 when K is scattered, and K' is infinite. Note first that $K' \setminus K''$ must be infinite. Indeed, otherwise any point of $K'' = K' \setminus (K' \setminus K'')$ will be an accumulation point of the same set, and K'' will be perfect, which is impossible.

Observe also that any $t \in K' \setminus K''$ is an accumulation point of $K \setminus K'$. Indeed, suppose otherwise, for the sake of contradiction. Then t has an open neighborhood W , disjoint from $K \setminus K'$. If U is another open neighborhood of t , then so is $U \cap W$. As t is an accumulation point of K , $U \cap W$ must meet K , hence also K' . This implies $t \in K''$, providing us with the desired contradiction.

Find distinct points $t_1, t_2, \dots \in K' \setminus K''$. For each i find an open set $A_i \ni t_i$ so that $t_j \notin A_i$ for $j \neq i$. Lemma 6.3 permits us to find an open set U_i so that $t_i \in U_i \subseteq \overline{U_i} \subseteq A_i$. Replacing U_2 by $U_2 \setminus \overline{U_1}$, U_3 by $U_3 \setminus \overline{U_1 \cup U_2}$, and so on, we can assume that the sets U_i are disjoint. Lemma 6.3 guarantees the existence of open sets V_i so that, for every i , $t_i \in V_i \subseteq \overline{V_i} \subseteq U_i$.

As noted above, each t_i is an accumulation point of $K \setminus K'$. Therefore, we can find distinct points $(s_{ji})_{j=1}^{\infty} \subseteq (K \setminus K') \cap V_i$. For each n , let S_n be the closure of $\{s_{j,2n} : j \in \mathbb{N}\}$ (note $S_n \subseteq \overline{V_{2n}} \subseteq U_{2n}$). Note that there exists $x^{(n)} \in C(K)$ such that:

- (i) $0 \leq x^{(n)} \leq 1$ everywhere.
- (ii) $x^{(n)}|_{S_n} = 1/2$.
- (iii) $x^{(n)}(s_{1,2n-1}) = 1$.
- (iv) $x^{(n)}(s_{n,2i}) = 1/2$ for $1 \leq i \leq n-1$.
- (v) $x^{(n)} = 0$ on $(K \setminus U_{2n}) \setminus \{s_{1,2n-1}, s_{n,2}, s_{n,4}, \dots, s_{n,2n-2}\}$.

To construct such an $x^{(n)}$, recall that $s_{1,2n-1}, s_{n,2}, s_{n,4}, \dots, s_{n,2n-2}$ are isolated points of K , hence the function g , defined by $g(s_{1,2n-1}) = 1$, $g(s_{n,2i}) = 1/2$ for $1 \leq i \leq n-1$, and $g = 0$ everywhere else, is continuous. Further, by Urysohn's Lemma, there exists $h \in C(K)$ so that $0 \leq h \leq 1/2 = h|_{S_n}$, vanishing outside of U_{2n} . Then $x^{(n)} = g + h$ has the desired properties.

We claim that $(x^{(n)})$ is equivalent to the standard c_0 -basis. Indeed, suppose $(\alpha_n) \in c_{00}$, with $\vee_n |\alpha_n| = 1$. We need to show $\|\sum_n \alpha_n x^{(n)}\| = 1$. The lower estimate on the norm is clear, since $x = \sum_n \alpha_n x^{(n)}$ attains the value of α_n at $s_{1,2n-1}$.

For an upper estimate, note that x vanishes outside of $\cup_m U_m$, and on U_m if m is large enough. If m is odd ($m = 2n-1$), then the only point of U_m where x does not vanish is $s_{1,2n-1}$, which we have already discussed. If m is even ($m = 2n$), then $|x| \leq 1/2$ except for the points $s_{i,2n}$ ($i > n$); at these points, x equals $(\alpha_n + \alpha_i)/2$, which has absolute value not exceeding 1.

It remains to show that $E = \text{span}[x^{(n)} : n \in \mathbb{N}]$ does SPR. In light of Proposition 3.4, it suffices to prove that $\| |x| \wedge |y| \| \geq 1/3$ for any norm one $x, y \in E$. Write $x = \sum_n \alpha_n x^{(n)}$ and $y = \sum_n \beta_n x^{(n)}$. Find n and m so that $|\alpha_n| = 1 = |\beta_m|$. If $n = m$, then both $|x|$ and $|y|$ equal 1 at $s_{1,2n-1}$, so $\| |x| \wedge |y| \| = 1$.

Otherwise, assume, by relabeling, that $n < m$. If $|\alpha_m| \geq 1/3$, then

$$\| |x| \wedge |y| \| \geq |x(s_{1,2m-1})| \wedge |y(s_{1,2m-1})| = |\alpha_m| \wedge |\beta_m| \geq \frac{1}{3}.$$

The case of $|\beta_m| \geq 1/3$ is treated similarly. If $|\alpha_m|, |\beta_n| < 1/3$, then $|x(s_{m,2n})| = |\alpha_n + \alpha_m|/2 > 1/3$, and similarly, $|y(s_{m,2n})| > 1/3$, which again gives us $\| |x| \wedge |y| \| \geq 1/3$. \square

Question 6.4 The proof of Theorem 6.1 shows that K' is infinite iff $C(K)$ contains an SPR copy of c_0 . If K is “large” enough (in terms of the smallest ordinal α for which $K^{(\alpha)}$ is finite), what SPR subspaces (other than c_0) does $C(K)$ have? Note that c_0 is isomorphic to $c = C[0, \omega]$ (ω is the first infinite ordinal). If $K^{(\alpha)}$ is infinite, does $C(K)$ contain an SPR copy of $C[0, \omega^\alpha]$?

In the spirit of Proposition 4.1, it is natural to ask which (isometric) subspaces of $C(K)$ are necessarily SPR. Below we give a “very local” condition on a Banach space E (finite or infinite dimensional) which guarantees that any isometric embedding of E into $C(K)$ has SPR.

Recall (see [48]) that a Banach space E is called *uniformly non-square* if there exists $\varepsilon > 0$ so that, for any norm one $f, g \in E$ we have $\min\{\|f + g\|, \|f - g\|\} < 2 - \varepsilon$.

Note that E fails to be uniformly non-square iff for every $\varepsilon > 0$ there exist norm one $f, g \in E$ so that $\|f + g\|, \|f - g\| > 2 - \varepsilon$. In the real case, this means that E contains ℓ_1^2 (equivalently, ℓ_∞^2) with arbitrarily small distortion. This is incompatible with uniform convexity or uniform smoothness.

Proposition 6.5 *Any uniformly non-square subspace of $C(K)$ does SPR.*

Proof Suppose E is a non-SPR subspace of $C(K)$; we shall show that it fails to be uniformly non-square. To this end, fix $\varepsilon \in (0, 1/2)$; by Theorem 3.4, there exist norm one $f, g \in E$ with $\| |f| \wedge |g| \| < \varepsilon$. Pointwise evaluation shows that

$$|f| \vee |g| + |f| \wedge |g| \geq |f + g| \geq |f| \vee |g| - |f| \wedge |g|.$$

As the ambient lattice is an M-space, we have $\| |f| \vee |g| \| = 1$, hence

$$1 - \varepsilon < \| |f| \vee |g| \| - \| |f| \wedge |g| \| \leq \|f + g\| \leq \| |f| \vee |g| \| + \| |f| \wedge |g| \| < 1 + \varepsilon.$$

Replacing g by $-g$, we conclude that $1 - \varepsilon < \|f - g\| < 1 + \varepsilon$.

Let $u = (f + g)/\|f + g\|$ and $v = (f - g)/\|f - g\|$. Then

$$\|u - (f + g)\| = |1 - \|f + g\|\| < \varepsilon,$$

and similarly, $\|v - (f - g)\| < \varepsilon$. Then

$$\|u + v\| \geq \|(f + g) + (f - g)\| - \|u - (f + g)\| - \|v - (f - g)\| > 2 - 2\varepsilon,$$

and likewise, $\|u - v\| > 2 - 2\varepsilon$. As ε is arbitrary, E fails to be uniformly non-square. \square

For infinite dimensional subspaces, Proposition 6.5 is only meaningful when K is not scattered. Indeed, if K is scattered, then $C(K)$ is c_0 -saturated [36, Theorem 14.26], hence any infinite dimensional subspace of $C(K)$ contains an almost isometric copy of c_0 [60, Proposition 2.e.3]. In particular, such subspaces contain almost isometric copies of ℓ_1^2 , hence they cannot be uniformly non-square.

In light of Proposition 6.5, we ask:

Question 6.6 Which Banach spaces E isometrically embed into $C(K)$ in a non-SPR way?

Note that containing an isometric copy of ℓ_∞^2 (and consequently, failing to be uniformly non-square) does not automatically guarantee the existence of a non-SPR embedding into $C(K)$ (in this sense, the converse to Proposition 6.5 fails). In the following example we look at isometric embeddings only; one can modify this example to allow for sufficiently small distortions.

Proposition 6.7 *There exists a 3-dimensional space E , containing ℓ_∞^2 isometrically (and consequently, failing to be uniformly non-square), so that, if K is a Hausdorff compact, and $J : E \rightarrow C(K)$ is an isometric embedding, then $\| |Jx| \wedge |Jy| \| \geq 1/3$ for any norm one $x, y \in E$.*

The following lemma is needed for the proof, and may be of interest in its own right.

Lemma 6.8 *Suppose K is a Hausdorff compact, E is a Banach space, and $J : E \rightarrow C(K)$ is an isometric embedding. Denote by \mathcal{F} the set of all extreme points of the unit ball of E^* . Then, for any $x, y \in E$, $\| |Jx| \wedge |Jy| \| \geq \sup_{e^* \in \mathcal{F}} |e^*(x)| \wedge |e^*(y)|$.*

Proof Standard duality considerations tell us that $J^* : M(K) \rightarrow E^*$ ($M(K)$ stands for the space of Radon measures on K) is a strict quotient—that is, for any $e^* \in E^*$ there exists $\mu \in M(K)$ so that $\|\mu\| = \|e^*\|$ and $J^*\mu = e^*$. Further, we claim that, for any $e^* \in \mathcal{F}$, there exists $t \in K$ so that $J^*\delta_t \in \{e^*, -e^*\}$. Indeed, the set $S = \{\mu \in M(K) : \|\mu\| \leq 1, J^*\mu = e^*\}$ is weak*-compact, hence it is the weak*-closure of the convex hull of its extreme points. We claim that any such extreme point is also an extreme point of $\{\mu \in M(K) : \|\mu\| \leq 1\}$. Indeed, suppose $\mu = (\mu_1 + \mu_2)/2$, with $\|\mu_1\|, \|\mu_2\| \leq 1$. Then $e^* = (J^*\mu_1 + J^*\mu_2)/2$, which guarantees that $e^* = J^*\mu_1 = J^*\mu_2$, so $\mu_1, \mu_2 \in S$, and therefore, they coincide with μ .

To finish the proof, recall that the extreme points of $\{\mu \in M(K) : \|\mu\| \leq 1\}$ are point evaluations and their opposites. \square

Proof of Proposition 6.7 To obtain E , equip \mathbb{R}^3 with the norm

$$\|(x, y, z)\| = \max \left\{ |x|, |y|, \frac{1}{2}(|x| + |y| + |z|) \right\}. \quad (6.1)$$

Clearly $\{(x_1, x_2, 0) : x_1, x_2 \in \mathbb{R}\}$ gives us an isometric copy of ℓ_∞^2 in E . Note that the unit ball of E^* is a polyhedron with vertices $(\pm 1, 0, 0)$, $(0, \pm 1, 0)$, and $(\pm 1/2, \pm 1/2, \pm 1/2)$; we denote this set of vertices by \mathcal{F} . In light of Lemma 6.8, we have to show that, for any norm one $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$ in E , there exists $e^* \in \mathcal{F}$ so that $|e^*(x)| \wedge |e^*(y)| \geq 1/3$.

In searching for e^* , we deal with several cases separately. Note first that, if $|x_1| \wedge |y_1| \geq 1/3$, then $e^* = (1, 0, 0)$ has the desired properties. The case of $|x_2| \wedge |y_2| \geq 1/3$ is treated similarly. Henceforth we assume $|x_1| \wedge |y_1|, |x_2| \wedge |y_2| < 1/3$. In light of (6.1), we need to consider three cases:

- (i) $|x_1| = 1 = |y_2|$ or $|x_2| = 1 = |y_1|$.
- (ii) Either $|x_1| \vee |x_2| = 1$ and $|y_1| + |y_2| + |y_3| = 2$, or $|y_1| \vee |y_2| = 1$ and $|x_1| + |x_2| + |x_3| = 2$.
- (iii) $|x_1| + |x_2| + |x_3| = 2 = |y_1| + |y_2| + |y_3|$.

In all the three cases, we look for $e^* = (\varepsilon_1, \varepsilon_2, \varepsilon_3)/2$, with $\varepsilon_1, \varepsilon_2, \varepsilon_3 = \pm 1$ selected appropriately.

Case (i). We shall assume $x_1 = 1 = y_2$, as other permutations of indices and choices of sign are handled similarly. Select $\varepsilon_1 = 1$, and take ε_3 so that $\varepsilon_3 x_3 \geq 0$. Pick $\varepsilon_2 = 1$ if $\varepsilon_1 y_1 + \varepsilon_3 y_3 \geq 0$ and $\varepsilon_2 = -1$ otherwise. Then $|x_2| < 1/3$, hence

$$e^*(x) = \frac{1}{2}(\varepsilon_1 + \varepsilon_2 x_2 + \varepsilon_3 x_3) \geq \frac{1 - |x_2|}{2} > \frac{1 - 1/3}{2} = \frac{1}{3}.$$

Further,

$$|e^*(y)| = \frac{|\varepsilon_1 y_1 + \varepsilon_2 + \varepsilon_3 y_3|}{2} \geq \frac{1}{2}.$$

Case (ii). We deal with $x_1 = 1$ (and consequently, $|y_1| < 1/3$) and $|y_1| + |y_2| + |y_3| = 2$, as other possible settings can be treated similarly. Let $\varepsilon_1 = 1$. If $|x_2| < 1/3$, select ε_3 so that $\varepsilon_3 x_3 \geq 0$. Pick ε_2 so that $\varepsilon_2 y_2$ and $\varepsilon_3 y_3$ have the same sign. Then

$$|e^*(x)| \geq \frac{1 + |x_3| - |x_2|}{2} \geq \frac{1 - |x_2|}{2} \geq \frac{1 - 1/3}{2} = \frac{1}{3},$$

and

$$|e^*(y)| \geq \frac{|y_2| + |y_3| - |y_1|}{2} = \frac{2 - 2|y_1|}{2} \geq \frac{2 - 2 \cdot 1/3}{2} = \frac{2}{3}.$$

Suppose, conversely, that $|x_2| \geq 1/3$, hence $|y_2| < 1/3$. Let $\varepsilon_2 = \text{sign } x_2$. Select ε_3 so that $\varepsilon_1 y_1$ and $\varepsilon_3 y_3$ are of the same sign. Then $|x_3| \leq 2 - (1 + |x_2|) = 1 - |x_2|$, hence

$$|e^*(x)| \geq \frac{1 + |x_2| - |x_3|}{2} \geq \frac{2|x_2|}{2} \geq \frac{1}{3}.$$

On the other hand, $2 - |y_2| = |y_1| + |y_3|$ and

$$|e^*(y)| \geq \frac{|y_1| + |y_3| - |y_2|}{2} = \frac{2 - 2|y_2|}{2} \geq \frac{2 - 2 \cdot 1/3}{2} = \frac{2}{3}.$$

Case (iii). If $|x_1|, |x_2| < 1/3$, let $\varepsilon_3 = \text{sign } x_3$, and select $\varepsilon_1, \varepsilon_2$ so that both $\varepsilon_1 y_1$ and $\varepsilon_2 y_2$ have the same sign as $\varepsilon_3 y_3$. Then

$$|e^*(x)| \geq \frac{|x_3| - |x_1| - |x_2|}{2} = \frac{2 - 2(|x_1| + |x_2|)}{2} \geq \frac{2 - 4 \cdot 1/3}{2} = \frac{1}{3},$$

and

$$|e^*(y)| = \frac{|y_1| + |y_2| + |y_3|}{2} = 1.$$

The case of $|y_1|, |y_2| < 1/3$ is handled similarly.

Now suppose neither of the above holds. Up to a permutation of indices, we assume that $|x_1| \geq 1/3$ (hence $|y_1| < 1/3$), and $|y_2| \geq 1/3$ (hence $|x_2| < 1/3$). Then let $\varepsilon_1 = \text{sign } x_1$ and $\varepsilon_3 = \text{sign } x_3$. Pick ε_2 so that $\text{sign } \varepsilon_2 y_2 = \text{sign } \varepsilon_3 y_3$, then

$$|e^*(x)| \geq \frac{|x_1| + |x_3| - |x_2|}{2} = \frac{2 - 2|x_2|}{2} \geq \frac{2 - 2 \cdot 1/3}{2} = \frac{2}{3},$$

and likewise,

$$|e^*(y)| \geq \frac{|y_2| + |y_3| - |y_1|}{2} \geq \frac{2}{3}.$$

□

7 Open problems

We now list some open questions and directions for further research. The reader can find additional questions embedded throughout the paper.

Question 7.1 (Classification of SPR subspaces): Given a Banach lattice X , it is of interest to classify the closed subspaces of X that do SPR. This question, of course, can be interpreted in various ways. Possibly the crudest of these is to classify the closed subspaces of X doing SPR up to Banach space isomorphism. One can then refine this classification by tracking the optimal SPR and isomorphism constants. On the other hand, one can ask about the “structure” of the collection of SPR subspaces of X . For example, whether certain natural candidates do SPR, or whether they have a further subspace/perturbation which does SPR. Compare with [24, Theorem 1.1], which, within a restricted class of subspaces of $L_2(\mathbb{R})$, is able to classify those that do SPR.

Phase retrieval is most often studied in terms of recovering a function f from $|Tf|$ where T is a linear transformation, such as the Fourier transform or Gabor transform. However, any use of phase retrieval in applications requires sampling at only finitely many points. Gabor frames are constructed by sampling the short-time Fourier transform at a lattice; however, any frame constructed by sampling the Gabor transform at an integer lattice cannot do phase retrieval. There has been significant recent interest in determining which sampling points allow for constructing frames which do phase retrieval [3, 43, 44].

The problem of sampling continuous frames which do stable phase retrieval to obtain frames which do stable phase retrieval was introduced in [37] and was shown to be connected to important integral norm discretization problems in approximation theory (such as in [29, 30, 52, 59]). In [38] it is proven that if $(x_t)_{t \in \Omega}$ is a bounded continuous frame of a separable Hilbert space H then there exist sampling points $(t_j)_{j \in J}$ in Ω such that $(x_{t_j})_{j \in J}$ is a frame of H . The corresponding quantitative and finite dimensional theorem in [59] gives that for each $\beta > 0$ there are universal constants $B > A > 0$ so that if $(x_t)_{t \in \Omega}$ is a continuous Parseval frame of an n -dimensional Hilbert space H and $\|x_t\| \leq \beta n^{1/2}$ for all $t \in \Omega$ then there exists m on the order of n sampling points $(t_j)_{j=1}^m$ in Ω so that $(m^{-1/2}x_{t_j})_{j=1}^m$ is a frame of H with lower frame bound A and upper frame bound B . The proof of the above theorem relies on the celebrated solution to the Kadison–Singer Problem and its connection to frame partitioning [62, 64]. It is natural to consider if this discretization theorem holds as well for stable phase retrieval, and the following question is stated in [37].

Question 7.2 Let $C, \beta > 0$. Do there exist constants $D, \kappa > 0$ so that for all $n \in \mathbb{N}$ there exists $m \leq Dn$ so that the following is true: Let H be an n -dimensional Hilbert space, (Ω, μ) a probability space, and $(x_t)_{t \in \Omega}$ a continuous Parseval frame of H which

does C -stable phase retrieval such that $\|x_t\| \leq \beta\sqrt{n}$ for all $t \in \Omega$. Then there exists a sequence of sampling points $(t_j)_{j=1}^m \subseteq \Omega$ such that $(m^{-1/2}x_{t_j})_{j=1}^m$ is a frame of H which does κ -stable phase retrieval.

Note that if $(x_t)_{t \in \Omega}$ is a continuous Parseval frame of H over a probability space Ω then the analysis operator $\Theta(x) = (\langle x, x_t \rangle)_{t \in \Omega}$ is an isometric embedding of H into $L_2(\Omega)$. We have that the continuous frame $(x_t)_{t \in \Omega}$ does C -stable phase retrieval if and only if the range of the analysis operator $\Theta(H)$ does C -stable phase retrieval as a subspace of $L_2(\Omega)$. In Theorem 5.3 we prove that there exists a subspace $E \subseteq L_2(\Omega)$ such that E does stable phase retrieval as a subspace of $L_2(\Omega)$ but that E does not do stable phase retrieval as a subspace of $L_1(\Omega)$. As shown by Theorem 5.6, doing stable phase retrieval in $L_1(\Omega)$ gives a lot of useful additional structure. This motivates the following problem.

Question 7.3 Let $C, \beta > 0$. Do there exist constants $D, \kappa > 0$ so that for all $n \in \mathbb{N}$ there exists $m \leq Dn$ so that the following is true: Let H be an n -dimensional Hilbert space, (Ω, μ) a probability space, and $(x_t)_{t \in \Omega}$ a continuous Parseval frame of H with analysis operator Θ such that $\Theta(H)$ does C -stable phase retrieval as a subspace of $L_1(\Omega)$ and as a subspace of $L_2(\Omega)$, and $\|x_t\| \leq \beta\sqrt{n}$ for all $t \in \Omega$. Then there exists a sequence of sampling points $(t_j)_{j=1}^m \subseteq \Omega$ such that $(m^{-1/2}x_{t_j})_{j=1}^m$ is a frame of H which does κ -stable phase retrieval.

The previous two questions on constructing frames by sampling continuous frames relate to discretizing the L_2 -norm on a subspace of $L_2(\Omega)$. There is significant interest in approximation theory on discretizing the L_p -norm on finite dimensional subspaces of $L_p(\Omega)$ which are called Marcinkiewicz-type discretization problems [29, 30, 52, 59]. For $p \neq 2$, it is too much to ask for the number of sampling points to be on the order of the dimension of the subspace. This leads to the following general problem on discretizing stable phase retrieval.

Question 7.4 Let $E \subseteq L_p(\Omega)$ be an n -dimensional subspace for some $1 \leq p < \infty$ and probability space Ω . Let $C > 0$ and let $f : \mathbb{N} \rightarrow \mathbb{N}$ be strictly increasing. What properties on E imply that there exists $m \leq f(n)$ and sampling points $(t_j)_{j=1}^m \subseteq \Omega$ so that the subspace $\{(m^{-1/p}x(t_j))_{j=1}^m : x \in E\}$ does C -stable phase retrieval in ℓ_p^m ? What properties on E imply that there exists $m \leq f(n)$, sampling points $(t_j)_{j=1}^m \subseteq \Omega$, and weights $(w_j)_{j=1}^m$ with $\sum_{j=1}^m |w_j|^p = 1$ so that the subspace $\{(w_jx(t_j))_{j=1}^m : x \in E\}$ does C -stable phase retrieval in ℓ_p^m ?

Data availability No data accompanies this paper.

Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

1. Abramovich, Y.A., Aliprantis, C.D.: An invitation to operator theory. Graduate Studies in Mathematics, vol. 50. American Mathematical Society, Providence (2002)
2. Alaiifari, R., Grohs, P.: Phase retrieval in the general setting of continuous frames for Banach spaces. *SIAM J. Math. Anal.* **49**(3), 1895–1911 (2017)
3. Alaiifari, R., Grohs, P.: Gabor phase retrieval is severely ill-posed. *Appl. Comput. Harmon. Anal.* **50**, 401–419 (2021)
4. Albiac, F., Kalton, N.J.: Topics in Banach space theory. Graduate Texts in Mathematics, vol. 233, 2nd edn. Springer, Berlin (2016). With a foreword by Gilles Godefroy
5. Alharbi, W., Alshabhi, S., Freeman, D., Ghoreishi, D.: Locality and stability for phase retrieval. *Sampling Theory, Signal Processing, and Data Analysis* (2023) **(to appear)**
6. Aliprantis, C.D., Burkinshaw, O.: Locally solid Riesz spaces with applications to economics, Mathematical Surveys and Monographs, vol. 105, 2nd edn. American Mathematical Society, Providence (2003)
7. Aliprantis, C.D., Burkinshaw, O.: Positive Operators. Springer, Dordrecht (2006). Reprint of the 1985 original
8. Andreys, S., Jaming, P.: Zak transform and non-uniqueness in an extension of Pauli's phase retrieval problem. *Anal. Math.* **42**(3), 185–201 (2016)
9. Astashkin, S.V.: $\Lambda(p)$ -spaces. *J. Funct. Anal.* **266**(8), 5174–5198 (2014)
10. Avilés, A., Tradacete, P.: Amalgamation and injectivity in Banach lattices. *Int. Math. Res. Not. IMRN* **2**, 956–997 (2023)
11. Bachelis, G.F., Ebenstein, S.E.: On $\Lambda(p)$ sets. *Pac. J. Math.* **54**(1), 35–38 (1974)
12. Balan, R.: Stability of phase retrievable frames. SPIE Optical Engineering + Applications, International Society for Optics and Photonics (2013)
13. Balan, R., Dock, C.B.: Lipschitz analysis of generalized phase retrievable matrix frames. *SIAM J. Matrix Anal. Appl.* **43**(3), 1518–1571 (2022)
14. Bandeira, A.S., Cahill, J., Mixon, D.G., Nelson, A.A.: Saving phase: injectivity and stability for phase retrieval. *Appl. Comput. Harmon. Anal.* **37**(1), 106–125 (2014)
15. Bilokopytov, E.: Disjointly non-singular operators on order continuous Banach lattices complement the unbounded norm topology. *J. Math. Anal. Appl.* **506**(1), 125556 (2022)
16. Bilokopytov, E.: Private communication (2023)
17. Borwein, J.M., Yost, D.T.: Absolute norms on vector lattices. *Proc. Edinb. Math. Soc.* (2) **27**(2), 215–222 (1984)
18. Botelho-Andrade, S., Casazza, P.G., Ghoreishi, D., Jose, S., Tremain, J.C.: Weak phase retrieval. *Compressed Sensing and its Applications*, pp. 221–234 (2017)
19. Bourgain, J.: Bounded orthogonal systems and the $\Lambda(p)$ -set problem. *Acta Math.* **162**(3–4), 227–245 (1989)
20. Bourgain, J.: Λ_p -sets in analysis: results, problems and related aspects. In: *Handbook of the Geometry of Banach Spaces*, vol. I, pp. 195–232. North-Holland, Amsterdam (2001)
21. Braga, B.M.: On the Hausdorff distance and some openings between Banach spaces as Borel functions. *Quaest. Math.* **38**(3), 403–411 (2015)
22. Cahill, J., Casazza, P.G., Peterson, J., Woodland, L.: Using projections for phase retrieval. In: *Proc. SPIE 8858, Wavelets and Sparsity XV* (2013)
23. Cahill, J., Casazza, P.G., Daubechies, I.: Phase retrieval in infinite-dimensional Hilbert spaces. *Trans. Amer. Math. Soc. Ser. B* **3**, 63–76 (2016)
24. Calderbank, R., Daubechies, I., Freeman, D., Freeman, N.: Stable phase retrieval for infinite dimensional subspaces of $L_2(\mathbb{R})$. arXiv preprint [arXiv:2203.03135](https://arxiv.org/abs/2203.03135) (2022)
25. Candès, E.J., Li, X.: Solving quadratic equations via PhaseLift when there are about as many equations as unknowns. *Found. Comput. Math.* **14**(5), 1017–1026 (2014)
26. Candès, E.J., Strohmer, T., Voroninski, V.: PhaseLift: exact and stable signal recovery from magnitude measurements via convex programming. *Commun. Pure Appl. Math.* **66**(8), 1241–1274 (2013)
27. Casazza, P.G., Ghoreishi, D., Jose, S., Tremain, J.C.: Norm retrieval and phase retrieval by projections. *Axioms* **6**(1), 6 (2017)
28. Christ, M., Pineau, B., Taylor, M.A.: Examples of Hölder-stable phase retrieval. *Math. Res. Lett.* (2023) **(to appear)**

29. Dai, F., Prymak, A., Shadrin, A., Temlyakov, V., Tikhonov, S.: Sampling discretization of integral norms. *Constr. Approx.* **54**(3), 455–471 (2021)
30. Dai, F., Temlyakov, V.: Universal sampling discretization. *Constr. Approx.*, 1–25 (2023)
31. Deng, Y., O'Brien, M., Troitsky, V.G.: Unbounded norm convergence in Banach lattices. *Positivity* **21**(3), 963–974 (2017)
32. Diestel, J., Jarchow, H., Tonge, A.: *Absolutely summing operators*. Cambridge Studies in Advanced Mathematics, vol. 43. Cambridge University Press, Cambridge (1995)
33. Dilworth, S.J.: Special Banach lattices and their applications. In: *Handbook of the Geometry of Banach Spaces*, vol. I, pp. 497–532. North-Holland, Amsterdam (2001)
34. Domel-White, D., Bodmann, B.G.: Phase retrieval by random binary questions: which complementary subspace is closer? *Constr. Approx.* **56**, 1–33 (2022)
35. Eldar, Y.C., Mendelson, S.: Phase retrieval: stability and recovery guarantees. *Appl. Comput. Harmon. Anal.* **36**(3), 473–494 (2014)
36. Fabian, M., Habala, P., Hájek, P., Montesinos, V., Zizler, V.: *Banach space theory. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC*. Springer, New York (2011). The basis for linear and nonlinear analysis
37. Freeman, D., Ghoreishi, D.: Discretizing L_p norms and frame theory. *J. Math. Anal. Appl.* **519**(2), 126846 (2023)
38. Freeman, D., Speegle, D.: The discretization problem for continuous frames. *Adv. Math.* **345**, 784–813 (2019)
39. Gao, N., Troitsky, V.G., Xanthos, F.: Uo-convergence and its applications to Cesàro means in Banach lattices. *Isr. J. Math.* **220**(2), 649–689 (2017)
40. González, M., Martínez-Abejón, A., Martínón, A.: Disjointly non-singular operators on Banach lattices. *J. Funct. Anal.* **280**(8), 108944 (2021)
41. González, M., Martínón, A.: A quantitative approach to disjointly non-singular operators. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM* **115**(4), 185 (2021)
42. Grohs, P., Koppenssteiner, S., Rathmair, M.: Phase retrieval: uniqueness and stability. *SIAM Rev.* **62**(2), 301–350 (2020)
43. Grohs, P., Liehr, L.: On foundational discretization barriers in STFT phase retrieval. *J. Fourier Anal. Appl.* **28**(2), 39 (2022)
44. Grohs, P., Liehr, L.: Stable Gabor phase retrieval in Gaussian shift-invariant spaces via biorthogonality. *Constr. Approx.*, 1–51 (2023)
45. Hare, K.E.: An elementary proof of a result on $\Lambda(p)$ sets. *Proc. Amer. Math. Soc.* **104**(3), 829–834 (1988)
46. Heinrich, S.: Ultraproducts in Banach space theory. *J. Reine Angew. Math.* **313**, 72–104 (1980)
47. Jagannathan, K., Eldar, Y.C., Hassibi, B.: Phase retrieval: an overview of recent developments. In: *Optical Compressive Imaging*, Ser. Opt. Optoelectron., pp. 263–296. CRC Press, Boca Raton (2017)
48. James, R.C.: Uniformly non-square Banach spaces. *Ann. Math.* **2**(80), 542–550 (1964)
49. Jaming, P.: Phase retrieval techniques for radar ambiguity problems. *J. Fourier Anal. Appl.* **5**(4), 309–329 (1999)
50. Jaming, P., Rathmair, M.: Uniqueness of phase retrieval from three measurements. *Adv. Comput. Math.* **49**(4), 47 (2023)
51. Johnson, W.B., Lindenstrauss, J.: Basic concepts in the geometry of Banach spaces. In: *Handbook of the Geometry of Banach Spaces*, vol. I, pp. 1–84. North-Holland, Amsterdam (2001)
52. Kashin, B., Kosov, E., Limanova, I., Temlyakov, V.: Sampling discretization and related problems. *J. Complex.* **71**, 101653 (2022)
53. Kechris, A.S.: *Classical descriptive set theory*. Graduate Texts in Mathematics, vol. 156. Springer, New York (1995)
54. Kraemer, F., Liu, Y.-K.: Phase retrieval without small-ball probability assumptions. *IEEE Trans. Inform. Theory* **64**(1), 485–500 (2018)
55. Kraemer, F., Stöger, D.: Complex phase retrieval from subgaussian measurements. *J. Fourier Anal. Appl.* **26**(6), 89 (2020)
56. Kreuter, M.: Sobolev spaces of vector-valued functions. Ulm University Faculty of Mathematics and Economics (2015)
57. Krivine, J.-L., Maurey, B.: Espaces de Banach stables. *Isr. J. Math.* **39**(4), 273–295 (1981)
58. Lacey, H.E.: *The isometric theory of classical Banach spaces*. Die Grundlehren der mathematischen Wissenschaften, Band 208. Springer, New York (1974)

59. Limonova, I., Temlyakov, V.: On sampling discretization in L_2 . *J. Math. Anal. Appl.* **515**(2), 126457 (2022)
60. Lindenstrauss, J., Tzafriri, L.: Classical Banach spaces. I. *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Band 92. Springer, Berlin (1977). Sequence spaces
61. Lindenstrauss, J., Tzafriri, L.: Classical Banach spaces. II, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, vol. 97. Springer, Berlin (1979). Function spaces
62. Marcus, A.W., Spielman, D.A., Srivastava, N.: Interlacing Families II: mixed characteristic polynomials and the Kadison–Singer problem. *Ann. Math.* **182**(1), 327–350 (2015)
63. Munkres, J.R.: *Topology*, 2nd edn. Prentice Hall, Inc., Upper Saddle River (2000)
64. Nitzan, S., Olevskii, A., Ulanovskii, A.: Exponential frames on unbounded sets. *Proc. Amer. Math. Soc.* **144**(1), 109–118 (2016)
65. Pauli, W.: Die allgemeinen prinzipien der wellenmechanik. In: *Quantentheorie*, pp. 83–272. Springer, Berlin (1933)
66. Popovici, I.M., Vuza, D.T.: Factoring compact operators and approximable operators. *Z. Anal. Anwendungen* **9**(3), 221–233 (1990)
67. Rosenthal, H.P.: On subspaces of L^p . *Ann. Math.* **2**(97), 344–373 (1973)
68. Rudin, W.: Trigonometric series with gaps. *J. Math. Mech.* **9**, 203–227 (1960)
69. Ryou, D.: A variant of the $\Lambda(p)$ -set problem in Orlicz spaces. *Math. Z.* **302**(4), 2545–2566 (2022)
70. Schaefer, H.H.: Banach lattices and positive operators. *Die Grundlehren der mathematischen Wissenschaften*, Band 215. Springer, New York (1974)
71. Tomczak-Jaegermann, N.: Banach–Mazur distances and finite-dimensional operator ideals, *Pitman Monographs and Surveys in Pure and Applied Mathematics*, vol. 38. Longman Scientific & Technical, Harlow; copublished in the United States with Wiley, New York (1989)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.