

NONVARYING, AFFINE, AND EXTREMAL GEOMETRY OF STRATA OF DIFFERENTIALS

DAWEI CHEN

ABSTRACT. We prove that the nonvarying strata of abelian and quadratic differentials in [CM1, CM2] have trivial tautological rings and are affine varieties. We also prove that strata of k -differentials of infinite area are affine varieties for all k . Vanishing of homology in degree higher than the complex dimension follows as a consequence for these affine strata. Moreover we prove that the stratification of the Hodge bundle for abelian and quadratic differentials of finite area is extremal in the sense that merging two zeros in each stratum leads to an extremal effective divisor in the boundary. A common feature throughout these results is a relation of divisor classes in strata of differentials as well as its incarnation in Teichmüller dynamics.

CONTENTS

1. Introduction	1
2. Triviality criteria	4
3. Nonvarying strata	5
4. Differentials of infinite area	7
5. Extremal stratification of the Hodge bundle	8
References	9

1. INTRODUCTION

For $\mu = (m_1, \dots, m_n)$ with $\sum_{i=1}^n m_i = k(2g - 2)$, let $\mathcal{P}_g^k(\mu)$ be the projectivized stratum of k -differentials ω on smooth and connected complex curves of genus g that have labeled zeros and poles whose orders are specified by μ . Equivalently, $\mathcal{P}_g^k(\mu)$ parameterizes k -canonical divisors of prescribed type μ .

The study of differentials is important in surface dynamics and moduli theory. We refer to [Zo, W, C2] for an introduction to this subject. Despite various known results, the global geometry of $\mathcal{P}_g^k(\mu)$ remains quite mysterious, e.g., the full structure of the tautological ring of $\mathcal{P}_g^k(\mu)$ is largely unknown. Moreover, it is generally unclear whether the underlying coarse moduli space of $\mathcal{P}_g^k(\mu)$ is an affine variety. The birational geometry of $\mathcal{P}_g^k(\mu)$ is also less studied. Such questions are meaningful for understanding a moduli space, e.g., many exciting ideas and results have been discovered in the study of these questions for the moduli space $\mathcal{M}_g$ of curves of genus g .

Date: September 7, 2023.

Research partially supported by National Science Foundation Grants DMS-2301030, DMS-2001040, and Simons Foundation Travel Support for Mathematicians.

There has been some expectation among experts that the (rational) Picard group of $\mathcal{P}_g^k(\mu)$ should behave similarly comparing to that of $\mathcal{M}_g$, which is of rank one (for $g \geq 3$) generated by the tautological divisor class λ . However, our result below shows that such an expectation should at least exclude the strata whose Teichmüller curves have nonvarying sums of Lyapunov exponents (as listed in [CM1, CM2] for about 30 of them). For simplicity we call them nonvarying strata. Note that a stratum can be disconnected due to spin and hyperelliptic structures ([KZ, L]). Since different connected components can have distinct properties, when speaking of a stratum we mean a specific connected component according to the hyperelliptic and spin parity labelings.

The tautological ring of a stratum is defined in [C4] to be generated by the tautological classes pulled back from $\mathcal{M}_{g,n}$ together with the tautological line bundle class of the projectivized Hodge bundle. We say that the tautological ring of a stratum is trivial if it is isomorphic to $\mathbb{Q}$ generated by the fundamental class, i.e., if every tautological class of positive codimension is zero over $\mathbb{Q}$. Here a cycle class on a stratum is treated in the stacky sense when we perform intersection calculations, just like doing calculations on $\mathcal{M}_{g,n}$ as a moduli stack. In contrast, when speaking of affine, birational, or topological properties, we refer to the underlying coarse moduli space of a stratum.

Theorem 1.1. *The nonvarying strata of abelian and quadratic differentials listed in [CM1, CM2] have trivial tautological rings and their coarse moduli spaces are affine varieties.*

In contrast if a stratum is varying, then the tautological classes should not be all trivial (see Remarks 3.1 and 3.2).

Next we consider k -differentials of infinite area, i.e., when μ contains at least one entry $\leq -k$ (see [BCGGM2] for an introduction to k -differentials).

Theorem 1.2. *The coarse moduli spaces of strata of k -differentials of infinite area are affine varieties.*

Affine varieties are Stein spaces in analytic geometry. By [AF, N] we thus obtain the following result about the homology of an affine stratum of differentials.

Corollary 1.3. *Let $\mathcal{P}_g^k(\mu)$ be one of the above affine strata of differentials. Then $H_d(\mathcal{P}_g^k(\mu), \mathbb{Z}) = 0$ if $d > \dim_{\mathbb{C}} \mathcal{P}_g^k(\mu)$ and $H_d(\mathcal{P}_g^k(\mu), \mathbb{Z})$ is torsion free if $d = \dim_{\mathbb{C}} \mathcal{P}_g^k(\mu)$.*

Remark 1.4. For strata of holomorphic differentials with $\mu = (m_1, \dots, m_n)$ and $m_i \geq 0$ for all i , a (strongly) $(g+1)$ -convex exhaustion function was constructed on $\mathcal{P}_g^1(\mu)$ by using the area and length functions on flat surfaces (see [M, Proposition 3.17]), i.e., $\mathcal{P}_g^1(\mu)$ is a $(g+1)$ -complete complex space. It follows that $H_d(\mathcal{P}_g^1(\mu), \mathbb{Z})$ is torsion and $H_d(\mathcal{P}_g^1(\mu), \mathbb{C}) = 0$ for $d > \dim_{\mathbb{C}} \mathcal{P}_g^1(\mu) + g = 3g - 2 + n$ (see [BB, Theorem 0.3]).¹

In general little is known about the homology of strata of differentials (nevertheless see [CMZ] for computing the Euler characteristics and [Zy] for the L^∞ -isodelaunay decomposition). Even if one is interested in strata of holomorphic differentials in high genus, differentials in low genus and meromorphic differentials

¹Note that the q -complete index is defined in [M] one higher than in [BB].

naturally appear in the boundary of compactified strata ([BCGGM1, BCGGM3]). Therefore, we expect that the above results can be useful for inductively computing the homology of strata of differentials.

Finally we turn to the stratification of the (ordinary) Hodge bundle of holomorphic differentials (up to scale) as the disjoint union of $\mathcal{P}_g^1\{\mu\}$, where μ runs over all partitions of $2g - 2$ and $\{\mu\}$ denotes the version of strata with unlabeled zeros. Note that $\mathcal{P}_g^1\{\mu\}$ is a finite quotient of $\mathcal{P}_g^1(\mu)$ under the group action permuting the zeros that have the same order. We also denote by $\overline{\mathcal{P}}_g^1\{\mu\}$ the closure of $\mathcal{P}_g^1\{\mu\}$ in the extended Hodge bundle over the Deligne–Mumford moduli space $\overline{\mathcal{M}}_g$ of stable curves. We use similar notations for the stratification of the (quadratic) Hodge bundle of quadratic differentials with at worst simple poles and the extension over $\overline{\mathcal{M}}_{g,s}$, where s is a given number of simple poles to start with.

Theorem 1.5. *Let $\mu = (m_1, \dots, m_n)$ be a signature of abelian differentials with $m_i \geq 1$ for all i , and let $\mu' = (m_1 + m_2, m_3, \dots, m_n)$. Then the stratum $\overline{\mathcal{P}}_g^1\{\mu'\}$ is an extremal effective divisor in $\overline{\mathcal{P}}_g^1\{\mu\}$.*

Let $\mu = (m_1, \dots, m_n)$ be a signature of quadratic differentials with $m_i \geq -1$ for all i , and let $\mu' = (m_1 + m_2, m_3, \dots, m_n)$, where at least one of m_1, m_2 is positive. Then the stratum $\overline{\mathcal{P}}_g^2\{\mu'\}$ is an extremal effective divisor in $\overline{\mathcal{P}}_g^2\{\mu\}$.

Differentials of type μ' as degeneration of differentials of type μ can be realized geometrically by merging the two zeros of order m_1 and m_2 . Theorem 1.5 thus has an amusing interpretation that the stratification of the Hodge bundle (for abelian and quadratic differentials of finite area) is extremal, where merging two zeros leads to an extremal effective divisor in the boundary. We remark that if a stratum of type μ is disconnected, then Theorem 1.5 holds for each connected component of type μ and the component of type μ' obtained by merging the zeros in the prescribed component of type μ .

Strategies of the proofs. To prove Theorem 1.1 we show that the nonvarying property yields an extra relation between tautological divisor classes which forces them to be trivial. To prove Theorem 1.2 we observe the sign change of $m_i + k$ for entries $m_i \leq -k$ and $m_i > -k$ in μ , and use it to exhibit an ample divisor class which is trivial on strata of k -differentials of infinite area. To prove Theorem 1.5 we show that Teichmüller curves contained in $\mathcal{P}_g^k\{\mu'\}$ have negative intersection numbers with the divisor class of $\overline{\mathcal{P}}_g^k\{\mu'\}$ in $\overline{\mathcal{P}}_g^k\{\mu\}$ for $k = 1, 2$.

Related works. Previously in [C3] the author showed that the nonvarying strata of abelian differentials for $\mu = (4)^{\text{odd}}, (3, 1), (2, 2)^{\text{odd}}$, and $(6)^{\text{even}}$ are affine by using the geometry of canonical curves in low genus (see also [LM] for a related discussion). This is now completed for all known nonvarying strata (including for quadratic differentials). Moreover in [C3] the author also showed that there is no complete curve in any stratum of k -differentials of infinite area, which now follows from Theorem 1.2. Note that in general the implication does not work the other way around, e.g., $\mathbb{A}^2$ removing a point contains no complete curves but it is not affine. For unprojectivized strata of holomorphic differentials, nonexistence of complete curves was shown in [Ge] and [C5]. Finally the beginning case of Theorem 1.5 for the principal strata, i.e., for all zeros equal to one, was previously established in [Gh]. Now this extremal behavior is shown to hold for all zero types. Therefore,

our current results can further enhance the understanding of the tautological ring, affine geometry, topology, and birational geometry of the strata. We hope that the ideas employed in this paper can shed some light on related questions.

Acknowledgements. The author would like to thank Frederik Benirschke, Matteo Costantini, Iulia Gheorghita, Samuel Grushevsky, Richard Hain, Martin Möller, Gabriele Mondello, and Scott Mullane for helpful discussions on relevant topics. The author also sincerely thanks the referee for very useful comments.

2. TRIVIALITY CRITERIA

Marking the zeros and poles of differentials, $\mathcal{P}_g^k(\mu)$ can be viewed as a subvariety in the moduli space $\mathcal{M}_{g,n}$ of curves of genus g with n marked points. We introduce the following tautological divisor classes that will be used throughout the paper. Let η be the first Chern class of the tautological line bundle $\mathcal{O}(-1)$ on $\mathcal{P}_g^k(\mu)$ whose fibers are spanned by the underlying k -differentials ω . Let λ be the first Chern class of the Hodge bundle, κ be the Miller–Morita–Mumford class, and ψ_i be the cotangent line bundle class associated to the i -th marked point. When $m_i \neq -k$ for all i , we will often use the following quantity

$$\kappa_\mu = \frac{1}{k}(2g - 2 + n) - \sum_{i=1}^n \frac{1}{m_i + k}.$$

Recall that the tautological ring of $\mathcal{P}_g^k(\mu)$ is defined to be generated by the tautological classes pulled back from $\mathcal{M}_{g,n}$ together with η . If μ does not contain $-k$ as an entry, then it was shown in [C4, Theorem 1.1] that in this case η can generate the entire tautological ring of $\mathcal{P}_g^k(\mu)$.

Proposition 2.1. *Suppose $m_i \neq -k$ for all the entries of μ . Let D be an effective divisor in $\mathcal{M}_{g,n}$ with divisor class $a\lambda + \sum_{i=1}^n b_i\psi_i$. Then the pullback of D to $\mathcal{P}_g^k(\mu)$ has divisor class*

$$\frac{1}{12} \left(\frac{(2g - 2 + n)a}{k} + \sum_{i=1}^n \frac{12b_i - a}{m_i + k} \right) \eta.$$

Moreover if the above coefficient of η is nonzero and if D and $\mathcal{P}_g^k(\mu)$ are disjoint, then η is trivial on $\mathcal{P}_g^k(\mu)$ and the tautological ring of $\mathcal{P}_g^k(\mu)$ is trivial.

For the nonvarying strata of abelian and quadratic differentials, the desired divisors D in Proposition 2.1 arise from those used in [CM1, CM2] which are disjoint from the respective strata. Occasionally we will use some variants of $\mathcal{M}_{g,n}$ by marking fewer points or lifting to the spin moduli spaces, for which the above reasoning still works as we only consider the interior of these moduli spaces.

Proof. The following relations hold on $\mathcal{P}_g^k(\mu)$ (see [C4, Proposition 2.1]):

$$\eta = (m_i + k)\psi_i, \quad 12\lambda = \kappa = \kappa_\mu \eta.$$

Then the claim on the pullback divisor class follows from these relations.

For the other claim, we only need to show that in this case η being trivial implies that the tautological ring of $\mathcal{P}_g^k(\mu)$ is trivial, which indeed follows from the fact that η generates the tautological ring when μ contains no entries equal to $-k$ (see [C4, Theorem 1.1]). $\square$

Proposition 2.2. *If all tautological divisor classes are trivial on $\mathcal{P}_g^k(\mu)$, then the coarse moduli space of $\mathcal{P}_g^k(\mu)$ is an affine variety.*

Proof. The tautological divisor class $\kappa + \sum_{i=1}^n \psi_i$ is ample on $\overline{\mathcal{M}}_{g,n}$ (see [Co, Theorem (2.2)]) and is trivial on $\mathcal{P}_g^k(\mu)$ by the assumption. Let δ be the total boundary of $\overline{\mathcal{M}}_{g,n}$ and $0 < \epsilon \ll 1$. The divisor class $\kappa + \sum_{i=1}^n \psi_i + \epsilon\delta$ remains ample and its restriction to the closure of $\mathcal{P}_g^k(\mu)$ can be represented by a divisor fully supported on the boundary. This proves the claim. $\square$

3. NONVARYING STRATA

In this section we prove Theorem 1.1. We use the effective divisors described in [CM1, CM2] and then apply Propositions 2.1 and 2.2.

3.1. The hyperelliptic strata. Note that λ is trivial on the hyperelliptic locus (see [CH]), which implies that the hyperelliptic strata have trivial tautological rings and are affine varieties.

3.2. $\mathcal{P}_3^1(4)^{\text{odd}}$ and $\mathcal{P}_3^1(3,1)$. The divisor H parameterizing hyperelliptic curves in $\mathcal{M}_3$ is disjoint from these strata and has divisor class 9λ (see [CM1, Section 2.5, Equation (1)]).

3.3. $\mathcal{P}_3^1(2,2)^{\text{odd}}$. For $(X, p_1, p_2) \in \mathcal{P}_3^1(2,2)^{\text{odd}}$, by definition $p_1 + p_2$ is an odd theta characteristic, which maps $\mathcal{P}_3^1(2,2)^{\text{odd}}$ to the odd spin moduli space $\mathcal{S}_3^-$. The divisor Z_3 in $\mathcal{S}_3^-$ parameterizes odd theta characteristics with a double zero. It is disjoint from $\mathcal{P}_3^1(2,2)^{\text{odd}}$ and has divisor class 11λ (see [CM1, Section 5.3]).

3.4. $\mathcal{P}_3^1(2,1,1)$. Marking the first two zeros maps $\mathcal{P}_3^1(2,1,1)$ to $\mathcal{M}_{3,2}$. The Brill–Noether divisor $BN_{3,(1,2)}^1$ in $\mathcal{M}_{3,2}$ parameterizes curves that admit a g_3^1 given by $p_1 + 2p_2$. It is disjoint from $\mathcal{P}_3^1(2,1,1)$ and has divisor class $-\lambda + \psi_1 + 3\psi_2$ (see [CM1, Section 2.5, Equation (5)]).

3.5. $\mathcal{P}_4^1(6)^{\text{even}}$. For $(X, p) \in \mathcal{P}_4^1(6)^{\text{even}}$, by definition $3p$ is an even theta characteristic. The theta-null divisor Θ_4 in $\mathcal{M}_{4,1}$ parameterizes curves that admit an odd theta characteristic whose support contains the marked point. It is disjoint from $\mathcal{P}_4^1(6)^{\text{even}}$ and has divisor class $30\lambda + 60\psi_1$ (see [CM1, Section 2.5, Equation (3)]).

3.6. $\mathcal{P}_4^1(6)^{\text{odd}}$ and $\mathcal{P}_4^1(5,1)$. Marking the zero of the largest order, these strata map to $\mathcal{M}_{4,1}$. The Brill–Noether divisor $BN_{3,(2)}^1$ in $\mathcal{M}_{4,1}$ parameterizes curves that admit a g_3^1 ramified at the marked point. It is disjoint from both strata and has divisor class $8\lambda + 4\psi_1$ (see [CM1, Section 2.5, Equation (7)]).

3.7. $\mathcal{P}_4^1(3,3)^{\text{nonhyp}}$. The divisor Lin_3^1 in $\mathcal{M}_{3,2}$ parameterizes curves that admit a g_3^1 with a fiber containing both marked points. It is disjoint from $\mathcal{P}_4^1(3,3)^{\text{nonhyp}}$ and has divisor class $8\lambda - \psi_1 - \psi_2$ (see [CM1, Section 2.5, Equation (8)]).

3.8. $\mathcal{P}_4^1(2,2,2)^{\text{odd}}$. For $(X, p_1, p_2, p_3) \in \mathcal{P}_4^1(2,2,2)^{\text{odd}}$, by definition $p_1 + p_2 + p_3$ is an odd theta characteristic. The divisor Z_4 in the odd spin moduli space $\mathcal{S}_4^-$ parameterizes odd theta characteristics with a double zero. It is disjoint from $\mathcal{P}_4^1(2,2,2)^{\text{odd}}$ and has divisor class 12λ (see [CM1, Section 5.3]).

3.9. $\mathcal{P}_4^1(3, 2, 1)$. The Brill–Noether divisor $BN_{4,(1,1,2)}^1$ in $\mathcal{M}_{4,3}$ parameterizes curves that admit a g_4^1 given by $p_1 + p_2 + 2p_3$. It is disjoint from $\mathcal{P}_4^1(3, 2, 1)$ and has divisor class $-\lambda + \psi_1 + \psi_2 + 3\psi_3$ (see [CM1, Section 2.5, Equation (6)]).

3.10. $\mathcal{P}_5^1(8)^{\text{even}}$. The Brill–Noether divisor BN_3^1 in $\mathcal{M}_{5,1}$ parameterizes curves with a g_3^1 . It is disjoint from $\mathcal{P}_5^1(8)^{\text{even}}$ and has divisor class 8λ (see [CM1, Section 2.5, Equation (4)]).

3.11. $\mathcal{P}_5^1(8)^{\text{odd}}$. The divisor $\text{Nfold}_{5,4}^1(1)$ in $\mathcal{M}_{5,1}$ parameterizes curves that admit a g_4^1 with a fiber containing $3p$. It is disjoint from $\mathcal{P}_5^1(8)^{\text{odd}}$ and has divisor class $7\lambda + 15\psi_1$ (see [CM1, Section 2.5, Equation (9)]).

3.12. $\mathcal{P}_5^1(5, 3)$. The Brill–Noether divisor $BN_{4,(1,2)}^1$ in $\mathcal{M}_{5,2}$ parameterizes curves that admit a g_4^1 with a fiber containing $p_1 + 2p_2$. It is disjoint from $\mathcal{P}_5^1(5, 3)$ and has divisor class $7\lambda + 7\psi_1 + 2\psi_2$ (see [CM1, Section 2.5, Equation (10)]).

3.13. $\mathcal{P}_4^1(4, 2)^{\text{even}}$, $\mathcal{P}_4^1(4, 2)^{\text{odd}}$, and $\mathcal{P}_5^1(6, 2)^{\text{odd}}$. The nonvarying property of these strata was proved in [YZ] by using the Harder–Narasimhan filtration of the Hodge bundle (see also [CM2, Appendix A]). Note that the filtration holds not only on a Teichmüller curve but also on (the interior of) each stratum. Let L_μ be the sum of (nonnegative) Lyapunov exponents which satisfies that $\lambda = L_\mu \eta$ from the Harder–Narasimhan filtration on these strata. In these cases we know that $L_\mu \neq \kappa_\mu/12$. Then the triviality of η on these strata follows from the two distinct relations $\lambda = (\kappa_\mu/12)\eta$ and $\lambda = L_\mu \eta$.

3.14. **Nonvarying strata of quadratic differentials in genus one and two.** Strata in genus ≤ 2 can be generally dealt with as in Sections 4.2 and 4.3 below.

3.15. $\mathcal{P}_3^2(9, -1)^{\text{irr}}$. This stratum is disjoint from the hyperelliptic divisor H in $\mathcal{M}_3$ whose divisor class is 9λ (see [CM2, Section 3.1, Equation (6)]).

3.16. $\mathcal{P}_3^2(8)$, $\mathcal{P}_3^2(7, 1)$, $\mathcal{P}_3^2(8, 1, -1)$, $\mathcal{P}_3^2(10, -1, -1)^{\text{nonhyp}}$, and $\mathcal{P}_3^2(9, -1)^{\text{reg}}$. Marking the zero of the largest order, these strata map to $\mathcal{M}_{3,1}$. The divisor W of Weierstrass points in $\mathcal{M}_{3,1}$ is disjoint from these strata and has divisor class $-\lambda + 6\psi_1$ (see [CM2, Section 3, Equation (5)]).

3.17. $\mathcal{P}_3^2(6, 2)^{\text{nonhyp}}$, $\mathcal{P}_3^2(6, 1, 1)^{\text{nonhyp}}$, $\mathcal{P}_3^2(5, 3)$, $\mathcal{P}_3^2(5, 2, 1)$, $\mathcal{P}_3^2(4, 4)$, $\mathcal{P}_3^2(4, 3, 1)$, $\mathcal{P}_3^2(5, 4, -1)$, $\mathcal{P}_3^2(5, 3, 1, -1)$, $\mathcal{P}_3^2(7, 2, -1)$, $\mathcal{P}_3^2(7, 3, -1, -1)$, and $\mathcal{P}_3^2(6, 3, -1)^{\text{reg}}$. Marking the first two zeros, these strata map to $\mathcal{M}_{3,2}$. The Brill–Noether divisor $BN_{3,(2,1)}^1$ in $\mathcal{M}_{3,2}$ parameterizes curves that admit a g_3^1 given by $2p_1 + p_2$. It is disjoint from these strata and has divisor class $-\lambda + 3\psi_1 + \psi_2$ (see [CM2, Section 3.1, Equation (8)]).

3.18. $\mathcal{P}_3^2(4, 2, 2)$, $\mathcal{P}_3^2(3, 3, 2)^{\text{nonhyp}}$, $\mathcal{P}_3^2(4, 3, 2, -1)$, $\mathcal{P}_3^2(3, 2, 2, 1)$, $\mathcal{P}_3^2(3, 3, 1, 1)^{\text{nonhyp}}$, and $\mathcal{P}_3^2(3, 3, 3, -1)^{\text{reg}}$. Marking the first three zeros, these strata map to $\mathcal{M}_{3,3}$. The Brill–Noether divisor $BN_{3,(1,1,1)}^1$ in $\mathcal{M}_{3,3}$ parameterizes curves that admit a g_3^1 given by $p_1 + p_2 + p_3$. It is disjoint from these strata and has divisor class $-\lambda + \psi_1 + \psi_2 + \psi_3$ (see [CM2, Section 3.1, Equation (7)]).

3.19. $\mathcal{P}_4^2(13, -1)$, $\mathcal{P}_4^2(11, 1)$, and $\mathcal{P}_4^2(12)^{\text{reg}}$. Marking the zero of the largest order, these strata map to $\mathcal{M}_{4,1}$. The divisor W of Weierstrass points in $\mathcal{M}_{4,1}$ is disjoint from these strata and has divisor class $-\lambda + 10\psi_1$ (see [CM2, Lemma 3.2]).

3.20. $\mathcal{P}_4^2(10, 2)^{\text{nonhyp}}$, $\mathcal{P}_4^2(8, 4)$, $\mathcal{P}_4^2(8, 3, 1)$, **and** $\mathcal{P}_4^2(9, 3)^{\text{reg}}$. Marking the first two zeros, these strata map to $\mathcal{M}_{4,2}$. The Brill–Noether divisor $BN_{4,(3,1)}^1$ in $\mathcal{M}_{4,2}$ parameterizes curves that admit a g_4^1 given by $3p_1 + p_2$. It is disjoint from these strata and has divisor class $-\lambda + 6\psi_1 + \psi_2$ (see [CM2, Lemma 3.2]).

3.21. $\mathcal{P}_4^2(7, 5)$ **and** $\mathcal{P}_4^2(6, 6)^{\text{reg}}$. The Brill–Noether divisor $BN_{4,(2,2)}^1$ in $\mathcal{M}_{4,2}$ parameterizes curves that admit a g_4^1 given by $2p_1 + 2p_2$. It is disjoint from these strata and has divisor class $-\lambda + 3\psi_1 + 3\psi_2$ (see [CM2, Lemma 3.2]).

3.22. $\mathcal{P}_4^2(7, 3, 2)$, $\mathcal{P}_4^2(5, 4, 3)$, **and** $\mathcal{P}_4^2(6, 3, 3)^{\text{reg}}$. The Brill–Noether divisor $BN_{4,(2,1,1)}^1$ in $\mathcal{M}_{4,3}$ parameterizes curves that admit a g_4^1 given by $2p_1 + p_2 + p_3$. It is disjoint from these strata and has divisor class $-\lambda + 3\psi_1 + \psi_2 + \psi_3$ (see [CM2, Lemma 3.2]).

3.23. $\mathcal{P}_4^2(3, 3, 3, 3)^{\text{reg}}$. It is disjoint from the Brill–Noether divisor $BN_{4,(1,1,1,1)}^1$ in $\mathcal{M}_{4,4}$ whose divisor class is $-\lambda + \psi_1 + \psi_2 + \psi_3 + \psi_4$ (see [CM2, Lemma 3.2]).

3.24. $\mathcal{P}_3^2(6, 3, -1)^{\text{irr}}$, $\mathcal{P}_4^2(12)^{\text{irr}}$, **and** $\mathcal{P}_4^2(9, 3)^{\text{irr}}$. We can use the Harder–Narasimhan filtration of the quadratic Hodge bundle for these strata (see [CM2, Appendix A] and [CY, Section 4.2]) and argue as in Section 3.13.

Remark 3.1. We say that a stratum of abelian differentials $\mathcal{P}_g^1(\mu)$ is varying, if it contains two Teichmüller curves that have distinct sums of Lyapunov exponents. In this case we claim that η is nontrivial on $\mathcal{P}_g^1(\mu)$.

To see this, let Δ_0 be the locus of stable differentials of type μ on curves with separating nodes only, where the differentials have simple poles with opposite residues at the nodes. If $\mathcal{P}_g^1(\mu)$ is irreducible, then Δ_0 is also irreducible and modeled on $\mathcal{P}_{g-1}^1(\mu, \{-1, -1\})$. Moreover, every Teichmüller curve in $\mathcal{P}_g^1(\mu)$ only intersects Δ_0 in the boundary (see [CM1, Corollary 3.2]).

Suppose that η is trivial on $\mathcal{P}_g^1(\mu)$. Then $\eta = e\delta_0$ on the partial compactification $\mathcal{P}_g^1(\mu) \cup \Delta_0$ for some constant e . Since $\lambda = (\kappa_\mu/12)\eta$ on $\mathcal{P}_g^1(\mu)$, we have $\lambda = \ell\delta_0$ on $\mathcal{P}_g^1(\mu) \cup \Delta_0$ for some constant ℓ . Then the sum of Lyapunov exponents of a Teichmüller curve C is given by $(\lambda \cdot C)/(\eta \cdot C) = \ell/e$ which is independent of C , contradicting that $\mathcal{P}_g^1(\mu)$ is a varying stratum.

Note that the above criterion is practically checkable for arithmetic Teichmüller curves generated by square-tiled surfaces by using the combinatorial description of area Siegel–Veech constants and sums of Lyapunov exponents (see [EKZ, Section 2.5.1] and also [C1, Theorem 1.8] for the relation with slopes of Teichmüller curves).

The same conclusion would hold for a varying stratum of quadratic differentials, assuming that all Teichmüller curves contained in the stratum intersect only one irreducible boundary divisor (see [CM2, Remark 4.7] for this assumption).

Remark 3.2. We speculate that the rational Picard group (and the rational second cohomology group) of any varying stratum of holomorphic differentials $\mathcal{P}_g^1(\mu)$ should be of rank one generated by η .

4. DIFFERENTIALS OF INFINITE AREA

In this section we prove Theorem 1.2 for strata of k -differentials of infinite area. For completeness we also include the discussion for some strata in low genus.

4.1. Strata in genus zero. Every stratum in genus zero is isomorphic to the corresponding moduli space of pointed smooth rational curves, which has a trivial Chow ring and is affine.

4.2. Strata in genus one. Since $\psi_i = 0$ on $\mathcal{M}_{1,n}$ for all i (see [AC, Theorem 2.2 c)]), every stratum $\mathcal{P}_1^k(\mu)$ in genus one (including the case of some $m_i = -k$ in μ) has a trivial tautological ring and is affine by Proposition 2.2. The affinity also follows from the fact that the ambient space $\mathcal{M}_{1,n}$ is affine (see [C3, Theorem 3.1]).

4.3. Strata in genus two. Since λ is trivial on $\mathcal{M}_2$, every stratum $\mathcal{P}_2^k(\mu)$ in genus two with $m_i \neq -k$ for all i has a trivial tautological ring and is affine by Proposition 2.2.

4.4. Strata with poles of order $\leq -k$. Suppose $m_1 \leq -k$. By the relations $\eta = (m_i + k)\psi_i$ for all i and $(2g - 2)\eta = k\kappa - \sum_{i=1}^n m_i\psi_i$, we obtain that

$$\begin{aligned} k\left(\kappa + \sum_{i=1}^n \psi_i\right) &= (2g - 2 + n)\eta \\ &= (2g - 2 + n)(m_1 + k)\psi_1. \end{aligned}$$

It implies that $\kappa + \sum_{i=1}^n \psi_i + a\psi_1$ is trivial on $\mathcal{P}_g^k(\mu)$ where $a = -(2g - 2 + n)(m_1 + k)/k \geq 0$. Since $\kappa + \sum_{i=1}^n \psi_i + a\psi_1$ (as ample plus nef) remains ample on $\overline{\mathcal{M}}_{g,n}$, it follows that $\mathcal{P}_g^k(\mu)$ is affine and Theorem 1.2 is justified.

5. EXTREMAL STRATIFICATION OF THE HODGE BUNDLE

In this section we prove Theorem 1.5. Recall the notation $\mu = (m_1, \dots, m_n)$, $\mu' = (m_1 + m_2, m_3, \dots, m_n)$, and we work with unordered zeros. The idea is to show that Teichmüller curves in $\mathcal{P}_g^k\{\mu'\}$ has negative intersection numbers with the divisor class of $\overline{\mathcal{P}}_g^k\{\mu'\}$ in $\overline{\mathcal{P}}_g^k\{\mu\}$ for $k = 1, 2$. In order to verify that, we make use of the aforementioned fact that degenerate differentials in the boundary of a Teichmüller curve can have only simple polar nodes (after taking a local square root in the case of quadratic differentials).

Proof of Theorem 1.5. First consider the case of abelian differentials. The following relation of divisor classes holds on $\overline{\mathcal{P}}_g^1\{\mu\}$:

$$12\lambda - D_h - \kappa_\mu\eta = (m_1 + m_2 + 1)\left(1 - \frac{1}{m_1 + 1} - \frac{1}{m_2 + 1} + \frac{1}{m_1 + m_2 + 1}\right)\overline{\mathcal{P}}_g^1\{\mu'\}$$

modulo the other boundary divisors, where D_h is the boundary divisor generically parameterizing stable differentials with simple polar nodes. This relation follows from [CCM, Proposition 6.3], where $\overline{\mathcal{P}}_g^1\{\mu\}$ can be identified with the boundary divisor D_Γ having $\Gamma_\perp$ as a rational vertex containing the two merged zeros and $\ell_\Gamma = m_1 + m_2 + 1$ therein. Note that the coefficient on the right of the above equality is positive for $m_1, m_2 \geq 1$.

Let C be the closure of a Teichmüller curve in $\mathcal{P}_g^1\{\mu'\}$. Since C does not intersect the part of the boundary of $\overline{\mathcal{P}}_g^1\{\mu\}$ away from D_h and $\overline{\mathcal{P}}_g^1\{\mu'\}$, we obtain

$$\begin{aligned} \frac{(12\lambda - D_h - \kappa_\mu\eta) \cdot C}{\eta \cdot C} &= \kappa_{\mu'} - \kappa_\mu \\ &= -1 + \frac{1}{m_1 + 1} + \frac{1}{m_2 + 1} - \frac{1}{m_1 + m_2 + 1}, \end{aligned}$$

where $12\lambda - D_h = \kappa_{\mu'}\eta$ holds on the stratum $\overline{\mathcal{P}}_g^1\{\mu'\}$ (modulo the other irrelevant boundary divisors of $\overline{\mathcal{P}}_g^1\{\mu'\}$ which do not meet C). Consequently, this implies

$$\frac{C \cdot \overline{\mathcal{P}}_g^1\{\mu'\}}{C \cdot \eta} = -\frac{1}{m_1 + m_2 + 1} < 0.$$

Since η has positive degree on every Teichmüller curve (e.g., from the positive sign of the area Siegel–Veech constant), it follows that $C \cdot \overline{\mathcal{P}}_g^1\{\mu'\} < 0$ in $\overline{\mathcal{P}}_g^1\{\mu\}$. Since Teichmüller curves are dense in $\overline{\mathcal{P}}_g^1\{\mu'\}$ and the negative ratio above is independent of each individual Teichmüller curve, the desired claim follows by using the same argument as in [Gh, Section 4].

The case of quadratic differentials is similar. The following relation of divisor classes holds on $\overline{\mathcal{P}}_g^2\{\mu\}$:

$$12\lambda - D_h - \kappa_\mu\eta = (m_1 + m_2 + 2) \left(\frac{1}{2} - \frac{1}{m_1 + 2} - \frac{1}{m_2 + 2} + \frac{1}{m_1 + m_2 + 2} \right) \overline{\mathcal{P}}_g^2\{\mu'\}$$

modulo the other boundary divisors, where the coefficient on the right is nonzero by the assumption that $m_1, m_2 \geq -1$ and at least one of them is positive. Let C be the closure of a Teichmüller curve in $\mathcal{P}_g^2\{\mu'\}$. Since C does not intersect the part of the boundary of $\overline{\mathcal{P}}_g^2\{\mu\}$ away from D_h and $\overline{\mathcal{P}}_g^2\{\mu'\}$, we obtain

$$\frac{(12\lambda - D_h - \kappa_\mu\eta) \cdot C}{\eta \cdot C} = -\frac{1}{2} + \frac{1}{m_1 + 2} + \frac{1}{m_2 + 2} - \frac{1}{m_1 + m_2 + 2}.$$

It follows that

$$\frac{C \cdot \overline{\mathcal{P}}_g^2\{\mu'\}}{C \cdot \eta} = -\frac{1}{m_1 + m_2 + 2} < 0.$$

The rest of the argument is the same as before. $\square$

Remark 5.1. If a stratum in Theorem 1.5 is disconnected, since each of its connected components contains a dense collection of Teichmüller curves, the conclusion of Theorem 1.5 still holds for each connected component.

Remark 5.2. For k -differentials with $k > 2$, since there is no meaningful way to define Teichmüller curves in this case, our method cannot be directly adapted. We leave it as an interesting question to determine whether the stratification of the k -th Hodge bundle is extremal for $k > 2$.

REFERENCES

- [AC] E. Arbarello and M. Cornalba, Calculating cohomology groups of moduli spaces of curves via algebraic geometry, *Inst. Hautes Études Sci. Publ. Math.* No. 88 (1998), 97–127.
- [AF] A. Andreotti and T. Frankel, The Lefschetz Theorem on Hyperplane Sections, *Ann. of Math. (2)* 69 (1959), 713–717.

- [BCGGM1] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, and M. Möller, Compactification of strata of Abelian differentials, *Duke Math. J.* 167 (2018), no. 12, 2347-2416.
- [BCGGM2] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, and M. Möller, Strata of k -differentials, *Algebr. Geom.* 6 (2019), no. 2, 196-233.
- [BCGGM3] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, and M. Möller, The moduli space of multi-scale differentials, *arXiv:1910.13492*.
- [BB] E. Ballico and G. Bolondi, On the homology groups of q -complete spaces, *Rend. Sem. Mat. Univ. Padova* 69 (1983), 19-25.
- [C1] D. Chen, Square-tiled surfaces and rigid curves on moduli spaces, *Adv. Math.* 228 (2011), no. 2, 1135-1162.
- [C2] D. Chen, Teichmüller dynamics in the eyes of an algebraic geometer, *Surveys on recent developments in algebraic geometry*, 171-197, Proc. Sympos. Pure Math., 95, Amer. Math. Soc., Providence, RI, 2017.
- [C3] D. Chen, Affine geometry of strata of differentials, *J. Inst. Math. Jussieu* 18 (2019), no. 6, 1331-1340.
- [C4] D. Chen, Tautological ring of strata of differentials, *Manuscripta Math.* 158 (2019), no. 3-4, 345-351.
- [C5] D. Chen, Complete curves in the strata of differentials, *C. R. Math. Acad. Sci. Paris* 361 (2023), 371-373.
- [CCM] D. Chen, M. Costantini, and M. Möller, On the Kodaira dimension of moduli spaces of Abelian differentials, *arXiv:2204.11943*.
- [CM1] D. Chen and M. Möller, Nonvarying sums of Lyapunov exponents of Abelian differentials in low genus, *Geom. Topol.* 16 (2012), no. 4, 2427-2479.
- [CM2] D. Chen and M. Möller, Quadratic differentials in low genus: exceptional and non-varying strata, *Ann. Sci. Éc. Norm. Supér. (4)* 47 (2014), no. 2, 309-369.
- [CY] D. Chen and F. Yu, Filtration and splitting of the Hodge bundle on the non-varying strata of quadratic differentials, *arXiv:2306.06702*.
- [Co] M. Cornalba, On the projectivity of the moduli spaces of curves, *J. Reine Angew. Math.* 443 (1993), 11-20.
- [CH] M. Cornalba and J. Harris, Divisor classes associated to families of stable varieties, with applications to the moduli space of curves, *Ann. Sci. École Norm. Sup. (4)* 21 (1988), no. 3, 455-475.
- [CMZ] M. Costantini, M. Möller, and J. Zachhuber, The Chern classes and the Euler characteristic of the moduli spaces of abelian differentials, *Forum Math. Pi* 10 (2022), Paper No. e16, 55 pp.
- [EKZ] A. Eskin, M. Kontsevich, and A. Zorich, Sum of Lyapunov exponents of the Hodge bundle with respect to the Teichmüller geodesic flow, *Publ. Math. Inst. Hautes Études Sci.* 120 (2014), 207-333.
- [Ge] Q. Gendron, Les strates ne possèdent pas de variétés complètes, *C. R. Math. Acad. Sci. Paris* 358 (2020), no. 2, 197-200.
- [Gh] I. Gheorghita, Effective divisors in the projectivized Hodge bundle, *Int. Math. Res. Not. IMRN* (2021), no. 1, 303-325.
- [KZ] M. Kontsevich and A. Zorich, Connected components of the moduli spaces of Abelian differentials with prescribed singularities, *Invent. Math.* 153 (2003), no. 3, 631-678.
- [L] E. Laneeau, Connected components of the strata of the moduli spaces of quadratic differentials, *Ann. Sci. Éc. Norm. Supér. (4)* 41 (2008), no. 1, 1-56.
- [M] G. Mondello, On the cohomological dimension of the moduli space of Riemann surfaces, *Duke Math. J.* 166 (2017), no. 8, 1463-1515.
- [LM] E. Looijenga and G. Mondello, The fine structure of the moduli space of abelian differentials in genus 3, *Geom. Dedicata* 169 (2014), 109-128.
- [N] R. Narasimhan, On the homology groups of Stein spaces, *Invent. Math.* 2 (1967), 377-385.
- [W] A. Wright, Translation surfaces and their orbit closures: an introduction for a broad audience, *EMS Surv. Math. Sci.* 2 (2015), no. 1, 63-108.
- [YZ] F. Yu and K. Zuo, Weierstrass filtration on Teichmüller curves and Lyapunov exponents, *J. Mod. Dyn.* 7 (2013), no. 2, 209-237.
- [Zo] A. Zorich, Flat surfaces, *Frontiers in number theory, physics, and geometry. I*, 437-583, Springer, Berlin, 2006.

- [Zy] B. Zykoski, *The L^∞ -isodelaunay decomposition of strata of abelian differentials*, *arXiv:2206.04143*.

DEPARTMENT OF MATHEMATICS, BOSTON COLLEGE, CHESTNUT HILL, MA 02467, USA
Email address: `dawei.chen@bc.edu`