

Copper-Catalyzed Hydroamination: Enantioselective Addition of Pyrazoles to Cyclopropenes

Minghao Wang, Julie C. Simon,[§] Mengfei Xu,[§] Stephanie A. Corio, Jennifer S. Hirschi,*
and Vy M. Dong*



Cite This: *J. Am. Chem. Soc.* 2023, 145, 14573–14580



Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: Chiral *N*-cyclopropyl pyrazoles and structurally related heterocycles are prepared using an earth-abundant copper catalyst under mild reaction conditions with high regio-, diastereo-, and enantiocontrol. The observed N²:N¹ regioselectivity favors the more hindered nitrogen of the pyrazole. Experimental and DFT studies support a unique mechanism that features a five-centered aminocupration.


Catalytic functionalization of heterocycles presents a challenge with implications for the discovery and preparation of medicines.^{1,2} Pyrazoles, a type of nitrogen-containing heterocycle,³ rank fourth in occurrence among the most recent FDA-approved drugs (Figure 1A).^{2d} Given that hydroamination is an efficient approach to form C–N bonds,^{4,5} variants that feature nitrogen-containing heterocycles warrant development.⁶ To date, pyrazoles undergo addition to

allenes,^{6a,c} alkynes,^{6b} and dienes,^{6d,e} albeit using precious metals (namely, Rh and Pd). As a promising alternative, Cu–amido complexes⁷ (first isolated and characterized by Gunnoe^{8a}) catalyze the addition of amines to electron-deficient olefins,^{8c,d} allenes,^{9b} nitrostyrenes,^{9c} and azabenzonornorbornadienes.^{9d} Despite this reactivity, no asymmetric variants were yet achieved (Figure 1B). In this study, we report a Cu-catalyzed hydroamination with pyrazoles that provides chiral cyclopropyl motifs with high regio-, diastereo-, and enantiocontrol (Figure 1C).


The hydrofunctionalization of cyclopropenes represents a versatile strategy for accessing chiral cyclopropanes.^{10–12} Most relevant to our current study, Hou and co-workers disclosed the asymmetric coupling of cyclopropenes with secondary amines (e.g., morpholine, pyrrolidine, and dibenzylamines) by using a rare-earth metal catalyst (Sm).^{12d} Buchwald and co-workers developed an enantioselective hydroamination of 1-silyl- or 1-aryl-substituted cyclopropenes using Cu–H catalysis,^{12h} with *O*-benzoylhydroxylamines as oxidants¹³ and silanes as the stoichiometric reductant.¹⁴ We hypothesized that the deprotonation of pyrazole with a catalytic amount of base would generate a Cu–pyrazolate catalyst, which would undergo aminocupration to cyclopropenes (Figure 1C).¹⁵ Subsequent protodemetalation would produce cyclopropyl pyrazoles. If successful, this method would enable a novel and late-stage¹⁶ cyclopropylation of pyrazoles with high atom economy.¹⁷

To begin this study, we focused on the desymmetrization of achiral cyclopropene **1a** with pyrazole (**2a**) to generate cyclopropyl pyrazole **3a**, which bears two stereogenic centers. An initial experiment with Cu in the absence of ligands

A. Inspiration: Bioactive molecules with pyrazole motifs



B. Background: Towards Cu-amido hydroamination



C. This work: Enantioselective Cu–N catalysis





Figure 1. Asymmetric hydroamination of cyclopropenes with pyrazoles.


Received: March 21, 2023

Published: June 30, 2023



resulted in the formation of allylic pyrazole **5a** (Figure 2A). Achiral ligands, such as *rac*-BINAP, led to the exclusive

A. Undesired reactivity identified



B. Effect of electron-rich chiral ligands




Figure 2. Reaction optimization using bisphosphine ligands. ^aReaction conditions: **1a** (0.12 mmol), **2** (0.10 mmol), Cu(CH₃CN)₄PF₆ (5 mol%), chiral ligand (6 mol%), toluene (0.4 mL), 30 °C, 6–24 h. Yields of isolated products are given. Enantiomeric ratios (*er*) were determined by SFC analysis on a chiral stationary phase. ^bThe reaction was performed using CH₃CN at 30 °C for 6 h. ^cThe reaction was performed using CH₃CN at 0 °C for 4 days.

formation of the same undesired isomer **5a**, likely via a ring-opening pathway^{12h,18} involving N–H bond insertion into allylic carbene **5a'**.¹⁹ Using commercial (IPr)CuCl offered high chemoselectivity to **3a** (see the [Supporting Information](#) (SI)).^{9d} However, efforts to generate chiral NHC–Cu complexes *in situ* from imidazolium salts provided **5a** more favorably; we presume that the ring-opening pathway with CuCl outcompetes NHC carbene formation and ligation. Gunnoe observed enhanced Cu–N nucleophilicity with bulky electron-rich phosphine ligands compared to NHC ligands.^{8b,d} We wondered whether bulky phosphine ligands would favor Cu–amido insertion over ring opening (Figure 2B). We found that bulky chiral phosphines (**L1**–**L7**) gave promising results. Among them, high enantioselectivity (90:10 *er*) and yield (90%) of **3a** were observed using (*R,R*)-*i*-Pr-Duphos (**L3**). We observed improved enantioselectivity (94:6 *er*) with CH₃CN as the solvent (see the SI). Although lowering the temperature to 0 °C improved the enantiomeric ratio, a longer reaction time (4 days) was necessary. Therefore, we chose to explore the substrate scope at 30 °C.

With this mild protocol in hand, we evaluated the enantioselective coupling of various pyrazoles to cyclopropene **1a** (Table 1). In general, high diastereoselectivity (>20:1) was observed, likely due to the large steric difference between methyl and phenyl substituents.²⁰ Symmetric pyrazoles afforded **3b**–**3i** with high enantioselectivity (91:9–99:1 *er*) in 40–94% yield. Introducing a methyl substituent on the

Table 1. Scope of Pyrazoles and Other N-Heterocycles.

Reaction conditions: **1a** (0.12 mmol), **2** (0.10 mmol), Cu(CH₃CN)₄PF₆ (5 mol%), **L3** (6 mol%), CH₃CN (0.4 mL), 30 °C, 6–12 h. Yields of isolated products are given. Nitrogen regioisomeric ratios (N²:N¹) were determined based on isolated yields and NMR experiments. Diastereomeric ratios (*dr*) were determined from ¹H NMR analysis of the reaction mixtures. Enantiomeric ratios (*er*) were determined by SFC analysis on a chiral stationary phase. ^b24 h. ^cThe reaction was performed at 60 °C.

^aReaction conditions: **1a** (0.12 mmol), **2** (0.10 mmol), Cu(CH₃CN)₄PF₆ (5 mol%), **L3** (6 mol%), CH₃CN (0.4 mL), 30 °C, 6–12 h. Yields of isolated products are given. Nitrogen regioisomeric ratios (N²:N¹) were determined based on isolated yields and NMR experiments. Diastereomeric ratios (*dr*) were determined from ¹H NMR analysis of the reaction mixtures. Enantiomeric ratios (*er*) were determined by SFC analysis on a chiral stationary phase. ^b24 h. ^cThe reaction was performed at 60 °C.


pyrazole showed no significant effect on reactivity (**3b**), whereas more hindered dimethylated pyrazoles gave decreased yields (**3h**, **3i**). Electron-withdrawing groups on pyrazoles were accommodated (**3c**–**3f**), although we observed more undesired allylic pyrazole with cyano substitution (**3e**). In the case of unsymmetric pyrazoles, 65–90% yield, 92:8–99:1 *er*, and nitrogen regioselectivity (N²:N¹ > 20:1) were observed (**3j**–**3o**). X-ray crystallographic analysis of **3j** confirmed the coupling of cyclopropene with the *more* sterically hindered nitrogen of the pyrazole; this regioselectivity is rare in pyrazole functionalization.^{9d,21,22} Further NOE experiments confirmed similar regiocontrol for related pyrazole substrates (see the SI). Despite the presence of a competing amino group, **3l** was isolated exclusively, showing a highly chemoselective cyclopropylation for pyrazole nitrogens.

Next, we investigated hydroamination using other heterocycles under the standard conditions. Pyridazinone (**2p**), a nitrogen-rich and medically relevant heterocycle,²³ provided **3p** in 55% yield, 87:13 *er*, and >20:1 nitrogen regioselectivity. Indazole, a heterocycle used as an indole bioisostere,²⁴ showed promising results. Chiral indazoles (**3q**–**3t**) were prepared in up to 66% yield with 89:11–92:8 *er* and nitrogen regioselectivity (3.1:1 to >20:1 *rr*).²⁵ The coupling of **1a** with indazoles required higher temperatures and resulted in

lower regioselectivity, except for electron-withdrawing ester substitution (**3t**). Other nitrogen nucleophiles, such as imidazole, triazole, and aniline, exhibited no desired reactivity under standard conditions and warrant further development.^{9a}

In addition, we studied the enantioselective hydroamination of pyrazole **2a** with various cyclopropenes **1** (Table 2). The

Table 2. Scope of Cyclopropenes



Cyclopropene 1	Product 4	Yield (%)	er
1a	4a	72%	94:6
1b	4b	75%	94:6
1c	4c	77%	94:6
1d	4d	89%	93:7
1e	4e	60%	92:8
1f	4f	68%	92:8

Substituent R	Product	Yield (%)	er
4-Me	4b	72%	94:6
4-OMe	4c	75%	94:6
4-Cl	4d	77%	94:6
4-CF ₃	4e	89%	93:7
2-Me	4f	60%	92:8
3-Me	4g	68%	92:8

Cyclopropene 1	Product 4	Yield (%)	er
1h	4h	96%	93:7
1i	4i	74%	93:7
1j	4j	94%	96:4
1k	4k	51%	99:1
1l	4l	80%	95:5
1m	4m	63%	94:6
1n	4n	80%	89:11

^aReaction conditions: **1** (0.12 mmol), **2a** (0.10 mmol), Cu(CH₃CN)₄PF₆ (5 mol%), L3 (6 mol%), CH₃CN (0.4 mL), 30 °C, 6–12 h. Yields of isolated products are given. Diastereomeric ratios (*dr*) were determined from ¹H NMR analysis of the unpurified reaction mixtures. Enantiomeric ratios (*er*) were determined by SFC analysis on a chiral stationary phase. ^b1:1 *dr*.

electronic properties of the phenyl ring on **1** have a negligible impact on the enantioselectivity and reactivity. Good yields (60–89%) and enantioselectivity (92:8–94:6 *er*) were observed with both electron-rich (**4b**, **4c**, **4f**, and **4g**) and electron-deficient (**4d**, **4e**) substrates. Replacing the phenyl ring on **1** with other aromatic rings afforded the desired results. Both thiophenyl- (**1h**) and naphthyl-substituted (**1i**) cyclopropenes resulted in high yields (74 and 96%, respectively) and enantioselectivities (93:7 *er*). The absolute configuration of compound **4i** was confirmed by X-ray crystallographic analysis. An enhancement in enantioselectivity was observed with the incorporation of a methoxy substituent (**1j**, 96:4 *er*) or an amide substituent (**1k**, 99:1 *er*) on the cyclopropene. X-ray crystallographic analysis of **4k** suggested a potential directing group effect because pyrazole added *cis* to the amide substituent.^{12a} Cyclopropenes with a spirocycle (**1l**) and an ethyl substituent (**1m**) both exhibited high reactivity (63–80%) and enantioselectivity (94:6–95:5 *er*). Furthermore, a

dialkyl-substituted cyclopropene (**1n**) provided product **4n** in 80% yield with 89:11 *er*, although as a 1:1 mixture of diastereomers.^{12c,d,i}

On the basis of our own observations and literature precedent, we propose a mechanism for this Cu-catalyzed hydroamination (Figure 3). The catalyst resting state is




Figure 3. Proposed catalytic cycle.

inactive off-cycle copper dipyrazolate V. Dissociation of one pyrazolate releases active copper–amido catalyst I, which enters the catalytic cycle and binds to cyclopropene **1a**. A subsequent *cis*-aminocupration of π complex II forges the key C–N bond and provides cyclopropylcopper intermediate III.^{12e,15} Protodemetalation of III with DBU–H⁺ affords copper complex IV,²⁶ which then undergoes ligand exchange with pyrazole **2a** to restart the catalytic cycle. Mechanistic studies that led to the proposed mechanism are discussed below.

To probe the mechanism, we studied the kinetic profile using variable-time normalization analysis (VTNA) (see the SI).²⁷ We observed a first-order dependence on both cyclopropene (**1a**) and the copper catalyst. We found an inverse-first-order dependence on pyrazole (**2a**), and a fractional order (0.5) for DBU. Our lab has previously identified negative fractional orders of thiols in Rh-catalyzed hydrothiolations, which we attributed to the coordination of multiple thiols to an off-cycle catalyst resting state.^{12g,28} Given the coordinating ability of pyrazoles, we propose that pyrazole (**1a**) is involved in the formation of the off-cycle copper resting state with a 2:1 relative ratio of pyrazole per copper center. In the Heck coupling, Blackmond and co-workers observed that the order in Pd catalyst varied between first order and fractional order (0.5), depending on the amount of catalyst monomer released from an off-cycle dimer.²⁹ By performing VTNA at higher copper loadings, we observed an apparent fractional order (0.5) in copper (Figure 4A), which suggests the possibility of a Cu dimer off-cycle resting state.³⁰

We then performed ³¹P NMR studies to study the catalyst resting state (see the SI). Through monitoring the chemical shift of L3 in the reaction of cyclopropene **1a** with pyrazole **2a**,




Figure 4. Mechanistic studies.

we identified a plausible catalyst resting state at -2.4 ppm, which was replaced by another species bearing a singlet at -4.7 ppm when the transformation was near completion. NMR titrations³¹ using stoichiometric catalyst, DBU, and pyrazole suggest the former resonance to be a Cu–Duphos–pyrazolate species³² and the latter to be a Cu–Duphos–DBU complex. These NMR studies provide evidence for the dual role of DBU, acting as a base to deprotonate the pyrazole and as a ligand to copper.²⁶ The apparent fractional order of DBU may arise from its multiple roles in proton transfer and ligation.

Efforts to characterize the resting state under high copper concentrations led us to the serendipitous observation of trimeric copper species VI. X-ray crystallographic analysis revealed the unique structure where the central copper bridges two neighboring Cu–Duphos complexes via four pyrazolates (Figure 4B). The reactivity of this crystal was then tested under otherwise standard conditions, where the desired product was isolated in 20% yield with 94:6 *er*. Due to an unexpected partial oxidation during crystallization, the central copper appears divalent, which accounts for the lowered reactivity. The structure of VI supports the feasibility of copper–pyrazolates.³²

A deuterium labeling experiment was conducted under the standard conditions using deuterated cyclopropene *d*-1a and indazole 2q (Figure 4C). Analysis of *d*-3q shows exclusively *syn* proton incorporation relative to indazole. The results support the idea that C–N bond formation is an inner-sphere *cis*-aminocupration, as opposed to an outer-sphere nucleophilic addition,^{7e} followed by a retentive protodemetalation.

Although a four-centered 1,2-migratory insertion of cyclopropene into the Cu–N bond was initially envisioned, the N²:N¹ regioselectivity observed suggests that aminocupration may occur via a 1',6'-migratory insertion (Figure 5A). In the proposed five-centered mechanism,³³ the less-hindered nitro-



A. Potential *cis*-aminocupration pathwaysB. Transition states for *cis*-aminocupration pathways

Figure 5. Proposed C–N bond formation pathways and TSs for the favored five-centered 1',6'-migratory insertion of N² (TS_{CA-N2}) and the higher energy, disfavored four-centered 1,2-migratory insertion of N² (TS_{MI-N2}).

gen (N¹) coordinates with copper, and the C–N bond is forged between the cyclopropane and the more-hindered nitrogen atom (N²). Moreover, this mechanistic rationale is in line with Lee's observations on pyrazole hydroamination.^{9d}

To explore the unique regioselectivity, we performed a density functional theory (DFT) analysis of the Cu–L3-catalyzed coupling of cyclopropene 1a and pyrazole 2j to yield 3j. DFT calculations were performed at the M06-2X/6-311+G**/PCM(MeCN)//M06-2X/6-31G* level of theory,^{34–36} as implemented in Gaussian 16.³⁷ The transition structures (TSs) for aminocupration, namely, the five-centered TS versus the four-centered TS, were pursued for formation of both the N² and N¹ isomers of 3j (Figure 5B). We discovered that the five-centered aminocupration leading to addition at N² represents the lowest-energy TS (Figure 5B, TS_{CA-N2}). This favorable transition structure (TS_{CA-N2}) has a C–Cu bond forming at 2.22 Å, a C–N bond occurring at 2.31 Å, and a Cu–N bond remaining intact at 1.99 Å. An analogous TS leading to the minor N¹ regioisomer shows the coordination of N² to copper and C–N bond formation from N¹ (TS_{CA-N1}); this pathway is 2.7 kcal/mol higher in energy than TS_{CA-N2}. These predictions are consistent with the experimentally

observed N²:N¹ regioselectivity. Finally, addition of pyrazole **2j** into cyclopropane **1a** via a 1,2-migratory insertion mechanism is disfavored for the TSs leading to both regioisomers of **3j**; the TSs for insertion of N¹ and N² (TS_{MI-N1} and TS_{MI-N2}) are higher in energy than TS_{CA-N2} by 13.6 and 16.1 kcal/mol, respectively.

In summary, hydroamination presents an attractive approach for the enantioselective coupling of cyclopropenes and pyrazoles. Chiral *N*-cyclopropyl pyrazoles and structurally related heterocycles are prepared using an earth-abundant copper catalyst under mild reaction conditions with high regio-, diastereo-, and enantiocontrol. Mechanistic studies suggest a unique 1',6'-migratory insertion. This Cu–amido strategy complements the Cu–hydride approach to hydroamination and will guide future studies of N-heterocycle functionalization.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.3c02971>.

Cartesian coordinates (XYZ)

Experimental procedures, computational details, and spectroscopic data for all new compounds (PDF)

Accession Codes

CCDC 2250602–2250605 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, U.K.; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Authors

Jennifer S. Hirschi – Department of Chemistry, Binghamton University, Binghamton, New York 13902, United States; orcid.org/0000-0002-3470-0561; Email: jhirschi@binghamton.edu

Vy M. Dong – Department of Chemistry, University of California, Irvine, Irvine, California 92697, United States; orcid.org/0000-0002-8099-1048; Email: dongv@uci.edu

Authors

Minghao Wang – Department of Chemistry, University of California, Irvine, Irvine, California 92697, United States

Julie C. Simon – Department of Chemistry, University of California, Irvine, Irvine, California 92697, United States

Mengfei Xu – Department of Chemistry, University of California, Irvine, Irvine, California 92697, United States

Stephanie A. Corio – Department of Chemistry, Binghamton University, Binghamton, New York 13902, United States

Complete contact information is available at: <https://pubs.acs.org/10.1021/jacs.3c02971>

Author Contributions

[§]J.C.S. and M.X. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

V.M.D. acknowledges the National Institutes of Health (R35GM127071) and National Science Foundation (CHE-2751956457). J.S.H. acknowledges the National Institute of General Medical Sciences, National Institutes of Health (R35GM147183) and the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) Program, which is supported by the National Science Foundation (Grants 2138259, 2138286, 2138307, 2137603, and 2138296). We acknowledge Sovias AG for the generous donation of phosphine ligands (L4 and L5) and the Jarvo Lab for the use of their SFC instrument. We thank Dr. Shaozhen Nie (GSK) for helpful discussions.

■ REFERENCES

- (1) Ertl, P.; Schuhmann, T. A Systematic Cheminformatics Analysis of Functional Groups Occurring in Natural Products. *J. Nat. Prod.* **2019**, *82*, 1258–1263.
- (2) For reviews, see: (a) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals: Miniperspective. *J. Med. Chem.* **2014**, *57*, 10257–10274. (b) Kalaria, P. N.; Karad, S. C.; Raval, D. K. A Review on Diverse Heterocyclic Compounds as the Privileged Scaffolds in Antimalarial Drug Discovery. *Eur. J. Med. Chem.* **2018**, *158*, 917–936. (c) Kerru, N.; Gummidi, L.; Maddila, S.; Gangu, K. K.; Jonnalagadda, S. B. A Review on Recent Advances in Nitrogen-Containing Molecules and Their Biological Applications. *Molecules* **2020**, *25*, 1909. (d) Bhutani, P.; Joshi, G.; Raja, N.; Bachhav, N.; Rajanna, P. K.; Bhutani, H.; Paul, A. T.; Kumar, R. U.S. FDA Approved Drugs from 2015–June 2020: A Perspective. *J. Med. Chem.* **2021**, *64*, 2339–2381.
- (3) Karrouchi, K.; Radi, S.; Ramli, Y.; Taoufik, J.; Mabkhot, Y.; Al-aizari, F.; Ansar, M. Synthesis and Pharmacological Activities of Pyrazole Derivatives: A Review. *Molecules* **2018**, *23*, 134.
- (4) For reviews, see: (a) Beller, M.; Seayad, J.; Tillack, A.; Jiao, H. Catalytic Markovnikov and Anti-Markovnikov Functionalization of Alkenes and Alkynes: Recent Developments and Trends. *Angew. Chem., Int. Ed.* **2004**, *43*, 3368–3398. (b) Müller, T. E.; Hultsch, K. C.; Yus, M.; Foubelo, F.; Tada, M. Hydroamination: Direct Addition of Amines to Alkenes and Alkynes. *Chem. Rev.* **2008**, *108*, 3795–3892. (c) Huang, L.; Arndt, M.; Gooßen, K.; Heydt, H.; Gooßen, L. J. Late Transition Metal-Catalyzed Hydroamination and Hydroamidation. *Chem. Rev.* **2015**, *115*, 2596–2697. (d) Colonna, P.; Bezenine, S.; Gil, R.; Hannedouche, J. Alkene Hydroamination via Earth-Abundant Transition Metal (Iron, Cobalt, Copper and Zinc) Catalysis: A Mechanistic Overview. *Adv. Synth. Catal.* **2020**, *362*, 1550–1563.
- (5) For selected examples of recent hydroamination, see: (a) Pronin, S. V.; Tabor, M. G.; Jansen, D. J.; Shenvi, R. A. A Stereoselective Hydroamination Transform To Access Polysubstituted Indolizidines. *J. Am. Chem. Soc.* **2012**, *134*, 2012–2015. (b) Yang, Y.; Shi, S.-L.; Niu, D.; Liu, P.; Buchwald, S. L. Catalytic Asymmetric Hydroamination of Unactivated Internal Olefins to Aliphatic Amines. *Science* **2015**, *349*, 62–66. (c) Gurak, J. A.; Yang, K. S.; Liu, Z.; Engle, K. M. Directed, Regiocontrolled Hydroamination of Unactivated Alkenes via Protodepalladation. *J. Am. Chem. Soc.* **2016**, *138*, 5805–5808. (d) Musacchio, A. J.; Lainhart, B. C.; Zhang, X.; Naguib, S. G.; Sherwood, T. C.; Knowles, R. R. Catalytic Intermolecular Hydroaminations of Unactivated Olefins with Secondary Alkyl Amines. *Science* **2017**, *355*, 727–730. (e) Xi, Y.; Ma, S.; Hartwig, J. F. Catalytic Asymmetric Addition of an Amine N-H Bond across Internal Alkenes. *Nature* **2020**, *588*, 254–260. (f) Chinn, A. J.; Sedillo, K.; Doyle, A. G. Phosphine/Photoredox Catalyzed Anti-Markovnikov Hydroamination of Olefins with Primary Sulfonamides via α -Scission from Phosphoranyl Radicals. *J. Am. Chem. Soc.* **2021**, *143*, 18331–18338.
- (6) For selected examples of hydroamination with heterocycles, see: (a) Haydl, A. M.; Xu, K.; Breit, B. Regio- and Enantioselective Synthesis of N-Substituted Pyrazoles by Rhodium-Catalyzed Asym-

metric Addition to Allenes. *Angew. Chem., Int. Ed.* **2015**, *54*, 7149–7153. (b) Haydl, A. M.; Hilpert, L. J.; Breit, B. Regioconvergent and Enantioselective Rhodium-Catalyzed Hydroamination of Internal and Terminal Alkynes: A Highly Flexible Access to Chiral Pyrazoles. *Chem. - Eur. J.* **2016**, *22*, 6547–6551. (c) Hilpert, L. J.; Sieger, S. V.; Haydl, A. M.; Breit, B. Palladium- and Rhodium-Catalyzed Dynamic Kinetic Resolution of Racemic Internal Allenes Towards Chiral Pyrazoles. *Angew. Chem., Int. Ed.* **2019**, *58*, 3378–3381. (d) Jiang, W.; Ji, D.; Zhang, W.; Zhang, G.; Min, X.; Hu, Y.; Jiang, X.; Chen, Q. Orthogonal Regulation of Nucleophilic and Electrophilic Sites in Pd-Catalyzed Regiodivergent Couplings between Indazoles and Isoprene. *Angew. Chem., Int. Ed.* **2021**, *60*, 8321–8328. (e) Jiu, A. Y.; Slocumb, H. S.; Yeung, C. S.; Yang, X.; Dong, V. M. Enantioselective Addition of Pyrazoles to Dienes. *Angew. Chem., Int. Ed.* **2021**, *60*, 19660–19664.

(7) For selected examples of copper-amido implicated in cross-coupling, see: (a) Strieter, E. R.; Bhayana, B.; Buchwald, S. L. Mechanistic Studies on the Copper-Catalyzed N-Arylation of Amides. *J. Am. Chem. Soc.* **2009**, *131*, 78–88. (b) Mankad, N. P.; Antholine, W. E.; Szilagy, R. K.; Peters, J. C. Three-Coordinate Copper(I) Amido and Aminyl Radical Complexes. *J. Am. Chem. Soc.* **2009**, *131*, 3878–3880. (c) Giri, R.; Hartwig, J. F. Cu(I)-Amido Complexes in the Ullmann Reaction: Reactions of Cu(I)-Amido Complexes with Iodoarenes with and without Autocatalysis by CuI. *J. Am. Chem. Soc.* **2010**, *132*, 15860–15863. (d) Creutz, S. E.; Lotito, K. J.; Fu, G. C.; Peters, J. C. Photoinduced Ullmann C–N Coupling: Demonstrating the Viability of a Radical Pathway. *Science* **2012**, *338*, 647–651. (e) Kainz, Q. M.; Matier, C. D.; Bartoszewicz, A.; Zultanski, S. L.; Peters, J. C.; Fu, G. C. Asymmetric Copper-Catalyzed C–N Cross-Couplings Induced by Visible Light. *Science* **2016**, *351*, 681–684. (f) Liang, Y.; Zhang, X.; MacMillan, D. W. C. Decarboxylative sp^3 C–N Coupling via Dual Copper and Photoredox Catalysis. *Nature* **2018**, *559*, 83–88. (g) Chen, C.; Peters, J. C.; Fu, G. C. Photoinduced Copper-Catalyzed Asymmetric Amidation via Ligand Cooperativity. *Nature* **2021**, *596*, 250–256.

(8) For selected examples, see: (a) Blue, E. D.; Davis, A.; Conner, D.; Gunnoe, T. B.; Boyle, P. D.; White, P. S. Synthesis, Solid-State Crystal Structure, and Reactivity of a Monomeric Copper(I) Anilido Complex. *J. Am. Chem. Soc.* **2003**, *125*, 9435–9441. (b) Goj, L. A.; Blue, E. D.; Munro-Leighton, C.; Gunnoe, T. B.; Petersen, J. L. Cleavage of X–H Bonds (X = N, O, or C) by Copper(I) Alkyl Complexes To Form Monomeric Two-Coordinate Copper(I) Systems. *Inorg. Chem.* **2005**, *44*, 8647–8649. (c) Munro-Leighton, C.; Blue, E. D.; Gunnoe, T. B. Anti-Markovnikov N–H and O–H Additions to Electron-Deficient Olefins Catalyzed by Well-Defined Cu(I) Anilido, Ethoxide, and Phenoxide Systems. *J. Am. Chem. Soc.* **2006**, *128*, 1446–1447. (d) Goj, L. A.; Blue, E. D.; Delp, S. A.; Gunnoe, T. B.; Cundari, T. R.; Pierpont, A. W.; Petersen, J. L.; Boyle, P. D. Chemistry Surrounding Monomeric Copper(I) Methyl, Phenyl, Anilido, Ethoxide, and Phenoxide Complexes Supported by N-Heterocyclic Carbene Ligands: Reactivity Consistent with Both Early and Late Transition Metal Systems. *Inorg. Chem.* **2006**, *45*, 9032–9045.

(9) For selected examples, see: (a) Kim, S.; Kang, S.; Kim, G.; Lee, Y. Copper-Catalyzed Aza-Michael Addition of Aromatic Amines or Aromatic Aza-Heterocycles to α,β -Unsaturated Olefins. *J. Org. Chem.* **2016**, *81*, 4048–4057. (b) Perego, L. A.; Blicek, R.; Groué, A.; Monnier, F.; Taillefer, M.; Ciofini, I.; Grimaud, L. Copper-Catalyzed Hydroamination of Allenes: From Mechanistic Understanding to Methodology Development. *ACS Catal.* **2017**, *7*, 4253–4264. (c) Park, S.; Kang, S.; Lee, Y. Copper-Catalyzed Intermolecular Hydroamination of Arylamines or Aza-Heterocycles with Nitrostyrene Derivatives. *Adv. Synth. Catal.* **2019**, *361*, 1071–1083. (d) Kim, K.; Lee, Y. Copper-Catalyzed Hydroamination of Oxa- and Azabenzonorbornadienes with Pyrazoles. *J. Org. Chem.* **2022**, *87*, 569–578.

(10) For reviews on the importance of cyclopropane moiety in medicinal chemistry, see: (a) Talele, T. T. The “Cyclopropyl Fragment” Is a Versatile Player That Frequently Appears in Preclinical/Clinical Drug Molecules. *J. Med. Chem.* **2016**, *59*, 8712–

8756. (b) Shearer, J.; Castro, J. L.; Lawson, A. D. G.; MacCoss, M.; Taylor, R. D. Rings in Clinical Trials and Drugs: Present and Future. *J. Med. Chem.* **2022**, *65*, 8699–8712.

(11) Dian, L.; Marek, I. Asymmetric Preparation of Polysubstituted Cyclopropanes Based on Direct Functionalization of Achiral Three-Membered Carbocycles: Focus Review. *Chem. Rev.* **2018**, *118*, 8415–8434.

(12) For selected examples of the hydrofunctionalization of cyclopropenes, see: (a) Tarwade, V.; Liu, X.; Yan, N.; Fox, J. M. Directed Carbozincation Reactions of Cyclopropene Derivatives. *J. Am. Chem. Soc.* **2009**, *131*, 5382–5383. (b) Phan, D. H. T.; Kou, K. G. M.; Dong, V. M. Enantioselective Desymmetrization of Cyclopropenes by Hydroacylation. *J. Am. Chem. Soc.* **2010**, *132*, 16354–16355. (c) Parra, A.; Amenós, L.; Guisán-Ceinos, M.; López, A.; García Ruano, J. L.; Tortosa, M. Copper-Catalyzed Diastereo- and Enantioselective Desymmetrization of Cyclopropenes: Synthesis of Cyclopropylboronates. *J. Am. Chem. Soc.* **2014**, *136*, 15833–15836. (d) Teng, H.-L.; Luo, Y.; Wang, B.; Zhang, L.; Nishiura, M.; Hou, Z. Synthesis of Chiral Aminocyclopropanes by Rare-Earth-Metal-Catalyzed Cyclopropene Hydroamination. *Angew. Chem., Int. Ed.* **2016**, *55*, 15406–15410. (e) Li, Z.; Zhao, J.; Sun, B.; Zhou, T.; Liu, M.; Liu, S.; Zhang, M.; Zhang, Q. Asymmetric Nitron Synthesis via Ligand-Enabled Copper-Catalyzed Cope-Type Hydroamination of Cyclopropene with Oxime. *J. Am. Chem. Soc.* **2017**, *139*, 11702–11705. (f) Dian, L.; Marek, I. Rhodium-Catalyzed Arylation of Cyclopropenes Based on Asymmetric Direct Functionalization of Three Membered Carbocycles. *Angew. Chem., Int. Ed.* **2018**, *57*, 3682–3686. (g) Zhang, H.; Huang, W.; Wang, T.; Meng, F. Cobalt-Catalyzed Diastereo- and Enantioselective Hydroalkenylation of Cyclopropenes with Alkenylboronic Acids. *Angew. Chem., Int. Ed.* **2019**, *58*, 11049–11053. (h) Feng, S.; Hao, H.; Liu, P.; Buchwald, S. L. Diastereo- and Enantioselective CuH-Catalyzed Hydroamination of Strained Trisubstituted Allenes. *ACS Catal.* **2020**, *10*, 282–291. (i) Dian, L.; Marek, I. Pd-Catalyzed Enantioselective Hydroalkynylation of Cyclopropenes. *ACS Catal.* **2020**, *10*, 1289–1293. (j) Huang, W.; Meng, F. Cobalt-Catalyzed Diastereo- and Enantioselective Hydroalkylation of Cyclopropenes with Cobalt Homo-enolates. *Angew. Chem., Int. Ed.* **2021**, *60*, 2694–2698. (k) Nie, S.; Lu, A.; Kuker, E. L.; Dong, V. M. Enantioselective Hydrothiolation: Diverging Cyclopropenes through Ligand Control. *J. Am. Chem. Soc.* **2021**, *143*, 6176–6184. (l) Huang, M.-Y.; Zhao, Y.-T.; Chai, H.; Zhang, C.-D.; Zhu, S.-F. Copper-Catalyzed Ring-Opening/Borylation of Cyclopropenes. *CCS Chem.* **2022**, *4*, 1232–1237. (m) Yu, R.; Cai, S.; Li, C.; Fang, X. Nickel-Catalyzed Asymmetric Hydroaryloxy- and Hydroalkoxycarbonylation of Cyclopropenes. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202200733. (n) Huang, Q.; Chen, Y.; Zhou, X.; Dai, L.; Lu, Y. Nickel-Hydride-Catalyzed Diastereo- and Enantioselective Hydroalkylation of Cyclopropenes. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202210560.

(13) For reviews, see: (a) Hirano, K.; Miura, M. Hydroamination, Aminoboration, and Carboamination with Electrophilic Amination Reagents: Umpolung-Enabled Regio- and Stereoselective Synthesis of N-Containing Molecules from Alkenes and Alkynes. *J. Am. Chem. Soc.* **2022**, *144*, 648–661. (b) Mohite, S. B.; Bera, M.; Kumar, V.; Karpoomath, R.; Baba, S. B.; Kumbhar, A. S. O-Benzoylhydroxylamines: A Versatile Electrophilic Aminating Reagent for Transition Metal-Catalyzed C–N Bond-Forming Reactions. *Top. Curr. Chem.* **2023**, *381*, 4.

(14) (a) Zhu, S.; Niljianskul, N.; Buchwald, S. L. Enantio- and Regioselective CuH-Catalyzed Hydroamination of Alkenes. *J. Am. Chem. Soc.* **2013**, *135*, 15746–15749. (b) Miki, Y.; Hirano, K.; Satoh, T.; Miura, M. Copper-Catalyzed Intermolecular Regioselective Hydroamination of Styrenes with Polymethylhydrosiloxane and Hydroxylamines. *Angew. Chem., Int. Ed.* **2013**, *52*, 10830–10834. (c) Liu, R. Y.; Buchwald, S. L. CuH-Catalyzed Olefin Functionalization: From Hydroamination to Carbonyl Addition. *Acc. Chem. Res.* **2020**, *53*, 1229–1243.

(15) Paderes, M. C.; Belding, L.; Fanovic, B.; Dudding, T.; Keister, J. B.; Chemler, S. R. Evidence for Alkene Cis-Aminocupration, an

Aminooxygenation Case Study: Kinetics, EPR Spectroscopy, and DFT Calculations. *Chem. - Eur. J.* **2012**, *18*, 1711–1726.

(16) For selected examples of late-stage functionalization, see: (a) Wencel-Delord, J.; Glorius, F. C-H Bond Activation Enables the Rapid Construction and Late-Stage Diversification of Functional Molecules. *Nat. Chem.* **2013**, *5*, 369–375. (b) DiRocco, D. A.; Dykstra, K.; Krska, S.; Vachal, P.; Conway, D. V.; Tudge, M. Late-Stage Functionalization of Biologically Active Heterocycles Through Photoredox Catalysis. *Angew. Chem., Int. Ed.* **2014**, *53*, 4802–4806. (c) Sharma, A.; Hartwig, J. F. Metal-Catalyzed Azidation of Tertiary C-H Bonds Suitable for Late-Stage Functionalization. *Nature* **2015**, *517*, 600–604. (d) Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. W. The Medicinal Chemist's Toolbox for Late-Stage Functionalization of Drug-like Molecules. *Chem. Soc. Rev.* **2016**, *45*, 546–576. (e) Gianatassio, R.; Lopchuk, J. M.; Wang, J.; Pan, C.-M.; Malins, L. R.; Prieto, L.; Brandt, T. A.; Collins, M. R.; Gallego, G. M.; Sach, N. W.; Spangler, J. E.; Zhu, H.; Zhu, J.; Baran, P. S. Strain-Release Amination. *Science* **2016**, *351*, 241–246. (f) Shang, M.; Wang, M.; Saint-Denis, T. G.; Li, M.; Dai, H.; Yu, J. Copper-Mediated Late-Stage Functionalization of Heterocycle-Containing Molecules. *Angew. Chem., Int. Ed.* **2017**, *56*, 5317–5321. (g) Le, C.; Liang, Y.; Evans, R. W.; Li, X.; MacMillan, D. W. C. Selective sp^3 C-H Alkylation via Polarity-Match-Based Cross-Coupling. *Nature* **2017**, *547*, 79–83. (h) Uehling, M. R.; King, R. P.; Krska, S. W.; Cernak, T.; Buchwald, S. L. Pharmaceutical Diversification via Palladium Oxidative Addition Complexes. *Science* **2019**, *363*, 405–408. (i) Feng, K.; Quevedo, R. E.; Kohrt, J. T.; Oderinde, M. S.; Reilly, U.; White, M. C. Late-Stage Oxidative $\text{C}(\text{sp}^3)\text{-H}$ Methylation. *Nature* **2020**, *580*, 621–627. (j) Zhang, L.; Ritter, T. A Perspective on Late-Stage Aromatic C-H Bond Functionalization. *J. Am. Chem. Soc.* **2022**, *144*, 2399–2414.

(17) Trost, B. The Atom Economy—A Search for Synthetic Efficiency. *Science* **1991**, *254*, 1471–1477.

(18) For reviews, see: (a) Vicente, R. C-C Bond Cleavages of Cyclopropanes: Operating for Selective Ring-Opening Reactions. *Chem. Rev.* **2021**, *121*, 162–226. (b) Huo, H.; Gong, Y. Construction of Heterocyclic Rings from Cyclopropanes. *Org. Biomol. Chem.* **2022**, *20*, 3847–3869.

(19) For selected examples, see: (a) Lee, E. C.; Fu, G. C. Copper-Catalyzed Asymmetric N-H Insertion Reactions: Couplings of Diazo Compounds with Carbamates to Generate α -Amino Acids. *J. Am. Chem. Soc.* **2007**, *129*, 12066–12067. (b) Phan, D. T. H.; Dong, V. M. Silver-Catalyzed Ring-Opening of Cyclopropanes: Preparation of Tertiary α -Branched Allylic Amines. *Tetrahedron* **2013**, *69*, 5726–5731. (c) Hyde, S.; Veliks, J.; Liégault, B.; Grassi, D.; Taillefer, M.; Gouverneur, V. Copper-Catalyzed Insertion into Heteroatom-Hydrogen Bonds with Trifluorodiazalkanes. *Angew. Chem., Int. Ed.* **2016**, *55*, 3785–3789. (d) Arredondo, V.; Hiew, S. C.; Gutman, E. S.; Premachandra, I. D. U. A.; Van Vranken, D. L. Enantioselective Palladium-Catalyzed Carbene Insertion into the N-H Bonds of Aromatic Heterocycles. *Angew. Chem., Int. Ed.* **2017**, *56*, 4156–4159. (e) Li, M.-L.; Yu, J.-H.; Li, Y.-H.; Zhu, S.-F.; Zhou, Q.-L. Highly Enantioselective Carbene Insertion into N-H Bonds of Aliphatic Amines. *Science* **2019**, *366*, 990–994.

(20) Muller, P. Glossary of Terms Used in Physical Organic Chemistry (IUPAC Recommendations 1994). *Pure Appl. Chem.* **1994**, *66*, 1077–1184.

(21) For examples of regioselective functionalization of more sterically hindered pyrazole nitrogen, see: (a) Xu, D.; Frank, L.; Nguyen, T.; Stumpf, A.; Russell, D.; Angelaud, R.; Gosselin, F. Magnesium-Catalyzed N2-Regioselective Alkylation of 3-Substituted Pyrazoles. *Synlett* **2020**, *31*, 595–599. (b) Bengel, L. L.; Aberle, B.; Egler-Kemmerer, A.; Kienzle, S.; Hauer, B.; Hammer, S. C. Engineered Enzymes Enable Selective N-Alkylation of Pyrazoles with Simple Haloalkanes. *Angew. Chem., Int. Ed.* **2021**, *60*, 5554–5560. (c) Chen, S.-J.; Golden, D. L.; Krska, S. W.; Stahl, S. S. Copper-Catalyzed Cross-Coupling of Benzylic C-H Bonds and Azoles with Controlled N-Site Selectivity. *J. Am. Chem. Soc.* **2021**, *143*, 14438–14444.

(22) For selected examples of regioselective functionalization of pyrazoles, see: (a) Antilla, J. C.; Baskin, J. M.; Barder, T. E.; Buchwald, S. L. Copper-Diamine-Catalyzed N-Arylation of Pyrroles, Pyrazoles, Imidazoles, and Triazoles. *J. Org. Chem.* **2004**, *69*, 5578–5587. (b) Goikhman, R.; Jacques, T. L.; Sames, D. C-H Bonds as Ubiquitous Functionality: A General Approach to Complex Arylated Pyrazoles via Sequential Regioselective C-Arylation and N-Alkylation Enabled by SEM-Group Transposition. *J. Am. Chem. Soc.* **2009**, *131*, 3042–3048. (c) Ye, Y.; Kevlishvili, I.; Feng, S.; Liu, P.; Buchwald, S. L. Highly Enantioselective Synthesis of Indazoles with a C3-Quaternary Chiral Center Using CuH Catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 10550–10556. (d) Dow, N. W.; Cabré, A.; MacMillan, D. W. C. A General N-Alkylation Platform via Copper Metallaphotoredox and Silyl Radical Activation of Alkyl Halides. *Chem* **2021**, *7*, 1827–1842. (e) Das, M.; Zamani, L.; Bratcher, C.; Musacchio, P. Z. Azolation of Benzylic C-H Bonds via Photoredox-Catalyzed Carbocation Generation. *J. Am. Chem. Soc.* **2023**, *145*, 3861–3868.

(23) For reviews, see: (a) Singh, J.; Sharma, D.; Bansal, R. Pyridazinone: An Attractive Lead for Anti-Inflammatory and Analgesic Drug Discovery. *Future Med. Chem.* **2017**, *9*, 95–127. (b) Dubey, S.; Bhosle, P. A. Pyridazinone: An Important Element of Pharmacophore Possessing Broad Spectrum of Activity. *Med. Chem. Res.* **2015**, *24*, 3579–3598.

(24) For reviews, see: (a) Denya, I.; Malan, S. F.; Joubert, J. Indazole Derivatives and Their Therapeutic Applications: A Patent Review (2013–2017). *Expert Opin. Ther. Pat.* **2018**, *28*, 441–453. (b) Mal, S.; Malik, U.; Mahapatra, M.; Mishra, A.; Pal, D.; Paidesetty, S. K. A Review on Synthetic Strategy, Molecular Pharmacology of Indazole Derivatives, and Their Future Perspective. *Drug Dev. Res.* **2022**, *83*, 1469–1504.

(25) The low yields of **3s** and **3t** may arise from the electron deficiency of **2s** and **2t**.

(26) Zall, C. M.; Linehan, J. C.; Appel, A. M. Triphosphine-Ligated Copper Hydrides for CO_2 Hydrogenation: Structure, Reactivity, and Thermodynamic Studies. *J. Am. Chem. Soc.* **2016**, *138*, 9968–9977.

(27) For a given species A, VTNA experiments are conducted by plotting the product formation against time normalized to various orders of $[A]$ (0, 1, 2, fractional, etc.). Overlays are attained when the assumed order is accurate. For more details, see the SI and: Burés, J. Variable Time Normalization Analysis: General Graphical Elucidation of Reaction Orders from Concentration Profiles. *Angew. Chem., Int. Ed.* **2016**, *55*, 16084–16087.

(28) (a) Yang, X.-H.; Davison, R. T.; Dong, V. M. Catalytic Hydrothiolation: Regio- and Enantioselective Coupling of Thiols and Dienes. *J. Am. Chem. Soc.* **2018**, *140*, 10443–10446. (b) Yang, X.-H.; Davison, R. T.; Nie, S.-Z.; Cruz, F. A.; McGinnis, T. M.; Dong, V. M. Catalytic Hydrothiolation: Counterion-Controlled Regioselectivity. *J. Am. Chem. Soc.* **2019**, *141*, 3006–3013.

(29) (a) Rosner, T.; Le Bars, J.; Pfaltz, A.; Blackmond, D. G. Kinetic Studies of Heck Coupling Reactions Using Palladacycle Catalysts: Experimental and Kinetic Modeling of the Role of Dimer Species. *J. Am. Chem. Soc.* **2001**, *123*, 1848–1855. (b) Burés, J. A Simple Graphical Method to Determine the Order in Catalyst. *Angew. Chem., Int. Ed.* **2016**, *55*, 2028–2031.

(30) For examples of off-cycle catalysts, see: (a) Hein, J. E.; Armstrong, A.; Blackmond, D. G. Kinetic Profiling of Prolinate-Catalyzed α -Amination of Aldehydes. *Org. Lett.* **2011**, *13*, 4300–4303. (b) Bandar, J. S.; Pirnot, M. T.; Buchwald, S. L. Mechanistic Studies Lead to Dramatically Improved Reaction Conditions for the Cu-Catalyzed Asymmetric Hydroamination of Olefins. *J. Am. Chem. Soc.* **2015**, *137*, 14812–14818. (c) Blackmond, D. G. Kinetic Profiling of Catalytic Organic Reactions as a Mechanistic Tool. *J. Am. Chem. Soc.* **2015**, *137*, 10852–10866.

(31) Gibbons, S. K.; Valteau, C. R. D.; Peltier, J. L.; Cain, M. F.; Hughes, R. P.; Glueck, D. S.; Golen, J. A.; Rheingold, A. L. Diastereoselective Coordination of P-Stereogenic Secondary Phosphines in Copper(I) Chiral Bis(Phosphine) Complexes: Structure, Dynamics, and Generation of Phosphido Complexes. *Inorg. Chem.* **2019**, *58*, 8854–8865.

(32) For examples, see: (a) Titov, A. A.; Filippov, O. A.; Smol'yakov, A. F.; Baranova, K. F.; Titova, E. M.; Averin, A. A.; Shubina, E. S. Dinuclear Cu^I and Ag^I Pyrazolates Supported with Tertiary Phosphines: Synthesis, Structures, and Photophysical Properties. *Eur. J. Inorg. Chem.* **2019**, 2019, 821–827. (b) Watanabe, Y.; Washer, B. M.; Zeller, M.; Savikhin, S.; Slipchenko, L. V.; Wei, A. Copper(I)-Pyrazolate Complexes as Solid-State Phosphors: Deep-Blue Emission through a Remote Steric Effect. *J. Am. Chem. Soc.* **2022**, 144, 10186–10192.

(33) Luo, Y.; Hou, Z. A Five-Center Rather than a Four-Center Transition State for Alkene Insertion into the Metal-Alkyl Bond of a Cationic Binuclear Yttrium Complex. *Organometallics* **2006**, 25, 6162–6165.

(34) Zhao, Y.; Truhlar, D. G. The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, non-covalent interactions, excited states, and transition elements: two new functionals and systematic testing of four M06-class functionals and 12 other functionals. *Theor. Chem. Acc.* **2008**, 120, 215–241.

(35) Hehre, W. J.; Stewart, R. F.; Pople, J. A. Self-Consistent Molecular-Orbital Methods. I. Use of Gaussian Expansions of Slater-Type Atomic Orbitals. *J. Chem. Phys.* **1969**, 51, 2657–2664.

(36) Validation of the five-centered aminocupration transition state (TS_{CA-N2}) was achieved by locating the corresponding transition state at both the ω B97XD/def2-TZVP PCM(MeCN)//B97D/def2-SVP and B3LYP-D3BJ/6-311+G** PCM(MeCN)//B3LYP-D3BJ/6-31G* levels of theory.

(37) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Petersson, G. A.; Nakatsuji, H.; Li, X.; Caricato, M.; Marenich, A.; Bloino, J.; Janesko, B. G.; Gomperts, R.; Mennucci, B.; Hratchian, H. P.; Ortiz, J. V.; Izmaylov, A. F.; Sonnenberg, J. L.; Williams-Young, D.; Ding, F.; Lipparini, F.; Egidi, F.; Goings, J.; Peng, B.; Petrone, A.; Henderson, T.; Ranasinghe, D.; Zakrzewski, V. G.; Gao, J.; Rega, N.; Zheng, G.; Liang, W.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Throssell, K.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E. N.; Kudin, K. N.; Staroverov, V. N.; Keith, T. A.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A. P.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Millam, J. M.; Klene, M.; Adamo, C.; Cammi, R.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Farkas, O.; Foresman, J. B.; Fox, D. J. *Gaussian 16*, rev. C.01; Gaussian, Inc.: Wallingford, CT, 2016.