

2D Voigt Boussinesq Equations

Mihaela Ignatova

ABSTRACT. We consider a critical conservative Voigt regularization of the 2D incompressible Boussinesq system on the torus. We prove the existence and uniqueness of global smooth solutions and their convergence in the smooth regime to the Boussinesq solution when the regularizations are removed. We also consider a range of mixed (subcritical-supercritical) Voigt regularizations for which we prove the existence of global smooth solutions.

1. Introduction

The Boussinesq equations are basic models of incompressible fluids in which density variations are due to variations of the temperature [4]. They form the basis of studies of thermally generated turbulence, with applications to geophysics and theoretical physics. In addition, in the absence of molecular friction due to viscosity and neglecting thermal diffusivity, in two spatial dimensions, the Boussinesq equations are similar to the 3D axisymmetric incompressible Euler equations, and as such they have been extensively studied numerically and theoretically in the context of the famous problems of finite time singularities. This connection between the 2D Boussinesq and 3D axisymmetric Euler singularities dates back at least as far as December 1991 [26], but it had a significant resurgence in recent years [5, 6, 9, 10, 11].

Fractional dissipative Boussinesq equations generalize the original dissipative system. The equations have the form

$$\partial_t u + \nu \Lambda^\alpha u + u \cdot \nabla u + \nabla p = Kg\theta e_2, \quad (1)$$

$$\nabla \cdot u = 0, \quad (2)$$

$$\partial_t \theta + \kappa \Lambda^\beta \theta + u \cdot \nabla \theta = 0 \quad (3)$$

where u is the two dimensional divergence-free velocity, θ is the temperature, p is the pressure. Above $\Lambda = (-\Delta)^{\frac{1}{2}}$ is the square root of the periodic Laplacian. For $s \in \mathbb{R}$, the periodic fractional Laplacian Λ^s applied to a mean zero function f is a Fourier multiplier with symbol $|k|^s$, that is, for f given by

$$f = \sum_{k \in \mathbb{Z}^2 \setminus \{0\}} f_k e^{ik \cdot x}, \quad (4)$$

and obeying

$$\sum_{k \in \mathbb{Z}^2 \setminus \{0\}} |k|^{2s} |f_k|^2 < \infty, \quad (5)$$

we have

$$\Lambda^s f = \sum_{k \in \mathbb{Z}^2 \setminus \{0\}} |k|^s f_k e^{ik \cdot x}. \quad (6)$$

In (1), the vector e_2 is the unit vector pointing in the direction opposite to gravity. When $\alpha = \beta = 2$, then we have the classical system, and $\nu > 0$ is the kinematic viscosity, $\kappa > 0$ is the thermal diffusivity, g is the constant of gravitational acceleration, and $K > 0$ is a constant thermal expansion coefficient. The reference constant density is taken to be 1.

Global existence and uniqueness of regular solutions have been proved for the fractional dissipative Boussinesq system, when $\alpha = 2, \nu > 0, \kappa = 0$ or when $\nu = 0, \beta = 2, \kappa > 0$ in [2] and in [15] for smooth

Key words and phrases. Boussinesq, Voigt regularization, global existence.

MSC Classification: 35Q30, 35Q35, 35Q92.

enough initial data. See also [18] for persistence of regularity with less smooth initial data. Several works [8, 21] treat anisotropic viscosity cases and [16] considers regularity in bounded domains.

Smaller powers of the fractional Laplacians have been successfully considered in [13, 14] with $\alpha = 1, \nu > 0, \kappa = 0$ or $\nu = 0, \beta = 1, \kappa > 0$. Global regularity for the 2D Boussinesq equations with $\nu\kappa > 0$ and fractional dissipation was obtained in [28] when $\alpha + \beta = 1, \alpha > 0.798103\dots$ and $\beta > 0$ and $(u_0, \theta_0) \in (H^s(\mathbb{R}^2))^2$ for $s > 2$ (see also [12]).

In this paper, we consider a Voigt regularization of the Boussinesq system. Voigt (or Kelvin-Voigt) equations have been introduced to model complex fluids with polymeric interactions [25]. Mathematically, they have been widely studied as Voigt regularizations in the context of incompressible fluid dynamics and magneto-hydrodynamics [1, 19, 20, 23]. Equations with Voigt regularizations have been used to obtain statistical solutions [22, 27], in the limit of the regularization parameter tending to zero. More recently, a Voigt regularization approach was used in [7] to tackle the problem of magnetic reconnection (topological change of magnetic field lines) and asymptotic approach to equilibrium.

As opposed to viscous regularizations, the Voigt regularizations are not dissipative, rather they are conservative. They work by regularizing the conserved energy. They have the advantage of preserving the steady states of the system, and are hence well suited for long time behavior studies, and also, they do not introduce spurious boundary layers. On the other hand, the Voigt regularizations do not respect the transport structure of the equations.

We consider the 2D Voigt Boussinesq equation,

$$(I + \epsilon\Lambda)\partial_t u + u \cdot \nabla u + \nabla p = \theta e_2, \quad (7)$$

$$\nabla \cdot u = 0, \quad (8)$$

$$(I + \epsilon\Lambda)\partial_t \theta + u \cdot \nabla \theta = 0 \quad (9)$$

on the torus $\mathbb{T}^2$, where $u = (u_1, u_2): \mathbb{T}^2 \times [0, T] \rightarrow \mathbb{R}^2$ is the fluid velocity, $\theta: \mathbb{T}^2 \times [0, T] \rightarrow \mathbb{R}$ is the temperature, I is the identity operator, $\Lambda = (-\Delta)^{\frac{1}{2}}$, $\epsilon > 0$ is a parameter, and $e_2 = (0, 1)$. We study the initial value problem for the system (7)–(9) with initial data

$$u(\cdot, 0) = u_0 \quad \text{and} \quad \theta(\cdot, 0) = \theta_0. \quad (10)$$

The curl $\omega = \nabla^\perp \cdot u$, with $\nabla^\perp = (-\partial_2, \partial_1)$ obeys the vorticity equation

$$(I + \epsilon\Lambda)\partial_t \omega + u \cdot \nabla \omega = \partial_1 \theta.$$

Thus, the Voigt Boussinesq equations (7)–(9) can be closed in terms of vorticity and temperature (ω, θ) ,

$$(I + \epsilon\Lambda)\partial_t \omega + u \cdot \nabla \omega = \partial_1 \theta, \quad (11)$$

$$u = \nabla^\perp \Delta^{-1} \omega, \quad (12)$$

$$(I + \epsilon\Lambda)\partial_t \theta + u \cdot \nabla \theta = 0 \quad (13)$$

with initial conditions

$$\omega(\cdot, 0) = \omega_0 = \nabla^\perp \cdot u_0 \quad \text{and} \quad \theta(\cdot, 0) = \theta_0. \quad (14)$$

Our main result is the global regularity of the Voigt Boussinesq equations.

THEOREM 1. *Let $s > 1$ and let $(\omega_0, \theta_0) \in (H^s(\mathbb{T}^2))^2$. Let T be arbitrary. Then, there exists a unique solution of 2D Voigt Boussinesq equations $(\omega, \theta) \in C([0, T]; (H^s(\mathbb{T}^2))^2)$ with initial data (ω_0, θ_0) .*

The proof is based on the following elements. We first prove the solutions (ω, θ) exist and are unique locally in time in the framework of Sobolev spaces $H^s \times H^s$, as soon as $s > 1$ (Theorem 6). Then, for $s > 1$, for any local solution defined on some time interval, we exploit the fundamental conservation property of Voigt regularizations. This conservation is shown to imply a priori information on the velocity $u \in L^\infty(0, T, H^{\frac{3}{2}})$, with bounds that depend only on the initial data and grow at most linearly in T . This is used, in conjunction with the local existence and uniqueness theorem to obtain global unique solutions in the Sobolev phase space $H^s \times H^s$ for $1 < s \leq \frac{3}{2}$. Next, we prove a natural Beale-Kato-Majda-type theorem

(Theorem 7) that gives sufficient conditions for persistence of regularity which do not involve the Lipschitz norm of the temperature. Finally, based on the Theorem 7, we prove that global regularity holds for any $s > \frac{3}{2}$.

The inviscid 2D Boussinesq system has a local existence theory which is similar to that of the 3D incompressible axisymmetric Euler equations.

THEOREM 2. *Let $s > 1$ and let $(\omega_0, \theta_0) \in (H^s(\mathbb{T}^2))^2$. There exists a time $T_0 > 0$ and a unique solution of the Boussinesq system (corresponding to $\epsilon = 0$ in (11)–(13)) $(\omega, \theta) \in C([0, T_0]; (H^s(\mathbb{T}^2))^2)$ with initial data (ω_0, θ_0) .*

The proof of this result is done by energy methods [24, Chapter 3, Section 2]. The fundamental open problem of blow up is whether or not a finite maximal time of existence $T_0 < \infty$ can arise from some smooth initial data.

In view of the interest in the blow up problem, it is important to study various relaxations of the regularization. In this paper we prove that the limit of vanishing regularization is the original equation, in a smooth enough regime.

THEOREM 3. *Let $(\omega_0, \theta_0) \in H^s(\mathbb{T}^2) \times H^{s+1}(\mathbb{T}^2)$, $s > 1$ and let $(\omega_B, \theta_B) \in L^\infty(0, T; H^s(\mathbb{T}^2) \times H^{s+1}(\mathbb{T}^2))$ be a solution of the 2D Boussinesq system on $[0, T]$. Then the solutions $(\omega_\epsilon, \theta_\epsilon)$ of the 2D Voigt Boussinesq equations with the same initial data converge as $\epsilon \rightarrow 0$ in $L^\infty(0, T; H^{-1}(\mathbb{T}^2) \times L^2(\mathbb{T}^2))$ to the solution (ω_B, θ_B) of the 2D Boussinesq system.*

REMARK 1. *The $s + 1$ regularity requirement for θ is needed because of the conservative nature of the Boussinesq equations, to ensure $\nabla \theta \in L^\infty$ uniformly in ϵ .*

We consider also fractional Voigt Boussinesq equations and prove global regularity for certain cases whereby lower powers of the fractional Laplacian are used for the temperature field.

THEOREM 4. *Let $s > 1$. Let $\alpha, \beta \geq 0$ with $\alpha + \beta \geq 2$, $\alpha > 1$, $\beta \geq \frac{2}{3}$. Let $(\omega_0, \theta_0) \in (H^s(\mathbb{T}^2))^2$. Let $T > 0$ be arbitrary. Then there exists a unique solution $(\omega, \theta) \in L^\infty(0, T; (H^s(\mathbb{T}^2))^2)$ of the fractional Voigt Boussinesq system*

$$\begin{cases} (I + \epsilon \Lambda)^\alpha \partial_t \omega + u \cdot \nabla \omega = \partial_1 \theta, \\ u = \nabla^\perp \Delta^{-1} \omega, \\ (I + \epsilon \Lambda)^\beta \partial_t \theta + u \cdot \nabla \theta = 0. \end{cases} \quad (15)$$

The proof of this result is based on commutator estimates and on bounds on θ in H^1 . We obtain also

THEOREM 5. *Let $\alpha > 2$, and $\beta = 0$. Let $(\omega_0, \theta_0) \in (H^s(\mathbb{T}^2))^2$. Let $T > 0$ be arbitrary. Then there exists a unique solution $(\omega, \theta) \in L^\infty(0, T; (H^s(\mathbb{T}^2))^2)$ of the fractional Voigt Boussinesq system (15)*

REMARK 2. *Theorem 4 and Theorem 5 imply regularity for $\beta \geq \frac{2}{3}$ when $\alpha \geq \frac{4}{3}$ and for $\beta = 0$ when $\alpha > 2$. It would be natural to conjecture global regularity for the whole range $\alpha + \beta \geq 2$, but at present we do not know how to prove this result.*

2. Proof of Theorem 1

2.1. Local existence and uniqueness in H^s , $s > 1$. The 2D Voigt Boussinesq equations can be written in divergence form as

$$\begin{cases} \partial_t \omega = -(I + \epsilon \Lambda)^{-1} \nabla \cdot (u \omega) + (I + \epsilon \Lambda)^{-1} \partial_1 \theta \\ \partial_t \theta = -(I + \epsilon \Lambda)^{-1} \nabla \cdot (u \theta), \end{cases} \quad (16)$$

where we used that u is divergence-free.

THEOREM 6. *Let $(\omega_0, \theta_0) \in (H^s(\mathbb{T}^2))^2$ with $s > 1$. There exists a time T depending only on $\epsilon > 0$ and the norms of ω_0 and θ_0 in $H^s(\mathbb{T}^2)$ and a unique solution (ω, θ) of the 2D Voigt Boussinesq equations, with initial data (ω_0, θ_0) and with $(\omega, \theta) \in L^\infty(0, T; (H^s(\mathbb{T}^2))^2)$.*

Proof of Theorem 6. The nonlinearity in the 2D Voigt Boussinesq equations is

$$\begin{pmatrix} \omega \\ \theta \end{pmatrix} \mapsto \begin{pmatrix} -(I + \epsilon \Lambda)^{-1} \nabla \cdot (u\omega) + (I + \epsilon \Lambda)^{-1} \partial_1 \theta \\ -(I + \epsilon \Lambda)^{-1} \nabla \cdot (u\theta) \end{pmatrix} = \begin{pmatrix} N_1(\omega, \theta) \\ N_2(\omega, \theta) \end{pmatrix}. \quad (17)$$

This follows from (16) by using that u is divergence-free. For $s > 1$, we have that $H^s(\mathbb{T}^2)$ is a Banach algebra, and that $H^s(\mathbb{T}^2) \subset L^\infty(\mathbb{T}^2)$. Moreover, the map $(I + \epsilon \Lambda)^{-1} \nabla$ is bounded in $H^s(\mathbb{T}^2)$. We deduce

$$\begin{aligned} \|N_1(\omega, \theta)\|_{H^s} &\leq C \|\omega\|_{H^s}^2 + \|\theta\|_{H^s} \\ \|N_2(\omega, \theta)\|_{H^s} &\leq C \|\omega\|_{H^s} \|\theta\|_{H^s} \end{aligned} \quad (18)$$

by using also the Poincaré inequality $\|u\|_{H^s} \leq C \|\omega\|_{H^s}$. The local existence and uniqueness of solutions follows from a fixed point argument and the fact that the nonlinearity (bilinear continuous plus linear continuous) is Lipschitz in the function space $(H^s(\mathbb{T}^2))^2$:

$$\|N(w_1) - N(w_2)\|_{(H^s(\mathbb{T}^2))^2} \leq L_B \|w_1 - w_2\|_{(H^s(\mathbb{T}^2))^2}, \quad w_1, w_2 \in B,$$

where $w_i = (\omega_i, \theta_i)$ and B is a ball in $(H^s(\mathbb{T}^2))^2$. We omit further details. $\square$

From now on, until we discuss convergence as $\epsilon \rightarrow 0$, for simplicity of exposition we set $\epsilon = 1$.

2.2. Basic Voigt energy bounds on H^s , $s > 1$ solutions of the 2D Voigt Boussinesq equations.

Multiplying the temperature equation (13) by θ and integrating, we obtain

$$\|\theta\|_{L^2}^2 + \|\theta\|_{H^{\frac{1}{2}}}^2 = \|\theta_0\|_{L^2}^2 + \|\theta_0\|_{H^{\frac{1}{2}}}^2 = A_0. \quad (19)$$

Multiplying the vorticity equation (11) by ω and integrating, we have

$$\frac{d}{dt} (\|\omega\|_{L^2}^2 + \|\omega\|_{H^{\frac{1}{2}}}^2) = 2 \int_{\mathbb{T}^2} (\partial_1 \theta) \omega dx \leq 2 \|\omega\|_{H^{\frac{1}{2}}} \sqrt{A_0}. \quad (20)$$

It follows that

$$\frac{d}{dt} \|\omega\|_{H^{\frac{1}{2}}} \leq \sqrt{A_0}, \quad (21)$$

$$\|\omega(\cdot, t)\|_{H^{\frac{1}{2}}} \leq \|\omega_0\|_{H^{\frac{1}{2}}} + t \sqrt{A_0}, \quad (22)$$

and thus $(\omega, \theta) \in L^\infty(0, T; (H^{\frac{1}{2}}(\mathbb{T}^2))^2)$ with a priori bounds in terms of initial data that grow at most linearly in T .

2.3. Proof of global existence and uniqueness for $1 < s \leq \frac{3}{2}$. The aim is to show that local solutions in H^s for $1 < s \leq \frac{3}{2}$ satisfy H^s bounds

$$\frac{d}{dt} (\|\theta\|_{H^s} + \|\omega\|_{H^s}) \leq C (\|u\|_{H^s} + 1) (\|\theta\|_{H^s} + \|\omega\|_{H^s}). \quad (23)$$

Then, because of the a priori bound due to the basic energy structure (22), specifically that $\omega \in L^\infty(H^{\frac{1}{2}})$, it follows that $u \in L^\infty(H^{\frac{3}{2}}) \subset L^\infty(H^s)$ with bounds that depend only on initial data, and grow at most linearly in time. The inequality (23) implies that the H^s norms of ω and θ are finite. Thus, by the local existence and uniqueness in H^s , obtained in Theorem 6, the solution can be continued indefinitely.

In order to obtain (23) we use the calculus inequality

$$\|fg\|_{H^s} \leq C (\|f\|_{L^\infty} \|g\|_{H^s} + \|f\|_{H^s} \|g\|_{L^\infty})$$

valid for any $s > 0$, to obtain from (16), using the boundedness of Riesz transforms in H^s spaces,

$$\frac{d}{dt} (\|\omega\|_{H^s} + \|\theta\|_{H^s}) \leq \|u\|_{L^\infty} (\|\omega\|_{H^s} + \|\theta\|_{H^s}) + (\|\omega\|_{L^\infty} + \|\theta\|_{L^\infty}) \|u\|_{H^s} + \|\theta\|_{H^s}. \quad (24)$$

We then apply the embedding $H^s \subset L^\infty$ to estimate $\|\omega\|_{L^\infty} + \|\theta\|_{L^\infty}$ by $\|\theta\|_{H^s} + \|\omega\|_{H^s}$ and $\|u\|_{L^\infty}$ by $\|u\|_{H^s}$. $\square$

2.4. A Beale-Kato-Majda-type theorem. We give here a regularity criterion which can be used to uniquely continue solutions.

THEOREM 7. *Let $s > 1$ and let $(\omega_0, \theta_0) \in (H^s(\mathbb{T}^2))^2$. We define the maximum existence time $T^* > 0$ to be*

$$T^* = \sup\{T > 0 \mid (\omega, \theta) \in C([0, T]; (H^s(\mathbb{T}^2))^2)\}$$

where (ω, θ) is the solution of the 2D Voigt-Boussinesq equations with initial data (ω_0, θ_0) . Then $T^ < \infty$ implies*

$$\lim_{T \rightarrow T^*} \int_0^T (\|\omega(\cdot, t)\|_{L^\infty} + \|\theta(\cdot, t)\|_{L^\infty}) dt = \infty. \quad (25)$$

Proof of Theorem 7. We return to the inequality (24). From the basic energy estimates, (ω, θ) are bounded $L^\infty(0, T; H^{\frac{1}{2}}(\mathbb{T}^2)^2)$ in terms of the initial data and T . The Sobolev embedding $H^{\frac{3}{2}} \subset L^\infty$ implies in particular that

$$\int_0^T \|u\|_{L^\infty} dt < \infty$$

is bounded a priori in terms of initial data and T , so that the only amplification factor that remains to be controlled in (24) is $\|\omega\|_{L^\infty} + \|\theta\|_{L^\infty}$, which is then controlled by the assumption (25). We omit further details. $\square$

REMARK 3. *We note that for the Voigt-Boussinesq system the Beale-Kato-Majda criterion does not require the finiteness of $\|\nabla\theta\|_{L^1(0, T; L^\infty)}$, as opposed to the criterion for the Boussinesq system [3].*

2.5. Proof of global existence and uniqueness for $s > \frac{3}{2}$. We use the result for $s = \frac{3}{2}$ which gives a priori bounds for (ω, θ) in $L^\infty(0, T; (H^{\frac{3}{2}}(\mathbb{T}^2))^2)$. Then the embedding $H^{\frac{3}{2}} \subset L^\infty$ implies that the BKM quantity (25) is controlled, and higher regularity is propagated. $\square$

3. Proof of Theorem 3

Denoting by θ_B and u_B the solutions of the 2D Boussinesq equations, by θ_V and u_V the solutions of the 2D Voigt Boussinesq equation, we set

$$u = u_V - u_B, \quad \theta = \theta_V - \theta_B,$$

and we have the equations

$$\begin{aligned} \partial_t(I + \epsilon\Lambda)\theta + (u + u_B) \cdot \nabla\theta + u \cdot \nabla\theta_B &= -\epsilon\Lambda\partial_t\theta_B, \\ \partial_t(I + \epsilon\Lambda)u + (u + u_B) \cdot \nabla u + u \cdot \nabla u_B - \theta e_2 &= -\epsilon\Lambda\partial_t u_B, \\ \nabla \cdot u &= 0, \end{aligned} \quad (26)$$

with vanishing initial data. We multiply the first equation by θ , the second equation by u , integrate and add. Because $\omega_B \in L^\infty(H^s)$ and $\theta_B \in L^\infty(H^{s+1})$, $s > 1$, we have $\nabla u_B \in L^\infty$ and $\nabla\theta_B \in L^\infty$ bounded uniformly by a constant C . Hence

$$\left| \int_{\mathbb{T}^2} ((u \cdot \nabla\theta_B)\theta + (u \cdot \nabla u_B)u) dx \right| \leq C\|u\|_{L^2}(\|\theta\|_{L^2} + \|u\|_{L^2}).$$

We consider the quantity

$$E = \|u\|_{L^2}^2 + \|\theta\|_{L^2}^2 + \epsilon \left(\|u\|_{H^{\frac{1}{2}}}^2 + \|\theta\|_{H^{\frac{1}{2}}}^2 \right)$$

and note that we have

$$\frac{d}{dt} E \leq C_1(\|u\|_{L^2}^2 + \|\theta\|_{L^2}^2) + \epsilon \left[\|\partial_t\theta_B\|_{H^{\frac{1}{2}}} \|\theta\|_{H^{\frac{1}{2}}} + \|\partial_t u_B\|_{H^{\frac{1}{2}}} \|u\|_{H^{\frac{1}{2}}} \right]$$

Using the Boussinesq equation, the assumptions $\omega_B \in L^\infty(H^s)$, $\theta_B \in L^\infty(H^{s+1})$, $s > 1$, ensure that $\|\partial_t \theta_B\|_{H^{\frac{1}{2}}}$ and $\|\partial_t u_B\|_{H^{\frac{1}{2}}}$ are bounded uniformly by a constant. From Young's inequality we have

$$\frac{d}{dt}E \leq C_2 E + \epsilon C_3$$

with initial data $E(0) = 0$, and thus we have the convergence $\lim_{\epsilon \rightarrow 0} E(t) = 0$ uniformly on finite time intervals. $\square$

4. Proof of Theorem 4

We consider the 2D fractional Voigt Boussinesq

$$\partial_t \omega + (I + \Lambda)^{-\alpha} (u \cdot \nabla \omega) = (I + \Lambda)^{-\alpha} \partial_1 \theta, \quad (27)$$

$$u = \nabla^\perp \Delta^{-1} \omega, \quad (28)$$

$$\partial_t \theta + (I + \Lambda)^{-\beta} (u \cdot \nabla \theta) = 0 \quad (29)$$

written in divergence form with initial conditions

$$\omega(\cdot, 0) = \omega_0 = \nabla^\perp \cdot u_0 \quad \text{and} \quad \theta(\cdot, 0) = \theta_0. \quad (30)$$

We have set the parameter $\epsilon = 1$ for simplicity.

Multiplying equation (29) by $(I + \Lambda)^\beta \theta$ and integrating,

$$\frac{1}{2} \frac{d}{dt} \|(I + \Lambda)^{\beta/2} \theta\|_{L^2}^2 = 0, \quad (31)$$

multiplying equation (27) by $(I + \Lambda)^\alpha \omega$ and integrating,

$$\frac{1}{2} \frac{d}{dt} \|(I + \Lambda)^{\alpha/2} \omega\|_{L^2}^2 = \int \partial_1 \theta \omega \leq C \|\theta\|_{H^{\beta/2}} \|\omega\|_{H^{\alpha/2}} \quad (32)$$

if $\alpha/2 + \beta/2 \geq 1$. By (31), we get $\theta \in L^\infty(H^{\beta/2})$ with $\|\theta\|_{H^{\beta/2}} \leq \|\theta_0\|_{H^{\beta/2}}$. By (32), we have $\omega \in L^\infty(H^{\alpha/2})$. Now we consider $s > 1$ and look at the evolution of the H^s norm of ω . We use the equivalent form $\|\omega\|_s \sim \|(I + \Lambda)^s \omega\|_{L^2}$. We have from (32)

$$\frac{1}{2} \frac{d}{dt} \|\omega\|_s^2 = A + B \quad (33)$$

with

$$A = - \int (I + \Lambda)^{s-\frac{\alpha}{2}} (u \cdot \nabla \omega) (I + \Lambda)^{s-\frac{\alpha}{2}} \omega dx \quad (34)$$

and

$$B = \int (I + \Lambda)^{s-\alpha} \partial_1 \theta (I + \Lambda)^s \omega dx \quad (35)$$

Now we use incompressibility, integration by parts and the commutator estimate

$$\|[(I + \Lambda)^\sigma, u \cdot \nabla] \omega\|_{L^2} \leq C \|\nabla u\|_{L^\infty} \|\omega\|_\sigma \quad (36)$$

valid for $\sigma > 0$ [17], together with the embedding inequality

$$\|\nabla u\|_{L^\infty} \leq C \|\omega\|_s \quad (37)$$

which is true because $s > 1$, to conclude that

$$|A| \leq C \|\omega\|_s \|\omega\|_{s-\frac{\alpha}{2}}^2. \quad (38)$$

We choose first $s = \alpha > 1$ and deduce in view of (31) and (32) that

$$|A| \leq C \|\omega\|_s \|\omega\|_{\frac{\alpha}{2}}^2 \leq (\|\omega_0\|_{\frac{\alpha}{2}} + C_\alpha t \|\theta_0\|_{\frac{\beta}{2}})^2 \|\omega\|_s = (a + bt)^2 \|\omega\|_s \quad (39)$$

holds for all $t \geq 0$. When $s = \alpha$ we see that

$$|B| \leq C \|\theta\|_1 \|\omega\|_s \quad (40)$$

and we have to estimate the H^1 norm of θ . This evolves according to

$$\frac{1}{2} \frac{d}{dt} \|\theta\|_1^2 = - \int (I + \Lambda)^{1-\frac{\beta}{2}} (u \cdot \nabla \theta) (I + \Lambda)^{1-\frac{\beta}{2}} \theta dx \quad (41)$$

and using the commutator estimate (36), we obtain

$$\frac{1}{2} \frac{d}{dt} \|\theta\|_1^2 \leq C \|\omega\|_s \|\theta\|_{1-\frac{\beta}{2}}^2. \quad (42)$$

Interpolating we may write

$$\|\theta\|_{1-\frac{\beta}{2}}^2 \leq \|\theta\|_1 \|\theta\|_{1-\beta} \leq C \|\theta\|_1 \|\theta_0\|_{\frac{\beta}{2}} \quad (43)$$

where we used $\beta \geq \frac{2}{3}$ and (31). We have thus from (42) and (43),

$$\frac{1}{2} \frac{d}{dt} \|\theta\|_1^2 \leq C_\beta \|\omega\|_s \|\theta\|_1. \quad (44)$$

The ODE inequality system in $X = \|\omega\|_s$ (with $s = \alpha$) and $Y = \|\theta\|_1$,

$$\begin{cases} \frac{1}{2} \frac{d}{dt} X^2 \leq (a + bt)^2 X + CXY \\ \frac{1}{2} \frac{d}{dt} Y^2 \leq C_\beta XY \end{cases} \quad (45)$$

follows from (33), (39), (40), (44). From this ODE we deduce a priori bounds for the quantities X and Y which are finite for all t . Once we know these bounds, we know that $\|\nabla u\|_{L^\infty}$ is controlled. The evolution of $\|\theta\|_s$ then obeys, using the commutator estimate (36),

$$\frac{d}{dt} \|\theta\|_s^2 \leq C \|\nabla u\|_{L^\infty} \|\theta\|_{s-\frac{\beta}{2}}^2, \quad (46)$$

and it implies a priori bounds on $\|\theta\|_s$. The bound (38) is replaced by

$$|A| \leq C \|\nabla u\|_{L^\infty} \|\omega\|_{s-\frac{\alpha}{2}}^2 \quad (47)$$

and (40), is replaced by

$$|B| \leq C \|\theta\|_s \|\omega\|_s \quad (48)$$

where we used $\alpha > 1$. Then from (33) it follows that $\|\omega\|_s$ is controlled. $\square$

5. Proof of Theorem 5

We start by noticing that because $\beta = 0$ we have that

$$\|\theta(t)\|_{L^p} \leq \|\theta_0\|_{L^p} \quad (49)$$

holds for all $t \geq 0$ and all $1 \leq p \leq \infty$. Then multiplying the u equation (29) by $(I + \Lambda)^\alpha u$ and integrating, we obtain

$$\frac{1}{2} \frac{d}{dt} \|u\|_{\frac{\alpha}{2}}^2 = \int \theta u_2 dx \leq \|\theta_0\|_{L^2} \|u\|_{L^2} \quad (50)$$

from whence we deduce that

$$\|u(t)\|_{\frac{\alpha}{2}} \leq C_D t. \quad (51)$$

Let us consider first $s = 1 + \epsilon$ where $0 < \epsilon = \alpha - 2$. Multiplying the ω equation (27) by $(I + \Lambda)^{2s} \omega$ and integrating, we have

$$\frac{1}{2} \frac{d}{dt} \|\omega\|_s^2 = - \int (I + \Lambda)^{s-\alpha} (\nabla(u\omega)) (I + \Lambda)^s \omega dx + \int (I + \Lambda)^{s-\alpha} \partial_1 \theta (I + \Lambda)^s \omega dx. \quad (52)$$

We note that $s - \alpha + 1 = 0$ by our choice of s , so we obtain that

$$\frac{1}{2} \frac{d}{dt} \|\omega\|_s^2 \leq C \|u\|_{L^4} \|\omega\|_{L^4} \|\omega\|_s + C \|\theta_0\|_{L^2} \|\omega\|_s \quad (53)$$

Thus we have that $\|\omega\|_s$ is bounded on $[0, T]$. Because $s > 1$, implies that $\|\nabla u\|_{L^\infty}$ is bounded on $[0, T]$ and the regularity for arbitrary $s > 1$ follows as in the proof of Theorem 4 above. $\square$

Conflict of interest The author declares that she has no conflicts of interest.

Acknowledgments. We acknowledge discussions with Jingyang Shu. This work was partially supported by NSF grant DMS-1713985.

References

- [1] Y. Cao, E. M. Lunasin, E. S. Titi. Global well-posedness of the three-dimensional viscous and inviscid simplified Bardina turbulence models. *Commun. Math. Sci.*, 4(2006), 823–848.
- [2] D. Chae, Global regularity for the 2D Boussinesq equations with partial viscosity terms, *Adv. Math.* 203(2) (2006), 497–513.
- [3] D. Chae, S.-K. Kim, and H.-S. Nam. Local existence and blow-up criterion of Hölder continuous solutions of the Boussinesq equations. *Nagoya Math. J.*, 155:55–80, 1999.
- [4] S. Chandrasekhar, *Hydrodynamic and Hydromagnetic Stability*, Oxford University Press (1961).
- [5] J. Chen, T. Y. Hou. Finite time blowup of 2D Boussinesq and 3D Euler equations with $C^{1,\alpha}$ velocity and boundary. *Comm. Math. Phys.*, 383:1559–1667, 2021.
- [6] J. Chen, T. Y. Hou. Stable nearly self-similar blowup of the 2D Boussinesq and 3D Euler equations with smooth data. 2022, arXiv:2210.07191.
- [7] P. Constantin, F. Pasqualotto. Magnetic Relaxation of a Voigt–MHD System. *Communications in Mathematical Physics*. 402(2):1931–1952, 2023.
- [8] R. Danchin, M. Paicu. Global existence results for the anisotropic Boussinesq system in dimension two. *Mathematical Models and Methods in Applied Sciences* 21(3), (2011) 421–457.
- [9] T. Elgindi. Finite-time singularity formation for solutions to the incompressible Euler equations on $\mathbb{R}^3$. *Annals of Mathematics* 194(3),(2021), 647–727.
- [10] T. Elgindi, I.-J. Jeong. Finite-Time Singularity Formation for Strong Solutions to the Axisymmetric 3D Euler Equations. *Annals of PDE* 5(2), 16 (2019).
- [11] T. Elgindi, I.-J. Jeong. Finite-time singularity formation for strong solutions to the Boussinesq system. *Annals of PDE* 6(1), 5 (2020).
- [12] F. Hadadifard and A. Stefanov. On the global regularity of the 2D critical Boussinesq system with $\alpha > 2/3$. *Comm. Math. Sci.* 15(5) (2017), 1325–1351.
- [13] T. Hmidi, S. Keraani, and F. Rousset. Global well-posedness for a Boussinesq-Navier-Stokes system with critical dissipation. *J. Differential Equations* 249 (2010), 2147–2174.
- [14] T. Hmidi, S. Keraani, and F. Rousset. Global well-posedness for Euler-Boussinesq system with critical dissipation. *Comm. Partial Differential Equations* 36 (2011), 420–445.
- [15] T.Y. Hou and C. Li. Global well-posedness of the viscous Boussinesq equations. *Discrete Contin. Dyn. Syst.* 12(1) (2005), 1–12.
- [16] W. Hu, I. Kukavica, and M. Ziane. On the regularity for the Boussinesq equations in a bounded domain. *J. Math. Phys.* 54(8) (2013), 081507, 10.
- [17] T. Kato and G. Ponce. Commutator estimates and the Euler and Navier-Stokes equations. *Comm. Pure Appl. Math.* 41(7) (1988), 891–907.
- [18] I. Kukavica, F. Wang and M. Ziane. Persistence of regularity for solutions of the Boussinesq equations in Sobolev spaces. *Adv. Differential Equations*. 21(1/2) (2016), 85–108.
- [19] A. Larios, E. Titi. On the higher-order global regularity of the inviscid Voigt-regularization of three-dimensional hydrodynamic models. *Discrete and Continuous Dynamical Systems B*, 14 (2010) 603–627.
- [20] A. Larios, E. Titi. Higher-order global regularity of an inviscid Voigt-regularization of the three-dimensional inviscid resistive magnetohydrodynamic equations. *J. Math. Fluid Mech.*, 16(1) (2014), 59–76.
- [21] A. Larios, E. Lunasin, and E.S. Titi. Global well-posedness for the 2D Boussinesq system with anisotropic viscosity and without heat diffusion. *J. Differential Equations* 255(9) (2013), 2636–2654.
- [22] B. Levant, F. Ramos, E. Titi. On the statistical properties of the 3D incompressible Navier-Stokes-Voigt model. *Commun. Math. Sci.*, 8(1) (2010) 277–293.
- [23] J. Linshiz E.Titi. Analytical study of certain magnetohydrodynamic models. *J. Math. Phys.*, 48(6) (2007) 065504, 28.
- [24] A. J. Majda, A. L. Bertozzi, *Vorticity and Incompressible Flow*, Cambridge University Press 2002.
- [25] A. P. Oskolkov. The uniqueness and solvability in the large of boundary value problems for the equations of motion of aqueous solutions of polymers. *Zap. Nauch. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 38 (1973) 98–136.
- [26] A. Pumir, E.D. Siggia. Finite-Time Singularities in the Axisymmetric Three-Dimensions Euler Equations. *Phys. Rev. Lett.* 68(10) (March 1992), 1511–1514
- [27] F. Ramos, E. Titi. Invariant measures for the 3D Navier-Stokes-Voigt equations and their Navier-Stokes limit. *Discrete Contin. Dyn. Syst.* 28(1) (2010) 375–403.
- [28] A. Stefanov and J. Wu. A global regularity result for the 2D Boussinesq equation with critical dissipation. *Journal d’Analyse Mathématique*, 137 (2019).

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA, PA 19122
Email address: `ignatova@temple.edu`