GALAXY CLUSTERS DISCOVERED VIA THE THERMAL SUNYAEV-ZEL'DOVICH EFFECT IN THE 500-SQUARE-DEGREE SPTPOL SURVEY

L. E. Bleem 1,2, M. Klein 3,4, T. M. C. Abbott 5, P. A. R. Ade 6, M. Aguena 7, O. Alves 8, A. J. Anderson 9, F. Andrade-Oliveira 8, B. Ansarinejad 10, M. Archipley 11,12, M. L. N. Ashby 13, J. E. Austermann 14,15, D. Bacon 16, J. A. Beall 14, A. N. Bender 1,2,17, B. A. Benson 9,2,17, F. Bianchini 18,19,20, S. Bocquet 3, D. Brooks 21, D. L. Burke M. Calzadilla 22, J. E. Carlstrom 2,23,1,17,24, A. Carnero Rosell 25,7,26, J. Carretero 27, C. L. Chang 2,11,17, P. Chaubal 10, H. C. Chiang 28,29, T-L. Chou 2,23, R. Citron 30, C. Corbett Moran 31, M. Costanzi 32,33,34, T. M. Crawford 2,17, A. T. Crites 35, L. N. da Costa 7, T. de Haan 36,37, J. De Vicente 38, S. Desai 39, M. A. Dobbs 24,41, J. Garc'ia-Bellido 45, M. Gatti 46, E. M. George 47, G. Giannini 27, S. Grandis 48, D. Gruen 3, R. A. Gruendl 12,11, N. Gupta 10, G. Gutierrez 9, N. W. Halverson 41,15, S. R. Hinton 49, G. P. Holder 11,50,40, D. L. Hollowood 51, W. L. Holzapfel 52, K. Honscheid 53,54, J. D. Hrubes 30, N. Huang 52, J. Hubmayr 14, K. D. Irwin 20,19 J. Mena-Fern'andez 55, D. J. James 56, F. K. éruzor é1, L. Knox 57, K. Kuehn 58,59, O. Lahav 21, A. T. Lee 52,60, S. Lee 31, D. Li 14,20, A. Lowitz 17, J. L. Marshal 61, M. McDonald 22, J. J. McMahon 2,23,17, F. Menanteau 12,11, S. S. Meyer 2,23,17,24, R. Miquel 67, B. J. J. Mohr 3,4,63, J. Montgomery 28, J. Myles 64, T. Natoli 17,2, J. P. Nibarger 14, G. I. Noble 65,66, V. Novosad 67, R. L. C. Ogando 68, S. Padin 69, S. Patil 10, M. E. S. Pereira 70, A. Pieres 7,68, A. A. Plazas Malag'on 18,20, C. Pryke 71, C. L. Reichardt 10, M. Rodr'iguez-Monroy 72, A. K. Romer 73, J. E. Ruhl 74, R. Sevilla-Noarbe 38, C. Sievers 30, G. Smecher 28,79, M. Smith 60, T. Somboonpanyakul 18,81, B. Stalder 82, A. A. Stark 56, E. Suchyta 83, M. E. C. Swanson 12, G. Tarle 85, P. Wiseman 80, W. L. K. Wu 20, V. Yefremenko 1, A. Zebrowski 2,17,9, and Y. Zhang 5 Version February 9, 2024

ABSTRACT

We present a catalog of 689 galaxy cluster candidates detected at significance $\xi > 4$ via their thermal Sunyaev-Zel'dovich (SZ) effect signature in 95 and 150 GHz data from the 500-square-degree SPTpol survey. We use optical and infrared data from the Dark Energy Camera and the Wide-field Infrared Survey Explorer (WISE) and Spitzer satellites, to confirm 544 of these candidates as clusters with ~ 94% purity. The sample has an approximately redshift-independent mass threshold at redshift z > 0.25. The confirmed sample spans $1.5 \times 10^4 < M_{500c} < 9 \times 10^{14} M_{\odot}/h_{70}$ and $0.03 < z \le 1.6$ in mass and redshift, respectively, with a median mass of $2.5 \times 10^{10} M_{\odot}/h_{70}$ and median redshift z = 0.7; 21% of the confirmed clusters are at z > 1. We use external radio data from the Sydney University Molonglo Sky Survey (SUMSS) to estimate contamination to the SZ signal from synchrotron sources. The contamination reduces the recovered ξ by a median value of 0.032, or \sim 0.8% of the ξ = 4 threshold value, and ~7% of candidates have a predicted contamination greater than $\Delta \xi = 1$. With the exception of a small number of systems (< 1%), an analysis of clusters detected in single-frequency 95 and 150 GHz data shows no significant contamination of the SZ signal by emission from dusty or synchrotron sources. This cluster sample, representing the deepest SZ-selected cluster sample todate, will be a key component in upcoming astrophysical and cosmological analyses of clusters. In addition to the cluster catalog, we also release the millimeter-wave maps and associated data products used to produce this sample. These maps have depths of 5.3 (11.7) µK_{CMB} -arcmin at 150 (95) GHz and an effective angular resolution of 1.2 (1.7). The SPTpol products are available at https:// pole.uchicago.edu/public/data/sptpol_500d_clusters/index.html, and the NASA LAMBDA website. An interactive sky server with the SPTpol maps and Dark Energy Survey data release 2 images is also available at NCSA https://skyviewer.ncsa.illinois.edu.

Keywords: Large-Scale Structure of the Universe, Galaxy Clusters

E-mail:lbleem@anl.gov High-Energy Physics Division, Argonne National Laboratory, 9700 South Cass Avenue., Lemont, IL, 60439, USA

² Kavli Institute for Cosmological Physics, University of Chicago, 5640 South Ellis Avenue, Chicago, IL, 60637, USA ³ University Observatory, Faculty Maximilians-Universit at, Scheinerstr. of Physics, Ludwig-1, 81679 Munich, Germany
⁴ Max-Planck-Institut für extraterrestrische Physik, Giessenbachstr. 85748 Garching, Germany Cerro Tololo Inter-American Observatory, NSF's National Optical-Infrared Astronomy Research Laboratory, Casilla 603,

La Serena, Chile ⁶ School of Physics and Astronomy, Cardiff University, Cardiff

CF24 3YB, United Kingdom

⁷ Laborat orio Interinstitucional de e-Astronomia - LIneA Jos'e Cristino 77, Rio de Janeiro, RJ - 20921-400,

Brazil

8 Department of Physics, University of Michigan, 450 Church

Street, Ann Arbor, MI, 48109, USA

⁹ Fermi National Accelerator Laboratory, MS209, P.O. Box 500, Batavia, IL, 60510, USA

10 School of Physics, University of Melbourne, Parkville, VIC

3010, Australia

¹¹ Department of Astronomy, University of Illinois Urbana-Champaign, 1002 West Green Street, Urbana, IL, 61801, USA ¹² Center for AstroPhysical Surveys, National Center for

Supercomputing Applications, Urbana, IL, 61801, USA

13 Center for Astrophysics | Harvard & Smithsonian,

and Infrared Astronomy Division, Cambridge, MA 01238, USA NIST Quantum Devices Group, 325 Broadway Mailcode

817.03, Boulder, CO, 80305, USA

15 Department of Physics, University of Colorado, Boulder,

CO, 80309, USA

16 Institute of Cosmology and Gravitation,
Portsmouth, Portsmouth, PO1 3FX, UK University of

17 Department of Astronomy and Astrophysics, University of Chicago, 5640 South Ellis Avenue, Chicago, IL, 60637, USA

18 Kavli Institute for Particle Astrophysics and Cosmology

Stanford University, 452 Lomita Mall, Stanford, CA,

USA

19 Department of Physics, Stanford University, 382 Via
Pueblo Mall, Stanford, CA, 94305, USA

SLAC National Accelerator Laboratory, 2575 Sand Hill
 Road, Menlo Park, CA, 94025, USA
 Department of Physics & Astronomy, University College
 London, Gower Street, London, WC1E 6BT, UK

University College

²² Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

²³ Department of Physics,

University of Chicago, 5640 South

Ellis Avenue, Chicago, IL, 60637, USA

²⁴ Enrico Fermi Institute, University of Chicago, 5640 South Ellis Avenue, Chicago, IL, 60637, USA

²⁵ Instituto de Astrofisica de Canarias. E-38205 La Laguna.

Tenerife, Spain

²⁶ Universidad de La Laguna, Dpto. Astrof'ısica, E-38206 La

Laguna, Tenerife, Spain

Institut de F´ısica d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, 08193 Bellaterra (Barcelona) Spain

²⁸ Department of Physics and McGill Space Institute, McGill University, 3600 Rue University, Montreal, Quebec H3A 2T8,

Canada
²⁹ School of Mathematics, Statistics & Computer Science University of KwaZulu-Natal, Durban, South Africa

³⁰ University of Chicago, 5640 South Ellis Avenue, Chicago. IL, 60637, USA

³¹ Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91011, USA
 ³² Astronomy Unit, Department of Physics, University

Department of Physics, University of Trieste, via Tiepolo 11, I-34131 Trieste, Italy

³³ INAF-Osservatorio Astronomico di Trieste, via G. B. Tiepolo 11, I-34143 Trieste, Italy

Institute for Fundamental Physics of the Universe, Via Beirut 2, 34014 Trieste, Italy

35 Cornell University, Ithaca, NY 14853, USA

³⁶ Institute of Particle and Nuclear Studies (IPNS), High Energy Accelerator Research Organization (KEK), Tsukuba.

Ibaraki 305-0801, Japan

37 International Center for Quantum-field Measurement Systems for Studies of the Universe and Particles (QUP-WPI), High Energy Accelerator Research Organization (KEK), Ibaraki 305-0801, Japann

38 Centro de Investigaciones Energ´eticas, Medioambientales y

Tecnol'ogicas (CIEMAT), Madrid, Spain

39 Department of Physics, IIT Hyderabad, Kandi, Telangana 502285, India

⁴⁰ Canadian Institute for Advanced Research, CIFAR Program in Gravity and the Extreme Universe, Toronto, ON, M5G 1Z8, Canada ´
⁴¹ Department of Astrophysical and Planetary Sciences,

University of Colorado, Boulder, CO, 80309, USA

42 Institute of Theoretical Astrophysics, University of Oslo.
 P.O. Box 1029 Blindern, NO-0315 Oslo, Norway
 43 Faculty of Physics and Astronomy, University of Missouri–

Kansas City, 5110 Rockhill Road, Kansas City, MO 64110, USA

44 Harvey Mudd College, 301 Platt Boulevard, Claremont, CA, 91711, USA

de Fisica Teorica UAM/CSIC, Universidad

Autonoma de Madrid, 28049 Madrid, Spain

46 Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, USA

European Southern Observatory, Karl-Schwarzschild-

Str..DE-85748 Garching b. Munchen, Germany

48 Universit at Innsbruck, Institut für Astro- und Teilchenphysik, Technikerstr. 25/8, 6020 Innsbruck, Austria

⁴⁹ School of Mathematics and Physics, University of Queens-

land, Brisbane, QLD 4072, Australia

50 Department of Physics, University of Illinois Urbana-Champaign, 1110 West Green Street, Urbana, IL, 61801, USA 51 Santa Cruz Institute for Particle Physics, Santa Cruz, CA

95064, USA
52 Department of Physics, University of California,

CA, 94720, USA

53 Center for Cosmology and Astro-Particle Physics. The Ohio State University, Columbus, OH 43210, USA

54 Department of Physics, The Ohio State University, Colum-

bus, OH 43210, USA

55 LPSC Grenoble - 53, Avenue des Martyrs 38026 Grenoble,

France
56 Harvard-Smithsonian Center for Astrophysics, 60 Garden

Street, Cambridge, MA, 02138, USA

57 Department of Physics, University of California, One Shields Avenue, Davis, CA, 95616, USA

Australian Astronomical Optics, Macquarie University, North Ryde, NSW 2113, Australia

⁵⁹ Lowell Observatory, 1400 Mars Hill Rd, Flagstaff, AZ 86001, USA

60 Physics Division, Lawrence Berkeley National Laboratory,
Berkeley, CA, 94720, USA
61 George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, and Department of Physics and Astronomy, Texas A&M University, College Station. 77843, USA

62 Instituci´o Catalana de Recerca i Estudis Avan cats, E-08010

Barcelona, Spain

63 Excellence Cluster Universe, Boltzmannstr. 2, 85748 Garching, Germany

Department of Astrophysical Sciences, Princeton Univer-

sity, Peyton Hall, Princeton, NJ 08544, USA

65 Dunlap Institute for Astronomy & Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON, M5S 3H4, Canada ⁶⁶ David A.

Dunlap Department of Astronomy & Astrophysics, University of Toronto, 50 St. George Street, Toronto,

ON, M5S 3H4, Canada

67 Materials Sciences Division, Argonne National Laboratory,

9700 South Cass Avenue, Lemont, IL, 60439, USA
⁶⁸ Observat'orio Nacional, Rua Gal. Jos'e Cristino 77, Rio de

Janeiro, RJ - 20921-400, Brazil
⁶⁹ California Institute of Technology, 1200 East California

Boulevard., Pasadena, CA, 91125, USA

⁷⁰ Hamburger Sternwarte, Universit at Hamburg, Gojen-bergsweg 112, 21029 Hamburg, Germany

71 School of Physics and Astronomy, University of Minnesota, 116 Church Street SE Minneapolis, MN, 55455, USA

⁷² Laboratoire de physique des 2 infinis Ir`ene Joliot-Curie, CNRS Universit'e Paris-Saclay, Bat. 100, Facult'e des sciences, F-91405 Orsay Cedex, France

1. INTRODUCTION

The advent of high-resolution cosmic microwave background (CMB) surveys has revolutionized the field of galaxy cluster science by enabling the identification of approximately mass-limited samples of massive galaxy clusters (Vanderlinde et al. 2010; Bleem et al. 2015; Planck Collaboration et al. 2016b; Hilton et al. 2018; Bleem et al. 2020) via the thermal Sunyaev-Zel'dovich (SZ) effect (Sunyaev & Zel'dovich 1972). These SZ samples have been used to place stringent constraints on cosmological models (Planck Collaboration et al. 2016a; Bocquet et al. 2019; Zubeldia & Challinor 2019; Salvati et al. 2022), the evolution of the intracluster medium (McDonald et al. 2017; Ghirardini et al. 2021; Flores et al. 2021; CHEX-MATE Collaboration et al. 2021; Anbajagane et al. 2022; Ruppin et al. 2023; Olivares et al. 2022) and the evolution of galaxies residing in clusters (Hennig et al. 2017; Chiu et al. 2018; Strazzullo et al. 2019; Khullar et al. 2022; Kim et al. 2023; Somboonpanyakul et al. 2022). The next generation of CMB surveys, with significantly increased sensitivity, is continuing this revolution by probing lower-mass and correspondingly higher-redshift systems (Huang et al. 2020; Hilton et al. 2021).

With this increased sensitivity comes both new opportunities (Simons Observatory Collaboration 2019; Chaubal et al. 2022) and challenges (Melin et al. 2018; Zubeldia et al. 2023) for cosmological and astrophysical analyses with SZ clusters. The enlarged sample sizes will also strengthen cross-wavelength analyses, for which comparisons of cluster samples selected via different techniques provide powerful ways to validate the robustness of analysis inferences and can signal the presence of unmitigated systematic errors (Grandis et al. 2021; Costanzi et al. 2021; Orlowski-Scherer et al. 2021).

In this work we continue the push to discover SZ clusters in low-noise and high-resolution CMB survey data by presenting a galaxy cluster sample constructed using the 500-square-degree SPTpol survey. his survey, which covers five times the area used in the 100-square-degree SPTpol field (hereafter SPTpol 100d) cluster analysis (Huang et al. 2020), benefits from approximately two-to-three times lower noise than the survey data used

⁷³ Department of Physics and Astronomy, Pevensey Building,
 University of Sussex, Brighton, BN1 9QH, UK
 ⁷⁴ Department of Physics, Case Western Reserve University,

Cleveland, OH, 44106, USA

To Brookhaven National Laboratory, Upton, NY 11973, USA

⁷⁵ Brookhaven National Laboratory, Upton, NY 11973, USA ⁷⁶ Universit'e Paris-Saclay, CNRS, Institut d'Astrophysique Spatiale, 91405, Orsay, France

⁷⁷ Liberal Arts Department, School of the Art Institute of Chicago, 112 South Michigan Avenue, Chicago, IL,60603, USA ⁷⁸ Argelander-Institut für Astronomie, Auf dem H¨ugel 71, D-53121 Bonn, Germany

⁷⁹ Three-Speed Logic, Inc., Victoria, B.C., V8S 3Z5, Canada ⁸⁰ School of Physics and Astronomy, University of Southampton, SQ17 1B.L. LIK

ton, Southampton, SO17 1BJ, UK

81 Department of Physics, Faculty of Science, Chulalongkorn University, 254 Phayathai Road, Pathumwan, Bangkok Thailand 10330

land. 10330

82 Vera C. Rubin Observatory Project Office, 933 North
Cherry Avenue Tucson AZ 85721 USA

Cherry Avenue, Tucson, AZ 85721, USA

83 Computer Science and Mathematics Division, Oak Ridge
National Laboratory, Oak Ridge, TN 37831

84 Space Science and Engineering Division, Southwest Research Institute, San Antonio, TX 78238

⁸⁵ Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA to construct the published cluster samples from SPT-SZ (Bleem et al. 2015) and ACTpol (Hilton et al. 2021). The new cluster sample is composed of 689 cluster candidates detected at significance $\xi > 4$, and we have confirmed 544 of these candidates as galaxy clusters using optical and infrared imaging data. Here we provide the complete cluster candidate list and—where available—redshifts, estimated masses, and select optical properties.

We organize our description of the identification and characterization of this new sample as followsIn Section 2 we describe the collection of the millimeter-wave data and its processing into the maps used for cluster identification. We detail the procedure by which we identify SZ cluster candidates in these maps in Section 3. Section 4 we describe the optical and infrared data sets we use to confirm candidates as galaxy clusters by the techniques described in Section 5. In Section 6 we describe how we estimate cluster masses and estimate the purity of our cluster sample. In Section 7 we describe a number of systematic checks to explore possible astrophysical contamination of the SZ signal. In Section 8 we present the properties of the resulting SZ cluster catalog and compare select properties of this catalog to other SZ- and optically selected cluster samples. Finally, we summarize our results and conclusions in Section 9.

In addition to the cluster sample we also release the SPTpol maps and associated data products used in the construction of this catalog. These products can be found on both the NASA Legacy Archive for Microwave Background Data (LAMBDA) ¹ and the South Pole Telescope (SPT) collaboration's website². Sky maps for the 500d region from SPTpol single-frequency data as well as images from the Dark Energy Survey (DES)-DR2 (Abbott et al. 2021) dataset can be accessed for exploration at the National Center for Supercomputing Applications (NCSA) Skyviewer.³

Finally we note that, where applicable, we assume a fiducial Λ CDM cosmology with $\sigma_8 = 0.80$, $\Omega_b = 0.046$, $\Omega_m = 0.30$, h = 0.70, $n_s(k_s = 0.002) = 0.972$, and $\Sigma m_v = 0.06$ eV. Magnitudes are reported in the AB system (Oke 1974).Cluster masses are reported in terms of M_{500c} , which is defined as the mass enclosed within a radius, r_{500c} , at which the average enclosed density is $500\times$ the critical density at the cluster redshift.

2. OBSERVATIONS AND DATA REDUCTION

The cluster catalog presented in this work was constructed via an analysis of 95 and 150 GHz data from the primary SPTpol (Austermann et al. 2012) survey field. This 500-square-degree field is centered at right ascension and declination (α =0 h , δ = –57.5 $^\circ$) and is hereafter referred to as the SPTpol 500d field; see Figure 1. The maps have depths of 11.7 and 5.3 μ K-arcmin at 95 and 150 GHz, respectively, and effective resolutions of 1.7 and 1.2 set by the instrumental response (i.e., telescope beam). The SPTpol data in this field have been the focus of a number of previous analyses including characterizations of low- and high- ℓ CMB temperature and polarization power spectra (Henning et al. 2018; Sayre

https://lambda.gsfc.nasa.gov/product/spt/sptpol_prod_table.html

² https://pole.uchicago.edu/public/Data%20Releases.html

https://skyviewer.ncsa.illinois.edu




Figure 1. Sequentially zoomed-in images of the SPTpol 500d field, with the full field shown in inset at top. (*Top panel*) 24 deg 2 of 95 GHz data (*Middle panel*) The same region at 150 GHz. The full field temperature maps and associated data products are released along with this work. (*Bottom panel*) To better illustrate the sensitivity of these maps to small scales, we further zoom in to a 1.25 × 1.25 degree region (outlined in black in above panels) centered on SPT-CL J2341-5724, a ξ = 14.3 cluster at z = 1.259. Plotted from left-to-right are the 95 and 150 GHz temperature maps and the ξ -map constructed for optimal detection of clusters with θ core = 0.25. Sources brighter than 6 mJy at 150 GHz were masked in the cluster detection step, with two examples visible in the bottom right panel.

et al. 2020; Reichardt et al. 2021), the CMB gravitational lensing signal (Wu et al. 2019; BICEP/Keck Collaboration and SPTpol Collaboration et al. 2021; Raghunathan et al. 2019), and the properties of emissive sources (Gupta et al. 2019). The overall map-making and data processing procedures utilized here closely follow those of previous efforts. In this section we provide a brief summary of these procedures, highlighting changes specific to this work, and refer readers to prior publications for more details.

2.1. Data Processing

The maps presented here are the weighted sum of > 4000 individual observations (~10,200 hours) of the SPTpol 500d field acquired between 2013 April 30 and 2016 Aug 20. The majority of the observations were obtained by scanning the telescope in azimuth back and forth across the field, stepping in elevation, and repeating this process until the full field was covered. As detailed in Henning et al. (2018), most of the data obtained before 2014 May 29 was acquired instead in "lead-trail" mode. In this mode, the field is split into two equal halves via an equal division in right ascension. These two subfields were then sequentially scanned, with starting times offset owing to sky rotation, such that the same azimuthal range was covered by the telescope in each subfield For this analysis we combine these "lead" and "trail" observations into full-field observations to match the rest of the 500d dataset.

The SPTpol data processing pipeline converts timeordered electrical signals recorded by the camera into calibrated maps of the millimeter-wave (mm-wave) sky. We apply standard detector quality cuts (Crites et al. 2015) and electrical cross talk corrections to the timeordered data (TOD). Following Bleem et al. (2020), to minimize the impact of low-frequency noise from the atmosphere and detector readout, both a common mode filter that removes the mean of all detectors in each frequency band and a seventh-order Legendre polynomial were fit to and subtracted from the TOD for each azimuthal scan. A scan-direction high-pass filter at angular multipole ℓ = 300 and similar low-pass filter at ℓ = 20, 000 were then applied. Bright emissive sources detected at 150 GHz at > 6 mJy were masked with masks of 4 radii during these filtering steps so as to not bias the fits or imprint artifacts in the resulting maps.

Using the telescope pointing model and weights for the individual bolometers based on their noise in the 1-3 Hz band, the TOD was binned into 0. 25 pixels in the Sanson-Flamsteed projection (Calabretta & Greisen 2002). The noise properties of each single-observation map were characterized and, following removal of a small number of maps with anomalous noise behavior, the observations were then combined via inverse noise-variance weighting to produce the final coadded maps of the field.

2.2. Removal of Emissive Sources that Cause Spurious Cluster Candidates

As discussed in Huang et al. (2020), unmasked emissive sources in the maps can lead to spurious cluster candidates owing to decrements produced by the sources "ringing" when the maps are high-pass filtered. To mitigate the number of such spurious detections, we identify and

remove moderate signal-to-noise (S/N) sources below our masking threshold from the maps before cluster detection occurs.

We first filter each individual frequency map with a filter optimized to detect point sources and construct a catalog of emissive sources detected at S/N > 6 and below the masking threshold of 6 mJy at 150 GHz used in the map construction. This corresponds to thresholds of \sim 4 mJy (3.5 mJy) at 95 GHz (150 GHz). A total of 348 sources were detected at 95 GHz and 382 at 150 GHz (245 at S/N> 6 in both maps). When sources were only detected in one of the two frequency maps the other frequency map was forced-photometered to recover the missing flux.

An empirical template of the 2D source profile, incorporating the effects of the beam and transfer function, was constructed by creating a flux-normalized median stack of the raw unfiltered maps at the locations of the \sim 200 brightest sources at each frequency (see Figure 2). A flux-scaled copy of this template was then subtracted from the raw maps at the location of each of the detected sources. Analysis of the cluster catalog (whose construction is detailed in the next section) showed this simple procedure was sufficient to remove the spurious candidates associated with these emissive sources. Artifacts from remaining lower flux sources contribute $<2\sigma$ SZ-like signals in the cluster detection maps. We use these source subtracted maps in all of our cluster identification steps.

3. IDENTIFICATION OF CLUSTER CANDIDATES

We closely follow the procedure utilized in previous SPT publications (most recently Huang et al. 2020; Bleem et al. 2020) to identify cluster candidates in the SPTpol 500d field with some small changes we explain below. As in these works, we filter the maps with a spatial-spectral filter that has been optimized to isolate cluster signals in the presence of frequency- and scale-dependent sources of noise (Melin et al. 2006). The spatial profile of clusters are modeled as a series of projected spherical β profiles (Cavaliere & Fusco-Femiano 1976)

$$\Delta T = \Delta T_{0} (1 + \theta^{2}/\theta_{c}^{2})^{-(3\beta-1)/2}$$
 (1)

with $\beta = 1$, a free normalization, ΔT_0 , and with core radii, θ_c , that are allowed to vary in twelve equally spaced steps from 0.25 to 3. The frequency dependence of the temperature signal in each map, ΔT , is given by the thermal SZ effect (Sunyaev & Zel'dovich 1972). At the SPTpol effective band centers of 95.9 and 148.5 GHz for a non-relativistic thermal SZ spectrum, this results in the thermal SZ signal being 1.6× brighter (in CMB temperature units) in the 95 GHz than 150 GHz data. This enhancement of the signal amplitude largely offsets the higher instrumental noise in the 95 GHz data, and—combined with the impacts of astrophysical noise discussed below—results in the two channels contributing close to equal weight in the construction of the cluster catalog. The noise in the SPTpol maps is broadly composed of two components(1) astrophysical/cosmological "noise" arising from fluctuations in the primary CMB and emission from extragalactic sources and (2) noise specific to our observations arising from the SPTpol instrument as well as residual atmospheric contamination.




Figure 2. 40 $^{'}$ × 40 $^{'}$ normalized median stack of sources at 95 GHz used to make the source subtraction template; a similar template was constructed at 150 GHz. The negative filtering artifacts along the scan (horizontal) direction are clearly visible even in these stacks which have significant residual large scale noise contributions from the primary CMB.

The SPTpol data are deep enough that—for identical reasons as in the case of moderate signal-to-noise emissive sources discussed in Section 2.2—the Fourier filtering induces spurious candidates around massive clusters along the telescope scan direction. As the sky density of such clusters is significantly lower than that of the emissive sources—0.09 per degat the conservative threshold we apply below—and they have a less well defined shape, we do not attempt to subtract a model of these clusters from the map. Instead we use a two-step procedure to construct a clean cluster candidate list that also recovers the properties of the most massive systems.

In the first step, we perform a cluster detection run on the maps masking both emissive sources with flux > 6 mJy at 150 GHz and massive clusters previously detected in the SPT-SZ survey 1 at significance 2 $\xi \ge 6$. This corresponds to a masking threshold of $M_{00c} \ge 4.5 \times 10^{14} M_{\odot}/h_{70}$. We use 4 radii masks and further exclude cluster candidates detected within 8 of these objects. Cluster candidates identified with $\xi > 4$ in this step from the SPTpol data form the core of the new cluster sample.

In the second step, to include the massive clusters masked in the previous step in our catalog, we perform a second filtering of the maps in which only the emissive sources > 6 mJy at 150 GHz are masked. We then add only the new detections of previously masked clusters and their properties to our cluster candidate list. There were no clusters masked in the first step that did not exceed our $\xi \ge 4$ cut. We validated through comparison of the noise properties in the two filtering steps that the presence of the most massive systems does not impact the estimated noise. We additionally visually inspected cutouts of the raw and filtered maps around all

 ξ > 4 cluster candidates to validate our treatment of both emissive sources and the highest signficance clusters. In total, after all masking is accounted for, 460 of the 498 deg uniform depth region (92%) of the SPTpol 500d field was included in the cluster search.

4. FOLLOW-UP OBSERVATIONS

SZ galaxy cluster candidates are confirmed and redshifts obtained via the identification of significant galaxy overdensities in optical and/or infrared imaging data. These data are drawn from both wide-field imaging surveys and targeted photometric and spectroscopic follow-up observations. The majority of our candidates are confirmed using data from the DES (Flaugher et al. 2015) and the highest-redshift clusters are confirmed using observations from the Wide-field Infrared Survey Explorer (WISE; Wright et al. 2010) and Spitzer/IRAC (Fazio et al. 2004). A small number of candidates at the borders of the SPTpol survey do not overlap with DES; these candidates are characterized using data from the DECam Legacy Survey (DECaLS; Dey et al. 2019).

4.1. Optical/Near-Infrared Imaging from the Dark Energy Camera

The DES is a ~5000 deg 2 5 band grizY optical and near-infrared imaging survey that was conducted using the 4 m Blanco telescope at Cerro Tololo Inter-American Observatory in Chile. Data were acquired over 760 nights between August 2013 and January 2019 (Abbott et al. 2021). In this analysis, we make use of photometric catalogs extracted from the full 6 year coadds. These data reach median 10σ coadded depths in 95 apertures of [24.7, 24.4, 23.8, 23.1, 21.7] magnitude in the grizY bands. The DES data cover almost the entire SPTpol 500d field and are typically deep enough to robustly confirm SPTpol cluster candidates to redshift $z \sim 1.1$.

Eighteen candidates at the edges of the SPTpol 500d field fall outside the DES coverage region. We use *griz* photometry data from the tenth data release of DECaL\$ to confirm and obtain redshifts for 11 of these candidates; the remaining 7 candidates are unconfirmed after optical/infrared analysis. Tests of cluster redshift estimates using systems in regions where DES and DECaLS overlap show good agreement.

4.2. Wide Field Infrared Explorer

To detect candidates at higher redshifts, we analyze data from the all-sky WISE dataset. Specifically we use 3.4 μ m and 4.6 μ m data (W1 and W2 band, respectively) from the "unWISE" analysis of Schlafly et al. (2019) which combined five years of data from WISE and NEOWISE (Mainzer et al. 2014) data at native WISE resolution.

4.3. Spitzer/IRAC

A subset of high-redshift cluster candidates were also targeted with Spitzer/IRAC at 3.6 and 4.5 μ m ([3.6] and [4.5] bands). These deeper data are significantly higher resolution than WISE ([3.6] $_{\rm FWHM}$ ~ 2 compared to 6 from WISE W1), allowing us to confirm additional high-redshift clusters as well as to validate our WISE analysis. Spitzer/IRAC observations of cluster candidates are

¹ The SPT-SZ survey covers all but 1 deg ² of the SPTpol 500d

 $^{^2}$ In this work, as in all previous SPT cluster publications, we define significance, ξ , as the maximum detection significance across the 12 matched filter scales.

https://www.legacysurvey.org/dr10/description/

drawn from three sources:50 candidates were previously targeted for follow-up observations in the production of the SPT-SZ cluster sample (PI: Brodwin, see details in Bleem et al. 2015), 84 cluster candidates were imaged in new targeted observations in *Spitzer* cycles 11, 12, and 14 (Program IDs 11096, 12073, 14096; PI: Bleem), and 163 cluster candidates (at any redshift) were within the footprint of the *Spitzer* South Pole Telescope Deep Field (SSDF; Ashby et al. 2013), with 31 of these candidates also having deeper targeted follow-up from the earlier SPT-SZ work.

The SPTpol candidates imaged in Cycles 11-14 were initially detected in preliminary versions of the SPTpol 500d cluster sample, and were selected for infrared follow-up based on the absence of galaxy overdensity counterparts in then existing DES data (DES Year 1 and DES Year 3, depending on the cycle; Abbott et al. 2018). In the new *Spitzer* observations, candidates were observed in each band in 6 × 30 s exposures and the data were reduced following the procedures detailed in Ashby et al. (2009). This exposure time is sufficient to reach a 5σ sensitivity of $4.8\mu\rm Jy$ at $3.6\mu\rm m$ in an aperture-corrected 4 aperture. Here we use 4 diameter aperture-corrected magnitudes.

5. CLUSTER CONFIRMATION AND REDSHIFT ESTIMATION

The majority of newly confirmed cluster candidates in this work were confirmed using the multi-component matched filter cluster confirmation tool (MCMF) previously described in detail in Klein et al. (2018, 2019).

5.1. MCMF

As explained in Klein et al. (2018, 2019), the MCMF algorithm was designed to provide—amongst other properties—cluster confirmation, redshift, and cluster galaxy richness information for samples of X-ray and SZ clusters by robustly identifying associated cluster galaxy counterparts. We provide a brief overview of the method here, including choices specific to this analysis, and refer readers to previous works for more details. The MCMF algorithm works as follows:

- At the location of each cluster candidate, local background-corrected cluster richnesses, λ_{MCMF} , are estimated as a function of redshift from 0.01 < z < 2, in steps of δ_z = 0.005.
- These richnesses are computed as the sum of galaxy weights within a projected radius of r _{500c} centered on the SZ candidate location where r _{500c} is determined from the ξ – M _{500c} relation (see Section 6) at the redshift of interest.
- Galaxy weights are computed for galaxies brighter than $i \le m^*(z)+1.25^1$ and are based on (1) the consistency of galaxy colors and magnitudes with a redshift-dependent cluster population model and (2) a radial weight from the SZ center based on a normalized Navarro, Frenk and White (NFW) profile (Navarro et al. 1997) with a scale radius

of $R_{\rm S}$ = $r_{\rm 500c}/3$ (Hennig et al. 2017). Following e.g., Gladders & Yee (2000), a passive red-sequence population model is used to describe the colormagnitude relation of cluster galaxies. This model was empirically calibrated using ~ 2, 500 clusters with spectroscopic redshifts from the SPT-SZ cluster catalog (Bleem et al. 2015; Bayliss et al. 2016; Khullar et al. 2019), the redMaPPer (RM) Y1 catalog (Rykoff et al. 2016; McClintock et al. 2019), and the MCXC cluster catalog (Piffaretti et al. 2011). At low redshift, the 4000Å break drives redshift determination, while at high redshift (z > 1.1) where WISE data are used, the "1.6 μ m Stellar Bump" feature (see Section 5.2) provides discriminating power.

- A correction is applied as needed to the richness at high redshift where the data are not complete to m*(z)+1.25.
- Random sight lines are used to determine the probability of false associations as a function of redshift (z_i) and richness (λ), with this contamination fraction given by:

$$f_{\text{cont}}(\lambda_i, Z_i) = \frac{R_{\infty}^{\infty}}{R_{\infty}^{i}} \frac{f_{\text{rand}}(\lambda, z_i) d\lambda}{f_{\text{obs}}(\lambda, z_i) d\lambda}.$$
 (2)

where $f_{\rm rand}$ is the richness distributions along random lines-of-sight and $f_{\rm obs}$ is the richness distributions along candidate lines of sight.

- Up to three peaks per candidate are analyzed in richness-redshift space, with the peak associated with the lowest chance of being contamination assigned as the most likely counterpart to the SPT cluster candidate.
- Below a threshold of f cont = 0.2 we denote candidates as "confirmed". The expected net contamination in the confirmed sample is a combination of the intrinsic purity of the SZ sample (see Section 6.4) and the optical/IR contamination. Under the assumption that the follow-up data is sufficiently deep to detect all real associations, the contamination of the optically confirmed sample is given by (see also Klein et al. 2023):

contamination =
$$f_{\text{cont}}^{\text{max}} * (1 - p(\xi > \xi_{\text{min}}))$$
 (3)

where $p(\xi > \xi_{\min})$ is the purity of the complete SZ candidate sample at $\xi > \xi_{\min}$, and we adopt $f_{\text{cont}}^{\max} = 0.2$.

In Table 1 we provide MCMF summary statistics for the cluster candidates (except when confirmed by *Spitzer*, see below) including their redshifts and optical richnesses.

5.2. Spitzer Confirmation

For those cluster candidates with available higherresolution and deeper *Spitzer* observations, we adopt a simpler, though related approach, to measuring the cluster redshifts, richnesses, and false associations. Given the small field-of-view of the targeted observations it is not possible to apply local background corrections.

 $^{^{1}}$ Here m^{\ast} corresponds to the apparent magnitude of L * galaxies, modeled as described in Klein et al. (2019).




Figure 3. (Left) Example of Spitzer redshift determination on two high-z SPTpol clusters. Plotted are IR richness (λ) versus redshift. In the inset we plot the median [3.6]–[4.5] color for SPT clusters with spectroscopic redshifts (green stars) and in brown we overplot our synthetic model calibrated using these systems. As can be seen, the Spitzer colors lose redshift discrimination power at z > 1.6 and so we assign all systems with higher best-fit redshifts to z = 1.6 and note this is a lower limit. (Right) SPT-CL J0151–5332 at $z \ge 1.6$ with identified cluster members marked with white squares. The RGB image is constructed from Spitzer [3.6] and DES *i*- and *g*- band data.

Clusters are confirmed in Spitzer data via the identification of excess galaxies at the cluster candidate locations as a function of [3.6]-[4.5] color via a variation of the "1.6 μm Stellar Bump method" previously employed in e.g., Papovich (2008); Muzzin et al. (2013); Gonzalez et al. (2019). As detailed in these previous works, after excluding low-redshift galaxies, there is a close mapping between the Spitzer color and redshift. We generate our model for this relation using the GALAXEV package (Bruzual & Charlot 2003) assuming that the cluster galaxy population was formed by a single starburst at z = 3 with a Salpeter initial mass function (Salpeter 1955) and then followed the MILES (Vazdekis et al. 2010) evolutionary tracks thereafter. As noted in e.g., Sorba & Sawicki (2010), the 1.6 μ m feature is a robust feature immune to the details of the star formation history for all but the youngest stellar populations.

We match sources selected from the *Spitzer* fields to optically selected counterparts from the DES using a 1 matching radius. Following Muzzin et al. (2013); Gonzalez et al. (2019), to reduce the number of low-z interloper galaxies, we remove galaxies with z = [3.6] < 1.6 and i < 21.3 from our catalogs. We next run the single-color λ -richness estimator introduced in Rykoff et al. (2012) modified to use our 1.6 μ m bump redshift model in place of the red-sequence model. We adopt an intrinsic color-spread in the model at fixed redshift of σ [3.6-4.5] = 0.07 (Muzzin et al. 2013). Background galaxy densities are estimated from either 25 blank-field pointings of unconfirmed cluster candidates from the SPT-SZ sample (Bleem et al. 2015) for the deep *Spitzer* observations or from the full SSDF field for the shallower observations.

Following MCMF (Klein et al. 2023), we also modify the radial extent of the richness aperture to extend to what would be r_{500c} for a cluster detected at significance ξ at the redshift of interest. Richnesses were computed from 0.8 < z < 2 with the candidate assigned the redshift and richness corresponding to the maximum λ value. The [3.6]–[4.5] color loses redshift discrimination power at z > 1.6 so systems with solutions at higher redshift are assigned z = 1.6 and flagged in the catalog. An example of this fitting procedure for two different high-z SPTpol clusters is shown in Figure 3.

While we do apply a color cut to remove low-redshift galaxies, faint galaxies below the DES detection limit can enter our catalog. As the 1.6 μ m bump redshift model above $z \sim 0.7$ is degenerate with solutions at lowredshift (see e.g., Figure 1 in Muzzin et al. 2013), we also apply a secondary run estimating λ versus redshift from 0.1 < z < 2 and visually inspect all outputs, flagging and removing cases where low-redshift interlopers are biasing our results. In the vast majority of cases, these contaminants do indeed correspond to galaxies in the faint end of the luminosity function of rich low-redshift clusters confirmed by MCMF in DES optical data alone. A future improvement to our Spitzer confirmation work will incorporate additional optical information beyond our simple color cut.

We also follow a similar convention to MCMF to estimate the contamination of the confirmed sample at fixed λ but here restrict ourselves to a single wide redshift bin from 1 < z < 2. To compute this contamination fraction we first select ~ 800 random locations in the SSDF (with galaxies also matched to DES and the color cut to remove low-z galaxies applied) that were screened to be > 5 from any SPTpol cluster candidate and not in highly masked regions of DES. We randomly assigned each location a ξ value from the SPTpol catalog and then

 $^{^2}$ Which we convert from Vega to AB magnitudes using the offsets of $m_{\rm AB}$ = $m_{\rm Vega}$ + 2.79 (3.26) for the [3.6]([4.5]) bands (Papovich et al. 2016).

estimated λ as for the real—candidates to generate our random distribution. We denote fractional—contamination determined using *Spitzer* data as f_{Scont} . We adopt a somewhat more conservative threshold ($f_{Scont} < 0.1$ versus $f_{Scont} < 0.2$ from MCMF) for our *Spitzer* confirmation thresholds given the steepness of this contamination fraction with declining richness and our limited ability to sample blind *Spitzer* fields across the full 500d survey.

5.3. Comparison Between WISE and Spitzer Observations

We can compare the performance of MCMF as applied to the DES-WISE dataset on high-z clusters that were also observed in higher-resolution Spitzer/IRAC data. In total we have 108 candidates observed with Spitzer with estimated redshifts $z \ge 0.85$ and Spitzer-derived contamination $f_{Scont} < 0.1$. Of these systems, 78 are also confirmed via MCMF and have redshifts in decent agreement ($|\delta z|$ < 0.2) with the *Spitzer* estimates. For the remaining 30 systems, differences in redshift/confirmation estimates between the MCMF and Spitzer analyses fall in two (expected) categories (1) detection in Spitzer data of higher-z systems not well detected by MCMF in DES+WISE data, (2) identification of different galaxy over-densities along the line of sight, leading to different redshift estimates. In the first scenario, 14 high-z systems confirmed by Spitzer are not confirmed by MCMF $(f_{cont} > 0.2)$; 5 of these systems have MCMF redshifts approximately consistent with those obtained by Spitzer, but with richnesses up to a factor of 2× smaller, and 9 are solely detected in Spitzer data. The remaining 16 systems have significant differences in identified counterparts/redshifts between the algorithms. After further inspection, for 9 of these systems we selected the Spitzeridentified over-density as the primary counterpart as it is better centered on the SZ detection or a richer detection and comparably centered (see Figure 4 and the middle panel of Figure 10 for two examples). For the remaining 7 systems the MCMF detection was selected as the primary counterpart. We flag systems with multiple significant over-densities along the line of sight in the cluster candidate table, Table 1.

In summary, we find generally good agreement between the MCMF/WISE and Spitzer analyses and that the WISE analysis provides an excellent addition to the analysis of DES data when deeper higher-resolution IR imaging is not available. The lower-resolution WISE data naturally has some limitations, and we find that the Spitzer data provide surer cluster confirmations (5 systems) or enables new detections not found in the WISE analysis (15 additional systems), impacting 20/108 clusters (16/51 at z > 1.1) for which we had both datasets. the near future data from the Euclid mission (Laureijs et al. 2011) will enable us to reach beyond the limits of WISE confirmations for the full sample. These data, expected to allow detections of high-z clusters to z ~ 2 (Euclid Collaboration et al. 2019), will enable confirmations of additional high-z systems from this SPTpol sample as well as future cluster samples from SPT-3G (Sobrin et al. 2018).

6. MASS ESTIMATION AND SAMPLE PURITY

In this section, we detail how we connect our SPT observable, ξ , to mass.We also explain how we use realistic




Figure 4. Example of a system with multiple significant overdensities along the line-of-sight. SPT-CL J2331–5737 at ξ = 7.6 has potential counterparts located at (redshift, richness) Z = 0.3, λ = 65 and z = 1.5, λ Spitzer = 25. We adopt the higher-z counterpart as our primary identification given its excellent alignment with the SPT position. While richer, the lower-z counterpart is centered ≥ 70 away from SPT detection; the statistical uncertainty on the SPT position is 11. The RGB image is constructed from *Spitzer* [3.6] and DES r - and g- band data.

simulations of the SPTpol 500d field to explore the expected behavior of our observable-mass scaling relation and to estimate the purity of the SZ candidate list.

6.1. The ζ -Mass Scaling Relation

As in previous SPT cluster catalogs, we use an observable-mass scaling relation to relate our detection significances to mass. This is done through a two step process. First, as discussed in Vanderlinde et al. (2010), to account for the bias owing to the maximization of the cluster detection algorithm over position and filter scales, we relate our observable ξ to ζ , an "unbiased" detection significance via

$$P(\xi|\zeta) = N(^{\mathsf{p}}\overline{\zeta^2 + 3}, 1) \tag{4}$$

for $\zeta > 2$ (see also a formal derivation of this maximization correction in Zubeldia et al. 2021).

Next, this unbiased significance is connected to mass via the relation:

$$\langle \ln \zeta \rangle = \ln A_{\rm SZ} \frac{M_{500c}}{3 \times 10^{14} M_{\odot} h^{-1}} \stackrel{B_{\rm SZ}}{\longrightarrow} \frac{H(z)}{H(0.6)} \stackrel{C_{\rm SZ}}{,}$$
 (5)

and

$$P\left(\ln \zeta | M, z\right) = N\left[\langle \ln \zeta \rangle (M, z), \sigma_{\ln \zeta}\right] \tag{6}$$

where $A_{\rm SZ}$ is the normalization, $B_{\rm SZ}$ the slope, $C_{\rm SZ}$ the redshift evolution, $\sigma_{\ln \zeta}$ the log-normal scatter on ζ , and H(z) is the Hubble parameter. As in Reichardt et al. (2013) and other SPT cluster works thereafter, we rescale the normalization to account for varying depth in the different SPT survey regions

$$A_{SZ} \rightarrow Y_{field} A_{SZ}.$$
 (7)

6.2. Simulations of the SPTpol field

We make use of simulations tailored to mimic the SPTpol observations to measure the rescaling factor, y, for the 500d field as well as to estimate the sample purity. Following Bleem et al. (2020); Huang et al. (2020), we have constructed five realizations of the 500d field by combining realizations of the CMB (Keisler et al. 2011) cosmic infrared background (CIB; Reichardt et al. 2021) discrete radio sources (De Zotti et al. 2005) with spectral indices consistent with Everett et al. (2020), thermal SZ, and instrumental/residual atmospheric noise (the latter hereafter referred to as "instrumental noise" for brevity). The thermal SZ maps are created by applying the methods of Flender et al. (2016) to halo lightcones from the Outer Rim simulation (Heitmann et al. 2019) and the instrumental noise maps are constructed from jackknife coadds of the SPT maps in which half of the observations were multiplied by -1 in order to produce maps with no sky signal. The SPT beam and transfer function are then applied to the simulated maps. The emissive source subtraction on the real maps (Section 2.2) reduces both the noise during cluster detection (by $\sim 3\%$ in the filtered maps) and the number of spurious cluster candidates. To mimic the effect of this subtraction in the simulations, we do not include radio sources brighter than the flux above which the source density of our radio source model matches the density of sources subtracted from the maps.

Following Reichardt et al. (2013), to measure y, we first filter the maps with the same optimal filters used in the cluster identification and extract the ζ values at the known location and optimal filter scales of the simulated clusters. We then fit for the scaling relation parameters (Eq. 5 above). We measure y = 2.23 for the SPTpol 500d field. This value implies the SPTpol 500d field depth should be roughly comparable to SPTpol 100d, and that clusters should be detected at ~1.8× higher significance in SPTpol than SPT-SZ (see e.g., Table 1 in de Haan et al. 2016).

We also tested the consistency of the B_{SZ} and C_{SZ} parameters measured in the SPTpol 500d simulations with those measured in simulated fields from the SPT-SZ (Bleem et al. 2015) and SPT-ECS (Bleem et al. 2020) surveys. While we find B_{SZ} to be consistent with previous values at better than the 1σ level, the best fit C_{SZ} values have increased by $\Delta G_{ZZ} = 0.26$, corresponding to a 7σ shift in the simulations.

This change in $C_{\rm SZ}$ arises from a combination of the reduced noise level and the larger contribution of the 95 GHz data to cluster detection in SPTpol compared to SPT-SZ. Decreasing the noise levelincreases sensitivity to higher-redshift clusters more than lower-redshift ones because of the fixed low- ℓ "noise" contribution from the primary CMB. Meanwhile, the increased weight at 95 GHz improves sensitivity to low-redshift clusters. We have confirmed both of these trends in simulations: The values for $C_{\rm SZ,95~only}$ and $C_{\rm SZ,150~only}$ determined by running the cluster finder on single frequency maps are below

and above the combined joint analysis value by 0.2, respectively, meanwhile excluding instrumental noise from simulations increases $\mathbb{Q}_{\mathbb{Z}}$. We surmise that, in the data, the effect of the noise reduction is larger, resulting in an increase in C_{SZ} .

6.3. Scaling Relation Parameters

In recent works, we have used scaling-relation parameters based on best-fit weighted averages from a Monte Carlo Markov chain (MCMC) analysis of the abundance of clusters in the SPT-SZ dataset as a function of ξ and redshift at our fiducial cosmology (Bocquet et al. 2019). This analysis was conducted using the $\xi \geq 5$ sample for which we had complete optical/IR follow-up and assuming a fixed scatter of $\sigma_{\ln \zeta} = 0.2$ whose value was motivated by comparisons with X-ray observables for a large sample of SPT clusters. Motivated by observed changes in $C_{\rm SZ}$ in our simulations, here we explore the validity of these scaling parameters for the SPTpol 500d sample using the data itself.

In this test we replace the SPT-SZ sample above with the confirmed SPTpol 500d clusters at $\xi > 5$ and run a new abundance analysis at fixed cosmology to estimate the SPTpol 500d mass- ζ scaling relation parameters. The simulated normalization rescaling factors are applied for each survey field to make the normalizations directly comparable. We plot the results in Figure 5 along with the best fit parameters derived for SPT-SZ and SPT-ECS; the latter two results were derived using the $\xi > 5$ confirmed samples from the respective surveys. We indeed observe a shift in the redshift evolution parameter of the scaling relation $C_{\rm SZ} = 0.87 \pm 0.17$ (compared to e.g., 0.64 ± 0.14 in SPT-SZ) but find better than 0.5σ consistency between the two surveys for the amplitude parameter $A_{\rm SZ} = 3.95 \pm 0.23$ (4.08 ± 0.1 in SPT-SZ) and mass slope $B_{\rm SZ} = 1.69 \pm 0.09$ (1.65 ± 0.08 for SPT-SZ).

Given the consistency in $A_{\rm SZ}$ and $B_{\rm SZ}$ for the 3 SPT surveys of significantly different depths, and the increased degeneracy between these parameters in the SPTpol 500d field given the smaller number of massive clusters observed owing to its smaller survey volume, we continue to use the best fit values from SPT-SZ for these parameters. Given the significant shift in the $C_{\rm SZ}$, we adopt the newly derived value from SPTpol 500d data when reporting our cluster masses for the SPTpol 500d sample.

6.4. Expected Purity of the SPTpol 500d Cluster Sample

In this work, we apply a new method to estimate the cluster sample purity that significantly improves the accuracy at lower detection significances. In previous SPT works, the number of false candidates as a function of significance was estimated by running the cluster detection algorithm on sky simulations with no tSZ signal. This is sufficient for characterizing the properties of high-significance samples drawn from maps in which significant instrumental and residual atmospheric noise is present (such as the SPT-SZ sample presented in Bleem et al. 2015 and used in the cosmological analyses of de Haan et al. 2016; Bocquet et al. 2019). However, it is known that astrophysical foregrounds and the tSZ itself are not necessarily Gaussian (see e.g., recent measurements in Crawford et al. 2014; Coulton et al. 2018). The

¹ We use Keisler et al. (2011) instead of more updated results from Planck Collaboration et al. (2020) to maintain consistency with previous generations of SPT cluster simulations, but do note such a change would have negligible impact on the results presented here.




Figure 5. Best fit mass- ζ scaling relation parameters (see Eq. 5) at our fiducial cosmology for the SPT-SZ, SPT-ECS, and SPTpol 500d cluster samples at $\xi > 5$. There is good consistency between the normalization and slope parameters (A $_{SZ}$ and B $_{SZ}$, respectively) in all three surveys; additionally the shift in the redshift evolution parameter (C $_{SZ}$) between SPTpol and the shallower surveys is captured in our simulations discussed in Section 6.2. The smaller area surveyed in SPTpol 500d results in fewer massive clusters and increases the degeneracy between A $_{SZ}$ and B $_{SZ}$; this increases the uncertainty on the recovered A $_{SZ}$ parameter.

impact of this non-Gaussianity on the expected number of false candidates becomes more pronounced as cluster samples are produced from lower-noise data and to lower detection significance.

To illustrate this effect, we measure the number of false candidates in a simulated 500 deg 2 region varying the amplitude of the tSZ signal. The process by which we identify false candidates is discussed below.We plot the results of these simulations in Figure 6. As can be seen, the relative amplitude of the tSZ to the total noise can have a significant impact on the expected number of false detections, especially at lower ξ values; using tSZ-free simulations this quantity can be overestimated by factors of \sim 1.4 at $\xi_{\rm min}$ = 4. This tension with estimations from tSZ-free simulations at low ξ is also seen empirically using optical follow-up observations as discussed in Klein et al.

In light of this effect, we update our method of prediction for the expected number of false candidates (and correspondingly sample purity) to use simulations that are statistically as close to the data as possible. We run the cluster finder on maps with tSZ included and make use of the fact that our tSZ maps are constructed with perfect correspondence to massive halos in the OuterRim Simulation. This provides us with a complete listing of all real possible cluster detections. We then estimate the purity of our cluster sample at $\xi > \xi_{\rm min}$ as follows:

 For each of 5 independent simulated realizations of the SPTpol 500d field (2500 square-degrees in total) we run the cluster detection algorithm. Each realization is drawn from a non-overlapping region of the simulated Compton-y map and has different




Figure 6. Variations in expected number of false detections based on differences in assumed tSZ amplitude for a 500-square-degree SPTpol-like field. In the (top) panel we show the expected number of false detections at $\xi > \xi$ min and in the (bottom) panel the ratio in the number of spurious detections compared to our reference assumptions. We vary the relative amplitude of the tSZ signal in the sky simulations, where "100%" corresponds to the maps having tSZ power which matches SPT power spectrum data (Reichardt et al. 2021) at $\ell = 3000$. Estimating the expected number of false detections through tSZ-free simulations can overestimate their counts by factors of 1.4 at low- ξ .

CMB, foreground, and noise realizations. Starting with the highest-significance cluster detection, candidates are matched to the most massive unmatched simulation halo within 2 arcmin. Matched halos are prevented from being matched with lower-significance candidates.

A similar matching is done for 5000 random sight-lines within each field footprint which determines the probability of randomly associating with a halo of a given mass within the association radius, f_R(M). Because of the strong dependence of number density on halo mass owing to the steepness of the halo mass function, it is important to include mass information when quantifying random associations.

• Next, using the observed number of candidates at $\xi > \xi_{\rm min}$ as well as this random association probability as a function of halo mass, we can estimate the number of false candidates above $\xi_{\rm min}$. Starting at the highest mass (with lowest chance of random association), we calculate the fraction of true associations to total candidates, $p(M, \xi_{\rm min})$, given $N_{\rm obs}$ associations for $N_{\rm cand}$ candidates at significance $> \xi_{\rm min}$ as

$$p(M, \, \xi_{\text{min}}) = \frac{N_{\text{obs}} - N_{\text{cand}} f_{R}(M)}{N_{\text{cand}} (1 - f_{R}(M))}$$

The number of false associations, N_{A} , at $\xi > \xi_{min}$ and mass is then simply

$$N_{\text{FA}}(M, \xi_{\text{min}}) = N_{\text{obs}} - p(M, \xi_{\text{min}})N_{\text{cand}}$$

We integrate with decreasing mass, reducing the number of candidates available for random associations by the true associations calculated in previous steps, to compute the total number of false candidates. This value is given by the sum of cluster candidates unassociated with simulated halos and the number candidates falsely associated with such halos.

 We then estimate the purity of the sample in the real survey by dividing this number of false candidates by the observed number of candidates in the SPTpol sample.

We have estimated this purity under two different assumptions that we plot in Figure 7. The first uses simulations in which we have normalized the tSZ power spectrum at ℓ = 3000 to match previous SPT results (George et al. 2015; Reichardt et al. 2021) and we adopt this as our baseline model. This tSZ model leads to approximately twice as many clusters at $\xi \ge 4$ in the simulated SPTpol field as compared to observation (likely owing to some combination of different cosmological parameters between the simulations and real data, differences from reality of the assumed tSZ profile as a function of mass and redshift, and lack of simulated correlated emission from cluster members with the tSZ signals, though we find no strong evidence for the latter in our data—see Section 7). As an alternative, we estimate the purity using simulations where we have scaled the tSZ amplitude so that the candidate count matches the observed SPTpol results. These two models agree with each other within 1 σ at ξ > 4.5. For reference we also plot the "tSZ-free" simulation case, which shows lower purity estimates at low ξ than the other two models.

Using Eq. 3, our baseline SZ purity model, and the subset of cluster candidates with contamination < 0.2, we estimate an overall purity of our confirmed cluster sample of 93%. We can omit use of the simulations, and estimate a purity such that Eq. 3 is satisfied and we observe consistency between the number of observed unassociated detections and predictions (as was done in Klein et al. 2023). This estimate raises the expected purity value slightly to 95% which corresponds to a $\sim 2.5\sigma$ difference between the two estimates of the purity of the full SZ candidate sample $\xi > 4$. Given the good agreement between the estimates for the confirmed portion, we quote $\sim 94\%$ as the expected purity of our confirmed

sample. To further refine our purity estimates and cluster selection modeling we are in the process of improving our simulations of cluster gas properties (K´eruzor´e et al. 2023) and correlated sources of mm-wavelength contamination. This work will be important to develop in parallel to keep pace with the upcoming cluster samples from e.g., SPT-3G (Sobrin et al. 2018), and Simons Observatory (Simons Observatory Collaboration et al. 2019).

7. SYSTEMATIC EXPLORATIONS

We conduct several tests to search for potential ases in the SZ signal we use to identify SPTpol clusters and estimate their masses. Such biases may arise from correlated emission from cluster galaxies, namely synchrotron from radio sources or emission from dust associated with star formation in cluster members. Based on multi-wavelength studies of the cluster galaxy population (see e.g., Gralla et al. 2011; Alberts et al. 2016; Melin et al. 2018), synchrotron contamination is expected to arise from bright discrete sources such as active galactic nuclei in cluster central galaxies, while contamination correlated with star formation is predominately sourced through the integrated emission from a large number of cluster members. Here we use external radio imaging data as well as internal comparisons of data from the two SPTpol frequencies to search for and estimate the level of bias from these sources in our sample. We note that as this study is being conducted using an SZ-selected sample, the impact of very strong contamination (such as from a powerful radio source) that completely fills in the SZ decrement will not be accounted for in this test. Based on previous studies of optical-, IR-, and X-rayselected systems (e.g., Lin et al. 2009; Gralla et al. 2011, 2014; Gupta et al. 2017; Mo et al. 2020; Dicker et al. 2021), such extreme sources are expected to be rare and not have a large impact on the completeness of SZ surveys at redshifts z > 0.25.

7.1. Radio Source Check

To assess potential contamination from radio sources below the detection threshold of SPTpol, we follow a procedure very similar to that used in Bleem et al. (2020)In this work, we use publicly available thumbnail maps from the Sydney University Molonglo Sky Survey (SUMSS, Mauch et al. 2003) at 843 MHz, while Bleem et al. (2020) used the 1.4 GHz National Radio Astronomy Observatory (NRAO) Very Large Array (VLA) Sky Survey (NVSS, Condon et al. 1998) which unfortunately does not extend as far south as the SPTpol field. Otherwise the process is identical; we summarize the analysis briefly here and refer the reader to Bleem et al. (2020) for more details.

First, we download all SUMSS postage-stamp maps that overlap with the SPTpol 500d field and reproject them onto the same pixel grid as the SPTpol maps. We make beam- and transfer-function-matched SUMSS maps for each of the SPTpol observing frequencies and scale the intensity of the maps assuming a single spectral index of -0.7 (Coble et al. 2003), and we convert the result to CMB fluctuation temperature. We produce maps of contamination to the cluster-finding by combining the single-frequency maps with the same weights as used in the cluster-finding and then filtering the result with each




Figure 7. Expected intrinsic purity of SPTpol 500d candidate list (without considering optical/IR confirmation) under different modeling assumptions for the tSZ contribution to simulated survey maps. Overall our purity for the SPTpol sample agree within 1 σ at ξ > 4.5 for our two bounding assumptions of either (1) matching the tSZ amplitude at ℓ = 3000 to recent power spectrum constraints (Reichardt et al. 2021) or (2) matching the identified candidate density in the 500-sq-degree footprint to that of the observed SPTpol sample. Estimating the number of false detections (and hence purity) of the sample using tSZ-free simulations significantly underestimates the sample purity at lower ξ values.

of the 12 matched filters. We estimate the contamination to ξ for each cluster by taking the value of the appropriate contamination map at the cluster center and dividing by the noise in the actual SPTpol combined, cluster-filtered map at that location. Because of artifacts in the SUMSS map around the bright radio source PKS 2356-61, we are not able to perform this calculation within approximately 1.5 degrees of that source. Eight of the 689 candidates in the SPTpol 500d catalog lie in this area.

The median contamination calculated in this way is $\Delta\xi_{\rm med}=0.032$ (where $\Delta\xi$ is defined such that $\xi_{\rm bserved}=\xi_{\rm true}-\Delta\xi$), or 0.8% of the $\xi=4$ threshold value for inclusion in the catalog. Of the 681 candidates in the catalog for which we are able to perform this test, 47 (~7%) have a predicted contamination of greater than $\Delta\xi=1$, and 18 (~2.5%) have a predicted contamination of greater than $\Delta\xi=2$. Of the 18 candidates with a predicted contamination greater than $\Delta\xi=2$, three are low-redshift (z < 0.25) systems, and two have $\xi<4.25$, leaving only thirteen such candidates that would be included in a cosmological analysis. We flag candidates with predicated contamination $\Delta\xi>2$ in the candidate Table 1.

7.2. 95/150 GHz Internal Consistency Test

We can use cluster samples selected independently from the SPTpol 95 and 150 GHz data to provide an independent test of predictions from our simulations as well as to look for signatures of potential contamination to the SZ signal. This test is based on the premise that




Figure 8. Masses estimated from 95 and 150 GHz data alone for 250 clusters individually detected in each band. Overall we find excellent agreement between the two, with the largest outlier, SPT-CL J2332-5358 (5.4 σ , at a ratio of M $_{95~\rm GHz}$ /M $_{150~\rm GHz}$ = 7/5) having previously been identified as a cluster-scale strong lens of a distant star forming galaxy whose emission results in significantly reduced tSZ signal in 150 GHz data.

the two key expected sources of correlated contaminating emission—synchrotron and thermal dust radiation from cluster member galaxies—should impact the recovered SZ differently at each frequency given their differing spectral energy distributions. Here we compare end-to-end predictions of the masses estimated from each frequency as cluster mass is what we are most concerned about for cosmological analyses.

Following the cluster identification procedure of Section 3 but now run on the single band maps, we detect 540 candidates (514 candidates) at $\xi > 4$ in the 95 (150 GHz) data with 250 common detections representing 221/544 confirmed systems with redshifts.Repeating the simulations above to standardize the extracted ξ values we expect the amplitude normalization factors, y, to be 0.85 (0.78) at 95 (150) GHz relative to the full depth field, with changes in $G_{\rm SZ}$ between the bands as discussed above.

In Figure 8 we plot the masses estimated from each individual frequency for the matched clusters. We find the median ratio in the masses estimated from 95 and 150 GHz alone to be 0.968 ± 0.005We attribute this offset from unity to a small misestimation of the foreground levels in the simulations, due to uncertainty in the amplitude of the uncorrelated CIB at 95 GHz (Reichardt et al. 2021), as well as our simplified modeling of the applied source cleaning, which might affect the bands differently due to the source populations in each map.For example, if we split the sample into halves based on the estimated level of radio contamination from the SUMSS data, the 95/150 GHz mass ratio is only 1% different between the two halves. Regardless,as the amplitude and other parameters of our observable-mass scaling relation will be

determined via weak lensing (e.g., Dietrich et al. 2019; Schrabback et al. 2018, 2021), small shifts of this order in our simulation calibration have negligible impact.

We next check for a redshift evolution in this relation. Splitting this joint sample at its median redshift of z = 0.66, we find the median ratio of the masses to be consistent, $0.97^{+0.01}_{-0.02}$ for the lower redshift half and 0.97 ± 0.01 for the higher redshift half. Restricting ourselves to the highest redshift clusters in the common sample at z > 1 (35 systems) we may see some evolution in the CIB contribution to the 150 GHz band, with the median ratio of 95/150 GHz masses shifting to 1.04 $^{+0.02}_{-0.03}$, indicating that the 150 GHz signal may be being partially filled in. This change would result in a ~ 3.5% change in the estimated masses of these highest redshift systems in our multi-band mass estimates, and is much smaller than our current best fit uncertainty in the C SZ relation calibrated with weak lensing (Schrabback et al. 2021; Zohren et al. 2022) or constrained by our joint lensing+cosmological analyses (Bocquet et al. 2019).

To check for outliers in the mass comparison, we difference the masses determined by data from each frequency and divide by the mass uncertainty determined by the statistical uncertainty only (as the intrinsic Comptony mass scatter should be the same for a given cluster measured at two different frequencies). To account for correlated scatter in the noise at each frequency from common foreground/atmospheric residuals we normalize this distribution to be a Gaussian of unit width by reducing the statistical uncertainty by 0.57, an empirical factor measured by taking a robust measure of the standard deviation of the distribution. 1 We find three clusters where the 95 and 150 GHz mass estimates differ at $> 3\sigma$. The largest discrepancy is SPT-CL J2332-5358 (5.4 σ) which was previously discussed in Vanderlinde et al. (2010); Andersson et al. (2011) as a cluster that is lensing a distant star-forming galaxy. The emission from this galaxy significantly reduces the SZ signal measured at 150 GHz which lowers its 150 GHz mass estimate. This cluster is noticeable as the biggest outlier in Figure 8.

Conversely, there are two systems (SPT-CL J2337-5942, SPT-CL J0154-5619) both with 95 GHz mass estimates less than the 150 GHz mass estimates by ~3.1 σ . These differences might be caused by co-located radio sources. Indeed for SPT-CL J2337-5942 there are 3 low-brightness SUMSS sources (7-10 mJy) within 2 . However, for SPT-CL J0154-5619, there is no SUMSS source within 2 of the SPT location. Given the sample size of 221 clusters, one would expect two systems as \geq 3.1 σ outliers ~ 7% of the time.

If we look instead at clusters detected significantly in one channel but not the other (> 7σ , or 3σ above our search threshold), we find only one system, SPT-CL J2240–6117 at z=0.95, detected in 95 GHz data at $\xi_{95}=8$. Using forced-photometry on the 150 GHz detection map, we find the cluster detected at $\xi_{150}=2.2$. Examination of a preliminary version of the first SPT-3G point source catalog (Archipley et al., in prep), reveals a dusty source detected at 220 GHz at 15 mJy within 0. 2 of the 95 GHz centroid, making this system potentially

a cluster lens similar to SPT-CL J2332-5358 discussed above. The high sensitivity of the SPT-3G at 220 GHz, in combination with significantly deeper data 95 and 150 GHz, will offer more opportunities to identify instances of high-z dusty sources lensed by massive clusters.

8. THE 500-SQUARE-DEGREE SPTPOL SURVEY SZ CLUSTER CATALOG

The SPTpol 500d cluster catalog consists of 689 SZ candidates detected at $\xi > 4$. Using optical and infrared observations we have confirmed 544 of these candidates as galaxy clusters with an expected contamination of our confirmed sample of less than ~ 6%. The redshifts of the confirmed sample are in the range $0.03 < z \le 1.6$ and the masses are in the range $1.5 \times 10^{14} < M_{500c} < 9 \times 10^{14} M_{\odot}/h_{70}$. The sample has a median redshift of 0.7, median mass of $2.5 \times 10^{14} M_{\odot}/h_{70}$, and a spatial density of 1.18 confirmed clusters/deg². We provide the complete cluster candidate list as well as redshifts, estimated masses, and select optical properties for confirmed clusters in Table 1.

In Figure 9 we plot the mass-redshift distribution of the SPTpol 500d sample as compared to several other SZ-selected cluster samples As seen in this plot, the high sensitivity of SPTpol and our extensive follow-up efforts have enabled us to confirm a large number of low-mass and high-redshift clusters; 114 of the SZ clusters are at z > 1 (21% of our total sample). This is a much higher fraction than the SPT-SZ (42/516 or 8%, Bleem et al. 2015) and ACT samples (222/4195, 5%, Hilton et al. 2021). We further compare select properties of the SPTpol sample to these other samples in Section 8.1. The redshift and mass reach of the SPTpol sample will enable a number of exciting studies of both cosmology (e.g., Bocquet et al. 2023) and astrophysics through studies of the mass and redshift evolution of clusters. In Figure 10 we highlight several of the new high-z clusters reported in this work.

In Figure 11 we plot the estimated selection function of the sample at z > 0.25. This selection function is empirically derived via propagating the $\xi \ge 4$ sample cut into mass- and redshift-space using the ξ -mass relation discussed in Section 6. The SPTpol sample is highly (> 90%) complete at masses $M_{500c} > 3.5 \times 10^{14} M_{\odot} / h_{70}$ and z > 0.25. The SPTpol survey, like previous SPT works, has an increasing sensitivity to lower-mass clusters as a function of increasing redshift. As described in Huang et al. (2020), this trend arises mainly from two effects: (1) residual fluctuations from the CMB and atmosphere increase the noise in the maps at larger angular scales (2) self-similar evolution of clusters leads to hotter clusters at fixed mass at higher redshifts and hence makes higher-z systems easier to detect.

8.1. Comparison to Other Cluster Surveys

In this section, we compare the properties of the SPT-pol 500d cluster sample to SZ and optical samples in the same survey region. For the SZ samples we focus on catalogs from ACT and SPT ¹ and check the consistency of mass and redshift estimates for the clusters in common

¹ This robust standard deviation is computed using the median absolute deviation as an initial estimate, and then using Tukey's biweight to weight points, see discussion in e.g., Beers et al. (1990).

¹ While there is overlap of 22 systems with *Planck*, the typically higher redshift and lower mass SPTpol 500d sample adds no significant new information to the SPT-*Planck* comparisons previously




Figure 9. The mass-redshift distribution of the SPTpol from *Planck* (Planck Collaboration et al. 2016b), ACT (Hilton et al. 2021), and SPT-SZ/SPT-ECS (Bleem et al. 2015, 2020). The SPTpo sample consists of 544 clusters, with 21% of the sample at z > 1. As discussed in Section 5.2, redshifts for clusters confirmed in *Spitzer* and WISE data at $z \ge 1.6$ are considered lower limits. These systems are plotted as right-facing triangles in the plot.

between the samples. For our comparisons to the optically selected systems from DES and WISE data we check whether the SZ systems were detected and compare estimated redshifts. Detailed comparisons that more fully leverage the constraining power offered by the lower-mass SPTpol systems, such as exploring the λ – M 500c relation and contamination of the optical samples (see, e.g., Bleem et al. 2020; Grandis et al. 2021), are reserved for future work.

8.1.1. SPT-SZ

There are 118 SPT-SZ cluster candidates detected at $\xi > 4.5$ that fall within the SPTpol 500d footprint and 113 of these candidates in unmasked regions (with masking owing to nearby bright point sources as discussed in Section 2.1). 2 Using an association radius of 3 to identify matching detections, we find that 97 of these candidates have matches in the SPTpol catalog, with median ratio of the significance of cluster detection between SPTpol/SPT-SZ being 1.7, and a median spatial separation of 0.26. As discussed in Klein et al. (2023), for 5 of these matched systems, the improved DES and WISE follow-up data allow us to confirm them and provide redshifts. Of the 16 unmatched SPT-SZ candidates ranging in $4.5 < \xi < 5.2$ (median $\xi = 4.7$), only one-SPT-CL J2232-6151, $\xi = 5.04$, z = 0.79—was reported confirmed by follow-up optical and infrared observations in Bleem et al. (2015). This is consistent with the es-

undertaken in (Bleem et al. 2020) for the SPT-SZ and SPT-ECS samples.

timate for the purity of the optically confirmed sample, using the framework developed in this paper. There are 597 cluster candidates at $\xi > 4$ in SPTpol—that are not found in SPT-SZ, with 448 (385) of these having optical contamination values less than 0.2 (0.05). Of these systems, 123 are at $\xi > 5.5$ where the raw SPTpol candidate list is > 99% pure (Section 6.4).

Comparing the estimated masses we find the median ratio of the masses of common systems in SPTpol/SPT-SZ to be $1.025^{+0.01}_{-0.04}$. A plot showing the masses of the common systems within $\delta_Z=0.1$ (this cut removes 5 systems at z>0.85) is shown in Figure 12.We apply the redshift cut here and in the other SZ sample comparisons in this section to avoid highlighting differences in mass that would purely arise from the masses being estimated at significantly different redshifts.

8.1.2. SPTpol 100d

Next, we compare the SPTpol 500d sample to the sample produced from the SPTpol 100d survey (Huang et al. 2020). This sample consists of 89 clusters detected at $\xi > 4.6$. Of these 89 systems, 86 are in the non-point source masked region and 73 are in the SPTpol sample. The matched candidates have a median spatial separation of 0.22. Using our updated scaling-relation parameters discussed in Section 6, we find the median ratio of the masses of clusters in SPTpol 500d/SPTpol 100d to be 1.02 \pm 0.02 for systems within $\delta = 0.1$. This comparison was conducted using 66 systems, which excludes from the 73 matches above 1 unconfirmed candidate, 2 newly confirmed systems in this work, and 4 systems with updated Spitzer redshifts that shifted them outside this range. We plot the SPTpol 500d and 100d masses against each other in the middle panel of Figure

 $^{^2}$ We note the point source mask between SPT-SZ and SPTpol differs slightly owing to survey noise and source variability as each was constructed using sources brighter than ${\sim}6$ mJy as measured in the maps used for cluster identification.




Figure 10. Three of the high-z clusters discovered in the SPTpol 500d survey. From left to right, SPT-CL J0128-5222 at ξ = 6 and z = 1.4 ± 0.07, SPT-CL J0122-5801 at ξ = 9 and z ≥ 1.6, and SPT-CL J0116-5039 at ξ = 6 and z ≥ 1.6. The SZ detection contours are overlaid on RGB images from *Spitzer* [3.6] and DES *i*- and *g*- band data. Star formation—as traced by bluer emission correlated with the galaxy overdensities—is prominently visible in some of these high-z cluster galaxies, highlighting the ability of this sample to probe clusters in the high-z transitional era between active star formation and passive galaxy evolution. SPT-CL J0122-5801 is one of the systems tagged as having multiple overdensities along the line-of-sight, with an additional foreground system of λ = 16, z = 0.6, and f cont = 0.05. The deep high-resolution *Spitzer* data allow us to detect the more significant distant background cluster.




Figure 11. Expected completeness of the $\xi > 4$ SPTpol sample as a function of M_{500c} at several redshifts. The sample is expected to be > 90% complete at $M_{500c} > 3.5 \times 10^{-14} \ M_{\odot}/h_{70}$ at z > 0.25. At lower redshifts the atmospheric filtering removes larger angular-scale cluster signals leading the sample to be increasingly incomplete at z < 0.25.

12. The ξ values of unmatched SPTpol 100d cluster candidates (8/13 confirmed with redshifts) range from 4.6 to 6.2.

8.1.3. ACT

Finally, repeating the matching exercise with the ACT cluster catalog of Hilton et al. (2021), we find 141 ACT cluster candidates detected at ACT signal-to-noise > 4 fall within the SPTpol 500d footprint and 132 in the non-

masked region. Using the same 3 matching radius as for SPT-SZ, we find 108 of these candidates have matches in the SPTpol sample, with a median ratio of SPTpol/ACT detection signal-to-noise of 1.66 and median spatial separation of 0.33. There are 24 candidates not matched, ranging in ACT detection significance from 4.1 to 5.4σ and having a median reported mass of M_{500c} = 2 × 10 ¹⁴ M_{\odot}/h_{70} .

Reversing the question, there are 389 SPTpol candidates at $\xi > 4$ and north of $\delta = -60$ (the southernmost extent of the ACT sample) not found in the ACT sample, with 290 having been confirmed by optical/IR follow-up observations. The ξ values of these unmatched confirmed clusters lie in the range $4 < \xi < 9$ with a median mass of $M_{500c} = 2.4 \times 10^{14}~M_{\odot}/h_{70}$. Of these systems, 64 are at $\xi > 5.5$, a significance above which candidates are highly pure, independent of optical follow-up (see Figure 7).

A more detailed comparison of the relative completeness of the ACT and SPTpol samples will require careful investigations of the cluster confirmation procedure as well as masking and spatial variations in the noise levels in the ACT maps. Hilton et al. (2021) also did not report a full candidate list, rather only cluster candidates confirmed by their optical/infrared analysis, which further complicates this comparison. The dominant source of follow-up confirmation and redshifts for ACT clusters in the SPTpol region was the redMaPPer algorithm run on DES data (which has excellent agreement with MCMF at z < 1, see Figure 14), followed by the zCluster algorithm run on DESI Legacy Survey data (Dey et al. 2019) which includes optical grz photometry combined with WISE W1 and W2 channel data, and then previously reported cluster redshifts from SPT and the literature. Given the challenges of confirming high-redshift clusters, we would naturally expect most variation in the confirmation fraction of this portion of the sample.

Moving on from the identification of systems, we now




Figure 12. Comparisons of mass estimates for clusters from SPTpol 500d that are in common with systems in SPT-SZ (Bleem et al. 2015), SPTpol 100d (Huang et al. 2020), and ACT (Hilton et al. 2021). Over-plotted is a line representing a one-to-one relationship. As discussed in Section 8.1, we find good agreement between the 3 SPT surveys, with the median ratio of masses of SPTpol 500d/SPT-SZ to be $1.02^{+0.02}_{-0.04}$ and SPTpol 500d/SPTpol 100d to be 1.02 ± 0.02 . We do note a small (2.3 σ) difference in the normalization of the ACT and SPT masses, finding the ratio of masses from SPTpol 500d/ACT = 1.07 \pm 0.03 (differing from the consistency shown in Hilton et al. 2021), but a stronger test of any cross experiment mass calibration differences will come when comparing weak-lensing calibrated samples in future works.

compare the estimated masses. The ACT collaboration uses a different procedure than SPT for estimating the masses of clustersFollowing initial cluster identification, a central Compton-y parameter, y₀, is measured for each candidate using a 24 matched filter. This central value is then corrected for the mismatch as a function of cluster mass and redshift between the expected cluster profile and that adopted for the matched filter. The Arnaud et al. (2010) pressure-mass scaling relation, statistical and assumed intrinsic scatter in 16, and the Tinker et al. (2008) mass function (computed at the same fiducial cosmology we adopt in this work) are then used to compute the probability distribution, $P(M_{500c} | y_o, z)$. The maximum in this distribution is assigned as the cluster mass. This mass is further corrected by a value of 0.71 ± 0.07 using the weak lensing λ - mass calibration of McClintock et al. (2019); these masses are reported as $M_{000c}^{\rm Cal}$ in the ACT catalog.

A plot showing the masses of the common SPTpol 500d/ACT systems within $\delta_z=0.1$ (100 clusters, 8 z>0.96 systems excluded for redshift differences) is shown in Figure 12. Comparing these masses, we find the ratio of SPTpol/ACT masses to be 1.07 ± 0.03 . We find this ratio to be consistent with the results comparing the median mass ratio of 362 common clusters between SPT-SZ and ACT ($1.06^{+0.02}_{-0.006}$). This difference in normalization is larger than the 1.027 ± 0.012 reported in Hilton et al. (2021). That work measured this ratio using 228 common clusters with ACT signal-to-noise > 6 and the full SPT-SZ sample at $\xi>4.5$; we recover the same mass ratio when applying ACT's thresholds. We note given the significant differences in how the mass-scaling relation parameters are determined—SPT's via

the abundance matching of our sample to a fiducial cosmology versus ACT's adoption of a weak lensing calibration via cross matching with the DES redMaPPER calibration (the latter scaling known to be biased by projection effects, especially at lower masses Bleem et al. 2020; Grandis et al. 2021)— that the agreement here between SPT and ACT masses is not a strong cross check of the mass estimation techniques.

8.1.4. Optically Selected Cluster Samples

We compare the SPTpol 500d sample to two optical samples produced by algorithms that adopt complementary approaches to finding clusters in photometric data. The first sample, the DES redMaPPer 1 sample (hereafter RM), was constructed by running the RM algorithm (Rykoff et al. 2016) on DES Y3 data (Abbott et al. 2018). RM identifies clusters by locating spatial over-densities of red-sequence galaxies. The full Y3 RM sample consists of 869,335 systems with a weighted galaxy-count mass proxy $\lambda > 5$ and redshifts 0.1 < z < 0.95. The second sample, produced by Wen & Han (2022, hereafter WH), was constructed using DES and WISE (Wright et al. 2010) data; clusters in this sample were identified as over-densities of stellar mass in photometric redshift slices. This sample contains 151,244 clusters from 0.1 < z < 1.5 above a detection signal-to-noise of 5. The inclusion of IR data from WISE greatly extends the redshift reach of the catalog compared to that of RM.

Here, as the spatial density of the optical systems is much greater than that of the SZ surveys, we define a cluster-match between samples as having a counterpart

¹ Version 6.4.22+2

18 ///, ///, Bleem et al.




Figure 13. Redshift distribution of optical/IR cluster sample matches from RM and WH to the full SPTpol 500d sample Matches are plotted at the SPTpol redshifts. Overall, the RM sample recovers 70% of the SPTpol sample and the WH sample recovers 81%; many of the lower-redshift clusters not recovered in RM are near regions masked in their analysis.

within 2 $^{'}$ and within $\delta_{z}=0.1\times(1+z)$. When multiple systems are found, we adopt the closest match in redshift as the cluster counterpart. In total we match 379/544 (70%) of the SZ clusters using RM and 441/544 (81%) using WH. We plot the redshift distribution of the SPTpol 500d sample and its matched counterparts in the wide-field optical/IR cluster searches in Figure 13.

We next compare the redshift consistency for estimates from the different samples. We relax our matching criteria to include optical systems matched within 2 for candidates which did not have a good spatial and redshift match to allow us to check for instances where RM and WH may have detected clusters not already captured in our analysis or identified the same optical system at a discrepant redshift. To account for the fact that there are candidates for which we have identified lower-significance galaxy excesses along the line-of-sight below our optical confirmation threshold, we assign these unconfirmed systems the redshifts of these over-densities for this exercise only. Finally, it is important to remember that the redshift estimates we are comparing here were conducted using similar or identical datasets (DES Years 3 or 6, with WISE for some a subset of the high-z clusters), though with different methods (photo-z versus red-sequence) and with different spectroscopic calibration samples.

As shown in Figure 14, overall there is good agreement in the measured redshifts for matching systems. We find $\sigma_{\delta_z}/(1+z)=0.009$ (0.014) for RM (WH) at z<1 for clusters we deem a spatial/redshift match, with the blind RM search systematically underestimating matched redshifts at z>0.9 (beyond its stated redshift reliability). At z>1, the scatter broadens for the difference between SPTpol 500d and WH to $\sigma_{\delta_z}/(1+z)=0.04$. Given their small number of high-z spectroscopic calibrators, WH do not report redshift uncertainties for these highest-z systems. The optical/IR systems from the wide-field




Figure 14. Comparison of redshifts of systems from wide-field cluster searches associated with SPTpol 500d candidates by a (1) redshift/spatial proximity match or (2) only a spatial-proximity match if no systems fell within the redshift association cut (for more details of the matching process see Section 8.1.4). In top panel are shown matches to the Wen & Han (2022) optical/IR sample and in the *bottom* panel to the DES RM sample constructed using the DES Y3 dataset. Note that redshift uncertainties are not provided at z > 1 for the WH sample. Plotted in gray on each panel is a line representing a one-to-one relationship. (RM) sample has good spatial and redshift matches for 81% (70%) of the SPTpol sample. Systems with redshifts that significantly differ between SPTpol and the wide-field optical/IR searches have typically low wide-field survey richnesses, and are likely spurious associations with the SZ detections. This highlights the importance of accounting for chance associations when matching galaxy overdensities to SZ detections.

searches that have discrepant redshifts are typically lowrichness and thus likely spurious associations; such discrepant associations are particularly noticeable for high-z SPT clusters at redshifts higher than the RM search. This highlights the importance of quantifying such false association probabilities (Klein et al. 2018; Bleem et al. 2020; Hilton et al. 2021) when mapping associations between optical galaxy over-densities and SZ clusters.

9. CONCLUSIONS

In this work we described the production of a new sample of galaxy clusters detected by their SZ signature in the SPTpol 500d survey. We implemented several important improvements to our cluster catalog production and characterization processes. These improvements include the subtraction of moderate signal-to-noise emissive sources from the temperature maps before cluster finding to reduce the number of spurious candidates, improved cluster simulations and sample purity estimation techniques, and new tests to identify potential biases to our recovered SZ signal.

The new SPTpol 500d sample consists of 689 galaxy cluster candidates detected at $\xi > 4$. These candidates

were selected from coadded maps of depths 5.3 (11.7) μK_{CMB} -arcmin at 150 (95) GHz that were created from > 4000 observations of the SPTpol 500d fieldTo confirm candidates as clusters, we search optical and IR observations from DECam, WISE, and *Spitzer* with the MCMF algorithm and a related IR-based code to identify significant red-sequence and/or IR galaxy over-densities at the candidate locations. We probabilistically confirm 544 of these candidates as galaxy clusters, with an expected confirmed sample purity of ~ 94%. The SPTpol 500d cluster sample has a median mass of 2.5 × 10⁴ M_{\odot}/h_{70} , a median redshift of z = 0.7, and is expected to be > 90% complete at M_{500c} > 3.5 × 10¹⁴ M_{\odot}/h_{70} at z > 0.25. A significant fraction of the systems are at high z, with 114 clusters at z > 1.

Masses are estimated for confirmed clusters via a ξ -mass scaling relation whose parameters are determined by matching the abundance of observed clusters to a fixed ΛCDM cosmology. Using simulations to relatively calibrate this relation for three independently extracted cluster catalogs from data of variable survey depth—the SPT-SZ (Bleem et al. 2015), SPTpol 100d (Huang et al. 2020), and SPTpol 500d cluster samples—we find excellent agreement for the masses of clusters in common between SPTpol 500d and the other surveys. We have used both external datasets and internal checks to test for contamination of the SZ signal that could bias these mass estimates. These investigations include an assessment of potential radio contamination via extrapolating data from the 843 MHz Sydney University Molonglo Sky Survey at cluster locations. This analysis predicts a median contamination of $\Delta \xi_{\text{med}} = 0.032$ and that $\sim 7\%$ of candidates could have a predicted contamination of greater than $\Delta \xi = 1$. However, this prediction requires a significant extrapolation from low-frequency data to the SPT bands, and may be an overestimate. Internal tests comparing masses for clusters detected at 95 and 150 GHz alone show—with the exception of a small number of clusters—insignificant radio or CIB contamination to the SZ signal.

Matching to previous high-resolution SZ surveys in the region, we find 73, 97, and 141 clusters in common with the SPTpol 100d, SPT-SZ, and ACT (Hilton et al. 2021) cluster samples, respectively. These clusters are detected at ~ 1.7× higher signal-to-noise in SPTpol data than SPT-SZ or ACT data. We find good agreement between the masses of systems in common between the SPT surveys and that the SPT masses are ~1.07× higher than the ACT reported masses for the common clusters. A spatial-redshift match to the wide-field optical/IR cluster samples of DES Y3 redMaPPer (Rykoff et al. 2016) and the sample produced by Wen & Han (2022) finds matches for 70-81% of the SPTpol sample, but also highlights the importance of probabilistic confirmation techniques to avoid spurious associations.

While this is the final cluster sample from the SPT-pol collaboration, a number of ongoing astrophysical and cosmological studies are underway that take advantage of its unique low-mass and high-redshift capabilities. Although surveys like SPT-3G (Sobrin et al. 2018), Simons Observatory (Simons Observatory Collaboration 2019), and CMB-S4 (CMB-S4 Collaboration et al. 2019) will in the future surpass the SPTpol survey with deeper and wider data sets, the work here—particularly the explo-

rations of emissive sources, simulation modeling of SZ cluster samples, and internal and external contamination tests—pave the way for maximizing the potential of the SPTpol and these future surveys for SZ cluster science.

As a final note — We also release the coadded mm-wave maps and associated data products used to produce this sample. The maps and supporting products are available at https://pole.uchicago.edu/public/data/sptpol_500d_clusters/index.html, and the NASA LAMBDA website. An interactive sky server with the SPTpol maps and Dark Energy Survey data release 2 images is also available at NCSA https://skyviewer.ncsa.illinois.edu.

ACKNOWLEDGEMENTS

The South Pole Telescope program is supported by the National Science Foundation (NSF) through award OPP-1852617. Partial support is also provided by the Kavli Institute of Cosmological Physics at the University of Chicago. Work at Argonne National Lab is supported by UChicago Argonne LLC, Operator of Argonne National Laboratory (Argonne). Argonne, a U.S. Department of Energy Office of Science Laboratory, is operated under contract no. DE-AC02-06CH11357. This research used resources of the Argonne Leadership Computing Facility, which is supported by DOE/SC under contract DE-AC02-06CH11357. TS acknowledges support from the German Federal Ministry for Economic Affairs and Energy (BMWi) provided through DLR under projects 50OR2002 and 50OR2302, from the German Research Foundation (DFG) under grant 415537506, and the Austrian Research Promotion Agency (FFG) the Federal Ministry of the Republic of Austria for Climate Action, Environment, Mobility, Innovation and Technology (BMK) via grants 899537 and 900565. The Melbourne group acknowledges support from the Australian Research Council's Discovery Projects scheme (No. DP200101068). This work is based in part on observations made with the Spitzer Space Telescope, which was operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA.

Funding for the DES Projects has been provided by the U.S. Department of Energy, the U.S. National Science Foundation, the Ministry of Science and Education of Spain, the Science and Technology Facilities Council of the United Kingdom, the Higher Education Funding Council for England, the National Center for Supercomputing Applications at the University of Urbana-Champaign, the Kavli Institute of Cosmological Physics at the University of Chicago, the Center for Cosmology and Astro-Particle Physics at the Ohio State University, the Mitchell Institute for Fundamental Physics and Astronomy at Texas A&M University, Financiadora de Estudos e Projetos, Funda cão Carlos Chagas Filho de Amparo `a Pesquisa do Estado do Rio de Janeiro, Conselho Nacional de Desenvolvimento Cient'ifico e Tecnol'ogico and the Minist'erio da Ci^encia, Tecnologia e Inovação, the Deutsche Forschungsgemeinschaftand the Collaborating Institutions in the Dark Energy Survey.

The Collaborating Institutions are Argonne National Laboratory, the University of California at Santa Cruz, the University of Cambridge, Centro de Investigaciones Energ´eticas, Medioambientales y Tecnol´ogicas-Madrid,

the University of Chicago, University College London, the DES-Brazil Consortium, the University of Edinburgh, the Eidgen ossische Technische Hochschule (ETH) Zürich, Fermi National Accelerator Laboratory, the University of Illinois at Urbana-Champaign, the Institut de Ci`encies de l'Espai (IEEC/CSIC), the Institut de F'isica d'Altes Energies, Lawrence Berkeley National Laboratory, the Ludwig-Maximilians Universit M unchen and the associated Excellence Cluster Universe, the University of Michigan, the National Optical Astronomy Observatory, the University of Nottingham, The Ohio State University, the University of Pennsylvania, the University of Portsmouth, SLAC National Accelerator Laboratory, Stanford University, the University of Sussex, Texas A&M University, and the OzDES Membership Consortium.

Based in part on observations at Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation

The DES data management system is supported by the National Science Foundation under Grant Numbers AST-1138766 and AST-1536171. The DES participants from Spanish institutions are partially supported by MINECO under grants AYA2015-71825, ESP2015-66861, FPA2015-68048, SEV-2016-0588, SEV-2016-0597, and MDM-2015-0509, some of which include ERDF funds from the European Union. IFAE is partially funded by the CERCA program of the Generalitat de Catalunya. Research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Program (FP7/2007-2013) including ERC grant agreements 240672, 291329, and 306478. We acknowledge support from the Brazilian Instituto Nacional de Ci[^]encia e Tecnologia (INCT) e-Universe (CNPg grant 465376/2014-2).

This manuscript has been authored by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

This research has made use of the VizieR catalogue access tool, CDS, Strasbourg, France (DOI: 10.26093/cds/vizier). The original description of the VizieR service was published in 2000, A&AS 143, 23.

This research made use of APLpy, an open-source plotting package for Python (Robitaille & Bressert 2012; Robitaille 2019).

Facilities: Blanco (DECAM), NSF/US Department of Energy, 10m South Pole Telescope (SPTpol), Spitzer (IRAC), WISE

REFERENCES

Abbott, T. M. C., Abdalla, F. B., Allam, S., et al. 2018, The Astrophysical Journal Supplement Series, 239, 18 Abbott, T. M. C., Adam'ow, M., Aguena, M., et al. 2021, ApJS, 255, 20

Alberts, S., Pope, A., Brodwin, M., et al. 2016, The Astrophysical Journal 829 Anbajagane, D., Chang, C., Jain, B., et al. 2022, MNRAS, 514, Andersson, K., Benson, B. A., Ade, P. A. R., et al. 2011, ApJ, Arnaud, M., Pratt, G. W., Piffaretti, R., et al. 2010, A&A, 517, Ashby, M. L. N., Stern, D., Brodwin, M., et al. 2009, ApJ,, 701, Ashby, M. L. N., Stanford, S. A., Brodwin, M., et al. 2013, ApJS, Austermann, J. E., Aird, K. A., Beall, J. A., et al. 2012, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 8452, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Bayliss, M. B., Ruel, J., Stubbs, C. W., et al. 2016, ApJS, 227, 3 Beers, T. C., Flynn, K., & Gebhardt, K. 1990, AJ, 100, 32 BICEP/Reck Collaboration and SPTpol Collaboration, Ade, P. A. R., Ahmed, Z., et al. 2021, Phys. Rev. D, 103, 022004 Bleem, L. E., Stalder, B., de Haan, T., et al. 2015, ApJS, 216, 27 Bleem, L. E., Bocquet, S., Stalder, B., et al. 2020, ApJS, 247, 25 Bocquet, S., Dietrich, J. P., Schrabback, T., et al. 2019, ApJ, 878, 556 Bocquet, S., Grandis, S., Bleem, L. E., et al. 2023, arXiv e-prints, Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000 Calabretta, M. R., & Greisen, E. W. 2002, A&A, 395, 1077 Cavaliere, A., & Fusco-Femiano, R. 1976, A&A, 49, 137 Chaubal, P. S., Reichardt, C. L., Gupta, N., et al. 2022, ApJ, 931, CHEX-MATE Collaboration, Arnaud, M., Ettori, S., et al. 2021, A&A, 650, A104 Chiu, I., Mohr, J. J., McDonald, M., et al. 2018, MNRAS, 478, CMB-S4 Collaboration, Abazajian, K., Addison, G., et al. 2019, arXiv e-prints, arXiv:1907.04473
Coble, K., Ade, P. A. R., Bock, J. J., et al. 2003, ArXiv Astrophysics e-prints, arXiv:astro-p Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, AJ, Costanzi, M., Saro, A., Bocquet, S., et al. 2021, Phys. Rev. D, Coulton, W. R., Aiola, S., Battaglia, N., et al. 2018, J. Cosmology Astropart. Phys., 2018, 022 Crawford, T. M., Schaffer, K. K., Bhattacharya, S., et al. 2014, Crites, A. T., Henning, J. W., Ade, P. A. R., et al. 2015, ApJ, de Haan, T., Benson, B. A., Bleem, L. E., et al. 2016, ApJ, 832, De Zotti, G., Ricci, R., Mesa, D., et al. 2005, A&A, 431, 893 Dey, A., Schlegel, D. J., Lang, D., et al. 2019, AJ, 157, 168 Dicker, S. R., Battistelli, E. S., Bhandarkar, T., et al. 2021, Dietrich, J. P., Bocquet, S., Schrabback, T., et al. 2019, MNRAS, Euclid Collaboration, Adam, R., Vannier, M., et al. 2019, Astronomy and Astrophysics, 627, A23
Everett, W. B., Zhang, L., Crawford, T. M., et al. 2020, ApJ, Fazio, G. G., Hora, J. L., Allen, L. E., et al. 2004, ApJS, 154, 10 Flaugher, B., Diehl, H. T., Honscheid, K., et al. 2015, AJ, 150, Flender, S., Bleem, L., Finkel, H., et al. 2016, ApJ, 823, 98 Flores, A. M., Mantz, A. B., Allen, S. W., et al. 2021, MNRAS, George, E. M., Reichardt, C. L., Aird, K. A., et al. 2015, ApJ, Ghirardini, V., Bulbul, E., Kraft, R., et al. 2021, ApJ, 910, 14 Gladders, M. D., & Yee, H. K. C. 2000, AJ, 120, 2148 Gonzalez, A. H., Gettings, D. P., Brodwin, M., et al. 2019, The Journal Gralla, M. B., Gladders, M. D., Yee, H. K. C., & Barrientos, L. F. 2011, ApJ, 734, 103 Gralla, M. B., Crichton, D., Marriage, T. A., et al. 2014, Grandis, S., Mohr, J. J., Costanzi, M., et al. 2021, MNRAS, 504, Gupta, N., Saro, A., Mohr, J. J., et al. 2017, MNRAS, 467, 3737 Gupta, N., Reichardt, C. L., Ade, P. A. R., et al. 2019, MNRAS, Heitmann, K., Finkel, H., Pope, A., et al. 2019, ApJS, 245, 16 Hennig, C., Mohr, J. J., Zenteno, A., et al. 2017, MNRAS, 467,

Henning, J. W., Sayre, J. T., Reichardt, C. L., et al. 2018, ApJ, 852, 97

```
Hilton, M., Hasselfield, M., Sif´on, C., et al. 2018, ApJS, 235, 20 Hilton, M., Sif´ on, C., Naess, S., et al. 2021, ApJS, 253, 3 Huang, N., Bleem, L. E., Stalder, B., et al. 2020, AJ, 159, 110 Keisler, R., Reichardt, C. L., Aird, K. A., et al. 2011, ApJ, 743, 28 K´eruzor'e, F., Bleem, L. E., Buehlmann, M., et al. 2023, arXiv
Khullar, G., Bleem, L. E., Bayliss, M. B., et al. 2019, ApJ, 870, 7
Khullar, G., Bayliss, M. B., Gladders, M. D., et al. 2022, ApJ,
Kim, K. J., Bayliss, M. B., Noble, A. G., et al. 2023, ApJ, 955, 32
Klein, M., Mohr, J. J., Desai, S., et al. 2018, MNRAS, 474, 3324
Klein, M., Grandis, S., Mohr, J. J., et al. 2019, MNRAS, 488, 739
Klein, M., Mohr, J. J., Bocquet, S., et al. 2023, arXiv e-prints,
Laureijs, R., Amiaux, J., Arduini, S., et al. 2011, arXiv e-prints,
Lin, Y., Partridge, B., Pober, J. C., et al. 2009, ApJ, 694, 992
Mainzer, A., Bauer, J., Cutri, R. M., et al. 2014, ApJ, 792, 30
Mauch, T., Murphy, T., Buttery, H. J., et al. 2003, MNRAS, 342,
McClintock, T., Varga, T. N., Gruen, D., et al. 2019, MNRAS
McDonald, M., Allen, S. W., Bayliss, M., et al. 2017, ApJ, 843, 28 Melin, J.-B., Bartlett, J. G., Cai, Z.-Y., et al. 2018, A&A, 617,
Melin, J.-B., Bartlett, J. G., & Delabrouille, J. 2006, A&A, 459,
Mo, W., Gonzalez, A., Brodwin, M., et al. 2020, ApJ, 901, 131
Muzzin, A., Wilson, G., Demarco, R., et al. 2013, ApJ, 767, 39
Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490,
Oke, J. B. 1974, ApJS, 27, 21
Olivares, V., Su, Y., Nulsen, P., et al. 2022, MNRAS, 516, L101
Orlowski-Scherer, J., Di Mascolo, L., Bhandarkar, T., et al. 2021,
A&A, 653, A135
Papovich, C. 2008, ApJ, 676, 206
Papovich, C., Shipley, H. V., Mehrtens, N., et al. 2016, ApJS,
Piffaretti, R., Arnaud, M., Pratt, G. W., Pointecouteau, E., &
Melin, J.-B. 2011, A&A, 534, A109
Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016a,
     2016b, A&A, 594, A27
Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2020,
Raghunathan, S., Patil, S., Baxter, E., et al. 2019,
Reichardt, C. L., Stalder, B., Bleem, L. E., et al. 2013, ApJ, 763,
Reichardt, C. L., Patil, S., Ade, P. A. R., et al. 2021, ApJ, 908,
```

```
Robitaille, T. 2019, APLpy v2.0: The Astronomical Plotting
Library in Python
Robitaille, T., & Bressert, E. 2012, APLpy: Astronomical
Plotting Library in Python, Astrophysics Source Code Library,
Ruppin, F., McDonald, M., Hlavacek-Larrondo, J., et al. 2023,

Ruppin, F., McDonaid, M., Hiavacek-Larrondo, J., et al. 2023, ApJ, 948, 49
Rykoff, E. S., Koester, B. P., Rozo, E., et al. 2012, ApJ, 746, 178
Rykoff, E. S., Rozo, E., Hollowood, D., et al. 2016, The Astrophysical Journal Supplement Series, 224, 1
Salpeter, E. E. 1955, ApJ, 121, 161
Salvati, L., Saro, A., Bocquet, S., et al. 2022, ApJ, 934, 129
Sayre, J. T., Reichardt, C. L., Henning, J. W., et al. 2020, Phys. Rev. D. 101, 122003

Phys. Rev. D, 101, 122003
Schlafly, E. F., Meisner, A. M., & Green, G. M. 2019, The
     Astrophysical Journal Sup
 Schrabback, T., Applegate, D., Dietrich, J. P., et al. 2018,
 Schrabback, T., Bocquet, S., Sommer, M., et al. 2021, MNRAS,
Simons Observatory Collaboration. 2019, Journal of Cosmology
    and Astro-Particle
Simons Observatory Collaboration, Ade, P., Aguirre, J., et al. 2019, Journal of Cosmology and Astro-Particle Physics, 2019,
Sobrin, J. A., Ade, P. A. R., Ahmed, Z., et al. 2018, in Society of
Photo-Optical Instrumentation Engineers (SPIE) Conference
Series, Vol. 10708, Proc. SPIE, 107081H
Somboonpanyakul, T., McDonald, M., Noble, A., et al. 2022, AJ,
Sorba, R., & Sawicki, M. 2010, ApJ, 721, 1056
Strazzullo, V., Pannella, M., Mohr, J. J., et al. 2019, A&A, 622,
Sunyaev, R. A., & Zel'dovich, Y. B. 1972, Comments on
Astrophysics and Space Physics, 4, 173
Tinker, J., Kravtsov, A. V., Klypin, A., et al. 2008, ApJ, 688, 709
Vanderlinde, K., Crawford, T. M., de Haan, T., et al. 2010, ApJ,
Vazdekis, A., S'anchez-Bl'azquez, P., Falc'on-Barroso, J., et al.
    2010, MNR
Wen, Z. L., & Han, J. L. 2022, MNRAS, 513, 3946
Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010,
Wu, W. L. K., Mocanu, L. M., Ade, P. A. R., et al. 2019, ApJ,
Zohren, H., Schrabback, T., Bocquet, S., et al. 2022, A&A, 668,
Zubeldia, I., & Challinor, A. 2019, MNRAS, 489, 401
Zubeldia, İ., Chluba, J., & Battye, R. 2023, MNRAS, 522, 5123
Zubeldia, Í., Rotti, A., Chluba, J., & Battye, R. 2021, MNRAS,
```

507, 4852

SPT ID	R.A. (J2000)	Decl. (J2000)	Be ξ	st $\theta_{ extsf{c}}$	Redshift	M_{500c} $(10^{14} h_{70}^{-1} M_{\odot})$	λ	Spitzer	Notes
SPT-CL J0000-6020	0.0348	-60.3367	7.15	0.25	0.76±0.02	3.15 ^{+0.43} _{-0.52} 2.14 ^{+0.33} _{-0.47}	51±8	0	-
SPT-CL J0000-5825	0.0855	-58.4250	4.03	0.50	0.74±0.03	2.14 ^{+0.33} _{-0.47}	30±7	0	-
SPT-CL J0000-6301	0.1269	-63.0190	4.44	0.25	0.245±0.008	2.67 ^{+0.44} _{-0.57}	55±9	0	-
SPT-CL J0000-6121	0.1890	-61.3538	4.01	0.25	-	_	-	0	-
SPT-CL J0000-5748	0.2479	-57.8083	14.58	0.25	0.702	5.12 ^{+0.56} _{-0.69}	85±10	0	RC
SPT-CL J0001-5917	0.3674	-59.2986	4.09	0.25	0.57±0.03	2.27+0.36	17±5	0	-
SPT-CL J0001-5552	0.3867	-55.8667	4.10	0.50	0.28±0.01	2.51+0.41	28±6	0	-
SPT-CL J0001-6258	0.4021	-62.9795	4.86	0.50	0.22±0.01	2 02+0.49	26±6	0	_
SPT-CL J0001-5440	0.4146	-54.6679	8.34	0.25	0.92±0.03	2 22+0.42	47±8	0	_
SPT-CL J0001-5614	0.4807	-56.2437	4.96	0.75	0.41±0.01	0.31	41±6	0	_
SPT-CL J0002-5557	0.4007	-55.9623	7.86	0.73	1.60±0.08	2.71 _{-0.56} 2.56 ^{+0.33} -0.39		1	LL
							18±3		LL
SPT-CL J0002-5017	0.6415	-50.2894	5.18	0.25	0.97±0.03	2.30 ^{+0.39} _{-0.46}	35±7	0	-
SPT-CL J0003-5938 SPT-CL J0004-5814	0.8124 1.0230	-59.6381 -58.2430	4.06 4.06	1.00 1.00	-	-	-	0 0	-
SPT-CL J0004 5614 SPT-CL J0005-5415	1.2639	-54.2519	7.99	0.25	0.52±0.01	3.70 ^{+0.48} _{-0.59}	56±8	0	_
		-58.8215			0.64±0.03	2.27 ^{+0.37}			_
SPT-CL J0005-5849	1.2785		4.25	0.25		2.27-0.50	24±5	0	-
SPT-CL J0005-5426	1.4344	-54.4359	4.28	0.75	0.85±0.03	2.12-0.47	39±7	0	-
SPT-CL J0005-5758	1.4655	-57.9698	5.17	0.25	0.37±0.01	2.82_0.57	44±7	0	-
SPT-CL J0006-5143	1.5704	-51.7297	4.52	1.25	0.40±0.01	2.55 ^{+0.43} _{-0.55}	49±7	0	LOS
SPT-CL J0007-5710	1.7826	-57.1680	4.48	0.25	0.58±0.02	2.55 ^{+0.43} 2.38 ^{+0.40} 2.38 ^{+0.40}	50±7	0	LOS
SPT-CL J0007-5719	1.9268	-57.3277	5.95	0.50	0.60±0.02	2.38 _{-0.51} 2.90 _{-0.51}	47±7	0	_
SPT-CL J0007-5605	1.9781	-56.0897	4.60	0.50	0.95±0.03	2 13+0.36	33±6	0	_
SPT-CL J0008-6247	2.0091	-62.7914	4.25	0.25	0.97±0.03	2 02+0.32	31±6	0	RC
SPT-CL J0008-5334	2.2486	-53.5782	4.52	0.50	-	2.02 _{-0.44}	-	0	-
SPT-CL J0009-4949	2.4618	-49.8267	4.08	1.00	-	-	-	Ö	_
SPT-CL J0010-6422	2.5448	-64.3674	4.16	0.50	-	-	-	0	-
SPT-CL J0010-6042	2.6230	-60.7137	5.43	0.50	0.80±0.02	2.53 ^{+0.43} _{-0.50}	38±7	0	-
SPT-CL J0010-5450	2.6790	-54.8345	4.29	0.25	0.52±0.02	2 27+0.38	23±5	0	_
SPT-CL J0010-5202	2.7092	-52.0437	4.27	0.25	0.96±0.04	2 03+0.34	21±6	0	_
SPT-CL J0010-5112	2.7398	-51.2038	6.83	1.50	0.210±0.007	IN-43	56±8	0	_
SPT-CL J0011-5541	2.9803	-55.6944	4.24	0.25	0.94±0.05	2 03+0.32	15±5	0	_
SPT-CL J0011 5341	3.0747	-53.8692	4.96	1.25	0.34±0.03	2 76+0.49	46±7	0	_
SPT-CL J0012-5552 SPT-CL J0012-4959	3.2111	-49.9852	5.22	0.50	0.34±0.01	2.70_0.57	40±1	0	_
SPT-CL J0012 4959 SPT-CL J0012-5959	3.2475	-59.9861	5.08	0.30	0.72±0.03	2.48 ^{+0.43}	24±6	0	_
		-57.2429				2.48 _{-0.50} 2.45 ^{+0.42} 2.52			_
SPT-CL J0013-5714	3.2946		4.73	1.50	0.61±0.02		45±7	0	-
SPT-CL J0013-5310	3.4148	-53.1815	5.56	0.50	0.89±0.03	2.40_0.49	45±8	0	LOS
SPT-CL J0013-5926	3.4787	-59.4499	4.62	0.25	0.98±0.06	2.11-0.46	29±6	0	-
SPT-CL J0014-5142	3.5477	-51.7065	4.03	0.75	-	- +0 54	-	0	-
SPT-CL J0014-4952	3.7042	-49.8776	14.48	0.25	0.752	5.02 ^{+0.54} _{-0.68}	98±11	0	-
SPT-CL J0014-6446	3.7288	-64.7675	4.21	0.25	0.78±0.04	2.15 ^{+0.35} _{-0.47}	17±5	0	-
SPT-CL J0015-5004	3.7826	-50.0757	9.45	0.25	0.96±0.02	3.54 ^{+0.43} _{-0.52}	61±9	0	-
SPT-CL J0015-5604	3.8840	-56.0809	4.21	0.25	1.31±0.07	5.02 ⁺ 0.68 2.15 ⁺ 0.35 2.15 ⁻ 0.47 3.54 ⁺ 0.43 1.79 ⁺ 0.27	24±6	0	_
SPT-CL J0015-5146	3.8981	-51.7727	7.62	0.50	0.414±0.009	1.79 ^{+0.50} _{-0.39} 3.72 ^{+0.50} _{-0.60} 2.35 ^{+0.37} _{-0.51}	84±9	0	_
SPT-CL J0016-5650	4.2114	-56.8405	4.02	0.75	0.47±0.02	2.35 +0.37	31±6	0	_
SPT-CL J0018-5504	4.5592	-55.0697	5.20	0.75	0.42±0.01	2.81 ^{+0.48} _{-0.56}	53±8	0	RC
SPT-CL J0018-5304 SPT-CL J0018-6148	4.5878	-61.8119	4.02	0.75	-	2.01-0.56	-	0	-
SPT-CL J0018-5700	4.6800	-57.0118	4.11	0.25	-	_	-	0	_
SPT-CL J0019-5527	4.8269	-55.4525	12.94	0.25	0.93±0.03	4.38 ^{+0.52} _{-0.61}	53±8	0	_
SPT-CL J0019-5128	4.8894	-51.4723	5.12	0.50	0.62±0.02	2.58 ^{+0.45} _{-0.52}	40±7	0	_
SPT-CL J0020-5612	5.2074	-56.2100	5.68	0.25	1.17±0.07	2.00-0.52	29±4	1	_
						2.29 ^{+0.40} 2.99 ^{+0.51} 2.99 ^{+0.51}	!		_
SPT-CL J0021-5900	5.3445 5.5924	-59.0070 -50.7104	5.56	0.75	0.36±0.01	2.99_0.57	46±7	0	_
SPT-CL J0022-5043 SPT-CL J0023-6037	5.5924 5.7988	-50.7194 -60.6255	4.70 4.38	2.00 0.25	_		_	0 0	_
SPT-CL J0023-6037 SPT-CL J0023-5450	5.9387	-54.8430	5.10	0.25	0.41±0.01	2 76 ^{+0.49}	- 44±7	0	_
						2.70-0.56			_
SPT-CL J0023-5905	5.9869	-59.0897	5.74	0.25	0.58±0.02	2.01 _{-0.53}	48±8	0	_
SPT-CL J0024-6001	6.0309	-60.0210	4.27	0.25	1.03±0.04	1.97-0.44	22±6	0	-
SPT-CL J0024-6338	6.0549	-63.6336	4.43	0.25	1.28±0.07	1.86-0.40	20±5	0	-
SPT-CL J0024-5752	6.0798	-57.8703	4.03	0.25	0.85±0.04	2.05 ^{+0.32} _{-0.44}	21±6	0	-
SPT-CL J0024-5151	6.1538	-51.8648	5.61	0.50	0.75±0.02	2.76 ^{+0.49} _{-0.56} 2.87 ^{+0.47} _{-0.53} 1.97 ^{+0.33} _{-0.44} 1.86 ^{+0.30} _{-0.40} 2.05 ^{+0.32} _{-0.44} 2.63 ^{+0.40} _{-0.50} 2.44 ^{+0.44} _{-0.51}	60±9	0	LOS
SPT-CL J0024-6301	6.1704	-63.0222	4.88	0.25	0.67±0.02	2 44+0.44	45±7	0	l _

Table 1 — Continued

SPT ID	R.A. (J2000)	Decl. (J2000)	Β ε	est $ heta_{ m c}$	Redshift	$M_{500c} (10^{14} h_{70}^{-1} M_{\odot})$	λ	Spitzer	Notes
SPT-CL J0024-5147	6.2343	-51.7968	4.37	0.25	0.86±0.04	2.12 ^{+0.35} _{-0.47}	15±5	0	_
SPT-CL J0025-5619	6.3622	-56.3232	6.79	0.50	0.81±0.02	2.99 ^{+0.45}	38±7	0	LOS
SPT-CL J0025-5034	6.3676	-50.5791	9.26	0.50	0.36±0.01	4.30+0.53	55±7	0	-
SPT-CL J0025-5120	6.4535	-51.3395	4.44	0.25	0.35±0.01	2.56 ^{+0.44}	41±6	0	-
SPT-CL J0026-5855	6.5763	-58.9184	6.07	1.00	0.59±0.02	2.98+0.47	44±8	0	-
SPT-CL J0026-5640	6.6204	-56.6818	4.81	0.25	0.47±0.02	2.59+0.46	30±6	0	-
SPT-CL J0026-5700	6.7270	-57.0123	4.88	0.25	0.78±0.02	2.35+0.40	45±7	0	LOS
SPT-CL J0027-5456	6.8825	-54.9483	5.00	0.25	0.83±0.03	2.35+0.40	29±6	0	-
SPT-CL J0027-6326	6.9483	-63.4353	9.67	0.25	0.86±0.03	3.75+0.45	24±7	0	_
SPT-CL J0027-5313	6.9607	-53.2293	6.29	0.75	0.53±0.02	3.73 _{-0.55} 3.11 ^{+0.48}	47±7	0	RC
SPT-CL J0028-6426	7.1692	-64.4468	5.35	0.25	0.61±0.02	2.67+0.46	45±7	0	_
SPT-CL J0030-5213	7.5117	-52.2299	6.74	0.25	0.52±0.02	3.27+0.49	45±8	0	LOS
SPT-CL J0030-5615	7.5495	-56.2654	4.56	0.25	0.52±0.03	2.45 ^{+0.42}	10±4	0	_
SPT-CL J0030-6134	7.7407	-61.5720	5.79	0.25	0.85±0.02	2.60+0.45	45±7	0	_
SPT-CL J0031-5508	7.8273	-55.1439	4.90	0.25	0.81±0.02	2 33+0.40	37±7	0	LOS
SPT-CL J0031-6433	7.9829	-64.5542	4.70	0.25	1.10±0.06	2 02+0.36	31±4	1	_
SPT-CL J0031-5712	7.9970	-57.2113	4.05	0.25	0.90±0.02	2.03 _{-0.45} 2.02 ^{+0.31}	50±7	0	_
SPT-CL J0032-5746	8.1115	-57.7734	6.99	0.50	0.67±0.01	3.21+0.47	74±10	0	_
SPT-CL J0032-5211	8.1165	-52.1948	4.12	0.25	0.51±0.03	2.34 +0.36	14±4	0	_
SPT-CL J0032 5211	8.2167	-55.6693	5.09	1.00	0.202±0.008	2 04+0.50	39±6	0	_
SPT-CL J0032=5340 SPT-CL J0033=6117	8.2608	-61.2901	4.03	0.25	0.82±0.003			0	_
SPT-CL J0033-6117 SPT-CL J0033-5027	8.2927	-50.4538	4.03	0.25	0.02±0.03	2.08 ^{+0.32}	23±5	0	_
SPT-CL J0033-6326	8.4711	-63.4456	13.46	0.50	0.597	5.03 ^{+0.58} _{-0.70}	68±8	0	_
SPT-CL J0034-5906	8.6528	-59.1130	4.92	0.50	0.59±0.02	2 52+0.45	36±7	0	LOS
SPT-CL J0035-5455	8.8176	-54.9295	6.58	0.75	0.30±0.01	2 47+0.49	42±7	0	_
SPT-CL J0035-6030	8.8704	-60.5161	5.11	0.50	0.92±0.03	2.32 ^{+0.41}	37±7	0	_
SPT-CL J0036-6241	9.1737	-62.6913	5.53	0.25	0.202±0.007	2 14+0.51	63±8	0	LOS
SPT-CL J0036-6000	9.2339	-60.0136	5.78	0.25	0.89±0.03	2 57+0.43	46±7	0	_
SPT-CL J0037-6327	9.2882	-63.4666	5.14	0.25	0.85±0.03	10.40	37±7	0	_
SPT-CL J0037 -5047	9.4429	-50.7934	8.26	0.25	1.026	2.39 ^{+0.42} _{-0.48} 3.18 ^{+0.40} _{-0.50}	44±0	0	_
SPT-CL J0037 = 3047 SPT-CL J0038 = 6215	9.5065	-62.2590	6.41	1.25	0.231±0.007	3.47 ^{+0.53}	68±8	0	_
SPT-CL J0038-6436	9.5726	-64.6005	6.18	0.25	0.231±0.007 0.61±0.02	0.00	60±8	0	_
SPT-CL J0038-5040	9.6165	-50.6735	4.34	0.25	1.22±0.06	-0.55	16±5	0	_
		-52.7439	7.87	0.73	0.43±0.01	1.87 ^{+0.30} _{-0.42} 3.78 ^{+0.49} _{-0.59}		0	_
SPT-CL J0038-5244 SPT-CL J0039-5007	9.7196 9.8005	-52.7439 -50.1267	4.01	0.50	0.43±0.01	3.70 _{-0.59}	67±8	0	_
SPT-CL J0040-6351	10.0648	-63.8555	4.09	0.75	1.01±0.05	1.95 ^{+0.30}	19±6	0	_
SPT-CL J0040-6347	10.1694	-63.7859	5.27	0.50	-	- 1.95_0.44	-	0	_
SPT-CL J0041-6030	10.2714	-60.5131	4.09	0.25	-		-	0	-
SPT-CL J0041-5343	10.2810	-53.7291	4.79	0.25	0.62±0.02	$2.47^{+0.42}_{-0.52}$	34±6	0	-
SPT-CL J0041-5107	10.3028	-51.1186	6.07	0.25	0.51±0.01	3.06+0.47	61±8	0	-
SPT-CL J0041-5119	10.4302	-51.3207	4.25	0.25	-			0	-
SPT-CL J0042-6244	10.5539	-62.7433	5.30	0.25	1.60±0.08	1.87-0.35	32±4	1	LL
SPT-CL J0043-5314	10.8292	-53.2362	4.22	0.25	0.70±0.02	2.22-0.35	70±9	0	-
SPT-CL J0043-6241	10.8382	-62.6836	4.10	0.50	0.55±0.02	2.30 -0.50	47±8	0	-
SPT-CL J0043-6433	10.9242	-64.5563	4.02	0.25	0.44±0.02	1.87 ^{+0.33} 2.22 ^{+0.35} 2.30 ^{+0.36} 2.30 ^{+0.36} 2.36 ^{+0.37} 2.66 ^{+0.40} 2.16 ^{+0.40} 2.16 ^{+0.40} 2.23 ^{+0.39} 2.23 ^{+0.39}	21±5	0	-
SPT-CL J0044-6415	11.0309	-64.2565	6.76	0.25	1.13±0.06	2.66 ^{+0.40} _{-0.45}	29±4	1	-
SPT-CL J0044-5737	11.0507	-57.6234	4.05	0.50	0.70±0.02	2.16 ^{+0.34} _{-0.48}	54±8	0	-
SPT-CL J0044-5157	11.2408	-51.9643	4.91	0.25	0.93±0.06	2.23 ^{+0.39} _{-0.47}	34±6	0	-
SPT-CL J0045-6254	11.2543	-62.9128	4.00	0.25	-	_	-	0	-
SPT-CL J0045-6039	11.2607	-60.6517	4.96	0.25	0.61±0.02	2.52-0.43	29±6	0	-
SPT-CL J0045-6236	11.4931	-62.6089	5.57	0.25	0.54±0.02	2.82-0.53	40±7	0	-
SPT-CL J0046-5239	11.5438	-52.6639	4.24	0.25	0.54±0.02	2.35-0.59	30±6	0	-
SPT-CL J0046-5537	11.6335	-55.6312	4.23	1.50	0.164±0.009	2.65-0.58	30±6	0	-
SPT-CL J0046-6456	11.6568	-64.9391	4.42	0.25	0.79±0.03	2.21 +0.30	22±5	0	LOS
SPT-CL J0046-5413	11.6642	-54.2276	4.19	0.50	0.44±0.01	2.41 -0.53	42±6	0	LOS
SPT-CL J0046-6502	11.6843	-65.0433	4.17	1.50	0.032±0.005	$2.71^{+0.45}_{-0.60}$	19±3	0	-
SPT-CL J0046-5825	11.7044	-58.4228	5.34	0.25	0.65±0.02	2.63 ^{+0.46} _{-0.51}	42±7	0	-
SPT-CL J0047-6428	11.7508	-64.4765	4.75	0.25	1.22±0.06	2.52 ^{+0.45} 2.82 ^{+0.47} 2.82 ^{+0.47} 2.35 ^{+0.53} 2.35 ^{-0.51} 2.65 ^{+0.44} 2.65 ^{+0.46} 2.21 ^{+0.36} 2.21 ^{+0.46} 2.41 ^{+0.45} 2.71 ^{+0.45} 2.63 ^{+0.46} 2.63 ^{+0.46} 1.97 ^{+0.44} 2.43 ^{+0.39}	44±8	0	-
SPT-CL J0047-6057	11.9330	-60.9513	4.24	0.50	0.43±0.01	2.43 ^{+0.39} _{-0.53}	40±7	0	-

Table 1 — Continued

SPT ID	R.A. (J2000)	Decl. (J2000)	Β e	st $\theta_{ extsf{c}}$	Redshift	M_{500c} $(10^{14}h_{70}^{-1}M_{\odot})$	λ	Spitzer	Notes
SPT-CL J0048-5244	12.0884	-52.7478	15.45	0.25	1.03±0.02	4.74 ^{+0.51}	72±10	0	LOS
SPT-CL J0048-6416	12.2362	-64.2709	14.32	0.25	0.92±0.02	4.74 ^{+0.51} 4.70 ^{+0.52} 4.70 ^{+0.52}	61±9	0	_
SPT-CL J0049-5615	12.2950	-56.2536	6.09	0.25	0.40±0.01	4.70 ^{+0.52} 	56±7	0	LOS
SPT-CL J0049-5315	12.3777	-53.2527	12.84	0.25	0.72±0.01	4.70 +0.52	87±10	0	_
SPT-CL J0049-5247	12.4467	-52.7946	6.65	0.25	0.60±0.02	3.16-0.54	43±7	0	LOS
SPT-CL J0049-6200	12.4603	-62.0003	4.65	0.25	0.57±0.02	2.45 ^{+0.43}	45±7	0	_
SPT-CL J0049-5709	12.4690	-57.1522	5.17	0.50	0.87±0.02	2.37 ^{+0.42} _{-0.49}	61±9	0	LOS
SPT-CL J0050-5646	12.6118	-56.7808	4.17	0.25	-	-0.49	-	0	-
SPT-CL J0050-5817	12.6648	-58.2841	4.00	0.25	0.88±0.03	2.03 ^{+0.32} 2.75 ^{+0.45}	38±7	0	LOS
SPT-CL J0050-5018	12.6693	-50.3017	4.49	0.50	0.160±0.009	2.75 ^{+0.45} _{-0.59}	27±5	0	-
SPT-CL J0050-5944	12.7493	-59.7402	4.03	0.25	-	-	-	0	-
SPT-CL J0051-5153	12.8530	-51.8936	6.49	0.50	0.42±0.01	3.31 ^{+0.48} _{-0.57}	60±8	0	-
SPT-CL J0051-6318	12.8557	-63.3085	4.13	0.25	-			0	-
SPT-CL J0051-5033	12.9869	-50.5516	4.18	0.25	0.90±0.03	2.05 ^{+0.33} 3.56 ^{+0.51} 3.56 ^{+0.50}	42±7	0	
SPT-CL J0052-5657	13.1433	-56.9604	6.84	0.50	0.314±0.010	3.56_0.60	52±8	0	LOS
SPT-CL J0052-5746	13.1456	-57.7698	7.05	0.50	0.256±0.009	3.71-0.62	50±7	0	-
SPT-CL J0052-5942	13.2060	-59.7116	9.88	0.25	1.13±0.06	3.44 ^{+0.42} _{-0.51}	51±5	1	-
SPT-CL J0053-5944	13.3739	-59.7374	4.13	0.25	-	- 4 00+0 33	-	0	-
SPT-CL J0054-6023 SPT-CL J0054-6104	13.6430 13.6782	-60.3881 -61.0817	5.06 4.19	0.25 0.75	1.47±0.07	1.89 ^{+0.33} _{-0.40}	37±4	1 0	-
SPT-CL J0054-6104 SPT-CL J0054-5821	13.6947	-58.3654	4.19	0.75	-	-	_	0	_
SPT-CL J0054-5731	13.7290	-57.5277	6.01	0.50	0.77±0.02	2.77 ^{+0.44} 0.50	55±8	0	_
SPT-CL J0054-5710	13.7362	-57.1737	4.09	2.75	-	-	-	0	-
SPT-CL J0055-6310	13.9758	-63.1752	4.30	0.25	0.54±0.02	2.37 ^{+0.39} _{-0.52}	46±7	0	-
SPT-CL J0057-5054	14.3376	-50.9053	4.47	1.25	0.54±0.02	2.43 ^{+0.39} _{-0.52}	43±7	0	_
SPT-CL J0058-6146	14.5858	-61.7667	16.29	0.25	0.87±0.02	5.17 ^{+0.56}	93±10	0	-
SPT-CL J0059-5105	14.7743	-51.0960	7.80	0.75	0.60±0.02	3.54+0.47	58±8	0	LOS
SPT-CL J0059-5539	14.7967	-55.6599	4.39	0.25	0.79±0.05	2.18+0.37	11±4	0	_
SPT-CL J0059-5436	14.8002	-54.6089	4.74	0.50	0.41±0.01	2.62 ^{+0.46} _{-0.56}	36±6	0	_
SPT-CL J0059-4955	14.9454	-49.9198	5.00	0.25	0.97±0.05	2.23+0.39	14±5	0	_
SPT-CL J0059-5217	14.9651	-52.2940	4.13	0.25	-	-	-	0	-
SPT-CL J0100-5359	15.0208	-53.9865	6.39	0.25	1.04±0.05	2.64 ^{+0.40} _{-0.45}	20±6	0	-
SPT-CL J0100-5021	15.0554	-50.3508	4.07	0.25	-	_	-	0	-
SPT-CL J0100-4958	15.0573	-49.9736	7.65	0.25	1.33±0.07	2.73 ^{+0.36} 2.46 ^{+0.43}	37±4	1	-
SPT-CL J0100-5910	15.2263	-59.1668	5.58	0.25	0.92±0.03	2.40_0.48	36±7	0	LOS
SPT-CL J0101-5332	15.2549	-53.5473	5.29	0.75	1.10±0.06	2.22 ^{+0.37} _{-0.45}	32±4	1	LOS
SPT-CL J0101-6047	15.2808	-60.7845	6.12	0.50	0.207±0.007	3.39 ^{+0.51} _{-0.61}	55±8	0	-
SPT-CL J0101-6007	15.3214	-60.1231	4.52	1.25	-	+0.38		0	-
SPT-CL J0101-5219	15.4187	-52.3228	4.13	0.50	0.45±0.02	2.38 ^{+0.38} -0.52	18±4	0	-
SPT-CL J0102-5028	15.7048	-50.4805	4.43	0.50	0.76±0.02	2.38 _{-0.52} 2.23 ^{+0.37} _{-0.48}	46±8	0	-
SPT-CL J0103-5242	15.8405	-52.7120	4.95	0.50	0.61±0.02	$2.52^{+0.44}_{-0.52}$	26±6	0	-
SPT-CL J0103-5516	15.9610	-55.2769	4.04	0.25		- - - - - - - - - - - - - - - - - - -	- 07.7	0	-
SPT-CL J0104-5706	16.1265	-57.1155	4.56	0.25	0.53±0.02	2.47 _{-0.53}	37±7	0	-
SPT-CL J0104-6219	16.1963	-62.3214	5.75	0.25	0.61±0.02	2.47 ^{+0.42} 2.81 ^{+0.47} 2.81 ^{+0.47} 2.11 ^{+0.35} 2.11 ^{+0.35} 2.02 ^{+0.33}	36±6	0	-
SPT-CL J0104-6226	16.2477	-62.4357	4.51	0.50	0.93±0.03	2.11_0.46	42±7	0	-
SPT-CL J0105-6257	16.3001	-62.9645	4.52	0.25	1.06±0.03	2.02 _{-0.45} 2.39 ^{+0.41} 2.39 _{-0.48}	52±9	0	-
SPT-CL J0105-6302	16.4095	-63.0373	5.33	0.25	0.92±0.03	2.39_0.48	37±7	0	-
SPT-CL J0106-5355	16.5707	-53.9277	11.63	0.25	0.48±0.01	4.80-0.68	64±8	0	-
SPT-CL J0106-5229	16.6116	-52.4875	4.67	0.25	1.52±0.08	1.76-0.39	15±3	1	-
SPT-CL J0106-5943	16.6119	-59.7195	16.02	0.50	0.348	6.09-0.81	93±10	0	-
SPT-CL J0106-5800	16.6646	-58.0071	7.09	1.25	0.280±0.008	4.80 ^{-0.48} 4.80 ^{-0.68} 1.76 ^{-0.29} 6.09 ^{-0.81} 3.69 ^{-0.80} 2.01 ^{-0.81}	64±8	0	RC
SPT-CL J0106-6233	16.7239	-62.5628	4.08	0.25	0.92±0.04	2.01+0.31	19±6	0	-
SPT-CL J0107-6324	16.8100	-63.4145	4.58	0.50	0.97±0.04	2.11 ^{+0.34} 1.90 ^{+0.42} 3.07 ^{+0.55} 3.73 ^{+0.42}	26±6	0	-
SPT-CL J0107-5242	16.8464	-52.7014	4.19	0.50	1.14±0.06	1.90 ^{+0.29} _{-0.42}	29±6	0	-
SPT-CL J0107-6332	16.8527	-63.5376	5.49	0.25	0.246±0.010	3.07 ^{+0.55} _{-0.57}	36±6	0	LOS
SPT-CL J0107-5833	16.9150	-58.5511	11.11	0.25	1.13±0.06	$3.73^{+0.42}_{-0.53}$	45±8	0	-
SPT-CL J0109-6028	17.3170	-60.4832	4.18	0.25	-	-	-	0	-
SPT-CL J0109-5132	17.3319	-51.5410	4.60	0.25	0.33±0.01	2.65 ^{+0.44} 2.23 ^{+0.36} 2.23 ^{+0.48}	34±6	0	-
SPT-CL J0109-5839	17.3347	-58.6522	4.32	0.50	0.72±0.02	2.23+0.36	40±7	0	-
SPT-CL J0109-5629	17.4039	-56.4875	5.12	0.25	0.49±0.01	2.70 ^{+0.47} 2.70 ^{+0.45} 2.84 ^{+0.45}	52±8	0	-
SPT-CL J0110-5121	17.5213	-51.3590	6.32	0.75	0.79±0.02	2.84 ^{+0.45}	64±8	0	_

Table 1 — Continued

	Table 1 — Continued									
SPT ID	R.A. (J2000)	Decl. (J2000)	Β ε	est $ heta_{ extsf{c}}$	Redshift	M_{500c} $(10^{14}h_{70}^{-1}M_{\odot})$	λ	Spitzer	Notes	
SPT-CL J0110-5857	17.5342	-58.9656	4.81	0.25	0.34±0.02	2.73 0.46	18±5	0	LOS	
SPT-CL J0110-6045	17.5847	-60.7615	5.79	1.00	0.72±0.03	2.73 ^{+0.46} 2.73 ^{+0.47} 2.73 ^{+0.48}	29±7	0	-	
SPT-CL J0110-5333	17.5874	-53.5659	4.44	0.75	0.93±0.03	2 00-0.33	29±6	0	_	
SPT-CL J0110-5622	17.6798	-56.3795	5.71	0.25	1.52±0.03	2.09 _{-0.46} 2.04 ^{+0.38}	40±6	0	LOS	
SPT-CL J0111-6336	17.7603	-63.6044	4.03	0.25	-	_	-	0	-	
SPT-CL J0111-5941	17.7612	-59.6971	4.86	0.25	0.97±0.06	2.18 ^{+0.38} _{-0.47}	18±3	1	LOS	
SPT-CL J0111-5425	17.7712	-54.4186	12.11	0.25	0.48±0.01	2.18 ^{+0.36} _{-0.47} 4.93 ^{+0.56} _{-0.68}	73±10	0	-	
SPT-CL J0111-5818	17.8279	-58.3026	7.81	0.25	0.70±0.03	$3.42^{+0.43}_{-0.54}$	24±6	0	-	
SPT-CL J0111-5519	17.8592	-55.3181	7.64	1.00	0.50±0.01	3.62 ^{+0.50} _{-0.58}	51±8	0	LOS	
SPT-CL J0111-5428 SPT-CL J0112-5942	17.9422 18.0257	-54.4753 -59.7081	4.54 5.39	0.75 0.25	- 1.20±0.07	2.18 ^{+0.39} _{-0.43}	- 34±4	0 1	-	
SPT-CL J0112-5942 SPT-CL J0112-6439	18.1680	-64.6598	4.45	0.23	1.20±0.07	2.10_0.43	J4 <u>1</u> 4	0	_	
SPT-CL J0113-5011	18.2646	-50.1954	4.11	0.25	_	-	-	ŏ	-	
SPT-CL J0113-5550	18.3560	-55.8433	4.80	0.75	0.360±0.010	2.69 ^{+0.48} _{-0.56}	61±8	0	RC,LOS	
SPT-CL J0113-6105	18.3854	-61.0885	6.69	0.75	0.44±0.01	$3.36^{+0.47}_{-0.57}$	67±8	0	-	
SPT-CL J0113-5502	18.4476	-55.0395	4.08	0.25	0.25±0.01	0 - 4+0 39	31±6	0	-	
SPT-CL J0114-5244	18.5018	-52.7395	5.95	0.75	0.43±0.01	2.54 _{-0.56} 3.09 ^{+0.48} _{-0.56}	64±8	0	-	
SPT-CL J0114-6144	18.5294	-61.7363	5.06	3.00	1.57±0.08	$1.82^{+0.32}_{-0.39}$	28±4	1	-	
SPT-CL J0115-5730	18.7995	-57.5107	4.40	0.25	0.20±0.01	2.67 ^{+0.45} _{-0.58}	16±4	0	-	
SPT-CL J0115-6516	18.8105	-65.2826	5.01	0.25	0.68±0.04	2.49 ^{+0.45} _{-0.51}	14±5	0	-	
SPT-CL J0115-5517	18.9960	-55.2845	4.08	0.75	0.49±0.02	2.34+0.37	26±6	0	-	
SPT-CL J0116-5039	19.0948	-50.6624	5.86	0.25	1.60±0.08	2.03+0.36	22±4	1	LL	
SPT-CL J0116-5045	19.1481	-50.7652	5.47	0.75	0.40±0.01	2.92 ^{+0.48} _{-0.56}	56±8	0	-	
SPT-CL J0117-5053	19.2821	-50.8946	7.15	1.75	0.205±0.007	2.92 _{-0.56} 3.80 ^{+0.53} -0.62 2.66 ^{+0.43} 2.66 _{-0.56}	63±8	0	RC	
SPT-CL J0117-5050	19.2897	-50.8481	4.36	0.25	0.206±0.009	2.66 ^{+0.43} _{-0.56}	35±6	0	-	
SPT-CL J0117-6032	19.3721	-60.5406	11.26	0.50	1.04±0.03	3.89+0.46	55±9	0	-	
SPT-CL J0118-5637	19.5198	-56.6332	6.75	1.00	0.205±0.007	3.64 ^{+0.51} _{-0.61}	59±8	0	-	
SPT-CL J0118-5156	19.5989	-51.9418	11.51	0.50	0.705	4.42 ^{+0.51} _{-0.61}	76±9	0	-	
SPT-CL J0118-5402	19.6647	-54.0450	5.27	1.00	0.236±0.008	2.99+0.49	51±7	0	-	
SPT-CL J0118-5826	19.7030	-58.4391	4.16	0.25	0.27±0.01	2.54+0.41	38±6	0	LOS	
SPT-CL J0118-6520	19.7399	-65.3345	9.54	0.50	0.083±0.008	4.79 ^{+0.56}	19±6	0	-	
SPT-CL J0118-5644	19.7483	-56.7487	4.65	0.25	1.47±0.07	1.79 +0.30	10±3	1	-	
SPT-CL J0119-5750	19.8480	-57.8488	4.92	0.25	0.95±0.03	1.79 _{-0.39} 2.23 ^{+0.38} 2.23 _{-0.48} 3.29 ^{+0.44}	40±7	0	-	
SPT-CL J0119-5919	19.9157	-59.3281	6.95	0.50	0.59±0.02		43±7	0	-	
SPT-CL J0119-5838	19.9825	-58.6462	5.19	0.25	0.56±0.02	2.65 ^{+0.47} _{-0.53}	20±5	0	-	
SPT-CL J0120-5435 SPT-CL J0120-6522	20.0146 20.0564	-54.5869 -65.3780	4.04 4.12	1.00 0.25	- 0.54±0.03	2.31 ^{+0.37}	- 15±4	0 0	-	
SPT-CL J0120-6522 SPT-CL J0121-5432	20.0304	-54.5366	4.12	0.25	0.54±0.05	2.31-0.51	13±4	0	_	
SPT-CL J0121-5547	20.3367	-55.7882	4.62	0.25	1.09±0.04	2.02 ^{+0.34} _{-0.44}	31±7	0	LOS	
SPT-CL J0121-5121	20.3561	-51.3578	4.14	0.50	0.24±0.01	2.56+0.41	23±5	0	_	
SPT-CL J0121-5458	20.3704	-54.9673	5.73	0.50	0.83±0.02	2.56 ^{+0.41} 2.60 ^{+0.43} 2.60 ^{+0.43}	58±8	0	LOS	
SPT-CL J0121-5804	20.4650	-58.0800	4.05	0.25	-	_	-	0	-	
SPT-CL J0122-5801	20.5492	-58.0218	8.98	0.25	1.60±0.08	2.79 ^{+0.36} 1.79 ^{+0.29} 1.79 ^{+0.29} 2.79	25±3	1	LOS,LL	
SPT-CL J0122-5241	20.5959	-52.6896	4.23	0.25	1.29±0.07	1.79 ^{+0.29} _{-0.40}	19±5	0	-	
SPT-CL J0122-5554	20.6445	-55.9067	4.36	0.50	0.69±0.03	2.26 ^{+0.37} _{-0.49}	25±6	0	LOS	
SPT-CL J0123-5112	20.8847	-51.2131	4.17	1.75		_		0	-	
SPT-CL J0123-6159	20.8886	-61.9975	8.53	0.25	0.87±0.02	3.44 ^{+0.43} 4.15 ^{+0.53} 4.15 ^{+0.63}	57±8	0	-	
SPT-CL J0124-5937	21.1861	-59.6276	8.13	0.75	0.204±0.006	4.15-0.64	76±9	0	-	
SPT-CL J0125-5637	21.2960	-56.6279	5.30	0.75	1.47±0.07	1.97-0.40	9±3	1	_	
SPT-CL J0126-6143	21.7079	-61.7230	4.34	0.50	0.78±0.03	1.97 ^{+0.34} 2.19 ^{+0.35} 2.19 ^{+0.36} 2.41 ^{+0.36}	21±6	0	-	
SPT-CL J0128-6506	22.0118	-65.1107	6.27	0.25	1.28±0.07	2.41_0.42	25±4	1	-	
SPT-CL J0128-5714	22.0502	-57.2363	4.08	1.00	0.42±0.02	2.39 ^{+0.32} _{-0.52} 2.27 ^{+0.36}	17±5	0	-	
SPT-CL J0128-5222 SPT-CL J0128-5058	22.0507 22.2053	-52.3827 -50.9773	6.12 4.80	0.25 2.00	1.37±0.07	2.27 ^{+0.36} _{-0.41}	28±4	1 0	_	
SPT-CL J0126-5056 SPT-CL J0129-5329	22.2053	-50.9773 -53.4900	4.80	0.50	-		-	0	_	
SPT-CL J0129-6442	22.3702	-64.7091	4.15	0.25	-	0.70	-	0	-	
SPT-CL J0129-6432	22.4317	-64.5462	21.42	0.50	0.326	7.26 ^{+0.78} _{-0.94}	118±11	0	-	
SPT-CL J0130-6357	22.6164	-63.9585	4.14	0.25	0.51±0.02	2.34+0.39	47±7	0	LOS	
SPT-CL J0130-6314	22.6190	-63.2355	6.01	0.25	0.77±0.02	2.75 ^{+0.45}	45±8	0	-	
SPT-CL J0131-5849	22.8516	-58.8206	4.76	0.50	0.48±0.01	2.57 ^{+0.46} _{-0.54} 2.47 ^{+0.41} _{-0.50}	67±8	0	-	
SPT-CL J0131-6248	22.8519	-62.8015	4.98	0.25	0.69±0.03	2.47 ^{+0.41} _{-0.50}	23±6	0	LOS	

Table 1 — Continued

			Iak	ne i –	- Conunuea					
SPT ID	R.A. (J2000)	Decl. (J2000)	Β ε	est $\theta_{ extsf{c}}$	Redshift	M_{500c} $(10^{14}h_{70}^{-1}M_{\odot})$	λ	Spitzer	Notes	
SPT-CL J0131-5921	22.8565	-59.3589	10.72	0.25	0.99±0.06	3.81 ^{+0.44} _{-0.55}	51±7	0	-	
SPT-CL J0131-5604	22.9319	-56.0802	11.68	0.25	0.80±0.02	4.30 ^{+0.51} _{-0.60}	86±10	0	_	
SPT-CL J0132-5535	23.0601	-55.5896	6.07	0.25	0.67±0.02	2 00+0.46	61±8	0	_	
SPT-CL J0132-5712	23.1202	-57.2154	5.24	0.25	1.21±0.05	2 11+0.38	18±6	0	_	
SPT-CL J0132-5312	23.2140	-53.2039	4.33	0.25	-	0.44	-	Ö	-	
SPT-CL J0133-5945	23.2542	-59.7631	4.09	0.75	-		-	0	-	
SPT-CL J0133-5821	23.2738	-58.3511	4.93	1.00	0.199±0.008	2.89 ^{+0.50} _{-0.59}	40±7	0	-	
SPT-CL J0133-6417	23.4020	-64.2980	4.21	0.50	-	5.57 ^{+0.62}	-	0	-	
SPT-CL J0133-6434	23.4076	-64.5699	13.68	1.00	0.319±0.007	5.57 _{-0.77} 2.22 _{-0.47}	113±11	0	-	
SPT-CL J0134-5318	23.7405	-53.3138	4.95	0.25	0.96±0.03		31±6	0	-	
SPT-CL J0135-5902	23.7892	-59.0340	7.40	0.25	0.51±0.01	3.52 ^{+0.48} _{-0.57}	57±8	0	-	
SPT-CL J0135-6139	23.8714	-61.6504	4.02	0.25	0.5010.04	2.04+0.48	-	0	-	
SPT-CL J0135-5904	23.9792	-59.0810	8.66	0.75	0.50±0.01	3.94 ^{+0.48} _{-0.59}	99±11	0	LOS	
SPT-CL J0136-6226	24.0412	-62.4467	6.02	0.50	0.89±0.03	2.65 ^{+0.42}	26±6	0	LOS	
SPT-CL J0137-6105	24.3434	-61.0908	6.03	0.25	0.42±0.01	3.13 ^{+0.48} _{-0.57}	45±7	0	-	
SPT-CL J0137-6044	24.4366	-60.7464	4.25	0.25	4 20 10 07	4 00+0.29	25.5	0	-	
SPT-CL J0138-6340 SPT-CL J0139-6135	24.6007 24.7755	-63.6745 -61.5980	4.26 4.56	0.25 0.50	1.29±0.07	1.80 ^{+0.29} _{-0.40}	25±5	0 0	-	
SPT-CL J0139-0133 SPT-CL J0139-5749	24.7733	-57.8267	5.04	0.30	0.76±0.02	2.44 ^{+0.41} _{-0.50}	49±8	0	_	
SPT-CL J0139 5749 SPT-CL J0139 5204	24.8873	-52.0786	9.37	0.25	0.70±0.02 0.93±0.02	2.44 _{-0.50} 3.56 ^{+0.44}	61±8	0	_	
SPT-CL J0139-5804	24.8881	-58.0673	4.80	0.25	0.93±0.02 0.77±0.03	3.56 ^{+0.44} _{-0.53} 2.33 ^{+0.42} _{-0.49}		0	-	
SPT-CL J0139-5604 SPT-CL J0140-4957	25.0294	-49.9538	4.20	0.23	0.77±0.03	2.33_0.49	33±6	0	_	
SPT-CL J0140 4937 SPT-CL J0140 5603	25.1185	-56.0547	4.20	0.50	_	_	_	0	_	
SPT-CL J0140-5718	25.1340	-57.3155	4.04	0.25	-	-	-	0	-	
SPT-CL J0140-6234	25.2163	-62.5690	4.04	0.25	1.06±0.05	1.91 ^{+0.30}	20±6	0	-	
SPT-CL J0140-5453	25.2227	-54.8877	4.02	0.75	-	-	-	0	-	
SPT-CL J0140-6129	25.2469	-61.4852	4.16	0.25	-	- a+0.61	-	0	-	
SPT-CL J0142-5032	25.5442	-50.5386	15.94	0.25	0.679	5.43 ^{+0.61} _{-0.72}	105±10	0	-	
SPT-CL J0142-5441	25.5801	-54.6979	4.13	0.25	0.20010.007	2.00+0.53	-	0	-	
SPT-CL J0143-5828	25.7887	-58.4755	5.40	0.50	0.200±0.007	3.09 ^{+0.53} 4.00 ^{+0.29}	56±8	0	-	
SPT-CL J0144-5035 SPT-CL J0144-5018	26.0838 26.1727	-50.5912 -50.3070	4.01 4.22	1.50 1.75	1.07±0.06	1.90-0.41	17±5	0 0	-	
SPT-CL J0144-5815	26.1727	-58.2593	4.53	1.75	0.84±0.05	2.18 ^{+0.37} _{-0.48}	10±4	0	_	
SPT-CL J0145-5301	26.2552	-53.0259	12.21	1.50	0.04±0.03	5.53 ^{+0.65}	52±7	0	_	
SPT-CL J0145-5627	26.2753	-56.4506	4.74	0.25	0.117 0.210±0.010	10.50	30±6	0	-	
					1	7 F0+0.80			_	
SPT-CL J0145-6033	26.3021	-60.5587	21.08	0.50	0.179	2.16 ^{+0.36} _{-0.47}	89±10	0	-	
SPT-CL J0145-5643 SPT-CL J0145-6044	26.3099 26.3217	-56.7209 -60.7346	4.57 4.21	0.25 1.00	0.89±0.06	2.16_0.47	9±5	0 0	-	
SPT-CL J0145-5641	26.3532	-56.6947	4.06	0.25	_	_	_	0	_	
SPT-CL J0145-5926	26.4847	-59.4435	4.44	0.25	0.52±0.02	2.43 ^{+0.41} _{-0.52}	25±5	0	_	
SPT-CL J0146-5956	26.5663	-59.9490	4.59	0.25	1.60±0.08	1.69 ^{+0.28} _{-0.37}		0	LL	
SPT-CL J0146-5002	26.5737	-50.0418	4.09	0.75	-	l -	-	Ö	-	
SPT-CL J0146-5320	26.6424	-53.3412	4.21	0.25	0.80±0.02	$2.14^{+0.34}_{-0.47} \\ 3.51^{+0.44}_{-0.53} \\ 2.20^{+0.38}_{-0.47}$	50±8	0	-	
SPT-CL J0146-6126	26.6588	-61.4365	9.17	0.25	0.93±0.03	3.51+0.44	49±8	0	_	
SPT-CL J0147-6330	26.9083	-63.5155	4.80	0.25	0.93±0.03	2.20+0.38	28±6	0	LOS	
SPT-CL J0147-5017	26.9284	-50.2965	4.33	0.75	_	-	-	0	-	
SPT-CL J0147-5150	26.9521	-51.8358	8.58	0.25	0.79±0.03	3.53 ^{+0.44} _{-0.54} 4.00 ^{+0.48} _{-0.58}	32±6	0	-	
SPT-CL J0147-5622	26.9634	-56.3758	9.63	0.50	0.66±0.01	4.00+0.48	83±10	0	LOS	
SPT-CL J0148-6525	27.0799	-65.4230	4.33	1.00	-	_	-	0	-	
SPT-CL J0148-5034	27.2160	-50.5780	4.00	0.25	1.20±0.07	1.80 ^{+0.27} _{-0.39}	13±3	1	-	
SPT-CL J0148-5500	27.2379	-55.0059	4.97	0.25	0.93±0.03	2.25+0.40	52±8	0	-	
SPT-CL J0148-6124	27.2474	-61.4148	4.08	0.75	0.92±0.05	1.80 ^{+0.27} -0.39 2.25 ^{+0.40} 1.99 ^{+0.31} 2.36 ^{+0.40} 2.36 ^{+0.40} 2.36-0.51	14±4	0	-	
SPT-CL J0149-5947	27.2665	-59.7894	4.58	0.25	0.65±0.02	2.36+0.40	30±5	0	_	
SPT-CL J0149-5127	27.3409	-51.4503	4.18	0.25	0.98±0.04	1.98+0.31	23±6	0	_	
SPT-CL J0149-6311	27.3928	-63.1919	4.04	0.75	_		-	Ö	-	
SPT-CL J0149-5840	27.3979	-58.6792	5.31	0.25	0.41±0.01	2.86+0.48	29±6	0	-	
SPT-CL J0149-5608	27.4563	-56.1483	4.88	0.50	0.95±0.03	2.21+0.40	32±6	0	LOS	
SPT-CL J0150-5224	27.5406	-52.4142	4.43	0.50	0.64±0.02	2.33+0.39	41±6	0	-	
SPT-CL J0150-5003	27.6380	-50.0563	4.77	0.25	0.82±0.02	2.29+0.41	53±8	0	_	
SPT-CL J0150-5215	27.6833	-52.2602	4.34	0.50	0.97±0.06	2.03+0.34	14±4	0	_	
SPT-CL J0150-5215 SPT-CL J0151-5654	27.7780	-56.9147	8.08	0.50	0.97±0.00 0.293±0.008	2.86 ^{+0.48} _{-0.56} 2.21 ^{+0.40} _{-0.47} 2.33 ^{+0.39} _{-0.50} 2.29 ^{+0.41} _{-0.48} 2.03 ^{+0.45} _{-0.45} 4.02 ^{+0.52} _{-0.63}	67±8	0	_	
SPT-CL J0151-6526	27.7780	-65.4403	4.40	0.75	-	-	-	0	_	
SPT-CL J0151-5848	27.8448	-58.8057	6.24	0.75	0.78±0.03	2.84 ^{+0.44} _{-0.50}	31±7	0	LOS	
	_,.5++0	33.3301	J.∠¬	3.20	3 0_0.00					

Table 1 — Continued

Table 1—Continued										
SPT ID	R.A. (J2000)	Decl. (J2000)	Β ε	est $\theta_{ extsf{c}}$	Redshift	M_{500c} $(10^{14}h_{70}^{-1}M_{\odot})$	λ	Spitzer	Notes	
SPT-CL J0151-5332	27.8514	-53.5388	4.67	0.25	1.60±0.08	1.70+0.28	23±3	1	LL	
SPT-CL J0151-5954	27.8559	-59.9075	13.71	0.25	1.07±0.02	1.70 ^{+0.28} 4.34 ^{+0.47} 4.35	100±11	0	-	
SPT-CL J0151-6145	27.8957	-61.7535	4.16	0.25	-	-0.56	-	0	-	
SPT-CL J0152-5133	28.1295	-51.5510	4.08	0.75	-	-	-	0	-	
SPT-CL J0152-5829	28.2060	-58.4845	5.90	0.25	0.73±0.02	2.77 ^{+0.45}	57±8	0	-	
SPT-CL J0152-6437	28.2378	-64.6272	4.01	0.25	0.39±0.01	2.40+0.38	40±6	0	RC	
SPT-CL J0152-5303	28.2379	-53.0567	12.60	0.50	0.58±0.01	4.88 ^{+0.54} _{-0.67} 2.08 ^{+0.35} _{-0.44}	107±11	0	-	
SPT-CL J0153-6059	28.2670	-60.9929	4.68	0.25	1.06±0.06	2.08 ^{+0.35} _{-0.44} 2.03 ^{+0.35} _{-0.45}	22±5	0	-	
SPT-CL J0153-5939	28.3719	-59.6523	4.69	0.25	1.12±0.06	2.03-0.45	39±7	0	-	
SPT-CL J0153-5827	28.4229	-58.4541	4.13	0.25	0.67±0.02	2.21 ^{+0.36} _{-0.49}	29±6	0	-	
SPT-CL J0153-6525	28.4928	-65.4211	5.55	0.25	-	0.00+0.34	-	0	-	
SPT-CL J0154-5843 SPT-CL J0154-6504	28.5128 28.5660	-58.7193 -65.0788	4.12 4.44	0.50 0.25	0.67±0.02	2.20 ^{+0.34} _{-0.48}	38±7	0 0	-	
SPT-CL J0154-5619	28.6214	-56.3236	7.86	0.25	0.79±0.02	3 34+0.44	51±8	0	_	
SPT-CL J0154-6409	28.6916	-64.1539	6.70	0.25	1.07±0.06	3.34 ^{+0.44} _{-0.52} 2.70 ^{+0.39} _{-0.46}	41±4	1	_	
SPT-CL J0154-5201	28.6989	-52.0286	4.05	0.25	-	2.70_0.46		Ö	_	
SPT-CL J0155-6105	28.7878	-61.0845	4.12	0.25	-	-	-	Ö	-	
SPT-CL J0155-5321	28.9014	-53.3514	4.29	0.50	0.34±0.01	2.52 ^{+0.42} _{-0.55}	39±8	0	-	
SPT-CL J0155-5218	28.9896	-52.3088	5.55	0.50	0.70±0.02		63±8	0	-	
SPT-CL J0156-5541	29.0426	-55.6992	18.69	0.25	1.288	2.67 ^{+0.43} _{-0.51} 4.87 ^{+0.52} _{-0.63}	50±0	0	RC	
SPT-CL J0156-5112	29.0510	-51.2090	4.74	0.75	0.65±0.02	2.40 ^{+0.42} _{-0.52}	31±6	0	-	
SPT-CL J0156-6224	29.0598	-62.4107	4.02	0.25	-	_	-	0	-	
SPT-CL J0156-5424	29.0937	-54.4062	4.53	0.50	0.41±0.01	2.55 ^{+0.44} _{-0.55}	44±7	0	LOS	
SPT-CL J0156-5003	29.1486	-50.0538	4.03	0.25	-	58		0	-	
SPT-CL J0157-5820	29.3060	-58.3476	9.81	0.50	0.217±0.006	4.67 ^{+0.58} _{-0.68} 3.00 ^{+0.50} _{-0.57}	69±8	0	-	
SPT-CL J0157-6117	29.3653	-61.2964	5.53	0.75	0.35±0.01	3.00-0.57	51±7	0	-	
SPT-CL J0157-6425	29.4117	-64.4229	4.35	0.50	0.073±0.006	2.76 ^{+0.45} _{-0.60}	33±6	0	-	
SPT-CL J0157-6442	29.4742	-64.7037	8.89	0.25	0.91±0.03	$3.47^{+0.46}_{-0.52}$	26±6	0	-	
SPT-CL J0157-5917 SPT-CL J0158-5505	29.4835 29.5343	-59.2867 -55.0964	4.27 4.10	0.25 1.25	-	-	-	0 0	-	
SPT-CL J0158-5438	29.5691	-54.6350	4.40	1.50	_	_	_	0	-	
SPT-CL J0158-5705	29.7120	-57.0980	4.19	1.00	-	-	-	0	-	
SPT-CL J0159-5947	29.7881	-59.7922	4.09	0.25	-	- 2 27+0.41	-	0	-	
SPT-CL J0159-6437	29.8483	-64.6238	4.83	0.25	0.73±0.02	2.37 ^{+0.41} _{-0.50}	39±7	0	-	
SPT-CL J2201-6105	330.3259	-61.0908	4.08	0.75	1 60 10 00	1 77+0.31	- 0.2	0 1	-	
SPT-CL J2201-5620	330.3831	-56.3441 50.0405	4.89	2.00	1.60±0.08	1.77-0.39	9±2		LL	
SPT-CL J2201-5956 SPT-CL J2202-5524	330.4675 330.5098	-59.9495 -55.4102	26.11 4.22	1.00 0.25	0.097	8.69 ^{+0.92} _{-1.12}	142±13	0 0	_	
SPT-CL J2202 5324 SPT-CL J2202-5936	330.5514	-59.6020	5.67	0.25	0.44±0.01	2.97+0.49	45±8	0	_	
SPT-CL J2202-5603	330.6143	-56.0649	4.06	0.25	1.14±0.06	1 97+0.28	19±5	0	_	
SPT-CL J2202 5005	330.6872	-63.7758	4.56	0.50	-	1.07 -0.41	-	0	_	
SPT-CL J2202-6308	330.7357	-63.1481	4.01	0.25	-	-	-	Ö	-	
SPT-CL J2203-6055	330.7531	-60.9186	5.53	0.25	0.413±0.009	2.93 ^{+0.49} _{-0.57}	83±9	0	-	
SPT-CL J2203-5735	330.7619	-57.5977	4.62	0.25	0.95±0.03	2.13 ^{+0.37} _{-0.47}	53±8	0	-	
SPT-CL J2203-5047	330.8042	-50.7861	10.68	0.25	0.94±0.02	2.93 ^{+0.49} _{-0.57} 2.13 ^{+0.37} 3.88 ^{+0.46} _{-0.55}	56±9	0	-	
SPT-CL J2203-5641	330.8305	-56.6834	4.05	0.50	-	-	-	0	-	
SPT-CL J2203-5042	330.9236	-50.7044	4.04	0.25	-	- 0.04+0.31	-	0	-	
SPT-CL J2204-5300	331.1467	-53.0118	4.03	0.50	0.87±0.03	2.04 ^{+0.31} 2.73 ^{+0.44} 2.73 ^{+0.44}	31±6	0	-	
SPT-CL J2204-5059	331.1949	-50.9870	4.28	0.25	0.084±0.008	2.73_0.58	19±5	0	-	
SPT-CL J2204-5340	331.2069	-53.6722	6.33	0.25	0.93±0.03	2.69 ^{+0.45} _{-0.47}	34±7	0	LOS	
SPT-CL J2204-6223	331.2076	-62.3834	6.34	0.50	0.279±0.009	3.41 ^{+0.50} -0.60	56±8	0	-	
SPT-CL J2205-5927	331.2582	-59.4525	11.47	0.50	0.349±0.008	4.96 ^{+0.58} 4.96 ^{+0.69} 2.50 ^{+0.41} 2.24 ^{+0.38}	84±9	0	LOS	
SPT-CL J2205-5705	331.3290	-57.0884	4.05	1.25	0.28±0.01	2.50 -0.55	40±6	0	-	
SPT-CL J2205-6310	331.3809	-63.1708	4.66	0.25	0.83±0.04	2.24+0.38	18±5	0	-	
SPT-CL J2205-5142	331.4881	-51.7090	4.17	0.25		-	- 20:7	0	-	
SPT-CL J2206-5528	331.5163	-55.4724	5.00	0.25	0.57±0.02	2.56 ^{+0.46} _{-0.53} 2.47 ^{+0.43}	36±7	0	-	
SPT-CL J2206-5820	331.6166	-58.3429	5.00	0.25	0.69±0.04	L 2.71_0.51	12±4	0	-	
SPT-CL J2206-5808	331.6486	-58.1347	11.65	0.75	0.59±0.01	4.63 ^{+0.52} -0.65 2.44 ^{+0.39} -0.44	90±10	0	LOS	
SPT-CL J2206-6407	331.6509	-64.1270	6.00	0.25	1.13±0.06	2.44_0.44	26±6	0	-	
SPT-CL J2206-5919	331.6748	-59.3248	5.60	0.25	0.77±0.02	2.62-0.50	48±8	0	-	
SPT-CL J2206-5558	331.7366	-55.9730	4.52	0.25	0.40±0.02	2.62 ^{+0.44} 2.62 ^{+0.44} 2.57 ^{+0.43} 1.96 ^{+0.32}	15±4	0	-	
SPT-CL J2207-5105	331.8744	-51.0958	4.28	0.25	1.04±0.06	1.96-0.44	11±6	0	-	
SPT-CL J2207-5522	331.8976	-55.3685	5.48	0.50	0.87±0.03	2.49 ^{+0.42} _{-0.48}	26±6	0	-	

Table 1 — Continued

Table 1 — Conunacu									
SPT ID	R.A. (J2000)	Decl. (J2000)	Β ε	est $\theta_{ extsf{c}}$	Redshift	M_{500c} $(10^{14}h_{70}^{-1}M_{\odot})$	λ	Spitzer	Notes
SPT-CL J2207-5056	331.9763	-50.9445	5.22	0.25	0.38±0.01	2.84 ^{+0.49} _{-0.56}	37±6	0	_
SPT-CL J2207-5136	331.9937	-51.6078	4.06	0.25		_	-	0	-
SPT-CL J2209-6350	332.2544	-63.8423	4.11	1.25	0.47±0.02	2.37 ^{+0.38} _{-0.52}	31±6	0	-
SPT-CL J2209-5807	332.3182	-58.1237	4.88	0.25	1.04±0.06	2.14 ^{+0.38} 2.14 ^{+0.46} 2.58 ^{+0.42} -0.56	10±3	1	-
SPT-CL J2209-5940	332.3285	-59.6756	4.26	0.25	0.264±0.009	2.58 _{-0.56}	48±7	0	-
SPT-CL J2209-5148	332.3422	-51.8033	7.69	1.25	0.110	4.11-0.67	41±7	0	-
SPT-CL J2209-5622	332.3739	-56.3690	4.05	0.25	1.10±0.06	1.07 _{-0.41}	29±6	0	-
SPT-CL J2209-5801 SPT-CL J2209-5002	332.4042	-58.0305 -50.0455	4.04	0.75	1.04±0.05	1.92 ^{+0.29} -0.42	17±6	0	-
SPT-CL J2209-5002 SPT-CL J2210-6347	332.4509 332.5159	-50.0455 -63.7939	4.23 4.00	0.50 0.25	1.44±0.07	1.71 ^{+0.26} -	22±6 -	0 0	_
SPT-CL J2210-6500	332.5375	-65.0160	6.04	0.25	1.52±0.08	2.14 ^{+0.36} -0.40 3.81 ^{+0.52}	45±4	1	LOS
SPT-CL J2210-5709	332.6261	-57.1640	7.52	0.50	0.301±0.008	3.81+0.52	71±9	0	_
SPT-CL J2210-6046	332.7304	-60.7747	4.18	0.25	0.49±0.03	2 26+0.38	10±4	0	_
SPT-CL J2212-6249	333.1006	-62.8198	5.12	0.50	0.84±0.04	2 37,0.45	20±5	0	_
SPT-CL J2212-5010	333.1814	-50.1772	4.28	0.25	0.78±0.04	2 17+0.36	14±5	0	_
SPT-CL J2212-5228	333.2036	-52.4717	4.68	0.75	0.58±0.03	2 45+0.43	12±4	0	_
SPT-CL J2212-5335	333.2299	-53.5938	4.60	1.25	0.205±0.010	2 75+0.48	30±6	0	_
SPT-CL J2212-5434	333.2302	-54.5769	4.22	0.50	0.67±0.02	2 24+0.37	53±8	0	_
SPT-CL J2212-5616	333.2488	-56.2754	4.65	2.00	-	2.24 _{-0.49} -	-	Ö	_
SPT-CL J2213-5024	333.4050	-50.4051	4.55	0.50	-	-	-	0	-
SPT-CL J2213-5150	333.4391	-51.8407	6.78	0.25	0.65±0.02	3.16 ^{+0.43} _{-0.53}	54±8	0	-
SPT-CL J2214-5501	333.5288	-55.0293	5.17	0.25	1.24±0.07	2.07 ^{+0.36}	49±4	1	-
SPT-CL J2215-5204	333.7659	-52.0807	5.35	0.75	0.51±0.01	2.77 ^{+0.48} _{-0.54}	54±8	0	-
SPT-CL J2215-5252	333.8019	-52.8702	4.03	0.25	-	_	-	0	-
SPT-CL J2215-5119	333.9097	-51.3244	4.54	1.00	0.49±0.02	2.49 ^{+0.42}	30±6	0	-
SPT-CL J2216-5705	334.0237	-57.0925	4.38	0.25	0.65±0.02	2.30 ^{+0.38} _{-0.50} 2.05 ^{+0.34} _{-0.45}	58±8	0	-
SPT-CL J2216-6421	334.1965	-64.3610	4.47	0.50	1.00±0.06	2.05-0.45	13±6	0	-
SPT-CL J2216-6422 SPT-CL J2217-5431	334.2234 334.3092	-64.3709 -54.5240	4.26 6.22	0.25 0.25	- 1.06±0.03	2.57 ^{+0.40}	- 68±10	0 0	-
SPT-CL J2217-5431	334.4391	-55.4083	6.00	0.25	1.00±0.03 1.02±0.06	2 55+0.39	35±4	1	-
	334.4880	-65.1749	9.46	1.25	0.095	4.74 ^{+0.59}		0	-
SPT-CL J2217-6510 SPT-CL J2217-5652	334.4942	-56.8718	4.25	2.25	0.095	4.74_0.70	66±9	0	_
SPT-CL J2218-5533	334.5298	-55.5525	8.42	0.25	0.98±0.03	3.28 ^{+0.41} 1.03 ^{+0.35}	39±8	0	_
SPT-CL J2218-5008	334.5761	-50.1496	5.35	0.25	1.52±0.08	1.93 ^{+0.35} _{-0.40}	18±3	1	_
SPT-CL J2218-5232	334.5780	-52.5424	4.40	0.25	-		-	0	_
SPT-CL J2218-5402	334.6165	-54.0460	4.29	0.75	-	-	-	0	-
SPT-CL J2218-5405	334.7446	-54.0929	4.69	0.25	1.28±0.07	1.90 ^{+0.33}	39±4	1	-
SPT-CL J2219-5239	334.8425	-52.6662	4.33	1.50	-	1.96 ^{+0.31}	-	0	-
SPT-CL J2219-6005	334.8464	-60.0984	4.12	1.00	1.00±0.04	1.96 ^{+0.43} 5.57 ^{+0.61} 5.57 _{-0.76}	26±10	0	LOS
SPT-CL J2219-5708	334.9564	-57.1431	13.46	0.50	0.296±0.007	5.57-0.76	106±10	0	-
SPT-CL J2219-5816	334.9954	-58.2698	4.84	0.25	0.29±0.01	2.77 ^{+0.46}	31±6	0	-
SPT-CL J2220-5228	335.0381	-52.4786	4.77	1.25	0.092±0.005	2.91-0.47	55±8	0	-
SPT-CL J2220-5159	335.1961	-51.9931	6.94	0.25	0.49±0.02	2.77 _{-0.57} 2.91 ^{-0.49} 3.39 ^{-0.47} 2.62 ^{-0.56} 2.62 ^{-0.56} 1.77 ^{-0.31} 2.53 ^{-0.39} 1.80 ^{-0.39} 4.36 ^{-0.54}	39±7	0	-
SPT-CL J2221-5631	335.4261	-56.5322	4.91	0.25	0.48±0.02	2.62-0.56	31±6	0	
SPT-CL J2221-5345	335.4376	-53.7550	4.90	0.25	1.60±0.08	1.77-0.31	12±3	1	LL
SPT-CL J2222-6106	335.5123	-61.1092	4.18	0.50	0.30±0.01	2.53-0.55	37±6	0	-
SPT-CL J2222-5105	335.7226	-51.0882	5.02	0.25	1.57±0.08	1.80 0.32	22±3	1	-
SPT-CL J2223-5228	335.8490	-52.4684	9.04	1.25	0.265±0.007	4.36-0.66	85±9	0	-
SPT-CL J2223-5604 SPT-CL J2224-5039	335.8969 336.0727	-56.0719 -50.6509	4.33 4.24	0.25 0.75	- 0.33±0.01	_	- 21±6	0 0	-
SPT-CL J2224-5039 SPT-CL J2224-5420	336.1979	-50.6509 -54.3351	4.24	1.00	0.55±0.01	2.52 ^{+0.42} -0.55	31±6	0	_
SPT-CL J2225-6526	336.2697	-65.4345	10.09	0.75	0.66±0.01	4.11 ^{+0.51} _{-0.60}	91±10	0	_
SPT-CL J2225-6120	336.3334	-61.3421	4.62	0.50	-	_	-	Ö	-
SPT-CL J2225-6131	336.4801	-61.5171	4.43	0.75	0.175±0.009	2.71 ^{+0.45} 2.61 ^{+0.42} 2.61 ^{+0.42}	28±5	0	-
SPT-CL J2226-5714	336.5119	-57.2414	4.01	1.50	0.128±0.007	2.61 ^{+0.42} _{-0.56}	33±6	0	-
SPT-CL J2226-5214	336.6484	-52.2420	4.18	0.75	-	_	-	0	-
SPT-CL J2227-6220	336.9977	-62.3345	7.37	0.25	0.63±0.02	$3.37^{+0.47}_{-0.55}$	64±8	0	-
SPT-CL J2228-6513	337.1396	-65.2240	5.78	0.25	0.59±0.02	2.85 ^{+0.48} _{-0.52}	35±6	0	LOS
SPT-CL J2228-5158	337.1944	-51.9822	4.14	0.75	0.65±0.02	2.24 ^{+0.36} _{-0.49}	29±6	0	-
SPT-CL J2228-5828	337.2109	-58.4731	9.92	1.00	0.74±0.02	3.37 ^{+0.47} _{-0.55} 2.85 ^{+0.48} _{-0.49} 2.24 ^{+0.36} _{-0.49} 3.95 ^{+0.50} _{-0.58}	86±10	0	-
SPT-CL J2230-6300	337.5477	-63.0111	5.04	0.25	1.10±0.06	2.15 ^{+0.38} _{-0.44}	34±4	1	_

Table 1 — Continued

SPT-LD_12230-5048 337.5656 -56.8067 4.11 0.75 0.58.00.2 4.47.08 3.416 0		Table 1 — Continued								
SPT-CL, J2230-6340 337,6880 -68,6800 10,95 25 0.6840,02 4,34,492 34,468 0 LOS SPT-CL, J2230-5834 337,6822 -68,6765 4,18 10,00 1,2640,07 182,039 29,66 0 - SPT-CL, J2230-5399 337,7122 -58,6678 4,18 0,50 - - 0 - 0 - SPT-CL, J2230-5399 338,1419 -99,9972 17,15 0.25 0.595 5,866,93 122,311 0 - SPT-CL, J2232-5599 338,1419 -99,9972 17,15 0.25 0.595 5,866,93 122,311 0 - SPT-CL, J2233-5391 338,154 -49,896 162 0.50 0.532,002 2,52,444 4356 0 - SPT-CL, J2233-5894 338,3154 -49,8862 16,20 0.25 0.522,023 482,7 0 - SPT-CL, J2233-5894 338,3767 78,082,23 4,10 0.0 0.440 0 2.52,433 <	SPT ID					Redshift	M_{500c} $(10^{14}h_{70}^{-1}M_{\odot})$	λ	Spitzer	Notes
SPT-CL J2230-5829 337.6775 -52.6575	SPT-CL J2230-5048	337.5504	-50.8037	4.11	0.75	-	-	-	0	-
SPT-CL J2230-5829 337.6775 -52.6575	SPT-CL J2230-6340	337.6580	-63.6806	10.95	0.25	0.65±0.02	4.34 ^{+0.52} _{-0.63}	34±6	0	LOS
SPT-CL J2230-5826 337.6930	SPT-CL J2230-5239	337.6775	-52.6575	6.13	0.75	0.58±0.02	3 01+0.45	49±8	0	-
SPT-CL J2230-5339 337.7122 53.6578 4.59 0.50	SPT-CL J2230-5634	337.6822	-56.5756	4.18	1.00	1.25±0.07	1.82 ^{+0.27} _{-0.39}	29±6		-
SPT-CL J2230-5598 338.1555 -52.2691 5.33 0.50 1.37±0.07 2.04 ^{-1.5} 21±3 1 LOS SPT-CL J2232-5216 338.1555 -52.2691 5.33 0.50 1.37±0.07 2.04 ^{-1.5} 21±3 1 LOS SPT-CL J2232-5516 338.1555 -52.2691 4.76 0.25 0.53±0.02 2.57±0.44 43:6 0 - SPT-CL J2232-5518 388.2656 -53.9616 1.608 0.50 0.4±0.01 3.12±0.02 4.3±0.02 0.5±0.02 2.57±0.44 43:6 0 - SPT-CL J2233-5512 338.352 -49.8562 1.430 1.00 0.440 0.557±0.02 3.07±0.02						-	-	-		-
SPIF-CL J2232-59599 338.1419 -59.9972 17.15 0.25 0.595 0.595 0.25 0.595 0.25 0.595 0.25 0.595 0.25 0.595 0.25 0.25 0.25 0.25 0.25 0.25 0.25 0.2						-	- 0.07+0.40	-		-
SPT-CL 12232-5216 338 1555 522691 5.38 0.50 0.374:007 2.04 ^{0.515} 348:7 0.5 0.5							2.31_049			-
SPT-CL J2232-6301 338.1787 -63.0171 4.76 0.25 0.532-0.02 2.25 34 4366 0 -										100
SPT-CL J2233-5358 338,2050 -53,9881 6.08 0.50 0.44±.001 31,276.27 482±.7 0 - SPT-CL J2233-5539 338,3272 -53,6512 14.30 1.00 0.440 5.22.37 107±11 0 - SPT-CL J2233-5539 338,3272 -53,6512 14.30 1.00 0.440 5.22.37 107±11 0 - SPT-CL J2233-5539 338,3272 -53,6512 4.40 0.25 - SPT-CL J2234-5639 338,676 -58,8223 4.40 0.25 - SPT-CL J2234-6029 338,7367 -60,4399 5.02 0.7±20.02 2.67.38 39±7 0 - SPT-CL J2234-5740 338,7367 -60,4399 5.02 0.25 0.47±20.02 2.67.38 39±7 0 - SPT-CL J2235-5625 338,7981 -52,8709 4.82 1.75 0.53±0.02 2.67.38 33±7 0 - SPT-CL J2235-5625 338,7981 -52,8709 4.82 1.75 0.53±0.02 2.67.38 36±7 0 - SPT-CL J2235-5137 338,6578 -51,9534 5.55 0.25 1.04±0.04 1.97.38 36±7 0 - SPT-CL J2236-5344 339,0388 -53,7471 5.89 1.25 0.45±0.01 3.05.38 43±8 0 - SPT-CL J2236-5344 339,0388 -53,7471 5.89 1.25 0.45±0.01 3.05.38 43±8 0 - SPT-CL J2237-5406 339,4800 -54,1010 7.42 0.75 0.75±0.01 2.70.38 13±5 0 SPT-CL J2238-6724 339,680 -54,1010 7.42 0.75 0.75±0.01 2.70.38 13±5 0 SPT-CL J2238-6349 339,9601 -64,262 4.45 0.50 0.18±0.01 2.70.38 2.8±6 0 SPT-CL J2238-6409 339,9601 -64,262 4.45 0.50 0.18±0.01 2.70.38 2.8±6 0 SPT-CL J2238-6409 339,9601 -64,262 4.45 0.50 0.18±0.01 2.72.38 4.3±8 0 SPT-CL J2238-6409 339,9601 -64,262 4.45 0.50 0.18±0.01 2.72.38 4.3±8 0 SPT-CL J2238-6409 339,9601 -64,262 4.45 0.50 0.18±0.01 2.72.38 4.3±8 0 SPT-CL J2238-6409 339,9601 -64,262 4.45 0.50 0.18±0.01 2.72.38 4.3±8 0 SPT-CL J2238-6409 339,9601 -64,262 4.45 0.50 0.18±0.01 2.72.38 4.3±8 0 SPT-CL J2238-6409 339,9601 -64,262 4.45 0.50 0.18±0.01 2.72.38 4.3±8 0 SPT-CL J2238-6409 339,9601 -64,262 4.45 0.50 0.18±0.01 2.72.38 4.3±8 0 SPT-CL J2245-6538 341,3774 -53,9784 4.3							0.50+0.44			LUS
SPT-CL J2233-4951 338.3154 -49.8562 6.62 0.25 0.72±0.02 3.02**\(\frac{0.25}{0.75} \)							TN.34			-
SPT-CL J2233-5539 338.3272 -53.6512 14.30 1.00 0.440 5.52***\(\frac{6.5}{0.45}\) 107:±11 0 -SPT-CL J2234-5552 338.6366 -58.8223 4.40 0.25 -1.06±0.06 2.58***\(\frac{6.45}{0.45}\) 39±7 0 - SPT-CL J2234-6549 338.7367 -60.4893 5.38 0.25 0.67±0.02 2.67**\(\frac{0.45}{0.45}\) 33±7 0 - SPT-CL J2234-55493 338.7367 -52.8709 4.16 0.25 0.74±0.02 2.67**\(\frac{0.45}{0.45}\) 36±7 0 - - 0 - - 0 - - 0 - - 0 - - 0 - - 0 - - 0 - - 0 - - 0 - - 0 - - 0 - - - 0 - - - - - - 0 - - - - - - - <t< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>-</td></t<>										-
SPT-CL J2234-5849 338,4056 -55.8770 6.26 0.25 1.06±0.06 2.56***\u00e44** 39±7 0 - SPT-CL J2234-5849 338,6376 -58.8223 4.40 0.25 0.67±0.02 2.63**\u00e5							5.02 _{-0.51} 5.52+0.62			-
SPT-CL J2234-5049 338.6376 -58.8223 4.40 0.25										-
SPT-CL J2234-6029 338.7367 -60.4839 5.38 0.25 0.67t.002 2.63**_0.65**_0.						1.00±0.06	2.56_0.45	39±1		_
SPT-CL J2235-5252 387.98						0.67+0.02	2 63+0.45	33+7		- -
SPT-CL J2235-5252 338.7981 -52.8709							2 67+0.46			_
SPT-CL J2235-5157 338.8578 -51.9534 5.54 0.25 1.03±0.03 2.36*0.34 5.0±8 0 LOS SPT-CL J2236-5344 339.0381 -53.2596 4.25 0.25 1.04±0.04 1.97*0.34 27±6 0 LOS SPT-CL J2236-5044 339.0388 -53.7471 5.89 1.25 0.45±0.01 3.05*0.32 13±5 0 - SPT-CL J2236-6041 339.1794 -60.6969 4.08 0.75 1.55±0.07 1.60*0.32 13±5 0 - SPT-CL J2237-5406 339.4693 -54.4351 4.39 0.25 0.17±0.01 2.70*0.32 20±5 0 - SPT-CL J2238-5720 339.5596 -67.3707 4.36 0.25 0 - SPT-CL J2238-6049 339.5591 -60.3229 4.00 0.25 0.50±0.03 2.35*0.32 20±5 0 - SPT-CL J2238-6018 339.5691 -60.3229 4.00 0.25 0.50±0.03 2.35*0.32 24±5 0 - SPT-CL J2238-6018 339.7697 -62.7595 4.10 0.25 1.24±0.07 1.79*0.44 28±6 0 - SPT-CL J2239-5409 339.7978 -54.1509 4.33 0.25 0 0 - SPT-CL J2239-5409 339.7978 -54.1509 4.33 0.25 0 0 - SPT-CL J2239-5409 339.7978 -64.1509 4.33 0.25 0 0 - SPT-CL J2239-5404 339.9902 -50.7053 4.31 0.50 0.69±0.02 2.24*0.34 29±6 0 - SPT-CL J2239-5415 339.9902 -50.7053 4.31 0.50 0.69±0.02 2.24*0.34 43±8 0 - SPT-CL J2241-5338 340.3322 -53.6443 7.64 0.25 0.99±0.02 2.24*0.34 66±9 0 - SPT-CL J2243-5358 340.3322 -53.6443 7.64 0.25 0.99±0.02 2.24*0.34 69±9 0 - SPT-CL J2243-5158 340.6446 -51.8453 4.58 0.50 0.75±0.03 2.27*0.39 24±6 0 - SPT-CL J2243-5518 340.7862 -51.9818 4.52 0.50 0.75±0.03 2.27*0.39 24±6 0 - SPT-CL J2243-5518 340.7862 -51.9818 4.52 0.50 0.75±0.03 2.27*0.39 24±6 0 - SPT-CL J2244-5508 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.08*0.44 0±7 0 LOS SPT-CL J2244-5503 341.2584 -57.1960 4.9 0.25 0.88±0.03 2.08*0.44 0±7 0 LOS SPT-CL J2244-5503 341.2584 -57.1960 4.9 0.25 0.88±0.03 2.08*0.44 0±7 0 - SPT-CL J2245-5043 341.2584 -57.1960 4.9 0.25 0.95±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341.3798 -53.9734 5.36 0.25 0.95±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341.3798 -53.9734 5.36 0.25 0.95±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341.3798 -53.9734 5.30 0.25 0.75±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341.4567 -50.3560 4.27 0.50 0.566 0.88*0.03 2.08*0.44 0±7 0 - SPT-CL J2245-5049 341.4567 -50.5040 4.99 0.25 0.75±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341						-	-0.56	-		-
SPT-CL J2235-5157 338.8578 -51.9534 5.54 0.25 1.03±0.03 2.36*0.34 5.0±8 0 LOS SPT-CL J2236-5344 339.0381 -53.2596 4.25 0.25 1.04±0.04 1.97*0.34 27±6 0 LOS SPT-CL J2236-5044 339.0388 -53.7471 5.89 1.25 0.45±0.01 3.05*0.32 13±5 0 - SPT-CL J2236-6041 339.1794 -60.6969 4.08 0.75 1.55±0.07 1.60*0.32 13±5 0 - SPT-CL J2237-5406 339.4693 -54.4351 4.39 0.25 0.17±0.01 2.70*0.32 20±5 0 - SPT-CL J2238-5720 339.5596 -67.3707 4.36 0.25 0 - SPT-CL J2238-6049 339.5591 -60.3229 4.00 0.25 0.50±0.03 2.35*0.32 20±5 0 - SPT-CL J2238-6018 339.5691 -60.3229 4.00 0.25 0.50±0.03 2.35*0.32 24±5 0 - SPT-CL J2238-6018 339.7697 -62.7595 4.10 0.25 1.24±0.07 1.79*0.44 28±6 0 - SPT-CL J2239-5409 339.7978 -54.1509 4.33 0.25 0 0 - SPT-CL J2239-5409 339.7978 -54.1509 4.33 0.25 0 0 - SPT-CL J2239-5409 339.7978 -64.1509 4.33 0.25 0 0 - SPT-CL J2239-5404 339.9902 -50.7053 4.31 0.50 0.69±0.02 2.24*0.34 29±6 0 - SPT-CL J2239-5415 339.9902 -50.7053 4.31 0.50 0.69±0.02 2.24*0.34 43±8 0 - SPT-CL J2241-5338 340.3322 -53.6443 7.64 0.25 0.99±0.02 2.24*0.34 66±9 0 - SPT-CL J2243-5358 340.3322 -53.6443 7.64 0.25 0.99±0.02 2.24*0.34 69±9 0 - SPT-CL J2243-5158 340.6446 -51.8453 4.58 0.50 0.75±0.03 2.27*0.39 24±6 0 - SPT-CL J2243-5518 340.7862 -51.9818 4.52 0.50 0.75±0.03 2.27*0.39 24±6 0 - SPT-CL J2243-5518 340.7862 -51.9818 4.52 0.50 0.75±0.03 2.27*0.39 24±6 0 - SPT-CL J2244-5508 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.08*0.44 0±7 0 LOS SPT-CL J2244-5503 341.2584 -57.1960 4.9 0.25 0.88±0.03 2.08*0.44 0±7 0 LOS SPT-CL J2244-5503 341.2584 -57.1960 4.9 0.25 0.88±0.03 2.08*0.44 0±7 0 - SPT-CL J2245-5043 341.2584 -57.1960 4.9 0.25 0.95±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341.3798 -53.9734 5.36 0.25 0.95±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341.3798 -53.9734 5.36 0.25 0.95±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341.3798 -53.9734 5.30 0.25 0.75±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341.4567 -50.3560 4.27 0.50 0.566 0.88*0.03 2.08*0.44 0±7 0 - SPT-CL J2245-5049 341.4567 -50.5040 4.99 0.25 0.75±0.03 2.28*0.44 0±7 0 - SPT-CL J2245-5049 341	SPT-CL J2235-5213	338.8268	-52.2202	4.82	1.75	0.53±0.02	2.54 ^{+0.43} _{-0.54}	36±7	0	-
SPT-CL J2236-6341 339.0138	SPT-CL J2235-5157	338.8578	-51.9534	5.54	0.25	1.03±0.03	2.36+0.43	50±8	0	-
SPT-CL J2236-6044 339,0388 -53,7471 5.89 1.25 0.45t0.01 3.05,048 4318 0 - SPT-CL J2237-5426 339,4693 -54,4351 4.39 0.25 0.17±0.01 2.70,38 20±5 0 - SPT-CL J2237-5406 339,4800 -54,1010 7.42 0.75 0.40±0.01 3.68,058 5628 0 - SPT-CL J2238-6722 339,5596 -57,3707 4.36 0.25 - SPT-CL J2238-6019 339,5591 -60,3289 4.20 0.25 0.50±0.03 2.55,058 16±5 0 - SPT-CL J2238-6338 339,6819 -61,6425 4.45 0.50 0.168±0.010 2.72,048 2±45 0 - SPT-CL J2238-6138 339,7367 -54,1509 4.33 0.25 1.24±0.07 1.79,041 2±45 0 - SPT-CL J2239-5409 339,7978 -54,1509 4.33 0.25 1.24±0.07 1.79,041 2±66 0 - SPT-CL J2239-5415 339,9061 -54,2562 4.39 1.25 0.45±0.02 2.66,054 2±6 0 - SPT-CL J2239-545 339,9061 -54,2562 4.39 1.25 0.45±0.02 2.66,054 2±6 0 - SPT-CL J2239-545 339,902 -61,2557 0.50 0.69±0.02 2.34,048 65±9 0 - SPT-CL J2240-6117 340,0204 -61,2957 7.05 0.50 0.95±0.02 2.33,048 65±9 0 - SPT-CL J2241-5338 340,3322 -53,6443 7.64 0.25 0.93±0.02 2.34,048 65±9 0 - SPT-CL J22425-5150 340,5482 -51,9818 4.42 0.50 0.47±0.02 2.46,054 2±6 0 - SPT-CL J2243-5518 340,7582 -51,9818 4.42 0.50 0.47±0.02 2.46,054 2±6 0 - SPT-CL J2243-5523 340,7582 -51,9818 4.22 0.50 0.47±0.02 2.46,054 2±6 0 - SPT-CL J2243-5623 341,2564 -57,1960 4.19 0.25 0.88±0.03 2.87,038 43±7 0 - SPT-CL J2245-5723 341,2584 -57,1960 4.19 0.25 0.88±0.03 2.87,038 43±7 0 - SPT-CL J2245-5723 341,2584 -57,1960 4.19 0.25 0.50±0.02 3.82,044 40±7 0 - SPT-CL J2245-5723 341,2584 -57,1960 4.19 0.25 0.50±0.02 3.82,044 40±7 0 - SPT-CL J2245-5723 341,2584 -53,9734 5.36 0.25 0.95±0.02 3.82,044 40±7 0 - SPT-CL J2245-5648 341,378 -53,9734 5.36 0.25 0.95±0.02 3.82,044 40±7 0 - SPT-CL J2245-5648 341,3929 -64,9787 5.52 0.25 0.40±0.01 2.89,043 30±4 1 - SPT-CL J2245-6548 341,3878 -53,9748 5.20 0.25 0.40±0.01 2.89,044 40±7 0 - SPT-CL J2245-6548 341,3878 -53,9748 5.20 0.25 0.40±0.01 2.85,044 40±7 0 - SPT-CL J2245-6548 341,3878 -53,9748 5.20 0.25 0.50±0.02 3.82,044 40±7 0 - SPT-CL J2245-6548 341,3878 -53,9748 5.20 0.25 0.50±0.02 3.82,044 40±7 0 - SPT-CL J2245-6548 341,3874 -53,9748 5.20 0.25 0.50±0.02 3.82,044 40±7 0 - SPT-CL J2245	SPT-CL J2236-5315	339.0131	-53.2596	4.25	0.25	1.04±0.04	1.97 0.31	27±6	0	LOS
SPT-CL J2237-5426 339,4680 -54,4351 4.39 0.25 0.17±0.01 2.70; 0.40±0.01 5.6±8 0 - SPT-CL J2238-6722 339,5596 -57,3707 4.36 0.25 - 0.25 0.50±0.03 2.35; 0.38 0.25 - 0.25 SPT-CL J2238-6191 339,5691 -60,3289 4.20 0.25 0.50±0.03 2.35; 0.38 0.25 SPT-CL J2238-6193 339,6819 -61,6425 4.45 0.50 0.168±0.010 2.72; 0.40 0.25 0.50±0.03 2.35; 0.38 0.25 SPT-CL J2238-6245 339,7367 -62,7595 4.10 0.25 1.24±0.07 1.75; 0.41 28±6 0 - SPT-CL J2239-5409 339,7367 -62,7595 4.10 0.25 1.24±0.07 1.75; 0.41 28±6 0 - SPT-CL J2239-5409 339,901 -54,1509 4.33 0.25 SPT-CL J2239-5402 339,902 -50,7053 4.31 0.50 0.69±0.02 2.46; 0.42 29±6 0 - SPT-CL J2239-5404 339,9061 -54,2562 4.39 1.25 0.45±0.02 2.46; 0.42 29±6 0 - SPT-CL J2241-5338 340,3322 -53,6443 7.64 0.25 0.93±0.02 2.93; 0.44 0.95 0 - SPT-CL J2241-5338 340,3322 -53,6443 7.64 0.25 0.93±0.02 3.12; 0.45 0.0 2 2.46; 0.42 0.95 0 - SPT-CL J2242-5150 340,6446 -51,8453 4.58 0.50 0.75±0.03 3.12; 0.45 0.0 2 2.46; 0.45	SPT-CL J2236-5344	339.0388	-53.7471	5.89	1.25	0.45±0.01	3.05+0.48	43±8	0	_
SPT-CL J2237-5426 339,4680 -54,4351 4.39 0.25 0.17±0.01 2.70; 0.40±0.01 5.6±8 0 - SPT-CL J2238-6722 339,5596 -57,3707 4.36 0.25 - 0.25 0.50±0.03 2.35; 0.38 0.25 - 0.25 SPT-CL J2238-6191 339,5691 -60,3289 4.20 0.25 0.50±0.03 2.35; 0.38 0.25 SPT-CL J2238-6193 339,6819 -61,6425 4.45 0.50 0.168±0.010 2.72; 0.40 0.25 0.50±0.03 2.35; 0.38 0.25 SPT-CL J2238-6245 339,7367 -62,7595 4.10 0.25 1.24±0.07 1.75; 0.41 28±6 0 - SPT-CL J2239-5409 339,7367 -62,7595 4.10 0.25 1.24±0.07 1.75; 0.41 28±6 0 - SPT-CL J2239-5409 339,901 -54,1509 4.33 0.25 SPT-CL J2239-5402 339,902 -50,7053 4.31 0.50 0.69±0.02 2.46; 0.42 29±6 0 - SPT-CL J2239-5404 339,9061 -54,2562 4.39 1.25 0.45±0.02 2.46; 0.42 29±6 0 - SPT-CL J2241-5338 340,3322 -53,6443 7.64 0.25 0.93±0.02 2.93; 0.44 0.95 0 - SPT-CL J2241-5338 340,3322 -53,6443 7.64 0.25 0.93±0.02 3.12; 0.45 0.0 2 2.46; 0.42 0.95 0 - SPT-CL J2242-5150 340,6446 -51,8453 4.58 0.50 0.75±0.03 3.12; 0.45 0.0 2 2.46; 0.45	SPT-CL J2236-6041	339.1794	-60.6969	4.08	0.75	1.55±0.07	1.60+0.25	13±5	0	-
SPT-CL J2238-5722 339.5966 -57.3707	SPT-CL J2237-5426	339.4693	-54.4351	4.39	0.25	0.17±0.01	2.70 +0.46	20±5	0	-
SPT-CL J2238-6019 393.95601 -50.3707 4.36 0.25	SPT-CL J2237-5406	339.4800	-54.1010	7.42	0.75	0.40±0.01	3.68+0.49	56±8	0	-
SPT-CL J2238-6138	SPT-CL J2238-5722	339.5596	-57.3707	4.36	0.25	-	_	-	0	-
SPT-CL J2238-6409 339,7367 -62,7595 4.10 0.25 1.24±0.07 1.79+0.41 -0.0	SPT-CL J2238-6019	339.5601	-60.3289	4.20	0.25	0.50±0.03	2.35 ^{+0.38} _{-0.52}	16±5	0	-
SPT-CL J2239-5405 339.7978 -54.1509 4.33 0.25 - - 0 - SPT-CL J2239-5415 339.9061 -54.2562 4.39 1.25 0.450.02 2.4610.40 2.24610.40 2.24610.40 - <td< td=""><td>SPT-CL J2238-6138</td><td>339.6819</td><td>-61.6425</td><td>4.45</td><td>0.50</td><td>0.168±0.010</td><td>$2.72^{+0.46}_{-0.60}$</td><td>24±5</td><td>0</td><td>-</td></td<>	SPT-CL J2238-6138	339.6819	-61.6425	4.45	0.50	0.168±0.010	$2.72^{+0.46}_{-0.60}$	24±5	0	-
SPT-CL J2239-5042 339.9061 -54.2562 4.39 1.25 0.45±0.02 2.46±0.49 29±6 0 - SPT-CL J2239-5042 339.9902 -50.7053 4.31 0.50 0.69±0.02 2.24±0.49 43±8 0 - SPT-CL J2241-65338 340.3322 -53.6443 7.64 0.25 0.93±0.02 2.93±0.49 43±8 0 - SPT-CL J2242-5150 340.6446 -51.8453 4.58 0.50 0.75±0.03 2.27±0.50 69±9 0 - SPT-CL J2243-5158 340.7582 -51.9818 4.42 0.50 0.75±0.03 2.27±0.50 2±±6 0 - SPT-CL J2243-5535 340.7780 -53.5881 6.52 0.75 0.44±0.01 3.29±0.45 67±8 0 - SPT-CL J2244-5908 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.05±0.45 40±7 0 LOS SPT-CL J2245-6207 341.2584 -62.1178 18.70 0.50 0.5886 6.18±0.61				_		1.24±0.07	1.79 ^{+0.29} _{-0.41}	28±6		-
SPT-CL J2239-5042 339.9902 -50.7053 4.31 0.50 0.69±0.02 2.24 ^{10.36} -036 43±8 0 - SPT-CL J2240-6117 340.0204 -61.2957 7.05 0.50 0.95±0.02 2.93 ^{-0.38} -0.98 65±9 0 - SPT-CL J2242-5150 340.6446 -51.8453 4.58 0.50 0.93±0.02 3.12 ^{-0.50} -050 96±9 0 - SPT-CL J2243-5158 340.7882 -51.9818 4.42 0.50 0.47±0.02 2.46 ^{-0.54} -0.54 36±6 0 LOS SPT-CL J2243-5535 340.7780 -53.5881 6.52 0.75 0.44±0.01 3.29 ^{-0.48} -0.56 67±8 0 - SPT-CL J2244-5908 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.08 ^{-0.55} -0.45 59±5 1 - SPT-CL J2244-5908 341.2584 -62.178 18.70 0.50 0.586 6.18 ^{-0.65} -0.45 40±7 0 - SPT-CL J2245-5003 341.2584 -62.1178 18.70 0.50 <t< td=""><td></td><td></td><td></td><td></td><td></td><td>-</td><td></td><td>-</td><td></td><td>-</td></t<>						-		-		-
SPT-CL J2241-5338 340.3322 -53.6443 7.64 0.25 0.93±0.02 3.12*0.50 69±9 0 - SPT-CL J2242-5150 340.6446 -51.8453 4.58 0.50 0.75±0.03 2.27*0.30 24±6 0 - SPT-CL J2243-5158 340.7580 -51.8818 4.42 0.50 0.47±0.02 2.46*0.02 36±6 0 LOS SPT-CL J2243-5835 340.7780 -53.5881 6.52 0.75 0.44±0.01 3.29*0.48 67±8 0 - SPT-CL J2243-5821 340.9549 -58.3610 7.48 0.75 0.44±0.01 3.29*0.48 67±8 0 - SPT-CL J2244-5908 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.08*0.45 40±7 0 LOS SPT-CL J2244-6920 341.2584 -62.1178 18.70 0.50 0.586 6.18*0.01 112±11 0 RC SPT-CL J2245-5023 341.2681 -50.3950 4.21 1.50 - - 0<							2.46 -0.54			-
SPT-CL J2241-5338 340.3322 -53.6443 7.64 0.25 0.93±0.02 3.12*0.50 69±9 0 - SPT-CL J2242-5150 340.6446 -51.8453 4.58 0.50 0.75±0.03 2.27*0.30 24±6 0 - SPT-CL J2243-5158 340.7580 -51.8818 4.42 0.50 0.47±0.02 2.46*0.02 36±6 0 LOS SPT-CL J2243-5835 340.7780 -53.5881 6.52 0.75 0.44±0.01 3.29*0.48 67±8 0 - SPT-CL J2243-5821 340.9549 -58.3610 7.48 0.75 0.44±0.01 3.29*0.48 67±8 0 - SPT-CL J2244-5908 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.08*0.45 40±7 0 LOS SPT-CL J2244-6920 341.2584 -62.1178 18.70 0.50 0.586 6.18*0.01 112±11 0 RC SPT-CL J2245-5023 341.2681 -50.3950 4.21 1.50 - - 0<							2.24-0.30			-
SPT-CL J2245-5150							TU. 11			-
SPT-CL J2243-5158							3. IZ _{-0.50}			-
SPT-CL J2243-5335 340.7780 -53.5881 6.52 0.75 0.44±0.01 3.29±0.57 67±8 0 - SPT-CL J2243-5821 340.9549 -58.3610 7.48 0.75 1.13±0.06 2.87±0.35 59±5 1 - SPT-CL J2244-5908 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.08±0.34 40±7 0 LOS SPT-CL J2244-6420 341.2154 -64.3362 4.03 0.25 - - 0 - SPT-CL J2245-5711 341.2580 -57.1960 4.19 0.25 0.48±0.01 2.39±0.38 43±7 0 - SPT-CL J2245-6207 341.2681 -50.3950 4.21 1.50 - - 0 - SPT-CL J2245-5023 341.3018 -57.3899 6.67 0.25 0.50±0.02 3.28±0.47 29±6 0 - SPT-CL J2245-5358 341.3774 -53.9734 5.36 0.25 0.92±0.03 2.39±0.45 45±8 0 - <t< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td>2.27_0.50</td><td></td><td></td><td>-</td></t<>							2.27_0.50			-
SPT-CL J2243-5821 340.9549 -58.3610 7.48 0.75 1.13±0.06 2.87+0.35 59±5 1 - SPT-CL J2244-6908 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.08±0.34 40±7 0 LOS SPT-CL J2244-6420 341.2154 -64.3362 4.03 0.25 - - 0 - SPT-CL J2245-5711 341.2580 -57.1960 4.19 0.25 0.48±0.01 2.39±0.38 43±7 0 - SPT-CL J2245-6207 341.2584 -62.1178 18.70 0.50 0.586 6.18±0.61 112±11 0 RC SPT-CL J2245-5023 341.2081 -50.3950 4.21 1.50 - - - 0 - SPT-CL J2245-5353 341.3018 -57.3899 6.67 0.25 0.50±0.02 3.28±0.67 29±6 0 - SPT-CL J2245-5358 341.3774 -53.974 4.97 0.25 0.95±0.03 2.24±0.47 34±6 0 - <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>2.46-0.54</td> <td></td> <td></td> <td>LOS</td>							2.46-0.54			LOS
SPT-CL J2244-5908 341.1715 -59.1403 4.22 0.25 0.88±0.03 2.0±0.34 40±7 0 LOS SPT-CL J2244-6420 341.2154 -64.3362 4.03 0.25 0 - 0 SPT-CL J2245-5711 341.2580 -57.1960 4.19 0.25 0.48±0.01 2.39±0.52 43±7 0 - SPT-CL J2245-5023 341.2681 -50.3950 4.21 1.50 0 - SPT-CL J2245-5023 341.2681 -50.3950 4.21 1.50 0 - SPT-CL J2245-5723 341.3018 -57.3899 6.67 0.25 0.50±0.02 3.28±0.47 2.9±6 0 - SPT-CL J2245-5358 341.3774 -53.9734 5.36 0.25 0.92±0.03 2.39±0.45 2.9±6 0 - SPT-CL J2245-5354 341.3798 -53.9158 4.97 0.25 0.95±0.03 2.24±0.40 34±6 0 - SPT-CL J2245-6458 341.3929 -64.9787 5.52 0.25 1.47±0.07 2.03±0.34 30±4 1 - SPT-CL J2245-5254 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85±0.55 40±7 0 - SPT-CL J2245-6113 341.4864 -61.2316 4.59 0.25 0.77±0.02 2.26±0.39 35±7 0 - SPT-CL J2246-5243 341.5638 -52.7302 8.41 2.25 0.102±0.003 4.38±0.67 96±10 0 RC SPT-CL J2246-6046 341.5810 -60.7755 4.10 0.50 0.41±0.01 2.40±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.34 41±7 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.34 41±7 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.35 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21±0.03 30±6 0							. 8.85			-
SPT-CL J22245-6420 341.2154 -64.3362 4.03 0.25 - - 0 - SPT-CL J2245-5711 341.2580 -57.1960 4.19 0.25 0.48±0.01 2.39 ^{+0.38} / _{-0.52} 43±7 0 - SPT-CL J2245-6207 341.2584 -62.1178 18.70 0.50 0.586 6.18 ^{+0.67} / _{-0.60} 112±11 0 RC SPT-CL J2245-5023 341.3681 -57.3899 6.67 0.25 0.50±0.02 3.28 ^{+0.47} / _{-0.56} 29±6 0 - SPT-CL J2245-5358 341.3774 -53.9734 5.36 0.25 0.92±0.03 2.39 ^{+0.45} / _{-0.47} 45±8 0 - SPT-CL J2245-5354 341.3798 -53.9158 4.97 0.25 0.95±0.03 2.24 ^{+0.47} / _{-0.47} 34±6 0 - SPT-CL J2245-6458 341.3929 -64.9787 5.52 0.25 1.47±0.07 2.03 ^{+0.46} / _{-0.47} 30±4 1 - SPT-CL J2245-6009 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85 ^{+0.47} / _{-0.40}							2.07 _{-0.45}			-
SPT-CL J2245-5711 341.2580 -57.1960 4.19 0.25 0.48±0.01 2.39±0.52 43±7 0 - SPT-CL J2245-6207 341.2584 -62.1178 18.70 0.50 0.586 6.18±0.67 112±11 0 RC SPT-CL J2245-5023 341.3018 -57.3899 6.67 0.25 0.50±0.02 3.28±0.47 29±6 0 - SPT-CL J2245-5358 341.3774 -53.9734 5.36 0.25 0.92±0.03 2.39±0.45 45±8 0 - SPT-CL J2245-5354 341.3798 -53.9158 4.97 0.25 0.95±0.03 2.39±0.49 34±6 0 - SPT-CL J2245-6458 341.3929 -64.9787 5.52 0.25 1.47±0.07 2.03±0.44 30±4 1 - SPT-CL J2245-6458 341.4516 -60.1568 4.73 0.25 1.47±0.07 1.80±0.49 30±4 1 - SPT-CL J2245-62543 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85±0.55						0.88±0.03	2.08-0.45	40±7		LOS
SPT-CL J2245-5023						0.49±0.01	2 20+0.38	- 12±7	-	-
SPT-CL J2245-5023							2.39 _{-0.52}			- DC
SPT-CL J2245-5723 341.3018 -57.3899 6.67 0.25 0.50±0.02 3.28 ^{-0.47} _{-0.56} 29±6 0 - SPT-CL J2245-5358 341.3774 -53.9734 5.36 0.25 0.92±0.03 2.39 ^{-0.45} _{-0.47} 45±8 0 - SPT-CL J2245-5354 341.3798 -53.9158 4.97 0.25 0.95±0.03 2.24 ^{-0.47} _{-0.47} 34±6 0 - SPT-CL J2245-6458 341.3929 -64.9787 5.52 0.25 1.47±0.07 2.03 ^{-0.40} _{-0.40} 30±4 1 - SPT-CL J2245-6009 341.4516 -60.1568 4.73 0.25 1.47±0.07 1.80 ^{-0.30} _{-0.40} 20±3 1 - SPT-CL J2245-5254 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85 ^{-0.55} _{-0.45} 40±7 0 - SPT-CL J2246-6113 341.4864 -61.2316 4.59 0.25 0.77±0.02 2.26 ^{-0.39} _{-0.49} 35±7 0 - SPT-CL J2246-5308 341.5670 -53.1338 4.96 0.2						0.566	0.10_0.81			RC -
SPT-CL J2245-5354 341.3798 -53.9158 4.97 0.25 0.95±0.03 2.24*0.47 (0.47) 34±6 0 - SPT-CL J2245-6458 341.3929 -64.9787 5.52 0.25 1.47±0.07 2.03*0.34 30±4 1 - SPT-CL J2245-6009 341.4516 -60.1568 4.73 0.25 1.47±0.07 1.80*0.30 20±3 1 - SPT-CL J2245-5254 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85*0.47 0 - SPT-CL J2245-6113 341.4864 -61.2316 4.59 0.25 0.77±0.02 2.26*0.39 35±7 0 - SPT-CL J2246-5243 341.5638 -52.7302 8.41 2.25 0.102±0.003 4.38*0.65 96±10 0 RC SPT-CL J2246-5308 341.5670 -53.1338 4.96 0.25 1.43±0.07 1.89*0.34 41±7 0 - SPT-CL J2246-6046 341.5810 -60.7755 4.10 0.50 0.41±0.01 2.40*0.40 0						0.50±0.02	3.28 +0.47	29±6		_
SPT-CL J2245-5354 341.3798 -53.9158 4.97 0.25 0.95±0.03 2.24*0.47 (0.47) 34±6 0 - SPT-CL J2245-6458 341.3929 -64.9787 5.52 0.25 1.47±0.07 2.03*0.34 30±4 1 - SPT-CL J2245-6009 341.4516 -60.1568 4.73 0.25 1.47±0.07 1.80*0.30 20±3 1 - SPT-CL J2245-5254 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85*0.47 0 - SPT-CL J2245-6113 341.4864 -61.2316 4.59 0.25 0.77±0.02 2.26*0.39 35±7 0 - SPT-CL J2246-5243 341.5638 -52.7302 8.41 2.25 0.102±0.003 4.38*0.65 96±10 0 RC SPT-CL J2246-5308 341.5670 -53.1338 4.96 0.25 1.43±0.07 1.89*0.34 41±7 0 - SPT-CL J2246-6046 341.5810 -60.7755 4.10 0.50 0.41±0.01 2.40*0.40 0							2.39+0.45			_
SPT-CL J2245-5254 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85-0.47 (10.55) 40±7 0 - SPT-CL J2245-6113 341.4864 -61.2316 4.59 0.25 0.77±0.02 2.26-0.39 (10.55) 35±7 0 - SPT-CL J2246-5243 341.5638 -52.7302 8.41 2.25 0.102±0.003 4.38-0.55 (10.67) 96±10 0 RC SPT-CL J2246-5308 341.5670 -53.1338 4.96 0.25 1.43±0.07 1.89-0.41 (10.67) 41±7 0 - SPT-CL J2246-6046 341.5810 -60.7755 (10.50) 4.10 0.50 (0.41±0.01) 2.40-0.40 (10.53) 30±6 (10.55) 0 - SPT-CL J2246-6234 341.5916 (10.56) -62.5697 (10.56) 4.35 (10.55) 0.25 (10.55) 0.75±0.03 (10.55) 2.21-0.48 (10.55) 10.50 (10.55) 42±7 (10.55) 0 SPT-CL J2246-5925 341.7126 (10.59) -59.4303 (10.56) 4.49 (10.55) 1.01±0.03 (10.55) 2.05-0.45 (10.55) 43±8 (10.55) 0 RC SPT-CL J2247-5715 341.7897 (10.55) -57.2546 (10.55) 5.21 (10.50) 1.28±0.07 (10.55) 2.05-0.45 (10.55) 43±4 (10.55) 1 - SPT-CL J2247-57139 341.8433 (10.56) -57.2546 (10.55) <td< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td>2.24+0.40</td><td></td><td></td><td>_</td></td<>							2.24+0.40			_
SPT-CL J2245-5254 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85-0.47 (10.55) 40±7 0 - SPT-CL J2245-6113 341.4864 -61.2316 4.59 0.25 0.77±0.02 2.26-0.39 (10.55) 35±7 0 - SPT-CL J2246-5243 341.5638 -52.7302 8.41 2.25 0.102±0.003 4.38-0.55 (10.67) 96±10 0 RC SPT-CL J2246-5308 341.5670 -53.1338 4.96 0.25 1.43±0.07 1.89-0.41 (10.67) 41±7 0 - SPT-CL J2246-6046 341.5810 -60.7755 (10.50) 4.10 0.50 (0.41±0.01) 2.40-0.40 (10.53) 30±6 (10.55) 0 - SPT-CL J2246-6234 341.5916 (10.56) -62.5697 (10.56) 4.35 (10.55) 0.25 (10.55) 0.75±0.03 (10.55) 2.21-0.48 (10.55) 10.50 (10.55) 42±7 (10.55) 0 SPT-CL J2246-5925 341.7126 (10.59) -59.4303 (10.56) 4.49 (10.55) 1.01±0.03 (10.55) 2.05-0.45 (10.55) 43±8 (10.55) 0 RC SPT-CL J2247-5715 341.7897 (10.55) -57.2546 (10.55) 5.21 (10.50) 1.28±0.07 (10.55) 2.05-0.45 (10.55) 43±4 (10.55) 1 - SPT-CL J2247-57139 341.8433 (10.56) -57.2546 (10.55) <td< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td>2.03+0.34</td><td></td><td></td><td>_</td></td<>							2.03+0.34			_
SPT-CL J2245-5254 341.4637 -52.9109 5.42 0.25 0.46±0.01 2.85-0.47 (10.55) 40±7 0 - SPT-CL J2245-6113 341.4864 -61.2316 4.59 0.25 0.77±0.02 2.26-0.39 (10.55) 35±7 0 - SPT-CL J2246-5243 341.5638 -52.7302 8.41 2.25 0.102±0.003 4.38-0.55 (10.67) 96±10 0 RC SPT-CL J2246-5308 341.5670 -53.1338 4.96 0.25 1.43±0.07 1.89-0.41 (10.67) 41±7 0 - SPT-CL J2246-6046 341.5810 -60.7755 (10.50) 4.10 0.50 (0.41±0.01) 2.40-0.40 (10.53) 30±6 (10.55) 0 - SPT-CL J2246-6234 341.5916 (10.56) -62.5697 (10.56) 4.35 (10.55) 0.25 (10.55) 0.75±0.03 (10.55) 2.21-0.48 (10.55) 10.50 (10.55) 42±7 (10.55) 0 SPT-CL J2246-5925 341.7126 (10.59) -59.4303 (10.56) 4.49 (10.55) 1.01±0.03 (10.55) 2.05-0.45 (10.55) 43±8 (10.55) 0 RC SPT-CL J2247-5715 341.7897 (10.55) -57.2546 (10.55) 5.21 (10.50) 1.28±0.07 (10.55) 2.05-0.45 (10.55) 43±4 (10.55) 1 - SPT-CL J2247-57139 341.8433 (10.56) -57.2546 (10.55) <td< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td>1.80+0.30</td><td></td><td></td><td>_</td></td<>							1.80+0.30			_
SPT-CL J2246-6046 341.5810 -60.7755 4.10 0.50 0.41±0.01 2.40_0.53 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21_0.48 19±5 0 - SPT-CL J2246-5130 341.6475 -51.5103 7.08 0.75 0.66±0.02 3.25_0.53 42±7 0 - SPT-CL J2246-5925 341.7126 -59.4303 4.49 0.25 1.01±0.03 2.05_0.45 43±8 0 RC SPT-CL J2247-5715 341.7897 -57.2546 5.21 0.50 1.28±0.07 2.05_0.36 43±4 1 - SPT-CL J2247-5139 341.8433 -51.6533 4.36 0.25 0 -							2.85+0.47			_
SPT-CL J2246-6046 341.5810 -60.7755 4.10 0.50 0.41±0.01 2.40_0.53 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21_0.48 19±5 0 - SPT-CL J2246-5130 341.6475 -51.5103 7.08 0.75 0.66±0.02 3.25_0.53 42±7 0 - SPT-CL J2246-5925 341.7126 -59.4303 4.49 0.25 1.01±0.03 2.05_0.45 43±8 0 RC SPT-CL J2247-5715 341.7897 -57.2546 5.21 0.50 1.28±0.07 2.05_0.36 43±4 1 - SPT-CL J2247-5139 341.8433 -51.6533 4.36 0.25 0 -							2 26+0.39			_
SPT-CL J2246-6046 341.5810 -60.7755 4.10 0.50 0.41±0.01 2.40_0.53 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21_0.48 19±5 0 - SPT-CL J2246-5130 341.6475 -51.5103 7.08 0.75 0.66±0.02 3.25_0.53 42±7 0 - SPT-CL J2246-5925 341.7126 -59.4303 4.49 0.25 1.01±0.03 2.05_0.45 43±8 0 RC SPT-CL J2247-5715 341.7897 -57.2546 5.21 0.50 1.28±0.07 2.05_0.36 43±4 1 - SPT-CL J2247-5139 341.8433 -51.6533 4.36 0.25 0 -							4.38+0.55			RC
SPT-CL J2246-6046 341.5810 -60.7755 4.10 0.50 0.41±0.01 2.40_0.53 30±6 0 - SPT-CL J2246-6234 341.5916 -62.5697 4.35 0.25 0.75±0.03 2.21_0.48 19±5 0 - SPT-CL J2246-5130 341.6475 -51.5103 7.08 0.75 0.66±0.02 3.25_0.53 42±7 0 - SPT-CL J2246-5925 341.7126 -59.4303 4.49 0.25 1.01±0.03 2.05_0.45 43±8 0 RC SPT-CL J2247-5715 341.7897 -57.2546 5.21 0.50 1.28±0.07 2.05_0.36 43±4 1 - SPT-CL J2247-5139 341.8433 -51.6533 4.36 0.25 0 -							1.89+0.34			_
SPT-CL J2247-5715 341.7897 -57.2546 5.21 0.50 1.28±0.07 2.05 _{-0.43} 43±4 1 - SPT-CL J2247-5139 341.8433 -51.6533 4.36 0.25 0 -							2.40+0.40			_
SPT-CL J2247-5715 341.7897 -57.2546 5.21 0.50 1.28±0.07 2.05 _{-0.43} 43±4 1 - SPT-CL J2247-5139 341.8433 -51.6533 4.36 0.25 0 -							2.21+0.37			_
SPT-CL J2247-5715 341.7897 -57.2546 5.21 0.50 1.28±0.07 2.05 _{-0.43} 43±4 1 - SPT-CL J2247-5139 341.8433 -51.6533 4.36 0.25 0 -							3.25+0.45			_
SPT-CL J2247-5715 341.7897 -57.2546 5.21 0.50 1.28±0.07 2.05 _{-0.43} 43±4 1 - SPT-CL J2247-5139 341.8433 -51.6533 4.36 0.25 0 -							2.05+0.35			RC
SPT-CL J2247-5139 341 8433 -51.6533 4.36 0.25 - - - 0 -							2.05 ^{+0.36}			-
SPT-CL J2247-5537 341.8992 -55.6304 7.10 0.25 0.93±0.05 2.96 ^{+0.40} _{-0.48} 13±5 0 SPT-CL J2248-5425 342.1721 -54.4325 4.68 0.25 0.85±0.03 2.23 ^{+0.39} _{-0.48} 33±6 0						-	_	-		_
SPT-CL J2248-5425 342.1721 -54.4325 4.68 0.25 0.85±0.03 2.23 ^{+0.39} _{-0.48} 33±6 0 -						0.93±0.05	2.96 ^{+0.40} _{-0.48}	13±5		-
_U.₩0	SPT-CL J2248-5425	342.1721	-54.4325	4.68	0.25	0.85±0.03	2.23 ^{+0.39} _{-0.48}	33±6	0	-

Table 1 — Continued

SPT ID	R.A. (J2000)	Decl. (J2000)	Β ε	est $\theta_{ extsf{c}}$	Redshift	M_{500c} $(10^{14} h_{70}^{-1} M_{\odot})$	λ	Spitzer	Notes		
SPT-CL J2249-6426	342.4579	-64.4379	8.18	1.25	0.094	4.31 ^{+0.56} _{-0.68}	78±9	0	-		
SPT-CL J2249-5528	342.4697	-55.4831	5.04	0.25	1.20±0.07	2.07 ^{+0.37} _{-0.43} 2.37 ^{+0.39} _{-0.52}	23±3	1	-		
SPT-CL J2250-5345	342.6720	-53.7555	4.26	0.25	0.52±0.01	2.37+0.39	52±8	0	-		
SPT-CL J2251-5257	342.8053	-52.9563	5.12	0.25	1.12±0.06	2.16 ^{+0.39}	37±6	0	-		
SPT-CL J2251-5246	342.8300	-52.7781	5.20	0.50	0.93±0.03	2.34 ^{+0.40} _{-0.47}	36±7	0	-		
SPT-CL J2251-5403	342.8682	-54.0533	5.91	0.25	0.49±0.02	$3.00^{+0.47}_{-0.56}$	24±5	0	LOS		
SPT-CL J2251-5049	342.9807	-50.8188	5.66	0.25	0.86±0.03	2.54 ^{+0.44} _{-0.49}	30±6	0	_		
SPT-CL J2252-6011	343.0828	-60.1969	5.97	0.25	1.10±0.06	0.45+0.40	39±7	0	_		
SPT-CL J2252-5720	343.1045	-57.3373	4.38	1.00	1.43±0.07	2.45 _{-0.45} 1.74 ^{+0.28} -0.39	15±4	0	_		
SPT-CL J2252-5737	343.1610	-57.6258	4.45	0.25	1.01±0.04	2.05 ^{+0.34} _{-0.44}	32±7	0	_		
SPT-CL J2253-5247	343.3036	-52.7870	4.45	0.25	0.79±0.03	2.21 ^{+0.36} _{-0.49}	19±5	0	_		
SPT-CL J2253-5408	343.3323	-54.1419	5.99	0.50	0.78±0.03	2.77+0.43	21±6	0	_		
SPT-CL J2254-6314	343.5099	-63.2484	15.57	0.75	0.211	6 22 +0.70	80±10	0	_		
SPT-CL J2254-6142	343.5547	-61.7067	4.21	0.25	0.92±0.04	2.04 ^{+0.34}	23±6	0	_		
SPT-CL J2254-5804	343.5842	-58.0804	5.74	2.25	0.153	3 27+0.53	32±6	0	_		
SPT-CL J2255-5747	343.8958	-57.7935	4.55	0.75	1.05±0.05	2.04+0.34	21±5	0	_		
SPT-CL J2256-6149	344.0089	-61.8259	4.48	0.25	0.34±0.01	2 50+0.45	34±6	0	_		
SPT-CL J2256-5821	344.1214	-58.3647	4.24	0.25	-	2.36 _{-0.56}	-	Ö	_		
SPT-CL J2256-5102	344.1636	-51.0495	4.05	0.25	-		-	0	-		
SPT-CL J2256-6246	344.1898	-62.7703	5.07	0.25	1.13±0.06	2.13 ^{+0.37} _{-0.44}	36±4	1	-		
SPT-CL J2257-5111	344.4303	-51.1863	4.05	0.25		+0.40		0	-		
SPT-CL J2258-5443	344.5558	-54.7271	4.21	2.50	0.30±0.02	2.54 ^{+0.40}	13±4	0			
SPT-CL J2258-6057	344.7486	-60.9582	19.71	0.25	0.79±0.01	5.94 ^{+0.62} _{-0.76}	114±11	0	RC,LOS		
SPT-CL J2259-5232	344.7604	-52.5360	4.68	0.50	0.97±0.04	2.13 ^{+0.38} _{-0.47}	19±6	0	-		
SPT-CL J2259-5349	344.7875	-53.8219	5.88	0.75	0.271±0.009	3.21-0.59	47±8	0	LOS		
SPT-CL J2259-5957	344.9537	-59.9554	4.10	1.50	-	4.15 ^{+0.52}	-	0	-		
SPT-CL J2259-5431	344.9798	-54.5280	9.00	0.50	0.42±0.01	4.15 ^{+0.52} _{-0.63} 4.50 ^{+0.56} _{-0.67}	39±7	0	-		
SPT-CL J2300-5617	345.0042	-56.2869	8.98	0.50	0.153	4.50_0.67	56±8	0	RC		
SPT-CL J2300-5331	345.1699	-53.5200	10.27	1.00	0.262	4.77 ^{+0.54} -0.69 3.05 ^{+0.50} -0.56	76±9	0	-		
SPT-CL J2301-5316	345.3374	-53.2793	5.72	0.75	0.37±0.01	3.05 _{-0.56} 3.08 _{-0.52}	39±7	0	-		
SPT-CL J2301-6055	345.4512	-60.9230	6.78	0.25	0.72±0.02	TX.32	68±8	0	-		
SPT-CL J2301-5546 SPT-CL J2302-5117	345.4595 345.6072	-55.7786 -51.2907	5.68 4.19	0.50 1.00	0.748	2.69-0.50	55±8	0 0	RC		
SPT-CL J2302-5405	345.6581	-51.2907 -54.0881	4.19	0.25	0.76±0.03	2.16 ^{+0.35}	24±6	0	_		
SPT-CL J2303-6335	345.9664	-63.5889	4.39	0.50	0.70±0.03 0.72±0.03	2.10 _{-0.48} 2.24 ^{+0.38} -0.49	29±6	0			
SPT-CL J2303-5600	345.9695	-56.0053	4.22	0.75	0.72±0.03	-	2910	0	_		
SPT-CL J2303-5006	345.9992	-50.1107	6.08	0.50	0.64±0.02	2.92 ^{+0.46} _{-0.53}	36±7	0	_		
SPT-CL J2304-5149	346.0677	-51.8328	4.02	0.25	1.52±0.08	2.92 _{-0.53} 1.63 ^{+0.25} _{-0.35}	14±3	1	_		
SPT-CL J2304-5718	346.0967	-57.3094	5.71	0.25	0.92±0.03	2.52 ^{+0.42} _{-0.47}	30±7	0	_		
SPT-CL J2305-6524	346.3119	-65.4084	4.26	0.50	-	-0.47	-	Ö	_		
SPT-CL J2305-5130	346.4023	-51.5099	4.42	0.25	0.66±0.02	$2.30^{+0.39}_{-0.50}$	36±7	0	LOS		
SPT-CL J2306-5120	346.6130	-51.3421	7.56	0.25	1.17±0.07	2.86+0.36	36±4	1	_		
SPT-CL J2306-6505	346.7210	-65.0939	13.66	0.75	0.530	2.86 ^{+0.36} 2.86 ^{+0.36} 5.21 ^{+0.58} 5.21 ^{+0.58} 2.46 ^{+0.43}	121±11	0	_		
SPT-CL J2307-5440	346.7911	-54.6689	5.01	0.50	0.70±0.02	$2.46^{+0.43}_{-0.51}$	51±8	0	LOS		
SPT-CL J2307-6521	346.8902	-65.3591	4.33	0.50	0.53±0.01	2.39+0.38	53±12	0	_		
SPT-CL J2307-5412	346.9352	-54.2038	4.30	0.25	0.76±0.03	2.19 ^{+0.36} _{-0.48}	26±6	0	_		
SPT-CL J2308-5133	347.0677	-51.5547	4.17	0.25	-	-	-	0	-		
SPT-CL J2308-5413	347.1451	-54.2220	4.41	0.75	0.66±0.02	$2.30^{+0.38}_{-0.51}$	58±8	0	-		
SPT-CL J2308-5710	347.2451	-57.1794	5.34	0.50	0.40±0.02	2.88 ^{+0.47} _{-0.55}	23±5	0	-		
SPT-CL J2309-5301	347.3109	-53.0172	4.98	0.75	-	-	-	0	-		
SPT-CL J2309-5532	347.3208	-55.5467	4.01	3.00	-	-	-	0	-		
SPT-CL J2309-6048 SPT-CL J2310-4951	347.3955 347.5532	-60.8018 -49.8563	4.53 4.96	0.25 0.50	- 0.57±0.02	2.56 ^{+0.45} _{-0.53}	- 25±6	0 0	_		
SPT-CL J2310-4951 SPT-CL J2310-5919	347.5604	-49.6563 -59.3194	5.73	0.50	0.57±0.02 0.75±0.02	2.50 _{-0.53} 2.60 ^{+0.44}	42±7	0			
SPT-CL J2310-5919 SPT-CL J2311-6225			5.73	0.50	0.75±0.02 0.77±0.02	2.09_0.50			_		
	347.7535	-62.4243				2.47_0.50	37±7	0	1.00		
SPT-CL J2311-5549	347.8682	-55.8172	4.02	1.00	0.57±0.02	2.69 ^{+0.44} 2.69 ^{+0.50} 2.47 ^{+0.50} 2.25 ^{+0.35} 2.25 ^{+0.35} 3.68 ^{+0.52}	34±6	0	LOS		
SPT-CL J2311-5522 SPT-CL J2311-5411	347.8924 347.9754	-55.3691 -54.1902	6.81 4.03	0.75 1.75	0.203±0.008	3.08_0.62	38±6	0 0	_		
SPT-CL J2311-5411 SPT-CL J2311-5820	347.9754	-54.1902 -58.3402	4.49	0.25	0.84±0.03	2.18 ^{+0.35}	39±6	0	[
SPT-CL J2311-3020 SPT-CL J2312-6117	348.1590	-61.2917	4.49	0.25	0.84±0.03	2.10 _{-0.48} 2.22 ^{+0.36} -0.48 2.40 ^{+0.42} 2.40 _{-0.43}	30±6	0			
	JTU. 1JJU	01.2317	7.55	0.23	0.01±0.00	_ _0.48	00±0	U	1 -		

Table 1 — Continued

$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	0 0 0	Notes
SPT-CL J2313-6428 348.4898 -64.4746 4.06 1.25 - - - SPT-CL J2314-5747 348.5009 -57.7929 4.41 0.25 0.87±0.03 2.14 ^{+0.35} _{-0.46} 25±6	0	-
SPT-CL J2314-5747 348.5009 -57.7929 4.41 0.25 0.87±0.03 2.14 ^{+0.35} 25±6		
SPT-CL J2314-5747 348.5009 -57.7929 4.41 0.25 0.87±0.03 2.14 ^{-0.36} 25±6	0	-
CDT CL 19944 COOC 940 E440 CO C4E0 444 0.75		-
SPT-CL J2314-6036 348.5118 -60.6158 4.11 0.75 - - - - - - SPT-CL J2314-5553 348.5243 -55.8984 4.90 0.25 0.71±0.04 2.42 ^{+0.41} _{-0.51} 13±4	0 0	-
SPT-CL J2314-5245 348.5272 -52.7514 4.53 0.25 1.07±0.05 2.02±0.45 20±6	0	_
SPT-CL J2314-5245 348.5572 -52.7514 4.53 0.25 1.07±0.05 2.02±0.01 2.02±0.34 20±6 SPT-CL J2316-5535 349.0204 -55.5974 4.17 0.75 0.22±0.01 2.58±0.42 20±5	0	_
SPT-CL J2316-5534 349 1689 -55 5684 4 26 0 75 -	0	_
SPT-CL J2316-5907 349.1772 -59.1173 6.67 0.50 0.84±0.03 2.92 ^{+0.42} _{-0.50} 29±6 SPT-CL J2316-5027 349.1901 -50.4517 5.44 0.50 1.28±0.07 2.14 ^{+0.37} _{-0.42} 26±4	0	-
SPT-CL J2316-5027 349.1901 -50.4517 5.44 0.50 1.28±0.07 2.14 ^{+0.37} 26±4	1	-
SPT-CL J2316-5454 349.2141 -54.9028 10.16 0.75 0.37±0.01 4.55 ^{+0.54} _{-0.67} 52±7	0	-
SPI-CL J2316-5822 349.2216 -58.3675 4.19 0.50 - - -	0	-
SPT-CL J2317-4959 349.3195 -49.9977 5.02 0.50 1.05±0.03 2.17 ^{+0.39} 41±8	0	LOS
3F1-CL J2317-3632 349.4413 -36.6799 4.40 0.30 0.36±0.02 2.30 _{-0.51} 32±0	0	-
SPT-CL J2318-5058 349.5349 -50.9796 4.52 0.75 0.37±0.02 2.58 ^{+0.43} _{-0.55} 18±5	0	-
SPT-CL J2318-5617 349.7071 -56.2862 6.47 0.25 0.55±0.01 3.15 ^{+0.45} _{-0.55} 54±8 SPT-CL J2319-5404 349.8139 -54.0784 4.10 0.25 0.276±0.009 2.52 ^{+0.39} _{-0.55} 47±7	0	-
	0	-
SPT-CL J2319-6404 349.8674 -64.0730 4.17 0.25 SPT-CL J2319-5842 349.8745 -58.7103 7.11 0.50 0.30±0.01 3.67 ^{+0.50} _{-0.64} 28±6	0 0	-
SPT-CL J2319-5842 349.8745 -58.7103 7.11 0.50 0.30±0.01 3.67 ^{+0.50} _{-0.61} 28±6 SPT-CL J2320-6430 350.0423 -64.5146 5.24 0.25 0.88±0.05 2.39 ^{+0.42} _{-0.48} 12±5	0	-
SPT-CL J2320-0430	0	_
SPT-CL J2320-6443 350.0567 -64.7285 4.88 0.25 0.80±0.03 2.33 ^{+0.41} _{-0.50} 32±7	0	LOS
CDT CL 12220 5000 250 0755 50 1227 6 00 0 50 0 54 0 02 2 24+0.46 25 17	0	_
SPT-CL J2320-5806 350.0755 -58.1337 6.96 0.50 0.54±0.02 3.54 _{-0.56} 35±7 SPT-CL J2320-5233 350.1251 -52.5614 6.53 0.25 0.76±0.02 2.94 ^{+0.45} _{-0.50} 66±8	0	_
SPT-CL J2321-5120 350.2922 -51.3484 4.77 0.50 0.62±0.02 2.44 ^{+0.42} 42±7	0	-
SPT-CL J2321-5419 350.4100 -54.3229 4.08 0.25 0.80±0.04 2.10 ^{+0.34} 14±5	0	-
SPT-CL J2321-5558 350.4379 -55.9800 4.32 1.00 0.72±0.02 2.23 ^{+0.38} _{-0.48} 34±6	0	-
SPT-CL J2322-5708 350 5600 -57.1426 4 06 0 25 - - -	0	-
SPT-CL J2323-5752 350.8837 -57.8770 5.99 0.25 1.20±0.07 2.38 ^{+0.38} 27±4	1	-
3F1-CL J2323-3037 330.0906 -30.9393 4.03 0.25 0.41±0.01 2.41 _{-0.52} 43±7	0	-
SPT-CL J2323-6529 350.9346 -65.4885 4.28 0.25 0.88±0.03 2.11 ^{+0.33} _{-0.45} 33±7	0	-
SPT-CL J2323-6529 350.9346 -65.4885 4.28 0.25 0.88±0.03 2.11 _{-0.45} 33±7 SPT-CL J2323-5126 350.9521 -51.4432 4.36 0.25 0.71±0.03 2.25 ^{+0.37} _{-0.49} 16±5	0	-
SPT-CL J2323-5453 350.9582 -54.8953 4.01 0.25 0.83±0.02 2.06 ^{+0.33} 45±7	0	-
SPT-CL J2324-6354 351.2028 -63.9152 4.47 0.25 0.88±0.03 2.15 ^{+0.36} 23±6	0	-
SPT-CL J2325-5129 351.3398 -51.4984 4.28 3.00 - - - SPT-CL J2325-5815 351.3437 -58.2548 4.88 0.25 0.56±0.02 2.54±0.43 48±8	0 0	-
SPT-CL J2325-6215 351.3732 -62.2548 4.58 1.00 0.55±0.02 2.44 ^{+0.41} _{-0.53} 36±7	0	_
SPT_CI_12325_5116	0	_
SPT-CL J2325-5316 351.4227 -53.2811 5.13 0.25 0.37±0.01 2.80 ^{+0.48} _{-0.56} 32±6	0	LOS
SPT-CL J2326-5027 351.5127 -50.4534 4.28 1.25	0	-
SPT-CL J2326-5949 351 7325 -59 8215 4 09 0 25 - - -	0	-
SPT-CL J2327-5137 351.7831 -51.6244 5.03 1.00 0.36±0.01 2.78 ^{+0.48} _{-0.57} 50±7	0	-
SPT-CL J2327-6225 351.9144 -62.4196 6.11 0.75 0.190±0.007 3.39 ^{+0.52} 56±8	0	LOS
SPT-CL J2327-5800 351.9221 -58.0158 4.15 0.75	0	-
SPT-CL J2328-5456 352.0115 -54.9349 4.47 0.25 1.02±0.06 2.04 ^{+0.34} _{-0.45} 19±3 SPT-CL J2328-5533 352.1774 -55.5602 8.84 0.25 0.81±0.02 3.59 ^{+0.44} _{-0.45} 49±7	1 0	-
SPT-CL J2328-5533 352.1774 -55.5602 8.84 0.25 0.81±0.02 3.59 ^{+0.47} _{-0.55} 49±7 SPT-CL J2329-5630 352.2591 -56.5072 4.14 1.00 1.09±0.04 1.91 ^{+0.30} _{-0.42} 27±6	0	-
SPT-CL J2329-5849 352.2920 -58.8313 4.15 0.75	0	_
SPT-CL J2329-5230 352.3115 -52.5014 4.32 0.25 0.70±0.02 2.25 ^{+0.36} _{-0.40} 45±8	0	_
SPT-CL J2329-5230 352.3115 -52.5014 4.32 0.25 0.70±0.02 2.25 ^{+0.36} _{-0.49} 45±8 SPT-CL J2329-5831 352.4743 -58.5267 11.75 0.25 0.80±0.02 4.32 ^{+0.50} _{-0.61} 80±10	0	LOS
SPT-CL J2329-5958 352.4795 -59.9741 4.11 1.25	0	-
SPT-CL J2329-5030 352.4943 -50.5052 5.12 0.25	0	-
SPT-CL J2330-5035 352.6022 -50.5884 4.21 0.25 1.07±0.06 1.94±0.30 -0.43 22±4 SPT-CL J2330-5129 352.6727 -51.4974 4.19 0.25 0.55±0.02 2.31±0.51 -0.51 33±7 SPT-CL J2331-5737 352.8963 -57.6175 7.65 0.50 1.57±0.08 2.53±0.33 -0.39 20±3	1	LOS
SPT-CL J2330-5129 352.6727 -51.4974 4.19 0.25 0.55±0.02 2.31 ^{+0.37} _{-0.51} 33±7	0	-
	1 0	LOS
SPT-CL J2331-5131 352.9506 -51.5162 4.26 0.50	0	_
SPT-CL J2332-5358 353.1092 -53.9761 18.08 1.25 0.402 6.43 ^{+0.70} 83±10	0	_
SPT-CL J2332–5220 353.1376 –52.3389 4.23 0.75 0.45±0.01 2.42 ^{+0.38} 46±7	0	LOS
SPT-CL J2332-6112 353.1431 -61.2127 5.12 0.25 1.04±0.06 2.21 ^{+0.39} 35±4	1	-
SPT-CL 32331-5131 332.9300 -51.3162 4.26 0.30 -50.8707 19.91 0.50 0.576 6.42+0.68 / 0.84 126±12 SPT-CL J2332-5358 353.1092 -53.9761 18.08 1.25 0.402 6.43+0.70 / 0.83 83±10 SPT-CL J2332-5220 353.1376 -52.3389 4.23 0.75 0.45±0.01 2.42+0.54 / 0.38 46±7 SPT-CL J2332-6112 353.1431 -61.2127 5.12 0.25 1.04±0.06 2.21+0.46 / 0.49 35±4 SPT-CL J2333-6501 353.4703 -65.0234 7.35 0.25 0.57±0.02 3.44+0.44 / 0.56 / 0.54 54±8 SPT-CL J2334-5308 353.5206 -53.1403 5.20 0.25 1.17±0.07 2.14+0.44 / 0.44 39±4	0	-
SPT-CL J2334-5308 353.5206 -53.1403 5.20 0.25 1.17±0.07 2.14 ^{+0.37} 39±4	1	_

Table 1 — Continued

					Continued				
SPT ID	R.A. (J2000)	Decl. (J2000)	Β ε	est $\theta_{ m c}$	Redshift	M_{500c} $(10^{14}h_{70}^{-1}M_{\odot})$	λ	Spitzer	Notes
SPT-CL J2334-5131	353.5897	-51.5236	4.18	0.25	0.68±0.02	2.22+0.35	39±7	0	-
SPT-CL J2334-5939	353.6880	-59.6503	6.70	0.25	0.39±0.01	$3.42^{+0.48}_{-0.58}$	32±6	0	-
SPT-CL J2335-6135	353.7724	-61.5859	4.24	0.75	0.35±0.01	2 50+0.41	39±6	0	-
SPT-CL J2335-5434	353.8726	-54.5789	5.27	0.25	1.03±0.04	2.27 ^{+0.40}	28±6	0	-
SPT-CL J2336-5352	354.0267	-53.8779	6.05	0.25	0.51±0.01	3.04+0.46	67±9	0	-
SPT-CL J2336-5252	354.0842	-52.8714	5.88	0.25	1.24±0.07	2.30 ^{+0.38} _{-0.43}	22±3	1	-
SPT-CL J2337-5942	354.3517	-59.7066	40.30	0.25	0.775	8.98+0.96	165±14	0	-
SPT-CL J2337-5912	354.3998	-59.2052	5.79	0.75	0.60±0.02	2.85 ^{+0.46} _{-0.52}	49±7	0	-
SPT-CL J2338-6246	354.5253	-62.7783	5.26	0.25	1.13±0.06	2.19 ^{+0.39} _{-0.44}	25±4	1	-
SPT-CL J2338-6527	354.5501	-65.4620	4.28	1.00	-	2.15 ^{+0.34}	-	0	-
SPT-CL J2338-6311	354.6854	-63.1970	4.10	0.25	0.73±0.03	10.10	31±6	0	LOS
SPT-CL J2339-5550	354.8613	-55.8347	5.08	0.25	0.39±0.01 0.98±0.04	2.76 ^{+0.46} 2.09 ^{+0.36} 2.09 ^{+0.46}	37±6	0	-
SPT-CL J2339-5008	354.9460	-50.1387	4.56	0.50		3.21 ^{+0.45}	22±6	0	-
SPT-CL J2339-6332 SPT-CL J2340-5719	354.9715	-63.5348 57.2214	7.18	0.25 0.25	0.71±0.02 0.82±0.03	1 '8.33	70±9	0 0	-
SPT-CL J2340-5719 SPT-CL J2340-5251	355.0673 355.1293	-57.3314 -52.8618	4.05 4.02	0.25	1.60±0.08	2.07 ^{+0.32} _{-0.46} 1.58 ^{+0.24} _{-0.34}	20±6 14±3	1	LL
SPT-CL J2340-5251 SPT-CL J2340-6045	355.2486	-60.7505	4.02	0.25	0.50±0.06	2.35 ^{+0.38}		0	
SPT-CL J2340-6045 SPT-CL J2341-5716	355.2636	-60.7505 -57.2719	5.95	1.00	0.50±0.02 0.44±0.01	3.07 ^{+0.49}	22±5 60±8	0	LOS
SPT-CL J2341-5141	355.2955	-51.6939	4.63	0.25	0.44±0.01 -	3.07 -0.56	- 0010	0	-
SPT-CL J2341-5119	355.3000	-51.3306	20.01	0.25	1.003	5.57 ^{+0.61}	51±0	0	_
SPT-CL J2341-5221	355.3005	-52.3619	4.09	0.25	0.83±0.04	5.57 ^{+0.61} -0.73 2.09 ^{+0.32} -0.46	18±5	0	_
SPT-CL J2341-5724	355.3506	-57.4156	14.32	0.50	1.259	, X-79	53±0	0	_
SPT-CL J2341-5639	355.4417	-56.6655	4.08	0.75	0.48±0.02	4.18 ^{+0.47} -0.57 2.36 ^{+0.37} -0.51	38±6	0	RC
SPT-CL J2341-5138	355.4460	-51.6389	4.71	0.25	0.55±0.02	דדיטיטא כי	36±6	0	_
SPT-CL J2341-5308	355.4778	-53.1371	5.34	0.75	0.54±0.01	$2.74^{+0.46}_{-0.54}$	59±8	0	RC
SPT-CL J2342-6030	355.5051	-60.5084	4.66	0.25	0.82±0.03	2.24+0.39	25±6	0	-
SPT-CL J2342-5621	355.6175	-56.3593	4.11	1.00	0.42±0.01	2.24 _{-0.49} 2.40 ^{+0.39} 2.53	44±7	0	-
SPT-CL J2342-5715	355.6404	-57.2509	5.49	0.50	0.83±0.03		27±6	0	-
SPT-CL J2342-6041	355.6921	-60.6995	5.10	0.25	0.80±0.03	2 42*0.41	31±7	0	-
SPT-CL J2342-5411	355.6928	-54.1864	12.54	0.25	1.075	4.12 ^{+0.45} _{-0.57}	32±0	0	-
SPT-CL J2342-5522	355.7060	-55.3771	4.58	2.50	-	-	-	0	-
SPT-CL J2343-5158	355.8285	-51.9801	4.22	0.50	0 80±0 03 -	2 70+0.44	36±7	0 0	-
SPT-CL J2343-5023 SPT-CL J2344-6251	355.8355	-50.3999 -62.8611	6.43 4.09	0.25 0.25	0.89±0.03 1.49±0.07	2.78 ^{+0.44} 1.65 ^{+0.25}	27±6	0	-
SPT-CL J2344-6251 SPT-CL J2344-6000	356.0183 356.0720	-60.0131	4.09	0.25	0.50±0.01	2.24+0.37	50±7	0	_
SPT-CL J2344-0000 SPT-CL J2344-5656	356.0758	-56.9349	4.10	0.50	1.60±0.01	1 50+0.23	13±3	1	LOS,LL
SPT-CL J2344-5233	356.0879	-52.5562	4.12	1.25	-	1.30 _{-0.34}	-	Ö	-
SPT-CL J2344-6004	356.2391	-60.0740	4.58	0.25	1.31±0.07	1.86 ^{+0.31} 5.63 ^{+0.59}	16±5	0	_
SPT-CL J2345-6405	356.2500	-64.0991	20.43	0.25	1.01±0.02	5.63 ^{+0.59} _{-0.72}	97±11	0	LOS
SPT-CL J2345-6218	356.3397	-62.3035	4.52	1.75	-	-	-	0	-
SPT-CL J2346-5740	356.6714	-57.6774	4.72	0.25	1.17±0.07	$2.00^{+0.35}_{-0.44}$	23±3	1	LOS
SPT-CL J2347-6008	356.8653	-60.1489	4.03	0.50	-	- 40+0.40	-	0	-
SPT-CL J2347-5532	356.8878	-55.5455	4.24	0.25	0.37±0.02	$\begin{array}{c} 2.48^{+0.40}_{-0.54} \\ 2.36^{+0.38}_{-0.52} \\ 2.54^{+0.44}_{-0.53} \end{array}$	22±5	0	-
SPT-CL J2347-6120	356.8988	-61.3361	4.19	0.50	0.50±0.02	2.36 _{-0.52}	27±6	0	-
SPT-CL J2347-5128 SPT-CL J2347-6530	356.9049 356.9071	-51.4828 -65.5137	4.84 4.53	0.75 0.25	0.55±0.02	2.54_0.53	20±5	0 0	_
SPT-CL J2347-5158	356.9431	-51.9802	4.23	0.25	0.84±0.03	2.10 ^{+0.33} _{-0.46}	38±7	0	_
SPT-CL J2348-5601	357.0260	-56.0316	4.18	0.50	0.42±0.01	2.44 ^{+0.39} 2.44 ^{-0.54}	39±7	0	_
SPT-CL J2348-5715	357.1200	-57.2629	4.40	0.25	0.60±0.02	2.33+0.40	28±6	0	LOS
SPT-CL J2349-5113	357.3861	-51.2285	5.60	0.75	0.42±0.01	2.94+0.50	58±9	0	LOS
SPT-CL J2349-6506	357.4426	-65.1067	4.77	0.25	0.57±0.02	2.33 ^{+0.40} 2.33 ^{+0.40} 2.94 ^{+0.57} 2.94 ^{+0.44} 2.50 ^{+0.44} 2.51 ^{+0.42} 2.51 ^{+0.42}	35±6	0	
SPT-CL J2349-5710	357.4939	-57.1692	4.66	0.50	0.51±0.02	2.51+0.42	19±5	0	_
SPT-CL J2350-5635	357.6085	-56.5973	4.79	0.25	1.33±0.07	1.90+0.35	13±3	1	_
SPT-CL J2350-5434	357.6180	-54.5812	4.11	0.25	0.55±0.02	1.90 ^{+0.35} 1.90 ^{+0.35} 2.29 ^{+0.37} 4.64 ^{+0.52}	27±5	0	_
SPT-CL J2350-5301	357.7247	-53.0222	11.42	0.50	0.54±0.01	4.64+0.52	68±9	0	_
SPT-CL J2351-5240	357.8173	-52.6769	4.36	0.25	-	-	-	Ö	-
SPT-CL J2351-5005	357.8295	-50.0952	5.03	0.25	0.56±0.02	2.59 ^{+0.45} 5.14 ^{+0.58} 5.14 ^{-0.71}	22±6	0	LOS
SPT-CL J2351-5452	357.9053	-54.8827	12.33	0.50	0.384	5.14 ^{+0.58}	102±10	0	RC
SPT-CL J2352-5755	358.1347	-57.9245	4.28	0.75	-	-		0	-
SPT-CL J2352-6122	358.1420	-61.3692	4.04	0.25	1.26±0.07	1.77 ^{+0.27} _{-0.39}	18±5	0	-

Table 1 — Continued

SPT ID	R.A. (J2000)	Decl. (J2000)	Be ξ	st $ heta_{ m c}$	Redshift	$M_{500c} (10^{14} h_{70}^{-1} M_{\odot})$	λ	Spitzer	Notes
SPT-CL J2352-6134	358.1941	-61.5687	11.81	0.25	0.92±0.02	4.18+0.48	72±9	0	_
SPT-CL J2352-6249	358.2094	-62.8287	4.55	0.25	0.93±0.04	2.11+0.37	24±6	0	LOS
SPT-CL J2352-5251	358.2117	-52.8648	4.19	1.00	0.48±0.02	0.00+0.40	38±7	0	-
SPT-CL J2353-6517	358.3013	-65.2921	4.91	0.25	0.68±0.02	0.40+0.43	40±8	0	-
SPT-CL J2354-5106	358.5422	-51.1048	6.62	0.50	0.33±0.01	2.43 _{-0.52} 3.46 ^{+0.50} 3.46 _{-0.59}	29±6	0	-
SPT-CL J2354-6518	358.5952	-65.3084	4.16	1.00	-	-	-	0	-
SPT-CL J2354-5633	358.7207	-56.5514	9.76	0.50	0.56±0.01	4.18 ^{+0.50} _{-0.62}	90±11	0	-
SPT-CL J2355-5156	358.8514	-51.9483	8.43	0.25	0.75±0.02	3.55 ^{+0.44} _{-0.55}	48±7	0	LOS
SPT-CL J2355-5029	358.8681	-50.4945	4.22	0.25	0.93±0.04	2.05 ^{+0.31} -0.45 1.79 ^{+0.28}	24±5	0	-
SPT-CL J2355-5515	358.8715	-55.2539	4.06	0.25	1.24±0.07	1.79 ^{+0.28} _{-0.40}	26±4	1	-
SPT-CL J2355-5259	358.9368	-52.9843	6.31	0.25	0.77±0.02	2.87 ^{+0.45}	58±8	0	-
SPT-CL J2355-5055	358.9492	-50.9297	10.36	0.75	0.320	2.87 _{-0.50} 4.70 ^{+0.54} -0.67	85±10	0	-
SPT-CL J2356-4957	359.0792	-49.9554	4.00	1.50	-	-	-	0	-
SPT-CL J2357-5421	359.2628	-54.3604	6.28	0.25	0.89±0.03	2.74 ^{+0.45}	25±6	0	LOS
SPT-CL J2358-5229	359.5293	-52.4846	7.15	0.25	0.67±0.01	2.74 ^{+0.45} 3.25 ^{+0.45} 3.25 ^{+0.45}	79±9	0	LOS
SPT-CL J2358-6129	359.7146	-61.4950	11.97	0.50	0.405±0.010	5.00+0.59	67±8	0	-
SPT-CL J2359-5023	359.7829	-50.3958	4.26	0.50	1.33±0.07	1.78 ^{+0.29} -0.40	11±3	1	-
SPT-CL J2359-5010	359.9318	-50.1703	10.86	0.50	0.775	4.14+0.50	72±9	0	_

Note. — Galaxy cluster candidates detected at $\xi > 4$ in the SPTpol 500d survey. We first report properties derived solely from the SPT data, namely the candidate names, positions, detection-significances (ξ) and corresponding β -model core-radii at the cortical $||\Omega||$ core-radii at which the candidates were detected. Next, using the optical/IR confirmation processes discussed in Section 5, we provide redshifts of associated galaxy counterparts to the SZ detections for systems with f_{cont} < 0.2. These redshifts are combined with the ξ values to estimate cluster masses (Section 6) for systems with optical/IR contamination fractions such that the overall sample is expected to be 94% pure. We provide the optical richness (λ) and mark whether these quantities were derived with our MCMF or Spitzer analysis. Finally we flag LL confirmed clusters with redshift lowerlimits of z = 1.6 (see Section 5.2), **RC** clusters in which potentially $\Delta \xi > 2$ is filled in by synchrotron emission (see Section 7) or LOS for which a secondary optical structure along the line-of-sight met our confirmation criteria. The richnesses, redshifts, and contamination fractions of both primary and secondary systems are reported in the online version of the cluster sample. provide in the online file information on line-of-sight structures for unconfirmed clusters.

This paper was built using the Open Journal of Astrophysics LaTeX template. The OJA is a journal which provides fast and easy peer review for new papers in the astro-ph section of the arXiv, making the reviewing process simpler for authors and referees alike. Learn more at http://astro.theoj.org.