



OPEN Proposal for a Lorenz qubit

Michael R. Geller

Nonlinear qubit master equations have recently been shown to exhibit rich dynamical phenomena such as period doubling, Hopf bifurcation, and strange attractors usually associated with classical nonlinear systems. Here we investigate nonlinear qubit models that support tunable Lorenz attractors. A Lorenz qubit could be realized experimentally by combining qubit torsion, generated by real or simulated mean field dynamics, with linear amplification and dissipation. This would extend engineered Lorenz systems to the quantum regime, allowing for their direct experimental study and possible application to quantum information processing.

Several recent papers^{1–7} have considered nonlinear generalizations of the Gorini-Kossakowski-Sudarshan-Lindblad (GKSL) master equation^{8,9} for qudits. The superoperators resulting from these evolutions each take the form of a positive trace-preserving (PTP) channel^{10,11} $X \mapsto \phi(X)/\text{tr}[\phi(X)]$, with X a density matrix and ϕ a positive map. The positivity of this *normalized* PTP channel follows from the positivity of ϕ and $\text{tr}[\phi(X)] > 0$. Its trace preservation property is actually a trace fixing one, but these are physically equivalent when applied to normalized initial states. Kowalski and Rembieliński¹, and also Rembieliński and Caban³, considered cases with linear ϕ and $\text{tr}[\phi(X)] \neq 1$, extending Gisin's 1981 model¹² to mixed states. We call these channels *nonlinear in normalization only* (NINO) to emphasize that the nonlinearity in this case serves only to conserve trace. We might think of NINO channels as being “mildly” nonlinear. In particular, they satisfy a convex quasilinear property¹³, preventing superluminal signaling^{14–18}. The main difference between linear completely positive trace-preserving (CPTP) and NINO channels are that the generators of linear CPTP evolution are negative definite, leading to strictly nonexpansive dynamics, whereas NINO channels support non-CP^{19–22} and entropy decreasing¹ processes that amplify the Bloch vector^{1,6}. Hence we can interpret the NINO master equation as extending the GKSL equation to non-Hermitian Hamiltonians. Fernengel and Drossel² studied cases where ϕ is nonlinear and $\text{tr}[\phi(X)] = 1$, a family of state-dependent CPTP channels obtained by adding state-dependence to a Hamiltonian and set of Lindblad jump operators. This is a stronger form of nonlinearity, supporting rich dynamical phenomena such as such Hopf bifurcations and strange attractors usually associated with classical nonlinear systems². State-dependent CPTP channels also support Bloch-ball *torsion*. Torsion can be created from the product of an $\text{SO}(3)$ rotation generator J_μ with the projection of the Bloch vector along the twist axis. Abrams and Lloyd²³ and Childs and Young²⁴ investigated state discrimination with z -axis torsion. Klobus et al.⁷ observed Feigenbaum's universal period doubling in a mean field model simulating torsion. Torsion also arises in a qubit friendly extension²⁵ of a rigorous duality between nonlinear mean field theory and the BBGKY hierarchy for n interacting bosons in the $n \rightarrow \infty$ limit^{26–30}. Many of these nonlinear models come from mean field theory.

Methods

In this paper we investigate qubit PTP channels with both nonlinear ϕ and $\text{tr}[\phi(X)] \neq 1$ that support generalized Lorenz attractors. The first version, which we call *Lor63*, implements Lorenz's 1963 model³¹

$$\frac{dx}{dt} = \sigma(y - x), \quad (1)$$

$$\frac{dy}{dt} = \rho x - y - gxz, \quad (2)$$

$$\frac{dz}{dt} = -\beta z + gxy, \quad (3)$$

where $\mathbf{r} = (x, y, z) = \text{tr}(X\sigma)$ is the Bloch vector. However here we increase the nonlinearity by a factor of $g \gg 1$ to shrink the attractor sufficiently as to contain it within the Bloch sphere. The master equation for the *Lor63* qubit in the Pauli basis is

$$\frac{dX}{dt} = \frac{\sigma^a}{2} \left(\frac{dr^a}{dt} \right), \quad \frac{dr^a}{dt} = \text{tr} \left(\frac{dX}{dt} \sigma^a \right) = G^{ab}(\mathbf{r}) r^b = (L + g \mathbf{J}_x)^{ab} r^b, \quad (4)$$

where $a, b \in \{1, 2, 3\}$ and

$$L = \begin{pmatrix} -\sigma & \sigma & 0 \\ \rho & -1 & 0 \\ 0 & 0 & -\beta \end{pmatrix} = L_+ + L_-, \quad L_+ = \begin{pmatrix} -\sigma & \frac{\rho+\sigma}{2} & 0 \\ \frac{\rho+\sigma}{2} & -1 & 0 \\ 0 & 0 & -\beta \end{pmatrix} = \left(\frac{\rho+\sigma}{2} \right) \lambda_1 - D, \quad (5)$$

$$L_- = \left(\frac{\rho-\sigma}{2} \right) J_z, \quad \lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad D = \begin{pmatrix} \sigma & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \beta \end{pmatrix}, \quad J_x = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad J_z = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (6)$$

$X \in \mathbb{C}^{2 \times 2}$ is a Hermitian positive-semidefinite matrix with unit trace. Model parameters ρ, σ, β, g are given in Table 1. The nonlinear generator $G^{ab}(\mathbf{r})$ is a 3×3 real matrix that depends on the Bloch vector \mathbf{r} . We decompose it into a linear (\mathbf{r} independent) operator L plus x -axis torsion. The J 's are SO(3) generators: $(J_a)_{bc} = -\varepsilon_{abc}$ with ε the Levi-Civita symbol. L is decomposed into symmetric and antisymmetric parts implementing a non-Hermitian Hamiltonian iL . λ_1 is a Gell-Mann matrix. Note that λ_1 has a positive eigenvalue corresponding to an amplifying and entropy *decreasing* non-CP process¹. Techniques for constructing Gell-Mann matrices and other symmetric generators from jump operators are given in⁶.

Result and discussion

The *Lor63* qubit is simulated in Fig. 1. The blue points indicate random initial conditions. Trajectories rapidly approach one of the two disc-shaped sets (pink or cyan) and bounce back and forth between them in an unpredictable manner, mirroring the aperiodic reversals of the Malkus waterwheel lying in its Fourier representation³².

A Lorenz-like attractor can also be created from the z -axis torsion coming from the Gross-Pitaevskii (GP) equation^{24,33,34}, leading to an aesthetic attracting set shown in Fig. 2. We call this channel the *GP butterfly*. The *GP butterfly* qubit has an especially simple master equation:

$$\frac{dr^a}{dt} = \text{tr} \left(\frac{dX}{dt} \sigma^a \right) = G^{ab}(\mathbf{r}) r^b = (m \lambda_4 + g z J_z)^{ab} r^b, \quad (7)$$

where $m = 10$, $g = 40$, and

	Original	Here
ρ	28	28
σ	10	10
β	8/3	8/3
g	1	80

Table 1. *Lor63* model parameters.




Figure 1. Bloch ball dynamics of the *Lor63* qubit. The faint yellow wireframe shows the Bloch sphere. Equator states $|\pm\rangle = 2^{-1/2}(|0\rangle \pm |1\rangle)$ and $|\pm i\rangle = 2^{-1/2}(|0\rangle \pm i|1\rangle)$ are also indicated with black dots and cyan lines. The model parameters used in the simulation are given in Table 1.




Figure 2. Bloch ball dynamics of the *GP butterfly* qubit. Blue dots indicate random initial conditions.

$$\lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \quad (8)$$

is another Gell–Mann matrix. The symmetric generator λ_4 can be implemented with Lindblad jump operators⁶.

Conclusion

In conclusion, we have proposed nonlinear PTP channels for the generation of Lorenz-like attractors in the Bloch ball. Despite its early prominence the Lorenz system defied rigorous analysis until rather recently when, in 2002, Tucker³⁵ established the existence of a strange attractor. Classical electrical circuits have been used to implement the Lorenz attractor and other chaotic and hyperchaotic attractors^{36–39}, which might find cryptographic application^{40,41}. It is tempting to speculate that chaotic attractors will find application in quantum technology as well. However it is important to recognize the very large nonlinear coupling strengths required, making experimental realization especially challenging.

Data availability

All data generated or analysed during this study are included in this published article.

Received: 22 November 2022; Accepted: 17 August 2023

Published online: 29 August 2023

References

1. Kowalski, K. & Rembieliński, J. Integrable nonlinear evolution of the qubit. *Ann. Phys.* **411**, 167955. <https://doi.org/10.1016/j.aop.2019.167955> (2019).
2. Fernengel, B. & Drossel, B. Bifurcations and chaos in nonlinear Lindblad equations. *J. Phys. A: Math. Theor.* **53**, 385701. <https://doi.org/10.1088/1751-8121/abaa85> (2020).
3. Rembieliński, J. & Caban, P. Nonlinear extension of the quantum dynamical semigroup. *Quantum* **5**, 420 (2020). <https://doi.org/10.22331/q-2021-03-23-420>
4. Kowalski, K. Linear and integrable nonlinear evolution of the qutrit. *Quant. Inf. Proc.* **19**, 145. <https://doi.org/10.1007/s11128-020-02640-6> (2020).
5. Buks, E. & Schwartz, D. Stability of the Grabert master equation. *Phys. Rev. A* **103**, 052217. <https://doi.org/10.1103/PhysRevA.103.052217> (2021).
6. Geller, M.R. Fast quantum state discrimination with nonlinear PTP channels. *Adv. Quantum Technol.* **2200156** (2023). [arXiv: 2111.05977](https://arxiv.org/abs/2111.05977). <https://doi.org/10.1002/qute.202200156>.
7. Kłobus, W., Kurzyński, P., Kuś, M., Laskowski, W., Przybycień, R. & Życzkowski, K. Transition from order to chaos in reduced quantum dynamics. [arXiv: 2111.13477](https://arxiv.org/abs/2111.13477).
8. Gorini, V., Kossakowski, A. & Sudarshan, E. C. G. Completely positive dynamical semigroups of N-level systems. *J. Math. Phys.* **17**, 821. <https://doi.org/10.1063/1.522979> (1976).
9. Lindblad, G. On the generators of quantum dynamical semigroups. *Commun. Math. Phys.* **48**, 119 (1976).
10. Sudarshan, E. C. G., Mathews, P. M. & Rau, J. Stochastic dynamics of quantum-mechanical systems. *Phys. Rev.* **121**, 920. <https://doi.org/10.1103/PhysRev.121.920> (1961).
11. Kraus, K. General state changes in quantum theory. *Ann. Phys.* **64**, 311. [https://doi.org/10.1016/0003-4916\(71\)90108-4](https://doi.org/10.1016/0003-4916(71)90108-4) (1971).
12. Gisin, N. A simple nonlinear dissipative quantum evolution equation. *J. Phys. A: Math. Gen.* **14**, 2259 (1981).
13. Rembieliński, J. & Caban, P. Nonlinear evolution and signaling. *Phys. Rev. Res.* **2**, 012027. <https://doi.org/10.1103/PhysRevRes.2.012027> (2020).
14. Gisin, N. Weinberg's nonlinear quantum mechanics and supraluminal communication. *Phys. Lett. A* **143**, 1 (1990).
15. Polchinski, J. Weinberg's nonlinear quantum mechanics and the Einstein–Podolsky–Rosen paradox. *Phys. Rev. Lett.* **66**, 397. <https://doi.org/10.1103/PhysRevLett.66.397> (1991).
16. Czachor, M. Mobility and non-separability. *Found. Phys. Lett.* **4**, 351 (1991).
17. Gisin, N. & Rigo, M. Relevant and irrelevant nonlinear Schrödinger equations. *J. Phys. A* **28**, 7375. <https://doi.org/10.1088/0305-4470/28/24/030> (1995).
18. Kent, A. Nonlinearity without superluminality. *Phys. Rev. A* **72**, 012108. <https://doi.org/10.1103/PhysRevA.72.012108> (2005).

19. Pechukas, P. Reduced dynamics need not be completely positive. *Phys. Rev. Lett.* **73**, 1060. <https://doi.org/10.1103/PhysRevLett.73.1060> (1994).
20. Shaji, A. & Sudarshan, E. C. G. Who's afraid of not completely positive maps?. *Phys. Lett. A* **341**, 48 (2005).
21. Carteret, H. A., Terno, D. R. & Życzkowski, K. Dynamics beyond completely positive maps: Some properties and applications. *Phys. Rev. A* **77**, 042113. <https://doi.org/10.1103/PhysRevA.77.042113> (2008).
22. Dominy, J. M., Shabani, A. & Lidar, D. A. A general framework for complete positivity. *Quantum Inf. Process.* **15**, 465. <https://doi.org/10.1007/s11128-015-1148-0> (2016).
23. Abrams, D. S. & Lloyd, S. Nonlinear quantum mechanics implies polynomial-time solution for NP-complete and #P problems. *Phys. Rev. Lett.* **81**, 3992. <https://doi.org/10.1103/PhysRevLett.81.3992> (1998).
24. Childs, A. M. & Young, J. Optimal state discrimination and unstructured search in nonlinear quantum mechanics. *Phys. Rev. A* **93**, 022314. <https://doi.org/10.1103/PhysRevA.93.022314> (2016).
25. Geller, M.R. *The Universe as a Nonlinear Quantum Simulation*. arXiv: [2112.09005](https://arxiv.org/abs/2112.09005)
26. Nachtergaelie, B., Ogata, Y. & Sims, R. Propagation of correlations in quantum lattice systems. *J. Stat. Phys.* **124**, 1. <https://doi.org/10.1007/s10955-006-9143-6> (2006).
27. Fröhlich, J., Graffi, S. & Schwarz, S. Mean-field and classical limit of many-body Schrödinger dynamics for bosons. *Commun. Math. Phys.* **271**, 681. <https://doi.org/10.1007/s00220-007-0207-5> (2007).
28. Rodnianski, I. & Schlein, B. Quantum fluctuations and rate of convergence towards mean field dynamics. *Commun. Math. Phys.* **291**, 31. <https://doi.org/10.1007/s00220-009-0867-4> (2009).
29. Erdős, L. & Schlein, B. Quantum dynamics with mean field interactions: A new approach. *J. Stat. Phys.* **134**, 859. <https://doi.org/10.1007/s10955-008-9570-7> (2009).
30. Gokler, C. *Mean Field Limit for Many-particle Interactions*. arXiv: [2006.05486](https://arxiv.org/abs/2006.05486)
31. Lorenz, E. N. Deterministic nonperiodic flow. *J. Atmos. Sci.* **20**, 130 (1963).
32. Sparrow, C. *The Lorenz Equations: Bifurcations, Chaos, and Strange Attractors* (Springer, 1982).
33. Meyer, D. A. & Wong, T. G. Nonlinear quantum search using the Gross–Pitaevskii equation. *New J. Phys.* **15**, 063014 (2013).
34. Meyer, D. A. & Wong, T. G. Quantum search with general nonlinearities. *Phys. Rev. A* **89**, 012312. <https://doi.org/10.1088/1367-2630/15/6/063014> (2014).
35. Tucker, W. A rigorous ODE solver and Smale's 14th problem. *Found. Comput. Math.* **2**, 52 (2002).
36. Cuomo, K. M. & Oppenheim, A. V. Circuit implementation of synchronized chaos with applications to communications. *Phys. Rev. Lett.* **71**, 65. <https://doi.org/10.1103/PhysRevLett.71.65> (1993).
37. Qi, G., van Wyk, M. A., van Wyk, B. J. & Chen, G. A new hyperchaotic system and its circuit implementation. *Chaos Solitons Fractals* **40**, 2544 (2009).
38. Liu, L., Du, C., Zhang, X., Li, J. & Shi, S. Dynamics and entropy analysis for a new 4-D hyperchaotic system with coexisting hidden attractors. *Entropy* **21**, 287. <https://doi.org/10.3390/e21030287> (2019).
39. Tian, H., Wang, Z., Zhang, P., Chen, M. & Wang, Y. Dynamic analysis and robust control of a chaotic system with hidden attractor. *Complexity* **2021**, 8865522 (2021).
40. Al-Maadeed, T.A., Hussain, I., Anees, A. & Mustafa, M.T. An image encryption algorithm based on chaotic Lorenz system and novel primitive polynomial S-boxes. arXiv: [2006.11847](https://arxiv.org/abs/2006.11847).
41. Mafco, A.G., Martinez, A.S. & Bruno, O.M. Fast, parallel, and secure cryptography algorithm using Lorenz's attractor. *Int. J. Mod. Phys. C* **21**, 365 (2010). arXiv: [1201.3114](https://arxiv.org/abs/1201.3114). <https://doi.org/10.1142/S0129183110015166>.

Author contributions

M.G. conceived of the project and wrote the manuscript.

Funding

This study was funded by National Science Foundation (No. DGE-2152159).

Competing interests

The author declares no competing interests.

Additional information

Correspondence and requests for materials should be addressed to M.R.G.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023