



Gorenstein Braid Cones and Crepant Resolutions

Joshua Hallam¹ · John Machacek²

Received: 14 January 2022 / Revised: 18 May 2023 / Accepted: 2 July 2023 /

Published online: 5 October 2023

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2023

Abstract

To any poset P , we associate a convex cone called a braid cone. We also associate a fan and study the toric varieties the cone and fan define. The fan always defines a smooth toric variety X_P , while the toric variety U_P of the cone may be singular. We show that $X_P \dashrightarrow U_P$ is a crepant resolution of singularities if and only if P is bounded. Next, we aim to determine when U_P is \mathbb{Q} -Gorenstein or Gorenstein. We prove that whether or not U_P is \mathbb{Q} -Gorenstein or Gorenstein depends only on the biconnected components of the Hasse diagram of P . In the case that P has a minimum or maximum element, we show that the Gorenstein property of U_P is completely determined by the Möbius function of P . We also provide a recursive method that determines if U_P is \mathbb{Q} -Gorenstein or Gorenstein in this case. We conjecture that U_P is Gorenstein if and only if it is \mathbb{Q} -Gorenstein. We verify this conjecture for posets of length 1 and also for posets with a minimum or maximum element.

Keywords Toric variety · Gorenstein variety · Braid cone · Poset · Möbius function

Mathematics Subject Classification Primary 06A07 · Secondary 05E14 · 14M25 · 52B05

Editor in Charge: János Pach

John Machacek acknowledges support of NSF grant DMS-2039316.

Joshua Hallam
joshua.hallam@lmu.edu

John Machacek
johnmach@uoregon.edu

¹ Department of Mathematics, Loyola Marymount University, Los Angeles, CA 90045, USA

² Department of Mathematics, University of Oregon, Eugene, OR 97403, USA

1 Introduction

The correspondence between convex cones and toric varieties is well established. We consider posets as an additional combinatorial layer. Every poset naturally gives rise to a cone as well as a fan that refines the cone. The cones that arise here are the unions of certain chambers of the braid arrangement and thus called *braid cones*. We use the combinatorics of the underlying posets and their Hasse diagrams to study affine toric varieties associated to braid cones. We also study a resolution of singularities for such a toric variety coming from another toric variety associated to a fan.

Our main focus is to be able to efficiently decide when the variety associated to a braid cone is Gorenstein and when the aforementioned resolution is crepant. Since we are working with affine toric varieties, this algebro-geometric decision problem ends up being equivalent to the discrete geometry problem of deciding if there exists a hyperplane containing certain lattice points. We find and use properties of posets which are equivalent to the desired resolutions of singularities being crepant and the Gorenstein property.

Crepant resolutions and the Gorenstein property are important in studying singularities in toric geometry (see e.g. [4]). Crepant resolutions have no discrepancy in the canonical class and play a role in the minimal model program. The class of Gorenstein varieties consists of varieties that are at worse “not too singular”. It properly sits between the class of smooth varieties and Cohen–Macaulay varieties. That is,

$$\text{smooth varieties} \subsetneq \text{Gorenstein varieties} \subsetneq \text{Cohen–Macaulay varieties}.$$

All toric varieties coming from fans are normal and hence by [11] are Cohen–Macaulay. Moreover, in [18], a classification¹ of smoothness for toric varieties arising from braid cones is given.

Recent works in a similar spirit to what we study here include [13, 15], and [14]. The smoothness of generic torus orbits inside Schubert varieties is studied in [15], and the maximal cones involved are each made from chambers of the braid arrangement fan. In [10] the authors give graph-theoretic characterizations of Gorensteinness for the toric variety associated to the base polytope of a graphic matroid arising from (simple) graphs. The work in [13] does the same in the case of graphic matroids coming from multigraphs. In [14], a matroid-theoretic characterization for Gorensteinness of the varieties associated to the base polytopes of general matroids is given. Also, we refer the reader to [25] which studies singularities for another construction from posets where one obtains a projective toric variety from a distributive lattice.

Cones corresponding to posets have been considered in a generalized setting of root systems for Coxeter groups under the names *parsets* [19] and *Coxeter cones* [24]. The case of posets is the Type *A* case. The cones we consider recently appeared in [16] where star subdivisions which correspond to toric blow-ups were studied. Recent work on closely related cones include [5], which focuses on their Whitney numbers.

¹ This characterization can be obtained by combining Proposition 3.5 (9), which characterizes which cones are simplicial, along Corollary 3.10 of [18], which shows that simplicial implies smoothness. Simplicial is a necessary condition for smoothness.

In the remainder of this section, we review posets and braid cones. Afterwards, we discuss the necessary background on toric varieties. In Sect. 2 we introduce labelings of posets which will determine when a poset defines a \mathbb{Q} -Gorenstein or Gorenstein toric variety as well as when a particular fan associated to a poset gives a crepant resolution of singularities. Our main result on crepant resolutions is in Sect. 3. In Theorem 3.3, we show there is a certain crepant resolution of singularities respecting Weyl chambers precisely for toric varieties coming from bounded posets (i.e. posets with both a minimum and maximum). In Sect. 4 we deal with the Gorenstein property and find, in Theorem 4.3, that Gorensteinness depends only on the biconnected components of the Hasse diagram. We consider posets with a minimum or a maximum element in Sect. 5. There, we are able to characterize when such posets give rise to a Gorenstein variety (Theorem 5.19) in a way which gives a recursive algorithm (Algorithm 5.20) to do this. In terms of parameterized complexity, this algorithm is an XP-algorithm (i.e. “slicewise polynomial”) implying it runs efficiently for small values of the parameter. The parameter used is a new parameter on posets which depends only on the minimal elements as well as those elements which cover only minimal elements. The Möbius function plays a significant role in developing the ideas in this section. We then look at posets of length 1 in Sect. 6 and conclude with a discussion on some open problems in Sect. 7.

Our main result on crepant resolutions is stated as follows

Theorem (Theorem 3.3) *Let P be a poset. $X_P \dashrightarrow U_P$ is crepant if and only if P is bounded.*

Here U_P is an affine toric variety coming from a braid cone defined in the next subsection while X_P is an abstract toric variety coming from the fan obtain by subdividing the aforementioned cone with Weyl chambers. This result implies bounded posets give Gorenstein toric varieties. Table 1 contains a summary of our results giving various sufficient conditions for a toric variety from a braid cone to be Gorenstein. In the table, when we say a poset is Gorenstein, we mean the corresponding variety is Gorenstein. Moreover, the “also necessary condition” column refers to if the conditions are also necessary when restricting to the type of poset described in that row. In all the cases described in the table, Gorenstein can also be replaced with \mathbb{Q} -Gorenstein without changing the results.

1.1 Posets, Connectedness, and Braid Cones

A much fuller treatment of braid cones can be found in [18]. Here we only discuss what is needed for our purposes. We write $[n] = \{1, 2, \dots, n\}$ and have the lattice $\mathbb{Z}^n \subseteq \mathbb{R}^n$ with standard basis denoted $\{e_i \mid i \in [n]\}$ and coordinates (x_1, \dots, x_n) . For each $A \subseteq [n]$, we define $e_A = \sum_{i \in A} e_i$. Our focus will be on the lattice $N = \mathbb{Z}^n / \mathbb{Z}e_{[n]}$ with corresponding vector space $N_{\mathbb{R}} = \mathbb{R} \otimes_{\mathbb{Z}} N$. The dual lattice and vector space will be denoted by M and $M_{\mathbb{R}}$ respectively.

We will assume that the reader is familiar with basic properties of posets. For more information on posets or any undefined terms, the reader may consult [20, Chap. 5] and [23, Chap. 3]. We will often need to refer to graph-theoretic aspects of the Hasse diagram of a poset. Unless otherwise noted, when we apply graph-theoretic adjectives

Table 1 A brief overview of the main results of the paper

Type of Poset	Sufficient Condition to be Gorenstein	Also Necessary Condition	Theorem or Proposition Number
Connected	Order dual is Gorenstein	Yes	Proposition 3.2
Connected	Biconnected components of Hasse diagram are Gorenstein	Yes	Theorem 4.3
Contains a min and max	None (they are always Gorenstein)	Yes	Corollary 3.4
Contains a min or max	Successive quotients of poset behave nicely ²	Yes	Theorem 5.19 (see also Algorithm 5.20)
Contains max and the Hasse diagram of bottom two levels ³ is connected	The Möbius function is zero outside of bottom two levels	Yes	Theorem 5.23
Has length 1	Hasse diagram is 2-regular and 2-connected	No	Proposition 6.4
Has length 1	Hasse diagram is complete bipartite	No	Proposition 6.5

²More explicitly, in the quotients, the Hasse diagram of the bottom two levels must be acyclic and the quotient must satisfy the tree-downset condition (see Definition 5.17)
³If the poset is ranked, this is just the bottom two rank levels of the poset. In the more general case, it is all the elements that are either minimal or cover only minimal elements




Fig. 1 The Hasse diagram of a poset P along with the cone σ_P . The values in red give a chamber crepant labeling of P

to a poset or its Hasse diagram, we are viewing the Hasse diagram as an undirected graph. For example, when we say a poset P is *connected* or *biconnected*, we mean that the underlying undirected Hasse diagram of P is connected or biconnected.

The *length* of a chain in a poset is one less the number of elements in the chain. For example, the chain $x_0 < x_1 < \dots < x_\ell$ has length ℓ . The *length* of a poset P is the maximum among all lengths of chains in P . We will use $\ell(P)$ to denote the length of a poset. Note that in the case that P is ranked, the length is the rank of the poset.

Let P be a connected poset with underlying set $[n]$. The *braid cone* associated to P , denoted by σ_P , is given by the intersection of the half-spaces $x_i \leq x_j$ for each pair (i, j) with $i <_P j$. Note that, by [18, Prop. 3.5 (8)], it suffices to only use the half-spaces $x_i \leq x_j$ when $i \ll_P j$. In Fig. 1 we have an example of a poset P and the cone σ_P . In this example, the cone is the intersection of the four half-spaces $x_1 \leq x_2$, $x_1 \leq x_3$, $x_2 \leq x_4$, and $x_3 \leq x_4$. One can check that σ_P is also given by the positive span of the vectors $e_{\{4\}}$, $e_{\{2,4\}}$, $e_{\{3,4\}}$, and $e_{\{2,3,4\}}$. We will give a general way to find these generating vectors in Lemma 2.3.

In [18, Prop. 3.5] it is shown that for any poset P , the cone σ_P is full dimensional and rational. The connectedness of P guarantees that σ_P is strongly convex (which the authors call *pointed* in [18]). We will assume throughout the article that all posets are connected and also assume that all cones encountered are strongly convex, rational polyhedral cones. We will also assume that $|P| \geq 2$ to avoid issues with triviality. It is often convenient to assume our posets have $[n]$ as their underlying set. However, if P is any poset with an underlying set of cardinality n , we may pick some bijection to $[n]$ and use this bijection to define σ_P . None of the properties of σ_P that we are interested in are influenced in any way by this choice of bijection. This will allow us to consider multiple posets with disjoint underlying sets.

The *braid arrangement* is the hyperplane arrangement in $N_{\mathbb{R}}$ consisting of the hyperplanes $x_i = x_j$ for $1 \leq i < j \leq n$. This is the Coxeter arrangement of type A_{n-1} . The regions of the braid arrangement are the Weyl chambers of type A_{n-1} and are σ_L for each linear order L on the elements of $[n]$. We use $|\Sigma|$ to denote the *support* of a fan Σ which consists of all points of $N_{\mathbb{R}}$ that are contained in any cone the fan Σ . A single cone together with its faces can be considered as a fan. In general $|\sigma_P|$ is the

union of $|\sigma_L|$ over all linear extensions L of P . For any poset P , we write Σ_P for the fan consisting of the cones σ_L and its faces for each linear extension L of P . Thus $|\Sigma_P| = |\sigma_P|$ as subsets of $N_{\mathbb{R}}$ though generally Σ_P contains additional faces which subdivide σ_P .

Returning to our example of P in Fig. 1, we can obtain Σ_P by adding a wall on the hyperplane $x_2 = x_3$ which is spanned by $e_{\{2,3,4\}}$ and $e_{\{4\}}$. The fan will then have two maximal cones, each corresponding to one of the two linear extensions of P .

1.2 The Corresponding Toric Varieties

For an in-depth introduction to toric varieties and the Gorenstein property, the reader can see the text [3], particularly Chapter 8 of the book. Given a cone σ , which is minimally generated by the positive span of finitely many vectors $\{v_i\}_i$, the set of *ray generators* of σ is $\{w_i\}_i$ where w_i is the first nonzero lattice point on the ray $\mathbb{R}_{\geq 0}v_i$. Note that the ray generators of a cone are unique. If σ is strongly convex with ray generators from some lattice, we get an affine toric variety we denote by U_{σ} . Similarly, given any fan of such cones Σ , it defines an abstract toric variety denoted X_{Σ} . We will write U_P for U_{σ_P} and X_P for X_{Σ_P} .

A variety is called \mathbb{Q} -*Gorenstein* if some multiple of its canonical divisor is Cartier. A variety is *Gorenstein* if it is Cohen–Macaulay and its canonical divisor is Cartier. When the variety is toric and defined by a cone, there is an equivalent definition of being \mathbb{Q} -Gorenstein or Gorenstein as we explain next. Let $v_1, v_2, \dots, v_k \in N$ be the ray generators of σ , then U_{σ} is \mathbb{Q} -Gorenstein if and only if the exists $u \in M$ and a positive integer r such that $\langle u, v_i \rangle = r$ for all $1 \leq i \leq k$. The *index* of a \mathbb{Q} -Gorenstein toric variety is the minimal possible r which can be taken. If the index is $r = 1$, then the toric variety is Gorenstein since toric varieties arising from fans are always Cohen–Macaulay [11].

A cone is *smooth* if its ray generators can be extended to a basis for the lattice, and a fan is smooth if it consists of smooth cones. If Σ is a smooth fan refining a cone σ then $X_{\Sigma} \dashrightarrow U_{\sigma}$ gives a resolution of singularities. Take a \mathbb{Q} -Gorenstein toric variety U_{σ} of index r and $u \in M$ such that $\langle u, v_i \rangle = r$ for all ray generators v_1, v_2, \dots, v_k of σ . If Σ is a smooth fan refining σ such that any ray generator v of a cone in Σ also has $\langle u, v \rangle = r$, then this resolution is called *crepant*. A crepant resolution does not change the canonical class.

As we saw earlier, the cone in Fig. 1 is generated by $e_{\{4\}}$, $e_{\{2,4\}}$, $e_{\{3,4\}}$, and $e_{\{2,3,4\}}$. In fact, these are the ray generators of the cone. If we take $u = e_{\{4\}}^* - e_{\{1\}}^*$, then

$$\langle u, e_{\{4\}} \rangle = \langle u, e_{\{2,4\}} \rangle = \langle u, e_{\{3,4\}} \rangle = \langle u, e_{\{2,3,4\}} \rangle = 1.$$

Moreover,

$$\langle u, e_{\{1,2,3,4\}} \rangle = 0$$

which must hold since $e_{\{1,2,3,4\}} = 0$ in $N_{\mathbb{R}}$. We conclude that U_P is Gorenstein. Because all the ray generators of the fan Σ_P are also ray generators of U_P , $X_P \dashrightarrow U_P$ is crepant.

At this point the reader may be wondering how one can find a $u \in M$ to determine if U_P is Gorenstein or if $X_P \dashrightarrow U_P$ is crepant. In the next section we introduce poset labelings to do this exactly. This allows us to translate our algebraic problem to a purely combinatorial one.

2 Ray Generators, Lattice Points, and Labelings

In this section we describe the ray generators of a braid cone as well as the lattice points of the form e_A for $A \subseteq [n]$ which are contained in the cone. We use this description to define labelings of the poset P which capture when U_P is \mathbb{Q} -Gorenstein or Gorenstein as well as when $X_P \dashrightarrow U_P$ is crepant.

For a binary relation R , let R^{op} denote the opposite binary relation where $(i, j) \in R^{op}$ if and only if $(j, i) \in R$. We define a *contraction* of a poset P to be the transitive closure of $P \cup R^{op}$ for some $R \subseteq P$. For example, taking the poset $P = \{2 <_P 1, 2 <_P 3\}$ and $R = \{2 <_R 1\}$ we find the transitive closure of $P \cup R^{op}$ to be $\{1 < 2, 2 < 1, 1 < 3, 2 < 3\}$ which is now not a poset, but is a preposet. A key fact is that if a poset P corresponds to a cone σ , then preposets P' that are contractions of P are in bijection with faces $\tau \subseteq \sigma$ [18, Prop. 3.5 (2)]. We use the notation $\tau = \sigma_{P'}$ where the face $\sigma_{P'}$ is the face obtained by intersecting σ_P with the hyperplanes $x_i = x_j$ for $(i, j) \in P'$ such that we also have $(j, i) \in P'$.

Remark 2.1 A related notion to the opposite binary relation, is the *order dual* of a poset P . This is given by reversing all the inequalities defining P . To avoid confusion, we will use the notation P^* when we are dealing with taking the order dual of an entire partial order P whereas R^{op} can be used for any binary relation.

Remark 2.2 It can be helpful to visualize the contraction of posets that was just defined as edge contraction in the Hasse diagram viewed as a graph. For example, this is done in [16] for the special case of tree posets. However, one needs to keep in mind there may be extra steps beyond just contracting edges accounting for the transitive closure. Also, R may include a poset relation which is not a cover relation and hence there would not be an edge draw in the Hasse diagram.

Given an upset A of P , we define

$$R_A = \{(i, j) \in P \mid i, j \in A \text{ or } i, j \notin A\}$$

and let the contraction P_A be the transitive closure of $P \cup R_A^{op}$. The fact that P_A is a preposet means that P_A is reflexive and transitive. In particular, P_A need not be antisymmetric. So, P_A defines an equivalence relation where i is equivalent to j if $(i, j) \in P_A$ and $(j, i) \in P_A$. We also define the *dimension* of A , denoted $\dim(A)$, to be one less than the number of equivalence classes determined by the preposet P_A . Note the $\dim(A)$ depends on the poset P , but A is always taken to be an upset of a

poset P so the data on the poset is present. The dimension, $\dim(A)$, can be computed by

$$\dim(A) = cc(A) + cc(\overline{A}) - 1$$

where $cc(A)$ and $cc(\overline{A})$ are the number of connected components of the Hasse diagram of P restricted to A and \overline{A} respectively. Here \overline{A} denotes the complement of A in $[n]$. For the poset depicted in Fig. 2, we have that $\dim(\{1, 3\}) = 1$ since $\{1, 3\}$ and $\overline{\{1, 3\}} = \{2\}$ are both connected, whereas $\dim(\{3\}) = 2$ since $\{3\}$ is connected and removing it disconnects the graph into two parts.


Since we are interested in the Gorenstein property and crepant resolutions, we need to know what the ray generators of a cone σ_P are as well as which ray generators are added when refining σ_P to Σ_P . The next lemma will describe primitive lattice points on these rays. Let us briefly recall a few definitions from convex geometry. A *face* of a cone is a subset of the cone obtained by intersection with a supporting hyperplane. The *dimension* of a face is the dimension of its linear span. Lastly, a point is said to be in the *relative interior* of a face τ if it is not contained in any proper face $\tau' \subsetneq \tau$.

Lemma 2.3 *If P is a poset, then the lattice point e_A is contained in σ_P if and only if A is an upset. Moreover, for an upset A the lattice point e_A is contained in the relative interior of a face $\sigma_{P_A} \subseteq \sigma_P$ with $\dim(\sigma_{P_A}) = \dim(A)$. In particular, e_A is a ray generator of σ_P if and only if A is an upset of dimension 1.*

Proof It follows immediately from the definitions that $e_A \in \sigma_P$ if and only if A is an upset. Now assume that A is an upset so that $e_A \in \sigma_P$. Then there is a face $\sigma_{P_A} \subseteq \sigma_P$ with $\dim(\sigma_{P_A}) = \dim(A)$ indexed by the contraction P_A . Furthermore, we have $e_A \in \sigma_{P_A}$ by construction. It only remains to show that e_A is in the relative interior of σ_{P_A} . That is, we must show e_A is not in any face properly contained in σ_{P_A} . Such faces properly contained in σ_{P_A} will correspond to contractions of P_A . Take any $(i, j) \in P_A$. If $(j, i) \in P_A$ this relation will not matter in any further contractions. If $(j, i) \notin P_A$, then since A is an upset we have that $j \in A$ and $i \notin A$. For any contraction which is the transitive closure of $P_A \cup R^{op}$ with $(i, j) \in R$, we must have $x_j \leq x_i$ in the corresponding cone. However, since $j \in A$ and $i \notin A$ we see that e_A does not satisfy this inequality. Therefore e_A is in the relative interior of σ_{P_A} as desired.

The final assertion now follows since the ray generators that show up in cones of the braid arrangement fan are precisely all 0-1 vectors except the all 0 vector e_\emptyset and all the 1 vector $e_{[n]}$ (which is equal to zero is our setting). \square

Remark 2.4 One can compare our cone σ_P and Lemma 2.3 with the cone denoted K_P^{wt} in [1, Prop. 5.1]. For the cone K_P^{wt} , the ray generators are the indicator vectors of all downsets which are connected (the complement need not be connected). This difference compared to our case arises because the cone K_P^{wt} lives in \mathbb{R}^n for a poset on n elements as opposed to $\mathbb{R}^n / e_{[n]}$ along with conventions which replace upsets with downsets. Also [7, Prop. 6.6] shows that for certain choices of P the affine semigroup rings of K_P^{wt} are the rings studied in [9]. These latter rings are the ones which give rise to the toric varieties in [25] mentioned in the introduction.



$$Q$$

Fig. 2 A poset together and its Gorenstein labeling given in red

Let us now determine if U_Q is Gorenstein for Q in Fig. 2. Recall that U_Q is Gorenstein if and only if there is a $u \in M$ such that $\langle u, e_A \rangle = 1$ for all ray generators e_A . Let $u = y_1 e_{\{1\}}^* + y_2 e_{\{2\}}^* + y_3 e_{\{3\}}^*$. Using Lemma 2.3, one can verify that the ray generators of σ_Q are $e_{\{1,3\}}$ and $e_{\{2,3\}}$. This together with the fact that $e_{\{1,2,3\}} = 0$ in $N_{\mathbb{R}}$ implies that if U_Q is Gorenstein, then the following system has integer solutions.

$$\begin{aligned} y_1 + y_3 &= 1 \\ y_2 + y_3 &= 1 \\ y_1 + y_2 + y_3 &= 0 \end{aligned}$$

The system does have a solution namely $y_1 = -1$, $y_2 = -1$, $y_3 = 2$, and so U_Q is Gorenstein. Since 1, 2, and 3 are elements of Q , we can “label” the element i of Q with the value of y_i to get a labeling that completely determines if U_Q is Gorenstein. Motivated by the ideas presented in this example, we introduce the following poset labelings.

Definition 2.5 Given a poset P on $[n]$, an r -Gorenstein labeling is a function $\phi : P \rightarrow \mathbb{Z}$ such that

$$\sum_{i \in P} \phi(i) = 0$$

and

$$\sum_{i \in A} \phi(i) = r$$

for all upsets A with $\dim(A) = 1$. We call a 1-Gorenstein labeling a *Gorenstein labeling*.

Remark 2.6 A poset admitting an r -Gorenstein labeling is unrelated to a poset being Gorenstein*. As the next proposition shows, an r -Gorenstein labeling of a poset P coincides with the toric variety U_P being \mathbb{Q} -Gorenstein. Whereas the Gorenstein* property is related to Gorensteinness of the order complex of P [22].

One can verify that the labelings in Fig. 1 and Fig. 2 are Gorenstein labelings. As we saw earlier, the varieties associated to these posets are both Gorenstein. This is no coincidence as we see next.

Proposition 2.7 Let P be a poset. U_P is \mathbb{Q} -Gorenstein with index r if and only if P has an r -Gorenstein labeling but has no s -Gorenstein labeling for any $0 < s < r$.

Proof Given a poset P , let

$$\{A_1, A_2, \dots, A_k\} = \{A \mid A \text{ is an upset and } \dim(A) = 1\}.$$

So by Lemma 2.3, the ray generators of σ_P are $e_{A_1}, e_{A_2}, \dots, e_{A_k}$. We observe that $u \in M$ with $\langle u, e_{A_j} \rangle = r$ for $1 \leq j \leq k$ is equivalent to an r -Gorenstein labeling ϕ where $\phi(i) = \langle u, e_i \rangle$ for $1 \leq i \leq n$. \square

Definition 2.8 Let P be a poset for which U_P is \mathbb{Q} -Gorenstein with index r . A *chamber crepant labeling* of P is a function $\phi : P \rightarrow \mathbb{Z}$ such that

$$\sum_{i \in P} \phi(i) = 0$$

and

$$\sum_{i \in A} \phi(i) = r$$

for all upsets A with $\dim(A) > 0$.

One can check that the labeling of the poset in Fig. 1 is a chamber crepant labeling. Indeed, every upset except the full poset only contains a single element with a nonzero label and the total sum of values is 0. On the other hand, the labeling of the poset in Fig. 2 is not a chamber crepant labeling. For example, the sum of the values in the upset $\{1, 3\}$ is 1 and the sum of values for $\{3\}$ is 2. As we will see in Sect. 3, there are no chamber crepant labelings of the poset in Fig. 2 because it is not bounded.

Proposition 2.9 Let P be a poset for which U_P is \mathbb{Q} -Gorenstein. The poset P has a chamber crepant labeling if and only if $X_P \dashrightarrow U_P$ is crepant.

Proof Having a chamber crepant labeling means there is a $u \in M$ such that $\langle u, e_A \rangle = r$ for some positive integer r and all upsets A with $\dim(A) > 0$. By Lemma 2.3, we see that the rays added when refining σ_P to Σ_P will be exactly e_A such that A is an upset with $\dim(A) > 0$. So, having a chamber crepant labeling is equivalent to U_P being \mathbb{Q} -Gorenstein and $X_P \dashrightarrow U_P$ being crepant. \square

Before we move on, let us note that if P has a r -Gorenstein labeling, then it only has one such labeling. Indeed, by [18, Prop. 3.5 (4)], σ_P is a full-dimensional cone. Thus the set $\{e_A \mid \dim(A) = 0, 1\}$ has full rank and so the linear equations given in Definition 2.5 have at most one solution. Since the chamber crepant labelings are special cases of r -Gorenstein labeling (where r is the index of the variety), we see that if there is a chamber crepant labeling, it is also unique.

As one can see from Lemma 2.3, once we know the upsets of a poset we can determine the ray generators of the cone. Then determining whether the corresponding

toric variety is Gorenstein or if it has a crepant resolution amounts to solving a system of linear equations. Proposition 2.7 and Proposition 2.9 show we can alternatively work with labelings of the poset to solve this system. Of course solving a linear system is easy, the real difficulty in these problems stems from the fact that determining the upsets of a general poset is known to be difficult. The problem of counting upsets has been shown to be #P-complete [17].

Since the ray generators correspond to dimension 1 upsets, we do not need to actually know all the upsets to determine if the variety is Gorenstein. However, even finding the ray generators does not have an easy known solution in general. The cover relations of P , which are the edges of the Hasse diagram, give the half-space representation of polyhedron σ_P . The problem of finding the ray generators of σ_P is equivalent to finding the vertex representation of a certain polytope. This polytope is the projectivization that one obtains by adding the “far face” at infinity to σ_P . Vertex enumeration of general polyhedra given by half-space representation is known to be a hard problem [12].

Despite the difficulties outlined above, we are able to show that under certain circumstances, there are efficient methods to solve these problems. This is what we explore in the remainder of the paper.

3 Chamber Crepant Labelings and Bounded Posets

In [16] it was shown that, for any poset P , Σ_P can always be obtained from σ_P by a sequence of star subdivisions. This means that X_P and U_P are related by a sequence of blow-ups. In this section we further look at the birational map $X_P \dashrightarrow U_P$ and determine when it is crepant. First, we explore how the Gorenstein and crepant property behave when taking order duals.

If A is an upset of P , then \overline{A} is a downset of P . Since the total sum of labels in a r -Gorenstein labeling must be 0, this means that the sum of the labels of A is completely determined by the sum of labels of \overline{A} when A and \overline{A} are both connected. As a result, we get the following.

Proposition 3.1 *Let P be a poset. The map $\phi : P \rightarrow \mathbb{Z}$ is a r -Gorenstein (resp. chamber crepant) labeling if and only if*

$$\sum_{i \in P} \phi(i) = 0$$

and

$$\sum_{i \in A} \phi(i) = -r$$

for all downsets A such $\dim(A) = 1$ (resp. $\dim(A) > 0$).

Upsets and downsets reverse roles when taking the order dual of a poset. Thus the previous proposition implies that if ϕ is an r -Gorenstein (resp. chamber crepant)




Fig. 3 Posets P_1 , P_2 and $(P_1, c) \star (P_2, x)$ along with Gorenstein labelings in red

labeling of P , then $-\phi$ is a r -Gorenstein (resp. chamber crepant) labeling of the order dual P^* . This gives the following.

Proposition 3.2 *Let P be a poset and let P^* be its order dual. U_P is \mathbb{Q} -Gorenstein if and only if U_{P^*} is \mathbb{Q} -Gorenstein. Similarly, U_P is Gorenstein if and only if U_{P^*} is Gorenstein. Moreover, $X_P \dashrightarrow U_P$ is crepant if and only if $X_{P^*} \dashrightarrow U_{P^*}$ is crepant.*

Recall a poset P is *bounded* if P has a minimal element (denoted by $\hat{0}$) and a maximal element (denoted by $\hat{1}$).

Theorem 3.3 *Let P be a poset. $X_P \dashrightarrow U_P$ is crepant if and only if P is bounded.*

Proof (\Rightarrow) By Definition 2.5, if ϕ is a chamber crepant labeling of P and m is a maximal element of P , then $\phi(m) = r$. Now let A be the set of maximal elements of P . Then A is an upset and since P is connected and non-trivial, $A \neq P$. So if $X_P \dashrightarrow U_P$ is crepant, the sum of the labels of A must be r and so $|A| = 1$. It follows that P has a $\hat{1}$. By Proposition 3.2, the crepant property is preserved by taking order duals. Using the fact that a poset has a $\hat{1}$ if and only if its order dual has a $\hat{0}$, we get this direction.

(\Leftarrow) Now suppose that P is bounded. Define $\phi : P \rightarrow \mathbb{Z}$ by $\phi(\hat{0}) = -1$, $\phi(\hat{1}) = 1$, and $\phi(x) = 0$ for all $x \in P \setminus \{\hat{0}, \hat{1}\}$. It is straightforward to show that ϕ is a chamber crepant labeling and so we have this direction. \square

The above theorem immediately gives us the following corollary.

Corollary 3.4 *Let P be a bounded poset. Then U_P is Gorenstein.*

4 Gorensteinness and Biconnected Components

Given any two disjoint posets P_1 and P_2 with Hasse diagrams H_1 and H_2 respectively along with $x_1 \in P_1$ and $x_2 \in P_2$ we define

$$(P_1, x_1) \star (P_2, x_2) = (P_1 \uplus P_2) / (x_1 \sim x_2)$$

which can be gotten by taking disjoint copies of H_1 and H_2 and identifying x_1 with x_2 . Informally, we can think of this as “gluing” together the posets along x_1 and x_2 . See Fig. 3 for an example.

In the example depicted in Fig. 3, the labelings of P_1 and P_2 are Gorenstein labelings. The labeling of $(P_1, c) \star (P_2, x)$ is obtained by summing the values of the elements that were identified and keeping all the other labels the same. One can verify that this new labeling is a Gorenstein labeling. As we show next, this is no coincidence.

Lemma 4.1 *Let P_1 and P_2 be posets with $x_1 \in P_1$ and $x_2 \in P_2$. If both P_1 and P_2 admit r -Gorenstein labelings, then $(P_1, x_1) \star (P_2, x_2)$ admits an r -Gorenstein labeling.*

Proof Let ϕ_1 and ϕ_2 be r -Gorenstein labelings for P_1 and P_2 respectively. Also, let $P = (P_1, x_1) \star (P_2, x_2)$. Let $\phi : P \rightarrow \mathbb{Z}$ given by

$$\phi(z) = \begin{cases} \phi_1(z) & \text{if } z \in P_1 \setminus \{x_1\}, \\ \phi_2(z) & \text{if } z \in P_2 \setminus \{x_2\}, \\ \phi_1(z) + \phi_2(z) & \text{if } z = x_1 = x_2. \end{cases}$$

We claim that ϕ is an r -Gorenstein labeling for P . Let $A \subseteq P$ be an upset with $\dim(A) = 1$. First suppose that A does not contain $x_1 = x_2$. Then either $A \subseteq P_1$ and

$$\sum_{z \in A} \phi(z) = \sum_{z \in A} \phi_1(z) = r$$

or else $A \subseteq P_2$ and the result follows by same calculation with ϕ_2 in place of ϕ_1 . Otherwise $x_1 = x_2$ is in A . Then it must be that $P_1 \subseteq A$ or $P_2 \subseteq A$. If not $\bar{A} \cap P_1$ and $\bar{A} \cap P_2$ are nonempty and disjoint. However this would imply that $\dim(A) > 1$. When $P_1 \subseteq A$ we have that

$$\begin{aligned} \sum_{z \in A} \phi(z) &= \sum_{z \in A \cap (P_1 \setminus \{x_1\})} \phi(z) + \phi(x_1 = x_2) + \sum_{z \in A \cap (P_2 \setminus \{x_2\})} \phi(z) \\ &= \sum_{z \in A \cap (P_1 \setminus \{x_1\})} \phi_1(z) + \phi_1(x_1) + \phi_2(x_2) + \sum_{z \in A \cap (P_2 \setminus \{x_1\})} \phi_2(z) \\ &= \sum_{z \in P_1} \phi_1(z) + \sum_{z \in A \cap P_2} \phi_2(z) \\ &= 0 + r \\ &= r \end{aligned}$$

where the second to last line holds since $A \cap P_2$ is an upset of dimension 1 in P_2 . A similar calculation also holds in the case that $P_2 \subseteq A$. Thus, the lemma is proven. \square

Recall, a graph is biconnected if it is connected and remains connected after the removal of any single vertex. A *biconnected component* of a graph is a maximal biconnected subgraph. For any graph its biconnected components and cut vertices form the vertex set of the *block-cut tree* (see e.g. [8, p. 36]). In this tree, we have an edge between a biconnected component and a cut vertex if and only if the cut vertex is contained in the biconnected component. The next lemma will be proven by induction using biconnected components and the block-cut tree of a Hasse diagram.

Lemma 4.2 *If P admits an r -Gorenstein labeling, then each biconnected component of P admits an r -Gorenstein labeling.*

Proof If P is biconnected, we are done. So we may assume it is not biconnected. Let x_0 be a cut vertex such that $X \cup Y = P$, $X \cap Y = \{x_0\}$, and $P|_X$ is a leaf of the block-cut tree. Let $P_X = P|_X$ and $P_Y = P|_Y$ both of which are connected posets. We show both P_X and P_Y have r -Gorenstein labelings.

By assumption P has an r -Gorenstein labeling, $\phi : P \rightarrow \mathbb{Z}$. Define $\phi_X : P_X \rightarrow \mathbb{Z}$ by

$$\phi_X(x) = \begin{cases} \phi(x) & \text{if } x \neq x_0, \\ \sum_{y \in Y} \phi(y) & \text{if } x = x_0. \end{cases}$$

We claim this is an r -Gorenstein labeling of P_X . Indeed, suppose that $A \subseteq P_X$ is an upset of dimension 1. If $x_0 \notin A$, then

$$\sum_{x \in A} \phi_X(x) = \sum_{x \in A} \phi(x) = r$$

since A is also an upset of P with $\dim(A) = 1$. In the case where $x_0 \in A$, we let $B = A \cup Y$, which is an upset of P with $\dim(B) = 1$. It follows that

$$\begin{aligned} \sum_{x \in A} \phi_X(x) &= \phi_X(x_0) + \sum_{x \in A \setminus \{x_0\}} \phi_X(x) \\ &= \sum_{x \in Y} \phi(x) + \sum_{x \in A \setminus \{x_0\}} \phi(x) \\ &= \sum_{x \in B} \phi(x) \\ &= r \end{aligned}$$

and so P_X has a r -Gorenstein labeling.

The same argument shows that the function $\phi_Y : P_Y \rightarrow \mathbb{Z}$ defined by

$$\phi_Y(x) = \begin{cases} \phi(x) & \text{if } x \neq x_0, \\ \sum_{y \in X} \phi(y) & \text{if } x = x_0. \end{cases}$$

is an r -Gorenstein labeling of P_Y . The result now holds by induction on the number of biconnected components of P . \square

Theorem 4.3 *A poset P has an r -Gorenstein labeling if and only if each biconnected component of P has an r -Gorenstein labeling.*

Proof The forward direction of the theorem is exactly Lemma 4.2. The backwards direction follows from Lemma 4.1, noting that P can be built from the operation \star applied to the biconnected components and cut vertices. \square

5 Posets with a Minimum Element or a Maximum Element

In this section we explore when a poset with a $\hat{0}$ or a $\hat{1}$ has a Gorenstein labeling. We show that if P has a $\hat{0}$ or a $\hat{1}$, then U_P is Gorenstein if and only if it is \mathbb{Q} -Gorenstein (Theorem 5.2). Moreover, we will see that whether U_P is Gorenstein, depends on the Möbius function of P . For posets P with a $\hat{0}$ or a $\hat{1}$, we are able to give a characterization of when U_P is Gorenstein (Theorem 5.19). Using this characterization, we give an algorithm to check if U_P is Gorenstein in this case (Algorithm 5.20). The characterization and the algorithm both make use of quotient posets.

A poset has a $\hat{0}$ if and only if its order dual has a $\hat{1}$. By Proposition 3.2, we know that U_P is Gorenstein if and only if U_{P^*} is Gorenstein. Therefore we only need to prove the results when either P has a $\hat{0}$ or has a $\hat{1}$. We believe the presentation of the proofs is simpler in the case that the poset has a $\hat{1}$ and thus we have chosen to present the proofs this way.

Throughout this section, we will often need to consider downsets generated by a single element. We will use the notation $\langle x \rangle_\downarrow$ to denote the downset generated by x . Similarly we will use $\langle S \rangle_\downarrow$ for the downset generated by the set S .

5.1 \mathbb{Q} -Gorenstein Implies Gorenstein

For any element x of a poset P we have the *principal downset generated by x* denoted $\langle x \rangle_\downarrow$ which contains all elements y so that $y \leq x$. If P has a $\hat{1}$ then for any proper downset $A \neq P$, \overline{A} is connected. Since any principal downset is connected, the next lemma will show that this forces any possible Gorenstein labeling to be completely determined by principal downsets. This allows us to give a recursive definition for an r -Gorenstein labeling.

Lemma 5.1 *Let P be a poset with a $\hat{1}$. If P has a r -Gorenstein labeling, then this labeling is given by*

$$\sum_{y \leq x} \phi(y) = r(\delta_{x, \hat{1}} - 1)$$

where $\delta_{x, \hat{1}}$ is the Kronecker delta function.

Proof Since P has a $\hat{1}$, if A is a proper downset, then \overline{A} is connected. Now suppose that $A = \langle x \rangle_\downarrow$ with $x \neq \hat{1}$. Then since x is the unique maximum element of A , A is connected. Thus, $\dim(A) = 1$.

Now $\langle x \rangle_\downarrow = P$ if and only if $x = \hat{1}$. It follows that from Proposition 3.1 that

$$\sum_{y \in \langle x \rangle_\downarrow} \phi(y) = r(\delta_{x, \hat{1}} - 1)$$

or equivalently,

$$\sum_{y \leq x} \phi(y) = r(\delta_{x, \hat{1}} - 1).$$

Thus, the result holds. \square

We are now ready to prove one of the main theorems in this section.

Theorem 5.2 *Let P be a poset with a $\hat{0}$ or a $\hat{1}$. Then U_P is Gorenstein if and only if U_P is \mathbb{Q} -Gorenstein.*

Proof We prove the result when P has a $\hat{1}$ which suffices by Proposition 3.2. The forward direction is immediate given the definitions of Gorenstein and \mathbb{Q} -Gorenstein.

For the backwards direction, it suffices to show that if ϕ is the unique r -Gorenstein labeling given in Lemma 5.1, then $r \mid \phi(x)$ for all $x \in P$. We do this by inducting on the maximum length of a chain from a minimal element to x . If the length is 0, then x must be a minimal element. So by Lemma 5.1, $\phi(x) = -r$ and so the base case holds.

Now suppose that the maximum length of a chain from x to a minimal element of P is k and that $x \neq \hat{1}$. Then by Lemma 5.1,

$$\phi(x) = - \sum_{y < x} \phi(y) - r.$$

By the inductive hypothesis, we have that $r \mid \phi(y)$ for all $y < x$. Thus, $r \mid \phi(x)$ in this case. Finally, by Lemma 5.1,

$$\phi(\hat{1}) = - \sum_{y \neq \hat{1}} \phi(y).$$

We have already shown that $r \mid \phi(y)$ for all $y \neq \hat{1}$ and so $r \mid \phi(\hat{1})$. It follows that $\frac{1}{r}\phi(x) \in \mathbb{Z}$ for all x and so $\frac{1}{r}\phi(x)$ is a Gorenstein labeling of P . \square

5.2 Relationship with the Möbius Function

Given the previous theorem, we will only focus on Gorenstein labelings (as opposed to r -Gorenstein labelings) in the remainder of this section. In the next proposition, we show that the labeling described in Lemma 5.1 (when $r = 1$) is related to the Möbius function. In this proposition and elsewhere in this section, we will be considering a new poset, namely, the poset obtained by adding a minimum element $\hat{0}$ to P . We note that we add a $\hat{0}$ even if P already has a minimum element. The poset obtained by adding a $\hat{0}$ to P will be denoted by \hat{P} . We now give the definition of the Möbius function.

Definition 5.3 Let P be a poset with a $\hat{0}$. The (*one-variable*) Möbius function, $\mu_P(x) : P \rightarrow \mathbb{Z}$, is defined recursively by

$$\sum_{y \leq x} \mu_P(y) = \delta_{\hat{0},x}$$

where $\delta_{\hat{0},x}$ is Kronecker delta function.




Fig. 4 Posets P and Q with the elements of M_P and M_Q colored blue. The values next to an element x given in red correspond to $\mu_{\widehat{P}}(x)$

We will use the notation $\mu_{\widehat{P}}(x)$ for the Möbius function of x in \widehat{P} . See Fig. 4 for examples of the Möbius function. From time to time, we will need to consider Möbius functions of different posets. In these cases, we will use the subscripts to distinguish them.

Lemma 5.4 *Let P be a poset with $\widehat{1}$ and let ϕ be the labeling described in Lemma 5.1 when $r = 1$. Then for all $x \in P$,*

$$\phi(x) = \begin{cases} \mu_{\widehat{P}}(x) & \text{if } x \neq \widehat{1}, \\ \mu_{\widehat{P}}(x) + 1 & \text{if } x = \widehat{1}. \end{cases}$$

Proof We prove the result for $x \neq \widehat{1}$, the argument for $x = \widehat{1}$ is similar.

We induct on the maximum length of a chain from a minimal element of P to x . If the length is 0, then x is minimal. Using the definition of the Möbius function, we have that $\mu_{\widehat{P}}(x) = -1$. This agrees with the value given in Lemma 5.1 (with $r = 1$) and so the base case holds.

Now suppose that the maximum length is more than 0. Then, by Lemma 5.1 (with $r = 1$), we have that

$$\sum_{y \leq x} \phi(y) = -1.$$

Rearranging gives us that

$$\phi(x) = - \sum_{y < x} \phi(y) - 1.$$

By the inductive hypothesis, $\phi(y) = \mu_{\hat{P}}(y)$ for all $y < x$. Moreover, by definition, $\mu_{\hat{P}}(\hat{0}) = 1$. Thus the previous displayed equation becomes

$$\phi(x) = - \sum_{y < x} \mu_{\hat{P}}(y) - \mu_{\hat{P}}(\hat{0}).$$

Again using the definition of the Möbius function, we see that the righthand side is exactly $\mu_{\hat{P}}(x)$. Thus the result holds by induction. \square

Combining Lemma 5.1 and Lemma 5.4, we get the following characterization of when U_P is Gorenstein in terms of the Möbius function.

Theorem 5.5 *Let P be a poset with a $\hat{1}$. If ϕ is a Gorenstein labeling of P , then*

$$\phi(x) = \begin{cases} \mu_{\hat{P}}(x) & \text{if } x \neq \hat{1}, \\ \mu_{\hat{P}}(x) + 1 & \text{if } x = \hat{1}. \end{cases}$$

Consequently, U_P is Gorenstein if and only if

$$\sum_{x \in A} \mu_{\hat{P}}(x) = -1$$

for all downsets A of dimension 1.

5.3 A Characterization and Algorithmic Considerations

As we will see when P has a $\hat{1}$, the minimal elements of P and the elements covering them play a crucial role in determining if U_P is Gorenstein. Because of this, we introduce notation for these elements. We will use M_P to denote the induced subposet of elements that are either minimal or cover only minimal elements. That is,

$$M_P = \{x \in P \mid \text{if } y \lessdot x, \text{ then } y \text{ is minimal}\}.$$

See Fig. 4 for two examples. Note that in the example, $e \notin M_Q$. This is because while e does cover b , it also covers c and d which are not minimal.

If U_P is Gorenstein, we must have that the sum of the Möbius values of any connected downset contained in M_P is -1 . Thus it is important for us to understand when this happens. As we show in the next lemma, this forces the connected downsets of M_P to be trees.

Lemma 5.6 *If C is a connected downset of M_P , then*

$$\sum_{x \in C} \mu_{\hat{P}}(x) = -1$$

if and only if C is a tree.

Proof A direct calculation using the definition of the Möbius function shows that for $x \in M_P$,

$$\mu_{\hat{P}}(x) = \begin{cases} -1 & \text{if } x \text{ is minimal in } P \\ \deg_{M_P}(x) - 1 & \text{otherwise} \end{cases}$$

where $\deg_{M_P}(x)$ refers to the degree of x considered as a vertex in the Hasse diagram of M_P . Let C' be the minimal elements of C and let C'' be the non-minimal elements of C . Then

$$\begin{aligned} \sum_{x \in C} \mu_{\hat{P}}(x) &= \sum_{x \in C'} \mu_{\hat{P}}(x) + \sum_{x \in C''} \mu_{\hat{P}}(x) \\ &= \sum_{x \in C'} (-1) + \sum_{x \in C''} (\deg_{M_P}(x) - 1) \\ &= -|C'| + \sum_{x \in C''} \deg_{M_P}(x) - |C''| \\ &= -|C| + \sum_{x \in C''} \deg_{M_P}(x) \end{aligned}$$

Now C is bipartite with partite sets C' and C'' . It follows that $\sum_{x \in C''} \deg_{M_P}(x)$ is the number of edges in C . Combining this fact with the above equations shows that

$$\sum_{x \in C} \mu_{\hat{P}}(x) = |E(C)| - |V(C)|$$

where $E(C)$ is the set of edges of C and $V(C)$ is the set of vertices in C . Since C is connected, $|E(C)| - |V(C)| = -1$ if and only if C is tree and so the result holds. \square

Lemma 5.6 along with Theorem 5.5, imply that M_P must be acyclic for U_P to be Gorenstein. However, this is not enough to guarantee that U_P is Gorenstein. For example, consider the poset P in Fig. 4. The sum of the Möbius values of M_P is -1 and the same is true for $\langle g \rangle \downarrow$. However, if we consider the downset $R \cup \langle g \rangle \downarrow$, where $R = \{a, b, d\}$ is the path in M_P connecting a and b , we see that the sum of the Möbius values is 0 . However, it needs to be -1 for U_P to be Gorenstein. It turns out the reason this arises is because while a and b are in the same tree of M_P , they are not in the same tree of $M_P \cap \langle g \rangle \downarrow$. This allows one to add in the vertices of this “missing” $a - b$ path in $M_P \cap \langle g \rangle \downarrow$ and keep the downset connected. However, by adding this path, we

change the sum of the Möbius values. It turns out that this issue is not unique to this example. To describe what is happening, we need a definition. In the next definition and elsewhere, we will need to count the number of connected components in a poset. As before, we will use the notation $cc(P)$ for the number of connected components of the Hasse diagram of P .

Definition 5.7 Let P be a poset with a $\hat{1}$ and let C be a connected downset of P . We say C satisfies the *connected component condition* (or simply the *cc condition*) if the number of connected components of M_P that have nonempty intersection with C is equal to $cc(C \cap M_P)$. We say P satisfies the cc condition if all of its connected downsets satisfy it.

The downset $\langle g \rangle_\downarrow$ in the poset P of Fig. 4 does not satisfy the cc condition. Indeed, the number of connected components of M_P that have nonempty intersection with $\langle g \rangle_\downarrow$ is 1 since M_P is connected. However, $\langle g \rangle_\downarrow \cap M_P$ has two connected components, namely $\{a, c\}$ and $\{b, e\}$. On the other hand, one can check that all connected downsets in Q of Fig. 4 satisfy the cc condition. And so Q satisfies the cc condition.

Lemma 5.8 Let P be a poset with a $\hat{1}$. If U_P is Gorenstein, then the following hold.

- (1) M_P is acyclic.
- (2) P satisfies the cc condition.

Proof First note that if (1) did not hold, then there would be a connected downset containing a cycle. But then by Lemma 5.6, the sum of the Möbius values in this downset would not be -1 , contradicting Theorem 5.5.

Now suppose that (2) did not hold and let C be a connected downset that does not satisfy the cc condition. Each connected component of $C \cap M_P$ must be contained in a connected component of M_P . Thus, the number of components of M_P that have nonempty intersection with C is less than $cc(C \cap M_P)$. It follows that there is a path completely contained in M_P that connects two connected components in $C \cap M_P$. Let C' be the downset generated by C and this path. Then C' is a connected downset. By construction, $C \setminus M_P = C' \setminus M_P$ and $cc(C \cap M_P) \neq cc(C' \cap M_P)$. But then by Lemma 5.6, C and C' have different sums of Möbius values since each connected component of $C \cap M_P$ and $C' \cap M_P$ contributes -1 to the total sum. This is impossible as $\dim(C) = 1 = \dim(C')$ and U_P is Gorenstein. \square

While the two conditions in Lemma 5.8 are necessary for U_P to be Gorenstein, they are not sufficient. To see why, consider the poset P in Fig. 5. Both conditions of Lemma 5.8 hold. However, U_P is not Gorenstein since the sum of the Möbius values of the downset $\{a, b, c, d, e, f, g\}$ is 0. Even though the two conditions in Lemma 5.8 are not sufficient, we will see that when they hold, we can modify P in a way that reduces the number of elements of P and does not change the Gorensteinness of the corresponding variety. Before we explain this in detail, let us return to P given in Fig. 5. As noted before, M_P is acyclic. Consider a new poset obtained by collapsing the elements of each tree of M_P to a single element. That is, collapse the elements, a, b and d to a single element and c and e to a single element. This new poset we get is labeled as P/\sim in Fig. 5. Note that $M_{P/\sim}$ is not acyclic and so by Lemma 5.8, $U_{P/\sim}$




Fig. 5 Posets P and P/\sim with the elements of M_P and $M_{P/\sim}$ colored blue. The values next to an element x given in red correspond to $\mu_{\hat{P}}(x)$

is not Gorenstein. As we will see, the fact that $U_{P/\sim}$ is not Gorenstein can be used to explain why U_P is not Gorenstein.

We now turn our attention to providing a characterization and an algorithm to determine when a poset with a $\hat{0}$ or $\hat{1}$ has variety that is Gorenstein. This will formalize the ideas we discussed in the previous paragraph. We have already discussed one of the main tools in this characterization, the Möbius function. The other main tool is quotient posets, which we now examine.

Definition 5.9 Let P be a poset and let \sim be an equivalence relation on P . The *quotient*, P/\sim , is the set of equivalence classes with the binary relation \leq given by $X \leq Y$ in P/\sim if and only if $x \leq y$ in P for some $x \in X$ and $y \in Y$. If P/\sim is a poset, we call P/\sim the *quotient poset*.

Remark 5.10 Whenever we are using quotient posets, we will use uppercase letters for elements of the quotient poset and lowercase letters for the original poset. We will use \leq for both orders, despite the fact that they are not the same.

In general, the quotient described in Definition 5.9 need not be a poset. For example, take a chain of length 2 and identify the bottom and top elements. The corresponding \leq relation defined for the quotient is reflexive and is transitive, but it is not antisymmetric. However, the relation that we use here will produce a poset.

Definition 5.11 Let P be a poset such that M_P is acyclic. The *tree relation* \sim is the relation such that $x \sim y$ if and only if $x = y$ or x and y are in the same connected component of M_P .

See Fig. 5 for an example of a poset P and the quotient poset P/\sim where \sim is the tree relation. We will now show the tree relation produces a quotient that is indeed a poset. Note that we assume that M_P is acyclic. Although this is not necessary for the quotient to be a poset, whenever we apply the relation, it will always be acyclic. This also explains the use of the word “tree” in “tree relation”.

Lemma 5.12 *Let P be a poset such that M_P is acyclic. Let \sim be the tree relation. Then P/\sim is a poset.*

Proof First note that it is trivial to verify that \leq is reflexive.

To show that \leq is antisymmetric, suppose on the contrary that there exists $X, Y \in P/\sim$ such that $X \leq Y$ and $Y \leq X$, but $X \neq Y$. Since $X \leq Y$, there is a $x \in X$ and a $y \in Y$ such that $x \leq y$. Since $X \neq Y$, we know that x and y are not in the same equivalence class. We claim that this means that $y \notin M_P$. Indeed, if $y \in M_P$, then $x \leq y$ would imply that $x \in M_P$. However, if $x, y \in M_P$ and $x \leq y$, then x and y are in the same tree of M_P and this would imply that $X = Y$. Now since $Y \leq X$, we have that there is an element in Y below an element of X . Since $y \notin M_P$, Y only has one element and this element must be y . So there exists a $x' \in X$ such that $y \leq x'$. The fact that $y \notin M_P$ and $y \leq x'$ implies that $x' \notin M_P$. But then this means that X must contain a single element and so $x = x'$. This together with $x \leq y \leq x'$ implies that $x = y$ and so $X = Y$, a contradiction.

Finally, we show that \leq is transitive. Suppose that $X \leq Y$ and $Y \leq Z$. If $X = Y$ or $Y = Z$, we immediately get that $X \leq Z$. So we may assume that $X < Y$ and $Y < Z$. Since $X < Y$, there is a $x \in X$, $y \in Y$ such that $x < y$. Using the same argument from the previous paragraph, we can see that $y \notin M_P$. This implies that $Y = \{y\}$. So $Y < Z$ implies that there is a $z \in Z$ such that $y < z$. So we have $x < y < z$, implying that $x < z$. Thus $X < Z$ and so \leq is transitive. \square

By Theorem 5.5 whether or not a poset with a $\hat{1}$ corresponds to a Gorenstein variety depends on the Möbius function. Because of this, it is prudent to understand how the Möbius function behaves when using the tree relation. If we return to our example poset P in Fig. 5, we see that when we took the quotient, none of the Möbius values of the elements outside of M_P changed. This is because when the trees collapsed to a single element, the Möbius value (in the quotient) of the single element is -1 which is the same as the sum of the Möbius values in the tree before it was collapsed. As we see in the next lemma, this property is not unique to our example.

Lemma 5.13 *Let P be a poset with a $\hat{1}$. Suppose that*

- (1) M_P is acyclic.
- (2) P satisfies the cc condition.

Let \sim be the tree relation on P . Then for all $x \in P \setminus M_P$

$$\mu_{\hat{P}}(x) = \mu_{\widehat{P/\sim}}(X),$$

where X is the equivalence class containing x .

Proof We induct on the number of elements strictly below x not in M_P . Suppose that this number is 0. In other words, x is only above elements of M_P . Let T_1, T_2, \dots, T_k be the connected components of $\langle x \rangle \downarrow \cap M_P$. By Lemma 5.6 the sum of Möbius values of a tree is -1 and so

$$\mu_{\widehat{P}}(x) = k - 1.$$

Since $x \notin M_P$, we know that x is in its own equivalence class, X . In P/\sim , each tree T_i has been collapsed to a single element and these elements are minimal in P/\sim . Moreover because of the assumption (2) applied to the connected downset $\langle x \rangle \downarrow$ and the definition of a quotient poset, $\langle X \rangle \downarrow \setminus X = \{T_1, T_2, \dots, T_k\}$. Since the T_i 's are minimal in P/\sim , we have that

$$\mu_{\widehat{P}/\sim}(T_i) = -1$$

and so

$$\mu_{\widehat{P}/\sim}(X) = k - 1$$

Thus, in this case, $\mu_{\widehat{P}}(x) = \mu_{\widehat{P}/\sim}(X)$ and so the base case holds.

Now suppose that the number of elements below x outside of M_P is nonzero. As before, let $X = \{x\}$. By the inductive hypothesis, if $Y < X$ and Y contains no elements of M_P , then

$$\mu_{\widehat{P}}(y) = \mu_{\widehat{P}/\sim}(Y)$$

where $Y = \{y\}$. Moreover, by the definition of the quotient poset, if $Y = \{y\}$ then $Y < X$ if and only if $y < x$. Using these facts, the definition of the Möbius function, and condition (2), we see that

$$\mu_{\widehat{P}}(x) = \mu_{\widehat{P}/\sim}(X)$$

and so the result holds by induction. \square

By Lemma 5.6, we know that when T is a tree, the sum of the Möbius values is -1 . This agrees with the Möbius value of the tree (as a single element) in the P/\sim . In other words, the sum of the Möbius values in that equivalence class is the Möbius value of the equivalence class in the quotient. Every other equivalence class only contains a single element. By Lemma 5.13, under the conditions given in that lemma, the same summation property holds for these other elements in the quotient. Thus, we have the following.

Corollary 5.14 *Let P be a poset with a $\hat{1}$. Suppose that*

- (1) M_P is acyclic.
- (2) P satisfies the cc condition.

Let \sim be the tree relation on P . Then for all $X \in P/\sim$,

$$\widehat{\mu_{P/\sim}}(X) = \sum_{x \in X} \mu_{\widehat{P}}(x).$$

Now that we have an understanding of how the Möbius function behaves when taking quotients, we move to understanding how connected downsets behave. As we will see in the following technical lemma, there is a mapping between connected downsets of P and P/\sim that preserves the sum of the Möbius values.

Lemma 5.15 *Let P be a poset with a $\widehat{1}$. Suppose that*

- (1) M_P is acyclic.
- (2) P satisfies the cc condition.

Let \sim be the tree relation on P and let $C(P)$ and $C(P/\sim)$ be the set of connected downsets of P and P/\sim . Define a map $\psi : C(P) \rightarrow C(P/\sim)$ by $\psi(A) = \{X \in P/\sim \mid X \cap A \neq \emptyset\}$. Then the following hold.

- (a) ψ is a well-defined surjection.
- (b) If $A, B \in \psi^{-1}(D)$, then

$$\sum_{x \in A} \mu_{\widehat{P}}(x) = \sum_{x \in B} \mu_{\widehat{P}}(x).$$

- (c) For all $A \in C(P)$,

$$\sum_{x \in A} \mu_{\widehat{P}}(x) = \sum_{X \in \psi(A)} \widehat{\mu_{P/\sim}}(X).$$

Proof First, let us prove (a). We start by showing ψ is well-defined. Suppose that A is a connected downset of P . Let $X, Y \in \psi(A)$ and let $x \in X \cap A$, $y \in Y \cap A$. Since $x, y \in A$ and A is connected, there exists an $x - y$ path in A , say $w_0 w_1 w_2 \dots w_k w_{k+1}$ where $x = w_0$ and $y = w_{k+1}$. Let W_i be the equivalence class containing w_i . Since $w_i \in A$, $W_i \in \psi(A)$. Since w_i is comparable to w_{i+1} in P , W_i and W_{i+1} are comparable or equal in P/\sim . In any case, after removing repeated vertices, we see that $X W_1 W_2 \dots W_k Y$ is a path in P/\sim whose vertices are in $\psi(A)$. It follows that $\psi(A)$ is connected.

Next, let us show that $\psi(A)$ is a downset. Suppose that $Y \in \psi(A)$ and let $X < Y$ in P/\sim . Then since $X \not\leq Y$, it must be the case that Y is not minimal in P/\sim . Then by definition of the tree relation, $Y = \{y\}$ for some $y \in P$. Since $Y \in \psi(A)$ and Y only has one element, this element must be in A . That is, $y \in A$. Now $X < Y$ implies that there is a $x \in X$ below an element of Y . Since Y only has one element, we have that $x < y$. Since A is a downset, $x \in A$. Thus, $X \in \psi(A)$ and so $\psi(A)$ is a downset. We conclude that ψ is well-defined.

Let us now show that ψ is a surjection. Let D be a connected downset of P/\sim . Let $A = \{a \in P \mid a \in Z \text{ for some } Z \in D\}$. In other words, A is the union of the equivalence classes contained in D . Clearly, if $A \in C(P)$, then $\psi(A) = D$, so it suffices to

show A is connected and a downset. First, we show A is connected. Let $x, y \in A$ and let X, Y be the corresponding equivalence classes containing x and y . Then we have that $X, Y \in D$. Since D is connected, there exists a path $W_0 W_1 W_2 \dots W_k W_{k+1}$ in D where $X = W_0$ and $Y = W_{k+1}$. For each i , if W_i and W_{i+1} are both not minimal in P/\sim , then $W_i = \{w_i\}$ and $W_{i+1} = \{w_{i+1}\}$ for some elements w_i and w_{i+1} . The fact that W_i and W_{i+1} are comparable in P/\sim implies that w_i and w_{i+1} are comparable in P . Thus there is a path from w_i to w_{i+1} . If W_j is minimal in P/\sim , then W_j corresponds to a tree in M_P . Moreover, the fact that $W_{j-1} W_j W_{j+1}$ is a path in D implies that W_{j-1} and W_{j+1} are not minimal. It follows that $W_{j-1} = \{w_{j-1}\}$ and $W_{j+1} = \{w_{j+1}\}$ for some elements w_{j-1} and w_{j+1} . Since W_{j-1} and W_j are comparable in P/\sim , we know that there is an element $w \in W_j$ comparable to w_{j-1} . Similarly, there is an element $w' \in W_j$ comparable to w_{j+1} . Now all the elements of W_j are in A and since W_j is a tree and thus connected, there is a path from w to w' in A . Thus, there is a path from w_{j-1} to w_{j+1} in A . By applying these ideas to the elements in $W_0 W_1 W_2 \dots W_k W_{k+1}$, we see that we can find a path from x to y in A . Thus A is connected.

Next, we show that A is a downset. Let $y \in A$ and let $x < y$ with $x \in X, y \in Y$. Then $Y \in D$. Since $x < y$ in P , $X \leq Y$ in P/\sim . Since D is a downset this implies that $X \in D$. It follows that $x \in A$ and so A is a downset. Thus ψ is a surjection and so (a) holds.

Now we prove (b). Let $A, B \in \psi^{-1}(D)$. We claim that A and B can only differ in values of M_P . To see why, suppose that there is an $x \notin M_P$ such that $x \in A$. Let X be the equivalence class containing x . Then $X \in \psi(A)$ and so $\psi(A) = \psi(B)$, implies that $X \in \psi(B)$. Since $x \notin M_P$, $X = \{x\}$ and so x must be in B .

We now break into two cases depending on if $A \subseteq M_P$ or not. If $A \subseteq M_P$, then since A is connected, A is a tree completely contained in M_P and so by Lemma 5.6 the sum of the Möbius values of A is -1 . Since the A and B agree except possibly for elements of M_P and since B is connected, the sum of the Möbius values of B is -1 as well. Thus (b) holds in this case.

Now suppose that A is not contained in M_P and suppose that there are k connected components of M_P that have nontrivial intersection with A . Since A is connected, for each such connected component, there is an element of $A \setminus M_P$ above some element of the component. Since A and B agree except possibly for elements of M_P , we see that there must also be k connected components of M_P that intersect nontrivially with B . By the cc condition, $cc(A \cap M_P) = k = cc(B \cap M_P)$. By Lemma 5.6 each connected component of $A \cap M_P$ and $B \cap M_P$ contributes -1 to the total sum of Möbius values for A and B . Since the two sets agree except possibly at M_P , we see that (b) holds in this case too.

Lastly let us prove (c). By (b), the sum of Möbius values of any preimage of $\psi(A)$ is the same. Thus it suffices to show the result when $A = \{a \in P \mid a \in X \text{ for some } X \in \psi(A)\}$. Then A contains all the elements of the equivalence classes of $\psi(A)$. Thus,

$$\sum_{x \in A} \mu_{\widehat{P}}(x) = \sum_{X \in \psi(A)} \sum_{x \in X} \mu_{\widehat{P}}(x) = \sum_{X \in \psi(A)} \mu_{\widehat{P}/\sim}(X)$$

where the last equality holds by Corollary 5.14. \square

The surjection ψ described in Lemma 5.15, provides a mapping between connected downsets of P and P/\sim that preserves the sum of the Möbius values. Thus we immediately get the following.

Lemma 5.16 *Let P be a poset with a $\hat{1}$. Suppose that*

- (1) M_P is acyclic.
- (2) P satisfies the cc condition.

Let \sim be the tree relation on P . Then U_P is Gorenstein if and only if $U_{P/\sim}$ is Gorenstein.

We will use the previous lemma to provide a characterization and an algorithm to determine when U_P is Gorenstein provided P has a $\hat{0}$ or $\hat{1}$. However, before we do that, we need to further discuss the cc condition. Clearly we do not want to actually check all connected downsets for the cc condition, since if we already knew all the connected downsets, we could just sum the Möbius values in these downsets to determine if U_P is Gorenstein. Instead, we will show the cc condition is equivalent to a condition that, when the number of connected components of M_P is small compared to the number of elements of P , can be checked efficiently.

Definition 5.17 Let P be a poset with a $\hat{1}$ and suppose that M_P is acyclic and contains k connected components. For all sets $S \subseteq P \setminus M_P$ such that $1 \leq |S| \leq k$, let T_1, T_2, \dots, T_m be the connected components of M_P such that $\langle S \rangle \downarrow \cap T_i \neq \emptyset$. Let $A_j = \langle S, T_1, T_2, \dots, T_{j-1}, T_{j+1}, \dots, T_m \rangle \downarrow$. We say that P satisfies the tree downset condition if whenever A_j is connected we have that $cc(A_j \cap M_P) = m$.

Clearly, the cc condition implies the tree downset condition. However, as we will see, the tree downset condition is actually equivalent to it. This equivalence means that if M_P has k connected components, we only need to look at downsets generated by k elements and the connected components of M_P below them. This can significantly improve the efficiency of checking the cc condition. For example, if M_P is connected, the tree downset condition says we need to only check principal downsets for the cc condition.

In the proof of the next lemma we will need some new terminology. Given a path in the Hasse diagram of P , $v_1 v_2 \dots v_\ell$, we say v_i is a *peak* if $v_i > v_{i-1}, v_{i+1}$. We say that a v_j is a *valley* if $v_j < v_{j-1}, v_{v_j+1}$. See Fig. 6 for an example of peaks and valleys.

Lemma 5.18 *Let P be a poset with a $\hat{1}$ such that M_P is acyclic. Then P satisfies the cc condition if and only if P satisfies the tree downset condition.*

Proof First, note that the forward direction is immediate given the definition of the cc condition and the tree downset condition. Thus, we may now focus on the backwards direction. Suppose that P satisfies the tree downset condition, but there is a connected downset C where the number of connected components of M_P that intersect non-trivially with C is not $cc(C \cap M_P)$. Then it must be the case that there are elements $a, b \in C \cap M_P$ which are connected in M_P , but not connected in $C \cap M_P$.

Since C is connected, we know that there is an $a - b$ path in C . Among all such $a - b$ paths, let Q be a path which uses the minimal number of peaks outside of M_P .




Fig. 6 Peaks are given in blue and valleys are given in red

We claim that Q has at most $k - 1$ peaks that are not in M_P . To see why, suppose this was not the case and let w_1, w_2, \dots, w_k be the first k peaks of Q not in M_P .

For each i , let T_i be a tree of M_P that has an element below the left valley of the peak w_i . Moreover, let T_{k+1} be a tree of M_P that has an element below the right valley of w_k . Since M_P has k trees, the sequence $T_1, T_2, \dots, T_k, T_{k+1}$ has a repeated element. Call this repeated element T . By definition, T is below a left valley of a peak in $\{w_1, w_2, \dots, w_k\}$. Moreover, we claim that T is also below a right valley of a peak in $\{w_1, w_2, \dots, w_k\}$. If $T = T_{k+1}$, this is true by definition of T_{k+1} . If not, then T is below a left valley of a peak say v . We claim that T must be below the right valley of the peak immediately preceding v in the list w_1, w_2, \dots, w_k . Call this peak u . Clearly if the left valley of v is the right valley of u this is true. If not, it is because the left valley of v and the right valley of u are in M_P and the portion of the path between these valleys stay in M_P . But then this means that both these valleys are in T . See Fig. 6 for a visualization. We conclude that T is below the right valley of some peak in w_1, w_2, \dots, w_k .

Let w_i and w_j be such that T is below the left valley of w_i and T is below the right valley of w_j and such that w_i and w_j chosen so that they are closest in the list $w_1, w_2, \dots, w_k, w_{k+1}$ with this property. Let A be downset generated by $\{w_i, w_{i+1}, \dots, w_j\}$ together with all the trees of M_P below an element of $\{w_i, w_{i+1}, \dots, w_j\}$ except for T . We will show that all the peaks (including those in M_P) of the portion of Q between w_i and w_j are in A . Since the path between w_i and w_j is connected, the downset the path generates is connected. Moreover since the path is generated by its peaks and since the additional trees added to A are below elements of the path and the trees are connected themselves, we see that A will be connected if we can show all the peaks are in A .

By definition, all the peaks outside of M_P on the portion of Q from w_i to w_j are in A . Now suppose that p is a peak in M_P in the portion of Q between w_i and w_j . Any time the path goes out of M_P , it creates a peak outside of M_P to go back down into M_P . Since the trees of M_P are connected components this means that once the path goes into M_P it stays in the same tree of M_P until the path leaves M_P . Thus any peak in M_P is in a tree that was below a peak outside of M_P . Since all the trees except for T are completely contained in A , as long as these peaks in M_P are not in T , they are in A . Because we chose w_i and w_j to be as close as possible, the peaks in M_P on this portion of Q cannot be in T . Thus, all the peaks of Q between w_i and w_j are in A and so A is connected.

Let t_1 be an element of T that is below the left valley of w_i and let t_2 be an element of T that is below the right valley of w_j . A is generated by no more than k elements together with all but one tree of M_P below it. We claim that there is a $t_1 - t_2$ path completely contained in $\langle w_i, w_{i+1}, \dots, w_j \rangle \downarrow \cap M_P$. If this was not the case, then this would contradict the tree downset condition. Since $w_i, w_{i+1}, \dots, w_j \in C$, this path is completely contained in $C \cap M_P$. However, we claim this allows us to modify Q to get a new path from a to b that has less peaks outside of M_P . Let Q' be the path from a to b which follows along Q until we reach the left valley of w_i , then travels down to t_1 , takes the path in M_P from t_1 to t_2 , then goes up to the right valley of w_j . Note that this new path Q' does not contain any of the peaks in w_i, w_{i+1}, \dots, w_j . When we descend from the left valley of w_i to t_1 we do not create any new peaks. Similarly we do not create new peaks when ascending from t_2 to the right valley of w_j . We may create new peaks between t_1 and t_2 , but these peaks are wholly contained in M_P . As a result, Q' has less peaks outside of M_P than Q does, contradicting the minimality of peaks outside of M_P of Q . We conclude that Q has at most $k - 1$ peaks.

To finish, apply the same reasoning we did before to Q . Since Q has at most $k - 1$ peaks outside of M_P , we see that this allows us to find an $a - b$ path in $C \cap M_P$, contradicting our original assumption. \square

In the next theorem, we provide a characterization of when the variety associated to a poset with a $\hat{1}$ is Gorenstein. To do this, we need some new notation. Let \sim be the tree relation on P . We define P/\sim^k recursively as

$$P/\sim^k = \begin{cases} P & \text{if } k = 0, \\ (P/\sim^{k-1})/\sim & \text{for } k > 0. \end{cases}$$

In other words, P/\sim^k is the poset obtained by applying the tree relation k times to P . See Fig. 7 for an example of a poset and its successive quotients. We are now ready to give one of the main theorems of this section. Recall that $\ell(P)$ denotes the length of P .

Theorem 5.19 *Let P be a poset with a $\hat{1}$. U_P is Gorenstein if and only for all $0 \leq k \leq \ell(P) - 1$ the following hold*

- (1) M_{P/\sim^k} is acyclic.
- (2) P/\sim^k satisfies the tree downset condition.




Fig. 7 Poset P and its successive quotients

Proof (\Rightarrow) Suppose that U_P is Gorenstein. We prove assumptions (1) and (2) hold for all $0 \leq k \leq \ell(P) - 1$ by using strong induction on k . When $k = 0$, the result holds by Lemma 5.8 and Lemma 5.18.

Now suppose that $k > 0$. By the inductive hypothesis, assumptions (1) and (2) hold for P/\sim^j for all $0 \leq j \leq k - 1$. By Lemma 5.18, (2) implies that P/\sim^j satisfies the cc condition for all $0 \leq j \leq k - 1$. Thus, we can repeatedly apply Lemma 5.16 to get that U_P Gorenstein implies that $U_{P/\sim^{k-1}}$ is Gorenstein. Applying the lemma to P/\sim^{k-1} implies that U_{P/\sim^k} is Gorenstein. Then assumptions (1) and (2) hold for P/\sim^k by Lemma 5.8.

(\Leftarrow) Now suppose assumptions (1) and (2) hold for all $0 \leq k \leq \ell(P) - 1$. Then we can repeatedly apply Lemma 5.16 to see that U_P is Gorenstein if and only $U_{P/\sim^{\ell(P)-1}}$ is Gorenstein. Now every time we apply the tree relation to poset its length drops by 1. Thus, applying the quotient $\ell(P) - 1$ times, we are left with a poset consisting of a $\hat{1}$ and minimal elements. Clearly such a poset gives rise of a Gorenstein variety. Thus, $U_{P/\sim^{\ell(P)-1}}$ is Gorenstein and so U_P is Gorenstein as well. \square

The previous theorem allows us to give an algorithm to determine if a poset with a $\hat{0}$ or $\hat{1}$ gives rise to a Gorenstein variety.

Algorithm 5.20 *Algorithm to determine if U_P is Gorenstein*

Input: P a poset with a $\hat{0}$ or $\hat{1}$.

Output: A Yes/No decision if U_P is Gorenstein.

Method:

- (1) If P has a $\hat{0}$, set P to be P^* .
- (2) While $\ell(P) > 1$

- (a) Check the truth value of the following statement.

M_P is acyclic.

If the statement is false, terminate algorithm and return “No”.

- (b) Check the truth value of the following statement.

P satisfies the tree downset condition.

If the statement is false, terminate algorithm and return “No”.

(c) Set P to be P/\sim , where \sim is the tree relation.

(3) Return “Yes”.

At each step of the algorithm, we need to check the tree downset condition. Since this condition depends on the number of connected components in M_P , it is important to understand how the number of connected components change as we take quotients.

Lemma 5.21 *Let P be a poset with k trees in M_P . Then $M_{P/\sim}$ has at most k trees.*

Proof Let T_1, T_2, \dots, T_k be the trees of M_P . Then T_1, T_2, \dots, T_k are the minimal elements of P/\sim . Every connected component of $M_{P/\sim}$ must contain a minimum element of P/\sim . So every connected component of $M_{P/\sim}$ contains some tree in T_1, T_2, \dots, T_k . Moreover, each T_i is in exactly one connected component of $M_{P/\sim}$. It follows that there are at most k trees in $M_{P/\sim}$. \square

In the next theorem, we give a bound on the complexity of Algorithm 5.20. Note that, the brute force method which would require first enumerating all downsets is known to be $\#P$ -complete as we have previously remarked. So, if the number of connected components of M_P is small compared to $|P|$, the algorithm can be significantly quicker than the brute force approach (assuming $\#P$ -complete problems are not actually easy). Toward formalizing this, we analyze Algorithm 5.20 in terms of parameterized complexity [6]. The parameterized complexity class XP (see e.g. [2, p. 7]) consists of problems such that given a pair of a problem instance and a parameter k there is an algorithm solving the problem with time bounded by $f(k) \cdot N^{g(k)}$ where N is the size of the problem instance and both f and g are computable functions. These problems are sometimes called “slice-wise polynomial” since for a fixed k the runtime is polynomial in N . We call an algorithm demonstrating membership in XP an *XP-algorithm*. In our situation, a problem instance will be given by a poset P on n elements. The parameter k we use is the number of connected components of M_P . We note that this parameter can be efficiently computed from the poset.

Theorem 5.22 *Algorithm 5.20 is a valid XP-algorithm which can be made to run in $O(kn^{k+3})$ time where $n = |P|$ and k is the number of connected components of M_P .*

Proof The validity of the algorithm follows from Theorem 5.19 and the fact that if $P = M_P$, then U_P is Gorenstein.

Now we show the result concerning the time complexity. First, the algorithm needs to determine M_P . This can be done by first finding all the minimal elements of P which can be done in $O(n^2)$ time by comparing elements. Then one must find the elements that cover only minimal elements. This too can be done in $O(n^2)$ time. Because we will need it later, we can also find the connected components of M_P again in $O(n^2)$ time using a depth first search. Next, the algorithm needs to determine if M_P is acyclic. Since we know the number of connected components of M_P , we can just count the number of edges in M_P to decide this.

After determining if M_P is acyclic, the algorithm must check the tree downset condition of P (Definition 5.17). Let $S \subseteq P \setminus M_P$ with $|S| \leq k$ where k is the number of connected components of M_P . To check the tree downset condition for S , we must find all the connected components that intersect nontrivially with $\langle S \rangle_\downarrow$. We can compute $\langle S \rangle_\downarrow$ in $O(kn)$ time by comparing each element of P with each element of S and from these comparisons we can also determine which connected components of M_P intersect nontrivially with $\langle S \rangle_\downarrow$.

Next, for each j we will compute A_j given in Definition 5.17 and check if it is connected. Computing A_j can be done in linear time given that we have already computed $\langle S \rangle_\downarrow$. Again, using a depth first search, determining connectedness of A_j can be done in $O(n^2)$ time. For those that are connected, we need to count the number of connected components of $A_j \cap M_P$. This can be done in $O(n^2)$ time. Since there at most k possible A_j 's to check, we see that for each set S , checking the tree downset condition for S , can be done in $O(kn^2)$ time. Now there are $O(n^k)$ sets to check, so this step can be done in $O(kn^{k+2})$ time. Lastly, if the tree downset condition is satisfied we must compute P/\sim . This can be handled by recording some of the computations performed above since P/\sim is determined by comparing elements of M_P with elements of $P \setminus M_P$.

We conclude that one iteration of steps (1) and (2) of the algorithm can be run in $O(kn^{k+2})$ time. By Lemma 5.21, the number of connected components of $M_{P/\sim}$ is at most k and so each iteration can be run in $O(kn^{k+2})$ time. Since the algorithm runs no more than n times, we see that the full algorithm can run in $O(kn^{k+3})$ time as claimed. \square

We finish this section by noting that when M_P is connected, we can easily check if U_P is Gorenstein. The main reason for the simplicity is that if we quotient out by the tree relation and M_P is connected, P/\sim is bounded and so $U_{P/\sim}$ is Gorenstein. We leave the details of the proof for the following theorem to the reader.

Theorem 5.23 *Let P be a poset with a $\hat{1}$ such that M_P is connected. Then the following are equivalent.*

- (a) U_P is Gorenstein.
- (b) For all $x \in P$, $\langle x \rangle_\downarrow \cap M_P$ is a tree.
- (c) For all $x \in P \setminus M_P$, $\mu_{\hat{P}}(x) = 0$.

6 Length 1 Posets

In this section we restrict our attention to posets of length 1. That is, posets in which every element is either minimal or maximal.

6.1 \mathbb{Q} -Gorenstein Implies Gorenstein

We start with a lemma which determines the values an r -Gorenstein labeling must take on any biconnected length 1 poset.




Fig. 8 A biconnected poset of length 1 which has the same number of minimal and maximal elements but is not Gorenstein

Lemma 6.1 *If P is a length 1 biconnected poset and $\phi : P \rightarrow \mathbb{Z}$ is an r -Gorenstein labeling, then $\phi(x) = r$ for all maximal elements $x \in P$ and $\phi(y) = -r$ for all minimal elements $y \in P$.*

Proof For each maximal element $x \in P$ we have that $A = \{x\}$ is an upset. Since P is biconnected, $\dim(A) = 1$ so $\phi(x) = r$ because ϕ is an r -Gorenstein labeling. Similarly, by Proposition 3.1, for any minimal $y \in P$, we must have that $\phi(y) = -r$

□

Proposition 6.2 *If P is a length 1 biconnected poset with an r -Gorenstein labeling, then P has the same number of minimal and maximal elements.*

Proof For a length 1 poset each element is either a minimal or maximal element and cannot be both since the poset is connected. Thus if ϕ is an r -Gorenstein labeling its values are completely determined by Lemma 6.1. Let α be the number of maximal elements of P and β denote the number of minimal elements. Then we have that

$$0 = \sum_{z \in P} \phi(z) = \alpha r - \beta r.$$

So $\alpha = \beta$ and the result is proven. □

Let us give an example which shows that the converse of Proposition 6.2 is false. The length 1 poset in Fig. 8 is biconnected with the same number of minimal and maximal elements, but is not r -Gorenstein. To see why, note that all the maximal elements would have to be labeled with a value of r . Now let A be the upset generated by 2, then $cc(A) = 1 = cc(\bar{A})$ and so it follows that 2 must be labeled by $-2r$. Thus by Lemma 6.1, the poset cannot be r -Gorenstein.

We now give one of the main theorems of this section.

Theorem 6.3 *Let P be a length 1 poset. Then U_P is Gorenstein if and only if U_P is \mathbb{Q} -Gorenstein.*

Proof The backwards direction is trivial given the definitions of \mathbb{Q} -Gorenstein and Gorenstein. By Theorem 4.3 we may assume that P is biconnected. So by Lemma 6.1, the r -Gorenstein labeling is given by $\phi(x) = \pm r$. It follows that the labeling $\frac{1}{r}\phi(x)$ is a Gorenstein labeling of P . □

6.2 Some Infinite Families

By Theorem 6.3 we need to only look at Gorenstein labelings. We will now give some infinite families of length 1 posets admitting Gorenstein labelings.

Proposition 6.4 *Let P be a length 1 poset which is 2-regular and 2-connected. Then U_P is Gorenstein.*

Proof For such a poset P , it must be that P is a $2k$ -cycle. This is a balanced bipartite graph with k elements in each part of the bipartition. The bipartition splits P into its minimal elements and its maximal elements. Let $\phi : P \rightarrow \mathbb{Z}$ be defined by

$$\phi(x) = \begin{cases} 1 & \text{if } x \text{ is maximal} \\ -1 & \text{if } x \text{ is minimal} \end{cases}$$

which is the only possibility for a Gorenstein labeling by Lemma 6.1. First, note that since there are k minimal elements and k maximal elements, we have that

$$\sum_{x \in P} \phi(x) = k - k = 0.$$

Now suppose that A is an upset such that $A \neq P$ and $\dim(A) = 1$. Then A is connected. The only proper connected subgraphs of a cycle are paths and so A is a path. Since A is an upset and a path, it must be that there are j minimal elements in the path and $j + 1$ maximal elements. It follows that for any upset A which is connected,

$$\sum_{x \in A} \phi(x) = (j + 1) - j = 1.$$

Thus U_P is Gorenstein. \square

As one can see from the poset in Fig. 8, we cannot just replace 2-regular with 3-regular in the previous proposition. However, there are n -regular length 1 posets which give rise to Gorenstein varieties as we see next. Throughout the remainder of this section, we use $K_{n,n}$ to denote the poset whose Hasse diagram is the complete bipartite graph with n minimal elements and n maximal elements.

Proposition 6.5 *If P is $K_{n,n}$, then U_P is Gorenstein.*

Proof First note that P has n minimal elements each of which is less than each of the n maximal elements. Let the minimal elements of P be a_1, a_2, \dots, a_n . Also, let the maximal elements of P be b_1, b_2, \dots, b_n . Let $\phi : P \rightarrow \mathbb{Z}$ be given by $\phi(a_i) = -1$ and $\phi(b_i) = 1$ for all $1 \leq i \leq n$ which is the only possibility for a Gorenstein labeling by Lemma 6.1. It is clear that $\sum_{x \in P} \phi(x) = 0$. Now if $A \subseteq P$ is an upset with $\dim(A) = 1$, then either $A = \{b_i\}$ for some $1 \leq i \leq n$ or else $\overline{A} = \{a_i\}$ again for some $1 \leq i \leq n$. In any case $\sum_{x \in A} \phi(x) = 1$ for any upset $A \subseteq P$ with $\dim(A) = 1$. Therefore U_P is Gorenstein. \square

By Proposition 6.5, we know that there are n -regular, n -connected posets which have Gorenstein labelings, namely $K_{n,n}$. Additionally, by Proposition 6.4, we know that there are 2-regular, 2-connected posets which give rise to Gorenstein varieties other than $K_{2,2}$. This naturally brings up the question of if there are n -regular, n -connected posets, other than $K_{n,n}$, whose variety is Gorenstein when $n \geq 3$. We have been able verify that $K_{3,3}$ is the only 3-regular, 3-connected length 1 poset whose variety is Gorenstein. However, at this time our methods do not seem to generalize when $n \geq 4$ and thus this case remains open.

7 Conclusion

We now conclude with some discussion of open problems. First we have the following question on recognizing when an r -Gorenstein labeling exists.

Question 7.1 *Is there a good characterization in terms of P of when U_P is Gorenstein or \mathbb{Q} -Gorenstein?*

In Question 7.1 the word “good” means a characterization which is not simply the definition of being \mathbb{Q} -Gorenstein or Gorenstein and preferably gives an efficient means of recognizing the Gorenstein property from the poset like is done in Algorithm 5.20 for posets with $\hat{0}$ or $\hat{1}$ when the number of connected components of M_P is small. A braid cone is smooth if and only if the Hasse diagram of the poset is a tree [18, Prop. 3.5 (9) and Corollary 3.10]. Since smooth implies Gorenstein, tree posets will be included in any characterization that answers Question 7.1. Another easy to describe class of posets we know have a Gorenstein labeling is bounded posets from Theorem 3.3. The largest class we understand, which contains tree posets and bounded posets, comes from the characterization in Theorem 5.19 combined with the reduction to biconnected components from Theorem 4.3. A question, possibly more approachable than in the general case, is the following for which we treated some cases of in Sect. 6.

Question 7.2 *Is there an efficient way to decide if U_P is Gorenstein when P is a length 1 poset?*

By Proposition 6.2 and reduction to the biconnected case previously mentioned, we see that Question 7.2 becomes a problem about neighborhoods in balanced bipartite graphs with a “top” and “bottom” set of vertices corresponding to maximal and minimal elements respectively. Approaches to this question could be to look for a clever algorithm or look for a reduction with an NP-complete problem.

Whether it is the length 1 case or the general case, in order to decide if U_P is Gorenstein, we need understand the ray generators of σ_P . As we mentioned before, doing this may be difficult since the ray generators come from upsets which are known to be hard to enumerate [17]. This makes the problem of deciding Gorensteinness potentially difficult for generic posets.

In Sect. 5 and Sect. 6 we showed that \mathbb{Q} -Gorenstein implies Gorenstein for posets with a $\hat{0}$ or a $\hat{1}$ and for length 1 posets. Moreover, using SageMath [21], we have been able to verify that for all posets on up to 10 elements, \mathbb{Q} -Gorenstein implies Gorenstein. This supporting data is available from the corresponding author on reasonable request. This leads us to the following conjecture.

Conjecture 7.3 *Let P be a poset, then U_P is Gorenstein if and only if U_P is \mathbb{Q} -Gorenstein.*

We note that braid cones have a nice property related to smoothness of a similar flavor to this conjecture. In particular, a braid cone is simplicial (i.e. ray generators are a basis of $N_{\mathbb{R}}$) if and only if it is smooth (i.e. ray generators are a basis of N) [18, Cor. 3.10]. Geometrically this means that the corresponding toric variety is an orbifold if and only if it is smooth.

Acknowledgements The authors wish to thank Victor Reiner for helpful comments on this work.

References

1. Boussicault, A., Féray, V., Lascoux, A., Reiner, V.: Linear extension sums as valuations on cones. *J. Algebraic Combin.* **35**(4), 573–610 (2012). <https://doi.org/10.1007/s10801-011-0316-2>
2. Cygan, M., Fomin, F.V., Kowalik, Ł., Lokshtanov, D., Marx, D., Pilipczuk, M., Michał Pilipczuk, S.Sa.: Parameterized Algorithms. Springer, Cham (2015). <https://doi.org/10.1007/978-3-319-21275-3>
3. Cox, D.A., Little, J.B., Schenck, H.K.: Toric Varieties. Graduate Studies in Mathematics, vol. 124. American Mathematical Society, Providence, RI (2011). <https://doi.org/10.1090/gsm/124>
4. Cox, D.A.: Update on Toric Geometry. In: Geometry of Toric Varieties, Volume 6 of Sémin. Congr., pp. 1–41. Soc. Math. France, Paris (2002)
5. Dorpalen-Barry, G., Kim, J.S., Reiner, V.: Whitney numbers for poset cones. *Order* (2021). <https://doi.org/10.1007/s11083-020-09541-4>
6. Downey, R.G., Fellows, M.R.: Parameterized Complexity. Monographs in Computer Science, Springer, New York (1999). <https://doi.org/10.1007/978-1-4612-0515-9>
7. Féray, V., Reiner, V.: P -partitions revisited. *J. Commut. Algebra* **4**(1), 101–152 (2012). <https://doi.org/10.1216/JCA-2012-4-1-101>
8. Harary, F.: Graph Theory. Addison-Wesley Publishing Co., Reading (1969)
9. Hibi, T.: Distributive lattices, affine semigroup rings and algebras with straightening laws. In: Commutative Algebra and Combinatorics (Kyoto, 1985), volume 11 of Adv. Stud. Pure Math., pp. 93–109. North-Holland, Amsterdam (1987). <https://doi.org/10.2969/aspm/01110093>
10. Hibi, T., Lason, M., Matsuda, K., Michałek, M., Vodicka, M.: Gorenstein graphic matroids. *Israel J. Math.* (2021). <https://doi.org/10.1007/s11856-021-2136-y>
11. Hochster, M.: Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes. *Ann. Math.* **2**(96), 318–337 (1972). <https://doi.org/10.2307/1970791>
12. Khachiyan, L., Boros, E., Borys, K., Elbassioni, K., Gurvich, V.: Generating all vertices of a polyhedron is hard. *Discrete Comput. Geom.* **39**(1–3), 174–190 (2008). <https://doi.org/10.1007/s00454-008-9050-5>
13. Kölbl, M.: Gorenstein graphic matroids from multigraphs. *Ann. Comb.* **24**(2), 395–403 (2020). <https://doi.org/10.1007/s00026-020-00495-3>
14. Lasoń, M., Michałek, M.: On algebraic properties of matroid polytopes (2020). [arXiv:2005.08359](https://arxiv.org/abs/2005.08359). <https://arxiv.org/abs/2005.08359>
15. Lee, E., Masuda, M.: Generic torus orbit closures in Schubert varieties. *J. Combin. Theory Ser. A* **170**, 105143, 44 (2020). <https://doi.org/10.1016/j.jcta.2019.105143>
16. Machacek, J.: Strong factorization and the braid arrangement fan. *Tohoku Math. J. (2)* **73**(1), 99–104 (2021). <https://doi.org/10.2748/tmj.20191223>
17. Provan, J.S., Ball, M.O.: The complexity of counting cuts and of computing the probability that a graph is connected. *SIAM J. Comput.* **12**(4), 777–788 (1983). <https://doi.org/10.1137/0212053>
18. Postnikov, A., Reiner, V., Williams, L.: Faces of generalized permutohedra. *Doc. Math.* **13**, 207–273 (2008)
19. Reiner, V.: Quotients of Coxeter complexes and P -partitions. *Mem. Am. Math. Soc.* **95**(460):vi+134 (1992). <https://doi.org/10.1090/memo/0460>
20. Sagan, B.E.: Combinatorics: The Art of Counting. Volume 210 of Graduate Studies in Mathematics, American Mathematical Society, Providence (2020)

21. The Sage Developers.: SageMath, The Sage Mathematics Software System (Version 9.4) (2021). <https://www.sagemath.org>
22. Stanley, R.P.: Flag f -vectors and the cd -index. *Math. Z.* **216**(3), 483–499 (1994). <https://doi.org/10.1007/BF02572336>
23. Stanley, R.P.: *Enumerative Combinatorics. Volume 1*, Volume 49 of Cambridge Studies in Advanced Mathematics, 2nd Edn. Cambridge University Press, Cambridge (2012)
24. Stembridge, J.R.: Coxeter cones and their h -vectors. *Adv. Math.* **217**(5), 1935–1961 (2008). <https://doi.org/10.1016/j.aim.2007.09.002>
25. Wagner, D.G.: Singularities of toric varieties associated with finite distributive lattices. *J. Algebraic Combin.* **5**(2), 149–165 (1996). <https://doi.org/10.1023/A:1022469218716>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.