

Manganese(III) Nitrate Complexes as Bench-Stable Powerful Oxidants

Ananya Saju, Matthew R. Crawley, Samantha N. MacMillan, and David C. Lacy*



Cite This: *J. Am. Chem. Soc.* 2024, 146, 11616–11621



Read Online

ACCESS |



Metrics & More



Article Recommendations

*

Supporting Information

ABSTRACT: We report herein a convenient one-pot synthesis for the shelf-stable molecular complex $[\text{Mn}(\text{NO})(\text{OPPh})_3]_2$ (**2**) and describe the properties that make it a powerful and selective one-electron oxidation (deelectronation) reagent. **2** has a high reduction potential of 1.02 V versus ferrocene (MeCN) (1.65 vs normal hydrogen electrode), which is one of the highest known among readily available redox agents used in chemical synthesis. **2** exhibits stability toward air in the solid state, can be handled with relative ease, and is soluble in most common laboratory solvents such as MeCN, dichloromethane, and fluorobenzene. **2** is substitutionally labile with respect to the coordinated (pseudo)halide ions enabling the synthesis of other new Mn(III) nitrate complexes also with high reduction potentials ranging from 0.6 to 1.0 V versus ferrocene.

Introduction. Single-electron oxidation, sometimes referred to as deelectronation,¹ is one of the most fundamental reactions in chemistry. Most very strong oxidants (>0.8 V; all potentials vs Fc/Fc^+ unless noted) such as main group fluorides (e.g., AsF_6^-) or transition-metal hexafluoride compounds (WF_6 , PtF_6 etc.) are impractical because they are difficult to synthesize and/or handle in standard laboratory settings.² Nitrosonium and Ag^+ salts of weakly coordinating anions (WCAs) are also established reagents for deelectronation,³ but, as with most strong oxidants, oxidation is often accompanied by side reactions in addition to electron removal. Some recent progress toward more selective single-electron oxidants such as $[\text{TCNQ} \cdot 4\text{B}(\text{C}_6\text{F}_5)_3]$ (TCNQ = tetracyanoquinodimethane), $[\text{phenazine}^+][\text{Al}(\text{OR}^F)_4]$ (phenazine^F = perfluoro-5,10-bis(perfluorophenyl)-5,10-dihydrophenazine), and aminium-cation carborane-anion pairs, like **T13**,⁴ have been reported, but the multistep syntheses and air and moisture sensitivities limit their utility.⁵ Hence, the search for selective single-electron oxidants continues, driven by the need for ease of synthesis and handling while maintaining high reduction potentials.

We noted that a characteristic of the Mn(III) ion is its high reduction potential, a property that is critical to life,⁶ battery materials,⁷ and oxidation in organic transformations such as alkene difunctionalization,⁸ alkene epoxidation,⁹ and free-radical additions and cyclizations.¹⁰ Within this framework of reactivity, we recently observed that the mononuclear Mn(III) complex $[\text{Mn}^{\text{III}}\text{Cl}_3(\text{OPPh}_3)_2]$ (**1**) is a potent Cl-atom transfer reagent¹¹ and mediates C–H chlorination in specific cases,¹² properties that appear to be intrinsic to the highly reactive MnCl_3 entity. We reasoned that the coordination of pnictogen-oxide ylide triphenylphosphine oxide (Ph_3PO) enabled the room-temperature stabilization of highly reactive MnCl_3 , which otherwise decomposes above -40°C .¹³ As an extension of this rationale, we hypothesized that analogous Mn(III) complexes with weakly coordinating pseudohalide ligands could likewise exhibit desirable bench-stability characteristics while maintain-

ing some of the intrinsic chemical properties, such as the high reduction potential.

Exploring this hypothesis with nitrate as the anion, reported herein, resulted in successful gram-scale synthesis of $[\text{Mn}^{\text{III}}(\text{NO}_3)_3(\text{OPPh}_3)_2]$ (**2**) and demonstration of desirable bench-stability and redox properties for oxidation. **2** has a strikingly high reduction potential of 1.02 V in MeCN (1.65 vs NHE) (Figure 1), which is one of the highest among available stoichiometric oxidants.¹⁴ While other reactants with higher potentials are known,⁵ **2** can be synthesized in a single step on the bench from commercially available reactants, is capable of open-to-air manipulations, and is soluble in common laboratory solvents. The byproducts of oxidation with **2** are $[\text{Mn}^{\text{II}}(\text{NO}_3)_2(\text{OPPh}_3)_2]$ (**9**) and non-nucleophilic nitrate ion.¹⁵ Except for use in exotic cases, where extremely electrophilic products are generated, **2** is an important addition to the growing class of powerful oxidants. **2** also is substitutionally labile with respect to halide/pseudohalide ion metathesis enabling the synthesis of new Mn(III) nitrate complexes, effectively allowing us to tune the reduction potential in a series of strong to very strong oxidants spanning 0.6 to 1.0 V.

Synthesis and Characterization of $[\text{Mn}^{\text{III}}(\text{NO}_3)_3(\text{OPPh}_3)_2]$. A procedure to prepare what is nominally $[\text{Mn}^{\text{III}}(\text{NO}_3)_3]$ (**3**) has been known since 1972,¹⁷ but it requires the use of highly toxic and inconvenient reagents like N_2O_5 and N_2O_4 and requires storage below -14°C (Figure 2). **2** was synthesized from $[\text{Mn}^{\text{III}}(\text{NO}_3)_3]$ via this cumbersome route and, aside from a solid-state magnetic

Received: March 8, 2024

Revised: April 12, 2024

Accepted: April 15, 2024

Published: April 19, 2024




E° (V vs. FeCp_2) in MeCN

Figure 1. Reduction potentials of common air-stable oxidants in MeCN (see the text for citations). (*) Potentials in DCM.




Figure 2. X-ray crystal structure of **2** (ellipsoids shown at 50% probability; H atoms omitted and only one component of disordered atoms for clarity) and material isolated from multigram scale synthesis.



measurement and select vibrational data, virtually nothing was known about it prior to this work.¹⁷ Motivated by the premise that **2** would exhibit a kind of tempered stability like **1**, we aimed to develop convenient ways of preparing **2** and new pnictogen-oxide stabilized complexes of Mn(III); pyridine-*N*-oxide (Opy), Ph_3PO , and triphenylarsine oxide (Ph_3AsO) were the chosen ligands for this study.

Two convenient routes to **2** were devised, both using an acetonitrile solution of trimethylsilylnitrate ($\text{Me}_3\text{SiONO}_2$) generated *in situ* from AgNO_3 and Me_3SiCl .¹⁸ In the first route, 36 equiv of $\text{Me}_3\text{SiONO}_2$ was added to $[\text{Mn}_{12}\text{O}_{12}(\text{OAc})_{16}(\text{H}_2\text{O})_4]\cdot 2\text{HOAc}\cdot 4\text{H}_2\text{O}$ (Mn_{12})¹⁹ and then treated with 24 equiv of Ph_3PO to give **2** in low yields (26% crystalline). The more efficient second route is analogous to the one-pot synthesis that we developed for **1** (Scheme 1),¹¹

Scheme 1. Synthesis of **2**

Previous Work

Can. J. Chem. 1972, 50, 3326



This Work




employing $\text{Me}_3\text{SiONO}_2$ instead of Me_3SiCl . Adding $\text{Me}_3\text{SiONO}_2$ to a mixture of $\text{Mn}(\text{OAc})_2$ and KMnO_4 in a 16:4:1 ratio formed a brown solution, presumably containing **3**, and upon addition of Ph_3PO **2** was obtained in good yields. The reaction is scalable; we routinely perform syntheses that furnish 6 g of crude product (73%, Figure 2) which after recrystallization in a glovebox furnishes 5 g (63%) of crystalline, analytically pure material. In the solid state, **2** is stable to air and can be stored for months without noticeable change on the bench (Figure S5). Performing an entire synthesis (8 g scale) on a Schlenk line under Ar followed by open-to-air filtration does not diminish the yield of the crude product.

X-ray diffraction (XRD) characterization of red-brown crystals of **2** revealed a pseudo-square-pyramidal geometry with the Ph_3PO ligands in the *cis* position (Figure 2), similar to **1**.¹¹ The solid-state ATR-FTIR spectrum of **2** contains a characteristic P–O stretch at 1436 cm^{-1} as well as nitrate stretches at 1528 , 1260 , and 951 cm^{-1} . The solution-state magnetic moment of **2** in DCM is consistent with a high-spin Mn(III) center ($\mu_{\text{eff}} = 5.46 \mu_{\text{B}}$), and the X-band EPR spectrum (77 K) is silent.

Synthesis of Other Mn(III) Nitrate Complexes. Mn(III) nitrate complexes with other pnictogen-oxide ligands were synthesized via **3** that was prepared directly from $\text{Mn}(\text{OAc})_2$ and KMnO_4 (Scheme 2). Addition of 2 equiv of Ph_3AsO to **3** produced $[\text{Mn}^{\text{III}}(\text{NO}_3)_3(\text{OAsPh}_3)_2]$ (**4**) in 61% crystalline yield. XRD analysis of deep purple crystals of **4** revealed a pseudo-trigonal-bipyramidal geometry with the Ph_3AsO ligands in the *trans* position (Figure 3). Addition of 2 equiv of Opy to **3** did not produce the expected $[\text{Mn}^{\text{III}}(\text{NO}_3)_3(\text{Opy})_2]$, and instead the mononitrate Mn(III) complex salt $[\text{Mn}^{\text{III}}(\text{NO}_3)(\text{MeCN})(\text{Opy})_2][\text{Mn}^{\text{II}}(\text{NO}_3)_4]$ (**5**) was obtained in 64% crystalline yield (Figure 3). The six-coordinate manganese center of **5** is in the +3 oxidation state.

Scheme 2. Synthesis of 3-5



The bond lengths and angles of the other manganese center are consistent with the known anion $[\text{Mn}^{\text{II}}(\text{NO}_3)_4]^{2-}$.²⁰ While 2 and 4 are bench stable in the solid state, 5 readily undergoes hydration when exposed to air. Furthermore, 2 and 4 are soluble in polar organic solvents like DCM, MeCN, and THF while 5 is only soluble in MeCN, consistent with its salt-like state in contrast to neutral 2. In fact, 2 is soluble in

fluorobenzene (2.8 mg/mL) and is sparingly soluble in toluene and benzene (<1 mg/mL).

We also prepared mixed nitrato dichlorido Mn^{III} complexes via halide/pseudohalide metathesis (Scheme 3). Reaction of 1 and 2 in a 2:1 ratio cleanly produces 6 (Figure 3). Similarly, complex $[\text{Mn}^{\text{III}}\text{Cl}_2(\text{NO}_3)(\text{OAsPh}_3)_2]$ (8) was synthesized from 4 and $[\text{Mn}^{\text{III}}\text{Cl}_3(\text{OAsPh}_3)_2]$ (7) (Figure 3).²¹

Electrochemical Investigation of the Mn^{III} Complexes. The complexes prepared here were analyzed with cyclic voltammetry (CV) and differential pulse voltammetry (DPV) in DCM, MeCN (Figure 4), and THF (Figure S47). The reduction potentials are remarkably high and span the range of 0.6 to 1.0 V (Table 1).

To confirm that 2 acts as a one-electron oxidant, we titrated a solution of ferrocene with 2 and found that one equivalent was required to cleanly generate FeCp_2^+ in DCM (Figure 4). The same titration was conducted for $[\text{Fe}^{\text{II}}(\text{bipy})_3](\text{PF}_6)_2$ (0.66 V in MeCN) and $\text{N}(4\text{-bromophenyl})_3$ (0.70 V in DCM), and likewise a clean one-electron transformation occurred to $[\text{Fe}^{\text{III}}(\text{bipy})_3]^{3+}$ ²² and $\text{N}(4\text{-bromophenyl})_3^{+}$, respectively.

The byproducts of oxidation with 2 are a weakly coordinating nitrate anion and 9, which are easy to separate





Figure 3. Molecular structures determined with XRD for 4-8 (ellipsoids shown at the 50% probability). H atoms are omitted, and only one component of disordered atoms is shown for clarity.

Scheme 3. Synthesis of 6-8



from reaction mixtures and will not likely interfere with subsequent chemistries if they are still present. The potentials of oxidants like KMnO_4 or ceric ammonium nitrate (CAN) in aqueous HClO_4 that are more powerful than 2 are only available in aqueous conditions, and such comparisons with aqueous oxidants are misleading. For instance, the $E_{1/2}$ for tetrabutylammonium ceric nitrate (CTAN)²³ is 0.6 V (Figure 1) in MeCN and DCM and requires addition of strong acids like HOTf to achieve higher oxidizing strength in organic solvent.²⁴

C–H Nitroxylation and Stability Limits of 2. Single-electron oxidation is not the only reaction that 2 mediates. We found that 2 and 4 react with certain benzylic C–H bonds to form nitrate esters (nitroxylation) (Scheme 4, Table 2).²⁵


Hexamethylbenzene (HMB) was chosen as model substrate to investigate this reactivity since it has a weak benzylic C–H bond dissociation free energy (BDFE) in MeCN of 81.0 kcal/mol²⁶ and low ionization potential (IP) of 7.85 eV.²⁷ Reaction of HMB with two equiv of 2 in DCM afforded quantitative conversion to 2,3,4,5,6-pentamethylbenzyl nitrate in 5 min (entry 1). The same reaction but with 4 gave only 58% conversion in 24 h (entry 2). Reaction of 1 equiv of 2 with HMB resulted in 50% conversion (entry 3) consistent with the proposed mechanism (*vide infra*). Due to the efficiency and short reaction time of 2 mediated nitroxylation of HMB, it can serve as an effective batch control for the purity of 2 (entry 4; see Supporting Information for details).

Table 1. Reduction Potentials for Mn(III) Complexes

Complex	$E_{1/2}$ (V vs Fc/Fc^+)	
	MeCN	DCM
$[\text{Mn}^{\text{III}}(\text{NO}_3)_3(\text{OPPh}_3)_2]$ (2) ^a	1.02	0.96
$[\text{Mn}^{\text{III}}(\text{NO}_3)_3(\text{OAsPh}_3)_2]$ (4)	0.95	0.91
$[\text{Mn}^{\text{III}}(\text{NO}_3)(\text{MeCN})(\text{Opy})_4][\text{Mn}^{\text{II}}(\text{NO}_3)_4]$ (5)	1.02	
$[\text{Mn}^{\text{III}}(\text{NO}_3)(\text{Cl})_2(\text{OPPh}_3)_2]$ (6)	0.72	0.68
$[\text{Mn}^{\text{III}}(\text{Cl})_3(\text{OAsPh}_3)_2]$ (7)	0.72	0.78
$[\text{Mn}^{\text{III}}(\text{NO}_3)(\text{Cl})_2(\text{OAsPh}_3)_2]$ (8)	0.68	0.68

^aPotentials of 2 in MeCN measured in triplicate were within 1 mV. Potentials in DCM measured in triplicate varied by ± 10 mV. Electrolyte: 0.5 M $[\text{nBu}_4\text{N}][\text{PF}_6]$. Analyte conc.: 6 mM.

Scheme 4. C–H Nitroxylation Mediated by 2

Table 2. Nitroxylation of HMB with $\text{Mn}(\text{NO}_3)_3$ Complexes^a

Entry	Complex	% yield
1	2	99
2	4 ^b	58
3	2 (1 equiv)	50
4	2 (crude)	82

^aConditions shown in Scheme 4. ^b24 h. Average yields for duplicated runs determined by ^1H NMR.

The reactivity of 2 with benzylic C–H bonds is limited, boding well for use of 2 as a selective single-electron oxidant. For example, 2 did not react with a range of benzylic substrates (e.g., aromatic aldehydes, ketones, or esters; Scheme 4), even under thermal (refluxing DCM) or photolytic (100 W Xe-arc




Figure 4. (Left) Cyclic voltammograms of 0.006 M 2 in MeCN with 0.500 M $[\text{nBu}_4\text{N}][\text{PF}_6]$ (see Supporting Information for details). (Middle & right) Titration of 2 with ferrocene in DCM showing one-to-one stoichiometry.

lamp) conditions. Other substrates affording clean nitroxylation were *tert*-butyldiphenylsilane (TBDPS) protected *p*-cresol and mesitylene (Scheme 4). One possible mechanism of the reaction with HMB is a C–H cleavage akin to what has been proposed for Ni(IV) nitrate-mediated C–H cleavage by Jackson and Wang,²⁸ chlorination of HMB using 1 by us,¹² and CF₃ functionalization of arenes using Ag²⁺ photooxidants by Nocera.²⁹ The first equivalent of 2 cleaves a C–H bond forming nitric acid and a benzylic radical, the latter of which couples with a second molecule of 2 explaining the need for two equivalents of 2 and observation of 9 as the byproduct in HMB nitroxylation. Given that 2 has an $E_{1/2, \text{MeCN}}$ of 1.02 V and HNO₃ has a $pK_{a, \text{MeCN}}$ of 9.4,³⁰ the H atom abstracting ability (i.e., effective BDFE) is estimated to be 89 kcal/mol.²⁶ This bond energy is high enough for C–H nitroxylation of HMB, but it is not so high as to interfere with selective single-electron oxidation under a wide range of conditions (most C–H strengths are above 89 kcal/mol). An alternative mechanism to C–H activation could initiate with single-electron oxidation of HMB to form the HMB^{•+} given its low IP.²⁷ 2 also reacts, albeit slowly, with mesitylene (IP = 8.39 eV) but is stable in dry fluorobenzene (IP = 9.19 eV), DCM (IP = 11.35 eV), and MeCN (IP = 12.20 eV),^{31,32} which puts an upper limit of single-electron oxidation of <9.2 eV.³³ This high IP and low BDFE_{eff} reflects the high stability of 2 toward air and common organic solvents while maintaining its potent oxidizing ability. In addition to this and the other properties of 2 that we have described herein, it is plain that 2 and other Mn(III) nitrate complexes are ideal powerful oxidants for organometallic chemists.

ASSOCIATED CONTENT

* Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.4c03411>.

Additional figures, experimental details, and NMR, FTIR, UV–vis spectra, and reaction data (PDF)

Accession Codes

CCDC 2338924–2338930 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

David C. Lacy – Department of Chemistry, University at Buffalo, State University of New York, Buffalo, New York 14260, United States; orcid.org/0000-0001-5546-5081; Email: DCLacy@Buffalo.edu

Authors

Ananya Saju – Department of Chemistry, University at Buffalo, State University of New York, Buffalo, New York 14260, United States

Matthew R. Crawley – Department of Chemistry, University at Buffalo, State University of New York, Buffalo, New York 14260, United States; orcid.org/0000-0002-2555-9543

Samantha N. MacMillan – Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853, United States; orcid.org/0000-0001-6516-1823

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/jacs.4c03411>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

NSF grant CHE-1847933, NIH-R21-GM141685-01, and the University at Buffalo (UB) provided support. X-ray diffraction (XRD) System Rigaku XtaLAB Synergy-S was purchased with NSF award CHE-2216151. A Q-exactive GC-Orbitrap tandem MS was purchased using NSF CHE-1919594 by the UB Chemistry Instrument Center (CIC). The Bruker Ascend-500 NMR spectrometer in the UB Magnetic Resonance Center was purchased with NSF CHE-2018160.

REFERENCES

- (1) (a) Himmel, D.; Radtke, V.; Butschke, B.; Krossing, I. Basic Remarks on Acidity. *Angew. Chem., Int. Ed.* 2018, 57, 4386. (b) Radtke, V.; Himmel, D.; Putz, K.; Goll, S. K.; Krossing, I. The Protoelectric Potential Map (PPM): An absolute two-dimensional chemical potential scale for a global understanding of chemistry. *Chem.-Eur. J.* 2014, 20, 4194.
- (2) (a) Malischewski, M.; Adelhardt, M.; Sutter, J.; Meyer, K.; Seppelt, K. Isolation and structural and electronic characterization of salts of the decamethylferrocene dication. *Science* 2016, 353, 678. (b) Boere, R. T.; Kacprzak, S.; Keßler, M.; Knapp, C.; Riebau, R.; Riedel, S.; Roemmele, T. L.; Ruhle, M.; Scherer, H.; Weber, S. Oxidation of closo-[B₁₂Cl₁₂]²⁻ to the radical anion [B₁₂Cl₁₂]^{•-} and to neutral B₁₂Cl₁₂. *Angew. Chem., Int. Ed.* 2011, 50, 549. (c) Seppelt, K. Molecular hexafluorides. *Chem. Rev.* 2015, 115, 1296–1306.
- (3) (a) Malinowski, P. J.; Krossing, I. Ag[Fe(CO)₅]²⁺: A Bare Silver Complex with Fe(CO)₅ as a Ligand. *Angew. Chem., Int. Ed.* 2014, 53, 13460. (b) Schmitt, M.; Mayländer, M.; Goost, J.; Richert, S.; Krossing, I. Chasing the Mond cation: synthesis and characterization of the homoleptic nickel tetracarbonyl cation and its tricarbonyl-nitrosyl analogue. *Angew. Chem., Int. Ed.* 2021, 60, 14800.
- (4) T13 is the aminium-cation carborane-anion pair derived from 2,4-dibromo-*N*-(2-bromo-4-(trifluoromethyl)phenyl)-*N*-(2,4-dibromophenyl)aniline. See ref 5c.
- (5) (a) Albrecht, P. A.; Rupf, S. M.; Sellin, M.; Schlögl, J.; Riedel, S.; Malischewski, M. Increasing the oxidation power of TCNQ by coordination of B(C₆F₅)₃. *Chem. Commun.* 2022, 58, 4958. (b) Schorpp, M.; Heizmann, T.; Schmucker, M.; Rein, S.; Weber, S.; Krossing, I. Synthesis and application of a perfluorinated ammoniumyl radical cation as a very strong deelectronator. *Angew. Chem., Int. Ed.* 2020, 59, 9453. (c) Davidson, J. J.; Gunther, S. O.; Leong, D. W.; Ozerov, O. V. Synthesis of fluorinated aminium cations coupled with carborane anions for use as strong one-electron oxidants. *Dalton Trans.* 2023, 52, 16027. (d) Sellin, M.; Friedmann, C.; Mayländer, M.; Richert, S.; Krossing, I. Towards clustered carbonyl cations [M₃(CO)₁₄]²⁺ (M = Ru, Os): the need for innocent deelectronation. *Chem. Sci.* 2022, 13, 9147–9158.
- (6) (a) Lingappa, U. F.; Monteverde, D. R.; Magyar, J. S.; Valentine, J. S.; Fischer, W. W. How manganese empowered life with dioxygen (and vice versa). *Free Radic. Biol. Med.* 2019, 140, 113. (b) Zhu, W.; Richards, N. G. J. Biological functions controlled by manganese redox changes in mononuclear Mn-dependent enzymes. *Essays Biochem.* 2017, 61, 259. (c) Mandal, M.; Kawashima, K.; Saito, K.; Ishikita, H. Redox potential of the oxygen-evolving complex in the electron transfer cascade of photosystem II. *J. Phys. Chem. Lett.* 2020, 11, 249.
- (7) Liu, C.; Neale, Z. G.; Cao, G. Understanding electrochemical potentials of cathode materials in rechargeable batteries. *Mater. Today* 2016, 19, 109.
- (8) (a) Fu, N.; Sauer, G. S.; Saha, A.; Loo, A.; Lin, S. Metal-catalyzed electrochemical diazidation of alkenes. *Science* 2017, 357, 575. (b) Sauer, G. S.; Lin, S. An Electrocatalytic approach to the radical

- difunctionalization of alkenes. *ACS Catal.* 2018, 8, 5175. (c) Dong, X.; Roeckl, J. L.; Waldvogel, S. R.; Morandi, B. Merging shuttle reactions and paired electrolysis for reversible vicinal dihalogenations. *Science* 2021, 371, 507.
- (9) Philip, R. M.; Radhika, S.; Abdulla, C. M. A.; Anilkumar, G. Recent trends and prospects in homogeneous manganese-catalyzed epoxidation. *Adv. Synth. Catal.* 2021, 363, 1272.
- (10) Demir, A.S.; Emrullahoglu, M. Manganese(III) acetate: a versatile reagent in organic chemistry. *Current Organic Synthesis* 2007, 4, 321.
- (11) Saju, A.; Griffiths, J. R.; MacMillan, S. N.; Lacy, D. C. Synthesis of a bench-stable manganese(III) chloride compound: coordination chemistry and alkene dichlorination. *J. Am. Chem. Soc.* 2022, 144, 16761.
- (12) Saju, A.; Gunasekera, P. S.; Morgante, P.; MacMillan, S. N.; Autschbach, J.; Lacy, D. C. Experimental and computational determination of a M–Cl homolytic bond dissociation free energy: Mn(III)Cl-mediated C–H cleavage and chlorination. *J. Am. Chem. Soc.* 2023, 145, 13384.
- (13) Nachtigall, O.; Pataki, A.; Molski, M.; Lentz, D.; Spandl, J. Solvates of manganese trichloride revisited - synthesis, isolation, and crystal structure of $\text{MnCl}_3(\text{THF})_3$. *Z. Anorg. Allg. Chem.* 2015, 641, 1164.
- (14) Connelly, N. G.; Geiger, W. E. Chemical redox agents for organometallic chemistry. *Chem. Rev.* 1996, 96, 877.
- (15) Pearson, R. G.; Sobel, H. R.; Songstad, J. Nucleophilic reactivity constants toward methyl iodide and trans-dichlorodi(pyridine)-platinum(II). *J. Am. Chem. Soc.* 1968, 90, 319.
- (16) Henceforth, when “nitrate” can be explicitly linked to a coordinated ligand it will be written as “nitrato” to differentiate it from nitrate salts, counterion, organo-nitrate, etc.
- (17) Johnson, D. W.; Sutton, D. The Preparation, properties, and some complexes of anhydrous manganese(III) nitrate. *Can. J. Chem.* 1972, 50, 3326.
- (18) Vankar, P. S.; Reddy, M. V. R.; Vankar, Y. D. Applications of trimethylsilyl halides-oxidants combinations in organic synthesis. a review. *Org. Prep. Proced. Int.* 1998, 30, 373.
- (19) Eppley, H. J.; Christou, G. Synthesis of dodecaoxohexadecarboxylatotetraquo-dodecamanganese $[\text{Mn}_{12}\text{O}_{12}(\text{O}_2\text{CR})_{16}(\text{H}_2\text{O})_4]$ (R = Me, Et, Ph, Cr) complexes. *Inorg. Syn.* 2002, 33, 61.
- (20) Drummond, J.; Wood, J. S. Crystal and molecular structure of tetraphenylarsonium tetranitratomanganate(II) and X-ray study of other $\text{M}(\text{NO}_3)_4^{2-}$ ions (M = Ni, Cu, or Zn). *J. Chem. Soc. A* 1970, 226, 226.
- (21) (a) A new synthesis for 7 was developed by us (see [Supporting Information](#)). (b) Uson, R.; Riera, V.; Ciriano, M. A.; Valderrama, M. Pentacoordinate neutral manganese(III) complexes. *Transition Met. Chem.* 1976, 1, 122.
- (22) Wong, C. L.; Kochi, J. K. Electron transfer with organometals. Steric effects as probes for outer-sphere and inner-sphere oxidations of homoleptic alkylmetals with iron(III) and iridate(IV) complexes. *J. Am. Chem. Soc.* 1979, 101, 5593.
- (23) Zhang, Y.; Raines, A. J.; Flowers, R. A., II Solvent-dependent chemoselectivities in Ce(IV)-mediated oxidative coupling reactions. *Org. Lett.* 2003, 5, 2363.
- (24) Jeong, D.; Lee, Y.; Lee, Y.; Kim, K.; Cho, J. Synthesis, characterization, and reactivity of a highly oxidative mononuclear manganese(IV)-bis(fluoro) Complex. *J. Am. Chem. Soc.* 2024, 146, 4172.
- (25) Examples of CH nitroxylation: (a) Li, B.; Han, Y.-Q.; Yang, X.; Shi, B.-F. Palladium-catalyzed $\text{C}(\text{sp}^3)\text{-H}$ nitroxylation with *tert*-butyl nitrate and molecular oxygen. *Org. Lett.* 2020, 22, 9719–9723. (b) Calvo, R.; Le Tellier, A.; Nauser, T.; Rombach, D.; Nater, D.; Katayev, D. Synthesis, characterization, and reactivity of a hyper-valent-iodine-based nitrooxylating reagent. *Angew. Chem., Int. Ed.* 2020, 59, 17162–17168.
- (26) Agarwal, R. G.; Coste, S. C.; Groff, B. D.; Heuer, A. M.; Noh, H.; Parada, G. A.; Wise, C. F.; Nichols, E. M.; Warren, J. J.; Mayer, J. M. Free energies of proton-coupled electron transfer reagents and their applications. *Chem. Rev.* 2022, 122, 1–49.
- (27) Schorpp, M.; Rein, S.; Weber, S.; Scherer, H.; Krossing, I. Guilty and charged: a stable solution of the hexamethylbenzene radical cation as a ligand forming oxidising agent. *Chem. Commun.* 2018, 54, 10036.
- (28) Kwon, Y. M.; Lee, Y.; Schmutz, A. K.; Jackson, T. A.; Wang, D. C–H bond activation by a mononuclear nickel(IV)- nitrate complex. *J. Am. Chem. Soc.* 2022, 144, 12072.
- (29) Campbell, B. M.; Gordon, J. B.; Raguram, E. R.; Gonzalez, M. I.; Reynolds, K. G.; Nava, M.; Nocera, D. G. Electrophotocatalytic perfluoroalkylation by LMCT excitation of Ag(II) perfluoroalkyl carboxylates. *Science* 2024, 383, 279.
- (30) Kutt, A.; Rodima, T.; Saame, J.; Raamat, E.; Mäemets, V.; Kaljurand, I.; Koppel, I. A.; Garlyauskayte, R. Y.; Yagupolskii, Y. L.; Yagupolskii, L. M.; Bernhardt, E.; Willner, H.; Leito, I. Equilibrium acidities of superacids. *J. Org. Chem.* 2011, 76, 391.
- (31) Watanabe, K. Ionization potentials of some molecules. *J. Chem. Phys.* 1957, 26, 542.
- (32) de Petris, G.; Fornarini, S.; Crestoni, M. E.; Troiani, A.; Mayer, P. M. What ion is generated when ionizing acetonitrile? *J. Phys. Chem. A* 2005, 109, 4425.
- (33) Malinowski, P. J.; Himmel, D.; Krossing, I. Coordination chemistry of diiodine and implications for the oxidation capacity of the synergistic Ag^+/X_2 (X = Cl, Br, I) system. *Angew. Chem., Int. Ed.* 2016, 55, 9262.