

Carboxylic Acids as Double Aryl Group Donors for Biaryl Synthesis

Wenzhi Zhang,^a Jie Ma,^a Fengyan Zhou,^a Michal Szostak^{b*} and Chengwei Liu^{c*}

Received 00th January 20xx,

Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

The synthesis of biaryl compounds by employing carboxylic acids as double aryl group donors is reported. Naturally present and benign carboxylic acids have been applied as double aryl group donors by sequential decarbonylations for the construction of unsymmetrical biaryls. This method represents an orthogonal approach for the synthesis of valuable biaryl compounds. Aryl carboxylic acids were first converted to aryl boronic esters via palladium-catalyzed decarbonylative borylation. Next, aryl boronic esters were transformed to aryl boronic acids via hydrolysis. Biaryl compounds were generated through decarbonylative coupling between aryl carboxylic acids and aryl boronic acids. Broad substrate scope and excellent functional group tolerance have been demonstrated. Furthermore, a range of pharmaceutical motifs can be readily engaged in excellent yields by this approach. The present method successfully achieves the synthesis of highly valuable biaryls using readily available and inexpensive carboxylic acids as a single class of precursors to access cross-coupling synthons of central importance to the synthetic community.

Introduction

The biaryl motif is a structural architecture of central importance in chemical synthesis.^{1–3} The importance of biaryl compounds is underscored by the fact that biaryls have found key applications in medicine, agriculture and materials science, affecting the quality of life and societal well-being.^{1–3} The synthesis of biaryls have represented an important endeavor in the development of chemical methodologies.^{1–5} Among methods available, at present, Suzuki–Miyaura cross-coupling represents the most powerful and reliable approach to the synthesis of biaryl compounds.

The two essential raw precursors for the Suzuki–Miyaura cross-coupling are aryl halides and aryl boronic acids. Aryl boronic acids for industrial synthesis are routinely prepared by the borylation of aryl halides to generate aryl boronic esters, which are then hydrolyzed to furnish aryl boronic acids.¹ Thus, aryl halides are two-fold essential raw precursors deployed for the Suzuki–Miyaura cross-couplings. However, aryl halides are not naturally present in nature, and are typically synthesized by electrophilic halogenation of aromatic hydrocarbons through the Friedel–Crafts reaction. Thus, the traditional sequence of preparing biaryls through the Suzuki–Miyaura cross-coupling involves four steps: 1) Friedel–Crafts halogenation of aromatic hydrocarbons to prepare aryl halides, 2) the borylation of aryl halides to furnish aryl boronic esters, 3) the hydrolysis of aryl boronic esters to afford aryl

boronic acids, 4) the reaction of aryl boronic acids with aryl halides to synthesize biaryls. Furthermore, there are safety concerns involving the use of halogens, which pose certain safety hazards. The prepared aryl halides also entail toxicity risks. The development of methods for the synthesis of biaryls using benign and naturally present substrates that can replace aryl halides and can be easily employed under the powerful Suzuki–Miyaura cross-coupling regimen is highly desirable from the societal, industrial and academic standpoints.

Carboxylic acids are readily available and inexpensive raw precursors, which are naturally present in nature.^{6–7} Following the wide use of aryl halides in cross-coupling reactions,^{1–5} chemists turned their attention to carboxylic acids as cross-coupling substrates, which led to the development of decarboxylative cross-coupling reactions of carboxylic acids.⁶ Compared to aryl halides, the advantages of carboxylic acids are plentiful, including their stability, non-toxicity and orthogonality.⁶ However, despite advantages of carboxylic acids over aryl halides, there are several major drawbacks of the decarboxylative cross-coupling reaction mode of carboxylic acids. Most notably, decarboxylative cross-couplings of carboxylic acids typically feature narrow substrate scope and poor functional group tolerance. In general, only ortho-functionalized carboxylic acid substrates or substrates containing strongly electron-withdrawing groups are compatible.⁶ In addition, decarboxylative cross-coupling reactions of carboxylic acids generate aryl nucleophiles and require oxidants in the cross-coupling with aryl boronic acids.⁸ At present, there are very few examples where carboxylic acids are used as double aryl group donors to achieve the synthesis of biaryl compounds.^{9,10} These methods are limited by the synthesis of symmetrical biaryls, specific substitution patterns, limited substrate scope and functional group tolerance, and are not viable for the synthesis of broadly useful biaryl compounds.

^a College of Chemistry, Chemical Engineering and Materials Science, Zaozhuang University, 1 Bei'an Road, Zaozhuang, Shandong 277160, China.

^b Department of Chemistry, Rutgers University, 73 Warren Street, Newark, NJ 07102, United States. *E-mail: michal.szostak@rutgers.edu.

^c Department of Chemistry, Shanghai University, 99 Shangda Road, Shanghai 200444, China.

* chengwei_liu@shu.edu.cn; https://www.x-mol.com/groups/chengwei_liu
Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x




Figure 1. The synthesis of biaryls using carboxylic acids.

In contrast, decarbonylative cross-coupling of carboxylic acids represents an exceedingly robust reaction mode, where the powerful Pd(0)/Pd(II) cycle shows the inherent broad substrate scope and excellent functional group tolerance. Most crucially, carboxylic acids applicable to decarbonylative cross-coupling are not limited to substrates with ortho-substituents or substrates with strongly electron-withdrawing groups, which results in broad compatibility of diverse aryl carboxylic acids to this reaction mode.¹¹

In 1998, de Vries and co-workers reported decarbonylative cross-coupling of carboxylic acid anhydrides with olefins establishing the first decarbonylative Heck reaction.^{12a} Afterwards, this reaction was extended to in situ activation of carboxylic acids.^{12b} These methods were based on precedents in decarbonylative β -hydride elimination of alkyl carboxylic acids.¹³⁻¹⁴ In 2013, the Shi group reported rhodium-catalyzed arylation of aromatic hydrocarbons via C-H bond activation using carboxylic acids as aryl donors.¹⁵ It was not until 2018 that decarbonylative borylation of carboxylic acids was reported, marking the first time carboxylic acids were used as coupling reagents for the formation of carbon-heteroatom bonds via decarbonylative pathway.¹⁶ In 2019, the first decarbonylative arylation of carboxylic acids was reported, which represented a general and practical method for the synthesis of biaryls using carboxylic acids as arylating reagents.¹⁷ Thereafter, a series of methods to generate carbon-carbon and carbon-heteroatom bonds through decarbonylative cross-coupling of carboxylic acids were developed.¹⁸⁻²⁴

Based on our interest in decarbonylative cross-coupling, we envisioned the use of carboxylic acids as double aryl donors to synthesize biaryl compounds. The specific steps of this approach involve 1) decarbonylative borylation of carboxylic acids to generate aryl borate esters, 2) hydrolysis of aryl borate esters to furnish aryl boronic acids, and 3) decarbonylative Suzuki-Miyaura cross-coupling of the generated aryl boronic acids with carboxylic acids to generate biaryls. Noteworthy features of this approach involve 1) the use of benign carboxylic acids that are naturally present and

orthogonal to aryl halides, 2) the first use of carboxylic acids as double aryl functional group donors for the synthesis of biaryl compounds, and 3) broad functional group tolerance, including functionalization of pharmaceutical motifs that can be readily synthesized via the present approach.

Results and Discussion

The decarbonylative borylation of carboxylic acids was first investigated using benzoic acid (**1a**) and bis(pinacolato)diborane (**2a**) as model substrates (Table 1). The use of base was first screened, and we found that 4-(dimethylamino)pyridine (DMAP) is the optimal base for this reaction (entries 1-3). Furthermore, various phosphine ligands were evaluated (entries 3-14), and we found that DPPP, XantPhos and DPPB gave the highest conversion under the tested conditions. The effect of temperature was also investigated (entries 15-16), and we determined that the reaction can still maintain 71% yield at 120 °C under these conditions.

Table 1. Optimization of borylation of carboxylic acids.^a


$\text{Ph-CO}_2\text{H}$ + B_2pin_2 $\xrightarrow[\text{base, Piv}_2\text{O, dioxane, 160 }^\circ\text{C, 15 h}]{[\text{Pd}] (3 \text{ mol\%}), \text{ligand} (6 \text{ mol\%})}$ Ph-Bpin				
entry	catalyst	ligand	base	yield
1	Pd(OAc) ₂	DPPB		77
2	Pd(OAc) ₂	DPPB	Et ₃ N	88
3	Pd(OAc) ₂	DPPB	DMAP	96
4	Pd(OAc) ₂	DPPP	DMAP	98
5	Pd(OAc) ₂	DPPPent	DMAP	90
6	Pd(OAc) ₂	DPPF	DMAP	90
7	Pd(OAc) ₂	BINAP	DMAP	41
8	Pd(OAc) ₂	XantPhos	DMAP	97
9	Pd(OAc) ₂	DavePhos	DMAP	13
10	Pd(OAc) ₂	XPhos	DMAP	<2
11	Pd(OAc) ₂	SPhos	DMAP	7
12	Pd(OAc) ₂	PCy ₃ HBF ₄	DMAP	29
13	Pd(OAc) ₂	PCyPh ₂	DMAP	37
14	Pd(OAc) ₂	PPh ₃	DMAP	53
15 ^b	Pd(OAc) ₂	DPPP	DMAP	88
16 ^c	Pd(OAc) ₂	DPPP	DMAP	71

^aConditions: **1a** (1.0 equiv), **2a** (1.5 equiv), Pd(OAc)₂ (3 mol%), ligand (6 mol%), base (1.5 equiv), Piv₂O (1.5 equiv), dioxane, 160 °C, 15 h; ^b140 °C; ^c120 °C.


With the optimized reaction conditions in hand, the scope of decarbonylative borylation of carboxylic acids was investigated (Scheme 1). As shown in Scheme 1, carboxylic acids bearing diverse electron-neutral (**3a-3c**), electron-donating (**3d**), and electron-deficient (**3e**) substituents are well tolerated in this approach. Furthermore, substrates containing sensitive cyano (**3f**), chloro (**3g**), ester (**3h**), ketone (**3i**) groups are compatible with this method. Naphthyl substrates (**3j-3k**) could also be converted to the desired borylation products in excellent yields. Furthermore, sterically-hindered (**3l**) and heterocyclic (**3m**) substrates could also be readily employed in this process. Notably, bioactive carboxylic acids, such as found in

pharmaceuticals (**3n–3o**), were also compatible with this reaction.

Scheme 1. Decarbonylative borylation of carboxylic acids.



Scheme 2. Hydrolysis of aryl boronic esters.



Having achieved the conversion of a series of carboxylic acids to aryl boronic esters, we next focused on the next step to hydrolyze aryl boronic esters to aryl boronic acids (Scheme 2). We established that these obtained aryl boronic esters could be hydrolyzed under mild conditions using NaIO₄/HCl.²⁵ As shown in Scheme 2, aryl boronic esters bearing electron-neutral (**4a–4c**), electron-rich (**4d**), and electron-withdrawing (**4e**) substituents were converted to aryl boronic acids in excellent yields. Substrates containing cyano (**4f**), chloro (**4g**), ester (**4h**), ketone (**4i**) groups were well-tolerated. Naphthyl

substrates (**4j–4k**), sterically-hindered (**4l**), and heterocyclic (**4m**) precursors were also well-suitable. Importantly, substrates derived from pharmaceuticals (**4n–4o**) were also compatible with this mild hydrolysis approach.

Table 2. Optimization of arylation of carboxylic acids.^a

entry	catalyst	ligand	base	yield
1 ^b	Pd(OAc) ₂	DPPB		14
2 ^b	Pd(OAc) ₂	DPPB	Et ₃ N	22
3	Pd(OAc) ₂	DPPB	Et ₃ N	92
4	Pd(OAc) ₂	DPPB	DMAP	36
5	Pd(OAc) ₂	DPPB	pyridine	74
6	Pd(OAc) ₂	DPPB	DIPEA	96
7 ^c	Pd(OAc) ₂	DPPB	DIPEA	75
8 ^d	Pd(OAc) ₂	DPPB	DIPEA	56
9	Pd(OAc) ₂	DPPP	DIPEA	47
10	Pd(OAc) ₂	DPPent	DIPEA	69
11	Pd(OAc) ₂	DPPF	DIPEA	35
12	Pd(OAc) ₂	BINAP	DIPEA	27
13	Pd(OAc) ₂	XantPhos	DIPEA	80
14	Pd(OAc) ₂	DavePhos	DIPEA	5
15	Pd(OAc) ₂	XPhos	DIPEA	7
16	Pd(OAc) ₂	SPhos	DIPEA	6
17	Pd(OAc) ₂	PCy ₃ HBF ₄	DIPEA	72
18	Pd(OAc) ₂	PCyPh ₂	DIPEA	67
19	Pd(OAc) ₂	PPh ₃	DIPEA	27


^aConditions: **1a** (1.0 equiv), **3b** (2.0 equiv), Pd(OAc)₂ (3 mol%), ligand (6 mol%), base (2.0 equiv), H₃BO₃ (2.0 equiv), Piv₂O (2.0 equiv), dioxane, 160 °C, 15 h; ^bwithout H₃BO₃; ^c**3b** (1.5 equiv), base (1.5 equiv), H₃BO₃ (1.5 equiv), Piv₂O (1.5 equiv); ^d**3b** (1.2 equiv), base (1.2 equiv), H₃BO₃ (1.2 equiv), Piv₂O (1.2 equiv).

Having secured access to aryl boronic acids, we then screened the conditions for the critical decarbonylative arylation of carboxylic acids using 3-py-CO₂H carboxylic acid as the model substrate (Table 2). We found that boric acid is an essential additive for this reaction (entries 1–3). *N,N*-Diisopropylethylamine (DIPEA) was identified as the optimal base for this decarbonylative arylation (entries 3–6). We established that stoichiometry is a crucial parameter for the arylation to ensure the optimal equivalency ratio (entries 6–8). Furthermore, different phosphine ligands were evaluated (entries 6, 9–19), and DPPB was identified as the optimal ligand. The first conditions (Scheme 1) are optimized for borylation, while the second are optimized for the Suzuki coupling (Table 2). The key difference is the use of triethylamine as a base to form more reactive acyl ammonium and boric acid.

Having identified the optimal conditions, the scope of the biaryl synthesis via decarbonylative arylation of carboxylic acids was next investigated (Scheme 3). As shown, aryl boronic acid substrates bearing electron-neutral (**5a–c**), electron-donating (**5d**), and electron-withdrawing (**5e**) substituents

were well compatible with this biaryl synthesis method, furnishing 3-pyridyl biaryls. Furthermore, aryl boronic acid substrates containing cyano (**5f**), chloro (**5g**), ester (**5h**), ketone (**5i**) functional groups were well-tolerated in this decarbonylative arylation. Moreover, naphthyl (**5j–5k**), sterically-hindered (**5l**), and heterocyclic (**5m**) aryl boronic acids were also well-compatible, affording the desired biaryl products in good to excellent yields. Interestingly, quinoline-6-carboxylic acid could also be applied to this reaction (**5n**), furnishing medicinally-relevant products. Next, we extended the scope of carboxylic acid substrates to benzoic acids. As shown in Scheme 3, carboxylic acid substrates containing sensitive ester (**5o**), ketone (**5p**), aldehyde (**5q**) functional groups were well-tolerated in this reaction, providing electrophilic handles for further functionalization. Moreover, carboxylic acids containing medicinally-relevant trifluoromethyl (**5r**), cyano (**5s**) and chloro (**5t**) substituents could be employed in good yields. Furthermore, naphthyl (**5u–5v**) substrates were readily amenable by this approach, furnishing conjugated biaryls. Finally, the sterically-hindered substrate (**5w**) was also well-tolerated in this method.

Scheme 3. Substrate scope of arylation of carboxylic acids.



Considering the benefits of carboxylic acids as double aryl donors, we next evaluated the rapid application of this method for the direct functionalization of bioactive molecules (Scheme 4A). As shown, this approach was successfully utilized for the arylation of pharmaceuticals and natural products, such as probenecid (**5x**, **5aa**), adapalene (**5y**), febuxostat (**5z**) and estrone (**5ab**). Crucially, this approach utilizes the inherent presence of the carboxylic acid moiety as a synthetic handle for decarbonylation. Furthermore, gram scale reaction has been implemented and a high conversion has been obtained (Scheme 4B).

Conclusions


In summary, we have reported the synthesis of biaryl compounds using carboxylic acids as double aryl group donors. This approach exploits sequential decarbonylations for the synthesis of unsymmetrical biaryls. The advantages of the method include the natural presence of carboxylic acids, their benign properties as well as orthogonal and ready availability, which provides an alternative approach to the traditional synthesis of biaryls using aryl halides. The approach features excellent functional group tolerance and broad substrate scope, demonstrating that this method represents an efficient pathway for the construction of biaryls using carboxylic acids as raw substrates. The utility has been further demonstrated in the direct functionalization of pharmaceutical molecules. This method provides new avenues for the synthesis of biaryl compounds using a single class of readily available and inexpensive precursors to access cross-coupling synthons of central importance to the synthetic community. Ongoing studies in our laboratories are focused on the development of one-pot decarbonylative processes of carboxylic acids and derivatives, and these results will be reported in due course.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank Shanghai University (C. L.), Overseas High-Level Talents Introduction Program of Shanghai (C. L.), Rutgers University (M. S.), the NSF (CAREER CHE-1650766, M. S.), Zaozhuang University (W. Z.) for generous support. Additional support was provided by the Rutgers Graduate School in the form of Dean's Dissertation Fellowship (C. L.).

Scheme 4. Versatile applications and gram scale reaction.

References

- (a) N. Miyaura, A. Suzuki, Palladium-Catalyzed Cross-Coupling Reactions of Organoboron Compounds, *Chem. Rev.* 1995, **95**, 2457-2483; (b) A. J. J. Lennox, G. C. Lloyd-Jones, Selection of Boron Reagents for Suzuki-Miyaura Coupling, *Chem. Soc. Rev.* 2014, **43**, 412-443.
- C. C. C. Johansson-Seechurn, M. O. Kitching, T. J. Colacot, V. Snieckus, Palladium-Catalyzed Cross-Coupling: A Historical Contextual Perspective to the 2010 Nobel Prize, *Angew. Chem. Int. Ed.* 2012, **51**, 5062-5085.
- (a) F. S. Han, Transition-Metal-Catalyzed Suzuki-Miyaura Cross-Coupling Reactions: a Remarkable Advance from Palladium to Nickel Catalysts, *Chem. Soc. Rev.* 2013, **42**, 5270-5298; (b) S. Z. Tasker, E. A. Standley, T. F. Jamison, Recent Advances in Homogeneous Nickel Catalysis, *Nature* 2014, **509**, 299-309; (c) V. P. Ananikov, Nickel: The "Spirited Horse" of Transition Metal Catalysis, *ACS Catal.* 2015, **5**, 1964-1971.
- (a) A. F. Littke, G. C. Fu, Palladium-Catalyzed Coupling Reactions of Aryl Chlorides, *Angew. Chem. Int. Ed.* 2002, **41**, 4176-4211; (b) T. E. Barder, S. D. Walker, J. R. Martinelli, S. L. Buchwald, Catalysts for Suzuki-Miyaura Coupling Processes: Scope and Studies of the Effect of Ligand Structure, *J. Am. Chem. Soc.* 2005, **127**, 4685-4696.
- (a) S. B. Blakey, D. W. C. MacMillan, The First Suzuki Cross-Couplings of Aryltrimethylammonium Salts, *J. Am. Chem. Soc.* 2003, **125**, 6046-6047; (b) M. Tobisu, T. Shimasaki, N. Chatani, Nickel-Catalyzed Cross-Coupling of Aryl Methyl Ethers with Aryl Boronic Esters, *Angew. Chem. Int. Ed.* 2008, **47**, 4866-4869; (c) B. T. Guan, Y. Wang, B. J. Li, D. G. Yu, Z. J. Shi, Biaryl Construction via Ni-Catalyzed C-O Activation of Phenolic Carboxylates, *J. Am. Chem. Soc.* 2008, **130**, 14468-14470.
- For reviews on decarboxylative cross-coupling of carboxylic acids, see: (a) L. J. Goossen, N. Rodriguez, K. Goossen,

- Carboxylic Acids as Substrates in Homogeneous Catalysis, *Angew. Chem. Int. Ed.* 2008, **47**, 3100-3120; (b) N. Rodriguez, L. J. Goossen, Decarboxylative Coupling Reactions: A Modern Strategy for C–C Bond Formation, *Chem. Soc. Rev.* 2011, **40**, 5030-5048; (c) Y. Wei, P. Hu, M. Zhang, W. Su, Metal-Catalyzed Decarboxylative C–H Functionalization, *Chem. Rev.* 2017, **117**, 8864-8907.
- 7 For reviews on carboxylic acids as directing group for C–H bond activation, see: (a) M. P. Drapeau, L. J. Goossen, Carboxylic Acids as Directing Groups for C–H Bond Functionalization, *Chem. Eur. J.* 2016, **22**, 18654-18677; (b) M. Font, J. M. Quibell, G. J. P. Perry, I. Larrosa, The Use of Carboxylic Acids as Traceless Directing Groups for Regioselective C–H Bond Functionalisation, *Chem. Commun.* 2017, **53**, 5584-5597.
 - 8 For a review on the synthesis of biaryls using carboxylic acids, see: G. J. P. Perry, I. Larrosa, Recent Progress in Decarboxylative Oxidative Cross-Coupling for Biaryl Synthesis, *Eur. J. Org. Chem.* 2017, 3517-3527.
 - 9 S. Kumagai, R. Yamasaki, T. Kameda, Y. Saito, A. Watanabe, C. Watanabe, N. Teramae, T. Yoshioka, Aromatic Hydrocarbon Selectivity as a Function of CaO Basicity and Aging during CaO-Catalyzed PET Pyrolysis using Tandem μ -Reactor-GC/MS, *Chem. Eng. J.* 2018, **332**, 169-173.
 - 10 F. Pu, L. Y. Zhang, Z. W. Liu, X. Y. Shi, Palladium (II)-Catalyzed Decarboxylative Cross-Dehydrogenative Coupling: Direct Synthesis of *meta*-Substituted Biaryls from Aromatic Acids, *Adv. Synth. Catal.* 2018, **360**, 2644-2649.
 - 11 (a) W. I. Dzik, P. P. Lange, L. J. Goossen, Carboxylates as Sources of Carbon Nucleophiles and Electrophiles: Comparison of Decarboxylative and Decarbonylative Pathways, *Chem. Sci.* 2012, **3**, 2671-2678.
 - 12 For studies on decarbonylative cross-coupling of aryl anhydrides, see: (a) M. S. Stephan, A. J. J. M. Teunissen, G. K. M. Verzijl, J. G. de Vries, Heck Reactions without Salt Formation: Aromatic Carboxylic Anhydrides as Arylating Agents, *Angew. Chem. Int. Ed.* 1998, **37**, 662-664; for preliminary exploration of decarbonylative cross-coupling of carboxylic acids, see: (b) L. J. Goossen, J. Paetzold, L. Winkel, Pd-Catalyzed Decarbonylative Heck Olefination of Aromatic Carboxylic Acids Activated in situ with Di-*tert*-butyl Dicarboxylate, *Synlett.* 2002, **10**, 1721-1723.
 - 13 For a review on β -hydride elimination of alkyl carboxylic acids, see: X. Zhang, F. Jordan, M. Szostak, Transition-Metal-Catalyzed Decarbonylation of Carboxylic Acids to Olefins: Exploiting Acyl C–O Activation for the Production of High Value Products, *Org. Chem. Front.* 2018, **5**, 2515-2521.
 - 14 For representative studies on β -hydride elimination of alkyl carboxylic acids, see: (a) J. A. Miller, J. A. Nelson, M. P. Byrne, A Highly Catalytic and Selective Conversion of Carboxylic Acids to 1-Alkenes of One Less Carbon Atom, *J. Org. Chem.* 1993, **58**, 18-20; (b) L. J. Goossen, N. Rodriguez, A Mild and Efficient Protocol for the Conversion of Carboxylic Acids to Olefins by A Catalytic Decarbonylative Elimination Reaction, *Chem. Commun.* 2004, 724-725; (c) J. L. Notre, E. L. Scott, M. C. R. Franssen, J. P. M. Sanders, Selective Preparation of Terminal Alkenes from Aliphatic Carboxylic Acids by A Palladium-Catalysed Decarbonylation-Elimination Reaction, *Tetrahedron Lett.* 2010, **51**, 3712-3715; (d) Y. Liu, K. E. Kim, M. B. Herbert, A. Fedorov, R. H. Grubbs, B. M. Stoltz, Palladium-Catalyzed Decarbonylative Dehydration of Fatty Acids for the Production of Linear Alpha Olefins, *Adv. Synth. Catal.* 2014, **356**, 130-136; (e) Y. Liu, S. C. Virgil, R. H. Grubbs, B. M. Stoltz, Palladium-Catalyzed Decarbonylative Dehydration for the Synthesis of α -Vinyl Carbonyl Compounds and Total Synthesis of (-)-Aspewentins A, B, and C, *Angew. Chem. Int. Ed.* 2015, **54**, 11800-11803.
 - 15 For selected Rh-catalyzed C–H activation using carboxylic acids as arylating reagents, see: (a) F. Pan, Z. Q. Lei, H. Wang, H. Li, J. Sun, Z. J. Shi, Rhodium(I)-Catalyzed Redox-Economic Cross-Coupling of Carboxylic Acids with Arenes Directed by N-Containing Groups, *Angew. Chem. Int. Ed.* 2013, **52**, 2063-2067; (b) L. Zhang, X. Xue, C. Xu, Y. Pan, G. Zhang, L. Xu, H. Li, Z. J. Shi, Rhodium-Catalyzed Decarbonylative Direct C2-Arylation of Indoles with Aryl Carboxylic Acids, *ChemCatChem* 2014, **6**, 3069-3074; (c) S. Kwon, D. Kang, S. Hong, Rh-I-Catalyzed Site-Selective Decarbonylative Alkenylation and Arylation of Quinolones under Chelation Assistance, *Eur. J. Org. Chem.* 2015, 3671-3678.
 - 16 C. Liu, C. L. Ji, X. Hong, M. Szostak, Palladium-Catalyzed Decarbonylative Borylation of Carboxylic Acids: Tuning Reaction Selectivity by Computation, *Angew. Chem. Int. Ed.* 2018, **57**, 16721-16726.
 - 17 (a) C. Liu, C. L. Ji, Z. X. Qin, X. Hong, M. Szostak, Synthesis of Biaryls via Decarbonylative Palladium-Catalyzed Suzuki-Miyaura Cross-Coupling of Carboxylic Acids, *iScience* 2019, **19**, 749-759. (b) A. Cervantes-Reyes, A. C. Smith, G. M. Chinigo, D. C. Blakemore, M. Szostak, Decarbonylative Pd-Catalyzed Suzuki Cross-Coupling for the Synthesis of Structurally Diverse Heterobiaryls, *Org. Lett.* 2022, **24**, 1678-1683.
 - 18 C. Liu, Z. X. Qin, C. L. Ji, X. Hong, M. Szostak, Highly-Chemoselective Step-Down Reduction of Carboxylic Acids to Aromatic Hydrocarbons via Palladium Catalysis, *Chem. Sci.* 2019, **10**, 5736-5742.
 - 19 For decarbonylative phosphorylation of carboxylic acids, see: (a) C. Liu, Y. Y. Xing, T. Zhou, T. Chen, X. Hong, M. Szostak, Carboxylic-Phosphoric Anhydrides as Direct Electrophiles for Decarbonylative Hirao Cross-Coupling of Carboxylic Acids: DFT Investigation of Mechanistic Pathway, *Chem. Asian J.* 2023, **18**, e202201262; (b) J. S. Zhang, T. Chen, L. B. Han, Palladium-Catalyzed Direct Decarbonylative Phosphorylation of Benzoic Acids with P(O)–H Compounds, *Eur. J. Org. Chem.* 2020, 1148-1153.
 - 20 W. Yu, L. Liu, T. Huang, X. Zhou, T. Chen, Palladium-Catalyzed Decarbonylative Heck Coupling of Aromatic Carboxylic Acids with Terminal Alkenes, *Org. Lett.* 2020, **22**, 7123-7128.
 - 21 For decarbonylative heteroarylation of carboxylic acids, see: (a) C. Liu, C. L. Ji, T. Zhou, X. Hong, M. Szostak, Bimetallic Cooperative Catalysis for Decarbonylative Heteroarylation of Carboxylic Acids via C–O/C–H Coupling, *Angew. Chem. Int. Ed.* 2021, **60**, 10690-10699; (b) K. Xiang, S. Zhang, L. Liu, T. Huang, Z. Tang, C. Li, K. Xu, T. Chen, Tunable C–H Arylation and Acylation of Azoles with Carboxylic Acids by Pd/Cu Cooperative Catalysis, *Org. Chem. Front.* 2021, **8**, 2543-2550.
 - 22 For decarbonylative alkylation of carboxylic acids, see: (a) X. Li, L. Liu, T. Huang, Z. Tang, C. Li, W. Li, T. Zhang, Z. Li, T. Chen, Palladium-Catalyzed Decarbonylative Sonogashira Coupling of Terminal Alkynes with Carboxylic Acids, *Org. Lett.* 2021, **23**, 3304-3309; (b) C. Liu, M. Szostak, Decarbonylative Sonogashira Cross-Coupling of Carboxylic Acids, *Org. Lett.* 2021, **23**, 4726-4730.
 - 23 For decarbonylative cyanation of carboxylic acids, see: (a) T. Xu, W. Li, K. Zhang, Y. Han, L. Liu, T. Huang, C. Li, Z. Tang, T. Chen, Palladium-Catalyzed Decarbonylative Cyanation of Carboxylic Acids with TMS-CN, *J. Org. Chem.* 2022, **87**, 11871-11879; (b) G. Zhang, H. Miao, C. Guan, C. Ding, Palladium-Catalyzed Direct Decarbonylative Cyanation of Aryl Carboxylic Acids, *J. Org. Chem.* 2022, **87**, 12791-12798.
 - 24 For decarbonylative thioetherification of carboxylic acids, see: (a) T. Xu, X. Zhou, X. Xiao, Y. Yuan, L. Liu, T. Huang, C. Li, Z. Tang, T. Chen, Nickel-Catalyzed Decarbonylative Thioetherification of Carboxylic Acids with Thiols, *J. Org. Chem.* 2022, **87**, 8672-8684; (b) H. Ji, H. Cao, G. Wang, F. Xing, M. Szostak, C. Liu, Predominant Intermolecular

- Decarbonylative Thioetherification of Carboxylic Acids using Nickel Precatalysts, *Org. Chem. Front.* 2023, **10**, doi: 10.1039/d3qo00744h.
- 25 A. G. Crawford, Z. Liu, I. A. I. Mkhalid, M. H. Thibault, N. Schwarz, G. Alcaraz, A. Steffen, J. C. Collings, A. S. Batsanov, J. A. K. Howard, T. B. Marder, Synthesis of 2- and 2,7-Functionalized Pyrene Derivatives: An Application of Selective C–H Borylation, *Chem. Eur. J.* 2012, **18**, 5022-5035.