

TEMPEREDNESS OF $L^2(\Gamma \backslash G)$ AND POSITIVE EIGENFUNCTIONS IN HIGHER RANK.

SAM EDWARDS AND HEE OH

ABSTRACT. Let $G = \mathrm{SO}^\circ(n, 1) \times \mathrm{SO}^\circ(n, 1)$ and $X = \mathbb{H}^n \times \mathbb{H}^n$ for $n \geq 2$. For a pair (π_1, π_2) of non-elementary convex cocompact representations of a finitely generated group Σ into $\mathrm{SO}^\circ(n, 1)$, let $\Gamma = (\pi_1 \times \pi_2)(\Sigma)$. Denoting the bottom of the L^2 -spectrum of the negative Laplacian on $\Gamma \backslash X$ by λ_0 , we show:

- (1) $L^2(\Gamma \backslash G)$ is tempered and $\lambda_0 = \frac{1}{2}(n-1)^2$;
- (2) There exists no positive Laplace eigenfunction in $L^2(\Gamma \backslash X)$.

In fact, analogues of (1)-(2) hold for any Anosov subgroup Γ in the product of at least two simple algebraic groups of rank one as well as for Hitchin subgroups $\Gamma < \mathrm{PSL}_d(\mathbb{R})$, $d \geq 3$. Moreover, if G is a semisimple real algebraic group of rank at least 2, then (2) holds for any Anosov subgroup Γ of G .

CONTENTS

1. Introduction	1
2. Preliminaries and notations	6
3. Positive joint eigenfunctions and conformal densities	10
4. Eigenvalues of positive eigenfunctions	14
5. Groups of the second kind and positive joint eigenfunctions	16
6. The L^2 -spectrum and uniqueness	19
7. Smearing argument in higher rank	26
8. Injectivity radius and $L^2(G) \propto L^2(\Gamma \backslash G)$	29
9. Temperedness of $L^2(\Gamma \backslash G)$	33
References	36

1. INTRODUCTION

Motivation and background. Locally symmetric spaces provide key examples of Riemannian manifolds for which there exist numerous tools for studying various aspects of spectral geometry. For example, properties of dynamical systems related to the manifold are closely connected to the spectral theory of the Laplace operator, as well as to representation theory. While

Edwards was supported by funding from the Heilbronn Institute for Mathematical Research and Oh was supported in part by NSF grant DMS-1900101.

the spectral theory of finite-volume locally symmetric spaces has been quite extensively developed, the infinite volume setting provides many examples of interesting phenomena that are less well understood. Nevertheless, for rank one locally symmetric spaces of infinite volume, a number of key facts about the spectrum have been established.

Let (\mathbb{H}^n, d) , $n \geq 2$, denote the n -dimensional hyperbolic space of constant curvature -1 , and let $G = \text{Isom}^+(\mathbb{H}^n) \simeq \text{SO}^\circ(n, 1)$ denote the group of all orientation preserving isometries of \mathbb{H}^n . Let $\Gamma < G$ be a torsion-free¹ discrete subgroup. The critical exponent $0 \leq \delta = \delta_\Gamma \leq n-1$ is defined as the abscissa of convergence of the Poincaré series $\sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)}$ for $o \in \mathbb{H}^n$. We denote by Δ the hyperbolic Laplacian and by $\lambda_0 = \lambda_0(\Gamma \backslash \mathbb{H}^n)$ the bottom of the L^2 -spectrum of the negative Laplace operator $-\Delta$, which is given as

$$\lambda_0 := \inf \left\{ \frac{\int_{\Gamma \backslash \mathbb{H}^n} \|\text{grad } f\|^2 d\text{vol}}{\int_{\Gamma \backslash \mathbb{H}^n} |f|^2 d\text{vol}} : f \in C_c^\infty(\Gamma \backslash \mathbb{H}^n) \right\} \quad (1.1)$$

(see [45, Theorem 2.2]). In a series of papers, Elstrodt ([11], [12], [13]) and Patterson ([33], [34], [35]) developed the relationship between δ and λ_0 , proving the following theorem for $n = 2$. The general case is due to Sullivan [45, Theorem 2.21].

Theorem 1.1 (Generalized Elstrodt-Patterson I). *For any discrete subgroup $\Gamma < \text{SO}^\circ(n, 1)$, the following are equivalent:*

- (1) $\delta \leq \frac{1}{2}(n-1)$;
- (2) $\lambda_0 = \frac{1}{4}(n-1)^2$.

The right translation action of G on the quotient space $\Gamma \backslash G$ equipped with a G -invariant measure gives rise to a unitary representation of G on the Hilbert space $L^2(\Gamma \backslash G)$, called a quasi-regular representation of G . If we set $K \simeq \text{SO}(n)$ to be a maximal compact subgroup of G and identify \mathbb{H}^n with G/K , the space of K -invariant functions of $L^2(\Gamma \backslash G)$ can be identified with $L^2(\Gamma \backslash \mathbb{H}^n)$. The bottom of the L^2 -spectrum λ_0 then provides information on which complementary series representation of G can occur in $L^2(\Gamma \backslash G)$. Indeed, it follows from the classification of the unitary dual of $\text{SO}^\circ(n, 1)$ that $\lambda_0 = (n-1)^2/4$ is equivalent to saying that the quasi-regular representation $L^2(\Gamma \backslash G)$ does not contain any complementary series representation (cf. [45], [10]), which is again equivalent to the *temperedness* of $L^2(\Gamma \backslash G)$. As first introduced by Harish-Chandra [18], a unitary representation (π, \mathcal{H}_π) of a semisimple real algebraic group G is tempered (Definition 2.6) if all of its matrix coefficients belong to $L^{2+\varepsilon}(G)$ for any $\varepsilon > 0$, or, equivalently, if π is weakly contained² in the regular representation $L^2(G)$ ([8], see Proposition 2.7).

¹all discrete subgroups in this paper will be assumed to be torsion-free

² π is weakly contained in a unitary representation σ of G if any diagonal matrix coefficients of π can be approximated, uniformly on compact sets, by convex combinations of diagonal matrix coefficients of σ .

Therefore Theorem 1.1 can be rephrased as follows:

Theorem 1.2 (Generalized Elstrodt-Patterson II). *For any discrete subgroup $\Gamma < G = \mathrm{SO}^\circ(n, 1)$, the following are equivalent:*

- (1) $\delta \leq \frac{1}{2}(n - 1)$;
- (2) $L^2(\Gamma \backslash G)$ is tempered.

The size of the critical exponent δ is also related to the existence of a square-integrable positive Laplace eigenfunction on $\Gamma \backslash \mathbb{H}^n$. A discrete subgroup $\Gamma < G$ is called convex cocompact if there exists a convex subspace of \mathbb{H}^n on which Γ acts co-compactly. For convex cocompact subgroups of G (more generally for geometrically finite subgroups), Patterson and Sullivan showed the following using their theory of conformal measures on the boundary $\partial \mathbb{H}^n$ ([36], [46], [45, Theorem 2.21]):

Theorem 1.3 (Sullivan). *For a convex cocompact subgroup $\Gamma < \mathrm{SO}^\circ(n, 1)$, the following are equivalent:*

- (1) $\delta \leq \frac{1}{2}(n - 1)$;
- (2) *There exists no positive Laplace eigenfunction in $L^2(\Gamma \backslash \mathbb{H}^n)$.*

Since λ_0 divides the positive spectrum and the L^2 -spectrum on $\Gamma \backslash \mathbb{H}^n$ by Sullivan's theorem [45, Theorem 2.1] (see Theorem 4.1), (2) is equivalent to saying that any λ_0 -harmonic function (i.e., $-\Delta f = \lambda_0 f$) on $\Gamma \backslash \mathbb{H}^n$ is not square-integrable.

Main results. The main aim of this article is to discuss analogues of Theorems 1.1, 1.2, and 1.3 for a certain class of discrete subgroups of a connected semisimple real algebraic group of higher rank, i.e., rank at least 2.

We begin by describing a special case of our main theorem when $G = \mathrm{SO}^\circ(n_1, 1) \times \mathrm{SO}^\circ(n_2, 1)$ with $n_1, n_2 \geq 2$. Let X be the Riemannian product $\mathbb{H}^{n_1} \times \mathbb{H}^{n_2}$ and Δ the Laplace-Beltrami operator on X . For a torsion-free discrete subgroup $\Gamma < G$, a smooth function f on $\Gamma \backslash X$ is called λ -harmonic if $-\Delta f = \lambda f$. The number $\lambda_0 = \lambda_0(\Gamma \backslash X)$ is given in the same way as (1.1) replacing $\Gamma \backslash \mathbb{H}^n$ by $\Gamma \backslash X$.

Theorem 1.4. *Let*

$$\Gamma = (\pi_1 \times \pi_2)(\Sigma) = \{(\pi_1(\sigma), \pi_2(\sigma)) \in G : \sigma \in \Sigma\} \quad (1.2)$$

where $\pi_i : \Sigma \rightarrow \mathrm{SO}^\circ(n_i, 1)$ is a non-elementary convex cocompact representation of a finitely generated group Σ for $i = 1, 2$. Then

- (1) $L^2(\Gamma \backslash G)$ is tempered and $\lambda_0 = \frac{1}{4}((n_1 - 1)^2 + (n_2 - 1)^2)$;
- (2) *There exists no positive Laplace eigenfunction in $L^2(\Gamma \backslash X)$, or equivalently, no λ_0 -harmonic function is square-integrable.*

Remark 1.5. Theorem 1.4 does not hold for a general subgroup $\Gamma < G$ of infinite co-volume. For example, if $\Gamma < \mathrm{SO}^\circ(n_1, 1) \times \mathrm{SO}^\circ(n_2, 1)$ is the product of two convex cocompact subgroups, each of which having critical exponent greater than $\frac{1}{2}(n_i - 1)$, then $L^2(\Gamma \backslash G)$ is not tempered and $L^2(\Gamma \backslash X)$ possesses a positive Laplace eigenfunction.

We now discuss a general setting. Let G be a connected semisimple real algebraic group and X the associated Riemannian symmetric space. In the rest of the introduction, we assume that $\Gamma < G$ is a torsion-free Zariski dense discrete subgroup. We let $\psi_\Gamma : \mathfrak{a} \rightarrow \mathbb{R} \cup \{-\infty\}$ denote the *growth indicator function* of Γ as defined in (2.4), where \mathfrak{a} is the Lie algebra of a maximal real split torus of G . The function ψ_Γ can be regarded as a higher rank generalization of the critical exponent of Γ . Let ρ denote the half sum of all positive roots for $(\mathfrak{g}, \mathfrak{a})$, counted with multiplicity. Analogous to the fact that the critical exponent δ is always bounded above by $n - 1$ for a discrete subgroup $\Gamma < \mathrm{SO}^\circ(n, 1)$, we have the upper bound $\psi_\Gamma \leq 2\rho$ for any discrete subgroup Γ of G [40].

The following Theorem 1.6 generalizes Theorems 1.1, 1.2, and 1.3 to Anosov subgroups of G (with respect to a minimal parabolic subgroup of G) which are regarded as higher rank generalizations of convex cocompact subgroups. For $G = \mathrm{SO}^\circ(n_1, 1) \times \mathrm{SO}^\circ(n_2, 1)$, they are precisely given by the class of subgroups considered in Theorem 1.4. We refer to Definition 2.4 for a general case. We mention that they were first introduced by Labourie [27] for surface groups and then generalized by Guichard and Wienhard for hyperbolic groups [17] (see also [16], [21]).

In the following theorem, the norm $\|\rho\|$ is defined via the identification \mathfrak{a}^* and \mathfrak{a} using the Killing form on \mathfrak{g} . Denote by $\sigma(\Gamma \backslash X)$ the L^2 -spectrum of $-\Delta$ on $\Gamma \backslash X$.

Theorem 1.6. *Let G be a connected semisimple real algebraic group and Γ a Zariski dense Anosov subgroup of G . The following (1)-(3) are equivalent, and imply (4):*

- (1) $\psi_\Gamma \leq \rho$;
- (2) $L^2(\Gamma \backslash G)$ is tempered and $\lambda_0(\Gamma \backslash X) = \|\rho\|^2$;
- (3) $L^2(G)$ and $L^2(\Gamma \backslash G)$ are weakly contained in each other and $\sigma(\Gamma \backslash X) = \sigma(X) = [\|\rho\|^2, \infty)$;
- (4) *There exists no positive Laplace eigenfunction in $L^2(\Gamma \backslash X)$.*

Moreover, if $\mathrm{rank} G \geq 2$, then (4) always holds for any Anosov subgroup $\Gamma < G$.

Our proof of the implication (1) \Rightarrow (2) is based on the asymptotic behavior of the Haar matrix coefficients for Anosov subgroups obtained in [9] and [6] as well as Harish-Chandra's Plancherel formula (see Theorems 6.4 and 9.4). The implication (2) \Rightarrow (1) is true for a general discrete subgroup (see the proof of Theorem 9.4). The equivalence (2) \Leftrightarrow (3) uses the observation that $L^2(G)$ is weakly contained in $L^2(\Gamma \backslash G)$ whenever the injectivity radius of $\Gamma \backslash G$ is infinite, and that $\Gamma \backslash G$ has infinite injectivity radius for any Anosov subgroup $\Gamma < G$, except for cocompact lattices of a rank one Lie group (see Section 8). For (4), we first prove that any positive Laplace eigenfunction in $L^2(\Gamma \backslash X)$ is indeed a joint eigenfunction for the whole ring of G -invariant differential operators, which then can be studied via Γ -conformal measures on the Furstenberg boundary of G (see Sections 3 and 6). We establish

a higher rank version of Sullivan-Thurston's smearing theorem (Theorem 7.4) from which we deduce the non-existence of square-integrable positive Laplace eigenfunctions for any higher rank Anosov subgroup (see Section 7 and Corollary 7.2). When rank $G = 1$, Anosov subgroups are convex cocompact groups and the implication $(1) + (2) \Rightarrow (4)$ is obtained in [45] (see also [42, Theorem 3.1]) for $X = \mathbb{H}^n$ and in [50] in general.

Although the condition $\psi_\Gamma \leq \rho$ may appear quite strong, it was verified in a recent work of Kim-Minsky-Oh [23] for Anosov subgroups in the following setting, and hence we deduce from Theorem 1.6:

Theorem 1.7. *Let Γ be a Zariski dense Anosov subgroup of the product of at least two simple real algebraic groups of rank one, or a Zariski dense Anosov subgroup of a Hitchin subgroup of $\mathrm{PSL}_d(\mathbb{R})$ for $d \geq 3$. Then (1)-(4) of Theorem 1.6 hold.*

It is conjectured in [23] that any Anosov subgroup of a higher rank semisimple real algebraic group satisfies the condition $\psi_\Gamma \leq \rho$. This conjecture suggests that Anosov subgroups in higher rank groups are more like generalizations of convex cocompact subgroups of *small* critical exponent.

Groups of the second kind and positive joint eigenfunctions. For any discrete subgroup Γ which is not cocompact in G and for any $\lambda \leq \lambda_0(\Gamma \backslash X)$, Sullivan proved the existence of a positive λ -harmonic function. We prove a higher-rank strengthening of this result: for any discrete subgroup of the *second kind* (see Definition 5.1) whose limit cone is contained in the interior of \mathfrak{a}^+ and for any linear form $\psi \geq \psi_\Gamma$, we construct a positive joint eigenfunction with character corresponding to ψ (Theorem 5.2).

Organization: In section 2, we review the basic notions and notations which will be used throughout the paper. In section 3, we show that any positive joint eigenfunction on $\Gamma \backslash X$ (i.e., an eigenfunction for the whole ring of G -invariant differential operators) arises from a (Γ, ψ) -conformal density (Proposition 3.7). In section 4, we compute the Laplace eigenvalue of a positive joint eigenfunction associated to a (Γ, ψ) -conformal measure (Proposition 4.2). In section 5, we introduce the notion of subgroups of the second kind. We then construct positive joint eigenfunctions for any $\psi \geq \psi_\Gamma$ for any subgroup of the second kind with its limit cone contained in $\mathrm{int} \mathfrak{a}^+ \cup \{0\}$ (Theorem 5.2). In section 6, we compute the L^2 -spectrum of X (Theorem 6.3) and show that $\lambda_0 = \|\rho\|^2$ if $L^2(\Gamma \backslash G)$ is tempered (Theorem 6.4). We show that a positive Laplace eigenfunction in $L^2(\Gamma \backslash X)$ is necessarily a joint eigenfunction (Corollary 6.6) and a spherical vector of a unique irreducible subrepresentation of $L^2(\Gamma \backslash G)$ (Theorem 6.8). In section 7, we prove a higher rank version of Sullivan-Thurston's smearing theorem (Theorem 7.4) to obtain the non-existence theorem of L^2 -positive Laplace eigenfunctions in higher rank. In section 8, we prove the weak containment $L^2(G) \propto L^2(\Gamma \backslash G)$ for all Anosov subgroups Γ in higher rank groups. In

section 9, we prove the equivalence of the temperedness of $L^2(\Gamma \backslash G)$ and $\psi_\Gamma \leq \rho$ (Theorem 9.4). We also deduce Theorem 1.6.

Acknowledgements: We would like to thank Marc Burger for bringing the reference [45] to our attention. We would also like to thank Dick Canary, Francois Labourie, Curt McMullen and Dennis Sullivan for useful conversations.

2. PRELIMINARIES AND NOTATIONS

Let G be a connected semisimple real algebraic group, i.e., the identity component of the group of real points of a semisimple algebraic group defined over \mathbb{R} . Let $\Gamma < G$ be a torsion-free discrete subgroup. Let P be a minimal parabolic subgroup of G with a fixed Langlands decomposition $P = MAN$ where A is a maximal real split torus of G , M is the maximal compact subgroup of P , which commutes with A , and N is the unipotent radical of P . We denote by $\mathfrak{g}, \mathfrak{a}, \mathfrak{n}$ respectively the Lie algebras of G, A, N . We fix a positive Weyl chamber $\mathfrak{a}^+ \subset \mathfrak{a}$ so that \mathfrak{n} consists of positive root subspaces. Let Σ^+ denote the set of all positive roots for $(\mathfrak{g}, \mathfrak{a}^+)$. We also write $\Pi \subset \Sigma^+$ for the set of all simple roots. We denote by

$$\rho = \frac{1}{2} \sum_{\alpha \in \Sigma^+} \alpha$$

the half sum of the positive roots for $(\mathfrak{g}, \mathfrak{a}^+)$, counted with multiplicity. We denote by $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$ the inner product and norm on \mathfrak{g} respectively, induced from the Killing form: $B(x, y) = \text{Tr}(\text{ad } x \text{ ad}(y))$ for $x, y \in \mathfrak{g}$.

We fix a maximal compact subgroup K of G so that the Cartan decomposition $G = K(\exp \mathfrak{a}^+)K$ holds, that is, for any $g \in G$, there exists a unique element $\mu(g) \in \mathfrak{a}^+$ such that $g \in K \exp \mu(g)K$. We call the map $\mu : G \rightarrow \mathfrak{a}^+$ the Cartan projection map.

The Riemannian symmetric space (X, d) can be identified with the quotient space G/K with the metric d induced from $\langle \cdot, \cdot \rangle$. We denote by $d \text{ vol}$ the Riemannian volume form on X or on $\Gamma \backslash X$. We also use dx to denote this volume form as well as the Haar measure on G , or on $\Gamma \backslash G$. In particular, $d(\cdot, \cdot)$ will denote both the left G -invariant Riemannian distance function on X , as well as the left G -invariant and right K -invariant distance on G . We set $o = [K] \in X$. We then have $\|\mu(g)\| = d(go, o)$ for $g \in G$. We do not distinguish a function on X and a right K -invariant function on G .

Let $w_0 \in K$ be an element of the normalizer of A so that $\text{Ad}_{w_0} \mathfrak{a}^+ = -\mathfrak{a}^+$. The opposition involution $i : \mathfrak{a} \rightarrow \mathfrak{a}$ is defined by

$$i(u) = -\text{Ad}_{w_0}(u) \quad \text{for all } u \in \mathfrak{a}. \quad (2.1)$$

Let $\mathcal{F} := G/P$ denote the Furstenberg boundary of G . We define the following *visual* maps $G \rightarrow \mathcal{F}$: for each $g \in G$,

$$g^+ := gP \in \mathcal{F} \quad \text{and} \quad g^- := gw_0P \in \mathcal{F}. \quad (2.2)$$

The unique open G -orbit $\mathcal{F}^{(2)}$ in $\mathcal{F} \times \mathcal{F}$ under the diagonal G -action is given by:

$$\mathcal{F}^{(2)} = G(e^+, e^-) = \{(g^+, g^-) \in \mathcal{F} \times \mathcal{F} : g \in G\}.$$

Two points ξ, η in \mathcal{F} are said to be in general position if $(\xi, \eta) \in \mathcal{F}^{(2)}$.

Conformal measures. Let $G = KAN$ be the Iwasawa decomposition, $\kappa : G \rightarrow K$ the K -factor projection of this decomposition, and $H : G \rightarrow \mathfrak{a}$ be the Iwasawa cocycle defined by the relation: for $g \in G$,

$$g \in \kappa(g) \exp(H(g))N.$$

Note that K acts transitively on \mathcal{F} and $K \cap P = M$, and hence we may identify \mathcal{F} with K/M . The Iwasawa decomposition can be used to describe both the action of G on $\mathcal{F} = K/M$ and the \mathfrak{a} -valued Busemann map as follows: for all $g \in G$ and $[k] \in \mathcal{F}$ with $k \in K$,

$$g \cdot [k] = [\kappa(gk)],$$

and the \mathfrak{a} -valued Busemann map is defined by

$$\beta_{[k]}(g(o), h(o)) := H(g^{-1}k) - H(h^{-1}k) \in \mathfrak{a} \quad \text{for all } g, h \in G.$$

We denote by $\mathfrak{a}^* = \text{Hom}_{\mathbb{R}}(\mathfrak{a}, \mathbb{R})$ the space of all linear forms on \mathfrak{a} .

Definition 2.1. Let $\psi \in \mathfrak{a}^*$.

(1) A finite Borel measure ν_o on $\mathcal{F} = K/M$ is said to be a (Γ, ψ) -conformal measure (with respect to $o = [K]$) if for all $\gamma \in \Gamma$ and $\xi = [k] \in K/M$,

$$\frac{d\gamma_*\nu_o}{d\nu_o}(\xi) = e^{-\psi(\beta_\xi(\gamma o, o))} = e^{-\psi(H(\gamma^{-1}k))},$$

or equivalently

$$d\nu_o([k]) = e^{\psi(H(\gamma k))} d\nu_o(\gamma \cdot [k]),$$

where $\gamma_*\nu_o(Q) = \nu_o(\gamma^{-1}Q)$ for any Borel subset $Q \subset \mathcal{F}$. Unless mentioned otherwise, all conformal measures in this paper are assumed to be with respect to o .

(2) A collection $\{\nu_x : x \in X\}$ of finite Borel measures on \mathcal{F} is called a (Γ, ψ) -conformal density if for all $x, y \in X$, $\xi \in \mathcal{F}$ and $\gamma \in \Gamma$,

$$\frac{d\nu_x}{d\nu_y}(\xi) = e^{-\psi(\beta_\xi(x, y))} \quad \text{and} \quad d\gamma_*\nu_x = d\nu_{\gamma(x)}. \quad (2.3)$$

A (Γ, ψ) -conformal measure ν_o defines a (Γ, ψ) -conformal density $\{\nu_x : x \in X\}$ by the formula:

$$d\nu_x(\xi) = e^{-\psi(\beta_\xi(x, o))} d\nu_o(\xi),$$

and conversely any (Γ, ψ) -conformal density $\{\nu_x\}$ is uniquely determined by its member ν_o by (2.3). For this reason, by abuse of terminology, we sometimes do not distinguish conformal measures and conformal densities.

Growth indicator function. Let $\Gamma < G$ be a Zariski dense discrete subgroup. Following Quint [40], let $\psi_\Gamma : \mathfrak{a} \rightarrow \mathbb{R} \cup \{-\infty\}$ denote the growth indicator function of Γ : for any non-zero $v \in \mathfrak{a}$,

$$\psi_\Gamma(v) := \|v\| \inf_{v \in \mathcal{C}} \tau_{\mathcal{C}}, \quad (2.4)$$

where the infimum is over all open cones \mathcal{C} containing v and $\tau_{\mathcal{C}}$ denotes the abscissa of convergence of the series $\sum_{\gamma \in \Gamma, \mu(\gamma) \in \mathcal{C}} e^{-s\|\mu(\gamma)\|}$. For $v = 0$, we let $\psi_\Gamma(0) = 0$. We note that ψ_Γ does not change if we replace the norm $\|\cdot\|$ by any other norm on \mathfrak{a} . For any discrete subgroup $\Gamma < G$, we have the upper bound $\psi_\Gamma \leq 2\rho$ [40]. On the other hand, when Γ is of infinite co-volume in a simple real algebraic group of rank at least 2, Quint deduced from [32] that $\psi_\Gamma \leq 2\rho - \eta_G$, where η_G is the half sum of a maximal strongly orthogonal subset of the root system of G ([41], see also [31, Theorem 7.1]).

Limit cone and limit set. The limit cone $\mathcal{L} = \mathcal{L}_\Gamma$ of Γ is defined as the asymptotic cone of $\mu(\Gamma)$, i.e.,

$$\mathcal{L} = \{\lim t_i \mu(\gamma_i) \in \mathfrak{a}^+ : t_i \rightarrow 0, \gamma_i \in \Gamma\}.$$

Benoist showed that for Γ Zariski dense, \mathcal{L} is a convex cone with non-empty interior [2]. Quint [40] showed that ψ_Γ is a concave and upper-semicontinuous function such that $\psi_\Gamma \geq 0$ on \mathcal{L} , $\psi_\Gamma > 0$ on $\text{int } \mathcal{L}$ and $\psi_\Gamma = -\infty$ outside \mathcal{L} .

For a sequence $g_i \in G$, we write $g_i \rightarrow \infty$ regularly if $\alpha(\mu(g_i)) \rightarrow \infty$ for all $\alpha \in \Pi$. For $g \in G$, we write $g = \kappa_1(g) \exp(\mu(g)) \kappa_2(g) \in KA^+K$; if $\mu(g) \in \text{int } \mathfrak{a}^+$, then $[\kappa_1(g)] \in K/M = \mathcal{F}$ is well-defined.

Definition 2.2. A sequence $p_i \in X$ is said to converge to $\xi \in \mathcal{F}$ and we write $\lim_{i \rightarrow \infty} p_i = \xi$ if there exists a sequence $g_i \rightarrow \infty$ regularly in G with $p_i = g_i(o)$ and $\lim_{i \rightarrow \infty} [\kappa_1(g_i)] = \xi$.

We denote by $\Lambda \subset \mathcal{F}$ the limit set of Γ , which is defined as

$$\Lambda = \{\lim \gamma_i(o) \in \mathcal{F} : \gamma_i \in \Gamma\}. \quad (2.5)$$

For $\Gamma < G$ Zariski dense, this is the unique Γ -minimal subset of \mathcal{F} ([2], [30]).

Tangent linear forms. We set

$$D_\Gamma = \{\psi \in \mathfrak{a}^* : \psi \geq \psi_\Gamma\}. \quad (2.6)$$

A linear form $\psi \in \mathfrak{a}^*$ is said to be tangent to ψ_Γ at $u \in \mathfrak{a}$ if $\psi \in D_\Gamma$ and $\psi(u) = \psi_\Gamma(u)$. We denote by D_Γ^* the set of all linear forms tangent to ψ_Γ at $\mathcal{L} \cap \text{int } \mathfrak{a}^+$, i.e.,

$$D_\Gamma^* := \{\psi \in D_\Gamma : \psi(u) = \psi_\Gamma(u) \text{ for some } u \in \mathcal{L} \cap \text{int } \mathfrak{a}^+\}. \quad (2.7)$$

For $\Gamma < \text{SO}^\circ(n, 1)$ and δ its critical exponent, we have $D_\Gamma^* = \{\delta\}$ and $D_\Gamma = \{s \geq \delta\}$.

Extending the construction of Patterson [36] and Sullivan [44], Quint [39] showed the following:

Theorem 2.3. *For any $\psi \in D_\Gamma^*$, there exists a (Γ, ψ) -conformal measure supported on Λ .*

Anosov subgroups. Let Σ be a finitely generated group. For $\sigma \in \Sigma$, let $|\sigma|$ denote the word length of σ for some fixed symmetric generating set of Σ .

Definition 2.4. ([17], [21], [16], [3]) A representation $\pi : \Sigma \rightarrow G$ is Anosov with respect to P if there exist a constant $c > 0$ such that for all $\sigma \in \Sigma$ and $\alpha \in \Pi$,

$$\alpha(\mu(\pi(\sigma))) \geq c|\sigma| - c^{-1}. \quad (2.8)$$

A discrete subgroup $\Gamma < G$ is called an Anosov subgroup (with respect to P) if Γ can be realized as the image $\pi(\Sigma)$ of an Anosov representation $\pi : \Sigma \rightarrow G$. If $\Gamma = \pi(\Sigma)$ is Anosov, then Σ is a Gromov hyperbolic group ([21], [3]). As mentioned in the introduction, Anosov subgroups of G were first introduced by Labourie for surface groups [27], and then extended by Guichard and Wienhard [17] to general word hyperbolic groups. Several equivalent characterizations have been established, one of which is the above definition (see [16], [21]). When G has rank one, the class of Anosov subgroups coincides with that of convex cocompact subgroups, and when G is a product of two rank one simple algebraic groups, any Anosov subgroup arises in a similar fashion to (1.2). Examples of Anosov subgroups include Schottky groups (cf. [9, Def. 7.1]), as well as Hitchin subgroups defined as follows. Let ι_d denote the irreducible representation $\mathrm{PSL}_2(\mathbb{R}) \rightarrow \mathrm{PSL}_d(\mathbb{R})$, which is unique up to conjugations. A Hitchin subgroup is the image of a representation $\pi : \Sigma \rightarrow \mathrm{PSL}_d(\mathbb{R})$ of a uniform lattice $\Sigma < \mathrm{PSL}_2(\mathbb{R})$, which belongs to the same connected component as $\iota_d|\Sigma$ in the character variety $\mathrm{Hom}(\Sigma, \mathrm{PSL}_d(\mathbb{R}))/\sim$ where the equivalence is given by conjugations.

One of the important features of an Anosov subgroup is the following:

Theorem 2.5. [38] *For any Anosov subgroup $\Gamma < G$, we have*

$$\mathcal{L} \subset \mathrm{int} \, \mathfrak{a}^+ \cup \{0\}.$$

Tempered representations. By definition, a unitary representation of G is a Hilbert space \mathcal{H}_π equipped with a strongly continuous homomorphism π from G to the group of unitary operators on \mathcal{H}_π . Given two unitary representations π and σ of G , π is said to be weakly contained in σ if any diagonal matrix coefficients of π can be approximated, uniformly on compact sets, by convex combinations of diagonal matrix coefficients of σ . We use the notation $\pi \propto \sigma$ for the weak containment.

The Harish-Chandra function $\Xi_G : G \rightarrow (0, \infty)$ is a bi- K -invariant function defined via the formula

$$\Xi_G(g) = \int_K e^{-\rho(H(gk))} dk \quad \text{for all } g \in G$$

where dk denotes the probability Haar measure on K . The following estimate is well-known, cf. e.g. [24]: for any $\varepsilon > 0$, there exist $C, C_\varepsilon > 0$ such that for any $g \in G$,

$$Ce^{-\rho(\mu(g))} \leq \Xi_G(g) \leq C_\varepsilon e^{-(1-\varepsilon)\rho(\mu(g))}. \quad (2.9)$$

Definition 2.6. A unitary representation (π, \mathcal{H}_π) of G is called *tempered* if for any K -finite unit vectors $v, w \in \mathcal{H}_\pi$ and any $g \in G$,

$$|\langle \pi(g)v, w \rangle| \leq (\dim \langle Kv \rangle \dim \langle Kw \rangle)^{1/2} \Xi_G(g),$$

where $\langle Kv \rangle$ denotes the linear subspace of \mathcal{H}_π spanned by Kv .

Proposition 2.7. [8] *The following are equivalent for a unitary representation (π, \mathcal{H}_π) of G :*

- (1) π is tempered;
- (2) $\pi \propto L^2(G)$;
- (3) for any vectors $v, w \in \mathcal{H}_\pi$, the matrix coefficient $g \mapsto \langle \pi(g)v, w \rangle$ lies in $L^{2+\varepsilon}(G)$ for any $\varepsilon > 0$;
- (4) for any $\varepsilon > 0$, π is strongly $L^{2+\varepsilon}$, i.e., there exists a dense subset of \mathcal{H}_π whose matrix coefficients all belong to $L^{2+\varepsilon}(G)$.

In the whole paper, the notation $f(v) \asymp g(v)$ means that the ratio $f(v)/g(v)$ is bounded uniformly between two positive constants, and $f \ll g$ means that $|f| \leq c|g|$ for some $c > 0$.

3. POSITIVE JOINT EIGENFUNCTIONS AND CONFORMAL DENSITIES

Let G be a connected semisimple real algebraic group and $\Gamma < G$ be a Zariski dense discrete subgroup. The main goal of this section is to obtain Proposition 3.7, which explains the relationship between positive joint eigenfunctions on $\Gamma \backslash X$ and Γ -conformal measures on the Furstenberg boundary of G .

Joint eigenfunctions on X . Let $\mathcal{D} = \mathcal{D}(X)$ denote the ring of all G -invariant differential operators on X . We call a real valued function on X a *joint eigenfunction* if it is an eigenfunction for all operators in \mathcal{D} . For each joint eigenfunction f , there exists an associated character $\chi_f : \mathcal{D} \rightarrow \mathbb{R}$ such that

$$Df = \chi_f(D)f$$

for all elements $D \in \mathcal{D}$. The ring \mathcal{D} is generated by $\text{rank}(G)$ elements, and the set of all characters of \mathcal{D} is in bijection with the space $\mathfrak{a}^* = \text{Hom}_{\mathbb{R}}(\mathfrak{a}, \mathbb{R})$ modulo the action of the Weyl group, as we now explain. Denote by $Z(\mathfrak{g}_{\mathbb{C}})$ the center of the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ of $\mathfrak{g}_{\mathbb{C}}$. Recall the well-known fact that the joint eigenfunctions on X can be identified with the right K -invariant real-valued $\mathcal{Z}(\mathfrak{g}_{\mathbb{C}})$ -eigenfunctions on G (cf. [19]).

Letting T be a maximal torus in M with Lie algebra \mathfrak{t} , set $\mathfrak{h} = (\mathfrak{a} \oplus \mathfrak{t})$. Then $\mathfrak{h}_{\mathbb{C}} := (\mathfrak{a} \oplus \mathfrak{t})_{\mathbb{C}}$ is a Cartan subalgebra of $\mathfrak{g}_{\mathbb{C}}$. We let

$$\iota : \mathcal{Z}(\mathfrak{g}_{\mathbb{C}}) \rightarrow \mathcal{S}^W(\mathfrak{h}_{\mathbb{C}})$$

denote the Harish-Chandra isomorphism from $Z(\mathfrak{g}_{\mathbb{C}})$ to the Weyl group-invariant elements of the symmetric algebra $\mathcal{S}(\mathfrak{h}_{\mathbb{C}})$ of $\mathfrak{h}_{\mathbb{C}}$ [24, Theorem 8.18].

For any $\psi \in \mathfrak{a}^*$, we can extend it to \mathfrak{h} by letting $\psi(J) = 0$ for all $J \in \mathfrak{m}$, and then to $\mathcal{S}(\mathfrak{h}_{\mathbb{C}})$ polynomially. This lets us define a character χ_{ψ} on $\mathcal{Z}(\mathfrak{g}_{\mathbb{C}})$ by

$$\chi_{\psi}(Z) := \psi(\iota(Z)) \quad (3.1)$$

for all $Z \in \mathcal{Z}(\mathfrak{g}_{\mathbb{C}})$. Conversely, if f is a right K -invariant $\mathcal{Z}(\mathfrak{g}_{\mathbb{C}})$ -eigenfunction, then, since \mathfrak{t} acts trivially on f , the associated character χ_f must arise as $\psi \circ \iota$ for some $\psi \in \mathfrak{a}^*$.

Example 3.1.

- Consider the hyperbolic space $\mathbb{H}^n = \{(x_1, \dots, x_{n-1}, y) \in \mathbb{R}^n : y > 0\}$ with the metric $\frac{\sqrt{\sum_{i=1}^{n-1} dx_i^2 + dy^2}}{y}$. The Laplacian Δ on \mathbb{H}^n is given as $\Delta = -y^2(\sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} + \frac{\partial^2}{\partial y^2}) + (n-2)y \frac{\partial}{\partial y}$ and the ring of $\mathrm{SO}^\circ(n, 1)$ -invariant differential operators is generated by Δ , i.e., a polynomial in Δ . If $\psi \in \mathfrak{a}^*$ is given by $\psi(v) = \delta v$ for some $\delta \in \mathbb{R}$ under the isomorphism $\mathfrak{a} = \mathbb{R}$, then $\chi_{\psi}(-\Delta) = \delta(n-1-\delta)$.
- Let $G = \mathrm{SO}^\circ(n_1, 1) \times \mathrm{SO}^\circ(n_2, 1)$ and X be the Riemannian product $\mathbb{H}^{n_1} \times \mathbb{H}^{n_2}$ for $n_1, n_2 \geq 2$. Then $\mathcal{D}(X)$ is generated by the hyperbolic Laplacians Δ_1, Δ_2 on each factor \mathbb{H}^{n_1} and \mathbb{H}^{n_2} . If we identify \mathfrak{a} with \mathbb{R}^2 and if a linear form $\psi \in \mathfrak{a}^*$ is given by $\psi(v) = \langle v, (\delta_1, \delta_2) \rangle$ for some vector $(\delta_1, \delta_2) \in \mathbb{R}^2$, then $\chi_{\psi}(-\Delta_i) = \delta_i(n_i - 1 - \delta_i)$ for $i = 1, 2$.

Joint eigenfunctions on $\Gamma \backslash X$. We now consider joint eigenfunctions on $\Gamma \backslash X$ or, equivalently, Γ -invariant joint eigenfunctions on X .

Definition 3.2. Let $\psi \in \mathfrak{a}^*$. Associated to a (Γ, ψ) -conformal density $\nu = \{\nu_x : x \in X\}$ on \mathcal{F} , we define the following function E_{ν} on G : for $g \in G$,

$$E_{\nu}(g) := |\nu_{g(o)}| = \int_{\mathcal{F}} e^{-\psi(H(g^{-1}k))} d\nu_o([k]). \quad (3.2)$$

Since $|\nu_{\gamma(x)}| = |\nu_x|$ for all $\gamma \in \Gamma$ and $x \in X$, the left Γ -invariance and right K -invariance of E_{ν} are clear. Hence we may consider E_{ν} as a K -invariant function on $\Gamma \backslash G$, or, equivalently, as a function on $\Gamma \backslash X$.

Proposition 3.3. For each (Γ, ψ) -conformal density ν on \mathcal{F} , E_{ν} is a positive joint eigenfunction on $\Gamma \backslash X$ with character $\chi_{\psi-\rho}$. Conversely, any positive joint eigenfunction on $\Gamma \backslash X$ arises in this way for some $\psi \geq \rho$ and a (Γ, ψ) -conformal density ν with (ψ, ν) uniquely determined.

In order to prove this proposition, we consider the following right K -invariant function on G for each $\psi \in \mathfrak{a}^*$ and $h \in G$:

$$\varphi_{\psi, h}(g) = e^{-\psi(H(g^{-1}h))} \quad (3.3)$$

so that

$$E_{\nu}(g) = \int_{\mathcal{F}} \varphi_{\psi, k}(g) d\nu_o([k]).$$

We may also consider $\varphi_{\psi,h}$ as a function on X . Hence the first part of Proposition 3.3 is a consequence of the following:

Lemma 3.4. ([24, Propositions 8.22 and 9.9]) *For any $\psi \in \mathfrak{a}^*$ and $h \in G$, the function $\varphi_{\psi,h}$ is a joint eigenfunction on X with character $\chi_{\psi-\rho}$.*

Proof. While we refer to [24] for the full proof, we outline some of the key points below, as we will use some part of this proof later. Since the elements of $\mathcal{Z}(\mathfrak{g}_{\mathbb{C}})$ commute with translation, we simply need to prove that

$$[Z\varphi_{\psi,e}](e) = \chi_{\psi-\rho}(Z)\varphi_{\psi,e}(e) \quad \text{for any } Z \in \mathcal{Z}(\mathfrak{g}_{\mathbb{C}});$$

the same identity will then hold for the function $g \mapsto \varphi_{\psi,e}(h^{-1}g)$, and thus also for $\varphi_{\psi,h}$ for any $h \in G$. Following [24, Chapter VII], we define the (non-unitary) principal series representation U^{ψ} : for all $g \in G$ and $f \in C(K)$

$$[U^{\psi}(g)f](k) := e^{-\psi(H(g^{-1}k))} f(\kappa(g^{-1}k))$$

for all $k \in K$. This extends to a representation dU^{ψ} of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ on the right M -invariant functions in $C^{\infty}(K)$ by way of the formula

$$[dU^{\psi}(X)f](k) = \frac{d}{dt} \bigg|_{t=0} [U^{\psi}(\exp(tX))f](k) \quad \text{for any } X \in \mathfrak{g}.$$

Observe that $[Z\varphi_{\psi,e}](e) = [dU^{\psi}(Z)](e)$, so in order to prove the proposition, it suffices to show that $dU^{\psi}(Z) = \chi_{\psi-\rho}(Z)$ for all $Z \in \mathcal{Z}(\mathfrak{g}_{\mathbb{C}})$.

The next key observation is that

$$Z(\mathfrak{g}_{\mathbb{C}}) \subset \mathcal{U}(\mathfrak{h}_{\mathbb{C}}) \oplus \mathfrak{n}\mathcal{U}(g_{\mathbb{C}}).$$

We thus write

$$Z = Y + \sum_i X_i U_i,$$

where $Y \in \mathcal{U}(\mathfrak{h}_{\mathbb{C}})$, $X_i \in \mathfrak{n}$, and $U_i \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}})$. Note that in this decomposition, Y is uniquely defined. Now, for arbitrary $X \in \mathfrak{n}$ and f ,

$$\begin{aligned} [dU^{\psi}(X)f](e) &= \frac{d}{dt} \bigg|_{t=0} [U^{\psi}(\exp(tX))f](e) = \frac{d}{dt} \bigg|_{t=0} [U^{\psi}(\exp(tX))f](e) \\ &= \frac{d}{dt} \bigg|_{t=0} e^{-\psi(H(\exp(-tX)))} f(\kappa(\exp(-tX))) = \frac{d}{dt} \bigg|_{t=0} f(e) = 0, \end{aligned}$$

so applying this to the X_i and functions $dU^{\psi}(U_i)f$ gives

$$[dU^{\psi}(X_i U_i)f](e) = [dU^{\psi}(X_i)(dU^{\psi}(U_i)f)](e) = 0,$$

hence $[dU^{\psi}(Z)f](e) = [dU^{\psi}(Y)f](e)$. For $L \in \mathfrak{m}$, we have $f(\exp(-L)) = f(e)$, so $[dU^{\psi}(J)f](e) = 0$ for all $J \in \mathfrak{t}$. Thus, it is only the \mathfrak{a} component of Y that contributes to $[dU^{\psi}(Y)f](e)$. Finally, note that for $X \in \mathfrak{a}$, we have

$$\begin{aligned} [dU^{\psi}(X)f](e) &= \frac{d}{dt} \bigg|_{t=0} e^{-\psi(H(\exp(-tX)))} f(\kappa(\exp(-tX))) \\ &= \frac{d}{dt} \bigg|_{t=0} e^{t\psi(X)} f(e) = \psi(X)f(e). \end{aligned}$$

Since the Harish-Chandra isomorphism consists of projection onto $\mathcal{U}(\mathfrak{h}_{\mathbb{C}})$ and then composition with the “ δ -shift” $H \mapsto H + \delta(H)1 = H + \rho(H)1$, where $\delta \in \mathfrak{h}_{\mathbb{C}}^*$ is the half-sum of the positive roots for $\mathfrak{g}_{\mathbb{C}}$, this shows that $dU^{\psi}(Z) = \chi_{\psi-\rho}(Z)$. \square

Letting $h = kan \in KAN$, we see that for any $g \in G$,

$$\varphi_{\psi,h}(g) = e^{-\psi(H(g^{-1}h))} = e^{-\psi(H(g^{-1}kan))} = e^{-\psi(H(g^{-1}k))} \cdot e^{-\psi(\log(a))},$$

i.e., the function $\varphi_{\psi,h}$ is a scalar multiple of $\varphi_{\psi,k(h)}$. In fact, the functions $\varphi_{\psi,k}$, $k \in K$ form a complete set of minimal positive joint eigenfunctions with character $\chi_{\psi-\rho}$ with $\psi \geq \rho$, in the sense that if f is a positive joint eigenfunction on X with character $\chi_{\psi-\rho}$ such that $f \leq \varphi_{\psi,k}$ for some $k \in K$, then

$$f = c \cdot \varphi_{\psi,k}$$

for some $c > 0$ (cf. [15, 22], see also [27, Theorem 1]).

As a consequence, we have the following (cf. [27, Theorem 3]):

Theorem 3.5. *For any positive joint eigenfunction f on X , there exist $\psi \in \mathfrak{a}^*$ with $\psi \geq \rho$ and a Borel measure ν_o on $\mathcal{F} = K/M$ such that for all $g \in G$,*

$$f(g) = \int_{\mathcal{F}} \varphi_{\psi,k}(g) d\nu_o([k]).$$

Moreover, the pair (ψ, ν_o) is uniquely determined by f .

Proof of the second part of Proposition 3.3: Let f be a Γ -invariant joint eigenfunction on X . By Theorem 3.5, there exist unique $\psi \in \mathfrak{a}^*$ and a Borel measure ν_o on \mathcal{F} so that for all $g \in G$,

$$f(g) = \int_{\mathcal{F}} \varphi_{\psi,k}(g) d\nu_o([k]).$$

Since f is Γ -invariant, for any $\gamma \in \Gamma$,

$$\begin{aligned} f(g) &= f(\gamma g) = \int_{\mathcal{F}} \varphi_{\psi,k}(\gamma g) d\nu_o([k]) \\ &= \int_{\mathcal{F}} \varphi_{\psi,\kappa(\gamma^{-1}k)}(g) e^{-\psi(H(\gamma^{-1}k))} d\nu_o([k]) \\ &= \int_{\mathcal{F}} \varphi_{\psi,\tilde{k}}(g) e^{\psi(H(\gamma\tilde{k}))} d\nu_o(\gamma \cdot [\tilde{k}]). \end{aligned}$$

By the uniqueness of ν_o in the integral representation of f ,

$$d\nu_o([k]) = e^{\psi(H(\gamma k))} d\nu_o(\gamma \cdot [k]).$$

Hence $\nu = \{\nu_x\}$ is a (Γ, ψ) -conformal density on \mathcal{F} , finishing the proof.

We denote by $\psi_{\Gamma} : \mathfrak{a} \rightarrow \mathbb{R} \cup \{-\infty\}$ the growth indicator function of Γ as defined in (2.4).

Theorem 3.6. [39, Theorem 8.1]. *Let $\Gamma < G$ be Zariski dense. If there exists a (Γ, ψ) -conformal measure on \mathcal{F} for some $\psi \in \mathfrak{a}^*$, then*

$$\psi \geq \psi_\Gamma.$$

Therefore Proposition 3.3 and Theorem 3.6 yield the following:

Proposition 3.7. *Let $\Gamma < G$ be a Zariski dense discrete subgroup. If ν is a (Γ, ψ) -conformal density for some $\psi \in \mathfrak{a}^*$, then E_ν is a positive joint eigenfunction on $\Gamma \backslash X$ with character $\chi_{\psi-\rho}$. Conversely, any positive joint eigenfunction on $\Gamma \backslash X$ is of the form E_ν for some (Γ, ψ) -conformal density ν with $\psi \geq \max(\rho, \psi_\Gamma)$, where (ψ, ν) is uniquely determined.*

4. EIGENVALUES OF POSITIVE EIGENFUNCTIONS

Let Γ be a torsion-free discrete subgroup of a connected semisimple real algebraic group G . Let Δ denote the Laplace-Beltrami operator on X or on $\Gamma \backslash X$. Since Δ is an elliptic differential operator, an eigenfunction is always smooth. We call a smooth function λ -harmonic if

$$-\Delta f = \lambda f.$$

Let $\mathcal{C} \in \mathcal{Z}(\mathfrak{g}_\mathbb{C})$ denote the Casimir operator on $C^\infty(G)$ (or on $C^\infty(\Gamma \backslash G)$) whose restriction to K -invariant functions coincides with Δ . Then K -invariant \mathcal{C} -eigenfunctions on $\Gamma \backslash G$ correspond to Laplace eigenfunctions on $\Gamma \backslash X$. In particular, a joint eigenfunction of $\Gamma \backslash X$ is a Laplace eigenfunction.

Define the real number $\lambda_0 = \lambda_0(\Gamma \backslash X) \in [0, \infty)$ as follows:

$$\lambda_0 := \inf \left\{ \frac{\int_{\Gamma \backslash X} \|\operatorname{grad} f\|^2 d\operatorname{vol}}{\int_{\Gamma \backslash X} |f|^2 d\operatorname{vol}} : f \in C_c^\infty(\Gamma \backslash X), f \neq 0 \right\}. \quad (4.1)$$

Positive Laplace eigenfunctions.

Theorem 4.1. [45, Theorem 2.1, 2.2] *Suppose that $\Gamma \backslash X$ is not compact.*

- (1) *For any $\lambda \leq \lambda_0$, there exists a positive λ -harmonic function on $\Gamma \backslash X$;*
- (2) *For any $\lambda > \lambda_0$, there is no positive λ -harmonic function on $\Gamma \backslash X$.*

We identify \mathfrak{a}^* with \mathfrak{a} via the inner product on \mathfrak{a} induced by the Killing form on \mathfrak{g} . This endows an inner product on \mathfrak{a}^* . More precisely, for each $\psi \in \mathfrak{a}^*$, there exist a unique $v_\psi \in \mathfrak{a}$ such that $\psi = \langle v_\psi, \cdot \rangle$. Then $\langle \psi_1, \psi_2 \rangle = \langle v_{\psi_1}, v_{\psi_2} \rangle$. Equivalently, fixing an orthonormal basis $\{H_i\}$ of \mathfrak{a} , we have $\langle \psi_1, \psi_2 \rangle = \sum_i \psi_1(H_i) \psi_2(H_i)$.

For $\psi \in \mathfrak{a}^*$, we set

$$\lambda_\psi := (\|\rho\|^2 - \|\psi - \rho\|^2). \quad (4.2)$$

Proposition 4.2. (1) *A positive joint eigenfunction on X with character $\chi_{\psi-\rho}$, $\psi \in \mathfrak{a}^*$, is λ_ψ -harmonic.*

- (2) *A positive Laplace eigenfunction on X is λ_ψ -harmonic for some $\psi \in \mathfrak{a}^*$ with $\psi \geq \rho$.*

Proof. Let $\psi \in \mathfrak{a}^*$. Recall the functions $\varphi_{\psi,h}$ in (3.3). By Theorem 3.5, (1) follows if we show that for any $h \in G$,

$$-\mathcal{C}\varphi_{\psi,h} = \lambda_\psi \varphi_{\psi,h}. \quad (4.3)$$

Let $\{H_i\}$ be an orthonormal basis of \mathfrak{a} . To each $\alpha \in \Sigma$, let $H_\alpha \in \mathfrak{a}$ be the unique vector such that $\alpha(x) = B(x, H_\alpha) = \langle x, H_\alpha \rangle$ for all $x \in \mathfrak{a}$, and choose a unit root vector $E_\alpha \in \mathfrak{n}$ so that $[x, E_\alpha] = \alpha(x)E_\alpha$ for all $x \in \mathfrak{a}$. We may write

$$\mathcal{C} = \sum_i H_i^2 + \sum_{\alpha \in \Sigma^+} (E_\alpha E_{-\alpha} + E_{-\alpha} E_\alpha) + J,$$

where $J \in \mathcal{U}(\mathfrak{m}_{\mathbb{C}})$ (cf. [25, Proposition 5.28]). Now using $E_{-\alpha} E_\alpha = E_\alpha E_{-\alpha} - H_\alpha$ gives

$$\mathcal{C} = \sum_i H_i^2 - \sum_{\alpha \in \Sigma^+} H_\alpha + \sum_{\alpha \in \Sigma^+} 2E_\alpha E_{-\alpha} + J.$$

As in the proof of Lemma 3.4, $[J\varphi_{\psi,h}](e) = 0$, and $[E_\alpha E_{-\alpha} \varphi_{\psi,h}](e) = 0$. Applying $-\mathcal{C}$ to $\varphi_{\psi,h}$ gives

$$\begin{aligned} -\mathcal{C}\varphi_{\psi,h} &= - \left(\sum_i \psi(H_i)^2 - \sum_{\alpha \in \Sigma^+} \psi(H_\alpha) \right) \varphi_{\psi,h} \\ &= -(\|\psi\|^2 - 2\langle \rho, \psi \rangle) \varphi_{\psi,h} \\ &= (\|\rho\|^2 - \|\psi - \rho\|^2) \varphi_{\psi,h}, \end{aligned}$$

proving (4.3). Let f be a positive λ -harmonic function on X , which we consider as a K -invariant function on G . By [27, Theorem 2], f is of the form: for any $g \in G$,

$$f(g) = \int_{K/M \times \{\psi \geq \rho : \lambda_\psi = \lambda\}} \varphi_{\psi,k}(g) d\mu([k], \psi)$$

for some Borel measure μ on $K/M \times \{\psi \geq \rho : \lambda_\psi = \lambda\}$. By (4.3), this implies (2). \square

Corollary 4.3. *For any Zariski dense discrete subgroup $\Gamma < G$,*

$$\sup\{\lambda_\psi : \psi \in D_\Gamma^*\} \leq \lambda_0.$$

Proof. If Γ is cocompact in G , then $\psi_\Gamma = 2\rho$ and hence $D_\Gamma^* = \{2\rho\}$. Since $\lambda_0 = 0 = \lambda_{2\rho}$ in this case, the claim follows. In general, it follows from Theorem 2.3 and Proposition 3.7 that for any $\psi \in D_\Gamma^*$, there exists a positive joint eigenfunction on $\Gamma \backslash X$ with character $\chi_{\psi-\rho}$. Hence the claim for the case when Γ is not cocompact in G follows from Theorem 4.1 and Proposition 4.2. \square

5. GROUPS OF THE SECOND KIND AND POSITIVE JOINT EIGENFUNCTIONS

When G has rank one (in which case the Furstenberg boundary is same as the geometric boundary of X), a discrete subgroup $\Gamma < G$ is said to be of the second kind if $\Lambda \neq \mathcal{F}$. We extend this definition to higher rank groups as follows:

Definition 5.1. A discrete subgroup $\Gamma < G$ is of *the second kind* if there exists $\xi \in \mathcal{F}$ which is in general position with all points of Λ , i.e., $(\xi, \Lambda) \subset \mathcal{F}^{(2)}$.

Theorem 4.1 provides a positive λ -harmonic function for any $\lambda \leq \lambda_0$, when $\Gamma \backslash X$ is not compact. The following theorem can be viewed as a higher rank strengthening of this result.

Theorem 5.2. *Let $\Gamma < G$ be of the second kind with $\mathcal{L} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$. For any $\psi \in D_\Gamma$, there exists a positive joint eigenfunction on $\Gamma \backslash X$ with character $\chi_{\psi-\rho}$.*

By Proposition 3.7, we get the following immediate corollary:

Corollary 5.3. *Let $\Gamma < G$ be of the second kind with $\mathcal{L} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$. Then for any $\psi \geq \max(\psi_\Gamma, \rho)$, there exists a (Γ, ψ) -conformal density.*

Remark 5.4. (1) Let $\Gamma_0 < G$ be an Anosov subgroup. Then any Anosov subgroup $\Gamma < \Gamma_0$ with $\Lambda_{\Gamma_0} \neq \Lambda_\Gamma$ is of the second kind. To see this, choose any $\xi \in \Lambda_{\Gamma_0} - \Lambda_\Gamma$, and note that $(\Lambda_\Gamma, \xi) \subset \mathcal{F}^{(2)}$, since any two distinct points of Λ_{Γ_0} are in general position by the Anosov assumption on Γ_0 .

- (2) If $\Lambda \subset gNw_0P$ for some $g \in G$, then $(\Lambda, g^+) \subset \mathcal{F}^{(2)}$. One can construct many Schottky groups with $\Lambda \subset Nw_0P$, which would then be of the second kind.
- (3) Let $G = \prod_{i=1}^k G_i$ be a product of simple algebraic groups G_i of rank one. Then $\mathcal{F} = \prod_i \mathcal{F}_i$ where $\mathcal{F}_i = G_i/P_i$. Let $\pi_i : \mathcal{F} \rightarrow \mathcal{F}_i$ denote the canonical projection. Then any discrete subgroup $\Gamma < G$ such that $\pi_i(\Lambda) \neq \mathcal{F}_i$ for all i is of the second kind. To see this, it suffices to note that $(\Lambda, \xi) \subset \mathcal{F}^{(2)}$ for any $\xi = (\xi_i)_i \in \mathcal{F}$ with $\xi_i \in \mathcal{F}_i - \pi_i(\Lambda)$.
- (4) The well-known properties of the limit set of a Hitchin subgroup of $\text{PSL}_d(\mathbb{R})$ imply that Hitchin groups are not of the second kind for any even $d \geq 4$ or $d = 3$; we thank Canary and Labourie for communicating this with us.

For $q \in X$ and $r > 0$, we set

$$B(q, r) = \{x \in X : d(x, q) \leq r\}.$$

For $p = g(o) \in X$, the shadow of the ball $B(q, r)$ viewed from p is defined as

$$O_r(p, q) := \{(gk)^+ \in \mathcal{F} : k \in K, gk \text{int } A^+ o \cap B(q, r) \neq \emptyset\}.$$

Similarly, for $\xi \in \mathcal{F}$, the shadow of the ball $B(q, r)$ viewed from ξ is defined by

$$O_r(\xi, q) := \{h^+ \in \mathcal{F} : h \in G \text{ satisfies } h^- = \xi, ho \in B(q, r)\}.$$

We will use the following shadow lemma to prove Theorem 5.2:

Lemma 5.5. [30, Lemma 5.6, 5.7]

(1) *If a sequence $q_i \in X$ converges to $\eta \in \mathcal{F}$, then for any $q \in X$, $r > 0$ and $\varepsilon > 0$,*

$$O_{r-\varepsilon}(q_i, q) \subset O_r(\eta, q) \subset O_{r+\varepsilon}(q_i, q)$$

for all sufficiently large i .

(2) *There exists $\kappa > 0$ such that for any $g \in G$ and $r > 0$,*

$$\sup_{\xi \in O_r(g(o), o)} \|\beta_\xi(g(o), o) - \mu(g^{-1})\| \leq \kappa r.$$

Lemma 5.6. *If $\mathcal{L} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$, then the union $\Gamma(o) \cup \Lambda$ is compact in the topology given in Definition 2.2.*

Proof. The hypothesis implies that any sequence $\gamma_i \rightarrow \infty$ in Γ tends to ∞ regularly, and hence has a limit in \mathcal{F} . Moreover the limit belongs to Λ by its definition. \square

Lemma 5.7. *Suppose that $\mathcal{L} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$. If $\xi \in \mathcal{F}$ satisfies that $(\xi, \Lambda) \subset \mathcal{F}^{(2)}$, then there exists $R > 0$ such that*

$$\xi \in \bigcap_{\gamma \in \Gamma} O_R(\gamma(o), o).$$

Proof. We first claim that $\xi \in \bigcap_{\eta \in \Lambda} O_R(\eta, o)$ for some $R > 0$. Note that $\lim_{R \rightarrow \infty} O_R(\eta, o) = \{z \in \mathcal{F} : (z, \eta) \in \mathcal{F}^{(2)}\}$. Hence for each $\eta \in \Lambda$, we have

$$R_\eta := \inf\{R + 1 : \xi \in O_R(\eta, o)\} < \infty.$$

It suffices to show that $R := \sup_{\eta \in \Lambda} R_\eta < \infty$. Suppose not; then $R_{\eta_i} \rightarrow \infty$ for some sequence $\{\eta_i\} \subset \Lambda$. By passing to a subsequence if necessary, we may assume that the η_i converge to some η . From this it follows that $O_{R_{\eta_i}+1}(\eta, o) \subset O_{R_{\eta_i}+2}(\eta_i, o)$ for all sufficiently large i . Therefore $R_{\eta_i} \leq R_\eta + 3$, yielding a contradiction.

We now claim that $\xi \in \bigcap_{\gamma \in \Gamma} O_{R'}(\gamma(o), o)$ for some $R' > 0$. Suppose not; then there exist sequences $\gamma_i \rightarrow \infty$ in Γ and $R_i \rightarrow \infty$ such that $\xi \notin O_{R_i}(\gamma_i(o), o)$. By Lemma 5.6, after passing to a subsequence, we may assume that $\gamma_i(o)$ converges to some $\eta \in \Lambda$. By the first claim, we have $\xi \in O_R(\eta, o)$. By Lemma 5.5, we have $\xi \in O_R(\eta, o) \subset O_{R+1}(\gamma_i(o), o)$ for all sufficiently large i . This is a contradiction, since for i large enough so that $R_i > R + 1$, we have $\xi \notin O_{R+1}(\gamma_i(o), o)$. This proves the claim. \square

As an immediate corollary of Lemmas 5.5 and 5.7, we obtain:

Corollary 5.8. *If $\mathcal{L} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$ and $\xi \in \mathcal{F}$ satisfies that $(\xi, \Lambda) \subset \mathcal{F}^{(2)}$, then*

$$\sup_{\gamma \in \Gamma} \|\beta_\xi(\gamma^{-1}o, o) - \mu(\gamma)\| < \infty.$$

Proof of Theorem 5.2: If $\psi \in D_\Gamma^*$, this follows from Theorem 2.3. Hence we assume $\psi \in D_\Gamma - D_\Gamma^*$; this implies that

$$\sum_{\gamma \in \Gamma} e^{-\psi(\mu(\gamma))} < \infty \quad (5.1)$$

by [40, Lem. III. 1.3]. As Γ is of the second kind, there exists $\xi \in \mathcal{F}$ such that $(\xi, \eta) \in \mathcal{F}^{(2)}$ for all $\eta \in \Lambda$. By Corollary 5.8, we have $\sup_{\gamma \in \Gamma} \|\beta_\xi(\gamma^{-1}o, o) - \mu(\gamma)\| < \infty$. Therefore (5.1) implies that

$$\sum_{\gamma \in \Gamma} e^{-\psi(\beta_\xi(\gamma^{-1}o, o))} < \infty. \quad (5.2)$$

For any fixed $x \in X$, we have $\beta_\xi(\gamma^{-1}x, o) = \beta_\xi(\gamma^{-1}o, o) + \beta_{\gamma\xi}(x, o)$ and $\|\beta_{\gamma\xi}(x, o)\| \leq d(x, o)$. Hence $e^{-\psi(\beta_\xi(\gamma^{-1}o, o))} \asymp e^{-\psi(\mu(\gamma))}$ with implied constant uniform for all $\gamma \in \Gamma$.

Therefore, by (5.1) the following function $F_{\psi, \xi}$ on X is well-defined:

$$F_{\psi, \xi}(x) := \sum_{\gamma \in \Gamma} e^{-\psi(\beta_\xi(\gamma^{-1}x, o))} \quad \text{for } x \in X. \quad (5.3)$$

If we write $\xi = [k_0] \in K/M = \mathcal{F}$, then for any $g \in G$,

$$\beta_\xi(\gamma^{-1}go, o) = \beta_M(k_0^{-1}\gamma^{-1}go, o) = H(g^{-1}\gamma k_0)$$

and hence $e^{-\psi(\beta_\xi(\gamma^{-1}go, o))} = \varphi_{\psi, \gamma k_0}(g)$. Therefore

$$F_{\psi, \xi} = \sum_{\gamma \in \Gamma} \varphi_{\psi, \gamma k_0}.$$

It now follows from Lemma 2.2 that $F_{\psi, \xi}$ is a positive Γ -invariant joint eigenfunction on X with eigenvalue $\chi_{\psi-\rho}$. This finishes the proof.

Remark 5.9. In the above proof, for any $\psi \in D_\Gamma - D_\Gamma^*$ and any $\xi \in \mathcal{F}$ with $(\Lambda, \xi) \subset \mathcal{F}^{(2)}$, we have constructed a positive joint eigenfunction $F_{\psi, \xi}$ on $\Gamma \backslash X$ of eigenvalue $\chi_{\psi-\rho}$.

Hence we get the following strengthened version of Corollary 4.3:

Corollary 5.10. *If $\Gamma < G$ is of the second kind with $\mathcal{L} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$, then*

$$\sup\{\lambda_\psi : \psi \in D_\Gamma\} \leq \lambda_0. \quad (5.4)$$

If $\Gamma < \text{SO}^\circ(n, 1)$ is a discrete subgroup with $\Lambda \neq \partial \mathbb{H}^n$, we have equality in (5.4), as was proved by Sullivan [45, Theorem 2.17].

6. THE L^2 -SPECTRUM AND UNIQUENESS

Let Γ be a torsion-free discrete subgroup of a connected semisimple real algebraic group G . The space $L^2(\Gamma \backslash X)$ consists of square-integrable functions together with the inner product $\langle f_1, f_2 \rangle = \int_{\Gamma \backslash X} f_1 \bar{f}_2 d\text{vol}$.

Let $W^1(\Gamma \backslash X) \subset L^2(\Gamma \backslash X)$ denote the closure of $C_c^\infty(\Gamma \backslash X)$ with respect to the norm $\|\cdot\|_{W^1}$ induced by the inner product

$$\langle f_1, f_2 \rangle_{W^1} := \int_{\Gamma \backslash X} f_1 \bar{f}_2 d\text{vol} + \int_{\Gamma \backslash X} \langle \text{grad } f_1, \text{grad } f_2 \rangle d\text{vol}$$

for any $f_1, f_2 \in W^1(\Gamma \backslash X)$.

As $\Gamma \backslash X$ is complete, there exists a unique self-adjoint operator on the space $W^1(\Gamma \backslash X)$ extending the Laplacian Δ on $C_c^\infty(\Gamma \backslash X)$, which we also denote by Δ . The L^2 -spectrum of $-\Delta$, which we denote by

$$\sigma(\Gamma \backslash X),$$

is the set of all $\lambda \in \mathbb{C}$ such that $\Delta + \lambda$ does not have a bounded inverse $(\Delta + \lambda)^{-1} : L^2(\Gamma \backslash X) \rightarrow W^1(\Gamma \backslash X)$. The self-adjointness of Δ and the fact that $\langle -\Delta f, f \rangle = \int_X \|\text{grad } f\|^2 d\text{vol}$ for all $f \in C_c^\infty(\Gamma \backslash X)$ imply $\sigma(\Gamma \backslash X) \subset [0, \infty)$.

We will be using Weyl's criterion to determine $\sigma(\Gamma \backslash X)$:

Theorem 6.1. (cf. [48, Lemma 2.17]) *For $\lambda \in \mathbb{R}$, we have $\lambda \in \sigma(\Gamma \backslash X)$ if and only if there exists a sequence of unit vectors $F_n \in W^1(\Gamma \backslash X)$ such that*

$$\lim_{n \rightarrow \infty} \|(\Delta + \lambda)F_n\| = 0.$$

The number $\lambda_0 = \lambda_0(\Gamma \backslash X)$ defined in (4.1) is the bottom of the L^2 -spectrum $\sigma(\Gamma \backslash X)$:

Theorem 6.2. [45, Theorem 2.1, 2.2] *We have*

$$\lambda_0 \in \sigma(\Gamma \backslash X) \subset [\lambda_0, \infty).$$

Using Harish-Chandra's Plancherel formula, we can identify $\lambda_0(X)$ and $\sigma(X)$ for the symmetric space $X = G/K$:

Proposition 6.3. *We have $\lambda_0(X) = \|\rho\|^2$ and $\sigma(X) = [\|\rho\|^2, \infty)$.*

Proof. It is shown in [22] that there are no positive Laplace eigenfunctions on X with eigenvalue strictly bigger than $\|\rho\|^2$; hence the inequality $\lambda_0(X) \leq \|\rho\|^2$ follows from Theorem 4.1 for $\Gamma = \{e\}$. On the other hand, as seen in the proof of (1), $\varphi_{\rho, h}$ is a positive $\|\rho\|^2$ -harmonic function (for any $h \in G$), hence $\lambda_0(X) = \|\rho\|^2$ by Theorem 4.1. We now deduce the second claim $\sigma(X) = [\|\rho\|^2, \infty)$ from Harish-Chandra's Plancherel theorem (cf. e.g. [43]). For $\psi \in \mathfrak{a}^*$, define $\Phi_\psi \in C^\infty(K \backslash G/K)$ by

$$\Phi_\psi(g) = \int_K \varphi_{\rho+i\psi, k}(g) dk.$$

where $\varphi_{\rho+i\psi,k}(g) = e^{-(\rho+i\psi)(H(g^{-1}k))}$.

Then by the same computation as (4.3), we have

$$-\mathcal{C}\Phi_\psi = -\Delta\Phi_\psi = (\|\rho\|^2 + \|\psi\|^2)\Phi_\psi.$$

Given any $f \in C_c^\infty(\mathfrak{a}^*)$, we can define a function $F \in L^2(X)$ by the formula

$$F(g) = \int_{\mathfrak{a}^*} f(\psi)\Phi_\psi(g) \frac{d\psi}{|\mathbf{c}(\psi)|^2};$$

here $d\psi$ denotes the Lebesgue measure on \mathfrak{a}^* and $\mathbf{c}(\psi)$ denotes the Harish-Chandra \mathbf{c} -function. The Plancherel formula says

$$\|F\|_{L^2(X)}^2 = \int_{\mathfrak{a}^*} |f(\psi)|^2 \frac{d\psi}{|\mathbf{c}(\psi)|^2}$$

(see [43]). Let $\lambda \in [\|\rho\|^2, \infty)$ be any number. Choose $\psi_0 \in \mathfrak{a}^*$ so that $\lambda = \|\rho\|^2 + \|\psi_0\|^2$. We then choose a sequence of non-negative functions $\{f_n\} \subset C_c^\infty(\mathfrak{a}^*)$ with $\text{supp } f_n \subset B_{1/n}(\psi_0)$ and $\|F_n\|_{L^2(X)} = 1$.

Then

$$\begin{aligned} (\Delta + \lambda)F_n &= \int_{\mathfrak{a}^*} f_n(\psi)(\Delta + \lambda)\Phi_\psi(g) \frac{d\psi}{|\mathbf{c}(\psi)|^2} \\ &= \int_{\mathfrak{a}^*} f_n(\psi)(\lambda - \|\rho\|^2 - \|\psi\|^2)\Phi_\psi(g) \frac{d\psi}{|\mathbf{c}(\psi)|^2}. \end{aligned}$$

This gives

$$\begin{aligned} \|(\Delta + \lambda)F_n\|_{L^2(X)}^2 &= \int_{\mathfrak{a}^*} |(\lambda - \|\rho\|^2 - \|\psi\|^2)f_n(\psi)|^2 \frac{d\psi}{|\mathbf{c}(\psi)|^2} \\ &\leq \max_{\psi \in B_{1/n}(\psi_0)} \|\psi_0\|^2 - \|\psi\|^2. \end{aligned}$$

Consequently,

$$\lim_{n \rightarrow \infty} \|(\Delta + \lambda)F_n\|_{L^2(X)} = 0.$$

By Weyl's criterion (Theorem 6.1), this implies that $\lambda \in \sigma(X)$. This proves the claim. \square

Theorem 6.4. *If $L^2(\Gamma \backslash G)$ is tempered, then*

$$\lambda_0(\Gamma \backslash X) = \|\rho\|^2.$$

Proof. Note that $\lambda_0 = \lambda_0(\Gamma \backslash X) \leq \lambda_0(X) = \|\rho\|^2$ by Proposition 6.3. Assume that $\lambda_0 < \|\rho\|^2$. By Theorem 6.1, we can then find a K -invariant unit vector $f \in L^2(\Gamma \backslash G)_K$ such that

$$\|(\Delta - \lambda_0)f\| < \frac{\|\rho\|^2 - \lambda_0}{2}.$$

This gives

$$\|\mathcal{C}f\| = \|\Delta f\| \leq \|(\Delta - \lambda_0)f\| + \lambda_0 < \frac{\|\rho\|^2 + \lambda_0}{2} < \|\rho\|^2.$$

On the other hand, consider the direct integral representation of $L^2(\Gamma \backslash G) = \int_{\mathbb{Z}}^{\oplus} (\pi_{\zeta}, \mathcal{H}_{\zeta}) d\mu(\zeta)$ into irreducible unitary representations of G which are tempered, by the hypothesis on the temperedness of $L^2(\Gamma \backslash G)$. Hence

$$\|\mathcal{C}f\|^2 = \int_{\mathbb{Z}} \|d\pi_{\zeta}(\mathcal{C})f_{\zeta}\|_{\zeta}^2 d\mu(\zeta) \geq \left(\min_{\pi \text{ spherical tempered}} |d\pi(\mathcal{C})|^2 \right),$$

where $d\pi$ denotes the derived representation of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ induced by π . By Schur's lemma, there exists a character χ_{π} of $\mathcal{Z}(\mathfrak{g}_{\mathbb{C}})$ such that $d\pi(Z) = \chi_{\pi}(Z)$ for all $Z \in \mathcal{Z}(\mathfrak{g}_{\mathbb{C}})$. Moreover, for any spherical π , there exists $\psi_{\pi} \in \mathfrak{a}_{\mathbb{C}}^*$ such that $\chi_{\pi} = \chi_{\psi_{\pi}}$ (cf. (3.1)). Now, by Harish-Chandra's Plancherel formula (cf. e.g. [20]), for any tempered spherical representation, we have

$$\psi_{\pi} = \rho + i \operatorname{Im}(\psi_{\pi}),$$

where $\operatorname{Im}(\psi_{\pi}) \in \mathfrak{a}^*$. As in the proof of Proposition 6.3, we then obtain

$$\chi_{\pi}(-\mathcal{C}) = \|\rho\|^2 + \|\operatorname{Im}(\psi_{\pi})\|^2.$$

Thus for any spherical tempered representation (π, \mathcal{H}) , we have $d\pi(\mathcal{C}) \in \sigma(X)$ and hence, by Proposition 6.3,

$$\min_{\pi \text{ spherical tempered}} |d\pi(\mathcal{C})| \geq \|\rho\|^2,$$

giving a contradiction. \square

Theorem 6.5. [45, Theorem 2.8 and Corollary 2.9]

- (1) *Any positive Laplace eigenfunction in $L^2(\Gamma \backslash X)$ is λ_0 -harmonic.*
- (2) *If there exists a λ_0 -harmonic function in $L^2(\Gamma \backslash X)$, then the space of λ_0 -harmonic functions in $\Gamma \backslash X$ is one-dimensional and generated by a positive function.*

Proof. Sullivan's proof in [45] uses the heat operator and superharmonic functions. We provide a more direct proof here.

Note that if $f \in L^2(\Gamma \backslash X) \cap C^{\infty}(\Gamma \backslash X)$ is a real-valued λ -harmonic function, then $f \in W^1(\Gamma \backslash X)$, since

$$\int_{\Gamma \backslash X} \|\operatorname{grad} f\|^2 d\operatorname{vol} = - \int_{\Gamma \backslash X} f \Delta f d\operatorname{vol} = \lambda \int_{\Gamma \backslash X} f^2 d\operatorname{vol}.$$

The key fact for us is that λ_0 may also be expressed as an infimum over real-valued functions in $W^1(\Gamma \backslash X)$; for $f \neq 0$ in $W^1(\Gamma \backslash X)$, define $R(f)$ by

$$R(f) = \frac{\|f\|_{W^1}^2}{\|f\|^2} - 1 \geq 0$$

where $\|\cdot\|$ denotes the $L^2(\Gamma \backslash X)$ norm. For any $f \neq 0 \in W^1(\Gamma \backslash X)$, and all φ with $\|f - \varphi\|_{W^1}$ small enough, we have

$$\frac{\|\varphi\|_{W^1} - \|f - \varphi\|_{W^1}}{\|\varphi\| + \|f - \varphi\|_{W^1}} - 1 \leq R(f) \leq \frac{\|\varphi\|_{W^1} + \|f - \varphi\|_{W^1}}{\|\varphi\| - \|f - \varphi\|_{W^1}} - 1,$$

i.e. $f \mapsto R(f)$ is continuous at each $f \neq 0 \in W^1(\Gamma \setminus X)$. The density of $C_c^\infty(\Gamma \setminus X)$ in $W^1(\Gamma \setminus X)$ then gives

$$\lambda_0 = \inf_{\substack{f \in C_c^\infty(\Gamma \setminus X) \\ f \neq 0}} R(f) = \inf_{\substack{f \in W^1(\Gamma \setminus X) \\ f \neq 0}} R(f).$$

Now suppose that $\phi \in L^2(\Gamma \setminus X)$ is a positive λ -harmonic function; so $\phi \in W^1(\Gamma \setminus X)$. We claim that $\lambda = \lambda_0$. By Green's identity, we have

$$\lambda_0 \leq R(\phi) = \frac{\int_{\Gamma \setminus X} \|\operatorname{grad} \phi\|^2 d\operatorname{vol}}{\int_{\Gamma \setminus X} |\phi|^2 d\operatorname{vol}} = \frac{\int_{\Gamma \setminus X} \phi(-\Delta\phi) d\operatorname{vol}}{\int_{\Gamma \setminus X} |\phi|^2 d\operatorname{vol}} = \lambda$$

(cf. Proposition 4.2). On the other hand, since $\phi > 0$, we have that for any $\varphi \in C_c^\infty(\Gamma \setminus X)$,

$$\frac{\int_{\Gamma \setminus X} \|\operatorname{grad} \varphi\|^2 d\operatorname{vol}}{\int_{\Gamma \setminus X} |\varphi|^2 d\operatorname{vol}} = \frac{\int_{\Gamma \setminus X} \|\operatorname{grad}(\phi \cdot \frac{\varphi}{\phi})\|^2 d\operatorname{vol}}{\int_{\Gamma \setminus X} |\varphi|^2 d\operatorname{vol}}.$$

By Barta's identity [1],

$$\int_{\Gamma \setminus X} \|\operatorname{grad}(\phi \cdot \frac{\varphi}{\phi})\|^2 d\operatorname{vol} = \int_{\Gamma \setminus X} \phi^2 \|\operatorname{grad} \frac{\varphi}{\phi}\|^2 d\operatorname{vol} - \int_{\Gamma \setminus X} \left(\frac{\varphi}{\phi}\right)^2 \phi \Delta\phi d\operatorname{vol},$$

so

$$\int_{\Gamma \setminus X} \|\operatorname{grad} \varphi\|^2 d\operatorname{vol} \geq \int_{\Gamma \setminus X} \left(\frac{\varphi}{\phi}\right)^2 \phi(-\Delta\phi) d\operatorname{vol} = \lambda \int_{\Gamma \setminus X} \varphi^2 d\operatorname{vol},$$

i.e.

$$\lambda \leq \frac{\int_{\Gamma \setminus X} \|\operatorname{grad} \varphi\|^2 d\operatorname{vol}}{\int_{\Gamma \setminus X} |\varphi|^2 d\operatorname{vol}},$$

showing that $\lambda_0 \geq \lambda$. Hence $\lambda = \lambda_0$.

In order to prove (2), we first claim that $f \in W^1(\Gamma \setminus X)$ satisfies $-\Delta f = \lambda_0 f$ if and only if $R(f) = \lambda_0$. Suppose that $R(f) = \lambda_0$. We will then show that for any $\varphi \in C_c^\infty(\Gamma \setminus X)$, we have

$$\langle f, -\Delta\varphi \rangle = \lambda_0 \langle f, \varphi \rangle; \quad (6.1)$$

this implies that f is λ_0 -harmonic. Let $\varphi \in C_c^\infty(\Gamma \setminus X)$. Since $R(f) = \lambda_0$, f minimizes R . So for any $\varphi \in C_c^\infty(\Gamma \setminus X)$, the function $F : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ defined by $F(x) = R(f + x\varphi)$ has a local minimum at $x = 0$, hence $F'(0) = 0$. Now computing $F'(0)$ gives

$$F'(0) = \frac{2\langle f, \varphi \rangle_{W^1} \|f\|^2 - 2\langle f, \varphi \rangle \|f\|_{W^1}^2}{\|f\|^4} = 0.$$

From $R(f) = \lambda_0$, we obtain $\|f\|_{W^1}^2 = (\lambda_0 + 1)\|f\|^2$, which, when entered into the identity above, gives

$$\langle f, \varphi \rangle_{W^1} = (\lambda_0 + 1)\langle f, \varphi \rangle. \quad (6.2)$$

Letting $\{f_i\}_{i \in \mathbb{N}} \subset C_c^\infty(\Gamma \setminus X)$ be a sequence converging to f in $W^1(\Gamma \setminus X)$, Green's identity again gives

$$\begin{aligned} \langle f, \varphi \rangle_{W^1} &= \lim_{i \rightarrow \infty} \langle f_i, \varphi \rangle_{W^1} = \lim_{i \rightarrow \infty} \int_{\Gamma \setminus X} f_i \varphi + \langle \operatorname{grad} f_i, \operatorname{grad} \varphi \rangle d\operatorname{vol} \\ &= \lim_{i \rightarrow \infty} \int_{\Gamma \setminus X} f_i \varphi + f_i(-\Delta \varphi) d\operatorname{vol} = \langle f, \varphi \rangle + \langle f, -\Delta \varphi \rangle. \end{aligned} \quad (6.3)$$

Combined with (6.2), this gives $\langle f, -\Delta \varphi \rangle = \lambda_0 \langle f, \varphi \rangle$ as in (6.1).

Conversely, if $f \in W^1(\Gamma \setminus X)$ satisfies $-\Delta f = \lambda_0 f$, then for any $\varphi \in C_c^\infty(\Gamma \setminus X)$, we have (as in (6.3))

$$\langle f, \varphi \rangle_{W^1} = \langle f, \varphi \rangle + \langle f, -\Delta \varphi \rangle = (\lambda_0 + 1) \langle f, \varphi \rangle,$$

hence

$$\|f\|_{W^1}^2 = \sup_{\varphi \in C_c^\infty(\Gamma \setminus X)} \langle f, \varphi \rangle_{W^1} = \sup_{\varphi \in C_c^\infty(\Gamma \setminus X)} (\lambda_0 + 1) \langle f, \varphi \rangle = (\lambda_0 + 1) \|f\|^2,$$

giving $R(f) = \lambda_0$. This proves the claim.

Let $f \in W^1(\Gamma \setminus X) \cap C^\infty(\Gamma \setminus X)$ now be a real-valued λ_0 -harmonic function. Then $|f| \in W^1(\Gamma \setminus X)$ and $R(|f|) = \lambda_0$. As shown above, $|f|$ is also a λ_0 -harmonic function. Hence either f is a constant multiple of $|f|$ or f must change sign at some point x_0 , hence $|f(x)| \geq |f(x_0)| = 0$ for all $x \in \Gamma \setminus X$. However, since $\Delta|f| = -\lambda_0|f| \leq 0$, the strong minimum principle (cf. e.g. [37, Theorem 66, p. 280]) gives that if $|f|$ attains its infimum, then $|f|$ is in fact constant (in this case equal to zero). We therefore conclude that any λ_0 -harmonic function in $L^2(\Gamma \setminus X)$ is a constant multiple of a positive function. This then implies that the space of λ_0 -harmonic functions must be one-dimensional as two positive functions cannot be orthogonal to each other. \square

The uniqueness in the above theorem has the following implications for joint eigenfunctions:

Corollary 6.6. (1) *There exists at most one positive joint eigenfunction in $L^2(\Gamma \setminus X)$ up to a constant multiple.*
 (2) *If there exists a positive joint eigenfunction in $L^2(\Gamma \setminus X)$ with character $\chi_{\psi-\rho}$, $\psi \in \mathfrak{a}^*$, then*

$$\lambda_0 = \lambda_\psi.$$

(3) *There exists a positive Laplace eigenfunction in $L^2(\Gamma \setminus X)$ if and only if there exists a positive joint eigenfunction in $L^2(\Gamma \setminus X)$ of character $\chi_{\psi-\rho}$ with $\lambda_\psi = \lambda_0$.*

Proof. We only need to verify the third claim. Suppose that $\phi \in L^2(\Gamma \setminus X)$ is a positive Laplace eigenfunction. Via the identification $L^2(\Gamma \setminus X) = L^2(\Gamma \setminus G)_K$, we may consider $\phi \in L^2(\Gamma \setminus G)_K$ as a positive \mathcal{C} -eigenfunction for the Casimir operator \mathcal{C} . By Theorem 6.5, $\mathcal{C}\phi = -\lambda_0\phi$. Let $D \in \mathcal{Z}(\mathfrak{g}_\mathbb{C})$. Then $\mathcal{C} \circ D\phi = D \circ \mathcal{C}\phi = -\lambda_0 D\phi$. By the uniqueness in Theorem 6.5, it follows that $D\phi$

is a constant multiple of ϕ ; and hence ϕ is an eigenfunction for D as well. Therefore ϕ is a joint eigenfunction. \square

Spherical unitary representations contained in $L^2(\Gamma \backslash G)$. We let $C_c(G//K)$ denote the Hecke algebra of G , i.e.

$$C_c(G//K) = \{f \in C_c(G) : f(k_1 g k_2) = f(g) \quad \text{for all } g \in G, k_1, k_2 \in K\}.$$

Each element of $C_c(G//K)$ acts on $C(G)$ via right convolution $*$.

Lemma 6.7. *A positive K -invariant joint eigenfunction on G is an eigenfunction for the action of the Hecke algebra. More precisely, if*

$$\phi(g) = \int_{\mathcal{F}} \varphi_{\psi, k}(g) d\nu_o([k]), \quad g \in G, \quad (6.4)$$

for some $\psi \in \mathfrak{a}^*$ and a (Γ, ψ) -conformal measure ν_o on $\mathcal{F} = K/M$, then for all $f \in C_c(G//K)$,

$$(\phi * f)(g) = \left(\int_G f(h) e^{-\psi(H(h))} dh \right) \phi(g).$$

Proof. Given $f \in C_c(G//K)$, we have

$$\begin{aligned} (\phi * f)(g) &= \int_G \phi(gh^{-1}) f(h) dh = \int_G \int_{\mathcal{F}} \varphi_{\psi, k}(gh^{-1}) f(h) d\nu_o([k]) dh \\ &= \int_{\mathcal{F}} \int_G f(h) e^{-\psi(H(hg^{-1}k))} dh d\nu_o([k]). \end{aligned}$$

Now using $H(hg^{-1}k) = H(h\kappa(g^{-1}k)) + H(g^{-1}k)$ and then the change of variables $h' = h\kappa(g^{-1}k)$ gives

$$\begin{aligned} (\phi * f)(g) &= \int_{\mathcal{F}} \left(\int_G f(h\kappa(g^{-1}k)^{-1}) e^{-\psi(H(h))} dh \right) e^{-\psi(H(g^{-1}k))} d\nu_o([k]) \\ &= \int_{\mathcal{F}} \left(\int_G f(h) e^{-\psi(H(h))} dh \right) e^{-\psi(H(g^{-1}k))} d\nu_o([k]) \\ &= \left(\int_G f(h) e^{-\psi(H(h))} dh \right) \phi(g), \end{aligned}$$

since $f \in C(G//K)$, and is thus right K -invariant. In total, we have shown that ϕ is an eigenfunction of the f -action, with eigenvalue $\int_G f(h) e^{-\psi(H(h))} dh$. \square

Theorem 6.8. *If $\phi \in L^2(\Gamma \backslash G)_K$ is a positive Laplace eigenfunction of norm one, there exists a unique irreducible spherical unitary subrepresentation (π, \mathcal{H}_ϕ) of $L^2(\Gamma \backslash G)$, and ϕ is the unique K -invariant unit vector in \mathcal{H}_ϕ .*

Proof. By Corollary 6.6, ϕ is given by (6.4) for some $\psi \in \mathfrak{a}^*$. Define $\Phi : G \rightarrow \mathbb{C}$ by

$$\Phi(g) := \langle g.\phi, \phi \rangle$$

for all $g \in G$ where the g action on $L^2(\Gamma \backslash G)$ is via the translation action of G on $\Gamma \backslash G$ from the right. Given $f \in C_c(G//K)$, we then have, using Lemma 6.7,

$$\begin{aligned} (\Phi * f)(g) &= \int_G \Phi(gh^{-1})f(h) dh = \int_G \langle (gh^{-1}).\phi, \phi \rangle f(h) dh \\ &= \int_G \langle f(h)h^{-1}.\phi, g^{-1}.\phi \rangle dh = \langle \phi * f, g^{-1}.\phi \rangle \\ &= \left(\int_G f(h)e^{-\psi(H(h))} dh \right) \Phi(g), \end{aligned}$$

i.e. Φ is also a $C_c(G//K)$ -eigenfunction. Also note that $\Phi(e) = 1$, and since ϕ is right K -invariant, Φ is *bi*- K -invariant. Moreover, being the matrix coefficient of a unitary representation, Φ is also positive definite, i.e., for any $g_1, \dots, g_n \in G$ and $z_1, \dots, z_n \in \mathbb{C}$,

$$\sum_{1 \leq i, j \leq n} z_i \bar{z}_j \Phi(g_j^{-1} g_i) \geq 0.$$

We have thus shown that Φ is a positive definite *spherical* function. Letting \mathcal{H}_ϕ denote the closure of $\text{span}\{g.\phi : g \in G\}$ in $L^2(\Gamma \backslash G)$, by [28, Chapter IV§5, Corollary of Theorem 9], \mathcal{H}_ϕ is an irreducible (spherical) unitary sub-representation of the quasi-regular representation $L^2(\Gamma \backslash G)$. The uniqueness follows from Corollary 6.6. \square

We require the following lemma in the proof of Theorem 6.10:

Lemma 6.9. *Let $\psi \geq \rho$ and $\psi \notin \mathbb{R}\rho$. Denote by ψ' be the element of the line $\mathbb{R}\psi$ closest to ρ . Then $\psi' \not\geq \rho$.*

Proof. Let $\phi := \psi - \rho$. Note that $\phi \geq 0$ on \mathfrak{a} by the hypothesis. Then

$$\psi' = \frac{\langle \psi, \rho \rangle}{\|\psi\|^2} \psi = \frac{\langle \rho + \phi, \rho \rangle}{\|\rho + \phi\|^2} \psi = \left(1 - \frac{\|\phi\|^2 + \langle \rho, \phi \rangle}{\|\rho + \phi\|^2} \right) \psi,$$

i.e. $\psi' = t\psi$ with $0 < t < 1$. Now, if $\psi' \geq \rho$, we could repeat the process with ψ' in place of ψ to find another, *different*, closest vector in $\mathbb{R}\psi$ to ρ , which is not possible. \square

Theorem 6.10. *Let $\Gamma < G$ be of the second kind with $\mathcal{L} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$. If there exists a λ_0 -harmonic function in $L^2(\Gamma \backslash X)$, then*

$$\lambda_0 = \lambda_\psi$$

for some $\psi \in D_\Gamma^* \cup \{\rho\}$.

Proof. Suppose that $\psi \in D_\Gamma \setminus (\{\rho\} \cup D_\Gamma^*)$ and that $\psi \geq \rho$. Assume that there exists a positive joint eigenfunction $\phi \in L^2(\Gamma \backslash X)$ with character $\chi_{\psi-\rho}$. By Corollary 6.6,

$$\lambda_0 = \lambda_\psi = \|\rho\|^2 - \|\psi - \rho\|^2. \tag{6.5}$$

Since ψ_Γ is concave, there exists $0 < c \leq 1$ such that $c\psi(u) = \psi_\Gamma(u)$ for some $u \in \mathcal{L}$. So $\psi_0 := c\psi \in D_\Gamma^*$. Since $\psi \notin D_\Gamma^*$, we have $0 < c < 1$. There exists a unique $s_0 \in \mathbb{R}$ such that

$$\|s_0\psi_0 - \rho\| = \min\{\|s\psi - \rho\| : s \in \mathbb{R}\}, \quad (6.6)$$

that is, $s_0\psi_0$ be the element on the line $\mathbb{R}\psi$ that is closest to ρ .

We claim that $s_0c \leq 1$; since $0 < c < 1$, this implies that $\max\{1, s_0\} < c^{-1}$. If $\psi \in \mathbb{R}\rho$, then $s_0\psi_0 = \rho$. Since $\psi_0 = c\psi$, we get $s_0c\psi = \rho$. By the hypothesis $\rho \leq \psi$, $s_0c \leq 1$. Now suppose $\psi \notin \mathbb{R}\rho$. Assume that $s_0c > 1$. Then $s_0\psi_0 = s_0c\psi > \psi$. Hence $s_0c\psi \in D_\Gamma$. By Corollary 5.10 and (6.5), we get

$$\|s_0c\psi - \rho\| \geq \|\psi - \rho\|.$$

By the choice of s_0 in (6.6), it follows that $\|s_0c\psi - \rho\| = \|\psi - \rho\|$. Since $s_0c\psi > \psi \geq \rho$, this yields a contradiction. Therefore the claim $s_0c \leq 1$ follows.

We now choose t so that $\max\{1, s_0\} < t < c^{-1}$. Since $t > 1$ and $\psi_0 \in D_\Gamma^*$, $t\psi_0 \in D_\Gamma$. Note also that $s \mapsto \lambda_{s\psi_0}$ is strictly decreasing on the interval $[s_0, \infty)$. Since $s_0 < t < c^{-1}$ and $c^{-1}\psi_0 = \psi$, we get

$$\lambda_0 = \lambda_\psi < \lambda_{t\psi_0}.$$

This contradicts Corollary 5.10. This implies the claim by Corollary 6.6. \square

If we use the norm on $\mathfrak{so}(n, 1)$ which endows the constant curvature -1 metric on \mathbb{H}^n , then for any non-elementary discrete subgroup $\Gamma < \mathrm{SO}^\circ(n, 1)$, $D_\Gamma^* = \{\delta\}$ and hence the above theorem says that if a λ_0 -harmonic function belongs to $L^2(\Gamma \backslash \mathbb{H}^n)$, then λ_0 must be given by either $\delta(n-1-\delta)$ or $\frac{1}{4}(n-1)^2$.

7. SMEARING ARGUMENT IN HIGHER RANK

Let Γ be a torsion-free discrete subgroup of a connected semisimple real algebraic group G . The goal of this section is to prove the following:

Theorem 7.1. *If $\mathcal{L} \neq \mathfrak{a}^+$, then no positive joint eigenfunction belongs to $L^2(\Gamma \backslash X)$.*

Combined with Corollary 6.6, we get the following corollary which implies Theorem 1.6(4) in higher rank.

Corollary 7.2. *If $\mathcal{L} \neq \mathfrak{a}^+$, there exists no positive Laplace eigenfunction in $L^2(\Gamma \backslash X)$. In particular, if $\mathrm{rank} G \geq 2$ and $\Gamma < G$ is Anosov, no positive Laplace eigenfunction belongs to $L^2(\Gamma \backslash X)$.*

The second part follows from the first by Theorem 2.5. Theorem 7.1 will be deduced from Theorem 7.4, the proof of which is based on the smearing argument of Thurston and Sullivan (see [46], [7] and also [47] for historical remarks and the origin of the name “smearing argument”). We also refer to [42, Theorem 3.1].

Definition 7.3 (Hopf parameterization). The homeomorphism $G/M \rightarrow \mathcal{F}^{(2)} \times \mathfrak{a}$ given by $gM \mapsto (g^+, g^-, b = \beta_{g^-}(e, g))$ is called the Hopf parameterization of G/M .

Fix a pair of linear forms $\psi_1, \psi_2 \in \mathfrak{a}^*$. For $x \in X$ and $(\xi, \eta) \in \mathcal{F}^{(2)}$, let

$$\phi_x(\xi, \eta) = e^{\psi_1(\beta_\xi(x, go)) + \psi_2(\beta_\eta(x, go))}, \quad (7.1)$$

where $g \in G$ is such that $g^+ = \xi$ and $g^- = \eta$.

Let $\nu = \{\nu_x : x \in X\}$ and $\bar{\nu} = \{\bar{\nu}_x : x \in X\}$ be respectively (Γ, ψ_1) and (Γ, ψ_2) -conformal densities on \mathcal{F} . Using the Hopf parametrization, we define the following locally finite Borel measure $\tilde{m}_{\nu, \bar{\nu}}$ on G/M : for $(\xi, \eta, v) \in \mathcal{F}^{(2)} \times \mathfrak{a}$,

$$d\tilde{m}_{\nu, \bar{\nu}}(\xi, \eta, v) = \phi_x(\xi, \eta) d\nu_x(\xi) d\bar{\nu}_x(\eta) dv \quad (7.2)$$

where dv is the Lebesgue measure on \mathfrak{a} and $x \in X$ is any element; it follows from the Γ -conformality of $\{\nu_x\}$ and $\{\bar{\nu}_x\}$ that this definition is independent of $x \in X$. The measure $\tilde{m}_{\nu, \bar{\nu}}$ is left Γ -invariant and right A -semi-invariant: for all $a \in A$,

$$a_* \tilde{m}_{\nu, \bar{\nu}} = e^{(-\psi_1 + \psi_2 \circ i)(\log a)} \tilde{m}_{\nu, \bar{\nu}}. \quad (7.3)$$

Note that $\psi_2 = \psi_1 \circ i$ if and only if $\tilde{m}_{\nu, \bar{\nu}}$ is A -invariant. We denote by $m_{\nu, \bar{\nu}}$ the M -invariant Borel measure on $\Gamma \backslash G$ induced by $\tilde{m}_{\nu, \bar{\nu}}$; this measure is called the (generalized) Bowen-Margulis-Sullivan measure associated to the pair $(\nu, \bar{\nu})$ [9].

Theorem 7.4 (Smearing theorem). *For any pair $(\nu, \bar{\nu})$ of Γ -conformal densities on \mathcal{F} , there exists $c > 0$ such that*

$$m_{\nu, \bar{\nu}}(\Gamma \backslash G) \leq c \int_{\text{one-neighborhood of } \text{supp } m_{\nu, \bar{\nu}}} E_\nu(x) E_{\bar{\nu}}(x) d\text{vol}(x).$$

Proof. Let $Z = G/K \times \mathcal{F}^{(2)}$. For any $(\xi, \eta) \in \mathcal{F}^{(2)}$, we write $[\xi, \eta] = gAo \subset X$ for any $g \in G$ such that $g^+ = \xi$ and $g^- = \eta$; $[\xi, \eta]$ is a maximal flat in X defined independently of the choice of $g \in G$. Let $\psi_1, \psi_2 \in \mathfrak{a}^*$ be linear forms such that ν and $\bar{\nu}$ are respectively (Γ, ψ_1) and (Γ, ψ_2) -conformal densities. Let ϕ_x be defined as in (7.1) for all $x \in X$. We also denote by $W_{\xi, \eta} \subset X$ the one neighborhood of $[\xi, \eta]$. Consider the following locally finite Borel measure α on Z defined as follows: for any $f \in C_c(Z)$,

$$\alpha(f) = \int_{(\xi, \eta) \in \mathcal{F}^{(2)}} \int_{z \in W_{\xi, \eta}} f(z, \xi, \eta) dz dm(\xi, \eta),$$

where dz is the G -invariant measure on X , and

$$dm(\xi, \eta) = \phi_x(\xi, \eta) d\nu_x(\xi) d\bar{\nu}_x(\eta)$$

(observe that this definition is independent of x).

Consider the natural diagonal action of Γ on Z . Since dz and dm are both left Γ -invariant, α is also left Γ -invariant and hence induces a measure on the quotient space $\Gamma \backslash Z$, which we also denote by α by abuse of notation.

Define the projection $\pi' : Z \rightarrow G/M$ as follows: for $(x, \xi, \eta) \in X \times \mathcal{F}^{(2)}$, choose $g \in G$ so that $g^+ = \xi$ and $g^- = \eta$. Then there exists a unique element $a \in A$ such that

$$d(x, gao) = d(x, gAo) = \inf_{b \in A} d(x, gbo);$$

this follows from [4, Proposition 2.4] since X is a CAT(0) space and $gA(o)$ is a convex complete subspace of X . In other words, the point gao is the orthogonal projection of x to the flat $[\xi, \eta] = gAo$. We then set

$$\pi'(x, \xi, \eta) = gAM \in G/M;$$

this is well-defined independent of the choice of $g \in G$. Noting that π' is Γ -equivariant, we denote by

$$\pi : \text{supp}(\alpha) \subset \Gamma \setminus Z \rightarrow \text{supp}(m_{\nu, \bar{\nu}}) \subset \Gamma \setminus G/M$$

the map induced by π' . Fixing $[ga] \in \Gamma \setminus G/M$, the fiber $\pi^{-1}[ga]$ is of the form $[(gaD_0, g^+, g^-)]$, where

$$D_0 = \{s \in X : d(s, o) \leq 1, \text{ the geodesic connecting } s \text{ and } o \text{ is orthogonal to } Ao \text{ at } o\}.$$

Noting that each fiber $\pi^{-1}(v)$, $v \in \text{supp } m_{\nu, \bar{\nu}}$, is isometric to D_0 , we have for any Borel subset $S \subset \text{supp } m_{\nu, \bar{\nu}}$,

$$\alpha(\pi^{-1}(S)) = \text{Vol}(D_0) \cdot m_{\nu, \bar{\nu}}(S); \quad (7.4)$$

the volume of D_0 being computed with respect to the volume form induced by the G -invariant measure on X . Consider now the map $p : \text{supp}(\alpha) \rightarrow \Gamma \setminus X$ defined by $p([(z, \xi, \eta)]) = [z]$ for any $(z, \xi, \eta) \in \text{supp}(\alpha)$. Let $F = \pi^{-1}(\text{supp } m_{\nu, \bar{\nu}}) \subset \text{supp}(\alpha)$. We write

$$\alpha(F) = \int_{\Gamma \setminus X} \alpha_x(p^{-1}(x) \cap F) dx,$$

where α_x is a conditional measure on the fiber $p^{-1}(x)$. We claim that there exists a constant $c > 0$ such that for any $x \in \Gamma \setminus X$,

$$\alpha_x(p^{-1}(x)) \leq c E_\nu(x) \cdot E_{\bar{\nu}}(x). \quad (7.5)$$

Since $p^{-1}(x) \cap F = \emptyset$ for x outside of the one neighborhood of $\text{supp}(m_{\nu, \bar{\nu}})$, this together with (7.4), implies that

$$\text{Vol}(D_0) \cdot |m_{\nu, \bar{\nu}}| = \alpha(F) \leq c \cdot \int_{\text{one neighborhood of } \text{supp}(m_{\nu, \bar{\nu}})} E_\nu(x) E_{\bar{\nu}}(x) dx$$

finishing the proof. Note that for any $h \in G$,

$$V_{ho} := \{(\xi, \eta) \in \mathcal{F}^{(2)} : [\xi, \eta] \cap B(ho, 1) \neq \emptyset\}$$

is a compact subset of $\mathcal{F}^{(2)}$; if $\{g_i\} \subset G$ and $\{a_i\} \subset A$ are sequences such that $d(g_i a_i o, ho) \leq 1$, then (by passing to a subsequence) we may assume that $g_i a_i$ converges to some $g_0 \in G$. This implies $(g_i^+, g_i^-) \rightarrow (g_0^+, g_0^-) \in \mathcal{F}^{(2)}$

as $i \rightarrow \infty$ and $d(g_0 o, ho) \leq 1$, from which the compactness of V_{ho} follows. It follows that

$$c := \sup\{\phi_o(\xi, \eta) : (\xi, \eta) \in V_o\} < \infty.$$

By the equivariance $\phi_{ho}(\xi, \eta) = \phi_o(h^{-1}\xi, h^{-1}\eta)$, we have for any $h \in G$,

$$\sup\{\phi_{ho}(\xi, \eta) : (\xi, \eta) \in V_{ho}\} = c.$$

Note that if $x = [ho] \in \Gamma \setminus X$ for $h \in G$, then

$$p^{-1}(x) = \{[(ho, \xi, \eta)] \in \text{supp}(\alpha) : [\xi, \eta] \cap B(ho, 1) \neq \emptyset\} \simeq V_{ho}.$$

Therefore for any $x = [ho] \in \Gamma \setminus X$,

$$\begin{aligned} \alpha_x(p^{-1}(x)) &= \alpha_x(V_{ho}) \\ &= \int_{(\xi, \eta) \in V_{ho}} \phi_{ho}(\xi, \eta) d\nu_{ho}(\xi) d\bar{\nu}_{ho}(\eta) \\ &\leq c \int_{(\xi, \eta) \in V_{ho}} d\nu_{ho}(\xi) d\bar{\nu}_{ho}(\eta) \\ &\leq c \cdot |\nu_{ho}| \cdot |\bar{\nu}_{ho}| = c \cdot E_\nu(x) \cdot E_{\bar{\nu}}(x). \end{aligned}$$

This proves (7.5), and hence finishes the proof. \square

Proof of Theorem 7.1. Suppose that $\phi \in L^2(\Gamma \setminus X)$ is a positive joint eigenfunction. By Proposition 3.7, $\phi = E_\nu$ for some (Γ, ψ) -conformal density ν . We may form the MA -semi-invariant measure $m_{\nu, \nu}$, and apply Theorem 7.4. Since $E_\nu \in L^2(\Gamma \setminus G)$, it follows that $m_{\nu, \nu}(\Gamma \setminus G) < \infty$. The finiteness of $|m_{\nu, \nu}|$ implies that $m_{\nu, \nu}$ is indeed MA -invariant by (7.3) and it is conservative for any one-parameter subgroup of A . In particular, for any non-zero $v \in \mathfrak{a}^+$, there exist $g \in G$, sequences $t_i \rightarrow +\infty$ and $\gamma_i \in \Gamma$ such that the sequence $\gamma_i g \exp(t_i v)$ is convergent. This implies that $\sup_i \|t_i v - \mu(\gamma_i^{-1})\| < \infty$ and hence $t_i^{-1} \mu(\gamma_i^{-1})$ converges to v , and hence $v \in \mathcal{L}$. Therefore $\mathcal{L} = \mathfrak{a}^+$. This finishes the proof.

Remark 7.5. If $\Gamma < G$ is Zariski dense and $\psi > \psi_\Gamma$, then for any (Γ, ψ) -conformal density ν , $E_\nu \notin L^2(\Gamma \setminus X)$. To see this, note that by [40, Lem. III. 1.3], the condition $\psi > \psi_\Gamma$ implies that

$$\sum_{\gamma \in \Gamma} e^{-\psi(\mu(\gamma))} < \infty.$$

On the other hand, by Theorem 1.4 of [5], the finiteness of $m_{\nu, \nu}$ implies that $\sum_{\gamma \in \Gamma} e^{-\psi(\mu(\gamma))} = \infty$. Hence we must have $|m_{\nu, \nu}| = \infty$. Then the claim follows from Theorem 7.4.

8. INJECTIVITY RADIUS AND $L^2(G) \otimes L^2(\Gamma \setminus G)$

As before let G be a connected semisimple real algebraic group. Recall from Proposition 6.3 that $\sigma(X) = [\|\rho\|^2, \infty)$. In this section, we prove the following:

Theorem 8.1. *Let $\Gamma < G$ be an Anosov subgroup. We suppose that Γ is not a cocompact lattice in a rank one group G . Then*

$$L^2(G) \propto L^2(\Gamma \backslash G) \quad \text{and} \quad \sigma(X) = [\|\rho\|^2, \infty) \subset \sigma(\Gamma \backslash X).$$

Note that if $\Gamma < G$ is Anosov, $\Gamma \backslash G$ has infinite volume except when Γ is a cocompact lattice in a rank one group G . The latter case has to be ruled out from the above theorem since the conclusions are not true in that case; $L^2(\Gamma \backslash G)$ contains the constant function and $\sigma(\Gamma \backslash X)$ is countable. When $G = \mathrm{SO}^\circ(n, 1)$, an Anosov subgroup $\Gamma < G$ is simply a convex cocompact subgroup, in which case this theorem is well-known due to the work of Lax and Phillips [29].

We will need the following lemma: when G is of rank one, we may write $A = \{a_t : t \in \mathbb{R}\}$ as a one-parameter subgroup, and a loxodromic element $g \in G$ is of the form $g = ha_tmh^{-1}$ for some $t \neq 0$, $m \in M$ and $h \in G$. The translation axis of g is then given by $hA(o)$.

Lemma 8.2. *Let G be a simple real algebraic group of rank one. For any loxodromic element $g \in G$ with translation axis L and any sequence $x_i \in X$ such that $d(x_i, L) \rightarrow \infty$, we have $d(x_i, gx_i) \rightarrow \infty$.*

Proof. Without loss of generality, we may assume $g = m^{-1}a_{-s_0} \in MA$ with $s_0 \neq 0$ so that $L = A(o)$. Let $x_i \in X$ be a sequence such that $d(x_i, A(o)) \rightarrow \infty$ as $i \rightarrow \infty$. Write $x_i = n_i a_{-t_i}(o)$ with $n_i \in N$ and $t_i \in \mathbb{R}$.

We may then write

$$d(gx_i, x_i) = d(a_{t_i}h_i n_i a_{-t_i}, a^{-1}o).$$

where $h_i = m a_{s_0} n_i^{-1} a_{-s_0} m^{-1} \in N$. As $d(x_i, A(o)) \rightarrow \infty$, we have $a_{t_i} n_i a_{-t_i} \rightarrow \infty$. It suffices to show $a_{t_i} h_i n_i a_{-t_i} \rightarrow \infty$.

By the assumption that G has rank one, there is only one simple root, say α , and \mathfrak{n} is the sum of at most two root subspaces $\mathfrak{n} = \mathfrak{n}_\alpha + \mathfrak{n}_{2\alpha}$ where $[\mathfrak{n}, \mathfrak{n}] = \mathfrak{n}_{2\alpha}$. Note that when N is abelian, $\mathfrak{n}_{2\alpha} = \{0\}$. Hence we have that for any $X, Y \in \mathfrak{n}$,

$$\log(\exp(X) \exp(Y)) = X + Y + \frac{1}{2}[X, Y]. \quad (8.1)$$

Write $\log n_i = Y_i + Z_i$ with $Y_i \in \mathfrak{n}_\alpha$ and $Z_i \in \mathfrak{n}_{2\alpha}$. Since Ad_m preserves \mathfrak{n}_α and $\mathfrak{n}_{2\alpha}$, we have

$$\log h_i = -\mathrm{Ad}_{m a_{s_0}} \log n_i = -e^{\alpha(s_0)} \mathrm{Ad}_m Y_i - e^{2\alpha(s_0)} \mathrm{Ad}_m Z_i.$$

Therefore by (8.1), we get

$$\log h_i n_i = (1 - e^{\alpha(s_0)} \mathrm{Ad}_m) Y_i + (1 - e^{2\alpha(s_0)} \mathrm{Ad}_m) Z_i - \frac{1}{2}[e^{\alpha(s_0)} \mathrm{Ad}_m Y_i, Y_i].$$

Hence

$$\begin{aligned} \mathrm{Ad}_{a_{t_i}} \log h_i n_i &= (1 - e^{\alpha(s_0)} \mathrm{Ad}_m) e^{\alpha(t_i)} Y_i + \\ &\quad (1 - e^{2\alpha(s_0)} \mathrm{Ad}_m) e^{2\alpha(t_i)} Z_i - [e^{\alpha(s_0)} \mathrm{Ad}_m e^{\alpha(t_i)} Y_i, e^{\alpha(t_i)} Y_i]. \end{aligned}$$

Now suppose that $a_{t_i}h_i n_i a_{-t_i}$ does not go to infinity as $i \rightarrow \infty$. By passing to a subsequence, we may assume that $\text{Ad}_{a_{t_i}} \log h_i n_i$ is uniformly bounded. It follows that both sequences $(1 - e^{\alpha(s_0)} \text{Ad}_m) e^{\alpha(t_i)} Y_i$ and $(1 - e^{2\alpha(s_0)} \text{Ad}_m) e^{2\alpha(t_i)} Z_i - [e^{\alpha(s_0)} \text{Ad}_m e^{\alpha(t_i)} Y_i, e^{\alpha(t_i)} Y_i]$ are uniformly bounded. Since $\alpha(s_0) \neq 0$, we have $e^{\alpha(t_i)} Y_i$ is uniformly bounded, which then implies that $e^{2\alpha(t_i)} Z_i$ is uniformly bounded. This implies that $\text{Ad}_{a_{t_i}} \log n_i = e^{\alpha(t_i)} Y_i + e^{2\alpha(t_i)} Z_i$ is uniformly bounded, contradicting the hypothesis that $d(a_{t_i} n_i a_{-t_i}) \rightarrow \infty$ as $i \rightarrow \infty$. This proves the claim. \square

Let $\Gamma < G$ be a discrete subgroup. For $x = [g] \in \Gamma \backslash G$, the injectivity radius $\text{inj } x$ is defined as the supremum $r > 0$ such that the ball $B_r(g) = \{h \in G : d(h, g) < r\}$ injects to $\Gamma \backslash G$ under the canonical quotient map $G \rightarrow \Gamma \backslash G$. The injectivity radius of $\Gamma \backslash G$ is defined as $\text{inj}(\Gamma \backslash G) = \sup_{x \in \Gamma \backslash G} \text{inj}(x)$.

Proposition 8.3. *For any Anosov subgroup $\Gamma < G$ which is not a cocompact lattice in a rank one group G , we have $\text{inj}(\Gamma \backslash G) = \infty$.*

Proof. If G has rank one, Γ is a convex cocompact subgroup which is not a cocompact lattice. In this case, take any $\xi \in \partial X$ which is not a limit point, and any $g_i \in G$ such that $g_i(o) \rightarrow \xi$. Then $\text{inj}(g_i(o)) \rightarrow \infty$ as $i \rightarrow \infty$.

Now suppose rank $G \geq 2$. We first observe that $\text{Vol}(\Gamma \backslash G) = \infty$; otherwise, $\Gamma < G$ is a co-compact lattice, as Anosov subgroups consist only of loxodromic elements. Since any Anosov subgroup Γ is a Gromov hyperbolic group as an abstract group ([21], [3]), it follows that G is a Gromov hyperbolic space and hence must be of rank one, which contradicts the hypothesis.

If every simple factor of G has rank at least 2, the claim $\text{inj}(\Gamma \backslash G) = \infty$ follows from a more general result of Fraczyk and Gelander [14] which applies to all discrete subgroups of infinite co-volume. Therefore it remains to consider the case where $G = G_1 \times G_2$ where G_1 and G_2 are respectively semisimple real algebraic subgroups of rank at least one and of rank precisely one. Let Σ be a finitely generated group and $\pi : \Sigma \rightarrow G$ be an Anosov representation with $\Gamma = \pi(\Sigma)$ as in Definition 2.4. Let $\pi_i : \Sigma \rightarrow G_i$ be the composition of π and the projection $G \rightarrow G_i$ for each i . It follows from (2.8) that $\pi_i(\Sigma)$ is a discrete subgroup of G_i for each $i = 1, 2$. Let X_i denote the rank one symmetric space associated to G_i and let X denote the Riemannian product $X = X_1 \times X_2$. Let $R > 0$ be an arbitrary number. We will find a point $x \in X$ with $\text{inj}(x) \geq R$, i.e., $d(x, \gamma x) > R$ for all non-trivial $\gamma \in \Gamma$; this implies the claim. Choose any $x_1 \in X_1$. By the discreteness of $\pi_1(\Sigma)$, the set $\{\sigma \in \Sigma - \{e\} : d_1(\pi_1(\sigma)x_1, x_1) < R\}$ is finite, which we write as $\{\sigma_1, \dots, \sigma_m\}$. For each $\sigma \in \Sigma \setminus \{e\}$, define a subset $T_2(\sigma) \subset X_2$ by

$$T_2(\sigma) = \{z \in X_2 : d_2(\pi_2(\sigma)z, z) < R\}.$$

Note that $\pi_2(\sigma)$ is a loxodromic element of G_2 and $T_2(\sigma)$ is contained in a bounded neighborhood of the translation axis of $\pi_2(\sigma)$ by Lemma 8.2. In particular, the symmetric space X_2 is not covered by the finite union $\bigcup_{j=1}^m T_2(\sigma_j)$. Hence we may choose $x_2 \in X_2$ outside of $\bigcup_{j=1}^m T_2(\sigma_j)$. We

now claim that the injectivity radius at $x := (x_1, x_2)$ is at least R ; suppose not. Then for some $\sigma \in \Sigma - \{e\}$, $d((\pi_1(\sigma)x_1, \pi_2(\sigma)x_2), x) < R$. In particular, for $i = 1, 2$, $d_i(\pi_i(\sigma)x_i, x_i) < R$. It follows that $\sigma = \sigma_j$ for some $1 \leq j \leq m$ and $x_2 \in T_2(\sigma_j)$, contradicting the choice of x_2 . This proves the claim. \square

Theorem 8.1 follows from Proposition 8.3 and the following proposition, which was suggested by C. McMullen.

Proposition 8.4. *Let $\Gamma < G$ be a discrete subgroup with $\text{inj}(\Gamma \backslash G) = \infty$. Then*

$$L^2(G) \propto L^2(\Gamma \backslash G) \quad \text{and} \quad \sigma(X) \subset \sigma(\Gamma \backslash X).$$

Proof. To prove the first claim, we need to show that the diagonal matrix coefficients of $L^2(G)$ can be approximated by the diagonal matrix coefficients of $L^2(\Gamma \backslash G)$ uniformly on compact subsets of G .

Let v be any element of $L^2(G)$ and $\mathcal{K} \subset G$ a compact subset containing e . We will use the fact that $\text{inj}(\Gamma \backslash G) = \infty$ to construct a sequence of functions $\{F_i\} \subset C_c(\Gamma \backslash G)$ such that

$$\lim_{i \rightarrow \infty} \max_{g \in \mathcal{K}} \left| \langle g.v, v \rangle_{L^2(G)} - \langle g.F_i, F_i \rangle_{L^2(\Gamma \backslash G)} \right| = 0,$$

as required. By the density of $C_c(G)$ in $L^2(G)$, there exists a sequence $\{f_i\} \subset C_c(G)$ such that $\lim_{i \rightarrow \infty} \|f_i - v\|_{L^2(G)} = 0$, hence

$$\lim_{i \rightarrow \infty} \max_{g \in \mathcal{K}} \left| \langle g.v, v \rangle_{L^2(G)} - \langle g.f_i, f_i \rangle_{L^2(G)} \right| = 0. \quad (8.2)$$

For each $i \geq 1$, we let $R_i > 0$ be such that $(\text{supp } f_i)\mathcal{K} \subset B_{R_i}(e)$. Since $\text{inj}(\Gamma \backslash G) = \infty$, there then exists a sequence $\{g_i\} \subset G$ such that $g_i B_{R_i}(e)$ injects to $\Gamma \backslash G$, i.e. the map $h \mapsto \Gamma h$ is injective on $g_i B_{R_i}(e)$.

For each i , consider the function $F_i \in C_c(\Gamma \backslash G)$ given by

$$F_i(x) = \sum_{\gamma \in \Gamma} f_i(g_i^{-1} \gamma h) \quad \text{for any } x = [h] \in \Gamma \backslash G.$$

We then have that for any $g \in G$,

$$\begin{aligned} \langle g.F_i, F_i \rangle_{L^2(\Gamma \backslash G)} &= \int_{\Gamma \backslash G} F_i(xg) F_i(x) dx \\ &= \int_{\Gamma \backslash G} F_i(\Gamma hg) \left(\sum_{\gamma \in \Gamma} f_i(g_i^{-1} \gamma h) \right) d(\Gamma h) = \int_G F_i(\Gamma hg) f_i(g_i^{-1} h) dh \\ &= \int_G \left(\sum_{\gamma \in \Gamma} f_i(g_i^{-1} \gamma g_i h g) \right) f_i(h) dh. \end{aligned}$$

We now observe that for $h \in G$, $g \in \mathcal{K}$, and $\gamma \in \Gamma$, $f_i(g_i^{-1} \gamma g_i h g) f_i(h) \neq 0$ implies that $g_i^{-1} \gamma g_i h g \in B_{R_i}(e)$ and $h \in \text{supp } f$. This in turn implies that

both $\gamma g_i hg$ and $g_i hg$ are in $g_i B_{R_i}(e)$. Since $g_i B_{R_i}(e)$ injects to $\Gamma \backslash G$, we must then have $\gamma = e$. Hence for all $i \geq 1$ and $g \in \mathcal{K}$,

$$\begin{aligned} \langle g.F_i, F_i \rangle_{L^2(\Gamma \backslash G)} &= \int_G \left(\sum_{\gamma \in \Gamma} f_i(g_i^{-1} \gamma g_i hg) \right) f_i(h) dh \\ &= \int_G f_i(g_i^{-1} e g_i hg) f_i(h) dh = \int_G f_i(hg) f_i(h) dh = \langle g.f_i, f_i \rangle_{L^2(G)}. \end{aligned}$$

Combined with (8.2), this proves the first claim.

In order to prove the second claim, let $W^1(\Gamma \backslash X) \subset L^2(\Gamma \backslash X)$ be as defined in the proof of Theorem 6.5. Let $\lambda \in \sigma(X)$. By Weyl's criterion (Theorem 6.1), there exists a sequence of $L^2(X)$ -unit vectors $\{u_n\}_{n \in \mathbb{N}} \subset W^1(X)$ such that

$$\lim_{n \rightarrow \infty} \|(\Delta + \lambda)u_n\|_{L^2(X)} = 0.$$

Since $C_c^\infty(X)$ is dense in $W^1(X)$ with respect to $\|\cdot\|_{W^1(X)}$, we may assume that $\{u_n\}_{n \in \mathbb{N}} \subset C_c^\infty(X)$. Since $\Gamma \backslash X$ has infinite injectivity radius, for each $n \in \mathbb{N}$, we can find $g_n \in G$ so that $g_n \text{supp}(u_n)$ injects to $\Gamma \backslash G$. We may therefore define $\{v_n\}_{n \in \mathbb{N}} \subset W^1(\Gamma \backslash X)$ by

$$v_n(\Gamma g_n x) = \begin{cases} u_n(x) & \text{if } x \in \text{supp}(u_n) \\ 0 & \text{otherwise.} \end{cases}$$

The G -invariance of Δ then gives

$$\lim_{n \rightarrow \infty} \|(\Delta + \lambda)v_n\|_{L^2(\Gamma \backslash X)} = \lim_{n \rightarrow \infty} \|(\Delta + \lambda)u_n\|_{L^2(X)} = 0;$$

and so using Weyl's criterion again yields $\lambda \in \sigma(\Gamma \backslash X)$. Hence $\sigma(X) \subset \sigma(\Gamma \backslash X)$, as claimed. \square

9. TEMPEREDNESS OF $L^2(\Gamma \backslash G)$

Let G be a connected semisimple real algebraic group and $\Gamma < G$ be a Zariski dense discrete subgroup. The goal of this section is to prove Theorem 9.4 and Corollary 9.6.

Burger-Roblin measures. We set $N^+ = w_0 N w_0^{-1}$ and $N^- = N$. For a (Γ, ψ) -conformal measure ν_o on \mathcal{F} , or equivalently for a (Γ, ψ) -conformal density $\nu = \{\nu_x : x \in X\}$, we denote by m_ν^{BR} and $m_\nu^{\text{BR}*}$ the associated N^+ and N^- -invariant Burger-Roblin measures on $\Gamma \backslash G$ respectively, as defined in [9]. By [9, Lem. 4.9], it can also be defined as follows: for any $f \in C_c(\Gamma \backslash G)$,

$$m_\nu^{\text{BR}}(f) = \int_{[k]m(\exp a)n \in K/M \times M\text{AN}^+} f([k]m(\exp a)n) e^{-\psi \circ i(a)} d\nu_o(k^-) dm da dn$$

and

$$m_\nu^{\text{BR}*}(f) = \int_{[k]m(\exp a)n \in K/M \times M\text{AN}^-} f([k]m(\exp a)n) e^{\psi(a)} d\nu_o(k^+) dm da dn$$

where dm, da, dn are Haar measures on M, \mathfrak{a}, N^\pm respectively.

Recall that dx denotes the G -invariant measure on $\Gamma \backslash G$ which is defined using the $(G, 2\rho)$ -conformal measure, that is, the K -invariant probability measure on \mathcal{F} (see [9, (3.11)]). For real-valued functions f_1, f_2 on $\Gamma \backslash G$, we write

$$\langle f_1, f_2 \rangle = \int_{\Gamma \backslash G} f_1(x) f_2(x) dx$$

whenever the integral converges. We write $C_c(\Gamma \backslash G)_K$ for the space of K -invariant compactly supported continuous functions on $\Gamma \backslash G$.

Lemma 9.1. *For a (Γ, ψ) -conformal density ν and any $f \in C_c(\Gamma \backslash G)_K$, we have*

$$m_\nu^{\text{BR}}(f) = \langle f, E_\nu \rangle = m_\nu^{\text{BR}*}(f).$$

Proof. If $g = (\exp b)nk \in AN^+K$, then

$$\beta_{e^-}(go, o) = \beta_{e^+}(\exp(-i(b)), o) = i(b).$$

Hence

$$\begin{aligned} m_\nu^{\text{BR}}(f) &= \int_{KAN^+} \int_K f(k \exp b n k_0) e^{-\psi \circ i(b)} dk_0 d\nu_o(k^-) db dn \\ &= \int_G \int_K f(kg) e^{-\psi(\beta_{e^-}(go, o))} d\nu_o(k^-) dg \\ &= \int_G f(g) \int_K e^{-\psi(\beta_{k^-}(go, o))} d\nu_o(k^-) dg = \langle f, E_\nu \rangle \end{aligned}$$

If $g = (\exp b)nk \in ANK$, then $\beta_{e^+}(go, o) = -b$ and using this, the second identity can be proved similarly. \square

Local matrix coefficients for Anosov subgroups. In the rest of this section, we assume that

$\Gamma < G$ is a Zariski dense Anosov subgroup.

Lemma 9.2. *For any $\psi \in D_\Gamma$, there exists a unique unit vector $u \in \mathfrak{a}^+$ and $0 < c \leq 1$ such that $c\psi(u) = \psi_\Gamma(u)$. Moreover $u \in \text{int } \mathcal{L}$.*

Proof. Since ψ_Γ is strictly concave [38, Propositions 4.6, 4.11], there exists $0 < c \leq 1$ and unique $u \in \mathcal{L}$ such that $c \cdot \psi(u) = \psi_\Gamma(u)$. Moreover there is no linear form tangent to ψ_Γ at $\partial \mathcal{L}$ [38], and hence $u \in \text{int } \mathcal{L}$. \square

For each $v \in \text{int } \mathcal{L}$, there exists a unique linear form $\psi_v \in D_\Gamma^*$ such that $\psi_v(v) = \psi_\Gamma(v)$ and a unique (Γ, ψ_v) -conformal density supported on Λ [9, Corollary 7.8 and Theorem 7.9], which we denote by ν_v . Hence [9, Theorem 7.12], together with Lemma 9.1, implies (let $r = \text{rank } G$):

Theorem 9.3. *For any $v \in \text{int } \mathcal{L}$, there exists $\kappa_v > 0$ such that for all $f_1, f_2 \in C_c(\Gamma \backslash G)_K$ and any $w \in \ker \psi_v$,*

$$\begin{aligned} \lim_{t \rightarrow +\infty} t^{(r-1)/2} e^{t(2\rho - \psi_v)(tv + \sqrt{t}w)} \langle \exp(tv + \sqrt{t}w) f_1, f_2 \rangle \\ = \kappa_v e^{-I(w)} \cdot \langle f_1, E_{\nu_{i(v)}} \rangle \cdot \langle f_2, E_{\nu_v} \rangle \end{aligned}$$

where $I(w) \in \mathbb{R}$ is given as in [9, 7.5]. Moreover, the left-hand side is uniformly bounded over all $(t, w) \in (0, \infty) \times \ker \psi_v$ such that $tv + \sqrt{t}w \in \mathfrak{a}^+$

Theorem 9.4. (1) We have $L^2(\Gamma \backslash G)$ is tempered if and only if $\psi_\Gamma \leq \rho$.
(2) If $L^2(\Gamma \backslash G)$ is tempered, then

$$\lambda_0(\Gamma \backslash X) = \|\rho\|^2 \quad \text{and} \quad \sigma(\Gamma \backslash X) = [\|\rho\|^2, \infty).$$

Proof. The second claim follows from Theorems 6.4 and 8.1. Suppose that $\psi_\Gamma \leq \rho$. In order to show that $L^2(\Gamma \backslash G)$ is tempered, by Proposition 2.7, it suffices to show that the matrix coefficients $g \mapsto \langle g.f_1, f_2 \rangle$ are in $L^{2+\varepsilon}(G)$ for all $\varepsilon > 0$ and for all $f_1, f_2 \in C_c(\Gamma \backslash G)$, since $C_c(\Gamma \backslash G)$ is dense in $L^2(\Gamma \backslash G)$. Without loss of generality, we may just consider non-negative functions $f_1, f_2 \in C_c(\Gamma \backslash G)$. Fix any $\varepsilon > 0$. Then using the Cartan decomposition $G = KA^+K$, we have

$$\int_G \langle g.f_1, f_2 \rangle^{2+\varepsilon} dg = \int_K \int_{\mathfrak{a}^+} \int_K \langle k_1 \exp(v) k_2 \cdot f_1, f_2 \rangle^{2+\varepsilon} \Xi(v) dk_1 dv dk_2,$$

where $\Xi(v) \asymp e^{2\rho(v)}$ (cf. [24]). Denoting $F_i(\Gamma g) = \max_{k \in K} f_i(\Gamma g k) \in C_c(\Gamma \backslash G)_K$, we then have

$$\int_G \langle g.f_1, f_2 \rangle^{2+\varepsilon} dg \ll \int_{\mathfrak{a}^+} \langle \exp(v).F_1, F_2 \rangle^{2+\varepsilon} e^{2\rho(v)} dv.$$

Since $\psi_\Gamma \leq \rho$, we have $\rho \in D_\Gamma$. By Lemma 9.2, there exists $0 < c \leq 1$ such that $c\rho \in D_\Gamma^*$ and a unit vector $u_0 \in \text{int } \mathcal{L}$ such that

$$\psi_\Gamma(u_0) = c\rho(u_0).$$

We now parameterize \mathfrak{a}^+ as follows: for each $v \in \ker \rho$, define

$$t_v := \min\{t \in \mathbb{R}_{>0} : tu_0 + \sqrt{t}v \in \mathfrak{a}^+\}.$$

Substituting $u = tu_0 + \sqrt{t}v$ for $t \geq 0$ and $v \in \mathfrak{b} \cap \ker \rho$ gives $du = s \cdot t^{\frac{r-1}{2}} dt dv$ for some constant $s > 0$. Then (letting $r = \dim(\mathfrak{a})$)

$$\begin{aligned} & \int_{\mathfrak{a}^+} \langle \exp(u).F_1, F_2 \rangle^{2+\varepsilon} e^{2\rho(u)} du \\ & \ll \int_{\ker \rho} \int_{t_v}^\infty \langle \exp(tu_0 + \sqrt{t}v).F_1, F_2 \rangle^{2+\varepsilon} e^{2t\rho(u_0)} t^{(r-1)/2} dt dv. \end{aligned}$$

By Theorem 9.3, there exists $C = C(F_1, F_2) > 0$ such that

$$t^{(r-1)/2} e^{(2-c)t\rho(u_0)} \langle \exp(tu_0 + \sqrt{t}v).F_1, F_2 \rangle \leq C$$

for all $(v, t) \in \ker \rho \times [t_v, \infty)$. Combining this with the trivial bound

$$\langle g.F_1, F_2 \rangle \leq \|F_1\| \|F_2\|,$$

we have (again, for all $(v, t) \in \ker \rho \times [t_v, \infty)$),

$$\begin{aligned} & \langle \exp(tu_0 + \sqrt{t}v).F_1, F_2 \rangle^{2+\varepsilon} \\ & \leq (C + \|F_1\| \|F_2\|)^{2+\varepsilon} \left(\min \left\{ 1, t^{-(r-1)/2} e^{-(2-c)t\rho(u_0)} \right\} \right)^{2+\varepsilon} \\ & \ll \min \{1, e^{-\eta t\rho(u_0)}\} \leq e^{-\eta t\rho(u_0)}, \end{aligned}$$

where $\eta = (2 - c)(2 + \varepsilon) > 2$. This gives

$$\begin{aligned} \int_G \langle g \cdot f_1, f_2 \rangle^{2+\varepsilon} dg & \ll \int_{v \in \ker \rho} \int_{t_v}^{\infty} e^{-\eta t\rho(u_0)} e^{2t\rho(u_0)} t^{(r-1)/2} dt dv \\ & \ll \int_{\mathfrak{a}^+} e^{-(\eta-2)\rho(u)} du < \infty. \end{aligned}$$

Therefore $L^2(\Gamma \backslash G)$ is tempered.

The converse holds for a general discrete subgroup. Suppose now that $L^2(\Gamma \backslash G)$ is tempered. Then by the definition of temperedness and the estimate of $\Xi_G(g)$ in (2.9), it follows that for any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that for any $f_1, f_2 \in L^2(\Gamma \backslash G)_K$ and $u \in \mathfrak{a}^+$,

$$|\langle \exp(u) \cdot f_1, f_2 \rangle| \leq C_\varepsilon \|f_1\| \|f_2\| e^{-(1-\varepsilon)\rho(u)}. \quad (9.1)$$

Applying [31, Prop. 7.3], we get $\psi_\Gamma \leq \rho$. \square

Now recall the following recent theorem of Kim, Minsky, and Oh [23]:

Theorem 9.5. [23] *Let Γ be an Anosov subgroup of the product G of at least two simple real algebraic groups or $\Gamma < G = \mathrm{PSL}_d(\mathbb{R})$ be a Zariski dense Anosov subgroup of a Hitchin subgroup. Then*

$$\psi_\Gamma \leq \rho.$$

Hence by Theorem 9.4, we get:

Corollary 9.6. *Let $\Gamma < G$ be as in Theorem 9.5. Then $L^2(\Gamma \backslash G)$ is tempered.*

Proofs of Theorem 1.6. The equivalence (1) \Leftrightarrow (2) is proved in Theorem 9.4. The equivalence (2) \Leftrightarrow (3) follows from Theorems 8.1 and 9.4. When $\mathrm{rank} G \geq 2$, (4) holds for any Anosov subgroup by Corollary 7.2. When $\mathrm{rank} G = 1$, the implication (1) + (2) \Rightarrow (4) is due to Sullivan [45] (see also [42, Theorem 3.1]) when X is a real hyperbolic space and to [50, Theorem 1.1 and Proposition 5.1] in general.

REFERENCES

- [1] J. Barta. *Sur la vibration fondamentale d'une membrane.* C. R. Acad. Sci. Paris 204 (1937), 472–473
- [2] Y. Benoist. *Propriétés asymptotiques des groupes linéaires.* Geom. Funct. Anal. 7 (1997), no. 1, 1–47.
- [3] J. Bochi, R. Potrie, and A. Sambarino. *Anosov Representations and dominated splittings.* J. Eur. Math. Soc., 21(11):3343–3414, 2019

- [4] M. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*. Springer, Vol 319.
- [5] M. Burger, O. Landesberg, M. Lee, and H. Oh. *The Hopf-Tsuji-Sullivan dichotomy in higher rank and applications to Anosov subgroups*. Journal of Modern Dynamics, Vol 19 (2023), 301–330
- [6] M. Chow and P. Sarkar. *Local mixing of one-parameter diagonal flows on Anosov homogeneous spaces*. Preprint, arXiv:2105.11377, To appear in IMRN.
- [7] K. Corelette and A. Iozzi *Limit sets of discrete groups of isometries of exotic hyperbolic spaces*. Trans. Amer. Math. Soc. 351 (1999), 1507–1530
- [8] M. Cowling, U. Haggerup and R. E. Howe. *Almost L^2 matrix coefficients*. J. Reiner Angew., Math 387 (1988), 97–110.
- [9] S. Edwards, M. Lee and H. Oh. *Anosov groups:local mixing, counting and equidistribution*. Geometry & Topology, 27 (2023), 513–573
- [10] S. Edwards and H. Oh. *Spectral gap and exponential mixing on geometrically finite hyperbolic manifolds*. Duke Math. J, Vol 170 (2021), 3417–3458.
- [11] J. Elstrodt. *Die Resolvente zum Eigenwert problem der automorphen Formen in der hyperbolische Ebene I*. Math. Ann 203 (1975), 295–330.
- [12] J. Elstrodt. *Die Resolvente zum Eigenwert problem der automorphen Formen in der hyperbolische Ebene II*. Math. Z. 132 (1973), 99–134.
- [13] J. Elstrodt. *Die Resolvente zum Eigenwert problem der automorphen Formen in der hyperbolische Ebene III*. Math. Ann 208 (1974), 99–132.
- [14] M. Fraczyk and T. Gelander. *Infinite volume and injectivity radius*. Annals of Math., Vol 197 (2023), 389–421
- [15] H. Furstenberg. *Translation invariant cones of functions*. Bull. AMS 71 (1965), 272–326.
- [16] F. Guéritaud, O. Guichard, F. Kassel, and A. Wienhard. *Anosov representations and proper actions*. Geom. Topol., vol. 21 (2017), 485–584.
- [17] O. Guichard and A. Wienhard. *Anosov representations: Domains of discontinuity and applications*. Inventiones Math., Volume 190, Issue 2 (2012), 357–438.
- [18] Harish-Chandra. *Discrete series for semisimple Lie groups. II* Acta Math Vol. 116 (1966), 1–111.
- [19] S. Helgason. *Some Results on Invariant Differential Operators on Symmetric Spaces*. Amer. J. Math., Vol. 114, No. 4 (Aug., 1992), 789–811.
- [20] S. Helgason. *The Fourier Transform on Symmetric spaces*. Élie Cartan et les mathématiques d’aujourd’hui - Lyon 25–29 juin 1984, Astérisque S.131 (1985), 14p.
- [21] M. Kapovich, B. Leeb, and J. Porti. *Anosov subgroups: dynamical and geometric characterizations*. Eur. J. Math. 3 (2017), no. 4, 808 –898.
- [22] F. I. Karpelevič. *The geometry of geodesics and the eigenfunctions of the Beltrami-Laplace operator on symmetric spaces*. Trudy Moskov. Mat. Obšč. 14 48–185.
- [23] D. Kim, Y. Minsky and H. Oh. *Tent property of the growth indicator functions and applications*. Preprint, arXiv:2112.00877
- [24] A. Knapp. *Representation Theory of semisimple Lie Groups*. Princeton University Press (1986).
- [25] A. Knapp. *Lie Groups Beyond an introduction*. Birkhäuser Boston (1996).
- [26] F. Labourie. *Anosov flows, surface groups and curves in projective space*. Invent. Math. 165 (2006), no. 1, 51–114.
- [27] R. G. Laha. *Nonnegative eigen functions of Laplace-Beltrami operators on symmetric spaces*. Bull. Amer. Math. Soc. 74 (1968), 167–170.
- [28] S. Lang. *SL(2, \mathbb{R})*. GTM 105, Springer-Verlag.
- [29] P. Lax and R. Phillips. *The asymptotic distribution of lattice points in Euclidean and non-Euclidean spaces*. J. Funct. Anal. 46 (1982), 280–350.
- [30] M. Lee and H. Oh. *Invariant measures for horospherical actions and Anosov groups*. Preprint, arXiv:2008.05296, To appear in IMRN.

- [31] M. Lee and H. Oh. *Dichotomy and measures on limit sets of Anosov groups*. To appear in IMRN, (arXiv:2203.06794).
- [32] H. Oh. *Uniform pointwise bounds for matrix coefficients of unitary representations and applications to Kazhdan constants*. Duke Math. J. 113 (2002), no. 1, 133–192.
- [33] S. Patterson. *The Laplacian operator on a Riemann surface*. Composition Math. 31 (1975), 83–107
- [34] S. Patterson. *The Laplacian operator on a Riemann surface II*. Composition Math. 32 (1976), 71–112
- [35] S. Patterson. *The Laplacian operator on a Riemann surface III*. Composition Math. 31 (1976), 227–259
- [36] S. Patterson. *The limit set of a Fuchsian group*. Acta Math. 136 (1976), 241–273.
- [37] P. Petersen. *Riemannian Geometry* GTM 171, Second edition, Springer 2006
- [38] R. Potrie and A. Sambarino. *Eigenvalues and entropy of a Hitchin representation*. Invent. Math., Vol 209 (2017), 885–925
- [39] J.-F. Quint. *Mesures de Patterson-Sullivan en rang supérieur*. Geom. Funct. Anal. 12 (2002), 776–809.
- [40] J.-F. Quint. *Divergence exponentielle des sous-groupes discrets en rang supérieur*. Comment. Math. Helv. 77 (2002), no. 3, 563–608.
- [41] J.-F. Quint. *Propriété de Kazhdan et sous-groupes discrets de covolume infini*. Travaux mathématiques 14 (2003), 143–151.
- [42] T. Roblin and S. Tapie. *Critical exponent and bottom of the spectrum in pinched negative curvature*. Mathematical Research Letters, 22(3):929–944, 2015
- [43] J. Rosenberg. *A Quick Proof of Harish-Chandra’s Plancherel Theorem for Spherical Functions on a Semisimple Lie Group*. Proc. Am. Math. Soc. 63 (1977), 143–149.
- [44] D. Sullivan. *The density at infinity of a discrete group of hyperbolic motions*. Publ. IHES. No. 50 (1979), 171–202.
- [45] D. Sullivan. *Related aspects of positivity in Riemannian geometry*. J. Diff geometry. 25 (1987), 327–351
- [46] D. Sullivan. *Entropy, Hausdorff measures old and new, and limits of geometrically finite Kleinian groups* Acta Math. 153 (1984), 259–277.
- [47] D. Sullivan. *A Decade of Thurston Stories*. What’s next? The Mathematical legacy of William P. Thurston, edited by Dylan Thurston, Annals of Mathematics Studies, Number 205.
- [48] G. Teschl. *Mathematical Methods in Quantum Mechanics With Applications to Schrödinger Operators*. Graduate Studies in Mathematics, Volume 157, Amer. Math. Soc., Providence, 2014.
- [49] J. Tits. *Classification of algebraic semisimple groups*. Proc. Sympos. Pure. Math., Vol. 9, 33–62, Amer. Math. Soc., Providence, 1966.
- [50] T. Weich and L. Wolf. *Absence of principal eigenvalues for higher rank locally symmetric spaces*. Preprint, arXiv:2205.03167
- [51] R. Zimmer. *Ergodic theory and semisimple groups*. 1984, Birkhauser.

DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, LOWER MOUNTJOY, DH1 3LE DURHAM, UNITED KINGDOM

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06511 AND KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL

Email address: samuel.c.edwards@durham.ac.uk

Email address: hee.oh@yale.edu