

Dark Radiation from Neutrino Mixing after Big Bang Nucleosynthesis

Daniel Aloni^{1,2}, Melissa Joseph¹, Martin Schmaltz^{1,3}, and Neal Weiner³

¹*Physics Department, Boston University, Boston, Massachusetts 02215, USA*

²*Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA*

³*Center for Cosmology and Particle Physics, Department of Physics, New York University, New York, New York 10003, USA*



(Received 6 March 2023; revised 8 June 2023; accepted 30 October 2023; published 28 November 2023)

Light dark fermions can mass mix with the standard model (SM) neutrinos. As a result, through oscillations and scattering, they can equilibrate in the early universe. Interactions of the dark fermion generically suppress such production at high temperatures but enhance it at later times. We find that for a wide range of mixing angles and interaction strengths equilibration with SM neutrinos occurs at temperatures near the dark fermion mass. For masses below an MeV, this naturally occurs after nucleosynthesis and opens the door to a variety of dark sector dynamics with observable imprints on the CMB and large scale structure, and with potential relevance to the tensions in H_0 and S_8 .

DOI: [10.1103/PhysRevLett.131.221001](https://doi.org/10.1103/PhysRevLett.131.221001)

Introduction.—The range of redshifts between $z \sim 10^9$ and $z \sim 10^3$ correspond to a “desert” in Λ CDM. As the temperature cools below the MeV scale where big bang nucleosynthesis (BBN), neutrino decoupling, and e^+e^- annihilation take place, no new threshold is reached for almost 6 orders of magnitude until the eV scale where matter-radiation equality, CMB decoupling, and eventually the sum of the neutrino masses can be found. The Λ CDM desert originates from the coincidence of a large gap in the mass spectrum of the standard model between the electron mass and the scale of neutrino masses with an unrelated but perfectly overlapping gap between nuclear and atomic binding energies.

Additional dark sectors can have new particles with masses in or below these scales, possibly leading to a rich phenomenology in the desert. A minimal extension of the standard model that realizes this has one noninteracting neutral dark fermion ν_d , with mass m_{ν_d} in the desert, which mixes with the standard model (SM) neutrino via a small Dirac mass. A combination of oscillations and weak interaction scattering can easily populate this species for large enough mixing. The relevant rate of this process Γ/H peaks near $T \sim 100 \text{ MeV} [m_{\nu_d}/\text{keV}]^{1/3}$ [1], yielding a fully thermalized fermion $\Delta N_{\text{eff}} \approx 1$ at BBN for mixing angles $\sin \theta_0 \gtrsim 10^{-3}$ and a dark fermion mass m_{ν_d} anywhere in the desert. This additional radiation affects BBN and is excluded from the measurements of light element abundances which require $\Delta N_{\text{eff}}|_{T \sim 1 \text{ MeV}} \leq 0.407(95.45\%)$ [2]. Smaller mixing angles yield dark fermions which are unthermalized and cosmologically uninteresting as radiation (absent a population from pre-TeV processes) and highly constrained as dark matter.

However, this minimal picture raises many questions, in particular regarding the origin of the new particle’s mass.

A natural expectation would be that the mass arises from some dynamics, and there would be other particles and interactions, such as self-interactions, connected to it. The consequences of such an interaction can be significant. Light fermions with large mixings can have their oscillations suppressed in the early universe [3–7], changing the cosmological constraints significantly. In the presence of a self-interaction, regions of parameter space arise where a $\sim \text{keV}$ fermion with small mixings can be dark matter [8–11]. In contrast, absent self-interactions, direct production of such dark matter through weak interactions, is excluded by a combination of x-ray data and the presence of small scale structure [12] (a famous loophole exists when SM neutrinos have chemical potentials and a lepton asymmetry [13]). Thus it is clear that a dark fermion with interactions is qualitatively different from the “unnaturally minimal” scenario of an inert dark state. Upcoming CMB and large scale structure (LSS) observations will probe the Λ CDM desert, motivating a broader exploration of such models.

In this Letter, we study the equilibration of dark sectors with the SM neutrinos after BBN and neutrino decoupling. Equilibration relies on the dark sector containing at least one neutral fermion which can mix with SM neutrinos and has interactions in the dark sector. For concreteness, we consider a single dark fermion ν_d which mixes with a SM neutrino by an amount $\sin \theta_0$ in vacuum. We assume that ν_d has a self-interaction mediated by a force carrier ϕ with $m_\phi \ll m_{\nu_d}$ and coupling strength α_d . We find two important results: (i) The dark sector comes into equilibrium with the neutrinos over a very large parameter space roughly bounded only by $\theta_0^2 \alpha_d^2 M_{\text{Pl}} > m_{\nu_d}$, allowing mixing angles ranging from 1 to 10^{-13} . (ii) Over most of the parameter space the temperature at which ν_d equilibrates is α_d independent and given by




FIG. 1. Thermal history of a universe with dark sector thermalization from neutrino mixing after BBN. The dark sector initially has negligible energy density (dashed line). After neutrino decoupling and electron annihilation it equilibrates with the SM neutrinos at T_{equil} . After ν_d annihilation at $T \sim m_{\nu_d}$ the SM neutrinos rededcouple and free-stream.

$$T_{\text{equil}} \simeq m_{\nu_d} \left(\theta_0^2 \frac{M_{\text{Pl}}}{m_{\nu_d}} \right)^{1/5}. \quad (1)$$

Thus even though the range of allowed values of θ_0 and α_d is huge, ν_d naturally equilibrates at temperatures near m_{ν_d} , and at most a few orders of magnitude higher, because of the $1/5$ power. Consequently, dark sectors with light ($< \text{MeV}$) fermions often equilibrate after BBN and are therefore unconstrained by primordial light element abundances.

The simplest thermal history is sketched in Fig. 1. After neutrino decoupling and electron self-annihilation at $T \sim \text{MeV}$, the dark sector ϕ and ν_d come into equilibrium with the SM neutrinos. At the lower temperature $T \sim m_{\nu_d}$, the dark fermions ν_d annihilate away. This causes the SM neutrinos to decouple and become free-streaming again, and the entropy of ν_d is shared between ϕ and the SM neutrinos.

Importantly, dark sector equilibration with SM neutrinos *after* neutrino decoupling does not change the relativistic energy density because the total energy in neutrinos + dark sector is conserved in the equilibration process. Thus N_{eff} is unchanged during equilibration, and constraints on N_{eff} from the CMB and LSS do not *a priori* constrain it.

However, if equilibration occurs prior to 100 keV, BBN can be modified. If ν_e (rather than ν_μ or ν_τ) equilibrates with ν_d , then ν_e is cooled, suppressing $n \rightarrow p$ conversion. When $T \sim m_{\nu_d}$ there is a “step” [14–16] in the total relativistic energy density (i.e., N_{eff} increases) as ν_d annihilates away. This can affect BBN as well [17] if it occurs before 100 keV. We leave a detailed study of this for future work.

For later equilibration, BBN is unaffected. However, prior to $T \sim m_{\nu_d}$, the $\nu - \nu_d - \phi$ fluid is tightly coupled.

This, combined with the step in N_{eff} , leaves an inevitable imprint on the density perturbations of the universe.

Should other particles have couplings to ϕ and ν_d , they, too, will come into equilibrium with the SM neutrinos below $T \sim \text{MeV}$. As a result, there is a possibility for other interesting dynamics within a dark sector to affect cosmology, such as the thermalization and freeze-out of dark matter, the presence of a second “step” [14–16] in the energy density of the dark sector due to the annihilation of additional massive particles into lighter ones. Alternatively, in a minimal scenario with $m_{\nu_d} \lesssim \text{eV}$, self-interactions in (a portion of) the relativistic energy density may arise only at late times, near recombination. Neutrino-dark sector equilibration after BBN thus has a very interesting and model-dependent impact on the CMB and structure formation with possible implications for H_0 and S_8 , all of which will be probed by a wide range of upcoming experiments.

Interactions and dark sector equilibration.—A generic dark sector which contains a fermion ν_d that mixes with the SM neutrinos can equilibrate with the SM neutrinos very efficiently by the combined effect of $\nu - \nu_d$ oscillations and scattering. The relevant formalism is well developed; see Refs. [18–20]. For simplicity we consider the case of one dark fermion oscillating with one SM neutrino. The rate of conversion of a SM neutrino into a dark fermion can be written as

$$\Gamma(E) = \frac{1}{2} \sin^2 2\theta_m \frac{\Gamma_{\text{int}}}{2}, \quad (2)$$

where we assume averaging over many oscillations, Γ_{int} is the rate of scattering, θ_m is the in-medium mixing angle between the SM neutrino and the dark fermion, and both depend on the incoming neutrino energy E . The process of dark sector equilibration is the usual competition between the production rate in Eq. (2) and Hubble. The mixing angle is generally suppressed by the presence of large diagonal effective thermal masses and thus the overall conversion rate grows rapidly as T declines.

The in-medium mixing angle is given by

$$\sin^2 2\theta_m = \frac{\sin^2 2\theta_0}{(\cos 2\theta_0 - 2E\Delta V_{\text{eff}}/\Delta m^2)^2 + \sin^2 2\theta_0}, \quad (3)$$

where θ_0 is the in-vacuum angle that parametrizes the mixing between the SM neutrino and the dark fermion, $\Delta m^2 \simeq m_{\nu_d}^2$ is the mass-squared difference between the two mass eigenstates and is dominated by the dark fermion mass, and $\Delta V_{\text{eff}} = V_{\text{eff}}^{\text{SM}} - V_{\text{eff}}^{\text{DS}}$. The effective potential of ν from the SM weak interactions is well known [1] and given by $V_{\text{eff}}^{\text{SM}} \simeq -c_V G_F^2 T^4 E$ where $c_V \simeq 22$ (for mixing with ν_μ or ν_τ), and we assume vanishing lepton asymmetry [13]. The dark sector effective potential arises due to scattering with light particles and a light mediator in the dark thermal bath and can be parametrized as $2EV_{\text{eff}}^{\text{DS}} \equiv \alpha_d T_d^2$ [5]. In

what follows we take this as the definition of α_d . The expression for the effective potential (and dark interaction rate) assumes that the dark sector is self-equilibrated with temperature T_d and vanishing chemical potentials (see discussion below). The exact expression can vary with Dirac/Majorana, internal symmetries, and other model dependencies which amount to an overall $O(\text{few})$ rescaling of α_d . The precise mapping onto a specific model Lagrangian is straightforward and not important for our discussion. We ignore a possible shift of the scalar expectation value in the thermal background which would change the mass of ν_d .

The scattering rate is the sum of the SM weak interaction $\Gamma_{\text{SM}} = n_\nu \langle \sigma v \rangle_{\text{SM}} = c_\Gamma T_\nu^4 G_F^2 E$ with $c_\Gamma \simeq 0.92$ [1], and the scattering rate of the dark fermions which we parametrize as $\Gamma_{\text{DS}} = n_{\text{DS}} \langle \sigma v \rangle_{\text{DS}} \equiv \kappa \alpha_d^2 T_d^2 / E$. This assumes that the cross section scales as $\langle \sigma v \rangle_{\text{DS}} \simeq \langle \kappa \alpha_d^2 / E_{\text{CM}}^2 \rangle_{\text{DS}} \simeq \kappa \alpha_d^2 / (E T_d)$ and $n_{\text{DS}} \propto T_d^3$. Here κ is a number greater than one, which allows for the presence of additional dark states which scatter via ϕ exchange. For simplicity, we set $\kappa = 3$, and in general it would shift the precise region of parameter space but not make it much larger or smaller.

Finally, averaging the conversion rate Γ over the thermal distribution of the SM neutrinos approximately replaces $E \rightarrow 3T_\nu$ so that

$$\langle \Gamma \rangle = \frac{\frac{1}{4} \sin^2 2\theta_0 \left(3c_\Gamma T_\nu^5 G_F^2 + \alpha_d^2 \frac{T_d^2}{T_\nu} \right)}{\left(\cos 2\theta_0 + \alpha_d \frac{T_d^2}{m_{\nu d}^2} + 18c_V \frac{G_F^2 T_\nu^6}{m_{\nu d}^2} \right)^2 + \sin^2 2\theta_0}. \quad (4)$$

We can now determine if and when the dark sector equilibrates with the neutrinos by comparing Γ with the expansion rate, $H \simeq T_\nu^2 / M_{\text{Pl}}$. There are two important limits to consider. First, in the Dodelson-Widrow (DW) [1] limit of vanishing dark sector interactions, $\alpha_d = 0$, the maximum conversion rate occurs when $G_F T_\nu^3 / m_{\nu d} \sim 0.1$. This peak temperature is above an MeV so that full equilibration from DW would yield a thermalized dark sector before BBN which is excluded. The dark sector equilibrates if $\Gamma = H$ at the peak; therefore, we obtain the constraint (in the DW limit) that $\theta_0^2 m_{\nu d} M_{\text{Pl}} G_F \lesssim 100$.

A qualitatively different solution is obtained when the dark sector interactions dominate over the weak interactions. Then $\langle \Gamma \rangle / H$ grows monotonically with decreasing temperature, and we can solve for the equilibration temperature (when $T_d = T_\nu$) by setting

$$1 \simeq \frac{\langle \Gamma \rangle}{H} \simeq \frac{\theta_0^2 \alpha_d^2 T_\nu}{(1 + \alpha_d \frac{T_\nu^2}{m_{\nu d}^2})^2} \frac{M_{\text{Pl}}}{T_\nu^2} \simeq \theta_0^2 \frac{M_{\text{Pl}}}{m_{\nu d}} \frac{m_{\nu d}^5}{T_\nu^5}, \quad (5)$$

giving $T_{\text{equil}} = m_{\nu d} (\theta_0^2 M_{\text{Pl}} / m_{\nu d})^{1/5}$. It is remarkable both that this is independent of α_d and the dependence on $\theta_0^2 M_{\text{Pl}}$ is mild because of the $1/5$ power. Thus for a very broad




FIG. 2. The ratio $T_d/T_\nu^{\Lambda\text{CDM}}$ obtained from solving Eq. (6) as a function of $T_\nu^{\Lambda\text{CDM}}$ for an example point with $\alpha_d = 1$, $m_{\nu d} = 100$ eV, $g_*^{\text{DS}}/g_*^{\text{SM}} = 1$ and initial dark sector temperature, T_d , calculated from Higgs decay. Here $T_\nu^{\Lambda\text{CDM}}$ is the temperature of the active neutrinos in a reference ΛCDM with no dark sector, where we have neglected changes in $T_\nu^{\Lambda\text{CDM}}$ from the annihilation of SM particles as they become nonrelativistic. The dashed lines correspond to $T_\nu/T_\nu^{\Lambda\text{CDM}}$ where the small drop shows the approach to equilibrium with the dark sector. Equilibration between the sectors occurs when $T_d/T_\nu^{\Lambda\text{CDM}} \approx 1$. The dark (light) gray region shows where this occurs after BBN (neutrino decoupling). See text for details.

range in parameter space the dark sector equilibrates with the neutrinos, and it does so at a temperature which is at most a few orders of magnitude above the dark fermion mass. This yields the important qualitative result that in the presence of a light ($\ll$ MeV) fermion, the natural equilibration scale is *below* the BBN scale, but also above recombination (a similar phenomenology can be achieved in models of neutrinos which couple to a Majoron, and resonantly produce dark matter at late times [21]).

This intuition is borne out by a numerical calculation. Integrating the Boltzmann equations for the phase space distribution functions of dark sector particles against energy and summing over dark sector species, we obtain an evolution equation for the total energy density in the dark sector

$$\frac{d}{d \log a} (a^4 \rho_{\text{DS}}) = \frac{\langle \Gamma \rangle}{H} a^4 \left(\rho_\nu - \frac{\rho_\nu}{\rho_{\text{DS}}|_{\text{eq}}} \rho_{\text{DS}} \right), \quad (6)$$

where ρ_{DS} is the total energy density in the DS, which we solve numerically. The evolution of the dark sector temperature is shown in Fig. 2. Details on the calculation of the dark sector temperature evolution are found in the Supplemental Material [22].

Our primary result is contained in Fig. 3 which shows the large regions of parameter space where the dark sector comes into equilibrium with the SM neutrinos at some point before $T_\nu = m_{\nu d}$ and where equilibration is reached below $T_\nu = \text{MeV}$, i.e. after neutrino decoupling and BBN.




FIG. 3. Colored regions indicate the parameter space over which the dark sector comes into equilibrium with the SM neutrinos after BBN, for different values of α_d . The lower boundary of each region is determined by $T_{\text{equil}} = m_{\nu d}$, while the upper (right) boundary corresponds to equilibration after BBN (dark shaded) or neutrino decoupling (light shaded), i.e. $T_{\text{equil}} = 100 \text{ keV}$ or $= 1 \text{ MeV}$, respectively. Also shown are contours of fixed equilibration temperatures T_{equil} (dashed contours labeled 10 eV, 1 keV) for the $\alpha_d = 1$ case. The gray region shows the parameter space over which equilibration would occur above BBN in absence of dark interactions via Dodelson-Widrow production.

Note that the small “fin” regions on the right of Fig. 3 correspond to parameter space in which $\alpha_d T_{\text{equil}}^2 / m_{\nu d}^2 < 1$. For the purposes of this figure we define the equilibration temperature T_{equil} as the temperature at which $\rho_{\text{DS}} g_{\text{DS}}^{\text{DS}} / g_{\text{SM}}^{\text{SM}}$ crosses $\rho_{\text{DS}} g_{\text{SM}}^{\text{SM}} / g_{\text{SM}}^{\text{SM}}$ with ρ_{DS} obtained from solving Eq. (6) with the backreaction term omitted.

It is worth noting that because of mixing of the SM neutrinos, for most of parameter space all three SM neutrinos equilibrate with the DS in rapid succession. That only a single SM neutrino equilibrates with the DS can occur for special regions in parameter space. Either the couplings of ν_d are tuned such that it only couples to a single SM neutrino mass eigenstate, or the dark parameters are such that equilibration with the first of the SM neutrinos occurs at a temperature just above $m_{\nu d}$ so that $\nu - \nu_d$ conversion shuts off because $m_{\nu d}$ is reached before another SM neutrino can equilibrate.

Discussion.—One of the simplest extensions of the standard model is to include a massive neutral fermion that mixes with the SM neutrino. It is natural—perhaps expected—that it should come with its own interaction, as well. In the presence of such an interaction, we find that

even for very small couplings and mixings, a new eV–MeV mass fermion is equilibrated with the neutrino bath at a temperature within a few orders of magnitude of its mass, and often much less. Consequently, it typically equilibrates after BBN, leaving no imprint on light element abundances. Its implications for the CMB and LSS, however, can be significant. Once the dark fermion equilibrates at T_{equil} , a whole series of additional particles can come into equilibrium as well, including dark matter, which can have mass above T_{equil} , including above an MeV.

Although the equilibration of the dark sector does not immediately increase the energy density in radiation, it can transform some or all of the radiation into an interacting fluid. The associated mass threshold can change the relative amount of relativistic radiation, turn on or off interactions in a dark sector, and provide a basis for equilibrating a broader dark sector which may contain part or all of the dark matter.

At high values of $100 \text{ eV} \lesssim m_{\nu d} \lesssim \text{MeV}$, the dark sector equilibrates with neutrinos and then goes through the mass threshold of the dark fermion before the CMB is directly sensitive to the transition. One consequence is the increase in N_{eff} by $\Delta N_{\text{eff}} = ((g_{\text{SM}}^{\text{UV}} / g_{\text{SM}}^{\text{IR}})^{1/3} - 1) N_{\text{eq}}$, where N_{eq} is the number of neutrinos that come into equilibrium with the dark sector, and $g_{\text{SM}}^{\text{UV}}(g_{\text{SM}}^{\text{IR}})$ is the total number of effective degrees of freedom above (below) the mass threshold, including the thermalizing neutrinos. The relativistic energy below this threshold could be interacting, non-interacting, or a combination.

At intermediate values of $O(1) \text{ eV} \lesssim m_{\nu d} \lesssim 100 \text{ eV}$, equilibration typically happens before 100 eV, but the mass threshold occurs in a period which is directly probed by the CMB and LSS. This can have important implications for many observables, including H_0 [14,15] and S_8 [16].

At very low values of $m_{\nu d}$, the equilibration can happen below 100 eV, and the signal could appear as a transition of the relativistic energy from free-streaming to strongly interacting. This transition would occur sequentially for the three SM neutrino mass eigenstates and would lead to observable signals in the CMB if it occurred at times near recombination. These implications for the CMB are beyond our scope and warrant their own study.

It is interesting to consider what might be a minimal setup, where a single dark Majorana fermion comes into equilibrium with all three SM neutrinos after BBN, but then annihilates away into a real scalar ϕ before the CMB or LSS are directly sensitive. The late universe would have $N_{\text{eff}} \simeq 3.30$ with $(1 - f)N_{\text{eff}} = 2.78$ free-streaming neutrinos and $fN_{\text{eff}} = 0.53$ interacting particles (arising from ϕ). Even in this minimal model, the resulting radiation ($\Delta N_{\text{eff}} \simeq 0.26$) is within the bounds from Planck [23] but is well above the sensitivity of Simons Observatory [24] and CMB-S4 [25]; and the fraction $f = 1/(1 + 3 \cdot 7/4)$ of the “neutrinos” that is interacting can be measured from phase shifts of the CMB peaks [26–33].

If additional particles couple to ν_d or ϕ , they, too, will equilibrate at or after T_{equil} and the thermal history can be yet richer. If additional light particles are present, then the requirement that $m_\phi \ll m_{\nu_d}$ is no longer necessary for a viable cosmology. Instead only $m_\phi \ll T_{\text{equil}}$ is needed for our calculations to hold, and in this case the neutrinos would become free-streaming again at m_ϕ rather than m_{ν_d} . With additional stable particles, dark matter could be produced through thermal processes. For freeze-out, in particular, the dark matter can have masses which are above T_{equil} , and dark matter would have naturally strong couplings to a radiation bath, at least for some period. In all of these cases, ΔN_{eff} can be found simply by an appropriate counting of degrees of freedom in the UV and IR (and intermediate steps, if needed).

In summary, we have considered the thermal history of dark fermions which mix with the SM neutrinos and have self-interactions through a light ($m_\phi \ll m_{\nu_d}$) mediator. We find that such particles equilibrate at temperatures near their mass, and thus typically at late times. This implies that later universe observables, such as LSS and the CMB, are independent probes when compared to BBN for such models. This can have important implications for models attempting to address cosmological tensions. As we look forward to upcoming results from CMB telescopes such as SPT, ACT, Simons Observatory, CMB-S4 as well as studies from LSS measurements KiDS, DES, HSC, and future galaxy surveys with Rubin, Roman, and UNIONS, such models provide an example of natural late-universe phenomena which may have significant impact. Should such particles populate the Λ CDM desert, these upcoming studies may show striking deviations from Λ CDM expectations.

We thank Joshua Ruderman and David Dunskey for useful discussions and comments on an early draft. The work of D. A., M. J., and M. S. is supported by the U.S. Department of Energy (DOE) under Award No. DE-SC0015845. N. W. is supported by NSF under Award No. PHY-2210498, by the BSF under Grants No. 2018140 and 2022287, and by the Simons Foundation. M. S. thanks the CCPP at NYU for their hospitality and support.

-
- [1] S. Dodelson and L. M. Widrow, Sterile-neutrinos as dark matter, *Phys. Rev. Lett.* **72**, 17 (1994).
 - [2] T.-H. Yeh, J. Shelton, K. A. Olive, and B. D. Fields, Probing physics beyond the standard model: Limits from BBN and the CMB independently and combined, *J. Cosmol. Astropart. Phys.* **10** (2022) 046.
 - [3] B. Dasgupta and J. Kopp, Cosmologically safe eV-scale sterile neutrinos and improved dark matter structure, *Phys. Rev. Lett.* **112**, 031803 (2014).

- [4] S. Hannestad, R. S. Hansen, and T. Tram, How self-interactions can reconcile sterile neutrinos with cosmology, *Phys. Rev. Lett.* **112**, 031802 (2014).
- [5] X. Chu, B. Dasgupta, and J. Kopp, Sterile neutrinos with secret interactions—lasting friendship with cosmology, *J. Cosmol. Astropart. Phys.* **10** (2015) 011.
- [6] J. F. Cherry, A. Friedland, and I. M. Shoemaker, Short-baseline neutrino oscillations, Planck, and IceCube, [arXiv:1605.06506](https://arxiv.org/abs/1605.06506).
- [7] Y. Farzan, Ultra-light scalar saving the $3 + 1$ neutrino scheme from the cosmological bounds, *Phys. Lett. B* **797**, 134911 (2019).
- [8] R. S. L. Hansen and S. Vogl, Thermalizing sterile neutrino dark matter, *Phys. Rev. Lett.* **119**, 251305 (2017).
- [9] L. Johns and G. M. Fuller, Self-interacting sterile neutrino dark matter: The heavy-mediator case, *Phys. Rev. D* **100**, 023533 (2019).
- [10] A. De Gouvêa, M. Sen, W. Tangarife, and Y. Zhang, Dodelson-widrow mechanism in the presence of self-interacting neutrinos, *Phys. Rev. Lett.* **124**, 081802 (2020).
- [11] T. Bringmann, P. F. Depta, M. Hufnagel, J. Kersten, J. T. Ruderman, and K. Schmidt-Hoberg, Minimal sterile neutrino dark matter, *Phys. Rev. D* **107**, L071702 (2023).
- [12] K. N. Abazajian, Neutrinos in astrophysics and cosmology: Theoretical Advanced Study Institute (TASI) 2020 lectures, *Proc. Sci. TASI2020* (2021) 001 [[arXiv:2102.10183](https://arxiv.org/abs/2102.10183)].
- [13] X.-D. Shi and G. M. Fuller, A new dark matter candidate: Nonthermal sterile neutrinos, *Phys. Rev. Lett.* **82**, 2832 (1999).
- [14] D. Aloni, A. Berlin, M. Joseph, M. Schmaltz, and N. Weiner, A step in understanding the Hubble tension, *Phys. Rev. D* **105**, 123516 (2022).
- [15] N. Schöneberg and G. Franco Abellán, A step in the right direction? Analyzing the Wess Zumino dark radiation solution to the Hubble tension, *J. Cosmol. Astropart. Phys.* **12** (2022) 001.
- [16] M. Joseph, D. Aloni, M. Schmaltz, E. N. Sivarajan, and N. Weiner, A step in understanding the S_8 tension, *Phys. Rev. D* **108**, 023520 (2023).
- [17] A. Berlin, N. Blinov, and S. W. Li, Dark sector equilibration during nucleosynthesis, *Phys. Rev. D* **100**, 015038 (2019).
- [18] R. Barbieri and A. Dolgov, Bounds on sterile-neutrinos from nucleosynthesis, *Phys. Lett. B* **237**, 440 (1990).
- [19] G. Sigl and G. Raffelt, General kinetic description of relativistic mixed neutrinos, *Nucl. Phys.* **B406**, 423 (1993).
- [20] B. Dasgupta and J. Kopp, Sterile neutrinos, *Phys. Rep.* **928**, 1 (2021).
- [21] A. Berlin and N. Blinov, Thermal neutrino portal to sub-MeV dark matter, *Phys. Rev. D* **99**, 095030 (2019).
- [22] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.131.221001> for details related to solving for the temperature evolution of the dark sector including initial conditions and assumptions related to self-thermalization of the dark sector.

- [23] N. Aghanim *et al.* (Planck Collaboration), Planck 2018 results. VI. Cosmological parameters, *Astron. Astrophys.* **641**, A6 (2020); **652**, C4(E) (2021).
- [24] P. Ade *et al.* (Simons Observatory Collaboration), The Simons Observatory: Science goals and forecasts, *J. Cosmol. Astropart. Phys.* **02** (2019) 056.
- [25] K. Abazajian *et al.*, CMB-S4 science case, reference design, and project plan, [arXiv:1907.04473](https://arxiv.org/abs/1907.04473).
- [26] D. Baumann, D. Green, J. Meyers, and B. Wallisch, Phases of new physics in the CMB, *J. Cosmol. Astropart. Phys.* **01** (2016) 007.
- [27] Z. Pan, L. Knox, B. Mulroe, and A. Narimani, Cosmic microwave background acoustic peak locations, *Mon. Not. R. Astron. Soc.* **459**, 2513 (2016).
- [28] C.D. Kreisch, F.-Y. Cyr-Racine, and O. Doré, Neutrino puzzle: Anomalies, interactions, and cosmological tensions, *Phys. Rev. D* **101**, 123505 (2020).
- [29] N. Blinov and G. Marques-Tavares, Interacting radiation after Planck and its implications for the Hubble tension, *J. Cosmol. Astropart. Phys.* **09** (2020) 029.
- [30] T. Brinckmann, J.H. Chang, and M. LoVerde, Self-interacting neutrinos, the Hubble parameter tension, and the cosmic microwave background, *Phys. Rev. D* **104**, 063523 (2021).
- [31] T. Brinckmann, J.H. Chang, P. Du, and M. LoVerde, Confronting interacting dark radiation scenarios with cosmological data, *Phys. Rev. D* **107**, 123517 (2023).
- [32] S. Roy Choudhury, S. Hannestad, and T. Tram, Updated constraints on massive neutrino self-interactions from cosmology in light of the H_0 tension, *J. Cosmol. Astropart. Phys.* **03** (2021) 084.
- [33] S. Roy Choudhury, S. Hannestad, and T. Tram, Massive neutrino self-interactions and inflation, *J. Cosmol. Astropart. Phys.* **10** (2022) 018.