

Layer separation of the 3D incompressible Navier–Stokes equation in a bounded domain

Alexis F. Vasseur & Jincheng Yang

To cite this article: Alexis F. Vasseur & Jincheng Yang (2024) Layer separation of the 3D incompressible Navier–Stokes equation in a bounded domain, Communications in Partial Differential Equations, 49:4, 381-409, DOI: [10.1080/03605302.2024.2346146](https://doi.org/10.1080/03605302.2024.2346146)

To link to this article: <https://doi.org/10.1080/03605302.2024.2346146>



Published online: 25 Apr 2024.



Submit your article to this journal 



Article views: 124



View related articles 



View Crossmark data 



Layer separation of the 3D incompressible Navier–Stokes equation in a bounded domain

Alexis F. Vasseur^a and Jincheng Yang^b

^aDepartment of Mathematics, The University of Texas at Austin, Austin, TX, USA; ^bDepartment of Mathematics, The University of Chicago, Chicago, IL, USA

ABSTRACT

We provide an unconditional L^2 upper bound for the boundary layer separation of Leray–Hopf solutions in a smooth bounded domain. By layer separation, we mean the discrepancy between a (turbulent) low-viscosity Leray–Hopf solution u^ν and a fixed (laminar) regular Euler solution \bar{u} with similar initial conditions and body force. We show an asymptotic upper bound $C\|\bar{u}\|_{L^\infty}^3 T$ on the layer separation, anomalous dissipation, and the work done by friction. This extends the previous result when the Euler solution is a regular shear in a finite channel. The key estimate is to control the boundary vorticity in a way that does not degenerate in the vanishing viscosity limit.

ARTICLE HISTORY

Received 10 March 2023

Accepted 16 April 2024

KEY WORDS

Navier–Stokes equation; inviscid limit; boundary regularity; blow-up technique; layer separation

MSC SUBJECT

CLASSIFICATION (2010)

76D05; 35Q30

1. Introduction

Let $T > 0$ and let $\Omega \subset \mathbb{R}^3$ be a smooth bounded domain. Given a smooth solution $\bar{u} : (0, T) \times \Omega \rightarrow \mathbb{R}^3$ to the Euler equation with impermeability boundary condition $\bar{u}|_{\partial\Omega} \cdot n = 0$ and a regular external force $\bar{f} : (0, T) \times \Omega \rightarrow \mathbb{R}^3$:

$$\partial_t \bar{u} + \bar{u} \cdot \nabla \bar{u} + \nabla \bar{P} = \bar{f} \quad \text{div } \bar{u} = 0 \quad \text{in } \Omega, \quad (\text{EE})$$

we estimate the $L^2(\Omega)$ difference at time T between \bar{u} and any Leray–Hopf weak solution $u^\nu : (0, T) \times \Omega \rightarrow \mathbb{R}^3$ to the Navier–Stokes equation with kinematic viscosity $\nu > 0$, body force $f^\nu \in L^1(0, T; L^2(\Omega))$, and non-slip boundary condition $u^\nu|_{\partial\Omega} = 0$:

$$\partial_t u^\nu + u^\nu \cdot \nabla u^\nu + \nabla P^\nu = \nu \Delta u^\nu + f^\nu \quad \text{div } u^\nu = 0 \quad \text{in } \Omega. \quad (\text{NSE}_\nu)$$

By Leray–Hopf solutions, we mean distributional solutions u^ν in the space $C_w(0, T; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega))$ satisfying the following energy inequality for every $T' \in [0, T]$:

$$\begin{aligned} & \frac{1}{2} \|u^\nu\|_{L^2(\Omega)}^2(T') + \int_0^{T'} \int_{\Omega} \nu |\nabla u^\nu|^2 \, dx \, dt \\ & \leq \frac{1}{2} \|u\|_{L^2(\Omega)}^2(0) + \int_0^{T'} \int_{\Omega} u^\nu \cdot f^\nu \, dx \, dt. \end{aligned} \quad (1)$$

One of the fundamental questions in fluid dynamics is whether ideal fluids, governed by the Euler equation, can be used to model viscous fluids with sufficiently small viscosity ν . This can be formulated as the so-called inviscid limit problem, which questions whether the following limit of layer separation is zero:

$$\text{LS}(\bar{u}) := \limsup_{\nu \rightarrow 0} \left\{ \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2 (T) : \begin{array}{l} u^\nu(0) \rightarrow \bar{u}(0) \text{ in } L^2(\Omega) \\ f^\nu \rightarrow \bar{f} \text{ in } L^1(0, T; L^2(\Omega)) \end{array} \right\}.$$

To the best of our knowledge, this question remains open for Leray–Hopf solutions, even for dimension 2.

This paper aims to provide the following unconditional upper bound for layer separation.

Theorem 1. *There exists a universal constant $C > 0$ such that the following holds. Let $T > 0$ and let $\Omega \subset \mathbb{R}^3$ be a domain with compact, smooth boundary satisfying Assumption 1. Let $\bar{u} \in L^\infty(0, T; C^1(\Omega))$ be a solution of (EE) with a forcing term $\bar{f} \in L^1(0, T; L^2(\Omega))$. Let u^ν be a family of Leray–Hopf weak solutions to the Navier–Stokes equation (NSE _{ν}) with force $f^\nu \in L^1(0, T; L^2(\Omega))$. Then the layer separation is bounded by*

$$\text{LS}(\bar{u}) \leq CA^3 T |\partial\Omega| \exp \left(2 \int_0^T \|D\bar{u}(t)\|_{L^\infty(\Omega)} dt \right),$$

where $A = \|\bar{u}\|_{L^\infty((0,T) \times \partial\Omega)}$ is the maximum boundary velocity of the Euler solution, and $D\bar{u} = \frac{1}{2}(\nabla\bar{u} + \nabla\bar{u}^\top)$ is the symmetric velocity gradient, also known as the rate-of-strain tensor.¹

Assumption 1 will be discussed in Section 2.3. It guarantees that the boundary $\partial\Omega$, as a compact manifold, can be triangularized in a uniform way. We conjecture this assumption should be satisfied by all smooth domains.

We remark that the norm $\|D\bar{u}\|_{L^\infty(\Omega)}$ only measures the rate of strain in the interior of Ω . On the boundary $\partial\Omega$, \bar{u} can be nonzero, and as a distribution in \mathbb{R}^3 , $D\bar{u}$ can be a measure. Indeed, if \bar{u} vanishes on the boundary, then $A = 0$ and $\text{LS}(\bar{u}) = 0$, which can also be verified by elementary computation.

This result is a generalization of the previous work by the authors [1] which studied the setting when \bar{u} is a static shear flow in a finite channel without force.

1.1. Literature review

Boundary layer and the inviscid limit problem. The gap between the Euler solution \bar{u} and the low-viscosity Navier–Stokes solution u^ν is due to the “boundary layer”, which refers to a thin layer of fluid near the boundary $\partial\Omega$ that exhibits instability and turbulent structure, in contrast with the regular Euler solution \bar{u} whose behavior near the boundary is predicted to be laminar. This has been observed from physical experiments [2] and numerical simulations [3]. Using a singular asymptotic expansion, Prandtl [4] conducted an asymptotic analysis of the Navier–Stokes system near the boundary and suggested that the turbulent structure is supported in a boundary layer of width $O(\sqrt{\nu})$. Even though the width of the boundary layer converges to zero in the inviscid limit, it is unclear whether the energy inside the boundary layer always converges to zero. In fact, the Prandtl layer with a non-monotonic shear background flow is unstable and ill-posed in Sobolev spaces. See [5–8].

¹The norm of $D\bar{u}$ should be interpreted as its largest absolute eigenvalue, which corresponds to the maximum expansion/contraction rate.

An important positive result for the inviscid limit to hold is the celebrated work of Kato [9], where he proved $\text{LS}(\bar{u}) = 0$ if the energy dissipation in a boundary layer of width $\delta = cv$ vanishes.

Theorem A (Kato's Criterion [9]). *Let $\mathcal{U}_\delta(\partial\Omega, \Omega)$ be the δ -tubular neighborhood of $\partial\Omega$ in Ω with $\delta = cv$ for some $c > 0$. If the following limit holds:*

$$\lim_{\nu \rightarrow 0} \int_0^T \int_{\mathcal{U}_\delta(\partial\Omega, \Omega)} \nu |\nabla u^\nu|^2 \, dx \, dt = 0, \quad (2)$$

then $\text{LS}(\bar{u}) = 0$.

Notice that this width is thinner than the Prandtl layer. This indicates that the inviscid limit fails only when the velocity gradient near the boundary has order $\nabla u^\nu \sim O(\nu^{-1})$. There have also been unconditional results for the inviscid limit to hold when the solution and the domain enjoy additional structure, for instance, analyticity or symmetry [10–12].

Nonuniqueness and anomalous dissipation. One important piece of theoretical evidence that suggests the inviscid limit may fail for Leray–Hopf solutions is the nonuniqueness. The recent work of Albritton, Brûe and Colombo [13, 14] exhibits the nonuniqueness of Leray–Hopf solutions for the forced Navier–Stokes equation. Their construction is based on self-similar solutions [15] and Euler instability [16–18]. Moreover, at the Euler level, even near a constant plug flow $\bar{u} = Ae_1$, Székelyhidi [1, 19] constructed nonunique Euler solutions \tilde{u} using convex integrations with a layer separation of

$$\|\tilde{u}(T) - \bar{u}\|_{L^2(\Omega)}^2 = CA^3 T.$$

Note that this rate is consistent with our upper bound of layer separation. Using convex integration or self-similar solutions, there has been an extensive amount of work in the study of the nonuniqueness of the Euler and the Navier–Stokes equations in the past decades [20–23].

Moreover, the layer separation is closely related to anomalous dissipation, which we define under our context as

$$\text{AD}(\bar{u}) := \limsup_{\nu \rightarrow 0} \left\{ \int_0^T \int_{\Omega} \nu |\nabla u^\nu|^2 \, dx \, dt : \begin{array}{l} u^\nu(0) \rightarrow \bar{u}(0) \text{ in } L^2(\Omega) \\ f^\nu \rightarrow \bar{f} \text{ in } L^1(0, T; L^2(\Omega)) \end{array} \right\}.$$

Kato's criterion shows that if $\text{AD}(\bar{u}) = 0$ then $\text{LS}(\bar{u}) = 0$. It is also straightforward to see from the energy inequality (1) that $\text{LS}(\bar{u}) = 0$ would imply $\text{AD}(\bar{u}) = 0$ as well. From this perspective, the validity of the inviscid limit is equivalent to whether the Kolmogorov's *zeroth law of turbulence* [24–26] can hold in a neighborhood of regular solution \bar{u} . In the absence of boundary, Brûe and De Lellis [27] constructed examples of classical solutions to the forced Navier–Stokes equation with positive anomalous dissipation. However, both the nonunique Leray–Hopf solutions in [13] and the anomalous dissipation of [27] are away from a smooth Euler solution, so they do not fall into the scope of this paper. Nevertheless, we provide the same unconditional bound on the limiting energy dissipation as well.

Corollary 1. *Under the same assumptions of Theorem 1, the anomalous dissipation is also bounded by*

$$\text{AD}(\bar{u}) \leq CA^3 T |\partial\Omega| \exp\left(2 \int_0^T \|D\bar{u}(t)\|_{L^\infty(\Omega)} dt\right).$$

Work of boundary friction and Kato's criterion. Let us briefly discuss the main ideas of the proof. The crucial term when estimating layer separation and anomalous dissipation is the work done by the friction on the boundary. It is easy to see that the validity of the inviscid limit is equivalent to the vanishing of the negative work of boundary friction in \bar{u} direction:

$$W_{\text{fric}}(\bar{u}) := \limsup_{v \rightarrow 0} \left\{ -v \int_{(0,T) \times \partial\Omega} \partial_n u^v \cdot \bar{u} dx' dt : \begin{array}{l} u^v(0) \rightarrow \bar{u}(0) \text{ in } L^2(\Omega) \\ f^v \rightarrow \bar{f} \text{ in } L^1(0, T; L^2(\Omega)) \end{array} \right\},$$

where $\partial_n u^v \cdot \bar{u} = \omega^v \cdot (n \times \bar{u})$ on the boundary $\partial\Omega$.

Using energy inequality and Grönwall inequality, it is easy to see that

$$\text{LS}(\bar{u}) + \text{AD}(\bar{u}) \leq W_{\text{fric}}(\bar{u}) \exp\left(2 \int_0^T \|D\bar{u}\|_{L^\infty(\Omega)} dt\right). \quad (3)$$

Hence, measuring the layer separation and anomalous dissipation relies on the estimation of **boundary vorticity** ω^v . We will provide a uniform bound in $L^{\frac{3}{2}}$ weak space in [Theorem 3](#), using the energy dissipation in the Kato's layer. As a consequence of this vorticity estimate, we can control the total work of boundary friction force asymptotically by

$$W_{\text{fric}}(\bar{u}) \leq C \|\bar{u}\|_{L^{3,1}((0,T) \times \partial\Omega)} \text{AD}_{cv}(\bar{u})^{\frac{2}{3}} \leq CA(T|\partial\Omega|)^{\frac{1}{3}} \text{AD}_{cv}(\bar{u})^{\frac{2}{3}},$$

where $\text{AD}_{cv}(\bar{u})$ is the limiting energy dissipation in the boundary layer of width cv . Therefore, if Kato's criterion holds, the work of (3) implies both layer separation and anomalous dissipation are also zero. Otherwise, by absorbing the anomalous dissipation into the left side of (3) we show the layer separation $\text{LS}(\bar{u})$, anomalous dissipation $\text{AD}(\bar{u})$, and total friction work $W_{\text{fric}}(\bar{u})$ are all bounded by $CA^3 T |\partial\Omega|$, up to an exponential factor which depends on the largest absolute eigenvalue of the strain-rate tensor $D\bar{u}$:

$$\text{LS}(\bar{u}) + \text{AD}(\bar{u}) + W_{\text{fric}}(\bar{u}) \leq CA^3 T |\partial\Omega| \exp\left(2 \int_0^T \|D\bar{u}(t)\|_{L^\infty(\Omega)} dt\right).$$

See [Remark 2](#) for a further discussion on physical relevance.

1.2. Main results

Both [Theorem 1](#) and [Corollary 1](#) are the consequence of the following bound at the Navier-Stokes level.

Theorem 2. *Let $\Omega \subset \mathbb{R}^3$ be a bounded domain with compact, smooth boundary satisfying [Assumption 1](#) with width $\bar{\delta}$. There exists a constant $C(\Omega) > 0$ depending only on Ω and a universal constant C such that the following is true. Given $T > 0$, let \bar{u} be a regular solution to [\(EE\)](#) with maximum boundary velocity $A = \|\bar{u}\|_{L^\infty((0,T) \times \partial\Omega)}$, and let u^v be a Leray-Hopf weak solution to [\(NSE_v\)](#) with initial value $u^v(0) \in H^1(\Omega)$ and force $f^v \in L^1(0, T; L^2(\Omega)) \cap L^{\frac{4}{3}}((0, T) \times \Omega)$. Define the characteristic frequency and Reynolds number by*

$$\frac{A}{L} = A^{-1} \|\partial_t \bar{u}\|_{L^\infty((0,T) \times \partial\Omega)} + \|\nabla \bar{u}\|_{L^\infty((0,T) \times \mathcal{U}_\delta(\partial\Omega, \Omega))}, \quad \text{Re} = \frac{AL}{v}.$$

Here $\delta = \min \{\bar{\delta}, A^{-1}v\}$. Then

$$\begin{aligned} & \|u^v - \bar{u}\|_{L^2(\Omega)}^2(T) + \frac{v}{2} \|\nabla u^v\|_{L^2((0,T) \times \Omega)}^2 \\ & \leq \left(\|u^v - \bar{u}\|_{L^2(\Omega)}^2(0) + CA^3 T |\partial\Omega| + R_v(T) \right) \\ & \quad \times \exp \left(\int_0^T 2 \|D\bar{u}\|_{L^\infty(\Omega)}(t) + \|f^v - \bar{f}\|_{L^2(\Omega)}(t) dt \right), \end{aligned}$$

where the remainder term $R_v(T)$ is defined by

$$\begin{aligned} R_v(T) &= \|f^v - \bar{f}\|_{L^1(0,T; L^2(\Omega))} + v \|\nabla \bar{u}\|_{L^2((0,T) \times \Omega)}^2 \\ &\quad + v^{\frac{1}{3}} \|f^v\|_{L^{\frac{4}{3}}((0,T) \times \Omega)}^{\frac{4}{3}} + 2v^{\frac{4}{3}} \|u^v(0)\|_{H^1(\Omega)}^{\frac{2}{3}} \\ &\quad + 2 \left(4 \log \left(\frac{4AL}{v} \right) + \frac{vT}{\bar{\delta}^2} + \frac{C(\Omega)(1+v^2)E_v T}{AL^4} \right) A v |\partial\Omega|. \end{aligned}$$

As mentioned earlier, the crucial step is to bound the boundary vorticity in a way that does not degenerate as $v \rightarrow 0$. We show that the averaged boundary vorticity can be bounded in $L^{\frac{3}{2}}$ weak norm, up to a remainder, by the energy dissipation in the boundary layer (2).

Theorem 3. *There exists a universal constant C such that the following is true. Let $\Omega \subset \mathbb{R}^3$ be a smooth bounded domain satisfying Assumption 1 with $\bar{\delta}$. Let $u^v \in L^2(0, T; H^1(\Omega))$ be a weak solution to (NSE_v) with force $f^v \in L^{\frac{4}{3}}((0, T) \times \Omega)$. We denote $\omega^v = \operatorname{curl} u^v$ to be the vorticity field. For any $0 < \delta \leq \bar{\delta}$, there exists a σ -algebra \mathcal{F} of $(0, T) \times \partial\Omega$, depending on u^v and δ , such that*

1. \mathcal{F} is a sub σ -algebra of the Borel σ -algebra on $(0, T) \times \partial\Omega$. For every integer $l \geq 0$ with $4^{-l}T \leq \delta^2$, the set $(0, 4^{-l-1}T) \times \Omega$ is \mathcal{F} -measurable.
2. For $\varphi \in C^1((0, T) \times \partial\Omega)$, we have

$$\|\varphi - \mathbb{E}[\varphi | \mathcal{F}]\|_{L^\infty} \leq \delta \left(\frac{\delta}{v} \|\partial_t \varphi\|_{L^\infty} + \|\nabla \varphi\|_{L^\infty} \right). \quad (4)$$

3. Denote $\tilde{\omega}^v = \mathbb{E}[\omega^v | \mathcal{F}]$. Then for every $\gamma \leq 1$,

$$\begin{aligned} & \left\| v \tilde{\omega}^v \mathbf{1}_{\{v|\tilde{\omega}^v| > \gamma \max\left\{\frac{v}{t}, \frac{v^2}{\delta^2}\right\}\}} \right\|_{L^{\frac{3}{2}, \infty}((0, T) \times \partial\Omega)}^{\frac{3}{2}} \\ & \leq C \gamma^{-\frac{1}{2}} \int_0^T \int_{\mathcal{U}_\delta(\partial\Omega, \Omega)} v |\nabla u^v|^2 + v^{\frac{1}{3}} |f^v|^{\frac{4}{3}} dx dt. \end{aligned} \quad (5)$$

Remark 1. If we set the boundary layer to have the width $\delta = O(v)$ as in Kato's condition, then (4) will be of order $O(v)$. We will recover Kato's results if the right-hand side of (5) vanishes in the inviscid limit.

The σ -algebra is constructed by a partition of $(0, T) \times \partial\Omega$ in a dyadic way. Morally speaking, we ensure in each piece with size $r < v$ in space and length $v^{-1}r^2$ in time, the average energy dissipation near it is $\int v |\nabla u^v|^2 dx dt \sim c_0 v^3 r^{-4}$, from which we control the

average boundary vorticity by $\tilde{\omega}^v = \int \omega^v \, dx' \, dt \lesssim \nu r^{-2}$ via a linear Stokes estimate. After a Calderón–Zygmund argument, we can control $\tilde{\omega}^v$ in a weak norm by (5).

This paper is organized as follows. Section 2 introduces the technical tools that will help deal with the non-flatness of the boundary, especially the dyadic decomposition. In Section 3 we prove the boundary vorticity estimate in Theorem 3. The main results will be proven in Section 4.

2. Preliminary on curved boundary

In this section, we discuss issues that arise due to the non-flatness of the boundary. We first rigorously define the triangular decomposition of $\partial\Omega$. Then we recall some classical estimates with curved boundaries.

2.1. Notation

Let \mathcal{D}_2 denote the set of open triangles in \mathbb{R}^2 with barycenter at the origin and side lengths between $\frac{5}{3}$ and $\frac{7}{3}$. Define Ψ to be the set of diffeomorphisms between any such triangle $\Delta_2 \in \mathcal{D}_2$ and any piece of two-dimensional surface $T_2 \subset \mathbb{R}^3$ under the following restriction:

$$\Psi := \left\{ \psi \in \text{Diff}(\Delta_2; T_2) : \begin{array}{l} \Delta_2 \in \mathcal{D}_2, T_2 \subset \mathbb{R}^3, \psi(0) = 0 \\ \nabla \psi(0) = i_{\mathbb{R}^2}, \|\nabla^2 \psi\|_{L^\infty} \leq \frac{1}{9} \end{array} \right\}.$$

Here $\text{Diff}(\Delta_2; T_2)$ is the set of smooth diffeomorphisms between Δ_2 and T_2 , and $i_{\mathbb{R}^2}$ is the natural inclusion from \mathbb{R}^2 to \mathbb{R}^3 defined by $(x_1, x_2) \mapsto (x_1, x_2, 0)$. By translation, rotation, reflection and dilation/contraction, we define $\Psi^{(r)}$ to be the set of diffeomorphisms $\psi : \Delta_2 \rightarrow T_{2r} \subset \mathbb{R}^3$ using the following:

$$\Psi^{(r)} := \{rR \circ \psi : \psi \in \Psi, R \in E(3)\}, \quad r > 0.$$

$E(3)$ is the isometry group of \mathbb{R}^3 .

We could also extend it with width. Denote $\Delta_2 = \Delta_2 \times (0, 2)$, and for $\psi \in \Psi^{(r)}$, denote the unit normal vector by $n = \frac{\partial_1 \psi \times \partial_2 \psi}{|\partial_1 \psi \times \partial_2 \psi|}$. We can define the extended diffeomorphism $\tilde{\psi} \in C^\infty(\Delta_2; \mathbb{R}^3)$ by

$$\tilde{\psi}(\xi, z) = \psi(\xi) - rzn(\xi), \quad \xi \in \Delta_2, z \in (0, 2).$$

The first and second derivative constraint ensures that $\tilde{\psi}$ is also a homeomorphism.

By $r\Delta_2$ we mean the scaling of Δ_2 by a factor of $r > 0$, and $r\Delta_2$ refers to the scaling of Δ_2 by a factor of r . For $r > 0$, we define the set of *curved triangles with size r* by

$$\mathcal{T}^{(r)} := \left\{ \psi(\Delta_1) : \psi \in \Psi^{(r)}, \Delta_1 = \frac{1}{2} \text{Dom } \psi \right\},$$

and we define the set of *curved triangular cylinders with size r* by

$$\mathcal{C}^{(r)} := \left\{ \tilde{\psi}(\Delta_1) : \psi \in \Psi^{(r)}, \Delta_1 = \frac{1}{2} \text{Dom } \tilde{\psi} \right\}.$$

Therefore, each curved triangle (triangular cylinder) has a neighborhood that is diffeomorphic to a triangle (triangular cylinder) with a uniform bound on the second derivative of the diffeomorphism. If $T_1 = \psi(\Delta_1)$ and $C_1 = \tilde{\psi}(\Delta_1)$, we say C_1 is the *extension* of T_1 and T_1 is the *base* of C_1 . For $r \in (0, 2)$ we denote $rT_1 = \psi(r\Delta_1)$ and $rC_1 = \tilde{\psi}(r\Delta_1)$. Note that these definitions do not depend on the choice of diffeomorphisms up to the orientation.

2.2. Dyadic decomposition of boundary

Below we describe the dyadic decomposition of triangles, cylinders, and time. Recall \simeq means that two sets are equal up to zero measure sets, and \sqcup is the disjoint union operator.

1. For a triangle $\Delta_2 \in \mathcal{D}_2$, it can be decomposed into four similar sub-triangles with half the side lengths, by connecting mid-points. Denote the four sub-triangles by $\Delta_1^{(i)}, i = 1, \dots, 4$. Then

$$\Delta_2 \simeq \bigsqcup_{i=1}^4 \Delta_1^{(i)}.$$

2. For a triangular cylinder $\Delta_2 = \Delta_2 \times (0, 2)$, we decompose $\Delta_2 \times (0, 1)$ into four pieces $\Delta_1^{(i)} = \Delta_1^{(i)} \times (0, 1)$, by decompose the base Δ_2 . See Figure 1. The top part $\Delta_2^* = \Delta_1 \times (1, 2)$ will be discarded.

$$\Delta_2 \simeq \left(\bigsqcup_{i=1}^4 \Delta_1^{(i)} \right) \sqcup \Delta_2^*.$$




Figure 1. Dyadic decomposition of $\Delta_2 = \left(\bigsqcup_{i=1}^4 \Delta_1^{(i)} \right) \sqcup \Delta_2^*$.

3. For $T_2 \in \mathcal{T}^{(2)}$, it is diffeomorphic to Δ_2 via $\psi(\frac{1}{2} \cdot)$ for some $\psi \in \Psi^{(2)} : \frac{1}{2} \Delta_2 \rightarrow T_2$. We first decompose Δ_2 . Then we map the four pieces to T_2 via $\psi : T_1^{(i)} = \psi(\frac{1}{2} \Delta_1^{(i)})$. In this way, T_2 is decomposed into

$$T_2 \simeq \bigsqcup_{i=1}^4 T_1^{(i)}.$$

Each $T_1^{(i)}$ belongs to $\mathcal{T}^{(1)}$. This decomposition is not unique and depends on the choice of the diffeomorphism.

4. For $C_2 \in \mathcal{C}^{(2)}$, it is diffeomorphic to Δ_2 via $\tilde{\psi}(\frac{1}{2} \cdot)$ for some $\tilde{\psi} \in \Psi^{(2)} : \frac{1}{2} \Delta_2 \rightarrow C_2$. We first decompose Δ_2 . Then we map the four pieces into C_2 via $\tilde{\psi} : C_1^{(i)} = \tilde{\psi}(\Delta_1^{(i)})$. The

remaining part $C_2^* = \tilde{\psi}(\Delta_2^*)$ is discarded.

$$C_2 \simeq \left(\bigsqcup_{i=1}^4 C_1^{(i)} \right) \sqcup C_2^*.$$

Each $C_1^{(i)}$ belongs to $\mathcal{C}^{(1)}$. Moreover, the base of C_2 is correspondingly decomposed to the bases of $C_1^{(i)}$.

5. For $\bar{Q}_2 := (-4, 0) \times T_2$, it can be decomposed into 16 pieces in both time and space: $\bar{Q}_1^{(i,j)} = (-j, -j+1) \times T_1^{(i)}$, where $i, j = 1, \dots, 4$.

$$\bar{Q}_2 \simeq \bigsqcup_{i=1}^4 \bigsqcup_{j=1}^4 \bar{Q}_1^{(i,j)}.$$

6. If C_2 is the extension of some $T_2 \in \mathcal{C}^{(2)}$, we say $Q_2 := (-4, 0) \times C_2$ is the extension of $\bar{Q}_2 = (-4, 0) \times T_2$. Q_2 can be decomposed into 16 pieces in both time and space: $Q_1^{(i,j)} = (-j, -j+1) \times C_1^{(i)}$, where $i, j = 1, \dots, 4$. The remaining part $Q_2^* = (-4, 0) \times C_2^*$ is discarded.

$$Q_2 \simeq \left(\bigsqcup_{i=1}^4 \bigsqcup_{j=1}^4 Q_1^{(i,j)} \right) \sqcup Q_2^*.$$

By scaling, every $T \in \mathcal{T}^{(\delta)}$ can be decomposed into four curved triangles in $\mathcal{T}^{(\frac{\delta}{2})}$, and every $C \in \mathcal{C}^{(\delta)}$ can be decomposed into four curved triangular cylinders in $\mathcal{C}^{(\frac{\delta}{2})}$ with a remainder part.

2.3. Structural assumption on the domain

Let $\Omega \subset \mathbb{R}^3$ be a bounded domain, whose boundary $\partial\Omega$ is a compact smooth manifold. We assume the following geometric property for the set Ω .

Assumption 1. Assume there exists a constant $\bar{\delta}$, such that Ω has a $\bar{\delta}$ -tubular neighborhood

$$\mathcal{U}_{\bar{\delta}}(\partial\Omega, \Omega) := \{x \in \Omega : \text{dist}(x, \partial\Omega) < \bar{\delta}\},$$

and $(x', \varepsilon) \mapsto x' - \varepsilon n(x')$ is a diffeomorphism from $\partial\Omega \times (0, \bar{\delta})$ to $\mathcal{U}_{\bar{\delta}}(\partial\Omega, \Omega)$, where $n(x')$ is the outer normal vector of $\partial\Omega$ at x' . Moreover, we assume that for every $\delta \in (0, \bar{\delta})$, $\partial\Omega$ has a *curved* triangular decomposition:

$$\partial\Omega \simeq \bigsqcup_i T_{\delta}^{(i)}, \quad T_{\delta}^{(i)} \in \mathcal{T}^{(\delta)}.$$

Intuitively, the assumption should hold for any compact smooth manifold with $\bar{\delta} \ll \gamma_{\partial\Omega}^{-1}$, where $\gamma_{\partial\Omega}$ is the greatest sectional curvature of $\partial\Omega$. In the computer vision community, the “marching triangle” algorithm [28, 29] is used to generate a triangular mesh for two-dimensional manifolds (or in general Lipschitz surfaces [30]) with triangular patches uniform in shape and size, meaning that each patch is close to an equilateral triangle and has comparable edge lengths. However, the authors did not provide explicit estimates for the size δ and the angles of the triangulation (Figure 2).




Figure 2. Triangulation of a torus.

2.4. Sobolev, trace, and Stokes on a curved cylinder

This subsection includes basic analysis tools that will be used later in the proofs. The constants in the following estimates must be uniform in the geometry of the boundaries of our interest.

The first lemma contains the Sobolev embedding and the trace theorem. We remind the reader that their constants are uniform for all $C \in \mathcal{C}^{(1)}$. These results are well-known so we omit the proof.

Lemma 1. *Let $C_1 \in \mathcal{C}^{(1)}$ be a curved cylinder with base $T_1 \in \mathcal{T}^{(1)}$. Let $\nabla u \in L^p(C_1)$ with either $\int_{C_1} u \, dx = 0$ or $u|_{T_1} = 0$, and $p \in [1, 3)$. Then there exists a constant C_p depending only on p , such that*

$$\|u\|_{L^{p^*}(C_1)} \leq C_p \|\nabla u\|_{L^p(C_1)}.$$

Here $p^* = \frac{dp}{d-p}$. Note that C_p does not depend on C_1 . Moreover, with $p^* = \frac{(d-1)p}{d-p}$,

$$\|u\|_{L^{p^*}(T_1)} \leq C_p \|\nabla u\|_{L^p(C_1)}.$$

In this paper, $d = \dim C_1 = 3$.

The next lemma is for the local boundary linear Stokes estimate, which is an extension of [1, Corollary 2.3]. The only difference is that the boundary part T_2 is no longer flat, yet the bound is still uniform for all $C_1 \in \mathcal{C}^{(1)}$.

Lemma 2. *Let $C_1 \in \mathcal{C}^{(1)}$ be a curved cylinder with base $T_1 \in \mathcal{T}^{(1)}$. Let C_2 be the image of a diffeomorphism ψ associated with C_1 , and denote its base by T_2 . Let $1 < p_2 < p_1 < \infty$, $1 < q_1, q_2 < \infty$, $f \in L^{p_1}(-4, 0; L^{q_1}(C_2))$. If (u, P) solves the linear evolutionary Stokes system*

$$\begin{cases} \partial_t u + \nabla P = \Delta u + f & \text{in } (-4, 0) \times C_2 \\ \operatorname{div} u = 0 & \text{in } (-4, 0) \times C_2 \\ u = 0 & \text{on } (-4, 0) \times T_2, \end{cases}$$

then there exists a decomposition $u = u_1 + u_2$ such that for any $q' < \infty$, there exists a constant $C = C(p_1, p_2, q_1, q_2, q')$ such that

$$\begin{aligned} & \| |\partial_t u_1| + |\nabla^2 u_1| \|_{L^{p_1}(-1, 0; L^{q_1}(C_1))} + \| |\partial_t u_2| + |\nabla^2 u_2| \|_{L^{p_2}(-1, 0; L^{q'}(C_1))} \\ & \leq C \left(\|f\|_{L^{p_1}(-4, 0; L^{q_1}(C_2))} + \|u\| + |\nabla u| + |P|\|_{L^{p_2}(-4, 0; L^{q_2}(C_2))} \right). \end{aligned}$$

In particular, C does not depend on the geometry of C_1 .

The proof of this lemma relies on [Lemma 1](#), and also the corresponding uniform bound for boundary estimates and the Cauchy problem for the Stokes with curved boundary. See [\[31\]](#).

Proof Pick a set Ω with C^2 boundary such that $C_1 \subset \Omega \subset C_2$. Note that the C^2 norm of $\partial\Omega$ can be uniformly bounded for all $C_1 \in \mathcal{C}^{(1)}$. By [\[32, Theorem 4.5\]](#), there exists a unique solution u_1 to the initial-boundary value problem

$$\begin{cases} \partial_t u_1 + \nabla P_1 = \Delta u_1 + f & \text{in } (-4, 0) \times \Omega \\ \operatorname{div} u_1 = 0 & \text{in } (-4, 0) \times \Omega \\ u_1 = 0 & \text{on } (-4, 0) \times \partial\Omega \\ u_1|_{t=-4} = 0 & \text{in } \Omega \end{cases}$$

with bound

$$\begin{aligned} & \| |\partial_t u_1| + |\nabla P_1| \|_{L^{p_1}(-4, 0; L^{q_1}(\Omega))} + \| u_1 \|_{L^{p_1}(-4, 0; W^{2, q_1}(\Omega))} \\ & \leq C \| f \|_{L^{p_1}(-4, 0; L^{q_1}(\Omega))}, \end{aligned}$$

where $C = C(p_1, q_1, \Omega)$. Dependence on Ω can be dropped if the C^2 norm of $\partial\Omega$ is uniformly bounded (see [\[31, 33, Lemma 1.2\]](#)).

Let $u_2 = u - u_1$. Then u_2 is a solution to

$$\begin{cases} \partial_t u_2 + \nabla P_2 = \Delta u_2 & \text{in } (-4, 0) \times \Omega \\ \operatorname{div} u_2 = 0 & \text{in } (-4, 0) \times \Omega \\ u_2 = 0 & \text{on } (-4, 0) \times (\partial\Omega \cap T_2) \end{cases}.$$

The local boundary estimates of Stokes equation in [\[32\]](#) imply the following bound:

$$\begin{aligned} & \| |\partial_t u_2| + |\nabla^2 u_2| \|_{L^{p_2}(-1, 0; L^{q'}(C_1))} \\ & \leq C \| |u_2| + |\nabla u_2| + |P_2| \|_{L^{p_2}(-4, 0; L^{\min\{q_1, q_2\}}(\Omega))}, \end{aligned}$$

where $C = C(p_2, q_1, q_2, \Omega)$. Again, the dependence on Ω can be dropped. Combining with estimates of u_1 , we finish the proof of the lemma. \square

Finally, we quote the following global Stokes theorem in [\[34, Theorem 1.1\]](#).

Lemma 3. *Let $\Omega \subset \mathbb{R}^d$ be a bounded C^2 domain with $d \geq 2$, and $T > 0$. Let $f \in L^p(0, T; L^q(\Omega))$ and $u_0 \in B_{q,p}^{2-2/p}(\Omega)$, where $B_{q,p}^l(\Omega)$ is the Besov space, and $1 < p, q < \infty$ satisfy $2 - 2/p < 1/q$. Then the following linear evolutionary Stokes system*

$$\begin{cases} \partial_t u + \nabla P = \Delta u + f & \text{in } (0, T) \times \Omega \\ \operatorname{div} u = 0 & \text{in } (0, T) \times \Omega \\ u = 0 & \text{on } (0, T) \times \partial\Omega \\ u|_{t=0} = u_0 & \text{in } \Omega \end{cases}$$

has a unique solution $u \in L^p(0, T; W^{2,q}(\Omega))$ with $\nabla P, \partial_t u \in L^p(0, T; L^q(\Omega))$, and

$$\begin{aligned} & \| u \|_{L^p(0, T; W^{2,q}(\Omega))} + \| \partial_t u \|_{L^p(0, T; L^q(\Omega))} + \| \nabla P \|_{L^p(0, T; L^q(\Omega))} \\ & \leq C \left(\| f \|_{L^p(0, T; L^q(\Omega))} + \| u_0 \|_{B_{q,p}^{2-2/p}(\Omega)} \right), \end{aligned}$$

where $C = C(\Omega, p, q)$.

3. Boundary vorticity estimate

In this section, we provide several estimates for the boundary vorticity $\omega^v = \operatorname{curl} u^v$. We first use linear parabolic theory to directly derive a coarse estimate. This estimate will degenerate in the inviscid limit. We compensate with a refined estimate [Theorem 3](#), which is based on a new local boundary vorticity estimate for the linear Stokes system.

3.1. Naïve linear global estimate

By treating Navier–Stokes equation as a Stokes system with a forcing term, we can derive the following naïve bound using parabolic regularization.

Proposition 1. *Let $u^v \in C_w(0, T; L^2(\Omega)) \cap L^2(0, T; H_0^1(\Omega))$ be a weak solution to [\(NSE_v\)](#) with divergence-free initial value $u^v(0) \in H_0^1(\Omega)$ and force $f^v \in L^{\frac{4}{3}}(0, T; L^{\frac{6}{5}}(\Omega))$. There exists a universal constant $C(\Omega)$, independent of u^v and v , such that*

$$\int_{(0,T) \times \partial\Omega} |\nu \nabla u^v|^{\frac{4}{3}} dx' dt \leq C(\Omega) \left[\|f^v\|_{L^{\frac{4}{3}}(0,T;L^{\frac{6}{5}}(\Omega))}^{\frac{4}{3}} + \right. \\ \left. + \|u^v\|_{L^{\infty}(0,T;L^2(\Omega))}^{\frac{2}{3}} \left(\int_{(0,T) \times \Omega} |\nabla u^v|^2 dx dt + \nu^{\frac{1}{3}} \|u^v(0)\|_{H^1(\Omega)}^{\frac{2}{3}} \right) \right].$$

Proof Let $u^v(t, x) = \nu v(\nu t, x)$ and $f^v(t, x) = \nu^2 g(\nu t, x)$. Then v solves [\(NSE₁\)](#) in $(0, \nu T) \times \Omega$ with unit viscosity and force g . Treating the nonlinear term $v \cdot \nabla v$ as a force, [Lemma 3](#) implies

$$\|v\|_{L^{\frac{4}{3}}(0, \nu T; W^{2, \frac{6}{5}}(\Omega))} \leq C(\Omega) \left(\| -v \cdot \nabla v + g \|_{L^{\frac{4}{3}}(0, \nu T; L^{\frac{6}{5}}(\Omega))} + \|v_0\|_{B_{\frac{6}{5}, \frac{4}{3}}^{\frac{1}{2}}(\Omega)} \right).$$

Here $C(\Omega)$ represent general constants depending only on Ω , and $v_0 = v|_{t=0}$. For the forcing term,

$$\begin{aligned} \|v \cdot \nabla v\|_{L^{\frac{4}{3}}(0, \nu T; L^{\frac{6}{5}}(\Omega))} &\leq \|v\|_{L^4(0, \nu T, L^3(\Omega))} \|\nabla v\|_{L^2((0, \nu T) \times \Omega)} \\ &\leq \|v\|_{L^\infty(0, \nu T; L^2(\Omega))}^{\frac{1}{2}} \|v\|_{L^2(0, \nu T; L^6(\Omega))}^{\frac{1}{2}} \|\nabla v\|_{L^2((0, \nu T) \times \Omega)} \\ &\leq C(\Omega) \|v\|_{L^\infty(0, \nu T; L^2(\Omega))}^{\frac{1}{2}} \|\nabla v\|_{L^2((0, \nu T) \times \Omega)}^{\frac{3}{2}}, \end{aligned}$$

where in the last step we used Sobolev embedding in $\Omega \subset \mathbb{R}^3$. For the initial value, we use Besov embedding and interpolation so

$$\begin{aligned} \|v_0\|_{B_{\frac{6}{5}, \frac{4}{3}}^{\frac{1}{2}}(\Omega)} &\leq C(\Omega) \|v_0\|_{B_{\frac{4}{3}, \frac{4}{3}}^{\frac{1}{2}}(\Omega)} \leq C(\Omega) \|v_0\|_{H^{\frac{1}{2}}(\Omega)} \\ &\leq C(\Omega) \|v_0\|_{L^2(\Omega)}^{\frac{1}{2}} \|v_0\|_{H^1(\Omega)}^{\frac{1}{2}}. \end{aligned}$$

By the Sobolev embedding and the trace theorem in Ω ,

$$\begin{aligned} \|\nabla v\|_{L^{\frac{4}{3}}((0, \nu T) \times \partial\Omega)} &\leq C(\Omega) \|\nabla v\|_{L^{\frac{4}{3}}(0, \nu T; W^{\frac{1}{6}, \frac{6}{5}}(\partial\Omega))} \\ &\leq C(\Omega) \|\nabla v\|_{L^{\frac{4}{3}}(0, \nu T; W^{1, \frac{6}{5}}(\Omega))} \\ &\leq C(\Omega) \|v\|_{L^{\frac{4}{3}}(0, \nu T; W^{2, \frac{6}{5}}(\Omega))}. \end{aligned}$$

Combining the above estimates, we have

$$\begin{aligned} \|\nabla v\|_{L^{\frac{4}{3}}((0,\nu T)\times\partial\Omega)} &\lesssim \|v\|_{L^{\infty}(0,\nu T;L^2(\Omega))}^{\frac{1}{2}} \left(\|\nabla v\|_{L^2((0,\nu T)\times\Omega)}^{\frac{3}{2}} + \|v_0\|_{H^1(\Omega)}^{\frac{1}{2}} \right) \\ &\quad + \|g\|_{L^{\frac{4}{3}}(0,\nu T;L^{\frac{6}{5}}(\Omega))}. \end{aligned}$$

Noting the scaling of the v and g , we have for any $p \in [1, \infty]$ and any norm X

$$\|u^\nu\|_{L^p(0,T;X)} = \nu^{1-\frac{1}{p}} \|v\|_{L^p(0,\nu T;X)}, \quad \|f^\nu\|_{L^p(0,T;X)} = \nu^{2-\frac{1}{p}} \|g\|_{L^p(0,\nu T;X)}.$$

By this scaling, we have the corresponding estimates on u^ν as

$$\begin{aligned} \nu^{-\frac{1}{4}} \|\nabla u^\nu\|_{L^{\frac{4}{3}}((0,T)\times\partial\Omega)} &\lesssim \nu^{-\frac{5}{4}} \|f^\nu\|_{L^{\frac{4}{3}}(0,T;L^{\frac{6}{5}}(\Omega))} + \\ &\quad + \nu^{-\frac{1}{2}} \|u^\nu\|_{L^{\infty}(0,T;L^2(\Omega))}^{\frac{1}{2}} \left(\nu^{-\frac{3}{4}} \|\nabla u^\nu\|_{L^2((0,T)\times\Omega)}^{\frac{3}{2}} + \nu^{-\frac{1}{2}} \|u^\nu(0)\|_{H^1(\Omega)}^{\frac{1}{2}} \right). \end{aligned}$$

This completes the proof of the proposition. \square

In the inviscid limit $\nu \rightarrow 0$, the main term $\int_{(0,T)\times\Omega} |\nabla u^\nu|^2 \, dx \, dt$ cannot be uniformly bounded, and the force $\|f^\nu\|_{L^{\frac{4}{3}}(0,T;L^{\frac{6}{5}}(\Omega))}$ does not vanish. Therefore, we need to look for another bound that does not degenerate in the inviscid limit.

3.2. Local estimate for the linear Stokes system

To overcome the degeneracy of the naïve bound in the inviscid limit, we show an improved bound in the next subsection, which is based on the following linear estimates for the Stokes system at the unit scale and unit viscosity.

Proposition 2. *Let $C_2 \in \mathcal{C}^{(2)}$ with base T_2 , and denote $Q_2 = (-4, 0) \times C_2$, $\bar{Q}_2 = (-4, 0) \times T_2$. Suppose $u \in L^2(-4, 0; H^1(C_2))$ is a solution to the following Stokes system with forcing term $f \in L^1(-4, 0; L^{\frac{6}{5}}(C_2))$:*

$$\begin{cases} \partial_t u + \nabla P = \Delta u + f & \text{in } Q_2 \\ \operatorname{div} u = 0 & \text{in } Q_2 \\ u = 0 & \text{on } \bar{Q}_2. \end{cases} \quad (6)$$

Then the average vorticity on the boundary is bounded by

$$\begin{aligned} \left| \int_{\bar{Q}_1} \omega(t, x', 0) \, dx' \, dt \right| &\leq \int_{T_1} \left| \int_{-1}^0 \omega(t, x', 0) \, dt \right| \, dx' \\ &\leq C \left(\|\nabla u\|_{L_t^2 L_x^2(Q_2)} + \|f\|_{L_t^1 L_x^{\frac{6}{5}}(Q_2)} \right). \end{aligned}$$

Proof The proof is the same as the one in [1], with only some mild modifications to resolve the curved boundary issue. Without loss of generality, assume by linearity that $\|\nabla u\|_{L_t^2 L_x^2(Q_2)}, \|f\|_{L_t^1 L_x^{\frac{6}{5}}(Q_2)} \leq 1$.

For $t \in (-3, 0)$, $x \in C_2$, we define

$$U(t, x) = \int_{t-1}^t u(s, x) \, ds.$$

Denote $\rho(t) = \mathbf{1}_{[0,1]}(t)$. Then $U = u *_t \rho$, where $*_t$ stands for convolution in t variable only. If we denote $Q = P *_t \rho$, and $F = f *_t \rho$, then U satisfies the Stokes system:

$$\begin{cases} \partial_t U + \nabla Q = \Delta U + F & \text{in } (-3,0) \times C_2 \\ \operatorname{div} U = 0 & \text{in } (-3,0) \times C_2 \\ U = 0 & \text{on } (-3,0) \times T_2 \end{cases}.$$

We have via Sobolev embedding [Lemma 1](#) and $u|_{T_2} = 0$ that

$$\|u\|_{L_t^2 L_x^6(Q_2)} \leq C. \quad (7)$$

Since $\partial_t U(t, x) = u(t, x) - u(t-1, x)$, we have

$$\|\partial_t U\|_{L_t^2 L_x^6((-3,0) \times C_2)} \leq C.$$

On the other hand, the Laplacian of U is bounded by

$$\|\Delta U\|_{L_t^\infty H_x^{-1}((-3,0) \times C_2)} \leq C \|\Delta u\|_{L_t^2 H_x^{-1}(Q_2)} \leq C \|\nabla u\|_{L^2(Q_2)} \leq C.$$

Note that the Sobolev constants depend on the geometry of C_2 . However, they are uniformly bounded as long as $C_2 \in \mathcal{C}^{(2)}$, since the Lipschitz norms of the boundary are uniformly bounded. Again by convolution, we bound F by

$$\|F\|_{L_t^\infty H_x^{-1}((-3,0) \times C_2)} \leq C \|F\|_{L_t^\infty L_x^{\frac{6}{5}}((-3,0) \times C_2)} \leq C.$$

Next, we estimate Q . Using $\nabla Q = \Delta U + F - \partial_t U$ we have

$$\|\nabla Q\|_{L_t^2 H_x^{-1}((-3,0) \times C_2)} \leq C.$$

Without loss of generality, we assume that the average of Q is zero at every t . Then by Nečas theorem (see [\[32\]](#), Section 1.4),

$$\|Q\|_{L_{t,x}^2((-3,0) \times T_2)} \leq C.$$

Note that the constant of Nečas theorem also depends on the Lipschitz norm of ∂C_1 , which is uniform for all $C_1 \in \mathcal{C}^{(1)}$.

By [Lemma 2](#), we can split $U = U_1 + U_2$, where for any $p < \infty$, we have

$$\|\partial_t U_1 + |\nabla^2 U_1|\|_{L_t^p L_x^{\frac{6}{5}}(Q_1)} + \|\partial_t U_2 + |\nabla^2 U_2|\|_{L_t^2 L_x^p(Q_1)} \leq C(p).$$

Denote $\Omega(t, z) := \int_{T_1} |\nabla U(t, x' - zn(x'))| dx'$. Then

$$|\partial_z \Omega| \leq C \int_{T_1} |\nabla^2 U(t, x' - zn(x'))| dx'.$$

Since $\nabla^2 U$ is in $L_t^2 L_z^p + L_t^p L_z^{\frac{6}{5}}(Q_1)$, $\partial_z \Omega$ is bounded in

$$\partial_z \Omega \in L_t^2 L_z^p + L_t^p L_z^{\frac{6}{5}}((-1, 0) \times (0, 1))$$

for any $p < \infty$. Note that

$$|\partial_t \Omega| \leq C \int_{T_1} |\nabla u(t, x' - zn(x'))| dx' \in L_{t,z}^2((-1, 0) \times (0, 1)).$$

Since by interpolation, $L_t^1 L_z^\infty \cap L_t^\infty L_z^1 \subset L_{t,z}^2$, by duality $\partial_t \Omega$ is bounded in $L_{t,z}^2 \subset L_t^1 L_z^\infty + L_t^\infty L_z^1$. Similarly, $\partial_z \Omega$ is bounded in

$$\partial_z \Omega \in L_t^2 L_z^p + L_t^p L_z^{\frac{6}{5}}((-1, 0) \times (0, 1)) \subset L_t^r L_z^\infty + L_t^\infty L_z^r((-1, 0) \times (0, 1))$$

for some $p > 6$ with $r > 1$ sufficiently small. Now we can use [1, Lemma 2.4] to show Ω is continuous up to the boundary with oscillation bounded by

$$\|\Omega\|_{\text{osc}((-1, 0) \times (0, 1))} \leq C.$$

Since the average of Ω is also bounded as

$$\int \Omega \, dz \, dt \leq C \int_{Q_1} |\nabla u| \, dx \, dt \leq C,$$

we have Ω is bounded in L^∞ , in particular

$$\int_{T_1} \left| \int_{-1}^0 \nabla u(t, x', 0) \, dt \right| \, dx' = \Omega(0, 0) \leq C.$$

This concludes the proof of this proposition. \square

3.3. Refined global estimate

Now we are ready to prove the main boundary vorticity estimate.

Proof of Theorem 3 The proof can be divided into four steps. In the first step, we triangularize $\partial\Omega$ and obtain a coarse partition $(0, T) \times \partial\Omega$. Next, we construct σ -algebra \mathcal{F} , which is generated by a finer partition of $(0, T) \times \partial\Omega$, by introducing a suitability criterion. Then we verify that in each piece of the partition, average boundary vorticity is controlled by the maximal function of the energy dissipation and the external force. Finally, we estimate the $L^{\frac{4}{3}}$ weak norm of the averaged vorticity function.

Up to rescaling $u^\nu(t, x) = \nu u(\nu t, x)$ and $f^\nu = \nu^2 f(\nu t, x)$, we assume $\nu = 1$ first and drop the superscript for simplicity.

Step 1. First, we introduce an initial partition of $(0, T) \times \Omega$ as follows. Select $L_0 = 4^{-K}T$, where $K = (\lceil \log_4 \left(\frac{T}{\delta^2} \right) \rceil)_+$ is the smallest nonnegative integer such that $L_0 = 4^{-K}T \leq \delta^2$. Set $r_0 = \frac{1}{2}\sqrt{L_0} = 2^{-K-1}\sqrt{T} \leq \delta$. Then

$$r_0 \leq \frac{1}{2} \min \left\{ \delta, \sqrt{T} \right\} < 2r_0.$$

Let $\{\mathsf{T}_{r_0}^{(i)}\}_i \subset \mathcal{T}^{(r_0)}$ be a partition of $\partial\Omega$ with size r_0 , as specified in Assumption 1. Then

$$(0, T) \times \partial\Omega \simeq \bigsqcup_{j=1}^{4^K} \bigsqcup_i \bar{\mathsf{Q}}^{(i,j)} \quad \text{where } \bar{\mathsf{Q}}^{(i,j)} = ((j-1)L_0, jL_0) \times \mathsf{T}_{r_0}^{(i)}.$$

We denote $\bar{\mathcal{Q}}_0 = \{\bar{\mathsf{Q}}^{(i,j)}\}_{i,j}$. By part (5) of Section 2.2, each $\bar{\mathsf{Q}}^{(i,j)}$ admits a sequence of dyadic decomposition. For $k \geq 1$, denote $\bar{\mathcal{Q}}_k$ to be the set of dyadic decompositions of spacetime curved triangles in $\bar{\mathcal{Q}}_{k-1}$. Then any $\bar{\mathsf{Q}} \in \bar{\mathcal{Q}}_k$ is a Cartesian product of curved triangles of size $r_k := 2^{-k}r_0$ in space and length $r_k^2 = 4^{-k}r_0^2$ in time.

Step 2. The next goal is to find a partition of $(0, T) \times \Omega$ consisting of “suitable” cubes, defined as follows. Let $\bar{Q} = (\hat{t} - r_k^2, \hat{t}) \times T_{r_k} \in \bar{\mathcal{Q}}_k$ for some $\hat{t} \in (0, T]$ and $T_{r_k} \in \mathcal{T}^{(r_k)}$. Denote \hat{x} to be the barycenter of T_{r_k} . We say \bar{Q} is **suitable** if both $\hat{t} \geq 4r_k^2$ and

$$\int_{\hat{t}-4r_k^2}^{\hat{t}} \int_{\partial\Omega \cap B_{2r_k}(\hat{x})} \int_0^{2r_k} \left(|\nabla u|^2 + |f|^{\frac{4}{3}} \right) (t, x' - zn(x')) dz dx' dt \leq c_0 r_k^{-4}. \quad (\text{S})$$

for some c_0 to be determined. Recall $n(x')$ is the outer normal vector at $x' \in \partial\Omega$.

Now we construct a partition according to suitability. Denote $\mathcal{S}_0 \subset \bar{\mathcal{Q}}_0$ to be the set of suitable cubes, $\mathcal{N}_0 = \bar{\mathcal{Q}}_0 \setminus \mathcal{S}_0$ be the set of non-suitable cubes. For $k \geq 1$, we perform a dyadic decomposition on each cube $\bar{Q} \in \mathcal{N}_{k-1}$, then put the suitable ones in \mathcal{S}_k and non-suitable ones in \mathcal{N}_k . This process may continue indefinitely, and we define $\mathcal{S} = \bigcup_k \mathcal{S}_k$ to be the set of suitable cubes that we obtained from this process.

We claim that \mathcal{S} is a partition of $(0, T) \times \partial\Omega$. It is easy to see from our process that cubes in \mathcal{S} are mutually disjoint. Moreover, for almost every $(t, x') \in (0, T) \times \partial\Omega$, the cube whose closure contains (t, x') becomes suitable if the cube is sufficiently small, by a partial regularity argument. Indeed, denote the singular set $\text{Sing}(u, f)$ to be the complement of the closure of $\bigcup_{\bar{Q} \in \mathcal{S}} \bar{Q}$ in $(0, T) \times \partial\Omega$. For every $(\hat{t}, \hat{x}') \in \text{Sing}(u, f)$, for every $k > 0$, there exists a cube $\bar{Q}_k \in \mathcal{N}_k$ such that \bar{Q}_k fails the suitability condition (S). Then we find a neighborhood of (\hat{t}, \hat{x}') in $(0, T) \times \Omega$ which is

$$U = \{(t, x' - zn(x')) : t \in (\hat{t} - 4r_k^2, \hat{t}), x' \in \partial\Omega \cap B_{2r_k}(\hat{x}), z \in (0, 2r_k)\},$$

such that $\int_U |\nabla u|^2 + |f|^{\frac{4}{3}} dx dt \gtrsim r_k$. Moreover, this neighborhood is comparable with a parabolic cylinder of radius r_k . These neighborhoods form an open cover of $\text{Sing}(u, f)$. By Vitali covering lemma, we find a disjoint subcollection U_i which covers $\text{Sing}(u, f)$ if dilating by a factor of 5. The radii are summable because $\sum_i r_k \lesssim \sum_i \int_{U_i} |\nabla u|^2 + |f|^{\frac{4}{3}} dx dt < \infty$, so the parabolic Hausdorff dimension of $\text{Sing}(u, f)$ is at most 1.

Define $\mathcal{F} = \sigma(\mathcal{S})$ to be the σ -algebra generated by these countably many suitable cubes. Then the conditional expectation $\tilde{\omega} := \mathbb{E}[\omega | \mathcal{F}]$ is simply a piecewise function, taking the average value of ω on each $\bar{Q} \in \mathcal{S}$.

Next we prove claim (1) and (2). First, we show the set $A = (0, 4^{-l-1}) \times \partial\Omega$ is \mathcal{F} -measurable. If $4^{-l}T \leq \delta^2$ and $l \geq 0$, then $4^{-l-1}T \leq \frac{1}{4} \min\{\delta^2, T\} < 4r_0^2$. Hence $4^{-l-1}T = 4^{-l-1} \cdot 4^{K+1}r_0^2 = 4^{-(l+1-K)} \cdot 4r_0^2 = 4r_{k'}^2$ for some $k' > 0$, and $A = (0, 4r_{k'}^2) \times \partial\Omega$. On the one hand, for $k < k'$, \mathcal{S}_k only contains cubes of the form $(\hat{t} - r_k^2, \hat{t}) \times T_{r_k}$ with $\hat{t} - r_k^2 \geq 3r_k^2 > 4r_{k'}^2$, so A is disjoint from every cube in \mathcal{S}_k . On the other hand, for $k \geq k'$, $\mathcal{S}_k \subset \bar{\mathcal{Q}}_k$ only contains cubes of the form $(jr_k^2, (j+1)r_k^2) \times T_{r_k}$. Since $4r_{k'}^2 = 4^{k-k'+1}r_k^2$, each cube in \mathcal{S}_k is either contained in A or disjoint from A . In conclusion, every set in \mathcal{S} is either a subset of A or a subset of $(0, T) \times \partial\Omega \setminus A$, hence $A \in \sigma(\mathcal{S}) = \mathcal{F}$.

To prove (2), note that each \bar{Q} in \mathcal{S} has size at most $r_0 < \delta$ in space and $r_0^2 < \delta^2$ in time, so for $(t_1, x_1), (t_2, x_2) \in \bar{Q}$,

$$\begin{aligned} |\varphi(t_1, x_1) - \varphi(t_2, x_2)| &\leq |\varphi(t_1, x_1) - \varphi(t_2, x_1)| + |\varphi(t_2, x_1) - \varphi(t_2, x_2)| \\ &\leq \|\partial_t \varphi\|_{L^\infty} |t_1 - t_2| + \|\nabla \varphi\|_{L^\infty} |x_1 - x_2| \\ &\leq \delta^2 \|\partial_t \varphi\|_{L^\infty} + \delta \|\nabla \varphi\|_{L^\infty}. \end{aligned}$$

Step 3. Take any cube $\bar{Q} \in \mathcal{S}_k$. By using the canonical scaling of the Navier-Stokes equation $u_r(t, x) := ru(r^2t + \hat{t}, rx)$ and $f_r(t, x) := r^3f(r^2t + \hat{t}, rx)$ with size $r = r_k$, u_r solves the Stokes equation (6) in $(-4, 0) \times C_2$ with some $C_2 \in \mathcal{C}^{(2)}$ and force term $f_r - u_r \cdot \nabla u_r$, and (S) implies

$$\begin{aligned} \|\nabla u_r\|_{L^2((-4,0) \times C_2)} &\leq c_0^{\frac{1}{2}}, \\ \|f_r\|_{L^1(-4,0; L^{\frac{6}{5}}(C_2))} &\leq \|f_r\|_{L^{\frac{4}{3}}((-4,0) \times C_2)} \lesssim c_0^{\frac{3}{4}}, \\ \|u_r \cdot \nabla u_r\|_{L^1(-4,0; L^{\frac{6}{5}}(C_2))} &\lesssim \|\nabla u_r\|_{L^2((-4,0) \times C_2)} \|u_r\|_{L^2(-4,0; L^6(C_2))} \lesssim c_0. \end{aligned}$$

In the last step we used the Sobolev embedding

$$\|u_r\|_{L^2(-4,0; L^6(C_2))} \lesssim \|\nabla u_r\|_{L^2((-4,0) \times C_2)},$$

when $u_r = 0$ on the base T_2 . Therefore, [Proposition 2](#) implies that after scaling, the average vorticity is bounded by

$$|\tilde{\omega}|_{\bar{Q}} := \left| \fint_{\bar{Q}} \omega(t, x') \, dx' \, dt \right| \leq \frac{1}{16} \gamma r_k^{-2},$$

where we choose $c_0 = \frac{1}{C} \gamma^2 \leq 1$.

Next, we separate two scenarios, $k = 0$ and $k > 0$. If $k = 0$, then for any $\bar{Q} \in \mathcal{S}_0$, for any $0 < t < T$,

$$|\tilde{\omega}|_{\bar{Q}} \leq \frac{1}{16} \gamma r_0^{-2} < \gamma \max \{ \delta^{-2}, T^{-1} \}.$$

If $k > 0$, then $\bar{Q} \in \mathcal{S}_k$ has an antecedent cube $\bar{P} \in \mathcal{N}_{k-1}$. Cube $\bar{P} = (\hat{t} - r_{k-1}^2, \hat{t}) \times T_{r_{k-1}}$ is not suitable, so either of the following two cases must be true.

1. $\hat{t} < 4r_{k-1}^2$. In this case, for any $(t, x) \in \bar{Q} \subset \bar{P}$,

$$|\tilde{\omega}|_{\bar{Q}} \leq \frac{1}{16} \gamma r_k^{-2} = \frac{1}{4} \gamma r_{k-1}^{-2} \leq \gamma \hat{t}^{-1} \leq \frac{\gamma}{t}.$$

2. $\hat{t} \geq 4r_{k-1}^2$, but

$$\fint_{\hat{t}-4r_{k-1}^2}^{\hat{t}} \fint_{\partial\Omega \cap B_{2r_{k-1}}(\hat{x})} \fint_0^{2r_{k-1}} \left(|\nabla u|^2 + |f|^{\frac{4}{3}} \right) (t, x' - zn(x')) \, dz \, dx' \, dt > c_0 r_{k-1}^{-4}.$$

In the latter case, note that the integral region is comparable to Q , the extension of \bar{Q} , which is contained in $(0, T) \times \mathcal{U}_\delta(\partial\Omega, \Omega)$. We then know that for any $(t, x) \in Q$, the parabolic maximal function is bounded from below by

$$\begin{aligned} M(t, x) &:= \mathcal{M}((|\nabla u|^2 + |f|^{\frac{4}{3}}) \mathbf{1}_{[0, T] \times \mathcal{U}_\delta(\partial\Omega, \Omega)})(t, x) \\ &:= \sup_{r > 0} \fint_{t-r^2}^{t+r^2} \fint_{B_r(x)} \left(|\nabla u|^2 + |f|^{\frac{4}{3}} \right) (s, y) \mathbf{1}_{[0, T] \times \mathcal{U}_\delta(\partial\Omega, \Omega)}(s, y) \, dy \, ds \\ &\geq \frac{1}{C} c_0 r_k^{-4} = \frac{\gamma^2}{C} r_k^{-4}. \end{aligned}$$

Note that the parabolic maximal function \mathcal{M} is a bounded map from $L^1(\mathbb{R} \times \mathbb{R}^3)$ to $L^{1,\infty}(\mathbb{R} \times \mathbb{R}^3)$.

In summary, for any $\bar{Q} \in \mathcal{S}_k$, we have $|\tilde{\omega}|_{\bar{Q}} \leq \frac{1}{16}\gamma r_k^{-2} \leq \gamma r_k^{-2}$, and

$$\text{either } |\tilde{\omega}|_{\bar{Q}} \leq \gamma \max \left\{ \frac{1}{t}, \frac{1}{\delta^2} \right\} \quad \text{or } M|_{\bar{Q}} \geq \frac{\gamma^2}{C} r_k^{-4}.$$

Step 4. Denote $\mathcal{S} = \{\bar{Q}^{(i)}\}$ and let $\bar{Q}^{(i)}$ has size $r^{(i)}$. For any $r_\star = 2^l r_0$ with $l \in \mathbb{Z}$, we have

$$\begin{aligned} & \left\{ (t, x') \in (0, T) \times \partial\Omega : |\tilde{\omega}| > \gamma \max \{r_\star^{-2}, t^{-1}, \delta^{-2}\} \right\} \\ & \subset \bigcup_i \left\{ \bar{Q}^{(i)} : r^{(i)} < r_\star, M|_{\bar{Q}^{(i)}} \geq \frac{\gamma^2}{C} (r^{(i)})^{-4} \right\} \\ & \subset \bigcup_i \bigcup_{k=1}^{\infty} \left\{ \bar{Q}^{(i)} : r^{(i)} = 2^{-k} r_\star, M|_{\bar{Q}^{(i)}} \geq \frac{\gamma^2}{C} (2^{-k} r_\star)^{-4} \right\}. \end{aligned}$$

Therefore the measure of the upper level set is controlled by the total measure of these suitable cubes, that is

$$\begin{aligned} & \left| \left\{ |\tilde{\omega}| > \gamma \max \{r_\star^{-2}, t^{-1}, \delta^{-2}\} \right\} \right| \\ & \leq \sum_{k=1}^{\infty} \sum_i \left\{ \left| \bar{Q}^{(i)} \right| : r^{(i)} = 2^{-k} r_\star, M|_{\bar{Q}^{(i)}} \geq \frac{\gamma^2}{C} (2^{-k} r_\star)^{-4} \right\} \\ & = \sum_{k=1}^{\infty} \frac{2^k}{r_\star} \sum_i \left\{ \left| \bar{Q}^{(i)} \right| : r^{(i)} = 2^{-k} r_\star, M|_{\bar{Q}^{(i)}} \geq \frac{\gamma^2}{C} (2^{-k} r_\star)^{-4} \right\} \\ & \lesssim \sum_{k=1}^{\infty} \frac{2^k}{r_\star} \left| \left\{ (t, x) \in (0, T) \times \Omega : M(t, x) \geq \frac{\gamma^2}{C} (2^{-k} r_\star)^{-4} \right\} \right| \\ & \lesssim \sum_{k=1}^{\infty} \frac{2^k}{\frac{\gamma^2}{C} r_\star} \|M\|_{L^{1,\infty}((0,T) \times \Omega)} (2^{-k} r_\star)^4 \\ & \lesssim \gamma^{-2} \left\| |\nabla u|^2 + |f|^{\frac{4}{3}} \right\|_{L^1((0,T) \times \mathcal{U}_\delta(\partial\Omega, \Omega))} r_\star^3 \\ & = \gamma^{-\frac{1}{2}} \left\| |\nabla u|^2 + |f|^{\frac{4}{3}} \right\|_{L^1((0,T) \times \mathcal{U}_\delta(\partial\Omega, \Omega))} (\gamma r_\star^{-2})^{-\frac{3}{2}}. \end{aligned}$$

This is true for any $r_\star = 2^l r_0$. By the definition of Lorentz space, for every $\gamma \leq 1$ we have

$$\begin{aligned} & \left\| \tilde{\omega} \mathbf{1}_{\{|\tilde{\omega}| > \gamma \max \left\{ \frac{1}{t}, \frac{1}{\delta^2} \right\} \}} \right\|_{L^{\frac{3}{2}, \infty}((0,T) \times \partial\Omega)}^{\frac{3}{2}} \\ & \lesssim \gamma^{-\frac{1}{2}} \left(\|\nabla u\|_{L^2((0,T) \times \mathcal{U}_\delta(\partial\Omega, \Omega))}^2 + \|f\|_{L^{\frac{4}{3}}((0,T) \times \mathcal{U}_\delta(\partial\Omega, \Omega))}^{\frac{4}{3}} \right). \end{aligned}$$

This completes the proof of the theorem with $\nu = 1$, and for general $\nu > 0$ the conclusion follows by scaling. \square

4. Proof of the main result

In this section, we first derive an estimate for the pairing between boundary vorticity with any C^1 vector field, which is the work done by the friction force, then apply this to estimate the layer separation.

Corollary 2. Let $\Omega \subset \mathbb{R}^3$ be a smooth bounded domain satisfying [Assumption 1](#) with $\bar{\delta}$. There exists a constant $C(\Omega) > 0$ depending only on Ω and a universal constant C such that the following holds. Given $T > 0$, $A > 0$, $L > 0$, suppose φ is a C^1 velocity field defined on $(0, T) \times \partial\Omega$, satisfying

$$\|\varphi\|_{L^\infty((0,T)\times\partial\Omega)}, \frac{L}{A} \|\partial_t \varphi\|_{L^\infty((0,T)\times\partial\Omega)}, L \|\nabla \varphi\|_{L^\infty((0,T)\times\partial\Omega)} \leq A. \quad (8)$$

Given any weak solution $u^\nu \in C_w(0, T; L^2(\Omega)) \cap L^2(0, T; H_0^1(\Omega))$ to [\(NSE_v\)](#) with initial value $u^\nu(0) \in H^1(\Omega)$ and force $f^\nu \in L^{\frac{4}{3}}((0, T) \times \Omega)$, denote

$$\begin{aligned} E_\nu &:= \|u^\nu\|_{L^\infty(0,T;L^2(\Omega))}^2, & D_\nu &:= \nu \|\nabla u^\nu\|_{L^2((0,T)\times\Omega)}^2, \\ H_\nu &:= \|u^\nu(0)\|_{H^1(\Omega)}^2, & F_\nu &:= \nu^{\frac{1}{3}} \|f^\nu\|_{L^{\frac{4}{3}}((0,T)\times\Omega)}^{\frac{4}{3}}. \end{aligned}$$

Then the vorticity ω^ν satisfies

$$\begin{aligned} \left| \nu \int_0^T \int_{\partial\Omega} \omega^\nu \cdot \varphi \, dx' \, dt \right| &\leq CA^3 T |\partial\Omega| + \frac{1}{4} D_\nu + \frac{1}{4} F_\nu + \nu^{\frac{4}{3}} H_\nu^{\frac{1}{3}} \\ &+ \left(4 \log \left(\frac{4AL}{\nu} \right)_+ + \frac{\nu T}{\bar{\delta}^2} + \frac{C(\Omega)(1+\nu^2)E_\nu T}{AL^4} \right) A \nu |\partial\Omega|. \end{aligned}$$

Proof For some $\delta \leq \bar{\delta}$ to be determined later, let \mathcal{F} be the σ -algebra introduced in [Theorem 3](#). For some $T_\nu = 4^{-k}T$ with k to be determined later, we compute the integral by

$$\begin{aligned} \nu \int_0^T \int_{\partial\Omega} \omega^\nu \cdot \varphi \, dx' \, dt &= \nu \int_0^{T_\nu} \int_{\partial\Omega} \omega^\nu \cdot \varphi \, dx' \, dt \\ &+ \nu \int_{T_\nu}^T \int_{\partial\Omega} (\omega^\nu - \mathbb{E}[\omega^\nu | \mathcal{F}]) \cdot \varphi \, dx' \, dt \\ &+ \nu \int_{T_\nu}^T \int_{\partial\Omega} \mathbb{E}[\omega^\nu | \mathcal{F}] \cdot \varphi \, dx' \, dt \\ &= \text{I} + \text{II} + \text{III}. \end{aligned}$$

We start with the second term. Note that since $T_\nu = 4^{-k}T$, $(T_\nu, T) \times \partial\Omega$ is a \mathcal{F} -measurable set, so

$$\int_{T_\nu}^T \int_{\partial\Omega} (\omega^\nu - \mathbb{E}[\omega^\nu | \mathcal{F}]) \cdot \varphi \, dx' \, dt = \int_{T_\nu}^T \int_{\partial\Omega} \omega^\nu \cdot (\varphi - \mathbb{E}[\varphi | \mathcal{F}]) \, dx' \, dt.$$

Therefore,

$$\begin{aligned} |\text{I} + \text{II}| &= \int_0^T \int_{\partial\Omega} \nu \omega^\nu \cdot (\varphi \mathbf{1}_{\{t \leq T_\nu\}} + (\varphi - \mathbb{E}[\varphi | \mathcal{F}]) \mathbf{1}_{\{t \geq T_\nu\}}) \, dx' \, dt \\ &\leq \|\nu \omega^\nu\|_{L^{\frac{4}{3}}((0,T)\times\partial\Omega)} \|\varphi \mathbf{1}_{\{t \leq T_\nu\}} + (\varphi - \mathbb{E}[\varphi | \mathcal{F}]) \mathbf{1}_{\{t \geq T_\nu\}}\|_{L^4((0,T)\times\partial\Omega)}. \end{aligned}$$

By assumption (8) on φ and (4) of [Theorem 3](#),

$$\begin{aligned} \|\varphi\|_{L^4((0,T_\nu)\times\partial\Omega)}^4 &\leq A^4 T_\nu |\partial\Omega|, \\ \|\varphi - \mathbb{E}[\varphi | \mathcal{F}]\|_{L^4((0,T)\times\partial\Omega)}^4 &\leq \left[\delta \left(\frac{\delta A^2}{\nu} + \frac{A}{L} \right) \right]^4 T |\partial\Omega|. \end{aligned}$$

Hence, by choosing $\delta = \min\{\bar{\delta}, \frac{\nu}{A}\}$ and choosing T_ν to satisfy

$$\frac{1}{4} T \min\left\{\frac{\nu}{AL}, 1\right\}^4 \leq T_\nu \leq T \min\left\{\frac{\nu}{AL}, 1\right\}^4,$$

we can bound

$$\|\varphi \mathbf{1}_{\{t \leq T_\nu\}} + (\varphi - \mathbb{E}[\varphi | \mathcal{F}]) \mathbf{1}_{\{t \geq T_\nu\}}\|_{L^4((0, T) \times \partial\Omega)} \leq \frac{\nu}{L} (T|\partial\Omega|)^{\frac{1}{4}}.$$

As for the $L^{\frac{4}{3}}$ norm of $\nu\omega^\nu$, we use the global linear estimate [Proposition 1](#):

$$\|\nu\omega^\nu\|_{L^{\frac{4}{3}}((0, T) \times \partial\Omega)}^{\frac{4}{3}} \leq C(\Omega) \left[F_\nu + E_\nu^{\frac{1}{3}} \left(\nu^{-1} D_\nu + \nu^{\frac{1}{3}} H_\nu^{\frac{1}{3}} \right) \right].$$

Here we used $\|f^\nu(t)\|_{L^{\frac{6}{5}}(\Omega)} \leq \|f^\nu(t)\|_{L^{\frac{4}{3}}(\Omega)} |\Omega|^{\frac{1}{12}}$. Combined we can bound the first two terms by

$$\begin{aligned} |\text{I} + \text{II}| &\leq C(\Omega) \frac{\nu}{L} \left[\nu^{-\frac{1}{3}} F_\nu + E_\nu^{\frac{1}{3}} \left(\nu^{-1} D_\nu + \nu^{\frac{1}{3}} H_\nu^{\frac{1}{3}} \right) \right]^{\frac{3}{4}} (T|\partial\Omega|)^{\frac{1}{4}} \\ &\leq C(\Omega) \left[\nu^{\frac{1}{2}} F_\nu^{\frac{3}{4}} + E_\nu^{\frac{1}{4}} \left(D_\nu^{\frac{3}{4}} + \nu H_\nu^{\frac{1}{4}} \right) \right] \frac{(\nu T|\partial\Omega|)^{\frac{1}{4}}}{L}. \end{aligned}$$

For the third term, denote $\tilde{\omega}^\nu = \mathbb{E}[\omega^\nu | \mathcal{F}]$. Then

$$\begin{aligned} |\text{III}| &\leq \left| \int_{(T_\nu, T) \times \partial\Omega} \nu \tilde{\omega}^\nu \mathbf{1}_{\{\nu|\tilde{\omega}^\nu| > \gamma \max\left\{\frac{\nu}{t}, \frac{\nu^2}{\delta^2}\right\}\}} \cdot \varphi \, dx' \, dt \right| \\ &\quad + \gamma \int_{(T_\nu, T) \times \partial\Omega} \frac{\nu}{t} |\varphi| \, dx' \, dt + \gamma \int_{(T_\nu, T) \times \partial\Omega} \frac{\nu^2}{\delta^2} |\varphi| \, dx' \, dt \\ &\leq \left\| \nu \tilde{\omega}^\nu \mathbf{1}_{\{\nu|\tilde{\omega}^\nu| > \gamma \max\left\{\frac{\nu}{t}, \frac{\nu^2}{\delta^2}\right\}\}} \right\|_{L^{\frac{3}{2}, \infty}((0, T) \times \partial\Omega)} \|\varphi\|_{L^{3,1}((0, T) \times \partial\Omega)} \\ &\quad + A\gamma \left(\nu|\partial\Omega| \log\left(\frac{T}{T_\nu}\right) + \nu^2 \delta^{-2} T|\partial\Omega| \right). \end{aligned}$$

Recalling the choice of T_ν and δ , we have

$$\begin{aligned} \nu|\partial\Omega| \log\left(\frac{T}{T_\nu}\right) + \nu^2 \delta^{-2} T|\partial\Omega| \\ \leq 4\nu|\partial\Omega| \log\left(\frac{4AL}{\nu}\right)_+ + \nu^2 \bar{\delta}^{-2} T|\partial\Omega| + A^2 T|\partial\Omega|. \end{aligned}$$

Moreover, by [Theorem 3](#) we control the $L^{\frac{3}{2}}$ weak norm by

$$\left\| \nu \tilde{\omega}^\nu \mathbf{1}_{\{\nu|\tilde{\omega}^\nu| > \gamma \max\left\{\frac{\nu}{t}, \frac{\nu^2}{\delta^2}\right\}\}} \right\|_{L^{\frac{3}{2}, \infty}((0, T) \times \partial\Omega)} \leq C\gamma^{-\frac{1}{3}} (D_\nu + F_\nu)^{\frac{2}{3}}. \quad (9)$$

And $\|\varphi\|_{L^{3,1}((0, T) \times \partial\Omega)} \leq A(T|\partial\Omega|)^{\frac{1}{3}}$. Hence

$$\begin{aligned} |\text{III}| &\leq CA(T|\partial\Omega|)^{\frac{1}{3}} \gamma^{-\frac{1}{3}} (D_\nu + F_\nu)^{\frac{2}{3}} \\ &\quad + \left(4 \log\left(\frac{4AL}{\nu}\right)_+ + \frac{\nu T}{\bar{\delta}^2} \right) \gamma A \nu |\partial\Omega| + \gamma A^3 T |\partial\Omega|. \end{aligned} \quad (10)$$

In conclusion, we have shown that

$$|\text{I} + \text{II} + \text{III}| \leq CA(T|\partial\Omega|)^{\frac{1}{3}}\gamma^{-\frac{1}{3}}(D_v + F_v)^{\frac{2}{3}} + \gamma A^3 T|\partial\Omega| + R_v,$$

with a remainder

$$\begin{aligned} R_v &= C(\Omega) \left[\nu^{\frac{1}{2}} F_v^{\frac{3}{4}} + E_v^{\frac{1}{4}} \left(D_v^{\frac{3}{4}} + \nu H_v^{\frac{1}{4}} \right) \right] \frac{(\nu T|\partial\Omega|)^{\frac{1}{4}}}{L} \\ &\quad + \left(4 \log \left(\frac{4AL}{\nu} \right)_+ + \frac{\nu T}{\delta^2} \right) A\nu|\partial\Omega|. \end{aligned}$$

Next, we use Young's inequality on each product, so

$$\begin{aligned} CA(T|\partial\Omega|)^{\frac{1}{3}}\gamma^{-\frac{1}{3}}(D_v + F_v)^{\frac{2}{3}} &\leq \frac{1}{8}(D_v + F_v) + \frac{C}{\gamma} A^3 T|\partial\Omega|, \quad (11) \\ C(\Omega)\nu^{\frac{1}{2}} F_v^{\frac{3}{4}} \frac{(\nu T|\partial\Omega|)^{\frac{1}{4}}}{L} &\leq \frac{1}{8}F_v + C(\Omega) \frac{\nu^3 T|\partial\Omega|}{L^4}, \\ C(\Omega)E_v^{\frac{1}{4}} D_v^{\frac{3}{4}} \frac{(\nu T|\partial\Omega|)^{\frac{1}{4}}}{L} &\leq \frac{1}{8}D_v + C(\Omega) \frac{\nu T|\partial\Omega|}{L^4} E_v, \\ C(\Omega)E_v^{\frac{1}{4}} \nu H_v^{\frac{1}{4}} \frac{(\nu T|\partial\Omega|)^{\frac{1}{4}}}{L} &\leq \nu^{\frac{4}{3}} H_v^{\frac{1}{3}} + C(\Omega) \frac{\nu T|\partial\Omega|}{L^4} E_v. \end{aligned}$$

Hence for every $\gamma \leq 1$, we have

$$\begin{aligned} |\text{I} + \text{II} + \text{III}| &\leq \left(\frac{C}{\gamma} + \gamma \right) A^3 T|\partial\Omega| + \frac{1}{4}D_v + \frac{1}{4}F_v + \nu^{\frac{4}{3}} H_v^{\frac{1}{3}} \\ &\quad + \left(4 \log \left(\frac{4AL}{\nu} \right)_+ + \frac{\nu T}{\delta^2} + \frac{C(\Omega)(1 + \nu^2)E_v T}{AL^4} \right) A\nu|\partial\Omega|. \end{aligned}$$

This finishes the proof of the corollary by selecting $\gamma = 1$. \square

To prove the main theorem, we will use the following elementary lemma, which computes the evolution of L^2 distance between a Navier–Stokes weak solution and a smooth vector field.

Lemma 4. *Let $u = u^\nu \in C_w(0, T; L^2(\Omega)) \cap L^2(0, T; H_0^1(\Omega))$ be a weak solution to (NSE_v) with force $f = f^\nu \in L^1(0, T; L^1(\Omega))$, and let v be any C^1 divergence-free flow with $v \cdot n = 0$ on $\partial\Omega$. Then the L^2 inner product (u, v) has the following time derivative:*

$$\frac{d}{dt}(u, v) = \int_{\Omega} u \cdot (\partial_t v + v \cdot \nabla v) + [(u - v) \otimes (u - v)] : Dv \, dx + (\nu \Delta u + f, v)$$

where

$$(\Delta u, v) = \int_{\partial\Omega} \partial_n u \cdot v \, dx' - \int_{\Omega} \nabla u : \nabla v \, dx.$$

If $v = \bar{u}$ solves the Euler equation (EE) with force $\bar{f} \in L^1(0, T; L^2(\Omega))$, then

$$\begin{aligned} \frac{d}{dt}(u, \bar{u}) &= \int_{\Omega} [(u - \bar{u}) \otimes (u - \bar{u}) - v \nabla u] : \nabla \bar{u} + u \cdot \bar{f} + \bar{u} \cdot f \, dx \\ &\quad + v \int_{\partial\Omega} \partial_n u \cdot \bar{u} \, dx'. \end{aligned}$$

In addition, if u is a Leray–Hopf solution with $f \in L^1(0, T; L^2(\Omega))$, then

$$\begin{aligned} & \frac{1}{2} \|u - \bar{u}\|_{L^2(\Omega)}^2(T) - \frac{1}{2} \|u - \bar{u}\|_{L^2(\Omega)}^2(0) + \frac{\nu}{2} \int_{(0,T) \times \Omega} |\nabla u|^2 - |\nabla \bar{u}|^2 \, dx \, dt \\ & \leq \int_0^T \|u - \bar{u}\|_{L^2(\Omega)}^2 \|D\bar{u}\|_{L^\infty(\Omega)} \, dt - \nu \int_{(0,T) \times \partial\Omega} \partial_n u \cdot \bar{u} \, dx' \, dt \\ & \quad + \int_{(0,T) \times \Omega} (u - \bar{u}) \cdot (f - \bar{f}) \, dx \, dt. \end{aligned}$$

Proof For $u \in L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H_0^1(\Omega))$, $\nu \in L^\infty(0, T; L^3(\Omega)) \cap L^2(0, T; H^1(\Omega)) \cap L^1(0, T; W^{1,\infty}(\Omega))$ with $\nu \cdot n = 0$ on $\partial\Omega$, we have

$$\begin{aligned} (v, u \cdot \nabla u) + (u, v \cdot \nabla v) &= (v, u \cdot \nabla(u - v)) + (u - v, v \cdot \nabla v) \\ &= (v, u \cdot \nabla(u - v)) - (v \cdot \nabla(u - v), v) \\ &= (v, (u - v) \cdot \nabla(u - v)) \\ &= (v, \operatorname{div}[(u - v) \otimes (u - v)]) \\ &= - \int_{\Omega} [(u - v) \otimes (u - v)] : \nabla v \, dx \\ &= - \int_{\Omega} [(u - v) \otimes (u - v)] : Dv \, dx. \end{aligned}$$

In the last step, we can replace ∇v by its symmetric part Dv because $(u - v) \otimes (u - v)$ is symmetric.

If $v = \bar{u}$ solves the Euler equation, then $\partial_t v + v \cdot \nabla v = -\nabla p + \bar{f}$, so

$$\begin{aligned} \frac{d}{dt} (u, \bar{u}) &= \int_{\Omega} u \cdot \bar{f} + [(u - \bar{u}) \otimes (u - \bar{u})] : D\bar{u} \, dx \\ & \quad + \nu \int_{\partial\Omega} \partial_n u \cdot v \, dx' - \nu \int_{\Omega} \nabla u : \nabla v \, dx + \int_{\Omega} u \cdot \bar{f} \, dx. \end{aligned}$$

Integrate between 0 and T :

$$\begin{aligned} (u, \bar{u})(T) - (u, \bar{u})(0) &= \int_0^T \int_{\Omega} [(u - \bar{u}) \otimes (u - \bar{u}) - v \nabla u] : \nabla \bar{u} \, dx \, dt \\ & \quad + \nu \int_0^T \int_{\partial\Omega} \partial_n u \cdot \bar{u} \, dx' + \int_0^T \int_{\Omega} \bar{u} \cdot f + u \cdot \bar{f} \, dx \, dt. \end{aligned}$$

Recall the energy inequality of the Leray–Hopf solutions to the Navier–Stokes equation and energy conservation for the Euler equation:

$$\begin{aligned} \frac{1}{2} \|u\|_{L^2(\Omega)}^2(T) + \int_0^T \int_{\Omega} \nu |\nabla u|^2 &\leq \frac{1}{2} \|u\|_{L^2(\Omega)}^2(0) + \int_0^T \int_{\Omega} u \cdot f \, dx \, dt, \\ \frac{1}{2} \|\bar{u}\|_{L^2(\Omega)}^2(T) &= \frac{1}{2} \|\bar{u}\|_{L^2(\Omega)}^2(0) + \int_0^T \int_{\Omega} \bar{u} \cdot \bar{f} \, dx \, dt. \end{aligned}$$

Combined we have

$$\begin{aligned} & \frac{1}{2} \|u - \bar{u}\|_{L^2(\Omega)}^2(T) - \frac{1}{2} \|u - \bar{u}\|_{L^2(\Omega)}^2(0) + \nu \|\nabla u\|_{L^2((0,T) \times \Omega)}^2 \\ & \leq - \int_0^T \int_{\Omega} [(u - \bar{u}) \otimes (u - \bar{u}) - v \nabla u] : \nabla \bar{u} \, dx - \nu \int_0^T \int_{\partial\Omega} \partial_n u \cdot \bar{u} \, dx' \end{aligned}$$

$$\begin{aligned}
& + \int_0^T \int_{\Omega} u f + \bar{u} \bar{f} - u \bar{f} - \bar{u} f \, dx \, dt \\
& \leq \int_0^T \|u - \bar{u}\|_{L^2(\Omega)}^2 \|D\bar{u}\|_{L^\infty(\Omega)} \, dt + \frac{\nu}{2} \|\nabla u\|_{L^2(\Omega)}^2 + \frac{\nu}{2} \|\nabla \bar{u}\|_{L^2(\Omega)}^2 \\
& \quad + \int_0^T \int_{\Omega} (u - \bar{u}) \cdot (f - \bar{f}) \, dx \, dt - \nu \int_0^T \int_{\partial\Omega} \partial_n u \cdot \bar{u} \, dx' \cdot
\end{aligned}$$

This completes the proof of the lemma. \square

Proof of Theorem 2 For any $0 < t < T$, by Lemma 4,

$$\begin{aligned}
& \frac{1}{2} \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(t) + \frac{\nu}{2} \|\nabla u^\nu\|_{L^2((0,t) \times \Omega)}^2 - \frac{\nu}{2} \|\nabla \bar{u}\|_{L^2((0,t) \times \Omega)}^2 \\
& \leq \frac{1}{2} \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(0) + \int_0^t \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2 \|D\bar{u}\|_{L^\infty(\Omega)} \, ds \\
& \quad - \nu \int_0^t \int_{\partial\Omega} \omega^\nu \cdot J[\bar{u}] \, dx' \, ds + \int_{(0,t) \times \Omega} (u^\nu - \bar{u}) \cdot (f^\nu - \bar{f}) \, dx \, ds.
\end{aligned}$$

Here $J[\bar{u}] = n \times u$. Using Corollary 2, we can control the total work of the friction force by

$$\begin{aligned}
& \left| \nu \int_0^t \int_{\partial\Omega} \omega^\nu \cdot J[\bar{u}] \, dx' \, dt \right| \leq CA^3 t |\partial\Omega| + \frac{\nu}{4} \|\nabla u^\nu\|_{L^2((0,t) \times \Omega)}^2 \\
& \quad + \frac{1}{4} \nu^{\frac{1}{3}} \|f^\nu\|_{L^{\frac{4}{3}}((0,t) \times \Omega)}^{\frac{4}{3}} + \nu^{\frac{4}{3}} \|u^\nu(0)\|_{H^1(\Omega)}^{\frac{2}{3}} \\
& \quad + \left(4 \log \left(\frac{4AL}{\nu} \right)_+ + \frac{\nu T}{\bar{\delta}^2} + \frac{C(\Omega)(1 + \nu^2)E_\nu T}{AL^4} \right) A\nu |\partial\Omega|.
\end{aligned}$$

Using Cauchy–Schwartz inequality, the forcing term can be controlled by

$$\begin{aligned}
& \int_{(0,t) \times \Omega} (u - \bar{u}) \cdot (f - \bar{f}) \, dx \, dt \\
& \leq \int_0^t \frac{\|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(s) + 1}{2} \|f^\nu - \bar{f}\|_{L^2(\Omega)}(s) \, ds.
\end{aligned}$$

By absorbing the dissipation term, we have

$$\begin{aligned}
& \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(t) + \frac{\nu}{2} \|\nabla u^\nu\|_{L^2((0,t) \times \Omega)}^2 - \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(0) \\
& \leq \int_0^t \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2 \left(2 \|D\bar{u}\|_{L^\infty(\Omega)} + \|f^\nu - \bar{f}\|_{L^2(\Omega)} \right) \, ds \\
& \quad + CA^3 t |\partial\Omega| + R_\nu(t),
\end{aligned} \tag{12}$$

where the remainder is

$$\begin{aligned}
R_\nu(t) & = \|f^\nu - \bar{f}\|_{L^1(0,t;L^2(\Omega))} + \nu \|\nabla \bar{u}\|_{L^2((0,t) \times \Omega)}^2 \\
& \quad + \nu^{\frac{1}{3}} \|f^\nu\|_{L^{\frac{4}{3}}((0,t) \times \Omega)}^{\frac{4}{3}} + 2\nu^{\frac{4}{3}} \|u^\nu(0)\|_{H^1(\Omega)}^{\frac{2}{3}} \\
& \quad + 2 \left(4 \log \left(\frac{4AL}{\nu} \right)_+ + \frac{\nu T}{\bar{\delta}^2} + \frac{C(\Omega)(1 + \nu^2)E_\nu T}{AL^4} \right) A\nu |\partial\Omega|.
\end{aligned}$$

By Grönwall inequality, we conclude that

$$\begin{aligned} & \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(T) + \frac{\nu}{2} \|\nabla u^\nu\|_{L^2((0,T) \times \Omega)}^2 \\ & \leq \left(\|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(0) + CA^3 T |\partial\Omega| + R_\nu(T) \right) \\ & \quad \times \exp \left(\int_0^T 2 \|D\bar{u}\|_{L^\infty(\Omega)} + \|f^\nu - \bar{f}\|_{L^2(\Omega)} dt \right). \end{aligned}$$

Note that

$$R_\nu(0) = 8(\log 4\text{Re})_+ A\nu |\partial\Omega| + 2\nu^{\frac{4}{3}} \|u^\nu(0)\|_{H^1(\Omega)}^{\frac{2}{3}}, \quad R_\nu(T) \rightarrow 0 \text{ as } \nu \rightarrow 0$$

provided $\nu^2 \|u^\nu(0)\|_{H^1(\Omega)} + \nu^{\frac{1}{4}} \|f^\nu\|_{L^{\frac{4}{3}}((0,T) \times \Omega)} + \|f^\nu - \bar{f}\|_{L^1(0,T;L^2(\Omega))} \rightarrow 0$. \square

Theorem 1 and **Corollary 1** are the consequence of **Theorem 2**.

Proof of Theorem 1 and Corollary 1 We first prove these results with an additional assumption that

$$f^\nu \in L^{\frac{4}{3}}((0, T) \times \Omega) \text{ and } \nu^{\frac{1}{4}} \|f^\nu\|_{L^{\frac{4}{3}}((0,T) \times \Omega)} \rightarrow 0 \text{ as } \nu \rightarrow 0. \quad (13)$$

For each ν we pick some $T_\nu > 0$ to be determined. By the energy inequality, it holds that

$$\nu \int_0^{T_\nu} \|\nabla u^\nu(t)\|_{L^2(\Omega)}^2 dt \leq \frac{1}{2} \|u^\nu(0)\|_{L^2(\Omega)}^2.$$

Therefore, there exists some time $\xi^\nu \in (0, T_\nu)$ such that

$$\nu T_\nu \|\nabla u^\nu(\xi^\nu)\|_{L^2(\Omega)}^2 \leq \frac{1}{2} \|u^\nu(0)\|_{L^2(\Omega)}^2.$$

Moreover, we know $\|u^\nu(\xi^\nu)\|_{L^2(\Omega)}^2 \leq \|u^\nu(0)\|_{L^2(\Omega)}^2$ due to energy inequality. Therefore

$$\nu^4 \|u^\nu(\xi^\nu)\|_{H^1(\Omega)}^2 \leq \left(\frac{\nu^3}{2T_\nu} + \nu^4 \right) \|u^\nu(0)\|_{L^2(\Omega)}^2 \rightarrow 0 \quad \text{as } \nu \rightarrow 0 \quad (14)$$

provided $\nu^3 T_\nu^{-1} \rightarrow 0$. Picking $T_\nu = \nu^2$ will work, for instance.

We claim that the work of the friction force between 0 and ξ^ν is negligible:

$$\liminf_{\nu \rightarrow 0} \left| \nu \int_0^{\xi^\nu} \int_{\partial\Omega} \omega^\nu \cdot J[\bar{u}] dx' dt \right| = 0.$$

This is because by **Lemma 4**, we integrate from 0 to ξ^ν :

$$\begin{aligned} & (u^\nu, \bar{u})(\xi^\nu) - (u^\nu, \bar{u})(0) \\ & = \int_0^{\xi^\nu} \int_{\Omega} [(u^\nu - \bar{u}) \otimes (u^\nu - \bar{u}) - \nu \nabla u^\nu] : \nabla \bar{u} dx dt \\ & \quad + \nu \int_0^{\xi^\nu} \int_{\partial\Omega} \omega^\nu \cdot J[\bar{u}] dx' dt + \int_0^{\xi^\nu} \int_{\Omega} \bar{u} \cdot f^\nu + u^\nu \cdot \bar{f} dx dt, \end{aligned}$$

in which as $\nu \rightarrow 0$, we establish the following convergences.

- $(u^\nu, \bar{u})(\xi^\nu) \rightarrow (\bar{u}, \bar{u})(0)$: $\bar{u}(\xi^\nu) \rightarrow \bar{u}(0)$ strongly in $L^2(\Omega)$, while $u^\nu(\xi^\nu) \rightharpoonup \bar{u}(0)$ weakly in $L^2(\Omega)$ up to a subsequence. This is because $u^\nu \rightarrow \bar{u}$ up to a subsequence in $C(0, T; H^{-1}(\Omega))$ using Aubin–Lions lemma, and u^ν are uniformly bounded in $L^\infty(0, T; L^2(\Omega))$, hence $u^\nu \rightarrow \bar{u}$ in $C_w(0, T; L^2(\Omega))$. Thus as $\nu \rightarrow 0$,

$$(u^\nu, \bar{u})(\xi^\nu) = (u^\nu(\xi^\nu), \bar{u}(\xi^\nu) - \bar{u}(0)) + (u^\nu(\xi^\nu) - \bar{u}(\xi^\nu), \bar{u}(0)) \\ + (\bar{u}(\xi^\nu) - \bar{u}(0), \bar{u}(0)) + \|\bar{u}(0)\|_{L^2(\Omega)}^2 \rightarrow \|\bar{u}(0)\|^2.$$

- $(u^\nu, \bar{u})(0) \rightarrow (\bar{u}, \bar{u})(0)$: this is simply because $u^\nu(0) \rightarrow \bar{u}(0)$ in $L^2(\Omega)$.
- $\iint [(u^\nu - \bar{u})^{\otimes 2} - \nu \nabla u^\nu] : \nabla \bar{u} \rightarrow 0$: $u^\nu - \bar{u}$ is uniformly bounded in $L^\infty(0, T; L^2(\Omega))$, and $\nu^{\frac{1}{2}} \nabla u^\nu$ is uniformly bounded in $L^2((0, T) \times \Omega)$.
- $\iint \bar{u} \cdot f^\nu \rightarrow 0$: this is because

$$\left| \int_0^{\xi^\nu} \int_{\Omega} \bar{u} \cdot f^\nu \, dx \, dt \right| \leq \|\bar{u}\|_{L^\infty(0, T; L^2(\Omega))} \|f^\nu - \bar{f}\|_{L^1(0, T; L^2(\Omega))} \\ + \left| \int_0^{\xi^\nu} \int_{\Omega} \bar{u} \cdot \bar{f} \, dx \, dt \right| \rightarrow 0.$$

- $\iint u^\nu \cdot \bar{f} \rightarrow 0$: u^ν is uniformly bounded in $L^\infty(0, T; L^2(\Omega))$.

These convergences prove the claim. Since this claim holds for any sequence of u^ν , it must hold that

$$R_\nu^{(1)} = \nu \int_0^{\xi^\nu} \int_{\partial\Omega} \omega^\nu \cdot J[\bar{u}] \, dx' \, dt \rightarrow 0 \quad \text{as } \nu \rightarrow 0.$$

Next, by [Corollary 2](#), we can control the work of the friction force from ξ^ν to t whenever $\xi^\nu < t \leq T$:

$$\left| \nu \int_{\xi^\nu}^t \int_{\partial\Omega} \omega^\nu \cdot J[\bar{u}] \, dx' \, dt \right| \leq CA^3 t |\partial\Omega| + \frac{\nu}{4} \|\nabla u^\nu\|_{L^2((0, t) \times \Omega)} \\ + \frac{1}{4} \nu^{\frac{1}{3}} \|f^\nu\|_{L^{\frac{4}{3}}((0, t) \times \Omega)}^{\frac{4}{3}} + \nu^{\frac{4}{3}} H_\nu^{\frac{1}{3}}(\xi^\nu) \\ + \left(4 \log \left(\frac{4AL}{\nu} \right)_+ + \frac{\nu T}{\delta^2} + \frac{C(\Omega)(1 + \nu^2)E_\nu T}{AL^4} \right) A\nu |\partial\Omega|.$$

Here $H_\nu(\xi^\nu) = \|u^\nu(\xi^\nu)\|_{H^1(\Omega)}^2$, and $\nu^{\frac{4}{3}} H_\nu^{\frac{1}{3}}(\xi^\nu) \rightarrow 0$ as $\nu \rightarrow 0$ by [\(14\)](#). Together with the energy inequalities, we have for every $0 < t < T$:

$$\|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(t) + \frac{\nu}{2} \|\nabla u^\nu\|_{L^2((0, t) \times \Omega)}^2 - \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(0) \\ \leq \int_0^t \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2 \left(2 \|D\bar{u}\|_{L^\infty(\Omega)} + \|f^\nu - \bar{f}\|_{L^2(\Omega)} \right) ds \\ + CA^3 t |\partial\Omega| + R_\nu(t),$$

where

$$R_\nu(t) = R_\nu^{(1)} + \|f^\nu - \bar{f}\|_{L^1(0, t; L^2(\Omega))} + \nu \|\nabla \bar{u}\|_{L^2((0, t) \times \Omega)}^2 \\ + \nu^{\frac{1}{3}} \|f^\nu\|_{L^{\frac{4}{3}}((0, t) \times \Omega)}^{\frac{4}{3}} + 2\nu^{\frac{4}{3}} H_\nu^{\frac{1}{3}}(\xi^\nu)$$

$$+ 2 \left(4 \log \left(\frac{4AL}{\nu} \right)_+ + \frac{\nu T}{\bar{\delta}^2} + \frac{C(\Omega)(1+\nu^2)E_\nu T}{AL^4} \right) A\nu |\partial\Omega|.$$

From our assumptions, we know $R_\nu(t) \rightarrow 0$ as $\nu \rightarrow 0$. By Grönwall inequality, we conclude

$$\begin{aligned} & \|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(T) + \frac{\nu}{2} \|\nabla u^\nu\|_{L^2((0,T) \times \Omega)}^2 \\ & \leq \left(\|u^\nu - \bar{u}\|_{L^2(\Omega)}^2(0) + CA^3 T |\partial\Omega| + R_\nu(T) \right) \\ & \quad \times \exp \left(\int_0^T 2 \|D\bar{u}\|_{L^\infty(\Omega)} + \|f^\nu - \bar{f}\|_{L^2(\Omega)} dt \right). \end{aligned}$$

Theorem 1 and **Corollary 1** are proven by sending $\nu \rightarrow 0$.

Finally, let us drop the assumption (13). Similar as before, we may assume $u^\nu \rightarrow \bar{u}$ in $C_w(0, T; L^2(\Omega))$. When f is not $L^{\frac{4}{3}}$ in time and space, we can take an average in time as follows. Let $\rho_\nu(t) = \frac{1}{\varepsilon_\nu} \mathbf{1}_{\{0 \leq t \leq \varepsilon_\nu\}}$, for some $\varepsilon_\nu > 0$ depending on ν to be determined, with $\varepsilon_\nu \rightarrow 0$ as $\nu \rightarrow 0$. Define $\tilde{u}^\nu = u^\nu *_t \rho^\nu$, $\tilde{f}^\nu = f^\nu *_t \rho^\nu$ by

$$\tilde{u}^\nu(t, x) = \int_{t-\varepsilon_\nu}^t u^\nu(s, x) ds, \quad \tilde{f}^\nu(t, x) = \int_{t-\varepsilon_\nu}^t f^\nu(s, x) ds$$

for $t \in [\varepsilon_\nu, T]$. We extend our definition by $\tilde{u}^\nu(t) = \tilde{u}^\nu(\varepsilon_\nu)$ and $\tilde{f}^\nu(t) = \tilde{f}^\nu(\varepsilon_\nu)$ for $0 < t < \varepsilon_\nu$. Then \tilde{u}^ν solves the Navier–Stokes equation in (ε_ν, T) :

$$\partial_t \tilde{u}^\nu + \tilde{u}^\nu \cdot \nabla \tilde{u}^\nu + \nabla \tilde{P}^\nu = \Delta \tilde{u}^\nu + \tilde{f}^\nu + f_1^\nu,$$

where

$$f_1^\nu = \tilde{u}^\nu \cdot \nabla \tilde{u}^\nu - \tilde{u}^\nu \cdot \nabla u^\nu := \tilde{u}^\nu \cdot \nabla \tilde{u}^\nu - \int_{t-\varepsilon_\nu}^t u^\nu \cdot \nabla u^\nu ds.$$

Then $\tilde{u}^\nu - u^\nu \rightarrow 0$ in $C_w(0, T; L^2(\Omega))$, $\tilde{f}^\nu - f^\nu \rightarrow 0$ in $L^1(0, T; L^2(\Omega))$, $f_1^\nu \rightarrow 0$ in $L^1(0, T; L^{\frac{3}{2}}(\Omega))$, and thus

$$\begin{aligned} \nu^{\frac{1}{4}} \|\tilde{f}^\nu\|_{L^{\frac{4}{3}}((0,T) \times \Omega)} & \leq C(\Omega) \nu^{\frac{1}{4}} \varepsilon_\nu^{-\frac{1}{4}} \|f^\nu\|_{L^1(0,T;L^2(\Omega))}, \\ \nu^{\frac{1}{4}} \|f_1^\nu\|_{L^{\frac{4}{3}}((0,T) \times \Omega)} & \leq C(\Omega) \nu^{\frac{1}{4}} \varepsilon_\nu^{-\frac{1}{4}} \|u^\nu\|_{L^2(0,T;L^6(\Omega))} \|\nabla u^\nu\|_{L^2(0,T;L^2(\Omega))}. \end{aligned}$$

If we set, for instance, $\varepsilon_\nu = \nu^{\frac{1}{2}}$, then $\nu^{\frac{1}{4}} \|\tilde{f}^\nu + f_1^\nu\|_{L^{\frac{4}{3}}((0,T) \times \Omega)} \rightarrow 0$ as $\nu \rightarrow 0$.

By **Lemma 4**, we can estimate the inner product of \tilde{u}^ν and \bar{u} :

$$\begin{aligned} & (\tilde{u}^\nu, \bar{u})(T) - (\tilde{u}^\nu, \bar{u})(\varepsilon_\nu) \\ & = \int_{\varepsilon_\nu}^T \int_\Omega [(\tilde{u}^\nu - \bar{u}) \otimes (\tilde{u}^\nu - \bar{u}) - \nu \nabla \tilde{u}^\nu] : \nabla \bar{u} dx dt \\ & \quad + \nu \int_{\varepsilon_\nu}^T \int_{\partial\Omega} \partial_n \tilde{u}^\nu \cdot \bar{u} dx' + \int_{\varepsilon_\nu}^T \int_\Omega \bar{u} \cdot \tilde{f}^\nu + \tilde{u}^\nu \cdot \bar{f} dx dt. \end{aligned}$$

Due to convergence $\tilde{u}^\nu - u^\nu \rightarrow 0$ in $C_w(0, T; L^2(\Omega))$, $\nabla \tilde{u}^\nu - \nabla u^\nu \rightarrow 0$ in $L^2((0, T) \times \Omega)$, and $\tilde{f}^\nu + f_1^\nu \rightarrow f^\nu$ in $L^1(0, T; L^{\frac{3}{2}}(\Omega))$, we conclude

$$\begin{aligned}
& (u^\nu, \bar{u})(T) - (u^\nu, \bar{u})(0) \\
&= \int_0^T \int_{\Omega} [(u^\nu - \bar{u}) \otimes (u^\nu - \bar{u}) - \nu \nabla u^\nu] : \nabla \bar{u} \, dx \, dt \\
&\quad + \nu \int_{\varepsilon_\nu}^T \int_{\partial\Omega} \partial_n \tilde{u}^\nu \cdot \bar{u} \, dx' + \int_0^T \int_{\Omega} \bar{u} \cdot f^\nu + u^\nu \cdot \bar{f} \, dx \, dt + R_\nu^{(2)}.
\end{aligned}$$

for some $R_\nu^{(2)} \rightarrow 0$ as $\nu \rightarrow 0$. Using [Corollary 2](#), we can bound the boundary term by

$$\begin{aligned}
\left| \nu \int_{\varepsilon_\nu}^T \int_{\partial\Omega} \partial_n \tilde{u}^\nu \cdot \bar{u} \, dx' \, dt \right| &\leq CA^3 T |\partial\Omega| + \frac{1}{4} \nu \int_0^T \int_{\Omega} |\nabla u^\nu|^2 \, dx \, dt \\
&\quad + \frac{1}{4} F_\nu + R_\nu,
\end{aligned}$$

where $R_\nu \rightarrow 0$ as $\nu \rightarrow 0$, and $F_\nu = \nu^{\frac{1}{3}} \|\tilde{f}^\nu + f_1^\nu\|_{L^{\frac{4}{3}}((0,T) \times \Omega)}^{\frac{4}{3}} \rightarrow 0$ as $\nu \rightarrow 0$ as well. Combining with the energy inequality and Grönwall inequality, we finish the proof of [Theorem 1](#) and [Corollary 1](#) for general force without assumption (13). \square

Finally, we recover the result of Kato from our analysis.

Proof of Theorem A The term $CA^3 T |\partial\Omega|$ of [Theorem 2](#) is due to the integral III in (10) in the proof of [Corollary 2](#), which comes from two sources: $\frac{C}{\gamma} A^3 T |\partial\Omega|$ in (11) can be traced back to the boundary vorticity estimate (9), and the other was $\gamma A^3 T |\partial\Omega|$. For the former, if the Kato's condition (2) holds for $\delta = \nu/A$, then by [Theorem 3](#) we have for any $\gamma < 1$,

$$\lim_{\nu \rightarrow 0} \left\| \nu \tilde{\omega}^\nu \mathbf{1}_{\{\nu |\tilde{\omega}^\nu| > \gamma \max\left\{\frac{\nu}{T}, \frac{\nu^2}{\delta^2}\right\}\}} \right\|_{L^{\frac{3}{2}, \infty}((0, T) \times \partial\Omega)}^{\frac{3}{2}} = 0,$$

thus $\limsup_{\nu \rightarrow 0} |\text{III}| \leq \gamma A^3 T |\partial\Omega|$. Consequently, $\text{LS}(\bar{u}) \leq \gamma A^3 T |\partial\Omega|$. This is true for any $\gamma \in (0, 1]$, therefore $\text{LS}(\bar{u}) = 0$. The general case $\delta = cv$ for $c > 0$ is a simple consequence of the rescaling of time. \square

Remark 2. The main part of the proof of [Theorem 2](#) is to use [Corollary 2](#) to bound the work of boundary friction toward the Euler flow. One could also study the work of fluid toward the boundary. This is related to the well-known d'Alembert's paradox: for an object traveling at a constant speed in a steady potential flow, there is no drag force, so the ambient fluid does zero work toward the object. However, in reality, an object moving in a fluid experiences a drag force, no matter how small the viscosity is or how fast the speed is.

To be more precise, imagine that an object K is moving at a constant velocity Ue_1 in a low-viscosity incompressible fluid in a large periodic domain. Sending the period to infinity is another nontrivial task, but we ignore it here. Then in the reference frame of K , the fluid around it solves the Navier-Stokes equation in $\Omega = \mathbb{T}^3 \setminus K$, with a background flow $\bar{u} \approx -Ue_1$ away from the object. Denote $\Sigma = -P^\nu \text{Id} + 2\nu Du^\nu$ to be the stress tensor of the fluid. Then the total force exerted on the object by the fluid at a given time is

$$\int_{\partial\Omega} -\Sigma n \, dx' = \int_{\partial\Omega} P^\nu n - \nu \partial_n u^\nu \, dx'. \tag{15}$$

Here n is the outer normal of $\partial\Omega$, i.e. the inward normal of ∂K . This force contains two parts: the first is due to pressure, and the second is due to friction. In e_1 direction, the former is

called “form drag”, whereas the latter is called “skin drag”. The work done on the object in the static frame of reference is

$$\int_{(0,T) \times \partial\Omega} -\Sigma n \cdot Ue_1 \, dx' \, dt = \int_{(0,T) \times \partial\Omega} \Sigma n \cdot (-Ue_1) \, dx' \, dt.$$

Recall that the work done on the Euler solution in the object’s frame of reference is

$$\nu \int_{(0,T) \times \partial\Omega} \partial_n u^\nu \cdot \bar{u} \, dx' \, dt = \int_{(0,T) \times \partial\Omega} \Sigma n \cdot \bar{u} \, dx' \, dt.$$

If $\bar{u} \approx -Ue_1$ on the boundary, then they are approximately the same. In particular, they are the same when K is a flat plate moving at a constant velocity tangential to its surface.

The drag force experienced by the object has the following empirical formula, which is derived from dimensional analysis by Lord Rayleigh:

$$F_{\text{drag}} = \frac{1}{2} \rho c_d(\text{Re}) U^2 S.$$

Here ρ is the density of the fluid, $c_d(\text{Re})$ is a dimensionless parameter called *drag coefficient*, depending on the shape of the object, and the *Reynolds number* $\text{Re} = \frac{UL}{\nu}$, where L is the characteristic length, and S is the reference area. One may choose L to be the diameter of the object K . It is customary to choose S as the cross-sectional area, but for wings it should be chosen as the lifting area. It has been observed experimentally that the drag coefficient $c_d(\text{Re})$ has a finite limit as $\text{Re} \rightarrow \infty$, i.e. $\nu \rightarrow 0$. For instance, a rigorous analysis shows the drag coefficient of a flat plate can be bounded by approximately 295.49 [35] as $\text{Re} \rightarrow \infty$. Upon fixing a unit system such that $\rho = 1$, the work done by the drag force from time 0 to T is exactly $c_d(\text{Re}) U^3 T S$. In the case of a flat plate, $\bar{u} = -Ue_1$, $A = U$, and $S = \frac{1}{2} |\partial\Omega|$. Our work then shows that even for weak solutions, the inviscid limit of drag coefficient has an upper bound:

$$\limsup_{\text{Re} \rightarrow \infty} c_d(\text{Re}) \leq C \left(\frac{A}{U} \right)^3 \cdot \frac{|\partial\Omega|}{S} \leq C.$$

For a general object K , we can set $\varphi = -Ue_1$ in Corollary 2 to provide a constant upper bound for the limiting skin drag coefficient

$$\limsup_{\text{Re} \rightarrow \infty} c_{d,\text{skin}}(\text{Re}) \leq \frac{C U^3 T |\partial\Omega|}{\frac{1}{2} U^2 S \cdot U T} \leq C \frac{|\partial\Omega|}{S},$$

where $c_{d,\text{skin}}$ is defined similarly as c_d but considering only the skin friction drag, neglecting the form drag component.

Acknowledgments

The authors would like to thank American Institute of Mathematics for the workshop “Criticality and stochasticity in quasilinear fluid systems”, where this project was initiated.

Funding

The first author was partially supported by the NSF grant: DMS 2306852. The second author was partially supported by the NSF grant: DMS 2054888.

References

- [1] Vasseur, A. F., Yang, J. (2023). Boundary vorticity estimates for Navier–Stokes and application to the inviscid limit. *SIAM J. Math. Anal.* 55(4):3081–3107.
- [2] Lee, J., de Silva, C., Monty, J., Hutchins, N. (2014). Video: Spatially developing turbulent boundary layers: The return of the plate. In: *67th Annual Meeting of the APS Division of Fluid Dynamics - Gallery of Fluid Motion*, DFD 2014. American Physical Society.
- [3] Deck, S. (2018). The spatially developing flat plate turbulent boundary layer. In: Mockett, C., Haase, W., Schwamborn, D., eds. *Go4Hybrid: Grey Area Mitigation for Hybrid RANS-LES Methods*. Cham: Springer, pp. 109–121.
- [4] Prandtl, L. (1904). Über flüssigkeitsbewegung bei sehr kleiner reibung. In: *Actes du 3me Congrès International des Mathématiciens, Heidelberg, Teubner, Leipzig*, pp. 484–491.
- [5] Grenier, E. (2000). On the nonlinear instability of Euler and Prandtl equations. *Commun. Pure Appl. Math.* 53(9):1067–1091.
- [6] Weinan, E. (2000). Boundary layer theory and the zero-viscosity limit of the Navier–Stokes equation. *Acta Math. Sin. (Engl. Ser.)* 16(2):207–218.
- [7] Gérard-Varet, D., Dormy, E. (2010). On the ill-posedness of the Prandtl equation. *J. Amer. Math. Soc.* 23(2):591–609.
- [8] Gérard-Varet, D., Nguyen, T. T. (2012). Remarks on the ill-posedness of the Prandtl equation. *Asymptot. Anal.* 77(1–2):71–88.
- [9] Kato, T. (1984). Remarks on zero viscosity limit for nonstationary Navier–Stokes flows with boundary. In: Chern, S. S., ed. *Seminar on Nonlinear Partial Differential Equations*. New York: Springer, pp. 85–98.
- [10] Maekawa, Y. (2014). On the inviscid limit problem of the vorticity equations for viscous incompressible flows in the half-plane. *Commun. Pure Appl. Math.* 67(7):1045–1128.
- [11] Maekawa, Y., Mazzucato, A. L. (2018). The inviscid limit and boundary layers for Navier–Stokes flows. In: *Handbook of Mathematical Analysis in Mechanics of Viscous Fluids*, pp. 781–828. Cham: Springer.
- [12] Fei, M., Tao, T., Zhang, Z. (2018). On the zero-viscosity limit of the Navier–Stokes equations in \mathbb{R}_+^3 without analyticity. *J. Math. Pures Appl.* (9) 112:170–229.
- [13] Albritton, D., Bruè, E., Colombo, M. (2022). Non-uniqueness of Leray solutions of the forced Navier–Stokes equations. *Ann. Math.* (2) 196(1):415–455.
- [14] Albritton, D., Bruè, E., Colombo, M. (2023). Gluing non-unique Navier–Stokes solutions. *Ann. PDE* 9(2):17.
- [15] Jia, H., Sverak, V. (2015). Are the incompressible 3D Navier–Stokes equations locally ill-posed in the natural energy space? *J. Funct. Anal.* 268(12):3734–3766.
- [16] Vishik, M. (2018). Instability and non-uniqueness in the Cauchy problem for the Euler equations of an ideal incompressible fluid. Part I. arXiv:1805.09426.
- [17] Vishik, M. (2018). Instability and non-uniqueness in the Cauchy problem for the Euler equations of an ideal incompressible fluid. Part II. arXiv:1805.09440.
- [18] De Lellis, C., Bruè, E., Albritton, D., Colombo, M., Giri, V., Janisch, M., Kwon, H. (2024). *Instability and Non-uniqueness for the 2D Euler Equations, after M. Vishik*, Princeton, NJ: Princeton University Press.
- [19] Székelyhidi, L. (2011). Weak solutions to the incompressible Euler equations with vortex sheet initial data. *Comptes Rendus Mathématique* 349(19):1063–1066.
- [20] Buckmaster, T., De Lellis, C., Székelyhidi, L., Vicol, V. (2019). Onsager’s conjecture for admissible weak solutions. *Commun. Pure Appl. Math.* 72(2):229–274.
- [21] Buckmaster, T., Vicol, V. (2019). Nonuniqueness of weak solutions to the Navier–Stokes equation. *Ann. Math.* (2) 189(1):101–144.
- [22] De Lellis, C., Székelyhidi, L. (2010). On admissibility criteria for weak solutions of the Euler equations. *Arch. Rational Mech. Anal.* 195(1):225–260.
- [23] Isett, P. (2018). A proof of Onsager’s conjecture. *Ann. Math.* (2) 188(3):871–963.
- [24] Kolmogoroff, A. N. (1941). The local structure of turbulence in incompressible viscous fluid for very large Reynold’s numbers. *C. R. (Doklady) Acad. Sci. URSS (N. S.)* 30:301–305.

- [25] Kolmogoroff, A. N. (1941). On degeneration of isotropic turbulence in an incompressible viscous liquid. *C. R. (Doklady) Acad. Sci. URSS (N. S.)* 31:538–540.
- [26] Kolmogoroff, A. N. (1941). Dissipation of energy in the locally isotropic turbulence. *C. R. (Doklady) Acad. Sci. URSS (N. S.)* 32:16–18.
- [27] Bruè, E., De Lellis, C. (2023). Anomalous dissipation for the forced 3D Navier–Stokes equations. *Commun. Math. Phys.* 400:1507–1533.
- [28] Hilton, A., Illingworth, J., et al. (1997). Marching triangles: Delaunay implicit surface triangulation. University of Surrey.
- [29] Hartmann, E. (1998). A marching method for the triangulation of surfaces. *Visual Comput.* 14(3):95–108.
- [30] McCormick, N. H., Fisher, R. B. (2002). Edge-constrained marching triangles. In *Proceedings. First International Symposium on 3D Data Processing Visualization and Transmission*, pp. 348–351.
- [31] Maremonti, P., Solonnikov, V. A. (1997). Estimates for solutions of the nonstationary Stokes problem in anisotropic sobolev spaces with mixed norm. *J. Math. Sci.* 87(5):3859–3877.
- [32] Seregin, G. (2014). *Lecture Notes on Regularity Theory for the Navier–Stokes Equations*. Singapore: World Scientific.
- [33] Maremonti, P., Solonnikov, V. A. (1995). On estimates for the solutions of the nonstationary Stokes problem in S. L. Sobolev anisotropic spaces with a mixed norm. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 222(Issled. po Linein. Oper. i Teor. Funktsii. 23):124–150, 309.
- [34] Solonnikov, V. A. (2002). Estimates of solutions of the Stokes equations in S. L. Sobolev spaces with a mixed norm. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 288(Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funkts. 32):204–231, 273–274.
- [35] Kumar, A., Garaud, P. (2020). Bound on the drag coefficient for a flat plate in a uniform flow. *J. Fluid Mech.* 900:A6.