

Brane wrapping, Alexandrov-Kontsevich-Schwarz-Zaboronsky sigma models, and QP manifolds

Alex S. Arvanitakis^{1,*} and David Tennyson^{2,†}

¹Theoretische Natuurkunde, Vrije Universiteit Brussel, and the International Solvay Institutes, Pleinlaan 2, B-1050 Brussels, Belgium

²Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, Texas 77843, USA



(Received 9 April 2023; accepted 7 September 2023; published 18 October 2023)

We introduce a technique to realize brane wrapping and double dimensional reduction in the context of Alexandrov-Kontsevich-Schwarz-Zaboronsky (AKSZ) topological sigma models and also in their target spaces, which are symplectic L_n algebroids (i.e., QP manifolds). Our procedure involves a novel coisotropic reduction combined with an AKSZ transgression that realizes degree shifting; the reduced QP manifold depends on topological data of the “wrapped” cycle. We check our procedure against the known rules for fluxes under wrapping in the context of M-theory/type IIA duality, and we also find a new relation between Courant algebroids and Poisson manifolds.


DOI: 10.1103/PhysRevD.108.086024

I. INTRODUCTION

In a series of recent papers [1–3] we have been establishing a correspondence between Bogomol'nyi-Prasad-Sommerfield p -branes in string/M-theory on the one hand and *symplectic L_{p+1} algebroids* on the other hand. The latter can be thought of as appropriate generalizations of (exact) *Courant* algebroids, and they are relevant for describing generic flux backgrounds of string theory/M-theory (see [4] for this point of view, and [5] for an alternative). The algebroid description of these backgrounds has been used to understand properties of reductions such as Kaluza-Klein (KK) spectra [6], consistent truncations [7], and marginal deformations [8,9]. In the aforementioned brane-algebroid

correspondence, the exact Courant algebroid (which is classified by the de Rham cohomology class of H [10]) is associated with the fundamental string (which couples to H electrically via a Wess-Zumino term). For other branes, such as the M2- and M5-branes in M-theory, or even/odd D-branes in type II string theory, the corresponding algebroids are roughly speaking the ones that are classified by whichever fluxes couple *electrically* to the brane in question.

In general, the correspondence is between a “physical” p -brane (an F1, M2, M5, …), a symplectic L_n algebroid for $n = p + 1$, and a *topological* Alexandrov-Kontsevich-Schwarz-Zaboronsky (AKSZ) brane sigma model of dimension $p + 2$. This can be schematically summarized in the following diagram:



Less tersely, the symplectic L_n algebroid—that is classified by a certain collection of fluxes—determines a topological

* alex.s.arvanitakis@vub.be
† dtennyson@tamu.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

n -brane sigma model via the AKSZ construction [11]. When the n -brane has an $(n - 1)$ -brane boundary, an inflow-type argument with an appropriate boundary condition produces the Wess-Zumino (WZ) term that couples those same fluxes to the $(n - 1)$ -brane [1,3].¹ The algebroid also determines the corresponding $(n - 1)$ -brane more directly via the *brane*

¹A slightly different boundary condition for the AKSZ sigma model can produce the entire $(n - 1)$ -brane Lagrangian, including kinetic terms. This was done for the fundamental string by Severa [12]. The other cases have not yet been considered in the literature.

phase space construction that yields the Poisson algebra of brane currents on phase space [2] (i.e., in the Hamiltonian formulation of brane dynamics).

This correspondence between branes and algebroids motivates the question: given that the string/M-theory duality web acts on the branes, *how is the duality web realized on the algebroid side?* Heretofore this was only known for dualities that preserve the world volume dimension; see [13] for T-duality and [3] for M-theory/type IIA duality along a transverse M-theory circle. [An example of the latter is the emergence of a D2-brane given an M2brane that does not wrap the M-theory circle, whose algebroid avatar is symplectic reduction modulo the U(1) action.] This “algebroid duality web” has already found applications in physics, including, e.g., to spacetime topology change induced by Poisson-Lie T-duality [13].

In this paper we provide an algebroid realization for the *brane wrapping* operation. In the string theory picture, this sends a p -brane to the $(p-d)$ -brane found by wrapping the original brane around a d -dimensional cycle on target space and then shrinking the volume of the cycle to zero. (Since both the dimensionality of the brane and that of the target space are reduced in this way, this is also known as *double dimensional reduction*.) The most basic example is M-theory/IIA duality, where M2-branes wrapped around the compactified 11th dimension give rise to fundamental strings in ten dimensions [14]. This already poses a puzzle: the corresponding algebroids are of degree $n = p + 1 = 3$ (for the M2-brane) and $n = 2$ (for the F1); *what is the mathematical operation that accounts for this degree shift?*

The mystery is resolved in the supergeometric formulation of symplectic L_n algebroids, defined by the data of a *QP manifold* (\mathcal{M}, ω, Q) where \mathcal{M} is a non-negatively graded manifold, ω a symplectic form of degree n , and Q a nilpotent vector field of degree 1, Hamiltonian for ω . Given a compact manifold X of dimension d —to be identified with the cycle to be “wrapped”—the odd tangent bundle $\mathcal{X} \equiv T[1]X$ possesses an integration measure $\int_{\mathcal{X}} : C^\infty(\mathcal{X}) \rightarrow \mathbb{R}$ of degree $-d$, namely the integral of differential forms. Then the mapping space

$$\mathcal{M}^{\mathcal{X}} \equiv \text{maps}(\mathcal{X} \rightarrow \mathcal{M}) \quad (1.2)$$

possesses a P structure of degree $(n-d)$, provided by the AKSZ construction. This is the correct degree shift; however, this manifold is infinite dimensional, and its structure sheaf is not non-negatively graded, so it cannot be the sought-after symplectic L_{n-d} algebroid.

A. A “brane wrapping” for QP manifolds

We introduce a coisotropic reduction of the space $\mathcal{M}^{\mathcal{X}}$ to a finite-dimensional QP manifold that resolves both issues. This resolution is heavily motivated by the intuitive




FIG. 1. The wrapping map specification for $N = \mathbb{R}$, $X = S^1$.

string-theoretic picture of brane wrapping. We deal with the case where the body of \mathcal{M} is a product $N \times X$, seen as a trivial bundle with fiber X , and we select a map $N \hookrightarrow \text{maps}(X \rightarrow N \times X)$, as in Fig. 1. The idea is that each point $n \in N$ is mapped to the cycle of $N \times X$ that shrinks to zero size in the double dimensional reduction procedure. Since $\text{maps}(X \rightarrow N \times X)$ is disconnected, with connected components corresponding to different winding sectors (as they would be called in physics), the choice of map $N \hookrightarrow \text{maps}(X \rightarrow N \times X)$ includes a choice of winding. On the string theory side, double dimensional reduction indeed depends on winding: for instance, an M2-brane wound w times around the M-theory circle yields a fundamental string coupled to the H -flux wH . Since the algebroids corresponding to these branes via the diagram (1.1) are defined by the same fluxes, we expect winding dependence in the obtained algebroid, and we will indeed find it.

In more detail: we start with the data of an NQP—“N” for non-negatively graded—manifold \mathcal{M} with body M and a “source” Q manifold $\mathcal{X} = T[1]X$ as above, along with a *wrapping map* $\mathfrak{w} : X \rightarrow M$ that defines a degree-0 submanifold $N \hookrightarrow \text{maps}(X \rightarrow M)$. We then produce a *finite-dimensional, non-negatively graded QP manifold* \mathcal{W} , whose P structure has degree $n-d$; we will call \mathcal{W} the *wrapped algebroid*, and we will call our procedure *(brane) wrapping*. The wrapping of QP manifolds/symplectic L_n algebroids is then a reduction of $\mathcal{M}^{\mathcal{X}}$ with respect to a coisotropic submanifold \mathcal{C} which may be thought of as the lift of $N \hookrightarrow \text{maps}(X \rightarrow M)$ to a graded submanifold of $\text{maps}(\mathcal{X} \rightarrow \mathcal{M}) = \mathcal{M}^{\mathcal{X}}$. The output QP manifold \mathcal{W} depends on the choice of wrapping map \mathfrak{w} only up to homotopy.

In fact, we were able to generalize beyond the case $M = N \times X$ (that was pictorially outlined above) to the case $M = N \times Y$, with Y and X not necessarily of the same dimension, even; then the wrapping is a map $\mathfrak{w} : X \rightarrow N \times Y$, and $d = \dim X$ controls the degree/dimensionality shifts as before. This generalization allows us to accommodate at least one example which might be of interest outside of string theory, namely the wrapping of a Courant algebroid into a Poisson manifold discussed in Sec. IV B, which has $\dim Y = 0$. When $\dim X = n+1$ in

addition to $\dim Y = 0$ (so that $\mathcal{M}^{\mathcal{X}}$ has a degree -1 P structure) our wrapping procedure agrees with that of [15]. Our approach gives a complementary perspective to the Losev-trick based wrapping-style reductions of [16,17] and to that of [18,19].

B. Brane wrapping and AKSZ sigma models

Our brane wrapping reduction—from a QP manifold \mathcal{M} to a QP manifold \mathcal{W} —also induces a reduction of the corresponding AKSZ topological field theories. Essentially, the two reductions commute, as in the schematic diagram

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{\text{wrapping}} & \mathcal{W} \\ \downarrow \text{AKSZ} & & \downarrow \text{AKSZ} \\ \mathcal{M}^{\mathcal{X} \times \mathcal{S}} & \xrightarrow{\text{dotted}} & \mathcal{W}^{\mathcal{S}} \end{array} \quad (1.3)$$

Here $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$ and $\mathcal{W}^{\mathcal{S}}$ are P manifolds of degree -1 created by the AKSZ construction for \mathcal{S} of appropriate dimension. The dotted arrow corresponds to a reduction of $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$ with respect to the coisotropic submanifold $\mathcal{C}^{\mathcal{S}} \equiv \text{maps}(\mathcal{S}, \mathcal{C})$, for \mathcal{C} the coisotropic submanifold that appears in the wrapping reduction $\mathcal{M} \rightarrow \mathcal{W}$. This “dotted” reduction always exists and is compatible with the AKSZ/BV (Batalin-Vilkovisky)master actions if the wrapping reduction does.

We provide the argument for the reduction of AKSZ sigma models in Sec. VI, along with an example: the reduction of a topological 3-brane sigma model (corresponding to the M2-brane symplectic L_3 algebroid) to a Courant sigma model (corresponding to the fundamental string symplectic L_2 algebroid). This provides an important consistency check: if we were to derive the corresponding physical brane sigma models, e.g., by introducing boundaries and using an inflow-type argument as in [1,12], we would find that the electric Wess-Zumino flux coupling has the correct winding dependence.

C. Structure of the paper

In Sec. II, we describe the general procedure for wrapping QP manifolds. We provide the conditions required of the QP structure on \mathcal{M} and define the coisotropic ideal $\mathcal{I} \subset C^\infty(\mathcal{M}^{\mathcal{X}})$ (that defines the coisotropic submanifold \mathcal{C}) in general. We show that it is well-defined and perform the reduction. The next three sections provide a multitude for examples. (If the reader finds the notation of Sec. II too terse, they may find it useful to first work their way through the examples before coming back to the general procedure.) Section III covers the case where $\dim X = 0$. In this case, we do not get any wrapping and our reduction is very similar to conventional dimensional reduction [20]. In Sec. IV we

consider examples where $\dim X \neq 0$, but the wrapping map \mathfrak{w} is trivial in homotopy. These provide examples which are simple but still present some of the main features of the reduction. Among these is the reduction of a Courant algebroid to a Poisson manifold given in Sec. IV B. In Sec. V, we consider examples relevant for physics and wrap string/M-theory branes on various manifolds. In Sec. VI we show how our procedure naturally lifts to a reduction of the AKSZ theory from $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$ to $\mathcal{W}^{\mathcal{S}}$. Section VII is left for comments and outlook. The appendixes cover our notation (Appendix A), some key properties and conventions of QP manifolds (Appendix B), and a review of coisotropic reduction in the graded context (Appendix C).

II. WRAPPING QP MANIFOLDS

We will describe a process of creating new QP manifolds from old, which effectively generalizes the notion of dimensional reduction, that we describe as wrapping QP manifolds. The nomenclature arises due to the consistency of this process with the AKSZ construction [11]—that is, one can reduce the AKSZ theory from the original QP manifold to that of the new manifold. Solutions of this reduced AKSZ theory will look like branes wrapping cycles of the target space. We will describe the relation to AKSZ sigma models in a later section and will describe the wrapping procedure here.

We start from the following ingredients:

- (i) An NQP manifold $\mathcal{M} = \mathcal{N} \times \mathcal{Y}$ of degree $n \geq 2$ where

$$\mathcal{Y} = T^*[n]T[1]Y \quad (2.1)$$

and \mathcal{N} is otherwise generic, with underlying commutative manifold² N . The underlying commutative manifold for \mathcal{M} is $M = N \times Y$, a direct product manifold. The symplectic form will be written $\omega_{\mathcal{M}} = d\vartheta_{\mathcal{M}}$, where $\vartheta_{\mathcal{M}}$ is the canonical symplectic potential. The induced Poisson bracket on \mathcal{M} will be written $(\cdot, \cdot)_{\mathcal{M}}$.

- (ii) The Q structure of \mathcal{M} should be a *lift of the de Rham differential of Y* , seen as the vector field $d_Y \equiv \xi^m \partial/\partial y^m$ on $T[1]Y$, with respect to the bundle projection p that is the composition $\mathcal{N} \times \mathcal{Y} \xrightarrow{\pi_{\mathcal{Y}}} \mathcal{Y} \xrightarrow{\pi_{T[1]Y}} T[1]Y$. Explicitly this lift condition means $Q_{\mathcal{M}} p^* = p^* d_Y$, which partially determines the form of the Hamiltonian $\Theta_{\mathcal{M}}$ in local coordinates:

²By “underlying commutative manifold” we mean the commutative manifold M whose structure sheaf is the sheaf of degree 0 functions on M , i.e., $C^\infty(M) = C_0^\infty(M)$. This is well defined since we are working on graded-commutative manifolds with a non-negative grading. We will also refer to this as the manifold in degree 0.

$$\Theta_{\mathcal{M}} = -\xi^m q_m + \sum_{k=0}^{n+1} \frac{1}{k!} \theta_{m_1 \dots m_k}(z, y) \xi^{m_1} \dots \xi^{m_k}, \quad (2.2)$$

where q are the degree n conjugate momenta to y on $T^* [n] T[1] Y$ and z are generic homogeneous coordinates on \mathcal{N} . The $\theta_k = \theta_k(z, y) \xi^k$ can be viewed as $[C^\infty(\mathcal{N})\text{-valued}]$ differential forms on Y , and we demand that they must be d_Y -closed differential forms.

- (iii) A Q manifold $\mathcal{X} = (T[1]X, d)$ where X is compact, without boundary, and has dimension $d < n$. d is the de Rham differential.
- (iv) A choice of “wrapping map” $\mathbf{w}: X \rightarrow Y$, defined up to homotopy.

We aim to produce a new NQP manifold \mathcal{W} from \mathcal{M}, \mathcal{X} , which describes a brane where X has been wrapped over Y and both cycles have been shrunk. The resulting QP manifold should therefore have degree $n - d$ and underlying commutative manifold N . There is a natural choice of manifold of degree $n - d$ given by the mapping space $\mathcal{M}^\mathcal{X} := \text{maps}(\mathcal{X} \rightarrow \mathcal{M})$. However, this manifold is infinite dimensional. We will see that we can define a coisotropic reduction of $\mathcal{M}^\mathcal{X}$ that produces a finite dimensional NQP manifold which only depends on the topology of X and the homotopy class of \mathbf{w} .

A. Properties of the mapping space

The infinite dimensional space $\mathcal{M}^\mathcal{X}$ consists of maps f which are defined by their pullback action on the coordinates on \mathcal{M} . Using generic homogeneous coordinates Z^A for \mathcal{M} and coordinates $(\sigma^\alpha, d\sigma^\alpha)$ for \mathcal{X} adapted to d $[d(\sigma^\alpha) = d\sigma^\alpha, dd\sigma^\alpha = 0]$ we have

$$\begin{aligned} f^* Z^A &= \mathbf{Z}^A(\sigma, d\sigma) = Z_0^A(\sigma) + Z_{1\alpha}^A(\sigma)d\sigma^\alpha + \dots \\ &+ \frac{1}{d!} Z_{d\alpha_1 \dots \alpha_d}^A(\sigma)d\sigma^{\alpha_1} \dots d\sigma^{\alpha_d}. \end{aligned} \quad (2.3)$$

Defining the components Z_k is equivalent to defining the map f . To interpret the Z_k we consider a change of coordinates on \mathcal{M} given by $\tilde{Z}^A = \tilde{Z}^A(Z)$ and note that

$$\begin{aligned} f^* \tilde{Z}^A(Z) &= \tilde{Z}^A(f^* Z) \\ &= \tilde{Z}^A(Z_0) + Z_{1\alpha}^B d\sigma^\alpha \frac{\partial \tilde{Z}^A}{\partial Z^B}(Z_0) \\ &+ \frac{1}{2} d\sigma^\alpha d\sigma^\beta \left(Z_{2\alpha\beta}^B \frac{\partial \tilde{Z}^A}{\partial Z^B}(Z_0) \right. \\ &\left. + Z_{1\alpha}^B Z_{1\beta}^C \frac{\partial^2 \tilde{Z}^A}{\partial Z^B \partial Z^C}(Z_0) \right) + \dots \end{aligned} \quad (2.4)$$

Therefore, in spite of the index structure, these in general are *not* vector-bundle-valued differential forms, with the

exception of Z_1 which is an $f_0^* T\mathcal{M}$ -valued one-form for the map $f_0 = f \circ s_0$, where $s_0: X \rightarrow \mathcal{X}$ is the zero section of $\mathcal{X} = T[1]X$. Of the other components, Z_0^A defines the map $f_0: X \rightarrow \mathcal{M}$, while the Z_k^A for $k > 1$ transform “affinely” whenever $Z_{k'}^A \neq 0$ for any $0 < k' < k$.³ Since we may not set $Z_k^A = 0$ consistently in general, this introduces a subtlety for our reduction procedure that we will discuss later in this section.

The QP structure on the mapping space is induced by that on \mathcal{M} through transgression. The symplectic structure is given by

$$\omega_{\mathcal{M}^\mathcal{X}} = \int_{\mathcal{X}} \frac{1}{2} \delta \mathbf{Z}^A (\omega_{\mathcal{M}})_{AB} \delta \mathbf{Z}^B = \sum_k \int_{\mathcal{X}} \frac{1}{2} \delta Z_k^A (\omega_{\mathcal{M}})_{AB} \delta Z_{d-k}^B, \quad (2.5)$$

which induces a Poisson bracket $[\cdot, \cdot]$ on $\mathcal{M}^\mathcal{X}$. This Poisson bracket can be conveniently expressed in terms of “test functions” as in [2]. Given arbitrary functions ϵ, η on \mathcal{X} —which correspond to differential forms on X since $\mathcal{X} = T[1]X$ —they write

$$\left[\int_{\mathcal{X}} \mathbf{Z}^A \epsilon, \int_{\mathcal{X}} \mathbf{Z}^B \eta \right] = (-1)^{(B+n)\epsilon+d} \int_{\mathcal{X}} (Z^A, Z^B)_{\mathcal{M}} \epsilon \eta, \quad (2.6)$$

where in the exponent we have used the shorthand B, ϵ for the degrees of the respective functions. From (2.5) and (2.6) we can see that if Z^A is dual to Z^B on \mathcal{M} , then Z_k^A will be dual to Z_{d-k}^B on $\mathcal{M}^\mathcal{X}$. Furthermore, if we are working in Darboux coordinates, so that components of $\omega_{\mathcal{M}}$ are constant, then by performing a Hodge decomposition

$$\Omega^k(X) = \mathcal{H}^k \oplus d\Omega^{k-1} \oplus d^\dagger \Omega^{k+1} \quad (2.7)$$

with respect to some arbitrary metric, exact forms Z_k^A will be dual to coexact Z_{d-k}^B and harmonic forms will be dual to harmonic forms. For convenience we introduce orthogonal projectors

$$P_{\mathcal{H}}, \quad P_{\text{ex}}, \quad P_{\text{co}} \quad (2.8)$$

onto harmonic, exact, and coexact forms, respectively.

The Q structure D on $\mathcal{M}^\mathcal{X}$ is defined as the Hamiltonian vector field

$$D = d + Q_{\mathcal{M}}, \quad D = [\Theta_{\mathcal{M}^\mathcal{X}}, \cdot], \quad (2.9)$$

where the Hamiltonian is

$$\Theta_{\mathcal{M}^\mathcal{X}} = (-1)^d \int_{\mathcal{X}} \Theta_{\mathcal{M}} + (-1)^{d+n+1} \int_{\mathcal{X}} \iota_d \vartheta_{\mathcal{M}}, \quad (2.10)$$

³Exploiting Batchelor’s theorem to write \mathcal{M} as a graded vector bundle only improves this situation in that some Z_0 take values in a vector bundle as well.

where each term generates the lift of $Q_{\mathcal{M}}$ and d to $\mathcal{M}^{\mathcal{X}}$, respectively. Note that implicit in this formula is the fact that we have pulled back/transgressed $\Theta_{\mathcal{M}}, \vartheta_{\mathcal{M}}$ to objects on \mathcal{X} ; we have used boldface to highlight this. The signs are such that $D = d_{\mathcal{X}} + Q_{\mathcal{M}}$.

B. The coisotropic

We need to perform a coisotropic reduction to obtain a finite dimensional NQP manifold. This is a generalization of symplectic reduction for Poisson manifolds which requires a coisotropic ideal $\mathcal{I} \subset C^{\infty}(\mathcal{M}^{\mathcal{X}})$, i.e., an ideal that satisfies

$$[\mathcal{I}, \mathcal{I}] \subseteq \mathcal{I}. \quad (2.11)$$

The description of the quotient manifold is given in two equivalent ways. In one description, we take the submanifold $\mathcal{C} \subset \mathcal{M}^{\mathcal{X}}$ defined by the vanishing of \mathcal{I} and quotient by transformations generated by \mathcal{I} . Alternatively, we can describe the structure sheaf of the quotient manifold as the normalizer $N(\mathcal{I})$ of \mathcal{I} , quotiented by \mathcal{I} . That is,

$$\mathcal{W} = \mathcal{C}/[\mathcal{I}, \cdot] \Leftrightarrow C^{\infty}(\mathcal{W}) = N(\mathcal{I})/\mathcal{I}. \quad (2.12)$$

Such a manifold has a natural Poisson structure induced from that on the mapping space; see Appendix C for a review. Further, provided the ideal is closed with respect to the Q structure, i.e., $D\mathcal{I} = [\Theta_{\mathcal{M}^{\mathcal{X}}}, \mathcal{I}] \subseteq \mathcal{I}$, the reduced space has a Q structure induced from the image of the Hamiltonian function under the quotient map:

$$\Theta_{\mathcal{W}} = \Pi(\Theta_{\mathcal{M}^{\mathcal{X}}}), \quad \Pi: N(\mathcal{I}) \rightarrow N(\mathcal{I})/\mathcal{I}. \quad (2.13)$$

This closure is precisely the statement that $\Theta_{\mathcal{M}^{\mathcal{X}}} \in N(\mathcal{I})$.

We build our ideal $\mathcal{I} = \langle \mathcal{I}_{\mathcal{N}}, \mathcal{I}_{\mathcal{Y}} \rangle$ in two parts, each defining a restriction to some submanifold of $\mathcal{M}^{\mathcal{X}} = \mathcal{N}^{\mathcal{X}} \times \mathcal{Y}^{\mathcal{X}}$. This factorization is convenient because \mathcal{Y} may be thought of as “longitudinal” to the cycle to be wrapped, while \mathcal{N} is “transverse.”

On $\mathcal{Y}^{\mathcal{X}}$, we would like the maps in degree 0 to restrict to the fixed wrapping map $\mathfrak{w}: X \rightarrow Y$. This restriction is naturally given by the zero locus of the ideal generated by $y - \mathfrak{w}$ and its closure under D . Using (2.2) and (2.10), we find

$$\mathcal{I}_{\mathcal{Y}} = \langle y - \mathfrak{w}, \xi + d\mathfrak{w} \rangle. \quad (2.14)$$

This is clearly coisotropic in the coordinates on \mathcal{Y} . The angled brackets $\langle \cdot \cdot \cdot \rangle$ will always denote the ideal generated by $\cdot \cdot \cdot$.

On $\mathcal{N}^{\mathcal{X}}$, we follow [15] and take the coisotropic submanifold to consist—in the first instance—of closed maps under the transgressed differential \mathbf{d} on $\mathcal{N}^{\mathcal{X}}$. In degree 0 we realize this via a choice of degree preserving

embedding $N \hookrightarrow \mathcal{N}^{\mathcal{X}}$. By degree counting this is a map of (ordinary) manifolds $N \hookrightarrow \mathcal{N}^{\mathcal{X}}$, and we choose this to be the map sending each $n \in N$ to the constant map $X \rightarrow \{n\}$ (which is \mathbf{d} -closed). Beyond degree 0, we simply set the coisotropic part of each coordinate in the superfield expansion (2.3) to zero (using the Hodge decomposition). Therefore we define $\mathcal{I}_{\mathcal{N}}$ such that⁴

$$\mathcal{I}_{\mathcal{N}} \supset \langle P_{\text{co}} z_k^A \rangle \quad (2.15)$$

for all values of k in the expansion (2.3), where z^A is a generic coordinate on \mathcal{N} . If we consider the vanishing locus of $\mathcal{I}_{\mathcal{N}}$ and $\mathcal{I}_{\mathcal{Y}}$ simultaneously, we see that we are restricted to $x_0 = \text{const}$ and $y_0 = \mathfrak{w}$. This gives an embedding $N \hookrightarrow \mathcal{M}^{\mathcal{X}}$. Similarly, a choice of degree preserving embedding $N \hookrightarrow \mathcal{M}^{\mathcal{X}}$ defines our ideal in degree 0.

This alone is not enough as we would like the reduced manifold \mathcal{W} to be an N manifold, i.e., a graded manifold with non-negative coordinates, such that in degree 0 the structure sheaf is that of an ordinary manifold.⁵ To remove these, we include harmonic generators of the maps z_k^A for maps such that $\deg z^A - k \leq 0$. The exception to this is the maps x_0 for which we do not include the harmonic (i.e., constant map) representatives. We therefore have

$$\mathcal{I}_{\mathcal{N}} = \left\langle \begin{cases} P_{\text{co}} z_k^A, P_{\mathcal{H}} z_{k'}^A & \forall k' \geq \deg z^A \quad \text{if } \deg z^A > 0 \\ P_{\text{co}} z_k^A, P_{\mathcal{H}} z_k^A & \forall k' > 0 \quad \text{if } \deg z^A = 0 \end{cases} \right\rangle. \quad (2.16)$$

To see that this is coisotropic, we use (2.6) and the surrounding discussion to note that the coexact generators are dual to exact generators. Hence, these terms are coisotropic with respect to all of $\mathcal{I}_{\mathcal{N}}$. The harmonic generators could be dual to some other harmonic generator in $\mathcal{I}_{\mathcal{N}}$. However, since we only include harmonic z_k^A for $0 \geq \deg z^A - k$, the dual coordinate $z_{k'}^B$ on $\mathcal{M}^{\mathcal{X}}$ has

$$\begin{aligned} \deg z^B - k' &= n - \deg z^A - (d - k) \\ &= (n - d) - (\deg z^A - k) \\ &> 0 \end{aligned} \quad (2.17)$$

so is not included in $\mathcal{I}_{\mathcal{N}}$. The total ideal $\mathcal{I} = \langle \mathcal{I}_{\mathcal{N}}, \mathcal{I}_{\mathcal{Y}} \rangle$ is coisotropic, as required.

C. Metric and coordinate independence

The construction of the ideal \mathcal{I} appears to rely on a choice of metric on X but we claim that the resulting reduced manifold depends on only the topological

⁴The reduction by such a coisotropic ideal is related to the reduction by “contractible pairs” in the BV formalism [21].

⁵There are issues not just with negative-graded but also with degree 0 “formal” coordinates; see, e.g., [22, Sec. 2].

data of X . This can be best seen from the vanishing locus $\mathcal{C} \subset \mathcal{M}^{\mathcal{X}}$. This is given by maps which are either d-closed (for $\deg z^A = 0$ and some values of k, k') or ones which are also d-exact (in all other cases), as specified in (2.16). The metric only appears in the specification of the vanishing ideal that represents \mathcal{C} but \mathcal{C} does not in itself depend on metric data. (This apparent metric dependence thus may perhaps be seen as due to “gauge-fixing.”)

Our construction also appears to depend on a choice of coordinates on \mathcal{N} . To see that this is well defined, we will show that the submanifold \mathcal{C} is invariant under a change of coordinates $\tilde{z}^A = \tilde{z}^A(z)$. Using formula (2.4), we can write the k -form component of the transgressed \tilde{z} as

$$\tilde{z}_k^A \sim \sum_j C^A_{A_1 \dots A_j}(z_0) z_{k_1}^{A_1} \dots z_{k_j}^{A_j} \quad (2.18)$$

such that $k_1 + \dots + k_j = k$ and $\deg z^{A_1} + \dots + \deg z^{A_j} \leq \deg \tilde{z}_k^A$; here we emphasize that the last inequality holds true because \mathcal{M} —and thus \mathcal{N} —was assumed to be an N manifold (its structure sheaf is non-negatively graded). Restricting to \mathcal{C} , the coordinates $z_{k_i}^{A_i}$ are all closed under d and hence so is \tilde{z}_k^A . Further, if $\deg \tilde{z}^A - k \leq 0$, then at least one of $\deg z^{A_i} - k_i \leq 0$. This means that $z_{k_i}^{A_i}$ is exact. A product of closed and exact forms is exact and hence so is \tilde{z}_k^A as required.

D. Closure under D

The final condition to check is that the ideal \mathcal{I} is closed under the Q structure $D = [\Theta_{\mathcal{M}^{\mathcal{X}}}, \cdot]$ on $\mathcal{M}^{\mathcal{X}}$. We have already checked that $\mathcal{I}_{\mathcal{Y}}$ is D -closed, and so we need only check how D acts on the generating coordinates of $\mathcal{I}_{\mathcal{N}}$. We can use formula (2.6) with $P_{\text{co}}\epsilon = 0$. This choice of epsilon selects out the harmonic and coexact generators z_k^A , respectively. We have

$$\begin{aligned} \left[\Theta_{\mathcal{M}^{\mathcal{X}}}, \int_{\mathcal{X}} z^A \epsilon \right] &= (-1)^d \left[\int_{\mathcal{X}} \Theta_{\mathcal{M}}, \int_{\mathcal{X}} z^A \epsilon \right] \\ &\quad + (-1)^{d+n+1} \left[\int_{\mathcal{X}} l_d \vartheta_{\mathcal{M}}, z^A \epsilon \right] \\ &= \int_{\mathcal{X}} (\Theta_{\mathcal{M}}, z^A)_{\mathcal{M}} \epsilon + \int_{\mathcal{X}} dz^A \epsilon. \end{aligned} \quad (2.19)$$

The second term vanishes when ϵ is closed. We therefore need to consider only the first term. We can see whether this term is contained within \mathcal{I} by transgressing the function $(\Theta_{\mathcal{M}}, z^A)_{\mathcal{M}}$ to the mapping space and evaluating it over \mathcal{C} . If the integral vanishes when integrated against all closed ϵ , then the ideal is closed under D . Using the form of the Hamiltonian function we find

$$(\Theta_{\mathcal{M}}, z^A)_{\mathcal{M}} \propto (\omega_{\mathcal{M}})^{AB} \sum_k \frac{\partial \theta_k(z, y)}{\partial z^B} \xi^k. \quad (2.20)$$

Transgressing this to the mapping space and restricting to \mathcal{C} , we replace $y \rightarrow \mathbf{w}$, $\xi \rightarrow d\mathbf{w}$, $z \rightarrow z$ for z -closed (or exact). Integrating this against ϵ we get

$$\left[\Theta_{\mathcal{M}^{\mathcal{X}}}, \int_{\mathcal{X}} z^A \epsilon \right]_{\mathcal{C}} \propto \int_{\mathcal{X}} \left((\omega_{\mathcal{M}})^{AB} \mathbf{w}^* \sum_k \frac{\partial \theta_k(z)}{\partial z^B} \right) \epsilon. \quad (2.21)$$

We require this to vanish for the above ϵ . When ϵ is exact, this indeed vanishes if we impose $d_y \theta_k = 0$. If ϵ is harmonic, however, we find constraints on the coefficients θ_k that we address case-by-case, in general.

E. The reduction, metric independence, and homotopy invariance

Given the coisotropic reduction \mathcal{I} , we consider the reduction given in (2.12). We will consider the structure sheaf construction of the reduced manifold. The normalizer $N(\mathcal{I})$ of the ideal is generated by

$$N(\mathcal{I}) \sim \{ P_{\mathcal{H}} x_0, P_{\mathcal{H}} z_k^A, \mathcal{I} | 0 < \deg z^A - k \leq n - d \}. \quad (2.22)$$

We can expand the $P_{\mathcal{H}} z_k^A = z_k^{A,a} e_a$ in some basis $\{e_a\}$ of \mathcal{H}^k , so the $z_k^{A,a}$ are constant parameters of degree $\deg z^A - k$. In the case that the respective cohomology group is one dimensional (e.g., for $\mathcal{H}^0, \mathcal{H}^d$) we will omit the a index and simply identify, e.g., $z_d^A = z_d^A \text{vol}_X$. We see that the structure sheaf $C^{\infty}(\mathcal{W}) = N(\mathcal{I})/\mathcal{I}$ is given therefore generated by

$$C^{\infty}(\mathcal{W}) = N(\mathcal{I})/\mathcal{I} \sim \{ x_0, z_k^{A,a} | 0 < \deg z^A - k \leq n - d \}. \quad (2.23)$$

That is, the structure sheaf is given by all smooth functions in the $z_k^{A,a}$ (and x_0).

The Hamiltonian function on \mathcal{W} is given by the projection $\Pi: N(\mathcal{I}) \rightarrow N(\mathcal{I})/\mathcal{I}$ of $\Theta_{\mathcal{M}^{\mathcal{X}}}$. We will confirm in the examples that the final result is given by

$$\Theta_{\mathcal{W}} = \Pi(\Theta_{\mathcal{M}^{\mathcal{X}}}) = \int_{\mathcal{X}} \sum_k (-1)^k \mathbf{w}^* \theta_k(z), \quad (2.24)$$

where the z are now the harmonic representatives of the cohomology groups on X . Expanding the harmonic z_k^A in terms of the constant coordinates $z_k^{A,a}$, we can perform the integral over X with the convention that the volume form is on the right of the integrand, so we pull out constants from the left. Once this is done, the final result will no longer be an integral but will be a function in the $z_k^{A,a}$ which will involve, in general, a sum over cohomology groups, which will be discrete in all cases (we only consider wrapping over compact cycles).

Formula (2.24) seems to depend on some metric to choose the harmonic representatives for the z . However, under a change of metric, the harmonic representatives change by a d-exact term, and since we have assumed that the forms θ_k are closed, this shift will not change the integral. Furthermore, since the forms θ_k are closed, the evaluation of the integral only depends on the homotopy class of $\mathfrak{w}: X \rightarrow Y$. Therefore the construction is metric-independent and homotopy invariant.

Now that we have defined the reduction in complete generality, we will see many examples of how this works in practice. There are three interesting cases to consider:

- (1) $\dim X = 0$ —The process effectively shrinks Y to a point.
- (2) $\dim Y = 0$ —We produce a QP manifold with the same underlying commutative manifold but with a different degree.

\mathcal{N}				
coord	x^μ	ψ^μ	χ_μ	p_μ
deg	0	1	$n-1$	n

- (3) $X = Y$ —We produce a QP manifold which corresponds to a brane wrapping the internal manifold.

III. EXAMPLE— $\dim X = 0$

We consider first a simple example to show that in the simple case that $\dim X = 0$, our procedure effectively reduces to dimensional reduction on Y . Consider the ingredients

$$\mathcal{M} = T^*[n]T[1](N \times Y), \quad X = \text{pt}. \quad (3.1)$$

Taking $\mathcal{N} = T^*[n]T[1]N$, $\mathcal{Y} = T^*[n]T[1]Y$, and $\mathcal{X} = T[1]X = \text{pt}$, we introduce the Darboux coordinates

\mathcal{Y}				
coord	y^m	ξ^m	ϕ_m	q_m
deg	0	1	$n-1$	n

We will take the QP structure to be given by the symplectic form and Hamiltonian function

$$\omega_{\mathcal{M}} = dpdx + dqdy - d\psi d\chi - d\xi d\phi, \quad (3.3)$$

$$\Theta_{\mathcal{M}} = -\psi p - \xi q + \frac{1}{n!} F_n \psi^n + \frac{1}{(n-1)!} F_{n-1} \psi^{n-1} \xi + \cdots + \frac{1}{d!(n-d)!} F_{n-d} \psi^{n-d} \xi^d. \quad (3.4)$$

We have suppressed all indices but they should be read as being contracted in the natural way. The coefficients F_k can be thought of as elements of $\Omega^k(N) \times \Omega^{n-k}(Y)$. These should be closed under the differential d_Y on Y . So, for example, $F_n \psi^n = F_n(x, y)_{\mu_1 \dots \mu_n} \psi^{\mu_1} \dots \psi^{\mu_n}$ should be viewed as a differential n -form on N , but a constant function on Y . In the ansatz above, we have assumed a trivial connection on the bundle. We can easily reintroduce it by making the replacement $\xi \rightarrow \mathcal{A} = \xi + A\psi$, where A is the connection; however, it will not change our final result, so we omit it for simplicity.

The first step in the reduction process is to transgress the QP structure to $\mathcal{M}^{\mathcal{X}}$. But since \mathcal{X} is zero dimensional, we have $\mathcal{M}^{\mathcal{X}} \simeq \mathcal{M}$. Next, we need to choose a wrapping map $\mathfrak{w}: X = \text{pt} \rightarrow Y$ or, equivalently, a (degree preserving) embedding $N \hookrightarrow \mathcal{M}^{\mathcal{X}} \simeq \mathcal{M}$. This is equivalent to choosing some point $\hat{y} \in Y$ and defining the embedding $N \rightarrow (N, \hat{y}) \subset M$. This is described by the ideal

$$\mathcal{I}_0 = \langle y^m - \hat{y}^m \rangle. \quad (3.5)$$

We then want to form the closure of this ideal with respect to differential Q on \mathcal{M} . We get

$$\mathcal{I}_Y = \langle \mathcal{I}_0, Q\mathcal{I}_0 \rangle = \langle y^m - \hat{y}^m, \xi^m \rangle. \quad (3.6)$$

It is easy to check from (3.3) that this is indeed coisotropic with respect to the Poisson bracket on \mathcal{M} . In principal, we also need to restrict the maps into \mathcal{N} to those that are closed/exact with respect to d on X . However, since $\dim X = 0$, this is a trivial constraint and so we just have $\mathcal{I} = \mathcal{I}_Y$.

To perform the coisotropic reduction, we need to go to first find the normalizer $N(\mathcal{I})$ of \mathcal{I} , which can easily be verified to be generated by the coordinates

$$N(\mathcal{I}) \sim \{x^\mu, \psi^\mu, \chi_\mu, p_\mu, y^m - \hat{y}^m, \xi^m\}. \quad (3.7)$$

The structure sheaf of the new QP manifold \mathcal{W} is then defined to be the quotient of this by the ideal \mathcal{I} . That is, $C^\infty(\mathcal{W}) = N(\mathcal{I})/\mathcal{I}$, which is generated by

$$N(\mathcal{I})/\mathcal{I} \sim \{x^\mu, \psi^\mu, \chi_\mu, p_\mu\} \Rightarrow \mathcal{W} = T^*[n]T[1]N. \quad (3.8)$$

Note that by construction $\Theta_{\mathcal{M}} \in N(\mathcal{I})$, and so we can find the new Hamiltonian function through the natural projection $\Pi: N(\mathcal{I}) \rightarrow N(\mathcal{I})/\mathcal{I}$, which gives

$$\Theta_{\mathcal{W}} = \Pi(\Theta_{\mathcal{M}}) = -\psi p + \frac{1}{n!} F_n(x, \hat{y}) \psi^n, \quad (3.9)$$

and the final symplectic form is

$$\omega_{\mathcal{W}} = dpdx - d\psi d\chi. \quad (3.10)$$

We see that this procedure has produced a new QP manifold with the same degree but with underlying commutative manifold N . We see that we have effectively collapsed Y to the point \hat{y} . In the case where Y is a Lie group, we find the same result as in symplectic reduction modulo $T[1]Y$ [13]. If we were to choose a different wrapping map $\mathfrak{w}' : X \mapsto \hat{y}'$ that is homotopic to $\mathfrak{w} : X \mapsto \hat{y}$, then we end up with the same graded manifold where the Q structure is evaluated for $F_n(x, \hat{y}')$. However, the condition that the F_n is closed on Y says that it is constant, and hence the Q structures are the same. This demonstrates the homotopy invariance of our construction.

With regards to applications to AKSZ sigma models, which feature prominently in the rest of the paper, this example is less interesting on account of the following: the AKSZ sigma model is based on the space of (super)fields $\mathcal{M}^{\mathcal{X}}$, which in this case is isomorphic to \mathcal{M} itself, since $\mathcal{X} = \text{point}$. This example retains physical significance in general, because it captures *direct* dimensional reduction (where there is no wrapping along the Y manifold).

IV. EXAMPLES— $\dim Y = 0$

A. n -Brane $\rightarrow (n-1)$ -brane

Let us now consider the same example as above, but instead of having $\dim X = 0$, we will take the dimension of the fiber $\dim Y = 0$ and take X to be nontrivial. We will take the ingredients

$$\mathcal{M} = T^*[n]T[1]M, \quad X = S^1. \quad (4.1)$$

We will use the homogeneous coordinates

\mathcal{M}				
coord	x^μ	ψ^μ	χ_μ	p_μ
deg	0	1	$n-1$	n

and use the coordinates $\sigma, d\sigma$ on $\mathcal{X} = T[1]S^1$. The Hamiltonian function and symplectic form are given by

$$\omega_{\mathcal{M}} = dpdx - d\psi d\chi, \quad (4.3)$$

$$\Theta_{\mathcal{M}} = -\psi p + \frac{1}{n!} F_n \psi^n. \quad (4.4)$$

Since $Y = \text{pt}$ in this example, we do not need to impose any constraints on the coefficients F_n .

We need to transgress this structure to the mapping space $\mathcal{M}^{\mathcal{X}}$. This is now an infinite dimensional graded manifold

whose points $f \in \mathcal{M}^{\mathcal{X}}$ can be described by their pullback action on coordinates on \mathcal{M} . That is, we have

$$f^* Z^A = Z^A(\sigma, d\sigma) = Z_0^A(\sigma) + Z_1^A(\sigma) d\sigma. \quad (4.5)$$

The transgressed Hamiltonian function is given by

$$\begin{aligned} \Theta_{\mathcal{M}^{\mathcal{X}}} &= (-1)^1 \int_{\mathcal{X}} \Theta_{\mathcal{M}} + (-1)^{n-1+1} \int_{\mathcal{X}} \iota_d \mathbf{Z}^A \\ &= - \int_{T[1]S^1} -\psi p + \frac{1}{n!} F_n(x) \psi^n \\ &\quad + (-1)^n \int_{T[1]S^1} p dx - \frac{1}{n} (\psi d\chi + (n-1)\chi d\psi). \end{aligned} \quad (4.6)$$

The boldfaced letters in the expression correspond to functions pulled back to functions on \mathcal{X} as in (4.5). The Berezin integral over $T[1]S^1$ selects the maximal degree component of the integrand (i.e., the one-form components) and integrates it over S^1 . Our convention is that we normalize with an overall factor of $\text{vol}(S^1)$, and so for the flat metric on S^1 we have

$$\int_{T[1]S^1} \dots = \frac{1}{2\pi} \int_{S^1} (\dots)_1. \quad (4.7)$$

The next step is to define the coisotropic ideal $\mathcal{I} = \langle \mathcal{I}_{\mathcal{N}}, \mathcal{I}_{\mathcal{Y}} \rangle$. Since Y is trivial, so is the ideal $\mathcal{I}_{\mathcal{Y}}$, and hence we need only determine $\mathcal{I}_{\mathcal{N}}$. Following Sec. II, we first start by restricting to all closed maps. That is, we take

$$\mathcal{I}_{\mathcal{N}} \supset \langle P_{\text{co}} x_k, P_{\text{co}} \psi_k, P_{\text{co}} \chi_k, P_{\text{co}} p_k \rangle. \quad (4.8)$$

To define this ideal we choose some arbitrary metric on S^1 , and for simplicity we can take the flat metric. We then also add the harmonic representatives for Z_k^A such that $\deg Z^A - k \leq 0$ (except for x_0). This gives

$$\mathcal{I}_{\mathcal{N}} = \langle P_{\text{co}} x_0, P_{\text{co}} \psi_0, P_{\text{co}} \chi_0, P_{\text{co}} p_0, P_{\mathcal{H}} x_1, P_{\mathcal{H}} \psi_1 \rangle. \quad (4.9)$$

Again, in degree 0, the vanishing locus of this ideal restricts us to maps $x_0 = \text{const}$ and hence defines a natural embedding $M \hookrightarrow \mathcal{M}^{\mathcal{X}}$. It is a quick check using (2.6) that this ideal is coisotropic. Indeed, the Poisson bracket of the coexact generators with any other generators will vanish, as they are dual to exact maps. The harmonic x_1, ψ_1 representatives are dual to $p_0, \chi_0 \in \mathcal{H}^0$, respectively, and these do not appear in the generating set of $\mathcal{I}_{\mathcal{N}}$.

We will also verify that this ideal is closed with respect to the Q structure (4.6). Using the test function form of the Poisson bracket (2.6), we can calculate D acting on the generators by calculating

$$\begin{aligned} \left[\Theta_{\mathcal{M}^x}, \int_{T[1]S^1} \mathbf{Z}^A \epsilon \right] &= - \left[\int_{T[1]S^1} \Theta_{\mathcal{M}}, \int_{T[1]S^1} \mathbf{Z}^A \epsilon \right] \\ &+ (-1)^n \left[\int_{T[1]S^1} \iota_d \vartheta_{\mathcal{M}}, \int_{T[1]S^1} \mathbf{Z}^A \epsilon \right], \end{aligned} \quad (4.10)$$

where ϵ is a function on \mathcal{N} that is closed under d . Taking $\epsilon \in \mathcal{H}^k$ selects the harmonic representative $Z_{1-k} \in \mathcal{H}^{1-k}$, while taking ϵ to be exact selects the coexact representative of Z_0^A . The second term gives us

$$\left[\int_{T[1]S^1} \iota_d \vartheta_{\mathcal{M}}, \int_{T[1]S^1} \mathbf{Z}^A \epsilon \right] \propto \int_{T[1]S^1} d\mathbf{Z} \epsilon = \frac{1}{2\pi} \int_{S^1} dZ_0^A \epsilon_0. \quad (4.11)$$

Taking ϵ to be closed tells us that ϵ_0 is constant. The integrand on the right-hand side is therefore exact and so the integral vanishes. The Poisson bracket is then determined by the first term alone which is proportional to

$$\left[\Theta_{\mathcal{M}^x}, \int_{T[1]S^1} \mathbf{Z}^A \epsilon \right] \propto \int_{T[1]S^1} (\Theta_{\mathcal{M}}, \mathbf{Z}^A)_M \epsilon, \quad (4.12)$$

where the function $(\Theta_{\mathcal{M}}, Z^A)$ is transgressed to the mapping space. We can use these results to confirm that $\Theta_{\mathcal{M}^x}$ lies always in $\mathcal{I}_{\mathcal{N}}$ as outlined in Sec. II. The only nontrivial checks are for the harmonic generators x_1, ψ_1 , for which we take $\epsilon = \epsilon_0$ to be constant. We have

$$(\Theta_{\mathcal{M}}, x)_M = \psi, \quad (\Theta_{\mathcal{M}}, \psi)_M = 0. \quad (4.13)$$

Transgressing these functions and evaluating on \mathcal{C} , we take ψ_1 to be exact. Hence, both vanish under the integral (4.12) when $\epsilon = \epsilon_0$ is constant. This proves that the ideal is closed under D .

Now that we have our coisotropic ideal, we perform the coisotropic reduction. The normalizer of \mathcal{I} is generated by all the coordinates that are not dual to those in \mathcal{I} ,

$$N(\mathcal{I}) \sim \{P_{\mathcal{H}}x_0, P_{\mathcal{H}}\psi_0, P_{\mathcal{H}}\chi_1, P_{\mathcal{H}}p_1, \mathcal{I}\}. \quad (4.14)$$

The structure sheaf for \mathcal{W} is then $N(\mathcal{I})/\mathcal{I}$, which is generated by

$$N(\mathcal{I})/\mathcal{I} \sim \{x_0, \psi_0, \chi_1, p_1\} \Rightarrow \mathcal{W} = T^*[n-1]T[1]M. \quad (4.15)$$

(Note in the expression above we are now working in the coordinates $z_k^{A,a}$ described in Sec. II: $P_{\mathcal{H}}x_0 = x_0 \cdot 1$ and $P_{\mathcal{H}}p_1 = p_1 \text{vol.}$) Thus we restrict to harmonic functions for x, ψ —so they retain their original degrees—while we restrict to harmonic one-forms for χ, p ; hence, they have their degrees shifted down by 1. We therefore end up with the manifold $T^*[n-1]T[1]M$.

To find the symplectic form, we use the Poisson brackets (2.6) with the ϵ, η appropriate harmonic representatives. We find that

$$\omega_{\mathcal{W}} = -dp_1 dx_0 - d\psi_0 d\chi_1. \quad (4.16)$$

To find the form of the Hamiltonian function we project $\Theta_{\mathcal{M}^x}$ under $\Pi: N(\mathcal{I}) \rightarrow N(\mathcal{I})/\mathcal{I}$. By restricting all coordinates to the harmonic representatives on which $d = 0$, we find $\Pi(\iota_d \bar{\vartheta}) = 0$. The term $\frac{1}{n!} \eta^n$ gets projected to $\frac{1}{n!} F_n(x_0) \psi_0^n$ which is a function on S^1 and hence vanishes under the Berezin integral. We find that we are left with⁶

$$\Theta_{\mathcal{W}} = \Pi(\Theta_{\mathcal{M}^x}) = \psi_0 p_1. \quad (4.17)$$

Making the change of coordinates $p_1 \rightarrow -p_1$ puts the QP manifold in the canonical form for a $(n-1)$ -brane. Interestingly, all flux twisting drops out of the Hamiltonian function in this case. This is what happens in the zero-wrapping sector of wrapped branes where physically one ends up with a tensionless brane [23]. These are somewhat pathological, and hence the physical interpretation of such reductions is less clear. We will see that one can get more interesting reductions if one allows X to wrap some part of M .

B. From Courant to Poisson

Using the formulation set out, we can already find novel relations between QP manifolds, and their associated AKSZ sigma models. Suppose M is a Poisson manifold with Poisson bivector π . There are at least two distinct ways to realize this structure as a QP structure.⁷ First, we can take the straight cotangent lift of π to obtain the following QP manifold:

$$\mathcal{W} = T^*[1]M \quad \begin{array}{c|cc} \text{coord} & \tilde{x} & \tilde{p} \\ \hline \text{deg} & 0 & 1 \end{array} \quad \begin{array}{l} \omega_{\mathcal{W}} = d\tilde{p} d\tilde{x}, \\ \Theta_{\mathcal{W}} = \frac{1}{2} \pi \tilde{p}^2. \end{array} \quad (4.18)$$

A quick calculation shows that $(\Theta_{\mathcal{W}}, \Theta_{\mathcal{W}}) = 0$ if and only if π is Poisson.

Alternatively, we can consider the Lie algebroid structure on T^*M whose anchor map is given by the bivector $\pi: T^*M \rightarrow TM$ and whose bracket is given by

$$[\alpha, \beta] = \mathcal{L}_{\pi(\alpha)} \beta - \iota_{\pi(\beta)} d\alpha. \quad (4.19)$$

This Lie algebroid can be lifted to a Dirac structure $L = \{\lambda + \pi(\lambda) | \lambda \in T^*M\}$ within the Courant algebroid $TM \oplus T^*M$. The Courant algebroid and the differential d_L associated with the Dirac structure can be lifted to a QP manifold structure via

⁶Our conventions are that we integrate with the volume form on the right of the integrand, and so we pull constants out from the left. This gives the overall sign.

⁷See, e.g., examples in the work of Voronov [24].

$$\mathcal{M} = T^*[2]T[1]M \quad \begin{array}{c|ccccc} \text{coord} & x & \psi & \chi & p \\ \hline \text{deg} & 0 & 1 & 1 & 2 \end{array} \quad \begin{aligned} \omega_{\mathcal{M}} &= dp dx - d\psi d\chi, \\ \Theta_{\mathcal{M}} &= -\pi p \chi + \frac{1}{2} \partial \pi \psi \chi^2. \end{aligned} \quad (4.20)$$

Once again $(\Theta_{\mathcal{M}}, \Theta_{\mathcal{M}}) = 0$ if and only if π is Poisson. We have suppressed indices for convenience. We want to see if we can pass from (4.20) to (4.18) via our brane wrapping procedure.

Let us perform a circle reduction of \mathcal{M} as above. We transgress the structure to $\mathcal{M}^{\mathcal{X}}$ where $\mathcal{X} = T[1]S^1$. As before, we define $\mathcal{I} = \mathcal{I}_{\mathcal{N}}$ by first including all coexact generators

$$\mathcal{I} \supset \langle P_{\text{co}}x_0, P_{\text{co}}\psi_0, P_{\text{co}}\chi_0, P_{\text{co}}p_0 \rangle. \quad (4.21)$$

Then we include harmonic representatives to remove coordinates of zero or negative degree. We will slightly relax the construction set out in Sec. II by allowing some new coordinates of degree 0.⁸ We will define

$$\mathcal{I} = \langle P_{\text{co}}x_0, P_{\text{co}}\psi_0, P_{\text{co}}\chi_0, P_{\text{co}}p_0, P_{\mathcal{H}}x_1, P_{\mathcal{H}}\chi_1 \rangle. \quad (4.22)$$

As before, this ideal is coisotropic.

We will check the closure of this ideal with respect to the Q structure D on $\mathcal{M}^{\mathcal{X}}$. The transgressed Hamiltonian function is

$$\begin{aligned} \Theta_{\mathcal{M}^{\mathcal{X}}} &= - \int_{T[1]S^1} \Theta_{\mathcal{M}} + \int_{T[1]S^1} \iota_d \Theta_{\mathcal{M}} \\ &= - \int_{T[1]S^1} -\pi(x)p\chi + \frac{1}{2} \partial \pi(x)\psi\chi^2 \\ &\quad + \int_{T[1]S^1} p dx - \frac{1}{2} (\psi d\chi + \chi d\psi). \end{aligned} \quad (4.23)$$

We then act with this on $\int \mathbf{Z}^A \epsilon$ for some test function ϵ that must be harmonic or exact. As in (4.12), the only nontrivial constraint to check is for the harmonic representatives. We need to check if the following vanishes:

$$\int_{T[1]S^1} (\Theta_{\mathcal{M}}, \mathbf{Z}^A)_{\mathcal{M}} \epsilon, \quad (4.24)$$

whenever the function $(\Theta_{\mathcal{M}}, \mathbf{Z}^A)$ is transgressed and evaluated on \mathcal{C} , and if ϵ is harmonic. Since the only harmonic generators of \mathcal{I} are x_1, χ_1 , we calculate

⁸The construction, as set out previously, would still work in this case but we would end up with a trivial Q structure. To result in a QP manifold with a nontrivial Q structure, we will need to perform an intermediate step before removing the additional degree 0 coordinates.

$$(\Theta_{\mathcal{M}}, x) = \pi(x)\chi, \quad (\Theta_{\mathcal{M}}, \chi_{\mu}) = \frac{1}{2} \partial \pi(x)\chi^2. \quad (4.25)$$

We transgress these functions to the mapping space and evaluate on the vanishing locus \mathcal{C} of \mathcal{I} . Noting that these are functions of x, χ alone, evaluating them on \mathcal{C} means that the zero-form component must be constant functions on X , while the one-form component must be an exact form. Integrating these against a constant function $\epsilon = \epsilon_0$ selects the one-form component, which is exact, and hence the integral vanishes as required.

The next step is to perform the coisotropic reduction with respect to this ideal. The normalizer is generated by all coordinates not dual to those in \mathcal{I} ,

$$N(\mathcal{I}) \sim \{P_{\mathcal{H}}x_0, P_{\mathcal{H}}\chi_0, P_{\mathcal{H}}\psi_1, P_{\mathcal{H}}p_1, \mathcal{I}\}, \quad (4.26)$$

and so we obtain the structure sheaf $C^{\infty}(\widetilde{\mathcal{M}}) = N(\mathcal{I})/\mathcal{I}$ which is generated by

$$N(\mathcal{I})/\mathcal{I} \sim \{x_0, \psi_1, \chi_0, p_1\} \Rightarrow \widetilde{\mathcal{M}} = T^*[1]TM. \quad (4.27)$$

This time, we restrict to harmonic functions for x, χ so they retain their degree, while we take harmonic one-forms for ψ, p , and hence their degree is shifted down by 1. The resulting Hamiltonian function is $\Pi(\Theta_{\mathcal{M}^{\mathcal{X}}})$ and the symplectic form is derived from the Poisson brackets (2.6) with harmonic representatives for ϵ, η ,

$$\Theta_{\widetilde{\mathcal{M}}} = -\pi p_1 \chi_0 - \frac{1}{2} \partial \pi \psi_1 \chi_0^2, \quad (4.28)$$

$$\omega_{\widetilde{\mathcal{M}}} = -dp_1 dx_0 + d\psi_1 d\chi_0. \quad (4.29)$$

We performed the change of coordinates $p_1 \rightarrow -p_1, \chi_0 \rightarrow -\chi_0$ to remove minus signs.

We have arrived at a “halfway house” QP manifold $\widetilde{\mathcal{M}}$. Interestingly, this is the cotangent lift of the complete lift of the Poisson structure π on M to the tangent bundle (TM, π^c) [25]. That is, given any Poisson structure (M, π) we define a Poisson structure (TM, π^c) by

$$\pi^c = \pi^{\mu\nu} \frac{\partial}{\partial x_0^{\mu}} \frac{\partial}{\partial \psi_1^{\nu}} + \frac{1}{2} \psi_1^{\rho} \partial_{\rho} \pi^{\mu\nu} \frac{\partial}{\partial \psi_1^{\mu}} \frac{\partial}{\partial \psi_1^{\nu}}, \quad (4.30)$$

where x_0 are coordinates on M and ψ_1 are coordinates along the vector bundle fibers. We can reduce the QP manifold $\widetilde{\mathcal{M}}$ further by following [25]. Given any (torsionless) connection on M , we can define a global vector field on TM given by the geodesic spray

$$s = \psi_1^\mu \frac{\partial}{\partial x_0^\mu} - \psi_1^\mu \psi_1^\nu \Gamma_{\mu\nu}^\rho \frac{\partial}{\partial \psi_1^\rho}. \quad (4.31)$$

This has a cotangent lift to $T^*[1]TM$ whose Hamiltonian is

$$S = \psi_1 p_1 - \Gamma \psi_1^2 \chi_0. \quad (4.32)$$

From this, we define a new Hamiltonian function

$$\Theta'_{\mathcal{M}} = -\frac{1}{2}(S, \Theta_{\mathcal{M}}) = \frac{1}{2}\pi p_1^2 + \psi_1 f(x_0, \psi_1, \chi_0, p_1), \quad (4.33)$$

where f is some function of the coordinates whose precise form is not important. All we will need is that besides the first term, $\frac{1}{2}\pi p_1^2$, each term is at least linear in the coordinate ψ_1 .

Consider the ideal generated by the single coordinate $\mathcal{I} = \langle \psi_1 \rangle$. This ideal is automatically closed under the Q structure since

$$\begin{aligned} (\Theta'_{\mathcal{M}}, \psi_1) &= \left(\frac{1}{2}\pi p_1^2 + \psi_1 f(x_0, \psi_1, \chi_0, p_1), \psi_1 \right) \\ &= (\psi_1 f(x_0, \psi_1, \chi_0, p_1), \psi_1) \\ &\propto \psi_1 (f, \psi_1) \\ &\in \mathcal{I}. \end{aligned} \quad (4.34)$$

Performing the coisotropic reduction with respect to this ideal we obtain the structure sheaf

$$N(\mathcal{I})/\mathcal{I} \sim \{x_0, p_1\} \Rightarrow \mathcal{W} = T^*[1]M. \quad (4.35)$$

That is, we reproduce the graded-commutative manifold \mathcal{W} . Further, the symplectic form and Hamiltonian function are easily shown to be the following:

$$\omega_{\mathcal{W}} = dp_1 dx_0, \quad (4.36)$$

$$\Theta_{\mathcal{W}} = \frac{1}{2}\pi p_1^2. \quad (4.37)$$

We see then that we precisely reproduce the QP manifold associated with the cotangent lift of the Poisson bivector that we described at the beginning of this section. Using the results of Sec. VI, this lifts to an association between the Courant and Poisson sigma models themselves. Note that if we had instead started with an H -twisted Courant sigma model, then our procedure would have resulted in the QP manifold associated with the H -twisted Poisson sigma model [26].

This construction provides new links between the Courant sigma models and the Poisson sigma models which are physically and mathematically distinct from previously found associations. Previous work [26–28] found that if one considers the Courant sigma model on a manifold with a boundary, then by studying the consistent

boundary conditions one finds that one can couple the (H -twisted) Poisson sigma model at the boundary.⁹ Our construction, however, requires no boundary and finds that the two theories—described in the QP language by (4.18) and (4.20), respectively—are also related via brane wrapping. (We note here that a similar relation was exhibited in a recent work [29] using a different reduction procedure.) Physically, our procedure should involve a compact cycle within the brane shrinking to zero size. One way to see this is by analogy with Kaluza-Klein theory; we retain the zero modes of superfields, which is precisely analogous to what happens in a KK scenario. This might provide a useful heuristic for understanding the link between these two theories in our construction.

V. EXAMPLES— $X = Y$

We will now generalize the previous two sections to allow for cases where the source manifold X wraps the target space fiber Y . In particular, we will be interested in the case where $X = Y$. We will see that the reduction procedure requires us to choose some self-wrapping map $\mathbf{w}: X \rightarrow X$. The examples we choose are physically motivated and fill our understanding of how brane dualities in M-theory/IIA arise in the QP setting. In particular, when $X = S^1$, we will see that our procedure produces the known relations from M-theory/type IIA duality. We will also see that this procedure reproduces other interesting relations between the M5-brane and the heterotic string [30,31].

A. M2 on S^1

Our first example will be wrapping the M2-brane on an S^1 . This will be very similar to the n -brane example in Sec. IV A, except in this case the wrapping will allow for more interesting Hamiltonian functions to be produced.

We start with the QP manifold \mathcal{M} associated with the M2-brane and a source manifold X :

$$\mathcal{M} = T^*[3]T[1](N \times S^1), \quad X = S^1. \quad (5.1)$$

Writing $\mathcal{N} = T^*[3]T[1]N$, $\mathcal{Y} = T^*[3]T[1]S^1$, and $\mathcal{X} = T[1]S^1$, we will introduce the coordinates

coord	\mathcal{N}				\mathcal{Y}			
	x^μ	ψ^μ	χ_μ	p_μ	coord	y	ξ	ϕ
deg	0	1	2	3	deg	0	1	2

(5.2)

and use coordinates $(\sigma, d\sigma)$ on \mathcal{X} . The Hamiltonian function and symplectic form are

⁹The coupled bulk and boundary theory was called WZW-Poisson theory in [27], or simply WZ-Poisson in [28].

$$\omega_{\mathcal{M}} = dp dx + dq dy - d\psi d\chi - d\xi d\phi, \quad (5.3)$$

$$\Theta_{\mathcal{M}} = -\psi p - \xi q + \frac{1}{4!} F_4 \psi^4 + \frac{1}{3!} H_3 \psi^3 \xi. \quad (5.4)$$

We require F_4 to be d_{S^1} -closed; we will also use the fact that $H_3 \xi$ is d_{S^1} -closed (which is automatic).

We transgress this to the mapping space $\mathcal{M}^{\mathcal{X}}$ and choose an ideal whose vanishing locus describes, in degree 0, some embedding $\iota: N \hookrightarrow \mathcal{M}^{\mathcal{X}}$. As explained in Sec. II, this depends on the choice of some wrapping map $\mathfrak{w}: S^1 \rightarrow S^1$. In fact, as stated, the final result only depends on the homotopy class of w , and hence we can take, for some $w \in \mathbb{Z}$,

$$\begin{aligned} \mathfrak{w}: S^1 &\longrightarrow S^1, \\ \sigma &\longmapsto w\sigma. \end{aligned} \quad (5.5)$$

To restrict to this wrapping sector of $\mathcal{M}^{\mathcal{X}}$, we define a coisotropic ideal $\mathcal{I} = \langle \mathcal{I}_{\mathcal{N}}, \mathcal{I}_{\mathcal{Y}} \rangle$ with

$$\mathcal{I}_{\mathcal{Y}} = \langle y - w\sigma, \xi + wd\sigma \rangle. \quad (5.6)$$

As explained in Sec. II, this is coisotropic and closed under the Q structure $D = [\Theta_{\mathcal{M}^{\mathcal{X}}}, \cdot]$. The ideal $\mathcal{I}_{\mathcal{N}}$ restricts all maps into \mathcal{N} to closed maps. That is, we take

$$\mathcal{I}_{\mathcal{N}} \supset \langle P_{\text{co}}x_0, P_{\text{co}}\psi_0, P_{\text{co}}\chi_0, P_{\text{co}}p_0 \rangle. \quad (5.7)$$

For any coordinate z_k^A with $\deg z_k^A - k \leq 0$, we need to further restrict to exact maps by including the harmonic representative in the ideal (except for x_0). Hence, we have

$$\mathcal{I}_{\mathcal{N}} = \langle P_{\text{co}}x_0, P_{\text{co}}\psi_0, P_{\text{co}}\chi_0, P_{\text{co}}p_0, P_{\mathcal{H}}x_1, P_{\mathcal{H}}\psi_1 \rangle. \quad (5.8)$$

The ideal $\mathcal{I} = \langle \mathcal{I}_{\mathcal{N}}, \mathcal{I}_{\mathcal{Y}} \rangle$ is clearly coisotropic.

We need to check that $\mathcal{I}_{\mathcal{N}}$ is closed under $D = [\Theta_{\mathcal{M}^{\mathcal{X}}}, \cdot]$. The transgressed Hamiltonian is

$$\begin{aligned} \Theta_{\mathcal{M}^{\mathcal{X}}} &= - \int_{\mathcal{X}} \Theta_{\mathcal{M}} - \int_{\mathcal{X}} \iota_d \vartheta_{\mathcal{M}} \\ &= - \int_{\mathcal{X}} -\psi p - \xi q + \frac{1}{4!} F_4(x, y) \psi^4 + \frac{1}{3!} H_3(x, y) \psi^3 \xi \\ &\quad - \int_{\mathcal{X}} p dx + q dy - \frac{1}{3} (\psi d\chi + 2\chi d\psi + \xi d\phi + 2\phi d\xi). \end{aligned} \quad (5.9)$$

As in the previous cases, the only nontrivial constraint comes from the Poisson bracket between the first term and the harmonic generators of $\mathcal{I}_{\mathcal{N}}$. We calculate

$$(\Theta_{\mathcal{M}}, x) = \psi, \quad d(\Theta_{\mathcal{M}}, \psi) = 0, \quad (5.10)$$

and hence we have

$$\left[\Theta_{\mathcal{M}^{\mathcal{X}}}, \int_{\mathcal{X}} \psi \epsilon \right] \propto \int_{\mathcal{X}} (\Theta_{\mathcal{M}}, \psi)_{\mathcal{M}} \epsilon = 0, \quad (5.11)$$

$$\left[\Theta_{\mathcal{M}^{\mathcal{X}}}, \int_{\mathcal{X}} x \epsilon \right] \propto \int_{\mathcal{X}} (\Theta_{\mathcal{M}}, x)_{\mathcal{M}} \epsilon = \int_{\mathcal{X}} \psi \epsilon. \quad (5.12)$$

Evaluating this on \mathcal{C} , we take ψ_1 to be exact, and so the integral vanishes when integrated over a constant $\epsilon = \epsilon_0$. This shows that the Poisson brackets with the harmonic generators x_1, ψ_1 vanish when evaluated on \mathcal{C} ; i.e., they are in \mathcal{I} .

To perform the coisotropic reduction we find the normalizer is generated by

$$N(\mathcal{I}) \sim \{P_{\mathcal{H}}x_0, P_{\mathcal{H}}\psi_0, P_{\mathcal{H}}\chi_1, P_{\mathcal{H}}p_1, \mathcal{I}\}, \quad (5.13)$$

and hence the structure sheaf is generated by

$$\begin{aligned} C^{\infty}(\mathcal{W}) &= N(\mathcal{I})/\mathcal{I} \sim \{x_0, \psi_0, \chi_1, p_1\} \Rightarrow \mathcal{W} \\ &= T^*[2]T[1]N, \end{aligned} \quad (5.14)$$

where the coordinates represent harmonic maps. The symplectic form can be derived from the Poisson brackets on $\mathcal{M}^{\mathcal{X}}$, as in Sec. IV A, and we find

$$\omega_{\mathcal{W}} = -dp_1 dx_0 - d\psi_0 d\chi_1. \quad (5.15)$$

The Hamiltonian function is given by

$$\Theta_{\mathcal{W}} = \Pi(\Theta_{\mathcal{M}^{\mathcal{X}}}) = \Pi \left(- \int_{\mathcal{X}} \Theta_{\mathcal{M}} - \int_{\mathcal{X}} \iota_d \vartheta_{\mathcal{M}} \right). \quad (5.16)$$

The second term vanishes when evaluated on harmonic maps where d annihilates the maps, except for the term $q dy$. We also get a piece $-\xi q$ from the first term. We find

$$\Pi \left(\int_{\mathcal{X}} \xi q - q dy \right) = \int_{\mathcal{X}} -dy q - q dy = 0, \quad (5.17)$$

where we pick up a minus sign from commuting dy (degree 1) through q (degree 3). This verifies the statement made in Sec. II about this cancellation. We then have

$$\begin{aligned} \Theta_{\mathcal{W}} &= \Pi \left(\int_{T[1]S^1} \psi p - \frac{1}{4!} F_4(x, y) \psi^4 - \frac{1}{3!} H_3(x, y) \psi^3 \xi \right) \\ &= \int_{\mathcal{X}} \psi_0 p_1 + -\frac{1}{4!} F_4(x_0, w\sigma) \psi_0^4 + \frac{1}{3!} H_3(x_0, w\sigma) \psi_0^3 w d\sigma \\ &= \frac{1}{2\pi} \int_{\mathcal{X}} \left(\psi_0 p_1 + \frac{w}{3!} H_3(x_0, w\sigma) \psi_0^3 \right) d\sigma \\ &= \psi_0 p_1 + \frac{w}{3!} \tilde{H}_3 \psi_0^3, \end{aligned} \quad (5.18)$$

where \tilde{H}_3 is the average of H_3 over the fiber.

Under the change of coordinates $p_1 \rightarrow -p_1$, we see that we recover the QP manifold associated with the F1 string with w units of \tilde{H}_3 flux, as we would expect from our

intuition of M-theory/IIA duality. Note that in the case that $w = 0$, the physical interpretation seems to break down—we find a string that does not couple to the NS three-form. However, as is noted in [23], this zero winding case corresponds to a scenario in which the original world volume is “collapsed.” This means that the map from the world volume to the target space is not an embedding. From the IIA perspective, the resulting string is tensionless, and thus the M2-brane must somehow be tensionless. We should discard that case on account of such objects appear not to exist on physical grounds; nevertheless, the QP procedure is well defined.

\mathcal{N}					
coord	x^μ	ψ^μ	ζ	χ_μ	p_μ
deg	0	1	3	5	6

\mathcal{Y}				
coord	y	ξ	ϕ	q
deg	0	1	5	6

and use coordinates $(\sigma, d\sigma)$ for \mathcal{X} . We write the symplectic form and Hamiltonian function as

$$\omega_{\mathcal{M}} = dp dx + dq dy - d\psi d\chi - d\xi d\phi - \frac{1}{2} d\zeta d\zeta, \quad (5.21)$$

$$\Theta_{\mathcal{M}} = -\psi p - \xi q + \frac{1}{7!} (H_7 + A \wedge F_6) \psi^7 + \frac{1}{6!} F_6 \psi^6 \xi + \frac{1}{4!} (F_4 - A \wedge H_3) \psi^4 \zeta + \frac{1}{3!} H_3 \psi^3 \xi \zeta. \quad (5.22)$$

We included, in this example, a nontrivial connection on the fiber bundle $N \times S^1$ which we will assume to be S^1 invariant. As previously, we can interpret the coefficients to be elements of $\Omega^i(N) \times \Omega^j(S^1)$, and we require that they are closed under the d_{S^1} on S^1 .

We then transgress the structure to the mapping space $\mathcal{M}^{\mathcal{X}}$ and aim to define a suitable ideal $\mathcal{I} = \langle \mathcal{I}_{\mathcal{N}}, \mathcal{I}_{\mathcal{Y}} \rangle$ with respect to which we perform the coisotropic reduction. The ideal $\mathcal{I}_{\mathcal{Y}}$ is taken as in the previous section

$$\mathcal{I}_{\mathcal{Y}} = \langle y - w\sigma, \xi + wd\sigma \rangle. \quad (5.23)$$

The ideal $\mathcal{I}_{\mathcal{N}}$ is also taken as in the previous section, but now with the additional constraints on the ζ coordinates, restricting them to closed maps. That is, we take

$$\mathcal{I}_{\mathcal{N}} = \langle P_{\text{co}}x_0, P_{\text{co}}\psi_0, P_{\text{co}}\zeta_0, P_{\text{co}}\chi_0, P_{\text{co}}p_0, P_{\mathcal{H}}x_1, P_{\mathcal{H}}\psi_1 \rangle. \quad (5.24)$$

B. M5 on S^1

The next case of interest is wrapping the M5 QP manifold on a circle. The M5 QP manifold was written down in [1], and our expectation is that we should recover that of the D4-brane [3]. We start with the following manifolds:

$$\mathcal{M} = T^*[6]T[1](N \times S^1) \times \mathbb{R}[3], \quad X = S^1. \quad (5.19)$$

Writing $\mathcal{N} = T^*[6]T[1]N \times \mathbb{R}[3]$, $\mathcal{Y} = T^*[6]T[1]S^1$, and $\mathcal{X} = T[1]S^1$, we introduce the homogeneous coordinates

\mathcal{N}					
coord	x^μ	ψ^μ	ζ	χ_μ	p_μ
deg	0	1	3	5	6

\mathcal{Y}				
coord	y	ξ	ϕ	q
deg	0	1	5	6

Since we have only added coexact generators to the ideal, the proof of coisotropy and closure under D goes exactly as in the previous case.

Performing the coisotropic reduction, we find the structure sheaf is generated by

$$C^\infty(\mathcal{W}) = N(\mathcal{I})/\mathcal{I} \sim \{x_0, \psi_0, \zeta_0, \zeta_1, \chi_1, p_1\}, \quad (5.25)$$

which gives

$$\mathcal{W} = T^*[5]T[1]N \times \mathbb{R}[2] \times \mathbb{R}[3]. \quad (5.26)$$

To find the symplectic form, we use the Poisson brackets on $\mathcal{M}^{\mathcal{X}}$ given by (2.6) with appropriate insertions of harmonic test functions ϵ, η and find

$$\omega_{\mathcal{W}} = -dp_1 dx_0 - d\psi_0 d\chi_1 - d\zeta_1 d\zeta_0, \quad (5.27)$$

and the Hamiltonian function is given by¹⁰

$$\begin{aligned} \Theta_{\mathcal{W}} &= \Pi(\Theta_{\mathcal{M}^{\mathcal{X}}}) \\ &= \Pi \left(\int_{T[1]S^1} \psi p - \frac{1}{7!} (H_7 + A \wedge F_6) \psi^7 - \frac{1}{6!} F_6 \psi^6 \xi - \frac{1}{4!} (F_4 - A \wedge H_3) \psi^4 \zeta - \frac{1}{3!} H_3 \psi^3 \xi \zeta \right) \\ &= \psi_0 p_1 + \frac{w}{6!} \tilde{F}_6 \psi_0^6 - \frac{1}{4!} (\tilde{F}_4 - A \wedge \tilde{H}_3) \psi_0^4 \zeta_1 - \frac{w}{3!} \tilde{H}_3 \psi_0^3 \zeta_0, \end{aligned} \quad (5.28)$$

¹⁰We are using the fact that the $\iota_d \theta_{\mathcal{M}}$ term vanishes, apart from the qdy term, which cancels against the ξq term in $\Theta_{\mathcal{M}}$.

where the tilde denotes the average over the S^1 fiber. For $w \neq 0$, we perform a canonical transformation generated by the function $-\frac{1}{2w}A\psi\zeta_1^2$ to obtain the Hamiltonian

$$\Theta_{\mathcal{W}} = \psi_0 p_1 + \frac{1}{4w} F_2 \psi_0^2 \zeta_1^2 - \frac{w}{3!} \tilde{H}_3 \psi_0^3 \zeta_0 - \frac{1}{4!} \tilde{F}_4 \psi_0^4 \zeta_1 + \frac{w}{6!} \tilde{F}_6 \psi_0^6, \quad (5.29)$$

where $F_2 = dA$. Making the change of coordinates $p_1 \rightarrow -p_1$, $\zeta_i \rightarrow -\zeta_i$ puts the QP manifold in the canonical form of that associated with the D4-brane [3].

C. M5 on X_4

The next example will be to wrap the M5-brane over a 4-manifold X_4 . In [30,31] it was shown that one could reproduce the noncritical heterotic string through such a reduction, where the dimension of the gauge group was related to the cohomology of the wrapping manifold. We will start with the manifolds

$$\mathcal{M} = T^*[6]T[1](N \times X_4) \times \mathbb{R}[3], \quad X = X_4. \quad (5.30)$$

Writing $\mathcal{N} = T^*[6]T[1]N \times \mathbb{R}[3]$, $\mathcal{Y} = T^*[6]T[1]X_4$, $\mathcal{X} = T[1]X_4$, we introduce the homogeneous coordinates as in the previous section

		\mathcal{N}					\mathcal{Y}				
coord		x^μ	ψ^μ	ζ	χ_μ	p_μ	coord	y^m	ξ^m	ϕ_m	q_m
deg		0	1	3	5	6	deg	0	1	5	6

where now $\alpha = 1, \dots, 4$, and we use the differential-graded (DG) coordinates $(\sigma^\alpha, d\sigma^\alpha)$ on \mathcal{X} . In these coordinates the symplectic form and Hamiltonian function take the form

$$\omega_{\mathcal{M}} = dp dx + dq dy - d\psi d\chi - d\xi d\phi - \frac{1}{2} d\zeta d\zeta, \quad (5.32)$$

$$\begin{aligned} \Theta_{\mathcal{M}} = & -\psi p - \xi q + \frac{1}{7!} H_7 \psi^7 + \frac{1}{6!} H_6 \psi^6 \xi + \frac{1}{25!} H_5 \psi^5 \xi^2 + \frac{1}{3!4!} H_4 \psi^4 \xi^3 + \frac{1}{4!3!} H_3 \psi^3 \xi^4 \\ & + \frac{1}{4!} F_4 \psi^4 \zeta + \frac{1}{3!} F_3 \psi^3 \xi \zeta + \frac{1}{22} F_2 \psi^2 \xi^2 \zeta + \frac{1}{3!} F_1 \psi \xi^3 \zeta + \frac{1}{4!} F_0 \xi^4 \zeta, \end{aligned} \quad (5.33)$$

where we have taken a trivial connection on the X_4 bundle again. As before, we can view the coefficients as differential forms on Y valued in $\Omega^k(N)$ that we take to be d_Y -closed.

We transgress this structure to $\mathcal{M}^{\mathcal{X}}$ and define a coisotropic ideal $\mathcal{I} = \langle \mathcal{I}_{\mathcal{N}}, \mathcal{I}_{\mathcal{Y}} \rangle$. To define the ideal $\mathcal{I}_{\mathcal{Y}}$ we need to choose some wrapping map $\mathfrak{w}: X_4 \rightarrow X_4$. Restriction to this winding sector of $\mathcal{M}^{\mathcal{X}}$ is given by

$$\mathcal{I}_{\mathcal{Y}} = \langle y - \mathfrak{w}, \xi + d\mathfrak{w} \rangle. \quad (5.34)$$

It is easy to verify that this is coisotropic and closed under D . The ideal $\mathcal{I}_{\mathcal{N}}$ is similar to that for the circle reduction done in the previous section, except now our transgressed coordinates are k -forms¹¹ for $k = 0, \dots, 4$. This means that we need to include more coexact generators and harmonic generators to remove unwanted coordinates. We take

$$\mathcal{I}_{\mathcal{N}} = \langle P_{\text{co}} x_k, P_{\text{co}} \psi_k, P_{\text{co}} \zeta_k, P_{\text{co}} \chi_k, P_{\text{co}} p_k, P_{\mathcal{H}} x_i, P_{\mathcal{H}} \psi_i, P_{\mathcal{H}} \zeta_j | i > 0, j > 2 \rangle. \quad (5.35)$$

We need to check whether this is closed under D . As in previous cases, the only nontrivial checks come from the harmonic generators. The Q structure D acting on the harmonic generators x_i and ψ_i return an element of $\mathcal{I}_{\mathcal{N}}$ precisely as in previous cases so we need only check the closure of $D\zeta_3$ and $D\zeta_4$. Once again, this can be done by calculating

$$(\Theta_{\mathcal{M}}, \zeta)_{\mathcal{M}} = \frac{1}{4!} F_4 \psi^4 + \frac{1}{3!} F_3 \psi^3 \xi + \frac{1}{4} F_2 \psi^2 \xi^2 + \frac{1}{3!} F_1 \psi \xi^3 + \frac{1}{4!} F_0 \xi^4. \quad (5.36)$$

We then transgress this function to $\mathcal{M}^{\mathcal{X}}$ and evaluate it on the vanishing locus \mathcal{C} of \mathcal{I} . We then check whether the following vanishes:

¹¹As noted in Sec. II, the transgressed coordinates z_k^A for $k \geq 2$ should be viewed as differential forms evaluated in some affine bundle. Our construction is still well defined so for simplicity we will ignore this subtlety here.

$$\int_{\mathcal{X}} (\Theta_{\mathcal{M}}, \zeta)_{\mathcal{M}} \epsilon \quad (5.37)$$

for suitable harmonic test functions ϵ . To determine the conditions coming from ζ_4 , we take $\epsilon = \epsilon_0$ a constant function. We then get the constraint

$$\int_X \mathfrak{w}^*(F_0) \epsilon_0 \stackrel{!}{=} 0, \quad (5.38)$$

where we are using the fact that F_0 is a four-form on Y which we pull back to X via the wrapping map. Similarly, the conditions coming from ζ_3 are given by choosing an arbitrary harmonic one-form $\epsilon = \epsilon_1$,

$$\cdot \int_X \mathfrak{w}^*(F_1) \wedge \epsilon_1 \stackrel{!}{=} 0. \quad (5.39)$$

This puts constraints on the coefficients F_0 and F_1 , which can be most easily satisfied if they vanish; i.e., they act as obstructions to the reduction. Note that in some cases, e.g., for $X = \text{K3}$, there are no nontrivial harmonic one-forms, and so (5.39) gives no constraints.

Assuming these constraints are satisfied, the coisotropic reduction with respect to $\mathcal{I} = \langle \mathcal{I}_{\mathcal{N}}, \mathcal{I}_{\mathcal{Y}} \rangle$ gives that the structure sheaf is generated by

$$\begin{aligned} C^\infty(\mathcal{W}) &= N(\mathcal{I})/\mathcal{I} \sim \{x_0, \psi_0, \zeta_2^a, \chi_4, p_4\} \Rightarrow \mathcal{W} \\ &= T^*[2]T[1]N \times H^2(X)[1]. \end{aligned} \quad (5.40)$$

We have introduced an index a parametrising a basis $\{e_a\}$ of $\mathcal{H}^2(X_4)$, and have expanded $\zeta_2 \in \mathcal{H}^2$ as $\zeta_2^a e_a$. Using the Poisson brackets on $\mathcal{M}^{\mathcal{X}}$, we get the symplectic form and the Hamiltonian function on \mathcal{W} to be

$$\omega_{\mathcal{W}} = dp_4 dx_0 - d\psi_0 d\chi_4 - \frac{1}{2} \kappa_{ab} d\zeta_2^a d\zeta_2^b, \quad (5.41)$$

$$\Theta_{\mathcal{W}} = \Pi(\Theta_{\mathcal{M}^{\mathcal{X}}}) = -\psi_0 p_4 + \frac{1}{3!} \tilde{H}_3 \psi^3 + \frac{1}{2} \tilde{F}_a \psi^2 \zeta_2^a, \quad (5.42)$$

where

$$\tilde{H}_3 = \int_X \mathfrak{w}^*(H_3), \quad \tilde{F}_a = \int_X \mathfrak{w}^*(F_2) \wedge e_a, \quad \kappa_{ab} = \int_X e_a \wedge e_b. \quad (5.43)$$

We get the canonical form of the QP manifold associated with a heterotic string with an Abelian gauge group of dimension $b_2(X_4)$. The Killing form on the gauge group is also given by the symmetric form κ_{ab} on $H^2(X_4)$. For example, if $X_4 = T^4$, we get an Abelian gauge group of dimension $b_2(T^4)$ with a Killing form of signature $(3,3)$. If $X_4 = \text{K3}$, then we get a gauge group of dimension $b_2(\text{K3}) = 22$ with a Killing form of signature $(3,19)$.

This matches the results of [30,31]. The fact that we can only obtain Abelian gauge groups arises because we are assuming that we are reducing on smooth manifolds. Degenerations of X_4 to some singular space should lead to gauge enhancement and non-Abelian groups.

VI. AKSZ SIGMA MODELS AND BRANE WRAPPING

In previous sections we obtained an NQP manifold \mathcal{W} from a coisotropic reduction of the mapping space $\mathcal{M}^{\mathcal{X}}$ with respect to a coisotropic submanifold \mathcal{C} that is invariant with respect to the Q structure $D = Q_{\mathcal{M}} + d_{\mathcal{X}}$ on $\mathcal{M}^{\mathcal{X}}$. (In the expression for D we have the lifts of vector fields on the target and source to the mapping space.) We will now point out that these data give rise—essentially trivially—to a reduction of AKSZ sigma models from an AKSZ model with target \mathcal{M} to an AKSZ model with target \mathcal{W} .

We start with the AKSZ sigma model with target \mathcal{M} where the source takes the form $\mathcal{X} \times \mathcal{S}$. The N -manifold \mathcal{S} is taken to be $T[1]\mathcal{S}$ where the (bosonic) manifold \mathcal{S} has dimension $\dim \mathcal{S} = n + 1 - \dim X$ (n being the degree of the target P structure). Then the BV master action is the Hamiltonian corresponding to the Q structure on $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$ given by

$$Q_{\text{BV}} \equiv Q_{\mathcal{M}} + d_{\mathcal{X} \times \mathcal{S}}, \quad (6.1)$$

where again $Q_{\mathcal{M}}$ denotes the lift to $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$ of the target space \mathcal{M} Q structure of the same name and $d_{\mathcal{X} \times \mathcal{S}}$ is the lift of the source $\mathcal{X} \times \mathcal{S}$ de Rham differential again to $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$. Since the source is a product, we can write $d_{\mathcal{X} \times \mathcal{S}} = d_{\mathcal{X}} + d_{\mathcal{S}}$.

The key point that leads to reduction is that we can write

$$\mathcal{M}^{\mathcal{X} \times \mathcal{S}} = (\mathcal{M}^{\mathcal{X}})^{\mathcal{S}}, \quad (6.2)$$

which is known as the product-exponential adjunction. Explicitly, this corresponds to interpreting a function $f \in \mathcal{M}^{\mathcal{X} \times \mathcal{S}}$, which is a function $f(x, s)$ of two arguments, as a function $s \rightarrow f(\bullet, s)$ where $f(\bullet, s)$ is a function of $x \in \mathcal{X}$ for each $s \in \mathcal{S}$.¹² Since $\mathcal{M}^{\mathcal{X}}$ is a QP manifold and \mathcal{S} is an NQ manifold with an integral measure we can consider the BV structure on $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$ as arising from an AKSZ construction with source \mathcal{S} and target $\mathcal{M}^{\mathcal{X}}$. If \mathcal{C} is coisotropic in $\mathcal{M}^{\mathcal{X}}$, then the mapping space $\mathcal{C}^{\mathcal{S}}$ will be a coisotropic submanifold in $(\mathcal{M}^{\mathcal{X}})^{\mathcal{S}} \cong \mathcal{M}^{\mathcal{X} \times \mathcal{S}}$.

The reduced AKSZ sigma model will be given by the coisotropic reduction of $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$ with respect to $\mathcal{C}^{\mathcal{S}}$. We need to confirm that $\mathcal{C}^{\mathcal{S}}$ is invariant with respect to Q_{BV} , so that the BV master action reduces. We rewrite Q_{BV} as

¹²The definition of mapping spaces for graded manifolds is such that this property is true; see, e.g., [32].

$$Q_{\text{BV}} = (Q_{\mathcal{M}} + d_{\mathcal{X}}) + d_{\mathcal{S}} = \hat{D} + \bar{d}_{\mathcal{S}}, \quad (6.3)$$

where in the last formula \hat{D} is the lift from $\mathcal{M}^{\mathcal{X}}$ to $(\mathcal{M}^{\mathcal{X}})^{\mathcal{S}}$ of the vector field D on $\mathcal{M}^{\mathcal{X}}$, while $\bar{d}_{\mathcal{S}}$ is the lift of $d_{\mathcal{S}}$ from \mathcal{S} to $(\mathcal{M}^{\mathcal{X}})^{\mathcal{S}}$. We denote these lifts explicitly now because it is the properties of these lifts that guarantee the reduction: if $V_{\mathcal{S}}$ is any vector field on \mathcal{S} , then the lift $\bar{V}_{\mathcal{S}}$ always leaves $\mathcal{C}^{\mathcal{S}}$ invariant (for any submanifold \mathcal{C} of $\mathcal{M}^{\mathcal{X}}$); $\mathcal{C}^{\mathcal{S}}$ is invariant for \hat{D} if \mathcal{C} is invariant for D . Therefore Q_{BV} gives rise to a homological (and Hamiltonian) vector field on the coisotropic reduction of $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$, which is simply $\mathcal{W}^{\mathcal{S}}$. Using the results of Appendix C we find that the new BV master action is given by evaluating the original action on $\mathcal{C}^{\mathcal{S}}$. In all examples we have investigated the result is another topological field theory of AKSZ type.

In summary, the brane wrapping of QP manifolds that we already discussed always leads to a brane wrapping procedure that takes the BV master action associated with an AKSZ topological field theory and produces the BV master action of another topological field theory.

A. AKSZ 3-brane to membrane example

To illustrate, we will treat the reduction of the AKSZ sigma model corresponding to the wrapping of an M2 algebroid (see Sec. V A) on a circle that we discussed in Sec. V A. This is a reduction of the four-dimensional (4D) topological field theory of Ikeda and Uchino [33] to a (3D) Courant sigma model.

This example thus has $\mathcal{X} = T[1]S^1$, and the coisotropic submanifold $\mathcal{C} \subset \mathcal{M}^{\mathcal{X}}$ is given by

$$\begin{aligned} d_{\mathcal{X}} p_0 &= 0, & d_{\mathcal{X}} x_0^{\mu} &= 0, & P_{\mathcal{H}} x_1^{\mu} &= P_{\text{co}} x_1^{\mu} = 0, & y_0 &= w\sigma, & y_1 &= 0, \\ d_{\mathcal{X}} \chi_0 &= 0, & d_{\mathcal{X}} \psi_0^{\mu} &= 0, & P_{\mathcal{H}} \psi_1^{\mu} &= P_{\text{co}} \psi_1^{\mu} = 0, & \xi_0 &= 0, & \xi_1 &= -w. \end{aligned} \quad (6.4)$$

We have used the superfield expansion of $\mathbf{Z}^{\bar{A}} = \{x^{\mu}, y, \psi^{\mu}, \xi, \dots\}$ in form degree [so $x(\sigma, d\sigma) = x_0(\sigma) + x_1(\sigma)d\sigma$, etc.]

For the original (4D) AKSZ theory degree counting to work we set $\mathcal{S} = T[1]S$ where S can be any 3-manifold, so that $X \times S = S^1 \times S$ is the four-dimensional world volume. Using the product-exponential adjunction to write $\mathcal{M}^{\mathcal{X} \times \mathcal{S}} \cong (\mathcal{M}^{\mathcal{X}})^{\mathcal{S}}$ amounts to promoting the components $Z_k^{\bar{A}}$ of the superfields $\mathbf{Z}^{\bar{A}}$ defining a map $\mathcal{M}^{\mathcal{X}}$ to superfields $Z_k^{\bar{A}}$ that now depend on the \mathcal{S} coordinates $\{s, ds\}$ as well as the \mathcal{X} coordinates $(\{\sigma, d\sigma\}$ in this case). Then the coisotropic submanifold $\mathcal{C}^{\mathcal{S}}$ is the locus of functions $\mathcal{S} \rightarrow \mathcal{M}^{\mathcal{X}}$ such that

$$\begin{aligned} d_{\mathcal{X}} p_0 &= 0, & d_{\mathcal{X}} x_0^{\mu} &= 0, & P_{\mathcal{H}} x_1^{\mu} &= P_{\text{co}} x_1^{\mu} = 0, & y_0 &= w\sigma, & y_1 &= 0, \\ d_{\mathcal{X}} \chi_0 &= 0, & d_{\mathcal{X}} \psi_0^{\mu} &= 0, & P_{\mathcal{H}} \psi_1^{\mu} &= P_{\text{co}} \psi_1^{\mu} = 0, & \xi_0 &= 0, & \xi_1 &= -w, \end{aligned} \quad (6.5)$$

where all bolded expressions depend on $\{\sigma, s, ds\}$. (The projectors to coexact/harmonic pieces refer to the Hodge decomposition with respect to \mathcal{X} as above.)

We can explicitly check the claim that $\mathcal{C}^{\mathcal{S}}$ is invariant with respect to $Q_{\text{BV}} = \hat{D} + \bar{d}_{\mathcal{S}}$. For example,

$$\hat{D} \int_{\mathcal{S} \times \mathcal{X}} (\mathbf{y} - w\sigma) \epsilon = \int_{\mathcal{S} \times \mathcal{X}} (\boldsymbol{\xi} + d\sigma \partial_{\sigma} \mathbf{y}) \epsilon \stackrel{\text{mod } \mathcal{I}(\mathcal{C}^{\mathcal{S}})}{=} \int_{\mathcal{S} \times \mathcal{X}} (-w d\sigma + d\sigma w) \epsilon = 0. \quad (6.6)$$

[We smeared against $\epsilon \in C^{\infty}(\mathcal{S} \times \mathcal{X})$ and employed (2.19)]. The other differential $\bar{d}_{\mathcal{S}}$ leaves the ideal invariant independently. This way we may confirm explicitly that S_{BV} lies in $N(\mathcal{I}(\mathcal{C}^{\mathcal{S}}))$.

It remains to calculate the reduced BV master action, which amounts to calculating $\Pi(S_{\text{BV}})$ where Π implements the quotient modulo $\mathcal{I}(\mathcal{C}^{\mathcal{S}})$. S_{BV} is the Hamiltonian for $Q_{\text{BV}} = D + d_{\mathcal{S}} = Q_{\mathcal{M}} + d_{\mathcal{X}} + d_{\mathcal{S}} \equiv Q_{\mathcal{M}} + d$ which is explicitly given by formula (2.10), which is a linear combination of $\int_{\mathcal{X} \times \mathcal{S}} \Theta_{\mathcal{M}}$ and $\int_{\mathcal{X} \times \mathcal{S}} \iota_d \Theta_{\mathcal{M}}$, for $\Theta_{\mathcal{M}}$ the transgression of a symplectic potential on \mathcal{M} that satisfies $d_{\mathcal{M}} \Theta_{\mathcal{M}} = \omega_{\mathcal{M}}$, $\omega_{\mathcal{M}}$ being given in (5.3). The bolded quantities are superfields corresponding to $\mathcal{M}^{\mathcal{X} \times \mathcal{S}}$ now. We then calculate

$$\begin{aligned} \Pi \int_{\mathcal{X} \times \mathcal{S}} \iota_d \Theta_{\mathcal{M}} &= \Pi \int_{\mathcal{X} \times \mathcal{S}} \mathbf{p} d\mathbf{x} + \mathbf{q} d\mathbf{y} - \boldsymbol{\chi} d\boldsymbol{\psi} - \boldsymbol{\phi} d\boldsymbol{\xi} \\ &= \int_{\mathcal{S}} \left(\int_{\mathcal{X}} \mathbf{p}_1 d\sigma \right) d_{\mathcal{S}} \mathbf{x}_0 + w \left(\int_{\mathcal{X}} \mathbf{q}_0 d\sigma \right) - \left(\int_{\mathcal{X}} \boldsymbol{\chi}_1 d\sigma \right) d_{\mathcal{S}} \boldsymbol{\psi}_0. \end{aligned} \quad (6.7)$$

Note that terms involving \mathbf{x}_1 and $\boldsymbol{\psi}_1$ will generate $d_{\mathcal{X}}$ -exact terms which will vanish under the $\int_{\mathcal{X}}$ integral. Using (5.4),

$$\Pi \int_{\mathcal{X} \times \mathcal{S}} \Theta_{\mathcal{M}} = \int_{\mathcal{S}} -\psi_0 \left(\int_{\mathcal{X}} \mathbf{p}_1 d\sigma \right) - w \left(\int_{\mathcal{X}} d\sigma \mathbf{q}_0 \right) + 0 - w \left(\int_{\mathcal{X}} \frac{1}{3!} H_3(\psi_0)^3 d\sigma \right). \quad (6.8)$$

We then read off the sign factors from (2.10) to find

$$\begin{aligned} \Pi S_{\text{BV}} &= \Pi \left(- \int_{\mathcal{X} \times \mathcal{S}} \Theta_{\mathcal{M}} + \int_{\mathcal{X} \times \mathcal{S}} \iota_d \vartheta_{\mathcal{M}} \right) \\ &= \int_{\mathcal{S}} \psi_0 \left(\int_{\mathcal{X}} \mathbf{p}_1 d\sigma \right) + w \left(\int_{\mathcal{X}} \frac{1}{3!} H_3(\psi_0)^3 d\sigma \right) + \left(\int_{\mathcal{X}} \mathbf{p}_1 d\sigma \right) d_{\mathcal{S}} \mathbf{x}_0 - \left(\int_{\mathcal{X}} \mathbf{x}_1 d\sigma \right) d_{\mathcal{S}} \psi_0. \end{aligned} \quad (6.9)$$

The signs were such that the terms $w \int_{\mathcal{X}} \mathbf{q}_0 d\sigma$ canceled.

In the above expression we can identify the integrated expressions $(\int_{\mathcal{X}} \mathbf{p}_1 d\sigma)$ and $(\int_{\mathcal{X}} \mathbf{x}_1 d\sigma)$ as the conjugate momenta superfields (with degrees 2 and 1 respectively) that appear in the Courant sigma model for an exact Courant algebroid structure defined by the three-form wH_3 . The result we calculated via coisotropic reduction of the original (four-dimensional) AKSZ topological sigma model is identical to the AKSZ sigma model constructed directly from the wrapped QP manifold \mathcal{W} with source manifold \mathcal{S} [see (5.14)].

Therefore we have recovered the correct relation between the M-theory fluxes, the M2-brane winding w , and the type IIA NS-flux wH_3 seen by the fundamental strings that arise as the M-theory circle $X = S^1$ is shrunk to zero, all at the level of the corresponding topological sigma models.

VII. CONCLUSIONS

We defined a reduction procedure of NQP manifolds $\mathcal{M} \rightarrow \mathcal{W}$ which encompasses the properties of wrapped branes. This is consistent with the AKSZ procedure in the sense that the reduction naturally lifts to a reduction of the AKSZ theory with target \mathcal{M} to the AKSZ theory with target \mathcal{W} . We applied this to many examples, including many physically motivated examples of wrapped branes, and we saw that it reproduced the known M-theory/IIA dualities. We also were able to find a novel relation between the Courant algebroid and the Poisson algebroid through this reduction.

As mentioned in the Introduction, there is a correspondence between branes, QP manifolds and higher Leibniz algebroids $E \rightarrow M$, which can be used to describe the geometry of string backgrounds. This geometry is described through the generalized metric [34–37]. While our procedure describes how the algebroids reduce under brane wrapping (captured by the graded-commutative manifold \mathcal{M}), determining how the full string background geometry reduces would likely require more input than the procedure outlined in this paper. In particular, one would likely have to select some precise representative of the cohomology classes we have constructed, breaking the topological nature. Nonetheless, in certain cases one may be able to obtain nontrivial physical data about the reduction from our construction as follows. In many cases, including many

supersymmetric backgrounds, the generalized metric is described, in part, via a choice of subalgebroid $L \rightarrow M$ which has a Lie algebroid structure [2,38–43]. This Lie algebroid structure has an associated differential d_L , which in the QP language is captured by the Hamiltonian vector field Q . We see from our construction that we define a reduction not only of the Leibniz algebroid E but also of the differential $Q = (\Theta, \cdot)$ and hence of the associated Lie algebroid L . Despite only describing the background in part, the L structure contains nontrivial physical data. For example, for reductions to Minkowski space, the L -bundle can determine certain massless moduli in the effective theory. For Anti-de Sitter reductions, the bundle L captures the holomorphic data of the associated Superconformal field theory [8]. An application of our reduction reasoning would be to find the bundle L (called the exceptional complex structure in [41]) associated with a IIA background with nontrivial Ramond-Ramond flux associated with branes arising from wrapped M2- and M5-branes in M-theory.

Beyond the application to the geometry of supersymmetric backgrounds, we expect that our work will have many interesting applications to other topological AKSZ theories. One can ask how general our procedure is, or whether it is possible to relax some of the assumptions made in Sec. II. For example, can we relax the trivial bundle condition $M = N \times Y$, perhaps by introducing some flat connection similar to [15]? We can also ask whether we can extend our construction to manifolds X with boundary. We can also relax the constraint on $\mathcal{X} = T[1]X$, and instead just take \mathcal{X} to be some DG manifold with some invariant measure of degree $n + 1$. For example, we can try to extend the reduction procedure to $\mathcal{X} = T^{1,0}[1]X$ for some complex manifold X with $\dim_{\mathbb{C}} X = n + 1$. We could then apply the reduction to, say, the work of [44].

In Sec. IV B, we found an interesting relation between the Courant algebroid and the Poisson algebroid QP structures. This was based on the embedding of the Poisson differential d_{π} into $T \oplus T^*$. There are other interesting differentials that can appear in these Courant algebroids [45] that are associated with topological theories on G_2 and $\text{Spin}(7)$ manifolds. One can try to embed these differentials in the language of QP structures and perform the reduction to get new topological models associated with these special holonomy manifolds.

ACKNOWLEDGMENTS

We thank Marco Zambon, Chris Blair, Dan Thompson, Ondrej Hulik, and Maxime Grigoriev for very helpful discussions during this project. The authors are grateful to the Mainz Institute for Theoretical Physics (MITP) of the DFG Cluster of Excellence PRISMA* (Project ID 39083149), for its hospitality and its partial support during the completion of this work. A. S. A. is supported by the FWO-Vlaanderen through the Project No. G006119N, as well as by the Vrije Universiteit Brussel through the Strategic Research Program “High-Energy Physics.” He is also supported by an FWO Senior Postdoctoral Fellowship (No. 1265122N). D. T., is supported by the NSF Grant No. PHYS-2112859. Part of the research for this project was performed while D. T. was supported by the EPSRC New Horizons Grant “New geometry from string dualities” EP/V049089/1.

APPENDIX A: NOTATION

1. Commutative manifolds

M	Starting/parent commutative manifold which is always a product manifold of a base and a fiber to be wrapped.
N	The commutative manifold which is the base of the trivial fiber bundle M .
Y	The fiber of the trivial bundle M . This is the manifold over which we wrap the branes.
X	The fiber of the brane that is wrapped over Y .

2. Graded-commutative manifolds

\mathcal{M}	Starting/parent QP manifold.
\mathcal{N}	A submanifold of \mathcal{M} which is the natural QP manifold restricted to the base of the fibration.
\mathcal{Y}	A submanifold of \mathcal{M} which is the natural QP manifold restricted to the fiber; usually $\mathcal{Y} = T^*[n]T[1]Y$.
\mathcal{X}	The shifted tangent bundle $T[1]X$; the source of the mapping space $\mathcal{M}^{\mathcal{X}}$.
\mathcal{W}	Final wrapped QP manifold.
$\mathcal{M}^{\mathcal{X}}$	maps($\mathcal{X} \rightarrow \mathcal{M}$).
\mathcal{S}	A DG manifold with invariant measure of degree $n + 1 - \dim X$.

3. Indices

A, B, C, \dots	Indices along \mathcal{M}, \mathcal{N} .
μ, ν, ρ, \dots	Indices along N .
m, n, p, \dots	Indices along Y .
$\alpha, \beta, \gamma, \dots$	Indices along X .
r, s, t, \dots	Indices corresponding to degree shifted real lines $\mathbb{R}[n_r]$.
a, b, c, \dots	Indices for a basis of differential forms on X .

4. Coordinates

Z^A	Homogeneous coordinates on \mathcal{M} .
z^A	Homogeneous coordinates on \mathcal{N} .
x^μ	Degree 0 coordinates on \mathcal{N} .
ψ^μ	Degree 1 coordinates on \mathcal{N} parametrizing the fiber of $T[1]N$.
p_μ	Coordinate dual to x^μ .
χ_μ	Coordinate dual to ψ^μ .
y^m	Degree 0 coordinates on \mathcal{Y} parametrizing the fiber of $T[1]Y$.
ξ^m	Degree 1 coordinates on \mathcal{Y} .
q_m	Coordinate dual to y^m .
ϕ_m	Coordinate dual to ξ^m .
$(\sigma^\alpha, d\sigma^\alpha)$	Coordinates for the DG manifold (\mathcal{X}, d) such that $d(\sigma^\alpha) = d\sigma^\alpha$.
ζ^r	Homogeneous coordinates corresponding to degree shifted real lines $\mathbb{R}[n_r]$.
Z^A	Transgressed coordinates of $\mathcal{M}^{\mathcal{X}}$.
Z_k^A	An expansion of the transgressed coordinates Z^A into differential k -forms.
$Z_k^{A,a}$	A coordinate labeling the harmonic k -forms, labeled by a , associated with the transgressed coordinate Z^A .

5. Functions and differential forms

Ω^k	The space of differential k -forms.
\mathcal{H}^k	Harmonic k -forms.
$e_{k,a}$	A basis of harmonic k -form(s) (occasionally the k is dropped).
$\Theta_{\mathcal{M}}$	The Hamiltonian function of \mathcal{M} (similarly for $\mathcal{N}, \mathcal{W}, \dots$).
$\omega_{\mathcal{M}}$	The symplectic form of \mathcal{M} (similarly for $\mathcal{N}, \mathcal{W}, \dots$).
$\vartheta_{\mathcal{M}}$	The canonical symplectic potential of \mathcal{M} (similarly for $\mathcal{N}, \mathcal{W}, \dots$).
$(\cdot, \cdot)_{\mathcal{M}}$	The Poisson bracket for \mathcal{M} (similarly for $\mathcal{N}, \mathcal{W}, \dots$).
$[\cdot, \cdot]$	The Poisson bracket on $\mathcal{M}^{\mathcal{X}}$.

6. Miscellaneous

w	Wrapping map $X \rightarrow Y$.
w	Winding number/matrix of a circle/torus over itself.
\mathcal{I}	The coisotropic ideal within $\mathcal{M}^{\mathcal{X}}$.
C	The vanishing locus of \mathcal{I} within $\mathcal{M}^{\mathcal{X}}$.

APPENDIX B: QP MANIFOLDS

1. Graded manifolds

A graded manifold \mathcal{M} is a supermanifold whose coordinates come equipped with a \mathbb{Z} grading.¹³ One can always find homogeneous coordinates Z^A of definite

¹³From [46], the consistency of the \mathbb{Z} grading of coordinates comes from the existence of a global degree counting vector field ε and transition functions which preserve degree.

degree, where $\deg Z^A \bmod 2$ is the Grassman parity of the coordinate. We will denote by A the degree of Z^A , and so we have

$$Z^A Z^B = (-1)^{AB} Z^B Z^A. \quad (\text{B1})$$

The sheaf of functions on \mathcal{M} splits into subsheaves $C_n^\infty(\mathcal{M})$ of functions of definite degree. The degree of a homogeneous function f is measured by the degree counting vector field ε (the “Euler vector field”) via

$$\varepsilon(f) = \deg(f)f. \quad (\text{B2})$$

In local homogeneous coordinates Z^A , we have

$$\varepsilon = \sum_A \deg(Z^A) Z^A \frac{\partial}{\partial Z^A}. \quad (\text{B3})$$

Unless otherwise stated, all derivations are left derivations. Hence, the de Rham d is

$$df = dZ^A \partial_A f, \quad (\text{B4})$$

and any homogeneous (in degree) vector field X acts as

$$X(fg) = X(f)g + (-1)^{Xf} f X(g), \quad (\text{B5})$$

where we have used the shorthand X, f for the degree of the respective components. In local coordinates we can write $X = X(Z)^A \partial_A$, and so $\deg X = \deg X^A - \deg Z^A$. We also define

$$\deg(df) = \deg f + 1. \quad (\text{B6})$$

For this to be consistent with $\iota_A dZ^B = \delta^B_A$, where ι_A denotes contraction with the vector field ∂_A , we require that the interior product has degree

$$\deg \iota = -1. \quad (\text{B7})$$

2. Poisson and symplectic structures

A graded Poisson structure of degree $-n$ is defined to satisfy

$$(f, g) = (-1)^{1+(f+n)(g+n)} (g, f) \quad (\text{B8})$$

and the graded Jacobi identity

$$(f, (g, h)) = ((f, g), h) + (-1)^{(f+n)(g+n)} (g, (f, h)) \quad (\text{B9})$$

for all homogeneous functions f, g, h . It also acts as a left derivation on the right-hand arguments, but a right derivation on the left-hand arguments. That is,

$$\begin{aligned} (f, gh) &= (f, g)h + (-1)^{(f+n)g} g(f, h), \\ (fg, h) &= f(g, h) + (-1)^{(h+n)g} (f, h)g. \end{aligned} \quad (\text{B10})$$

If the Poisson structure is induced from a symplectic structure ω , we have that

$$\iota_{X_f} = (-1)^f df, \quad X_f := (f, \cdot). \quad (\text{B11})$$

In local homogeneous coordinates we can write

$$\omega = \frac{1}{2} dZ^A \omega_{AB} dZ^B, \quad (\text{B12})$$

which implies the symmetry

$$\omega_{AB} = (-1)^{1+AB+n(A+B)} \omega_{BA}. \quad (\text{B13})$$

If we define ω^{AB} via $\omega^{AB} \omega_{BC} = \delta^A_C$, then (B11) implies

$$(f, g) = (-1)^f \partial_A^R f \omega^{AB} \partial_B g, \quad (\text{B14})$$

where ∂_A^R is defined by $df = dZ^A \partial_A f = \partial_A^R dZ^A$. Note that it is not a right derivation by itself, but the combination $(-1)^f \partial_A^R f$ is a right derivation. This is consistent with (B10). Note that this implies

$$(Z^A, Z^B) = (-1)^A \omega^{AB}. \quad (\text{B15})$$

The symplectic potential is defined such that $d\vartheta = \omega$, and can be defined canonically through the Euler vector field ε . We have that¹⁴

$$n\omega = \mathcal{L}_\varepsilon \omega = \iota_\varepsilon d\omega + d(\iota_\varepsilon \omega) = d(\iota_\varepsilon \omega), \quad (\text{B16})$$

where we have used $d\omega = 0$. This implies we can take

$$\vartheta = \frac{1}{n} \iota_\varepsilon \omega = (\deg Z^A) Z^A \omega_{AB} dZ^B. \quad (\text{B17})$$

3. Transgressed QP structure on $\mathcal{M}^\mathcal{X}$

Let $(\mathcal{X} = T[1]X, d)$ be a DG manifold with homogeneous coordinates $\sigma, d\sigma$. A point $f \in \mathcal{M}^\mathcal{X}$ can be defined by how it pulls back the coordinates on \mathcal{M} . We have

$$\begin{aligned} f^* Z^A &= Z^A(\sigma, d\sigma) = Z_0^A(\sigma) + Z_{1\alpha}^A(\sigma) d\sigma^\alpha + \dots \\ &\quad + \frac{1}{d!} Z_{d\alpha_1 \dots \alpha_d}^A(\sigma) d\sigma^{\alpha_1} \dots d\sigma^{\alpha_d}. \end{aligned} \quad (\text{B18})$$

We use the shorthand $Z_k^A = \frac{1}{k!} Z_{k\alpha_1 \dots \alpha_k}^A(\sigma) d\sigma^{\alpha_1} \dots d\sigma^{\alpha_k}$, where $Z_{k\alpha_1 \dots \alpha_k}^A(\sigma)$ is a function of degree $\deg Z^A - k$. These act as coordinates on $\mathcal{M}^\mathcal{X}$. Our conventions are always that the form components come to the right of the function. So, e.g.,

¹⁴More generally, the Lie derivative on any graded differential form along a vector field X is given by $\mathcal{L}_X = \iota_X d + (-1)^X d \iota_X$. The Euler vector field is degree 0, hence the expression given.

$$Z_1^A = Z_{1a}^A(\sigma) d\sigma^\alpha = (-1)^{A-1} d\sigma^\alpha Z_{1a}^A(\sigma). \quad (\text{B19})$$

We can always define an evaluation map

$$\begin{aligned} \text{ev}: \mathcal{M}^{\mathcal{X}} \times \mathcal{X} &\longrightarrow \mathcal{M}, \\ (f, \sigma, d\sigma) &\longmapsto f(\sigma, d\sigma). \end{aligned} \quad (\text{B20})$$

We also have the chain map defined by

$$\begin{aligned} \mu_*: \Omega^*(\mathcal{M}^{\mathcal{X}} \times \mathcal{X}) &\longrightarrow \Omega^*(\mathcal{M}^{\mathcal{X}}), \\ \alpha &\longmapsto \int_{\mathcal{X}} \alpha. \end{aligned} \quad (\text{B21})$$

The combination $\mu_* \text{ev}^*: \Omega^*(\mathcal{M}) \rightarrow \Omega^*(\mathcal{M}^{\mathcal{X}})$ is called the transgression map. The QP structure on the mapping space is defined by

$$\begin{aligned} \omega_{\mathcal{M}^{\mathcal{X}}} &= \mu_* \text{ev}^* \omega_{\mathcal{M}}, \\ \Theta_{\mathcal{M}^{\mathcal{X}}} &= (-1)^d \mu_* \text{ev}^* \Theta_{\mathcal{M}} + (-1)^{n+d+1} \iota_{d\mu_* \text{ev}^* \vartheta}, \end{aligned} \quad (\text{B22})$$

where we use the same symbol d for the lift of the vector field on \mathcal{X} to $\mathcal{M}^{\mathcal{X}}$.

This can be given more explicitly in the coordinates (B18). We will use the boldface notation to denote a function, differential form, or coordinate on \mathcal{M} that is pulled back to \mathcal{X} via some function $f \in \mathcal{M}^{\mathcal{X}}$. That is, we effectively take $f = \text{ev}^* f$. We can then write

$$\omega_{\mathcal{M}^{\mathcal{X}}} = \frac{1}{2} \int_{\mathcal{X}} \delta \mathbf{Z}^A (\omega_{\mathcal{M}})_{AB} \delta \mathbf{Z}^B. \quad (\text{B23})$$

Our convention for integrals is that constants are pulled out from the left. The symplectic form above gives rise to a Poisson bracket that takes the following form on homogeneous functionals F, G

$$[F, G] = \int_{\mathcal{X}} (-1)^F \frac{\delta^R F}{\delta \mathbf{Z}^A} (\omega_{\mathcal{M}})^{AB} \frac{\delta G}{\delta \mathbf{Z}^B}, \quad (\text{B24})$$

where

$$\delta F = \int_{\mathcal{X}} \delta \mathbf{Z}^A \frac{\delta F}{\delta \mathbf{Z}^A} = \int_{\mathcal{X}} \frac{\delta^R F}{\delta \mathbf{Z}^A} \delta \mathbf{Z}^A. \quad (\text{B25})$$

We can define a functional F via some pulled-back function f by

$$F(\epsilon) = \int_{\mathcal{X}} f \epsilon, \quad \forall \epsilon \in C^\infty(\mathcal{X}). \quad (\text{B26})$$

Then the Poisson bracket (B24) can be expressed nicely as

$$\left[\int_{\mathcal{X}} f \epsilon, \int_{\mathcal{X}} g \eta \right] = (-1)^{(f+n)\epsilon+d} \int_{\mathcal{X}} (f, g)_{\mathcal{M}} \epsilon \eta, \quad (\text{B27})$$

where $(f, g)_{\mathcal{M}} = \text{ev}^*(f, g)_{\mathcal{M}}$.

We can use this to calculate the Poisson bracket on two harmonic generators Z_k^A and $Z_{k'}^B$. Let $e_{k,a}$ be a basis of harmonic k -forms and \tilde{e}_{d-k}^b be a dual basis of harmonic $d-k$ -forms. So

$$\delta^b_a = \int_X e_{k,a} \wedge \tilde{e}_{d-k}^b. \quad (\text{B28})$$

Noting that $\Omega^*(\mathcal{X}) \simeq C^\infty(T[1]X) = C^\infty(\mathcal{X})$, and by expanding $Z_k^A = Z_k^{A,a} e_{k,a}$ with $Z_k^{A,a}$ some constant coefficient, we have

$$Z_k^{A,a} = \int_{\mathcal{X}} Z_k^A \tilde{e}_{d-k}^a = \int_{\mathcal{X}} \mathbf{Z}^A \tilde{e}_{d-k}^a. \quad (\text{B29})$$

We then see that we get an induced Poisson bracket on the coefficients given by

$$\begin{aligned} [Z_k^{A,a}, Z_{k'}^{B,b}] &\equiv \left[\int_{\mathcal{X}} \mathbf{Z}^A \tilde{e}_{d-k}^a, \int_{\mathcal{X}} \mathbf{Z}^B \tilde{e}_{d-k'}^b \right] \\ &= (-1)^{(A+n)(d-k)+d} \int_{\mathcal{X}} (\mathbf{Z}^A, \mathbf{Z}^B)_{\mathcal{M}} \tilde{e}_{d-k}^a \wedge \tilde{e}_{d-k'}^b \\ &= (-1)^{(A+n)(d-k)+d} (-1)^A \omega^{AB} \int_{\mathcal{X}} \tilde{e}_{d-k}^a \wedge \tilde{e}_{d-k'}^b \\ &= (-1)^{(A+n)(d-k)+d} (-1)^A \omega^{AB} \kappa^{ab} \delta_{k+k',d}, \end{aligned} \quad (\text{B30})$$

where we have assumed Darboux coordinates, so the ω^{AB} are constant, and where

$$\kappa^{ab} = \int_X \tilde{e}_{d-k}^a \wedge \tilde{e}_k^b. \quad (\text{B31})$$

We use this to find the symplectic form of the reduced theory.

APPENDIX C: COISOSTROPIC REDUCTION OF GRADED POISSON ALGEBRAS

Let \mathcal{P} be a graded algebra with a graded Poisson bracket $[\bullet, \bullet]$ of degree $-P$ along with a left derivation \mathcal{V} of \mathcal{P} , possibly Hamiltonian (i.e., given by Poisson brackets, so $\mathcal{V} = [\mathcal{H}_{\mathcal{V}}, \bullet]$ for $\mathcal{H}_{\mathcal{V}} \in \mathcal{P}$). We will explain how all of these objects behave under *coisotropic reduction*. The derivation is as in the ungraded case considered originally by Śniatycki and Weinstein [47].

If \mathcal{I} is a (multiplicative, degree-homogeneous) ideal of \mathcal{P} , it is a *coisotropic* if it is a Poisson subalgebra, i.e., $[\mathcal{I}, \mathcal{I}] \subseteq \mathcal{I}$. Then the *coisotropic reduction* of \mathcal{P} with respect to \mathcal{I} is the quotient

$$\overline{\mathcal{P}} \equiv N(\mathcal{I})/\mathcal{I}, \quad (\text{C1})$$

where $N(\mathcal{I}) \equiv \{f \in \mathcal{P} | [f, \mathcal{I}] \subseteq \mathcal{I}\}$ is the Poisson normalizer of \mathcal{I} . Then the bracket on $\overline{\mathcal{P}}$ is defined in terms of the bracket $[\bullet, \bullet]$ via

$$[\Pi f, \Pi g]_{\bar{\mathcal{P}}} \equiv \Pi[f, g], \quad (C2)$$

where Πf is the equivalence class $f + \mathcal{I}$. For any \mathcal{P} derivation \mathcal{V} we define its reduction $\bar{\mathcal{V}}$ via

$$\bar{\mathcal{V}}(\Pi f) = \Pi \mathcal{V}(f), \quad f \in \mathcal{P}. \quad (C3)$$

Theorem 1. Given any coisotropic \mathcal{I} , the bracket $[\bullet, \bullet]_{\bar{\mathcal{P}}}$ is well defined. It is moreover a Poisson bracket of degree $-P$, and so $\bar{\mathcal{P}}$ is a graded Poisson algebra.

If the derivation \mathcal{V} on \mathcal{P} preserves the Poisson structure ($\mathcal{V}[f, g] = [\mathcal{V}f, g] \pm [f, \mathcal{V}g]$) and the coisotropic $[\mathcal{V}(\mathcal{I}) \subseteq \mathcal{I}]$, then the reduced derivation $\bar{\mathcal{V}}$ is well defined.

Finally, if \mathcal{V} is furthermore Hamiltonian with Hamiltonian $\mathcal{H}_{\mathcal{V}} \in \mathcal{P}$ (so $\mathcal{V} = [\mathcal{H}_{\mathcal{V}}, \bullet]$), then $\bar{\mathcal{V}}$ is Hamiltonian with Hamiltonian $\Pi(\mathcal{H}_{\mathcal{V}})$. [In this latter case \mathcal{V} automatically preserves the Poisson structure, but the condition $\mathcal{V}(\mathcal{I}) \subseteq \mathcal{I}$ implies $[\mathcal{H}_{\mathcal{V}}, \mathcal{I}] \subseteq \mathcal{I}$.]

If all derivations we are interested in are, in fact, Hamiltonian (which is the case in the main text), then we just need to check that the ideal \mathcal{I} is a coisotropic and that $[\mathcal{H}_{\mathcal{V}}, \mathcal{I}] \subseteq \mathcal{I}$.

Proof.—The bracket $[\bullet, \bullet]_{\bar{\mathcal{P}}}$ is well defined because

$$\begin{aligned} [\Pi f, \Pi g]_{\bar{\mathcal{P}}} &= \Pi[f + \mathcal{I}, g + \mathcal{I}] = \Pi([f, g] + [f, \mathcal{I}] \\ &\quad + [\mathcal{I}, g] + [\mathcal{I}, \mathcal{I}]) = \Pi[f, g], \end{aligned} \quad (C4)$$

where the last three terms in the second equality vanish because $f, g \in N(\mathcal{I})$ and $[\mathcal{I}, \mathcal{I}] \subseteq \mathcal{I}$. This new bracket inherits the antisymmetry and Jacobi identity properties from $[\bullet, \bullet]$. Since furthermore \mathcal{I} is homogeneous in degree, Πf will have a well-defined degree, and so the new bracket defines a graded Poisson algebra structure.

Similarly since $\mathcal{V}(f + \mathcal{I}) = \mathcal{V}(f) + \mathcal{V}(\mathcal{I})$ we have that $\bar{\mathcal{V}}$ is well defined on \mathcal{P}/\mathcal{I} when $\mathcal{V}(\mathcal{I}) \subseteq \mathcal{I}$. We then need to show that it preserves the subspace $N(\mathcal{I})/\mathcal{I} = \bar{\mathcal{P}}$. Since \mathcal{V} preserves the Poisson bracket, we have

$$[\mathcal{V}f, \mathcal{I}] = \mathcal{V}[f, \mathcal{I}] \pm [f, \mathcal{V}\mathcal{I}]. \quad (C5)$$

If $f \in N(\mathcal{I})$, this becomes $\mathcal{V}(\mathcal{I}) \pm [f, \mathcal{V}\mathcal{I}]$ which lies in the coisotropic when $\mathcal{V}(\mathcal{I}) \subseteq \mathcal{I}$. Therefore $\mathcal{V}(f)$ lies in $N(\mathcal{I})$.

Finally, if $\mathcal{V} = [\mathcal{H}_{\mathcal{V}}, \bullet]$, then $\Pi \mathcal{V}(\Pi f) = \Pi \mathcal{V}(f) = \Pi[\mathcal{H}_{\mathcal{V}}, f] = [\Pi \mathcal{H}_{\mathcal{V}}, \Pi f]_{\Pi \mathcal{P}}$, which completes the proof. \square

[1] A. S. Arvanitakis, Brane Wess-Zumino terms from AKSZ and exceptional generalised geometry as an L_∞ -algebroid, *Adv. Theor. Math. Phys.* **23**, 1159 (2019).

[2] A. S. Arvanitakis, Brane current algebras and generalised geometry from QP manifolds. Or, “when they go high, we go low”, *J. High Energy Phys.* **11** (2021) 114.

[3] A. S. Arvanitakis, E. Malek, and D. Tennyson, Romans massive QP manifolds, *Universe* **8**, 147 (2022).

[4] A. S. Arvanitakis, Generalising courant algebroids to M-theory, *Proc. Sci. CORFU2018* (2019) 127.

[5] R. Sun, U-duality and courant algebroid in exceptional field theory, [arXiv:2211.08286](https://arxiv.org/abs/2211.08286).

[6] E. Malek and H. Samtleben, Kaluza-Klein Spectrometry for Supergravity, *Phys. Rev. Lett.* **124**, 101601 (2020).

[7] D. Cassani, G. Josse, M. Petrini, and D. Waldram, Systematics of consistent truncations from generalised geometry, *J. High Energy Phys.* **11** (2019) 017.

[8] A. Ashmore, M. Petrini, E. L. Tasker, and D. Waldram, Exactly Marginal Deformations and Their Supergravity Duals, *Phys. Rev. Lett.* **128**, 191601 (2022).

[9] A. Ashmore, M. Gabella, M. Graña, M. Petrini, and D. Waldram, Exactly marginal deformations from exceptional generalised geometry, *J. High Energy Phys.* **01** (2017) 124.

[10] P. Ševera, Letters to Alan Weinstein about courant algebroids, [arXiv:1707.00265](https://arxiv.org/abs/1707.00265).

[11] M. Alexandrov, A. Schwarz, O. Zaboronsky, and M. Kontsevich, The geometry of the master equation and topological quantum field theory, *Int. J. Mod. Phys. A* **12**, 1405 (1997).

[12] P. Ševera, Poisson-Lie T-duality as a boundary phenomenon of Chern-Simons theory, *J. High Energy Phys.* **05** (2016) 044.

[13] A. S. Arvanitakis, C. D. A. Blair, and D. C. Thompson, A QP perspective on topology change in Poisson-Lie T-duality, *J. Phys. A* **56**, 255205 (2023).

[14] M. J. Duff, P. S. Howe, T. Inami, and K. S. Stelle, Superstrings in $D = 10$ from supermembranes in $D = 11$, *Phys. Lett. B* **191**, 70 (1987).

[15] F. Bonechi, P. Mnev, and M. Zabzine, Finite dimensional AKSZ-BV theories, *Lett. Math. Phys.* **94**, 197 (2010).

[16] Z. Kókényesi, A. S. Sinkovics, and R. J. Szabo, AKSZ constructions for topological membranes on G_2 -manifolds, *Fortschr. Phys.* **66**, 1800018 (2018).

[17] A. S. Cattaneo, J. Qiu, and M. Zabzine, 2D and 3D topological field theories for generalized complex geometry, *Adv. Theor. Math. Phys.* **14**, 695 (2010).

[18] A. Chatzistavrakidis, C. J. Grewcoe, L. Jonke, F. S. Khoo, and R. J. Szabo, BRST symmetry of doubled membrane sigma-models, *Proc. Sci. CORFU20182019* (2019) 147.

[19] Z. Kókényesi, A. S. Sinkovics, and R. J. Szabo, Double field theory for the A/B-models and topological S-duality in generalized geometry, *Fortschr. Phys.* **66**, 1800069 (2018).

[20] F. Bonechi, A. Cabrera, and M. Zabzine, AKSZ construction from reduction data, *J. High Energy Phys.* **07** (2012) 068.

[21] G. Barnich and M. Grigoriev, A Poincaré lemma for sigma models of AKSZ type, *J. Geom. Phys.* **61**, 663 (2011).

[22] A. S. Arvanitakis, Formal exponentials and linearisations of QP-manifolds, [arXiv:2204.12613](https://arxiv.org/abs/2204.12613).

[23] P. K. Townsend, Four lectures on M theory, in *Proceedings of the ICTP Summer School in High-energy Physics and Cosmology* (1996), pp. 385–438, [arXiv:hep-th/9612121](https://arxiv.org/abs/hep-th/9612121).

[24] T. T. Voronov, Q-manifolds and mackenzie theory, *Commun. Math. Phys.* **315**, 279 (2012).

[25] G. Mitric and I. Vaisman, Poisson structures on tangent bundles, *Differential Geometry and its Applications* **18**, 207 (2003).

[26] N. Ikeda and T. Strobl, BV and BFV for the h-twisted poisson sigma model, *Ann. Henri Poincaré* **22**, 1267 (2021).

[27] C. Klimcik and T. Strobl, WZW - Poisson manifolds, *J. Geom. Phys.* **43**, 341 (2002).

[28] N. Ikeda, Lectures on AKSZ sigma models for physicists, in *Proceedings of the Workshop on Strings, Membranes and Topological Field Theory* (World Scientific Publishing Company, Singapore, 2017), pp. 79–169, [10.1142/9789813144613_0003](https://doi.org/10.1142/9789813144613_0003).

[29] A. Cabrera and M. Cueca, Dimensional reduction of courant sigma models and lie theory of poisson groupoids, *Lett. Math. Phys.* **112**, 104 (2022).

[30] S. A. Cherkis and J. H. Schwarz, Wrapping the M theory five-brane on K3, *Phys. Lett. B* **403**, 225 (1997).

[31] J. Park and W. Sim, Supersymmetric heterotic action out of M5 brane, *J. High Energy Phys.* **08** (2009) 047.

[32] D. Roytenberg, AKSZ-BV formalism and courant algebroid-induced topological field theories, *Lett. Math. Phys.* **79**, 143 (2007).

[33] N. Ikeda and K. Uchino, QP-structures of degree 3 and 4D topological field theory, *Commun. Math. Phys.* **303**, 317 (2011).

[34] C. M. Hull, Generalised geometry for M-theory, *J. High Energy Phys.* **07** (2007) 079.

[35] P. Pires Pacheco and D. Waldram, M-theory, exceptional generalised geometry and superpotentials, *J. High Energy Phys.* **09** (2008) 123.

[36] O. Hohm, C. Hull, and B. Zwiebach, Generalized metric formulation of double field theory, *J. High Energy Phys.* **08** (2010) 008.

[37] M. Gualtieri, Generalized complex geometry, *Ann. Math.* **174**, 75 (2011).

[38] P. Severa, Some title containing the words “homotopy” and “symplectic”, e.g., this one, [arXiv:math/0105080](https://arxiv.org/abs/math/0105080).

[39] D. Tennyson and D. Waldram, Exceptional complex structures and the hypermultiplet moduli of 5d Minkowski compactifications of M-theory, *J. High Energy Phys.* **08** (2021) 088.

[40] A. Ashmore, C. Strickland-Constable, D. Tennyson, and D. Waldram, Heterotic backgrounds via generalised geometry: moment maps and moduli, *J. High Energy Phys.* **11** (2020) 071.

[41] A. Ashmore, C. Strickland-Constable, D. Tennyson, and D. Waldram, Generalising G_2 geometry: Involutivity, moment maps and moduli, *J. High Energy Phys.* **01** (2021) 158.

[42] H. Bursztyn, N. Martinez Alba, and R. Rubio, On higher dirac structures, *Int. Math. Res. Not.* **2019**, 1503 (2017).

[43] M. Cueca, The geometry of graded cotangent bundles, *J. Geom. Phys.* **161**, 104055 (2021).

[44] J. Qiu and M. Zabzine, On the AKSZ formulation of the Rozansky-Witten theory and beyond, *J. High Energy Phys.* **09** (2009) 024.

[45] A. Ashmore, A. Coimbra, C. Strickland-Constable, E. E. Svanes, and D. Tennyson, Topological G_2 and $Spin(7)$ strings at 1-loop from double complexes, *J. High Energy Phys.* **02** (2022) 089.

[46] M. Józwikowski and M. Rotkiewicz, A note on actions of some monoids, *Differential Geometry and its Applications* **47**, 212 (2016).

[47] J. Śniatycki and A. Weinstein, Reduction and quantization for singular momentum mappings, *Lett. Math. Phys.* **7**, 155 (1983).