

Looking for the sponge loop: analyses of detritus on a Caribbean forereef using stable isotope and eDNA metabarcoding techniques

Lauren K. Olinger^{1,2}, Beverly McClenaghan³, Mehrdad Hajibabaei^{3,4}, Nicole Fahner³, Lesley Berghuis³, Hoda Rajabi³, Patrick Erwin², Chad S. Lane⁵ and Joseph R. Pawlik²

¹ Center for Marine and Environmental Studies, University of the Virgin Islands, St Thomas, Virgin Islands, U.S. Virgin Islands, United States of America

² Department of Biology and Marine Biology, University of North Carolina Wilmington, Wilmington, NC, United States of America

³ eDNAtec Inc., Newfoundland and Labrador, St. John's, Canada

⁴ Department of Integrative Biology, University of Guelph, Guelph, Ontario, Canada

⁵ Department of Earth and Ocean Sciences, University of North Carolina Wilmington, Wilmington, NC, United States of America

ABSTRACT

Coral reefs are biodiverse ecosystems that rely on trophodynamic transfers from primary producers to consumers through the detrital pathway. The sponge loop hypothesis proposes that sponges consume dissolved organic carbon (DOC) and produce large quantities of detritus on coral reefs, with this turn-over approaching the daily gross primary production of the reef ecosystem. In this study, we collected samples of detritus in the epilithic algal matrix (EAM) and samples from potential sources of detritus over two seasons from the forereef at Carrie Bow Cay, Belize. We chose this location to maximize the likelihood of finding support for the sponge loop hypothesis because Caribbean reefs have higher sponge abundances than other tropical reefs worldwide and the Mesoamerican barrier reef is an archetypal coral reef ecosystem. We used stable isotope analyses and eDNA metabarcoding to determine the composition of the detritus. We determined that the EAM detritus was derived from a variety of benthic and pelagic sources, with primary producers (micro- and macroalgae) as major contributors and metazoans (Arthropoda, Porifera, Cnidaria, Mollusca) as minor contributors. None of the sponge species that reportedly produce detritus were present in EAM detritus. The cnidarian signature in EAM detritus was dominated by octocorals, with a scarcity of hard corals. The composition of detritus also varied seasonally. The negligible contribution of sponges to reef detritus contrasts with the detrital pathway originally proposed in the sponge loop hypothesis. The findings indicate a mix of pelagic and benthic sources in the calmer summer and primarily benthic sources in the more turbulent spring.

Submitted 27 November 2023
Accepted 29 January 2024
Published 23 February 2024

Corresponding author
Lauren K. Olinger,
l.olinger12993@gmail.com

Academic editor
Magnus Johnson

Additional Information and
Declarations can be found on
page 23

DOI 10.7717/peerj.16970

© Copyright
2024 Olinger et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Biodiversity, Ecology, Marine Biology, Biogeochemistry

Keywords Detritus, Coral reef, Trophodynamics, Porifera, eDNA, Stable isotope analyses

INTRODUCTION

Coral reefs are among the most biodiverse marine ecosystems despite occurring in oligotrophic tropical waters (Darwin & Bonney, 1896). The paradoxical survival of coral reefs in such marine deserts is made possible by high primary productivity and efficient nutrient cycles (Mumby & Steneck, 2018). However, the survival of coral reefs is at risk due to human-driven climate change, diseases, overfishing, and other stressors (Hughes, 1994; Hoegh-Guldberg *et al.*, 2007). These stressors are driving the replacement of hermatypic corals by non-reef building organisms, such as algae, sponges, and benthic cyanobacterial mats (BCM), reducing reef complexity, accretion, water quality, and overall habitat value for higher trophic levels (Done, 1992; Aronson *et al.*, 2002; Fabricius, 2005; Norström *et al.*, 2009; De Bakker *et al.*, 2017). Changes to benthic communities also cause cascading effects to the nutrient cycles on which these ecosystems rely, including how energy is transferred from primary producers to higher trophic levels through the detrital pathway (Begon, Harper & Townsend, 1986; Johnson *et al.*, 1995; Wilson *et al.*, 2003).

Detritus is a protein- and amino acid-rich mixture (Wilson, 2000; Crossman *et al.*, 2001) of non-living organic matter derived from non-fossil living sources and living organic matter from heterotrophic and autotrophic microbes (Begon, Harper & Townsend, 1986; Wilson *et al.*, 2003). The combination of detritus and inorganic carbonates and silicates forms benthic particulates on coral reefs (Tebbott & Bellwood, 2019). The accumulation of detritus-laden particulates in the turf-forming algae that covers dead coral surfaces is collectively known as the epilithic algal matrix (EAM). This amalgamation of nutritious detritus and algae is an important food resource for many fish and invertebrate detritivores (Hatcher, 1983; Purcell & Bellwood, 2001; Wilson *et al.*, 2003; Kramer, Bellwood & Bellwood, 2012).

Detrital organic matter on coral reefs is derived from a variety of autochthonous and allochthonous benthic and pelagic sources. Micro- and macro-algae are major sources of detritus that accumulate on the benthos following mechanical breakdown by physical forces or through grazing (Hatcher, 1983; Alongi, 1988; Wilson *et al.*, 2003). Algae can also produce detritus indirectly by generating photosynthate, forms of dissolved organic carbon (DOC) that can aggregate into non-cellular amorphous detritus by adsorbing to carbonate sediments or spontaneously forming polymer gels (Otsuki & Wetzel, 1973; Chin, Orellana & Verdugo, 1998; Wilson *et al.*, 2003). Another source of detritus is bacteria (Tebbott & Bellwood, 2019) that can form biofilms and act as nucleation sites for DOC aggregation, and heterotrophic bacteria can consume detritus and mediate its transformation to a form that can be consumed by metazoans (Biddanda, 1985; Alongi, 1988; Wilson *et al.*, 2003). Benthic cyanobacterial mats (BCM) are also becoming more abundant on Caribbean coral reefs (De Bakker *et al.*, 2017) and may be similar to benthic macroalgae with respect to their contribution to detritus, either directly through mechanical breakdown following physical detachment from the reef, or indirectly through production of large quantities DOC (Brocke *et al.*, 2015) that may aggregate and build up in the EAM.

Detritus may also be derived from the shedding of mucus or cells by benthic metazoans, namely corals and sponges. Coral mucus has long been considered an important source

of detritus on coral reefs ([Gottfried & Roman, 1983](#)). Sponges have more recently been implicated in detritus production, and sponge conversion of DOC to detritus may be an important mechanism recycling carbon back to the benthos before it can escape the reef, thereby aiding in tight carbon cycles that ensure reef survival under oligotrophic conditions ([De Goeij et al., 2013](#)). This conversion of DOC to detritus is the theorized sponge loop, named after the similar microbial loop which describes the return of DOC back into the grazing food web *via* predation on free-living microbes by protists ([Azam et al., 1983](#)). Since its proposal, the sponge loop hypothesis has been widely cited and has been featured in undergraduate marine biology textbooks ([Levinton, 2021](#)).

The sponge loop hypothesis was originally proposed based on incubation experiments using flow chambers or open pots spiked with isotopically enriched DOC containing thinly encrusting, cryptic (found in reef interstices) Caribbean reef sponge species, with the isotopic signature traced in detritus collected by filtration or by pipette ([De Goeij et al., 2013](#)). Four cryptic sponge species were used in these experiments (*Halisarca caerulea*, *Haliclona vandoesti*, *Chondrilla caribensis*, and *Scopalina ruetzleri*) with the mechanism of detritus production proposed as rapid cellular proliferation and choanocyte turnover associated with space-limitation and restricted sponge growth ([De Goeij et al., 2013](#)). Detritus production *via* the sponge loop was proposed to be very large, with sponges (both cryptic and massive, or emergent, species) transforming most of the DOC they consumed into detritus, estimated at 9.3% of sponge biomass per day ([Alexander et al., 2014](#)) and with the turnover of DOC to detritus approaching the daily gross primary production of the entire reef ecosystem ([De Goeij et al., 2013](#)). However, subsequent experiments with larger, emergent sponge species that grow on the reef surface failed to find these sponges to be net producers of detritus and instead suggested that they return DOC to the benthos as sponge biomass ([McMurray et al., 2018](#)). More recently, some emergent sponge species, including the Caribbean species *Aplysina archeri*, were observed “sneezing” mucus and particulates from near their incurrent ostia ([Kornder et al., 2022](#)), but it is not yet clear if this is a method of detritus production analogous to that of cryptic sponge species or whether this mechanism is common across the emergent sponge fauna. The discovery of the microbial loop accounted for a large fraction of missing oceanic carbon ([Fenchel, 2008](#)), but it is unclear whether the carbon transformation mediated by the sponge loop is as important a pathway.

Detritus collected from a fore reef off Palmyra Atoll showed abundant pelagic inputs ([Max et al., 2013](#)). Studies such as those conducted on the Palmyra Atoll by [Max et al. \(2013\)](#) used stable isotope analyses (SIA) to determine the sources contributing to benthic detritus. SIA is a popular technique for tracing energy flow through ecosystems, and it is based on the discrimination against heavier isotopes in metabolic processes, which separates organisms from different trophic levels along axes of $^{15}\text{N}/^{14}\text{N}$ and separates organisms with different types of primary productivity and carbon sources along axes of $^{13}\text{C}/^{12}\text{C}$ ([Post, 2002](#); [Michener & Kaufman, 2007](#); [Middelburg, 2014](#); [Zapata-Hernández et al., 2021](#)). SIA has been relatively underused to explore trophodynamics on coral reefs ([Greenwood, Sweeting & Polunin, 2010](#)).

While popular, SIA is not able to discern between species with similar trophic levels and carbon-fixing pathways. This limitation can be resolved with supplemental collection and metabarcoding of environmental DNA (eDNA) samples, an increasingly popular approach for characterizing biodiversity that does not require collection of whole biological specimens and instead relies on recovery and analysis of DNA from the physical environment in which they live (e.g., water, soil, sediment; [Thomsen & Willerslev, 2015](#)). This eDNA is released from organisms through various mechanisms, including cell shedding and the excretion of various bodily fluids and feces, and it can be isolated, amplified, and sequenced using high-throughput genomic sequencing platforms and then screened against publicly available databases (e.g., GenBank) to assign taxonomy to the organisms contributing to that environmental sample ([Baird & Hajibabaei, 2012](#); [Taberlet et al., 2012](#); [Thomsen & Willerslev, 2015](#)). Analyses of eDNA have been used to quantify biodiversity ([Thomsen & Willerslev, 2015](#)), determine species diets ([Boyi et al., 2022](#)), and identify elusive species ([McClanaghan et al., 2020](#)). Analyses of eDNA can also complement SIA-based estimations of source contributions to organic matter resources, such as detritus. A similar SIA/eDNA approach was recently used to quantify the relative contributions of different sources to sedimentary organic carbon in seagrass meadows ([Reef et al., 2017](#)).

In this study, we chose to test the sponge loop hypothesis from the “sink side”, focusing on the sponge signature in natural EAM detritus accumulating on a coral reef, rather than from the “source side” as detritus is produced by targeted sponges, whether in laboratory incubations or in the field ([De Goeij et al., 2013](#); [McMurray et al., 2018](#), reviewed in [Pawlik & McMurray, 2020](#)). Further, we chose a location for our study that would maximize the likelihood of confirming the sponge loop hypothesis in providing high sponge abundance. Caribbean reefs are distinct from others in the tropics in having higher sponge abundance ([Wilkinson & Cheshire, 1990](#)), with mean percentage cover of sponges on Caribbean fore reefs of ~16% ([Loh & Pawlik, 2014](#)). If, as proposed by the sponge loop hypothesis, >9% of sponge biomass per day is returned to the benthos as detritus and sponges cover >10% of the reef surface, we would anticipate that the sponge signal in natural EAM detritus would be very large. We collected samples of reef EAM detritus and samples from potential sources of detritus, including from organisms and from areas where detritus may originate (e.g., water column, fish feces) from a Caribbean fore reef within the archetypal Mesoamerican barrier reef ecosystem (Carrie Bow Cay, Belize) over two seasons (July 2018 and March 2019). Stable isotope and eDNA metabarcoding analyses were applied to the samples to explore the composition of detritus, with a particular focus on sponges to test claims that sponges are major sources of detritus on Caribbean coral reefs ([De Goeij et al., 2013](#)). In addition to testing the sponge loop hypothesis, this study provided an opportunity to further develop both SIA- and eDNA-based methodologies, allowing comparisons across sampling seasons for the same reef location and comparisons with other studies.

MATERIALS & METHODS

Sample collection

Samples were collected in July 2018 and March 2019 from the fore reef near the Smithsonian field station at Carrie Bow Cay, Belize. Carrie Bow Cay is situated on the Mesoamerican Barrier Reef, approximately 22 km from the city of Dangriga on mainland Belize. The characteristic spur-and-groove formation of the fore reef at 15–20 m depth is an *Orbicella annularis* framework with abundant octocorals, sponges, and algae, and with hard coral cover estimated at ~10% at the times of this study. The fore reef location (16°48'10.6"N, 88°4'43.9"W) approximately 500 m northeast of the island of Carrie Bow Cay drops off steeply at the outer ridge less than 100 m to the east. Cuts to the north and south of the island and reef isolate the complex from the rest of the Mesoamerican barrier reef and expose the area to currents that flow through the channels, including waters from the open ocean and from the shoreward lagoon with large mangrove islands and seagrass beds nearby. The reef complex at Carrie Bow Cay has been extensively studied since its founding as a research site by scientists in 1972 ([Rützler & Macintyre, 1982](#)). Field experiments were approved by the Belize Fisheries Department (permit number 00023-18).

Two categories of samples were collected from the fore reef: detritus and detritus sources, where the latter refers to organisms or materials that may be contributing to detritus. Samples also encompassed either tissues or composites, where tissues refer to organic material deliberately collected from a living organism and composites refer to organic remnants of multiple organisms that had been previously deposited to the water column or benthos as particulates or feces.

Two types of detritus were collected: detritus from the epilithic algal matrix (EAM detritus) and detritus from plastic trays placed under reef overhangs (tray detritus), with the latter added for the second sampling season. Seven types of source samples were collected: algae tissue, benthic cyanobacterial mat (BCM) tissue, feces from two species of herbivorous fish, feces from two species of spongivorous fish, tissues of three emergent sponge species, tissues of three cryptic sponge species, and water column particulates from sediment traps ([Table 1](#)) The difficulty in collecting some source samples (e.g., spongivores feces) is reflected in lower levels of sample replication.

EAM and tray detritus

Detrital material was suctioned from a 25 cm² area of epilithic algal matrix (EAM) using 50 ml polypropylene syringes. EAM samples (2018 $n = 5$, 2019 $n = 11$) were collected at least 10 m apart and from flat patches of dead rock covered with closely cropped turf and devoid of pits ([Purcell & Bellwood, 2001](#)). In 2019, trays (white plastic bins with dimensions 40 × 31.8 × 15.2 cm; $n = 6$) were deployed under reef overhangs to target detrital outfall from organisms, including cryptic sponges. After 72 h, the particulates and detritus that had accumulated in the trays were suctioned up using 50 ml polypropylene syringes. Syringes of EAM and tray detritus were transported back to the lab at Carrie Bow Cay, where their contents were filtered through a 150 µm mesh to isolate the smaller size fraction targeted by detritivores ([Wilson et al., 2003](#)). The mixture was divided, with approximately 5 ml

Table 1 Number of replicates of detritus and detritus sources collected in July 2018 and March 2019 and analyzed using stable isotope analyses (n_{SIA}) and genetic sequencing (n_{seq}).

Category	Sample type	Species	July 2018		March 2019	
			n_{SIA}	n_{seq}	n_{SIA}	n_{seq}
Detritus	EAM	Total	5	5	11	6
	Tray	Total	0	0	3	2
		<i>Dictyota</i> sp.	5	5	5	0
	Algae tissue	<i>Halimeda</i> sp.	0	0	3	0
		<i>Lobophora variegata</i>	5	5	6	0
		Total	10	10	14	0
Source	BCM tissue	Total	3	3	5	0
		<i>Acanthurus bahianus</i>	3	3	4	6
	Herbivore feces	<i>Acanthurus coeruleus</i>	2	2	1	1
		Total	5	5	5	7
		<i>Holacanthus ciliaris</i>	1	1	1	2
	Spongivore feces	<i>Pomacanthus paru</i>	0	0	2	2
Emergent sponge tissue		Total	1	1	3	4
		<i>Aplysina caerulea</i>	10	0	10	0
		<i>Niphates digitalis</i>	8	0	10	0
		<i>Xestospongia muta</i>	10	5	10	0
		Total	28	5	30	0
		<i>Chondrilla</i> sp.	2	2	3	0
Cryptic sponge tissue		<i>Halisarca caerulea</i>	3	3	4	0
		<i>Scopalina ruetzleri</i>	3	3	5	0
		Total	8	8	12	0
	Sediment trap	Total	5	5	14	7

Notes.

Composite samples are indicated in bold.

transferred to a microcentrifuge tube with 0.5 ml ethanol and the remaining filtered onto a 0.7 μ m filter (Watson GFF), and both samples were stored at -20°C .

Tissue samples

The tissues of organisms that are abundant and possible contributors to reef detritus were collected at the same location and time (July 2018, March 2019) as detritus collections. Divers sought out different individuals or patches of each targeted organism; species were visually identified on site, and identifications were later validated with genetic sequence data for all but the most conspicuous organisms (*Aplysina caerulea*, *Niphates digitalis*, and *Halimeda* sp.). When collecting the tissue samples, the sample quantity and collection method varied by species. The algae and benthic cyanobacterial mat (BCM) tissues were collected in handfuls, using snips to separate algae or bacterial tissues from each source patch, and transferred to the lab at Carrie Bow Cay in plastic bags. The tissues of emergent sponge species *Xestospongia muta*, *A. caerulea*, and *N. digitalis* were collected with a dive knife or cork borer (one cm diameter), and clean pieces of tissue (approximately eight cm³ or two cm cylinders) were transported to the lab at Carrie Bow Cay in plastic bags. Tissues

of cryptic sponge species *Chondrilla* sp., *Halisarca caerulea*, and *Scopalina ruetzleri* were removed from rock using a scalpel blade, producing tissue samples that were approximately four cm² and 1–2 mm thick. All tissue samples were sealed at depth in plastic bags and transported back to the lab, where a subsample of tissue (<1 cm³) from each organism was transferred to a microcentrifuge tube, rinsed twice with ethanol, and stored in 0.5 ml ethanol at –20 °C. The plastic bag with the remaining tissue sample was squeezed of excess water, resealed, and frozen at –20 °C.

Feces samples

Feces of two herbivorous fish species (*Acanthurus bahianus*, *Acanthurus coeruleus*) and two spongivorous fish species (*Holacanthus ciliaris*, *Pomacanthus paru*) were also collected. For these samples, divers observed fish, and when the fish was seen defecating, the diver used a turkey baster to collect the feces before it touched the benthos. The contents of the turkey baster were transferred to a plastic bag, sealed at depth, and transported back to the lab. The mixture was divided, with approximately 5 ml transferred to a microcentrifuge tube with 0.5 ml ethanol and the remaining filtered onto a 0.7 µm filter (Watson GFF), and both samples were stored at –20 °C.

Sediment trap samples

Sediment traps made of PVC pipe with diameter of 7.6 cm and length of 30.5 cm (4:1 aspect ratio) were deployed on the reef for 72 h to capture water column particulates that may be contributing to detritus. While seawater samples would have been a more representative sample of pelagic contributions to reef detritus, this was not feasible due to the volume of water (>>4 L) required for stable isotope analyses. Traps (2018 $n = 5$; 2019 $n = 15$; [Table 1](#)) were deployed at least 10 m apart on the reef and affixed with cable ties to a metal stake that had been securely driven into dead coral heads. After 72 h, it was determined that sufficient sediment had accumulated for analyses, and plastic bags and rubber bands were used to seal sediment traps at depth before being transported to the lab at Carrie Bow Cay for processing. The mixture was divided, with approximately 5 ml transferred to a microcentrifuge tube with 0.5 ml ethanol and the remaining filtered onto a 0.7 µm filter (Watson GFF), and both samples were stored at –20 °C.

Stable isotope analyses

Samples were transported on ice to University of North Carolina Wilmington (UNCW), where tissue samples and filters containing composite samples were transferred to combusted scintillation vials and lyophilized for 24 h. Dried tissue samples were ground into a powder using a mortar and pestle, and the powder was returned to the scintillation vials. Filters were cut in half using a scalpel. One half of each filter and approximately 10 mg of each tissue sample was transferred to a second combusted vials for acidification by exposure to vapors of 12 M hydrochloric acid (HCl) for 24 h. Acidified samples were then placed on a 60 °C hotplate for 48 h to boil off remaining HCl. Each of the non-acidified and acidified filter halves were halved again, and the quarter filter segments were transferred to foil capsules. For each non-acidified and acidified tissue sample, an ultra-microbalance was used to precisely measure (within 5%) a target weight of sample into foil capsules.

Target weights varied from 2 mg for *Dictyota* to 7 mg for *Lobophora*, and 4 mg for all other sampled species (six sponge species, BCM, and *Halimeda*).

Non-acidified and acidified samples were analyzed for nitrogen and carbon stable isotope compositions, respectively, using a Costech 4010 Elemental Analyzer interfaced with a Thermo Delta V Plus stable isotope mass spectrometer at UNCW. Isotopic results are expressed in standard delta (δ) units calculated as follows: $\delta^{13}\text{C}$ or $\delta^{15}\text{N}$ (‰) = $1000[(R_{\text{sample}}/R_{\text{standard}}) - 1]$, where R_{sample} and R_{standard} are the $^{13}\text{C}/^{12}\text{C}$ or $^{15}\text{N}/^{14}\text{N}$ ratios of samples and standards, respectively. Within-run standard deviations of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were $< 0.3\text{‰}$ across repeated analyses of USGS 40 and USGS 41a glutamic acid standards (average = 0.2‰ , min = 0.1‰). A total of 8 samples from 2019 were omitted from analyses due to insufficient acidification leading to remnant inorganic carbon species, as determined by $\delta^{13}\text{C}$ values that exceeded -10 (Schidlowski, 2001). For fish feces samples collected in 2019, the stable isotope composition of three feces samples ($n = 2$ from herbivore *Acanthurus bahianus* and $n = 1$ from spongivore *Holocanthus ciliarus*) were not measured due to insufficient material on the filter.

Metabarcoding analyses

The samples stored in microcentrifuge tubes in ethanol were shipped to the Center for Environmental Genomics Analysis (CEGA) for sequencing in 2019. CEGA designed and optimized a fully custom eDNA analysis. DNA extraction methods were optimized for each sample type to maximize DNA recovery, and DNA was extracted from all samples using the optimized protocols. Three DNA markers partially covering three genetic regions (cytochrome *c* oxidase I, 18S rRNA, 16S rRNA) were amplified from each sample using PCR (Tables S1–S2). These markers were chosen to provide comprehensive detection of metazoans, algae, and cyanobacteria. Negative controls were added during extraction and PCR steps and carried through to sequencing. The three amplicons from each sample were sequenced on an Illumina NovaSeq 6000 instrument with a target minimum sequencing depth of 250,000 sequences per sample per marker. See [Supplemental Information](#) for detailed DNA extraction and library preparation procedures. Raw sequence reads are available in NCBI's sequence read archive under project [PRJNA965826](#).

After DNA sequencing, base calling and demultiplexing were performed using Illumina's *bcl2fastq* software (v2.20), and primers were trimmed from sequences using *cutadapt* (v2.8; Martin, 2011). Using *DADA2* (v1.14; Callahan *et al.*, 2016) with default parameters, sequences were filtered for quality and length and then denoised to create exact sequence variants (ESVs), each of which represent a unique sequence from the sample. ESVs were assigned taxonomy using NCBI's BLASTN tool (v2.11; Altschul *et al.*, 1990) and the NT database (downloaded November 25, 2020) with an e-value cut-off of 0.001. The taxonomy presented here matches the naming conventions used in NCBI's taxonomy database and was assigned based on a sequence similarity score (the product of the percent sequence similarity and the percent query coverage). The minimum scores for taxonomic assignment at each level were as follows: phylum at 85%, class and order at 90%, family at 95%, genus at 98%, and species at 99%. The results from all markers were consolidated to create the taxonomic lists. Taxonomic identifications were verified against the World

Register of Marine Species (WoRMS), Global Biodiversity Information Facility (GBIF), and the Encyclopedia of Life (EOL) to ensure that spurious matches resulting from poor reference database coverage were not included in the list. As standard practice, ESVs assigned to humans and other common contaminants were removed from the sample data. Sequencing results from the negative controls generated during lab processing were screened for contamination, and any ESVs detected in the negative controls were removed from the associated samples. No ESVs were amplified from negative controls that were also detected in associated samples and could be assigned taxonomy based on the selection criteria described above.

Data analyses

Bayesian stable isotope mixing models (R package SIMMR; [Parnell, 2019](#)) were used for the two tracer isotopes (^{13}C , ^{15}N). For July 2018 samples, the mixing model was parameterized using the EAM detritus as the mixture and 11 sources: algae tissue (combined *Dictyota* and *Lobophora*), BCM tissue, herbivore feces (combined *A. bahianus* and *A. coeruleus*), spongivore feces (*H. ciliaris*), emergent sponge tissue from three species (*Aplysina cauliformis*, *Niphates digitalis*, and *Xestospongia muta*), cryptic sponge tissue from three species (*Chondrilla* sp., *Halisarca* sp., *Scopalina ruetzleri*), and sediment trap samples. The algae tissue types were initially pooled as a single source because of the large overlap between the two species in isotopic space, and the herbivore feces were combined for the same reason. However, sponge species were retained as separate sources in the mixing model due to little overlap in isotopic space. For March 2019 samples, two mixing models were parameterized (one each for EAM and tray detritus) with the same number and type of sources as the 2018 samples, with the addition of *Halimeda* sp. to the combined algae tissue category and feces of *P. paru* to the combined spongivore feces category. Each model was run with 10,000 iterations, thinned by 10, and with an initial discard of the first 1,000 iterations. We examined correlations among the posterior distributions in each model to assess the ability of the model to distinguish between different sources, pooled highly correlated sources, and re-ran the model with the pooled sources, keeping other settings the same.

For genetic sequencing data, exact sequence variant (ESV) identifications were tabulated across sample types and years to determine the frequency of occurrence at the level of taxonomic phyla and species, where frequency of occurrence was calculated as the number of samples in which the taxa (either phyla or species) was present, divided by the total number of sequenced samples of that sample type. Multivariate analyses were used to compare the detritus (EAM and tray) from 2018 and 2019. A non-metric multidimensional scaling (NMDS) biplot based on the Jaccard similarity measure was used to visualize differences among detritus types. ESV identifications were converted to presence/absence values at the level of taxonomic order. Ellipses generated from the standard error of the weighted average of the scores in each treatment habitat were superimposed on each biplot to visualize the dispersion of each detritus type. A permutational analysis of variance (PERMANOVA) was conducted to test the multivariate response of taxonomic order presence/absence across the detritus types. When PERMANOVAs were significant

($p < 0.05$) a similarity percentage analysis (SIMPER) was used to evaluate the contribution of each taxonomic order to differences between detritus types. All statistical analyses were performed in R ([R Core Team, 2020](#)).

RESULTS

Stable isotope analyses

Detritus $\delta^{13}\text{C}$ values ranged from $-19.5\text{\textperthousand}$ (2019 EAM detritus) to $-14.4\text{\textperthousand}$ (2019 tray detritus), and source $\delta^{13}\text{C}$ values ranged from $-27.7\text{\textperthousand}$ (2019 BCM tissue) to $-10.7\text{\textperthousand}$ (2019 *Halisarca caerulea* tissue; [Table S3](#)). Detritus $\delta^{15}\text{N}$ values ranged from $0.2\text{\textperthousand}$ to $2.7\text{\textperthousand}$ (both from 2019 EAM detritus), and source $\delta^{15}\text{N}$ values ranged from $-2.6\text{\textperthousand}$ (2018 sediment traps) to $5.5\text{\textperthousand}$ (2018 *Scopalina ruetzleri* tissue). In general, $\delta^{15}\text{N}$ values were greater in sponge tissue samples than algae and BCM tissue samples ([Fig. 1](#); [Table S4](#)). The isotope signatures of the feces of spongivorous fishes (*Holacanthus ciliaris*, *Pomacanthus paru*) were similar to sponge tissues, and the isotope signatures of the feces of herbivorous fishes (*Acanthurus bahianus*, *Acanthurus coeruleus*) were similar to algae tissues ([Fig. 1](#); [Tables S3–S4](#)).

There were notable seasonal changes in stable isotope compositions of some composite and tissue samples between July 2018 and March 2019. For the $\delta^{15}\text{N}$ values of sediment trap samples, there was a large increase and shift from negative to positive $\delta^{15}\text{N}$ values from 2018 (mean $\pm \text{SD} = -1.1 \pm 1.3\text{\textperthousand}$) to 2019 ($-1.7 \pm 0.5\text{\textperthousand}$; [Table S4](#)). The emergent sponge species *A. cauliformis* and cryptic sponge species *Chondrilla* sp. exhibited similar isotopic values, and both exhibited increasing $\delta^{13}\text{C}$ values and decreasing $\delta^{15}\text{N}$ values from 2018 to 2019 ([Fig. 1](#); [Tables S3–S4](#)). The $\delta^{13}\text{C}$ of the cryptic sponge species *Halisarca caerulea* also increased from $-17.9 \pm 1.4\text{\textperthousand}$ to $-12.6 \pm 2.8\text{\textperthousand}$, and the $\delta^{13}\text{C}$ of BCM decreased markedly from $-21.1 \pm 0.1\text{\textperthousand}$ to $-27.1 \pm 0.5\text{\textperthousand}$, and this was the only tissue sample type with reduced $\delta^{13}\text{C}$ values in 2019 ([Table S3](#)).

The Bayesian isotope mixing model based on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values was used to establish the relative contribution of each source to the detritus samples ([Fig. 1](#)). For 2018 samples, algae and herbivore feces were pooled after initial runs indicated a high correlation between the sample types (correlation = -0.72). The resulting model showed further correlations between the combined algae/herbivore feces source category and BCM tissue and sediment trap samples (correlation = -0.63 , -0.74 respectively); these were further pooled into a category “algae/BCM/herbivore feces/sediment trap” ([Fig. S1](#)). The resulting model indicated high contribution by the pooled four source mixture to the EAM detritus (mean $\pm \text{SD} = 58.3 \pm 11.1\text{\textperthousand}$), with the other source categories showing much lower contribution ($<8.5\%$; [Fig. 2A](#)). Earlier iterations of the model with unpooled sources showed high contribution by algal tissue and herbivore feces ($44.8 \pm 17\%$, $11.9 \pm 14.5\%$, respectively; [Fig. S2](#)), suggesting the disproportionate contribution was not merely a byproduct of the pooling of four sources.

The mixing model of 2019 EAM detritus samples showed no major correlations among detrital sources, thus analyses proceeded on the original model with unpooled sources. The mixing model revealed similarly low contributions with large confidence intervals for




Figure 1 Isotope biplots of samples collected in (A) July 2018 and (B) March 2019. Detritus samples, including EAM and tray detritus, are represented by open and filled circles, respectively, that denote the values of individual samples. All other symbols represent the average \pm SD of source samples.

[Full-size](#) DOI: [10.7717/peerj.16970/fig-1](https://doi.org/10.7717/peerj.16970/fig-1)

all sources, ranging from $4.5 \pm 3.3\%$ for BCM tissue to $16 \pm 13.5\%$ for algal tissue (Fig. 2B), and likely owing to the greater variability in EAM detritus isotopic values and tighter clustering of source samples around the EAM detritus (Fig. 1B). The tray detritus mixing model showed similarly low contributions and large confidence intervals for all sources (Fig. 2C).




Figure 2 Simulated contributions of each source to EAM detritus in (A) July 2018 and (B) March 2019, and to (C) tray detritus in March 2019. In the box plots, the boundary of the box closest to zero indicates the 25th percentile, a black line within the box marks the median, and the boundary of the box farthest from zero indicates the 75th percentile. Whiskers above and below the box indicate the 10th and 90th percentiles, and outermost points indicate outliers.

Full-size DOI: 10.7717/peerj.16970/fig-2

Metabarcoding analyses

A total of 119,012,810 sequence reads were generated with a mean of 577,732 ($\pm 302,189$ SD) DNA sequence reads retained per sample per marker after bioinformatic filtering. The 18S marker yielded a mean of 1,121 ($\pm 2,991$ SD) ESVs per sample, COI yielded a mean of

452 (± 390 SD) ESVs per sample, and 16S yielded a mean of 11,657 ($\pm 26,518$ SD) ESVs per sample. Approximately 37.5% of ESVs (295,497 ESVs) were assigned taxonomy based on the selection criteria outlined in the methods.

Genetic sequences of tissue samples largely matched the morphology-based taxonomy (Table S5). Three out of four sponge species analyzed were identified to the genus level, and the fourth sponge species (*Halisarca caerulea*) was identified at the family level (Halisarcidae), with the expected genus *Halisarca* scoring a 96%, just below the 98% threshold. The BCM was assigned the expected species *Moorea producens*, and *Lobophora* was assigned its proper genus. The tissue samples from the alga visually identified as *Dictyota* matched to multiple taxa, including phaeophyte genus *Dictyopteris* and rhodophytes in family Rhodomelaceae including *Lomentaria* and *Neosiphonia*.

The composite samples (EAM and tray detritus, as well as source samples from sediment traps and the feces of herbivores and spongivores) contained rich genetic diversity. For the following analyses, we retained eukaryotes and cyanobacteria and excluded Archaea and bacteria other than cyanobacteria. A total of 17 metazoan phyla and 20 phyla from other kingdoms were identified in the composite samples (Fig. 3). Of these phyla, 34 were found in detritus (four unique), 27 were found in sediment trap samples (0 unique), 23 were found in herbivore feces (0 unique) and 21 were found in spongivore feces (2 unique; Fig. 3). Metazoan phyla Arthropoda, Nematoda, Xenacoelomorpha, Annelida, Mollusca, and Porifera were found in over 50% of EAM detritus samples. The most frequent phylum, Arthropoda, was detected in 82% of EAM detritus samples, while Porifera was detected in 55% of EAM detritus samples. Metazoan phyla Arthropoda, Porifera, Chordata, Cnidaria, and Mollusca were found in over 50% of sediment trap samples, and the most frequent phyla were Porifera and Arthropoda (83% each). Metazoan phyla Chordata, Cnidaria, Porifera, and Platyhelminthes were found in over 50% of spongivore feces samples, and the most frequent phyla were Chordata and Cnidaria (80% each) while Porifera was detected in 60% of samples. No metazoan phyla were found in more than 50% of herbivore feces samples (Fig. 3).

On average, non-metazoan phyla had higher frequencies of occurrence than metazoan phyla (Fig. 3). Bacillariophyta, Rhodophyta, Cyanobacteria, and Ciliophora were detected in at least 90% of EAM detritus samples, and those four phyla as well as Chlorophyta, Cercozoa, and Haptista were detected in at least 90% of sediment trap samples. Bacillariophyta, Cyanobacteria and Rhodophyta were detected in at least 90% of herbivore feces samples, and Chlorophyta and Streptophyta were detected in 58% and 50% of herbivore feces samples, respectively. Streptophyta was detected in 100% of spongivore feces samples (Fig. 3).

The main taxa of interest in investigating the sponge-loop hypothesis were sponges, algae, and cyanobacteria. In total, we identified 36 sponge taxa spanning 14 families, 13 genera, and six species (Fig. 4). Multiple sponge taxa were detected in detritus, sediment trap, and spongivore feces samples, including all the sponge taxa identified from the tissue samples, except the family Halisarchidae. However, sponge taxa had very low frequencies of occurrence compared to other algae and metazoans (Fig. 4), and only two of the 36 taxa were detected in more than one of the 11 EAM detritus samples.




Figure 3 The frequencies of occurrence for all phyla in each composite sample in each year. Number of replicates is shown in parentheses. J18, July 2018; M19, March 2019; EAM, detritus suctioned from epilithic algal matrix; tray, detritus sampled from trays placed under reef overhangs; sed. trap, particulates captured in sediment traps; h'vore feces, feces collected from herbivorous fishes; s'vore feces, feces collected from spongivorous fishes.

Full-size DOI: 10.7717/peerj.16970/fig-3




Figure 4 The frequencies of occurrence in composite sample types for the 15 most abundant algae and non-sponge metazoans, and all 36 identified sponge taxa. Number of replicates is shown in parentheses. J18, July 2018; M19, March 2019; EAM, detritus suctioned from epilithic algal matrix; tray, detritus sampled from trays placed under reef overhangs; sed. trap, particulates captured in sediment traps; h'vore feces, feces collected from herbivorous fishes; s'vore feces, feces collected from spongivorous fishes. Classification includes phylum (P), class (C), order (O), family (F), genus (G), and species (S), where applicable. See Tables S6 and S7 for frequencies of all algae and metazoan taxa, respectively.

Full-size DOI: 10.7717/peerj.16970/fig-4

Among the algae (including Cyanobacteria), we identified 25 classes, 107 orders, 163 families, 202 genera, and 159 species. The most frequently detected taxa in detritus samples were also among the most frequently detected taxa in sediment trap samples as well as herbivore feces samples but were less frequent in spongivore feces samples (Fig. 4; Table S6). The cyanobacterium in the provided reference tissue sample was only detected in EAM and tray detritus, while the *Lobophora* algae identified from the tissue samples was detected

in all sample types (Table S6). The *Dictyota* algae tissue did not yield consistent taxonomic identifications and the best algal matches from two samples (family Rhodomelaceae and genus *Dictyopteris*) were detected in EAM detritus, sediment trap samples, and herbivore feces (Table S6).

Metazoans other than Porifera also contributed to detritus. The most frequent metazoan taxa were harpacticoid copepods (phylum Arthropoda), identified in all the EAM detritus samples collected in July 2018 and half of the EAM detritus samples collected in March 2019. Cyclopoid copepods were also well represented in the EAM detritus collected in both years (Fig. 4). Among macrofaunal metazoans, the most frequently occurring in EAM included cnidarians of class Anthozoa (octocorals; Fig. 4), while cnidarians of class Scleractinia (hard corals) were notably scarce, with only one identification in a single EAM sample collected in 2018 (Table S7). Meiofaunal and epifaunal metazoans were more frequent than macrofauna, and Xenacoelomorpha (class Acoela), Platyhelminthes, and Nematoda were frequent in EAM detritus collected in both years (Fig. 4).

There were some notable differences among 2018 and 2019 EAM and tray detritus. The most immediate difference was the number of detected ESVs, and the number of detected ESVs in 2018 EAM detritus (7135 ± 8220 ESVs per sample) outnumbered tray detritus (4150 ± 4164 ESVs per sample) and greatly outnumbered 2019 EAM detritus (466 ± 435 ESVs per sample). There was also relatively low overlap of individual ESVs across seasons, only 245 of the total 34,226 ESVs detected in EAM detritus were present in at least one sample from each season.

The NMDS based on presence-absence of taxonomic orders revealed separation of 2018 and 2019 EAM detritus (Fig. S3). Interestingly, tray detritus collected in 2019 was more similar to 2018 EAM detritus than 2019 EAM detritus, but this may be due to greater similarity in the number of detected ESVs in tray and 2018 EAM detritus. PERMANOVA revealed significant differences in the communities (PERMANOVA $F_{2,12} = 1.7636$, $p = 0.029$), and pairwise PERMANOVA indicated significant differences between the 2018 and 2019 EAM detritus (adjusted $p = 0.024$). The phylum with the greatest contribution to dissimilarity was Bacillariophyta (SIMPER: 11.1%).

Frequency of occurrence data revealed some additional potential drivers of dissimilarity between the 2018 and 2019 EAM detritus samples. For the algae and cyanobacterial taxa, 41 of the top 50 most frequent algae taxa were present in 100% of 2018 EAM detritus samples, while only three of the top 50 taxa were present in 100% of 2019 EAM detritus samples (Table S6). Additionally, sequences matching to collected algae tissues were more frequent in 2018 EAM detritus compared to 2019 EAM detritus, particularly *Lobophora* (Table S6).

DISCUSSION

Unlike studies that led to the proposal of the sponge loop hypothesis (De Goeij *et al.*, 2013) and those that subsequently tested it (McMurray *et al.*, 2018, reviewed in Pawlik & McMurray, 2020) by examining sponge detritus production on the “source side”, this study tested the “sink side” of the sponge loop, exploring the sponge signature in the

accumulating detritus on the reef surface. The detritus in the epilithic algal matrix (EAM) on the Carrie Bow Cay fore reef was derived from numerous benthic and pelagic sources. Primary producers were the dominant group of organisms contributing to detritus, consistent with previous studies (Hatcher, 1983; Alongi, 1988; Wilson, Burns & Codi, 2001; Wilson *et al.*, 2003). There was genetic signature of sponges (phylum Porifera) in detritus, confirming the validity of the eDNA methodology, but detritus contained low frequencies of Porifera sequences compared to primary producers and other metazoans, and detritus isotopic signatures were distinct from isotopic signatures of sponge tissues, indicating minimal sponge inputs to detritus. The use of complementary analyses and seasonal sampling herein yielded additional insights into how physical conditions (flow, currents, turbidity) may influence detritus composition.

Minimal sponge inputs to detritus

The results of this study are at odds with the detrital component of the sponge loop hypothesis, which proposed that “sponges transform a majority of DOM into particulate detritus” as shed sponge cells with “DOM turnover by sponges approaching the daily gross primary production of the entire reef ecosystem” (De Goeij *et al.*, 2013). Detritus contributions by sponges were proposed to be approximately 6-fold greater than contributions from other sources (De Goeij *et al.*, 2013). Considering these estimates and that the expected form of sponge-derived detrital material is cellular debris, the sponge signal from eDNA and stable isotopic signatures of fore reef detritus samples should have been substantial, but it was not.

The six sponges sampled here are common species that represented a cross section of sponge diversity on Caribbean reefs (Loh & Pawlik, 2014). This diversity was evident in the range of isotopic space that species occupied, and such isotopic differences can be attributed to species-specific diets, microbiomes, and nitrogen cycling and fractionation pathways (Freeman *et al.*, 2021; Van Duy, Mueller & Meesters, 2018). Despite a diverse sampling of sponge species, isotope mixing models indicated low contribution by any species, consistent with the much lower frequencies of occurrence of Porifera sequences in eDNA analyses compared to algae and metazoans, especially when higher-order taxonomic groupings are compared. While frequencies are not true measurements of relative abundance of these diverse organisms, previous studies have demonstrated that DNA frequencies can approximate community abundance (Yoccoz *et al.*, 2012), and low frequencies of Porifera sequences also paralleled differences between sponge and detritus isotopic signatures.

The deployment of trays under reef overhangs in 2019 was a generous test of the sponge loop hypothesis designed to further maximize the odds of capturing choanocytes shed by cryptic sponge species. The three cryptic sponge species targeted in the present study (*Halisarca caerulea*, *Chondrilla caribensis*, and *Scopalina ruetzleri*) were also the same species for which detritus production had been measured using incubation experiments and cited to support the sponge loop hypothesis, with *H. caerulea* being the most frequently used target species for these studies (De Goeij *et al.*, 2008a; De Goeij *et al.*, 2008b; De Goeij *et al.*, 2009; De Goeij *et al.*, 2013; Lesser *et al.*, 2020; Campana *et al.*, 2021; Hudspith *et al.*, 2021). Notably, none of the EAM or tray detritus samples collected in the present study contained

sequences matching to *H. caerulea* (or *Halisaeca*), despite the relative abundance of this species on the Carrie Bow Cay forereef. Sequences matching to *S. ruetzleri* and *Chondrilla* sp. were also absent from all detritus samples, except for one identification of *Chondrilla* sp. in one of the two sequenced tray detritus samples. The absence of these species in the EAM, where detritivore grazing occurs (Hatcher, 1983), calls into question the ecological relevance of any detritus they may produce.

Detritus inputs by algae and non-sponge metazoans

Isotope mixing models revealed a significant contribution of benthic macroalgae to EAM detritus collected in 2018, and genetic analyses revealed a high frequency of sequences from major plant phyla. These findings are congruent with reports that benthic and filamentous algae are important contributors to coral reef detritus (Wilson, Burns & Codi, 2001). The benthic cover of *Dictyota* experienced historical increases at Carrie Bow Cay following die-offs of *Diadema antillarum* in the 1980s and persistent overfishing, but a ban on the harvest of herbivorous fishes in Belize has led to the recovery of herbivore fish populations in recent years (Cox et al., 2013; Mumby & Steneck, 2018; De Pablo et al., 2021). Herbivores not only benefit the reef by minimizing algae cover, but their ceaseless grazing and frequent defecations act as a pathway to recycle algae tissue to the benthos as detritus (Wernberg et al., 2006), and fish egestion of organic matter contain more nutrients than their excretion of inorganic matter (Schiettekatte et al., 2023). A similar function is likely performed by spongivore grazers. Indeed, the feces of prominent spongivore species (*Holacanthus ciliaris* and *Pomacanthus paru*, Randall & Hartman, 1968) matched isotopic values of sponge tissues and contained Porifera sequences, validating the use of eDNA methods in attempting to find the sponge signature in reef detritus. However, spongivore feces were isotopically and genetically distinct from EAM detritus, suggesting minimal detritus contributions through the spongivore grazing pathway.

Pelagic sources were well represented in EAM detritus, consistent with the pelagic signature found in detritus collected from the forereef of Palmyra Atoll (Max et al., 2013). All EAM detritus samples contained Bacillariales and Naviculales, orders of diatoms (phylum Bacillariophyceae) known for being among the most abundant microalgal components of detritus, accounting for up to 14% of detrital organic material, and likely contributing important dietary resources such as fatty acids and proteins (reviewed in Wilson et al., 2003). Many EAM detritus samples also contained crustacean signatures at lower overall frequencies, which may represent zooplankton or their feces that have arrived at the EAM through sinking or grazing by the “wall of mouths” (Hamner et al., 1988). Meiofaunal and epifaunal sequences were also abundant in the detritus, likely from living inhabitants of the EAM that are not part of the detritus but may consume it. Sequences matching to copepod order Harpacticoida were notably abundant in EAM detritus, consistent with past observations that harpacticoid copepod nauplii were the most numerically dominant members of the zooplankton on the Carrie Bow Cay forereef (Fornshell, 1994). The xenacoelomorph order Acoela was also found in 73% of detritus samples, and a diversity of these meiofaunal flatworms are known to inhabit the shallow sediments of Carrie Bow Cay (Achatz, Hooge & Tyler, 2007; Hooge & Tyler, 2008).

Nematoda were also abundant, but are unlikely to interfere or compete with detritivore grazers (Leduc & Probert, 2009).

The most common macrofaunal metazoans in EAM detritus were Mollusca (55% frequency) and Cnidaria (45% frequency). Identified molluscs included cephalaspidean gastropods in the family Haminoeidae, which may have been a non-selective detritivore grazer akin to other species in this family (Malaquias et al., 2004). Scleractinia (hard corals) were largely absent from EAM detritus. This was unexpected considering longstanding assertions about the importance of coral mucus to reef detritus (Gottfried & Roman, 1983), but missing identifications may also be due to the lack of cellular material containing the targeted genetic sequences in coral mucus. Interestingly, most cnidarian identifications in detritus belonged to octocorals (Order Alcyonacea). Gorgonian octocorals are abundant on Carrie Bow Cay but have received relatively less research attention (Kupfner Johnson & Hallock, 2020). No reports or studies could be found addressing detrital contributions by octocorals, but further study is merited considering the findings herein of abundance of octocoral sequences in detritus.

Unusually negative $\delta^{15}\text{N}$ in water column particulates collected in sediment traps

The purpose of sediment traps was to capture the particulates sinking from the water column and distinguish them from particulates originating on the benthos. Lower $\delta^{13}\text{C}$ values of sediment trap particulates compared to benthic detritus were consistent with previous findings (Max et al., 2013) and in agreement with the pattern of generally lower $\delta^{13}\text{C}$ in phytoplankton compared to benthic autotrophs (Van Duy, Mueller & Meesters, 2018). Samples from the Palmyra atoll fore reef exhibited comparable $\delta^{15}\text{N}$ values between sediment trap and EAM detritus, indicating a similar pelagic origin and highlighting the importance of pelagic inputs to fore reef detritus (Max et al., 2013). In contrast, the isotopic signatures of Carrie Bow Cay fore reef detritus suggest a combination of pelagic and benthic inputs, as $\delta^{15}\text{N}$ of sediment trap particulates was highly reduced compared to EAM detritus, even taking on an unusually negative value in 2018. The observation of lower $\delta^{15}\text{N}$ in sediment traps (pelagic particulates) compared to EAM detritus (benthic particulates) contradicts other reports of elevated $\delta^{15}\text{N}$ in pelagic compared to benthic organic matter (Van Duy, Mueller & Meesters, 2018). This trend reversal may be due to unique circumstances driving negative $\delta^{15}\text{N}$ in waters around Carrie Bow Cay and reflected in the sediment trap samples in 2018, and two possible explanations for negative $\delta^{15}\text{N}$ are presented below.

First, uncharacteristically negative $\delta^{15}\text{N}$ values for detritus in sediment traps from July 2018 samples may be a result of grazing by zooplankton on the diazotrophic cyanobacterium *Trichodesmium*. Similarly negative $\delta^{15}\text{N}$ values (−1 to −2‰) of zooplankton collected from oligotrophic waters were attributed to nitrogen fixation by *Trichodesmium* that are consumed by zooplankton (Montoya, Carpenter & Capone, 2002). *Trichodesmium* is abundant during the summer at Carrie Bow Cay, and the fore reef may act as a sink of *Trichodesmium* (Villareal, 1995). We identified *Trichodesmium* in two EAM detritus samples from 2018 (40% frequency of occurrence) and zero sediment trap samples;

Trichodesmium colonies were likely either retained in the 150 μm prefilter or grazed, for example by cyanobacteria-grazing mixotrophic dinoflagellates such as *Alexandrium* (Jeong *et al.*, 2010). *Alexandrium* was found in 80% of 2018 sediment traps but missing from all other composite samples. This dinoflagellate could have contributed to uniquely negative $\delta^{15}\text{N}$ values of organic matter if it arrived in large quantities after grazing on *Trichodesmium*. We were unable to measure the relative abundance of *Alexandrium*, but this explanation is consistent with reports of sharply declining abundances of *Trichodesmium* across the Carrie Bow Cay fore reef for which zooplankton grazing was implicated as a likely cause (Villareal, 1995).

A second explanation could be that nitrogen in the atmosphere and rain is notably depleted in ^{15}N around Carrie Bow Cay, and the $\delta^{15}\text{N}$ values of nearby nutrient-starved mangroves can be as low as -17 (Fogel *et al.*, 2008). Currents may carry particulates from the ^{15}N -depleted mangroves from the lagoon, through the tidal cut, and onto the reef. However, July is marked by northeasterly trade winds and a net onshore current flow (Greer & Kjerfve, 1982; Koltos & Opishinski, 2009) that flushes the fore reef with more oceanic than lagoonal seawater. Irrespective of the origin of ^{15}N -depleted particles in sediment traps, whether open ocean or lagoon, the isotopic and genetic distinctions between sediment trap particulates and EAM detritus indicate separate origins and composition of the particulates in July 2018.

Seasonal differences in sources and composition of detritus

There were considerable seasonal shifts in the isotopic signatures of several detritus sources between July 2018 and March 2019. Tissue samples of the HMA sponge species *Aplysina cauliformis* and *Chondrilla* sp. exhibited a rise in $\delta^{13}\text{C}$, perhaps because higher irradiance and water clarity led to more CO_2 fixation by the sponge photosymbionts and consequently lower $\delta^{13}\text{C}$ of these sponge species in July 2018, compared to March 2019 when reduced irradiance may have favored heterotrophy. The $\delta^{13}\text{C}$ shifts in tissues of other sponge species reflected their different feeding strategies and dietary preferences. The overlap in $\delta^{13}\text{C}$ between EAM detritus and the tissues of *Niphates digitalis* is consistent with the detritus-dominated diet of this species (Freeman *et al.*, 2021). The DOC-dominated diets of *X. muta* and *H. caerulea* (De Goeij *et al.*, 2008a; McMurray *et al.*, 2018) may also explain the seasonal rise in $\delta^{13}\text{C}$ values in their tissues, for example if there was a shift from pelagic to benthic sources of DOC; *X. muta* consumes DOC of varying pelagic and benthic origins, depending on environmental availability (Van Duyl, Mueller & Meesters, 2018).

There was a collective decline in the $\delta^{15}\text{N}$ values of tissues of all six sponge species in March 2019, suggesting widespread changes to nitrogen taken up by sponges from their environment. The sponge microbiome can also influence $\delta^{15}\text{N}$ in complex ways (Van Duyl, Mueller & Meesters, 2018) and may explain why some species showed larger $\delta^{15}\text{N}$ reductions than others. For example, the connection between nitrogen fixation and reduced $\delta^{15}\text{N}$ (Southwell *et al.*, 2008) may explain larger $\delta^{15}\text{N}$ reductions in *A. cauliformis*, which relies on symbionts for nutrition, compared to the smaller $\delta^{15}\text{N}$ reductions in *N. digitalis*, which predominantly depends on external nitrogen sources (Weisz *et al.*, 2007).

Samples of tissues from benthic cyanobacterial mats (BCM) exhibited an unexpected drop in $\delta^{13}\text{C}$ values. This is the first known report of such a seasonal shift for BCM, and it may be due to seasonal variation in dissolved inorganic carbon (DIC) or BCM species composition. Sequencing was only performed on BCM samples from July 2018, precluding seasonal comparison of the species known to occur in the BCM consortia (e.g., *Moorea*, *Oscillatoria*, *Hydrocoleum*).

Although isotopic and genetic signatures of detritus sources varied from July 2018 to March 2019, isotopic signatures of the detritus itself remained relatively consistent between the seasons. EAM detritus contained an abundance of sequences matching to primary producers whose mechanical breakdown and predation can directly contribute to detritus (Wilson *et al.*, 2003). The abundant dissolved organic matter (DOM) produced by benthic algae and other autotrophs may also spontaneously form polymer gels or adsorb to carbonate sediments and become a form of amorphic detritus (Otsuki & Wetzel, 1973; Chin, Orellana & Verdugo, 1998; Wilson *et al.*, 2003). Amorphic detritus is a non-cellular composite undetectable by genetic sequencing methods but with an isotopic signature reflective of its source organism, and significant inputs of this form of detritus may explain similar isotopic signatures despite distinct genetic sequences in detritus from both seasons. Amorphic detritus is a form of molecularly uncharacterized materials, byproducts of complex biogeochemical processes that escape characterization by traditional chromatography-based techniques and are poorly understood across all aquatic ecosystems, especially coral reefs (Bowen, 1979; Wilson, 2002; Wilson *et al.*, 2003; Wakeham & Lee, 2019).

Multiple lines of evidence indicate that detritus composition was shaped by environmental conditions on the forereef, namely higher flow and greater turbulence in March 2019 compared to July 2018. First, detritus collected in March 2019 had more variable isotopic delta values and fewer genetic sequences compared to detritus collected in July 2018, indicating a more heterogeneous and scarce mixture that may result from greater turbulence and mixing. Detritus collected in March 2019 was also characterized by greater frequency of meiofauna and epifauna that likely inhabit the EAM, including Annelida, Nematoda, and Xenacoelomorpha, and lower frequencies of macrofauna that originated elsewhere on the benthos, pointing to surge-driven export of macrofaunal-derived organic matter before it could settle and get trapped in the EAM in March 2019. Stronger surge conditions may have also caused the resuspension of benthic particulates in the water column, which would account for the presence of sequences belonging to benthic metazoans (e.g., Porifera, Mollusca, Cnidaria) in sediment trap samples in March 2019. Sediment traps are effective water column samplers when conditions are calm (flow < 0.2 m/s), as was the case during July 2018 sampling, but sediment traps may capture resuspended benthic materials when conditions are turbulent as was the case during March 2019 sampling (Gardner, 1980; White, 1990; Wilson *et al.*, 2003). The more energetic environment in March 2019 may have also amplified production of amorphic detritus in the EAM, as DOM aggregation may be facilitated by more encounters with substrates and increased laminar shear and bubbles formed in the higher water motion (Wilson *et al.*, 2003).

Ecological implications

Organic matter cycling through the detrital food web is complicated by the number of organisms that may act as both sources and sinks of detritus, including bacteria, sponges, surgeonfishes, and parrotfishes (Choat, Clements & Robbins, 2002; Wilson et al., 2003; Crossman, Choat & Clements, 2005; Dromard et al., 2015; Mumby & Steneck, 2018). It is important to understand how detritus is processed and transformed by these groups. Certain detritivores can selectively reduce protein levels on the reef, resulting in higher carbon-to-nitrogen ratios and decreased nutritional value in nitrogen-limited systems, as observed in the backreef habitats of Palmyra Atoll in the Pacific Ocean (Max et al., 2013). Detritus is likely more abundant on Caribbean coral reefs (Mumby & Steneck, 2018), and there may be a surplus of nutritious detritus on the Carrie Bow Cay forereef due to the windward exposure of the reef and other favorable conditions driving import of particulate matter onto the reef.

The higher productivity in the Summer (July 2018) and potential influx of sinking phyto- and zooplankton may have supplemented benthic sources of detritus, while the more energetic early spring (March 2019) resulted in greater mixing and flushing and perhaps better conditions for formation and accumulation of amorphic detritus in the EAM. Seasonal conditions may bring a variety and surplus of detritus that is regularly replenished to the forereef and may even supply downstream backreef and lagoonal habitats (Hatcher, 1983). Despite an abundance of detritus, there may be an imbalance in the number of detritivores feeding on it, as detritivores are less diverse and abundant in the Caribbean compared to Pacific reefs (Bellwood et al., 2004; Roff & Mumby, 2012; Edwards et al., 2014). A shortage of detritivores in the Caribbean may leave this overflow detritus to get recycled into the microbial loop, returned to benthic suspension feeders (e.g., sponges), or lost from the reef system (Mumby & Steneck, 2018).

It may be argued that this “sink side” test of the sponge loop is based on a single coral reef location, but as indicated above, the location was chosen to maximize the likelihood of confirming the hypothesis, with among the highest sponge abundances reported for Caribbean coral reefs (Wilkinson & Cheshire, 1990; Loh & Pawlik, 2014) and the deployment of trays directly under reef overhangs that supported cryptic sponge species. Despite the foregoing, the sponge signature in reef detritus was negligible. The breadth of sampling conducted in the present study (16 tissue and composite sample types collected over two seasons) encompassed many common organisms and the most likely candidates for detritus production, although this sample breadth limited the number of replicates from each group that could feasibly be analyzed. The stable isotope analyses and eDNA metabarcoding analysis methods used herein each had limitations that the other method accounted for to some degree, but some gaps remained. Isotope mixing models can only estimate contributions of sampled sources, so while this was an expansive sampling effort, it was not exhaustive, and detritus isotopic signatures may have been influenced by diagenetic changes, especially of $\delta^{15}\text{N}$ values (Caraco et al., 1998). Genetic sequencing complemented the isotopic analyses by identifying all traces of organisms that are cataloged and with genetic material that resisted degradation, but some organisms could have escaped detection if they were novel or degraded (Reef et al., 2017). The omission of bacteria besides

cyanobacteria in the genetic sequence data also discounted a key component of detritus ([Biddanda, 1985](#); [Alongi, 1988](#); [Wilson et al., 2003](#)), but the bacterial signal would have been present in the isotopic signatures of detritus.

CONCLUSION

To better understand how reefs of the future will function, it is crucial to understand how energy is transferred throughout contemporary coral reefs, including the complex and poorly-elucidated detrital pathway. While sponges may not be major contributors through direct production of detritus, they may extend the residence time of organic matter on coral reefs through efficient uptake of DOC and accumulation of biomass that is eventually grazed or mechanically broken down and returned to the detrital pool. This carbon pathway may not be essential to the healthy functioning of Caribbean coral reefs, however, considering low demand by relatively few detritivorous grazers. As benthic communities continue to shift from coral- to algae- or sponge-dominated systems, this will alter trophodynamics and influence the detrital resources on which these systems rely.

ACKNOWLEDGEMENTS

Many thanks to Laura Gaitan Daza, Steve McMurray, Jim Evans, Mellissa Dionesotes, and Sasha Giametti for assistance in the field, the staff of Smithsonian Institution's Carrie Bow Cay Field Station in Belize for logistical support, Evan Heit and Kim Rosov for assistance with preparation and analyses of stable isotope samples, and Nicole Fogarty and Amy Grogan for additional assistance with reviewing and providing input on earlier drafts of this manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was funded by the National Science Foundation (OCE 2218863), the Atlantic Canada Opportunities Agency's Atlantic Innovation Fund (project number 781-37749-207993), and a grant from Energy Research and Innovation Newfoundland and Labrador (ERINL). Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of ERINL or its members. There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

National Science Foundation: OCE 2218863.

Atlantic Canada Opportunities Agency's Atlantic Innovation Fund: project number 781-37749-207993.

Energy Research and Innovation Newfoundland and Labrador (ERINL).

Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of ERINL or its members.

Competing Interests

Beverly McClenaghan, Lesley Berghuis, Nicole Fahner and Hoda Rajabi are employees of eDNAtec Inc. Mehrdad Hajibabaei is the founder and Chief Scientific Officer of eDNAtec Inc.

Author Contributions

- Lauren K. Olinger conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Beverly McClenaghan analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Mehrdad Hajibabaei analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Nicole Fahner analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Lesley Berghuis analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Hoda Rajabi analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Patrick Erwin conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Chad S. Lane conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Joseph R. Pawlik conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the article, and approved the final draft.

Field Study Permissions

The following information was supplied relating to field study approvals (*i.e.*, approving body and any reference numbers):

Field experiments were approved by the Belize Fisheries Department (permit number 00023-18).

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The DNA sequence data generated during and/or analyzed during the current study are available at GenBank: [PRJNA965826](#).

Data Availability

The following information was supplied regarding data availability:

The raw data is available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.16970#supplemental-information>.

REFERENCES

- Achatz JG, Hooge MD, Tyler S. 2007. Convolutidae (Acoela) from Belize. *Zootaxa* **1479**:35–66.
- Alexander BE, Liebrand K, Osinga R, HG van der Geest, Admiraal W, Cleutjens JPM, Schutte B, Verheyen F, Ribes M, E van Loon, JM de Goeij. 2014. Cell turnover and detritus production in marine sponges from tropical and temperate benthic ecosystems. *PLOS ONE* **9**:e109486 DOI [10.1371/journal.pone.0109486](https://doi.org/10.1371/journal.pone.0109486).
- Alongi DM. 1988. Detritus in coral reef ecosystems: fluxes and fates. In: *Proceedings of the sixth international coral reef symposium*. 29–36.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* **215**:403–410 DOI [10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).
- Aronson RB, Precht WF, Toscano MA, Koltes KH. 2002. The 1998 bleaching event and its aftermath on a coral reef in Belize. *Marine Biology* **141**:435–447 DOI [10.1007/s00227-002-0842-5](https://doi.org/10.1007/s00227-002-0842-5).
- Azam F, Fenchel T, Field J, Gray J, Meyer-Reil L, Thingstad F. 1983. The ecological role of water-column microbes in the sea. *Marine Ecology Progress Series* **10**:257–263 DOI [10.3354/meps010257](https://doi.org/10.3354/meps010257).
- Baird DJ, Hajibabaei M. 2012. Biomonitoring 2.0: a new paradigm in ecosystem assessment made possible by next-generation DNA sequencing. *Molecular Ecology* **21**:2039–2044 DOI [10.1111/j.1365-294X.2012.05519.x](https://doi.org/10.1111/j.1365-294X.2012.05519.x).
- Begon M, Harper JL, Townsend CR. 1986. *Ecology: individuals, populations and communities*. Oxford: Blackwell Scientific Publications.
- Bellwood DR, Hughes TP, Folke C, Nyström M. 2004. Confronting the coral reef crisis. *Nature* **429**:827–833 DOI [10.1038/nature02691](https://doi.org/10.1038/nature02691).
- Biddanda B. 1985. Microbial synthesis of macroparticulate matter. *Marine Ecology Progress Series* **20**:241–251 DOI [10.3354/meps020241](https://doi.org/10.3354/meps020241).
- Bowen SH. 1979. A nutritional constraint in detritivory by fishes: the stunted population of *Sarotherodon mossambicus* in Lake Sibaya, South Africa. *Ecological Monographs* **49**:17–31 DOI [10.2307/1942570](https://doi.org/10.2307/1942570).
- Boyi JO, Heße E, Rohner S, Säurich J, Siebert U, Gilles A, Lehnert K. 2022. Deciphering Eurasian otter (*Lutra lutra* L.) and seal (*Phoca vitulina* L.; *Halichoerus grypus* F.) diet: metabarcoding tailored for fresh and saltwater fish species. *Molecular Ecology* **31**:5089–5106 DOI [10.1111/mec.16635](https://doi.org/10.1111/mec.16635).
- Brocke HJ, Wenzhoefer F, De Beer D, Mueller B, Van Duyl FC, Nugues MM. 2015. High dissolved organic carbon release by benthic cyanobacterial mats in a Caribbean reef ecosystem. *Scientific Reports* **5**:8852 DOI [10.1038/srep08852](https://doi.org/10.1038/srep08852).

- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016.** DADA2: high-resolution sample inference from Illumina amplicon data. *Nature Methods* 13:581–583 DOI [10.1038/nmeth.3869](https://doi.org/10.1038/nmeth.3869).
- Campana S, Hudspith M, Lankes D, De Kluijver A, Demey C, Schoorl J, Absalah S, Van der Meer MTJ, Mueller B, De Goeij JM. 2021.** Processing of naturally sourced macroalgal- and coral-dissolved organic matter (DOM) by high and low microbial abundance encrusting sponges. *Frontiers in Marine Science* 8:640583 DOI [10.3389/fmars.2021.640583](https://doi.org/10.3389/fmars.2021.640583).
- Caraco N, Lampman G, Cole J, Limburg K, Pace M, Fischer D. 1998.** Microbial assimilation of DIN in a nitrogen rich estuary: implications for food quality and isotope studies. *Marine Ecology Progress Series* 167:59–71 DOI [10.3354/meps167059](https://doi.org/10.3354/meps167059).
- Chin W-C, Orellana MV, Verdugo P. 1998.** Spontaneous assembly of marine dissolved organic matter into polymer gels. *Nature* 391:568–572 DOI [10.1038/35345](https://doi.org/10.1038/35345).
- Choat J, Clements K, Robbins W. 2002.** The trophic status of herbivorous fishes on coral reefs. *Marine Biology* 140:613–623 DOI [10.1007/s00227-001-0715-3](https://doi.org/10.1007/s00227-001-0715-3).
- Cox CE, Jones CD, Wares JP, Castillo KD, McField MD, Bruno JF. 2013.** Genetic testing reveals some mislabeling but general compliance with a ban on herbivorous fish harvesting in Belize: genetic testing on coral reef fish in Belize. *Conservation Letters* 6:132–140 DOI [10.1111/j.1755-263X.2012.00286.x](https://doi.org/10.1111/j.1755-263X.2012.00286.x).
- Crossman D, Choat J, Clements K. 2005.** Nutritional ecology of nominally herbivorous fish on coral reefs. *Marine Ecology Progress Series* 296:129–142 DOI [10.3354/meps296129](https://doi.org/10.3354/meps296129).
- Crossman DJ, Choat HJ, Clements KD, Hardy T, McConochie J. 2001.** Detritus as food for grazing fishes on coral reefs. *Limnology and Oceanography* 46:1596–1605 DOI [10.4319/lo.2001.46.7.1596](https://doi.org/10.4319/lo.2001.46.7.1596).
- Darwin C, Bonney TG. 1896.** *The structure and distribution of coral reefs*. New York: D. Appleton and Company.
- De Bakker DM, Van Duyl FC, Bak RPM, Nugues MM, Nieuwland G, Meesters EH. 2017.** 40 Years of benthic community change on the Caribbean reefs of Curaçao and Bonaire: the rise of slimy cyanobacterial mats. *Coral Reefs* 36:355–367 DOI [10.1007/s00338-016-1534-9](https://doi.org/10.1007/s00338-016-1534-9).
- De Goeij JM, De Kluijver A, Van Duyl FC, Vacelet J, Wijffels RH, De Goeij AFPM, Cleutjens JPM, Schutte B. 2009.** Cell kinetics of the marine sponge *Halisarca caerulea* reveal rapid cell turnover and shedding. *Journal of Experimental Biology* 212:3892–3900 DOI [10.1242/jeb.034561](https://doi.org/10.1242/jeb.034561).
- De Goeij JM, Moodley L, Houtekamer M, Carballeira NM, Van Duyl FC. 2008b.** Tracing ¹³C-enriched dissolved and particulate organic carbon in the bacteria-containing coral reef sponge *Halisarca caerulea*: evidence for DOM-feeding. *Limnology and Oceanography* 53:1376–1386 DOI [10.4319/lo.2008.53.4.1376](https://doi.org/10.4319/lo.2008.53.4.1376).
- De Goeij JM, Van den Berg H, Van Oostveen M, Epping E, Van Duyl F. 2008a.** Major bulk dissolved organic carbon (DOC) removal by encrusting coral reef cavity sponges. *Marine Ecology Progress Series* 357:139–151 DOI [10.3354/meps07403](https://doi.org/10.3354/meps07403).

- De Goeij JM, Van Oevelen D, Vermeij MJA, Osinga R, Middelburg JJ, De Goeij AFPM, Admiraal W.** 2013. Surviving in a marine desert: the sponge loop retains resources within coral reefs. *Science* **342**:108–110 DOI [10.1126/science.1241981](https://doi.org/10.1126/science.1241981).
- De Pablo LX, Lefcheck JS, Harper L, Paul VJ, Jones S, Whippo R, Seemann J, Kline DI, Duffy JE.** 2021. A doubling of stony coral cover on shallow reefs at Carrie Bow Cay, Belize from 2014 to 2019. *Scientific Reports* **11**:19185 DOI [10.1038/s41598-021-96799-2](https://doi.org/10.1038/s41598-021-96799-2).
- Done TJ.** 1992. Phase shifts in coral reef communities and their ecological significance. *Hydrobiologia* **247**:121–132 DOI [10.1007/BF00008211](https://doi.org/10.1007/BF00008211).
- Dromard CR, Bouchon-Navaro Y, Harmelin-Vivien M, Bouchon C.** 2015. Diversity of trophic niches among herbivorous fishes on a Caribbean reef (Guadeloupe, Lesser Antilles), evidenced by stable isotope and gut content analyses. *Journal of Sea Research* **95**:124–131 DOI [10.1016/j.seares.2014.07.014](https://doi.org/10.1016/j.seares.2014.07.014).
- Edwards CB, Friedlander AM, Green AG, Hardt MJ, Sala E, Sweatman HP, Williams ID, Zgliczynski B, Sandin SA, Smith JE.** 2014. Global assessment of the status of coral reef herbivorous fishes: evidence for fishing effects. *Proceedings of the Royal Society B* **281**:20131835 DOI [10.1098/rspb.2013.1835](https://doi.org/10.1098/rspb.2013.1835).
- Fabricius KE.** 2005. Effects of terrestrial runoff on the ecology of corals and coral reefs: review and synthesis. *Marine Pollution Bulletin* **50**:125–146 DOI [10.1016/j.marpolbul.2004.11.028](https://doi.org/10.1016/j.marpolbul.2004.11.028).
- Fenchel T.** 2008. The microbial loop—25 years later. *Journal of Experimental Marine Biology and Ecology* **366**:99–103 DOI [10.1016/j.jembe.2008.07.013](https://doi.org/10.1016/j.jembe.2008.07.013).
- Fogel ML, Wooller MJ, Cheeseman J, Smallwood BJ, Roberts Q, Romero I, Meyers MJ.** 2008. Unusually negative nitrogen isotopic compositions ($\delta^{15}\text{N}$) of mangroves and lichens in an oligotrophic, microbially-influenced ecosystem. *Biogeosciences* **5**:1693–1704 DOI [10.5194/bg-5-1693-2008](https://doi.org/10.5194/bg-5-1693-2008).
- Fornshell JA.** 1994. Copepod nauplii from the barrier reef of Belize. *Hydrobiologia* **292**:295–301.
- Freeman CJ, Easson CG, Fiore CL, Thacker RW.** 2021. Sponge–microbe interactions on coral reefs: multiple evolutionary solutions to a complex environment. *Frontiers in Marine Science* **8**:705053 DOI [10.3389/fmars.2021.705053](https://doi.org/10.3389/fmars.2021.705053).
- Gardner W.** 1980. Sediment trap dynamics and calibration: a laboratory evaluation. *Journal of Marine Research* **38**:17–39.
- Gottfried M, Roman MR.** 1983. Ingestion and incorporation of coral-mucus detritus by reef zooplankton. *Marine Biology* **72**:211–218 DOI [10.1007/BF00396825](https://doi.org/10.1007/BF00396825).
- Greenwood NDW, Sweeting CJ, Polunin NVC.** 2010. Elucidating the trophodynamics of four coral reef fishes of the Solomon Islands using $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$. *Coral Reefs* **29**:785–792 DOI [10.1007/s00338-010-0626-1](https://doi.org/10.1007/s00338-010-0626-1).
- Greer JE, Kjerfve B.** 1982. Water currents adjacent to Carrie Bow Cay, Belize. *Smithsonian Contributions to the Marine Sciences* **12**:53–58.
- Hamner WM, Jones MS, Carleton JH, Hauri IR, Williams DM.** 1988. Zooplankton, planktivorous fish, and water currents on a windward reef face: great Barrier Reef, Australia. *Bulletin of Marine Science* **42**:459–479.

- Hatcher BG.** 1983. Role of detritus in the metabolism and secondary production of coral reef ecosystems. In: Baker JT, Carter RM, Sammarco PW, Stark KP, eds. *Proceedings of the inaugural great barrier reef conference, August 28-September 2, 1983*. Townsville: James Cook University, 317–325.
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A, Hatziolos ME.** 2007. Coral reefs under rapid climate change and ocean acidification. *Science* **318**:1737–1742 DOI [10.1126/science.1152509](https://doi.org/10.1126/science.1152509).
- Hooge MD, Tyler S.** 2008. Acoela (Acoelomorpha) from Bocas del Toro, Panama. *Zootaxa* **1719**:1–40.
- Hudspith M, Rix L, Achlatis M, Bougoure J, Guagliardo P, Clode PL, Webster NS, Muyzer G, Pernice M, De Goeij JM.** 2021. Subcellular view of host–microbiome nutrient exchange in sponges: insights into the ecological success of an early metazoan–microbe symbiosis. *Microbiome* **9**:44 DOI [10.1186/s40168-020-00984-w](https://doi.org/10.1186/s40168-020-00984-w).
- Hughes TP.** 1994. Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. *Science* **265**:1547–1551 DOI [10.1126/science.265.5178.1547](https://doi.org/10.1126/science.265.5178.1547).
- Jeong HJ, Yoo YD, Kim JS, Seong KA, Kang NS, Kim TH.** 2010. Growth, feeding and ecological roles of the mixotrophic and heterotrophic dinoflagellates in marine planktonic food webs. *Ocean Science Journal* **45**:65–91 DOI [10.1007/s12601-010-0007-2](https://doi.org/10.1007/s12601-010-0007-2).
- Johnson C, Klumpp D, Field J, Bradbury R.** 1995. Carbon flux on coral reefs: effects of large shifts in community structure. *Marine Ecology Progress Series* **126**:123–143 DOI [10.3354/meps126123](https://doi.org/10.3354/meps126123).
- Koltes KH, Opishinski TB.** 2009. Patterns of water quality and movement in the vicinity of Carrie Bow Cay, Belize. *Smithsonian Contributions to the Marine Sciences* **38**:379–390.
- Kornder NA, Esser Y, Stoupin D, Leys SP, Mueller B, Vermeij MJA, Huisman J, Goeij JM de.** 2022. Sponges sneeze mucus to shed particulate waste from their seawater inlet pores. *Current Biology* **32**:3855–3861 DOI [10.1016/j.cub.2022.07.017](https://doi.org/10.1016/j.cub.2022.07.017).
- Kramer MJ, Bellwood DR, Bellwood O.** 2012. Cryptofauna of the epilithic algal matrix on an inshore coral reef, Great Barrier Reef. *Coral Reefs* **31**:1007–1015 DOI [10.1007/s00338-012-0924-x](https://doi.org/10.1007/s00338-012-0924-x).
- Kupfner Johnson S, Hallock P.** 2020. A review of symbiotic gorgonian research in the western Atlantic and Caribbean with recommendations for future work. *Coral Reefs* **39**:239–258 DOI [10.1007/s00338-020-01891-0](https://doi.org/10.1007/s00338-020-01891-0).
- Leduc D, Probert PK.** 2009. The effect of bacterivorous nematodes on detritus incorporation by macrofaunal detritivores: a study using stable isotope and fatty acid analyses. *Journal of Experimental Marine Biology and Ecology* **371**:130–139 DOI [10.1016/j.jembe.2009.01.011](https://doi.org/10.1016/j.jembe.2009.01.011).
- Lesser MP, Mueller B, Pankey MS, Macartney KJ, Slattery M, De Goeij JM.** 2020. Depth-dependent detritus production in the sponge, *Halisarca caerulea*. *Limnology and Oceanography* **65**:1200–1216 DOI [10.1002/limo.11384](https://doi.org/10.1002/limo.11384).

- Levinton J.** 2021. *Marine biology*. Sixth Edition. Oxford: Oxford Univ. Press.
- Loh T-L, Pawlik JR.** 2014. Chemical defenses and resource trade-offs structure sponge communities on Caribbean coral reefs. *Proceedings of the National Academy of Sciences of the United States of America* **111**:4151–4156.
- Malaquias MAE, Condinho S, Cervera JL, Sprung M.** 2004. Diet and feeding biology of *Haminoea orbygniana* (Mollusca: Gastropoda: Cephalaspidea). *Journal of the Marine Biological Association of the United Kingdom* **84**:767–772 DOI [10.1017/S0025315404009890h](https://doi.org/10.1017/S0025315404009890).
- Martin M.** 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet Journal* **17**:10–12.
- Max L, Hamilton S, Gaines S, Warner R.** 2013. Benthic processes and overlying fish assemblages drive the composition of benthic detritus on a central Pacific coral reef. *Marine Ecology Progress Series* **482**:181–195 DOI [10.3354/meps10259](https://doi.org/10.3354/meps10259).
- McClanahan B, Fahner N, Cote D, Chawarski J, McCarthy A, Rajabi H, Singer G, Hajiabaei M.** 2020. Harnessing the power of eDNA metabarcoding for the detection of deep-sea fishes. *PLOS ONE* **15**:e0236540 DOI [10.1371/journal.pone.0236540](https://doi.org/10.1371/journal.pone.0236540).
- McMurray SE, Stubler AD, Erwin PM, Finelli CM, Pawlik JR.** 2018. A test of the sponge-loop hypothesis for emergent Caribbean reef sponges. *Marine Ecology Progress Series* **588**:1–14 DOI [10.3354/meps12466](https://doi.org/10.3354/meps12466).
- Michener RH, Kaufman L.** 2007. Stable isotope ratios as tracers in marine food webs: an update. In: Michener RH, Lajtha K, eds. *Stable isotopes in ecology and environmental science*. Malden: Blackwell Publishing Ltd, 238–282.
- Middelburg JJ.** 2014. Stable isotopes dissect aquatic food webs from the top to the bottom. *Biogeosciences* **11**:2357–2371 DOI [10.5194/bg-11-2357-2014](https://doi.org/10.5194/bg-11-2357-2014).
- Montoya JP, Carpenter EJ, Capone DG.** 2002. Nitrogen fixation and nitrogen isotope abundances in zooplankton of the oligotrophic North Atlantic. *Limnology and Oceanography* **47**:1617–1628 DOI [10.4319/lo.2002.47.6.1617](https://doi.org/10.4319/lo.2002.47.6.1617).
- Mumby PJ, Steneck RS.** 2018. Paradigm lost: dynamic nutrients and missing detritus on coral reefs. *BioScience* **68**:487–495 DOI [10.1093/biosci/biy055](https://doi.org/10.1093/biosci/biy055).
- Norström AV, Nyström M, Lokrantz J, Folke C.** 2009. Alternative states on coral reefs: beyond coral-macroalgal phase shifts. *Marine Ecology Progress Series* **376**:293–306.
- Otsuki A, Wetzel RG.** 1973. Interaction of yellow organic acids with calcium carbonate in freshwater. *Limnology and Oceanography* **18**:490–493 DOI [10.4319/lo.1973.18.3.0490](https://doi.org/10.4319/lo.1973.18.3.0490).
- Parnell A.** 2019. simmr: a stable isotope mixing model. R package version 0.5.0.9000. Available at <https://cran.r-project.org/web/packages/simmr/index.html>.
- Pawlik JR, McMurray SE.** 2020. The emerging ecological and biogeochemical importance of sponges on coral reefs. *Annual Review of Marine Science* **12**:315–337 DOI [10.1146/annurev-marine-010419-010807](https://doi.org/10.1146/annurev-marine-010419-010807).
- Post DM.** 2002. Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology* **83**:703–718 DOI [10.1890/0012-9658\(2002\)083\[0703:USITET\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[0703:USITET]2.0.CO;2).

- Purcell S, Bellwood D.** 2001. Spatial patterns of epilithic algal and detrital resources on a windward coral reef. *Coral Reefs* **20**:117–125 DOI [10.1007/s003380100150](https://doi.org/10.1007/s003380100150).
- Randall JE, Hartman WD.** 1968. Sponge-feeding fishes of the West Indies. *Marine Biology* **1**:216–225 DOI [10.1007/BF00347115](https://doi.org/10.1007/BF00347115).
- R Core Team.** 2020. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <https://www.r-project.org>.
- Reef R, Atwood TB, Samper-Villarreal J, Adame MF, Sampayo EM, Lovelock CE.** 2017. Using eDNA to determine the source of organic carbon in seagrass meadows. *Limnology and Oceanography* **62**:1254–1265 DOI [10.1002/limo.10499](https://doi.org/10.1002/limo.10499).
- Roff G, Mumby PJ.** 2012. Global disparity in the resilience of coral reefs. *Trends in Ecology & Evolution* **27**:404–413 DOI [10.1016/j.tree.2012.04.007](https://doi.org/10.1016/j.tree.2012.04.007).
- Rützler K, Macintyre IG.** 1982. The habitat distribution and community structure of the barrier reef complex at Carrie Bow Cay, Belize. *Smithsonian Contributions to the Marine Sciences* **12**:9–45.
- Schidlowski M.** 2001. Carbon isotopes as biogeochemical recorders of life over 3.8 Ga of Earth history: evolution of a concept. *Precambrian Research* **106**:117–134 DOI [10.1016/S0301-9268\(00\)00128-5](https://doi.org/10.1016/S0301-9268(00)00128-5).
- Schiettekatte NMD, Casey JM, Brandl SJ, Mercière A, Degregori S, Burkepile D, Van Wert JC, Ghilardi M, Villéger S, Parravicini V.** 2023. The role of fish feces for nutrient cycling on coral reefs. *Oikos* **n/a**:e09914.
- Southwell MW, Weisz JB, Martens CS, Lindquist N.** 2008. *In situ* fluxes of dissolved inorganic nitrogen from the sponge community on Conch Reef, Key Largo, Florida. *Limnology and Oceanography* **53**:986–996 DOI [10.4319/lo.2008.53.3.0986](https://doi.org/10.4319/lo.2008.53.3.0986).
- Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH.** 2012. Environmental DNA. *Molecular Ecology* **21**:1789–1793 DOI [10.1111/j.1365-294X.2012.05542.x](https://doi.org/10.1111/j.1365-294X.2012.05542.x).
- Tebbett SB, Bellwood DR.** 2019. Algal turf sediments on coral reefs: what's known and what's next. *Marine Pollution Bulletin* **149**:110542 DOI [10.1016/j.marpolbul.2019.110542](https://doi.org/10.1016/j.marpolbul.2019.110542).
- Thomsen PF, Willerslev E.** 2015. Environmental DNA—an emerging tool in conservation for monitoring past and present biodiversity. *Biological Conservation* **183**:4–18 DOI [10.1016/j.biocon.2014.11.019](https://doi.org/10.1016/j.biocon.2014.11.019).
- Van Duyl FC, Mueller B, Meesters EH.** 2018. Spatio-temporal variation in stable isotope signatures ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of sponges on the Saba Bank. *PeerJ* **6**:e5460 DOI [10.7717/peerj.5460](https://doi.org/10.7717/peerj.5460).
- Villareal TA.** 1995. Abundance and photosynthetic characteristics of *Trichodesmium* spp. along the Atlantic Barrier Reef at Carrie Bow Cay, Belize. *Marine Ecology* **16**:259–271 DOI [10.1111/j.1439-0485.1995.tb00410.x](https://doi.org/10.1111/j.1439-0485.1995.tb00410.x).
- Wakeham SG, Lee C.** 2019. Limits of our knowledge, part 2: selected frontiers in marine organic biogeochemistry. *Marine Chemistry* **212**:16–46 DOI [10.1016/j.marchem.2019.02.005](https://doi.org/10.1016/j.marchem.2019.02.005).
- Weisz JB, Hentschel U, Lindquist N, Martens CS.** 2007. Linking abundance and diversity of sponge-associated microbial communities to metabolic differences in host sponges. *Marine Biology* **152**:475–483 DOI [10.1007/s00227-007-0708-y](https://doi.org/10.1007/s00227-007-0708-y).

- Wernberg T, Vanderklift MA, How J, Lavery PS.** 2006. Export of detached macroalgae from reefs to adjacent seagrass beds. *Oecologia* 147:692–701 DOI [10.1007/s00442-005-0318-7](https://doi.org/10.1007/s00442-005-0318-7).
- White J.** 1990. The use of sediment traps in high-energy environments. *Marine Geophysical Research* 12:145–152 DOI [10.1007/BF00310569](https://doi.org/10.1007/BF00310569).
- Wilkinson C, Cheshire A.** 1990. Comparisons of sponge populations across the Barrier Reefs of Australia and Belize: Evidence for higher productivity in the Caribbean. *Marine Ecology Progress Series* 67:285–294 DOI [10.3354/meps067285](https://doi.org/10.3354/meps067285).
- Wilson SK.** 2000. Trophic status and feeding selectivity of blennies (Blenniidae: Salariini). *Marine Biology* 136:431–437 DOI [10.1007/s002270050702](https://doi.org/10.1007/s002270050702).
- Wilson S.** 2002. Nutritional value of detritus and algae in blenny territories on the Great Barrier Reef. *Journal of Experimental Marine Biology and Ecology* 271:155–169 DOI [10.1016/S0022-0981\(02\)00035-7](https://doi.org/10.1016/S0022-0981(02)00035-7).
- Wilson S, Bellwood D, Choat J, Furnas M.** 2003. Detritus in the epilithic algal matrix and its use by coral reef fishes. *Oceanography and Marine Biology: An Annual Review* 41:279–309.
- Wilson S, Burns K, Codi S.** 2001. Identifying sources of organic matter in sediments from a detritivorous coral reef fish territory. *Organic Geochemistry* 32:1257–1269 DOI [10.1016/S0146-6380\(01\)00084-5](https://doi.org/10.1016/S0146-6380(01)00084-5).
- Yoccoz NG, Bråthen KA, Gielly L, Haile J, Edwards ME, Goslar T, Von Stedingk H, Brysting AK, Coissac E, Pompanon F, Sønstebo JH, Miquel C, Valentini A, De Bello F, Chave J, Thuiller W, Wincker P, Cruaud C, Gavory F, Rasmussen M, Gilbert MTP, Orlando L, Brochmann C, Willerslev E, Taberlet P.** 2012. DNA from soil mirrors plant taxonomic and growth form diversity. *Molecular Ecology* 21:3647–3655 DOI [10.1111/j.1365-294X.2012.05545.x](https://doi.org/10.1111/j.1365-294X.2012.05545.x).
- Zapata-Hernández G, Sellanes J, Letourneur Y, Harrod C, Morales NA, Plaza P, Meerhoff E, Yannicelli B, Carrasco SA, Hinojosa I, Gaymer CF.** 2021. Tracing trophic pathways through the marine ecosystem of Rapa Nui (Easter Island). *Aquatic Conservation: Marine and Freshwater Ecosystems* 31:304–323 DOI [10.1002/aqc.3500](https://doi.org/10.1002/aqc.3500).