

Microlocal Morse theory of wrapped Fukaya categories

By SHEEL GANATRA, JOHN PARDON, and VIVEK SHENDE

Abstract

The Nadler–Zaslow correspondence famously identifies the finite-dimensional Floer homology groups between Lagrangians in cotangent bundles with the finite-dimensional Hom spaces between corresponding constructible sheaves. We generalize this correspondence to incorporate the infinite-dimensional spaces of morphisms “at infinity,” given on the Floer side by Reeb trajectories (also known as “wrapping”) and on the sheaf side by allowing unbounded infinite rank sheaves which are categorically compact. When combined with existing sheaf theoretic computations, our results confirm many new instances of homological mirror symmetry.

More precisely, given a real analytic manifold M and a subanalytic isotropic subset Λ of its co-sphere bundle S^*M , we show that the partially wrapped Fukaya category of T^*M stopped at Λ is equivalent to the category of compact objects in the unbounded derived category of sheaves on M with microsupport inside Λ . By an embedding trick, we also deduce a sheaf theoretic description of the wrapped Fukaya category of any Weinstein sector admitting a stable polarization.

Contents

1. Introduction	945
1.1. Weinstein manifolds and partially wrapped Fukaya categories	945
1.2. Topological and sheaf theoretic interpretations	948
1.3. Main results	949
Convention.	953
1.4. Acknowledgements	953
2. Stratifications	953
2.1. Generalities	953
2.2. Microlocal approximation of constructible sets	956

Keywords: Fukaya category, microlocal sheaf, Stein manifold, Weinstein manifold, mirror symmetry

AMS Classification: Primary: 53D37, 14J33, 53D40.

© 2024 Department of Mathematics, Princeton University.

2.3. Cornering and conormals of constructible open sets	959
3. Microlocal Morse categories	962
3.1. Strata poset categories and refinement functors	962
3.2. A category for any Λ	963
3.3. Open inclusions	966
4. Sheaf categories	967
4.1. Categories of sheaves and functors between them	967
4.2. Constructible sheaves	969
4.3. Microsupport	971
4.4. Microstalks	972
4.5. Compact objects	975
4.6. In conclusion	978
5. Wrapped Fukaya categories	980
5.1. Wrapped Floer cohomology	980
5.2. Wrapped Fukaya categories	981
5.3. Gradings and orientations	982
5.4. Wrapping exact triangle, stop removal, generation	985
5.5. Conormals and corners	986
5.6. Floer cohomology between conormals of balls and stable balls	987
5.7. Fukaya categories of conormals to stars	991
5.8. Functors from poset categories to Fukaya categories	994
5.9. In conclusion	998
6. Examples	1001
6.1. Cotangent bundles	1001
6.2. Plumblings	1001
6.3. Proper modules and infinitesimal Fukaya categories	1002
6.4. Legendrians and constructible sheaves	1005
6.5. Fukaya–Seidel categories of cotangent bundles	1007
7. Partially wrapped Fukaya categories and microlocal sheaves	1008
7.1. Homological cocores	1008
7.2. Liouville hypersurfaces	1010
7.3. Doubling I: Fully faithful embeddings of Fukaya categories	1012
7.4. A first comparison	1015
7.5. Doubling II: Microlocal sheaves and antimicrolocalization	1015
7.6. Proof of Theorem 1.4	1021
7.7. Making the core subanalytic	1025
Appendix A. Review of categorical notions	1027
A.1. Functors, modules, and bimodules	1027
A.2. Pre-triangulated, idempotent complete, and co-complete categories	1029
A.3. Compactly generated categories	1030

A.4. Morita equivalence	1031
A.5. Adjoints and compact objects	1031
A.6. Brown representability	1031
A.7. Quotients and localization	1032
A.8. Proper modules	1033
A.9. Smooth and proper categories	1034
A.10. Exceptional collections	1034
References	1035

1. Introduction

I saw the angel in the marble,
and carved until I set him free.

—Michelangelo (attributed)

The calculation of Fukaya categories of symplectic manifolds has emerged as a question of central interest in geometry. Within symplectic geometry itself, many questions may be phrased in terms of intersections of Lagrangian submanifolds; as the Fukaya category is built from these, it is a natural tool for their study. In low-dimensional topology, a number of invariants of smooth manifolds and smooth knots can be extracted from Fukaya categories of associated symplectic manifolds. Homological mirror symmetry, a largely conjectural correspondence arising from non-rigorous reasoning in mathematical physics, further predicts that many categories of interest in algebraic geometry and representation theory also arise as Fukaya categories.

Beyond the intrinsic interest in confirming or explaining these predictions, knowing that a category of interest arises as a Fukaya category suggests the existence of additional structures. For one example, morphism spaces in the Fukaya category are Floer homology chain complexes, and as such come with a natural basis; this often “explains” the existence of previously known “canonical” bases of these Hom spaces. For another, the relative ease of constructing symplectomorphisms (which act on the relevant Fukaya categories) gives a natural source of automorphisms of these categories that are far less apparent from other points of view. The difficulty in calculating Fukaya categories, which is present in all of the aforementioned settings, stems from the global and analytic nature of the pseudo-holomorphic disks appearing in the definition.

In this paper, we obtain a combinatorial description of the partially wrapped Fukaya categories of all stably polarized Weinstein manifolds (more generally, sectors), by showing that they are isomorphic to certain corresponding categories of microlocal sheaves.

1.1. Weinstein manifolds and partially wrapped Fukaya categories. A vector field Z on a symplectic manifold (X, ω) is said to be *Liouville* when

$L_Z\omega = \omega$. Recall that in this case, $\lambda := \omega(Z, \cdot)$ is a primitive for ω , and that such symplectic manifolds are necessarily non-compact. Such a triple (X, ω, Z) (equivalently (X, λ)) is called a *Liouville manifold* if, in addition, the non-compact ends of X are identified (necessarily uniquely) by Z with the positive end of the symplectization of a contact manifold [32]. The *core* of a Liouville manifold is the locus of points \mathfrak{c}_X which do not escape to infinity under the Liouville flow. The inclusion $\mathfrak{c}_X \subseteq X$ is a homotopy equivalence, and in some sense \mathfrak{c}_X carries all of the symplectic topology of X as well.

A Liouville manifold (X, ω, Z) is said to be *Weinstein* if Z is gradient-like [20]. The key feature of such manifolds is that the core \mathfrak{c}_X is a union of isotropic submanifolds, and moreover admits transverse Lagrangian disks (“cocores”) at its smooth Lagrangian points. Prototypical examples include cotangent bundles, affine algebraic varieties, and more generally (finite type) Stein complex manifolds. Many examples of interest in geometric representation theory, such as conical symplectic resolutions, quiver varieties, moduli of Higgs bundles, and cluster varieties are in this class. Moreover, any compact symplectic manifold whose symplectic form has rational periods can be presented as a compactification of a Weinstein manifold [24], [39], and hence, through a strategy introduced in [78], understanding the Fukaya categories of Weinstein manifolds serves as a stepping stone to studying the Fukaya categories of closed symplectic manifolds.

While various analytic difficulties in Floer theory are simplified in the Liouville setting (due to strong topological and geometric control on pseudo-holomorphic disks), there is a significant new layer of complexity possible, thanks to the non-compactness of the target space. Namely, as has been understood for some time, it is desirable in this context to enlarge the Fukaya category of compact Lagrangians by adding certain non-compact (properly embedded, conical at infinity) Lagrangians as well. These larger Fukaya categories often have better formal properties due to there being an ample supply of non-compact Lagrangians, and they are also required by mirror symmetry, where non-compact Lagrangians in non-compact targets arise as mirror objects to sheaves on non-compact or non-Calabi–Yau manifolds, whose Ext groups could be of infinite rank or fail to satisfy Poincaré duality. Fukaya categories of non-compact Lagrangians also have bearing on questions about the Reeb dynamics at infinity. There are many different ways to add non-compact Lagrangians to the Fukaya category, with substantially different results; the basic parameters are (1) in which directions at infinity to allow non-compact Lagrangians and (2) in what direction and by how much to perturb (“wrap”) Lagrangians at infinity when computing Floer homology.

The framework of *partially wrapped Fukaya categories* [14], [15], [88], [37], [38] has emerged as a way to describe and relate different prescriptions for asymptotics and wrapping. One specifies a subset at infinity which Lagrangians

cannot limit to or wrap past, called the *stop*. In the resulting category, Lagrangians which are isotopic in the complement of the stop induce isomorphic objects, and symplectomorphisms preserving the stop induce autoequivalences. The resulting category is also invariant under isotopies of the complement of the stop. A *stopped Liouville manifold* (X, Λ) consists of a Liouville manifold X and a stop Λ . The *relative core* $\mathfrak{c}_{X, \Lambda}$ of (X, Λ) is the set of those which do not escape to the complement of Λ at infinity under the positive Liouville flow.

The partially wrapped setting includes variants of many previous constructions:

- When the stop is empty, one obtains the (fully) wrapped Fukaya category of Abouzaid–Seidel [10].
- Given a smooth Legendrian Λ , there is a naive “infinitesimally wrapped” Fukaya category with asymptotics along Λ given equivalently by either (1) take the stop to be the complement of a small regular neighborhood of Λ , or (2) take the stop to be Λ and consider just the full subcategory of Lagrangians which admit wrappings converging to Λ ; see Section 6.3 for further discussion, including a comparison with [66].¹ More generally, we can take Λ to be the core of a Liouville hypersurface.
- The Fukaya category of a *Landau–Ginzburg model* $w : Y \rightarrow \mathbb{C}$, also known as the *Fukaya–Seidel category* when w is a Lefschetz fibration [79], can be modeled as the partially wrapped Fukaya category of Y stopped at (the core of) the Weinstein hypersurface $w^{-1}(-\infty)$.²

Of course, partially wrapped Fukaya categories form a significantly broader class than infinitesimal or Fukaya–Seidel categories (e.g., they can have infinite-dimensional morphism spaces). An illustrative example: categories of coherent complexes on arbitrary (not necessarily compact or smooth) n -dimensional toric stacks can be shown equivalent to partially wrapped Fukaya categories of $(\mathbb{C}^*)^n = T^*T^n$, by combining the sheaf theoretic work of [54] with the main theorem of the present article.

Of particular importance are (possibly singular) isotropic stops. Isotropicity of the stop plays the same role as the Weinstein condition on the symplectic manifold itself; for instance, the core of a *Weinstein hypersurface at infinity* is a typical singular isotropic stop of interest to mirror symmetry.

¹This naive category embeds fully faithfully into the category of “proper modules” over the partially wrapped Fukaya category stopped at Λ , which should be regarded as the more correct category. It is an open and likely hard geometric question to determine when this embedding is an equivalence, already for $\Lambda = \emptyset$.

²From our point of view, this should just be taken as the *definition* of the Fukaya–Seidel category. However, we note there are some technical differences between this definition and the standard definition, and a careful proof of their equivalence is, as far as we know, a folk result whose proof has no available reference (though a special case is treated in [38, §8.6]).

1.2. Topological and sheaf theoretic interpretations. Despite their analytic origins, Fukaya categories have often been found to admit topological interpretations. The first prototype is the fact that the Lagrangian Floer homology of an exact Lagrangian is nothing other than its ordinary cohomology. The work of Nadler and Zaslow [66], [62] provides a sweeping generalization of this, identifying infinitesimally wrapped Floer homologies of exact Lagrangians in cotangent bundles with morphisms of sheaves on the base. In a seemingly different direction, it was observed that *wrapped* Floer homology also has a topological interpretation. Indeed, work of Abbondandalo and Schwarz [2], [1], [3] (see also Cieliebak and Latschev [21]) found many instances where wrapped Floer homology is isomorphic to the homology of spaces of paths and loops. Building on these, Abouzaid showed that in fact the wrapped Fukaya category of a cotangent bundle is naturally identified with perfect modules over chains on the based loop space of the base [6] (see also [7]). This last result may be restated (by the ∞ version of the van Kampen theorem): the wrapped Fukaya category of a cotangent bundle is the global sections of the constant cosheaf of categories on the zero section with costalk $\text{Perf } \mathbb{Z}$. This formulation exhibits an instance of a more general conjecture of Kontsevich [51]: the wrapped Fukaya category of any Weinstein manifold X should be the global sections of a cosheaf of categories on its core \mathbf{c}_X .

Nadler’s work [63] unified these points of view, by proposing that while infinitesimally wrapped Fukaya categories are modeled by (micro)sheaves with perfect (micro)stalks (as in [66]), the partially wrapped category should be modeled by compact objects in the category of all (micro)sheaves with appropriate microsupport conditions. For essentially formal reasons, these categories of compact objects may be organized into a cosheaf of categories, so Nadler’s proposal is a strengthening of Kontsevich’s conjecture. At the time of Nadler’s original proposal, microsheaves were only defined for subsets of cotangent bundles, but the high codimension embedding trick from [81] has now defined a category of microsheaves on the core of any Liouville manifold [65].

Since microlocal sheaf categories are entirely combinatorial/topological in nature, Nadler’s proposal is a (conjectural) computation of (partially) wrapped Fukaya categories. An illustrative example calculation is given in [63], where it was shown that the relevant category of microlocal sheaves on the skeleton of the (higher dimensional) symplectic pairs of pants matched the mirror-symmetric prediction for the wrapped Fukaya category (which was more recently verified directly [55]).

Example. Remarkably, while wrapped and infinitesimal Floer homologies are rather different creatures, the same does not appear to be the case on the sheaf side of Nadler’s proposal. Notably, there is nothing like “wrapping” on

the sheaf side; instead, there are purely categorical operations. As it is difficult to appreciate the depth of the distinction at first, let us return to the example of the cotangent bundle, with empty stop. The core is just the zero section, and sheaves microsupported in the zero section are (almost by definition) nothing other than locally constant sheaves. If we require these to have finite stalks, then we are studying finite rank local systems (which coincides with the infinitesimally wrapped Fukaya category, i.e., in this case the Fukaya category of compact Lagrangians). On the other hand, a typical compact object in this category is the “tautological” local system, with fiber given by chains on the based loop space. Usually of infinite rank, this object is best understood in terms of the functor it co-represents: taking the stalk at a point. Note that no “wrapping” appears in this purely categorical procedure proposed by Nadler, yet it does in fact correctly recover the wrapped Fukaya category.

1.3. Main results. We now fix notation and state our main results more precisely. [Theorem 1.1](#) concerns the special case of cotangent bundles, and its proof comprises the bulk of the paper. [Theorem 1.4](#) is derived from [Theorem 1.1](#) and concerns more general stably polarized Liouville manifolds.

For a Liouville symplectic manifold X and closed subset $\Lambda \subseteq \partial_\infty X$, we write $\mathcal{W}(X, \Lambda)$ for the (partially) wrapped Fukaya category of X , stopped at Λ . Its objects are Lagrangians $L \subseteq X$ which are eventually conical and disjoint from Λ at infinity, and its morphism complexes are Floer cochains after wrapping Lagrangians in the complement of Λ . It is an A_∞ category defined in [\[37\]](#), [\[38\]](#) (see also [\[10\]](#), [\[11\]](#), [\[88\]](#)); we review its definition at the beginning of [Section 5](#). Particularly important objects of $\mathcal{W}(X, \Lambda)$ include: the Lagrangian linking disks to the smooth Legendrian points of Λ [\[38, §5.3\]](#), the Lagrangian cocore disks when X is Weinstein, and the cotangent fibers when $X = T^*M$ (which may be viewed as a special case of cocore disks). We write $\text{Perf } \mathcal{W}(X, \Lambda)$ for the idempotent-completed pre-triangulated closure of $\mathcal{W}(X, \Lambda)$. Whenever $\Lambda \subseteq \Lambda'$, there is a tautological functor $\mathcal{W}(X, \Lambda') \rightarrow \mathcal{W}(X, \Lambda)$.

For a smooth manifold M , we write $\text{Sh}(M)$ for the dg category of sheaves of dg \mathbb{Z} -modules on M . The microsupport of a sheaf \mathcal{F} is a closed conical locus $\text{ss}(\mathcal{F}) \subseteq T^*M$ whose role is to encode, infinitesimally, which restriction maps are quasi-isomorphisms. We write $\text{Sh}_\Lambda(M)$ for the full subcategory of $\text{Sh}(M)$ spanned by those sheaves whose microsupport at infinity is contained in Λ . Particularly nice functors $\text{Sh}_\Lambda(M) \rightarrow \text{Mod } \mathbb{Z}$ include the stalk functors at points of M and the microstalk functors at smooth Legendrian points of Λ . These definitions are reviewed in [Section 4](#). We denote by $\text{Sh}_\Lambda(M)^c$ the category of compact objects in $\text{Sh}_\Lambda(M)$. The reader is cautioned that the compact objects of $\text{Sh}_\Lambda(M)$ do not necessarily have perfect stalks or bounded homological degree; that is, they need not be constructible sheaves in the usual sense.

The relevance of such objects on the sheaf side was pointed out in [63], where [Theorem 1.1](#) was implicitly conjectured by the terminology “wrapped sheaves” for compact objects of $\text{Sh}_\Lambda(M)$ and “wrapped (microlocal) skyscrapers” for co-representatives of (micro)stalk functors.

THEOREM 1.1. *Let M be a real analytic manifold, and let $\Lambda \subseteq S^*M$ be a subanalytic closed isotropic subset. There there is a canonical equivalence of categories*

$$(1.1) \quad \text{Perf } \mathcal{W}(T^*M, \Lambda)^{\text{op}} = \text{Sh}_\Lambda(M)^c$$

which carries the linking disk at any smooth Legendrian point $p \in \Lambda$ to a co-representative of the microstalk functor at $p \in \Lambda$ and carries the cotangent fiber at a point $p \in M$ not in the image of Λ to a co-representative of the stalk functor.

Remark 1.2. Rather than passing to the opposite category of the Fukaya category, we could equivalently negate either Λ or the Liouville form on T^*M .

To prove [Theorem 1.1](#), we do not calculate a single $\mathcal{W}(T^*M, \Lambda)$ on its own. Instead, we calculate the functor $\Lambda \mapsto \text{Perf } \mathcal{W}(T^*M, \Lambda)^{\text{op}}$ (functoriality is with respect to inclusions $\Lambda \subseteq \Lambda'$) which is a much more rigid object. In fact, we formulate a pair of axioms which uniquely characterize such a system of categories $\Lambda \mapsto \mathcal{C}(\Lambda)$ ([Section 3](#)), so the proof of [Theorem 1.1](#) then reduces to verifying these axioms for both $\Lambda \mapsto \text{Perf } \mathcal{W}(T^*M, \Lambda)^{\text{op}}$ ([Section 5](#)) and $\Lambda \mapsto \text{Sh}_\Lambda(M)^c$ ([Section 4](#)).

The underlying reason this strategy can succeed is that there are special stops Λ (specifically $\Lambda = N_\infty^* \mathcal{S}$, the union of conormals to the strata of a Whitney triangulation \mathcal{S}) for which the Reeb dynamics in the complement of Λ are simple, thus making it tractable to show (1.1) by direct calculation. Since every Λ is a subset of some $N_\infty^* \mathcal{S}$, it then suffices to show that both sides of (1.1) transform in the same way when Λ gets smaller. On the Fukaya side, the functor $\mathcal{W}(T^*M, \Lambda') \rightarrow \mathcal{W}(T^*M, \Lambda)$ for $\Lambda' \supseteq \Lambda$ is the quotient by the linking disks to $\Lambda' \setminus \Lambda$; this was established recently in [38]. On the sheaf side, one quotients by co-representatives of microstalks; this is ultimately a consequence of co-isotropicity of the microsupport [49, Thm. 6.5.1]. The identification of linking disks with microstalks matches the wrapping exact triangle of [38] with the microlocal Morse description of sheaf cohomology from [41], [49]. The conclusion is then that choosing a Whitney triangulation \mathcal{S} of M whose conormal $N_\infty^* \mathcal{S}$ contains Λ yields a description of both categories $\text{Perf } \mathcal{W}(T^*M, \Lambda)^{\text{op}}$ and $\text{Sh}_\Lambda(M)^c$ as the same localization of the category $\text{Perf } \mathcal{W}(T^*M, N_\infty^* \mathcal{S})^{\text{op}} = \text{Perf } \mathcal{S} = \text{Sh}_{N_\infty^* \mathcal{S}}(M)^c$.

This approach to the proof of [Theorem 1.1](#) is rather different from the previous computations of Fukaya categories of cotangent bundles [66], [62], [6].

In particular, we rely on no results from these articles. In fact, our ability to add geometry to simplify the situation is such that the only Floer cohomology calculations which need to be made in this entire article are between Lagrangians which intersect in at most one point, obviating, in particular, the need to ever compute a holomorphic disc. We therefore expect that the proofs of the results in this paper would apply to the case of more general (e.g., sphere spectrum) coefficients, provided one has access to the definitions of the sheaf and Fukaya categories in these settings.

The equivalence of [Theorem 1.1](#) is also functorial under open inclusions:

PROPOSITION 1.3. *Let $M' \hookrightarrow M$ be an analytic open inclusion of analytic manifolds. For subanalytic isotropics $\Lambda' \subseteq S^*M'$ and $\Lambda \subseteq S^*M$ with $\Lambda' \supseteq \Lambda \cap S^*M'$, the following diagram commutes*

$$(1.2) \quad \begin{array}{ccc} \text{Perf } \mathcal{W}(M', \Lambda')^{\text{op}} & \longrightarrow & \text{Perf } \mathcal{W}(M, \Lambda)^{\text{op}} \\ \parallel \text{Theorem 1.1} & & \parallel \text{Theorem 1.1} \\ \text{Sh}_{\Lambda'}(M')^c & \longrightarrow & \text{Sh}_{\Lambda}(M)^c, \end{array}$$

where the bottom horizontal arrow is (the restriction to compact objects of) the left adjoint of restriction.

In [Section 6](#), we detail a number of applications and corollaries of [Theorem 1.1](#). These include a version of the original Nadler–Zaslow correspondence, translations of the microsheaf theoretic work on mirror symmetry for toric varieties and toric boundaries, and a sheaf theoretic description of the augmentation category of the partially wrapped Floer cochains of the linking disk to a Legendrian (expected to be equivalent to the Legendrian DGA), valid for the jet bundle of a manifold of any dimension. After the present work, a host of sheaf theoretic calculations [\[35\]](#), [\[84, 83\]](#), [\[82\]](#), [\[54\]](#), [\[64\]](#), [\[63\]](#), [\[36\]](#) can now be understood as computations of wrapped Fukaya categories.

Finally, in [Section 7](#), we turn from cotangent bundles to the general setting of stably polarized Weinstein manifolds. We proceed by combining the “doubling trick” of [\[38, Ex. 10.7 and 13.4\]](#) and the “antimicrolocalization” of [\[65, §7\]](#) to reduce to the cotangent bundle case. We arrive the following sheaf theoretic description of partially wrapped Fukaya categories:

THEOREM 1.4. *Let X be a real analytic Liouville manifold, and let $\Lambda \subseteq \partial_\infty X$ be a stop whose relative core $\mathfrak{c}_{X, \Lambda} := \mathfrak{c}_X \cup (\Lambda \times \mathbb{R}) \subseteq X$ is subanalytic singular isotropic. For any stable polarization³ of X , there is a fully faithful*

³A stable polarization of a symplectic manifold is the expression of its tangent bundle plus \mathbb{C}^k (some finite k) as the complexification of a real vector bundle; a choice of stable polarization controls the “twisting” of the categories on both sides of [\(1.3\)](#). Many examples

functor

$$(1.3) \quad \text{Perf } \mathcal{W}(X, \Lambda)^{\text{op}} \hookrightarrow \mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda})^c,$$

where $\mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda})$ denotes the category of microlocal sheaves on $\mathfrak{c}_{X, \Lambda}$. This functor sends a homological cocore at a smooth Lagrangian point p of $\mathfrak{c}_{X, \Lambda}$ to a co-representative of the microstalk at p .

In particular, if X is Weinstein, or more generally admits homological cocores, then (1.3) is an equivalence.

Let us comment on the hypotheses of [Theorem 1.4](#) and, in particular, argue that it applies to all stably polarized Weinstein sectors. The analyticity assumptions hold in most concrete cases of interest, and abstractly speaking, any Weinstein manifold (more generally, sector) may be perturbed so as to be real analytic and to have subanalytic relative core (see [Corollary 7.28](#)). A homological cocore at a smooth Lagrangian point $p \in \mathfrak{c}_{X, \Lambda}$ is an object of $\text{Perf } \mathcal{W}(X, \Lambda)$ whose image in $\text{Perf } \mathcal{W}((X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$ is the linking disk at $p \times \infty$. Admitting homological cocores means that every smooth Lagrangian point of $\mathfrak{c}_{X, \Lambda}$ has a homological cocore; this condition turns out to depend only on X up to deformation, and holds whenever X is Weinstein. For (X, Λ) as in [Theorem 1.4](#), the stabilization $X \times \mathbb{C}$ always admits homological cocores, so there is always an equivalence

$$\text{Perf } \mathcal{W}((X, \Lambda) \times (\mathbb{C}, \pm\infty))^{\text{op}} = \mu\text{sh}_{\mathfrak{c}_{X, \Lambda} \times \mathbb{R}}(\mathfrak{c}_{X, \Lambda} \times \mathbb{R})^c = \mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda})^c.$$

The embedding of [Theorem 1.4](#) depends a priori on a choice of analytic Liouville hypersurface embedding of $X \times \mathbb{C}^k$ into S^*M for some auxiliary analytic manifold M , compatible with stable polarizations. (Part of the proof is to show such data always exists.) We expect our methods could be extended to show that the embedding of [Theorem 1.4](#) is independent of this choice.

The embedding of [Theorem 1.4](#) associated to a given analytic Liouville hypersurface embedding $X \hookrightarrow S^*M$ is, by construction, compatible with the equivalence of [Theorem 1.1](#) in the sense that the following diagram commutes:

$$(1.4) \quad \begin{array}{ccc} \text{Perf } \mathcal{W}(X, \Lambda)^{\text{op}} & \longrightarrow & \text{Perf } \mathcal{W}(T^*M, \overline{\mathfrak{c}_{X, \Lambda}})^{\text{op}} \\ \text{Theorem 1.4} \downarrow & & \parallel \text{Theorem 1.1} \\ \mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda})^c & \xrightarrow{\mu^*} & \text{Sh}_{\overline{\mathfrak{c}_{X, \Lambda}}}(M)^c \end{array}$$

(see [Proposition 7.24](#)), where μ^* denotes the left adjoint to microlocalization. Using this compatibility, it is proven in [\[36\]](#) that for a Fano toric stack Y with

of interest are stably polarized, such as all cotangent bundles and all complete intersections in \mathbb{C}^n .

toric divisor D , there is a commutative diagram

$$(1.5) \quad \begin{array}{ccccccc} \text{Perf } \mathcal{W}(W^{-1}(-\infty)) & \longrightarrow & \text{Perf } \mathcal{W}(X, W) & \longrightarrow & \text{Perf } \mathcal{W}(X) & \longrightarrow & 0 \\ \parallel & & \parallel & & \parallel & & \\ \text{Coh}(D) & \longrightarrow & \text{Coh}(Y) & \longrightarrow & \text{Coh}(Y \setminus D) & \longrightarrow & 0, \end{array}$$

where $W : X \rightarrow \mathbb{C}$ is the mirror Landau–Ginzburg model (see [Example 7.25](#)).

Convention. Throughout this document, we work in the setting of dg and, equivalently, A_∞ categories over \mathbb{Z} (or more generally any commutative ring). We only ever consider “derived” functors, we only ever mean “homotopy” limits or colimits, and we systematically omit the word “quas.i” By modules, we mean dg or A_∞ modules; e.g., by \mathbb{Z} -modules we mean the category of chain complexes of abelian \mathbb{Z} -modules, localized at quasi-isomorphisms, except when, as in this sentence, we qualify it with the word “abelian.” In [Section A](#) we detail our assumptions about these categories and collect relevant categorical notions which will appear throughout the paper.

1.4. Acknowledgements. We thank Mohammed Abouzaid, Roger Casals, Alexander Efimov, Tobias Ekholm, Benjamin Gammage, David Nadler, Amnon Neeman, and Lenhard Ng for helpful discussions, some of which took place during visits to the American Institute of Mathematics.

S.G. was partially supported by NSF grant DMS–1907635 and would also like to thank the Simons Laufer Mathematical Sciences Institute (previously known as MSRI) for its hospitality during visits in Spring 2018 (supported by NSF grant DMS–1440140) and Fall 2022 (supported by NSF grant DMS–1928930) during which some of this work was completed. This research was conducted during the period J.P. served as a Clay Research Fellow and was partially supported by a Packard Fellowship and by the National Science Foundation under the Alan T. Waterman Award, Grant No. 1747553. V.S. was supported by the NSF CAREER grant DMS–1654545.

2. Stratifications

2.1. Generalities. Let X be a topological space. By a stratification \mathcal{S} of X , we mean a locally finite decomposition into disjoint locally closed subsets $\{X_\alpha\}_{\alpha \in \mathcal{S}}$, called strata, such that each boundary $\overline{X_\alpha} \setminus X_\alpha$ is a union of other strata X_β . The collection of strata \mathcal{S} is naturally a poset, in which there is a map $\beta \rightarrow \alpha$ if and only if $X_\alpha \subseteq \overline{X_\beta}$.

Remark 2.1. The poset \mathcal{S} does not generally capture the homotopy type of the space X . Conditions under which it does (contractibility of various strata/stars) are well known and recalled below.

We will say a subset $Y \subseteq X$ is \mathcal{S} -constructible when it is a union of strata of the stratification \mathcal{S} of X . We say that a stratification \mathcal{T} refines a stratification \mathcal{S} when the strata of \mathcal{S} are \mathcal{T} -constructible.

We recall that an abstract simplicial complex on a vertex set \mathcal{V} is a collection Σ of nonempty finite subsets of \mathcal{V} , containing all singletons and all subsets of elements of Σ . By a simplicial complex, we mean the geometric realization $|\Sigma|$ of an abstract simplicial complex Σ ; it comes with a stratification by the “open simplices” (which, of course, are locally closed, not necessarily open, subsets of $|\Sigma|$). We say a stratification \mathcal{S} on X is a triangulation when there exists a homeomorphism $|\Sigma| \xrightarrow{\sim} X$ identifying stratifications. We never impose any sort of regularity condition (differentiability, smoothness, analyticity, etc.) on this homeomorphism, even in the context of stratifications of a given regularity class. Note that the following are *not* triangulations: a stratification of a circle into single point and its complement, or into two points and their complement; the stratification into three points and their complement is a triangulation.

The open star of a stratum is the union of strata whose closures contain it. Taking stars reverses the inclusion: we have $X_\alpha \subseteq \overline{X_\beta} \iff \text{star}(X_\beta) \subseteq \text{star}(X_\alpha)$. Note that $\text{star}(X_\alpha) \cap \text{star}(X_\beta) = \bigcup_{\alpha \leftarrow \gamma \rightarrow \beta} \text{star}(X_\gamma)$. For triangulations, we can do better: $\text{star}(X_\alpha) \cap \text{star}(X_\beta) = \text{star}(X_\gamma)$, where γ is the simplex spanned by the vertices of α union the vertices of β (if this simplex is present), and otherwise $\text{star}(X_\alpha) \cap \text{star}(X_\beta) = \emptyset$.

For a C^p manifold M , we say a stratification \mathcal{S} is C^p if each stratum M_α is a (locally closed) C^p submanifold.

A C^1 stratification \mathcal{S} of a C^1 manifold M is called a *Whitney stratification* if and only if it satisfies Whitney’s conditions (a) and (b). These are usually stated as the following conditions on pairs of strata X and Y of \mathcal{S} :

- (a) For any sequence $x_i \in X$ converging to $y \in Y$ such that $T_{x_i}X$ converges to a subspace $V \subseteq T_yM$, we have $T_yY \subseteq V$.
- (b) For any pair of sequences $x_i \in X$ and $y_i \in Y$ both converging to $y \in Y$ such that $T_{x_i}X$ converges to a subspace $V \subseteq T_yM$ and the secant directions from y_i to x_i converge to a line $L \subseteq T_yM$, we have $L \subseteq V$.

By compactness of flag varieties, we may pass to convergent subsequences, and hence conditions (a) and (b) may be reformulated as follows:

- (a) For strata $Y \subseteq \overline{X}$, as $X \ni x \rightarrow y \in Y$, the tangent spaces T_xX become arbitrarily close to containing T_yY , uniformly over compact subsets of Y .
- (b) For strata $Y \subseteq \overline{X}$, as $X \ni x \rightarrow y \in Y$, the secant lines between x and y become arbitrarily close to being contained in T_xX , uniformly over compact subsets of Y .

Whitney’s condition (a) is equivalent to the assertion that the union of conormals $N^*\mathcal{S} := \bigcup_\alpha N^*M_\alpha \subseteq T^*M$ is closed. In fact, it is not hard to see that

Whitney's condition (b) implies Whitney's condition (a) [61, Prop. 2.4]. A Whitney stratification is, by definition, at least C^1 ; it makes sense to consider C^p Whitney stratifications for any $p \geq 1$, including $p = \infty$.

In order to guarantee the existence of Whitney stratifications, we will ultimately restrict to the setting of (real) analytic manifolds and subanalytic stratifications. We recall that a set is defined to be subanalytic when locally (i.e., in a neighborhood of every point of its closure) it is the analytic image of a relatively compact semianalytic set (i.e., locally defined by finitely many analytic inequalities). The canonical modern reference for subanalytic geometry is [18]. By a subanalytic stratification, we mean a stratification in which all strata are subanalytic. Every subanalytic stratification admits a subanalytic refinement in which all strata are locally closed analytic submanifolds. It is a fundamental result that for any locally finite collection of subanalytic subsets of an analytic manifold, there exists a subanalytic stratification with respect to which all the subsets are constructible. For proofs of these results, see [18], [85]. We also require the result that every subanalytic stratification admits a refinement to a subanalytic Whitney triangulation [86], [22], [23].

Remark 2.2. Wherever we have written “subanalytic,” one could substitute “defineable” with respect to any fixed analytic-geometric category [26], [25]. Every defineable stratification has a defineable refinement to a C^p Whitney triangulation for any given $p < \infty$ [86], [22], [23]. The fact that this is not known to hold for $p = \infty$ does not create any difficulties.

LEMMA 2.3. *Let M be a manifold with Whitney stratification \mathcal{S} . If $N \subseteq M$ is a locally closed submanifold transverse to every stratum of \mathcal{S} , then the intersected stratification $\mathcal{S} \cap N$ is a Whitney stratification of N .*

Proof. It suffices to verify that $\mathcal{S} \cap N$ satisfies Whitney (b). Thus consider a pair of strata $X \cap N$ and $Y \cap N$ with $Y \subseteq \overline{X}$. Whitney (b) for the stratification \mathcal{S} guarantees that the secant line from $x \in X \cap N$ to $y \in Y \cap N$ becomes arbitrarily close to being contained in $T_x X$ as $x \rightarrow y$, uniformly over compact subsets of $Y \cap N$. On the other hand, Whitney (b) for the stratification for $\mathcal{S} \cap N$ requires this secant line to become arbitrarily close to being contained in $T_x X \cap T_x N$, a stronger condition. Since N is a submanifold, the secant line certainly becomes arbitrarily close to being contained in $T_x N$. Our task is thus to pass from being close to $T_x X$ and $T_x N$ to being close to their intersection $T_x X \cap T_x N$. It thus suffices to show that $T_x X$ and $T_x N$ are uniformly transverse as $x \rightarrow y$. Let us see how this follows from Whitney (a) for \mathcal{S} . Since N is transverse to Y , we have $T_y Y + T_y N = T_y M$. Whitney (a) for \mathcal{S} means that $T_x X$ is arbitrarily close to containing $T_y Y$ as $x \rightarrow y$. Thus $T_x X + T_x N = T_x M$ uniformly as $x \rightarrow y$, as desired. \square

Remark 2.4. Here is a typical application of [Lemma 2.3](#). Let M be a manifold with a Whitney stratification \mathcal{S} , and let $N \subseteq M$ be a closed (as a subset) submanifold transverse to every stratum of \mathcal{S} . Then the stratification \mathcal{S}_N of M with strata $M_\alpha \cap N$ and $M_\alpha \setminus N$ for $\alpha \in \mathcal{S}$ (the poset of strata \mathcal{S}_N is thus $\mathcal{S} \times \{(M \setminus N) > N\}$) is Whitney. Indeed, Whitney (b) for \mathcal{S}_N is a special case of Whitney (b) for \mathcal{S} and $\mathcal{S} \cap N$. A slight modification of this example will also come up later. Let $B \subseteq M$ be a closed ball whose boundary is transverse to a Whitney stratification \mathcal{S} . The stratification of M by $M_\alpha \cap \partial B$, $M_\alpha \cap B^\circ$, and $M_\alpha \setminus B$, for strata $M_\alpha \subseteq M$, is now Whitney by the same argument using [Lemma 2.3](#).

2.2. Microlocal approximation of constructible sets. A constructible set X with respect to a Whitney stratification is in general quite singular. Our goal in this section is to show how such sets can be *microlocally* approximated by manifolds-with-corners X^η parametrized by small $\eta > 0$, in the sense that $X^\eta \rightarrow X$ and the conormal to X^η converges to (being contained in) the conormal of X as $\eta \rightarrow 0$. This result will be used in the proof of [Proposition 4.8](#) and in [Section 5.7](#). We fix an integer $p \geq 1$, possibly $p = \infty$, and work throughout with stratifications of class C^p .

A C^p radius function for a locally closed C^p submanifold $Y \subseteq M$ is a pair (U, ρ) , where $U \subseteq M$ is an open set containing Y and $\rho : U \rightarrow \mathbb{R}_{\geq 0}$ is of class C^p on $U \setminus Y$ and satisfies the following three conditions:

- $\rho^{-1}(0) = Y$.
- ρ is Lipschitz on a neighborhood of any compact subset of Y .
- The \liminf of the evaluation of $d\rho(x)$ on the secant direction from $y \in Y$ to $x \in M$ is bounded below by some $\epsilon > 0$ as $x \rightarrow y$, provided y is constrained to a compact subset of Y and the ratio $\frac{d(x,y)}{d(x,Y)}$ is bounded by some fixed $N < \infty$. (This condition is well defined since ρ is assumed Lipschitz; it implies $|d\rho(x)|$ is bounded away from zero over neighborhoods of compact subsets of Y .)

The standard radius function for $\mathbb{R}^n \times 0 \subseteq \mathbb{R}^n \times \mathbb{R}^m$ is of course

$$(a_1, \dots, a_n, b_1, \dots, b_m) \mapsto (b_1^2 + \dots + b_m^2)^{1/2}.$$

Every locally closed C^p submanifold admits a C^p radius function, as can be seen by choosing a collection of local coordinate patches and summing together standard radius functions via a partition of unity. (A convex combination of radius functions is a radius function.)

The following (trivial) restriction property for radius functions will be important: if (U, ρ) is a radius function for $Y \subseteq M$, and $N \subseteq M$ is transverse to Y , then $(U \cap N, \rho|_{U \cap N})$ is a radius function for $Y \cap N \subseteq N$.

When $Y \subseteq M$ is relatively compact, a radius function (U, ρ) for Y will be called *proper* when for every open set V containing \overline{Y} , there exists $\epsilon > 0$ such

that $\rho^{-1}([0, \epsilon]) \subseteq V$. It is easy to produce proper radius functions: for any radius function (U, ρ) , there exists an open set $U' \subseteq U$ containing Y such that $(U', \rho|_{U'})$ is proper.

The purpose of a radius function is to define tubular neighborhoods $\rho^{-1}([0, \epsilon])$. The conditions in the definition of a radius function are chosen so as to be able to prove the following two key assertions:

LEMMA 2.5. *Let (U, ρ) be a radius function for $Y \subseteq M$. We have $d\rho(x) \rightarrow N^*Y$ as $x \rightarrow Y$, uniformly over compact subsets of Y . (Equivalently, $N^*\rho^{-1}(r)$ approaches being contained in N^*Y as $r \rightarrow 0$, uniformly over compact subsets of Y .)*

Proof. Suppose for sake of contradiction that there is a sequence $x_i \rightarrow y \in Y$ with $d\rho(x_i) \rightarrow \xi \in T_y^*M \setminus N_y^*Y$. Since $\xi \notin N_y^*Y$, there exist $y_i \in Y$ with $\frac{d(x_i, y_i)}{d(x_i, Y)}$ uniformly bounded and the secant direction from y_i to x_i converging to the kernel of ξ . This contradicts the final axiom of a radius function. \square

LEMMA 2.6. *Let $Y \subseteq M$ be a stratum of a Whitney stratification \mathcal{S} , and let (U, ρ) be a radius function for Y . There exists an open set $V \subseteq U$ containing Y such that the submanifold $\rho^{-1}(r)$ is transverse to \mathcal{S} over V for all $r > 0$.*

Proof. The secant line to $x \in \rho^{-1}(r)$ from nearby $y \in Y$ pairs positively with $d\rho(x)$, whereas Whitney (b) requires that this secant line approach the tangent space to the stratum containing x in the limit $x \rightarrow y$. Since $d\rho$ is bounded, this gives a positive lower bound on the restriction of $d\rho$ to any stratum of \mathcal{S} in a neighborhood of any compact subset of Y . \square




Figure 1. A compact constructible set X (left) and its outward cornering X^ϵ (right).

We now turn to the setting of a compact set $X \subseteq M$ constructible with respect to a chosen Whitney stratification \mathcal{S} . Given a proper radius function for each stratum of \mathcal{S} contained in X , we define the “outward cornering” of X with respect to \mathcal{S} (see Figure 1) to be

$$(2.1) \quad X^\epsilon = \bigcup_{M_\alpha \subseteq X} \rho_\alpha^{-1}([0, \epsilon_\alpha])$$

for $\epsilon = (\epsilon_\alpha)_{M_\alpha \subseteq X}$, where it is tacitly required that $\epsilon_\alpha > 0$ be sufficiently small as an unspecified function of $(\epsilon_\beta)_{M_\beta \subsetneq \overline{M_\alpha}}$. Note that, no matter our notion of sufficiently small, we can always find a parametrization $\underline{\epsilon}(\eta)$ where each $\epsilon_\alpha > 0$ is an increasing function of $\eta > 0$, limiting to zero as $\eta \rightarrow 0$, such that $\epsilon_\alpha(\eta) > 0$ is sufficiently small in terms of $(\epsilon_\beta(\eta))_{M_\beta \subsetneq \overline{M_\alpha}}$ for all $\eta > 0$ (proof: by induction on strata).

The significance of properness of the given radius functions is that it (along with compactness of X) ensures that the part of ∂X^ϵ coming from $\rho_\alpha^{-1}(\epsilon_\alpha)$ is contained in a neighborhood of a *compact* subset of $M_\alpha \subseteq X$ (depending on $(\epsilon_\beta)_{M_\beta \subsetneq \overline{M_\alpha}}$), hence falls within the scope of [Lemma 2.6](#).

Here is the first key property of X^ϵ :

COROLLARY 2.7. *Fix a Whitney stratification \mathcal{S} of M . Let $X \subseteq M$ be a compact constructible subset, with a choice of proper radius function for each of its strata. Then $M \setminus X^\epsilon$ is a manifold-with-corners, all of whose corner strata are transverse to \mathcal{S} .*

Proof. We proceed by induction on the number of strata of X . Thus let $X = X_0 \cup M_\alpha$, and suppose the result is known for X_0 . This means, in particular, that $M \setminus X_0^\epsilon$ is a manifold-with-corners, all of whose corner strata are transverse to M_α . It follows that further removing the small regular neighborhood $\rho_\alpha^{-1}([0, \epsilon_\alpha])$ produces a manifold-with-corners $M \setminus X^\epsilon$, provided $\epsilon_\alpha > 0$ is sufficiently small as a function of $(\epsilon_\beta)_{M_\beta \subsetneq \overline{M_\alpha}}$.

Now let us show that the corner strata of $M \setminus X^\epsilon$ are transverse to \mathcal{S} . The boundary stratum of $M \setminus X^\epsilon$ coming from $\rho_\alpha^{-1}(\epsilon_\alpha)$ is transverse to \mathcal{S} by [Lemma 2.6](#). A general corner stratum of $M \setminus X^\epsilon$ is the intersection $C \cap \rho_\alpha^{-1}(\epsilon_\alpha)$, where C is a corner stratum of $M \setminus X_0^\epsilon$. To see that such an intersection is also transverse to \mathcal{S} , apply [Lemma 2.6](#) to the stratum $C \cap M_\alpha \subseteq C$ of the restriction to C of \mathcal{S} (which is Whitney by [Lemma 2.3](#), locally uniformly in $\epsilon \setminus \epsilon_\alpha$) equipped with the restriction of ρ_α (which remains a radius function as noted earlier). \square

Note that the conclusion of [Corollary 2.7](#) (transversality of ∂X^ϵ and \mathcal{S}) is equivalent to saying that $N^*\mathcal{S}$ (the union of the conormals of all strata) and N^*X^ϵ (the union of the conormals of all corner strata) are disjoint at infinity.

Given transversality of ∂X^ϵ and \mathcal{S} , we can define the “big conormal” $N^*(\mathcal{S}|(M \setminus X^\epsilon))$ to be the union of conormals of intersections of strata of \mathcal{S} and corner strata of $M \setminus X^\epsilon$. The second key property of X^ϵ is the following convergence result:

COROLLARY 2.8. *Fix a Whitney stratification \mathcal{S} of M . Let $X \subseteq M$ be a compact constructible subset, with a choice of proper radius function for each of its strata. Then the big conormal $N^*(\mathcal{S}|(M \setminus X^\epsilon))$ lies in arbitrarily small neighborhoods of $N^*\mathcal{S}$ as $\epsilon \rightarrow 0$.*

Proof. We proceed by induction on the number of strata of X . Thus let $X = X_0 \cup M_\alpha$, and suppose the result is known for X_0 . It thus suffices to show that $N^*(\mathcal{S}|(M \setminus X^\epsilon))$ lies in an arbitrarily small neighborhood of $N^*(\mathcal{S}|(M \setminus X_0^\epsilon))$ as $\epsilon_\alpha \rightarrow 0$ (uniformly over compact subsets of $\epsilon \setminus \epsilon_\alpha$).

The conormal of the intersection of a stratum M_γ of \mathcal{S} with $\rho_\alpha^{-1}(\epsilon_\alpha)$ is the sum of the conormals of M_γ and $\rho_\alpha^{-1}(\epsilon_\alpha)$. These individually approach being contained in N^*M_α in the limit $\epsilon_\alpha \rightarrow 0$ (the first by Whitney (a) and the second by [Lemma 2.5](#)), and they are quantitatively transverse by the axioms of a radius function (as was the main point of the proof of [Lemma 2.6](#)). It follows that their sum also approaches N^*M_α as $\epsilon_\alpha \rightarrow 0$.

The general case is that of the conormal of $M_\gamma \cap \rho_\alpha^{-1}(\epsilon_\alpha) \cap C$, where C is a corner stratum of $M \setminus X_0^\epsilon$. It follows by applying the same argument to the intersection of the situation with C , as in the proof of [Corollary 2.7](#). \square

Remark 2.9. In the context of [Corollary 2.8](#), note that any subanalytic family of Legendrians inside S^*M , whose projections converge to X , will themselves converge to a subset of the conormal of X *with respect to some refinement of \mathcal{S}* . [Corollary 2.8](#) provides a stronger convergence result (we do not need to refine the stratification) for the particular family of outward cornerings defined in [\(2.1\)](#).

2.3. Cornering and conormals of constructible open sets. Here we develop some finer properties of the microlocal approximations constructed in the previous subsection. They will not be used until [Section 5.7](#).

Let \mathcal{S} be a Whitney stratification of M by locally closed smooth submanifolds. For any \mathcal{S} -constructible relatively compact open set $U \subseteq M$, we define its *inward cornering*

$$(2.2) \quad U^{-\epsilon} := U \setminus \overline{(\partial U)^\epsilon},$$

where $(\partial U)^\epsilon$ denotes the outward cornering ∂U defined in [\(2.1\)](#). Thus $U^{-\epsilon}$ is (the interior of) a codimension zero submanifold-with-corners ([Corollary 2.7](#)), depending smoothly on ϵ , such that as $\epsilon \rightarrow 0$, its conormal $N^*U^{-\epsilon}$ remains disjoint from $N^*\mathcal{S}$ at infinity ([Corollary 2.7](#)) yet limits inside it ([Corollary 2.8](#)). Strictly speaking, $U^{-\epsilon}$ also depends on the choices of tubular neighborhoods of the strata comprising ∂U , however we will leave this choice out of the notation as it is never particularly relevant. (Note that it is a convex, hence contractible, choice.) When even the choice of ϵ is not relevant, we will simply write U^- .

Taking $\epsilon \rightarrow 0$, we learn that

LEMMA 2.10. *The open sets U and U^- are diffeomorphic, and the diffeomorphism may be chosen to be the identity on any fixed compact subset of U .*

When \mathcal{S} is a triangulation, we may consider for any simplex $s \in \mathcal{S}$ its open star, $\text{star}(s)$.

LEMMA 2.11. *For \mathcal{S} a triangulation and any $s \in \mathcal{S}$, there is a homotopy equivalence $\partial \text{star}(s)^- \simeq \partial \text{star}(s)$.*

Proof. Both $\text{star}(s)$ and $\text{star}(s)^-$ are the interiors of compact (topological) manifolds-with-boundary (namely their closures). They are also diffeomorphic: $\text{star}(s) \cong \text{star}(s)^-$ by Lemma 2.10. It therefore suffices to recall the standard fact that the interior of a compact manifold-with-boundary remembers its boundary, up to homotopy equivalence.

Indeed, let \overline{M} be a compact manifold-with-boundary, and let $M = \overline{M} \setminus \partial \overline{M}$ denote its interior. The “end space” $e(M)$ is the space of proper maps $\mathbb{R}_{\geq 0} \rightarrow M$. (This is a model for the homotopy inverse limit of $M \setminus K$ over compact subsets $K \subseteq M$.) A choice of collar $\partial \overline{M} \times [0, 1] \hookrightarrow \overline{M}$ determines a homotopy equivalence

$$(2.3) \quad e(M) \xleftarrow{\sim} e(\partial \overline{M} \times (0, \frac{1}{2})) = C(\mathbb{R}_{\geq 0}, \partial \overline{M}) \times e((0, \frac{1}{2}]),$$

and we have homotopy equivalences $C(\mathbb{R}_{\geq 0}, \partial \overline{M}) \simeq \partial \overline{M}$ and $e((0, \frac{1}{2})) \simeq *$, so we have $e(M) \simeq \partial \overline{M}$. (Compare [45, §1].) \square

LEMMA 2.12. *For \mathcal{S} a triangulation and for simplices $s, t \in \mathcal{S}$ with $\text{star}(t) \cap \text{star}(s) \neq \emptyset$ and $t \not\rightarrow s$, the intersection $\text{star}(t)^{-\varepsilon} \cap \partial \text{star}(s)^{-\underline{\delta}}$ is contractible for ε sufficiently small and $\underline{\delta}$ sufficiently small in terms of ε .*

Proof. As in the proof of Lemma 2.11, the intersection

$$\text{star}(t)^{-\varepsilon} \cap \partial \text{star}(s)^{-\underline{\delta}}$$

is the end space of $\overline{\text{star}(t)^{-\varepsilon} \cap \text{star}(s)^{-\underline{\delta}}}$. As in Lemma 2.10, this intersection $\overline{\text{star}(t)^{-\varepsilon} \cap \text{star}(s)^{-\underline{\delta}}}$ is diffeomorphic to $\overline{\text{star}(t)^{-\varepsilon} \cap \text{star}(s)}$, whose end space is in turn given by $\text{star}(t)^{-\varepsilon} \cap \partial \text{star}(s)$. Now we may take $\varepsilon \rightarrow 0$ again mimicking the proof of Lemma 2.10 to see that this is homotopy equivalent to $\text{star}(t) \cap \partial \text{star}(s)$. The assumptions on s and t now imply that this is contractible. \square

LEMMA 2.13. *If U has smooth boundary, then there is a C^0 -small isotopy between U^- and a small inward pushoff of U .*

Proof. The definition of U^- depends on choice of radius functions for the strata comprising ∂U . Fix coordinates $\partial U \times \mathbb{R} \subseteq M$ near ∂U , and choose radius functions whose inward derivative in the \mathbb{R} -direction is positive inside U . Such radius functions exist locally, hence can be patched together using a partition of unity pulled back from ∂U (i.e., independent of the \mathbb{R} -coordinate), which preserves the property of having positive inward derivative inside U . Now using these radius functions, each vertical line $p \times \mathbb{R}$ for $p \in \partial U$ intersects ∂U^- exactly once, transversally, which provides the desired isotopy. \square

Remark 2.14. It is *not* asserted that the isotopy in [Lemma 2.13](#) will ensure that the conormal remains disjoint from N^*S at infinity. This will require us to exercise some care when applying it.

For the next result, consider a Whitney stratification S and a point $q \in M$ lying in a stratum M_χ . Let S_q denote the Whitney stratification obtained from S by replacing M_χ with $M_\chi \setminus q$ and $\{q\}$. Given a compact S -constructible set $X \subseteq M$ containing q , we can consider its outward cornerings X^ϵ and $X^{\epsilon,\delta}$ with respect to S and S_q , respectively, where $\epsilon = (\epsilon_\alpha)_{M_\alpha \subseteq X}$ and $\delta > 0$ is associated to q . Evidently,

$$(2.4) \quad X^{\epsilon,\delta} = X^\epsilon \cup B_\delta(q),$$

and according to the definition of “outward cornering” above, there is the implicit requirement that $\epsilon_\chi > 0$ be sufficiently small as a function of $(\epsilon_\beta)_{M_\beta \subsetneq M_\alpha}$ and δ . The next result concerns the behavior of $X^{\epsilon,\delta}$ when we *remove* the dependence of ϵ_χ on δ . The resulting neighborhoods are illustrated in [Figure 2](#), which should be contrasted with [Figure 1](#).




Figure 2. The neighborhoods $X^{\epsilon,\delta}$.

PROPOSITION 2.15. *Let $X \subseteq M$ be a compact S -constructible subset, and fix $q \in X$ living in stratum χ . Let $\tilde{X}^{\epsilon,\delta} = X^\epsilon \tilde{\cup} B_\delta(q)$, where X^ϵ is the outward cornering (2.1) and the notation $\tilde{\cup}$ indicates that the boundary of the union $\rho_\chi^{-1}([0, \epsilon_\chi]) \cup B_\delta(q)$ is smoothed along its potential corner locus $\rho_\chi^{-1}(\epsilon_\chi) \cup \partial B_\delta(q)$, and then the remaining tubes are added. Then for suitable choices of radius functions near q , these modified outward cornerings $X^{\epsilon,\delta}$ satisfy the conclusion of [Corollary 2.7](#).*

Proof. Choose a local Euclidean chart near q in which $q \in M_\chi \subseteq M$ is locally modelled on $0 \in \mathbb{R}^k \subseteq \mathbb{R}^n$, and let us choose the usual radius (i.e., distance) functions in this chart. Note that we thus have a completely explicit picture of how these tubes intersect near q ; in particular, their union with smoothed boundary is well behaved. We regard the union of tubes (with smoothed boundary) $\rho_\chi^{-1}([0, \epsilon_\chi]) \tilde{\cup} B_\delta(q)$ as a single object associated to the stratum χ , albeit depending on two parameters ϵ_χ and δ . This single object satisfies the conclusion of [Lemma 2.6](#) by Whitney (b), which is all that is used in the inductive proof of [Corollary 2.7](#). \square

3. Microlocal Morse categories

3.1. *Strata poset categories and refinement functors.* Let \mathcal{S} be a stratification. We fix the following notation for the Yoneda embedding:

$$(3.1) \quad \mathcal{S} \rightarrow \text{Fun}(\mathcal{S}^{\text{op}}, \text{Set}),$$

$$(3.2) \quad \alpha \mapsto \text{Hom}(\cdot, \alpha) =: 1_{\text{star}(\alpha)}.$$

Note that

$$(3.3) \quad \text{Hom}(1_{\text{star}(\alpha)}, 1_{\text{star}(\beta)}) = 1_{\text{star}(\beta)}(\alpha) = \text{Hom}(\alpha, \beta) = \begin{cases} \{1\} & \text{star}(\alpha) \subseteq \text{star}(\beta), \\ \emptyset & \text{otherwise.} \end{cases}$$

For any \mathcal{S} -constructible open set U , we introduce the functor $1_U \in \text{Fun}(\mathcal{S}^{\text{op}}, \text{Set})$ defined by the analogous formula

$$(3.4) \quad \text{Hom}(1_{\text{star}(\alpha)}, 1_U) = 1_U(\alpha) := \begin{cases} \{1\} & \text{star}(\alpha) \subseteq U, \\ \emptyset & \text{otherwise.} \end{cases}$$

(The action of 1_U on morphism sets is in fact uniquely determined by the above, since when $\alpha \rightarrow \beta$, the set $\text{Hom}_{\text{Set}}(1_U(\beta), 1_U(\alpha))$ always consists of one element.) Note that $\text{star}(\alpha) \subseteq U$ if and only if $\alpha \subseteq U$. More generally, for $U \subseteq V$, there is a unique natural transformation

$$(3.5) \quad 1_U \rightarrow 1_V.$$

It sends $1 \in 1_U(\alpha)$ to $1 \in 1_V(\alpha)$ for any $\text{star}(\alpha) \subseteq U \subseteq V$.

Now let \mathcal{S}' be a stratification refining \mathcal{S} . There is a natural map $r : \mathcal{S}' \rightarrow \mathcal{S}$, sending a stratum in \mathcal{S}' to the unique stratum in \mathcal{S} containing it. We write

$$(3.6) \quad r^* : \text{Fun}(\mathcal{S}^{\text{op}}, \text{Set}) \rightarrow \text{Fun}(\mathcal{S}'^{\text{op}}, \text{Set})$$

for the pullback of functors along this map r . For $\tau' \in \mathcal{S}'$ and an \mathcal{S} -constructible open set U , we have

$$(3.7) \quad \text{Hom}(1_{\text{star}(\tau')}, r^* 1_U) = (r^* 1_U)(\tau') = 1_U(r(\tau')) = \begin{cases} \{1\} & \text{star}(r(\tau')) \subseteq U, \\ \emptyset & \text{otherwise.} \end{cases}$$

Since U is open and \mathcal{S} -constructible, we have $\text{star}(r(\tau')) \subseteq U$ if and only if $\text{star}(\tau') \subseteq U$, so we conclude that $r^* 1_U = 1_U$.

We now linearize. We write $\mathbb{Z}[\mathcal{S}]$ for the linearization of a poset \mathcal{S} . We write $\text{Mod } \mathcal{S}$ for the category of modules $\text{Fun}(\mathcal{S}^{\text{op}}, \text{Mod } \mathbb{Z}) = \text{Fun}(\mathbb{Z}[\mathcal{S}]^{\text{op}}, \text{Mod } \mathbb{Z})$, and we use $r^* : \text{Mod } \mathcal{S} \rightarrow \text{Mod } \mathcal{S}'$ for pullback of modules as above. As with any pullback of modules, this functor has a left adjoint, typically termed extension of scalars or induction, which we write as $r_! : \text{Mod } \mathcal{S}' \rightarrow \text{Mod } \mathcal{S}$, which fits into

the commuting diagram

$$(3.8) \quad \begin{array}{ccc} \mathcal{S}' & \xrightarrow{s \mapsto 1_{\text{star}(s)}} & \text{Mod } \mathcal{S}' \\ r \downarrow & & \downarrow r_! \\ \mathcal{S} & \xrightarrow{s \mapsto 1_{\text{star}(s)}} & \text{Mod } \mathcal{S}. \end{array}$$

Restriction of scalars r^* is co-continuous, so its left adjoint $r_!$ extension of scalars preserves compact objects, giving a map $r_! : \text{Perf } \mathcal{S}' \rightarrow \text{Perf } \mathcal{S}$ (which can also be viewed as the canonical extension of $r : \mathcal{S}' \rightarrow \mathcal{S}$ to the idempotent-completed pre-triangulated hulls).

3.2. A category for any Λ . We now wish to define a *microlocal Morse category* $\mathcal{C}(\Lambda)$ for any subanalytic (possibly) singular isotropic $\Lambda \subseteq S^*M$, together with functors $\mathcal{C}(\Lambda') \rightarrow \mathcal{C}(\Lambda)$ for inclusions $\Lambda' \supseteq \Lambda$. We define this system of categories $\Lambda \mapsto \mathcal{C}(\Lambda)$, the *microlocal Morse theatre*, by formulating axioms which characterize it uniquely. (Recall that $S^*M := (T^*M \setminus M)/\mathbb{R}_{>0}$ denotes the co-sphere bundle of M , and a closed subanalytic set $\Lambda \subseteq S^*M$ is called isotropic if and only if for some, or, equivalently, every, cover of Λ by locally closed C^1 submanifolds, all of them are isotropic.)

The previous subsection defined categories $\text{Perf } \mathcal{S}$ together with functors $r_! : \text{Perf } \mathcal{S}' \rightarrow \text{Perf } \mathcal{S}$ whenever \mathcal{S}' is a refinement of \mathcal{S} . For our current purpose, these categories do not have the correct significance for general stratifications \mathcal{S} (compare [Remark 2.1](#)). As such, we will consider these categories only for triangulations \mathcal{S} .⁴ The microlocal Morse theatre is an extension of this functor $\mathcal{S} \mapsto \text{Perf } \mathcal{S}$ on triangulations.

Definition 3.1. A *microlocal Morse pre-theatre* $\Lambda \mapsto \mathcal{C}(\Lambda)$ is a functor from the category of subanalytic singular isotropics inside S^*M to the category of dg categories over \mathbb{Z} . A *normalized* microlocal Morse pre-theatre is one equipped with an isomorphism of functors $(\mathcal{S} \mapsto \mathcal{C}(N_\infty^* \mathcal{S})) = (\mathcal{S} \mapsto \text{Perf } \mathcal{S})$ on Whitney triangulations \mathcal{S} .

Remark 3.2. Any isomorphism of functors $(\mathcal{S} \mapsto H^* \mathcal{C}(N_\infty^* \mathcal{S})) = (\mathcal{S} \mapsto H^* \text{Perf } \mathcal{S})$ automatically lifts to an isomorphism $(\mathcal{S} \mapsto \mathcal{C}(N_\infty^* \mathcal{S})) = (\mathcal{S} \mapsto \text{Perf } \mathcal{S})$ by [Proposition 5.28](#). This will be crucial when discussing Fukaya categories (specifically [Theorem 5.35](#)).

We will characterize the microlocal Morse theatre in terms of microlocal Morse theory.⁵ Let $f : M \rightarrow \mathbb{R}$ be a function and \mathcal{S} a stratification. An

⁴In fact, there are weaker conditions on a stratification \mathcal{S} (which are satisfied if \mathcal{S} is a triangulation) implying that $\text{Perf } \mathcal{S}$ is the correct category to associate to \mathcal{S} .

⁵More conventionally [41], this is called stratified Morse theory. We find the term “microlocal” more descriptive, and also the word stratified would otherwise take on too many meanings in this article.

intersection of Γ_{df} with $N^*\mathcal{S}$ is called an \mathcal{S} -critical point, which is said to be Morse if it is a transverse intersection at a smooth point of $N^*\mathcal{S}$. The function f is said to be \mathcal{S} -Morse when all its \mathcal{S} -critical points are Morse. When \mathcal{S} is subanalytic, such functions are plentiful and can be chosen analytic. (See [41, Thm. 2.2.1] for this assertion, which is collected there from various results in the literature.)

More generally, for any singular isotropic $\Lambda \subseteq S^*M$, a Λ -critical point of f is by definition an intersection of Γ_{df} with the union of the zero section and $\mathbb{R}_{>0} \times \Lambda$. That is,

$$(3.9) \quad \text{crit}_\Lambda(f) = \Gamma_{df} \cap (0_M \cup (\mathbb{R}_{>0} \times \Lambda)) \subseteq T^*M.$$

Such a Λ -critical point is said to be Morse if the intersection is transverse and occurs at a smooth point of $0_M \cup (\mathbb{R}_{>0} \times \Lambda)$, and any f whose Λ -critical points are all Morse is called Λ -Morse.

Definition 3.3. In any normalized microlocal Morse pre-theatre $\Lambda \mapsto \mathcal{C}(\Lambda)$, the *Morse characters* $\mathcal{X}_{\Lambda,p}(f, \epsilon, \mathcal{S}) \in \mathcal{C}(\Lambda)$ are defined as follows for smooth Legendrian points $p \in \Lambda$.

Let $f : M \rightarrow \mathbb{R}$ be an analytic function with a Morse Λ -critical point at p (i.e., somewhere in $\mathbb{R}_{>0} \times \{p\} \subseteq T^*M$) with critical value 0, no other Λ -critical points with critical values in the interval $[-\epsilon, \epsilon]$, and with relatively compact sublevel set $f^{-1}(-\infty, \epsilon)$. Let \mathcal{S} be a Whitney triangulation for which $\Lambda \subseteq N_\infty^*\mathcal{S}$ and for which both $f^{-1}(-\infty, -\epsilon)$ and $f^{-1}(-\infty, \epsilon)$ are \mathcal{S} -constructible.

The Morse character $\mathcal{X}_{\Lambda,p}(f, \epsilon, \mathcal{S})$ is then defined as the image of

$$(3.10) \quad \text{cone}(1_{f^{-1}(-\infty, -\epsilon)} \rightarrow 1_{f^{-1}(-\infty, \epsilon)}) \in \text{Perf } \mathcal{S} = \mathcal{C}(N_\infty^*\mathcal{S}).$$

under the map $\mathcal{C}(N_\infty^*\mathcal{S}) \rightarrow \mathcal{C}(\Lambda)$, where $1_{f^{-1}(-\infty, -\epsilon)} \rightarrow 1_{f^{-1}(-\infty, \epsilon)}$ is (the linearization of) the unique map (3.5).

The Morse character $\mathcal{X}_{\Lambda,p}(f, \epsilon, \mathcal{S}) \in \mathcal{C}(\Lambda)$ depends a priori on the ‘‘casting directors’’ $(f, \epsilon, \mathcal{S})$. Casting directors (f, ϵ) exist at any smooth Legendrian point $p \in \Lambda$ by general position, and \mathcal{S} exists by the following argument. First, by [49, Prop. 8.3.10] every closed subanalytic singular isotropic $\Lambda \subseteq S^*M$ is contained in $N_\infty^*\mathcal{S}$ for some subanalytic stratification \mathcal{S} of M . Next, by refining \mathcal{S} the subanalytic subsets $f^{-1}(-\infty, \pm\epsilon)$ can be made constructible [18], [85]. Finally, \mathcal{S} can be made a subanalytic Whitney triangulation by [86], [22], [23].

Definition 3.4. A *microlocal Morse theatre* is a normalized microlocal Morse pre-theatre $\Lambda \mapsto \mathcal{C}(\Lambda)$ satisfying the *localization property*: for any inclusion $\Lambda \subseteq \Lambda'$ and any collection of Morse characters $\mathcal{X}_{\Lambda',p}(f, \epsilon, \mathcal{S}) \in \mathcal{C}(\Lambda')$ at smooth Legendrian points $p \in \Lambda' \setminus \Lambda$ with at least one in every component of the smooth Legendrian locus of $\Lambda' \setminus \Lambda$, the functor $\mathcal{C}(\Lambda') \rightarrow \mathcal{C}(\Lambda)$ is the idempotent-completed quotient by these Morse characters.

The definition of a microlocal Morse theatre allows one to readily compute any particular microlocal Morse category $\mathcal{C}(\Lambda)$: embed Λ into some $N_\infty^* \mathcal{S}$, cast Morse characters in $\mathcal{C}(N_\infty^* \mathcal{S}) = \text{Perf } \mathcal{S}$ for all Legendrian components of $N_\infty^* \mathcal{S} \setminus \Lambda$, and take the quotient of $\text{Perf } \mathcal{S}$ by these characters and idempotent complete. It follows that

PROPOSITION 3.5. *Any two microlocal Morse theatres $\Lambda \mapsto \mathcal{C}(\Lambda)$ are uniquely isomorphic.*

Proof. For any normalized microlocal Morse pre-theatre \mathcal{C} , let $\mathcal{X}_{\Lambda' \setminus \Lambda} \subseteq \mathcal{C}(\Lambda')$ denote the collection of all Morse characters at all smooth Legendrian points of $\Lambda' \setminus \Lambda$. Now for any microlocal Morse theatre \mathcal{C} , we have a canonical quasi-equivalence (functorial in Λ),

$$(3.11) \quad \varinjlim_{\substack{N^* \mathcal{S} \supseteq \Lambda \\ \mathcal{S} \text{ Whitney triangulation}}} (\mathcal{C}(N^* \mathcal{S}) / \mathcal{X}_{N^* \mathcal{S} \setminus \Lambda})^\pi \xrightarrow{\sim} \mathcal{C}(\Lambda),$$

and the left-hand side is independent of \mathcal{C} since \mathcal{C} is normalized. \square

A *dramatic realization* is a particular construction of the microlocal Morse theatre $\Lambda \mapsto \mathcal{C}(\Lambda)$. We give two dramatic realizations, namely via sheaves and via Lagrangians in [Sections 4](#) and [5](#), respectively. Both these dramatic realizations *cast* the Morse characters as certain familiar objects. They moreover show that the Morse characters in fact depend only on p (up to shifts) and are independent of the casting directors.

THEOREM 3.6. *The microlocal Morse theatre $\Lambda \mapsto \mathcal{C}(\Lambda)$ exists, and the Morse characters $\mathcal{X}_{\Lambda, p} \in \mathcal{C}(\Lambda)$ are independent of the casting directors and form a local system over the smooth Legendrian locus of Λ .*

Proof. This follows from either [Theorem 4.28](#) or [Theorem 5.36](#). \square

Proof of Theorem 1.1. Combine [Theorem 4.28](#) and [Theorem 5.36](#) with [Proposition 3.5](#). \square

In fact, both dramatic realizations show that $\mathcal{C}(\Lambda)$ is invariant under contact isotopy of $S^* M$, something which is not apparent from the present combinatorial prescription. This is immediate on the Fukaya side, and on the sheaf side it is “sheaf quantization” [\[43\]](#). In fact, there are even stronger invariance statements: it is shown in [\[38\]](#) that in fact $\mathcal{C}(\Lambda)$ is invariant under contact isotopy of $S^* M \setminus \Lambda$ inside $S^* M$; meanwhile, it is shown in [\[65\]](#) that $\mathcal{C}(\Lambda)$ is invariant under “gapped” deformations of Λ .

Remark 3.7. The construction of this subsection makes sense in any stable setting, e.g., over the sphere spectrum. To show existence of the microlocal Morse theatre in such a more general setting, one could set up either microlocal sheaf theory or the Fukaya category over the sphere spectrum. In principle, one could also show existence directly from the stratified Morse theory of [\[41\]](#),

as it already establishes results about homotopy types of spaces (not just their cohomologies). A more interesting question is whether any symplectically invariant statement can be made beyond the stable setting.

3.3. Open inclusions. We now discuss functoriality of the microlocal Morse theatre under open inclusions.

Given any analytic open inclusion of analytic manifolds $M' \hookrightarrow M$, a microlocal Morse pre-theatre \mathcal{C}' on M' determines a microlocal Morse pre-theatre $\Lambda \mapsto \mathcal{C}'(\Lambda \cap S^* M')$ on M ; let us call this the extension of \mathcal{C}' to M . For microlocal Morse pre-theatres \mathcal{C}' on M' and \mathcal{C} on M , a morphism $\mathcal{C}' \rightarrow \mathcal{C}$ means a morphism to \mathcal{C} from the extension of \mathcal{C}' to M . In other words, such a morphism consists of a coherent system of maps $\mathcal{C}'(\Lambda \cap S^* M') \rightarrow \mathcal{C}(\Lambda)$ for subanalytic singular isotropics $\Lambda \subseteq S^* M$. Equivalently, we may view such a morphism as a coherent system of maps

$$(3.12) \quad \mathcal{C}'(\Lambda') \rightarrow \mathcal{C}(\Lambda)$$

for subanalytic singular isotropics $\Lambda' \subseteq S^* M'$ and $\Lambda \subseteq S^* M$ with $\Lambda \cap S^* M' \subseteq \Lambda'$. If \mathcal{C} and \mathcal{C}' are normalized, then restricting these maps to the case that Λ and Λ' are both conormals of Whitney triangulations, we obtain a coherent collection of maps

$$(3.13) \quad \text{Perf } \mathcal{S}' = \mathcal{C}'(N_\infty^* \mathcal{S}') \rightarrow \mathcal{C}(N_\infty^* \mathcal{S}) = \text{Perf } \mathcal{S}$$

for every pair of Whitney triangulations \mathcal{S} of M and \mathcal{S}' of M' such that \mathcal{S}' refines $\mathcal{S} \cap M'$. A normalized morphism $\mathcal{C}' \rightarrow \mathcal{C}$ is one equipped with an isomorphism between (3.13) and the extension to Perf of the tautological maps of posets $\mathcal{S}' \rightarrow \mathcal{S}$ (sending a stratum of \mathcal{S}' to the unique stratum of \mathcal{S} containing it). A morphism of microlocal Morse theatres $\mathcal{C}' \rightarrow \mathcal{C}$ is, by definition, a normalized morphism of underlying normalized microlocal Morse pre-theatres.

We now have the following refinement of [Proposition 3.5](#):

PROPOSITION 3.8. *For an analytic open inclusion of analytic manifolds $M' \hookrightarrow M$ and microlocal Morse theatres \mathcal{C}' on M' and \mathcal{C} on M , there exists a unique morphism of microlocal Morse theatres $\mathcal{C}' \rightarrow \mathcal{C}$.*

Proof. It follows from (3.13) that the maps $\mathcal{C}'(\Lambda') \rightarrow \mathcal{C}(\Lambda)$ send Morse characters to Morse characters. (Choose casting directors on M which are “supported inside M' ” and appeal to the independence of Morse characters of the casting directors from [Theorem 3.6](#).) Now we have the following commutative diagram, functorial in Λ and Λ' :

$$(3.14) \quad \begin{array}{ccc} \varinjlim (\mathcal{C}(N_\infty^* \mathcal{S}') / \mathcal{X}_{N_\infty^* \mathcal{S}' \setminus \Lambda'})^\pi & \longrightarrow & \varinjlim (\mathcal{C}(N_\infty^* \mathcal{S}) / \mathcal{X}_{N_\infty^* \mathcal{S} \setminus \Lambda})^\pi \\ \downarrow \sim & & \downarrow \sim \\ \mathcal{C}'(\Lambda') & \longrightarrow & \mathcal{C}(\Lambda), \end{array}$$

where both direct limits take place over pairs of Whitney triangulations \mathcal{S} and \mathcal{S}' for which \mathcal{S}' refines $M' \cap \mathcal{S}$ and for which $\Lambda \subseteq N_\infty^* \mathcal{S}$ and $\Lambda' \subseteq N_\infty^* \mathcal{S}'$. Hence the maps $\mathcal{C}'(\Lambda') \rightarrow \mathcal{C}(\Lambda)$ are determined uniquely. \square

Proof of Proposition 1.3. Combine Propositions 4.29 and 5.37 with Proposition 3.8. \square

4. Sheaf categories

We recall the general formalism of sheaves and properties of stratifications. We then recall from [49] the notion of microsupport, and the category $\text{Sh}_\Lambda(M)$ of sheaves on M whose microsupport at infinity is contained in Λ . We show that the assignment $\Lambda \mapsto \text{Sh}_\Lambda(M)$ is a microlocal Morse theatre in the sense of Definition 3.4.

4.1. Categories of sheaves and functors between them. Here we give a brief review of the general formalism of sheaves. Our presentation is somewhat modern in that we *never* discuss sheaves of abelian groups, rather we work at the dg level and with unbounded complexes from the beginning, but it is essentially the same as any standard account such as [47], [49], [75], complemented by [87] in order to work with unbounded complexes and, in particular, for the proper base change theorem in this setting. Some discussion about working in the unbounded setting can be found in [50].

Given a topological space T , we write $\text{Op}(T)$ for the category whose objects are open sets and morphisms are inclusions. A (\mathbb{Z} -module valued) presheaf on T is by definition a functor $\text{Op}(T)^{\text{op}} \rightarrow \text{Mod } \mathbb{Z}$. In particular, a presheaf \mathcal{F} takes a value $\mathcal{F}(U) \in \text{Mod } \mathbb{Z}$ on an open set $U \subseteq T$, termed its sections; given open sets $U \subseteq V$, it gives a morphism $\mathcal{F}(V) \rightarrow \mathcal{F}(U)$, termed the restriction, etcetera. Given any subset $X \subseteq T$, we write $\mathcal{F}(X) = \varinjlim_{X \subseteq U} \mathcal{F}(U)$; when X is a point, this is termed the stalk and is written \mathcal{F}_x .

The category of sheaves is the full subcategory of presheaves on objects \mathcal{F} taking covers to limits:

$$(4.1) \quad \mathcal{F}\left(\bigcup_{i \in I} U_i\right) \xrightarrow{\sim} \varinjlim_{\emptyset \neq J \subseteq I} \mathcal{F}\left(\bigcap_{j \in J} U_j\right).$$

The inclusion of sheaves into presheaves has a left adjoint termed “sheafification,” giving, for any presheaf \mathcal{F} , a sheaf \mathcal{F}^{sh} such that any map from \mathcal{F} to a sheaf factors uniquely through \mathcal{F}^{sh} .

We write $\text{Sh}(T)$ for the (dg) category of sheaves of (dg) \mathbb{Z} -modules on T . It is complete and co-complete. Its homotopy category is what was classically called the unbounded derived category of sheaves on T .

For any continuous map $f : S \rightarrow T$, there is an adjoint pair $f^* : \text{Sh}(T) \leftrightarrow \text{Sh}(S) : f_*$. The pushforward f_* is given by the formula $(f_* \mathcal{F})(U) = \mathcal{F}(f^{-1}(U))$,

while the pullback f^* is the sheafification of the presheaf given by $(f^* \mathcal{G})(V) = \mathcal{G}(f(V))$.

Example 4.1. Consider $f : S \rightarrow \text{pt}$ and the constant sheaf $\mathbb{Z} := f^* \mathbb{Z}$. Note that in our conventions, $\mathbb{Z}(U)$ is a chain complex computing the cohomology of U . This should illustrate where, in this account of sheaf theory, the usual homological algebra of resolutions is hiding: it is in the sheafification.

Being a left adjoint, f^* is co-continuous (preserves colimits, in particular, sums). When $j : U \rightarrow T$ is the inclusion of an open set, j^* is given by the simpler formula $(j^* \mathcal{F})(V) = \mathcal{F}(V)$, no sheafification required, and hence preserves limits as well. In particular, it must also be a right adjoint. The corresponding left adjoint $j_!$ is easy to describe: it is the sheafification of

$$V \mapsto \begin{cases} \mathcal{F}(V) & V \subseteq U, \\ 0 & \text{otherwise.} \end{cases}$$

The sheaf $j_! \mathcal{F}$ is termed the extension by zero, since its stalks in U are isomorphic to the corresponding stalks of \mathcal{F} , and its stalks outside of U are zero. For a sheaf \mathcal{F} on T , we write $\mathcal{F}_U := j_! j^* \mathcal{F}$. By adjunction there is a canonical morphism $\mathcal{F}_U \rightarrow \mathcal{F}$. The object \mathbb{Z}_U co-represents the functor of sections over U , i.e., $\text{Hom}(\mathbb{Z}_U, \mathcal{F}) = \mathcal{F}(U)$.

Being a right adjoint, f_* is continuous. When f is proper, it is in addition co-continuous. More generally, for a morphism of locally compact spaces $f : S \rightarrow T$, one defines⁶

$$\begin{aligned} f_! : \text{Sh}(S) &\rightarrow \text{Sh}(T), \\ \mathcal{F} &\mapsto \varinjlim_{U \subset\subset S} f_* \mathcal{F}_U. \end{aligned}$$

Here the notation $U \subset\subset S$ means that the closure of U is compact. When S is an open subset, this recovers the original definition. When f is proper, then $f_! = f_*$. When f is the map to a point, then $f_! f^* \mathbb{Z}$ is the compactly supported cohomology.

As $f_!$ is built from colimits, left adjoints, and pushforwards from compact sets, it is co-continuous. As such it has a right adjoint, denoted $f^!$. When f is the inclusion of an open subset, we already had the right adjoint f^* , so in this case $f^* = f^!$.

For any locally closed subset $v : V \subseteq T$, we extend the notation $\mathcal{F}_V := v_! v^* \mathcal{F}$. This sheaf has the same stalks as \mathcal{F} at points in V and has vanishing stalks outside.

⁶This particular way of defining the $!$ pushforward is taken from [75]. It has the virtue of making the co-continuity of $f_!$ obvious.

For an open-closed decomposition $U \xrightarrow{j} T \xleftarrow{i} V$ (j open, i closed), the functors j_* , $j_!$ and i_* , $i_!$ are fully faithful, and there is an exact triangle

$$(4.2) \quad j_! j^! \rightarrow \text{id} \rightarrow i_* i^* \xrightarrow{[1]}.$$

Denoting by $\text{Op}(M)$ the poset of open sets, there are functors

$$(4.3) \quad \begin{aligned} \text{Op}(M) &\xrightarrow{!} \text{Sh}(M), & \text{Op}(M)^{\text{op}} &\xrightarrow{*} \text{Sh}(M), \\ U &\mapsto u_! \mathbb{Z}, & U &\mapsto u_* \mathbb{Z}, \end{aligned}$$

where $u : U \rightarrow M$ denotes the inclusion. We have the following criterion for when (pullbacks of) these functors are fully faithful:

LEMMA 4.2. *Let Π be a poset with a map to $\text{Op}(M)$, and let $\mathbb{Z}[\Pi]$ denote its dg linearization. The following are equivalent:*

- $H^*(U) \cong \mathbb{Z}$ for all $U \in \Pi$ and $H^*(U) \xrightarrow{\sim} H^*(U \setminus V)$ whenever $U \not\subseteq V$.
- The composition $\mathbb{Z}[\Pi] \rightarrow \mathbb{Z}[\text{Op}(M)] \xrightarrow{!} \text{Sh}(M)$ is fully faithful.

Proof. We have

$$(4.4) \quad \begin{aligned} \text{Hom}_M(\mathbb{Z}_U, \mathbb{Z}_V) &= \text{Hom}_M(u_! \mathbb{Z}, v_! \mathbb{Z}) = \text{Hom}_U(\mathbb{Z}, u^! v_! \mathbb{Z}) = \text{Hom}_U(\mathbb{Z}, u^* v_! \mathbb{Z}) \\ &= H^*(U, \mathbb{Z}_{V \cap U}) = \text{cone}(H^*(U) \rightarrow H^*(U \setminus V))[-1], \end{aligned}$$

where we have used the exact triangle (4.2). The second condition asks that this be \mathbb{Z} when $U \subseteq V$ and zero otherwise, which is exactly what is asserted in the first condition. \square

LEMMA 4.3. *Let Π be a poset with a map to $\text{Op}(M)$ satisfying the equivalent conditions in Lemma 4.2, and suppose that $W \subseteq M$ is an open set such that $H^*(U) \xrightarrow{\sim} H^*(U \setminus W)$ is an isomorphism whenever $U \not\subseteq W$. Then the pullback of the module $\text{Hom}(-, \mathbb{Z}_W)$ along $\mathbb{Z}[\Pi] \xrightarrow{!} \text{Sh}(M)$ is the indicator functor*

$$(4.5) \quad 1_W : U \mapsto \begin{cases} \mathbb{Z} & U \subseteq W, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. This is true by the same calculation as above. \square

4.2. *Constructible sheaves.* Let T be a topological space and $\mathcal{S} : T = \coprod T_\alpha$ a stratification. Write $T_{\mathcal{S}}$ for the topological space with underlying set T and base given by the stars of strata in \mathcal{S} . (Note that the intersection of any two stars is expressible as a union of stars.) Note the continuous map $T \rightarrow T_{\mathcal{S}}$.

Remark 4.4. Let $\pi : T \rightarrow T'$ be any continuous bijection. For any open set U of T' , and any sheaf \mathcal{F} on T , one has by definition $\pi_* \mathcal{F}(U) = \mathcal{F}(U)$. It follows that $\pi^* \mathbb{Z}_U = \mathbb{Z}_U$, as this sheaf co-represents the functor of sections over U .

LEMMA 4.5. *Pulling back sheaves under $\mathcal{S} \xrightarrow{\text{star}} \text{Op}(T_{\mathcal{S}})$ defines an equivalence*

$$(4.6) \quad \text{Sh}(T_{\mathcal{S}}) \xrightarrow{\sim} \text{Fun}(\mathcal{S}^{\text{op}}, \text{Mod } \mathbb{Z}) = \text{Mod } \mathcal{S},$$

$$(4.7) \quad \mathcal{F} \mapsto (s \mapsto \mathcal{F}(\text{star}(s))) = \text{Hom}_{T_{\mathcal{S}}}(\mathbb{Z}_{\text{star}(-)}, \mathcal{F})$$

which sends $\mathbb{Z}_{\text{star}(s)}$ to $\text{Hom}_{\mathcal{S}}(\cdot, s) = 1_{\text{star}(s)}$.

Proof. The functor in question is simply restricting a sheaf on $T_{\mathcal{S}}$ to the base consisting of stars of strata. This functor is fully faithful because a map of sheaves is determined by its restriction to a base for the topology. It is essentially surjective because there are no nontrivial covers of stars of strata by stars of strata. The behavior on objects is as asserted because $\mathbb{Z}_{\text{star}(s)}$ and s are the co-representatives of the functors of sections over s and the value of the module at s , respectively. \square

LEMMA 4.6. *If \mathcal{S}' refines \mathcal{S} , then the following diagram commutes:*

$$(4.8) \quad \begin{array}{ccc} \text{Sh}(T_{\mathcal{S}'}) & \longrightarrow & \text{Mod } \mathcal{S}', \\ \pi^* \uparrow & & \uparrow r^* \\ \text{Sh}(T_{\mathcal{S}}) & \longrightarrow & \text{Mod } \mathcal{S} \end{array}$$

where π^* denotes pullback of sheaves under the continuous map $\pi : T_{\mathcal{S}'} \rightarrow T_{\mathcal{S}}$ and $r^* : \text{Mod } \mathcal{S} \rightarrow \text{Mod } \mathcal{S}'$ denotes the pullback along the natural map $r : \mathcal{S}' \rightarrow \mathcal{S}$.

Proof. The proof follows by [Remark 4.4](#) and the characterization of the horizontal functors as $\mathbb{Z}_U \mapsto 1_U$. \square

A sheaf is said to be constant when it is isomorphic to the star pullback of a sheaf on a point, and locally constant when this is true after restriction to an open cover. For a stratification \mathcal{S} of M , we say a sheaf is \mathcal{S} -constructible⁷ if it is locally constant when star restricted to each stratum of \mathcal{S} . We write $\text{Sh}_{\mathcal{S}}(T)$ for the full subcategory of $\text{Sh}(T)$ on the \mathcal{S} -constructible sheaves.

Note that the image of the pullback map $\text{Sh}(T_{\mathcal{S}}) \rightarrow \text{Sh}(T)$ is contained in $\text{Sh}_{\mathcal{S}}(T)$.

LEMMA 4.7. *For a triangulation \mathcal{S} , the map $\text{Sh}(T_{\mathcal{S}}) \rightarrow \text{Sh}_{\mathcal{S}}(T)$ is an equivalence.*

⁷Some sources, such as [49], also ask that the word constructible should mean that sheaves should have perfect stalks and bounded cohomological degree. We do not.

Proof. To show full faithfulness, in view of the equivalence of [Lemma 4.5](#) it is enough to check that $\text{Hom}_{T_s}(\mathbb{Z}_{\text{star}(s)}, \mathbb{Z}_{\text{star}(t)}) = \text{Hom}_T(\mathbb{Z}_{\text{star}(s)}, \mathbb{Z}_{\text{star}(t)})$. The former is the indicator of $\text{star}(s) \subseteq \text{star}(t)$ again by [Lemma 4.5](#). To show that $\text{Hom}_T(\mathbb{Z}_{\text{star}(s)}, \mathbb{Z}_{\text{star}(t)})$ is as well, by [Lemma 4.2](#) it is enough to show that $H^*(\text{star}(s)) \rightarrow H^*(\text{star}(s) \setminus \text{star}(t))$ is an isomorphism for $\text{star}(s) \not\subseteq \text{star}(t)$. If $\text{star}(s) \not\subseteq \text{star}(t)$, then $\text{star}(s) \setminus \text{star}(t)$ is the join of something with s and is hence contractible.

Regarding essential surjectivity, note that the exact triangle of [\(4.2\)](#) serves to decompose any sheaf into an iterated extension of (extensions by zero of) sheaves on the strata, hence any constructible sheaf into (extensions by zero of) locally constant sheaves on the strata. Since the strata are all contractible, these sheaves are in fact constant. This shows that the \mathbb{Z}_s generate. To conclude that the $\mathbb{Z}_{\text{star}(s)}$ generate, use the exact triangle $\mathbb{Z}_{\text{star}(s) \setminus s} \rightarrow \mathbb{Z}_{\text{star}(s)} \rightarrow \mathbb{Z}_s \xrightarrow{[1]}$ and induction on dimension of strata (noting that the first term is in the span of \mathbb{Z}_t for $\dim(t) < \dim(s)$). \square

4.3. Microsupport. The notion of microsupport is developed in [\[49\]](#).⁸ We recall some basic facts here.

For what follows, let M denote an analytic manifold. Given a sheaf \mathcal{F} and a smooth function $\phi : M \rightarrow \mathbb{R}$, consider a point m in a level set $\phi^{-1}(t)$. We say that $m \in M$ is a cohomological \mathcal{F} -critical point of ϕ if, for inclusion of the superlevel set $i : \phi^{-1}(\mathbb{R}_{\geq t}) \hookrightarrow M$, one has $(i^! \mathcal{F})_m \neq 0$.

The *microsupport* $\text{ss}(\mathcal{F}) \subseteq T^*M$ is by definition the closure of the locus of differentials of functions at their cohomological \mathcal{F} -critical points [\[49\]](#). It is conical.

If \mathcal{F} is locally constant, then a cohomological \mathcal{F} -critical point can only occur where the function in question has zero derivative. Thus the microsupport of a locally constant sheaf is contained in the zero section (and is equal to it where the sheaf is not locally zero). If $U \subseteq M$ is an open set and m is a point in the smooth locus of ∂U , then over m , the locus $\text{ss}(\mathbb{Z}_U) = \text{ss}(u_! \mathbb{Z})$ is the half-line of outward conormals to ∂U . The locus $\text{ss}(u_* \mathbb{Z})$ is the inward conormal.

For a subset $X \subseteq T^*M$, we write $\text{Sh}_X(M)$ for the full subcategory of $\text{Sh}(M)$ spanned by objects with microsupport contained in X . Similarly, for $X \subseteq S^*M$, we write $\text{Sh}_X(M)$ for the full subcategory of $\text{Sh}(M)$ with microsupport at infinity contained in X . Evidently if $0_M \subseteq X$, then $\text{Sh}_X(M) = \text{Sh}_{\partial_\infty X}(M)$.

⁸In [\[49\]](#), the authors work in the bounded derived category. As noted in [\[73\]](#), the only real dependence on this was in the proof of one lemma, which is extended to the unbounded setting in that reference.

The following result is a strengthening of [49, Prop. 8.4.1]:

PROPOSITION 4.8. *For a Whitney stratification \mathcal{S} of a C^1 manifold M , we have $\mathrm{Sh}_{\mathcal{S}}(M) = \mathrm{Sh}_{N^*\mathcal{S}}(M)$ (i.e., having microsupport contained in $N^*\mathcal{S}$ is equivalent to being \mathcal{S} -constructible).*

Proof. We first show the inclusion $\mathrm{Sh}_{\mathcal{S}}(M) \subseteq \mathrm{Sh}_{N^*\mathcal{S}}(M)$. Let us first show that $\mathrm{ss}(\mathbb{Z}_X) \subseteq N^*\mathcal{S}$. When X is relatively compact, express X as the ascending union of locally closed submanifolds-with-corners $X_i = X \setminus (\partial X)^{\underline{\epsilon}_i}$ where $(\partial X)^{\underline{\epsilon}_i}$ denotes the outward cornering of $\partial X = \overline{X} \setminus X$ in the sense of (2.1) and $\underline{\epsilon}_i \rightarrow 0$. [Corollary 2.7](#) implies $X \setminus (\partial X)^{\underline{\epsilon}_i}$ are indeed locally closed submanifolds-with-corners, and [Corollary 2.8](#) implies that their conormals limit inside $N^*\mathcal{S}$ as $\underline{\epsilon}_i \rightarrow 0$, and hence that $\mathrm{ss}(\mathbb{Z}_X) = \mathrm{ss}(\varinjlim \mathbb{Z}_{X_i}) \subseteq N^*\mathcal{S}$. The case of general X may be reduced to the relatively compact case by refining the stratification as in [Remark 2.4](#) (the assertion $\mathrm{ss}(\mathbb{Z}_X) \subseteq N^*\mathcal{S}$ is local). The same argument shows that for any locally constant sheaf on X , its lower shriek pushforward to M has microsupport contained in $N^*\mathcal{S}$. Since any \mathcal{S} -constructible sheaf is (locally) a finite iterated extension of such sheaves, we conclude that $\mathrm{Sh}_{\mathcal{S}}(M) \subseteq \mathrm{Sh}_{N^*\mathcal{S}}(M)$.

We now show that the inclusion $\mathrm{Sh}_{\mathcal{S}}(M) \subseteq \mathrm{Sh}_{N^*\mathcal{S}}(M)$ implies the reverse inclusion $\mathrm{Sh}_{N^*\mathcal{S}}(M) \subseteq \mathrm{Sh}_{\mathcal{S}}(M)$ by a straightforward dévissage argument. Suppose $\mathrm{ss}(\mathcal{F}) \subseteq N^*\mathcal{S}$, and let us show that \mathcal{F} is \mathcal{S} -constructible. Let X be a maximal stratum over which \mathcal{F} is nonzero, and let $U \subseteq M$ be an open set containing X so that $X \subseteq U$ is the support of $\mathcal{F}|_U$ (hence, in particular, $X \subseteq U$ is closed). Since $\mathrm{ss}(\mathcal{F}) \subseteq N^*\mathcal{S}$, there exists a (derived) local system on X whose lower shriek pushforward \mathcal{F}_0 (which is \mathcal{S} -constructible) agrees with \mathcal{F} over U . Since $\mathcal{F}_0 \in \mathrm{Sh}_{\mathcal{S}}(M) \subseteq \mathrm{Sh}_{N^*\mathcal{S}}(M)$, it suffices to show that the cone of $\mathcal{F}_0 \rightarrow \mathcal{F}$ is \mathcal{S} -constructible. We have thus reduced to a sheaf with smaller support. Iterating, we eventually reduce to the case of $\mathcal{F} = 0$ which is obviously \mathcal{S} -constructible. \square

4.4. Microstalks. Recall that if \mathcal{F} is a sheaf and ϕ is a smooth function with $\phi(x) = t$ and $d\phi_x = \xi$, and we denote the inclusion $i : \phi^{-1}(\mathbb{R}_{\geq t}) \rightarrow M$, then if $(i^!F)_x \neq 0$, we have $(x, \xi) \in \mathrm{ss}(\mathcal{F})$ (though not conversely). Given this, one wants to assign the complex $(i^!F)_x$ itself as an invariant of \mathcal{F} at (x, ξ) . This is not generally possible, but it can be done when ξ is a point in the smooth Lagrangian locus of $\mathrm{ss}(\mathcal{F})$ [49, Prop. 7.5.3]. Namely, at any smooth Lagrangian point $(x, \xi) \in X \subseteq T^*M$, there is a “microstalk” functor

$$(4.9) \quad \mu_{(x, \xi)} : \mathrm{Sh}_X(M) \rightarrow \mathrm{Sh}(\mathrm{pt}).$$

It is given by a shift of $\mathcal{F} \mapsto (i^!F)_x$ for any ϕ with $d_x\phi = \xi$ with the graph of $d\phi$ transverse to X . The shift can be fixed using the index of the three transverse Lagrangians $(\mathrm{ss}(\mathcal{F}), T_x^*M, \Gamma_{d\phi})$. When $\xi = 0$, the microstalk functor is simply the stalk functor.

LEMMA 4.9. *The microstalk functors are co-continuous.*

Proof. Every stalk functor (i^* for i the inclusion of a point) is a left adjoint, hence is co-continuous. To show co-continuity of the microstalk at a point $(x, \xi) \in \Lambda$ with $\xi \neq 0$, argue as follows. By applying a contact transformation, we may reduce to the case that Λ is (locally near (x, ξ)) the conormal of a smooth hypersurface $N \subseteq M$. Let $B \subseteq M$ be an open ball with smooth boundary whose inward conormal at $x \in \partial B$ is ξ . Moreover, choose B so that $N^* \partial B$ and $\Lambda = N^* N$ intersect cleanly at (x, ξ) . (That is, ∂B and N are tangent at x , differing by a non-degenerate quadratic form.) Define B_- and B_+ from B by pushing ∂B inward/outward near x . Now the cone of the map $\Gamma_c(B_-, -) \rightarrow \Gamma_c(B_+, -)$ is (up to a shift) the microstalk functor $\mu_{(x, \xi)}$. The compactly supported sections functor Γ_c is co-continuous, since it is the composition of the restriction and lower shriek pushforward functors, both of which are co-continuous. \square

PROPOSITION 4.10. *Let $X \subseteq T^* M$ be closed and conical, and let $\Lambda \subseteq T^* M \setminus X$ be closed, conical, and stratified by isotropic submanifolds. Then $\mathrm{Sh}_X(M) \subseteq \mathrm{Sh}_{X \cup \Lambda}(M)$ is the kernel of all microstalks at Lagrangian points of Λ .*

Proof. If $\mathrm{ss}(\mathcal{F}) \subseteq X$, then the microstalks of \mathcal{F} at Lagrangian points of Λ vanish by definition of microsupport. To prove the converse, suppose that $\mathrm{ss}(\mathcal{F}) \subseteq X \cup \Lambda$ and that the microstalks of \mathcal{F} vanish at all Lagrangian points of Λ , and let us show that $\mathrm{ss}(\mathcal{F}) \subseteq X$. By the fundamental result [49, Thm. 6.5.4] that the microsupport is co-isotropic, it is enough to show that $p \notin \mathrm{ss}(\mathcal{F})$ for every Lagrangian point $p \in \Lambda$. It is not quite immediate from the definitions that vanishing of the microstalk implies there is no microsupport, since the microsupport is defined in terms of arbitrary test functions, whereas microstalks are defined in terms of microlocally transverse test functions. To see it is true, and that moreover the microstalk is locally constant along Λ , one can apply a contact transformation so that Λ becomes locally the conormal to a smooth hypersurface; for details, see [49, Chap. 7]. \square

It will be central to our discussion to find co-representatives of the microstalk functors. Here is a first step:

THEOREM 4.11 ([49, Cor. 5.4.19, Props. 5.4.20 and 7.5.3] or [41], [76]). *Let $X \subseteq T^* M$ be a closed conical subset, let $\phi : M \rightarrow \mathbb{R}$ be a proper function, and assume that over $\phi^{-1}([a, b])$, one has $\Gamma_{d\phi} \cap X = (x, \xi)$, where (x, ξ) is a smooth Lagrangian point of X .*

Let $A : \phi^{-1}((-\infty, a)) \rightarrow M$, $A' : \phi^{-1}((a, \infty)) \rightarrow M$, $B : \phi^{-1}((-\infty, b)) \rightarrow M$, and $B' : \phi^{-1}((b, \infty)) \rightarrow M$ be the inclusions. Then (up to a shift), the following functors $\mathrm{Sh}_X(M) \rightarrow \mathrm{Sh}(\mathrm{pt})$ are isomorphic:

- the microstalk functor $\mu_{(x,\xi)}$;
- $\text{Hom}(\text{cone}(A_! \mathbb{Z} \rightarrow B_! \mathbb{Z}), -)$;
- $\text{Hom}(\text{cone}(A'_* \mathbb{Z} \rightarrow B'_* \mathbb{Z}), -)$.

Here the maps are the canonical ones coming from restriction of sections.

We do not say that $\text{cone}(A_! \mathbb{Z} \rightarrow B_! \mathbb{Z})$ co-represents the microstalk because it is not an element of $\text{Sh}_X(M)$. As observed in [63], such co-representatives do exist, for categorical reasons, as we now explain. First, we need to know that the categories in question are well generated in the sense of Neeman [69, 52].

LEMMA 4.12. *The category $\text{Sh}_X(M)$ is well generated.*

Proof. The category of all sheaves $\text{Sh}(M)$ is the derived category of a Grothendieck abelian category, hence is well generated [68]. A sheaf \mathcal{F} having singular support inside X is equivalent to the restriction maps $\mathcal{F}(U_\alpha) \rightarrow \mathcal{F}(V_\alpha)$ being isomorphisms for some list of pairs $(U_\alpha, V_\alpha)_\alpha$ depending on X . This condition is equivalent to \mathcal{F} being right-orthogonal to the cone of the map $\mathbb{Z}_{V_\alpha} \rightarrow \mathbb{Z}_{U_\alpha}$. Now the right-orthogonal complement of a set of objects in a well generated category is well generated [69], [72, Thm. 4.9]. \square

Now note that the microsupport of a sum or product is contained in the closure of the union of the microsupports. Thus if X is closed, then the subcategory $\text{Sh}_X(M) \subseteq \text{Sh}(M)$ is closed under sums and products. In particular, $\text{Sh}_X(M)$ is complete and co-complete, and the inclusion $\text{Sh}_X(M) \rightarrow \text{Sh}(M)$ is continuous and co-continuous. More generally, if $X \subseteq X'$ are closed, then the inclusion $\iota : \text{Sh}_X(M) \rightarrow \text{Sh}_{X'}(M)$ is continuous and co-continuous. It follows that

LEMMA 4.13. *For closed $X \subseteq X' \subseteq T^*M$, the inclusion $\iota : \text{Sh}_X(M) \rightarrow \text{Sh}_{X'}(M)$ has both adjoints: $(\iota^*, \iota, \iota^!)$.*

Proof. Since $\text{Sh}_X(M)$ is well generated and ι is co-continuous, it has a right adjoint $\iota^!$ by Brown representability for well generated categories [69].

For the left adjoint ι^* , it would be sufficient to know Brown representability for the opposite of $\text{Sh}_X(M)$. However according to Neeman [70], it is an open problem to establish Brown representability for the opposites of well-generated categories. Instead, we may argue as follows. The categories in question are presentable (co-complete and accessible [58, Def. 5.4.2.1 and 5.5.0.1]; this is a version of well generation), and so a functor has a left adjoint if and only if it is continuous and accessible (preserves κ -filtered colimits [58, Def. 5.4.2.5 and 5.3.4.5]) by Lurie [58, Corollary 5.5.2.9].

Another proof of the existence of the left adjoint ι^* has been given by Efimov [28]. \square

For example, if $V \subseteq M$ is a closed subset, then taking $X = T^*M|_V$ and $X' = T^*M$ recovers the adjoint triple for the pushforward along $V \rightarrow M$, because $\mathrm{Sh}_{T^*M|_V}(M) = \mathrm{Sh}(V)$.

Using the left adjoint and [Theorem 4.11](#), we can obtain a co-representative for the microstalk as follows. Take any $X' \supseteq \mathrm{ss}(\mathrm{cone}(A_! \mathbb{Z} \rightarrow B_! \mathbb{Z}))$, e.g., $X' = T^*M$. Then $\iota^* \mathrm{cone}(A_! \mathbb{Z} \rightarrow B_! \mathbb{Z}) \in \mathrm{Sh}_X(M)$ co-represents the microstalk.

We do not generally have a good understanding of $(\iota^*, \iota, \iota^!)$, but when $X' \setminus X$ is isotropic we have the following (special cases of which have appeared in [\[63\]](#), [\[46\]](#)):

THEOREM 4.14. *Let $X \subseteq T^*M$ be closed and conical, and let $\Lambda \subseteq T^*M \setminus X$ be closed, conical, and stratified by isotropic submanifolds. The left adjoint ι^* to the inclusion $\iota : \mathrm{Sh}_X(M) \rightarrow \mathrm{Sh}_{X \cup \Lambda}(M)$ realizes the quotient*

$$(4.10) \quad \mathrm{Sh}_{X \cup \Lambda}(M)/\mathcal{D} \xrightarrow{\sim} \mathrm{Sh}_X(M),$$

where \mathcal{D} denotes co-representing objects for the microstalks at Lagrangian points of Λ .

Proof. According to [Proposition 4.10](#), the full subcategory $\mathrm{Sh}_X(M) \subseteq \mathrm{Sh}_{X \cup \Lambda}(M)$ is precisely the right-orthogonal to \mathcal{D} . The left adjoint ι^* to the inclusion ι is thus termed the quotient by \mathcal{D} . \square

Remark 4.15. For our purposes in this paper, we do not need [Lemma 4.13](#) and [Theorem 4.14](#) in their general formulations given above, rather only in the special case of subanalytic singular isotropic singular supports. In this setting, we give an elementary derivation (i.e., without appealing to general Brown representability type statements) in the next subsection.

4.5. Compact objects. Here we elaborate upon some assertions of [\[63\]](#).

We write $\mathrm{Sh}_X(M)^c$ for the compact objects in the category $\mathrm{Sh}_X(M)$. Be warned:

PROPOSITION 4.16 ([68]). *When M is connected and non-compact,*

$$\mathrm{Sh}(M)^c = 0.$$

There are not many more compact objects in the compact case. However, for sheaves with prescribed isotropic microsupport, the situation is different:

PROPOSITION 4.17. *For $\Lambda \subseteq T^*M$ a conic subset Whitney stratifiable by isotropics, the category $\mathrm{Sh}_\Lambda(M)$ is compactly generated by the (co-representatives of the) microstalk functors at the smooth Lagrangian points of Λ .*

Proof. It was shown immediately after the proof of [Lemma 4.13](#) that the microstalk functors at smooth Lagrangian points of Λ are co-represented by objects of $\mathrm{Sh}_\Lambda(M)$. Since the microstalk functors are co-continuous ([Lemma 4.9](#)), these co-representatives are compact. Any sheaf right-orthogonal to these

co-representatives has by definition vanishing microstalks at all smooth Lagrangian points of Λ , hence has microsupport contained in the complement of the smooth Lagrangian locus (see the proof of [Proposition 4.10](#)). This complement, being stratified by subcritical isotropics, has no co-isotropic subset; hence by the involutivity of microsupports [[49, Thm. 6.5.4](#)], the microsupport is in fact the empty set and the sheaf vanishes. \square

As promised in [Remark 4.15](#), we now give arguments avoiding the use of representability theorems in non-compactly-generated categories. This comes at the cost of assuming subanalyticity in order to ensure the existence of triangulations, but for the main results we will anyway need this hypothesis.

LEMMA 4.18. *For \mathcal{S} a triangulation, the category $\mathrm{Sh}_{\mathcal{S}}(M)$ is compactly generated, and the objects of $\mathrm{Sh}_{\mathcal{S}}(M)^c$ are the sheaves with perfect stalks and compact support.*

Proof. Under the identification ([Lemma 4.7](#)) $\mathrm{Sh}_{\mathcal{S}}(M) = \mathrm{Mod} \mathcal{S}$, the $\mathbb{Z}_{\mathrm{star}(s)}$ go to compact generators. The dévissage in the proof of the same lemma shows that \mathbb{Z}_s also generate, and can be expressed using finitely many $\mathbb{Z}_{\mathrm{star}(s)}$, hence are compact. The \mathbb{Z}_s evidently generate the sheaves with perfect stalks and compact support. \square

Remark 4.19. Note that while a non-compact manifold does not admit a finite triangulation, it can sometimes be a relatively compact constructible subset of a larger manifold.

Recall that $\mathrm{Sh}_{\mathcal{S}}(M) = \mathrm{Sh}_{N^*\mathcal{S}}(M)$ for any Whitney stratification by [Proposition 4.8](#).

PROPOSITION 4.20. *For any subanalytic Whitney triangulation \mathcal{S} , the category $\mathrm{Sh}_{\mathcal{S}}(M)$ is compactly generated by co-representatives of the microstalks at smooth points of $N^*\mathcal{S}$.*

Proof. Consider the microstalk at some smooth point $(x, \xi) \in N^*\mathcal{S}$. It is possible to choose real analytic ϕ as in [Theorem 4.11](#); see [[41, Thm. 2.2.1](#)] or [[49, Prop. 8.3.12](#)]. We keep the notation of [Theorem 4.11](#). Refine \mathcal{S} to a subanalytic Whitney triangulation \mathcal{S}' for which $A_!\mathbb{Z}$ and $B_!\mathbb{Z}$ are constructible.

By [Lemma 4.18](#), $\mathrm{cone}(A_!\mathbb{Z} \rightarrow B_!\mathbb{Z})$ is a compact object in $\mathrm{Sh}_{\mathcal{S}'}(M)$. [Lemmas 4.5](#) and [4.7](#) give $\mathrm{Sh}_{\mathcal{S}}(M) = \mathrm{Mod} \mathcal{S}$, and [Lemma 4.6](#) states that the inclusion $\iota : \mathrm{Sh}_{\mathcal{S}}(M) \rightarrow \mathrm{Sh}_{\mathcal{S}'}(M)$ corresponds to the map $r^* : \mathrm{Mod} \mathcal{S} \rightarrow \mathrm{Mod} \mathcal{S}'$ from [Section 3.1](#). It was observed in [Section 3.1](#) that r^* has a left adjoint $r_! : \mathrm{Mod} \mathcal{S}' \rightarrow \mathrm{Mod} \mathcal{S}$, thus giving us a left adjoint $\iota^* : \mathrm{Sh}_{\mathcal{S}'}(M) \rightarrow \mathrm{Sh}_{\mathcal{S}}(M)$. These left adjoints $r_!/\iota^*$ preserve compact objects since r^*/ι are co-continuous. Thus the object $\iota^* \mathrm{cone}(A_!\mathbb{Z} \rightarrow B_!\mathbb{Z}) \in \mathrm{Sh}_{\mathcal{S}}(M)$, which co-represents the desired microstalk, is compact.

Co-representatives of the microstalks at all smooth points of $N^*\mathcal{S}$ generate $\mathrm{Sh}_{\mathcal{S}}(M)$ by [Proposition 4.10](#). \square

Remark 4.21. A similar argument shows that the stalk at any point of M (not necessarily a smooth point of N^*S) is co-representable by a compact object of $\text{Sh}_S(M)$. Indeed, note that for any $x \in M$, the functor of taking stalks at x , which is by definition $\mathcal{F}_x := \varinjlim \mathcal{F}(B_\epsilon(x))$, is in fact computed by some fixed $\mathcal{F}_x = \mathcal{F}(B_{\epsilon_x}(x))$. Indeed, further shrinking of the ball will be non-characteristic with respect to N^*S , as follows from Whitney's condition (b) (or alternatively from microlocal Bertini–Sard [49, Prop. 8.3.12]). We may now argue as above, choosing any analytic function with sublevelset $B_{\epsilon_x}(x)$.

COROLLARY 4.22. *For any closed conical subanalytic isotropic $\Lambda \subseteq T^*M$, the category $\text{Sh}_\Lambda(M)$ is compactly generated by co-representatives of the microstalks at smooth points of Λ .*

Proof. Fix a subanalytic Whitney triangulation S for which $\Lambda \subseteq N^*S$. Denote by $\mathcal{D} \subseteq \text{Sh}_{N^*S}(M)^c$ the co-representatives of the microstalks at smooth points of $N^*S \setminus \Lambda$. By [Proposition 4.10](#), $\iota : \text{Sh}_\Lambda(M) \subseteq \text{Sh}_{N^*S}(M)$ is precisely the inclusion of the right-orthogonal to \mathcal{D} . Since the objects of \mathcal{D} are compact by [Proposition 4.20](#), [Lemma A.7](#) applies to show that $\text{Sh}_\Lambda(M) = \text{Sh}_{N^*S}(M)/\mathcal{D}$ is compactly generated by $\text{Sh}_\Lambda(M)^c = (\text{Sh}_{N^*S}(M)^c/\mathcal{D})^\pi$ and that the resulting functor $\iota^* : \text{Sh}_{N^*S}(M) \rightarrow \text{Sh}_\Lambda(M)$ is left adjoint to ι . \square

COROLLARY 4.23. *Let $X \subseteq T^*M$ and $\Lambda \subseteq T^*M \setminus X$ be closed conical subanalytic isotropics. The inclusion $\iota : \text{Sh}_X(M) \rightarrow \text{Sh}_{X \cup \Lambda}(M)$ has a left adjoint $\iota^* : \text{Sh}_{X \cup \Lambda}(M) \rightarrow \text{Sh}_X(M)$ whose restriction to compact objects defines an equivalence*

$$(4.11) \quad (\text{Sh}_{X \cup \Lambda}(M)^c/\mathcal{D})^\pi \xrightarrow{\sim} \text{Sh}_X(M)^c,$$

where \mathcal{D} denotes co-representing objects for the microstalks at Lagrangian points of Λ .

Proof. By [Corollary 4.22](#), $\text{Sh}_{X \cup \Lambda}(M)$ is compactly generated by the microstalks at smooth points of $X \cup \Lambda$. Now argue as in the proof of [Corollary 4.22](#). \square

The following result was shown in [63] using arborealization; here is a direct argument.

COROLLARY 4.24. *The Yoneda embedding induces an equivalence between the full subcategory of $\text{Sh}_\Lambda(M)$ of objects with perfect stalks and the category $\text{Prop Sh}_\Lambda(M)^c$.*

Proof. From the argument in [Proposition 4.20](#), we see that the microstalks are calculated by comparing sections over precompact sets; it follows that a sheaf microsupported in Λ (thus constructible) with perfect stalks has perfect microstalks. The microstalk functors split-generate $\text{Sh}_\Lambda(M)^c$ by [Corollary 4.22](#), so we see that a sheaf with perfect stalks defines a proper module over $\text{Sh}_\Lambda(M)^c$.

To see the converse, recall from [Remark 4.21](#) that the stalk functors can be expressed in terms of sections over open sets constructible with respect to some \mathcal{S} satisfying $N^*\mathcal{S} \supseteq \Lambda$. The left adjoint to $\mathrm{Sh}_\Lambda(M) \hookrightarrow \mathrm{Sh}_{\mathcal{S}}(M)$ preserves compact objects as observed previously, hence proper over $\mathrm{Sh}_\Lambda(M)^c$ implies perfect stalks. \square

For compact M , we establish smoothness and/or properness for some of these categories.

PROPOSITION 4.25. *If M is compact and \mathcal{S} is a triangulation, then $\mathrm{Sh}_{\mathcal{S}}(M)^c$ is smooth and proper.*

Proof. The $\mathbb{Z}_{\mathrm{star}(s)}$ give a finite generating exceptional collection which is proper, and this implies smoothness by [Lemma A.11](#). \square

More generally,

COROLLARY 4.26. *If M is compact and Λ is closed conical subanalytic singular isotropic, then $\mathrm{Sh}_\Lambda(M)^c$ is smooth, and hence*

$$\mathrm{Prop} \, \mathrm{Sh}_\Lambda(M)^c \subseteq \mathrm{Perf} \, \mathrm{Sh}_\Lambda(M)^c$$

and $\mathrm{Prop} \, \mathrm{Sh}_\Lambda(M)^c$ is proper.

Proof. By [Proposition 4.25](#) and [Corollary 4.23](#), the category $\mathrm{Sh}_\Lambda(M)^c$ is a quotient of a smooth category, hence smooth ([Lemma A.9](#)). Smoothness implies proper modules are perfect ([Lemma A.8](#)) and that the category of proper modules is proper. \square

Remark 4.27. When (M, Λ) are non-compact but finite-type in a suitable sense, the same result is true. One can prove it by embedding into a compact manifold as in [Remark 4.19](#).

4.6. *In conclusion.* Collecting the results of this section, we have shown

THEOREM 4.28. *The functor $\Lambda \mapsto \mathrm{Sh}_\Lambda(M)^c$ is a microlocal Morse theatre in the sense of [Definition 3.4](#), which casts the co-representatives of the microstalk functors at smooth points of Λ as the Morse characters.*

Proof. The most obvious functor $\Lambda \rightarrow \mathrm{Sh}_\Lambda(M)$ is the one which carries *inclusions* $\Lambda \subseteq \Lambda'$ to *inclusions* $\mathrm{Sh}_\Lambda(M) \hookrightarrow \mathrm{Sh}_{\Lambda'}(M)$; note that this is in fact a strict diagram of categories (as all are simply full subcategories of $\mathrm{Sh}(M)$) and takes values in the category whose objects are large dg categories and whose morphisms are continuous and co-continuous functors. Passing to left adjoints and taking compact objects (see [Corollary 4.23](#)), we obtain a microlocal Morse pre-theatre $\Lambda \mapsto \mathrm{Sh}_\Lambda(M)^c$.

For triangulations \mathcal{S} , the functors

$$(4.12) \quad \mathcal{S} \xrightarrow{s \mapsto \mathbb{Z}_{\text{star}(s)}} \text{Sh}_{\mathcal{S}}(M) \xrightarrow{\mathcal{F} \mapsto \text{Hom}(\mathbb{Z}_{\text{star}(-)}, \mathcal{F})} \text{Mod } \mathcal{S}$$

define an equivalence $\text{Perf } \mathcal{S} = \text{Sh}_{\mathcal{S}}(M)^c$ by [Lemmas 4.5](#) and [4.7](#). When \mathcal{S} is a Whitney stratification, we have $\text{Sh}_{\mathcal{S}}(M)^c = \text{Sh}_{N^* \mathcal{S}}(M)^c$ by [Proposition 4.8](#).

Taking the commutative diagram in [Lemma 4.6](#) and passing to the left adjoints of the vertical maps shows that this equivalence respects refinement of Whitney triangulations. This shows that $\Lambda \mapsto \text{Sh}_{\Lambda}(M)^c$ is normalized.

By [Theorem 4.11](#), the Morse characters in $\text{Perf } \mathcal{S}$ correspond, under this isomorphism, to co-representatives of the microstalks. According to [Corollary 4.23](#), the functor $\text{Sh}_{\Lambda'}(M)^c \rightarrow \text{Sh}_{\Lambda}(M)^c$ is the quotient by co-representatives of the microstalks. Thus $\Lambda \mapsto \text{Sh}_{\Lambda}(M)^c$ satisfies the localization property and is thus a microlocal Morse theatre. \square

PROPOSITION 4.29. *For any analytic open inclusion of analytic manifolds $M' \hookrightarrow M$, the restriction functors $\text{Sh}_{\Lambda}(M) \rightarrow \text{Sh}_{\Lambda'}(M')$ for subanalytic singular isotropics with $\Lambda' \supseteq \Lambda \cap S^* M'$ have left adjoints whose restrictions to compact objects form a morphism of microlocal Morse theatres.*

Proof. The categories $\text{Sh}_{\Lambda}(M)$ are compactly generated by [Corollary 4.22](#), and Brown representability holds for the opposites of compactly generated categories by [\[69\]](#), [\[53\]](#). Thus since the restriction functors $\text{Sh}_{\Lambda}(M) \rightarrow \text{Sh}_{\Lambda'}(M')$ are continuous, they admit left adjoints. Since restriction is co-continuous, these left adjoints preserve compact objects. Restricting these left adjoints to compact objects defines a morphism of microlocal Morse pre-theatres in the sense of [Section 3.3](#).

Let us show that this is a morphism of microlocal Morse theatres, i.e., that it is normalized. For a stratification \mathcal{S} of M and a stratification \mathcal{S}' refining $\mathcal{S} \cap M'$, we have the following commutative diagram:

$$(4.13) \quad \begin{array}{ccccc} \text{Sh}_{\mathcal{S}'}(M') & \longleftarrow & \text{Sh}(M'_{\mathcal{S}'}) & \xrightarrow{\sim} & \text{Mod } \mathcal{S}' \\ \uparrow & & \uparrow & & \uparrow \\ \text{Sh}_{\mathcal{S}}(M) & \longleftarrow & \text{Sh}(M_{\mathcal{S}}) & \xrightarrow{\sim} & \text{Mod } \mathcal{S} \end{array}$$

(compare [Lemmas 4.5](#) and [4.6](#)). When \mathcal{S} and \mathcal{S}' are triangulations, the left horizontal maps are also equivalences by [Lemma 4.7](#). Finally, when \mathcal{S} is Whitney, we have $\text{Sh}_{\mathcal{S}}(M) = \text{Sh}_{N^* \mathcal{S}}(M)$ (and the same for \mathcal{S}') by [Proposition 4.8](#). Thus passing to left adjoints of the vertical arrows and restricting to compact objects, we conclude. \square

5. Wrapped Fukaya categories

5.1. *Wrapped Floer cohomology.* Here we quickly fix notation and review basic facts (see, e.g., [37, §3] for more details). Fix a Liouville manifold or open Liouville sector X .

For a pair of exact Lagrangians $L, K \subseteq X$, conical and disjoint at infinity, we write $HF^*(L, K)$ for their Floer cohomology. We write $HF^*(L, L)$ to mean $HF^*(L^+, L)$, where L^+ denotes an (unspecified) small positive (meaning positive at infinity) pushoff of L . There is an isomorphism of groups $HF^*(L, L) = H^*(L)$, and the group $HF^*(L, L) = HF^*(L^+, L)$ is a unital algebra;⁹ its unit is termed the *continuation element*. Composition of continuation elements associated to small pushoffs defines more generally a continuation element in $HF^*(L^{++}, L)$ for L^{++} any (not necessarily small) positive wrapping (i.e., isotopy) of L . Composition with the continuation element associated to $L \rightsquigarrow L^{++}$ gives maps $HF^*(L, K) \rightarrow HF^*(L^{++}, K)$ and $HF^*(K, L^{++}) \rightarrow HF^*(K, L)$ for any K disjoint at infinity from L and L^{++} , which are termed *continuation maps*. If the entire positive isotopy $L \rightsquigarrow L^{++}$ takes place in the complement of $\partial_\infty K$, then these continuation maps are isomorphisms. More generally, if $L \rightsquigarrow L'$ is any isotopy taking place in the complement of $\partial_\infty K$ (for example, any compactly supported isotopy), then there is an induced identification $HF^*(L, K) = HF^*(L', K)$ (see [37, Lem. 3.21]) which coincides with the continuation isomorphism if $L \rightsquigarrow L'$ is positive at infinity (see [37, Lem. 3.26]). In particular, seeing as $HF^*(L, K) = 0$ tautologically when K and L are disjoint, Floer cohomology $HF^*(L, K)$ vanishes whenever L is disjoinable from K by an isotopy in the complement of $\partial_\infty K$.

The wrapped Floer cohomology $HW^*(L, K)_X$ is equivalently calculated by (5.1)

$$\varinjlim_{L \rightsquigarrow L^{++}} HF^*(L^{++}, K) = \varinjlim_{\substack{L \rightsquigarrow L^{++} \\ K^{--} \rightsquigarrow K}} HF^*(L^{++}, K^{--}) = \varinjlim_{K^{--} \rightsquigarrow K} HF^*(L, K^{--}).$$

Here, the direct limits are taken using the continuation maps over positive-at-infinity isotopies of L and negative-at-infinity isotopies of K . The freedom to wrap in only one factor is extremely useful in practice.

Given any closed subset $\Lambda \subseteq \partial_\infty X$, and L, K disjoint at infinity from Λ , we similarly define partially wrapped Floer cohomology $HW^*(L, K)_{X, \Lambda}$ by restricting wrappings to take place in the complement of Λ .

The following lemma allows one to explicitly describe some cofinal wrapping sequences in a given (X, Λ) . Its typical use is the following. To compute $HW^*(L, K)_{X, \Lambda}$, if one can find a cofinal sequence L_t such that the induced

⁹One expects (as is known for compact L) that the isomorphism $HF^*(L, L) = H^*(L)$ is further compatible with algebra structures; we are not aware of a reference for this.

maps $HF^*(L_t, K) \rightarrow HF^*(L_{t+1}, K)$ are eventually all isomorphisms, then $HW^*(L, K)_{X, \Lambda} = HF^*(L_t, K)$ for any L_t in this stable range.

LEMMA 5.1 ([37, Lem. 3.29] [38, Lem. 2.2]). *Let L_t be a positive isotopy of Lagrangians in X avoiding Λ at infinity. If $\partial_\infty L_t$ escapes to infinity in (i.e., is eventually disjoint from any given compact subset of) $\partial_\infty X \setminus \Lambda$ as $t \rightarrow \infty$, then it is a cofinal wrapping of L_0 in (X, Λ) .*

5.2. *Wrapped Fukaya categories.* In [37], [38], for any Liouville sector X and any closed subset $\Lambda \subseteq (\partial_\infty X)^\circ$, we constructed A_∞ categories $\mathcal{W}(X, \Lambda)$ whose objects are exact Lagrangians in $X \setminus \Lambda$, conical at infinity (by convention, $\mathcal{W}(X) := \mathcal{W}(X, \emptyset)$). The cohomology-level morphisms are simply the wrapped Floer cohomology groups as defined above: $H^* \mathcal{W}(L, K) = HW^*(L, K)_{X, \Lambda}$. For a compact manifold-with-boundary M , its cotangent bundle T^*M is a Liouville sector [37, Ex. 2.7].

One main point of [37] was the construction of a covariant functor $\mathcal{W}(X) \rightarrow \mathcal{W}(Y)$ for an inclusion of Liouville sectors $X \subseteq Y$. In [38] we remarked that the same construction gives a functor $\mathcal{W}(X, \Lambda \cap (\partial_\infty X)^\circ) \rightarrow \mathcal{W}(Y, \Lambda)$. This covariance is a nontrivial result having to do with the fact that holomorphic disks can be made to not cross the boundary of a Liouville sector (if the Lagrangian boundary conditions do not). By contrast, it is immediate from the definition that if $\Lambda \subseteq \Lambda'$, then there is a natural map $\mathcal{W}(X, \Lambda') \rightarrow \mathcal{W}(X, \Lambda)$: just wrap more. Both covariance statements allow one to calculate in a potentially simpler geometry and push forward the result.

Here we wish to consider categories $\mathcal{W}(T^*M, \Lambda)$ for (possibly non-compact) manifolds M without boundary and closed subsets $\Lambda \subseteq S^*M = \partial_\infty T^*M$. Such a cotangent bundle T^*M is an *open Liouville sector* in the sense of [37, Rem. 2.8] (meaning, concretely, it admits an exhaustion by Liouville sectors, in this case $T^*M_0 \subseteq T^*M_1 \subseteq \dots$, where $M_0 \subseteq M_1 \subseteq \dots$ is an exhaustion of M by compact codimension zero submanifolds-with-boundary).

The construction of the wrapped Fukaya category of an open Liouville sector is given in [37, §3.8]. The generalization to the case with a stop following [38, §2] is straightforward. The result is the following definition. We consider tuples $(P, \{M_p\}_{p \in P}, \{L_p\}_{p \in P}, \underline{J}, \underline{\xi})$ where

- (i) P is a partially ordered set;
- (ii) each $M_p \subseteq M$ is a compact codimension zero submanifold with smooth boundary, equipped with a choice of projection from T^*M_p to $\mathbb{C}_{\text{Re} \geq 0}$ as in [37, Def. 2.26] defined near the boundary;
- (iii) each $L_p \subseteq T^*M_p$ is an exact Lagrangian, cylindrical at infinity, disjoint from Λ at infinity, equipped with grading and orientation data as in Section 5.3 below, such that for every totally ordered subset $p_0 > \dots > p_k \in P$, the Lagrangians L_{p_0}, \dots, L_{p_k} are mutually transverse;

- (iv) the pair $(\underline{\xi}, \underline{J})$ is a choice of compatible Floer data (strip-like coordinates and almost complex structures) as in [37, Eq. (3.31)–(3.33)] for every totally ordered subset $p_0 > \dots > p_k \in P$ (so J_{p_0, \dots, p_k} is an almost complex structure on $T^*M_{p_0}$), such that all moduli spaces of Fukaya A_∞ disks are cut out transversally.

Any such tuple gives rise to a A_∞ category whose objects are the elements of P , whose morphism spaces from p to p' are $CF^*(L_p, L_{p'})$ for $p > p'$, are \mathbb{Z} for $p = p'$, and otherwise vanish. We may ask that such a tuple be *cofinite* (meaning $P^{\leq p}$ is finite for every $p \in P$) and *duplicate-free* (meaning $P^{\leq p}$ equipped with the restriction of the remaining data are pairwise non-isomorphic for $p \in P$). There is a *universal* cofinite duplicate-free tuple $(P, \{M_p\}_{p \in P}, \{L_p\}_{p \in P}, \underline{J}, \underline{\xi})$ [37, Lem. 3.42], which thus gives a canonically defined A_∞ category $\mathcal{O}(T^*M, \Lambda)$. The wrapped category $\mathcal{W}(T^*M, \Lambda)$ is defined as the localization $\mathcal{O}(T^*M, \Lambda)[C^{-1}]$ (refer to [37, §3.1.3] for localizations of A_∞ categories) at the class C of all continuation elements in $HF^0(L_p, L_{p'})$ for positive isotopies $L_{p'} \rightsquigarrow L_p$ inside T^*M_p disjoint at infinity from Λ . That this category deserves the name $\mathcal{W}(T^*M, \Lambda)$ is justified by [37, Prop. 3.43, Prop. 3.39, Lem. 3.37] and [38, §2]; in particular, these show that it has the correct cohomology category.

The resulting category $\mathcal{W}(T^*M, \Lambda)$ is moreover strictly functorial in M and Λ : for any open inclusion of manifolds $M \hookrightarrow M'$ such that Λ contains the inverse image of Λ' , there is an induced functor $\mathcal{W}(T^*M, \Lambda) \rightarrow \mathcal{W}(T^*M', \Lambda')$, and these functors respect compositions of inclusions $M \hookrightarrow M' \hookrightarrow M''$.

5.3. Gradings and orientations. We briefly review the setup for defining gradings and orientations in Floer theory; for more details, see Seidel [77] and [79, (11e)–(11l)]. Our Floer cohomology groups and Fukaya categories are all \mathbb{Z} -graded and with \mathbb{Z} coefficients.

Denote by $\text{LGr}(V)$ the Grassmannian of Lagrangian subspaces of a given symplectic vector space V . A map $\partial D^2 = S^1 \rightarrow \text{LGr}(V)$ defines elliptic boundary conditions for the $\bar{\partial}$ -operator on the trivial vector bundle with fiber V over D^2 (choosing also a compatible complex structure on V , which is a contractible choice), and hence a virtual vector space, namely the index (kernel minus cokernel) of this operator, thus giving a map

$$(5.2) \quad \mathcal{L}(\text{LGr}(V)) \rightarrow \mathbb{Z} \times BO.$$

Identifying $U/O = \varinjlim_n \text{LGr}(\mathbb{C}^n)$ and restricting to the based loop space, the resulting map $\Omega(U/O) \rightarrow \mathbb{Z} \times BO$ is (almost [79, Rmk. 11.8]) the Bott periodicity homotopy equivalence. For Floer theory with \mathbb{Z} -grading and \mathbb{Z} coefficients, we care just about the dimension and orientation, i.e., we compose the above map with $(\text{id}, w_1) : \mathbb{Z} \times BO \rightarrow \mathbb{Z} \times K(\mathbb{Z}/2, 1)$. Now restricting to the based loop space $\Omega \text{LGr}(V)$ and applying B , we obtain cohomology

classes on $\mathrm{LGr}(V)$, which are (see the related [79, Lem. 11.7] or [9, Prop. 4.2.8]) the Maslov class $\mu \in H^1(\mathrm{LGr}(V), \mathbb{Z})$ and $w_2 \in H^2(\mathrm{LGr}(V), \mathbb{Z}/2)$ (the second Stiefel–Whitney class of the tautological bundle $L \rightarrow \mathrm{LGr}(V)$). The class w_2 is represented by a map $\mathrm{LGr}(V) \rightarrow K(\mathbb{Z}/2, 2)$ given by the pullback of $w_2 : BO \rightarrow K(\mathbb{Z}/2, 2)$ under the map $\mathrm{LGr}(V) \rightarrow BO$ (which is *well defined up to contractible choice*) classifying the tautological bundle. In contrast, the map $\mathrm{LGr}(V) \rightarrow K(\mathbb{Z}, 1) = S^1$ classified by the Maslov class μ is *not* well defined up to contractible choice. Rather, given a compatible complex structure on V (a contractible choice), the Maslov class is represented by the canonical map

$$(5.3) \quad \mathrm{LGr}(V) \rightarrow ((\wedge_{\mathbb{C}}^{\mathrm{top}} V)^{\otimes 2} \setminus 0)/\mathbb{R}_{>0}$$

given by the composition of $\wedge_{\mathbb{R}}^{\mathrm{top}} : \mathrm{LGr}(V) \rightarrow \mathrm{LGr}(\wedge_{\mathbb{C}}^{\mathrm{top}} V) = \mathbb{R}P(\wedge_{\mathbb{C}}^{\mathrm{top}} V)$ with the squaring map $\mathbb{R}P(\wedge_{\mathbb{C}}^{\mathrm{top}} V) \rightarrow ((\wedge_{\mathbb{C}}^{\mathrm{top}} V)^{\otimes 2} \setminus 0)/\mathbb{R}_{>0}$. Given a “basepoint” $S \in \mathrm{LGr}(V)$ (so $V = S \otimes_{\mathbb{R}} \mathbb{C}$), we obtain canonical identifications $\mathrm{LGr}(V) = \mathrm{LGr}(S \otimes_{\mathbb{R}} \mathbb{C}) = U(S \otimes_{\mathbb{R}} \mathbb{C})/O(S)$ and $((\wedge_{\mathbb{C}}^{\mathrm{top}} V)^{\otimes 2})/\mathbb{R}_{>0} = (\mathbb{C} \setminus 0)/\mathbb{R}_{>0} = U(1)$, under which (5.3) is given by \det^2 . Given a map $\mathrm{LGr}(V) \rightarrow K(\mathbb{Z}, 1) \times K(\mathbb{Z}/2, 2)$ representing (μ, w_2) , we obtain a $(\mathbb{Z} \times \mathbb{R}P^\infty)$ -bundle $\mathrm{LGr}(V)^\# \rightarrow \mathrm{LGr}(V)$.

We now globalize. Let X be a symplectic manifold, and denote by $\mathrm{LGr}(X)$ the bundle of Lagrangian Grassmannians of TX over X . There is a canonical map $\mathrm{LGr}(X) \rightarrow K(\mathbb{Z}/2, 2)$ restricting to w_2 on each fiber, namely the pullback of $w_2 : BO \rightarrow K(\mathbb{Z}/2, 2)$ under the map classifying the tautological bundle over $\mathrm{LGr}(X)$. There need not be a map $\mathrm{LGr}(X) \rightarrow K(\mathbb{Z}, 1)$ whose restriction to each fiber represents μ ; the obstruction to the existence of such a map is given by $2c_1(TX) \in H^2(X, \mathbb{Z})$ and is represented geometrically by the complex line bundle $(\wedge_{\mathbb{C}}^{\mathrm{top}} TX)^{\otimes 2}$. *Grading/orientation data for X* is, by definition, a choice of map $X \rightarrow K(\mathbb{Z}/2, 2)$ and map $\mathrm{LGr}(X) \rightarrow K(\mathbb{Z}, 1)$ whose restriction to each fiber represents μ . The choice of map $X \rightarrow K(\mathbb{Z}/2, 2)$ induces a map $\mathrm{LGr}(X) \rightarrow K(\mathbb{Z}/2, 2)$ by pulling back and adding the canonical map $\mathrm{LGr}(X) \rightarrow K(\mathbb{Z}/2, 2)$ restricting to w_2 on each fiber. Grading/orientation data on X thus induces a map $\mathrm{LGr}(X) \rightarrow K(\mathbb{Z}, 1) \times K(\mathbb{Z}/2, 2)$ whose restriction to each fiber represents (μ, w_2) . The pullback of the tautological $(\mathbb{Z} \times \mathbb{R}P^\infty)$ -bundle over $K(\mathbb{Z}, 1) \times K(\mathbb{Z}/2, 2)$ thus defines a $(\mathbb{Z} \times \mathbb{R}P^\infty)$ -bundle $\mathrm{LGr}(X)^\# \rightarrow \mathrm{LGr}(X)$ associated to this choice of grading/orientation data.

We now introduce Lagrangians. Fix a choice of grading/orientation for X , giving $\mathrm{LGr}(X)^\# \rightarrow \mathrm{LGr}(X)$. Given a Lagrangian $L \subseteq X$, *grading/orientation data for L* means a lift of the canonical section of $\mathrm{LGr}(X)|_L$ to $\mathrm{LGr}(X)^\#|_L$. It is explained in Seidel [79, (11e)–(11l)] (also reviewed in [37, §3.2]) how such data determines graded orientation lines associated to transverse intersections of ordered pairs of Lagrangians, as well as a recipe for orienting moduli spaces of pseudo-holomorphic disks relative to these orientation lines.

For our purposes in this paper, we will induce grading/orientation data from Lagrangian polarizations. Recall that a (Lagrangian) *polarization* of a symplectic manifold X is a global section of $\mathrm{LGr}(X)$; equivalently (up to homotopy) it is a real vector bundle B with an isomorphism $B \otimes_{\mathbb{R}} \mathbb{C} = TX$. Given such a polarization, we obtain a map $\mathrm{LGr}(X) \rightarrow K(\mathbb{Z}, 1)$ using the section as the fiberwise basepoint, and we obtain a map $X \rightarrow K(\mathbb{Z}/2, 2)$ by pulling back $w_2 : BO \rightarrow K(\mathbb{Z}/2, 2)$ under the map classifying B . By this very definition, any Lagrangian which is everywhere tangent to the polarization admits canonical grading/orientation data (i.e., section of $\mathrm{LGr}(X)^\#|_L$). A stable polarization (a global section of $\mathrm{LGr}(TX \oplus \mathbb{C}^k)$ for some $k < \infty$) also induces grading/orientation data by restriction from $\mathrm{LGr}(TX \oplus \mathbb{C}^k)$ to $\mathrm{LGr}(X) = \mathrm{LGr}(TX)$.

In the specific case of cotangent bundles T^*M , there is a tautological polarization given by (the tangent space of) the tautological foliation by Lagrangian fibers of the projection $T^*M \rightarrow M$; the fibers are thus equipped with canonical grading/orientation data with respect to the grading/orientation data on T^*M induced by this polarization. Conormals to open sets with smooth (or cornered) boundary also have canonical grading/orientation data; see [Section 5.5](#). We will see in [Remark 5.27](#) and [Lemma 5.10](#) the point in the proof of [Theorem 1.1](#) where it matters to have chosen this particular grading/orientation data on T^*M .

Remark 5.2. The notion of grading/orientation data given above may be reformulated as follows, which connects it to the corresponding discussion of coefficient twisting in microlocal sheaf categories as it appears in [\[42\]](#), [\[48\]](#). The stable J -homomorphism sends a (stable) vector bundle to (the suspension spectrum of) its Thom space, which is a family of invertible modules over the sphere spectrum. Applying cochains, we may obtain a family of invertible dg \mathbb{Z} -modules. We thus have an infinite loop map

$$(5.4) \quad \mathbb{Z} \times BO \xrightarrow{J} \mathrm{Pic} \mathbb{S} \rightarrow \mathrm{Pic} \mathbb{Z}$$

sending a vector bundle V to the local system $C^*(V, V \setminus 0)$ (where Pic denotes the space of invertible modules). The invertible module $\mathbb{Z}[1]$ and the automorphism -1 of the invertible module \mathbb{Z} together define an isomorphism of infinite loop spaces $\mathbb{Z} \times B(\mathbb{Z}/2) \xrightarrow{\sim} \mathrm{Pic} \mathbb{Z}$. The map $\mathbb{Z} \times BO \rightarrow \mathrm{Pic} \mathbb{Z} = \mathbb{Z} \times B(\mathbb{Z}/2)$ is then the evident projection to \mathbb{Z} times the Stiefel–Whitney class w_1 , as considered above. Applying B as before, we obtain a map

$$(5.5) \quad U/O = B(\mathbb{Z} \times BO) \rightarrow B \mathrm{Pic} \mathbb{Z} = B\mathbb{Z} \times B^2(\mathbb{Z}/2).$$

Now the tangent bundle of a symplectic manifold X is classified by a map $X \rightarrow BU$, which we may compose with $BU \rightarrow B(U/O)$ to obtain a map $X \rightarrow B(U/O)$ which classifies the (stable) Lagrangian Grassmannian of X .

Composing this with $B(5.5)$ yields a map

$$(5.6) \quad X \rightarrow B^2\mathbb{Z} \times B^3(\mathbb{Z}/2).$$

Now grading/orientation data on X is equivalently a null-homotopy of this map. Indeed, the map to the second factor is canonically null-homotopic (since by definition it factors through $BU \rightarrow B(U/O) \rightarrow B^2O$ which is canonically null-homotopic), so a choice of null-homotopy of it is the same as a choice of map $X \rightarrow \Omega B^3(\mathbb{Z}/2) = B^2(\mathbb{Z}/2)$. The map to the first factor by definition classifies $(\wedge_C^{\text{top}} TX)^{\otimes 2}$, a trivialization of which is the same as a map $\text{LGr}(X) \rightarrow K(\mathbb{Z}, 1)$ whose restriction to each fiber represents μ .

On a Lagrangian $L \subseteq X$, there is a tautological section of $\text{LGr}(X)|_L$ given by the tangent space to the Lagrangian. That is, the restricted map $L \rightarrow B(U/O)$ has a canonical null-homotopy, inducing in turn a null-homotopy of the map $L \rightarrow B^2\mathbb{Z} \times B^3(\mathbb{Z}/2)$. Now given grading/orientation data for X , grading/orientation data on L is equivalently a homotopy between this null-homotopy and the restriction to L of the chosen null-homotopy of (5.6). Note that the space of such null-homotopies has the homotopy type of maps from L to $\Omega(B^2\mathbb{Z} \times B^3(\mathbb{Z}/2)) = B\mathbb{Z} \times B^2(\mathbb{Z}/2)$, the component group of which is $H^1(L, \mathbb{Z}) \oplus H^2(L, \mathbb{Z}/2)$. The obstruction to the existence of grading/orientation data for L thus lies in $H^1(L, \mathbb{Z}) \oplus H^2(L, \mathbb{Z}/2)$, and if this obstruction vanishes, the homotopy classes of choices of grading/orientation data for L form a torsor over $H^0(L, \mathbb{Z}) \oplus H^1(L, \mathbb{Z}/2)$.

A stable polarization of X gives a global section of the stable Lagrangian Grassmannian, hence a null-homotopy of $X \rightarrow B(U/O)$, hence of (5.6), which by definition agrees with the canonical homotopy of its restriction to any Lagrangian $L \subseteq X$ everywhere tangent to the polarization.

5.4. Wrapping exact triangle, stop removal, generation. The fundamental ingredients underlying our work in this section are the wrapping exact triangle and its consequence stop removal, both proved in [38]. The wrapping exact triangle can be thought of as quantifying the price of wrapping through a stop; it should be compared with [Theorem 4.11](#).

THEOREM 5.3 (Wrapping exact triangle [38, Thm. 1.10]). *Let (X, Λ) be a stopped Liouville sector, and let $p \in \Lambda$ be a point near which Λ is a Legendrian submanifold. If $L \subseteq X$ is an exact Lagrangian submanifold and $L^w \subseteq X$ is obtained from L by passing $\partial_\infty L$ through Λ transversally at p in the positive direction, then there is an exact triangle*

$$(5.7) \quad L^w \rightarrow L \rightarrow D_p \xrightarrow{[1]}$$

in $\mathcal{W}(X, \Lambda)$, where $D_p \subseteq X$ denotes the small Lagrangian disk linking Λ at p and the map $L^w \rightarrow L$ is the continuation map.

The following result about wrapped Fukaya categories is a consequence of the wrapping exact triangle and can be compared with [Theorem 4.14](#).

THEOREM 5.4 (Stop removal [[38](#), Thm. 1.20]). *Let (X, Λ') be a stopped Liouville sector. Let $\Lambda \subseteq \Lambda'$ be closed so that its complement $\Lambda' \setminus \Lambda \subseteq (\partial_\infty X)^\circ \setminus \Lambda$ is an isotropic submanifold. Then pushforward induces an equivalence*

$$(5.8) \quad \mathcal{W}(X, \Lambda')/\mathcal{D} \xrightarrow{\sim} \mathcal{W}(X, \Lambda),$$

where \mathcal{D} denotes the collection of small Lagrangian disks linking (Legendrian points of) $\Lambda' \setminus \Lambda$.

We will also need to know that

THEOREM 5.5. *The cotangent fibers split-generate $\mathcal{W}(T^*M)$.*

Proof. When M is compact (including the case with boundary), this is [[38](#), Thm. 1.14 and Ex. 1.15]. For a general possibly non-compact M , we observe that any Lagrangian $L \in \mathcal{W}(T^*M)$ is in the essential image of the pushforward functor $\mathcal{W}(T^*M_L) \rightarrow \mathcal{W}(T^*M)$ for some compact codimension zero submanifold-with-boundary $M_L \subseteq M$. Now push foward the fact that L is split-generated by a fiber in $\mathcal{W}(T^*M_L)$. \square

Remark 5.6. In fact, the argument above shows that the fibers generate $\mathcal{W}(T^*M)$, however we only need split-generation.

Another ingredient which proves useful in our computations is the Künneth theorem for Floer cohomology and wrapped Fukaya categories, also proved in [[38](#)].

5.5. Conormals and corners. Let $U \subseteq M$ be a relatively compact open set. When U has smooth boundary, we write $L_U \subseteq T^*M$ for (a smoothing of) the union of $U \subseteq M \subseteq T^*M$ with the *outward* conormal along its boundary. More generally, if \bar{U} is a compact manifold-with-corners with interior U , then L_U shall mean $L_{\tilde{U}}$, where \tilde{U} is obtained from U by smoothing out its boundary. We also allow the degenerate case that U is a point p (hence, in particular, not open), in which case L_p denotes the cotangent fiber over p . In all of the above cases, we could also equivalently say that L_U is a rounding of $\text{ss}(\mathbb{Z}_U)$ (compare [Section 4.3](#)).

Remark 5.7. Various natural constructions, such as the cornering operations of [Section 2.2–2.3](#) and taking products, introduce corners. By convention, we conflate such cornered objects with their smoothings, usually without comment. The choice of this smoothing is always a contractible choice which is ultimately irrelevant.

Recall that for a Lagrangian L , we write L^+ for an unspecified small positive Reeb pushoff of L , and L^- for a negative pushoff. Thus if U is a relatively compact open set with smooth boundary and U^+ denotes its ϵ neighborhood

in some metric, then $L_{U^+} = L_U^+$. That is, positive Reeb flow pushes outward conormals out. In particular, $(T_p^*M)^+ = L_{B_\epsilon(p)}$.

Each L_U is exact and possesses canonical grading/orientation data: the codimension zero inclusion $U \subseteq L_U$ is a homotopy equivalence, and U is a codimension zero submanifold of, and thereby inherits all of this data from, the zero section. The grading/orientation data for the zero section arises from the canonical homotopy from its tangent bundle $TM \subseteq T(T^*M)|_M$ to the family of tangent spaces of the cotangent fibers $T^*M \subseteq T(T^*M)|_M$ (which is the chosen polarization of T^*M) given by $e^{i\theta}T^*M$ for $\theta \in [0, \pi/2]$ (where J is chosen so that $J(T^*M) = TM$).

5.6. Floer cohomology between conormals of balls and stable balls. Here we study the Floer cohomology between conormals of open sets with smooth boundary (though recall [Remark 5.7](#) about implicit smoothing of corners). The assertion that an open set with smooth boundary is a ball shall mean that its *closure* is diffeomorphic to the standard closed unit ball. Note that a small positive pushoff of the cotangent fiber L_p over a point p is the conormal of a small open ball around p , so we may substitute “point” in place of “open ball” in many of the statements below.

LEMMA 5.8. *Let $U, V \subseteq M$ be balls with $\overline{U} \subseteq V$. Then $HF^*(L_V, L_U)$ equals \mathbb{Z} and is canonically generated by the continuation element, lying in degree zero.*

Proof. There is a positive isotopy from L_U^+ to L_V in the complement of $\partial_\infty L_U$. Hence $HF^*(L_V, L_U) = HF^*(L_U, L_U)$ (compare [\[37, Lem. 3.21\]](#)), but this latter group (which is isomorphic to $H^*(L_U) = H^*(U)$) is generated by its identity element. \square

LEMMA 5.9. *Let $U, V \subseteq M$ be balls with $\overline{U} \subseteq V$, and let $\overline{V} \subseteq W \subseteq M$. The continuation map $HF^*(L_W, L_V) \rightarrow HF^*(L_W, L_U)$ (i.e., multiplication by the continuation element in $HF^*(L_V, L_U)$) is an isomorphism.*

Proof. The positive isotopy $L_U \rightsquigarrow L_V$ takes place in the complement of $\partial_\infty L_W$, hence induces an isomorphism $HF^*(L_W, L_V) = HF^*(L_W, L_U)$ which agrees with multiplication by the continuation element by [\[37, Lem. 3.26\]](#). \square

LEMMA 5.10. *Let $U \subseteq M$ be a ball, and let $\overline{U} \subseteq W \subseteq M$. There is a canonical isomorphism $HF^*(L_W, L_U) = \mathbb{Z}$, with respect to which the continuation maps from [Lemma 5.9](#) act as the identity on \mathbb{Z} .*

Proof. The groups $HF^*(L_W, L_p)$ form a local system of $p \in W$ [\[37, Lem. 3.21\]](#). It suffices to show that this local system is canonically isomorphic to the constant local system \mathbb{Z} . Indeed, by [Lemma 5.9](#) we have a canonical isomorphism $HF^*(L_W, L_U) = HF^*(L_W, L_p)$ for any $p \in U$, which is compatible with the local system structure of $HF^*(L_W, L_p)$ by [\[37, Lem. 3.26\]](#).

The assertion that the local system $p \mapsto HF^*(L_W, L_p)$ is canonically trivialized is local on W , so let us consider p varying only in a small ball $U \subseteq W$. Now the isomorphism $HF^*(L_W, L_p) = HF^*(L_W, L_{p'})$ for nearby points p and p' from [37, Lem. 3.21] is induced by a diffeomorphism supported in U sending p to p' . As such, it is sent under the identifications $HF^*(L_W, L_p) = HF^*(L_U, L_p)$ and $HF^*(L_W, L_{p'}) = HF^*(L_U, L_{p'})$ (coming from the fact that, in both cases, they are generated by the “same” Lagrangian intersection and have no Floer differential, and we have chosen the “same” grading/orientation data on L_W and L_U in Section 5.5) to the corresponding isomorphism $HF^*(L_U, L_p) = HF^*(L_U, L_{p'})$. Now $HF^*(L_U, L_p)$ and $HF^*(L_U, L_{p'})$ are canonically \mathbb{Z} by Lemma 5.9, and the isomorphism between them acts as the identity on \mathbb{Z} by [37, Lem. 3.26]. \square

LEMMA 5.11. *Let V be an open set with smooth boundary, and let U be a ϵ -ball centered at a point on ∂V . Then $HF^*(L_U, L_V) = 0 = HF^*(L_V, L_U)$.*

Proof. During the obvious isotopy of U outward to become disjoint from V , their conormals never intersect at infinity. \square

By a *stable ball*, we mean a compact manifold-with-boundary which is contractible; the statement that an open set with smooth boundary is a stable ball shall mean its closure is a stable ball. The reason we study stable balls is that we do not know how to prove that for a subanalytic Whitney triangulation, the “inward cornering” in the sense of Section 2.3 of an open star is a ball; it is, however, obviously a stable ball.

To compute Floer cohomology between conormals of stable balls, we reduce to the case of conormals to balls by stabilizing (i.e., taking their product with conormals to standard balls in \mathbb{R}^k) and appealing to the Künneth theorem for Floer cohomology. We begin by showing that the stabilization of a stable ball is indeed a ball, thus justifying the name. This uses the following famous corollary of the h -cobordism theorem:

THEOREM 5.12. *A stable ball of dimension ≥ 6 with simply connected boundary is a ball.*

COROLLARY 5.13. *Let M be a stable ball. Then $M \times I^k$ is a ball provided $\dim M + k \geq 6$ and $k \geq 1$.*

Proof. We just need to check that the boundary of $M \times I^k$ is simply connected. It suffices to show that for any stable ball N of dimension ≥ 2 , the boundary of $N \times I$ is simply connected. The boundary of $N \times I$ is, up to homotopy, two copies of N glued along their common boundary. Since N is contractible, the fundamental group of this gluing vanishes provided ∂N is

connected. If ∂N were disconnected, then by Poincaré duality, the cohomology group $H^{\dim N-1}(N)$ would be nonzero, which contradicts contractibility as $\dim N \geq 2$. \square

PROPOSITION 5.14. *Let $U, V \subseteq M$ be stable balls with $\overline{U} \subseteq V$. Then $HF^*(L_V, L_U) = \mathbb{Z}$, and it is equipped with a canonical generator 1_{VU} which we call the pseudo-continuation element. (It coincides with the usual continuation map when U and V are balls.) The pseudo-continuation elements are closed under composition: for any triple of stable balls $U, V, W \subseteq M$ with $\overline{U} \subseteq V$ and $\overline{V} \subseteq W$, we have $1_{WV}1_{VU} = 1_{WU}$.*

Proof. We multiply L_U, L_V by $L_{B_1(0)}, L_{B_2(0)} \subseteq T^*\mathbb{R}^k$, where k is sufficiently large to guarantee that $U \times B_1(0)$ and $V \times B_2(0)$ are balls by [Corollary 5.13](#). By the Künneth formula for Floer cohomology (see, e.g., [38, Lem. 8.3]) and freeness of $HF^*(L_{B_2(0)}, L_{B_1(0)})$ (compare [38, Rmk. 8.4]), we have

$$\begin{aligned} HF^*(L_V \times L_{B_2(0)}, L_U \times L_{B_1(0)}) &= HF^*(L_V, L_U) \otimes HF^*(L_{B_2(0)}, L_{B_1(0)}) \\ &= HF^*(L_V, L_U). \end{aligned}$$

On the other hand, by the result for balls [Lemma 5.8](#), we have

$$HF^*(L_V \times L_{B_2(0)}, L_U \times L_{B_1(0)}) = HF^*(L_{V \times B_2(0)}, L_{U \times B_1(0)}) = \mathbb{Z}.$$

After arguing that the above identification is compatible with rounding of corners, this defines the canonical generator $1_{VU} \in HF^*(L_V, L_U)$. The proof that $1_{WV}1_{VU} = 1_{WU}$ is the same: stabilize to reduce to the corresponding fact for honest continuation maps. \square

In order to make sense of the next corollary, recall that [Lemma 5.8](#) and [Proposition 5.14](#) continue to apply in the limiting situation in which L_U is replaced by a cotangent fibre $L_p = T_p^*M$ for some $p \in V$.

COROLLARY 5.15. *Let $U \subseteq M$ be any stable ball. Then the pseudo-continuation element $L_U \rightarrow T_p^*M$ is an isomorphism in $\mathcal{W}(T^*M)$ for any point $p \in U$.*

Proof. Note that this corollary is tautologically true if U is a ball, as genuine continuation elements by definition are isomorphisms in the wrapped Fukaya category. By pushing forward, it suffices to treat the case $M = U^+$. Appealing to the fully faithful Künneth embedding (see [38, Thm. 1.5]) $\mathcal{W}(T^*U^+) \otimes \mathcal{W}(T^*I^k) \hookrightarrow \mathcal{W}(T^*(U^+ \times I^k))$, it further suffices to show the result after taking (the image under this embedding of) the product of this pseudo-continuation element with the continuation element $L_{I^k} \rightarrow [\text{fiber}]$ (which is an isomorphism in $\mathcal{W}(T^*I^k)$). The pseudo-continuation element $L_U \rightarrow [\text{fiber}]$ is, by definition, sent by this stabilization to the continuation element $L_{U \times I^k} \rightarrow$

[fiber] (which is defined since the stabilized stable ball $U^+ \times I^k$ is a ball). This latter map is an isomorphism, so we are done. \square

Here is an improved version of [Lemma 5.10](#):

LEMMA 5.16. *For any stable ball $U \subseteq M$ whose closure is contained in $W \subseteq M$, there is canonical isomorphism $HF^*(L_W, L_U) = \mathbb{Z}$, such that for an inclusion $\overline{U} \subseteq V$ of such stable balls, the pseudo-continuation map $\mathbb{Z} = HF^*(L_W, L_V) \rightarrow HF^*(L_W, L_U) = \mathbb{Z}$ (multiplication by the pseudo-continuation element in $HF^*(L_V, L_U)$) is the identity on \mathbb{Z} .*

Proof. Stabilize to reduce to [Lemma 5.10](#). \square

There is similarly an improved version of [Lemma 5.11](#):

LEMMA 5.17. *Let V be an open set with smooth boundary, and let U be stable ball such that $U \cap \partial V$ is also a stable ball. Then $HF^*(L_U, L_V) = 0 = HF^*(L_V, L_U)$.*

Proof. Stabilization (multiplying both U and V by I^k) and appealing to the Künneth formula for Floer cohomology (as in the proof of [Lemma 5.14](#)) reduces this proof to [Lemma 5.11](#). (Note that $U \cap \partial V$ necessarily divides U into two stable balls.) \square

A more subtle result about stable balls is the following, which will be important later:

PROPOSITION 5.18. *Let $X^m \subseteq Y^n$ be an inclusion of stable balls, with $\partial X \subseteq \partial Y$. Assume there exists another stable ball (with corners) $Z^{m+1} \subseteq Y^n$ such that ∂Z is the union of X with a smooth submanifold of ∂Y . Then the pseudo-continuation element $L_Y \rightarrow L_{B_\epsilon(x)}$ is an isomorphism in $\mathcal{W}(T^*Y, N_\infty^*X)$ for any $x \in X$.*

Proof. By stabilization, we reduce to the case that X , Y , and Z are all balls. This implies that, up to diffeomorphism, everything is standard: Y is the unit ball, X is the intersection of Y with a linear subspace, and Z is the intersection of Y with a linear halfspace. Indeed, since X and Z are balls, we can use Z to push X to $Z \cap \partial Y$, thus showing that X is simply a slight inward pushoff of the ball $Z \cap \partial Y \subseteq \partial Y$.

By definition, the pseudo-continuation element becomes the continuation element under stabilization (i.e., after multiplying by the continuation isomorphism from the conormal of a large ball to that of a small ball as in the proof of [Corollary 5.15](#)). Once everything is standard, the continuation map $L_Y \rightarrow L_{B_\epsilon(x)}$ is an isomorphism, since there is a positive isotopy $L_{B_\epsilon(x)} \rightsquigarrow L_Y$ disjoint from N_∞^*X at infinity. \square

We will apply [Proposition 5.18](#) when (before rounding) X is a simplex in a triangulation, Y is its star, and Z is any simplex containing X of dimension one larger.

5.7. Fukaya categories of conormals to stars. Let \mathcal{S} be a Whitney stratification of M by locally closed smooth submanifolds. For an \mathcal{S} -constructible open set U , we abuse notation and denote by L_U the conormal of the inward cornering of U with respect to \mathcal{S} in the sense of [Section 2.3](#). More precisely, we have $L_U := L_{U-\underline{\epsilon}}$ for $\underline{\epsilon} \in \mathbb{R}_{>0}^{\mathcal{S}}$ satisfying $\epsilon_\alpha \leq f((\epsilon_\beta)_{\beta \subsetneq \alpha})$. Recall that $L_{U-\underline{\epsilon}}$ is disjoint from $N^*\mathcal{S}$ at infinity but limits to it as $\underline{\epsilon} \rightarrow 0$.

LEMMA 5.19. *Let L be any Lagrangian disjoint at infinity from $N^*\mathcal{S}$. Then for all $\underline{\epsilon} > 0$ sufficiently small, $CF^*(L_{U-\underline{\epsilon}}, L) \xrightarrow{\sim} CW^*(L_{U-\underline{\epsilon}}, L)_{N_\infty^*\mathcal{S}}$.*

Proof. Taking $\underline{\epsilon} \rightarrow 0$ is a positive wrapping of $L_{U-\underline{\epsilon}}$ (compare [Section 5.5](#)) which converges to (while remaining disjoint from) $N_\infty^*\mathcal{S}$. It is thus cofinal by [Lemma 5.1](#). \square

Now assume further that \mathcal{S} is a triangulation, and let us consider the conormals to (the inward cornerings of) open stars $L_{\text{star}(s)} \in \mathcal{W}(T^*M, N_\infty^*\mathcal{S})$. Since $\text{star}(s)$ is contractible, $L_{\text{star}(s)}$ is the conormal to a stable ball ([Lemma 2.10](#)), and hence the results about stable balls from [Section 5.6](#) above apply, allowing us to deduce the following:

PROPOSITION 5.20. *We have*

$$(5.9) \quad HW^*(L_{\text{star}(s)}, L_{\text{star}(t)})_{N_\infty^*\mathcal{S}} = \begin{cases} \mathbb{Z} & t \rightarrow s, \\ 0 & \text{otherwise} \end{cases}$$

generated in the former case by the pseudo-continuation element.

Proof. Fix a small $\underline{\epsilon} > 0$, and let $\underline{\delta} \rightarrow 0$. By [Lemma 5.19](#), the wrapped Floer cohomology $HW^*(L_{\text{star}(s)}, L_{\text{star}(t)})$ is calculated by $HF^*(L_{\text{star}(s)-\underline{\delta}}, L_{\text{star}(t)-\underline{\epsilon}})$.

Now if $t \rightarrow s$, then $\text{star}(t)^{-\underline{\epsilon}} \subseteq \text{star}(s)^{-\underline{\delta}}$ is an inclusion of stable balls, so by [Proposition 5.14](#), $HF^*(L_{\text{star}(s)-\underline{\delta}}, L_{\text{star}(t)-\underline{\epsilon}}) = \mathbb{Z}$ is generated by the pseudo-continuation element.

Now suppose that $t \not\rightarrow s$. If $\text{star}(s) \cap \text{star}(t) = \emptyset$, then the desired vanishing is trivial. Otherwise, we have $\text{star}(s) \cap \text{star}(t) = \text{star}(r)$, where r is the simplex spanned by the union of the vertices of s and t . To show the desired vanishing, it suffices by [Proposition 5.17](#) to show that $\text{star}(t)^{-\underline{\epsilon}} \cap \partial \text{star}(s)^{-\underline{\delta}}$ is a stable ball, which is the content of [Lemma 2.12](#). \square

It will be convenient to have another perspective on the objects $L_{\text{star}(s)}$. Let L_s denote the conormal to a small ball centered at any point on the stratum s (this conormal is disjoint from $N_\infty^*\mathcal{S}$ at infinity by Whitney's condition

(b), compare with the proof of [Lemma 2.6](#)); this is well defined up to Lagrangian isotopy. One reason the L_s are nice to consider is the following calculation:

LEMMA 5.21. *For any \mathcal{S} -constructible open set U , we have*

$$(5.10) \quad HW^*(L_U, L_s)_{N_\infty^* \mathcal{S}} = \begin{cases} \mathbb{Z} & \text{star}(s) \subseteq U, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We calculate using [Lemma 5.19](#). If s is a stratum in the interior of U , then the ball centered at s is contained in U and, in particular, as L_s is a small positive pushoff of a cotangent fiber to a point in s , one can arrange for there to be a single intersection point between $L_{U-\epsilon}$ (for small ϵ) and L_s . Hence $HW^*(L_U, L_s)_{N_\infty^* \mathcal{S}} = HF^*(L_{U-\epsilon}, L_s) = \mathbb{Z}$. If s is a stratum not contained in the closure of U , then the morphism space obviously vanishes since the two Lagrangians ($L_{U-\epsilon}$ for any ϵ and L_s) are disjoint.




Figure 3. The isotopy from $U^{-\epsilon, \delta+\eta}$ to $U^{-\epsilon, \delta-\eta}$ to $U^{-\epsilon}$.

Finally, we claim that if s is a stratum on the boundary of U , the morphism space still vanishes. To prove this, it suffices to construct a cofinal wrapping of L_U which begins disjoint from L_s and remains forever disjoint from $\partial_\infty L_s$ (see [\[37, Lem. 3.26\]](#)). Such an isotopy is illustrated in [Figure 3](#), which we now define precisely. Fix a point $q \in s$, and consider the stratification \mathcal{S}_q obtained from \mathcal{S} by declaring $\{q\}$ to be its own stratum. Now the inward cornering with respect to \mathcal{S}_q may be denoted $U^{-\epsilon, \delta}$ for $\epsilon \in \mathbb{R}_{>0}^S$ and $\delta > 0$ the parameter associated to the new stratum $\{q\}$. Fix ϵ , and take δ to zero, and note that [Proposition 2.15](#) implies this gives an isotopy of $U^{-\epsilon, \delta}$ whose conormals remain disjoint at infinity from $N^* \mathcal{S}$. Once $\delta = 0$, we just have $U^{-\epsilon}$, whose conormal has cofinal wrapping by taking $\epsilon \rightarrow 0$. Now take L_s to be the conormal of $B_\delta(q)$ and take the isotopy of L_U to be given the conormal of the isotopy

$$(5.11) \quad U^{-\epsilon, \delta+\eta} \rightsquigarrow U^{-\epsilon, \delta-\eta} \rightsquigarrow U^{-\epsilon}$$

illustrated in [Figure 3](#), followed by isotoping $U^{-\epsilon}$ by taking $\epsilon \rightarrow 0$. [Corollary 2.7](#) implies that this isotopy remains disjoint at infinity from $L_s = N^* B_\delta(q)$ except possibly at $U^{-\epsilon, \delta}$, but these do not intersect at infinity due to their coorientations being opposite. \square

Another reason that the L_s are nice to consider is that we can show using the wrapping exact triangle and stop removal that they (split-)generate:

PROPOSITION 5.22. *The objects L_s associated to strata s split-generate $\mathcal{W}(T^*M, N_\infty^* \mathcal{S})$.*

Proof. Denote by $\mathcal{S}_{\leq k}$ the stratification where we keep all strata of dimension $\leq k$ and combine all other strata into a single top stratum. We consider the sequence of categories

$$(5.12) \quad \begin{aligned} \mathcal{W}(T^*M, N_\infty^* \mathcal{S}) &= \mathcal{W}(T^*M, N_\infty^* \mathcal{S}_{\leq n-1} \rightarrow \mathcal{W}(T^*M, N_\infty^* \mathcal{S}_{\leq n-2}) \\ &\rightarrow \cdots \rightarrow \mathcal{W}(T^*M, N_\infty^* \mathcal{S}_{\leq 0}) \rightarrow \mathcal{W}(T^*M). \end{aligned}$$

Each of these functors removes a locally closed Legendrian submanifold $N_\infty^* \mathcal{S}_{\leq k} \setminus N_\infty^* \mathcal{S}_{\leq k-1}$, and thus by the stop removal theorem, [Theorem 5.4](#), is the quotient by the corresponding linking disks.

The linking disk at a point on $N_\infty^* \mathcal{S}_{\leq k} \setminus N_\infty^* \mathcal{S}_{\leq k-1}$ can be described as follows. A point on $N_\infty^* \mathcal{S}_{\leq k} \setminus N_\infty^* \mathcal{S}_{\leq k-1}$ is simply a point x on a k -dimensional stratum together with a covector ξ at x conormal to the stratum. Consider a small ball B_a centered at x , and consider a smaller ball $B_b \subseteq B_a$ disjoint from the stratum containing x . There is a family of balls starting at B_a and shrinking down to B_b whose boundaries are tangent to the stratum containing x only at (x, ξ) . It follows from the wrapping exact triangle [Theorem 5.3](#) that the cone on the resulting continuation map $L_{B_a} \rightarrow L_{B_b}$ is precisely the linking disk at (x, ξ) .

We have thus shown that the linking disks to each locally closed Legendrian $N_\infty^* \mathcal{S}_{\leq k} \setminus N_\infty^* \mathcal{S}_{\leq k-1}$ are generated by the objects L_s . By [Theorem 5.5](#) above, these L_s also split-generate the final category $\mathcal{W}(T^*M)$. We conclude that the L_s split-generate $\mathcal{W}(T^*M, N_\infty^* \mathcal{S})$, as the quotient by all of them vanishes. \square

Remark 5.23. A small variation on the above proof and an appeal to [38, Thm. 1.14] shows that the objects L_s in fact generate $\mathcal{W}(T^*M, N_\infty^* \mathcal{S})$. We give the weaker argument above to minimize the results we need to appeal to.

PROPOSITION 5.24. *The pseudo-continuation element $L_{\text{star}(s)} \rightarrow L_s$ is an isomorphism in $\mathcal{W}(T^*M, N_\infty^* \mathcal{S})$.*

Proof. We proceed by induction on the codimension of s . When s has codimension zero, the desired statement follows from [Corollary 5.15](#).

Now suppose that s has positive codimension. For any t of strictly smaller codimension than s , we have $\text{Hom}(L_{\text{star}(t)}, L_{\text{star}(s)}) = 0$ by [Proposition 5.20](#) and $\text{Hom}(L_{\text{star}(t)}, L_s) = 0$ by [Lemma 5.21](#).

Now by the discussion in the proof of [Proposition 5.22](#), the functor

$$(5.13) \quad \mathcal{W}(T^*M, N_\infty^* \mathcal{S}) \rightarrow \mathcal{W}(T^*M, N_\infty^* \mathcal{S}_{\leq \dim s})$$

quotients by cones of L_t for t of strictly smaller codimension than s . By the induction hypothesis and the calculations of the previous paragraph, such cones are left-orthogonal to L_s and $L_{\text{star}(s)}$. Hence it suffices to check that $L_{\text{star}(s)} \rightarrow L_s$ is an isomorphism in $\mathcal{W}(T^*M, N_\infty^* \mathcal{S}_{\leq \dim s})$.

Finally, we observe that by [Proposition 5.18](#), $L_{\text{star}(s)} \rightarrow L_s$ is an isomorphism in $\mathcal{W}(T^*M, N_\infty^* \mathcal{S}_{\leq \dim s})$. Namely, we take $Y = \text{star}(s)^-$, $X = s \cap \text{star}(s)^-$, and $Z = t \cap \text{star}(s)^-$ for any simplex t containing s and of one higher dimension. \square

Remark 5.25. For a “smooth triangulation” \mathcal{S} , there is an obvious positive isotopy from L_s to $L_{\text{star}(s)}$ disjoint from $N_\infty^* \mathcal{S}$ (thus proving [Proposition 5.24](#) in this case), obtained by expanding a small ball centered at a point on s to $\text{star}(s)$, keeping the boundary transverse to the strata of \mathcal{S} . We do not know whether this proof can be generalized from smooth triangulations to subanalytic Whitney triangulations.

5.8. Functors from poset categories to Fukaya categories.

Definition 5.26. Let M be a manifold with Whitney stratification \mathcal{S} , and let $U : \Pi \rightarrow \text{Ops}_\mathcal{S}(M)$ be a map from a poset Π to the poset of \mathcal{S} -constructible open subsets of M . Suppose further that each $U(\pi)^-$ (from [Section 2.3](#)) is a stable ball. Define a functor on cohomology categories

$$(5.14) \quad H^*F_U : \mathbb{Z}[\Pi] \rightarrow H^*\mathcal{W}(T^*M, N_\infty^* \mathcal{S})^{\text{op}}$$

by $H^*F_U(\pi) := L_{U(\pi)}$ and $H^*F_U(1_{\pi, \pi'}) = 1_{U(\pi'), U(\pi)} \in HW^*(L_{U(\pi')}, L_{U(\pi)})$ (the pseudo-continuation element).

Remark 5.27. Note that the definition of H^*F_U depends on having defined pseudo-continuation elements with the compatibility properties from [Proposition 5.14](#), which in turn depends on having equipped the cotangent fibers of T^*M with continuously varying grading/orientation data (compare [Section 5.3](#)). For general grading/orientation data on T^*M , it may not be possible to define continuously varying grading/orientation data on the cotangent fibers, in which case we could only define H^*F_U to respect composition up to sign. The resulting 2-cocycle, or rather its class in $H^2(N\Pi, \mathbb{Z}/2)$, would represent (the pullback of) the obstruction in $H^2(M, \mathbb{Z}/2)$ to choosing continuously varying relative Pin-structures on the cotangent fibers.

PROPOSITION 5.28. *For any functor $f : \mathbb{Z}[\Pi] \rightarrow H^*\mathcal{C}$ such that*

$$H^*\mathcal{C}(f(x), f(y))$$

is concentrated in degree zero for every pair $x \leq y \in \Pi$, there exists an A_∞ functor $F : \mathbb{Z}[\Pi] \rightarrow \mathcal{C}$ with $H^*F = f$. Moreover, given any two A_∞ functors $F, G : \mathbb{Z}[\Pi] \rightarrow \mathcal{C}$ such that $H^*\mathcal{C}(F(x), G(y))$ is concentrated in degree zero for every pair $x \leq y \in \Pi$ and a natural transformation $t : f \rightarrow g$, the space of A_∞ natural transformations $T : F \rightarrow G$ with $H^*T = t$ is contractible.

Proof. We show existence of a lift F by induction. Lift the action on objects arbitrarily. Take F^1 to be any map in the correct cohomology class $H^*F^1 = f$. Having chosen F^1, \dots, F^{k-1} , the existence of an F^k satisfying the A_∞ functor equations is equivalent to a certain element of

$$(5.15) \quad \prod_{\pi_0, \dots, \pi_k \in \Pi} \text{Hom}(\mathbb{Z}[\Pi](\pi_0, \pi_1) \otimes \cdots \otimes \mathbb{Z}[\Pi](\pi_{k-1}, \pi_k), \mathcal{C}(F(\pi_0), F(\pi_k)))$$

(namely the sum of all the terms of the A_∞ functor equations with k inputs except for those involving F^k) being a coboundary. This element is always a cocycle due to F^1, \dots, F^{k-1} satisfying the A_∞ functor equations, so it suffices to show that its class in cohomology vanishes. The cohomology of (5.15) is of course simply

$$(5.16) \quad \prod_{\pi_0 \leq \cdots \leq \pi_k \in \Pi} H^*\mathcal{C}(F(\pi_0), F(\pi_k)),$$

which is concentrated in degree zero by hypothesis. The obstruction class thus vanishes for degree reasons for $k \geq 3$. For $k = 2$, the obstruction class measures the failure of H^*F to respect composition, so by hypothesis the obstruction vanishes in this case as well. We conclude that there always exists an F^k compatible with the previously chosen F^1, \dots, F^{k-1} . (Compare [79, Lem. 1.9], where a variant on this obstruction theory argument is explained in more detail.)

To construct a natural transformation [79, (1d)] $T : F \rightarrow G$ with $H^*T = t$, first pick some T^0 lifting t . Given T^0, \dots, T^{k-1} , the obstruction to the existence of T^k is a degree $1 - k$ cohomology class in

$$(5.17) \quad \prod_{\pi_0 \leq \cdots \leq \pi_k \in \Pi} \mathcal{C}(F(\pi_0), G(\pi_k)).$$

It hence vanishes for degree reasons for $k \geq 2$, and for $k = 1$, it measures the failure of H^*T to respect morphisms, hence vanishes in this case as well.

Contractibility of the space of natural transformations T with $H^*T = t$ is, concretely, the assertion that the complex of pre-natural transformations $\text{Hom}(F, G)$ is acyclic in negative (cohomological) degree and that any two natural transformations (i.e., degree zero cocycles) T and T' with $H^*T = t = H^*T'$ are cohomologous. In both cases, we should produce a pre-natural transformation Q of degree $-g < 0$ with prescribed value of dQ . Such $Q = (Q^0, Q^1, \dots)$ is again constructed by induction. The existence of Q^k then comes

down to the vanishing in degree $1 - k - g$ of the cohomology of (5.17) for all $g \geq 1$ and $k \geq 0$ (except for $g = 1$ and $k = 0$, for which the relevant obstruction measures the failure of the desired value $T - T'$ of dQ to vanish on $H^* \mathcal{C}$, hence vanishes by the assumption $H^* T = t = H^* T'$). \square

Remark 5.29. The assertion of [Proposition 5.28](#) over a field k (instead of over \mathbb{Z}) is a straightforward consequence of the fact that any A_∞ category \mathcal{C} over k is quasi-isomorphic to an A_∞ category $\tilde{\mathcal{C}}$ with vanishing differential, sometimes called a *minimal model* of \mathcal{C} . In this case, the essential image of f inside $\tilde{\mathcal{C}}$ would be necessarily concentrated in degree zero on the chain level, hence have vanishing higher order A_∞ structure maps for degree reasons (and hence this essential image inside $\tilde{\mathcal{C}}$, equivalently \mathcal{C} , is *formal*). It follows that any functor on cohomology categories $k[\Pi] \rightarrow H^* \tilde{\mathcal{C}}$ would lift tautologically to an A_∞ functor by taking all higher operations to vanish. The proof above bypasses the question of the existence of minimal models over \mathbb{Z} .

COROLLARY 5.30. *There is a unique up to contractible choice A_∞ functor*

$$(5.18) \quad F_U : \mathbb{Z}[\Pi] \rightarrow \mathcal{W}(T^* M, N_\infty^* \mathcal{S})^{\text{op}}$$

lifting the functor on cohomology categories from [Definition 5.26](#).

Proof. By [Corollary 5.19](#), the wrapped Floer cohomology group

$$HW^*(L_{U(\pi)}, L_{U(\pi')})$$

is simply the Floer cohomology of two nested stable balls, which is \mathbb{Z} by [Proposition 5.14](#). Thus [Proposition 5.28](#) is applicable. \square

Remark 5.31. To extend [Corollary 5.30](#) to the Fukaya category with a \mathbb{Z}/N -grading, we would need to add to the requirement that F (and the natural transformations $F_1 \rightarrow F_2$) must lift to \mathbb{Z} -graded categories locally. (The \mathbb{Z} -grading is only defined locally, over any contractible open subset of M .)

For the next corollary, let us denote by \mathcal{H}^* the functor from $\text{Mod } \mathbb{Z}$ to the category of graded abelian groups given by taking the cohomology of *objects* of $\text{Mod } \mathbb{Z}$. The functor \mathcal{H}^* factors through, but does not coincide with, the functor $H^* : \text{Mod } \mathbb{Z} \rightarrow H^* \text{Mod } \mathbb{Z}$ which exists for any A_∞ category in place of $\text{Mod } \mathbb{Z}$ (and which takes cohomology of *morphisms*).

COROLLARY 5.32. *Consider functors $F : \mathbb{Z}[\Pi] \rightarrow \text{Mod } \mathbb{Z}$ such that $\mathcal{H}^* F(x)$ is free and concentrated in degree zero for all $x \in \Pi$. Given any two such functors F and G and a natural transformation $t : \mathcal{H}^* F \rightarrow \mathcal{H}^* G$, the space of natural transformations $T : F \rightarrow G$ with $\mathcal{H}^* T = t$ is contractible. In particular, any such functor F is quasi-isomorphic to $i \mathcal{H}^* F : \mathbb{Z}[\Pi] \rightarrow \text{Mod } \mathbb{Z}$, namely the composition of $\mathcal{H}^* F$ with the inclusion i of free abelian groups into $\text{Mod } \mathbb{Z}$ (as complexes concentrated in degree zero).*

Proof. We first argue that if $P, Q \in \text{Mod } \mathbb{Z}$ are such that \mathcal{H}^*P and \mathcal{H}^*Q are free and concentrated in degree zero, then the natural map

$$(5.19) \quad H^* \text{Hom}(P, Q) \rightarrow \text{Hom}(\mathcal{H}^*P, \mathcal{H}^*Q)$$

is an isomorphism. Since \mathcal{H}^*P is projective, there is a quasi-isomorphism $\mathcal{H}^*P \rightarrow P$ (and the same for Q). It follows that there is a quasi-isomorphism of chain complexes $\text{Hom}(P, Q) = \text{Hom}(\mathcal{H}^*P, \mathcal{H}^*Q)$ (homomorphisms in $\text{Mod } \mathbb{Z}$). Since \mathcal{H}^*P and \mathcal{H}^*Q are projective and concentrated in degree zero, the complex $\text{Hom}(\mathcal{H}^*P, \mathcal{H}^*Q)$ is (quasi-isomorphic to) homomorphisms of abelian \mathbb{Z} modules $\mathcal{H}^*P \rightarrow \mathcal{H}^*Q$ concentrated in degree zero; thus (5.19) is an isomorphism as desired.

Now suppose F and G are as in the statement and a natural transformation $t : \mathcal{H}^*F \rightarrow \mathcal{H}^*G$ is given. Since \mathcal{H}^*F and \mathcal{H}^*G are free and concentrated in degree zero, we see from (5.19) that the data of t is equivalent to the data of a natural transformation $\bar{t} : H^*F \rightarrow H^*G$. Now (5.19) also implies that the hypotheses of [Proposition 5.28](#) are satisfied, so the space of natural transformations T with $H^*T = \bar{t}$ (which is, as just noted, equivalent to $\mathcal{H}^*T = t$) is contractible.

It is immediate from the definition that $\mathcal{H}^*F = \mathcal{H}^*i\mathcal{H}^*F$, so the final statement follows from the first. \square

Definition 5.33. For a Whitney triangulation \mathcal{S} , let

$$(5.20) \quad F_{\mathcal{S}} : \mathbb{Z}[\mathcal{S}] \rightarrow \mathcal{W}(T^*M, N_{\infty}^*\mathcal{S})^{\text{op}}$$

denote the functor induced from [Definition 5.26](#) and [Corollary 5.30](#) by the map associating to each simplex of \mathcal{S} its open star.

THEOREM 5.34. *The functor $F_{\mathcal{S}}$ is a Morita equivalence.*

Proof. [Proposition 5.20](#) shows full faithfulness of $F_{\mathcal{S}}$. [Propositions 5.22](#) and [5.24](#) together show essential surjectivity of $F_{\mathcal{S}}$ (after passing to Perf). \square

We now show that $F_{\mathcal{S}}$ is compatible with refinement (compare [Lemma 4.6](#)):

THEOREM 5.35. *For \mathcal{S}' a refinement of \mathcal{S} , the following diagram commutes up to contractible choice:*

$$(5.21) \quad \begin{array}{ccc} \mathbb{Z}[\mathcal{S}'] & \xrightarrow{F_{\mathcal{S}'}} & \mathcal{W}(T^*M, N_{\infty}^*\mathcal{S}')^{\text{op}} \\ r \downarrow & & \downarrow \rho \\ \mathbb{Z}[\mathcal{S}] & \xrightarrow{F_{\mathcal{S}}} & \mathcal{W}(T^*M, N_{\infty}^*\mathcal{S})^{\text{op}}. \end{array}$$

Proof. There are two functors $\rho \circ F_{\mathcal{S}'}$ and $F_{\mathcal{S}} \circ r$ from $\mathbb{Z}[\mathcal{S}']$ to $\mathcal{W}(T^*M, N_{\infty}^*\mathcal{S})$. By [Proposition 5.28](#), it suffices to define a canonical natural isomorphism between the induced functors on cohomology categories. It is most natural to define this canonical natural isomorphism in the direction $F_{\mathcal{S}} \circ r \Rightarrow \rho \circ F_{\mathcal{S}'}$.

To a stratum s of \mathcal{S}' , the composition $F_{\mathcal{S}} \circ r$ associates the conormal of $\text{stars}_s(r(s))$, and the composition $\rho \circ F_{\mathcal{S}'}$ associates the conormal of $\text{stars}'_s(s)$. Since $\text{stars}_s(r(s)) \supseteq \text{stars}'_s(s)$ is an inclusion of stable balls, by [Proposition 5.14](#) we may consider the pseudo-continuation element from one to the other. Given that pseudo-continuation elements are closed under composition by [Proposition 5.14](#), it is easy to check that this defines a natural transformation $H^*(F_{\mathcal{S}} \circ r) \Rightarrow H^*(\rho \circ F_{\mathcal{S}'})$.

This natural transformation is in fact a natural isomorphism since the natural maps from both $L_{\text{stars}_s(r(s))}$ and $L_{\text{stars}'_s(s)}$ to $L_s = L_{r(s)}$ are isomorphisms by [Proposition 5.24](#). \square

5.9. In conclusion.

THEOREM 5.36. *The functor $\Lambda \mapsto \text{Perf } \mathcal{W}(T^*M, \Lambda)^{\text{op}}$ is a microlocal Morse theatre in the sense of [Definition 3.4](#), which casts the linking disks at smooth points of Λ as the Morse characters.*

Proof. [Definition 5.33](#) and [Theorems 5.34](#) and [5.35](#) give the identification between $\mathcal{S} \mapsto \text{Perf } \mathcal{S}$ and $\mathcal{S} \mapsto \text{Perf } \mathcal{W}(T^*M, N_{\infty}^* \mathcal{S})^{\text{op}}$ via the functors $F_{\mathcal{S}}$.

Stop removal [Theorem 5.4](#) says that $\mathcal{W}(T^*M, \Lambda') \rightarrow \mathcal{W}(T^*M, \Lambda)$ is the quotient by the linking disks at the smooth points of $\Lambda' \setminus \Lambda$. It therefore suffices to show that the Morse characters are precisely (isomorphic to) these linking disks.

Recall from [Definition 3.3](#) that a Morse character at a smooth point $p \in \Lambda$ is defined as follows. We choose a function $f : M \rightarrow \mathbb{R}$ and an $\epsilon > 0$ such that $f^{-1}(-\infty, \epsilon)$ is relatively compact, f has no critical values in $[-\epsilon, \epsilon]$ and df is transverse to $\mathbb{R}_{>0} \times \Lambda$ over $f^{-1}[-\epsilon, \epsilon]$, intersecting it only at p (where f vanishes). We also choose a subanalytic Whitney triangulation \mathcal{S} such that $\Lambda \subseteq N_{\infty}^* \mathcal{S}$ and $f^{-1}(-\infty, -\epsilon)$ and $f^{-1}(-\infty, \epsilon)$ are constructible. The Morse character associated to these choices is then defined as the image in $\mathcal{W}(T^*M, \Lambda)$ of

$$(5.22) \quad \text{cone}(1_{f^{-1}(-\infty, -\epsilon)} \rightarrow 1_{f^{-1}(-\infty, \epsilon)}) \in \text{Perf } \mathcal{S} = \text{Perf } \mathcal{W}(T^*M, N_{\infty}^* \mathcal{S})^{\text{op}},$$

where the morphism $1_{f^{-1}(-\infty, -\epsilon)} \rightarrow 1_{f^{-1}(-\infty, \epsilon)}$ is (the linearization of) the canonical one from [\(3.5\)](#). To show that this cone is indeed sent to the linking disk at p in $\mathcal{W}(T^*M, \Lambda)$, we will make use of the wrapping exact triangle [Theorem 5.3](#), which says that the linking disk at p is the cone of the continuation map associated to any positive isotopy of Lagrangians in T^*M which crosses Λ exactly once transversely at p . Specifically, there is an obvious positive isotopy from the conormal of $f^{-1}(-\infty, -\epsilon)$ to the conormal of $f^{-1}(-\infty, \epsilon)$, namely $f^{-1}(-\infty, t)$ for $t \in [-\epsilon, \epsilon]$, since f has no critical values in the interval $[-\epsilon, \epsilon]$; the cone of the associated continuation element in $\mathcal{W}(T^*M, \Lambda)$ is thus the desired linking disk.

The conormals of $f^{-1}(-\infty, \pm\epsilon)$ are not themselves objects of the wrapped Fukaya category associated to the triangulation $\mathcal{W}(T^*M, N_\infty^*\mathcal{S})$, seeing as they by definition touch the stop. However, in [Section 5.7](#) we studied the conormals of the inward cornerings of \mathcal{S} -constructible open sets U , which we denoted $L_U := L_{U-\epsilon}$. These Lagrangians $L_{f^{-1}(-\infty, \pm\epsilon)}$ are thus, in particular, objects of $\mathcal{W}(T^*M, N_\infty^*\mathcal{S})$, and [Lemma 2.13](#) provides an isotopy between them and the (usual) conormals of $f^{-1}(-\infty, \pm\epsilon)$ which takes place in the complement of Λ , thus inducing an isomorphism in $\mathcal{W}(T^*M, \Lambda)$. Therefore to complete the argument, it suffices to show that

- (i) $F_{\mathcal{S}}(1_{f^{-1}(-\infty, \pm\epsilon)}) \in \text{Perf } \mathcal{W}(T^*M, N_\infty^*\mathcal{S})$ is isomorphic to $L_{f^{-1}(-\infty, \pm\epsilon)}$, and
- (ii) the canonical morphism $1_{f^{-1}(-\infty, -\epsilon)} \rightarrow 1_{f^{-1}(-\infty, \epsilon)}$ is sent by $F_{\mathcal{S}}$ and the isomorphisms (i) to an element in $HW^*(L_{f^{-1}(-\infty, \epsilon)}, L_{f^{-1}(-\infty, -\epsilon)})_{N_\infty^*\mathcal{S}}$ whose image in $\mathcal{W}(T^*M, \Lambda)$ is the continuation element associated to the natural positive isotopy between the conormals of $f^{-1}(-\infty, \pm\epsilon)$.

Regarding (i), let us establish the more general assertion that $F_{\mathcal{S}}(1_W)$ is isomorphic to L_W for any relatively compact open \mathcal{S} -constructible set W . Since $F_{\mathcal{S}}$ is a Morita equivalence, it suffices by Yoneda to show that the pullback $F_{\mathcal{S}}^*L_W = CW^*(L_W, F_{\mathcal{S}}(-))$ is isomorphic in $\text{Mod } \mathcal{S}$ to 1_W . Using [Lemmas 5.19](#) and [5.16](#) (for the case $\text{star}(s) \subseteq W$) and [Proposition 5.24](#) and [Lemma 5.21](#) (for the case $\text{star}(s) \not\subseteq W$), we have canonical isomorphisms

$$(5.23) \quad HW^*(L_W, L_{\text{star}(s)})_{N_\infty^*\mathcal{S}} = \begin{cases} \mathbb{Z} & \text{star}(s) \subseteq W, \\ 0 & \text{otherwise} \end{cases} = 1_W(s).$$

This identifies $F_{\mathcal{S}}^*L_W = 1_W$ objectwise (i.e., identifies their evaluations at every $s \in \mathcal{S}$). To identify $F_{\mathcal{S}}^*L_W = 1_W$ as modules, it suffices by [Corollary 5.32](#) to compare the action of morphisms $t \rightarrow s$ for $\text{star}(t) \subseteq \text{star}(s) \subseteq W$ at the level of cohomology. (Since the category $\mathbb{Z}[\mathcal{S}]$ and both modules are cohomologically concentrated in degree zero, there is no room for any higher homotopies.) In other words, we should show that the map

$$(5.24) \quad HW^*(L_W, L_{\text{star}(s)})_{N_\infty^*\mathcal{S}} \rightarrow HW^*(L_W, L_{\text{star}(t)})_{N_\infty^*\mathcal{S}}$$

that is obtained by multiplication by the pseudo-continuation element living in $HW^*(L_{\text{star}(s)}, L_{\text{star}(t)})_{N_\infty^*\mathcal{S}}$ acts as the identity map on \mathbb{Z} under the isomorphism (5.23), and this is precisely what is stated in [Lemma 5.16](#). This completes the proof of (i).

Turning to (ii), first we note that, as $\partial_\infty L_{f^{-1}(-\infty, \epsilon)}$ falls immediately into the stop $N_\infty^*\mathcal{S}$ ([Lemma 5.19](#)), we have

$$(5.25) \quad HW^*(L_{f^{-1}(-\infty, \epsilon)}, L_{f^{-1}(-\infty, -\epsilon)})_{N_\infty^*\mathcal{S}} = HF^*(L_{f^{-1}(-\infty, \epsilon)}, L_{f^{-1}(-\infty, -\epsilon)}).$$

The isotopies of [Lemma 2.13](#) between the (usual) conormals of $f^{-1}(-\infty, \pm\epsilon)$ and the inward cornerings $L_{f^{-1}(-\infty, \pm\epsilon)}$ do not cross each other at infinity, hence

induce an isomorphism between HF^* of the conormals with the above HF^* group of their inward cornerings. These transfer the continuation element associated to the isotopy of conormals from $f^{-1}(-\infty, -\epsilon)$ to $f^{-1}(-\infty, \epsilon)$ to an element of (5.25), which we will call the *cornered continuation element* (multiplication by which is called the cornered continuation map). The image of the cornered continuation element in $\mathcal{W}(T^*M, \Lambda)$ is, by definition, the continuation element between the conormals of $f^{-1}(-\infty, \pm\epsilon)$ (as the isomorphisms between HF^* are compatible with the map to HW^*). It thus suffices to show that the cornered continuation element in (5.25) is the image of the canonical map $1_{f^{-1}(-\infty, -\epsilon)} \rightarrow 1_{f^{-1}(-\infty, \epsilon)}$ under F_S and the isomorphisms of (i). Equivalently, we are to show that the canonical map $1_{f^{-1}(-\infty, -\epsilon)} \rightarrow 1_{f^{-1}(-\infty, \epsilon)}$ agrees under the isomorphisms $HW^*(L_{f^{-1}(-\infty, \pm\epsilon)}, F_S(-)) = 1_{f^{-1}(-\infty, \pm\epsilon)}(-)$ from (i) with the pulled back cornered continuation map. Again by Corollary 5.32, it suffices to make this comparison at the level of cohomology. That is, we are to show that multiplication by the cornered continuation map

$$(5.26) \quad HW^*(L_{f^{-1}(-\infty, -\epsilon)}, L_{\text{star}(s)})_{N_\infty^* S} \rightarrow HW^*(L_{f^{-1}(-\infty, \epsilon)}, L_{\text{star}(s)})_{N_\infty^* S}$$

acts as $\text{id} : \mathbb{Z} \rightarrow \mathbb{Z}$ under the isomorphisms (5.23) (for $\text{star}(s) \subseteq f^{-1}((-\infty, -\epsilon))$). Now, the isomorphisms of both sides with \mathbb{Z} coming from Lemma 5.16 are compatible with pseudo-continuation elements, so by Proposition 5.24 it is equivalent to show that multiplication by the cornered continuation map

$$(5.27) \quad HF^*(L_{f^{-1}(-\infty, -\epsilon)}, L_s) \rightarrow HF^*(L_{f^{-1}(-\infty, \epsilon)}, L_s)$$

acts as the identity on \mathbb{Z} under the isomorphisms of Lemma 5.10 (for $s \subseteq f^{-1}((-\infty, -\epsilon))$). (We replaced HW^* with HF^* using Lemma 5.19.) By the definition given in Lemma 5.10, the “identity on \mathbb{Z} ” map (5.27) is simply the identity map on the single Floer generator we get when unperturbing L_s back to be the cotangent fiber of a point on s . Now the cornered continuation map (5.27) agrees by the last part of [37, Lem. 3.26] with the isomorphism (5.27) from [37, Lem. 3.21] associated to the isotopy from $L_{f^{-1}(-\infty, -\epsilon)}$ to $L_{f^{-1}(-\infty, \epsilon)}$. Since this isotopy takes place far away from L_s , by definition it also acts as the identity on the single Floer generator of both sides. \square

PROPOSITION 5.37. *For any analytic open inclusion of analytic manifolds $M' \hookrightarrow M$, the pushforward functor $\mathcal{W}(T^*M', \Lambda') \rightarrow \mathcal{W}(T^*M, \Lambda)$ for subanalytic singular isotropics with $\Lambda' \supseteq \Lambda \cap S^*M'$ defines a morphism of microlocal Morse theatres.*

Proof. The reasoning of Theorem 5.35 applies without change. \square

6. Examples

6.1. *Cotangent bundles.* Let M be a smooth manifold (assumed connected for sake of notation). The cotangent fiber $F_q \in \mathcal{W}(T^*M)$ generates by Abouzaid [5], [6] when M is closed and by [38, Thm. 1.14] in general.

When M is closed, Abbondandolo–Schwarz [3] and Abouzaid [8] calculated the endomorphism algebra of the fiber as $CW^*(F_q, F_q) = C_{-*}(\Omega_q M)$ (using relative Pin structures as in Section 5.3). The present Theorem 1.1 (which does not depend on any of [3], [8], [6], [5]) gives a proof of this fact for all (not necessarily closed) M :

COROLLARY 6.1. *There is a quasi-isomorphism $CW^*(F_q, F_q) = C_{-*}(\Omega_q M)$. Moreover if $M \subseteq N$ is a codimension zero inclusion, there is a commutative diagram*

$$(6.1) \quad \begin{array}{ccc} CW^*(F_q, F_q)_{T^*M} & \longrightarrow & C_{-*}(\Omega_q M) \\ \downarrow & & \downarrow \\ CW^*(F_q, F_q)_{T^*N} & \longrightarrow & C_{-*}(\Omega_q N), \end{array}$$

where the left-hand vertical arrow is covariant inclusion and the right-hand vertical arrow is induced by pushforward of loops.

Proof. Note that there exists a real analytic structure on M whose induced smooth structure agrees with the given one. Taking $\Lambda = \emptyset$ in Theorem 1.1 gives $\text{Perf } \mathcal{W}(T^*M) = \text{Sh}_\emptyset(M)^c$. It is well known that $\text{Sh}_\emptyset(M)^c = \text{Perf } C_{-*}(\Omega_q M)$, for example, because both are the global sections of the constant cosheaf of linear categories with costalk $\text{Perf } \mathbb{Z}$. Indeed, $U \mapsto U$ is a cosheaf of spaces, equivalently of ∞ -groupoids, which upon linearizing yields $U \mapsto \text{Perf } C_{-*}(\Omega U)$, and $U \mapsto \text{Sh}_\emptyset(U)^c$ is a cosheaf since $U \mapsto \text{Mod Sh}_\emptyset(U)^c = \text{Sh}_\emptyset(U)$ is a sheaf.

We may derive the more precise assertion that $C_{-*}(\Omega_q M)$ is endomorphisms of the cotangent fiber by following a fiber through the equivalence, e.g., by considering the inclusion of the cotangent bundle of a disk, or equivalently by introducing a stop along the conormal of the boundary of a disk and then removing it. \square

6.2. *Plumbings.* Many authors have studied Fukaya categories of plumbings [7], [12], [34] and their sheaf counterparts [17]. Here we compute the wrapped category of a plumbing.

Let Π_{2n} be the Liouville pair $(\mathbb{C}^n, \partial_\infty(\mathbb{R}^n \cup i\mathbb{R}^n))$; we term it the plumbing sector. Plumblings are formed by taking a manifold M (usually disconnected) with spherical boundary $\partial M = \coprod S^{n-1}$, and gluing the Liouville pair $(T^*M, \partial M)$ to some number of plumbing sectors along the spheres.

One can model the wrapped Fukaya category of the plumbing sector directly in sheaf theory: we can view it as the pair $(T^*\mathbb{R}^n, N_\infty^*\{0\})$, and the category $\text{Sh}_{N_\infty^*\{0\}}(\mathbb{R}^n)$ has a well-known description in terms of the Fourier

transform as described in [17]. This category is equivalent to $\mathcal{W}(\Pi_{2n})$ by [Theorem 1.1](#). To apply the gluing results of [38], however, we need to know how the wrapped Fukaya categories of the two boundary sectors include, which is slightly more than what [Theorem 1.1](#) tells us. Hence we give a direct computation of the wrapped Fukaya category of the plumbing sector. Take a positive Reeb pushoff of the boundary of a cotangent fiber in $T^*\mathbb{R}^n$, so it is now the outward conormal of a small ball. Deleting the original cotangent fiber, we obtain the Liouville sector $T^*S^{n-1} \times \mathcal{A}_2$, where \mathcal{A}_2 denotes the Liouville sector $(\mathbb{C}, \{e^{2\pi ik/3}\}_{k=0,1,2\infty})$. We can get back to the plumbing sector Π_{2n} by adding back the missing fiber, which amounts to attaching a Weinstein handle along one of the boundary sectors $T^*(S^{n-1} \times I)$. We may thus deduce from [38, Thms. 1.28, 1.5, and Cor. 1.18] that

LEMMA 6.2. *The category $\mathcal{W}(\Pi_{2n})$ is Morita equivalent to*

$$(6.2) \quad \text{colim}(\text{Perf}(\bullet) \leftarrow \text{Perf } C_*(\Omega S^{n-1}) \rightarrow \text{Perf}(\bullet \rightarrow \bullet) \otimes \text{Perf } C_*(\Omega S^{n-1})).$$

Gluing in the remaining manifolds, we conclude

COROLLARY 6.3. *The wrapped Fukaya category of a plumbing is Morita equivalent to the colimit of the diagram*

(6.3)

$$\begin{array}{c} \coprod \text{Perf}(\bullet) \\ \uparrow \\ \coprod \text{Perf } C_*(\Omega S^{n-1}) \rightarrow \coprod \text{Perf}(\bullet \rightarrow \bullet) \otimes \text{Perf } C_*(\Omega S^{n-1}) \\ \uparrow \\ \coprod \text{Perf } C_*(\Omega S^{n-1}) \longrightarrow \coprod \text{Perf } C_*(\Omega M_i), \end{array}$$

where M_i are the components of M .

6.3. Proper modules and infinitesimal Fukaya categories. Recall that for a dg or A_∞ category \mathcal{C} , we write $\text{Prop } \mathcal{C} := \text{Fun}(\mathcal{C}, \text{Perf } \mathbb{Z})$ for the category of proper (also known as pseudo-perfect) modules. It is immediate from our main result that $\text{Prop } \text{Sh}_\Lambda(M)^c = \text{Prop } \mathcal{W}(T^*M, \Lambda)^{\text{op}}$.

Recall from [Corollary 4.24](#) that any proper $\text{Sh}_\Lambda(M)^c$ -module is representable by an object of $\text{Sh}_\Lambda(M)$ with perfect stalks, i.e., a constructible sheaf in the classical sense. Let us describe some objects in the Fukaya category $\mathcal{W}(T^*M, \Lambda)$ which necessarily give rise to proper modules (and thus to sheaves on M with perfect stalks, microsupported inside Λ).

Definition 6.4. For any stopped Liouville manifold (X, Λ) , we define the *forward stopped subcategory* $\mathcal{W}^\epsilon(X, \Lambda)$ to be the full subcategory of $\mathcal{W}(X, \Lambda)$ generated by Lagrangians which admit a positive wrapping into Λ , meaning $\partial_\infty L$ becomes eventually contained in arbitrarily small neighborhoods of Λ . By [Lemma 5.1](#), such a wrapping is necessarily cofinal.

Example 6.5. If Λ admits a ribbon F (or, alternatively, is itself equal to a Liouville hypersurface F), then $\mathcal{W}^\epsilon(X, \Lambda)$ contains all Lagrangians whose boundary at infinity is contained in a neighborhood of a small negative Reeb pushoff of Λ (or F).

Example 6.6. All compact (exact) Lagrangians are contained in $\mathcal{W}^\epsilon(X, \Lambda)$, as their boundary at infinity \emptyset is wrapped into Λ by the trivial wrapping.

PROPOSITION 6.7. *All objects of $\mathcal{W}^\epsilon(X, \Lambda)$ co-represent proper modules over $\mathcal{W}(X, \Lambda)$; that is, the restriction of the Yoneda embedding $\mathcal{W}(X, \Lambda) \hookrightarrow \text{Mod } \mathcal{W}(X, \Lambda)^{\text{op}}$ to $\mathcal{W}^\epsilon(X, \Lambda)$ has image contained in $\text{Prop } \mathcal{W}(X, \Lambda)^{\text{op}}$.*

Proof. Morphisms in the wrapped category can be computed by cofinally positively wrapping the first factor. Any $L \in \mathcal{W}^\epsilon(X, \Lambda)$ admits such a wrapping $\{L_t\}_{t \geq 0}$ which converges at infinity to Λ . It follows that after some time t , its boundary at infinity stays disjoint at infinity from K , and hence $CW^*(L, K) = CF^*(L_t, K)$ for sufficiently large t . \square

COROLLARY 6.8. *The equivalence $\text{Perf } \mathcal{W}(T^*M, \Lambda)^{\text{op}} = \text{Sh}_\Lambda(M)^c$ sends $\mathcal{W}^\epsilon(T^*M, \Lambda)$ into $\text{Prop } \text{Sh}_\Lambda(M)^c$.*

Recall that for a Whitney triangulation \mathcal{S} , the category $\mathcal{W}(T^*M, N_\infty^* \mathcal{S})^{\text{op}}$ is Morita equivalent to $\mathbb{Z}[\mathcal{S}]$, hence smooth and proper. The generators $L_{\text{star}(s)}$ of $\mathcal{W}(T^*M, N_\infty^* \mathcal{S})$ used to prove this equivalence were shown in that proof to lie in $\mathcal{W}^\epsilon(T^*M, N_\infty^* \mathcal{S})$, so we have

PROPOSITION 6.9. *For a Whitney triangulation \mathcal{S} , the inclusion*

$$\mathcal{W}^\epsilon(T^*M, N_\infty^* \mathcal{S}) \subseteq \mathcal{W}(T^*M, N_\infty^* \mathcal{S})$$

is a Morita equivalence.

Remark 6.10. [Corollary 6.8](#) is very similar to the original Nadler–Zaslow correspondence [66], restricted to Lagrangians with fixed asymptotics. To be more precise, recall that Nadler–Zaslow wish to consider an infinitesimally wrapped Fukaya category $\mathcal{W}^{\text{inf}}(T^*M, \Lambda)$ of Lagrangians “asymptotic at infinity to Λ ” and then show it is equivalent to a category of sheaves on M with microsupport inside Λ .

If Λ is a smooth Legendrian and $\mathcal{W}^{\text{inf}}(T^*M, \Lambda)$ is defined to consist of Lagrangians which are conical at infinity, ending inside Λ , then there is a fully faithful embedding $\mathcal{W}^{\text{inf}}(T^*M, \Lambda) \hookrightarrow \mathcal{W}^\epsilon(T^*M, \Lambda)$, sending a Lagrangian ending inside Λ to its small negative pushoff, as this pushoff tautologically wraps positively back into Λ . Hence [Corollary 6.8](#) recovers a version of [66] when Λ is a smooth Legendrian. One can certainly imagine constructing such an embedding $\mathcal{W}^{\text{inf}}(T^*M, \Lambda) \hookrightarrow \mathcal{W}^\epsilon(T^*M, \Lambda)$ for more general (e.g., subanalytic isotropic) Λ , provided one is given a definition of $\mathcal{W}^{\text{inf}}(T^*M, \Lambda)$ for such Λ .

Remark 6.11. We *do not* know when $\mathcal{W}^\epsilon(T^*M, \Lambda)^{\text{op}} \hookrightarrow \text{Prop Sh}_\Lambda(M)^c$ is a Morita equivalence. Note that the assertion of such an equivalence (for $\mathcal{W}^{\text{inf}}(T^*M, \Lambda)^{\text{op}}$) is *not* made in [62], although that work is occasionally misquoted to suggest that it is. What is actually said is that one can get all objects of $\text{Prop Sh}_\Lambda(M)^c$ from twisted complexes of objects of $\mathcal{W}^{\text{inf}}(T^*M, \Lambda')^{\text{op}}$ for a *possibly larger* Λ' which, as twisted complexes, pair trivially with all Lagrangians contained in a neighborhood of $\Lambda' \setminus \Lambda$. Such Lagrangians might be said to be “Floer theoretically supported away from $\Lambda' \setminus \Lambda$.”

To make a precise statement along the lines of [Remark 6.11](#), realizing a version of the Nadler–Zaslow equivalence, we have

PROPOSITION 6.12. *If \mathcal{S} is any subanalytic Whitney triangulation of compact M with $\Lambda \subseteq N_\infty^*\mathcal{S}$, and \mathcal{D} denotes the collection of linking disks to smooth points of $N_\infty^*\mathcal{S} \setminus \Lambda$, then*

$$(6.4) \quad \text{Prop Sh}_\Lambda(M)^c = \text{Prop } \mathcal{W}(T^*M, \Lambda)^{\text{op}} = (\text{Tw } \mathcal{W}^\epsilon(T^*M, N_\infty^*\mathcal{S})^{\text{op}})_{\text{Ann}(\mathcal{D})},$$

where Tw denotes twisted complexes (i.e., any model for the the pre-triangulated, non idempotent-completed, hull), and the subscript $\text{Ann}(\mathcal{D})$ indicates taking the full subcategory of objects annihilated by $CW^*(-, D) = 0$ for all $D \in \mathcal{D}$.

Proof. For such an \mathcal{S} , the functor $j : \mathcal{W}(T^*M, N_\infty^*\mathcal{S}) \rightarrow \mathcal{W}(T^*M, \Lambda)$ is the quotient by \mathcal{D} by [Theorem 5.4](#). Pullback of modules under any localization is a fully faithful embedding, identifying the category of modules over the localized category with the full subcategory of modules over the original category which annihilate the objects quotiented by (see [Section A.7](#) and [37, Lem. 3.12 and 3.13]). Properness of a module is also clearly equivalent to properness of its pullback. We thus conclude that

$$(6.5) \quad j^* : \text{Prop } \mathcal{W}(T^*M, \Lambda)^{\text{op}} \hookrightarrow \text{Prop } \mathcal{W}(T^*M, N_\infty^*\mathcal{S})^{\text{op}}$$

embeds the former as the full subcategory of the latter annihilating \mathcal{D} .

Now $\mathcal{W}(T^*M, N_\infty^*\mathcal{S})$ (Morita equivalent to $\text{Perf } \mathcal{S}^{\text{op}}$ by [Proposition 5.34](#)) is smooth and proper by [Lemma A.11](#) (since M is compact and thus there are finitely many simplices). Hence $\text{Prop } \mathcal{W}(T^*M, N_\infty^*\mathcal{S})^{\text{op}} = \text{Perf } \mathcal{W}(T^*M, N_\infty^*\mathcal{S})^{\text{op}} = \text{Perf } \mathcal{W}^\epsilon(T^*M, N_\infty^*\mathcal{S})^{\text{op}}$ (by [Proposition 6.9](#)). Finally, we observe that idempotent completion is unnecessary by [Lemma A.10](#), as $\text{Perf } \mathcal{S}$ has a generating exceptional collection. \square

Remark 6.13. For non-compact M , the same proof implies that

$$(6.6) \quad \begin{aligned} \text{Prop Sh}_\Lambda(M)^c &= (\text{Prop } \mathcal{W}^\epsilon(T^*M, N_\infty^*\mathcal{S})^{\text{op}})_{\text{Ann}(\mathcal{D})} \\ &\supseteq (\text{Perf } \mathcal{W}^\epsilon(T^*M, N_\infty^*\mathcal{S})^{\text{op}})_{\text{Ann}(\mathcal{D})} \end{aligned}$$

but the inclusion is not generally an equality. Though, if at least Λ is compact, a similar argument should relate proper modules annihilating co-representatives of the stalks at infinity with the annihilator in $\text{Tw } \mathcal{W}^\epsilon(T^*M, N_\infty^*\mathcal{S})^{\text{op}}$ of \mathcal{D} and the fibers at infinity.

Example 6.14. Let us explain how our “stopped” setup can be used to make ordinary (not wrapped) Floer cohomology calculations using sheaves. Suppose we have two Lagrangians $L, K \subseteq T^*M$ for which $\Lambda := \partial_\infty L \cup \partial_\infty K$ is subanalytic. We are interested in computing $HF^*(L^+, K)$. Thus consider the wrapped category $\mathcal{W}(T^*M, \Lambda)$ and small negative pushoffs $L^-, K^- \in \mathcal{W}(T^*M, \partial_\infty L \cup \partial_\infty K)$, and observe that

$$HF^*(L^+, K) = HW^*(L^-, K^-)_\Lambda.$$

By our main result, the right-hand side can be computed as $\text{Hom}(\mathcal{F}_K, \mathcal{F}_L)$ in the sheaf category $\text{Sh}_\Lambda(M)$, provided we can determine the sheaves \mathcal{F}_L and \mathcal{F}_K to which L^- and K^- are sent by our [Theorem 1.1](#).

Here we make only a few observations regarding how to determine these sheaves. Because linking disks go to microstalks and L^-, K^- are forward stopped, we can see immediately that $\mathcal{F}_L, \mathcal{F}_K$ have microstalk \mathbb{Z} along the respective loci $\partial_\infty L, \partial_\infty K \subseteq \Lambda$. For the same reason, for p away from the front projection of $\Lambda = \partial_\infty L \cup \partial_\infty K$, we have

$$\mathcal{F}_L|_p \cong CF^*(L, T_p^*M), \quad \mathcal{F}_K|_p \cong CF^*(K, T_p^*M).$$

In some cases, e.g., in the case where L intersects every cotangent fiber either once or not at all, this data already suffices to determine \mathcal{F}_L . In particular, this situation occurs in [83], where sheaf calculations are made exhibiting cluster transformations arising from comparing different fillings of Legendrian knots. The present discussion suffices to translate those calculations into calculations in Lagrangian Floer theory.

6.4. Legendrians and constructible sheaves.

COROLLARY 6.15. *Let $\Lambda \subseteq J^1\mathbb{R}^n \subseteq S^*\mathbb{R}^{n+1}$ be a smooth compact Legendrian. Let $D = D_1 \sqcup \dots \sqcup D_n$ be a disjoint union of linking disks at distinct points of Λ , at least one on each connected component. Consider the algebra*

$$(6.7) \quad \mathcal{A}_\Lambda := CW^*(D, D)_{T^*\mathbb{R}^{n+1}, \Lambda} = \bigoplus_{i, j=1}^n CW^*(D_i, D_j)_{T^*\mathbb{R}^{n+1}, \Lambda}.$$

Then $\text{Mod } \mathcal{A}_\Lambda^{\text{op}}$ is equivalent to the category $\text{Sh}_\Lambda(T^\mathbb{R}^{n+1})_0$ of sheaves micro-supported inside Λ and with vanishing stalk at infinity.*

This equivalence identifies the microstalk along Λ near D_i with the direct summand of the forgetful functor $\text{Mod } \mathcal{A}_\Lambda^{\text{op}} \rightarrow \text{Mod } \mathbb{Z}$ corresponding to the

idempotent $e_i := \mathbf{1}_{D_i} \in CW^*(D_i, D_i) \subseteq \mathcal{A}_\Lambda$ (e_1, \dots, e_n are orthogonal idempotents summing to the identity). Hence $\text{Prop } \mathcal{A}_\Lambda^{\text{op}}$ is equivalent to the subcategory of $\text{Sh}_\Lambda(T^*\mathbb{R}^{n+1})_0$ of objects with perfect microstalks along Λ (or, equivalently, with perfect stalks).

Proof. Our generation results [38, Thm. 1.14] imply that $\mathcal{W}(T^*\mathbb{R}^{n+1}, \Lambda)$ is generated by D_1, \dots, D_n and a cotangent fiber F near infinity. Because we assume that $\Lambda \subseteq J^1\mathbb{R}^n$, the cotangent fiber at negative (in the last coordinate) infinity can be cofinally positively wrapped without intersecting Λ , and likewise the (isomorphic) cotangent fiber at positive infinity can be cofinally negatively wrapped without intersecting Λ . These large wrappings are conormals to large disks in \mathbb{R}^{n+1} containing the projection of Λ ; they thus have vanishing wrapped Floer cohomology (in both directions) with the linking disks D_i to Λ . Thus $D = D_1 \sqcup \dots \sqcup D_n$ and F are orthogonal objects of $\mathcal{W}(T^*\mathbb{R}^{n+1}, \Lambda)$.

Denote by $\mu = \mu_1 \oplus \dots \oplus \mu_n$ and $\sigma \in \text{Sh}_\Lambda(T^*\mathbb{R}^{n+1})^c$ the objects corresponding to $D = D_1 \oplus \dots \oplus D_n$ and F . They are orthogonal and have endomorphism algebras $\mathcal{A}_\Lambda^{\text{op}}$ and \mathbb{Z} , respectively.

We have $\text{Sh}_\Lambda(T^*\mathbb{R}^{n+1}) = \text{Mod } \text{Sh}_\Lambda(T^*\mathbb{R}^{n+1})^c = \text{Mod } \mathcal{W}(T^*\mathbb{R}^{n+1}, \Lambda)^{\text{op}} = \text{Mod } \mathcal{A}_\Lambda^{\text{op}} \oplus \text{Mod } \mathbb{Z}$, and this equivalence is given concretely by $\mathcal{F} \mapsto \text{Hom}(\mu, \mathcal{F}) \oplus \text{Hom}(\sigma, \mathcal{F})$. By Theorem 1.1, $\text{Hom}(\mu, \mathcal{F}) = \text{Hom}(\mu_1, \mathcal{F}) \oplus \dots \oplus \text{Hom}(\mu_n, \mathcal{F})$ is the direct sum of microstalks along Λ near D_1, \dots, D_n , and $\text{Hom}(\sigma, \mathcal{F})$ is the stalk at infinity.

To see that having perfect stalks is equivalent to having perfect microstalks along Λ for objects of $\text{Sh}_\Lambda(T^*\mathbb{R}^{n+1})_0$, argue as follows. Suppose microstalks are perfect. Stalks are computed by $\text{Hom}(\mathbb{Z}_{B_\epsilon(x)}, \mathcal{F})$ for some sufficiently small $\epsilon > 0$ (in terms of Λ), since changing ϵ is non-characteristic by Whitney's condition (b) for a subanalytic Whitney stratification \mathcal{S} whose conormal contains Λ . Now moving $B_\epsilon(x)$ generically to infinity picks up some number of microstalks when its conormal passes through Λ (transversally), and eventually gives zero since the stalk of \mathcal{F} near infinity vanishes. Thus perfect microstalks implies perfect stalks. That perfect stalks implies perfect microstalks was proven in Corollary 4.24. \square

Let us comment on the relation of the above result to the “augmentations are sheaves” statement in [84], [71] (and later developments such as [74], [13]). There is an evident similarity: both relate augmentations of an algebra associated to a Legendrian to categories of sheaves microsupported in that Legendrian. But they are not exactly the same: the algebra \mathcal{A}_Λ is not by definition the Chekanov–Eliashberg dga, and moreover in [71] the category of augmentations is defined by a somewhat complicated procedure, not just as proper modules over a dga. Also in [71], the authors restrict attention to augmentations, i.e., 1-dimensional representations of the dga, whereas the above

result concerns the entire representation category (the underlying \mathbb{Z} -module of the representation being the microstalk), specializing to a comparison of rank k representations with rank k microstalk sheaves for every k .

In fact, \mathcal{A}_Λ was conjectured by Sylvan to be a version of the Chekanov–Eliashberg dga with enhanced $C_*(\Omega\Lambda)$ coefficients. A precise statement comparing \mathcal{A}_Λ to such a generalized “loop space dga” can be found in [30, Conj. 3], where it is explained that the comparison should follow from a slight variant of the surgery techniques of [19], [29]. The relation between the multiple copy construction of [71] and the loop space dga can also be extracted from [30].

Finally we note that a version of the above discussion serves to translate between the arguments of [80] and [31].

6.5. Fukaya–Seidel categories of cotangent bundles. Let $W : T^*M \rightarrow \mathbb{C}$ be an exact symplectic fibration with singularities. The associated Fukaya–Seidel category is by (our) definition $\mathcal{W}(T^*M, W^{-1}(-\infty))$. According to [38, Cor. 3.9], retracting the stop to its core does not affect the category:

$$\mathcal{W}(T^*M, W^{-1}(-\infty)) = \mathcal{W}(T^*M, \mathfrak{c}_{W^{-1}(-\infty)}).$$

Thus if the fiber is Weinstein, then we may calculate the corresponding Fukaya–Seidel category using [Theorem 1.1](#) (provided the core is subanalytic).

In particular, the sheaf theoretic work on mirror symmetry for toric varieties may now be translated into assertions regarding the wrapped Fukaya category. Recall that for any n -dimensional toric variety \mathbf{T} , [35] introduced a certain Lagrangian $\Lambda_{\mathbf{T}} \subseteq T^*(S^1)^n$. They conjectured,¹⁰ and [54] proved, that $\mathrm{Sh}_{\partial_\infty \Lambda_{\mathbf{T}}}((S^1)^n)^c = \mathrm{Coh}(\mathbf{T})$, where we use Coh to denote the dg category of coherent complexes. By [Theorem 1.1](#), we may conclude

COROLLARY 6.16. *We have $\mathrm{Perf}(\mathcal{W}(T^*(S^1)^n, \partial_\infty \Lambda_{\mathbf{T}}))^{\mathrm{op}} = \mathrm{Coh}(\mathbf{T})$.*

When \mathbf{T} is smooth and Fano, it was expected that the $\mathrm{Coh}(\mathbf{T})$ should be equivalent to the Fukaya–Seidel category of the mirror Hori–Vafa superpotential [44]. To compare this expectation with [Corollary 6.16](#), it suffices to show that $\partial_\infty \Lambda_{\mathbf{T}}$ is in fact the core of the fiber of said superpotential in the Fano case. This is shown under certain hypotheses in [36] and in general in [91]. We summarize the above discussion in the right column of (7.31).

These results may be compared with [4], which for smooth projective \mathbf{T} gives a fully faithful embedding of $\mathrm{Perf}(\mathbf{T})$ into an infinitesimal Fukaya category for the superpotential. In the Fano case, this is recovered and upgraded to an equivalence by taking proper modules of the formulation in [36].

¹⁰Strictly speaking, they conjectured the proper module version of this statement.

Note that [Corollary 6.16](#) gives an equivalence in the general (non-Fano, non-compact, singular, and stacky) case, although this equivalence is not yet formulated in terms of a superpotential. Such a formulation is known to be somewhat subtle, requiring the exclusion of some critical values; see [\[16, §5\]](#) or [\[4\]](#). It may be interesting to explore this using the present methods.

7. Partially wrapped Fukaya categories and microlocal sheaves

The purpose of this section is to prove [Theorem 1.4](#). The reasoning in this section depends only on the statement of [Theorem 1.1](#), together with various results from [\[38\]](#) and [\[65\]](#); as such, it can be read independently of previous sections of this article.

The main point in the derivation of [Theorem 1.4](#) from [Theorem 1.1](#) is to properly exploit, on both the Fukaya side and the sheaf side, the “doubling trick,” which allows one to embed the category associated (on either side) to a Liouville manifold (possibly relative a singular isotropic stop) into the category associated to a cotangent bundle relative an appropriate “doubled” stop obtained from an embedding of (a stabilization of) the given Liouville manifold into the co-sphere bundle. The use of such an embedding to reduce to cotangent bundles was advocated for on the sheaf side in [\[81\]](#). The doubling trick has appeared in various forms on the Fukaya side [\[38\]](#), [\[89\]](#), and we develop it systematically below. It has been used on the sheaf side in [\[42\]](#), [\[65\]](#) to embed categories of *microlocal* sheaves into categories of (usual) sheaves, and we will use it for the same purpose here.

While our eventual appeal to [Theorem 1.1](#) will require the relative core in question to be subanalytic isotropic, most of the intermediate results of this section require much weaker assumptions. A subset Λ of a contact (resp. symplectic) manifold will be called *mostly Legendrian* (resp. *Lagrangian*) [\[38, Def. 1.7\]](#) if and only if the complement $\Lambda^{\text{subcrit}} := \Lambda \setminus \Lambda^{\text{crit}}$ of the open locus $\Lambda^{\text{crit}} \subseteq \Lambda$, where Λ is a smooth Legendrian (resp. Lagrangian), can be covered by the smooth image of a second countable smooth manifold of dimension strictly less than Legendrian (resp. Lagrangian). A conical mostly Lagrangian subset is (locally) the cone over a mostly Legendrian subset. (To show this, intersect with a generic contact type hypersurface). A subanalytic isotropic subset is mostly Legendrian/Lagrangian.

7.1. Homological cocores. We begin by introducing homological cocores, which are a simultaneous generalization of linking disks and cocores. They are analogous to co-representatives of microstalks in the sheaf theoretic context. At various points in our arguments below, it will be relevant to assume a given Liouville manifold X “admits homological cocores” in the sense defined below. (In fact, admitting homological cocores is most naturally a condition

on a pair (X, Λ) , which turns out to be independent of Λ and invariant under deformations of X .) Every Weinstein manifold admits homological cocores.

We begin by recalling a special case of the Künneth embedding from [38]. Let (X, Λ) be a stopped Liouville manifold. We have a Künneth functor [38, Thm. 1.5]

$$(7.1) \quad \mathcal{W}(X, \Lambda) \hookrightarrow \mathcal{W}((X, \Lambda) \times (\mathbb{C}, \infty \cup e^{i[\frac{\pi}{2}, \frac{3\pi}{2}]} \infty))$$

given by multiplication by the linking disk $D_\infty \in \mathcal{W}(\mathbb{C}, \infty \cup e^{i[\frac{\pi}{2}, \frac{3\pi}{2}]} \infty)$ at $\infty \in \partial_\infty \mathbb{C}$; it is fully faithful since the endomorphism algebra of D_∞ is \mathbb{Z} . We also have a fully faithful embedding

$$(7.2) \quad \mathcal{W}((X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty)) \subseteq \mathcal{W}((X, \Lambda) \times (\mathbb{C}, \infty \cup e^{i[\frac{\pi}{2}, \frac{3\pi}{2}]} \infty))$$

by [38, Lem. 3.7]. (To be explicit, $(X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty) = (X \times \mathbb{C}_{\text{Re} \geq 0}, (\mathfrak{c}_X \times \infty) \cup (\Lambda \times \mathbb{R}_{>0}))$.) The image of the Künneth functor (7.1) is evidently contained in this full subcategory (7.2), so we obtain a functor

$$(7.3) \quad \mathcal{W}(X, \Lambda) \hookrightarrow \mathcal{W}((X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$$

which will be used throughout this section.

Definition 7.1 (Homological cocore). Let (X, Λ) be a stopped Liouville manifold whose relative core $\mathfrak{c}_{X, \Lambda}$ is mostly Lagrangian. A *homological cocore* at a smooth Lagrangian point $p \in \mathfrak{c}_{X, \Lambda}$ is an object of $\text{Perf } \mathcal{W}(X, \Lambda)$ whose image under (Perf of) the Künneth embedding (7.3) is the linking disk at $p \times \infty \in \mathfrak{c}_{X, \Lambda} \times \infty$.

Recall from [38, Proof of Thm. 1.14] that if $L \subseteq X$ is exact, cylindrical at infinity, and intersects $\mathfrak{c}_{X, \Lambda}$ precisely once, transversely, at a smooth Lagrangian point $p \in \mathfrak{c}_{X, \Lambda}$, then L is a homological cocore at p . In particular, (properly embedded) cocores of critical Weinstein handles are homological cocores. Also recall from [38, §9.1] that the linking disk at a point of Λ is a homological cocore at the corresponding point of $\mathfrak{c}_{X, \Lambda}$.

Definition 7.2. We say that (X, Λ) (with $\mathfrak{c}_{X, \Lambda}$ mostly Lagrangian) *admits homological cocores* if and only if every smooth Lagrangian point of $\mathfrak{c}_{X, \Lambda}$ has a homological cocore.

It follows from stop removal and the vanishing of $\mathcal{W}(X \times \mathbb{C}_{\text{Re} \geq 0})$ [38, Lem. 9.1] that the linking disks to $\mathfrak{c}_{X, \Lambda} \times \infty$ split-generate $\mathcal{W}((X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$. Hence (X, Λ) admits homological cocores if and only if the Künneth embedding (7.3) is a Morita equivalence (and in this case $\text{Perf } \mathcal{W}(X, \Lambda)$ is split-generated by the homological cocores). In fact, we have the following equivalent characterizations of admitting homological cocores:

PROPOSITION 7.3. *For a stopped Liouville manifold (X, Λ) whose relative core $\mathfrak{c}_{X, \Lambda}$ is mostly Lagrangian, the following are equivalent:*

- (X, Λ) admits homological cocores;
- the Künneth embedding (7.3) is a Morita equivalence;
- the Künneth embedding is a Morita equivalence:

$$(7.4) \quad \mathcal{W}(X, \Lambda) \hookrightarrow \mathcal{W}((X, \Lambda) \times (\mathbb{C}, \pm\infty));$$

- the Künneth embedding is a Morita equivalence:

$$(7.5) \quad \mathcal{W}(X) \hookrightarrow \mathcal{W}(X \times (\mathbb{C}_{\text{Re} \geq 0}, \infty));$$

- the Künneth embedding is a Morita equivalence:

$$(7.6) \quad \mathcal{W}(X) \hookrightarrow \mathcal{W}(X \times (\mathbb{C}, \pm\infty)).$$

Proof. The equivalence of admitting homological cocores and the Künneth embedding (7.3) being a Morita equivalence was already argued for above.

We argue that (7.3) and (7.5) are equivalent. These are the statements that the categories $\mathcal{W}((X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$ and $\mathcal{W}(X \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$ (respectively) are split-generated by Lagrangians of the form $L \times [i\mathbb{R}]$. Taking L to be a linking disk of Λ , we see that the linking disks to $\Lambda \times \mathbb{R}_{>0}$ inside $\mathcal{W}((X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$ are of the desired form. Therefore it is split-generated by Lagrangians of the form $L \times [i\mathbb{R}]$ if and only if its quotient by the linking disks to $\Lambda \times \mathbb{R}_{>0}$ is split-generated by these objects, and this quotient is precisely $\mathcal{W}(X \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$ by stop removal. The same argument shows that (7.4) and (7.6) are equivalent.

We argue that (7.5) and (7.6) are equivalent. They are the statements that the categories $\mathcal{W}(X \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$ and $\mathcal{W}(X \times (\mathbb{C}, \pm\infty))$ (respectively) are split-generated by Lagrangians of the form $L \times [i\mathbb{R}]$. These statements are equivalent since the natural functor $\mathcal{W}(X \times (\mathbb{C}_{\text{Re} \geq 0}, \infty)) \rightarrow \mathcal{W}(X \times (\mathbb{C}, \pm\infty))$ is an equivalence by [38, Cor. 3.9]. \square

Note that condition (7.6) does not involve Λ and is invariant under deforming X ; it holds whenever X is Weinstein by [38, Cor. 1.18].

If $\mathfrak{c}_{X, \Lambda}$ is mostly Lagrangian, the stabilization $(X, \Lambda) \times (\mathbb{C}, \pm\infty)$ admits homological cocores since every component of the smooth Lagrangian locus of $\mathfrak{c}_{(X, \Lambda) \times (\mathbb{C}, \pm\infty)} = \mathfrak{c}_{X, \Lambda} \times \mathbb{R}$ is unbounded. It follows that the Künneth embedding

$$(7.7) \quad \mathcal{W}((X, \Lambda) \times (\mathbb{C}, \pm\infty)^k) \hookrightarrow \mathcal{W}((X, \Lambda) \times (\mathbb{C}, \pm\infty)^{k+1})$$

is a Morita equivalence for every $k > 0$.

7.2. Liouville hypersurfaces. Recall that a *Liouville hypersurface embedding* $X \hookrightarrow Y$ is a codimension one embedding of a Liouville domain (X, λ) into a contact manifold (Y, ξ) such that there exists a contact form α on (Y, ξ) whose restriction to X coincides with λ . A *Liouville pair* (Z, X) is a Liouville manifold Z together with a Liouville hypersurface embedding $X \hookrightarrow \partial_\infty Z$.

We will often abuse terminology and speak of a Liouville hypersurface embedding of a Liouville *manifold* into a contact manifold to mean a Liouville hypersurface embedding of a Liouville domain whose completion is the given Liouville manifold.

Here we record two real analytic approximation results for later use.

LEMMA 7.4. *Any codimension zero smooth embedding of real analytic contact manifolds $(U, \xi_U) \hookrightarrow (Y, \xi_Y)$ can be smoothly approximated over compact subsets of the domain by real analytic embeddings.*

Proof. First, we approximate the given embedding by a real analytic map f which does not necessarily respect contact structures. We now have two real analytic contact structures $f^*\xi_Y$ and ξ_U on U which are C^∞ -close. Interpolating linearly yields a real analytic family of real analytic contact structures ξ_t interpolating between $f^*\xi_Y$ and ξ_U . By Gray's theorem, we obtain a real analytic family of real analytic vector fields V_t defined uniquely by the properties $V_t \in \xi_t$ and $\mathcal{L}_{V_t}\xi_t = \frac{d}{dt}\xi_t$. The total flow of this family V_t thus defines a real analytic diffeomorphism of U (possibly defined only on a large compact subset due to lack of completeness) carrying ξ_U to $f^*\xi_Y$. Pre-composing f by this diffeomorphism gives the desired real analytic map. \square

COROLLARY 7.5. *Let (X, λ) be a real analytic Liouville domain, and let (Y, ξ) be a real analytic contact manifold. Any Liouville hypersurface embedding $X \hookrightarrow Y$ can be smoothly approximated by real analytic Liouville hypersurface embeddings.*

Proof. Given a Liouville hypersurface embedding $X \hookrightarrow Y$, there is an induced codimension zero inclusion of contact manifolds $X \times [0, 1] \hookrightarrow Y$ to which we may apply Lemma 7.4. \square

We now study the question of when a Liouville manifold X admits a Liouville hypersurface embedding $X \hookrightarrow S^*M$. Such an embedding determines three pieces of “formal” data:¹¹

- (i) a smooth map $f : X \rightarrow M$;
- (ii) a splitting $f^*TM = B \oplus \underline{\mathbb{R}}$;
- (iii) an isomorphism of complex vector bundles $TX = B \otimes_{\mathbb{R}} \mathbb{C}$.

The first two pieces of data are equivalent to a homotopy class of smooth maps $X \rightarrow S^*M$. Indeed, up to contractible choices, a lift of $f : X \rightarrow M$ to S^*M is the same as a non-vanishing section of f^*TM , which is the same as a trivialized subbundle $\underline{\mathbb{R}} \subseteq f^*TM$, which is the same as a splitting $f^*TM = \underline{\mathbb{R}} \oplus B$. The isomorphism $TX = B \otimes_{\mathbb{R}} \mathbb{C}$ comes from the derivative of the embedding, which

¹¹The term “formal” has a precise meaning in the context of the h -principle; see [33, §5.3].

identifies TX with the pullback of the contact distribution on S^*M . There is an existence h -principle for Liouville hypersurface embeddings (under a certain “half-dimensional” hypothesis on the core); namely,

LEMMA 7.6. *Let X be a Liouville manifold whose core \mathbf{c}_X is contained in a finite union of locally closed submanifolds of dimension at most half the dimension of X . Every triple of formal data as above comes from a Liouville hypersurface embedding $X \hookrightarrow S^*M$.*

Proof. The formal data is (homotopy) equivalent to a smooth map $p : X \rightarrow S^*M$ together with an isomorphism $q : TX = p^*\xi$, where ξ is the contact distribution of S^*M . Equivalently, it is the data of a smooth map $p : X \times [0, 1] \rightarrow S^*M$ together with an *isocontact isomorphism* $q : T(X \times [0, 1]) = p^*TS^*M$ (i.e., an isomorphism respecting contact distributions and their conformal symplectic structure). The h -principle [33, 16.1.1] now guarantees that the pair (p, q) is homotopic to an *isocontact immersion*, i.e., one for which p is an immersion which pulls back the contact structure on S^*M to the contact structure $\lambda + dt$ on $X \times [0, 1]$. The assumption on the core \mathbf{c}_X ensures that a generic perturbation of p is an embedding in a small neighborhood of $\mathbf{c}_X \times \{\frac{1}{2}\}$. \square

COROLLARY 7.7. *Let X be a Liouville manifold. For any stable polarization $TX = B \otimes_{\mathbb{R}} \mathbb{C}$, there is a Liouville hypersurface embedding of $X \times \mathbb{C}^k$ (some $k < \infty$) into some S^*M , compatible with stable polarizations.*

Proof. In view of [Lemma 7.6](#) (whose hypothesis is trivially satisfied for $X \times \mathbb{C}^k$ once $k \geq \frac{1}{2} \dim X$), it suffices to show that there exists a manifold N and a map $f : X \rightarrow N$ such that f^*TN and B are stably isomorphic (i.e., isomorphic after direct summing with some \mathbb{R}^m). To see that this is true, note that the tangent bundle to the Grassmannian of n -planes in \mathbb{R}^N is (stably) inverse to the tautological vector bundle. \square

Remark 7.8. [Corollary 7.7](#) concerns the product polarization of $X \times \mathbb{C}^k$. By contrast, there always exists a twisted stabilization, i.e., the total space of an arbitrary polarized symplectic vector bundle over X (equivalently, the complexification of a rank k real vector bundle) which embeds into $S^*\mathbb{R}^N$ as in [81], [65]. In the present article, we need to restrict to untwisted stabilization because in [38] we have only proven an untwisted Künneth theorem. Meanwhile in the sheaf-theoretic settings of [81], [65], the corresponding twisted Künneth result is a formality, and it is convenient to embed into $S^*\mathbb{R}^N$ rather than some S^*M in order to have (homotopical) uniqueness of embeddings.

7.3. Doubling I: Fully faithful embeddings of Fukaya categories. We first recall the doubling trick in the “absolute” (i.e., no stop) setting. Consider a

Liouville pair (Z, X) . Grading/orientation data on Z determines such data on X by restriction; when $Z = T^*M$, our primary case of interest, this means that X is equipped with the polarization induced from the Legendrian foliation of S^*M by co-spheres. With respect to such compatible data, there is a functor $\mathcal{W}(X) \rightarrow \mathcal{W}(Z, \mathfrak{c}_X)$ obtained by composing the Künneth map $\mathcal{W}(X) \rightarrow \mathcal{W}(X \times \mathbb{C}_{\text{Re} \geq 0}, \mathfrak{c}_X \times \{\infty\})$ with the canonical neighborhood $X \times \mathbb{C}_{\text{Re} \geq 0} \hookrightarrow Z$ of the Liouville hypersurface X (an embedding of Liouville sectors).

We now consider the double $D(\mathfrak{c}_X) := \mathfrak{c}_X \sqcup \mathfrak{c}_X^\epsilon$, where \mathfrak{c}_X^ϵ denotes a small positive pushoff of \mathfrak{c}_X . There is a functor

$$(7.8) \quad \mathcal{W}(X) \rightarrow \mathcal{W}(Z, \mathfrak{c}_X \sqcup \mathfrak{c}_X^\epsilon)$$

defined by including $(X \times \mathbb{C}_{\text{Re} \geq 0}, \mathfrak{c}_X \times \{\infty\})$ into $(T^*M, \mathfrak{c}_X \sqcup \mathfrak{c}_X^\epsilon)$ as the canonical neighborhood of the first copy of \mathfrak{c}_X inside $D(\mathfrak{c}_X)$. When \mathfrak{c}_X is mostly Lagrangian, this functor evidently sends a homological cocore at a point of \mathfrak{c}_X to the linking disk at the corresponding point of the first copy of \mathfrak{c}_X inside the double.

PROPOSITION 7.9. *The functor (7.8) is fully faithful.*

Proof. It is asserted in [38, Ex. 10.7] that the covariant pushforward $\mathcal{W}(X) \rightarrow \mathcal{W}(Z, X \sqcup X^\epsilon)$ is fully faithful, where X^ϵ denotes a small positive pushoff of X . Combining this with the fact that the functor $\mathcal{W}(Z, X \sqcup X^\epsilon) \rightarrow \mathcal{W}(Z, \mathfrak{c}_X \sqcup \mathfrak{c}_X^\epsilon)$ is an equivalence [38, Cor. 3.9], we conclude that (7.8) is fully faithful. \square

We now explain the doubling trick in the presence of a stop (“relative doubling”).

Construction 7.10 (Doubling the relative core of a Liouville hypersurface). Let (Z, X) be a Liouville pair, and let $\Lambda \subseteq \partial_\infty X$ be a stop. We will define the double $D(\mathfrak{c}_{X,\Lambda}) \subseteq \partial_\infty Z$. The double is contained in a small neighborhood of X , so it suffices to define it as a subset of $\partial_\infty(X \times \mathbb{C}_{\text{Re} \geq 0})$ (and then push forward under the standard neighborhood $X \times \mathbb{C}_{\text{Re} \geq 0} \hookrightarrow Z$ of X inside Z).

The double $D(\mathfrak{c}_{X,\Lambda}) \subseteq \partial_\infty(X \times \mathbb{C}_{\text{Re} \geq 0})$ is the stop of the product of stopped Liouville manifolds

$$(7.9) \quad (X, \Lambda) \times (\mathbb{C}, \{\pm i\infty\}) = (X \times \mathbb{C}, (\mathfrak{c}_X \times \{\pm i\infty\}) \cup (\Lambda \times i\mathbb{R})),$$

which indeed lies inside $\partial_\infty(X \times \mathbb{C}_{\text{Re} \geq 0}) \subseteq \partial_\infty(X \times \mathbb{C})$. The double $D(\mathfrak{c}_{X,\Lambda})$ is evidently comprised of a “first copy” of $\mathfrak{c}_{X,\Lambda}$, namely $(\mathfrak{c}_X \times \{-\infty\}) \cup (\Lambda \times i\mathbb{R}_{<0})$, and a “second copy” of $\mathfrak{c}_{X,\Lambda}$, namely $(\mathfrak{c}_X \times \{+\infty\}) \cup (\Lambda \times i\mathbb{R}_{>0})$, joined along their common boundary $\Lambda = \Lambda \times 0$.

The fact that $D(\mathfrak{c}_{X,\Lambda}) \subseteq \partial_\infty(X \times \mathbb{C}_{\text{Re} \geq 0})$ lies on the boundary poses no issue for defining $D(\mathfrak{c}_{X,\Lambda}) \subseteq \partial_\infty Z$ as its image under (the action on boundaries

at infinity of) $X \times \mathbb{C}_{\text{Re} \geq 0} \hookrightarrow Z$. We will, however, want to consider $\mathcal{W}(X \times \mathbb{C}_{\text{Re} \geq 0}, D(\mathfrak{c}_{X,\Lambda}))$, and for the purpose of defining this category, we implicitly push $D(\mathfrak{c}_{X,\Lambda})$ inward using a choice of contact vector field transverse to the boundary. Alternatively, we could use a different Liouville structure on $\mathbb{C}_{\text{Re} \geq 0}$ which is strictly isomorphic to $T^*[0, \epsilon)$ near the boundary, which makes pushing easy (simple translation).

Let us now generalize the functor (7.8) and [Proposition 7.9](#) to the relative setting. We consider the composition

$$(7.10) \quad \begin{aligned} \mathcal{W}(X, \Lambda) &\xrightarrow{(7.3)} \mathcal{W}((X, \Lambda) \times (\mathbb{C}_{\text{Im} \leq 0}, -i\infty)) \\ &\rightarrow \mathcal{W}(X \times \mathbb{C}, (\mathfrak{c}_X \times \{-\infty, \pm i\infty\}) \cup (\Lambda \times i\mathbb{R})). \end{aligned}$$

The target category is identified with $\mathcal{W}(X \times \mathbb{C}_{\text{Re} \geq 0}, D(\mathfrak{c}_{X,\Lambda}))$ by [\[38, Cor. 3.9\]](#), so we obtain a canonical functor

$$(7.11) \quad \mathcal{W}(X, \Lambda) \rightarrow \mathcal{W}(X \times \mathbb{C}_{\text{Re} \geq 0}, D(\mathfrak{c}_{X,\Lambda})),$$

and hence composing with any inclusion $X \times \mathbb{C}_{\text{Re} \geq 0} \hookrightarrow Z$, a functor

$$(7.12) \quad \mathcal{W}(X, \Lambda) \rightarrow \mathcal{W}(Z, D(\mathfrak{c}_{X,\Lambda})).$$

When $\mathfrak{c}_{X,\Lambda}$ is mostly Lagrangian, this functor evidently sends a homological cocore at a point of $\mathfrak{c}_{X,\Lambda}$ to the linking disk at the corresponding point of the first copy of $\mathfrak{c}_{X,\Lambda}$ inside the double.

PROPOSITION 7.11. *The functor (7.12) is fully faithful.*

Proof. Appealing to the definition of the functor (7.3), the functor (7.12) is the composition

$$(7.13) \quad \begin{aligned} \mathcal{W}(X, \Lambda) &\xrightarrow{(7.1)} \mathcal{W}((X, \Lambda) \times (\mathbb{C}, -i\infty \cup e^{i[0,\pi]}\infty)) \\ &\rightarrow \mathcal{W}(X \times \mathbb{C}, (\mathfrak{c}_X \times \{-\infty, \pm i\infty\}) \cup (\Lambda \times i\mathbb{R})) \rightarrow \mathcal{W}(Z, D(\mathfrak{c}_{X,\Lambda})) \end{aligned}$$

of Künneth, stop removal, and pushforward. Given that Künneth is fully faithful, it suffices to show that the composition of the latter two functors is fully faithful when restricted to product objects $L \times D_{-i\infty} \subseteq (X, \Lambda) \times (\mathbb{C}, -i\infty \cup e^{i[0,\pi]}\infty)$. In fact, we will show they are both fully faithful on such objects.

To show full faithfulness comes down to understanding cofinal wrappings. It was shown in [\[38, §7.4\]](#) that products of cofinal wrappings are cofinal. (This was the basis for full faithfulness of the Künneth functor.) But the results of [\[38, §7.4\]](#) are better: they in fact show that if wrappings of $L \subseteq (X, \Lambda)$ and $D_{-i\infty} \subseteq (\mathbb{C}, -i\infty \cup e^{i[0,\pi]}\infty)$ satisfy the cofinality criterion [\[38, Lem. 2.2\]](#), then so does their product inside $(X, \Lambda) \times (\mathbb{C}, -i\infty \cup e^{i[0,\pi]}\infty)$.

Now the cofinality criterion is robust in an important way. Choose wrappings of $L \subseteq (X, \Lambda)$ and $D_{-i\infty} \subseteq (\mathbb{C}, \{\pm i\infty\})$ satisfying the cofinality criterion.

Their product satisfies the cofinality criterion in $(X, \Lambda) \times (\mathbb{C}, \{\pm i\infty\})$, hence also in $(X \times \mathbb{C}, (\mathfrak{c}_X \times \{-\infty, \pm i\infty\}) \cup (\Lambda \times i\mathbb{R}))$, as it stays away from the additional stop $\mathfrak{c}_X \times \{-\infty\}$. Satisfaction of the cofinality criterion is also preserved under cutting out a neighborhood of this additional stop at $\mathfrak{c}_X \times \{-\infty\}$ and embedding into $\mathcal{W}(Z, D(\mathfrak{c}_{X, \Lambda}))$.

We have thus described cofinal wrappings of product objects in the three categories in (7.13) other than $\mathcal{W}(X, \Lambda)$. The desired full faithfulness results follow using [37, Lem. 3.20]. \square

7.4. A first comparison. We now combine the doubling trick embeddings with [Theorem 1.1](#) to arrive at a first sheaf theoretic description of some partially wrapped Fukaya categories. Note that the doubling construction works real analytically by appealing to [Corollary 7.4](#) to make the contactomorphisms involved in [Construction 7.10](#) real analytic. (In the below, we tacitly assume that doubling takes place real analytically in this sense.)

Combining [Proposition 7.11](#) and [Theorem 1.1](#), we obtain the following:

COROLLARY 7.12. *Let (X, Λ) be a stopped real analytic Liouville manifold whose relative core $\mathfrak{c}_{X, \Lambda}$ is subanalytic isotropic. Let M be a real analytic manifold and $X \hookrightarrow S^*M$ an analytic Liouville hypersurface embedding. There is a fully faithful embedding*

$$(7.14) \quad \mathcal{W}(X, \Lambda)^{\text{op}} \hookrightarrow \text{Sh}_{D(\mathfrak{c}_{X, \Lambda})}(M)^c$$

which sends homological cocores of (X, Λ) to co-representatives of microstalks at the corresponding points of the first copy of $\mathfrak{c}_{X, \Lambda}$ inside $D(\mathfrak{c}_{X, \Lambda})$.

In particular, if (X, Λ) admits homological cocores, then (7.14) is a Morita equivalence onto the full subcategory split-generated by co-representatives of the microstalks at smooth points of the first copy of $\mathfrak{c}_{X, \Lambda}$ inside $D(\mathfrak{c}_{X, \Lambda})$.

In order to bridge the gap between [Corollary 7.12](#) and [Theorem 1.4](#), note first that [Corollary 7.7](#) implies that there always exists some hypersurface embedding $X \hookrightarrow S^*M$ (which can be assumed real analytic by [Corollary 7.5](#)). The remaining work thus concerns only the sheaf side: we must relate the sheaf category in (7.14) (which is, in particular, not a priori independent of the choice of Liouville hypersurface embedding) to the microsheaf category defined in [63], [81], [65]. This is accomplished in [65], which we adapt to our purposes in the next subsection.

7.5. Doubling II: Microlocal sheaves and antimicrolocalization. Let us now recall the definition of the microlocal sheaf categories appearing in [Theorem 1.4](#). The category which sheaf theorists typically associate to a closed subset of S^*M is defined as follows. One forms the ‘‘Kashiwara–Schapira stack’’ by sheafifying the presheaf of categories on T^*M given by the formula

$\mu\text{sh}^{\text{pre}}(\Omega) := \text{Sh}(M)/\text{Sh}_{T^*M \setminus \Omega}(M)$. The presheaf $\mu\text{sh}^{\text{pre}}$ is already discussed in [49]; working with its sheafification is a more modern phenomenon; see, e.g., [42], [63], [65]. The notion of microsupport makes sense for a section of this sheaf, and we write μsh_Λ for the subsheaf of full subcategories of objects with microsupport inside Λ . The subsheaf $\mu\text{sh}_\Lambda \subseteq \mu\text{sh}$ is evidently supported on Λ .

The sheaf μsh is conic; in particular, given $(T^*M \setminus M) \xrightarrow{\pi} S^*M \xrightarrow{\iota} T^*M$ we have canonically $\pi^* \iota^* \mu\text{sh} = \mu\text{sh}|_{T^*M \setminus M}$. We also denote the sheaf $\iota^* \mu\text{sh}$ on S^*M by μsh . Likewise for $\Lambda \subseteq S^*M$ we have μsh_Λ . We will consider this sheaf for Λ locally closed and will be interested in the category $\mu\text{sh}_\Lambda(\Lambda)$.

By construction there are evident maps $\text{Sh}(M) \rightarrow \mu\text{sh}(\Omega)$ for any open $\Omega \subseteq T^*M$, and similarly $\text{Sh}_\Lambda(M) \rightarrow \mu\text{sh}_\Lambda(\Lambda \cap \Omega)$; in particular, $\text{Sh}_\Lambda(M) \rightarrow \mu\text{sh}_{\Lambda \setminus M}(\Lambda \setminus M) = \mu\text{sh}_{\Lambda_\infty}(\Lambda_\infty)$ (the last of which being in the cosphere bundle). We term all such maps “microlocalization functors.”

In fact, the category $\mu\text{sh}_\Lambda(\Lambda)$ is defined for any space Λ equipped with a germ of closed embedding into a contact manifold carrying a stable polarization [81]. Indeed, such a contact manifold admits a homotopically unique isocontact embedding into $S^*\mathbb{R}^N$ as $N \rightarrow \infty$ by the h -principle [33, 16.1.2]. The key insight of [81] is that, while the image of Λ under such an embedding would have vanishing microsheaf category, one can obtain the correct category by thickening Λ along the relevant Lagrangian polarization of the normal bundle. The role of these polarizations on the sheaf side is entirely parallel to the role of polarizations on the Fukaya side to determine grading/orientation data as discussed in [Section 5.3](#); also compare with [Corollary 7.7](#) and [Remark 7.8](#). Note that we may also define $\mu\text{sh}_\Lambda(\Lambda)$ by embedding into S^*M for any manifold M , since such an M admits an embedding into \mathbb{R}^N .

Remark 7.13. For Liouville manifolds X and X' satisfying the hypotheses of [Theorem 1.4](#), it follows from [Theorem 1.4](#) that if X and X' are within the same Liouville deformation class, then $\mu\text{sh}_{\mathfrak{c}_X}(\mathfrak{c}_X) = \mu\text{sh}_{\mathfrak{c}_{X'}}(\mathfrak{c}_{X'})$. This equivalence is highly non-obvious from the sheaf theoretic standpoint, but is proven directly in [65] under certain assumptions of isotropicity.

The doubling trick in the sheaf context is developed in [65], resulting in embeddings between categories of (microlocal) sheaves parallel to the embeddings between Fukaya categories discussed above. Let us now recall the precise definition of the doubling operation which is relevant in the sheaf context [65], so as to compare it with [Construction 7.10](#) from the Fukaya context.

We begin with a discussion of the contact manifold

$$(7.15) \quad (\mathbb{C} \times V, \lambda_{\mathbb{C}} + \alpha_V),$$

where $\lambda_{\mathbb{C}}$ is a Liouville form on \mathbb{C} (for the standard symplectic structure) and (V, α_V) is a contact manifold with choice of contact form. First, note that the

specific choice of Liouville form on \mathbb{C} is of no importance, as for $f : \mathbb{C} \rightarrow \mathbb{R}$ there is a strict contactomorphism

$$(7.16) \quad (\mathbb{C} \times V, \lambda_{\mathbb{C}} + \alpha_V) \rightarrow (\mathbb{C} \times V, \lambda_{\mathbb{C}} + df + \alpha_V),$$

$$(7.17) \quad (x, y) \mapsto (x, e^{-f(x)R_{\alpha_V}} y),$$

(at least, provided the Reeb flow on V is complete). Next, for a subset $\Lambda_0 \subseteq V$ and a smooth arc γ in \mathbb{C} , define

$$(7.18) \quad \gamma \tilde{\times} \Lambda_0 := \bigcup_{t \in \gamma} (\{t\} \times e^{-g(t)R_{\alpha_V}} \Lambda_0),$$

where $g : \gamma \rightarrow \mathbb{R}$ is a primitive for $\lambda_{\mathbb{C}}|_{\gamma}$, namely $dg = \lambda_{\mathbb{C}}|_{\gamma}$ (so g is well-defined up to adding a locally constant function). Note that the meaning of $\gamma \tilde{\times} \Lambda_0$ does not depend on the choice of Liouville form on \mathbb{C} , as the definition is compatible with the contactomorphisms (7.16). If Λ_0 is isotropic, then so is $\gamma \tilde{\times} \Lambda_0$.

Construction 7.14 (Doubling a subset with boundary coordinates). Begin with a contact manifold Y and a locally closed relatively compact $\Lambda \subseteq Y$. Also fix, in a neighborhood of $\overline{\Lambda} \setminus \Lambda$, coordinates on Y of the form (7.15) (regarded as a germ near $\{0\} \times \Lambda_0$ for compact $\Lambda_0 \subseteq V$) in which $\Lambda = \mathbb{R}_{>0} \tilde{\times} \Lambda_0$ (so Λ_0 is identified with $\overline{\Lambda} \setminus \Lambda$); we call these *boundary coordinates* for Λ . Now the double $D(\Lambda)$ is, near Λ_0 , defined to be $\gamma \tilde{\times} \Lambda_0$, where γ is the immersed arc obtained from two copies of $\mathbb{R}_{>0}$ by adding a small loop enclosing a sufficiently small positive area near the origin (a contractible choice). Away from $\{0\} \times V$, the double is thus $\Lambda \sqcup e^{\epsilon R_{\lambda_{\mathbb{C}} + \alpha_V}} \Lambda$ (note that $R_{\lambda_{\mathbb{C}} + \alpha_V} = R_{\alpha_V}$), which is extended globally by extending the contact form $\lambda_{\mathbb{C}} + \alpha_V$ globally (a contractible choice). The double $D(\Lambda)$ thus consists of Λ (the “first copy”) and a positive Reeb pushoff of Λ (the “second copy”) joined appropriately near their boundary.

Note that when Λ is subanalytic with C^r subanalytic boundary coordinates, then we may ensure that the double $D(\Lambda)$ is subanalytic by choosing both γ and the global extension of α to be C^r subanalytic.¹²¹³ We will tacitly assume that $D(\Lambda)$ is defined in this way whenever Λ is assumed to be subanalytic with C^r subanalytic boundary coordinates.

¹²A C^r subanalytic function $\mathbb{R}^n \rightarrow \mathbb{R}$ is one which is C^r and has subanalytic graph. This class of functions is closed under composition, hence gives rise to a notion of C^r subanalytic manifolds, etc.

¹³The integer r is tacitly assumed to be sufficiently large, and we make no attempt to determine the minimum value of r needed for our constructions to go through (though it will not be particularly large). Note that the tangent bundle of a C^r subanalytic manifold is a C^{r-1} subanalytic vector bundle, hence the highest regularity one can impose on a contact form α is C^{r-1} subanalytic. The exterior derivative $d\alpha$, hence also the Reeb vector field R_{α} , will then be C^{r-2} subanalytic. This would suggest that at a very minimum we must take $r \geq 3$ to ensure we can integrate R_{α} .

The relative core of a Liouville hypersurface always has boundary coordinates (of the same regularity as the hypersurface) in the sense of [Construction 7.14](#). Indeed, a neighborhood of the boundary of a Liouville hypersurface is given by $(Y \times \mathbb{R}_s \times \mathbb{R}_t, e^s \lambda + dt)$ (a germ near $(s, t) = (0, 0)$, with the hypersurface itself being the locus $Y \times \{s \leq 0\} \times \{t = 0\}$), and we can scale the contact form to be $\lambda + e^{-s} dt$, which has the desired form [\(7.15\)](#) near $(s, t) = (0, 0)$. In these coordinates, any relative core will have the desired form $\Lambda_0 \times \{s \leq 0\} \times \{t = 0\}$.

PROPOSITION 7.15. *For the relative core of a Liouville hypersurface, the doubles defined in [Constructions 7.10](#) and [7.14](#) are canonically isotopic.*

Proof. Fix a Liouville manifold X with a stop $\Lambda \subseteq \partial_\infty X$, and consider the stopped Liouville manifold

$$(7.19) \quad (X, \Lambda) \times (\mathbb{C}, \pm i\infty) = (X \times \mathbb{C}, (\Lambda \times i\mathbb{R}) \cup (\mathfrak{c}_X \times \{\pm i\infty\})).$$

The stop, as a subset of $\partial_\infty(X \times \mathbb{C}_{\text{Re} \geq 0})$, is the double of $\mathfrak{c}_{X, \Lambda}$ defined in [Construction 7.10](#). We will exhibit an isotopy from it to the double defined in [Construction 7.14](#).

We begin with the family of stops

$$(7.20) \quad (\Lambda \times \{e^{i\theta}, e^{-i\theta}\} \mathbb{R}_{\geq 0}) \cup (\mathfrak{c}_X \times \{e^{i\theta}, e^{-i\theta}\} \infty)$$

for θ from $\pi/2$ to 0. This family may be written as

$$(7.21) \quad (\gamma_\theta \tilde{\times} \Lambda) \cup (\partial_\infty \gamma_\theta \times \mathfrak{c}_X),$$

where $\gamma_\theta = \{e^{i\theta}, e^{-i\theta}\} \mathbb{R}_{\geq 0}$ (illustrated in the top row of [Figure 4](#)) and we have fixed a contact form on $V = \partial_\infty X$ to obtain coordinates $(\mathbb{C}, \lambda_{\mathbb{C}}) \times (V, \alpha_V) \subseteq \partial_\infty(X \times \mathbb{C})$.




Figure 4. The family of arcs γ_θ (top) and their smoothings $\tilde{\gamma}_\theta$ near the origin (bottom).

Now we smooth γ_θ near the origin to obtain a family of immersed arcs $\tilde{\gamma}_\theta$ (embedded except at $\theta = 0$) as in the bottom row of [Figure 4](#). We would like to consider (7.21) with $\tilde{\gamma}_\theta$ in place of γ_θ . Note, however, that while γ_θ and $\tilde{\gamma}_\theta$ agree outside a compact subset of \mathbb{C} , the same is not true of $\gamma_\theta \tilde{\times} \Lambda$ and $\tilde{\gamma}_\theta \tilde{\times} \Lambda$, due to the fact that the actions of γ_θ and $\tilde{\gamma}_\theta$ necessarily differ at $\theta = 0$ (this being the difference of areas enclosed). Thus while $\gamma_\theta \tilde{\times} \Lambda$ has two “arms” which coincide at $\theta = 0$, the twisted product $\tilde{\gamma}_\theta \tilde{\times} \Lambda$ has two “arms” which near infinity differ by a small positive Reeb pushoff at $\theta = 0$. This small positive isotopy extends to the ambient contact manifold $\partial_\infty(X \times \mathbb{C})$, hence we can, in particular, apply it to the part of (7.21) lying near infinity in the \mathbb{C} -coordinate. This defines the desired isotopy from the double in the sense of [Construction 7.10](#) (at $\theta = \pi/2$) to the double in the sense of [Construction 7.14](#) (at $\theta = 0$). \square

The following “stabilize and then double” construction will be crucial in what follows. Let $\Lambda \subseteq S^*M$ be equipped with boundary coordinates. We consider its “stabilization” $\Lambda \times (0, 1)$ inside $S^*(M \times \mathbb{R})$, where $(0, 1) \subseteq \mathbb{R} \subseteq T^*\mathbb{R}$ is contained in the zero section. This stabilization is naturally equipped with two boundary charts, one near $\partial\Lambda \times (0, 1)$ (obtained from the boundary coordinates for Λ by multiplying by $T^*(0, 1)$) and one near $\Lambda \times \partial(0, 1)$ (obtained from the trivial chart $(0, 1) \subseteq T^*(0, 1)$ by multiplying by $\Lambda \subseteq S^*M$). When M is real analytic and the boundary coordinates for Λ are C^r subanalytic, the first chart is also C^r subanalytic; the second chart is always analytic. These two boundary charts overlap near $\partial\Lambda \times \partial(0, 1)$ in a chart of the form $(\mathbb{C}^2, \mathbb{R}_{\geq 0}^2) \times W$. Viewing the first factor as $(T^*\mathbb{R}^2, \mathbb{R}_{\geq 0}^2)$, we may simply smooth the corner of $\mathbb{R}_{\geq 0}^2$ as in [\[38, §7.1\]](#) to obtain the smoothed product $(\Lambda \times (0, 1))^{\text{sm}}$. This splices together the charts near $\partial\Lambda \times (0, 1)$ and $\Lambda \times \partial(0, 1)$ to define boundary coordinates for $(\Lambda \times (0, 1))^{\text{sm}}$. This smoothing and splicing can be done in the C^r subanalytic category. The double $D((\Lambda \times (0, 1))^{\text{sm}})$ is thus defined, and we abuse notation by writing it as $D(\Lambda \times (0, 1))$.

The doubling trick for sheaf categories from [\[65\]](#) concerns $D(\Lambda \times (0, 1))$, and its proof relies on just a short list of its properties, which are easier to see from a different description of it, as a “movie of creation and destruction” of Λ , denoted $(\Lambda, \partial\Lambda) \prec \succ$ in [\[65, §7.4\]](#). Thus to apply the results of [\[65\]](#) we must give an isotopy $D(\Lambda \times (0, 1)) \sim (\Lambda, \partial\Lambda) \prec \succ$. Let us do this now:

LEMMA 7.16. *For $\Lambda \subseteq S^*M$ equipped with boundary coordinates, the double of the stabilization $D(\Lambda \times (0, 1)) \subseteq S^*(M \times \mathbb{R})$ is isotopic to the “movie of creation and destruction” $(\Lambda, \partial\Lambda) \prec \succ \subseteq S^*(M \times \mathbb{R})$ from [\[65, §7.4\]](#).*

Proof. The main point is to take the picture from [Construction 7.14](#) based on the Lagrangian projection and translate it into the front projection.

First we translate [Construction 7.14](#) itself into the front projection. We add an imaginary third coordinate \mathbb{R}_z to \mathbb{C} to form $(\mathbb{C} \times \mathbb{R}_z, \lambda_{\mathbb{C}} + dz)$. A

Legendrian curve γ in $(\mathbb{C} \times \mathbb{R}_z, \lambda_{\mathbb{C}} + dz)$ determines $\tilde{\gamma} \times \Lambda_0 \subseteq (\mathbb{C} \times V, \lambda_{\mathbb{C}} + \alpha_V)$ for $\Lambda_0 \subseteq V$ via (7.18). The “front projection” is the projection $(\mathbb{C}_{x+iy} \times \mathbb{R}_z, dz - y dx) \rightarrow \mathbb{R}_x \times \mathbb{R}_z$. The Legendrian curve relevant for [Construction 7.14](#) has front projection given by the two rays $\mathbb{R}_{x \geq 0} \times \{z = 0, \epsilon\}$ joined by a single cusp in the standard way illustrated in [Figure 5](#).



Figure 5. A standard cusp.

To understand the double of the stabilization $D(\Lambda \times (0, 1))$, we may look in the corner boundary coordinates $T^* \mathbb{R}_{\geq 0}^2 \times W$. Passing to the front projection and smoothing the corner, we see that in these coordinates, the double is given by two parallel copies of $\mathbb{R}_{\geq 0}^2$ (with its corner smoothed) joined by cusps (i.e., the standard cusp in [Figure 5](#) times a smoothing of the boundary of $\mathbb{R}_{\geq 0}^2$). Up to fixing a standard model of this cusped object (which some readers might call a “square-ish quarter of a flying saucer”), this is exactly the definition of $(\Lambda, \partial\Lambda)^{\prec\succ} \subseteq S^*(M \times \mathbb{R})$ from [\[65, §7.4\]](#). \square

We now state the doubling trick for sheaf categories from [\[65\]](#) (substituting $D(\Lambda \times (0, 1))$ in place of $(\Lambda, \partial\Lambda)^{\prec\succ}$ in accordance with [Lemma 7.16](#) and the preceding discussion). The crucial point for us is that it realizes a given category of *microlocal* sheaves as (a full subcategory of) a certain category of *sheaves* with a given singular support condition; this process is termed “antimicrolocalization” in [\[65\]](#).

THEOREM 7.17 ([\[65, Thm. 7.30\]](#)). *Let Λ be a Whitney stratifiable isotropic inside $S^* M$ with C^r boundary coordinates. The category $\text{Sh}_{D(\Lambda \times (0, 1))}(M \times \mathbb{R})$ is the orthogonal direct sum of its full subcategories $\text{Sh}_{\emptyset}(M \times \mathbb{R})$ (local systems) and $\text{Sh}_{D(\Lambda \times (0, 1))}(M \times \mathbb{R})_0$ (objects with vanishing stalk at infinity), and the microlocalization functor*

$$(7.22) \quad \text{Sh}_{D(\Lambda \times (0, 1))}(M \times \mathbb{R})_0 \xrightarrow{\sim} \text{ush}_{\Lambda \times (0, 1)}(\Lambda \times (0, 1)) = \text{ush}_{\Lambda}(\Lambda)$$

is an equivalence.

(The Künneth equivalence $\text{ush}_{\Lambda \times (0, 1)}(\Lambda \times (0, 1)) = \text{ush}_{\Lambda}(\Lambda)$ is standard.)

Remark 7.18. The actual hypothesis of [Theorem 7.17](#) in [\[65\]](#) (“sufficiently isotropic”) is somewhat weaker than being Whitney stratifiable isotropic. In our applications, we will in fact always have subanalyticity of Λ hence, in particular, Whitney stratifiability.

COROLLARY 7.19. *Let Λ be a locally closed relatively compact subanalytic isotropic inside S^*M with C^r subanalytic boundary coordinates. The category $\mu\text{sh}_\Lambda(\Lambda)$ is compactly generated by co-representatives of microstalks at the smooth points of Λ .*

Proof. By [Theorem 7.17](#), we have $\mu\text{sh}_\Lambda(\Lambda) = \text{Sh}_{D(\Lambda \times (0,1))}(M \times \mathbb{R})_0$. It thus suffices to show that the microstalk co-representatives $\mu_\xi \in \text{Sh}_{D(\Lambda \times (0,1))}(M \times \mathbb{R})$ for smooth points ξ of the first copy of $\Lambda \times (0, 1)$ inside $D(\Lambda \times (0, 1))$ lie in the full subcategory $\text{Sh}_{D(\Lambda \times (0,1))}(M \times \mathbb{R})_0$ and compactly generate it.

The category $\text{Sh}_{D(\Lambda \times (0,1))}(M \times \mathbb{R})$ is compactly generated by [Corollary 4.22](#) (which immediately implies its orthogonal full subcategories from [Theorem 7.17](#) are also compactly generated). The μ_ξ (which are compact) are by definition left orthogonal to local systems, so they lie in $\text{Sh}_{D(\Lambda \times (0,1))}(M \times \mathbb{R})_0$ by [Theorem 7.17](#). An object of $\text{Sh}_{D(\Lambda \times (0,1))}(M \times \mathbb{R})$ right-orthogonal to all these μ_ξ must have microsupport contained in $\overline{\Lambda} \times [0, 1]$ (the closure of the second copy), but this implies empty microsupport since $\overline{\Lambda} \times [0, 1]$ is isotropic and every smooth Legendrian component has boundary (so the relevant microstalk always vanishes, which is enough by [Proposition 4.10](#)). It follows that the μ_ξ compactly generate $\text{Sh}_{D(\Lambda \times (0,1))}(M \times \mathbb{R})_0$ as claimed. \square

Remark 7.20. One may eliminate the subanalyticity hypotheses from [Corollary 7.19](#) at the cost of appealing to more general representability theorems as in [Lemma 4.13](#). For example, for Λ stratifiable by isotropics, $\mu\text{sh}_\Lambda(\Lambda)$ is compactly generated by co-representatives of microstalks at the smooth points of Λ . Indeed, the arguments of [Lemma 4.13](#) imply that for any open $U \subseteq \Lambda$, the restriction $\mu\text{sh}_\Lambda(\Lambda) \rightarrow \mu\text{sh}_U(U)$ has a left adjoint which preserves compact objects. Taking U to be a contractible open subset near a given smooth Legendrian point of Λ produces a compact co-representative of the microstalk. These then compactly generate by [Proposition 4.10](#).

7.6. Proof of Theorem 1.4. We begin by deriving from [Theorem 7.17](#) a sheaf theoretic analogue of [Proposition 7.11](#). Although it is a purely sheaf theoretic statement, the proof we give passes through the Fukaya category and the results of [38].

COROLLARY 7.21. *Let (X, Λ) be a stopped Liouville manifold. Let M be a real analytic manifold and $X \hookrightarrow S^*M$ a Liouville hypersurface embedding such that the image of $\mathfrak{c}_{X, \Lambda}$ is subanalytic isotropic with C^r subanalytic boundary coordinates. The left adjoint μ^* of the microlocalization functor $\mu : \text{Sh}_{D(\mathfrak{c}_{X, \Lambda})}(M) \rightarrow \mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda})$ is fully faithful.*

We will see in the proof that μ^* exists and preserves compact objects for formal reasons.

Proof. We consider the commuting diagram

$$(7.23) \quad \begin{array}{ccccc} \mathrm{Sh}_{D(\mathfrak{c}_{X,\Lambda})}(M) & \xleftarrow{r} & \mathrm{Sh}_{D(\mathfrak{c}_{X,\Lambda}) \times (0,1)}(M \times (0,1)) & \xleftarrow{r} & \mathrm{Sh}_{D(\mathfrak{c}_{X,\Lambda} \times (0,1))}(M \times \mathbb{R}) \\ \downarrow \mu & & \downarrow \mu & & \swarrow \mu \\ \mu\mathrm{sh}_{\mathfrak{c}_{X,\Lambda}}(\mathfrak{c}_{X,\Lambda}) & \xleftarrow{r} & \mu\mathrm{sh}_{\mathfrak{c}_{X,\Lambda} \times (0,1)}(\mathfrak{c}_{X,\Lambda} \times (0,1)) & & \end{array}$$

where the functors are restriction r and microlocalization μ . It is a standard result of microlocal sheaf theory that the two leftmost restriction functors r are equivalences. We note that we may indeed choose the double of the stabilization $D(\mathfrak{c}_{X,\Lambda} \times (0,1))$ so that over $M \times (0,1)$ it coincides with $D(\mathfrak{c}_{X,\Lambda}) \times (0,1)$ (compare the picture from [Lemma 7.16](#)), so the upper right restriction functor r is defined.

Because microsupport respects limits and colimits, so too do the microlocalization functors μ (see, e.g., [\[65, Rem. 6.1\]](#) for more details) and restriction functors r . The domain sheaf categories are compactly generated by [Corollary 4.22](#), and Brown representability holds for the opposites of compactly generated categories by [\[69\]](#), [\[53\]](#), so all functors r and μ in (7.23) admit left adjoints r^* and μ^* .¹⁴ Being left adjoint to co-continuous functors, each r^* and μ^* preserves compact objects.

The microlocal sheaf categories appearing in (7.23) are compactly generated by co-representatives of microstalk functors by [Corollary 7.19](#). The images of these compact generators under μ^* are again co-representatives of the same microstalk functors. To show that a given μ^* is fully faithful, it suffices to check on compact objects.

By [Theorem 7.17](#), the diagonal μ in (7.23) is the projection onto an orthogonal direct summand of the domain. Its left adjoint μ^* is thus the inclusion of this orthogonal direct summand hence, in particular, is fully faithful. Thus to prove full faithfulness of the other vertical μ^* functors, it suffices to show full faithfulness of

$$(7.24) \quad r^* : \mathrm{Sh}_{D(\mathfrak{c}_{X,\Lambda}) \times (0,1)}(M \times (0,1)) \rightarrow \mathrm{Sh}_{D(\mathfrak{c}_{X,\Lambda} \times (0,1))}(M \times \mathbb{R})$$

restricted to co-representatives of the microstalk functors at the first copy of $\mathfrak{c}_{X,\Lambda} \times (0,1)$. By [Proposition 1.3](#), this functor (restricted to compact objects) corresponds under [Theorem 1.1](#) to the pushforward functor

$$(7.25) \quad \mathcal{W}(T^*(M \times (0,1)), D(\mathfrak{c}_{X,\Lambda}) \times (0,1)) \rightarrow \mathcal{W}(T^*(M \times \mathbb{R}), D(\mathfrak{c}_{X,\Lambda} \times (0,1))).$$

It thus suffices to show that the restriction of this functor to the linking disks of the first copy of $\mathfrak{c}_{X,\Lambda} \times (0,1)$ is fully faithful.

¹⁴In fact, r^* and μ^* exist in general by arguing as in [Lemma 4.13](#).

The inclusion

$$(7.26) \quad (X, \Lambda) \times (\mathbb{C}, \pm\infty) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty) \hookrightarrow (T^*(M \times \mathbb{R}), D(\mathfrak{c}_{X, \Lambda} \times (0, 1)))$$

around the first copy of $\mathfrak{c}_{X, \Lambda} \times (0, 1) = \mathfrak{c}_{(X, \Lambda) \times (\mathbb{C}, \pm\infty)}$ induces a fully faithful functor on \mathcal{W} by [Proposition 7.11](#) and the fact that $(X, \Lambda) \times (\mathbb{C}, \pm\infty)$ admits homological cocores. Now, after a deformation, the above inclusion factors through $(T^*(M \times (0, 1)), D(\mathfrak{c}_{X, \Lambda}) \times (0, 1))$ as the identity map on $T^*(0, 1)$ times the canonical inclusion

$$(7.27) \quad (X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty) \hookrightarrow (T^*M, D(\mathfrak{c}_{X, \Lambda}))$$

around the first copy of $\mathfrak{c}_{X, \Lambda}$. It thus suffices to show that the induced map on wrapped Fukaya categories is also fully faithful. To do this, we multiply the proof of [Proposition 7.11](#) by $T^*(0, 1)$. Namely, we consider Lagrangians inside $T^*(0, 1) \times (X, \Lambda)$ times the linking disk of $(\mathbb{C}_{\text{Re} \geq 0}, \infty)$ and consider product wrappings inside $T^*(0, 1) \times (X, \Lambda) \times (\mathbb{C}, \{\pm i\infty\})$ which we conclude satisfy the cofinality criterion, hence remain cofinal after removing a neighborhood of the additional stop at $\mathfrak{c}_X \times \{-\infty\}$ and gluing onto T^*M . \square

COROLLARY 7.22. *Let (X, Λ) be a stopped Liouville manifold. Let M be a real analytic manifold and $X \hookrightarrow S^*M$ a Liouville hypersurface embedding such that the image of $\mathfrak{c}_{X, \Lambda}$ is subanalytic isotropic with C^r subanalytic boundary coordinates. There is a fully faithful functor $\mathcal{W}(X, \Lambda)^{\text{op}} \hookrightarrow \mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda})^c$ characterized uniquely by commutativity of the diagram*

$$(7.28) \quad \begin{array}{ccc} \mathcal{W}(X, \Lambda)^{\text{op}} & \xhookrightarrow{\text{(7.12)}} & \text{Perf } \mathcal{W}(T^*M, D(\mathfrak{c}_{X, \Lambda}))^{\text{op}} \\ \downarrow & & \parallel \text{Theorem 1.1} \\ \mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda})^c & \xhookrightarrow{\mu^*} & \text{Sh}_{D(\mathfrak{c}_{X, \Lambda})}(M)^c, \end{array}$$

where μ^* denotes the restriction to compact objects of the left adjoint of the microlocalization functor.

Proof. [Proposition 7.11](#) and [Corollary 7.21](#) ensure that the horizontal arrows in (7.28) are fully faithful. The essential image of (7.12) is contained in the subcategory generated by linking disks of the first copy of $\mathfrak{c}_{X, \Lambda}$ since it factors through $\mathcal{W}((X, \Lambda) \times (\mathbb{C}_{\text{Re} \geq 0}, \infty))$, which is generated by linking disks. The functor μ^* obviously sends co-representatives of microstalks (which exist by [Corollary 7.19](#)) to co-representatives of microstalks, which are identified with linking disks under [Theorem 1.1](#). \square

Remark 7.23. The hypotheses of [Corollaries 7.21](#) and [7.22](#) may be ensured by assuming (X, Λ) is analytic, $\mathfrak{c}_{X, \Lambda}$ is subanalytic isotropic, and the embedding $X \hookrightarrow S^*M$ is analytic. This is how we proceed to prove [Theorem 1.4](#), relying on the abstract analytic approximation result of [Corollary 7.5](#). However when applying [Corollary 7.5](#) in practice (and in particular in [36]), it can be more

convenient to simply check subanalyticity of the core inside S^*M and existence of analytic boundary coordinates (which holds vacuously if $\Lambda = \emptyset$).

Now the embedding of [Theorem 1.4](#) is simply defined to be that of [Corollary 7.22](#) for a choice of Liouville hypersurface embedding, which is guaranteed to exist by [Corollary 7.7](#).

Proof of Theorem 1.4. By [Corollary 7.7](#), there is a Liouville hypersurface embedding $X \times \mathbb{C}^k \hookrightarrow S^*M$ compatible with polarizations for some manifold M . Equip M with a real analytic structure, and use [Corollary 7.5](#) to perturb the embedding to be analytic. Now apply [Corollary 7.22](#) to $(X, \Lambda) \times (\mathbb{C}, \pm\infty)^k$ and the embedding $X \times \mathbb{C}^k \hookrightarrow S^*M$ to obtain an embedding

$$(7.29) \quad \mathcal{W}((X, \Lambda) \times (\mathbb{C}, \pm\infty)^k) \hookrightarrow \mu\text{sh}_{\mathfrak{c}_{X, \Lambda} \times \mathbb{R}^k}(\mathfrak{c}_{X, \Lambda} \times \mathbb{R}^k),$$

which sends homological cocores to co-representatives of microstalks since [Theorem 1.1](#) sends linking disks to co-representatives of microstalks. Finally, combine this with the Künneth embedding $\mathcal{W}(X, \Lambda) \hookrightarrow \mathcal{W}((X, \Lambda) \times (\mathbb{C}, \pm\infty)^k)$ and the equivalence $\mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda}) = \mu\text{sh}_{\mathfrak{c}_{X, \Lambda} \times \mathbb{R}^k}(\mathfrak{c}_{X, \Lambda} \times \mathbb{R}^k)$. \square

While the equivalence of [Theorem 1.1](#) is canonical, the embedding of [Theorem 1.4](#) depends a priori on a choice of analytic hypersurface embedding $X \times \mathbb{C}^k \hookrightarrow S^*M$ compatible with polarizations. We do strongly expect that it is independent of these choices, and moreover that pursuing the present methods a bit further would show this.

In some instances, there is a particularly natural choice of Liouville hypersurface embedding for which the category $\text{Sh}_{\overline{\mathfrak{c}_{X, \Lambda}}}(M)$ is of interest. It is then of interest to know that the embedding of [Theorem 1.4](#) (associated to this particular hypersurface embedding) and the equivalence of [Theorem 1.1](#) intertwine pushforward on Fukaya categories and (the left adjoint of) microlocalization. We stated this compatibility in the introduction as (1.4). Here we make a stronger statement, relevant in applications, with [Corollary 7.22](#) in place of [Theorem 1.4](#). [Corollary 7.22](#) requires only that the image of the core be subanalytic and have subanalytic boundary coordinates, rather than requiring the hypersurface itself to be analytic.

PROPOSITION 7.24. *In the notation and assuming the hypotheses of [Corollary 7.22](#), the following diagram commutes:*

$$(7.30) \quad \begin{array}{ccc} \mathcal{W}(X, \Lambda)^{\text{op}} & \longrightarrow & \text{Perf } \mathcal{W}(T^*M, \overline{\mathfrak{c}_{X, \Lambda}})^{\text{op}} \\ \text{Corollary 7.22} \downarrow & & \parallel \text{Theorem 1.1} \\ \mu\text{sh}_{\mathfrak{c}_{X, \Lambda}}(\mathfrak{c}_{X, \Lambda})^c & \xrightarrow{\mu^*} & \text{Sh}_{\overline{\mathfrak{c}_{X, \Lambda}}}(M)^c. \end{array}$$

Proof. Append to the right side of (7.28) a square diagram forgetting down from $D(\mathfrak{c}_{X, \Lambda})$ to the first copy $\overline{\mathfrak{c}_{X, \Lambda}} \subseteq D(\mathfrak{c}_{X, \Lambda})$. \square

Example 7.25 (Mirror symmetry for very affine hypersurfaces). Let $W_{\mathbf{T}} : (\mathbb{C}^*)^n \rightarrow \mathbb{C}^*$ be the Hori–Vafa mirror superpotential to a smooth toric stack \mathbf{T} . In [Corollary 6.16](#), we discussed how the results of the present article allows one to translate the sheaf theoretic work of [\[35\]](#), [\[54\]](#) into a mirror symmetry statement equating the Fukaya–Seidel category of $W_{\mathbf{T}}$ with $\text{Coh}(\mathbf{T})$. This also depended on certain calculations of skeleta in [\[36\]](#), [\[91\]](#).

The main purpose of [\[36\]](#) was to provide the relevant skeletal calculations and microlocal sheaf theoretic results to prove the expected mirror symmetry between the wrapped Fukaya category of a generic fiber (which we denote $W_{\mathbf{T}}^{-1}(-\infty)$) and the category of coherent sheaves on the toric boundary $\text{Coh}(\partial\mathbf{T})$. [Theorem 7.22](#) provides the translation between microlocal sheaf theory and wrapped Fukaya categories. To summarize, we have the following commutative diagram:

$$(7.31) \quad \begin{array}{ccc} \text{Coh}(\partial\mathbf{T}) & \longrightarrow & \text{Coh } \mathbf{T} \\ \text{[36]} \parallel & & \parallel \text{[35], [54]} \\ \mu\text{sh}_{\Lambda_{\mathbf{T}}}(\Lambda_{\mathbf{T}})^c & \xrightarrow{\mu^*} & \text{Sh}_{\Lambda_{\mathbf{T}}}((S^1)^n)^c \\ \text{Theorem 7.22} \parallel & & \parallel \text{Theorem 1.1} \\ \text{Perf } \mathcal{W}(W_{\mathbf{T}}^{-1}(-\infty)) & \longrightarrow & \text{Perf } \mathcal{W}((\mathbb{C}^*)^n, W_{\mathbf{T}}^{-1}(-\infty)) \end{array}$$

in which the bottom square is [\(7.30\)](#), using the fact that $-\Lambda_{\mathbf{T}}$ is the core of $W_{\mathbf{T}}^{-1}(-\infty)$ from [\[36\]](#), [\[91\]](#). (The absence of an “op” is due to the appearance of the minus sign in $-\Lambda_{\mathbf{T}}$ and a corresponding use of an antipodal map.)

7.7. Making the core subanalytic. The goal of this subsection is to show that every Weinstein sector may be perturbed to be real analytic and have subanalytic relative core (and hence satisfy the hypotheses of [Theorem 1.4](#)).

PROPOSITION 7.26. *Let M be a real analytic manifold, and let V be a real analytic vector field on M which is convex and complete at infinity and which is gradient-like with respect to a proper Morse function with finitely many critical points. Suppose that a neighborhood of every zero of V has local analytic coordinates in which $V = \sum_i a_i x_i \frac{\partial}{\partial x_i}$ for some $a_i \in \mathbb{Q} \setminus \{0\}$. Then the union of all stable manifolds $C \subseteq M$ is a subanalytic subset of M . In fact, for any subanalytic subset $\Lambda \subseteq \partial_\infty M$, the union $C \cup (\Lambda \times \mathbb{R}) \subseteq M$ is subanalytic.*

Proof. Fix a proper smooth Morse function $\phi : M \rightarrow \mathbb{R}$ with respect to which V is gradient-like. There is no real need to make ϕ real analytic, though the usual real analytic approximation results allow us to do so if we like.

The core C is compact, so it is vacuously true that $C_\Lambda := C \cup (\Lambda \times \mathbb{R})$ is subanalytic over $\{\phi > T\}$ for some large $T < \infty$. By V -invariance of C_Λ , if an interval $[T', T]$ contains no critical values of ϕ , then C_Λ subanalytic over

$\{\phi > T\}$ implies C_Λ is subanalytic over $\{\phi > T'\}$. Thus the point is to understand what happens when we cross a critical value of ϕ . We may assume the critical values of ϕ are distinct.

Fix a critical value of ϕ , which by translating ϕ we may assume is zero. Supposing that C_Λ is subanalytic over $\{\phi > \epsilon\}$, let us show that C_Λ is subanalytic over $\{\phi > -\epsilon\}$. It is trivial that C_Λ is subanalytic away from the stable manifold of the critical point of ϕ in question. Thus let us work in local analytic coordinates $[-1, 1]^{n+m}$ near this critical point in which $V = \sum_i a_i x_i \frac{\partial}{\partial x_i} - \sum_j b_j y_j \frac{\partial}{\partial y_j}$ for $a_i, b_j \in \mathbb{Q}_{>0}$. We now consider the proper map

$$(7.32) \quad \left\{ \begin{array}{l} x_1^2 + \cdots + x_n^2 = 1 \\ y_1^2 + \cdots + y_m^2 = 1 \\ s, t \geq 0 \end{array} \right\} \xrightarrow{(s^{a_1} x_1, \dots, s^{a_n} x_n, t^{b_1} y_1, \dots, t^{b_m} y_m)} \mathbb{R}^{n+m},$$

which is analytically definable since $a_i, b_i \in \mathbb{Q}$. Note that for fixed values of $(x_1, \dots, x_n, y_1, \dots, y_m)$ and of the product st , the image is a flow line of V ; in fact, this identifies the space of broken flow lines of $\sum_i a_i x_i \frac{\partial}{\partial x_i} - \sum_j b_j y_j \frac{\partial}{\partial y_j}$ on \mathbb{R}^{n+m} with

$$(7.33) \quad \{x_1^2 + \cdots + x_n^2 = 1\} \times \{y_1^2 + \cdots + y_m^2 = 1\} \times \mathbb{R}_{\geq 0}.$$

We may now show that C_Λ is subanalytic in a neighborhood of the stable manifold $\{x_1 = \cdots = x_n = 0\} \times \mathbb{R}_y^m$ as follows. Choose a small real analytic hypersurface H transverse to V near

$$\{x_1^2 + \cdots + x_n^2 = 1\} \times \{y_1 = \cdots = y_m = 0\}.$$

Since H lies in the locus where ϕ is positive, the intersection $C_\Lambda \cap H$ is subanalytic. Now the image of $C_\Lambda \cap H$ under the backward flow of V may be described by projecting it to (7.33), taking its inverse image in the domain of (7.32), and taking its image under (7.32); the result is subanalytic since (7.32) is proper. Near the stable manifold, C_Λ is the union of this subanalytic set (the image of $C_\Lambda \cap H$ under the backward flow of V) with the stable manifold, hence is subanalytic. \square

COROLLARY 7.27. *Every Weinstein manifold can be perturbed to admit a real analytic structure such that for every subanalytic subset at infinity, the associated relative core is subanalytic.*

Proof. The standard Weinstein handle

$$(7.34) \quad \left(\mathbb{R}^{2k} \times \mathbb{R}^{2(n-k)}, \sum_{i=1}^n dx_i \wedge dy_i, \sum_{i=1}^k \frac{1}{2}(-x_i \partial_{x_i} + 3y_i \partial_{y_i}) + \sum_{i=k+1}^n \frac{1}{2}(x_i \partial_{x_i} + y_i \partial_{y_i}) \right)$$

is real analytic. Any critical point of a Weinstein manifold may be perturbed so as to coincide locally with (7.34) (see [20] and [40, Lemma 6.6]). We may thus

construct (after perturbation) any Weinstein manifold by iteratively attaching such standard handles. Now the attaching maps may be perturbed to be real analytic by [Lemma 7.4](#). We therefore obtain a real analytic Weinstein manifold (X, ω, Z) to which [Proposition 7.26](#) applies. \square

COROLLARY 7.28. *Every Weinstein sector is equivalent to a stopped Weinstein manifold with subanalytic isotropic relative core.*

Proof. A Weinstein sector is (equivalent to) a Liouville pair (X, F) where X and F are both Weinstein. Apply [Corollary 7.27](#) to X and F individually, and apply [Corollary 7.5](#) to the embedding $F \hookrightarrow \partial_\infty X$. \square

Appendix A. Review of categorical notions

We will assume the reader is familiar with the basic definitions of differential graded (dg) and/or A_∞ categories, functors between them, modules, and bimodules, for which there are many references. In this section we review notation, assumptions, and relevant notions/results.

All of our dg or A_∞ categories \mathcal{C} have morphism cochain complexes linear over a fixed commutative ring (which we take for simplicity of notation to be \mathbb{Z}), which are \mathbb{Z} -graded and cofibrant in the sense of [\[37, §3.1\]](#) (an assumption which is vacuous if working over a field). We further assume that all such \mathcal{C} are at least cohomologically unital, meaning that the underlying cohomology-level category $H^*(\mathcal{C})$ has identity morphisms. (This follows if \mathcal{C} itself is strictly unital, as is the case in the dg setting.) We say objects in \mathcal{C} are isomorphic if they are isomorphic in $H^*(\mathcal{C})$.

A.1. Functors, modules, and bimodules. For two (A_∞ or dg) categories \mathcal{C} and \mathcal{D} , we use the notation

$$(A.1) \quad \text{Fun}(\mathcal{C}, \mathcal{D})$$

to refer to the (A_∞) category of A_∞ functors from \mathcal{C} to \mathcal{D} . (Compare [\[79, §1d\]](#), noting that here we consider homologically unital functors.) Note that $\text{Fun}(\mathcal{C}, \mathcal{D})$ is in fact a dg category whenever \mathcal{D} is. The morphism space between $f, g \in \text{Fun}(\mathcal{C}, \mathcal{D})$ is the *derived* space of natural transformations (as opposed to the space of strict natural transformations, which can be defined in the dg setting but not in the more general A_∞ setting).

An A_∞ functor $f : \mathcal{C} \rightarrow \mathcal{D}$ is called fully faithful (essentially surjective, an equivalence) if the induced functor on cohomology categories $H^*(f) : H^*(\mathcal{C}) \rightarrow H^*(\mathcal{D})$ is. We freely use the similar notion of a bilinear A_∞ functor $\mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ (see [\[59\]](#)), which are themselves objects of an A_∞ category which is dg if \mathcal{E} is.

Denote by $\text{Mod}\mathbb{Z}$ the dg category of dg \mathbb{Z} -modules, i.e., the category of (implicitly \mathbb{Z} -graded) unbounded complexes of \mathbb{Z} -modules localized at acyclic

complexes. When relevant, we take cofibrant complexes of \mathbb{Z} -modules as our model of this category.

A left (respectively right) module over a category \mathcal{C} is, by definition, a functor from \mathcal{C}^{op} (respectively \mathcal{C}) to $\text{Mod } \mathbb{Z}$. More generally, a $(\mathcal{C}, \mathcal{D})$ bimodule is a bilinear functor $\mathcal{C}^{\text{op}} \times \mathcal{D} \rightarrow \text{Mod } \mathbb{Z}$; this notion specializes to the previous two notions by taking \mathcal{C} or $\mathcal{D} = \mathbb{Z}$ (meaning the category with one object $*$ and endomorphism algebra \mathbb{Z}); see [37, §3.1]. By the above discussion, left modules, right modules, and bimodules are each objects of dg categories, denoted

$$(A.2) \quad \text{Mod } \mathcal{C} = \text{Fun}(\mathcal{C}^{\text{op}}, \text{Mod } \mathbb{Z}),$$

$$(A.3) \quad \text{Mod } \mathcal{C}^{\text{op}} = \text{Fun}(\mathcal{C}, \text{Mod } \mathbb{Z}),$$

$$(A.4) \quad [\mathcal{C}, \mathcal{D}] = \text{Fun}(\mathcal{C}^{\text{op}} \times \mathcal{D}, \text{Mod } \mathbb{Z}),$$

respectively. We will most frequently discuss left modules, which we simply call modules. There are canonical fully faithful Yoneda embeddings (see, e.g., [79, §(11)] for a more detailed description on morphism spaces):

$$(A.5) \quad \mathcal{C} \hookrightarrow \text{Mod } \mathcal{C} \quad X \mapsto \text{hom}_{\mathcal{C}}(-, X),$$

$$(A.6) \quad \mathcal{C}^{\text{op}} \hookrightarrow \text{Mod } \mathcal{C}^{\text{op}} \quad Y \mapsto \text{hom}_{\mathcal{C}}(Y, -),$$

$$(A.7) \quad \mathcal{C} \times \mathcal{D}^{\text{op}} \hookrightarrow [\mathcal{C}, \mathcal{D}] \quad (X, Y) \mapsto \text{hom}_{\mathcal{C}}(-, X) \otimes_{\mathbb{Z}} \text{hom}_{\mathcal{D}}(Y, -),$$

and we call any (bi)module in the essential image of these embeddings representable. Recall that any \mathcal{C} possesses a canonical (not necessarily representable) $(\mathcal{C}, \mathcal{C})$ bimodule, the diagonal bimodule \mathcal{C}_{Δ} (defined on the level of objects by $\mathcal{C}_{\Delta}(-, -) = \text{hom}_{\mathcal{C}}(-, -)$).

A $(\mathcal{D}, \mathcal{C})$ bimodule \mathcal{B} induces, via convolution (also known as tensor product), a functor

$$(A.8) \quad \mathcal{B} \otimes_{\mathcal{C}} - : \text{Mod } \mathcal{C} \rightarrow \text{Mod } \mathcal{D},$$

$$(A.9) \quad \mathcal{M} \mapsto \mathcal{B}(-, -) \otimes_{\mathcal{C}} \mathcal{M}(-)$$

(note that this is a version of the derived tensor product), and more generally a functor $[\mathcal{C}, \mathcal{E}] \rightarrow [\mathcal{D}, \mathcal{E}]$ for any category \mathcal{E} . This functor always has a right adjoint, given by $\mathcal{N} \mapsto \text{hom}_{\text{Mod } \mathcal{D}}(\mathcal{B}, \mathcal{N})$.¹⁵ As one might expect, convolving with the diagonal bimodule is (isomorphic to) the identity. Not every functor $\text{Mod } \mathcal{C} \rightarrow \text{Mod } \mathcal{D}$ comes from a bimodule, however there is a characterization of those which do:

THEOREM A.1 (compare [90, Thm. 1.4]). *The convolution map $[\mathcal{D}, \mathcal{C}] \rightarrow \text{Fun}(\text{Mod } \mathcal{C}, \text{Mod } \mathcal{D})$ is fully faithful, and its essential image is precisely the co-continuous functors, i.e., those which preserve small direct sums.*

¹⁵We say $f : \mathcal{C} \rightarrow \mathcal{D}$ has right adjoint (or is the left adjoint of) $g : \mathcal{D} \rightarrow \mathcal{C}$ if there is an isomorphism in $[\mathcal{C}, \mathcal{D}]$ between $\text{hom}_{\mathcal{D}}(f(-), -)$ and $\text{hom}_{\mathcal{C}}(-, g(-))$.

(By “ F preserves small direct sums” we mean “the natural map $\bigoplus_\alpha F(X_\alpha) \rightarrow F(\bigoplus_\alpha X_\alpha)$ is an isomorphism.”)

Proof Sketch. If $\text{Fun}_{\text{co-continuous}}(\text{Mod } \mathcal{C}, \text{Mod } \mathcal{D})$ denotes the co-continuous functors, observe that restriction to (the Yoneda image of) \mathcal{C} induces tautologically a map (which is an equivalence) $\text{Fun}_{\text{co-continuous}}(\text{Mod } \mathcal{C}, \text{Mod } \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \text{Mod } \mathcal{D}) = \text{Fun}(\mathcal{C}, \text{Fun}(\mathcal{D}^{\text{op}}, \text{Mod } \mathbb{Z})) = [\mathcal{D}, \mathcal{C}]$; in other words, co-continuous functors from $\text{Mod } \mathcal{C}$ are determined by what they do on \mathcal{C} . One checks that this is a two-sided inverse to the convolution map, up to homotopy. \square

Given an A_∞ functor $f : \mathcal{C} \rightarrow \mathcal{D}$, there is a pair of (adjoint) induced functors on module categories: first, there is an induced restriction map

$$(A.10) \quad f^* : \text{Mod } \mathcal{D} \rightarrow \text{Mod } \mathcal{C}$$

given by pre-composing with f^{op} ; one can show this is isomorphic to tensoring with the graph $(\mathcal{C}, \mathcal{D})$ bimodule $(f^{\text{op}}, \text{id})^* \mathcal{D}_\Delta = \mathcal{D}_\Delta(f(-), -)$ (see [37, Lem. 3.7]). In particular, there is a natural functor $\mathcal{D} \rightarrow \text{Mod } \mathcal{C}$ given by composing (A.10) with the Yoneda embedding for \mathcal{D} . There is also (left adjoint to f^*) an induction map

$$(A.11) \quad f_! : \text{Mod } \mathcal{C} \rightarrow \text{Mod } \mathcal{D}$$

given by tensoring with the graph $(\mathcal{D}, \mathcal{C})$ bimodule $(\text{id}, f)^* \mathcal{D}_\Delta = \mathcal{D}_\Delta(-, f(-))$. One can directly compute that $f_!$ sends a representable over $X \in \mathcal{C}$ to an object isomorphic to the representable over $f(X)$. Conversely, we have

LEMMA A.2. *If a $(\mathcal{D}, \mathcal{C})$ bimodule \mathcal{B} has the property that $\mathcal{B}(-, c)$ is representable by an object $f(c) \in \mathcal{D}$ for each $c \in \mathcal{C}$, then convolving with \mathcal{B} is isomorphic in $\text{Fun}(\text{Mod } \mathcal{C}, \text{Mod } \mathcal{D})$ to the induction of a (unique up to isomorphism) A_∞ functor $f : \mathcal{C} \rightarrow \mathcal{D}$ sending c to $f(c)$. In particular, $f_! = \mathcal{B} \otimes_{\mathcal{C}} -$ admits a right adjoint, namely f^* .*

Note that f^* also admits a right adjoint f_* , called co-induction, induced by taking hom from $(f^{\text{op}}, \text{id})^* \mathcal{D}_\Delta$, by the earlier discussion.

A.2. Pre-triangulated, idempotent complete, and co-complete categories. A category \mathcal{C} is called pre-triangulated if and only if it is closed under taking mapping cones; in this case $H^0 \mathcal{C}$ is triangulated in the usual sense. Every category has a well-defined pre-triangulated closure $\text{Tw } \mathcal{C}$,¹⁶ which can be defined as the closure of the image of \mathcal{C} in $\text{Mod } \mathcal{C}$ under taking mapping cones (see [79, §3]). An object of $\text{Mod } \mathcal{C}$ which is in the closure of $\mathcal{A} \subseteq \mathcal{C}$ under taking mapping cones is said to be generated by \mathcal{A} .

¹⁶The notation $\text{Tw } \mathcal{C}$ is usually taken to mean the specific model of the pre-triangulated closure of \mathcal{C} given by the category of so-called “twisted complexes” of objects of \mathcal{C} .

A category \mathcal{C} is called idempotent complete if and only if $H^0\mathcal{C}$ is closed under retracts. (Usually this property is only considered when \mathcal{C} is already pre-triangulated.) Every category has a well-defined idempotent completion \mathcal{C}^π , which can be defined similarly as a full subcategory of $\text{Mod } \mathcal{C}$ (see [79, §4]). An object of $\text{Mod } \mathcal{C}$ is said to be split-generated by $\mathcal{A} \subseteq \mathcal{C}$ when it is in the closure of \mathcal{A} under the operations of taking mapping cones and retracts. The category of perfect modules $\text{Perf } \mathcal{C} \subseteq \text{Mod } \mathcal{C}$ is by definition the full subcategory spanned by objects split-generated by \mathcal{C} ; in other words, $\text{Perf } \mathcal{C} = (\text{Tw } \mathcal{C})^\pi$.

A category \mathbf{C} is called co-complete if and only if it is pre-triangulated and has all (small) direct sums. Equivalently, \mathbf{C} is co-complete if and only if it has all small colimits. In particular, a co-complete category is idempotent complete (compare [69, Prop. 1.6.8]). We will also call co-complete categories *large categories*.

A.3. Compactly generated categories. Any category of modules $\text{Mod } \mathcal{C}$ (or more generally bimodules, etc.) over a small category \mathcal{C} inherits from $\text{Mod } \mathbb{Z}$ the property of being co-complete. Large (i.e., co-complete) categories of the form $\text{Mod } \mathcal{C}$ may be characterized intrinsically as follows.

Let \mathbf{C} be a large category. We say an object $X \in \mathbf{C}$ is compact if $\text{hom}_{\mathbf{C}}(X, -)$ commutes with arbitrary direct sums (i.e., is co-continuous). Denoting by $\mathbf{C}^c \subseteq \mathbf{C}$ the full subcategory of compact objects, we say that a co-complete category \mathbf{C} is compactly generated if there is a small collection (i.e., a set) of compact objects $\mathcal{C} \subseteq \mathbf{C}^c$ satisfying the following equivalent conditions:

- An object $X \in \mathbf{C}$ is zero if and only if it is right-orthogonal to \mathcal{C} (meaning $\text{hom}_{\mathbf{C}}(-, X)$ annihilates \mathcal{C}).
- The natural map $\mathbf{C} \rightarrow \text{Mod } \mathcal{C}$ sending $Y \mapsto \text{hom}_{\mathbf{C}}(-, Y)$ is an equivalence.

Thus compactly generated categories \mathbf{C} are precisely those of the form $\text{Mod } \mathcal{C}$ for some small category \mathcal{C} .

It is natural to ask to what extent $\text{Mod } \mathcal{C}$ determines \mathcal{C} . This is answered by the following well-known fact:

LEMMA A.3. *The compact objects of $\text{Mod } \mathcal{C}$ are precisely $\text{Perf } \mathcal{C}$.*

Proof. Let $\mathcal{M} \in \text{Mod } \mathcal{C}$ be compact. There is a natural quasi-isomorphism $\mathcal{C}_\Delta \otimes_{\mathcal{C}} \mathcal{M} \xrightarrow{\sim} \mathcal{M}$ [37, Lem. 3.7], [38, Lem. A.1] which expresses \mathcal{M} as an infinite twisted complex of Yoneda modules. Since \mathcal{M} is compact, the inverse quasi-isomorphism factors through some finite subcomplex, so \mathcal{M} is a retract of a finite twisted complex of Yoneda modules. \square

In particular, the inclusion $\mathcal{C} \subseteq \text{Perf } \mathcal{C}$ induces an equivalence $\text{Mod } \text{Perf } \mathcal{C} = \text{Mod } \mathcal{C}$.

A.4. Morita equivalence. We say categories \mathcal{C} and \mathcal{D} are Morita equivalent if there exist a $(\mathcal{C}, \mathcal{D})$ bimodule \mathcal{P} and a $(\mathcal{D}, \mathcal{C})$ bimodule \mathcal{Q} inducing, via convolution, an inverse pair of equivalences

$$(A.12) \quad \text{Mod } \mathcal{C} \xrightleftharpoons{\sim} \text{Mod } \mathcal{D}.$$

Actually, every equivalence $\text{Mod } \mathcal{C} \xrightarrow{\sim} \text{Mod } \mathcal{D}$ is isomorphic to convolution by a bimodule by [Theorem A.1](#) (since an equivalence is necessarily co-continuous), so \mathcal{C} and \mathcal{D} are Morita equivalent if and only if there is an equivalence $\text{Mod } \mathcal{C} = \text{Mod } \mathcal{D}$.

LEMMA A.4. *Categories \mathcal{C} and \mathcal{D} are Morita equivalent if and only if there is an equivalence $\text{Perf } \mathcal{C} = \text{Perf } \mathcal{D}$.*

Proof. An equivalence preserves compact objects, so an equivalence between $\text{Mod } \mathcal{C}$ and $\text{Mod } \mathcal{D}$ restricts to an equivalence $\text{Perf } \mathcal{C} = \text{Perf } \mathcal{D}$. Conversely, any equivalence $\text{Perf } \mathcal{C} = \text{Perf } \mathcal{D}$ induces an equivalence $\text{Mod } \mathcal{C} = \text{Mod } \text{Perf } \mathcal{C} = \text{Mod } \text{Perf } \mathcal{D} = \text{Mod } \mathcal{D}$. \square

In particular, the canonical inclusion $\mathcal{C} \hookrightarrow \text{Perf } \mathcal{C}$ is a Morita equivalence. In light of the above lemma, we will also refer to an equivalence $\text{Perf } \mathcal{C} = \text{Perf } \mathcal{D}$ as a Morita equivalence between \mathcal{C} and \mathcal{D} . We say a property of \mathcal{C} is “a Morita-invariant notion” if its validity only depends on $\text{Perf } \mathcal{C}$ up to equivalence.

A.5. Adjoints and compact objects. The following is a useful criterion for when a functor preserves compact objects.

LEMMA A.5. *If a functor $f : \mathbf{C} \rightarrow \mathbf{D}$ has a co-continuous right adjoint g , then f sends compact objects to compact objects.*

Proof. For $c \in \mathbf{C}$ a compact object, we have

$$(A.13) \quad \begin{aligned} \text{hom}_{\mathbf{D}}\left(f(c), \bigoplus_{\alpha} d_{\alpha}\right) &= \text{hom}_{\mathbf{C}}\left(c, g\left(\bigoplus_{\alpha} d_{\alpha}\right)\right) = \text{hom}_{\mathbf{C}}\left(c, \bigoplus_{\alpha} g(d_{\alpha})\right) \\ &= \bigoplus_{\alpha} \text{hom}_{\mathbf{C}}(c, g(d_{\alpha})) = \bigoplus_{\alpha} \text{hom}_{\mathbf{D}}(f(c), d_{\alpha}) \end{aligned}$$

as desired. \square

For example, if $f : \mathcal{C} \rightarrow \mathcal{D}$ is a functor of small categories, the pullback on module categories $f^* : \text{Mod } \mathcal{D} \rightarrow \text{Mod } \mathcal{C}$ is co-continuous and has a left adjoint $f_! : \text{Mod } \mathcal{C} \rightarrow \text{Mod } \mathcal{D}$ extending f (see [Section A.1](#)) which thus preserves compact objects (a fact which can also be seen from [Lemma A.3](#)).

A.6. Brown representability. On the level of large categories, a version of Brown representability gives effective criteria for deducing the existence of adjoints to functors.

THEOREM A.6 (Compare [69, Thm. 8.4.4] or [58, Cor. 5.5.2.9]). *Let \mathbf{C} and \mathbf{D} be large categories with \mathbf{C} compactly generated. If an A_∞ functor $f : \mathbf{C} \rightarrow \mathbf{D}$ is co-continuous, then f admits a right adjoint.*

Sketch of proof. We suppose that \mathbf{D} is also compactly generated, so one can write $\mathbf{C} = \text{Mod } \mathcal{C}$, $\mathbf{D} = \text{Mod } \mathcal{D}$ with $\mathcal{C} = \mathbf{C}^c$ and $\mathcal{D} = \mathbf{D}^c$. Then we observe that if f is co-continuous, it comes (by [Theorem A.1](#)) from convolving with a bimodule, which always has a right adjoint as described above. \square

[Theorem A.6](#) also holds under the weaker hypothesis that \mathbf{C} is *well generated* rather than compactly generated, by work of Neeman adapted to the dg/ A_∞ case. (For a definition of this notion, see [69, §8], and for a proof of [Theorem A.6](#) in that setting, see [69, Prop. 8.4.2 and Thm. 8.4.4].)

A.7. Quotients and localization. Given a (small) A_∞ (or dg) category \mathcal{C} and a full subcategory $\mathcal{D} \subseteq \mathcal{C}$, there is a well-defined notion of the quotient (dg or A_∞) category \mathcal{C}/\mathcal{D} which comes equipped with a functor

$$(A.14) \quad q : \mathcal{C} \rightarrow \mathcal{C}/\mathcal{D}.$$

(See [27], [60] for an explicit model in the dg and A_∞ cases respectively, also discussed in [37, §3.1.3].) The pair \mathcal{C}/\mathcal{D} and q satisfy the following universal property: any functor $\mathcal{C} \rightarrow \mathcal{E}$ which sends \mathcal{D} to 0 factors essentially uniquely through \mathcal{C}/\mathcal{D} via q ; more precisely, the pre-composition $q^* : \text{Fun}(\mathcal{C}/\mathcal{D}, \mathcal{E}) \hookrightarrow \text{Fun}(\mathcal{C}, \mathcal{E})$ fully faithfully embeds the former category as the full subcategory $\text{Fun}_{\text{Ann}(\mathcal{D})}(\mathcal{C}, \mathcal{E})$ of the latter consisting of functors from \mathcal{C} to \mathcal{E} which annihilate \mathcal{D} . Taking \mathcal{E} to be $(\text{Mod } \mathbb{Z})^{\text{op}}$, we note, in particular, that the pullback map

$$(A.15) \quad q^* : \text{Mod}(\mathcal{C}/\mathcal{D}) \rightarrow \text{Mod } \mathcal{C}$$

is a fully faithful embedding whose essential image is the \mathcal{C} modules which annihilate \mathcal{D} (see [37, Lem. 3.12 and 3.13]). It follows from these universal properties that the quotient \mathcal{C}/\mathcal{D} depends only on the full subcategory of \mathcal{C} split-generated by \mathcal{D} . If \mathcal{C} is pre-triangulated, then so is \mathcal{C}/\mathcal{D} , however be warned that \mathcal{C}/\mathcal{D} need not be idempotent complete even if \mathcal{C} is.

In light of (A.15), we have the following equivalent perspective on localization in terms of large categories. Let \mathbf{C} be a compactly generated large category, and let $\mathbf{D} \subseteq \mathbf{C}$ be a full subcategory closed under cones and arbitrary direct sums (hence itself a large category) which is compactly generated by a subset of \mathbf{C} 's compact objects $\mathcal{D} \subseteq \mathcal{C} := \mathbf{C}^c$. (Conversely, any full subcategory $\mathcal{D} \subseteq \mathcal{C}$ gives rise to such a $\mathbf{D} \subseteq \mathbf{C}$, namely the image of the induced functor $\text{Mod } \mathcal{D} \rightarrow \text{Mod } \mathcal{C}$, which is co-continuous since it is left adjoint to pull-back of modules, and is fully faithful because the unit of this adjunction is an isomorphism [37, Lem. 3.7].) The quotient of \mathbf{C} by \mathbf{D} , denoted \mathbf{C}/\mathbf{D} , is by definition the full subcategory of \mathbf{C} which is right-orthogonal to \mathbf{D} . (We

may also write this as \mathbf{C}/\mathbf{D} ; note that the universal property of direct sum implies that the right-orthogonal of \mathbf{D} is the same as the right-orthogonal of \mathbf{D} .) According to (A.15), this quotient \mathbf{C}/\mathbf{D} is precisely $\text{Mod}(\mathcal{C}/\mathcal{D})$. Thus the large quotient \mathbf{C}/\mathbf{D} is compactly generated, and passing to compact objects recovers the quotient of categories of compact objects, up to Morita equivalence. By the discussion in Section A.1, the embedding $q^* : \mathbf{C}/\mathbf{D} \hookrightarrow \mathbf{C}$ is right adjoint to a functor $q_! : \mathbf{C} \rightarrow \mathbf{C}/\mathbf{D}$ extending $q : \mathcal{C} \rightarrow \mathcal{C}/\mathcal{D}$. We thus conclude

LEMMA A.7 (Compatibility of large quotients with compact objects, compare [67, Thm. 2.1]). *Let $\mathbf{D} \subseteq \mathbf{C}$ be a co-continuous inclusion of compactly generated large categories which sends compact objects to compact objects. The large quotient \mathbf{C}/\mathbf{D} (by definition the right-orthogonal to $\mathbf{D} \subseteq \mathbf{C}$) is also a compactly generated large category with co-continuous inclusion into \mathbf{C} . The fully faithful inclusion $q^* : \mathbf{C}/\mathbf{D} \rightarrow \mathbf{C}$ is right adjoint to a “quotient functor” $q_! : \mathbf{C} \rightarrow \mathbf{C}/\mathbf{D}$ whose restriction to compact objects is the corresponding quotient functor on small categories $q : \mathcal{C} \rightarrow (\mathcal{C}/\mathcal{D})^\pi$ (with idempotent-completed target).*

If \mathcal{C} is a pre-triangulated dg/ A_∞ category and Z is a set of morphisms in $H^0(\mathcal{C})$, one can form the localization of \mathcal{C} with respect to Z by taking the quotient

$$(A.16) \quad \mathcal{C}[Z^{-1}] := \mathcal{C}/\text{cones } Z,$$

where $\text{cones } Z$ denotes any set of cones of morphisms in \mathcal{C} representing the elements in Z . (Regardless of how one chooses such a subset, one notices that $\text{cones } Z$ is a well-defined full subcategory of \mathcal{C} and, in particular, $\mathcal{C}[Z^{-1}]$ is unaffected by the choice.) If \mathcal{C} is not pre-triangulated, one can still define this localization by taking the essential image of \mathcal{C} under

$$(A.17) \quad \mathcal{C} \rightarrow \text{Tw } \mathcal{C} \rightarrow \text{Tw } \mathcal{C}/(\text{cones } Z).$$

The tautological localization map $\mathcal{C} \rightarrow \mathcal{C}[Z^{-1}]$ possesses a host of nice properties, simply as a special case of the properties of quotients discussed above; we leave it to the reader to spell out the details.

A.8. *Proper modules.* Recall that $\text{Perf } \mathbb{Z} \subseteq \text{Mod } \mathbb{Z}$ is the subcategory of perfect \mathbb{Z} -linear chain complexes, namely those chain complexes which are quasi-isomorphic to a bounded complex of finite projective \mathbb{Z} -modules.

We say a module or bimodule is proper (sometimes called pseudo-perfect in the literature) if as a functor to $\text{Mod } \mathbb{Z}$, it takes values in the full subcategory $\text{Perf } \mathbb{Z}$ (i.e., for a module \mathcal{M} if $\mathcal{M}(X)$ is a perfect chain complex for every $X \in \mathcal{C}$). Denote the full subcategory of proper modules by

$$(A.18) \quad \text{Prop } \mathcal{C} := \text{Fun}(\mathcal{C}^{\text{op}}, \text{Perf } \mathbb{Z}) \subseteq \text{Mod } \mathcal{C}.$$

A.9. Smooth and proper categories. We say a category \mathcal{C} is smooth (sometimes called homologically smooth) if its diagonal bimodule \mathcal{C}_Δ is perfect. (A $(\mathcal{C}, \mathcal{D})$ bimodule is called perfect if it is split-generated by tensor products of representable bimodules $\text{hom}_{\mathcal{C}}(-, X) \otimes \text{hom}_{\mathcal{D}}(Y, -)$.)

We say \mathcal{C} is proper (sometimes called compact) if its diagonal bimodule \mathcal{C}_Δ is proper, or if equivalently $\text{hom}_{\mathcal{C}}(X, Y)$ is a perfect \mathbb{Z} -module for any two objects $X, Y \in \mathcal{C}$. Smoothness and properness are Morita-invariant notions; in particular, \mathcal{C} is smooth (resp. proper) if and only if $\text{Perf } \mathcal{C}$ is.

In general, the subcategories of modules $\text{Perf } \mathcal{C}$ and $\text{Prop } \mathcal{C}$ do not coincide,¹⁷ however they are related under the above finiteness assumptions on \mathcal{C} :

LEMMA A.8. *If \mathcal{C} is proper, then $\text{Perf } \mathcal{C} \subseteq \text{Prop } \mathcal{C}$, and if \mathcal{C} is smooth, then $\text{Prop } \mathcal{C} \subseteq \text{Perf } \mathcal{C}$. In particular, if \mathcal{C} is smooth and proper, then $\text{Prop } \mathcal{C} = \text{Perf } \mathcal{C}$.*

LEMMA A.9. *Properness is inherited by full subcategories, and smoothness passes to quotients/localizations.*

A.10. Exceptional collections. We say a (full) subcategory of finitely many objects $\mathcal{A} \subseteq \mathcal{C}$ is an exceptional collection if there exists a partial ordering of the objects of \mathcal{A} such that

$$(A.19) \quad \text{hom}(X, X) = \mathbb{Z}\langle \text{id}_X \rangle,$$

$$(A.20) \quad \text{hom}(X, Y) = 0 \quad \text{unless } X \leq Y.$$

LEMMA A.10. *If $\mathcal{N} \in \text{Mod } \mathcal{C}$ is split-generated by an exceptional collection $\mathcal{A} \subseteq \mathcal{C}$, then \mathcal{N} is generated by \mathcal{A} (i.e., it is not necessary to add direct summands).*

Proof. Let $X \in \mathcal{A}$ be any maximal (with respect to the given partial order) object. We consider the functor

$$(A.21) \quad F_X : \text{Mod } \mathcal{C} \rightarrow \text{Mod } \mathbb{Z},$$

$$(A.22) \quad \mathcal{M} \mapsto \mathcal{M}(X).$$

Certainly if \mathcal{M} is generated by \mathcal{A} (i.e., by the Yoneda modules $\text{hom}_{\mathcal{C}}(-, A)$ for $A \in \mathcal{A}$), then $F_X(\mathcal{M}) \in \text{Perf } \mathbb{Z}$ by maximality of X , as all of the Yoneda modules except $\text{hom}_{\mathcal{C}}(-, X)$ contribute trivially to F_X , and each $\text{hom}_{\mathcal{C}}(-, X)$ contributes a perfect \mathbb{Z} -module.

There is a tautological map of \mathcal{C} modules $\text{hom}_{\mathcal{C}}(-, X) \otimes F_X(\mathcal{M}) \rightarrow \mathcal{M}(-)$; denote its cone by $\mathcal{M}|_{\mathcal{A} - \{X\}}$. Now given any maximal object Y of $\mathcal{A} - \{X\}$,

¹⁷Rather, they are in some sense “Morita dual” in that $\text{Prop } \mathcal{C} = \text{Fun}(\text{Perf } \mathcal{C}^{\text{op}}, \text{Perf } \mathbb{Z})$.

we may define a functor

$$(A.23) \quad F_Y : \text{Mod } \mathcal{C} \rightarrow \text{Mod } \mathbb{Z},$$

$$(A.24) \quad \mathcal{M} \mapsto \mathcal{M}|_{A-\{X\}}(Y).$$

Again, if \mathcal{M} is generated by \mathcal{A} , then $F_Y(\mathcal{M}) \in \text{Perf } \mathbb{Z}$. To see this, simply note that given a twisted complex \mathcal{M} of objects of \mathcal{A} , the object $\mathcal{M}|_{A-\{X\}}$ is just the same twisted complex but with all instances of X deleted. We may now similarly define $\mathcal{M}|_{A-\{X,Y\}}$ to be the cone of $\text{hom}_{\mathcal{C}}(-, Y) \otimes F_Y(\mathcal{M}) \rightarrow \mathcal{M}|_{A-\{X\}}(-)$.

Iterating this procedure defines a sequence of functors $F_X : \text{Mod } \mathcal{C} \rightarrow \text{Mod } \mathbb{Z}$ for all $X \in \mathcal{A}$. (In fact, these are independent of the order in which we pick off maximal elements, however we will not use this.) The above arguments show that for any \mathcal{M} generated by \mathcal{A} , all $F_X(\mathcal{M})$ are in $\text{Perf } \mathbb{Z}$ (and hence the same holds for \mathcal{M} split-generated by \mathcal{A}). They also show (for arbitrary \mathcal{M}) that if all $F_X(\mathcal{M})$ are in $\text{Perf } \mathbb{Z}$, then there exists $\mathcal{M}' \in \text{Mod } \mathcal{C}$ generated by \mathcal{A} and a map $\mathcal{M}' \rightarrow \mathcal{M}$ which is an isomorphism in $\text{Mod } \mathcal{A}$.

We may now conclude: if \mathcal{N} is split-generated by \mathcal{A} , then $F_X(\mathcal{N}) \in \text{Perf } \mathbb{Z}$, so there is $\mathcal{N}' \in \text{Mod } \mathcal{C}$ generated by \mathcal{A} and a map $\mathcal{N}' \rightarrow \mathcal{N}$ which is an isomorphism in $\text{Mod } \mathcal{A}$, and since \mathcal{N}' and \mathcal{N} are split-generated by \mathcal{A} , an isomorphism in $\text{Mod } \mathcal{A}$ is an isomorphism in $\text{Mod } \mathcal{C}$. \square

LEMMA A.11. *If \mathcal{A} is an exceptional collection which is proper, then it is smooth.*

Proof. In the case where \mathcal{A} has one object, this is true because \mathbb{Z} is trivially smooth. Now inductively apply the following assertion: If \mathcal{C} and \mathcal{D} are both smooth, and \mathcal{E} denotes the semi-orthogonal gluing of \mathcal{C} with \mathcal{D} along a $(\mathcal{C}, \mathcal{D})$ bimodule \mathcal{B} which is perfect, then \mathcal{E} is smooth as well. (See [56, Prop. 3.11] and [57, Thm. 3.24] for the dg case, which immediately extends to this setting.) In the assertion, observe it suffices that \mathcal{B} be proper, since proper bimodules over smooth categories are automatically perfect (by the bimodule version of Lemma A.8). Hence, one can induct from \mathbb{Z} to any proper exceptional collection \mathcal{A} . \square

References

- [1] A. ABBONDANDOLO, A. PORTALURI, and M. SCHWARZ, The homology of path spaces and Floer homology with conormal boundary conditions, *J. Fixed Point Theory Appl.* **4** no. 2 (2008), 263–293. [MR 2465553](#). [Zbl 1171.53344](#). <https://doi.org/10.1007/s11784-008-0097-y>.
- [2] A. ABBONDANDOLO and M. SCHWARZ, On the Floer homology of cotangent bundles, *Comm. Pure Appl. Math.* **59** no. 2 (2006), 254–316. [MR 2190223](#). [Zbl 1084.53074](#). <https://doi.org/10.1002/cpa.20090>.

- [3] A. ABBONDANDOLO and M. SCHWARZ, Floer homology of cotangent bundles and the loop product, *Geom. Topol.* **14** no. 3 (2010), 1569–1722. [MR 2679580](#). [Zbl 1201.53087](#). <https://doi.org/10.2140/gt.2010.14.1569>.
- [4] M. ABOUZAID, Morse homology, tropical geometry, and homological mirror symmetry for toric varieties, *Selecta Math. (N.S.)* **15** no. 2 (2009), 189–270. [MR 2529936](#). [Zbl 1204.14019](#). <https://doi.org/10.1007/s00029-009-0492-2>.
- [5] M. ABOUZAID, A geometric criterion for generating the Fukaya category, *Publ. Math. Inst. Hautes Études Sci.* no. 112 (2010), 191–240. [MR 2737980](#). [Zbl 1215.53078](#). <https://doi.org/10.1007/s10240-010-0028-5>.
- [6] M. ABOUZAID, A cotangent fibre generates the Fukaya category, *Adv. Math.* **228** no. 2 (2011), 894–939. [MR 2822213](#). [Zbl 1241.53071](#). <https://doi.org/10.1016/j.aim.2011.06.007>.
- [7] M. ABOUZAID, A topological model for the Fukaya categories of plumbings, *J. Differential Geom.* **87** no. 1 (2011), 1–80. [MR 2786590](#). [Zbl 1228.57015](#). <https://doi.org/10.4310/jdg/1303219772>.
- [8] M. ABOUZAID, On the wrapped Fukaya category and based loops, *J. Symplectic Geom.* **10** no. 1 (2012), 27–79. [MR 2904032](#). [Zbl 1298.53092](#). <https://doi.org/10.4310/JSG.2012.v10.n1.a3>.
- [9] M. ABOUZAID, Symplectic cohomology and Viterbo’s theorem, in *Free Loop Spaces in Geometry and Topology, IRMA Lect. Math. Theor. Phys.* **24**, Eur. Math. Soc., Zürich, 2015, pp. 271–485. [MR 3444367](#). [Zbl 1385.53078](#). <https://doi.org/10.4171/153>.
- [10] M. ABOUZAID and P. SEIDEL, An open string analogue of Viterbo functoriality, *Geom. Topol.* **14** no. 2 (2010), 627–718. [MR 2602848](#). [Zbl 1195.53106](#). <https://doi.org/10.2140/gt.2010.14.627>.
- [11] M. ABOUZAID and P. SEIDEL, Lefschetz fibration techniques in wrapped Floer cohomology, in preparation.
- [12] M. ABOUZAID and I. SMITH, Exact Lagrangians in plumbings, *Geom. Funct. Anal.* **22** no. 4 (2012), 785–831. [MR 2984118](#). [Zbl 1266.53073](#). <https://doi.org/10.1007/s00039-012-0162-y>.
- [13] B. H. AN, Y. BAE, and T. SU, Augmentations are sheaves for Legendrian graphs, *J. Symplectic Geom.* **20** no. 2 (2022), 259–415. [MR 4537126](#). [Zbl 1511.53082](#). <https://doi.org/10.4310/JSG.2022.v20.n2.a1>.
- [14] D. AUROUX, Fukaya categories and bordered Heegaard-Floer homology, in *Proceedings of the International Congress of Mathematicians. Volume II*, Hindustan Book Agency, New Delhi, 2010, available at author’s website, pp. 917–941. [MR 2827825](#). [Zbl 1275.53082](#).
- [15] D. AUROUX, Fukaya categories of symmetric products and bordered Heegaard-Floer homology, *J. Gökova Geom. Topol. GGT* **4** (2010), 1–54. [MR 2755992](#). [Zbl 1285.53077](#). Available at <https://gokovagt.org/journal/2010/auroux.html>.
- [16] D. AUROUX, L. KATZARKOV, and D. ORLOV, Mirror symmetry for weighted projective planes and their noncommutative deformations, *Ann. of Math. (2)* **167** no. 3 (2008), 867–943. [MR 2415388](#). [Zbl 1175.14030](#). <https://doi.org/10.4007/annals.2008.167.867>.

- [17] R. BEZRUKAVNIKOV and M. KAPRANOV, Microlocal sheaves and quiver varieties, *Ann. Fac. Sci. Toulouse Math.* (6) **25** no. 2-3 (2016), 473–516. [MR 3530166](#). [Zbl 1370.14026](#). <https://doi.org/10.5802/afst.1502>.
- [18] E. BIERSTONE and P. D. MILMAN, Semianalytic and subanalytic sets, *Inst. Hautes Études Sci. Publ. Math.* no. 67 (1988), 5–42. [MR 0972342](#). [Zbl 0674.32002](#). Available at http://www.numdam.org/item?id=PMIHES_1988_67_5_0.
- [19] F. BOURGEOIS, T. EKHOLM, and Y. ELIASHBERG, Effect of Legendrian surgery, *Geom. Topol.* **16** no. 1 (2012), 301–389, With an appendix by Sheel Ganatra and Maksim Maydanskiy. [MR 2916289](#). [Zbl 1322.53080](#). <https://doi.org/10.2140/gt.2012.16.301>.
- [20] K. CIELIEBAK and Y. ELIASHBERG, *From Stein to Weinstein and Back: Symplectic Geometry of Affine Complex Manifolds*, Amer. Math. Soc. Colloq. Publ., Amer. Math. Soc., Providence, RI, 2012. [MR 3012475](#). [Zbl 1262.32026](#). <https://doi.org/10.1090/coll/059>.
- [21] K. CIELIEBAK and J. LATSCHEV, The role of string topology in symplectic field theory, in *New Perspectives and Challenges in Symplectic Field Theory*, CRM Proc. Lecture Notes **49**, Amer. Math. Soc., Providence, RI, 2009, pp. 113–146. [MR 2555935](#). [Zbl 1214.53067](#). <https://doi.org/10.1090/crmp/049/04>.
- [22] M. CZAPLA, Definable triangulations with regularity conditions, *Geom. Topol.* **16** no. 4 (2012), 2067–2095. [MR 3033513](#). [Zbl 1375.14188](#). <https://doi.org/10.2140/gt.2012.16.2067>.
- [23] M. CZAPLA and W. PAWŁUCKI, Strict C^1 -triangulations in o-minimal structures, *Topol. Methods Nonlinear Anal.* **52** no. 2 (2018), 739–747. [MR 3915661](#). [Zbl 1423.32011](#). <https://doi.org/10.12775/TMNA.2018.033>.
- [24] S. K. DONALDSON, Symplectic submanifolds and almost-complex geometry, *J. Differential Geom.* **44** no. 4 (1996), 666–705. [MR 1438190](#). [Zbl 0883.53032](#). <https://doi.org/10.4310/jdg/1214459407>.
- [25] L. VAN DEN DRIES, *Tame Topology and o-minimal Structures*, London Math. Soc. Lect. Note Ser. **248**, Cambridge Univ. Press, Cambridge, 1998. [MR 1633348](#). [Zbl 0953.03045](#). <https://doi.org/10.1017/CBO9780511525919>.
- [26] L. VAN DEN DRIES and C. MILLER, Geometric categories and o-minimal structures, *Duke Math. J.* **84** no. 2 (1996), 497–540. [MR 1404337](#). [Zbl 0889.03025](#). <https://doi.org/10.1215/S0012-7094-96-08416-1>.
- [27] V. DRINFELD, DG quotients of DG categories, *J. Algebra* **272** no. 2 (2004), 643–691. [MR 2028075](#). [Zbl 1064.18009](#). <https://doi.org/10.1016/j.jalgebra.2003.05.001>.
- [28] A. I. EFIMOV, Sheaves with singular support condition form a left admissible subcategory, 2019, unpublished manuscript.
- [29] T. EKHOLM, Holomorphic curves for Legendrian surgery, 2019. [arXiv 1906.07228](#).
- [30] T. EKHOLM and Y. LEKILI, Duality between Lagrangian and Legendrian invariants, *Geom. Topol.* **27** no. 6 (2023), 2049–2179. [MR 4634745](#). [Zbl 07781510](#). <https://doi.org/10.2140/gt.2023.27.2049>.

- [31] T. EKHOLM, L. NG, and V. SHENDE, A complete knot invariant from contact homology, *Invent. Math.* **211** no. 3 (2018), 1149–1200. [MR 3763406](#). [Zbl 1385.57015](#). <https://doi.org/10.1007/s00222-017-0761-1>.
- [32] Y. ELIASHBERG and M. GROMOV, Convex symplectic manifolds, in *Several Complex Variables and Complex Geometry, Part 2* (Santa Cruz, CA, 1989), *Proc. Sympos. Pure Math.* **52**, Amer. Math. Soc., Providence, RI, 1991, pp. 135–162. [MR 1128541](#). [Zbl 0742.53010](#). <https://doi.org/10.1090/pspum/052.2/1128541>.
- [33] Y. ELIASHBERG and M. GROMOV, *Introduction to the h-principle*, *Grad. Stud. Math.* **48**, Amer. Math. Soc., Providence, RI, 2002. [MR 1909245](#). [Zbl 1008.58001](#). <https://doi.org/10.1090/gsm/048>.
- [34] T. ETGÜ and Y. LEKILI, Koszul duality patterns in Floer theory, *Geom. Topol.* **21** no. 6 (2017), 3313–3389. [MR 3692968](#). [Zbl 1378.57041](#). <https://doi.org/10.2140/gt.2017.21.3313>.
- [35] B. FANG, C.-C. M. LIU, D. TREUMANN, and E. ZASLOW, A categorification of Morelli’s theorem, *Invent. Math.* **186** no. 1 (2011), 79–114. [MR 2836052](#). [Zbl 1250.14011](#). <https://doi.org/10.1007/s00222-011-0315-x>.
- [36] B. GAMMAGE and V. SHENDE, Mirror symmetry for very affine hypersurfaces, *Acta Math.* **229** no. 2 (2022), 287–346. [MR 4554224](#). [Zbl 07681949](#). <https://doi.org/10.4310/ACTA.2022.v229.n2.a2>.
- [37] S. GANATRA, J. PARDON, and V. SHENDE, Covariantly functorial wrapped Floer theory on Liouville sectors, *Publ. Math. Inst. Hautes Études Sci.* **131** (2020), 73–200. [MR 4106794](#). [Zbl 1508.53091](#). <https://doi.org/10.1007/s10240-019-00112-x>.
- [38] S. GANATRA, J. PARDON, and V. SHENDE, Sectorial descent for wrapped Fukaya categories, *J. Amer. Math. Soc.* **37** no. 2 (2024), 499–635. [MR 4695507](#). [Zbl 7798120](#). <https://doi.org/10.1090/jams/1035>.
- [39] E. GIROUX, Remarks on Donaldson’s symplectic submanifolds, *Pure Appl. Math. Q.* **13** no. 3 (2017), 369–388. [MR 3882202](#). [Zbl 1403.53071](#). <https://doi.org/10.4310/pamq.2017.v13.n3.a1>.
- [40] E. GIROUX and J. PARDON, Existence of Lefschetz fibrations on Stein and Weinstein domains, *Geom. Topol.* **21** no. 2 (2017), 963–997. [MR 3626595](#). [Zbl 1372.32035](#). <https://doi.org/10.2140/gt.2017.21.963>.
- [41] M. GORESKY and R. MACPHERSON, *Stratified Morse Theory*, *Ergeb. Math. Grenzgeb.* **14**, Springer-Verlag, Berlin, 1988. [MR 0932724](#). [Zbl 0639.14012](#). <https://doi.org/10.1007/978-3-642-71714-7>.
- [42] S. GUILLERMOU, Quantization of conic Lagrangian submanifolds of cotangent bundles, 2015. [arXiv 1212.5818](#).
- [43] S. GUILLERMOU, M. KASHIWARA, and P. SCHAPIRA, Sheaf quantization of Hamiltonian isotopies and applications to nondisplaceability problems, *Duke Math. J.* **161** no. 2 (2012), 201–245. [MR 2876930](#). [Zbl 1242.53108](#). <https://doi.org/10.1215/00127094-1507367>.
- [44] K. HORI, S. KATZ, A. KLEMM, R. PANDHARIPANDE, R. THOMAS, C. VAFA, R. VAKIL, and E. ZASLOW, *Mirror Symmetry*, *Clay Math. Monogr.* **1**, Amer. Math. Soc., Providence, RI; Clay Math. Inst., Cambridge, MA, 2003, with a preface by Vafa. [MR 2003030](#). [Zbl 1044.14018](#).

- [45] B. HUGHES and A. RANICKI, *Ends of Complexes*, Cambridge Tracts in Math. **123**, Cambridge Univ. Press, Cambridge, 1996. MR 1410261. Zbl 0876.57001. <https://doi.org/10.1017/CBO9780511526299>.
- [46] Y. IKE and T. KUWAGAKI, Categorical localization for the coherent-constructible correspondence, *Publ. Res. Inst. Math. Sci.* **55** no. 1 (2019), 1–24. MR 3898322. Zbl 1456.14047. <https://doi.org/10.4171/PRIMS/55-1-1>.
- [47] B. IVERSEN, *Cohomology of Sheaves*, Universitext, Springer-Verlag, Berlin, 1986. MR 0842190. Zbl 1272.55001. <https://doi.org/10.1007/978-3-642-82783-9>.
- [48] X. JIN, Microlocal sheaf categories and the J -homomorphism, 2020. arXiv 2004.14270.
- [49] M. KASHIWARA and P. SCHAPIRA, *Sheaves on Manifolds*, Grundlehren Math. Wissen. **292**, Springer-Verlag, Berlin, 1990, with a chapter in French by Christian Houzel. MR 1074006. Zbl 0709.18001. <https://doi.org/10.1007/978-3-662-02661-8>.
- [50] M. KASHIWARA and P. SCHAPIRA, *Categories and Sheaves*, Grundlehren Math. Wissens. **332**, Springer-Verlag, Berlin, 2006. MR 2182076. Zbl 1118.18001. <https://doi.org/10.1007/3-540-27950-4>.
- [51] M. KONTSEVICH, Symplectic geometry of homological algebra, 2009, available at author’s website.
- [52] H. KRAUSE, On Neeman’s well generated triangulated categories, *Doc. Math.* **6** (2001), 121–126. MR 1836046. Zbl 0987.18012. <https://doi.org/10.4171/DM/99>.
- [53] H. KRAUSE, A Brown representability theorem via coherent functors, *Topology* **41** no. 4 (2002), 853–861. MR 1905842. Zbl 1009.18010. [https://doi.org/10.1016/S0040-9383\(01\)00010-6](https://doi.org/10.1016/S0040-9383(01)00010-6).
- [54] T. KUWAGAKI, The nonequivariant coherent-constructible correspondence for toric stacks, *Duke Math. J.* **169** no. 11 (2020), 2125–2197. MR 4132582. Zbl 1461.53064. <https://doi.org/10.1215/00127094-2020-0011>.
- [55] Y. LEKILI and A. POLISHCHUK, Homological mirror symmetry for higher-dimensional pairs of pants, *Compos. Math.* **156** no. 7 (2020), 1310–1347. MR 4120165. Zbl 1467.14099. <https://doi.org/10.1112/s0010437x20007150>.
- [56] V. A. LUNTS, Categorical resolution of singularities, *J. Algebra* **323** no. 10 (2010), 2977–3003. MR 2609187. Zbl 1202.18006. <https://doi.org/10.1016/j.jalgebra.2009.12.023>.
- [57] V. A. LUNTS and O. M. SCHNÜRER, Smoothness of equivariant derived categories, *Proc. Lond. Math. Soc.* (3) **108** no. 5 (2014), 1226–1276. MR 3214679. Zbl 1300.14046. <https://doi.org/10.1112/plms/pdt053>.
- [58] J. LURIE, *Higher Topos Theory*, Ann. of Math. Stud. **170**, Princeton Univ. Press, Princeton, NJ, 2009. MR 2522659. Zbl 1175.18001. <https://doi.org/10.1515/9781400830558>.
- [59] V. LYUBASHENKO, A_∞ -morphisms with several entries, *Theory Appl. Categ.* **30** (2015), Paper No. 45, 1501–1551. MR 3421458. Zbl 1348.18013. Available at <http://www.tac.mta.ca/tac/volumes/30/45/30-45abs.html>.

- [60] V. LYUBASHENKO and S. OVSIENKO, A construction of quotient A_∞ -categories, *Homology Homotopy Appl.* **8** no. 2 (2006), 157–203. [MR 2259271](#). [Zbl 1134.18002](#). <https://doi.org/10.4310/HHA.2006.v8.n2.a9>.
- [61] J. MATHER, Notes on topological stability, *Bull. Amer. Math. Soc. (N.S.)* **49** no. 4 (2012), 475–506. [MR 2958928](#). [Zbl 1260.57049](#). <https://doi.org/10.1090/S0273-0979-2012-01383-6>.
- [62] D. NADLER, Microlocal branes are constructible sheaves, *Selecta Math. (N.S.)* **15** no. 4 (2009), 563–619. [MR 2565051](#). [Zbl 1197.53116](#). <https://doi.org/10.1007/s00029-009-0008-0>.
- [63] D. NADLER, Wrapped microlocal sheaves on pairs of pants, 2016. [arXiv 1604.00114](#).
- [64] D. NADLER, Mirror symmetry for the Landau-Ginzburg A -model $M = \mathbb{C}^n$, $W = z_1 \cdots z_n$, *Duke Math. J.* **168** no. 1 (2019), 1–84. [MR 3909893](#). [Zbl 1411.53075](#). <https://doi.org/10.1215/00127094-2018-0036>.
- [65] D. NADLER and V. SHENDE, Sheaf quantization in Weinstein symplectic manifolds, 2022. [arXiv 2007.10154v3](#).
- [66] D. NADLER and E. ZASLOW, Constructible sheaves and the Fukaya category, *J. Amer. Math. Soc.* **22** no. 1 (2009), 233–286. [MR 2449059](#). [Zbl 1227.32019](#). <https://doi.org/10.1090/S0894-0347-08-00612-7>.
- [67] A. NEEMAN, The connection between the K -theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel, *Ann. Sci. École Norm. Sup. (4)* **25** no. 5 (1992), 547–566. [MR 1191736](#). [Zbl 0868.19001](#). <https://doi.org/10.24033/asens.1659>.
- [68] A. NEEMAN, On the derived category of sheaves on a manifold, *Doc. Math.* **6** (2001), 483–488. [MR 1874232](#). [Zbl 1026.18006](#). <https://doi.org/10.4171/DM/111>.
- [69] A. NEEMAN, *Triangulated Categories*, *Ann. of Math. Stud.* **148**, Princeton Univ. Press, Princeton, NJ, 2001. [MR 1812507](#). [Zbl 0974.18008](#). <https://doi.org/10.1515/9781400837212>.
- [70] A. NEEMAN, Explicit cogenerators for the homotopy category of projective modules over a ring, *Ann. Sci. Éc. Norm. Supér. (4)* **44** no. 4 (2011), 607–629. [MR 2919978](#). [Zbl 1258.16013](#). <https://doi.org/10.24033/asens.2151>.
- [71] L. NG, D. RUTHERFORD, V. SHENDE, S. SIVEK, and E. ZASLOW, Augmentations are sheaves, *Geom. Topol.* **24** no. 5 (2020), 2149–2286. [MR 4194293](#). [Zbl 1457.53064](#). <https://doi.org/10.2140/gt.2020.24.2149>.
- [72] M. PORTA, The Popescu-Gabriel theorem for triangulated categories, *Adv. Math.* **225** no. 3 (2010), 1669–1715. [MR 2673743](#). [Zbl 1227.18011](#). <https://doi.org/10.1016/j.aim.2010.04.002>.
- [73] M. ROBALO and P. SCHAPIRA, A lemma for microlocal sheaf theory in the ∞ -categorical setting, *Publ. Res. Inst. Math. Sci.* **54** no. 2 (2018), 379–391. [MR 3784874](#). [Zbl 1403.35016](#). <https://doi.org/10.4171/PRIMS/54-2-5>.
- [74] D. RUTHERFORD and M. SULLIVAN, Sheaves via augmentations of Legendrian surfaces, *J. Homotopy Relat. Struct.* **16** no. 4 (2021), 703–752. [MR 4343078](#). [Zbl 1496.57028](#). <https://doi.org/10.1007/s40062-021-00292-6>.

- [75] P. SCHAPIRA, Abelian sheaves, 2014, available at author's website.
- [76] W. SCHMID and K. VILONEN, Characteristic cycles of constructible sheaves, *Invent. Math.* **124** no. 1-3 (1996), 451–502. [MR 1369425](#). [Zbl 0851.32011](#). <https://doi.org/10.1007/s002220050060>.
- [77] P. SEIDEL, Graded Lagrangian submanifolds, *Bull. Soc. Math. France* **128** no. 1 (2000), 103–149. [MR 1765826](#). [Zbl 0992.53059](#). <https://doi.org/10.24033/bsmf.2365>.
- [78] P. SEIDEL, Fukaya categories and deformations, in *Proceedings of the International Congress of Mathematicians, Vol. II* (Beijing, 2002), Higher Ed. Press, Beijing, 2002, pp. 351–360. [MR 1957046](#). [Zbl 1014.53052](#).
- [79] P. SEIDEL, *Fukaya Categories and Picard-Lefschetz Theory*, Zurich Lect. Adv. Math., European Math. Soc. (EMS), Zürich, 2008. [MR 2441780](#). [Zbl 1159.53001](#). <https://doi.org/10.4171/063>.
- [80] V. SHENDE, The conormal torus is a complete knot invariant, *Forum Math. Pi* **7** (2019), e6. [MR 4010558](#). [Zbl 1426.57021](#). <https://doi.org/10.1017/fmp.2019.1>.
- [81] V. SHENDE, Microlocal category for Weinstein manifolds via the h -principle, *Publ. Res. Inst. Math. Sci.* **57** no. 3-4 (2021), 1041–1048. [MR 4322006](#). [Zbl 1483.53103](#). <https://doi.org/10.4171/prims/57-3-10>.
- [82] V. SHENDE, D. TREUMANN, H. WILLIAMS, and E. ZASLOW, On the combinatorics of exact Lagrangian surfaces, 2016. [arXiv 1603.07449](#).
- [83] V. SHENDE, D. TREUMANN, H. WILLIAMS, and E. ZASLOW, Cluster varieties from Legendrian knots, *Duke Math. J.* **168** no. 15 (2019), 2801–2871. [MR 4017516](#). [Zbl 1475.53094](#). <https://doi.org/10.1215/00127094-2019-0027>.
- [84] V. SHENDE, D. TREUMANN, and E. ZASLOW, Legendrian knots and constructible sheaves, *Invent. Math.* **207** no. 3 (2017), 1031–1133. [MR 3608288](#). [Zbl 1369.57016](#). <https://doi.org/10.1007/s00222-016-0681-5>.
- [85] M. SHIOTA, *Geometry of Subanalytic and Semialgebraic Sets*, Progr. Math. **150**, Birkhäuser Boston, Inc., Boston, MA, 1997. [MR 1463945](#). [Zbl 0889.32006](#). <https://doi.org/10.1007/978-1-4612-2008-4>.
- [86] M. SHIOTA, Whitney triangulations of semialgebraic sets, *Ann. Polon. Math.* **87** (2005), 237–246. [MR 2208550](#). [Zbl 1091.14016](#). <https://doi.org/10.4064/ap87-0-20>.
- [87] N. SPALTENSTEIN, Resolutions of unbounded complexes, *Compositio Math.* **65** no. 2 (1988), 121–154. [MR 0932640](#). [Zbl 0636.18006](#). Available at http://www.numdam.org/item?id=CM_1988_65_2_121_0.
- [88] Z. SYLVAN, On partially wrapped Fukaya categories, *J. Topol.* **12** no. 2 (2019), 372–441. [MR 3911570](#). [Zbl 1430.53097](#). <https://doi.org/10.1112/topo.12088>.
- [89] Z. SYLVAN, Orlov and Viterbo functors in partially wrapped Fukaya categories, 2019. [arXiv 1908.02317](#).
- [90] B. TOËN, The homotopy theory of dg-categories and derived Morita theory, *Invent. Math.* **167** no. 3 (2007), 615–667. [MR 2276263](#). [Zbl 1118.18010](#). <https://doi.org/10.1007/s00222-006-0025-y>.

[91] P. ZHOU, Lagrangian skeleta of hypersurfaces in $(\mathbb{C}^*)^n$, *Selecta Math. (N.S.)* **26** no. 2 (2020), Paper No. 26, 33. [MR 4087022](#). [Zbl 1439.53077](#). <https://doi.org/10.1007/s00029-020-00555-9>.

(Received: October 2, 2018)

(Revised: August 27, 2023)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA,
3620 S. VERMONT AVE., KAP 104, LOS ANGELES, CA 90089, USA

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, FINE HALL,
WASHINGTON RD., PRINCETON, NJ 08544, USA

Current address: SIMONS CENTER FOR GEOMETRY AND PHYSICS, STATE UNIVERSITY
OF NEW YORK, STONY BROOK, NY 11794, USA

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY,
BERKELEY, CA 94720, USA AND CENTER FOR QUANTUM MATHEMATICS, UNIVERSITY
OF SOUTHERN DENMARK, CAMPUSVEJ 55, 5230 ODENSE M, DENMARK