

Search for Boosted Dark Matter in COSINE-100

G. Adhikari,¹ N. Carlin,² J. J. Choi,^{3,4} S. Choi,³ A. C. Ezeribe,⁵ L. E. França,² C. Ha,⁶ I. S. Hahn,^{7,8,9} S. J. Hollick,¹ E. J. Jeon,⁴ J. H. Jo,¹ H. W. Joo,³ W. G. Kang,⁴ M. Kauer,¹⁰ B. H. Kim,⁴ H. J. Kim,¹¹ J. Kim,⁶ K. W. Kim,^{4,*} S. H. Kim,⁴ S. K. Kim,³ W. K. Kim,^{9,4} Y. D. Kim,^{4,12,9} Y. H. Kim,^{4,13,9} Y. J. Ko,⁴ D. H. Lee,¹¹ E. K. Lee,⁴ H. Lee,^{9,4} H. S. Lee,^{4,9,†} H. Y. Lee,⁴ I. S. Lee,⁴ J. Lee,⁴ J. Y. Lee,¹¹ M. H. Lee,^{4,9} S. H. Lee,^{9,4} S. M. Lee,³ Y. J. Lee,⁶ D. S. Leonard,⁴ N. T. Luan,¹¹ B. B. Manzano,² R. H. Maruyama,¹ R. J. Neal,⁵ J. A. Nikkel,¹ S. L. Olsen,⁴ B. J. Park,^{9,4} H. K. Park,¹⁴ H. S. Park,¹³ K. S. Park,⁴ S. D. Park,¹¹ R. L. C. Pitta,² H. Prihadi,⁴ S. J. Ra,⁴ C. Rott,^{15,16} K. A. Shin,⁴ D. F. F. S. Cavalcante,² A. Scarff,⁵ N. J. C. Spooner,⁵ W. G. Thompson,¹ L. Yang,¹⁷ and G. H. Yu^{15,4}

(COSINE-100 Collaboration)

¹Department of Physics and Wright Laboratory, Yale University, New Haven, Connecticut 06520, USA²Physics Institute, University of São Paulo, 05508-090, São Paulo, Brazil³Department of Physics and Astronomy, Seoul National University, Seoul 08826, Republic of Korea⁴Center for Underground Physics, Institute for Basic Science (IBS), Daejeon 34126, Republic of Korea⁵Department of Physics and Astronomy, University of Sheffield, Sheffield S3 7RH, United Kingdom⁶Department of Physics, Chung-Ang University, Seoul 06973, Republic of Korea⁷Department of Science Education, Ewha Womans University, Seoul 03760, Republic of Korea⁸Center for Exotic Nuclear Studies, Institute for Basic Science (IBS), Daejeon 34126, Republic of Korea⁹IBS School, University of Science and Technology (UST), Daejeon 34113, Republic of Korea¹⁰Department of Physics and Wisconsin IceCube Particle Astrophysics Center, University of Wisconsin-Madison, Madison, Wisconsin 53706, USA¹¹Department of Physics, Kyungpook National University, Daegu 41566, Republic of Korea¹²Department of Physics, Sejong University, Seoul 05006, Republic of Korea¹³Korea Research Institute of Standards and Science, Daejeon 34113, Republic of Korea¹⁴Department of Accelerator Science, Korea University, Sejong 30019, Republic of Korea¹⁵Department of Physics, Sungkyunkwan University, Suwon 16419, Republic of Korea¹⁶Department of Physics and Astronomy, University of Utah, Salt Lake City, Utah 84112, USA¹⁷Department of Physics, University of California San Diego, La Jolla, California 92093, USA

(Received 1 June 2023; accepted 30 October 2023; published 16 November 2023)

We search for energetic electron recoil signals induced by boosted dark matter (BDM) from the galactic center using the COSINE-100 array of NaI(Tl) crystal detectors at the Yangyang Underground Laboratory. The signal would be an excess of events with energies above 4 MeV over the well-understood background. Because no excess of events are observed in a 97.7 kg · yr exposure, we set limits on BDM interactions under a variety of hypotheses. Notably, we explored the dark photon parameter space, leading to competitive limits compared to direct dark photon search experiments, particularly for dark photon masses below 4 MeV and considering the invisible decay mode. Furthermore, by comparing our results with a previous BDM search conducted by the Super-Kamionkande experiment, we found that the COSINE-100 detector has advantages in searching for low-mass dark matter. This analysis demonstrates the potential of the COSINE-100 detector to search for MeV electron recoil signals produced by the dark sector particle interactions.

DOI: [10.1103/PhysRevLett.131.201802](https://doi.org/10.1103/PhysRevLett.131.201802)

A number of astrophysical observations provide evidence that the dominant matter component of the Universe is not ordinary matter but rather non-baryonic dark matter [1,2]. Many searches for signs of dark matter have been pursued by direct detection experiments [3–8], indirect detection experiments [9–14], and collider experiments [15–17] without success [18]. It motivates searches for alternative types of dark matter that produce substantially

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

different signatures in detectors, such as light (mass) dark matter models that predict extremely low-energy signals [19–22] or relativistically boosted dark matter (BDM) that produce more energetic signals [23–25].

A relativistic dark matter particle, i.e., one that is boosted by interactions with cosmic rays in the galaxy [26–35] or produced by the decay [36–38] or annihilation [39–41] of heavier dark sector particles, can deposit signals with energies that are above MeV in detectors. Since typical direct detection experiments search for low-energy nuclear recoil signals, scenarios for such energetic events have not been very well studied. Here we consider a model in which a BDM is produced by heavier dark matter particles [42–44]. It would require at least two species of dark matter particles, denoted by χ_0 and χ_1 for the heavier and lighter dark matter particles, respectively [42,45]. The first direct search for BDM from annihilations of heavy dark matter particles in the galactic center was performed with the Super-Kamiokande detector that searched for energetic electron recoil signals above 100 MeV induced by BDM elastic scattering [46]. With COSINE-100 data, we searched for the inelastic scattering of BDM (IBDM) [24] induced by the existence of another dark sector particle [40,41]. Recently, searches for cosmic-ray BDM interacting with protons in dark matter detectors with energies of a few keV [47,48], as well as in neutrino detectors with energies between a few MeV [49] and a few GeV [50], were performed.

In this Letter, we report on a search for BDM that elastically interacts with electrons in the NaI(Tl) crystals of the COSINE-100 detector. Such interactions would produce energetic electrons in the NaI(Tl) crystals. Our region of interest for the BDM interaction consists of an energy deposition above 4 MeV, since radioactive γ or β particles primarily have energy less than 4 MeV.

COSINE-100 [51] is composed of an array of eight ultra-pure NaI(Tl) crystals, each coupled to two photomultiplier tubes (PMTs). Because of high background levels and low light yields of three crystals, this analysis only uses data from five crystals, corresponding to an effective mass of 61.3 kg [52,53]. The crystals are immersed in an active veto detector that is composed of 2200 L of linear alkylbenzene (LAB)-based liquid scintillator (LS) [54]. The LS is contained within a shield comprising a 3 cm thick layer of oxygen-free copper, a 20 cm thick layer of lead, and an array of plastic scintillation counters for cosmic-ray muon tagging [55,56].

We used data obtained between 21 October 2016 and 18 July 2018, corresponding to 1.7 years of effective live time, and a total exposure of 97.7 kg · yr for this search. The same dataset was already adopted for a precise understanding and modeling of the observed background between 1 keV and 3 MeV [57], as well as for a dark matter search that concentrated on the low-energy nuclear recoil spectrum [52].

The COSINE-100 data acquisition system recorded two different signals from the crystal PMTs, covering a wide dynamic range from single-photoelectron to 4 MeV high energy [58]. In addition to the low-energy (0–100 keV) anode readout, the 5th stage dynode readout was recorded by 500 MHz flash analog-to-digital converters (FADCs) for 8 μ s long waveforms. It provided sufficient energy resolution for events with energies between 50 keV and 3 MeV.

We have previously presented a background model for the COSINE-100 detectors that covered energies below 3 MeV [57,59]. However, events with energies greater than 4 MeV were above the limit of the FADC dynamic range and suffered from a saturated, nonlinear response. To address this issue, we developed an algorithm to detect the saturation of the recorded pulse and reconstruct the saturated event. A template from the unsaturated events at the 2–3 MeV energy was compared to the saturated pulse, and the reconstruction at the saturated region was performed, as shown in Fig. 1(a). The original energy spectrum as well as the recovered energy spectrum are shown in Fig. 1(b).

The energy scale above 4 MeV is calibrated with 7.6 and 7.9 MeV γ rays from ^{56}Fe and ^{63}Cu , respectively, that are produced by thermal neutron capture in the steel supporter of the lead shield and the copper encapsulation of the NaI(Tl) crystals [51]. Figure 1(b) shows the reconstructed energy spectrum of the single hit events, in which the spectrum above 6 MeV is well described by the neutron capture events from GEANT4-based [60] simulation.

Candidate events are selected if the reconstructed energy is greater than 4 MeV with no coincident muon candidate tracks in the muon detector [55]. We reject α -induced events in the crystals using a pulse shape discrimination method [61]. Selected candidate events are sorted into two different categories: single-hit and multiple-hit events. A multiple-hit event has accompanying crystal signals with more than four photoelectrons or has a liquid scintillator signal above 80 keV [54]. A single-hit event is classified as one where the other detectors do not meet these criteria. Although a BDM interaction with the NaI(Tl) crystal would generate an energetic single electron with energy between a few MeV and a few 100 MeV [40,41], this energetic electron could generate a number of Bremsstrahlung radiation-induced γ s that could convert and deposit energy in the other crystals or the LS. Therefore, we use both single and multiple-hit channels in this analysis. Although the single-hit channel exhibits dominant sensitivity, the multiple-hit channel yields a slight enhancement in final sensitivity.

Four different categories of events contribute to the background above 4 MeV. Internal or external β/γ radiation induced by environmental radioactivities were well understood by the background modeling of the COSINE-100 detector for energy below 3 MeV [57,59]. We extended this model to energies above 4 MeV. Here the main contribution




FIG. 1. (a) An example of a saturated event due to the limited dynamic range (12 bit, 4096 for 2.5 V) is presented as a black solid line. Reconstruction of the saturated event (red dashed line) is achieved by comparison with the template from unsaturated events (thin blue solid line). In this example, a 6.02 MeV energy is reconstructed. (b) The measured energy spectra before (black-solid line) and after (red dots) reconstruction of the saturated events are presented. The reconstructed energy spectrum is calibrated with 7.6 and 7.9 MeV γ -rays from the neutron capture of iron and copper, respectively, as shown in the blue dotted line.

is caused by internal ^{228}Th decay, especially sequenced ^{212}Bi (β -decay with a Q value of 2.25 MeV) and ^{212}Po (α decay with a Q value of 8.95 MeV) decays with a 300 ns half-life of ^{212}Po . Because of the short half-life, ^{212}Bi and ^{212}Po events pile up in the 8 μs event window. Based on their distinct pulse shapes, we can partially reject them, but their residual is the main contribution above 4 MeV from environmental β/γ radiation.

Although the muon veto detector tags a coincident event with muons [56], $2.14 \pm 0.21\%$ of the muons that transit the detector are mis-tagged due to gaps between plastic scintillator panels. We applied a data-driven method to estimate the muon mis-tag contribution in the signal region, as described in Ref. [24]. Because of the 4π solid-angle coverage of the LS active shield [54], almost no events can reach the NaI(Tl) crystals without hits on the LS detector. Therefore, we do not consider the mis-tagged muon contribution for the single-hit events.

Thermal neutron capture by copper or iron nuclei in the shielding materials produces γ rays with energies as high as 8 MeV via (n, γ) reactions. The thermal neutron and total

neutron flux measured at the Yangyang underground laboratory are $(1.44 \pm 0.15) \times 10^{-5}/(\text{cm}^2 \cdot \text{s})$ and $(4.46 \pm 0.66) \times 10^{-5}/(\text{cm}^2 \cdot \text{s})$, respectively [62]. The neutron-induced events shown in Fig. 1(b) were simulated based on this flux.

In addition, we estimate the expected background events from ^8B solar neutrino elastic scattering on electrons. Table I presents the expected backgrounds from the aforementioned contributions for the single-hit and multiple-hit channels, which are compared with the measured data. The measured data agree with the total expected backgrounds within their uncertainties.

In Ref. [42], it is proposed that the boosted, lighter χ_1 dark matter particles are produced in the pair annihilation of two heavier χ_0 with a total flux,

$$\mathcal{F} = 1.6 \times 10^{-4} \text{ cm}^{-2} \text{ s}^{-1} \left(\frac{\langle \sigma v \rangle_{0 \rightarrow 1}}{5 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}} \right) \left(\frac{\text{GeV}}{m_0} \right)^2, \quad (1)$$

where the reference value for $\langle \sigma v \rangle_{0 \rightarrow 1}$, which is the velocity-averaged annihilation cross section of $\chi_0 \chi_0 \rightarrow \chi_1 \chi_1$,

TABLE I. The expected number of background events and the observed events from the 1.7 years COSINE-100 dataset are shown for both the single-hit and the multiple-hit channels. The individual contributions from environmental β/γ , thermal neutron capture, muon mis-tag, and solar neutrino are also listed.

Energy (MeV)	Single hit					Multiple hit				
	β/γ	Neutron	Neutrino	Total	Data	β/γ	Neutron	Muon	Total	Data
4–6	172 ± 26	203 ± 30	0.039 ± 0.006	375 ± 40	322	12 ± 2	889 ± 91	15 ± 2	915 ± 91	873
6–8	0	592 ± 63	0.024 ± 0.004	592 ± 63	545	0	1165 ± 120	16 ± 2	1181 ± 120	1194
8–10	0	60 ± 25	0.011 ± 0.002	60 ± 25	78	0	30 ± 11	21 ± 3	51 ± 12	37
> 10	0	0	0.003 ± 0.001	0.003 ± 0.001	0	0	2 ± 1	211 ± 4	213 ± 4	218




FIG. 2. The summed energy spectra for the five crystals (black solid circles) and the best fit (green solid lines) with BDM signal of $m_1 = 6$, $m_X = 13$ MeV, $\gamma_1 = 20$, $\epsilon = 8 \times 10^{-4}$ are presented for the single-hit events (a) and the multiple-hit events (b). Fitted contributions to the background from β/γ radiation, neutron-capture, and muon mis-tag are indicated. The green bands are the 68% confidence level (C.L.) intervals of the uncertainties obtained from the likelihood fit. For presentation purposes, we draw the BDM signal shapes assuming BDM parameters of $m_1 = 10$ MeV, $m_X = 1$ MeV, $\gamma_1 = 50$, $\epsilon = 3 \times 10^{-4}$ and $m_1 = 6$ MeV, $m_X = 13$ MeV, $\gamma_1 = 20$, $\epsilon = 8 \times 10^{-4}$ with $\times 30$ amplification of the signal amplitude.

corresponds to a correct dark matter thermal relic density for χ_0 that is derived by a so-called “assisted” freeze-out mechanism [45], and m_0 denotes the mass of χ_0 . This production rate is subject to uncertainties in the dark matter halo models [63–65]. Here we assume the NFW halo profile [63,66] described in Ref. [42]. The relativistic χ_1 (mass m_1) travels and interacts with terrestrial detector elements either elastically or inelastically. We consider $\chi_1 e^-$ elastic scattering via a mediator X (mass m_X) exchange.

We generate expected signals for various values of BDM parameters ($\gamma_1 = m_0/m_1$, m_X , and ϵ , where ϵ is the coupling between the dark sector mediator X and the electron) based on Refs. [40,41]. The generated signal events undergo detector simulation [57,59] and event selection. To search for BDM-induced events, we use a Bayesian approach with a likelihood function based on Poisson probability [52]. We perform binned maximum likelihood fits to the measured energy spectra for two different channels of the single-hit and the multiple-hit events for each signal of the various BDM parameters. Each crystal for each channel is fitted with a crystal- and channel-specific background model and a crystal- and channel-correlated BDM signal for the combined fit by multiplying the ten likelihoods of the five crystals. We use evaluated background contributions to set Gaussian priors for the known background rates.

Figure 2 presents an example of the maximum likelihood fit for BDM signals with assumed parameters of $m_1 = 6$ MeV, $m_X = 13$ MeV, $\gamma_1 = 20$, $\epsilon = 8 \times 10^{-4}$. The summed event spectra for the five crystals in the single-hit

(a) and multiple-hit (b) events are shown together with the best-fit result. For comparison, the expected signals for the BDM parameters $m_1 = 10$ MeV, $m_X = 1$ MeV, $\gamma_1 = 50$, $\epsilon = 3 \times 10^{-4}$ and $m_1 = 6$ MeV, $m_X = 13$ MeV, $\gamma_1 = 20$, $\epsilon = 8 \times 10^{-4}$ are presented. No excess of events that could be attributed to BDM interaction is found for the considered BDM signals. The posterior probabilities of signals are consistent with zero in all cases, and 90% C.L. upper limits are determined.

We interpret this result in the context of dark photon phenomenology by assuming that the interaction between the standard model particles and the dark sector particles is mediated by a dark photon. It allows us to compare this result with other dark photon searches in terms of the parameters m_X and ϵ . A similar interpretation with 59.6 days of COSINE-100 data for IBDM was presented in Ref. [24]. In our analysis, we generate signals using different sets of model parameters, fixing m_1 and γ_1 while varying m_X . Figure 3 shows the measured 90% C.L. upper limits obtained from the 1.7 yr of COSINE-100 data for the aforementioned model parameters. We compare our results with those of direct dark photon searches for both the visible decay mode ($m_X < 2m_1$) and the invisible decay mode ($m_X \geq 2m_1$) in Figs. 3(a) and 3(b), respectively. (Note that additional constraints from cosmological and astrophysical observations, depending on the detailed model of the dark sector particles discussed in Ref. [67], need to be taken into account.) Notably, for the invisible mode, our analysis yields a competitive limit for the dark photon mass below 4 MeV, assuming parameters of




FIG. 3. Interpretation of the BDM searches with 1.7 years of COSINE-100 data in terms of the dark photon mass (m_X) and coupling (ϵ) parameters is presented. (a) When $m_X < 2m_1$, our limits are compared with the currently excluded parameter space (shaded region) for the visible decay mode, including results from E141 [68], NA48 [69], NA64 [70], Babar [71], as well as bounds from the electron anomalous magnetic moment [72] and our previous IBDM search with 59.5 days of COSINE-100 data [24]. (b) When $m_X \geq 2m_1$, our limits are compared with the currently excluded parameter space (shaded region) for the invisible decay mode from NA64 [73] and Babar [74] experiments.

$m_1 = 0.5$ MeV and $\gamma_1 = 10$. This result highlights the complementarity of our search for the dark photon, although the specific model discussed in this Letter has to be assumed.

An additional interpretation of the Super-Kamiokande (SK) IV search result [23] considers the relic dark matter




FIG. 4. Interpretation of the BDM searches with 97.7 kg · years of COSINE-100 data in terms of the relic dark matter mass (m_0) and ϵ parameters is conducted, assuming $m_1 = 20$ and $m_X = 2$ MeV. This result is then compared with limits from 161.9 kiloton · years of SK IV data, assuming $m_1 = 200$ and $m_X = 20$ MeV [23]. To ensure a proper comparison with the SK result, we assume the coupling constant between χ_1 and X through elastic scattering to be 0.5.

mass (m_0) and coupling (ϵ) parameter space, as shown in Fig. 4. Two results are obtained using the same NFW halo profile [63] with a 0.5 coupling constant between the BDM and the mediator through the elastic interaction. Despite using a much smaller dataset of 97.7 kg · yr compared to the 161.9 kton · yr SK exposure, the lowest bound for ϵ at m_0 of about 200 MeV is at a similar level as that of the SK search result for an m_0 of about 5 GeV. Because of the well understood backgrounds in the COSINE-100 detector above 4 MeV, the COSINE-100 data is complementary to results from the SK detector in searching unexplored parameter space at low-mass dark matter.

In summary, we searched for evidence of boosted dark matter (BDM) by observing energetic electron recoil events induced by the elastic scattering of the BDM. Based on 1.7 yr of COSINE-100 data, we found no evidence of BDM interaction, and we set 90% C.L. limits for various model parameters. Our investigation of dark photon interactions explored a parameter space that complements other dark photon search experiments. We also demonstrate that a small-scale dark matter search detector has some unique advantages for the low-mass dark matter in the BDM scenario compared to the much larger neutrino detectors. Although our results are interpreted in the context of the BDM model that elastically scatters electron, this search can apply to any theory that predicts an excess of events in electron recoil of a few MeV, for which the COSINE-100 detector has world-competitive sensitivity.

We thank Jong-Chul Park and Seodong Shin for insightful discussions. We thank the Korea Hydro and Nuclear Power (KHNP) Company for providing underground laboratory space at Yangyang and the IBS Research Solution Center (RSC) for providing high performance computing resources. This work is supported by the Institute for Basic Science (IBS) under project code IBS-R016-A1, NRF-2021R1A2C3010989, and NRF-2021R1A2C1013761, Republic of Korea; NSF Grants No. PHY-1913742, No. DGE-1122492, WIPAC, the Wisconsin Alumni Research Foundation, United States; STFC Grants No. ST/N000277/1 and No. ST/K001337/1, United Kingdom; Grants No. 2021/06743-1 and No. 2022/12002-7 FAPESP, CAPES Finance Code 001, CNPq No. 131152/2020-3 and No. 303122/2020-0, Brazil.

*kwkim@ibs.re.kr

†hyunsulee@ibs.re.kr

- [1] D. Clowe, M. Bradač, A. H. Gonzalez, M. Markevitch, S. W. Randall, C. Jones, and D. Zaritsky, A direct empirical proof of the existence of dark matter, *Astrophys. J.* **648**, L109 (2006).
- [2] N. Aghanim *et al.* (Planck Collaboration), Planck 2018 results. VI. Cosmological parameters, *Astron. Astrophys.* **641**, A6 (2020); **652**, C4(E) (2021).

- [3] E. Aprile *et al.* (XENON Collaboration), Dark matter search results from a one tonne $\times$ year exposure of XENON1T, *Phys. Rev. Lett.* **121**, 111302 (2018).
- [4] Y. Meng *et al.* (PandaX-4T Collaboration), Dark matter search results from the PandaX-4T commissioning run, *Phys. Rev. Lett.* **127**, 261802 (2021).
- [5] J. Aalbers *et al.* (LUX-ZEPLIN Collaboration), First dark matter search results from the LUX-ZEPLIN (LZ) experiment, *Phys. Rev. Lett.* **131**, 041002 (2023).
- [6] E. Aprile *et al.* (XENON Collaboration and XENON Collaboration), First dark matter search with nuclear recoils from the XENONnT experiment, *Phys. Rev. Lett.* **131**, 041003 (2023).
- [7] J. Billard *et al.*, Direct detection of dark matter—APPEC committee report, *Rep. Prog. Phys.* **85**, 056201 (2022).
- [8] D. S. Akerib *et al.*, Snowmass2021 cosmic frontier dark matter direct detection to the neutrino fog, [arXiv:2203.08084](https://arxiv.org/abs/2203.08084).
- [9] K. Abe *et al.* (Super-Kamiokande Collaboration), Indirect search for dark matter from the Galactic Center and halo with the Super-Kamiokande detector, *Phys. Rev. D* **102**, 072002 (2020).
- [10] A. Albert *et al.* (ANTARES and IceCube Collaborations), Combined search for neutrinos from dark matter self-annihilation in the Galactic Center with ANTARES and IceCube, *Phys. Rev. D* **102**, 082002 (2020).
- [11] R. Abbasi *et al.* (IceCube Collaboration), Search for GeV-scale dark matter annihilation in the Sun with IceCube DeepCore, *Phys. Rev. D* **105**, 062004 (2022).
- [12] H. Abdalla *et al.* (H.E.S.S. Collaboration), Search for dark matter annihilation signals in the H.E.S.S. inner galaxy survey, *Phys. Rev. Lett.* **129**, 111101 (2022).
- [13] J. Conrad and O. Reimer, Indirect dark matter searches in gamma and cosmic rays, *Nat. Phys.* **13**, 224 (2017).
- [14] D. Hooper, TASI lectures on indirect searches for dark matter, *Proc. Sci. TASI2018* (2019) 010.
- [15] A. M. Sirunyan *et al.* (CMS Collaboration), Search for dark matter in events with a leptoquark and missing transverse momentum in proton-proton collisions at 13 TeV, *Phys. Lett. B* **795**, 76 (2019).
- [16] G. Aad *et al.* (ATLAS Collaboration), Search for dark matter in events with missing transverse momentum and a Higgs boson decaying into two photons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **10** (2021) 013.
- [17] O. Buchmueller, C. Doglioni, and L.-T. Wang, Search for dark matter at colliders, *Nat. Phys.* **13**, 217 (2017).
- [18] R. L. Workman *et al.* (Particle Data Group), Review of particle physics, *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [19] G. Adhikari *et al.* (COSINE-100 Collaboration), Searching for low-mass dark matter via the Migdal effect in COSINE-100, *Phys. Rev. D* **105**, 042006 (2022).
- [20] L. Barak *et al.* (SENSEI Collaboration), SENSEI: Direct-detection results on sub-GeV dark matter from a new skipper-CCD, *Phys. Rev. Lett.* **125**, 171802 (2020).
- [21] P. Agnes *et al.* (DarkSide Collaboration), Search for dark-matter–nucleon interactions via Migdal effect with DarkSide-50, *Phys. Rev. Lett.* **130**, 101001 (2023).
- [22] I. Arnquist *et al.* (DAMIC-M Collaboration), First constraints from DAMIC-M on Sub-GeV dark-matter particles interacting with electrons, *Phys. Rev. Lett.* **130**, 171003 (2023).
- [23] C. Kachulis *et al.* (Super-Kamiokande Collaboration), Search for boosted dark matter interacting with electrons in Super-Kamiokande, *Phys. Rev. Lett.* **120**, 221301 (2018).
- [24] C. Ha *et al.* (COSINE-100 Collaboration), First direct search for inelastic boosted dark matter with COSINE-100, *Phys. Rev. Lett.* **122**, 131802 (2019).
- [25] D. Kim, P. A. N. Machado, J.-C. Park, and S. Shin, Optimizing energetic light dark matter searches in dark matter and neutrino experiments, *J. High Energy Phys.* **07** (2020) 057.
- [26] T. Bringmann and M. Pospelov, Novel direct detection constraints on light dark matter, *Phys. Rev. Lett.* **122**, 171801 (2019).
- [27] Y. Ema, F. Sala, and R. Sato, Light dark matter at neutrino experiments, *Phys. Rev. Lett.* **122**, 181802 (2019).
- [28] C. V. Cappiello and J. F. Beacom, Strong new limits on light dark matter from neutrino experiments, *Phys. Rev. D* **100**, 103011 (2019); **104**, 069901(E) (2021).
- [29] C. Xia, Y.-H. Xu, and Y.-F. Zhou, Production and attenuation of cosmic-ray boosted dark matter, *J. Cosmol. Astropart. Phys.* **02** (2022) 028.
- [30] Y. Jho, J.-C. Park, S. C. Park, and P.-Y. Tseng, Cosmic-neutrino-boosted dark matter (ν BDM), [arXiv:2101.11262](https://arxiv.org/abs/2101.11262).
- [31] A. Das and M. Sen, Boosted dark matter from diffuse supernova neutrinos, *Phys. Rev. D* **104**, 075029 (2021).
- [32] D. Ghosh, A. Guha, and D. Sachdeva, Exclusion limits on dark matter-neutrino scattering cross section, *Phys. Rev. D* **105**, 103029 (2022).
- [33] D. Bardhan, S. Bhowmick, D. Ghosh, A. Guha, and D. Sachdeva, Bounds on boosted dark matter from direct detection: The role of energy-dependent cross sections, *Phys. Rev. D* **107**, 015010 (2023).
- [34] A. Granelli, P. Ullio, and J.-W. Wang, Blazar-boosted dark matter at Super-Kamiokande, *J. Cosmol. Astropart. Phys.* **07** (2022) 013.
- [35] J. Alvey, T. Bringmann, and H. Kolesova, No room to hide: Implications of cosmic-ray upscattering for GeV-scale dark matter, *J. High Energy Phys.* **01** (2023) 123.
- [36] A. Bhattacharya, R. Gandhi, and A. Gupta, The direct detection of boosted dark matter at high energies and PeV events at IceCube, *J. Cosmol. Astropart. Phys.* **03** (2015) 027.
- [37] J. Kopp, J. Liu, and X.-P. Wang, Boosted dark matter in IceCube and at the Galactic Center, *J. High Energy Phys.* **04** (2015) 105.
- [38] L. Heurtier, D. Kim, J.-C. Park, and S. Shin, Explaining the ANITA anomaly with inelastic boosted dark matter, *Phys. Rev. D* **100**, 055004 (2019).
- [39] H. Alhazmi, K. Kong, G. Mohlabeng, and J.-C. Park, Boosted dark matter at the deep underground neutrino experiment, *J. High Energy Phys.* **04** (2017) 158.
- [40] D. Kim, J.-C. Park, and S. Shin, Dark Matter “collider” from inelastic boosted dark matter, *Phys. Rev. Lett.* **119**, 161801 (2017).
- [41] G. F. Giudice, D. Kim, J.-C. Park, and S. Shin, Inelastic boosted dark matter at direct detection experiments, *Phys. Lett. B* **780**, 543 (2018).
- [42] K. Agashe, Y. Cui, L. Necib, and J. Thaler, (In)direct detection of boosted dark matter, *J. Cosmol. Astropart. Phys.* **10** (2014) 062.

- [43] K. Kong, G. Mohlabeng, and J.-C. Park, Boosted dark matter signals uplifted with self-interaction, *Phys. Lett. B* **743**, 256 (2015).
- [44] J. Berger, Y. Cui, and Y. Zhao, Detecting boosted dark matter from the sun with large volume neutrino detectors, *J. Cosmol. Astropart. Phys.* **02** (2015) 005.
- [45] G. Belanger and J.-C. Park, Assisted freeze-out, *J. Cosmol. Astropart. Phys.* **03** (2012) 038.
- [46] C. Kachulis *et al.* (Super-Kamiokande Collaboration), Search for boosted dark matter interacting with electrons in Super-Kamiokande, *Phys. Rev. Lett.* **120**, 221301 (2018).
- [47] X. Cui *et al.* (PandaX-II Collaboration), Search for cosmic-ray boosted Sub-GeV dark matter at the PandaX-II experiment, *Phys. Rev. Lett.* **128**, 171801 (2022).
- [48] R. Xu *et al.* (CDEX Collaboration), Constraints on sub-GeV dark matter boosted by cosmic rays from the CDEX-10 experiment at the China Jinping Underground Laboratory, *Phys. Rev. D* **106**, 052008 (2022).
- [49] M. Andriamirado *et al.* (PROSPECT Collaboration), Limits on sub-GeV dark matter from the PROSPECT reactor antineutrino experiment, *Phys. Rev. D* **104**, 012009 (2021).
- [50] K. Abe *et al.* (Super-Kamiokande Collaboration), Search for cosmic-ray boosted Sub-GeV dark matter using recoil protons at Super-Kamiokande, *Phys. Rev. Lett.* **130**, 031802 (2023); **131**, 159903(E) (2023).
- [51] G. Adhikari *et al.* (COSINE-100 Collaboration), Initial performance of the COSINE-100 experiment, *Eur. Phys. J. C* **78**, 107 (2018).
- [52] G. Adhikari *et al.* (COSINE-100 Collaboration), Strong constraints from COSINE-100 on the DAMA dark matter results using the same sodium iodide target, *Sci. Adv.* **7**, abk2699 (2021).
- [53] G. Adhikari *et al.* (COSINE-100 Collaboration), Three-year annual modulation search with COSINE-100, *Phys. Rev. D* **106**, 052005 (2022).
- [54] G. Adhikari *et al.* (COSINE-100 Collaboration), The COSINE-100 liquid scintillator veto system, *Nucl. Instrum. Methods Phys. Res., Sect. A* **1006**, 165431 (2021).
- [55] H. Prihtiadi *et al.* (COSINE-100 Collaboration), Muon detector for the COSINE-100 experiment, *J. Instrum.* **13**, T02007 (2018).
- [56] H. Prihtiadi *et al.* (COSINE-100 Collaboration), Measurement of the cosmic muon annual and diurnal flux variation with the COSINE-100 detector, *J. Cosmol. Astropart. Phys.* **02** (2021) 013.
- [57] G. Adhikari *et al.* (COSINE-100 Collaboration), Background modeling for dark matter search with 1.7 years of COSINE-100 data, *Eur. Phys. J. C* **81**, 837 (2021).
- [58] G. Adhikari *et al.* (COSINE-100 Collaboration), The COSINE-100 data acquisition system, *J. Instrum.* **13**, P09006 (2018).
- [59] P. Adhikari *et al.* (COSINE-100 Collaboration), Background model for the NaI(Tl) crystals in COSINE-100, *Eur. Phys. J. C* **78**, 490 (2018).
- [60] S. Agostinelli *et al.* (GEANT4 Collaboration), GEANT4—a simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [61] B. J. Park *et al.*, Development of ultra-pure NaI(Tl) detectors for the COSINE-200 experiment, *Eur. Phys. J. C* **80**, 814 (2020).
- [62] Y. S. Yoon, J. Kim, and H. Park, Neutron background measurement for rare event search experiments in the YangYang underground laboratory, *Astropart. Phys.* **126**, 102533 (2021).
- [63] J. F. Navarro, C. S. Frenk, and S. D. M. White, The structure of cold dark matter halos, *Astrophys. J.* **462**, 563 (1996).
- [64] A. V. Kravtsov, A. A. Klypin, J. S. Bullock, and J. R. Primack, The cores of dark matter dominated galaxies: Theory versus observations, *Astrophys. J.* **502**, 48 (1998).
- [65] B. Moore, T. R. Quinn, F. Governato, J. Stadel, and G. Lake, Cold collapse and the core catastrophe, *Mon. Not. R. Astron. Soc.* **310**, 1147 (1999).
- [66] J. F. Navarro, C. S. Frenk, and S. D. M. White, A universal density profile from hierarchical clustering, *Astrophys. J.* **490**, 493 (1997).
- [67] A. Kamada, H. J. Kim, J.-C. Park, and S. Shin, Manifesting hidden dynamics of a sub-component dark matter, *J. Cosmol. Astropart. Phys.* **10** (2022) 052.
- [68] E. M. Riordan *et al.*, A search for short lived axions in an electron beam dump experiment, *Phys. Rev. Lett.* **59**, 755 (1987).
- [69] J. R. Batley *et al.* (NA48/2 Collaboration), Search for the dark photon in π^0 decays, *Phys. Lett. B* **746**, 178 (2015).
- [70] D. Banerjee *et al.* (NA64 Collaboration), Improved limits on a hypothetical X(16.7) boson and a dark photon decaying into e^+e^- pairs, *Phys. Rev. D* **101**, 071101 (2020).
- [71] J. P. Lees *et al.* (BABAR Collaboration), Search for a dark photon in e^+e^- collisions at BABAR, *Phys. Rev. Lett.* **113**, 201801 (2014).
- [72] H. Davoudiasl, H.-S. Lee, and W. J. Marciano, Muon g-2, rare kaon decays, and parity violation from dark bosons, *Phys. Rev. D* **89**, 095006 (2014).
- [73] Y. M. Andreev *et al.*, Improved exclusion limit for light dark matter from e^+e^- annihilation in NA64, *Phys. Rev. D* **104**, L091701 (2021).
- [74] J. P. Lees *et al.* (BABAR Collaboration), Search for invisible decays of a dark photon produced in e^+e^- collisions at BABAR, *Phys. Rev. Lett.* **119**, 131804 (2017).