

# Bis(N-xylyl-imino)phenyl “NCN” Iridium Pincer Complexes.

## Thermodynamics of Ligand Binding and C-C Bond Cleavage

Soumyadipa Das, Souvik Mandal, Santanu Malakar, Thomas J. Emge, Alan S. Goldman\*

*Department of Chemistry and Chemical Biology, Rutgers, The State University of New Jersey, New Brunswick, New Jersey 08854, United States*

Dedicated to Prof. Peter T. Wolczanski on the occasion of this 70th birthday

### Abstract

Iridium dibromide complexes of the phenyldiimine ligand 2,6-bis(1-((2,6-dimethylphenyl)imino)ethyl)phenyl, *trans*-(<sup>x</sup>PhDI)IrBr<sub>2</sub>L, have been synthesized, and relative Ir-L BDFEs have been experimentally determined for a wide range of corresponding adducts of ligands L. An estimate of the absolute enthalpy of Ir-L binding has been obtained from dynamic NMR measurements. The results of DFT calculations are in very good agreement with the relative and absolute experimental values. Computational studies were extended to the formation of adducts of (<sup>x</sup>PhDI)IrH<sub>2</sub> and (<sup>x</sup>PhDI)Ir<sup>I</sup>, as well as other (pincer)Ir<sup>I</sup> fragments, (Phebox)Ir<sup>I</sup> and (PCP)Ir<sup>I</sup>, to enable a comparison of electronic and steric effects with these archetypal pincer ligands. Attempts to reduce (<sup>x</sup>PhDI)IrBr<sub>2</sub>(MeCN) to a hydride or an Ir<sup>I</sup> complex yielded a dinuclear CN-bridged complex with a methyl ligand on the cyanide-C-bound Ir center (characterized by scXRD), indicating that C-CN bond cleavage took place at that Ir center. DFT calculations indicate that the C-CN bond cleavage occurs at one Ir center with strong assistance by coordination of the CN nitrogen to the other Ir center.


**Keywords:** C-C bond cleavage; metal-ligand bond strengths; iridium; pincer ligands; phenyldiimine

## 1. Introduction

Iridium complexes bearing 2,6-bisphosphinomethyl aryl (PCP motif) and many related pincer ligands, including those with PNP, PPP, and bis(NHC)aryl (CCC) motifs, have been explored and developed extensively in the past 25 years. In particular such complexes have seen great success in C-H bond activation including catalytic alkane dehydrogenation and tandem reactions based upon dehydrogenation<sup>1-11</sup>, as well as reactions involving cleavage and formation of C-O<sup>12, 13</sup>, N-H<sup>14-16</sup>, and other strong bonds<sup>17</sup>. Iridium complexes of pincer ligands with terminal N-coordinating groups (e.g. NCN-type, such as 2,6-bis-oxazolinephenyl, i.e. Phebox) have also seen development in catalysis and strong-bond activation. This chemistry, however, has typically not been analogous to that of the aforementioned ligands with phosphino- or carbene-coordinating “arms” (terminal groups) which largely operates via Ir(I) complexes. Instead, the chemistry of such NCN-iridium complexes has largely focused on Ir(III) carboxylate complexes that are believed to operate via concerted metalation-deprotonation (CMD) or other mechanisms involving high-oxidation complexes.<sup>18-25</sup>

N-coordinating groups potentially offer significant advantages over P-coordinating and other “soft” groups, including ease of ligand synthesis, cost, and resistance to oxidation. From a fundamental perspective, it is of interest to understand the effect of such variations on catalytic or stoichiometric reactivity. However, whereas PXP-type pincer ligands (typically, X = C or N) largely incorporate sterically demanding phosphino groups (e.g. P<sup>t</sup>Bu<sub>2</sub> or P<sup>i</sup>Pr<sub>2</sub>), the NXN pincer ligands explored in this context have largely made use of groups such as oxazolines, which are much less bulky, and in which the limited steric bulk is positioned very differently than in the PCP complexes. These factors obfuscate any meaningful comparisons.

In this context we wished to explore the chemistry of an NCN complex with a relatively bulky N-coordinating group, and particularly one in which the steric bulk is not positioned only near the coordination site trans to the coordinating aryl carbon as in the case of Phebox. Toward this end we have synthesized adducts, **1-L**, of the iridium dibromide complex of the phenyldiimine ligand, 2,6-bis(1-((2,6-dimethylphenyl)imino)ethyl)phenyl (<sup>xv</sup>PhDI). The complexes isolated represent somewhat unusual examples of late-metal pincer complexes bearing two electron-withdrawing low-field ancillary ligand (bromides). We have investigated the thermodynamics of the binding of the various ligands, L. DFT calculations are found to be in very good agreement with the experimental results. Encouraged by this agreement, we compare these results with binding thermodynamics calculated for the corresponding hydrides, and for the Ir(I) complexes of <sup>xv</sup>PhDI and other pincer ligands. In an effort to synthesize the dihydride, which is of particular interest as a precursor of the corresponding Ir(I) fragment, we treated the dibromide precursor **1-MeCN** with KO<sup>t</sup>Bu under H<sub>2</sub> atmosphere. This resulted in the formation of a bimetallic species, the structure of which was determined crystallographically. Remarkably the molecular structure revealed that one equivalent CH<sub>3</sub>CN had been hydrogenated to give ethylamine, and a second equivalent had undergone C-C bond cleavage to give an iridium center with a methyl and C-bound cyanide bridged to the second metal center.





## 2. Results and Discussion

### 2.1. Synthesis of (<sup>xy</sup>PhDI)Ir complexes


**Experimental results.** 1,1'-(2-bromo-1,3-phenylene)bis(N-(2,6-dimethylphenyl)ethan-1-imine)<sup>26</sup> (<sup>xy</sup>PhDI-Br) was metalated by the reaction with [Ir(COD)Cl]<sub>2</sub> (COD = 1,5-cyclooctadiene) (Scheme 1) following the procedure reported by Oakley *et al.* for synthesis of aldimine analogues.<sup>27</sup> Crystals of **1-MeCN** were grown by diffusion of pentane into a THF solution at room temperature, and the molecular structure was determined by scXRD (Figure 1).

#### Scheme 1. Synthesis of **1-MeCN**



**Figure 1.** Crystallographically determined molecular structure of **1-MeCN**


Acetonitrile was displaced from **1-MeCN** by reaction with 1.2 equiv of pyridine, PMe<sub>3</sub>, N-ethylamine, or P(OMe)<sub>3</sub>, or CO (1 atm), to give **1-py**, **1-PMe<sub>3</sub>**, **1-NH<sub>2</sub>Et**, **1-P(OMe)<sub>3</sub>** and **1-CO** respectively. Their molecular structures are shown in Figures 2a-e.



**Figure 2.** Crystallographically determined molecular structures of (a) **1-py** (b) **1-PMe<sub>3</sub>** (c) **1-NH<sub>2</sub>Et** (d) **1-CO** (e) **1-P(OMe)<sub>3</sub>** (f) **1-C<sub>2</sub>H<sub>4</sub>**

Bubbling ethylene through a toluene solution of **1-MeCN** to dryness yielded a solid that was redissolved in benzene under argon atmosphere. Crystals were obtained by slow evaporation and the molecular structure of the product, **1-C<sub>2</sub>H<sub>4</sub>**, was determined by scXRD (Figure 2f).

Addition of ethylene atmosphere to a toluene solution of **1-MeCN** (without bubbling to dryness) led to a mixture of **1-MeCN** and **1-C<sub>2</sub>H<sub>4</sub>**. The equilibrium of Scheme 2 was established with  $K_{eq} = 0.181$  ( $\Delta G^\circ = 1.0$  kcal/mol).

**Scheme 2.** Equilibrium between **1-MeCN** and **1-C<sub>2</sub>H<sub>4</sub>**.

Bubbling a solution of **1-C<sub>2</sub>H<sub>4</sub>** to dryness with N<sub>2</sub> gas resulted in no substitution, nor did the analogous approach lead to substitution with H<sub>2</sub> gas. It appears that these species bind much more weakly than C<sub>2</sub>H<sub>4</sub>, if at all. Likewise addition of 1-hexene to a solution of **1-C<sub>2</sub>H<sub>4</sub>** did not result in any observable substitution.

Addition of CO (1.9 atm) to a benzene-d<sub>6</sub> solution of **1-PMe<sub>3</sub>** did not result in any substitution or other reaction. Conversely, however, addition of 1.2 equiv PMe<sub>3</sub> to a solution of **1-CO** resulted in complete conversion to **1-PMe<sub>3</sub>**. Given that CO typically binds very strongly to iridium complexes, including Ir(III) complexes, we found this result to be somewhat intriguing.<sup>28, 29</sup>

## 2.2. Thermodynamics of ligand binding

### 2.2.1. Relative energies of ligand binding to (<sup>x</sup>PhDI)Ir: experimental and computational results.

Equilibrium constants were determined for various pairs of ligands. In all cases the equilibrium was reached from both directions, by starting with a given complex **1-L**, adding the complementary ligand L', waiting until equilibrium was apparently reached, and then adding an additional quantity of ligand L.  $K_{eq}$  was determined for the following ligand pairs at 25 °C: MeCN/C<sub>2</sub>H<sub>4</sub>,  $K_{eq}$  = 0.181 (Scheme 2); MeCN/*t*BuNH<sub>2</sub>,  $K_{eq}$  = 32.0; *t*BuNH<sub>2</sub>/*i*PrNH<sub>2</sub>,  $K_{eq}$  = 48.2; *i*PrNH<sub>2</sub>/py,  $K_{eq}$  = 3.71; py/EtNH<sub>2</sub>,  $K_{eq}$  = 2.63; py/CO,  $K_{eq}$  = 13.8; EtNH<sub>2</sub>/CO,  $K_{eq}$  = 4.95; EtNH<sub>2</sub>/PPh<sub>2</sub>Me,  $K_{eq}$  = 2.15. For PPh<sub>2</sub>Me/P(OMe)<sub>3</sub>,  $K_{eq}$  = 29.2 as determined at 80 °C. The corresponding relative free energies of binding are given in Table 1.

**Table 1.** Free energies of binding to **1** relative to **1-C<sub>2</sub>H<sub>4</sub>**

| Ligand                        | Experimental | Calculated | $\Delta(\text{Calc} - \text{Exptl})$ |
|-------------------------------|--------------|------------|--------------------------------------|
| H <sub>2</sub>                | > 0          | 12.28      | --                                   |
| N <sub>2</sub>                | > 0          | 7.57       | --                                   |
| 1-hexene                      | > 0          | 3.00       | --                                   |
| C <sub>2</sub> H <sub>4</sub> | [0.0]        | [0.0]      | [0.0]                                |
| MeCN                          | -1.01        | -0.82      | 0.19                                 |
| <i>t</i> -BuNH <sub>2</sub>   | -3.05        | -4.01      | -0.96                                |
| <i>i</i> -PrNH <sub>2</sub>   | -5.34        | -7.19      | -1.85                                |
| pyridine                      | -6.11        | -6.75      | -0.64                                |
| EtNH <sub>2</sub>             | -6.68        | -7.58      | -0.90                                |
| PPh <sub>2</sub> Me           | -7.13        | -8.69      | -1.56                                |
| CO                            | -7.64        | -8.52      | -0.88                                |
| P(OMe) <sub>3</sub>           | -9.49        | -12.11     | -2.62                                |
| PMe <sub>3</sub>              | << -9.5      | -13.72     | --                                   |

Electronic structure (DFT) calculations were then conducted, initially for those complexes that were studied experimentally. Geometries were optimized in the gas phase using the M06 functional<sup>30</sup> and split valence basis set 6-31G(d,p) for C, H, N, O and P<sup>31-35</sup>. For Ir, the Stuttgart-Dresden effective core potential was used for the 60 core electrons; the associated basis set was used for the 17 valence electrons (SDD).<sup>36</sup> All calculations were done at standard conditions of temperature (298.15K) and pressure (1 atm); full details are given in the SI. Ir-L binding energies were calculated and are given in Table 2 (absolute BDFEs)(see SI for electronic energy, enthalpy and entropy of binding). The calculated relative free energies of binding are in excellent agreement with experimentally obtained values (Table 1). For those ligands for which relative binding free energies were obtained experimentally, the root-mean-

square deviation of all calculated relative values compared with all relative values extrapolated from the experimental measurements is 0.8 kcal/mol<sup>37</sup>; we consider this to be very satisfactory agreement.


**Table 2.** Calculated free energies ( $\Delta G^\circ$ , kcal/mol) of binding of various ligands to *trans*-(<sup>Xy</sup>PhDI)IrBr<sub>2</sub> (**1**), *trans*-(<sup>Xy</sup>PhDI)IrH<sub>2</sub> (**2**)<sup>a</sup>, (<sup>Xy</sup>PhDI)Ir (**3**), (Phebox)Ir (**4**), (<sup>i</sup>PrPCP)Ir (**5**) and (<sup>t</sup>BuPCP)Ir (**6**)

| L                              | ( <sup>Xy</sup> PhDI)IrBr <sub>2</sub> | ( <sup>Xy</sup> PhDI)IrH <sub>2</sub> | ( <sup>Xy</sup> PhDI)Ir | (Phebox)Ir | ( <sup>i</sup> PrPCP)Ir | ( <sup>t</sup> BuPCP)Ir |
|--------------------------------|----------------------------------------|---------------------------------------|-------------------------|------------|-------------------------|-------------------------|
| H <sub>2</sub>                 | 2.7                                    | -5.5                                  | -6.7                    | -9.2       | -13.8                   | -9.5                    |
| N <sub>2</sub>                 | -2.0                                   | -10.3                                 | -16.1                   | -17.2      | -20.1                   | -19.2                   |
| 1-hexene                       | -6.6                                   | -12.9                                 | -20.1                   | -15.1      | -21.4                   | -8.0                    |
| C <sub>2</sub> H <sub>4</sub>  | -9.6                                   | -14.0                                 | -22.6                   | -16.4      | -23.7                   | -16.7                   |
| MeCN                           | -10.4                                  | -14.0                                 | -18.0                   | -16.4      | -18.4                   | -17.1                   |
| <sup>t</sup> BuNH <sub>2</sub> | -13.6                                  | -19.1                                 | -18.6                   | -9.1       | -12.8                   | -5.5                    |
| pyridine                       | -16.3                                  | -18.1                                 | -18.8                   | -16.4      | -17.2                   | -14.7                   |
| <sup>i</sup> PrNH <sub>2</sub> | -16.8                                  | -21.2                                 | -21.3                   | -11.3      | -16.3                   | -10.7                   |
| EtNH <sub>2</sub>              | -17.1                                  | -20.3                                 | -20.4                   | -13.2      | -15.8                   | -13.9                   |
| CO                             | -18.1                                  | -32.5                                 | -37.4                   | -43.0      | -49.3                   | -48.1                   |
| PPh <sub>2</sub> Me            | -18.3                                  | -25.6                                 | -23.9                   | -20.2      | -32.6                   | -14.6                   |
| P(OMe) <sub>3</sub>            | -21.7                                  | -32.2                                 | -33.1                   | -31.5      | -45.9                   | -35.0                   |
| PMe <sub>3</sub>               | -23.3                                  | -28.2                                 | -27.4                   | -17.8      | -29.4                   | -17.0                   |

(a) See reference 38

With potential ligands H<sub>2</sub>, N<sub>2</sub> and 1-hexene we were unable to observe any displacement of C<sub>2</sub>H<sub>4</sub>, this is in agreement with their calculated low energies of binding. For H<sub>2</sub> and N<sub>2</sub> in particular, the failure to observe substitution or loss of **1-C<sub>2</sub>H<sub>4</sub>**, even after bubbling solutions of **1-C<sub>2</sub>H<sub>4</sub>** to dryness with the respective gas (and thereby providing a very strong entropic driving force for substitution) indicates that these molecules bind particularly weakly to **1** (if at all). The very unfavorable calculated free binding energies are consistent with this result. At the other extreme, none of the ligands used in this study were able to displace PMe<sub>3</sub> to any observable extent, in accord with its calculated very high relative free binding energy (13.7 kcal/mol greater than ethylene, Table 1).

Having established the ability of the computational method to reliably calculate relative energies of binding to **1**, we used such calculation to study binding of the same ligands to related Ir fragments, specifically the corresponding dihydride (**2**), and the 14-electron Ir(I) fragment, (<sup>Xy</sup>PhDI)Ir (**3**; no ancillary ligands) (Figure 3). We calculate that for amines and phosphines, the variability of the energy of binding to these three fragments was fairly small, with bond dissociation free energies (BDFEs) for each ligand found to be within a range spanning 6 kcal/mol. In contrast, the BDFE of CO increased strongly over the series **1** < **2** < **3** with the free energy of CO binding to Ir(I) fragment **3** (37.4 kcal/mol) being 19.3 kcal/mol greater ( $\Delta G_1 - \Delta G_3$ ) than binding to Ir(III) fragment **1** (18.1 kcal/mol). Presumably this large variability reflects the degree of increasing  $\pi$ -donating ability among the various fragments, and commensurately increased metal-ligand  $\pi$ -backbonding.



<https://doi.org/10.26434/chemrxiv-2023-r4jt6> **ORCID:** <https://orcid.org/0009-0008-4667-930X> Content not peer-reviewed by ChemRxiv. License: CC BY-NC-ND 4.0

exception to that rule (e.g. the calculated free energy of N<sub>2</sub> binding to **1** is only 2.0 kcal/mol). But although N<sub>2</sub> is also generally considered to be a much weaker  $\pi$ -acceptor than CO<sup>39</sup> the variation of N<sub>2</sub> binding energies among the complexes is nearly as great as that found for CO; N<sub>2</sub> is calculated to bind 14.1 kcal/mol more strongly to fragment **3** than to fragment **1** ( $\Delta G_1 - \Delta G_3$ ). Ethylene and 1-hexene show sensitivity to the nature of the fragment that is approximately equal to that of N<sub>2</sub> ( $\Delta G_1 - \Delta G_3 = 13.0$  kcal/mol and 13.5 kcal/mol respectively) while trimethyl phosphite is comparably sensitive ( $\Delta G_1 - \Delta G_3 = 11.4$  kcal/mol). Acetonitrile ( $\Delta G_1 - \Delta G_3 = 7.6$  kcal/mol) shows variability somewhere in between that of the amine ligands and those ligands that are apparently very sensitive to  $\pi$ -donating ability (e.g. CO, N<sub>2</sub>, olefins).

H<sub>2</sub> is calculated to add to the three (<sup>x</sup>PhDI)Ir fragments investigated to give a dihydrogen complex with a relatively short H-H distance varying only from 0.81 Å (**1**-H<sub>2</sub>) to 0.85 Å (**3**-H<sub>2</sub>). Thermodynamically, the binding energy shows moderate sensitivity to the nature of the iridium fragment ( $\Delta G_1 - \Delta G_3 = 9.5$  kcal/mol).


Generally speaking, our calculations predict that  $\pi$ -acceptor ligands such as CO bind much more strongly to the (<sup>x</sup>PhDI)Ir species with greater  $\pi$ -electron-donating ability ((<sup>x</sup>PhDI)IrH<sub>2</sub> and (<sup>x</sup>PhDI)Ir), with a sensitivity to  $\pi$ -electron-donating ability that is much greater than calculated for ligands such as amines; this is consistent with the most fundamental organometallic precepts and classical organometallic bonding descriptions. In such descriptions, CO is presented as a strong  $\pi$ -acceptor while amines are pure sigma donors or are even considered to be  $\pi$ -donors as well as  $\sigma$ -donors. From that perspective, however, it is noteworthy that the binding energy of N-ethylamine for example is even slightly greater toward the more electron-rich iridium fragments; the model of simple donation seems inconsistent with these data. Moreover, the variation found for phosphines is very nearly equal to that for amines; this seems inconsistent with the former being considered even modest  $\pi$ -acceptors.<sup>40, 41</sup> A full analysis of the origin of these effect is beyond the scope of this paper, but we plan to address this question in future work.

### 2.2.2. Ligand binding to other (pincer)Ir(I) fragments.


We have also calculated, for comparison with (<sup>x</sup>PhDI)Ir, the energies of binding of monodentate ligands to the (pincer)Ir(I) fragments (Phebox)Ir, (<sup>i</sup>PrPCP)Ir, and (<sup>t</sup>BuPCP)Ir (Table 2 and Figure 3). Of these, the Phebox pincer ligand is of course the most closely related of these to <sup>x</sup>PhDI, having the same diaminoaryl NCN motif. van Koten and co-workers have shown that nickel complexes of these pincer ligands have fairly similar redox properties as might be expected.<sup>26</sup>

The sterically undemanding ligands H<sub>2</sub>, N<sub>2</sub>, and CO bind more strongly to the (Phebox)Ir fragment than to (<sup>x</sup>PhDI)Ir, while acetonitrile binds slightly less strongly. This could suggest that (Phebox)Ir is a more  $\pi$ -electron-donating fragment but a slightly poorer  $\sigma$ -acceptor. Larger differences between Ir-L BDFEs of (Phebox)IrL versus (<sup>x</sup>PhDI)IrL are seen in the case of larger ligands, with the binding to (Phebox)Ir being weaker in all cases. The respective complexes are four-coordinate d<sup>8</sup>, therefore approximately square planar, and therefore this indicates specifically that the binding site trans to the

Ir-bound aryl carbon is more crowded in (Phebox)Ir than in (<sup>xy</sup>PhDI)Ir. Inspection of the calculated structures of the <sup>xy</sup>PhDI complexes supports this conclusion. The N-xylyl groups are oriented so that this trans coordination site of the <sup>xy</sup>PhDI complexes is significantly more open than that of (Phebox)Ir. For example, in the respective PMe<sub>3</sub> complexes there are numerous close contacts (*d*<sub>H-H</sub> < 2.4 Å) between the Phebox methyl groups and the PMe<sub>3</sub> ligand (Figure 4a), but no close contacts between coordinated PMe<sub>3</sub> and the <sup>xy</sup>PhDI ligand (Figure 4b).




**Figure 4.** DFT-calculated structures of (Phebox)Ir(PMe<sub>3</sub>) and (<sup>xy</sup>PhDI)Ir(PMe<sub>3</sub>) with closest contacts between PMe<sub>3</sub> and pincer ligand indicated (Å) (illustrating much more severe crowding in the Phebox complex).



**Figure 5.** DFT-calculated structures (space-filling models at 100% van der Waals radii) of (Phebox)Ir(PMe<sub>3</sub>) (a and b) and (<sup>xy</sup>PhDI)Ir(PMe<sub>3</sub>) (c and d). Viewed down Ir-C axis (a and c) and axis perpendicular to the plane of the pincer backbone (b and d) to illustrate the greater crowding of (Phebox)Ir at the site trans to the Ir-bond C atom, in contrast with the greater steric hindrance of (<sup>xy</sup>PhDI)Ir(PMe<sub>3</sub>) at the sites cis to the Ir-bond C atom. PMe<sub>3</sub> methyl groups in green and non-coordinating atoms of the pincer backbones in light grey.

Notably, although the trans site of the (<sup>Xy</sup>PhDI)Ir fragment is much more open than that of (Phebox)Ir, the sites cis to the Ir-bound carbon are fairly crowded in the case of (<sup>Xy</sup>PhDI)Ir while extremely open in the case of (Phebox)Ir. This is illustrated in Figure 5 with space filling models of the respective (pincer)Ir(PMe<sub>3</sub>) complexes and the Buried Volume maps of the (pincer)Ir fragments of Figure 6.<sup>42, 43</sup> Qualitatively at least, the distribution of steric bulk in the (<sup>Xy</sup>PhDI)Ir fragment resembles that of the (PCP)Ir fragments more closely than that of (Phebox)Ir, in that the (PCP)Ir fragments are also more crowded at the coordination sites cis to the Ir-bound carbon than at the trans sites. The symmetry of the (<sup>tBu</sup>PCP)Ir fragment (Figure 6c) makes the qualitative resemblance to (<sup>Xy</sup>PhDI)Ir (Figure 6a) more apparent than for the (<sup>iPr</sup>PCP)Ir (Figure 6d) fragment because the two i-propyl groups on each P atom are generally oriented in opposite directions (one tertiary C-H bond toward the Ir center and the other pointed away). Quantitatively however the Buried Volume calculations indicate that the (<sup>Xy</sup>PhDI)Ir fragment (%V<sub>BUR</sub> = 68.2%) is much more similar to (<sup>iPr</sup>PCP)Ir (67.3%) than to (<sup>tBu</sup>PCP)Ir (78.4%).



**Figure 6.** Topographical ligand steric maps and percent Buried Volume (%V<sub>BUR</sub>)<sup>42, 43</sup> of (a) (<sup>Xy</sup>PhDI)Ir (68.2%) (b) (Phebox)Ir (64.0%) (c) (<sup>tBu</sup>PCP)Ir (78.4%) (d) (<sup>iPr</sup>PCP)Ir (67.3%)

All ligands that we have studied are calculated to bind much more strongly to (<sup>iPr</sup>PCP)Ir than to (Phebox)Ir (Figure 3). Compared with (<sup>Xy</sup>PhDI)Ir, however, the amines bind less strongly to (<sup>iPr</sup>PCP)Ir while the phosphines bind more strongly to (<sup>iPr</sup>PCP)Ir. The small  $\pi$ -accepting ligands bind more strongly to (<sup>iPr</sup>PCP)Ir than to (<sup>Xy</sup>PhDI)Ir, with CO in particular binding 12 kcal/more strongly, while P(OMe)<sub>3</sub> also binds much more strongly to (<sup>iPr</sup>PCP)Ir (by 12.8 kcal/mol). P(OMe)<sub>3</sub> is known to be a good  $\pi$ -acceptor, but presumably much less  $\pi$ -accepting than CO. The greater energy of binding of P(OMe)<sub>3</sub> to (<sup>iPr</sup>PCP)Ir versus (<sup>Xy</sup>PhDI)Ir may therefore be a combination of greater  $\pi$ -donating ability of the (<sup>iPr</sup>PCP)Ir fragment


combined with a greater tendency to bind to the P-donating ligands generally, perhaps related to the “softness” of these ligands in contrast with the “hard” N-donors.

### 2.2.3. Kinetics of exchange with ethylene: thermodynamic implications

At room temperature, the  $^1\text{H}$ NMR spectrum of **1**- $\text{C}_2\text{H}_4$  in the presence of free  $\text{C}_2\text{H}_4$  indicates rapid exchange of free and bound ethylene. Decoalescence of the respective  $^1\text{H}$  NMR signals was observed at slightly reduced temperature, and the individual signals, attributable to free and bound  $\text{C}_2\text{H}_4$  respectively, were sharp at 255 K. Dynamic NMR allowed determination of the rate constants for exchange, over the temperature range 255 K – 298 K (Table 3), by simulation using the dNMR feature in the program Topspin.<sup>44</sup> An Eyring plot (Figure 7) of the rates thus obtained yielded activation parameters  $\Delta H^\ddagger = 25.0$  kcal/mol and  $\Delta S^\ddagger = 36.0 \pm 1.0$  cal/mol•deg. The positive activation entropy indicates that the reaction proceeds via dissociation of ethylene, as would be expected of an 18-electron ethylene complex. The activation enthalpy is very close to the calculated *thermodynamic* value of the enthalpy of dissociation,  $\Delta H^\circ = 23.7$  kcal/mol. Taken at face value, this would imply, as might also be expected, that addition of ethylene to the 16-electron dissociation product, **1**, has a near-zero enthalpic barrier of approximately  $\Delta H^\ddagger = 1.3$  kcal/mol. These results offer experimental support for the DFT-calculated thermodynamic values for ethylene (albeit approximate). Accordingly, they also support the validity of the absolute values of the DFT-calculated binding free energies of those ligands for which the *relative* (to ethylene) binding free energies are in agreement with experimental values.

**Table 3.** Rates and free energy for exchange between free and **1**-bound  $\text{C}_2\text{H}_4$  obtained by dynamic NMR and simulations


| T (K) | k ( $\text{s}^{-1}$ ) | $\Delta G^\ddagger$ (kcal/mol) |
|-------|-----------------------|--------------------------------|
| 298   | 190.1                 | 14.34                          |
| 285   | 29.6                  | 14.74                          |
| 275   | 5.66                  | 15.11                          |
| 265   | 0.877                 | 15.52                          |
| 255   | 0.138                 | 15.85                          |



**Figure 7.** Eyring plot for exchange between free and **1**-bound  $\text{C}_2\text{H}_4$

### 2.3. Cleavage of the C-C bond of acetonitrile

In an attempt to reduce **1-MeCN** to the corresponding dihydride or Ir(I) complex, the complex was treated with KO<sup>t</sup>Bu (3 equiv) in benzene under H<sub>2</sub> atmosphere and was left to stir at room temperature overnight under the hydrogen atmosphere. Benzene was evaporated and the solution was extracted with pentane to remove excess base. Crystals were grown under inert atmosphere at room temperature by diffusion of pentane into a concentrated benzene solution. Unexpectedly, scXRD revealed the product to be a binuclear bridging cyanide complex with the molecular structure as shown in Figure 8. Although hydrides were not located unambiguously by crystallography, the <sup>1</sup>H NMR spectrum of the crystals, after dissolving in benzene-*d*<sub>6</sub>, indicated the presence of two equivalent hydride ligands (-26.5 ppm) in accord with formulation as [(<sup>xy</sup>PhDI)Ir(Me)(H<sub>2</sub>NEt)](μ-CN)[(<sup>xy</sup>PhDI)IrH<sub>2</sub>] (**7**).




**Figure 8.** Molecular structure of **7**. H atoms except hydrides and those of the methyl and ethylamine ligands omitted for clarity. Carbon atoms of methyl and ethylamine ligands and bridging cyanide shown in green for emphasis.

The net reaction to give **7** thus involves hydrogenation of one molecule of CH<sub>3</sub>CN (per molecule of binuclear product) and cleavage of the C-C bond of a second molecule of CH<sub>3</sub>CN. C-C bond cleavage generally, and cleavage of alkyl cyanides in particular, is a reaction of much interest.<sup>45-52</sup> The examples that are perhaps most closely related to the present work have been reported by Garcia and Jones.<sup>53-58</sup>


We investigated the reaction of the (<sup>xy</sup>PhDI)Ir fragment with CH<sub>3</sub>CN computationally and located a transition state for direct oxidative addition of the acetonitrile C-C bond. However the barrier of this reaction was calculated to be too high, at 33.1 kcal/mol, to account for the formation of **7** at room temperature. Lewis acids, however, have been reported to accelerate C-CN bond cleavage<sup>54, 59</sup> or the reverse, C-CN reductive elimination<sup>60, 61</sup>. (Notably, Garcia and Jones have reported an example where

Lewis acid binding actually *inhibited* C-C cleavage.<sup>53</sup>) Therefore, given that the nitrile group of **7** is bridging, we considered that the second (<sup>x</sup>PhDI)Ir unit played a role in promoting the C-C cleavage reaction. In accord with this hypothesis, the free energy barrier to C-C cleavage was calculated to be only 10.4 kcal/mol when the acetonitrile N atom was coordinated to a (<sup>Me</sup>PhDI)IrH<sub>2</sub> fragment (the N-coordinated (PhDI)Ir moiety was truncated, with xyl groups replaced by methyl groups for computational simplicity; **3**\*-H<sub>2</sub>; Figure 9). Although the free energy of this binding is positive,  $\Delta G^\circ = 7.7$  kcal/mol, due to a large unfavorable entropy term ( $T\Delta S = -16.7$  kcal/mol at 298 K), the overall calculated barrier of C-C cleavage,  $\Delta G^\ddagger = 18.1$  kcal/mol, is still dramatically lowered by assistance from the second (PhDI)Ir center.



**Figure 9.** Calculated energy profile (free energies in kcal/mol) for oxidative addition of the C-C bond of acetonitrile by **1**, assisted by **1**\*-H<sub>2</sub> and unassisted. (Selected distances in Å)

The calculations indicate, remarkably, that in the transition states for C-CN cleavage, both metal assisted and non-assisted, the Ir-CN bond is significantly shorter in the transition states (1.94 Å and 1.96 Å for Ir-bridged and non-bridged respectively) than in the products in which the Ir-CN bond is fully formed (2.10 Å and 2.07 Å) (Figure 10). Note, however, that as would generally be expected for an oxidative addition reaction, the Ir-CH<sub>3</sub> distance is significantly shorter in the products than in the transition states. Acetonitrile C-C oxidative addition by (dippe)Ni(0)<sup>55</sup> and Cp\*(PMe<sub>3</sub>)Rh(I)<sup>56</sup> (not assisted by a second metal) has been investigated computationally in detail by Jones. While our results are generally in agreement with those, in the Rh(I) case the M-CN bond in the TS (2.00 Å) was slightly longer<sup>56</sup> than in the C-C cleavage product (1.97 Å). In the case of Ni(0), however, as in the present systems the M-C bond was shorter in the TS (1.82 Å) than in the product (1.88 Å) although the difference was not as pronounced.



**Figure 10.** Calculated transition state (a) and product (b) of oxidative addition of the C-C bond of acetonitrile by **3**, assisted by **3**\*-H<sub>2</sub>. Transition state (c) and product (d) of “unassisted” C-CN addition by **3**. N-mesityl groups of **3**, and most H atoms, omitted for clarity

Alkyl cyanide elimination/C-C bond formation has been compared with alkyl migration to CO (i.e. CO insertion into M-alkyl bonds).<sup>61-63</sup> This perspective might help to rationalize this unusual example of a TS for C-C cleavage with a M-C distance shorter than that of the product with the fully formed M-C bond. For example, we have computationally studied alkyl migration to CO of Mn(CO)<sub>5</sub>(CH<sub>2</sub>Ar), and found that the M-CO bond in the migration transition state is shorter (1.82 Å) than in the carbonyl reactant (1.86 Å) or in the acyl product of migration (1.88 Å).<sup>64</sup> Note also that in the transition states calculated in this work, as well as in the systems studied by Jones,<sup>55, 56</sup> there is a significant agostic interaction with the acetonitrile methyl C-H bond. In our studies of alkyl migration to CO (or alkyl migration from acyl ligand to metal) it was shown that formation of an analogous agostic interaction played a significant role in the energy of the transition state<sup>64</sup>; these shared feature would seem to further support the proposed relationship between alkyl migration and alkyl-CN cleavage/elimination.

### 3. Conclusions

Iridium dibromide complexes of the phenyldiimine ligand 2,6-bis(1-((2,6-dimethylphenyl)imino)ethyl)phenyl have been synthesized, and the relative Ir-L BDFEs have been experimentally determined for a wide range of corresponding adducts of ligands L. An estimate of the absolute enthalpy of Ir-L binding has been obtained from dynamic NMR measurements. The results of DFT calculations are in very good agreement with the relative and absolute experimental values.

A computational study has been conducted, first comparing (<sup>Xy</sup>PhDI)Br<sub>2</sub>Ir-L BDFEs with Ir-L BDFEs of adducts *trans*-(<sup>Xy</sup>PhDI)H<sub>2</sub>IrL and (<sup>Xy</sup>PhDI)IrL. The Ir-L BDFEs are all greater for the dihydride than for the dibromide, with the difference being much more pronounced for  $\pi$ -accepting ligands. BDFEs of  $\pi$ -accepting ligands are even markedly greater still for the Ir(I) complexes (<sup>Xy</sup>PhDI)IrL versus *trans*-(<sup>Xy</sup>PhDI)H<sub>2</sub>IrL, while there is little difference for ligands that are less  $\pi$ -accepting.

We also compared (<sup>Xy</sup>PhDI)Ir-L BDFEs with BDFEs of (pincer)Ir-L complexes for pincers Phebox, <sup>i</sup>PrPCP, and <sup>t</sup>BuPCP. The (Phebox)Ir fragment forms slightly stronger bonds than (<sup>Xy</sup>PhDI)Ir with the smallest  $\pi$ -accepting ligands, but (Phebox)Ir-L bonds are significantly weaker with bulky ligands, particularly those that are not significantly  $\pi$ -accepting. Although Buried Volume calculations indicate that the (Phebox)Ir fragment has overall greater “unburied” volume than (<sup>Xy</sup>PhDI)Ir, the coordination site occupied by L in the four-coordinate d<sup>8</sup> (pincer)IrL complexes (i.e. the site *trans* to Ir-bound aryl C) is significantly more crowded in (Phebox)Ir. The motif of (<sup>Xy</sup>PhDI)Ir, with greater steric crowding at the *cis* sites and a more open site *trans* to the aryl C is also found for both (<sup>i</sup>PrPCP)Ir and (<sup>t</sup>BuPCP)Ir pincer ligands. The (<sup>R</sup>PCP)Ir fragments appear to be more  $\pi$ -donating than the Phebox or PhDI fragments, as illustrated by stronger binding to CO or N<sub>2</sub>, but with respect to sterically demanding ligands, the BDFEs of (<sup>Xy</sup>PhDI)IrL are somewhere between those of the very crowded (<sup>t</sup>BuPCP)Ir and the much less crowded (<sup>i</sup>PrPCP)Ir.

An attempt to convert (<sup>Xy</sup>PhDI)IrBr<sub>2</sub>(MeCN) to a hydride or possibly an Ir(I) complex yielded an unexpected dinuclear cyano-bridged complex with N-ethylamine (presumably the product of hydrogenation of one mol acetonitrile) coordinated to one metal center, and a methyl ligand on the cyanide carbon-bound iridium center, indicating the occurrence of C-CN bond cleavage at the latter Ir center. DFT calculations indicate that the C-CN bond cleavage occurs at one Ir center with strong assistance by coordination of the CN nitrogen to the other center. At the metal center effecting the C-CN addition the transition state is calculated to strongly resemble a transition state for alkyl migration to CO or the microscopic reverse, C-C bond cleavage of an acyl ligand. Similarities include a M-CX (X = N or O) bond distance in the TS that is shorter than that in the M-CX C-C cleavage product, and a strong agostic interaction with a C-H bond of the alkyl group being cleaved from CX.

The inference of C-C cleavage by the (<sup>Xy</sup>PhDI)Ir(I) fragment may suggest promising activity, related but distinct from the chemistry of (<sup>R</sup>PCP)Ir(I) fragments, if such a species can be generated in the absence of acetonitrile.

## Supplementary Data

Electronic Supplementary Information available:

Complete experimental details and synthetic procedures, NMR data, computational details and data, computed energies and thermodynamic quantities (**PDF**)

CCDC 2268483-226848, 2280217, 2303676

Supplementary data to this article can be found online at <https://doi.org/xxxxx>

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgments

This work was supported by the U. S. Department of Energy Office of Science (DE-SC0020139).

## AUTHOR INFORMATION

### Corresponding Author

**Alan S. Goldman** – *Department of Chemistry and Chemical Biology, Rutgers, The State University of New Jersey, New Brunswick, New Jersey 08854, United States; orcid.org/0000-0002-2774-710X; Email: alan.goldman@rutgers.edu*

### Authors

**Soumyadipa Das** – *Department of Chemistry and Chemical Biology, Rutgers, The State University of New Jersey, New Brunswick, New Jersey 08854, United States; orcid.org/0009-0008-4667-930X*

**Souvik Mandal** – *Department of Chemistry and Chemical Biology, Rutgers, The State University of New Jersey, New Brunswick, New Jersey 08854, United States; orcid.org/0000-0003-2402-6662*

**Santanu Malakar** – *Department of Chemistry and Chemical Biology, Rutgers, The State University of New Jersey, New Brunswick, New Jersey 08854, United States; orcid.org/0000-0002-3838-5095*

**Thomas J. Emge** – *Department of Chemistry and Chemical Biology, Rutgers, The State University of New Jersey, New Brunswick, New Jersey 08854, United States; orcid.org/0000-0003-4685-8419*

## ■ REFERENCES

- [1] Y. Wang, Z. Huang, G. Liu and Z. Huang, A New Paradigm in Pincer Iridium Chemistry: PCN Complexes for (De)Hydrogenation Catalysis and Beyond, *Acc. Chem. Res.* 55 (2022) 2148-2161. <https://doi.org/10.1021/acs.accounts.2c00311>.
- [2] B. M. Gordon, N. Lease, T. J. Emge, F. Hasanayn and A. S. Goldman, Reactivity of Iridium Complexes of a Triphosphorus-Pincer Ligand Based on a Secondary Phosphine. Catalytic Alkane Dehydrogenation and the Origin of Extremely High Activity, *J. Am. Chem. Soc.* 144 (9) (2022) 4133-4146. <https://doi.org/10.1021/jacs.1c13309>.
- [3] A. D. R. Shada, A. J. M. Miller, T. J. Emge and A. S. Goldman, Catalytic Dehydrogenation of Alkanes by PCP-Pincer Iridium Complexes Using Proton and Electron Acceptors, *ACS Catal.* 11 (2021) 3009-3016. <https://doi.org/10.1021/acscatal.0c05160>.
- [4] K. Das and A. Kumar, Alkane dehydrogenation reactions catalyzed by pincer-metal complexes, *Adv. Organomet. Chem.* 72 (2019) 1-57. <https://doi.org/10.1016/bs.adomc.2019.02.0041>.
- [5] X. Tang, X. Jia and Z. Huang, Thermal, Catalytic Conversion of Alkanes to Linear Aldehydes and Linear Amines, *J. Am. Chem. Soc.* 140 (11) (2018) 4157-4163. <https://doi.org/10.1021/jacs.8b01526>.
- [6] A. Kumar, T. M. Bhatti and A. S. Goldman, Dehydrogenation of Alkanes and Aliphatic Groups by Pincer-Ligated Metal Complexes, *Chem. Rev.* 117 (19) (2017) 12357-12384. <http://dx.doi.org/10.1021/acs.chemrev.7b00247>.
- [7] X. Jia and Z. Huang, Conversion of alkanes to linear alkylsilanes using an iridium-iron-catalyzed tandem dehydrogenation-isomerization-hydrosilylation, *Nat. Chem.* 8 (2) (2016) 157-161. <https://doi.org/10.1038/nchem.2417>.
- [8] W. Yao, X. Jia, X. Leng, A. S. Goldman, M. Brookhart and Z. Huang, Catalytic Alkane Transfer-Dehydrogenation by PSCOP Iridium Pincer Complexes, *Polyhedron* 116 (2016) 12-19. <https://doi.org/10.1016/j.poly.2016.02.044>.
- [9] M. C. Haibach, S. Kundu, M. Brookhart and A. S. Goldman, Alkane Metathesis by Tandem Alkane-Dehydrogenation-Olefin-Metathesis Catalysis and Related Chemistry, *Acc. Chem. Res.* 45 (2012) 947-958. <http://dx.doi.org/10.1021/ar3000713>.
- [10] J. Choi and A. S. Goldman, Ir-Catalyzed Functionalization of C-H Bonds, *Top. Organomet. Chem.* 34 (2011) 139-167. [http://dx.doi.org/10.1007/978-3-642-15334-1\\_6](http://dx.doi.org/10.1007/978-3-642-15334-1_6).
- [11] A. Ray, K. Zhu, Y. V. Kissin, A. E. Cherian, G. W. Coates and A. S. Goldman, Dehydrogenation of aliphatic polyolefins catalyzed by pincer-ligated iridium complexes, *Chem. Commun.* (2005) 3388-3390. <https://doi.org/10.1039/B502120K>.
- [12] J. Choi, Y. Choliy, X. Zhang, T. J. Emge, K. Krogh-Jespersen and A. S. Goldman, Cleavage of  $sp^3$  C-O Bonds via Oxidative Addition of C-H Bonds, *J. Am. Chem. Soc.* 131 (43) (2009) 15627-15629. <http://dx.doi.org/10.1021/ja906930u>.
- [13] M. C. Haibach, C. Guan, D. Y. Wang, B. Li, N. Lease, A. M. Steffens, K. Krogh-Jespersen and A. S. Goldman, Olefin Hydroaryloxylation Catalyzed by Pincer-Iridium Complexes, *J. Am. Chem. Soc.* 135 (40) (2013) 15062-15070. <http://dx.doi.org/10.1021/ja404566v>.
- [14] M. Kanzelberger, X. Zhang, T. J. Emge, A. S. Goldman, J. Zhao, C. Incarvito and J. F. Hartwig, Distinct Thermodynamics for the Formation and Cleavage of N-H Bonds in Aniline and Ammonia. Directly-Observed Reductive Elimination of Ammonia from an Isolated Amido Hydride Complex, *J. Am. Chem. Soc.* 125 (45) (2003) 13644-13645. <https://doi.org/10.1021/ja037363u>.
- [15] J. Zhao, A. S. Goldman and J. F. Hartwig, Oxidative Addition of Ammonia to Form a Stable Monomeric Amido Hydride Complex, *Science* 307 (5712) (2005) 1080-1082. <http://www.sciencemag.org/cgi/content/full/307/5712/1080>.
- [16] D. Y. Wang, Y. Choliy, M. C. Haibach, J. F. Hartwig, K. Krogh-Jespersen and A. S. Goldman, Assessment of the Electronic Factors Determining the Thermodynamics of "Oxidative Addition" of C-H and N-H Bonds to Ir(I) Complexes, *J. Am. Chem. Soc.* 138 (1) (2016) 149-163. <http://dx.doi.org/10.1021/jacs.5b09522>.
- [17] J. Choi, D. Y. Wang, S. Kundu, Y. Choliy, T. J. Emge, K. Krogh-Jespersen and A. S. Goldman, Net Oxidative Addition of  $C(sp^3)$ -F Bonds to Iridium via Initial C-H Bond Activation, *Science* 332 (2011) 1545-1548. <http://www.sciencemag.org/content/332/6037/1545>.

- [18] J.-i. Ito, T. Shiomi and H. Nishiyama, Efficient preparation of new rhodium- and iridium-[bis(oxazolynyl)-3,5-dimethylphenyl] complexes by C-H bond activation: applications in asymmetric synthesis, *Adv. Synth. Catal.* 348 (2006) 1235-1240. <https://doi.org/10.1002/adsc.200606049>.
- [19] J.-i. Ito, T. Kaneda and H. Nishiyama, Intermolecular C-H Bond Activation of Alkanes and Arenes by NCN Pincer Iridium(III) Acetate Complexes Containing Bis(oxazolynyl)phenyl Ligands, *Organometallics* 31 (2012) 4442-4449. <http://dx.doi.org/10.1021/om3002137>.
- [20] C. P. Owens, A. Varela-Alvarez, V. Boyarskikh, D. G. Musaev, H. M. L. Davies and S. B. Blakey, Iridium(III)-bis(oxazolynyl)phenyl catalysts for enantioselective C-H functionalization, *Chem. Sci.* 4 (6) (2013) 2590-2596. <http://dx.doi.org/10.1039/C3SC50886B>.
- [21] K. E. Allen, D. M. Heinekey, A. S. Goldman and K. I. Goldberg, Alkane Dehydrogenation by C-H Activation at Iridium(III), *Organometallics* 32 (6) (2013) 1579-1582. <http://dx.doi.org/10.1021/om301267c>.
- [22] D. R. Pahl, K. E. Allen, K. I. Goldberg and T. R. Cundari, Understanding the Effect of Ancillary Ligands on Concerted Metalation-Deprotonation by (dmPhebox)Ir(OAc)<sub>2</sub>(H<sub>2</sub>O) Complexes: A DFT Study, *Organometallics* 33 (22) (2014) 6413-6419. <http://dx.doi.org/10.1021/om500752m>.
- [23] M. Zhou, S. I. Johnson, Y. Gao, T. J. Emge, R. J. Nielsen, W. A. Goddard and A. S. Goldman, Activation and Oxidation of Mesitylene C-H Bonds by (Phebox)Iridium(III) Complexes, *Organometallics* 34 (12) (2015) 2879-2888. <http://pubs.acs.org/doi/abs/10.1021/acs.organomet.5b00200>.
- [24] N. Li, W.-J. Zhu, J.-J. Huang, X.-Q. Hao, J.-F. Gong and M.-P. Song, Chiral NCN Pincer Iridium(III) Complexes with Bis(imidazolynyl)phenyl Ligands: Synthesis and Application in Enantioselective C-H Functionalization of Indoles with  $\alpha$ -Aryl- $\alpha$ -diazoacetates, *Organometallics* 39 (12) (2020) 2222-2234. <https://doi.org/10.1021/acs.organomet.0c00174>.
- [25] X. Zhou, S. Malakar, T. Dugan, K. Wang, A. Sattler, D. O. Marler, T. J. Emge, K. Krogh-Jespersen and A. S. Goldman, Alkane Dehydrogenation Catalyzed by a Fluorinated Phebox Iridium Complex, *ACS Catal.* 11 (22) (2021) 14194-14209. <https://doi.org/10.1021/acscatal.1c03562>.
- [26] M. Stol, D. J. M. Snelders, M. D. Godbole, R. W. A. Havenith, D. Haddleton, G. Clarkson, M. Lutz, A. L. Spek, G. P. M. Van Klink and G. Van Koten, 2,6-Bis(oxazolynyl)phenylnickel(II) Bromide and 2,6-Bis(ketimine)phenylnickel(II) Bromide: Synthesis, Structural Features, and Redox Properties, *Organometallics* 26 (16) (2007) 3985-3994. <https://doi.org/10.1021/om061055y>.
- [27] S. H. Oakley, M. P. Coogan and R. J. Arthur, Synthesis of Bis(imino)aryl Iridium Pincer Complexes and Demonstration of Catalytic Hydrogen-Transfer Activity, *Organometallics* 26 (9) (2007) 2285-2290. <https://doi.org/10.1021/om070059f>.
- [28] G. P. Rosini, F. Liu, K. Krogh-Jespersen, A. S. Goldman, C. Li and S. P. Nolan, Thermodynamics of Addition of H<sub>2</sub>, CO, N<sub>2</sub>, and C-H Bonds to M(P<sup>i</sup>Pr<sub>3</sub>)<sub>2</sub>Cl (M = Ir, Rh). An Unprecedented Metal Carbonyl Bond Strength, *J. Am. Chem. Soc.* 120 (36) (1998) 9256-9266. <https://doi.org/10.1021/ja974100p>.
- [29] K. Wang, G. P. Rosini, S. P. Nolan and A. S. Goldman, Thermodynamics of Addition of CO, Isocyanide, and H<sub>2</sub> to Rh(PR<sub>3</sub>)<sub>2</sub>Cl, *J. Am. Chem. Soc.* 117 (18) (1995) 5082-5088. <https://doi.org/10.1021/ja00123a010>.
- [30] C. Y. Zhao and D. G. Truhlar, The M06 Suite of Density Functionals for Main Group Thermochemistry, Kinetics, Noncovalent Interactions, Excited States, and Transition Elements: Two New Functionals and Systematic Testing of Four M06 Functionals and Twelve Other Functionals, *Theo. Chem. Acc.* 120 (2008) 215-241. <https://doi.org/10.1007/s00214-007-0310-x>.
- [31] R. Ditchfield, W. J. Hehre and J. A. Pople, Self-consistent molecular-orbital methods. IX. Extended Gaussian-type basis for molecular-orbital studies of organic molecules, *J. Chem. Phys.* 54 (2) (1971) 724-728. <https://doi.org/10.1063/1.1674902>.
- [32] W. J. Hehre, R. Ditchfield and J. A. Pople, Self-Consistent Molecular Orbital Methods. XII. Further Extensions of Gaussian-Type Basis Sets for Use in Molecular Orbital Studies of Organic Molecules, *J. Chem. Phys.* 56 (5) (1972) 2257-2261. <https://doi.org/10.1063/1.1677527>.
- [33] P. C. Hariharan and J. A. Pople, The influence of polarization functions on molecular orbital hydrogenation energies, *Theor. Chim. Acta* 28 (3) (1973) 213-222. <https://doi.org/10.1007/BF00533485>.
- [34] P. C. Hariharan and J. A. Pople, Accuracy of AH<sub>n</sub> equilibrium geometries by single determinant molecular orbital theory, *Mol. Phys.* 27 (1) (1974) 209-214. <https://doi.org/10.1080/00268977400100171>.
- [35] M. M. Francl, W. J. Pietro, W. J. Hehre, J. S. Binkley, M. S. Gordon, D. J. DeFrees and J. A. Pople, Self-consistent molecular orbital methods XXIII A polarization-type basis set for second-row elements, *J. Chem. Phys.* 77 (7) (1982) 3654-3665. <https://doi.org/10.1063/1.444267>.

- [36] D. Andrae, U. Haeussermann, M. Dolg, H. Stoll and H. Preuss, Energy-adjusted ab initio pseudopotentials for the second and third row transition elements, *Theor. Chim. Acta* 77 (2) (1990) 123-141. <https://doi.org/10.1007/BF01114537>.
- [37] This root-mean-square value was not taken from the deviations between experiment and calculated values shown in Table 1, which are specifically relative to ethylene and therefore any such error determination would put undue weight on the values calculated and measure for ethylene in particular. Instead relative values were extrapolated for all L/L' couples and the root-mean-square calculation was based on those values.
- [38] The calculated global minimum for (<sup>x</sup>PhDI)IrH<sub>2</sub> is a dihydrogen complex (with the H<sub>2</sub> ligand coordinated trans to the Ir-bound carbon, oriented in the plane of the pincer backbone). For purposes of comparison to the addition of L to (<sup>x</sup>PhDI)IrBr<sub>2</sub>, in which the global minimum structure has mutually trans Br ligands (both cis to the Ir-bound carbon) for (<sup>x</sup>PhDI)IrH<sub>2</sub>L Ir-L BDEs we calculated addition to an analogous structure of trans-(<sup>x</sup>PhDI)IrH<sub>2</sub>, which is actually a local energy minimum, 8.4 kcal/mol in free energy above the global minimum (<sup>x</sup>PhDI)Ir(H<sub>2</sub>).
- [39] M. D. Fryzuk, Side-on End-on Bound Dinitrogen: An Activated Bonding Mode That Facilitates Functionalizing Molecular Nitrogen, *Acc. Chem. Res.* 42 (1) (2009) 127-133. <https://doi.org/10.1021/ar800061g>.
- [40] R. H. Crabtree, in *The Organometallic Chemistry of the Transition Metals*, John Wiley & Sons, Hoboken, NJ, 7th edn., 2019, p. 100.
- [41] J. F. Hartwig, in *Organotransition Metal Chemistry*, University Science Books, Sausalito, CA, 2010, p. 36.
- [42] L. Falivene, R. Credendino, A. Poater, A. Petta, L. Serra, R. Oliva, V. Scarano and L. Cavallo, SambVca 2. A Web Tool for Analyzing Catalytic Pockets with Topographic Steric Maps, *Organometallics* 35 (13) (2016) 2286-2293. <https://doi.org/10.1021/acs.organomet.6b00371>.
- [43] L. Falivene, Z. Cao, A. Petta, L. Serra, A. Poater, R. Oliva, V. Scarano and L. Cavallo, Towards the online computer-aided design of catalytic pockets, *Nat. Chem.* 11 (10) (2019) 872-879. <https://doi.org/10.1038/s41557-019-0319-5>.
- [44] Rohonczy, J; Loránd, E. DNMR Lineshape Analysis Software Manual Version 1.1 Rev. 071103, Bruker BioSpin GmbH, TopSpin 2.1, 2007.
- [45] Y. Nakao, Metal-mediated C–CN Bond Activation in Organic Synthesis, *Chem. Rev.* 121 (1) (2021) 327-344. <https://doi.org/10.1021/acs.chemrev.0c00301>.
- [46] D. Churchill, J. H. Shin, T. Hascall, J. M. Hahn, B. M. Bridgewater and G. Parkin, The Ansa Effect in Permethylmolybdenocene Chemistry: A [Me<sub>2</sub>Si] Ansa Bridge Promotes Intermolecular C–H and C–C Bond Activation, *Organometallics* 18 (13) (1999) 2403-2406. <https://doi.org/10.1021/om990195n>.
- [47] F. L. Taw, P. S. White, R. G. Bergman and M. Brookhart, Carbon–Carbon Bond Activation of R–CN (R = Me, Ar, iPr, tBu) Using a Cationic Rh(III) Complex, *J. Am. Chem. Soc.* 124 (16) (2002) 4192-4193. <https://doi.org/10.1021/ja0255094>.
- [48] F. L. Taw, A. H. Mueller, R. G. Bergman and M. Brookhart, A Mechanistic Investigation of the Carbon–Carbon Bond Cleavage of Aryl and Alkyl Cyanides Using a Cationic Rh(III) Silyl Complex, *J. Am. Chem. Soc.* 125 (32) (2003) 9808-9813. <https://doi.org/10.1021/ja034468o>.
- [49] H. Nakazawa, T. Kawasaki, K. Miyoshi, C. H. Suresh and N. Koga, C–C Bond Cleavage of Acetonitrile by a Carbonyl Iron Complex with a Silyl Ligand, *Organometallics* 23 (1) (2004) 117-126. <https://doi.org/10.1021/om0208319>.
- [50] T. Lu, X. Zhuang, Y. Li and S. Chen, C–C Bond Cleavage of Acetonitrile by a Dinuclear Copper(II) Cryptate, *J. Am. Chem. Soc.* 126 (15) (2004) 4760-4761. <https://doi.org/10.1021/ja031874z>.
- [51] H. Nakazawa, K. Kamata and M. Itazaki, Catalytic C–C bond cleavage and C–Si bond formation in the reaction of RCN with Et<sub>3</sub>SiH promoted by an iron complex, *Chem. Commun.* (31) (2005) 4004-4006. <http://dx.doi.org/10.1039/B504131G>.
- [52] F. Xu, W. Huang and X.-Z. You, Novel cyano-bridged mixed-valent copper complexes formed by completely in situ synthetic method via the cleavage of C–C bond in acetonitrile, *Dalton Trans.* 39 (44) (2010) 10652-10658. <http://dx.doi.org/10.1039/C0DT00345J>.
- [53] J. J. Garcia, A. Arevalo, N. M. Brunkan and W. D. Jones, Cleavage of Carbon–Carbon Bonds in Alkyl Cyanides Using Nickel(0), *Organometallics* 23 (16) (2004) 3997-4002. <https://doi.org/10.1021/om049700t>.

- [54] N. M. Brunkan, D. M. Brestensky and W. D. Jones, Kinetics, Thermodynamics, and Effect of BPh<sub>3</sub> on Competitive C-C and C-H Bond Activation Reactions in the Interconversion of Allyl Cyanide by [Ni(dippe)], *J. Am. Chem. Soc.* 126 (11) (2004) 3627-3641. <https://doi.org/10.1021/ja037002e>.
- [55] T. A. Atesin, T. Li, S. Lachaize, W. W. Brennessel, J. J. Garcia and W. D. Jones, Experimental and Theoretical Examination of C-CN and C-H Bond Activations of Acetonitrile Using Zerovalent Nickel, *J. Am. Chem. Soc.* 129 (24) (2007) 7562-7569. <https://doi.org/10.1021/ja0707153>.
- [56] M. E. Evans, T. Li and W. D. Jones, C-H vs. C-C Bond activation of acetonitrile and benzonitrile via oxidative addition: rhodium vs. nickel and Cp\* vs. Tp' (Tp' = hydrotris(3,5-dimethylpyrazol-1-yl)borate, Cp\* = h<sup>5</sup>-pentamethylcyclopentadienyl), *J. Am. Chem. Soc.* 132 (2010) 16278-16284. <https://doi.org/10.1021/ja107927b>.
- [57] T. Li, J. J. Garcia, W. W. Brennessel and W. D. Jones, C-CN Bond activation of aromatic nitriles and fluxionality of the h<sup>2</sup>-arene intermediates: experimental and theoretical investigations, *Organometallics* 29 (2010) 2430-2445. <https://doi.org/10.1021/om100001m>.
- [58] T. Tanabe, M. E. Evans, W. W. Brennessel and W. D. Jones, C-H and C-CN Bond Activation of Acetonitrile and Succinonitrile by [Tp'Rh(PR<sub>3</sub>)], *Organometallics* 30 (4) (2011) 834-843. <https://doi.org/10.1021/om101002m>.
- [59] L. Munjanja, C. Torres-López, W. W. Brennessel and W. D. Jones, C-CN Bond Cleavage Using Palladium Supported by a Dippe Ligand, *Organometallics* 35 (11) (2016) 2010-2013. <https://doi.org/10.1021/acs.organomet.6b00304>.
- [60] C. A. Tolman, W. C. Seidel, J. D. Druliner and P. J. Domaille, Catalytic hydrocyanation of olefins by nickel(0) phosphite complexes - effects of Lewis acids, *Organometallics* 3 (1) (1984) 33-38. <https://doi.org/10.1021/om00079a008>.
- [61] J. Huang, C. M. Haar, S. P. Nolan, J. E. Marcone and K. G. Moloy, Lewis Acids Accelerate Reductive Elimination of RCN from P<sub>2</sub>Pd(R)(CN), *Organometallics* 18 (3) (1999) 297-299. <https://doi.org/10.1021/om980897x>.
- [62] J. E. Marcone and K. G. Moloy, Kinetic Study of Reductive Elimination from the Complexes (Diphosphine)Pd(R)(CN), *J. Am. Chem. Soc.* 120 (33) (1998) 8527-8528. <https://doi.org/10.1021/ja980762i>.
- [63] J. Ledford, C. S. Shultz, D. P. Gates, P. S. White, J. M. DeSimone and M. Brookhart, Bond Angle Effects on the Migratory Insertion of Ethylene and Carbon Monoxide into Palladium(II)-Methyl Bonds in Complexes Bearing Bidentate Phosphine Ligands, *Organometallics* 20 (25) (2001) 5266-5276. <https://doi.org/10.1021/om010489k>.
- [64] T. Zhou, S. Malakar, S. L. Webb, K. Krogh-Jespersen and A. S. Goldman, Polar molecules catalyze CO insertion into metal-alkyl bonds through the displacement of an agostic C-H bond, *Proc. Natl. Acad. Sci.* 116 (9) (2019) 3419-3424. <https://doi.org/10.1073/pnas.1816339116>.