


Southern Giant Petrels *Macronectes giganteus* as indicators of ocean surface currents

Megan A Cimino & Carlos Moffat



Cimino MA, Moffat C 2024. Southern Giant Petrels *Macronectes giganteus* as indicators of ocean surface currents. Biodiversity Observations 14: 71–74.

10 June 2024

DOI: 10.15641/bo.1503

ORNITHOLOGY

Southern Giant Petrels *Macronectes giganteus* as indicators of ocean surface currents

Megan A Cimino^{1*} & Carlos Moffat²

¹*Institute of Marine Science, University of California Santa Cruz,
1156 High St, Santa Cruz, CA 95064, USA.*

²*School of Marine Science and Policy, University of Delaware,
210 South College Ave., Newark, DE 19716, USA.*

**mecimino@ucsc.edu*

Abstract

A Southern Giant Petrel *Macronectes giganteus* was satellite tracked during a long foraging trip. While at rest on the sea surface, the giant petrel drifted in a counterclockwise corkscrew pattern that is characteristic of an inertial oscillation in the Southern Ocean. This note demonstrates that tracking data from resting seabirds can be used as passive drifters to estimate ocean surface currents in a notoriously stormy environment where data near the air-sea interface are difficult to obtain.

Introduction

Seabirds are well-known ocean sentinels that can provide unique information on atmospheric, ocean, and food web processes, especially species capable of traveling long distances (Hazen et al. 2019, Ellis-Soto et al. 2023). An exemplary species is the Southern Giant Petrel *Macronectes giganteus*, which is a long-lived, widely distributed predator in the Southern Ocean. Giant petrels can have long foraging trips during both the breeding and non-breeding seasons, where they can be at sea for days to weeks and cover distances of 100s to 1000s of kilometres (e.g., Finger et al. 2023). Therefore, their foraging behaviour across multiple spatio-temporal scales can provide information on their surrounding environment in intriguing ways as shown here.

Observation

As part of the Palmer Long-Term Ecological Research (LTER) program, Southern Giant Petrels nesting on small islands near Anvers Island (64.076°W, 64.769°S), West Antarctic Peninsula, were satellite tracked using Axy-Trek Marine devices during the incubation phase in 2022. The trackers record GPS coordinates (longitude, latitude, and time) every 15 minutes, which can be used to estimate statistics such as distance traveled and speed. One male giant petrel traveled to Peter I Island in the Bellingshausen Sea (~1200 km away from its nest) before returning to its nest and finishing its ~3200 km journey in about 14 days (8-22 December) (Figure 1).

When we examined the petrel's trajectory, we found several instances of a counterclockwise corkscrew pattern (Figure 2A) that coincided with low speeds and displacements, indicating the bird was sitting on the sea surface, presumably foraging or resting, and thus tracking the movement of the surface ocean water. Counterclockwise surface circulation can be evident in the surface ocean after strong winds suddenly stop. The currents that are forced by the wind, and that continue after the wind stops, are influenced by the earth's rotation and friction, veering to the left and forming a characteristic counterclockwise corkscrew pattern called an inertial oscillation. In the northern hemisphere, this motion would be clockwise.




Figure 1: A Southern Giant Petrel foraging track coloured by speed. The track begins and ends near Anvers Island. The direction of travel is indicated by the blue arrows. The region within the red box is shown in Figure 2 .

If what we observed was indeed an inertial oscillation, theory predicts the time for the trajectory to complete a full circle to be $T = 2\pi/f$ (the inertial period), where $\pi=3.1416$ and f is the Coriolis parameter, which depends on the latitude. At the latitude of 70°S where the inertial oscillations appeared to have been tracked by the petrel, T is 12.7 hours. This expected value is broadly consistent with the observations, which showed the petrel finishing a complete circle every 12 hours or so several times (Figure 2AB). The radius of the circle scales like V/f , where V is the velocity of the surface current. Typical speeds are of the order of 0.15 m/s, which should result in “corkscrews” with a radius of 1 km or so, which is again broadly consistent with the observations (Figure 2B).




Figure 2: A) Zoomed-in giant petrel track within the red box in Fig. 1. Arrows represent the direction of drift. With the addition of flight at higher speeds, the bird travelled ~34 km north. The start and end points are marked, 13 December at 00:00 and 15 December at 13:00. B) Zoomed-in counterclockwise “corkscrew” located within the red box in (A) with the hour and radius of the corkscrew. The track started at approximately 00:00 and ended at ~14:00 .

Discussion

The Southern Ocean is a notorious harsh environment with strong winds and large waves. During rare and prolonged calm periods, seabirds may take advantage of these conditions for resting, sitting, or foraging for prey near the sea surface. Sleeping and resting are an important part of an animal's daily time budget and can provide an opportunity to digest food during long foraging trips, and these long resting bouts may be more common than realized (Ropert-Coudert et al. 2004). Additionally, the cost of flying without strong winds may make it energetically favourable for a bird to sit and wait for wind speeds to increase to continue its journey (Thorne et al. 2023).

Ocean surface winds are a critical factor for understanding ocean and atmospheric interactions. In polar environments, obtaining high-resolution and continuous in situ meteorological measurements or data at the air-sea interface can be challenging. It has been previously shown that flying birds take advantage of winds and vertical air motion (Treep et al. 2016) and tracking birds with instruments measuring elevation and air pressure allow for the calculation of wind speed and direction (Yonehara et al. 2016, Goto et al. 2017, Ellis-Soto et al. 2021). Seabirds resting on the sea surface can also be used as passive drifters to estimate ocean currents or tides (Yoda et al. 2014, Shamoun-Baranes et al. 2011, Cooper et al. 2018), and such drift observations that represent geostrophic currents (not wind-driven components) have been assimilated into ocean forecasts (Miyazawa et al. 2015).

The observations shown here suggest Southern Giant Petrels may also be an indicator of atmospheric and oceanic processes at the sea surface. It is possible that some biologists and ecologists have overlooked drift patterns in tracking data because these small-scale movements are not always noticeable when viewing tracking data over a large spatiotemporal scale and/or current/tidal patterns (including, inertial oscillations) may not be known or of interest when the goal is to answer major research questions on climate or human impacts, predator-prey interactions, or physiology (Watanabe, Papas-tamatiou 2023). With the availability of large, long-term tracking data-

bases on numerous seabird species (Hindell et al. 2020, Bernard et al. 2020), the drift patterns of resting seabirds could be used as another fine-scale biomonitoring tool by turning ocean wanderers into ocean drifters.

Acknowledgements

The authors acknowledge funding from the National Science Foundation, Office of Polar Programs (OPP – 2026045). We are grateful to the field technicians that deployed the tracker.

References

- Cooper M, Bishop C, Lewis M, Bowers D, Bolton M, Owen E, Dodd S** 2018. What can seabirds tell us about the tide? *Ocean Science* 14(6): 1483–1490.
- Bernard A, Rodrigues AS, Cazalis V, Grémillet D** 2021. Toward a global strategy for seabird tracking. *Conservation Letters* 14(3): p.e12804.
- Ellis-Soto D, Wikelski M, Jetz W** 2023. Animal-borne sensors as a biologically informed lens on a changing climate. *Nature Climate Change* 13(10): 1042–1054.
- Finger JVG, Krüger L, Corá DH, Petry MV** 2023. Habitat selection of southern giant petrels: potential environmental monitors of the Antarctic Peninsula. *Antarctic Science* 35(4): 256–269.
- Goto Y, Yoda K, Sato K** 2017. Asymmetry hidden in birds' tracks reveals wind, heading, and orientation ability over the ocean. *Science Advances* 3: e1700097.
- Hazen EL, Abrahms B, Brodie S, Carroll G, Jacox MG, Savoca MS, Scales KL, Sydeman WJ, Bograd SJ** 2019. Marine top predators as climate and ecosystem sentinels. *Frontiers in Ecology and the Environment* 17(10): 565–574.
- Hindell MA, Reisinger RR, Ropert-Coudert Y, Hückstädt LA, Trathan PN, Bornemann H, Charrassin JB, Chown SL, Costa DP, Danis B, Lea MA** 2020. Tracking of marine predators to protect Southern Ocean ecosystems. *Nature* 580(7801): 87–92.

Miyazawa Y, Guo X, Varlamov SM, Miyama T, Yoda K, Sato K, Kano T, Sato K 2015. Assimilation of the seabird and ship drift data in the north-eastern Sea of Japan into an operational ocean nowcast/forecast system. *Scientific Reports* 5(1): 17672.

Ropert-Coudert Y, Gremillet D, Kato A, Ryan PG, Naito Y, Le Maho Y 2004. A fine-scale time budget of Cape Gannets provides insights into the foraging strategies of coastal seabirds. *Animal Behavior* 67: 985–992.

Shamoun-Baranes JUDY, Bouten W, Camphuysen CJ, Baaij E 2011. Riding the tide: intriguing observations of gulls resting at sea during breeding. *Ibis* 153(2): 411–415.

Thorne LH, Clay TA, Phillips RA, Silvers LG, Wakefield ED 2023. Effects of wind on the movement, behavior, energetics, and life history of seabirds. *Marine Ecology Progress Series* 723: 73–117.

Treep J, Bohrer G, Shamoun-Baranes J, Duriez O, de Moraes Frasson RP, Bouten W 2016. Using high-resolution GPS tracking data of bird flight for meteorological observations. *Bulletin of the American Meteorological Society* 97(6): 951–961.

Watanabe YY, Papastamatiou YP 2023. Biologging and biotelemetry: tools for understanding the lives and environments of marine animals. *Annual Review of Animal Biosciences* 11: 247–267.

Yoda K, Shiomi K, Sato K 2014. Foraging spots of streaked shearwaters in relation to ocean surface currents as identified using their drift movements. *Progress in Oceanography* 122: 54–64.

Yonehara Y, Goto Y, Yoda K, Watanuki Y, Young LC, Weimerskirch H, Bost CA, Sato K 2016. Flight paths of seabirds soaring over the ocean surface enable measurement of fine-scale wind speed and direction. *Proceedings of the National Academy of Sciences* 113(32): 9039–9044.

Paper edited by René A. Navarro

FitzPatrick Institute of African Ornithology,
Department of Biological Sciences,
University of Cape Town



Biodiversity Observations is powered by [Open Journal Systems \(OJS\)](#) and is hosted by the [University of Cape Town Libraries](#). OJS is an open source software application for managing and publishing scholarly journals. Developed and released by the [Public Knowledge Project](#) in 2001, it is the most widely used open source journal publishing platform in existence, with over 30,000 journals using it worldwide.