



Local tropicalizations of splice type surface singularities

Maria Angelica Cueto¹ · Patrick Popescu-Pampu² · Dmitry Stepanov³

With an appendix by Jonathan Wahl (Department of Mathematics, The University of North Carolina at Chapel Hill, 120 E Cameron Avenue, CB #3250, Chapel Hill, NC 27599-3250, USA. Email: jmwahl@email.unc.edu)

Received: 30 May 2022 / Revised: 1 October 2023 / Accepted: 18 October 2023

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023

Abstract

Splice type surface singularities were introduced by Neumann and Wahl as a generalization of the class of Pham–Brieskorn–Hamm complete intersections of dimension two. Their construction depends on a weighted tree called a splice diagram. In this paper, we study these singularities from the tropical viewpoint. We characterize their local tropicalizations as the cones over the appropriately embedded associated splice diagrams. As a corollary, we reprove some of Neumann and Wahl’s earlier results on these singularities by purely tropical methods, and show that splice type surface singularities are Newton non-degenerate complete intersections in the sense of Khovanskii. We also confirm that under suitable coprimality conditions on its weights, the diagram can be uniquely recovered from the local tropicalization. As a corollary of the Newton non-degeneracy property, we obtain an alternative proof of a recent theorem of de Felipe, González Pérez and Mourtada, stating that embedded resolutions of any plane curve singularity can be achieved by a single toric morphism, after re-embedding the

Communicated by Jean-Yves Welschinger.

To Walter Neumann, on the occasion of his 75th birthday.

✉ Patrick Popescu-Pampu
patrick.popescu-pampu@univ-lille.fr

Maria Angelica Cueto
cueto.5@osu.edu

Dmitry Stepanov
stepanov.da@phystech.edu

¹ Mathematics Department, The Ohio State University, 231 W 18th Ave, Columbus, OH 43210, USA

² Univ. Lille, CNRS, UMR 8524 - Laboratoire Paul Painlevé, 59000 Lille, France

³ Department of Higher Mathematics and Center of Fundamental Mathematics, Moscow Institute of Physics and Technology, 9 Institutskiy per., Dolgoprudny, Moscow, Russia 141701

ambient smooth surface germ in a higher-dimensional smooth space. The paper ends with an appendix by Jonathan Wahl, proving a criterion of regularity of a sequence in a ring of convergent power series, given the regularity of an associated sequence of initial forms.

Mathematics Subject Classification Primary 14B05 · 14T90 · 32S05; Secondary 14M25 · 57M15

1 Introduction

Splice diagrams are finite trees with half-edges weighted by integers and with nodes (internal vertices) decorated by $\pm$ signs. If the half-edge weights around each node are pairwise coprime, we say that the splice diagram is *coprime*. This class of weighted trees was first introduced by Siebenmann [52] in 1980 to encode graph manifolds which are integral homology spheres. Coprime splice diagrams with only $+$ node decorations and positive half-edge weights were used by Eisenbud and Neumann in [10] to study special kinds of links in integral homology spheres, in particular those corresponding to curves on normal surface singularities with integral homology sphere links. One of the main theorems of [10] states that such integral homology spheres are described by positively-weighted coprime splice diagrams satisfying the *edge determinant condition*, namely, that the product of the two weights associated to any fixed internal edge must be greater than the product of the weights of the neighboring half-edges.

Interesting isolated surface singularities arise from splice diagrams. For example, complete intersections of *Pham–Brieskorn–Hamm* hypersurface singularities are associated to star splice diagrams (i.e., those with a single node). As recognized by Hamm in [19, Sect. 5] and [20], in order to determine an isolated singularity in $\mathbb{C}^n$, all maximal minors of the coefficient matrix $(c_{ij})_{i,j}$ of each polynomial $f_i := \sum_j c_{ij} z_j^{a_j}$ in the Brieskorn system must be non-zero. In turn, work of Neumann [35] shows that universal abelian covers of quasi-homogeneous complex normal surface singularities with rational homology sphere links are complete intersections of Pham–Brieskorn–Hamm hypersurface singularities.

In 2002, Neumann and Wahl [37] extended this family of complete intersections by defining *splice type surface singularities* associated to splice diagrams whose weights satisfy a special arithmetic property called the *semigroup condition*. These singularities are defined by explicit splice type systems of convergent power series near the origin, whose coefficients satisfy generalizations of Hamm’s maximal minors conditions. Splice type surface singularities and the related class of *splice quotients* (determined by diagrams subject to an additional congruence condition) have been further studied by both authors in [38–40], and by Lamberson, Némethi, Okuma and Pedersen in [27, 31–33, 42–44, 46–48]. For more details, we refer the reader to the surveys [36, 45, 58, 59].

The present paper uses tropical geometry techniques to study splice type systems with n leaves associated to splice diagrams satisfying the edge determinant and semi-

group conditions. Our first main result recovers and strengthens a central theorem from [38] (see Theorem 2.16). More precisely:

Theorem 1.1 *Splice type systems are Newton non-degenerate complete intersection systems of equations. The associated splice type singularities are isolated, irreducible and not contained in any coordinate subspace of the corresponding ambient space $\mathbb{C}^n$.*

This statement plays a major role in the proof of the Neumann–Wahl Milnor fiber conjecture for splice type singularities with integral homology sphere links [39] obtained by the present authors. For an overview of this proof, we refer the reader to [6]. Motivated by Theorem 1.1, we formulate Question 7.17.

The notion of Newton non-degeneracy (in the sense of Kouchnirenko [25] and Khovanskii [22]) is closely related to the notion of the initial form of a series relative to a weight vector, which lies at the core of tropical geometry (see Sect. 3, in particular Definition 3.1). A regular sequence of convergent power series $(f_1, \dots, f_s)$ in $\mathbb{C}\{z_1, \dots, z_n\}$ defining a germ $(X, 0) \hookrightarrow \mathbb{C}^n$ is a *Newton non-degenerate complete intersection system* if for each weight vector w with positive entries, the associated initial forms $(\text{in}_w(f_i))_i$ determine hypersurfaces of the algebraic torus $(\mathbb{C}^*)^n$ whose sum is a normal crossings divisor in the neighborhood of their intersection. This condition is automatically satisfied whenever the intersection is empty. Surprisingly, not many examples of Newton non-degenerate complete intersection systems are known in codimension two or higher. Theorem 1.1 contributes a large class of examples of such systems.

Newton non-degeneracy enables the resolution of the germ $(X, 0) \hookrightarrow \mathbb{C}^n$ by a single toric morphism. Indeed, works of Varchenko [57] (for hypersurfaces) and Oka [41, Chapter III, Theorem (3.4)] (for complete intersections) show that such a morphism may be defined by a regular subdivision of the positive orthant refining the dual fan of the Newton polyhedron of each function f_i . Furthermore, the complete dual fan is not needed to achieve a resolution. Indeed, [41, Theorem III.3.4] allows us to restrict to the subfan corresponding to the orbits intersecting the strict transform of the given germ.

The support of this subfan depends only on the ideal defining the germ $(X, 0)$ but not on the particular generators f_i . It is the finite part of the so-called *local tropicalization* of the embedding $(X, 0) \hookrightarrow \mathbb{C}^n$. This local version of the standard notion of tropicalization of a subvariety of an algebraic torus was first introduced by the last two authors in [49] as a tool to study arbitrary subgerms of $\mathbb{C}^n$ or, more generally, arbitrary morphisms from analytic or formal germs to germs of toric varieties.

As was mentioned earlier, splice diagrams record topological information about the link of splice type singularities. Indeed, starting from a normal surface singularity with a rational homology sphere link, Neumann and Wahl [37] build a splice diagram which determines the dual tree of the minimal normal crossings resolution (up to a finite ambiguity) whenever this graph is not a star tree. This diagram is homeomorphic to the dual tree: it is obtained by disregarding all bivalent vertices of the dual tree. It satisfies the edge determinant condition, but not necessarily the semigroup or the congruence conditions. When $(X, 0) \hookrightarrow \mathbb{C}^n$ has an integral homology sphere link,

the ambiguity disappears and the dual tree is completely determined by the splice diagram.

Given a splice diagram Γ with n leaves, the construction of Neumann and Wahl associates a weight vector $w_u \in (\mathbb{Z}_{>0})^n$ to each vertex u of Γ . These vectors induce a piecewise linear embedding of Γ into the standard simplex in $\mathbb{R}^n$ after appropriate normalization. Our second main result shows the close connection between Γ , the local tropicalization of the associated splice type system in $\mathbb{C}^n$ and its resolution diagrams:

Theorem 1.2 *Let Γ be a splice diagram satisfying the semigroup condition and let $(X, 0) \hookrightarrow \mathbb{C}^n$ be the germ defined by an associated splice type system. Then, the finite local tropicalization of X is the cone over an embedding of Γ in $\mathbb{R}^n$. Furthermore, in the coprime case, Γ can be uniquely recovered from this fan.*

Theorem 1.2 shows that the link at the origin of the local tropicalization of $(X, 0) \hookrightarrow \mathbb{C}^n$ (obtained by intersecting the fan with the $(n - 1)$ -dimensional sphere) is homeomorphic to the splice diagram Γ . To the best of our knowledge, *this is the first tropical interpretation of Siebenmann's splice diagrams*. In this spirit, we view Siebenmann's paper [52] as a precursor to tropical geometry (for others, see [28, Chapter 1]).

Our method to characterize the local tropicalization is different from the general one discussed in Oka's book [41] and described briefly above. Namely, we do not use the Newton polyhedra of the collection of series defining the germ. Instead, we use a “mine-sweeping” approach, using successive stellar subdivisions of the standard simplex in $\mathbb{R}^n$ dictated by the splice diagram, in order to remove relatively open cones in the positive orthant avoiding the local tropicalization.

Once the local tropicalization is determined (via Theorem 1.2), a simple computation confirms the Newton non-degeneracy of the system. In turn, by analyzing the local tropicalizations of the intersections of the germ with the coordinate subspaces of $\mathbb{C}^n$, we conclude that $(X, 0)$ is an isolated complete intersection surface singularity, thus completing the proof of Theorem 1.1.

As a consequence of Theorem 1.1, we provide an alternative proof of the main theorem of de Felipe, González Pérez and Mourtada [7], stating that *any germ of a reduced plane curve may be resolved by one toric modification after re-embedding its ambient smooth germ of surface into a higher-dimensional germ $(\mathbb{C}^n, 0)$* (see Corollary 7.14). The first theorem of this kind was proved by Goldin and Teissier [17] for irreducible germs of plane curves.

Our paper is organized as follows. In Sect. 2, we review the definitions and main properties of splice diagrams, splice type systems and end-curves associated to rooted splice diagrams, following [38, 39]. Sections 3 and 4 include background material about local tropicalizations and Newton non-degeneracy. In Sect. 5 we show how to embed a given splice diagram Γ with n leaves into the standard $(n - 1)$ -simplex in $\mathbb{R}^n$, and we highlight various convexity properties of this embedding. The proof of the first part of Theorem 1.2 is discussed in Sect. 6, while Theorem 1.1 is proven in Sect. 7. Section 8 characterizes local tropicalizations of splice type systems defined by a coprime splice diagram and shows how to recover the diagram from the tropical fan, thus yielding the second part of Theorem 1.2. Finally, Sect. 9 discusses the dependency

of the construction of splice type systems on the choice of admissible monomials for arbitrary splice diagrams, in the spirit of [38, Sect. 10].

Appendix A, written by Jonathan Wahl, includes a proof of [38, Lemma 3.3] that was absent from the literature. This result confirms that given a finite sequence $(f_1, \dots, f_s)$ in $\mathbb{C}\{z_1, \dots, z_n\}$ and a fixed positive integer vector $w \in (\mathbb{Z}_{>0})^n$, the regularity of the sequence $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ of initial forms ensures that the original sequence is regular, and furthermore, that the w -initial ideal must be generated by the sequence of initial forms. This statement can be used to determine if a given w lies in the local tropicalization of the germ defined by the vanishing of the input sequence (see Corollary 7.7), providing an alternative proof to part of Theorem 1.2.

2 Splice diagrams, splice type systems and end-curves from rooted splice diagrams

In this section, we recall the notions of splice diagram and splice type systems associated to them. The definitions follow closely the work of Neumann and Wahl [38, 39].

We start with some basic terminology and notations about trees:

Definition 2.1 A *tree* is a finite connected graph with no cycles and at least one vertex. The *star* of a vertex v of the tree Γ is the set $\text{Star}_\Gamma(v)$ of edges adjacent to v . The *valency* of v is the cardinality of $\text{Star}_\Gamma(v)$, which we denote by δ_v . A *node* of a tree is a vertex v whose valency is greater than one, whereas a *leaf* is a one-valent vertex. We denote the set of nodes of Γ by $V^\circ(\Gamma)$ and its set of leaves by $\partial\Gamma$.

When the ambient tree is understood from context, we remove it from the notation and simply write $\text{Star}(v)$.

Remark 2.2 Endowing a tree with a metric allows us to consider geodesics on it and distances between vertices. Whenever these notions are invoked, it is understood that each edge of the tree has length one.

Definition 2.3 Given a subset $W = \{p_1, \dots, p_k\}$ of vertices of the tree Γ , we denote by $[W]$ or $[p_1, \dots, p_k]$ the subtree of Γ spanned by these points. We call it the *convex hull* of the set $\{p_1, \dots, p_k\}$ inside Γ . For example, $\Gamma = [\partial\Gamma]$.

Splice diagrams are special kinds of trees enriched with weights around all nodes, as we now describe:

Definition 2.4 A *splice diagram* is a pair $(\Gamma, \{d_{v,e}\}_{v,e})$, where Γ is a tree without valency-two vertices, with at least one node, and decorated with a weight function on the star of each node v of Γ , denoted by

$$\text{Star}_\Gamma(v) \rightarrow \mathbb{Z}_{>0}: e \mapsto d_{v,e}.$$

We call $d_{v,e}$ the *weight of e at v* . If u is any other vertex of Γ such that e lies in the unique geodesic of Γ joining u and v , we write $d_{v,u} := d_{v,e}$. We view this as the

weight in the neighborhood of v pointing towards u . The *total weight* of a node v of Γ is the product $\boxed{d_v} := \prod_{e \in \text{Star}_\Gamma(v)} d_{v,e}$.

Remark 2.5 Let $(\Gamma, \{d_{v,e}\}_{v,e})$ be a splice diagram. For simplicity, we remove the collection of weights from the notation and simply use Γ to refer to the splice diagram. By a similar abuse of notation, we may view $\text{Star}_\Gamma(v)$ also as a splice diagram, whose weights around its unique node v are inherited from Γ . Splice diagrams with one node will be referred to as *star splice diagrams*, and the underlying graphs as *star trees*.

Definition 2.6 Let u and v be two distinct vertices of the splice diagram Γ . The *linking number* $\boxed{\ell_{u,v}}$ between u and v is the product of all the weights adjacent to, but not on, the geodesic $[u, v]$ joining u and v . Thus, $\ell_{v,u} = \ell_{u,v}$. We set $\boxed{\ell_{v,v}} := d_v$ for each node v of Γ . The *reduced linking number* $\boxed{\ell'_{u,v}}$ is defined via a similar product where we exclude the weights around u and v . In particular, $\ell'_{v,v} = 1$ for each node v of Γ .

Remark 2.7 Given a node v and a leaf λ of Γ , it is immediate to check that $\ell_{v,\lambda} d_{v,\lambda} = \ell'_{v,\lambda} d_v$.

Linking numbers satisfy the following useful identity, whose proof is immediate from Definition 2.6 (see [13, Proposition 69]):

Lemma 2.8 *If u, v, w are vertices of Γ with $u \in [v, w]$, then $\ell_{u,v} \ell_{u,w} = d_u \ell_{v,w}$.*

In [34, Theorem 1], Neumann gave explicit descriptions of integral homology spheres associated to star splice diagrams as links of Pham–Brieskorn–Hamm surface singularities. The following definition was introduced by Neumann and Wahl in [39, Sect. 1] to characterize which integral homology spheres may be realized as links of normal surface singularities. Its origins can be traced back to [10, p. 82].

Definition 2.9 Let Γ be a splice diagram. Given two adjacent nodes u and v of Γ , the *determinant* of the edge $[u, v]$ is the difference between the product of the two decorations on $[u, v]$ and the product of the remaining decorations in the neighborhoods of u and v , that is,

$$\boxed{\det([u, v])} := d_{u,v} d_{v,u} - \ell_{u,v}. \quad (2.1)$$

The splice diagram Γ satisfies the *edge determinant condition* if all edges have positive determinants.

The next result on integral homology sphere links is due to Eisenbud and Neumann (see [10, Theorem 9.4] for details and Definition 8.1 for the meaning of the coprimality condition). It was strengthened by Pedersen in [47, Theorem 1] to address the case of rational homology sphere links:

Theorem 2.10 *The integral homology sphere links of normal surface singularities are precisely the oriented 3-manifolds $\Sigma(\Gamma)$ associated to coprime splice diagrams Γ which satisfy the edge determinant condition.*

The construction of oriented 3-manifolds from splice diagrams is due to Siebenmann [52]. They are obtained from splicing Seifert-fibered oriented 3-manifolds $\Sigma(\text{Star}_\Gamma(v))$ associated to each node v of Γ along special fibers of their respective Seifert fibration corresponding to the edges of Γ . For each $\Sigma(\text{Star}_\Gamma(v))$, these special fibers are in bijection with the δ_v -many edges adjacent to v . Each edge $[u, v]$ induces a splicing of both $\Sigma(\text{Star}_\Gamma(u))$ and $\Sigma(\text{Star}_\Gamma(v))$ along the oriented fibers corresponding to the edge. These fibers are knots in both Seifert-fibered manifolds and their linking number is precisely $\ell_{u,v}$ (see [10, Theorem 10.1]).

The following result shows that the edge determinant condition yields Cauchy–Schwarz’ type inequalities:

Lemma 2.11 *Assume that the splice diagram Γ satisfies the edge determinant condition. Then,*

$$d_u d_v \geq \ell_{u,v}^2, \quad \text{for all nodes } u, v \in \Gamma. \quad (2.2)$$

Furthermore, equality holds if and only if $u = v$.

Proof The result follows by induction on the distance $\text{dist}_\Gamma(u, v)$ between u and v (see Remark 2.2). The base case corresponds to adjacent nodes. Lemma 2.8 is used for the inductive step. $\square$

If Γ satisfies the edge determinant condition, then the linking numbers verify the following inequality, which generalizes Lemma 2.8. This inequality is reminiscent of the ultrametric condition for dual graphs of arborescent singularities (see [16, Proposition 1.18]):

Proposition 2.12 *Assume that the splice diagram Γ satisfies the edge determinant condition. Then, for all nodes u, v and w of Γ , we have $\ell_{u,v} \ell_{u,w} \leq d_u \ell_{v,w}$. Furthermore, equality holds if and only if $u \in [v, w]$.*

Proof Consider the tree T spanned by u, v and w and let a be the unique node in the intersections of the three geodesics $[u, v]$, $[u, w]$ and $[v, w]$. We prove the inequality by a direct calculation. By Lemma 2.8 applied to the triples $\{a, u, v\}$, $\{a, u, w\}$ and $\{a, v, w\}$, we have:

$$\ell_{a,u} \ell_{a,v} = d_a \ell_{u,v}, \quad \ell_{a,u} \ell_{a,w} = d_a \ell_{u,w} \quad \text{and} \quad \ell_{a,w} \ell_{a,v} = d_a \ell_{v,w}.$$

These expressions combined with the inequality (2.2) applied to the pair $\{a, u\}$ yield:

$$\ell_{u,v} \ell_{u,w} = \frac{\ell_{a,v} \ell_{a,u}}{d_a} \frac{\ell_{a,w} \ell_{a,u}}{d_a} = \frac{\ell_{a,v} \ell_{a,w}}{d_a} \frac{(\ell_{a,u})^2}{d_a} = \ell_{v,w} \frac{(\ell_{a,u})^2}{d_a} \leq \ell_{v,w} d_u.$$

Furthermore, Lemma 2.11 confirms that equality is attained if and only if $a = u$, that is, if and only if u lies in the geodesic $[v, w]$. This concludes our proof. $\square$

Before introducing splice type systems as defined by Neumann and Wahl in [38, 39], we set up notation arising from toric geometry. We write $\boxed{n} := |\partial \Gamma|$ for the number of leaves of the splice diagram Γ (where $|\cdot|$ denotes the cardinality of a finite

set) and let $M(\partial \Gamma)$ be the free abelian group generated by all leaves of Γ . We denote by $N(\partial \Gamma)$ its dual lattice and write the associated pairing using dot product notation, i.e. $w \cdot m$ whenever $w \in N(\partial \Gamma)$ and $m \in M(\partial \Gamma)$. Fixing a basis $\{\overline{w_\lambda} : \lambda \in \partial \Gamma\}$ for $N(\partial \Gamma)$ and its dual basis $\{\overline{m_\lambda} : \lambda \in \partial \Gamma\}$ for $M(\partial \Gamma)$ identifies both lattices with $\mathbb{Z}^n$. To each m_λ , we associated a variable $\overline{z_\lambda}$. We view $M(\partial \Gamma)$ as the lattice of exponents of monomials in those variables and $N(\partial \Gamma)$ as the associated lattice of weight vectors.

In addition to defining weights for all leaves of Γ , each node u in Γ has an associated weight vector:

$$\overline{w_u} := \sum_{\lambda \in \partial \Gamma} \ell_{u,\lambda} w_\lambda \in N(\partial \Gamma). \quad (2.3)$$

As was mentioned in Sect. 1, star splice diagrams Γ with a unique node v produce Pham–Brieskorn–Hamm singularities using the monomials $\{z_\lambda^{d_{v,\lambda}} : \lambda \in \partial \Gamma\}$. Neumann and Wahl’s splice type systems [38, 39] generalize this construct to diagrams with more than one node. In addition to satisfying the edge determinant condition, Γ must have an extra arithmetic property that allows to replace each monomial $z_v^{d_{v,\lambda}}$ by a suitable monomial associated to the pair (v, e) where v is any node and e is an edge adjacent to it (see (2.7)). This property, which ensures that all monomials associated to a vertex v have the same w_v -degree, will automatically hold for star splice diagrams.

Definition 2.13 A splice diagram Γ satisfies the *semigroup condition* if for each node v and each edge $e \in \text{Star}(v)$, the total weight d_v of v belongs to the subsemigroup of $(\mathbb{N}, +)$ generated by the set of linking numbers between v and the leaves λ seen from v in the direction of e , that is, such that $e \subseteq [v, \lambda]$. Therefore, we may write:

$$d_v = \sum_{\lambda \in \partial_{v,e}\Gamma} m_{v,e,\lambda} \ell_{v,\lambda}, \quad \text{or equivalently} \quad d_{v,e} = \sum_{\lambda \in \partial_{v,e}\Gamma} m_{v,e,\lambda} \ell'_{v,\lambda}, \quad (2.4)$$

where $\overline{m_{v,e,\lambda}} \in \mathbb{N}$ for all λ and $\overline{\partial_{v,e}\Gamma}$ is the set of leaves λ of Γ with $e \subseteq [v, \lambda]$.

Assume that Γ satisfies the semigroup condition and pick coefficients $m_{v,e,\lambda}$ satisfying (2.4). Using these integers we define an exponent vector (i.e., an element of $M(\partial \Gamma)$) for each pair (v, e) as above:

$$\overline{m_{v,e}} := \sum_{\lambda \in \partial_{v,e}\Gamma} m_{v,e,\lambda} m_\lambda \in M(\partial_{v,e}\Gamma) \subset M(\partial \Gamma). \quad (2.5)$$

Following [39], we refer to it as an *admissible exponent* for (v, e) . Note that the relation (2.4) is equivalent to:

$$w_v \cdot m_{v,e} = d_v. \quad (2.6)$$

Each admissible exponent $m_{v,e}$ defines an *admissible monomial*, which was denoted by $M_{v,e}$ in [39]:

$$\overline{z^{m_{v,e}}} := \prod_{\lambda \in \partial_{v,e}\Gamma} z_\lambda^{m_{v,e,\lambda}}. \quad (2.7)$$

Definition 2.14 Let Γ be a splice diagram which satisfies both the edge determinant and the semigroup conditions of Definitions 2.9 and 2.13. We fix an order for its set $\partial \Gamma$ of n leaves.

- A *strict splice type system* associated to Γ is a finite family of $(n - 2)$ polynomials of the form:

$$\boxed{f_{v,i}(\underline{z})} := \sum_{e \in \text{Star}(v)} \boxed{c_{v,e,i}} \underline{z}^{m_{v,e}} \quad \text{for all } i \in \{1, \dots, \delta_v - 2\} \text{ and } v \text{ a node of } \Gamma, \quad (2.8)$$

where $m_{v,e} \in M(\partial \Gamma)$ are the admissible exponent vectors defined by (2.5) for each node $v \in \Gamma$ and each edge $e \in \text{Star}(v)$. We also require the coefficients $c_{v,e,i}$ to satisfy the *Hamm determinant conditions*. Namely, for any node $v \in \Gamma$, if we fix an ordering of the edges in $\text{Star}(v)$, then all the maximal minors of the matrix of coefficients $(c_{v,e,i})_{e,i} \in \mathbb{C}^{\delta_v \times (\delta_v - 2)}$ must be non-zero.

- A *splice type system* $\boxed{\mathcal{S}(\Gamma)}$ associated to Γ is a finite family of power series of the form

$$\boxed{F_{v,i}(\underline{z})} := f_{v,i}(\underline{z}) + g_{v,i}(\underline{z}) \quad \text{for all } i \in \{1, \dots, \delta_v - 2\} \text{ and } v \text{ a node of } \Gamma, \quad (2.9)$$

where the collection $(f_{v,i})_{v,i}$ is a strict splice type system associated to Γ and each $g_{v,i}$ is a convergent power series satisfying the following condition for each exponent m in the support of $g_{v,i}$:

$$w_v \cdot m > d_v \quad \text{and} \quad w_u \cdot m > \ell_{u,v} \quad \text{for each node } u \text{ of } \Gamma \text{ with } u \neq v. \quad (2.10)$$

- A *splice type singularity* associated to Γ is the subgerm of $(\mathbb{C}^n, 0)$ defined by $\mathcal{S}(\Gamma)$.

Remark 2.15 The inequalities in (2.10) should be compared with the equality imposed in (2.6). As was shown by Neumann and Wahl in [38, Lemma 3.2], the right-most inequality in (2.10) follows from the left-most one and the edge determinant condition. We choose to include both inequalities in (2.10) for mere convenience since we will need both of them for several arguments in Sect. 6.

The issue of dependency of the set of germs defined by splice type systems on the choice of admissible monomials is a subtle one. We postpone this discussion to Sect. 9.

As was mentioned in Sect. 1, splice type singularities satisfy the following crucial property, proved by Neumann and Wahl in [38, Theorem 2.6]. An alternative proof of this statement, using local tropicalization, will be provided at the end of Sect. 7.

Theorem 2.16 *Splice type singularities are isolated complete intersection surface singularities.*

Example 2.17 We let Γ be the splice diagram to the left of Fig. 1. Then, $d_u = 294$, $d_v = 770$ and $\ell_{u,v} = 420$ and so the edge determinant condition $d_u d_v > \ell_{u,v}^2$ holds for $[u, v]$.




Fig. 1 From left to right: a splice diagram and its associated rooted diagram obtained by fixing one of the leaves as its root r , and removing one weight from the star of each node

The semigroup condition is also satisfied, since

$$49 = 0 \cdot (2 \cdot 5) + 1 \cdot (2 \cdot 7) + 1 \cdot (5 \cdot 7) \quad \text{and} \quad 11 = 1 \cdot (3) + 4 \cdot (2) = 3 \cdot (3) + 1 \cdot (2).$$

Thus, we may take as exponents $m_{u,[u,v]} = (0, 0, 0, 1, 1)$ and $m_{v,[u,v]} = (1, 4, 0, 0, 0)$ or $(3, 1, 0, 0, 0)$ in $\mathbb{Z}^5$. A possible strict splice type system for Γ is:

$$\begin{cases} f_{u,1} := z_1^2 - 2 z_2^3 + z_4 z_5, \\ f_{v,1} := z_1 z_2^4 + z_3^7 + z_4^5 - 2155 z_5^2, \\ f_{v,2} := 33 z_1 z_2^4 + z_3^7 + 2 z_4^5 - 2123 z_5^2. \end{cases} \quad (2.11)$$

An alternative system is obtained by replacing the admissible monomial $z_1 z_2^4$ with $z_1^3 z_2$. The coefficients of the system were chosen to simplify the parameterization of the end-curve of the corresponding splice type surface singularity associated to the leaf λ_1 (see Example 2.22 for details).

A central role in this paper will be played by tropicalizations and weighted initial forms of series and ideals of $\mathbb{C}\{z_\lambda : \lambda \in \partial \Gamma\}$, which we discuss in Sect. 3. In Proposition 7.1, we determine the initial forms of the series $F_{v,i}$ of a splice type system $\mathcal{S}(\Gamma)$ with respect to each weight vector w_u from (2.3). Its proof is a consequence of the next two lemmas:

Lemma 2.18 *Assume that Γ is a splice diagram satisfying the edge determinant condition. Then, for any pair of adjacent nodes u, v of Γ we have:*

$$w_u \in \frac{\ell_{u,v}}{d_v} w_v + \mathbb{R}_{>0} \{w_\lambda : \lambda \in \partial_{v,[u,v]} \Gamma\}.$$

Proof We write $e = [u, v]$. The definition of linking numbers gives the following expressions for each $\lambda \in \partial \Gamma$:

$$\ell_{u,\lambda} = \begin{cases} (\ell_{v,\lambda} d_u) / (d_{v,u} d_{u,v}) & \text{if } \lambda \in \partial_{u,e} \Gamma, \\ (\ell_{v,\lambda} d_{v,u} d_{u,v}) / d_v & \text{if } \lambda \in \partial_{v,e} \Gamma. \end{cases}$$

The statement follows by substituting these expressions in the definition of w_u from (2.3) and by using the edge determinant condition, i.e.,

$$\begin{aligned} w_u &= \frac{d_u}{d_{v,u} d_{u,v}} \sum_{\lambda \in \partial_{u,e} \Gamma} \ell_{v,\lambda} w_\lambda + \frac{d_{v,u} d_{u,v}}{d_v} \sum_{\lambda \in \partial_{v,e} \Gamma} \ell_{v,\lambda} w_\lambda \\ &= \frac{\ell_{u,v}}{d_v} w_v + \sum_{\lambda \in \partial_{v,e} \Gamma} \underbrace{\frac{\det(e) \ell_{v,\lambda}}{d_v}}_{>0} w_\lambda. \end{aligned} \quad (2.12)$$

□

Lemma 2.19 *Assume that Γ satisfies the edge determinant and semigroup conditions. Then, the exponent vector $m_{v,e}$ from (2.5) satisfies $w_u \cdot m_{v,e} \geq \ell_{u,v}$ for all nodes u of Γ and each edge $e \in \text{Star}_\Gamma(v)$. Furthermore, equality holds if and only if $e \not\subseteq [u, v]$.*

Proof If $u = v$, then $w_v \cdot m_{v,e} = d_v = \ell_{v,v}$. If $u \neq v$, we argue by induction on the distance $\text{dist}_\Gamma(u, v) > 0$ between u and v in the tree Γ . If $\text{dist}_\Gamma(u, v) = 1$, we let $e' = [u, v]$. Expression (2.12) yields

$$\begin{aligned} w_u \cdot m_{v,e} &= \frac{\ell_{u,v}}{d_v} w_v \cdot m_{v,e} + \frac{\det(e')}{d_v} \left(\sum_{\lambda \in \partial_{v,e'} \Gamma} m_{v,e,\lambda} \ell_{v,\lambda} \right) \\ &= \ell_{u,v} + \frac{\det(e')}{d_v} \left(\sum_{\lambda \in \partial_{v,e'} \Gamma} m_{v,e,\lambda} \ell_{v,\lambda} \right). \end{aligned}$$

The second summand is always non-negative and it equals zero if and only if $e \neq e'$.

If $\text{dist}_\Gamma(u, v) > 1$, we let u' be the unique node adjacent to u in $[u, v]$ and set $e' := [u', u]$. Note that $e \in [v, u]$ if and only if $e \in [v, u']$. Expression (2.12) applied to $\{u, u'\}$, the non-negativity of each $m_{v,e,\lambda}$, the inductive hypotheses on $\{u', v\}$ and Proposition 2.12 yield

$$w_u \cdot m_{v,e} = \frac{\ell_{u,u'}}{d_{u'}} \underbrace{w_{u'} \cdot m_{v,e}}_{\geq \ell_{u',v}} + \frac{\det(e')}{d_{u'}} \sum_{\lambda \in \partial_{u',e'} \Gamma} \underbrace{m_{v,e,\lambda} \ell_{u',\lambda}}_{\geq 0} \geq \frac{\ell_{u,u'} \ell_{u',v}}{d_{u'}} = \ell_{u,v}.$$

By construction, equality is achieved if and only if $e \not\subseteq [u', v]$, which is equivalent to $e \not\subseteq [u, v]$. □

In Sect. 7, we will be interested in curves obtained from a given splice type system when we choose a leaf r of the corresponding splice diagram Γ to be its root. We orient the resulting rooted tree $[\Gamma_r]$ towards the root and remove one weight in the neighborhood of each node, namely the one pointing towards the root, as seen on the right of Fig. 1. We write $[\partial_r \Gamma]$ for the set of $(n-1)$ non-root leaves of Γ_r and assume it is ordered. The following definition was introduced in [38, Sect. 3] by the name of *splice diagram curves*.

Definition 2.20 Assume that the rooted splice diagram Γ_r satisfies the semigroup condition and consider a fixed strict splice type system $\mathcal{S}(\Gamma)$ associated to the (unrooted) splice diagram Γ . For each node v of Γ and each index $i \in \{1, \dots, \delta_v\}$, we let $\boxed{h_{v,i}(\underline{z})} \in \mathbb{C}[z_\lambda : \lambda \in \partial_r \Gamma]$ be the polynomial obtained from $f_{v,i}(\underline{z})$ by removing the term corresponding to the unique edge adjacent to v pointing towards r . The subvariety of $\mathbb{C}^{\partial_r \Gamma} \simeq \mathbb{C}^{n-1}$ defined by the vanishing of $(h_{v,i}(\underline{z}))_{v,i}$ is called the *end-curve* of $\mathcal{S}(\Gamma)$ relative to r . We denote it by $\boxed{C_r}$.

A planar embedding of Γ_r determines an ordering of the edges adjacent to a fixed node v that point away from r (for example, by reading them from left to right). Once this order is fixed, by the Hamm determinant conditions, each group of equations $(h_{v,i}(\underline{z}) = 0)_{i=1}^{\delta_v-2}$ becomes equivalent to a collection of w_v -homogeneous binomial equations of the form

$$\underline{z}^{m_{v,e_j}} - a_{v,j} \underline{z}^{m_{v,e_{\delta_v-1}}} = 0 \quad \text{for } j \in \{1, \dots, \delta_v - 2\},$$

with all $a_{v,j} \neq 0$. The next statement summarizes the main properties of C_r discussed in [38, Theorem 3.1]:

Theorem 2.21 *The subvariety $C_r \subseteq \mathbb{C}^{n-1}$ is a reduced complete intersection curve, smooth away from the origin, and meets any coordinate subspace of $\mathbb{C}^{n-1}$ only at the origin. It has g many components, where $g := \gcd\{\ell_{r,\lambda} : \lambda \in \partial_r \Gamma\}$. All of them are isomorphic to torus-translates of the monomial curve in $\mathbb{C}^{n-1}$ with parameterization $t \mapsto (t^{\ell_{r,\lambda_1}/g}, \dots, t^{\ell_{r,\lambda_{n-1}}/g})$.*

Example 2.22 We fix the splice diagram from Example 2.17 and consider its rooted analog obtained by setting the first leaf as its root r , as seen in the right of Fig. 1. By construction, $\ell_{r,2} = 49$, $\ell_{r,3} = 30$, $\ell_{r,4} = 42$, $\ell_{r,5} = 105$, $\ell_{r,u} = 147$, $\ell_{r,v} = 210$, $w_u = (49, 30, 42, 105)$ and $w_v = (70, 10, 14, 35)$. The equations defining this end-curve are obtained by removing the monomial indexed by the edge pointing towards r in each equation from (2.11). Since $g = 1$, the curve C_r is reduced and irreducible. It is defined as the solution set to

$$-2z_2^3 + z_4z_5 = z_3^7 + z_4^5 - 2155z_5^2 = z_3^7 + 2z_4^5 - 2123z_5^2 = 0.$$

Linear combinations of the last two expressions yield the equivalent binomial system:

$$-2z_2^3 + z_4z_5 = z_4^5 + 32z_5^2 = z_3^7 - 2187z_5^2 = 0.$$

An explicit parameterization is given by $(z_2, z_3, z_4, z_5) = (-t^{49}, 3t^{30}, -2t^{42}, t^{105})$.

The collection $h_{v,i}(\underline{z})$ of polynomials defining the end-curve C_r determines a map $G_r: \mathbb{C}^{n-1} \rightarrow \mathbb{C}^{n-2}$. Our next result, which we state for comparison's sake with Corollary 7.12, discusses the restriction of this map to each coordinate hyperplane of $\mathbb{C}^{n-1}$:

Corollary 2.23 *For every $\lambda \in \partial_r \Gamma$, the restriction of G_r to the hyperplane H_λ of $\mathbb{C}^{n-1}$ defined by the equation $z_\lambda = 0$ is dominant.*

Proof By Theorem 2.21, the fiber over the origin of the restricted map $G_{r|H_\lambda}$ is finite. Upper semicontinuity of fiber dimensions implies that the generic fiber is also 0-dimensional. Since $\dim H_\lambda = n - 2$, the map $G_{r|H_\lambda}$ must be dominant. $\square$

3 Local tropicalization

In [49], the last two authors developed a theory of *local tropicalizations* of algebraic, analytic or formal germs endowed with maps to (not necessarily normal) toric varieties, adapting the original formulation of global tropicalization (see, e.g. [28]) to the local setting. In this section, we recall the basics on local tropicalizations that will be needed in Sect. 6. We focus our attention on germs $(Y, 0) \hookrightarrow \mathbb{C}^n$ defined by ideals I of the ring of convergent power series $\boxed{\mathcal{O}} := \mathbb{C}\{z_1, \dots, z_n\}$ near the origin, rather than of its completion $\hat{\mathcal{O}} := \mathbb{C}[[z_1, \dots, z_n]]$. As Remark 3.5 confirms, both local tropicalizations yield the same set.

The notion of local tropicalization of an embedded germ is rooted on the construction of initial ideals associated to non-negative weight vectors, which we now describe. Any weight vector $w := (w_1, \dots, w_n) \in (\mathbb{R}_{\geq 0})^n$ induces a real-valued valuation on $\mathcal{O}$, known as the w -weight, as follows. Given a monomial $\boxed{z^\alpha} := z_1^{\alpha_1} \cdots z_n^{\alpha_n}$, we set $w(\boxed{z^\alpha}) := w \cdot \alpha = \sum_{i=1}^n w_i \alpha_i$. In turn, for each $f = \sum_{\alpha} c_{\alpha} \boxed{z^\alpha}$ with $f \neq 0$ we set

$$\boxed{w(f)} := \min\{w(\boxed{z^\alpha}) : c_{\alpha} \neq 0\} = \min\{w \cdot \alpha : c_{\alpha} \neq 0\}. \quad (3.1)$$

We define $w(0) := \infty$. The set $\{\alpha : c_{\alpha} \neq 0\}$ is called the *support* of f , and it is the basis of the construction of the *Newton polyhedron* and the *Newton fan* of f (see Definition 3.3).

Definition 3.1 Given $f \in \mathcal{O}$ and $w \in (\mathbb{R}_{\geq 0})^n$, the w -initial form $\boxed{\text{in}_w(f)} \in \mathcal{O}$ is the sum of the terms in the series f with minimal w -weight $w(f)$. In turn, given an ideal I of $\mathcal{O}$, the w -initial ideal $\boxed{\text{in}_w(I)\mathcal{O}}$ of I in $\mathcal{O}$ is generated by the w -initial forms of all elements of I . If $w \in (\mathbb{R}_{>0})^n$, the w -initial ideal $\boxed{\text{in}_w(I)}$ of I is the ideal of $\mathbb{C}[z_1, \dots, z_n]$ generated by the w -initial forms of all elements of I .

Example 3.2 Given the minimal splice type system from Example 2.17, we have $w_u = (147, 98, 60, 84, 210)$ and $w_v = (210, 140, 110, 154, 385)$. The polynomial $f_{u,1}$ is w_u -homogeneous, whereas $f_{v,1}$ and $f_{v,2}$ are w_v -homogeneous. Their initial forms relative to the weight vectors w_u and w_v are:

$$\begin{cases} \text{in}_{w_v}(f_{u,1}) = z_1^2 - 2z_2^3, \\ \text{in}_{w_u}(f_{v,1}) = z_3^7 + z_4^5 - 2155z_5^2, \\ \text{in}_{w_u}(f_{v,2}) = z_3^7 + 2z_4^5 - 2123z_5^2. \end{cases}$$

We will see in Lemma 7.4 that $\text{in}_w(S(\Gamma)) = \langle \text{in}_w(f_{u,1}), \text{in}_w(f_{v,1}), \text{in}_w(f_{v,2}) \rangle \mathcal{O}$ when w equals w_u or w_v .

The initial forms of a series determine its Newton fan as follows:

Definition 3.3 The *Newton polyhedron* $\boxed{\mathcal{NP}(f)}$ of a non-zero element $f \in \mathcal{O}$ is the convex hull of the Minkowski sum of $(\mathbb{R}_{\geq 0})^n$ and the support of f . Given a face K of $\mathcal{NP}(f)$, we let $\boxed{\sigma_K}$ be the closure of the set of weight vectors w in $(\mathbb{R}_{\geq 0})^n$ supporting K (that is, such that the convex hull of the support of $\text{in}_w(f)$ is K). The set $\{\sigma_K : K \text{ face of } \mathcal{NP}(f)\}$ is the *Newton fan* $\boxed{\mathcal{NF}(f)}$ of f .

The map $K \mapsto \sigma_K$ yields an inclusion-reversing bijection between the set of faces of $\mathcal{NP}(f)$ and the Newton fan $\mathcal{NF}(f)$. Furthermore, every face K of $\mathcal{NP}(f)$ satisfies

$$\dim K + \dim \sigma_K = n. \quad (3.2)$$

Definition 3.4 Let $(Y, 0) \subseteq \mathbb{C}^n$ be a germ defined by an ideal I of $\mathcal{O}$. The *local tropicalization of I or of the germ Y* , is the set of all vectors $w \in (\mathbb{R}_{\geq 0})^n$ such that the w -initial ideal $\text{in}_w(I)\mathcal{O} \subseteq \mathcal{O}$ of I is monomial-free. We denote it by $\boxed{\text{Trop } I}$ or $\boxed{\text{Trop } Y}$. In turn, the *positive local tropicalization of I or of the germ Y* , is the intersection of the local tropicalization with the positive orthant $(\mathbb{R}_{> 0})^n$. We denote it by $\boxed{\text{Trop}_{> 0} I}$ or $\boxed{\text{Trop}_{> 0} Y}$.

Even though Definition 3.4 depends heavily on the fixed embedding $(Y, 0) \subseteq \mathbb{C}^n$, we omit it from the notation for the sake of simplicity. The next remarks clarify some differences between the present approach and that of [49], which defines local tropicalizations using local valuation spaces (see [49, Definitions 5.13 and 6.7]). A recent extension of this construction to toric prevarieties by means of Berkovich analytification can be found in [26].

Remark 3.5 As shown in [49, Theorem 11.2] and [54, Corollary 4.3], local tropicalizations of ideals in either $\mathcal{O}$ or $\hat{\mathcal{O}}$ admit several equivalent characterizations analogous to the *Fundamental Theorem of Tropical Algebraic Geometry* [28, Theorem 3.2.3]. One of them is as Euclidean closures in $(\mathbb{R}_{\geq 0})^n$ of images of local valuation spaces. By [49, Corollary 5.17], the canonical inclusion $(\mathcal{O}, \mathfrak{m}) \hookrightarrow (\hat{\mathcal{O}}, \hat{\mathfrak{m}})$ induces an isomorphism of local valuation spaces. This implies that extending an ideal in $\mathcal{O}$ to the complete ring $\hat{\mathcal{O}}$ will yield the same local tropicalization. Therefore, we can define local tropicalizations for ideals of $\mathcal{O}$ rather than of $\hat{\mathcal{O}}$, in agreement with the setting of splice type systems.

As in the global case, local tropicalizations of hypersurface germs can be obtained from the corresponding Newton fans. Indeed, if the ideal I is principal, generated by $f \in \mathcal{O}$, and $w \in (\mathbb{R}_{\geq 0})^n$, then each initial ideal $\text{in}_w(I)\mathcal{O}$ is also principal, with generator $\text{in}_w(f)$. Equality (3.2) then yields the following statement (see [49, Proposition 11.8]):

Proposition 3.6 *The set $\text{Trop}(f)$ is the union of all cones of the Newton fan of f dual to bounded edges of the Newton polyhedron of f .*




Fig. 2 Splice diagram, Newton polyhedron and local tropicalization of the E_8 surface singularity




Fig. 3 From left to right: Splice diagram, and representations of the global and local tropicalizations of Example 3.9

Example 3.7 The E_8 surface singularity is the splice type surface singularity defined by the polynomial $z_1^2 + z_2^3 + z_3^5$. Its associated splice diagram, Newton polyhedron and local tropicalization are depicted in Fig. 2. Its local tropicalization is a 2-dimensional fan with four rays spanned by e_1, e_2, e_3 , and $w_u = (15, 10, 6)$. Its three maximal cones are spanned by the pairs $\{e_i, w_u\}$ for $i \in \{1, 2, 3\}$. These three cones are dual to the three bounded edges of the Newton polyhedron.

By contrast, if I has two or more generators, their w -initial forms need not generate $\text{in}_w(I)\mathcal{O}$. However, for the purpose of characterizing $\text{Trop}_{>0} I$, it is enough to have a *tropical basis* for I in the sense of [49, Definition 10.1], i.e., a finite set of generators $\{f_1, \dots, f_s\}$ of I which is a universal standard basis of I in the sense of [49, Definition 9.8] and such that for any $w \in (\mathbb{R}_{>0})^n$, the w -initial ideal $\text{in}_w(I)\mathcal{O}$ contains a monomial if and only if one of the initial forms $\text{in}_w(f_i)$ is a scalar multiple of a monomial.

Remark 3.8 Such tropical bases exist by [49, Theorem 10.3] and can be used to determine $\text{Trop} I$ by intersecting the local tropicalizations of the corresponding hypersurface germs. Furthermore, their existence ensures that local tropicalizations are supports of rational fans in $(\mathbb{R}_{\geq 0})^n$. Indeed, the corresponding fan is obtained by considering the common refinement of the intersection of the local tropicalization of each member of a tropical basis for I combined with Proposition 3.6. Furthermore, under the hypothesis that no irreducible component of $(Y, 0)$ is included in a coordinate subspace of $\mathbb{C}^n$, such fans and their refinements are *standard tropicalizing fans* of Y in the sense of Definition 3.17 below.

Example 3.9 We consider the germ of splice type surface singularity from [39, Example 2], given by

$$\begin{cases} f_u := z_1^2 + z_2^3 + z_3 z_4 = 0, \\ f_v := z_3^5 + z_4^2 + z_1 z_2^4 = 0. \end{cases} \quad (3.3)$$

associated to the splice diagram on the left of Fig. 3. A computation with the package `Tropical.m2` [1], available in `Macaulay2` [18], determines the global tropicalization of the system (see [28, Definition 3.2.1]). It is a 2-dimensional fan in $\mathbb{R}^4$ with f -vector $(1, 6, 9)$. Its six rays are generated by the primitive vectors

$$\begin{aligned} r_1 &:= (-28, -13, -16, -40), & r_3 &:= (-2, -7, -6, -15), & r_5 &:= (21, 14, 12, 30), \\ r_2 &:= (-15, -10, -11, -19), & r_4 &:= (-6, -4, -1, -11), & r_6 &:= (30, 20, 22, 55). \end{aligned}$$

Notice that $r_5 = w_u$ and $r_6 = w_v$. Its nine top-dimensional cones are encoded by the graph depicted at the center of the figure.

The local tropicalization of the germ defined by (3.3) is obtained by intersecting the global tropicalization with the positive orthant [49, Theorem 12.10]. We indicate the positions of the four canonical basis elements $\{e_1, \dots, e_4\}$ of $\mathbb{R}^4$ with unfilled dots inside the edges of the central graph of Fig. 3. As we will see in Theorem 6.2, the local tropicalization of this germ can be obtained as the cone over the red graph depicted in the standard tetrahedron seen in the right of the figure. Note that this graph is homeomorphic to the splice diagram. This fact is general, as confirmed by Theorem 5.11.

Remark 3.10 Our choice of terminology for local tropicalizations differs slightly from [49], as we now explain. As was shown in [49, Sect. 6], the local tropicalizations of the intersection of a germ $(Y, 0) \hookrightarrow \mathbb{C}^n$ with each coordinate subspace of $\mathbb{C}^n$ can be glued together to form an *extended fan* in $(\mathbb{R}_{\geq 0} \cup \{\infty\})^n$ called the *local nonnegative tropicalization* of Y in [49]. In the present paper, we refer to this structure as an *extended tropicalization* of the germ Y , in agreement with Kashiwara and Payne's constructions for global tropicalizations (see [28, Sect. 6.2]). The finite part of this extended tropicalization (i.e., its intersection with $(\mathbb{R}_{\geq 0})^n$) is the local tropicalization from Definition 3.4. A precise description of the boundary strata of the extended local tropicalization of splice type singularities is given in Sect. 6.2.

Remark 3.11 The local tropicalization of a germ $(Y, 0) \subseteq \mathbb{C}^n$ coincides with the local tropicalization of the associated reduced germ $(Y_{red}, 0) \subseteq \mathbb{C}^n$ since the defining ideal of $(Y_{red}, 0)$ is the radical $\sqrt{I} \subseteq \mathcal{O}$ and w -initial forms respect products. As a consequence, $\text{in}_w(\sqrt{I})$ is monomial-free if and only if the same is true for $\text{in}_w(I)$. Alternatively, the same statement can be obtained from the definition of local tropicalization as the image of the local valuation space of Y and the fact that the embedding $Y_{red} \hookrightarrow Y$ induces a homeomorphism of local valuation spaces (see [49, Lemma 5.18]).

Remark 3.12 The local tropicalization of a reduced germ $(Y, 0) \subseteq \mathbb{C}^n$ is equal to the union of the local tropicalizations of its irreducible components, and the same is true for their local positive tropicalizations. This is a direct consequence of the fact that the local valuation space of $(Y, 0)$ is the union of the local valuation spaces of its irreducible components (see [49, Lemma 5.18]).

The next result determines the local tropicalization of a germ $(Y, 0) \hookrightarrow \mathbb{C}^n$ from the positive one:

Proposition 3.13 *The local tropicalization $\text{Trop } Y$ is the closure of the positive local tropicalization $\text{Trop}_{>0} Y$ inside the cone $(\mathbb{R}_{\geq 0})^n$.*

Proof By Remarks 3.11 and 3.12, it suffices to consider the case where $(Y, 0)$ is irreducible. In this situation, [49, Theorem 11.9] shows that the extended local tropicalization of $(Y, 0)$ is the closure of the extended positive local tropicalization in the extended non-negative orthant $(\mathbb{R}_{\geq 0} \cup \{\infty\})^n$. Applying Lemma 3.14 below to the case when E is the extended local tropicalization of $(\mathbb{C}^n, 0)$, $U = (\mathbb{R}_{\geq 0})^n$ and A is the extended local tropicalization of $(Y, 0)$ confirms the claim about the closure of $\text{Trop}_{>0} Y$ in $(\mathbb{R}_{\geq 0})^n$. $\square$

The following lemma is a standard statement in general set topology, which may for instance be obtained as an immediate consequence of [30, Theorem 17.4]:

Lemma 3.14 *Let E be a topological space and U be an open subset. Then, for any subset A of E we have:*

$$\text{cl}_U(A \cap U) = \text{cl}_E(A) \cap U,$$

where $\text{cl}_E(A)$ is the closure of A in E and $\text{cl}_U(A \cap U)$ denotes the closure of $A \cap U$ in U .

In what follows, we restrict our attention to subgerms of $(\mathbb{C}^n, 0)$ with no components included in coordinate subspaces. Their local tropicalizations verify the following key property (see [49, Proposition 9.21, Theorems 10.3 and 11.9]):

Proposition 3.15 *Let $(Y, 0)$ be a subgerm of $(\mathbb{C}^n, 0)$ with no irreducible components contained in coordinate subspaces of $\mathbb{C}^n$, and let $I \subseteq \mathcal{O}$ be its defining ideal. Then, $\text{Trop } Y$ is the support of a rational polyhedral fan $\mathcal{F}$ which satisfies the following conditions:*

1. *the dimension of all the maximal cones of $\mathcal{F}$ agrees with the complex dimension of Y ;*
2. *the maximal cones of $\mathcal{F}$ have non-empty intersections with $(\mathbb{R}_{>0})^n$;*
3. *given any cone τ of $\mathcal{F}$, the w -initial ideal of I is independent of the choice of $w \in \tau^\circ$.*

Proof By Remarks 3.11 and 3.12, we may restrict to the case when $(Y, 0)$ is reduced and irreducible. The existence of $\mathcal{F}$ satisfying the first two conditions is a direct consequence of [49, Theorem 11.9], which ensures the analogous properties hold for the extended local tropicalization of $(Y, 0)$. Note that $\text{Trop } Y$ is non-empty, since $(Y, 0)$ meets the dense torus of $\mathbb{C}^n$ (see [49, Lemma 7.4]). Furthermore, the fan structure on $\text{Trop } Y$ induced from a tropical basis $\{f_1, \dots, f_r\}$ for I discussed in Remark 3.8 satisfies condition (3). In turn, any refinement will satisfy conditions (1) and (2) as well. $\square$

Remark 3.16 Proposition 3.15 allows us to recover the complex dimension of an irreducible germ $(Y, 0) \hookrightarrow \mathbb{C}^n$ meeting the dense torus from its positive tropicalization (see, for instance, the proofs of Corollaries 6.20 and 6.21). Indeed, its dimension agrees with the dimension of any of the top-dimensional cones in any fixed fan satisfying conditions (1)–(3) in Proposition 3.15. If $(Y, 0)$ is not irreducible, a similar procedure determines the maximal dimension of a component of the germ meeting the dense torus.

Definition 3.17 Let $(Y, 0)$ be a subgerm of $\mathbb{C}^n$ with no irreducible component contained in a coordinate subspace of $\mathbb{C}^n$. Let I be the ideal of $\mathcal{O}$ defining $(Y, 0)$. Any fan $\mathcal{F}$ satisfying all three conditions in Proposition 3.15 is called a *standard tropicalizing fan* for $(Y, 0)$ or for I . For every cone τ of $\mathcal{F}$ meeting $(\mathbb{R}_{>0})^n$ and any $w \in \tau^\circ$, we write $\boxed{\text{in}_\tau(I)} := \text{in}_w(I)$ for the associated initial ideal in the polynomial ring $\mathbb{C}[z_1, \dots, z_n]$ and $\boxed{\text{in}_\tau(Y)} := \text{in}_w(Y)$ for the associated subscheme of $\mathbb{C}^n$ (see Definition 3.1).

Remark 3.18 When Y is a hypersurface germ defined by a series $f \in \mathcal{O}$, the set of cones of codimension one of the Newton fan of f dual to bounded edges of the Newton polyhedron of f satisfies all three conditions listed in Proposition 3.15. Furthermore, it is the coarsest fan with these properties. However, for germs of higher codimension such canonical choice need not always exist. For an example in the global (i.e., polynomial) setting, we refer the reader to [28, Example 3.5.4].

The next proposition emphasizes the relevance of tropicalizing fans for producing birational models of irreducible germs with desirable geometric properties, in the spirit of Tevelev's construction of tropical compactifications of subvarieties of tori [56].

Proposition 3.19 Fix a rational polyhedral fan $\mathcal{F}$ contained in $(\mathbb{R}_{\geq 0})^n$ and let $\pi_{\mathcal{F}}: \mathcal{X}_{\mathcal{F}} \rightarrow \mathbb{C}^n$ be the associated toric morphism. Given an irreducible germ $(Y, 0) \hookrightarrow \mathbb{C}^n$ meeting the dense torus, let $Y_{\mathcal{F}}$ be the strict transform of Y under $\pi_{\mathcal{F}}$ and write $\pi: Y_{\mathcal{F}} \rightarrow Y$ for the restriction of the map $\pi_{\mathcal{F}}$ to $Y_{\mathcal{F}}$. Then, the following properties hold:

1. The restriction π is proper if and only if the support $|\mathcal{F}|$ contains the local tropicalization $\text{Trop } Y$.
2. Assume that π is proper. Then, the strict transform $Y_{\mathcal{F}}$ intersects every orbit S of $\mathcal{X}_{\mathcal{F}}$ along a non-empty pure-dimensional subvariety with $\text{codim}_{Y_{\mathcal{F}}}(Y_{\mathcal{F}} \cap S) = \text{codim}_{\mathcal{X}_{\mathcal{F}}}(S)$ if and only if $|\mathcal{F}| = \text{Trop } Y$.

Proof In what follows, we use standard terminology and notation from toric geometry, which can be found in Fulton's book [12]. The proof of (2) is similar to the global analog [28, Proposition 6.4.7 (2)], so we leave it to the reader.

It remains to prove assertion (1). To this end, we consider a fan Σ subdividing the non-negative cone $(\mathbb{R}_{\geq 0})^n$ and containing $\mathcal{F}$ as a subfan. Such a fan exists by [11, Theorem 2.8 (III.2)].

We consider the toric varieties $\mathcal{X}_{\mathcal{F}}$ and $\mathcal{X}_{\Sigma}$ associated to the fans $\mathcal{F}$ and Σ and the natural toric morphisms $\pi_{\mathcal{F}}: \mathcal{X}_{\mathcal{F}} \rightarrow \mathbb{C}^n$ and $\pi_{\Sigma}: \mathcal{X}_{\Sigma} \rightarrow \mathbb{C}^n$. We let Y_{Σ} and $Y_{\mathcal{F}}$ be the strict transforms of Y under these two maps. The aforementioned varieties and maps fit naturally into the commutative diagram

$$\begin{array}{ccccccc}
 Y_\Sigma & \hookrightarrow & \mathcal{X}_\Sigma & \xleftarrow{\quad} & \mathcal{X}_\mathcal{F} & \xleftarrow{\quad} & Y_\mathcal{F} \\
 & & \searrow \pi_\Sigma & & \swarrow \pi_\mathcal{F} & & \swarrow \pi \\
 & & & \mathbb{C}^n & \xleftarrow{\quad} & Y &
 \end{array}$$

where the central triangle involves toric morphisms and the horizontal arrows are embeddings. The vertical map π on the right is the restriction of $\pi_\mathcal{F}$ to $Y_\mathcal{F}$. In what follows we view $\mathcal{X}_\mathcal{F}$ as an open subvariety of $\mathcal{X}_\Sigma$. Note that the toric birational morphism π_Σ is proper by [12, Sect. 2.4], as the defining fans of its source and target have the same support, namely $(\mathbb{R}_{\geq 0})^n$.

By construction, π is proper if and only if Y_Σ is contained in $\mathcal{X}_\mathcal{F}$. Thus, claim (1) will follow if we show that for every cone τ of Σ we have the following equivalence:

$$\mathcal{O}_\tau \cap Y_\Sigma \neq \emptyset \iff \tau^\circ \cap \text{Trop } Y \neq \emptyset, \quad (3.4)$$

where $\mathcal{O}_\tau := \text{Hom}_{\text{monoid}}(\tau^\perp \cap M, \mathbb{C})$ is the corresponding toric orbit.

It remains to prove (3.4). We start with the forward implication and fix $y_0 \in \mathcal{O}_\tau \cap Y_\Sigma$. Then, there exists a holomorphic arc in $Y \cap (\mathbb{C}^*)^n$ parameterized as $t \mapsto y(t)$ such that the limit as $t \rightarrow 0$ of its strict transform in Y_Σ equals y_0 , i.e.,

$$\lim_{t \rightarrow 0} y(t) = y_0 \in \mathcal{O}_\tau \cap Y_\Sigma.$$

Such an arc can be built by choosing an irreducible subgerm of a curve C_Σ of the germ (Y_Σ, y_0) , not contained in the toric boundary, then by projecting it to a subgerm C of Y via π_Σ and, finally, by choosing a normalization of C , which we identify with $(\mathbb{C}, 0)$.

We consider the weight vector $w := \text{ord } y(t)$ in the dual lattice $N := M^\vee$ recording the orders of vanishing of the components of $y(t)$. We claim that w belongs to $\tau^\circ \cap \text{Trop } Y$, so $\tau^\circ \cap \text{Trop } Y \neq \emptyset$, as desired. To prove this claim, notice that the arc $t \mapsto y_{\text{in}}(t)$ of Y_Σ obtained by keeping the w -initial terms $y_{\text{in}}(t)$ of the components of $y(t)$ has the same limit when $t \rightarrow 0$ as $y(t)$ does. This fact can be checked by working in the affine toric variety associated to the cone $\mathbb{R}_{\geq 0}\langle w \rangle$. Properties of limits in toric varieties from [12, Sect. 2.3] ensure that $\lim_{t \rightarrow 0} y_{\text{in}}(t) \in \mathcal{O}_\tau$ is equivalent to the fact that the weight vector w lies in $\tau^\circ \cap N$. Therefore, we conclude that $\text{ord } y(t) \in \tau^\circ$.

It remains to verify that $w \in \text{Trop } Y$. To do so, it suffices to notice that $\text{Trop } C \subseteq \text{Trop } Y$ (since $C \subseteq Y$) and that $\text{Trop } C = \mathbb{R}_{\geq 0}\langle w \rangle$. The latter is a direct consequence of [54, Corollary 4.3 3'')] (see also Maurer's paper [29] which includes a precursor of local tropicalization for germs of space curves).

Finally, to prove the reverse implication of (3.4), we assume that $\tau^\circ \cap \text{Trop } Y \neq \emptyset$ and let $w \in \tau^\circ \cap \text{Trop } Y$ be a primitive lattice vector in $N \simeq \mathbb{Z}^n$. We consider a refinement Σ_w of Σ such that the ray $\tau_w = \mathbb{R}_{\geq 0}\langle w \rangle \in \Sigma_w$. By construction, the orbit $\mathcal{O}_{\tau_w}$ is mapped via the toric morphism $\pi_w: \mathcal{X}_{\Sigma_w} \rightarrow \mathcal{X}_\Sigma$ to the orbit $\mathcal{O}_\tau$.

The intersection of Y_{Σ_w} with the orbit $\mathcal{O}_{\tau_w}$ is determined by the w -initial ideal $\text{in}_w(I(Y))$ of the ideal $I(Y)$ defining Y , viewed in the Laurent polynomial ring. Since this initial ideal is monomial free because $w \in \text{Trop } Y$, this intersection is non-empty.

Since $\mathcal{O}_{\tau_w} \subseteq Y_{\Sigma_w}$, the map π_w ensures that $Y_{\Sigma} \cap \mathcal{O}_{\tau} \neq \emptyset$ as well. This concludes our proof. $\square$

One well-known feature of global tropicalizations of equidimensional subvarieties of toric varieties is the so-called *balancing condition* [28, Sect. 3.3]. To this end, tropical varieties must be endowed with positive integer weights (called *tropical multiplicities*) along their top-dimensional cones. Such multiplicities may be also defined in the local situation. We restrict the exposition to equidimensional germs, since this is sufficient for the purposes of this paper.

Definition 3.20 Let $(Y, 0) \hookrightarrow \mathbb{C}^n$ be an equidimensional germ meeting $(\mathbb{C}^*)^n$, defined by an ideal I of $\mathcal{O}$ and let $\mathcal{F}$ be a standard tropicalizing fan for it. Given a top-dimensional cone τ of $\mathcal{F}$, we define the *tropical multiplicity* of $\mathcal{F}$ at τ to be the number of irreducible components of $\text{in}_{\tau}(Y) \cap (\mathbb{C}^*)^n$, counted with multiplicity.

In order to state the balancing condition for local tropicalization, we first define the notion of a pure rational weighted balanced fan. Recall that a fan is pure if all its top-dimensional cells have the same dimension.

Definition 3.21 Consider a pure rational polyhedral fan $\mathcal{F}$ in $\mathbb{R}^n$ relative to the lattice $\mathbb{Z}^n$, with positive integer weights on its maximal cones. Fix a cone τ of $\mathcal{F}$ of codimension one in $\mathcal{F}$, and let $\sigma_1, \dots, \sigma_s$ be the maximal cones of $\mathcal{F}$ containing τ as a face. Denote by $m_1, \dots, m_s$ the corresponding weights. For each $i \in \{1, \dots, s\}$ consider a vector $\omega_{\sigma_i|\tau} \in \mathbb{Z}^n \cap \sigma_i$ generating the lattice

$$\Lambda_i := \frac{\mathbb{Z}^n \cap \mathbb{R}\langle \sigma_i \rangle}{\mathbb{Z}^n \cap \mathbb{R}\langle \tau \rangle}. \quad (3.5)$$

We say that $\mathcal{F}$ is *balanced at τ* if $\sum_{i=1}^s m_i \omega_{\sigma_i|\tau} \in \mathbb{Z}^n \cap \mathbb{R}\langle \tau \rangle$. The fan $\mathcal{F}$ is *balanced* if it is balanced at each of its codimension one cones.

Remark 3.22 By construction, the lattice Λ_i in (3.5) is free of rank one. It has one generator, up to sign. Even though there are many choices for $\omega_{\sigma_i|\tau}$, their projections onto Λ_i give the same generator of the lattice.

Our last statement in this section confirms that the balancing condition holds for local tropicalizations of equidimensional germs in $\mathbb{C}^n$ meeting the dense torus $(\mathbb{C}^*)^n$. Its validity follows from [49, Remark 11.3, Theorem 12.10]:

Theorem 3.23 Let $(Y, 0) \hookrightarrow \mathbb{C}^n$ be an equidimensional germ meeting $(\mathbb{C}^*)^n$ and let $\mathcal{F}$ be any refinement of a standard tropicalizing fan for Y . Then, $\mathcal{F}$ is a balanced fan when endowed with tropical multiplicities as in Definition 3.20.

Remark 3.24 Balancing for local tropicalizations of equidimensional germs of surfaces features in the proof of Proposition 6.26. This property will help us prove that the embedded splice diagrams in $\mathbb{R}^n$ are included in the local tropicalizations of the corresponding splice type systems (see Sect. 6.3). Tropical multiplicities will be also used in Sect. 8 to recover the edge weights on any coprime splice diagram Γ from the local tropicalization of any splice type surface singularity associated to it.

4 Newton non-degeneracy

In this section we discuss the notion of Newton non-degenerate complete intersection in the sense of Khovanskii [22], starting with the case of formal power series in $\hat{\mathcal{O}}$, as introduced by Kouchnirenko in [24, Sect. 8] and [25, Definition 1.19]. Kouchnirenko's definition was later extended by Steenbrink [53, Definition 5] to $\mathbb{C}$ -algebras of formal power series $\mathbb{C}[[P]]$ with exponents on an arbitrary saturated sharp toric monoid P . For precursors to this notion we refer the reader to Teissier's work [55, Sect. 5].

Definition 4.1 Given a series $f \in \hat{\mathcal{O}}$, we say that f is *Newton non-degenerate* if for any positive weight vector $w \in (\mathbb{R}_{>0})^n$, the subvariety of the dense torus $(\mathbb{C}^*)^n$ defined by $\text{in}_w(f)$ is smooth.

Example 4.2 We consider the E_8 singularity from Example 3.7, defined by $f = z_1^2 + z_2^3 + z_3^5$. Its local tropicalization is depicted in Fig. 2. Since $\text{in}_w(f)$ is a monomial whenever $w \notin \text{Trop}_{>0} Z(f)$, the Newton non-degeneracy condition only needs to be checked for $w \in \text{Trop}_{>0} Z(f)$.

The calculations are simplified since $\text{Trop}_{>0} Z(f)$ is a subfan of the normal fan to the Newton polyhedron of f . If $w \in \mathbb{R}_{>0}\langle w_u \rangle$, we have $\text{in}_w(f) = f$. In turn, weight vectors in the relative interiors of maximal cones of $\text{Trop}_{>0} Z(f)$ satisfy

$$\text{in}_w(f) = \begin{cases} z_2^3 + z_3^5 & \text{if } w \in \mathbb{R}_{>0}\langle e_1, w_u \rangle, \\ z_1^2 + z_3^5 & \text{if } w \in \mathbb{R}_{>0}\langle e_2, w_u \rangle, \\ z_1^2 + z_2^3 & \text{if } w \in \mathbb{R}_{>0}\langle e_3, w_u \rangle. \end{cases}$$

All four initial forms describe surfaces that are smooth when restricted to the torus $(\mathbb{C}^*)^3$. Thus, $Z(f)$ is Newton non-degenerate.

The notion of Newton non-degeneracy extends naturally to finite sequences of functions. For our purposes, it suffices to restrict ourselves to *regular sequences*, i.e., collections $(f_1, \dots, f_s)$ in $\hat{\mathcal{O}}$ where

1. f_1 is not a zero divisor of $\hat{\mathcal{O}}$, and
2. for each $i \in \{1, \dots, s-1\}$, the element f_{i+1} is not a zero divisor in the quotient ring $\hat{\mathcal{O}}/\langle f_1, \dots, f_i \rangle \hat{\mathcal{O}}$.

As $\hat{\mathcal{O}}$ is a regular local ring, the germs defined by regular sequences in $\hat{\mathcal{O}}$ are exactly formal complete intersections at the origin of $\mathbb{C}^n$.

Definition 4.3 Fix a positive integer s and a regular sequence $(f_1, \dots, f_s)$ in $\hat{\mathcal{O}}$. Consider the germ $(Y, 0) \hookrightarrow \mathbb{C}^n$ defined by the ideal $\langle f_1, \dots, f_s \rangle \hat{\mathcal{O}}$. The sequence $(f_1, \dots, f_s)$ is a *Newton non-degenerate complete intersection system* for $(Y, 0)$ if for any positive weight vector $w \in (\mathbb{R}_{>0})^n$, the hypersurfaces of $(\mathbb{C}^*)^n$ defined by each $\text{in}_w(f_i)$ form a normal crossings divisor in a neighborhood of their intersection. Equivalently, the differentials of the initial forms $\text{in}_w(f_1), \dots, \text{in}_w(f_s)$ must be linearly independent at each point of this intersection.

Remark 4.4 Notice that Definition 4.3 allows for the initial forms to be monomials. This would determine an empty intersection with the dense torus $(\mathbb{C}^*)^n$.

Our main result in this section is a useful tool for proving that a regular sequence in $\hat{\mathcal{O}}$ defines a Newton non-degenerate system. We make use of this statement in Theorem 7.3:

Proposition 4.5 *Let s be a positive integer, $(f_1, \dots, f_s)$ be a sequence in $\hat{\mathcal{O}}$ and w be a weight vector in $(\mathbb{R}_{>0})^n$. Assume that $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is a regular sequence in $\mathbb{C}[z_1, \dots, z_n]$ defining a subscheme Z_w of $\mathbb{C}^n$. If p is a smooth point of $Z_w \cap (\mathbb{C}^*)^n$, then the differentials at p of the initial forms $\text{in}_w(f_1), \dots, \text{in}_w(f_s)$ are linearly independent.*

Proof Fix any point $p \in Z_w \subseteq (\mathbb{C}^*)^n$. Our regularity hypothesis implies that $\dim(Z_w) = n - s$. Since p is a smooth point of Z_w , by [8, Lemma 4.3.5] we know that $\mathcal{O}_{Z_w, p}$ is a regular local ring of dimension $\dim(Z_w)$. Furthermore, this quantity equals the embedding dimension $\text{edim}(\mathcal{O}_{Z_w, p})$ of the local ring $\mathcal{O}_{Z_w, p}$.

In turn, the Jacobian criterion for smoothness (see, e.g., [8, Theorem 4.3.6]) enables us to compute the rank of the $n \times s$ Jacobian matrix of the initial forms $\{\text{in}_w(f_i)\}_{i=1}^s$ evaluated at p , in terms of n and this embedding dimension. More precisely, we have

$$\text{rk}_p(\text{Jac}(\text{in}_w(f_1), \dots, \text{in}_w(f_s))) = n - \text{edim}(\mathcal{O}_{Z_w, p}) = n - (n - s) = s.$$

This implies that the differentials of the initial forms $\text{in}_w(f_1), \dots, \text{in}_w(f_s)$ are linearly independent at p . $\square$

Example 4.6 We consider the splice type system from Example 3.9. As claimed by Theorem 1.1, the germ defined by it is Newton non-degenerate. By Remark 4.4 we need only focus on weights $w \in \text{Trop}_{>0}\langle f_u, f_v \rangle \mathcal{O}$. A direct computation using Lemma 2.19 shows that the initial forms of f_u and f_v are constant along the relative interiors of maximal cones of $\text{Trop}_{>0}\langle f_u, f_v \rangle \mathcal{O}$. We check that the Newton non-degeneracy condition is satisfied for w in two of the five maximal cones. The remaining three cases are similar.

Pick $w \in \mathbb{R}_{>0}\langle w_u, w_v \rangle$. Then, $\text{in}_w(f_u) = z_1^2 + z_2^3$ and $\text{in}_w(f_v) = z_3^5 + z_4^2$. Their differentials are $2z_1 dz_1 + 3z_2^2 dz_2$ and $5z_3^4 dz_3 + 2z_4 dz_4$, so they are linearly independent everywhere in the torus $(\mathbb{C}^*)^4$. Similarly, if $w \in \mathbb{R}_{>0}\langle e_1, w_u \rangle$, we have $\text{in}_w(f_u) = z_2^3 + z_3 z_4$ and $\text{in}_w(f_v) = z_3^5 + z_4^2$. Their differentials are $3z_2^2 dz_2 + z_4 dz_3 + z_3 dz_4$ and $5z_3^4 dz_3 + 2z_4 dz_4$. They are also linearly independent everywhere in $(\mathbb{C}^*)^4$.

Remark 4.7 Definition 4.3 modifies slightly Khovanskii's original definition from [22, Remark 4 of Sect. 2.4], by imposing the regularity of the sequence $(f_1, \dots, f_s)$. The generalization to formal power series $\mathbb{C}[[P]]$ associated to an arbitrary sharp toric monoid P is straightforward, and parallels that done by Steenbrink [53] for hyper-surfaces. A slightly different notion of *Newton non-degenerate ideals* was introduced by Saia in [50] for ideals of finite codimension in $\hat{\mathcal{O}}$. For a comparison with Khovanskii's approach we refer the reader to Bivià-Ausina's work [5, Lemma 6.8]. For a general perspective on Newton non-degenerate complete intersections, the reader can consult Oka's book [41]. A definition of *Newton non-degenerate algebraic subgerms* of $(\mathbb{C}^n, 0)$ for not necessarily complete intersections was given by Aroca, Gómez-Morales and Shabbir [2]. The results of their paper (and the proofs involving Gröbner

bases) can be extended to the analytic and formal contexts by working with standard bases, in the spirit of [49]. Recent work of Aroca, Gómez-Morales and Mourtada [3] generalize the constructions from [2] to subgerms of arbitrary normal affine toric varieties.

5 Embeddings and convexity properties of splice diagrams

In this section, we describe simplicial fans in the real weight space $N(\partial \Gamma) \otimes_{\mathbb{Z}} \mathbb{R} \simeq \mathbb{R}^n$ that arise from splice diagrams and appropriate subdiagrams. These constructions will play a central role in Sect. 6 when characterizing local tropicalizations of splice type surface singularities. Throughout this section we assume that the splice diagram Γ satisfies the edge determinant condition of Definition 2.9. The semigroup condition from Definition 2.13 plays no role.

We let $\boxed{\Delta_{n-1}}$ be the standard $(n - 1)$ -simplex of the real weight space $\mathbb{R}^n$, with vertices $\{w_\lambda : \lambda \in \partial \Gamma\}$. We start by defining a piecewise linear map from Γ to Δ_{n-1} :

$$\boxed{\iota} : \Gamma \rightarrow \Delta_{n-1} \quad \text{where} \quad \iota(v) = \frac{w_v}{|w_v|} \quad \text{for each vertex } v \text{ of } \Gamma. \quad (5.1)$$

Here, $|\cdot|$ denotes the L^1 -norm in $\mathbb{R}^n$. In particular, $|w_v| = \sum_{\lambda \in \partial \Gamma} \ell_{v,\lambda}$ for each node v of Γ . After identifying each edge e of Γ with the interval $[0, 1]$, the map ι on e is defined by convex combinations of the assignment at its endpoints. The injectivity of ι will be discussed in Theorem 5.11.

The following combinatorial constructions, in particular Definitions 5.1, 5.3 and 5.4, play a prominent role in proving Theorem 6.2. Stars of vertices (see Definition 2.1) and convex hulls of vertices (see Definition 2.3) are central to many arguments below.

Definition 5.1 A subtree T of the splice diagram Γ is *star-full* if $\text{Star}_\Gamma(v) \subseteq T$ for every node v of T . A node of T is called an *end-node* if it is adjacent to exactly one other node of T .

Every tree with two or more nodes contains at least two end-nodes. The following statement, illustrated in Fig. 4, describes a method to produce new star-full subtrees from old ones by *pruning from an end-node*. Its proof is straightforward, so we omit it.

Lemma 5.2 Let T be a star-full subtree of Γ with d leaves $\{u_1, \dots, u_d\}$. Fix an end-node v of T and assume that $u_1, \dots, u_{\delta_v-1}$ are the only leaves of T adjacent to v . Then, the convex hull $T' := [v, u_{\delta_v}, \dots, u_d]$ is also star-full.

Definition 5.3 A *branch* of a tree T adjacent to a node v is a connected component of

$$T \setminus \left(\{v\} \cup \bigcup_{e \in \text{Star}_\Gamma(v)} e^\circ \right),$$

where $\boxed{e^\circ}$ denotes the interior of the edge e .




Fig. 4 Pruning a star-full subtree T of Γ from an end-node v of T produces a new star-full subtree T' with one fewer node, as in Lemma 5.2. Here, $s = \delta_v - 1$ and $s' = \delta_{v'} - 1$

For example, $\{u_1\}, \dots, \{u_s\}$ and the convex hull $[v', T_1, \dots, T_{s'}]$ constructed inside the tree T on the left of Fig. 4 are the δ_v branches of T adjacent to the node v . Similarly, the branches of T' adjacent to v' are $T_1, \dots, T_{s'}$ and $\{v\}$.

Star-full subtrees of splice diagrams have a key convexity property rooted in barycentric calculus. This is the content of Proposition 5.10. The following definition plays a crucial role in its proof.

Definition 5.4 Let v be a node of Γ and let L be a set of leaves of Γ . We define $\text{Bar}(L; v)$ as the *barycenter* of the leaves in L with weights determined by w_v , that is:

$$\boxed{\text{Bar}(L; v)} := \sum_{\lambda \in L} \frac{\ell_{v,\lambda}}{\ell} w_\lambda \quad \text{where} \quad \ell := \sum_{\lambda \in L} \ell_{v,\lambda}. \quad (5.2)$$

In particular, $\text{Bar}(\{\lambda\}; v) = w_\lambda$ for any leaf λ .

Remark 5.5 Fix a node v of Γ with adjacent branches $\Gamma_1, \dots, \Gamma_{\delta_v}$. Then, the set $\{\text{Bar}(\partial \Gamma_i; v) : i = 1, \dots, \delta_v\}$ is linearly independent, and a direct computation gives

$$\iota(v) = \sum_{j=1}^{\delta_v} \left(\frac{1}{|w_v|} \sum_{\lambda \in \partial \Gamma_j} \ell_{v,\lambda} \right) \text{Bar}(\partial \Gamma_j; v). \quad (5.3)$$

In particular, $\iota(v)$ lies in the relative interior of the simplex $\text{conv}(\{\text{Bar}(\partial \Gamma_j; v) : j = 1, \dots, \delta_v\})$, where conv denotes the affine convex hull inside Δ_{n-1} .

Following the notation from Definition 2.13, we write $\boxed{\partial_a \Gamma} := \partial_{b,[a,b]} \Gamma$ and $\boxed{\partial_b \Gamma} := \partial_{a,[a,b]} \Gamma$ for each pair of adjacent nodes a, b of Γ . Barycenters determined by splitting Γ along the edge $[a, b]$ are closely related, as the following lemma shows:

Lemma 5.6 Let a, b be two adjacent nodes of Γ , with associated sets of leaves $\partial_a \Gamma$ and $\partial_b \Gamma$ on each side of Γ . Then:

1. $\text{Bar}(\partial_a \Gamma; a) = \text{Bar}(\partial_a \Gamma; b)$ and $\text{Bar}(\partial_b \Gamma; a) = \text{Bar}(\partial_b \Gamma; b)$ (which we denote by $\boxed{\text{Bar}(\partial_a \Gamma; [a, b])}$ and $\boxed{\text{Bar}(\partial_b \Gamma; [a, b])}$, respectively).
2. The points $\iota(a)$ and $\iota(b)$ lie in the line segment $[\text{Bar}(\partial_a \Gamma; [a, b]), \text{Bar}(\partial_b \Gamma; [a, b])]$.

3. $\text{Bar}(\partial_a \Gamma; [a, b]) < \iota(a) < \iota(b) < \text{Bar}(\partial_b \Gamma; [a, b])$, where $<$ is the order given by identifying the segment $[\text{Bar}(\partial_a \Gamma; [a, b]), \text{Bar}(\partial_b \Gamma; [a, b])]$ with $[0, 1]$.

Proof We start by showing (1). Definition 2.6 induces the following identities:

$$\ell_{b,\lambda} = \ell_{a,\lambda} \frac{\ell_{a,b}}{d_a} \quad \text{for all } \lambda \in \partial_a \Gamma, \quad \text{and} \quad \ell_{a,\mu} = \ell_{b,\mu} \frac{\ell_{a,b}}{d_b} \quad \text{for all } \mu \in \partial_b \Gamma. \quad (5.4)$$

Thus, a and b contribute proportional weights to $\partial_a \Gamma$ and $\partial_b \Gamma$, respectively. This implies that the corresponding barycenters agree, proving (1).

Next, we discuss (2). To simplify notation, we write $w = \text{Bar}(\partial_a \Gamma; [a, b])$ and $w' = \text{Bar}(\partial_b \Gamma; [a, b])$. Then, (5.4) yields:

$$\begin{aligned} w_a &= \left(\sum_{\lambda \in \partial_a \Gamma} \ell_{a,\lambda} \right) w + \left(\sum_{\mu \in \partial_b \Gamma} \ell_{a,\mu} \right) w' \\ \text{and} \quad w_b &= \left(\sum_{\lambda \in \partial_a \Gamma} \ell_{b,\lambda} \right) w + \left(\sum_{\mu \in \partial_b \Gamma} \ell_{b,\mu} \right) w'. \end{aligned}$$

The definition of ι then confirms that $\iota(a)$ and $\iota(b)$ are convex combinations of w and w' .

It remains to prove (3). The condition $w < \iota(a) < \iota(b) < w'$ claimed in (3) is equivalent to:

$$\left(\sum_{\lambda \in \partial_a \Gamma} \ell_{a,\lambda} \right) \left(\sum_{\mu \in \partial_b \Gamma} \ell_{a,\mu} \right)^{-1} > \left(\sum_{\lambda \in \partial_a \Gamma} \ell_{b,\lambda} \right) \left(\sum_{\mu \in \partial_b \Gamma} \ell_{b,\mu} \right)^{-1}.$$

Rearranging these expressions by sums with common indexing sets and simplifying further using (5.4) yields the equivalent identity:

$$1 > \left(\sum_{\lambda \in \partial_a \Gamma} \ell_{b,\lambda} \right) \left(\sum_{\lambda \in \partial_a \Gamma} \ell_{a,\lambda} \right)^{-1} \left(\sum_{\mu \in \partial_b \Gamma} \ell_{a,\mu} \right) \left(\sum_{\mu \in \partial_b \Gamma} \ell_{b,\mu} \right)^{-1} = \frac{\ell_{a,b}}{d_a} \frac{\ell_{a,b}}{d_b}. \quad (5.5)$$

The edge determinant condition for $[a, b]$ confirms the validity of (5.5), and so (3) holds. $\square$

Our next result shows that the image under ι of the vertices adjacent to a fixed node v satisfy a convexity property analogous to that of Remark 5.5:

Proposition 5.7 *Let v be a node of Γ , with adjacent vertices $u_1, \dots, u_{\delta_v}$. Then:*

1. $\{\iota(u_i) : i = 1, \dots, \delta_v\}$ is linearly independent;
2. $\iota(v) \in (\text{conv}(\{\iota(u_j) : j = 1, \dots, \delta_v\}))^\circ$.

Proof Both statements are clear if v is only adjacent to leaves of Γ (i.e., when Γ is a star splice diagram), so we may assume v is adjacent to some node of Γ . Thus, up to relabeling if necessary, we suppose $\{u_1, \dots, u_s\}$ are leaves of Γ and $\{u_{s+1}, \dots, u_{\delta_v}\}$ are nodes of Γ (we set $s = 0$ if v is only adjacent to nodes).

For each $j \in \{1, \dots, \delta_v\}$, we let Γ_j be the branch of Γ adjacent to v containing u_j , and set $w_j := \text{Bar}(\partial \Gamma_j; v)$. Lemma 5.6 (2) and the definition of barycenters ensures the existence of $\alpha_j \in (0, 1]$ satisfying

$$\iota(u_j) = \alpha_j w_j + (1 - \alpha_j) \iota(v) \quad \text{for each } j \in \{1, \dots, \delta_v\}, \quad (5.6)$$

with $\alpha_j = 1$ if and only if $j \in \{1, \dots, s\}$.

We start by proving (1). We fix a linear relation $\sum_{j=1}^{\delta_v} \beta_j \iota(u_j) = 0$. Substituting (5.6) into this dependency relation yields

$$\sum_{j=1}^{\delta_v} (\beta_j \alpha_j) w_j + B \iota(v) = 0 \quad \text{where} \quad B := \sum_{j=1}^{\delta_v} \beta_j (1 - \alpha_j). \quad (5.7)$$

We claim that $B = 0$. Indeed, assuming this is not the case, we use (5.7) to write $\iota(v)$ in terms of $w_1, \dots, w_{\delta_v}$. Comparing this expression with (5.3) and using the linear independence of $\{w_1, \dots, w_{\delta_v}\}$ gives:

$$\beta_j \alpha_j = -\frac{B}{|w_v|} \sum_{\lambda \in \partial T_j} \ell_{v,\lambda} \quad \text{for each } j \in \{1, \dots, \delta_v\}. \quad (5.8)$$

In particular, all β_j are non-zero and have the same sign, namely the opposite sign to B . Summing up the expressions (5.8) over all j yields $\sum_{j=1}^{\delta_v} \beta_j = 0$, which cannot happen due to the sign constraint on the β_j 's. From this it follows that $B = 0$.

Since $B = 0$, the linear independence of $\{w_1, \dots, w_{\delta_v}\}$ forces $\beta_j \alpha_j = 0$ for all j . Combining this with our assumption that $\alpha_j > 0$ for all j , gives $\beta_j = 0$ for all $j = 1, \dots, \delta_v$, thus confirms (1).

To finish, we discuss (2). We let $q_1, \dots, q_{\delta_v}$ be the coefficients used in (5.3) to write $\iota(v)$ as a convex combination of $w_1, \dots, w_{\delta_v}$. Substituting the value of each w_j obtained from (5.6) in expression (5.3) yields:

$$\iota(v) = \sum_{j=1}^s \frac{q_j}{A} \iota(u_j) + \sum_{j=s+1}^{\delta_v} \frac{q_j}{A \alpha_j} \iota(u_j) \quad \text{where} \quad A := 1 + \sum_{j=s+1}^{\delta_v} \frac{q_j (1 - \alpha_j)}{\alpha_j}. \quad (5.9)$$

The conditions $0 < \alpha_j < 1$ for $j \in \{s+1, \dots, \delta_v\}$, and the definition of $q_1, \dots, q_{\delta_v}$ ensure that the right-hand side of (5.9) is a positive convex combination of $\iota(u_1), \dots, \iota(u_{\delta_v})$, as we wanted to show. $\square$

Each subtree T of Γ determines a polytope via the map ι :

$$\boxed{\Delta_T} := \text{conv}(\{\iota(u) : u \in \partial T\}) \subseteq \Delta_{n-1}. \quad (5.10)$$

For example, Δ_Γ is the standard simplex Δ_{n-1} . We view the next result as a key convexity property of star-full subtrees of splice diagrams.

Lemma 5.8 *Fix a star-full subtree T of Γ . For every node u of T , $\iota(u)$ admits an expression of the form*

$$\iota(u) = \sum_{\mu \in \partial T} \alpha_\mu \iota(\mu) \quad \text{with} \quad \sum_{\mu \in \partial T} \alpha_\mu = 1 \quad \text{and} \quad \alpha_\mu > 0 \quad \text{for all } \mu \in \partial T. \quad (5.11)$$

In particular, $\iota(T) \subset \Delta_T$.

Proof We let p be the number of nodes of T and proceed by induction on p . The statement is vacuous for $p = 0$. If $p = 1$, then T is a star tree and the result follows by Proposition 5.7 (2). For the inductive step, we let $p \geq 2$ and suppose that the result holds for star-full subtrees with $(p - 1)$ nodes.

We fix $\partial T = \{u_1, \dots, u_d\}$ and we let v be an end-node of T . Following Fig. 4, we let v' be the unique node of T adjacent to v and assume that $u_1, \dots, u_{\delta_v-1}$ are adjacent to v . Proposition 5.7 (2) applied to $\text{Star}_\Gamma(v)$ gives

$$\iota(v) = \beta_0 \iota(v') + \sum_{j=1}^{\delta_v-1} \beta_j \iota(u_j) \quad \text{with} \quad \sum_{j=0}^{\delta_v-1} \beta_j = 1 \quad \text{and} \quad \beta_j > 0 \quad \text{for all } j. \quad (5.12)$$

Using Lemma 5.2, we let T' be the star-full subtree of Γ obtained by pruning T from v . By construction, T' has $(p - 1)$ nodes and its leaves are $\{v\} \cup \{u_{\delta_v}, \dots, u_d\}$. The inductive hypothesis yields the following expressions for v' and all other (potential) nodes $u \neq v'$ of T' :

$$\iota(v') = \gamma_0 \iota(v) + \sum_{j=\delta_v}^d \gamma_j \iota(u_j) \quad \text{and} \quad \iota(u) = \alpha_0 \iota(v) + \sum_{j=\delta_v}^d \alpha_j \iota(u_j), \quad (5.13)$$

with $\alpha_0 + \sum_{j=\delta_v}^d \alpha_j = \gamma_0 + \sum_{j=\delta_v}^d \gamma_j = 1$ and $\alpha_i, \gamma_i > 0$ for all $i \in \{0, \delta_v, \dots, d\}$. Since $0 < \gamma_0 \beta_0 < 1$, substituting the expression for $\iota(v')$ obtained from (5.13) into (5.12) produces the desired positive convex combination for $\iota(v)$:

$$\iota(v) = \sum_{j=1}^{\delta_v-1} \frac{\beta_j}{1 - \gamma_0 \beta_0} \iota(u_j) + \sum_{j=\delta_v}^d \frac{\beta_0 \gamma_j}{1 - \gamma_0 \beta_0} \iota(u_j).$$

In turn, substituting this identity in both expressions from (5.13) gives the positive convex combination statement for all remaining nodes of T . The inclusion $\iota(T) \subset \Delta_T$ follows by the convexity of Δ_T . $\square$

Proposition 5.7 and Lemma 5.8 combined have the following natural consequence:

Corollary 5.9 For each pair of star-full subtrees T, T' of Γ with $T' \subseteq T$, we have $\Delta_{T'} \subseteq \Delta_T$.

Our next result is a generalization of Proposition 5.7 and it highlights a key combinatorial property shared by Γ and all its star-full subtrees.

Proposition 5.10 Let T be a star-full subtree of Γ . Then:

1. the weights $\{\iota(u) : u \in \partial T\}$ are linearly independent;
2. Δ_T is a simplex of dimension $|\partial T| - 1$;
3. for each node v of T we have $\iota(v) \in \Delta_T^\circ$.

Proof Item (2) is a direct consequence of (1). In turn, item (3) follows from (2) and Lemma 5.8. Thus, it remains to prove (1). We distinguish two cases, depending on the number of nodes of T , denoted by p .

Case 1: If $p < 2$, then T is either a vertex, an edge of Γ , or a star tree. If T is a vertex of Γ , then the claim holds because $\iota(u) \neq 0$ for any vertex u of Γ . If T is a star tree, the statement agrees with Proposition 5.7 (1).

Next, assume T is an edge of Γ . We consider two scenarios. First, if T joins a leaf λ and a node u of Γ , then the result follows immediately since $\iota(u)$ and w_λ are linearly independent. On the contrary, assume T joins two adjacent nodes of Γ , say u and v . Pick two leaves λ, μ of Γ with $u \in [\lambda, v]$ and $v \in [u, \mu]$ (i.e., λ is on the u -side and μ is on the v -side of Γ , as seen from the edge $[u, v]$). Lemma 2.8 yields the following formula for the (λ, μ) -minor of the matrix $(w_u|w_v)$, involving the determinant $\det([u, v])$ of the edge $[u, v]$, as defined in (2.1):

$$\begin{pmatrix} \ell_{\lambda,u} & \ell_{\lambda,v} \\ \ell_{\mu,u} & \ell_{\mu,v} \end{pmatrix} = \ell_{\lambda,u} \ell_{\mu,v} \frac{d_{u,v} d_{v,u} - \ell_{u,v}}{d_{u,v} d_{v,u}} = \frac{\ell_{\lambda,u} \ell_{\mu,v} \det([u, v])}{d_{u,v} d_{v,u}}.$$

The last expression is positive by the edge determinant condition, so $\{\iota(u), \iota(v)\}$ is linearly independent.

Case 2: If $p \geq 2$, we know that $d := |\partial T| \geq 4$. We prove the result by reverse induction on d . When $d = n$, we have $T = \Gamma$ and there is nothing to prove since $\iota(\lambda) = w_\lambda$ for all $\lambda \in \partial \Gamma$. For the inductive step, we take $d < n$ and label the leaves of T by $u_1, u_2, \dots, u_d$. We assume that the linear independence holds for any star-full subtree with $k \geq d + 1$ leaves. Without loss of generality we assume u_1 is a node of Γ (one must exist since $T \neq \Gamma$ is star-full). Set $s = \delta_{u_1} - 1$.

Next, we define $T' := T \cup \text{Star}_\Gamma(u_1)$. By construction, T' is a star-full subtree of Γ with $(d + s - 1)$ leaves. Since u_1 is a node of T' , Lemma 5.8 applied to T' yields a positive convex combination:

$$\iota(u_1) = \sum_{v \in \partial T'} \beta_v \iota(v) \quad \text{with} \quad \sum_{v \in \partial T'} \beta_v = 1 \quad \text{and} \quad \beta_v > 0 \quad \text{for all } v \in \partial T'. \quad (5.14)$$

To prove the linear independence for the d points $\iota(u_1), \dots, \iota(u_d)$, we fix a potential dependency relation $\sum_{j=1}^d \alpha_j \iota(u_j) = 0$. Substituting (5.14) into it gives a linear

dependency relation for the leaves of T' :

$$\sum_{j=2}^d (\alpha_j + \alpha_1 \beta_{u_j}) \iota(u_j) + \sum_{j=d+1}^{d+s} (\alpha_1 \beta_{u_j}) \iota(u_j) = 0,$$

where $\{u_{d+1}, \dots, u_{d+s}\}$ are the leaves of T' adjacent to u_1 . The inductive hypothesis applied to T' and the positivity of each β_v with $v \in \partial T'$ forces $\alpha_j = 0$ for all $j = 1, \dots, d$. Thus, (1) holds. $\square$

Next, we state the main result in this section, which is a natural consequence of Lemma 5.8:

Theorem 5.11 *The map ι from (5.1) is injective.*

Proof We prove that the statement holds when restricted to any star-full subtree T of Γ . As in the proof of Lemma 5.8, we argue by induction on the number p of nodes of T . If $p = 0$, then T is either a vertex u or an edge $[u, v]$. The statement in the first case is tautological. The result for the second one holds by construction because $\{\iota(u), \iota(v)\}$ is linearly independent by Proposition 5.7 (1).

If $p = 1$, then T is a star tree. Let v be its unique node and $\{u_1, \dots, u_{\delta_v}\}$ be its leaves. Injectivity over T is a direct consequence of the following identity:

$$\iota([u_i, v]) \cap \iota([u_j, v]) = \{\iota(v)\} \quad \text{for all } i \neq j,$$

which we prove by a direct computation. Indeed, pick $0 \leq a \leq b \leq 1$ with

$$a \iota(u_i) + (1 - a) \iota(v) = b \iota(u_j) + (1 - b) \iota(v). \quad (5.15)$$

By Proposition 5.7 (2), $\iota(v)$ admits a unique expression:

$$\iota(v) = \sum_{k=1}^{\delta_v} \alpha_k \iota(u_k) \quad \text{with} \quad \sum_{k=1}^{\delta_v} \alpha_k = 1 \quad \text{and} \quad \alpha_k > 0 \quad \text{for all } k.$$

Substituting this identity in (5.15) yields the following affine dependency relation for $\iota(\partial T)$:

$$(a + (b - a)\alpha_i)\iota(u_i) + ((b - a)\alpha_j - b)\iota(u_j) + \sum_{k \neq i, j} \alpha_k(b - a)\iota(u_k) = 0$$

By Proposition 5.7 (1), we conclude that $a + (b - a)\alpha_i = (b - a)\alpha_j - b = 0$ and $\alpha_k(b - a) = 0$ for all $k \neq i, j$. Since $\alpha_k > 0$ for all $k \neq i, j$ and $\delta_v \geq 3$, it follows that $b - a = 0$ and $a = -b = 0$. Therefore, expression (5.15) represents $\iota(v)$.

Finally, pick $p \geq 2$ and assume the result holds for star-full subtrees with $p - 1$ nodes. Let T be a star-full subtree with p nodes and pick an end-node v of T . As in Fig. 4, write v' for the unique node of T adjacent to it and $\{u_1, \dots, u_{\delta_v-1}\}$ for the leaves of T adjacent to v .

As in Lemma 5.2, let T' be the star-full subtree obtained by pruning T from v . Our inductive hypothesis ensures that ι is injective when restricted to T' . By the $p = 1$ case we know that $\iota([v, u_i]) \cap \iota([v, u_j]) = \{\iota(v)\}$ if $i \neq j$. Thus, the injectivity of ι when restricted to T will be proven if we show:

$$\iota([v, u_i]) \cap \iota(T') = \{\iota(v)\} \quad \text{for all } i = 1, \dots, \delta_v - 1. \quad (5.16)$$

The identity follows from Proposition 5.10. Indeed, we write any w on the left-hand side of (5.16) as

$$w := at(u_i) + (1 - a)\iota(v) \in \iota([v, u_i]) \cap \iota(T') \quad \text{with } 0 \leq a \leq 1. \quad (5.17)$$

Recall that $w \in \iota(T') \subset \Delta_{T'}$ and $\Delta_{T'} \subset \Delta_T$ by Corollary 5.9. Since $\iota(v) \in (\Delta_T)^\circ$ as in Proposition 5.10 (3), substituting this expression into (5.17) and comparing it with the known expression for w as an element of Δ_T yields an affine dependency equation for $\{\iota(u) : u \in \partial T\}$. The positivity constraint on the coefficients used to write $\iota(v)$ as an element of $(\Delta_T)^\circ$ forces $a = 0$, and so (5.16) holds. $\square$

6 Local tropicalization of splice type systems

Let Γ be a splice diagram with n leaves and let $\mathcal{S}(\Gamma)$ be a splice type system associated to it, as in Definition 2.14. Fixing a total order on $\partial \Gamma$ yields an embedding of the corresponding splice type singularity $(X, 0)$ into $\mathbb{C}^n$. In this section we describe the local tropicalization of this embedded germ, following the characterization from Definition 3.4. As a byproduct, we confirm the first half of Theorem 1.1, namely that $(X, 0)$ is a complete intersection in $\mathbb{C}^n$ with no irreducible components contained in any coordinate subspace.

The injectivity of the map ι from (5.1), discussed in Theorem 5.11, fixes a natural simplicial fan structure on the cone over $\iota(\Gamma)$ in $\mathbb{R}^n$:

Definition 6.1 Let Γ be a splice diagram. Then, the set $\mathbb{R}_{\geq 0}\iota(\Gamma)$ has a natural fan structure, with top-dimensional cones

$$\{\mathbb{R}_{\geq 0}\iota([u, v]) : [u, v] \text{ is an edge of } \Gamma\}.$$

We call it the *splice fan* of Γ .

Here is the main result of this section:

Theorem 6.2 *The local tropicalization of $(X, 0) \hookrightarrow \mathbb{C}^n$ is supported on the splice fan of Γ .*

We prove Theorem 6.2 by a double inclusion argument, first restricting our attention to the positive local tropicalization. In Sect. 6.1 we show that the positive local tropicalization of $\mathcal{S}(\Gamma)$ is contained in the support of the splice fan of Γ . We prove this fact by working with simplices associated to star-full subtrees of Γ , which were

introduced in Definition 5.1. For clarity of exposition, we break the arguments into a series of combinatorial lemmas and propositions. These results allow us to certify that the ideal generated by the w -initial forms of all the series $F_{v,i}$ in $\mathcal{S}(\Gamma)$ always contains a monomial when w lies in the complement of the splice fan of Γ in $(\mathbb{R}_{>0})^n$.

In turn, showing that the support of the splice fan of Γ lies in the Euclidean closure of the local tropicalization of $(X, 0)$ involves the so-called balancing condition for pure-dimensional local tropicalizations. This is the subject of Sect. 6.3. An alternative proof will be given in Sect. 7 after proving the Newton non-degeneracy of the germ $(X, 0)$.

The fact that the positive local tropicalization of $(X, 0)$ is pure-dimensional is verified in an indirect way. Our proof technique relies on the explicit computation of the boundary components of the extended tropicalization, which is done in Sect. 6.2. This establishes the first half of Theorem 1.1 discussed above (see Corollary 6.20). As a consequence, we confirm by Corollary 6.22 that the local tropicalization is the Euclidean closure of the positive one. This result together with the findings in Sects. 6.1 and 6.3 complete the proof of Theorem 6.2.

Remark 6.3 Throughout the next subsections, we adopt the following convenient notation for the admissible exponent vectors $m_{v,e}$ from (2.5). Given a node v and a vertex u of Γ with $u \neq v$, we define $\boxed{m_{v,u}} := m_{v,e}$ where e is the unique edge adjacent to v and lying in the geodesic $[v, u]$. Similarly, given a star-full subtree T of Γ not containing v , we write $\boxed{m_{v,T}} := m_{v,u}$, where u is any vertex of T .

6.1 The positive local tropicalization is contained in the support of the splice fan

In this subsection we show that the only points in Δ_{n-1} contained in the positive local tropicalization $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ are included in $\iota(\Gamma)$. We exploit the terminology and convexity results stated in Sect. 5.

Our first technical result will be used extensively throughout this section to determine $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$. As the proof shows, the Hamm determinant conditions imposed on the system $\mathcal{S}(\Gamma)$ play a crucial role.

Lemma 6.4 Fix a node v of Γ and let e, e', e'' be three distinct edges of $\text{Star}_\Gamma(v)$. Fix $w \in (\mathbb{R}_{\geq 0})^n$ and suppose that the admissible exponent vectors $m_{v,e}, m_{v,e'}, m_{v,e''}$ from (2.5) satisfy:

$$w \cdot m_{v,e} < w \cdot m_{v,e'} \quad \text{and} \quad w \cdot m_{v,e} < w \cdot m_{v,e''}. \quad (6.1)$$

Then, $\underline{z}^{m_{v,e}} = \text{in}_w(f)$ for some f in the linear span of $\{f_{v,i} : i = 1 \dots, \delta_v - 2\}$. If, in addition, w satisfies

$$w \cdot m_{v,e} < w \cdot m \quad \text{for each } m \in \bigcup_{i=1}^{\delta_v-2} \text{Supp}(g_{v,i}), \quad (6.2)$$

then $\underline{z}^{m_{v,e}} = \text{in}_w(F)$ for some series F in the linear span of $\{F_{v,i} : i = 1 \dots, \delta_v - 2\}$. In particular, $w \notin \text{Trop}(\mathcal{S}(\Gamma))$.

Proof We let $e_1, \dots, e_{\delta_v}$ be the edges adjacent to v and assume that $e_{\delta_v-2} := e$, $e_{\delta_v-1} := e'$ and $e_{\delta_v} := e''$. Using the Hamm determinant conditions, we build a basis $\{f'_{v,i}\}_{i=1}^{\delta_v-2}$ for the linear span of $\{f_{v,i}\}_{i=1}^{\delta_v-2}$ where

$$f'_{v,i} := \underline{z}^{m_{v,e_i}} + a_i \underline{z}^{m_{v,e'}} + b_i \underline{z}^{m_{v,e''}} \quad \text{for each } i \in \{1, \dots, \delta_v - 2\}.$$

From (6.1) we conclude that $\text{in}_w(f'_{v,\delta_v-2}) = \underline{z}^{m_{v,e}}$. Taking $f := f'_{v,\delta_v-2}$ proves the first part of the statement.

For the second part, the technique yields a new basis $\{F'_{v,i}\}_{i=1}^{\delta_v-2}$ for the linear span of $\{F'_{v,i}\}_{i=1}^{\delta_v-2}$ with

$$F'_{v,i} = f'_{v,i} + g'_{v,i} \quad \text{for each } i \in \{1, \dots, \delta_v - 2\},$$

where each $g'_{v,i}$ is a linear combination of $\{g_{v,j}\}_{j=1}^{\delta_v-2}$. Condition (6.2) then ensures that

$$\text{in}_w(F'_{v,\delta_v-2}) = \text{in}_w(f'_{v,\delta_v-2}) = \underline{z}^{m_{v,e}}.$$

Thus, the series $F = F'_{v,\delta_v-2}$ satisfies the required properties. In particular, the ideal $\text{in}_w(\langle \mathcal{S}(\Gamma) \rangle)$ contains the monomial $\underline{z}^{m_{v,e}}$ and so $w \notin \text{Trop} \langle \mathcal{S}(\Gamma) \rangle$ by definition. $\square$

Next, we state the main theorem in this subsection, which yields one of the required inclusions in Theorem 6.2 when choosing $T = \Gamma$. More precisely:

Theorem 6.5 *For every star-full subtree T of Γ , we have $\Delta_T \cap \text{Trop}_{>0} \langle \mathcal{S}(\Gamma) \rangle \subseteq \iota(T)$.*

Proof Recall from (5.10) that Δ_T is the convex hull of the set of leaves ∂T of T , viewed in Δ_{n-1} via the map ι . We proceed by induction on the number of nodes of T , which we denote by p . If $p = 0$, then T is either a vertex or an edge of Γ , and $\Delta_T = \iota(T)$. For the inductive step, assume $p \geq 1$ and pick a node v of T . Let $T_1, \dots, T_{\delta_v}$ be the branches of T adjacent to v , as in Definition 5.3. We use the point $\iota(v)$ to perform a stellar subdivision of Δ_T , giving a decomposition $\Delta_T = \bigcup_{\lambda \in \partial T} \tau_{v,\lambda}$, where

$$\boxed{\tau_{v,\lambda}} := \text{conv}(\{\iota(v)\} \cup \iota(\partial T \setminus \{\lambda\})) \quad \text{for all } \lambda \in \partial T. \quad (6.3)$$

By Lemma 5.8, $\tau_{v,\lambda}$ is a simplex of dimension $|\partial T| - 1$. Proposition 6.7 below shows that $\tau_{v,\lambda} \cap \text{Trop}_{>0} \langle \mathcal{S}(\Gamma) \rangle$ lies in the boundary of $\tau_{v,\lambda}$. In turn, Proposition 6.10 below ensures that

$$\tau_{v,\lambda} \cap \text{Trop}_{>0} \langle \mathcal{S}(\Gamma) \rangle \subseteq \bigcup_{\substack{1 \leq i \leq \delta_v \\ i \neq j}} \Delta_{[T_i, v]},$$

where T_j is the unique branch of T adjacent to v that contains the leaf λ , and $\boxed{[T_i, v]}$ denotes the convex hull in Γ of $T_i \cup \{v\}$. Combining this fact with the inductive hypothesis applied to all star-full subtrees $[T_i, v]$ of Γ with $i \in \{1, \dots, \delta_v\}$ gives

$$\Delta_T \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma)) \subseteq \bigcup_{i=1}^{\delta_v} (\Delta_{[T_i, v]} \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma))) \subseteq \bigcup_{i=1}^{\delta_v} \iota([T_i, v]) \subseteq \iota(T). \quad (6.4)$$

□

As a natural consequence of this result, we deduce one of the two inclusions required to confirm Theorem 6.2:

Corollary 6.6 *The positive local tropicalization of $\mathcal{S}(\Gamma)$ is contained in the support of the splice fan of Γ .*

In the remainder of this subsection, we discuss the two key propositions used in the proof of Theorem 6.5. We start by showing that the relative interior of any of the top-dimensional simplices $\tau_{v,\lambda}$ from (6.3) obtained from the stellar subdivision of Δ_T induced by a node v of T does not meet $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$. Lemma 6.4 plays a central role. The task is purely combinatorial and the difficulty lies in how to select the triple of admissible exponent vectors required by the lemma. Throughout, we make use of branches of subtrees, which were introduced in Definition 5.3.

Proposition 6.7 *Let T be a star-full subtree of Γ and $\lambda \in \partial T$. Consider the simplex $\tau_{v,\lambda}$ defined by (6.3). Then, its relative interior $\tau_{v,\lambda}^\circ$ is disjoint from $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$.*

Proof Let u be the unique node of T adjacent to λ , and denote by $T_1, \dots, T_{\delta_u}$ the branches of T adjacent to u . We assume that $T_{\delta_u} = \{\lambda\}$ and fix any $w \in \tau_{v,\lambda}^\circ$. Since $\tau_{v,\lambda}$ is a simplex, we can write w uniquely as

$$w = \alpha_v \iota(v) + \sum_{j=1}^{\delta_u-1} w_j \quad \text{where} \quad [w_j] := \sum_{\mu \in \partial T_j} \alpha_{\mu,j} \iota(\mu) \quad \text{for all } j \in \{1, \dots, \delta_u-1\}, \quad (6.5)$$

and $[\alpha_v], [\alpha_{\mu,j}] > 0$ for all $\mu \in \partial T_j$.

In what follows we analyze the w -initial forms of the series $F_{u,i}$ from $\mathcal{S}(\Gamma)$ for $i \in \{1, \dots, \delta_u-2\}$ and use Lemma 6.4 to confirm that $w \notin \text{Trop}(\mathcal{S}(\Gamma))$. To this end, we compare the w -weights of $\underline{z}^{m_{u,\lambda}}$ and of the remaining monomials in $F_{u,i}$, for each $i \in \{1, \dots, \delta_u-2\}$. We treat the monomials appearing in $f_{u,i}$ and $g_{u,i}$ separately.

Lemma 6.8 discusses the monomials in $g_{u,i}$ and confirms that the required condition (6.2) of Lemma 6.4 holds for $m_{u,\lambda}$. In turn, Lemma 6.9 verifies that we can find two edges e', e'' of T adjacent to u satisfying the inequalities (6.1) for $e = [u, \lambda]$. Therefore, Lemma 6.4 applied to the node u confirms that $\underline{z}^{m_{u,\lambda}}$ is the w -initial form of a series in the linear span of $\{in_w(F_{u,i}) : i = 1, \dots, \delta_u-2\}$. Thus, $w \notin \text{Trop}(\mathcal{S}(\Gamma))$ as we wanted to show. □

In the next two lemmas we place ourselves in the setting of Proposition 6.7. In particular, we use the notations introduced in its proof.

Lemma 6.8 *Let w be a weight vector satisfying condition (6.5). Then, for each $i \in \{1, \dots, \delta_u-2\}$ and each monomial $\underline{z}^m$ appearing in $g_{u,i}$ we have*

$$w \cdot m_{u,\lambda} < w \cdot m. \quad (6.6)$$

Proof First, we compare the w -weights of $\underline{z}^{m_{u,\lambda}}$ and all the monomials $\underline{z}^m$ appearing in $g_{u,i}$. We do so by looking at the weight contributed by each summand of w in the decomposition (6.5). Lemma 2.19 and conditions (2.10) ensure that

$$\iota(v) \cdot m_{u,\lambda} = \frac{\ell_{u,v}}{|w_v|} < \iota(v) \cdot m. \quad (6.7)$$

In turn, to compare the w_j -weights of the monomials $\underline{z}^{m_{u,\lambda}}$ and $\underline{z}^m$, we notice that the only summands $\alpha_{\mu,j} \iota(\mu)$ of w_j contributing a positive weight to $\underline{z}^{m_{u,\lambda}}$ are the ones coming from those $\mu \in \partial T_j$ that are nodes in Γ . Again, Lemma 2.19 and the conditions on $g_{u,i}$ listed in (2.10) confirm that

$$\iota(\mu) \cdot m_{u,\lambda} = \frac{\ell_{u,\mu}}{|w_\mu|} < \iota(\mu) \cdot m \quad \text{for all } \mu \in \partial T_j \setminus \partial \Gamma.$$

Combining this inequality with the positivity of the coefficients $\alpha_{\mu,j}$ yields the inequality

$$w_j \cdot m_{u,\lambda} = \sum_{\mu \in \partial T_j \setminus \partial \Gamma} \alpha_{\mu,j} \frac{\ell_{u,\mu}}{|w_\mu|} \leq \sum_{\mu \in \partial T_j \setminus \partial \Gamma} \alpha_{\mu,j} (\iota(\mu) \cdot m) \leq w_j \cdot m \quad \text{for } 1 \leq j < \delta_u. \quad (6.8)$$

Furthermore, the leftmost inequality is strict if the set $\partial T_j \setminus \partial \Gamma$ is non-empty.

Finally, the positivity of α_v together with the inequalities (6.7) and (6.8) ensure that each m in the support of $g_{u,i}$ satisfies the inequality (6.6). This concludes our proof. $\square$

Lemma 6.9 *Let w be a weight vector verifying condition (6.5). Then, there exist two different edges e' , e'' adjacent to the node u of Γ satisfying*

$$w \cdot m_{u,\lambda} < w \cdot m_{u,e'} \quad \text{and} \quad w \cdot m_{u,\lambda} < w \cdot m_{u,e''}.$$

Proof We use the notation $m_{u,T}$ introduced in Remark 6.3. It is enough to find two different branches T_{j_1} and T_{j_2} of T adjacent to u with $j_1, j_2 < \delta_v$, and verifying

$$w \cdot m_{u,\lambda} < w \cdot m_{u,T_{j_1}} \quad \text{and} \quad w \cdot m_{u,\lambda} < w \cdot m_{u,T_{j_2}}. \quad (6.9)$$

Our choice will depend on the nature of u . If $u = v$ we pick any pair of distinct indices $j_1, j_2 < \delta_v$. On the contrary, if $u \neq v$ we take T_{j_1} to be the unique branch of T adjacent to u containing v , and let T_{j_2} be any other branch of T adjacent to u not containing λ with $j_2 < \delta_v$. After relabeling, we may assume that $j_1 = 1$ and $j_2 = 2$.

In the remainder, we confirm that these two branches satisfy the inequalities in (6.9) by analyzing the contributions of each summand of w in the decomposition (6.5) to the total weight of each of the three relevant monomials. We follow the same reasoning as in the proof of Lemma 6.8. Lemma 2.19 is again central to our computations.

We start with the contribution of $\iota(v)$. The lemma confirms that

$$\iota(v) \cdot m_{u,\lambda} = \frac{\ell_{u,v}}{|w_v|} = \min\{\iota(v) \cdot m_{u,T_1}, \iota(v) \cdot m_{u,T_2}\}. \quad (6.10)$$

To analyze the w_j -weight of the three monomials, we recall from the proof of Lemma 6.8 that we need only consider the contributions of those vertices $\mu \in \partial T_j$ that are nodes of Γ . Indeed, we have

$$w_j \cdot m_{u,\lambda} = \sum_{\mu \in \partial T_j \setminus \partial \Gamma} \alpha_{\mu,j} \frac{\ell_{u,\mu}}{|w_\mu|} \quad \text{for } j \in \{1, \dots, \delta_u - 1\}. \quad (6.11)$$

This expression agrees with the value of $w_j \cdot m_{u,T_1} = w_j \cdot m_{u,T_2}$ when $j > 2$ even if $\partial T_j \setminus \partial \Gamma$ is the empty-set.

If $j = 1$, Lemma 2.19 confirms that

$$w_1 \cdot m_{u,\lambda} = w_1 \cdot m_{u,T_2} = \sum_{\mu \in \partial T_1 \setminus \partial \Gamma} \alpha_{\mu,1} \frac{\ell_{u,\mu}}{|w_\mu|} < w_1 \cdot m_{u,T_1}. \quad (6.12)$$

The inequality is strict even if the set $\partial T_1 \setminus \partial \Gamma$ is empty. Indeed, the fact that the tree T is star-full ensures that when $\partial T_1 \subset \partial \Gamma$, any variable appearing in $\underline{z}^{m_{u,T_1}}$ will be indexed by an element in ∂T_1 . Therefore, the total w_1 -weight of m_{u,T_1} is positive no matter the choice of admissible exponent m_{u,T_1} .

By symmetry, we deduce that $w_2 \cdot m_{u,\lambda} = w_2 \cdot m_{u,T_1} < w_2 \cdot m_{u,T_2}$. Combining this fact, the positivity of α_v , and the expressions (6.10), (6.11) and (6.12) yields the inequalities in (6.9). This concludes our proof. $\square$

Our next result is central to the inductive step in the proof of Theorem 6.5. As with the previous two lemmas, the proof is combinatorial and the difficulty lies in how to select the triple of admissible exponent vectors required by Lemma 6.4 in a way that is compatible with a given input proper face of $\tau_{v,\lambda}$.

Proposition 6.10 *Let T be a star-full subtree of Γ , and let v be a node of T . Denote by $T_1, \dots, T_{\delta_v}$ the branches of T adjacent to v . Let L be a proper subset of ∂T which is not included in any ∂T_i , and set*

$$\boxed{\mathcal{P}} := \text{conv}(\{\iota(v)\} \cup \iota(L)). \quad (6.13)$$

Then, $\mathcal{P}$ is a simplex of dimension $|L|$ and its relative interior $\mathcal{P}^\circ$ is disjoint from $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$.

Proof By Lemma 5.8, we have $\iota(v) \in (\Delta_T)^\circ$. In addition, Proposition 5.10 (2) and (3) imply that $\mathcal{P}$ is a simplex of the expected dimension. It remains to show that $\mathcal{P}^\circ \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma)) = \emptyset$. For each $j = 1, \dots, \delta_v$, we set

$$\boxed{L_j} := L \cap \partial T_j \quad \text{and} \quad \boxed{\tau_j} := \mathbb{R}_{\geq 0} \langle \iota(L_j) \rangle. \quad (6.14)$$

By definition, we have $\tau_j = \{\mathbf{0}\}$ if $L_j = \emptyset$. Moreover, each τ_j is a simplicial cone of dimension $|L_j|$. Our assumptions on L and a suitable relabeling of the branches T_j (if necessary) ensure the existence of some $q \in \{2, \dots, \delta_v\}$ with $L_j \neq \emptyset$ for all $j \in \{1, \dots, q\}$ and $L_j = \emptyset$ for all $j \in \{q+1, \dots, \delta_v\}$.

We argue by contradiction and pick $w \in \mathcal{P}^\circ \cap \text{Trop}_{>0} \langle S(\Gamma) \rangle$. Since $\mathcal{P}$ is a simplex and w belongs to its relative interior, we may write w as

$$w = \alpha_v \iota(v) + \sum_{j=1}^q w_j \quad \text{with } \alpha_v > 0 \text{ and } w_j \in \tau_j^\circ \text{ for all } j. \quad (6.15)$$

Lemma 6.11 below implies that $L_j = \partial T_j$ for all $j \leq q$. Since $L \subsetneq \partial T$, it follows that $q < \delta_v$.

Next, we use Lemma 6.4 to confirm that $\underline{z}^{m_{v,T_{\delta_v}}}$ is the w -initial form of a series in the linear span of $\{\text{in}_w(F_{v,i})\}_{i=1}^{\delta_v-2}$, which contradicts our assumption $w \in \text{Trop}_{>0} \langle S(\Gamma) \rangle$. We use the admissible exponents m_{v,T_1} , m_{v,T_2} and $m_{v,T_{\delta_v}}$.

First, the star-full property of T combined with the condition that $L_j = \partial T_j$ for all $j \leq q$ ensures that

$$\begin{aligned} \iota(v) \cdot m_{v,T_{\delta_v}} &= \iota(v) \cdot m_{v,T_2} = \iota(v) \cdot m_{v,T_1} = \frac{d_v}{|w_v|} \quad \text{and} \\ w_j \cdot m_{v,T_{\delta_v}} &= w_j \cdot m_{v,T_1} = w_j \cdot m_{v,T_2} \quad \text{for all } j \in \{3, \dots, q\}. \end{aligned} \quad (6.16)$$

These identities follow directly from Lemma 2.19. In turn, the same arguments employed in the proof of Lemma 6.9, together with the equalities $L_1 = \partial T_1$ and $L_2 = \partial T_2$ give

$$w_1 \cdot m_{v,T_{\delta_v}} = w_1 \cdot m_{v,T_2} < w_1 \cdot m_{v,T_1} \quad \text{and} \quad w_2 \cdot m_{v,T_{\delta_v}} = w_2 \cdot m_{v,T_1} < w_2 \cdot m_{v,T_2}. \quad (6.17)$$

Expressions (6.16) and (6.17) combined yield $w \cdot m_{v,T_{\delta_v}} < w \cdot m_{v,T_1}$ and $w \cdot m_{v,T_{\delta_v}} < w \cdot m_{v,T_2}$. Thus, the first condition required by Lemma 6.4 is satisfied.

To finish, we must compare the w -weight of $\underline{z}^{m_{v,T_{\delta_v}}}$ with that of any exponent m in the support of a fixed series $g_{v,i}$. For each $j \in \{1, \dots, q\}$ we write $w_j := \sum_{\mu \in \partial T_j} \alpha_{\mu,j} \iota(\mu)$ with $\alpha_{\mu,j} > 0$ for all μ . Then, the defining properties (2.10) of $g_{v,i}$ and the reasoning followed in the proof of Lemma 6.8 imply that

$$\iota(v) \cdot m_{v,T_{\delta_v}} < \iota(v) \cdot m \quad \text{and} \quad w_j \cdot m_{v,T_{\delta_v}} = \sum_{\mu \in L_j \setminus \partial \Gamma} \alpha_{\mu,j} \frac{\ell_{v,\mu}}{|w_\mu|} \leq w_j \cdot m \quad \text{for } 1 \leq j \leq q. \quad (6.18)$$

Note that the right-most inequality for $j \in \{1, \dots, q\}$ is strict whenever $L_j \not\subseteq \partial \Gamma$.

Combining both parts of (6.18) and the positivity of α_v yields $w \cdot m_{v,T_{\delta_v}} < w \cdot m$ whenever $\underline{z}^m$ appears in $g_{v,i}$. This verifies the second hypothesis required for Lemma 6.4, contradicting our choice of $w \in \text{Trop}_{>0} \langle S(\Gamma) \rangle$. $\square$

Lemma 6.11 *Let L be a set as in Proposition 6.10 and $\mathcal{P}$ be the simplex defined by formula (6.13). For each $j \in \{1, \dots, \delta_v\}$, let L_j be the set of elements of L which are leaves of T_j , as in (6.14). If $\mathcal{P}^\circ \cap \text{Trop}_{>0} \langle S(\Gamma) \rangle \neq \emptyset$, then L_j is either empty or equal to ∂T_j .*




Fig. 5 Building a maximal sub-branch of T_j avoiding L_j starting from a suitable leaf $\lambda \in \partial T_j \setminus L_j$ and moving inwards towards v , as in the proof of Lemma 6.11. Here, $s \leq \delta_{v'} - 2$ and $r = \delta_{u''} - 1$

Proof We argue by contradiction and assume $\emptyset \subsetneq L_j \subsetneq \partial T_j$, so in particular $|\partial T_j| > 1$. We break the argument into four combinatorial claims, guided by Fig. 5. The left-most picture informs the discussion for Claims 1 and 2. The central picture refers to Claim 3, and the right-most picture illustrates Claim 4. Throughout, we fix $w \in \mathcal{P}^\circ \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma))$ and write w as in (6.15), where we write each $w_k \in \tau_k^\circ$ as $w_k := \sum_{\mu \in L_k} \alpha_{\mu,k} \iota(\mu)$ for each $k \in \{1, \dots, q\}$, with $\alpha_{\mu,k} > 0$ for all μ, k . $\square$

First, we pick $\lambda \in \partial T_j \setminus L_j$ furthest away from v in the geodesic metric on T (see Remark 2.2). Let v' be the unique node of T_j adjacent to λ . The condition $|\partial T_j| > 1$ ensures that $v' \neq v$. The maximality condition satisfied by λ restricts the nature of the node v' . More precisely:

Claim 1. *The node v' is an end-node of T_j .*

Proof We consider all branches T'_k of T_j adjacent to v' and containing neither v nor λ . Our goal is to show that $|\partial T'_k| = 1$ for all k . We argue by contradiction.

Assume that k is such that $|\partial T'_k| > 1$. Then, by the maximality of the distance between v and λ , we have $\partial T'_k \subseteq L_j$. We consider the series $F_{v',i}$ of $\mathcal{S}(\Gamma)$ for $i \in \{1, \dots, \delta_{v'} - 2\}$, and the admissible exponent vectors at v' associated to λ , v and T'_k . We claim that the weight w satisfies the inequalities

$$w \cdot m_{v',\lambda} < w \cdot m_{v',T'_k}, \quad w \cdot m_{v',\lambda} < w \cdot m_{v',v} \quad \text{and} \quad w \cdot m_{v',\lambda} < w \cdot m \quad (6.19)$$

for each m in the support of some $g_{v',i}$. This cannot happen by Lemma 6.4, since $w \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$.

To prove the inequalities (6.19), we analyze the contributions of each summand of w in the decomposition (6.15) to the total w -weight of each admissible monomial, as we did in the proofs of Lemmas 6.8 and 6.9. Here, the node u appearing in those lemmas is replaced by the node v' . As usual, Lemma 2.19 is central to our arguments. The contribution of each weight vector w_p for $p \neq j$ to the total weight of $\underline{z}^{m_{v',\lambda}}$ comes from elements in $L_p \setminus \partial \Gamma$.

We start by verifying the left-most inequality in (6.19). Since v is a node of Γ , Lemma 2.19 ensures that:

$$\iota(v) \cdot m_{v',\lambda} = \frac{\ell_{v',v}}{|w_v|} = \iota(v) \cdot m_{v',T'_k} < \iota(v) \cdot m_{v',v}. \quad (6.20)$$

In turn, for each $p \in \{1, \dots, q\}$ with $p \neq j$, the weight w_p satisfies

$$w_p \cdot m_{v', \lambda} = \sum_{\mu \in L_p \setminus \partial \Gamma} \alpha_{\mu, p} \frac{\ell_{v', \mu}}{|w_\mu|} = w_p \cdot m_{v', T'_k}. \quad (6.21)$$

On the contrary, if $p = j$, the condition $\partial T'_k \subseteq L_j$ ensures that $w_j \cdot m_{v', \lambda} < w_j \cdot m_{v', T'_k}$. Indeed, separating the w_j -weight of m_{v', T'_k} into the contributions of three groups of elements from L_j we have

$$\begin{aligned} w_j \cdot m_{v', T'_k} &= \sum_{\mu \in L_j \setminus (\partial T'_k \cup \partial \Gamma)} \alpha_{\mu, j} \frac{\ell_{v', \mu}}{|w_\mu|} + \sum_{\mu \in \partial T'_k \setminus \partial \Gamma} \alpha_{\mu, j} \frac{w_\mu \cdot m_{v', T'_k}}{|w_\mu|} \\ &\quad + \sum_{\mu \in \partial T'_k \cap \partial \Gamma} \alpha_{\mu, j} \frac{w_\mu \cdot m_{v', T'_k}}{|w_\mu|}. \end{aligned} \quad (6.22)$$

Notice that $w_\mu \cdot m_{v', T'_k} > \ell_{v', \mu}$ whenever $\mu \in \partial T'_k \setminus \partial \Gamma$ and $w_\mu \cdot m_{v', T'_k} \geq 0$ for $\mu \in \partial T'_k \cap \partial \Gamma$.

Similarly, the w_j -weight of $m_{v', \lambda}$ can be determined by separating the elements of L_j into two types:

$$w_j \cdot m_{v', \lambda} = \sum_{\mu \in L_j \setminus (\partial T'_k \cup \partial \Gamma)} \alpha_{\mu, j} \frac{\ell_{v', \mu}}{|w_\mu|} + \sum_{\mu \in \partial T'_k \setminus \partial \Gamma} \alpha_{\mu, j} \frac{\ell_{v', \mu}}{|w_\mu|}. \quad (6.23)$$

Comparing expressions (6.22) and (6.23), the positivity of all the coefficients $\alpha_{\mu, j}$ implies the inequality $w_j \cdot m_{v', \lambda} < w_j \cdot m_{v', T'_k}$ if the set $\partial T'_k \setminus \partial \Gamma$ is non-empty. In turn, if this last set is empty, the fact that T is star-full ensures that all variables featuring in the admissible monomial $\underline{z}^{m_{v', T'_k}}$ are indexed by elements of $\partial T'_k \subseteq \partial \Gamma$. Therefore, the last summand in (6.22) is strictly positive. Thus, the comparison of the same two expressions gives $w_j \cdot m_{v', \lambda} < w_j \cdot m_{v', T'_k}$ in this situation as well.

By adding up the strict inequality $w_j \cdot m_{v', \lambda} < w_j \cdot m_{v', T'_k}$ and the equalities $w_p \cdot m_{v', \lambda} = w_p \cdot m_{v', T'_k}$ from (6.21) for each $p \in \{1, \dots, q\} \setminus \{j\}$, we obtain:

$$\left(\sum_{k=1}^q w_k \right) \cdot m_{v', \lambda} < \left(\sum_{k=1}^q w_k \right) \cdot m_{v', T'_k}.$$

Combining this inequality with the decomposition (6.15) the equality $\iota(v) \cdot m_{v', \lambda} = \iota(v) \cdot m_{v', T'_k}$ from (6.20) yields the left-most strict inequality in (6.19).

Next, we confirm the central inequality in (6.19). By (6.20) and the positivity of α_v , it is enough to check that $w_p \cdot m_{v', \lambda} \leq w_p \cdot m_{v', v}$ for all $p \in \{1, \dots, q\}$. Indeed, Lemma 2.19 yields:

$$w_p \cdot m_{v',\lambda} = \sum_{\mu \in L_p \setminus \partial \Gamma} \alpha_{\mu,p} \frac{\ell_{v',\mu}}{|w_\mu|} \leq \sum_{\mu \in L_p} \alpha_{\mu,p} \frac{w_\mu \cdot m_{v',v}}{|w_\mu|} = w_p \cdot m_{v',v}, \quad (6.24)$$

since all coefficients $\alpha_{v,\mu}$ are positive, $\ell_{v',\mu} \leq w_\mu \cdot m_{v',v}$ for each $\mu \in L_p \setminus \partial \Gamma$ and $0 \leq w_\mu \cdot m_{v',v}$ if $\mu \in L_p \cap \partial \Gamma$.

To finish, we must certify the right-most inequality in (6.19). We use the same reasoning as in Lemma 6.8. The properties defining the series $g_{v',i}$ combined with the star-full condition of T , the inclusion $\partial T'_k \subseteq L_j$ and the expressions (6.20), (6.21) and (6.23) imply that for each monomial $\underline{z}^m$ appearing in a fixed $g_{v',i}$ we have the inequalities

$$\iota(v) \cdot m_{v',\lambda} < \iota(v) \cdot m \quad \text{and} \quad w_p \cdot m_{v',\lambda} \leq w_p \cdot m \quad \text{for all } p = 1, \dots, q.$$

The inequality $\alpha_v > 0$ then confirms the validity of the right-most inequality in (6.19). $\diamond$

Claim 2. *We have $\text{Star}_{T_j}(v') \cap L_j = \emptyset$. In particular, none of the leaves of T_j adjacent to v' can lie in L_j .*

Proof Since v is an end-node by Claim 1, any branch of T emanating from v' and not containing v is a singleton. The statement follows by the same line of reasoning as Claim 1, working with the exponents $m_{v',v}$, $m_{v',\lambda}$ and m_{v',T'_k} , where T'_k is any of the remaining branches of T_j adjacent to v' . Indeed, if $T'_k = \partial T'_k \subseteq L_j$, the inequalities in (6.19) will remain valid, and this will contradict $w \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$. $\diamond$

As a consequence of Claim 2 we conclude that λ is part of a branch of T_j which avoids L_j and has at least one node. Let T' be a maximal branch of T_j with this property and furthest away from v . We claim that $T' = T_j$. To prove this, we argue by contradicting the maximality of T' . We let u be the node of T_j adjacent to T' and u' be the node of T' adjacent to u , as seen in the center of Fig. 5. Note that $u \neq v$ since $T' \neq T_j$ because L_j is non-empty.

Next, we analyze the intersections between L_j and the leaves of all relevant branches of T_j adjacent to u . We treat two cases, depending on the size of each such branch, starting with singleton branches:

Claim 3. *None of the leaves of T_j adjacent to u belongs to L_j .*

Proof We argue by contradiction and pick a leaf μ of T_j adjacent to u satisfying $\mu \in L_j$. Then, a reasoning similar to that of Claim 1 for the series $F_{u,i}$ replacing λ by T' and T'_k by μ confirms that

$$w \cdot m_{u,T'} < w \cdot m_{u,\mu}, \quad w \cdot m_{u,T'} < w \cdot m_{u,v}, \quad \text{and} \quad w \cdot m_{u,T'} < w \cdot m \quad (6.25)$$

whenever $m \in \text{Supp}(g_{u,i})$. Note that the inequality $w_j \cdot m_{u,T'} < w_j \cdot m_{u,\mu}$ follows because T_j is star-full. However, Lemma 6.4 and the assumption $w \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$ refute the validity of (6.25), so $\mu \notin L_j$. $\diamond$

Claim 4. Given a non-singleton branch $T'' \neq T'$ of T_j adjacent to u and with $v \notin T''$, we have $\partial T'' \cap L_j = \emptyset$.

Proof First, we show that $\partial T'' \not\subseteq L_j$. We argue by contradiction and consider the series of $\mathcal{S}(\Gamma)$ determined by the node u . Replacing the roles of v' , λ and T'_k in (6.19) by u , T' and T'' , respectively, the same proof technique from Claim 1 yields

$$w \cdot m_{u,T'} < w \cdot m_{u,T''}, \quad w \cdot m_{u,T'} < w \cdot m_{u,v} \quad \text{and} \quad w \cdot m_{u,T'} < w \cdot m$$

for each m in the support of any fixed $g_{u,i}$. Lemma 6.4 then shows that $w \notin \text{Trop}_{>0} \mathcal{S}(\Gamma)$, contradicting our original assumption on w . From here it follows that $\partial T'' \not\subseteq L_j$.

Next, we pick some leaf $\lambda \in \partial T'' \setminus L_j$. Following the reasoning of Claims 1 and 2, we build a maximal branch of T'' not meeting L_j . If $\partial T'' \cap L_j \neq \emptyset$ this branch would be a maximal branch of T avoiding L_j and properly contained in T'' . Furthermore, following the notation of the right-most picture in Fig. 5, it would be contained in one of the branches T''_k for $k \in \{1, \dots, r\}$. As a result, the distance between v and this branch would be strictly larger than $\text{dist}_{T_j}(v, T')$, contradicting the maximality choice of T' . Thus, we conclude that $\partial T'' \cap L_j = \emptyset$, as we wanted to show. $\diamond$

To finish, we observe that Claims 3 and 4 combined contradict the maximality of T' , since the convex hull of all branches adjacent to u and not containing v will be a branch of T_j strictly containing T' and not meeting L_j . From here it follows that $T' = T_j$, which cannot happen since $L_j \neq \emptyset$. $\square$

6.2 Boundary components of the extended tropicalization

In this subsection, we characterize the boundary strata of the extended tropicalization of the germ $(X, 0)$ defined by $\mathcal{S}(\Gamma)$ (see Remark 3.10). These strata are determined by the positive local tropicalization of the intersection of X with each coordinate subspace of $\mathbb{C}^n$. This will serve two purposes. First, it will show by combinatorial methods that $(X, 0)$ is a two-dimensional complete intersection with no boundary components (that is, without irreducible components contained in some coordinate hyperplane). Second, it will help us prove the reverse inclusion to the one in Theorem 6.5. The latter is the subject of Sect. 6.3 (see Theorem 6.23).

We start by setting up notation. Throughout, we write $\boxed{\sigma} := (\mathbb{R}_{\geq 0})^n$ and fix a positive-dimensional proper face $\boxed{\tau}$ of σ . Since $N(\partial\Gamma) \simeq \mathbb{Z}^n$ by choosing the basis $\{w_\lambda : \lambda \in \partial\Gamma\}$ from Sect. 2, we define

$$\boxed{L_\tau} := \{\lambda \in \partial\Gamma : w_\lambda \text{ is a ray of } \tau\}. \quad (6.26)$$

We let $\boxed{k}$ be the dimension of τ and consider the natural projection of vector spaces

$$\boxed{p_\tau} : \mathbb{R}^n \rightarrow \mathbb{R}^n / \langle \tau \rangle \simeq \mathbb{R}^{n-k}. \quad (6.27)$$

By abuse of notation, whenever $w_\lambda \notin \tau$, we identify w_λ with its image in $\mathbb{R}^{n-k}$ under p_τ .

Definition 6.12 Given a series $f \in \mathbb{C}\{z_\lambda : \lambda \in \partial \Gamma\}$, we let $\boxed{f^\tau}$ be the series obtained from f by setting all z_λ with $\lambda \in L_\tau$ to be zero. We view $f^\tau \in \mathbb{C}\{z_\lambda : \lambda \in \tau^\perp \cap \mathbb{Z}^n\}$ as a series in the $n - k$ variables in $\{z_\lambda : \lambda \in \partial \Gamma \setminus L_\tau\}$. We call it the τ -truncation of f .

Definition 6.13 We let $\boxed{X_\tau}$ be the intersection of the germ $(X, 0)$ defined by $\mathcal{S}(\Gamma)$ with the dense torus in the $(n - k)$ -dimensional coordinate subspace of $\mathbb{C}^n$ associated to τ . This new germ is defined by the vanishing of the τ -truncations of all series in $\mathcal{S}(\Gamma)$. The *positive local tropicalization of $\mathcal{S}(\Gamma)$ with respect to τ* is defined as the positive local tropicalization of X_τ in $\mathbb{R}^{n-k}$. Following [49, Sect. 12], we denote it by $\boxed{\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)}$.

Our first result generalizes Lemma 6.4, when some, but not all, admissible monomials at a fixed node of Γ have trivial τ -truncations. It will simplify the computation of $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$ for each proper face τ of σ .

Lemma 6.14 Fix a positive-dimensional proper face τ of σ , a node v of Γ and some $w \in p_\tau(\sigma)$. Assume that some τ -truncated series in $\{f_{v,i}^\tau\}_{i=1}^{\delta_v-2}$ is not identically zero and that one of two conditions holds:

1. the system involves at most $\delta_v - 2$ admissible monomials and there exists an edge e adjacent to v with $m_{v,e} \notin \mathbb{N}\langle z_\lambda : \lambda \in L_\tau \rangle$ satisfying $w \cdot m_{v,e} < w \cdot m$ for each m in the support of some $(g_{v,i})^\tau$; or
2. the system involves exactly $\delta_v - 1$ monomials, and we have two distinct edges e, e' of $\text{Star}_\Gamma(v)$ with $m_{v,e}, m_{v,e'} \notin \mathbb{N}\langle z_\lambda : \lambda \in L_\tau \rangle$ such that $w \cdot m_{v,e} < w \cdot m_{v,e'}$ and $w \cdot m_{v,e} < w \cdot m$ for each m in the support of some $(g_{v,i})^\tau$.

Then, $\underline{z}^{m_{v,e}}$ is the w -initial form of a series in the linear span of $\{(F_{v,i})^\tau\}_{i=1}^{\delta_v-2}$ and $w \notin \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$.

Proof Recall from (2.9) that $F_{v,i} = f_{v,i} + g_{v,i}$ for each node v of Γ and each $i \in \{1, \dots, \delta_v - 2\}$. Our assumptions on $(f_{v,i})^\tau$ and the weight vector w ensure both that the τ -truncation $(F_{v,i})^\tau$ of $F_{v,i}$ is non-zero and that the w -initial form of $(F_{v,i})^\tau$ agrees with that of $(f_{v,i})^\tau$.

The proof follows the same reasoning as that of Lemma 6.4, considering the truncations of a new basis $\{F'_{v,i}\}_{i=1}^{\delta_v-2}$ obtained by suitable linear combinations of the original series $\{F_{v,i}\}_{i=1}^{\delta_v-2}$ and ordering the edges adjacent to v in a convenient way. Writing each element in the new basis as $F'_{v,i} = f'_{v,i} + g'_{v,i}$ in the spirit of (2.9), the Hamm determinant conditions on the original splice system allow us to choose a special form for the polynomials $\{f'_{v,i}\}_i$. Indeed, their coefficient matrix has the block form $(\text{Id}_{\delta_v-2} \mid * \mid *)$, where $*$ represents a column vector in $(\mathbb{C}^*)^{\delta_v-2}$. By construction, the new tails $g'_{v,i}$ will satisfy conditions (1), respectively (2), if and only if the old tails $g_{v,i}$ did.

If condition (1) holds, we pick any two edges e', e'' with $m_{v,e'}, m_{v,e''} \in \mathbb{N}\langle z_\lambda : \lambda \in L_\tau \rangle$ and order the edges adjacent to v so that $e_{\delta_v-2} := e, e_{\delta_v-1} := e'$ and $e_{\delta_v} := e''$. In

this situation, the statement follows by our assumptions on the basis $\{F'_{v,i}\}_i$, combined with Lemma 2.19. Indeed, we have

$$\text{in}_w((F'_{v,\delta_v-2})^\tau) = \text{in}_w((f'_{v,\delta_v-2})^\tau) = \underline{z}^{m_{v,e}}.$$

Similarly, if condition (2) holds, we pick e'' to be the unique edge with $m_{v,e''} \in \mathbb{N}\langle z_\lambda : \lambda \in L_\tau \rangle$. Indeed, $m_{v,e''}$ is the single monomial of the polynomials $\{f_{v,i}\}_i$ that drops when taking their τ -truncations. We set $e_{\delta_v} := e''$ and fix $e := e_{\delta_v-2}$ and $e' := e_{\delta_v-1}$. As in the previous case, we have $\text{in}_w((F'_{v,\delta_v-2})^\tau) = \underline{z}^{m_{v,e}}$. $\square$

Here is the main result of this section:

Theorem 6.15 *Let τ be a positive-dimensional proper face of σ with $\lambda \in L_\tau$. Let v be the unique node of Γ adjacent to λ . Then:*

1. *If $\dim \tau = 1$, then $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) \subseteq \mathbb{R}_{>0}\langle p_\tau(w_v) \rangle$.*
2. *If $\dim \tau \geq 2$, then $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) = \emptyset$.*

Proof We fix $k = \dim \tau$. Since $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$ is included in $(\mathbb{R}_{>0})^{n-k}$, we can determine the positive tropicalization by its intersection with the standard simplex Δ_{n-k-1} . We do this by following the proof strategy of Theorem 6.5. Since the vector $p_\tau(w_v)$ lies in $(\mathbb{R}_{>0})^{n-k}$, we set

$$\boxed{w'_v} := \frac{p_\tau(w_v)}{|p_\tau(w_v)|} \in (\Delta_{n-k-1})^\circ \quad (6.28)$$

and perform a stellar subdivision of this standard simplex using w'_v .

If $k = 1$, Proposition 6.16 below ensures that every positive-dimensional simplex ρ of the stellar subdivision that meets $(\Delta_{n-2})^\circ$ satisfies $\rho^\circ \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) = \emptyset$. The subdivision has only one zero-dimensional simplex, namely $\{p_\tau(w_v)\}$, so item (1) holds. In turn, when $\dim \tau = 2$, the same proposition ensures the intersection is empty even if ρ is a point. This forces $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) = \emptyset$, as stated in item (2). $\square$

Our next result is an adaptation of Proposition 6.10, needed to rule out points of the extended tropicalization $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$ in a given simplex $\rho \subseteq \Delta_{n-\dim \tau-1}$. It is central to proving Theorem 6.15.

Proposition 6.16 *Let τ , λ and v be as in Theorem 6.15 and L_τ be as in (6.26). Fix a proper subset L of $\partial \Gamma \setminus L_\tau$. Let w'_v be as in (6.28) and set $\boxed{\rho} := \text{conv}(\{w'_v\} \cup \{w_\mu : \mu \in L\}) \subseteq \Delta_{n-k-1}$, where $k = \dim \tau$. If $|L \cup L_\tau| > 1$, then*

$$\rho^\circ \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) = \emptyset. \quad (6.29)$$

Proof We follow the proof strategy of Proposition 6.10 for $T = \Gamma$ but now incorporating the set L_τ into the arguments. We let $T_1, \dots, T_{\delta_v}$ be the branches of Γ adjacent to v , with $T_1 = \{\lambda\}$. Given $j \in \{2, \dots, \delta_v\}$, we set

$$\boxed{L_{\tau,j}} := L_\tau \cap \partial T_j, \quad \boxed{L_j} := L \cap \partial T_j \quad \text{and} \quad \boxed{\rho_j} := \mathbb{R}_{\geq 0}\langle p_\tau(w_\mu) : \mu \in L_j \rangle \subset \mathbb{R}^{n-k}. \quad (6.30)$$

If $L_j = \emptyset$, we declare $\rho_j = \{0\}$. Since $1 < |L \cup L_\tau| < n$, upon relabeling the branches $T_2, \dots, T_{\delta_v}$ if necessary, there exists a unique $q \in \{2, \dots, \delta_v\}$ with $L_j \cup L_{\tau,j} \neq \emptyset$ for all $j \in \{1, \dots, q\}$, and $L_j \cup L_{\tau,j} = \emptyset$ for $j > q$.

We argue by contradiction and pick $w \in \rho^\circ \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$. Since ρ is a simplex and $T_1 \subset L_\tau$, we have

$$w = \alpha_v w'_v + \sum_{j=2}^q w_j \quad \text{with } \alpha_v > 0 \text{ and } w_j \in \rho_j^\circ \text{ for all } j \in \{2, \dots, q\}. \quad (6.31)$$

In particular, we know that $w_j = 0$ if and only if $\rho_j = \{0\}$.

Lemma 6.17 below ensures that $L_j \cup L_{\tau,j} = \partial T_j$ for all $j \leq q$. Since $1 < |L \cup L_\tau| < n$, we conclude that:

$$1 < q < \delta_v. \quad (6.32)$$

In particular, we see that $\partial T_{\delta_v} \cap (L \cup L_\tau) = \emptyset$, so the τ -truncation of $\underline{z}^{m_{v,T_{\delta_v}}}$ is non-zero.

By the Hamm conditions, the admissible monomial $\underline{z}^{m_{v,T_{\delta_v}}}$ must feature in some $f_{v,i}$. Therefore, the τ -truncated series $\{(f_{v,i})^\tau\}_{i=1}^{\delta_v-2}$ are not all identically zero. In turn, since $\lambda \in \partial T_1 \cap L_\tau$ and $\underline{z}^{m_{v,T_1}} = z_\lambda^{d_{v,\lambda}}$, we conclude that its τ -truncation is zero. Thus, the system of τ -truncations $\{(f_{v,i})^\tau\}_{i=1}^{\delta_v-2}$ involves at most $\delta_v - 1$ admissible monomials.

The proposition follows from our next claim, which contradicts our assumption that $w \in \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$.

Claim. *The monomial $\underline{z}^{m_{v,T_{\delta_v}}}$ is the w -initial form of a series in the linear span of $\{(F_{v,i})^\tau\}_{i=1}^{\delta_v-2}$.*

The statement of this claim matches the conclusion of Lemma 6.14. Recall that our earlier discussion confirmed that the τ -truncations of $f_{v,i}$ are not all zero, and there are two options for the number of monomials featured in these τ -truncations. We prove the claim via a case-by-case analysis, each one matching one of the two possible scenarios portrayed in the lemma.

First, we assume that the system of τ -truncations $\{(f_{v,i})^\tau\}_{i=1}^{\delta_v-2}$ involves exactly $\delta_v - 1$ monomials. In particular, $(\underline{z}^{m_{v,T_2}})^\tau \neq 0$, so no variable appearing in $\underline{z}^{m_{v,T_2}}$ can be indexed by an element of $L_{\tau,2}$. Since $L_2 \cup L_{\tau,2} = \partial T_2$, we conclude that $\rho_2 \neq \{0\}$ and $w_2 \cdot m_{v,T_2} > 0$.

Next, we confirm that the hypotheses of Lemma 6.14 (2) hold for the weight vector w and the edges $e = e_{\delta_v}$ and $e' = e_2$ giving rise to the admissible exponents $m_{v,T_{\delta_v}}$ and m_{v,T_2} , respectively. Notice that our earlier discussion ensures that these two vectors lie outside $\mathbb{N}\langle z_\lambda : \lambda \in L_\tau \rangle$. To check the required inequalities between the w -weights of these two admissible monomials and of any monomial $\underline{z}^m$ in the support of some $(g_{v,i})^\tau$, we compare the contribution of each summand of w featured in (6.31) to the total w -weight of each monomial. This was the same method used in the proofs of Lemmas 6.8 and 6.9.




Fig. 6 Given a branch T_j of Γ adjacent to v and a fixed proper subset of leaves L of $\partial \Gamma \setminus \{\lambda\}$ meeting ∂T_j properly, we find a node u in T_j with $\text{dist}_\Gamma(u, v)$ maximal so that the branches $T'_1, \dots, T'_r$ have a given property, but $T'_{r+1}, \dots, T'_{\delta_u-1}$ do not. This construction is the main ingredient in the proof of Lemma 6.17

First, we focus on w'_v . Lemma 2.19 and the conditions (2.10) on $g_{v,i}$ imply that

$$w'_v \cdot m_{v, T_{\delta_v}} = w'_v \cdot m_{v, T_2} = \frac{d_v}{|p_\tau(w_v)|} < w'_v \cdot m. \quad (6.33)$$

In turn, our earlier discussion and the inequality $q < \delta_v$ seen in (6.32) imply the following relations among the w_j -weights:

$$w_2 \cdot m_{v, T_{\delta_v}} = 0 < w_2 \cdot m_{v, T_2} \quad \text{and} \quad w_j \cdot m_{v, T_{\delta_v}} = w_j \cdot m_{v, T_2} = 0 \leq w_j \cdot m \\ \text{for } j \in \{3, \dots, q\}, \quad (6.34)$$

where $\underline{z}^m$ is any monomial in the support of $g_{v,i}$ surviving the τ -truncation. Combining the positivity of α_v with expressions (6.33) and (6.34) confirms the validity of the conditions required in Lemma 6.14 (2).

To finish proving the claim, we must analyze the case when the τ -truncated system $\{(f_{v,i})^\tau\}_{i=1}^{\delta_v-2}$ has at most $\delta_v - 2$ admissible monomials. In this case, we consider the edge $e = e_{\delta_v}$ and certify that the hypotheses imposed in Lemma 6.14 (1) hold. This is done by checking that the inequalities in (6.33) and (6.34) involving $m_{v, T_{\delta_v}}$ and m remain valid in this new setting. $\square$

Our next result is analogous to Lemma 6.11. We prove it by using the same techniques.

Lemma 6.17 Fix $L_{\tau,j}$, L_j and ρ_j as in (6.30). If $\rho^\circ \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) \neq \emptyset$, then the set $L_j \cup L_{\tau,j}$ is either empty or it equals ∂T_j .

Proof We use the notation established in the proof of Proposition 6.16 and pick $w \in \rho^\circ \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$. We decompose w as in (6.31). We argue by contradiction, assuming that $0 < |L_j \cup L_{\tau,j}| < |\partial T_j|$.

First, note that we can find a branch T' of Γ contained in T_j with $\partial T' \cap (L_j \cup L_{\tau,j}) = \emptyset$. For example, any leaf μ of ∂T_j not in $L \cup L_\tau$ will produce such a branch. Since T_j is finite, we can choose the branch T' to be maximal with respect to the condition $\partial T' \cap (L_j \cup L_{\tau,j}) = \emptyset$ and furthest away from v in the geodesic metric on Γ . Let u be the unique node of Γ adjacent to T' . Since $L_j \cup L_{\tau,j} \neq \emptyset$, we know that $u \neq v$.

Assume that there are r branches adjacent to u with this maximality property, and denote them by $T'_1 = T', \dots, T'_r$ as in Fig. 6. We let $T'_{r+1}, \dots, T'_{\delta_u-1}$ be the remaining branches of Γ adjacent to u and not containing λ . The maximality of both T' and $\text{dist}_\Gamma(u, v)$ combined with the condition $u \neq v$ implies that $r < \delta_u - 1$ and $\partial T'_i \subseteq L_j \cup L_{\tau,j}$ for each $i \in \{r+1, \dots, \delta_u-1\}$. Indeed, if the latter were not the case, we could find a branch T'' of Γ inside T'_i with the same properties as T' , but further away from v .

The construction of T'_1 ensures that $(\underline{z}^{m_{u,T'_1}})^\tau \neq 0$. Therefore, the collection $\{(f_{u,i})^\tau\}_{i=1}^{\delta_u-2}$ is not identically zero. Our original assumption that $w \in \text{ Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$ will be contradicted by the following assertion:

Claim. *The admissible monomial $\underline{z}^{m_{u,T'_1}}$ is the w -initial form of a series in the linear span of $\{(F_{u,i})^\tau\}_{i=1}^{\delta_u-2}$.*

We prove our claim by analyzing three cases (in decreasing order of difficulty), depending on whether the number of admissible monomials in the τ -truncation of the collection $\{f_{u,i}\}_{i=1}^{\delta_u-2}$ is exactly δ_u , at most $\delta_v - 2$, or exactly $\delta_u - 1$. The first case will make use of Lemma 6.4, whereas we will invoke Lemma 6.14 to establish the other two.

Case 1: We assume that the number of admissible monomials at u remaining after τ -truncation is δ_u . By the Hamm conditions, this forces $f_{u,i} = (f_{u,i})^\tau$ for each $i \in \{1, \dots, \delta_u-2\}$. After replacing each $g_{u,i}$ with its τ -truncation, we are in the setting of Lemma 6.4, where we view $(\mathbb{R}_{\geq 0})^{n-k} \subset \mathbb{R}_{\geq 0}^n$ by adding zeros as complementary coordinates. In this situation, the condition $\partial T'_1 \cap L = \emptyset$ and the defining properties of $g_{u,i}$ ensure that

$$w \cdot m_{u,T'_1} < w \cdot m_{u,v}, \quad w \cdot m_{u,T'_1} < w \cdot m_{u,T'_{\delta_u-1}} \quad \text{and} \quad w \cdot m_{u,T'_1} < w \cdot m \quad (6.35)$$

for each $\underline{z}^m$ appearing in $(g_{u,i})^\tau$. The aforementioned lemma then implies the validity of our claim.

To prove the inequalities in (6.35), we proceed by comparing the contributions of each summand of w in the decomposition (6.31) to the weight of each monomial. Lemma 2.19 and (2.10) ensure that

$$w'_v \cdot m_{u,T'_1} = w'_v \cdot m_{u,T'_{\delta_u-1}} = \frac{\ell_{u,v}}{|p_\tau(w_v)|} < \min\{w'_v \cdot m_{u,v}, w'_v \cdot m\}. \quad (6.36)$$

In turn, for each $p \in \{2, \dots, \delta_v\}$ with $p \neq j$, the fact that L contains no node of Γ gives

$$w_p \cdot m_{u,T'_1} = w_p \cdot m_{u,T'_{\delta_u-1}} = 0 \leq \min\{w_p \cdot m_{u,v}, w_p \cdot m\}. \quad (6.37)$$

Finally, since $L_j \cap \partial T'_1 = \emptyset$, $\partial T'_{\delta_v-1} \subseteq L_j \cap L_{\tau,j}$ and the admissible monomial $\underline{z}^{m_{u,T'_{\delta_v-1}}}$ has no variable indexed by an element in $L_{\tau,j}$, we see that

$$w_j \cdot m_{u,T'_1} = 0 \leq \min\{w_j \cdot m_{u,v}, w_j \cdot m\} \quad \text{whereas} \quad 0 < w_j \cdot m_{u,T'_{\delta_u-1}}. \quad (6.38)$$

Expressions (6.36), (6.37) and (6.38), combined with the positivity of α_v confirm (6.35).

Case 2: Assume that the τ -truncated series $\{(f_{u,i})^\tau\}_{i=1}^{\delta_u-2}$ involve at most $\delta_u - 2$ admissible monomials. In this situation the claim follows by Lemma 6.14 (1) since the inequalities involving m_{u,T'_1} and m in (6.35) remain valid in this scenario.

Case 3: Suppose that the τ -truncations of the collection $\{f_{u,i}\}_{i=1}^{\delta_u-2}$ involve exactly $\delta_u - 1$ admissible monomials. If so, up to relabeling of $T'_{r+1}, \dots, T'_{\delta_u-1}$ we may assume that exactly one of the admissible monomials with exponent $m_{u,v}$ or $m_{u,T'_{\delta_u-1}}$ vanishes under τ -truncation. The inequalities in (6.35) involving m_{u,T'_1} , m and the aforementioned surviving admissible exponent remain valid. Thus, the hypotheses required by Lemma 6.14 (2) hold, hence confirming the claim. $\square$

Theorem 6.15 has the following two important consequences:

Corollary 6.18 *The intersection of the germ $(X, 0)$ defined by the system $\mathcal{S}(\Gamma)$ with any coordinate subspace H of $\mathbb{C}^n$ of codimension at least two is just the origin.*

Proof Assume the contrary and let H be a coordinate subspace of $\mathbb{C}^n$ of maximal codimension such that $X \cap H \neq \{0\}$. Consider the face τ of $(\mathbb{R}_{\geq 0})^n$ associated to H and let X_τ be the intersection of X with the dense torus of H . The maximality condition on H implies that $X_\tau \neq \emptyset$. Since $\dim X_\tau = \dim \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$ by Proposition 3.15 (1), we conclude that $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) \neq \emptyset$.

Since the dimension of τ agrees with the codimension of H in $\mathbb{C}^n$, our hypothesis implies that $\dim \tau \geq 2$. Therefore, we have $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) = \emptyset$ by Theorem 6.15 (2). This contradicts our earlier observation. $\square$

Corollary 6.19 *The germ $(X, 0)$ defined by the system $\mathcal{S}(\Gamma)$ intersects each coordinate hyperplane along a germ of a curve. All its irreducible components meet the dense torus of the corresponding hyperplane.*

Proof We fix a coordinate hyperplane $\{z_\lambda = 0\}$ and let τ be the cone generated by w_λ in $\mathbb{R}^n$. Counting the number of equations defining $Y := X \cap \{z_\lambda = 0\}$ in $\mathbb{C}^n$, we see that $\dim Y \geq 1$ by Krull's principal ideal theorem. By construction, $X_\tau = Y \cap (\mathbb{C}^*)^{n-1}$. Since any face τ' of $(\mathbb{R}_{\geq 0})^n$ properly containing τ satisfies $X_{\tau'} = \emptyset$ by Corollary 6.18, it follows that no component of Y lies in the toric boundary of $\mathbb{C}^{n-1}$. Furthermore, Proposition 3.15 (1) implies that $\dim X_\tau = \dim \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$. Since Theorem 6.15 (1) ensures that $\dim \text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) \leq 1$, we conclude that $\dim X_\tau = 1$. $\square$

In turn, the last corollary has two consequences. First, it confirms that the germ defined by $\mathcal{S}(\Gamma)$ is a complete intersection, and second, it shows that equality must hold in Theorem 6.15 (1).

Corollary 6.20 *The germ $(X, 0)$ defined by the system $\mathcal{S}(\Gamma)$ is a two-dimensional complete intersection. Each of its irreducible components meets the dense torus $(\mathbb{C}^*)^n$ non-trivially in dimension two.*

Note that this result allows a priori for the germ $(X, 0)$ to have several irreducible components. We will see in Corollary 7.13 that $(X, 0)$ is in fact irreducible.

Corollary 6.21 *Let λ be a leaf of Γ and let v be the unique node of Γ adjacent to it. If $\tau = \mathbb{R}_{\geq 0}\langle w_\lambda \rangle$, then $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau) = \mathbb{R}_{>0}\langle p_\tau(w_v) \rangle$.*

Proof The result follows by combining Theorem 6.15 (1), Corollary 6.19, and the equality between the dimensions of X_τ and $\text{Trop}_{>0}(\mathcal{S}(\Gamma), \tau)$ stated in Proposition 3.15 (1). $\square$

Our last result in this subsection is a special case of Proposition 3.13. It will be used in the next subsection.

Corollary 6.22 *The local tropicalization of the germ defined by the splice type system $\mathcal{S}(\Gamma)$ is the Euclidean closure of $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ in $\mathbb{R}^n$.*

6.3 The support of the splice fan is contained in the local tropicalization

In this subsection, we prove the remaining inclusion in Theorem 6.2. Our arguments are purely combinatorial, and rely on the balancing condition for pure-dimensional local tropicalizations (see Theorem 3.23 and Remark 3.24). Corollary 6.20 confirms that such condition holds for the positive tropicalization of the germ defined by $\mathcal{S}(\Gamma)$. Furthermore, the proofs in this section imply that no proper two-dimensional subset of $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ is balanced.

We start by stating the main result in this section. Its proof will be broken into several lemmas and propositions for clarity of exposition.

Theorem 6.23 *For every splice diagram Γ , we have $\iota(\Gamma) \subseteq \text{Trop}(\mathcal{S}(\Gamma))$ in $(\mathbb{R}_{\geq 0})^n$.*

Proof By Lemma 6.24 below we know that $\Delta_{n-1} \cap \text{Trop}(\mathcal{S}(\Gamma))$ is a 1-dimensional polyhedral complex and $\iota(v) \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$ for some node v of Γ . We claim that, in fact, $\iota(u) \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$ for each node u .

We prove this claim by induction on the distance between u and v (recall that Γ is connected). If $u = v$ there is nothing to show. For the inductive step, pick a node u with $\text{dist}_\Gamma(u, v) = k \geq 1$ and assume that the claim holds for each node u' of Γ with $\text{dist}_\Gamma(u', v) = k - 1$. Let u' be the unique node of $[u, v]$ adjacent to u . Then, $\iota(u') \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$ by our inductive hypothesis. Proposition 6.26 below and Corollary 6.22 yield:

$$\iota(u) \in \iota(\text{Star}_\Gamma(u')) \cap (\mathbb{R}_{>0})^n \subseteq \text{Trop}(\mathcal{S}(\Gamma)).$$

The desired inclusion $\iota(\Gamma) \subseteq \overline{\text{Trop}_{>0}(\mathcal{S}(\Gamma))}$ follows by combining Proposition 6.26 with the identity

$$\iota(\Gamma) = \bigcup_{v \text{ node of } \Gamma} \iota(\text{Star}_\Gamma(v)).$$

$\square$

Our first lemma ensures that the image of some node of Γ lies in the positive tropicalization of $\mathcal{S}(\Gamma)$:

Lemma 6.24 *The intersection $\Delta_{n-1} \cap \text{Trop}(\mathcal{S}(\Gamma))$ is a 1-dimensional polyhedral complex. Furthermore, there exists a node v of Γ with $\iota(v) \in \Delta_{n-1} \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma))$.*

Proof By Corollary 6.20 and Proposition 3.15 (1) we know that $\text{Trop} \mathcal{S}(\Gamma)$ is a fan of pure dimension two, and so $\Delta_{n-1} \cap \text{Trop}(\mathcal{S}(\Gamma))$ is a pure 1-dimensional polyhedral complex. To conclude, we must find a node v with $\iota(v) \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$.

Since $\text{Trop}_{>0}(\mathcal{S}(\Gamma)) \subseteq \iota(\Gamma)$ by Corollary 6.6, we have two possibilities for any $w \in \Delta_{n-1} \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma))$: either $w = \iota(v)$ for some node v of Γ or $w \in \iota([u, u']^\circ)$ for two adjacent vertices u, u' of Γ . In the second situation, Lemma 6.25 below ensures that $\iota([u, u']) \subset \Delta_{n-1} \cap \text{Trop}(\mathcal{S}(\Gamma))$. Since one of u or u' must be a node of Γ and each node maps to $(\Delta_{n-1})^\circ$ under ι , the claim follows. $\square$

Our next lemma is central to the proof of both Theorem 6.23 and Lemma 6.24. It describes the possible intersections between the local tropicalization of the ideal $\langle \mathcal{S}(\Gamma) \rangle$ and the edges of Γ embedded in Δ_{n-1} via the map ι . The balancing condition for positive local tropicalizations plays a prominent role.

Lemma 6.25 *Let u, u' be two adjacent vertices of Γ . If $\iota([u, u']^\circ)$ intersects $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ non-trivially, then $\iota([u, u']) \subseteq \text{Trop}(\mathcal{S}(\Gamma))$.*

Proof It suffices to show that $\iota([u, u']^\circ) \subseteq \text{Trop}(\mathcal{S}(\Gamma))$. We argue by contradiction and fix a point $w \in \iota([u, u']) \setminus \text{Trop}_{>0}(\mathcal{S}(\Gamma))$. Let $\varphi: [0, 1] \rightarrow [u, u']$ be the affine map such that $\varphi(0) = u$ and $\varphi(1) = u'$. There exists a unique $t_0 \in [0, 1]$ such that $w = \iota(\varphi(t_0))$. Since $\text{Trop}(\mathcal{S}(\Gamma))$ is closed in $(\mathbb{R}_{\geq 0})^n$, we can find a pair $a, b \in [0, 1]$ satisfying $a < t_0 < b$ and such that the open segment $(\iota(\varphi(a)), \iota(\varphi(b)))$ avoids $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$. Furthermore, we pick (a, b) to be the maximal open interval in $[0, 1]$ containing t_0 with this property. A contradiction will arise naturally if we prove that $a = 0$ and $b = 1$.

By symmetry, it suffices to show that $a = 0$. We argue by contradiction, and assume $a > 0$, so $\iota(\varphi(a)) \in (\Delta_{n-1})^\circ$ by construction. The maximality of (a, b) combined with Corollary 6.22 ensures that $\iota(\varphi(a)) \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$. Recall from Corollary 6.6 that $\Delta_{n-1} \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma)) \subseteq \iota(\Gamma)$. Since ι is an injection by Theorem 5.11, any neighborhood of $\iota(\varphi(a))$ in $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ lies in the relative interior of the two-dimensional cone $\mathbb{R}_{\geq 0} \langle \iota([u, \varphi(a)]) \rangle$. In turn, since $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ is two-dimensional, and $\iota([\varphi(a), \varphi(b)]^\circ)$ avoids $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$, it follows that $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ is not balanced at $\iota(\varphi(a))$ when $a > 0$. Indeed, the balancing condition (see Theorem 3.23) at $\iota(\varphi(a))$ involves a single non-zero vector, namely, the image of $\iota(u) - \iota(\varphi(a))$ in the lattice $(\mathbb{Z}^n \cap \langle \iota(u), \iota(\varphi(a)) \rangle) / (\mathbb{Z}^n \cap \langle \iota(\varphi(a)) \rangle)$, and a non-zero scalar. Therefore, we must have $a = 0$. $\square$

Our next result plays a prominent role in the induction arguments used to prove Theorem 6.23. Once again, as in the proof of Lemma 6.25, the balancing condition for local tropicalizations is crucial in our reasoning.

Proposition 6.26 *Let v be a node of Γ with $\iota(v) \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$. Then, $\iota(\text{Star}_\Gamma(v)) \subseteq \text{Trop}(\mathcal{S}(\Gamma))$.*

Proof After refinement if necessary, we may assume that $\boxed{\tau} := \mathbb{R}_{\geq 0}\iota(v)$ is a ray of $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$. In order to exploit the balancing condition, we use analogous notation to that of Definition 3.21, simplified slightly by the fact that $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ is 2-dimensional.

Given a vertex u of Γ adjacent to v , we consider the 1-dimensional saturated lattice

$$\boxed{\Lambda} := \frac{\mathbb{Z}^n \cap \mathbb{R}\langle w_u, w_v \rangle}{\mathbb{Z}^n \cap \mathbb{R}\langle w_v \rangle}.$$

Fix a vector $w_{u|\tau} \in \mathbb{Z}^n \cap \mathbb{R}_{\geq 0}\iota([u, v])$ whose natural projection to Λ generates this lattice. In particular, we can write $w_{u|\tau}$ uniquely as

$$w_{u|\tau} = \alpha_u \iota(u) + \beta_u \iota(v) \quad (6.39)$$

for some $\boxed{\alpha_u}, \boxed{\beta_u} \in \mathbb{Q}$ and with $\alpha_u > 0$.

We let $u_1, \dots, u_{\delta_v}$ be the vertices of Γ adjacent to v . The balancing condition for $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ at τ combined with Theorem 6.5 ensures the existence of non-negative integers $\{k_1, \dots, k_{\delta_v}\}$ (i.e., the tropical multiplicities of Definition 3.20) satisfying:

$$\sum_{i=1}^{\delta_v} k_i w_{u_i|\tau} \in \mathbb{Z}^n \cap \mathbb{R}\langle w_v \rangle. \quad (6.40)$$

Moreover, by Definition 3.20, $(\iota([u_j, v]))^\circ$ intersects $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ non-trivially in a neighborhood of $\iota(v)$ if and only if $k_j \neq 0$.

Since $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ is pure of dimension two and $\iota(v) \in \text{Trop}_{>0}(\mathcal{S}(\Gamma))$, we know that $k_{j_0} > 0$ for some $j_0 \in \{1, \dots, \delta_v\}$. We claim that, furthermore, all $k_1, \dots, k_{\delta_v}$ are positive integers. The inclusion $\iota(\text{Star}_\Gamma(v)) \subset \text{Trop}(\mathcal{S}(\Gamma))$ follows by combining this statement with Lemma 6.25.

To prove our claim, we consider the following linear equation in $k_1, \dots, k_{\delta_v}$ that is equivalent to (6.40):

$$\sum_{j=1}^{\delta_v} (k_j \alpha_{u_j}) \iota(u_j) = \beta \iota(v) \quad \text{with } \beta \in \mathbb{Q}. \quad (6.41)$$

First, we argue that this system admits a unique solution $(k_1, \dots, k_{\delta_v}) \in \mathbb{Q}^{\delta_v}$ for all β . Uniqueness follows directly since $\alpha_{u_j} > 0$ for all j and the set $\{\iota(u_1), \dots, \iota(u_{\delta_v})\}$ is linearly independent by Proposition 5.10 (1).

Second, we claim that for $\beta \neq 0$, any solution to (6.41) has $k_j \neq 0$ for all j . By homogeneity, we may assume $\beta = 1$. Then, Proposition 5.7 (2) and the linear independence of $\{\iota(u_j)\}_{j=1}^{\delta_v}$ force the coefficients $k_j \alpha_{u_j}$ in (6.41) to be the ones used to write $\iota(v)$ as an element of $(\Delta_{\text{Star}_\Gamma(v)})^\circ$. Since $\alpha_{u_j} > 0$ for all j , we have $k_j > 0$ for all j , as we wanted.

To finish, we argue that (6.40) has a solution with $k_{j_0} > 0$ by the balancing condition. This forces $\beta \neq 0$ in (6.41), and so $k_j \neq 0$ for all j by the previous discussion. This concludes our proof. $\square$

7 Splice type singularities are Newton non-degenerate

In this section we discuss the Newton non-degeneracy of splice type systems (see Theorem 7.3), following the original framework introduced by Khovanskii (see Definition 4.3). This property only involves the initial forms of the generators of the system $\mathcal{S}(\Gamma)$, as opposed to the initial ideals of the ideal generated by the elements of the system $\mathcal{S}(\Gamma)$. In turn, Theorem A.1 ensures the w -initial forms of $(F_{v,i})_{v,i}$ generate the w -initial ideal of $\langle \mathcal{S}(\Gamma) \rangle$, for each rational weight vector w in the positive local tropicalization of $\mathcal{S}(\Gamma)$, thus giving an alternative proof of Theorem 6.23. Furthermore, Newton non-degeneracy implies that condition (3) of Proposition 3.15 holds for every cell in $\iota(\Gamma)$, thus showing that the splice fan of Γ is a standard tropicalizing fan for the germ defined by the system $\mathcal{S}(\Gamma)$, in the sense of Definition 3.17. This is the content of Corollary 7.10. In turn, Corollary 7.14 gives an alternative proof of a recent theorem of de Felipe, González Pérez and Mourtada [7], which resolves any germ of a reduced plane curve by one toric morphism after a reembedding in a higher dimensional complex affine space. The section concludes with an open question regarding embedding dimension of splice type surface singularities preserving the Newton non-degeneracy property (see Question 7.17).

Our first result is a direct consequence of Lemma 2.19 and (2.10). It allows us to determine the w_u -initial form of each series $F_{v,i}$ in the system $\mathcal{S}(\Gamma)$ for each node u of Γ . This computation is central to all arguments in this section:

Proposition 7.1 *For each pair of nodes u, v in Γ , and each $i \in \{1, \dots, \delta_v - 2\}$, we have:*

$$\text{in}_{w_u}(F_{v,i}) = \text{in}_{w_u}(f_{v,i}) = \begin{cases} f_{u,i} & \text{if } u = v, \\ f_{v,i} - c_{v,[u,v]} \underline{z}^{m_{v,e}} & \text{otherwise,} \end{cases}$$

where e is the unique edge adjacent to v with $e \subseteq [v, u]$.

The second formula, for the case where $u \neq v$, means that we remove from $f_{v,i}$ the term associated to the edge starting from v in the direction of u .

Remark 7.2 Recall from Definition 2.20 that rooted splice diagrams Γ_r with $n + 1$ leaves (including the root r) give rise to end-curves in $\mathbb{C}^n$. Consider a fixed strict splice type system $\mathcal{S}(\Gamma)$ associated to Γ giving rise to this curve and let u be the unique node of Γ_r adjacent to r . In view of Proposition 7.1, the curve $\mathcal{C}_r$ is defined by the vanishing set of z_r and the w_u -initial forms of the $n - 1$ polynomials in the system $\mathcal{S}(\Gamma)$, viewed in $\{0\} \times \mathbb{C}^n$.

Next, we set up notation that we will use throughout this section. For each weight vector $w \in (\mathbb{R}_{>0})^n$ we define:

$$J_w := (\text{in}_w(F_{v,i}) : v \text{ node of } \Gamma, i = 1, \dots, \delta_v - 2) \subset \mathbb{C}[z_{\lambda_1}, \dots, z_{\lambda_n}], \quad (7.1)$$

and let Z_w be the subscheme of $\mathbb{C}^n$ defined by J_w .

Next, we state the main result of this section. The rest of the section is devoted to its proof.

Theorem 7.3 *The splice type system $\mathcal{S}(\Gamma)$ is a Newton non-degenerate complete intersection system.*

Proof We certify the two conditions from Definition 4.3 starting with the regularity of the sequence defined by $\mathcal{S}(\Gamma)$. This property follows by Lemma 7.4 and Theorem A.1. Indeed, since the weight vector w_u associated to any node u of Γ lies in $(\mathbb{Z}_{>0})^n$, and Z_{w_u} is a complete intersection of dimension two with defining ideal J_w by the lemma, we conclude that the $n - 2$ generators of J_w form a regular sequence in $\mathcal{O}$. In turn, the theorem confirms that the series $\{F_{v,i}\}_{v,i}$ form a regular sequence in $\mathcal{O}$.

Recall from Definition 3.4 that $Z_w \cap (\mathbb{C}^*)^n$ is the empty-set whenever $w \notin \text{Trop}_{>0}(\mathcal{S}(\Gamma))$. Thus, it suffices to certify the linear independence condition of the differentials of $\text{in}_w(F_{v,i})$ when $w \in (\mathbb{R}_{>0})^n \cap \text{Trop}_{>0}(\mathcal{S}(\Gamma))$. In turn, by Theorem 6.2 it is enough to restrict our analysis to vectors $w \in (\mathbb{R}_{>0})^n \cap \iota(\Gamma)$.

Since the generators of J_w form a regular sequence in $\mathcal{O}$ by Corollary 7.7, Proposition 4.5 simplifies our task: we need only verify that $Z_w \cap (\mathbb{C}^*)^n$ is smooth for each such w . We analyze three cases, depending on the nature of the unique cell of $\iota(\Gamma)$ containing w in its relative interior. Each case matches the settings of Lemmas 7.4, 7.5 and 7.6 below.

First, we pick a node u of Γ and consider the weight vector $w = w_u/|w_u|$. By Lemma 7.4, the components of the scheme $Z_{w_u} \subseteq \mathbb{C}^n$ are Pham–Brieskorn–Hamm singularities, and they can only meet at coordinate subspaces of $\mathbb{C}^n$. Furthermore, the Hamm conditions ensure that these germs are smooth away from the origin. This fact confirms the smoothness of $Z_{w_u} \cap (\mathbb{C}^*)^n$.

Next, we consider a weight vector $w \in \iota([\lambda, u])^\circ$, where λ is a leaf of Γ and u is a node of Γ adjacent to λ . Then, by Lemma 7.5, Z_w is a cylinder over an end-curve $\mathcal{C}_\lambda$ associated to the rooted diagram Γ_λ . By Theorem 2.21, such curves are smooth away from the origin. This confirms that $Z_w \cap (\mathbb{C}^*)^n$ is smooth in this case as well.

Finally, we fix a weight vector $w \in \iota([u, u'])^\circ$ where u, u' are adjacent nodes of Γ . Then, Lemma 7.6 confirms that the scheme Z_w is the product of two end-curves $\mathcal{C}_{\Gamma'_u}$ and $\mathcal{C}_{\Gamma'_v}$ associated to two splice subdiagrams of Γ rooted at u and v , respectively. More precisely, we have $\Gamma' := [u, \partial_{u,[u,v]}\Gamma]$ and $\Gamma'' := [v, \partial_{v,[u,v]}\Gamma]$ in the notation of Definition 2.13. By Theorem 2.21, these curves are smooth in the corresponding ambient tori, so $Z_w \cap (\mathbb{C}^*)^n$ is smooth. This concludes our proof. $\square$

The next three lemmas characterize the ideal J_w associated to points in the relative interior of each cell in the piecewise-linearly embedded tree $\iota(\Gamma) \hookrightarrow \Delta_{n-1}$.

Lemma 7.4 *Let u be a node of Γ and let $w = w_u$. Let Γ_0 be the star of Γ at u , viewed as a splice diagram with inherited weights around u . Then:*




Fig. 7 Rooted splice diagrams used in the proofs of Lemmas 7.4, 7.6 and 7.11. The vertex u' on the right is allowed to be a leaf of Γ , whereas u is always a node

1. $J_w = \langle \text{in}_w(f_{v,i}) : v \text{ node of } \Gamma, i = 1, \dots, \delta_v - 2 \rangle$;
2. Z_w is a complete intersection of dimension two;
3. Z_w has no irreducible component contained in a coordinate hyperplane of $\mathbb{C}^n$, so J_w is monomial-free;
4. all components of Z_w are obtained as images of torus-translated monomial maps $\mathbb{C}^{\delta_u} \rightarrow \mathbb{C}^n$ and their preimages lie in Pham–Brieskorn–Hamm complete intersections in $\mathbb{C}^{\delta_u}$ determined by Γ_0 ;
5. the intersection of any two distinct components of Z_w lies outside of $(\mathbb{C}^*)^n$.

Proof Item (1) follows from the identity $\text{in}_{w_u}(F_{v,i}) = \text{in}_{w_u}(f_{v,i})$ for all v and i , seen in Proposition 7.1. To prove the remaining items, we let $\boxed{k}$ be the number of nodes of Γ adjacent to u and $\boxed{T_1}, \dots, \boxed{T_k}$ be the branches of Γ adjacent to u containing these nodes, as seen in Fig. 7. Set $\boxed{\Gamma_i} := [T_i, u]$ and express J_w as the sum of the following $k + 1$ ideals:

$$\begin{aligned} \boxed{J_0} &:= \langle \text{in}_w(f_{u,j}) : j = 1, \dots, \delta_u - 2 \rangle, \\ \boxed{J_i} &:= \langle \text{in}_w(f_{v,j}) : v \text{ node of } \Gamma_i, j = 1, \dots, \delta_v - 2 \rangle, \text{ for } i = 1, \dots, k. \end{aligned}$$

Note that the generators of each J_i with $i \neq 0$ lie in $\mathbb{C}[z_\lambda : \lambda \in \partial T_i]$. In particular, no variable z_λ with λ adjacent to u appears in them. We call these leaves $\boxed{\lambda_1}, \dots, \boxed{\lambda_l}$, with $k + l = \delta_u$.

To simplify notation, we write $\boxed{Z} := Z_w$. We start by characterizing the components of Z . Each ideal J_i with $i \in \{1, \dots, k\}$ defines an end-curve $\mathcal{C}_i$ determined by the splice diagram Γ_i rooted at u , as seen in the center of Fig. 7. In turn, Theorem 2.21 ensures that each of the $(g_i$ -many) components of $\mathcal{C}_i$ admits a (torus-translated) monomial parameterization of the form

$$\mathbb{C}[z_\lambda : \lambda \in \partial_u \Gamma_i] \rightarrow \mathbb{C}[t_i] \quad \text{where } z_\lambda \mapsto c_{\lambda,i}^{(u)} t_i^{\ell_{u,\lambda}/g_i}, \quad (7.2)$$

with $c_{\lambda,i}^{(u)} \neq 0$ for all λ, i . Thus, each component of $\mathcal{C}_1 \times \dots \times \mathcal{C}_k$ can be parameterized using $\mathbb{C}^k$ by combining these (torus-translated) monomial maps.

Substituting the expressions from (7.2) for the chosen component of each $\mathcal{C}_i$ into the generators of J_0 shows that the closure of each component of Z can be parameterized

using a component of a splice type singularity defined by the diagram Γ_0 . Note that the Hamm determinant conditions at u arising from $\mathcal{S}(\Gamma_0)$ agree with those determined by the generators of J_0 up to multiplying the columns corresponding to the branches Γ_i , $i = 1, \dots, k$ by non-zero constants. Therefore, each component of Z is the image of a (torus-translated) monomial map from $\mathbb{C}^{\delta_u} = \mathbb{C}^k \times \mathbb{C}^\ell$ restricted to a Pham–Brieskorn–Hamm complete intersection defined by Γ_0 . In particular, no component of Z lies in a coordinate subspace of $\mathbb{C}^n$, so J_w is monomial-free. This proves both (3) and (4).

Since Z is equidimensional of dimension two and it is defined by $n - 2$ polynomial equations, it is a complete intersection. This proves (2). It remains to address (5) when Z is not irreducible.

To determine the intersection of two distinct components (say, Z_1 and Z_2) of Z we exploit the parameterization described earlier. Let $I \subseteq \{1, \dots, k\}$ be the collection of indices i for which the projections of Z_1 and Z_2 to $\text{Spec}(\mathbb{C}[t_i])$ disagree. If $I = \emptyset$, then the two components are parameterized using the same Pham–Brieskorn–Hamm system of equations associated to Γ_0 and the same (torus-translated) monomial map. This cannot happen since such germs are irreducible and $Z_1 \neq Z_2$.

Next, assume $|I| \geq 1$. In this setting, the projection of $Z_1 \cap Z_2$ to $\mathbb{C}^{\partial_u \Gamma_i}$ for each $i \in I$ lies in the intersection of two components of the end-curve $\mathcal{C}_i$, which can only be the origin of $\mathbb{C}^{\partial_u \Gamma_i}$ by Theorem 2.21. This means that $Z_1 \cap Z_2$ lies in the coordinate subspace of $\mathbb{C}^n$ defined by the vanishing of all z_λ with $\lambda \in \bigcup_{i \in I} \partial \Gamma_i$. This concludes our proof. $\square$

Lemma 7.5 *Let u be a node of Γ adjacent to a leaf λ and pick $w \in [\iota(u), \iota(\lambda)]^\circ$. Then, Z_w is a cylinder over a monomial curve in $\mathbb{C}^{n-1}$ with $\gcd(\ell_{\lambda, \mu} : \mu \in \partial \Gamma \setminus \{\lambda\})$ many components and J_w is monomial-free.*

Proof We prove that J_w defines the cylinder over the end-curve $\mathcal{C}_\lambda$ associated to the rooted splice diagram Γ_λ . A simple inspection shows that for all nodes v of Γ and each $i \in \{1, \dots, \delta_v - 2\}$, the initial form $\text{in}_w(F_{v,i})$ agrees with the corresponding polynomial $h_{v,i}$ defining $\mathcal{C}_\lambda$ (see Definition 2.20). Indeed, if we write w as

$$w = \alpha \iota(\lambda) + (1 - \alpha) \iota(u) \quad \text{with} \quad 0 < \alpha < 1, \quad (7.3)$$

then the defining properties of $g_{v,i}$ ensure that $\text{in}_w(F_{v,i}) = \text{in}_w(f_{v,i})$ for all v, i . In turn, combining this fact with Lemma 2.19 and Proposition 7.1 yields

$$w \cdot m_{v,e} \geq (1 - \alpha) \frac{\ell_{u,v}}{|w_u|} \quad \text{for each edge } e \text{ in } \text{Star}_\Gamma(v).$$

Furthermore, equality holds if and only if $e \not\subseteq [v, \lambda]$. Thus, $\text{in}_w(f_{v,i})$ is obtained from $f_{v,i}$ by dropping the admissible monomial at v pointing towards λ . In particular, these initial forms do not involve z_λ .

The above discussion shows that $\boxed{J'_w} := J_w \cap \mathbb{C}[z_\mu : \mu \in \partial \Gamma \setminus \{\lambda\}]$ defines $\mathcal{C}_\lambda$.

The general point of this curve lies in $(\mathbb{C}^*)^{n-1}$ by Theorem 2.21, so J'_w is monomial-free. In turn, the same result confirms that J'_w defines a reduced complete intersection,

smooth outside the origin, and with $\gcd(\ell_{\lambda,\mu} : \mu \in \partial \Gamma \setminus \{\lambda\})$ many irreducible components. Since J_w is obtained from J'_w by base change to $\mathbb{C}[z_\lambda : \lambda \in \partial \Gamma]$, and both ideals admit a common generating set, the result follows. $\square$

Given two adjacent nodes u and u' of Γ , we let T' be the branch of Γ adjacent to u' containing u . Similarly, we let T be the branch of Γ adjacent to u and containing u' . Our final lemma is analogous to Lemma 7.5:

Lemma 7.6 *Let u and u' be two adjacent nodes of Γ and pick $w \in [\iota(u), \iota(u')]^\circ$. Then, Z_w is isomorphic to a product of two monomial curves in $\mathbb{C}^{|\partial T'|} \times \mathbb{C}^{|\partial T|}$. Furthermore, the number of irreducible components of Z_w equals $\gcd(\ell'_{u',\lambda} : \lambda \in \partial T') \gcd(\ell'_{u,\mu} : \mu \in \partial T)$ and J_w is monomial-free.*

Proof We write $w = \alpha w_u + (1 - \alpha) w_{u'}$ with $0 < \alpha < 1$ and follow the same proof-strategy as in Lemma 7.5. By Theorem 5.11, we can find a unique $r \in [u, u']^\circ$ with $\iota(r) = w/|w|$ (see the rightmost picture in Fig. 7).

The conditions on α together with Proposition 7.1 guarantee that for each node v in T' , $\text{in}_w(F_{v,i}) = \text{in}_w(f_{v,i})$. Furthermore, each form is obtained from $f_{v,i}$ by dropping the admissible monomial at v pointing towards u' . This observation and the symmetry between u and u' determine a partition of the generating set of J_w , where each initial form $\text{in}_w(F_{v,i})$ for any node v of T' (respectively, in T) only involves the leaves of T' (respectively, of T). Thus, each set determines an end-curve for the diagrams $[T', r]$ and $[T, r]$, respectively, both rooted at r .

By construction, J_w defines the product of these two monomial curves. Since each of them is a complete intersection in their respective ambient spaces, the same is true for J_w . The number of components is determined by Theorem 2.21. Since Z_w meets $(\mathbb{C}^*)^n$, we conclude that J_w is monomial-free. $\square$

The next result is a direct consequence of Theorem A.1 and the previous three lemmas. It gives an alternative proof of Theorem 6.23 since for any w in $\iota(\Gamma)^\circ$, these lemmas confirm that J_w is monomial-free.

Corollary 7.7 *For any strictly positive vector w in the splice fan of Γ , we have $\text{in}_w(\langle \mathcal{S}(\Gamma) \rangle) = J_w$, where $\langle \mathcal{S}(\Gamma) \rangle$ is the ideal generated by the splice type system $\mathcal{S}(\Gamma)$.*

Proof By construction, we have $J_w \subseteq \text{in}_w(\mathcal{S}(\Gamma))$ for each $w \in (\Delta_{n-1})^\circ \cap \iota(\Gamma)$. In addition, Lemmas 7.4, 7.5 and 7.6 imply that the generators of J_w form a regular sequence in the ring of convergent power series $\mathbb{C}\{z_\lambda : \lambda \in \partial \Gamma\}$ near the origin. Thus, Theorem A.1 implies that $J_w = \text{in}_w(\mathcal{S}(\Gamma))$ as ideals of $\mathbb{C}\{z_\lambda : \lambda \in \partial \Gamma\}$ for each $w \in \mathbb{Q}^n \cap \mathbb{R}_{>0} \langle \iota(\Gamma) \rangle$. Finally, since these ideals are constant when we consider all rational points in the relative interior of any fixed edge of $\iota(\Gamma)$, Lemma 7.8 below confirms that the same must be true for all points in these open segments. $\square$

Lemma 7.8 *Fix an edge $[u, v]$ of Γ . If J_w is constant for all $w \in \iota([u, v]) \cap (\mathbb{Q}_{>0})^n$, the same is true for all $w \in \iota([u, v])^\circ$.*

Proof We set $\boxed{J} := \langle \mathcal{S}(\Gamma) \rangle$, $\boxed{\sigma} := \mathbb{R}_{\geq 0} \langle \iota([u, v]) \rangle$, and fix a weight vector $w' \in \sigma^\circ \setminus \mathbb{Q}^n$. We wish to show that $\text{in}_{w'}(J) = \text{in}_w(J)$ for all $w \in \sigma^\circ \cap \mathbb{Q}^n$. We prove this equality by double-inclusion.

The inclusion $\text{in}_{w'}(J) \subseteq \text{in}_w(J)$ is a direct consequence of the following claim:

Claim. For each $f \in J$, we have $\text{in}_{w'}(f) \in \text{in}_w(J)$.

Proof Throughout our reasoning, we need the following auxiliary fact. By [28, Lemma 2.4.6], given w_1 and w_2 in $(\mathbb{R}_{>0})^n$, there exists $\varepsilon > 0$ with

$$\text{in}_{w_1}(\text{in}_{w_2}(f)) = \text{in}_{w_2 + \varepsilon' w_1}(f) \quad (7.4)$$

for all $0 < \varepsilon' < \varepsilon$ and for each $f \in \mathbb{C}[z_\lambda : \lambda \in \partial \Gamma]$. Restricting further the value of $\varepsilon > 0$ to require that $w_2 + \varepsilon' w_1 \in (\mathbb{R}_{>0})^n$ whenever $0 < \varepsilon' < \varepsilon$, extends the validity of (7.4) to convergent power series $f \in \mathbb{C}\{z_\lambda : \lambda \in \partial \Gamma\}$.

Since $w' \in (\mathbb{R}_{>0})^n$ by construction, the initial form $\text{in}_{w'}(f)$ is a polynomial. Its Newton fan is rational and complete, and we get an induced rational fan (denoted by Σ) on the cone σ by common refinement. Thus, w' cannot be a ray of Σ (the vector is not rational), so it lies in a 2-dimensional cone τ of Σ . By construction, $\tau \subseteq \sigma$. Picking any $w'' \in \tau \cap (\mathbb{Q}_{>0})^n$, our earlier discussion ensures the existence of $\varepsilon > 0$ with

$$\text{in}_{w'}(f) = \text{in}_{w''}(\text{in}_{w'}(f)) = \text{in}_{w' + \varepsilon' w''}(f) \quad \text{for all } 0 < \varepsilon' < \varepsilon.$$

Since the set $\{w' + \varepsilon' w'' : 0 < \varepsilon' < \varepsilon\}$ contains a point w in $\sigma^\circ \cap \mathbb{Q}^n$, we conclude that $\text{in}_{w'}(f) \in \text{in}_w(J)$. $\diamond$

For the reverse inclusion, we argue as follows. Direct calculations used in the proofs of Lemmas 7.5 and 7.6 confirm the identities

$$\text{in}_w(F_{v,i}) = \text{in}_{w'}(F_{v,i}) \quad \text{for all nodes } v \in \Gamma \text{ and each } i \in \{1, \dots, \delta_v - 2\}.$$

In turn, Theorem A.1 ensures that this collection generates $\text{in}_w(J)$. Thus, $\text{in}_w(J) \subseteq \text{in}_{w'}(J)$, as desired. $\square$

Remark 7.9 The geometric information collected in Lemmas 7.5 and 7.6 combined with Corollary 7.7 determines the tropical multiplicities of $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$. Using Remark 2.7 and Definition 3.20, we have:

1. if $\tau = \mathbb{R}_{\geq 0}\iota([\lambda, u])$ for a node u of Γ , then τ has multiplicity

$$\frac{1}{d_{u,\lambda}} \gcd(\ell_{u,\mu} : \mu \in \partial \Gamma \setminus \{\lambda\});$$

2. if $\tau = \mathbb{R}_{\geq 0}\iota([u, v])$ for two adjacent nodes u and v of Γ , then τ has multiplicity

$$\frac{1}{d_{u,v}d_{v,u}} \gcd(\ell_{u,\lambda} : \lambda \in \partial \Gamma, u \in [\lambda, v]) \gcd(\ell_{v,\lambda} : \lambda \in \partial \Gamma, v \in [\lambda, u]).$$

This information completes the characterization of $\text{Trop}_{>0}(\mathcal{S}(\Gamma))$ as a tropical object, i.e., as a weighted balanced polyhedral fan. We use this data in Sect. 8 when discussing how to recover Γ from its splice fan.

Our next statement follows naturally from Proposition 3.15 and Corollary 7.7:

Corollary 7.10 *The splice fan of Γ is a standard tropicalizing fan for the germ $(X, 0)$ defined by $\mathcal{S}(\Gamma)$, in the sense of Definition 3.17.*

Next, we show how we can use their local tropicalizations to recover some known facts about splice type systems and their associated end-curves from [38, 39]. Our first statement discusses the intersections of the initial degenerations of the germ defined by $\mathcal{S}(\Gamma)$ will suitable codimension-2 coordinate subspaces of $\mathbb{C}^n$:

Lemma 7.11 *Let w be a vector in $(\mathbb{R}_{>0})^n$ which lies in the embedded splice diagram $\iota(\Gamma)$, and let λ and μ be two leaves of Γ in different branches of $\iota(\Gamma)$ adjacent to w . Then, the system*

$$\begin{cases} \text{in}_w(F_{v,i}) = 0, & v \text{ is a node of } \Gamma, \quad i \in \{1, \dots, \delta_v - 2\}, \\ z_\lambda = z_\mu = 0 \end{cases} \quad (7.5)$$

has $0 \in \mathbb{C}^n$ as its only solution.

Proof If $w = \iota(u)$ is the (normalized) weight vector corresponding to a node u of Γ , this statement is proved as part of [38, Theorem 2.6], more precisely, on [38, p. 710]. For any point w in $\iota(\Gamma) \cap (\Delta_n)^\circ$, the claim follows by similar arguments, but for the reader's convenience, we give a complete proof that includes the node case. Our reasoning is guided by Fig. 7, and it is based on the techniques discussed in the proofs of Lemmas 7.4, 7.5 and 7.6.

By Theorem 5.11, we can find a unique point r in Γ with $w = \iota(r)$. As in the figure, we write $r = u$ if u is a node of Γ . Otherwise, r is in the relative interior of a unique edge $[u, u']$, where u is a node of Γ but u' is allowed to be a leaf of Γ . The proofs of the three aforementioned lemmas ensure that $\text{in}_w(F_{v,j}) = \text{in}_w(f_{v,j})$ for all nodes v in Γ and $j \in \{1, \dots, \delta_v - 2\}$. Thus, to prove the statement, we may assume that $g_{v,i} = 0$ for all v, i .

By abuse of notation, we think of the point r as a node of Γ (potentially, of valency two). We assume r is adjacent to k nodes and ℓ leaves of Γ , with $k + \ell = \delta_r$. We let $T_1, \dots, T_{\delta_r}$ be the branches of Γ adjacent to T . We assume $T_1, \dots, T_k$ contain nodes of Γ , whereas the remaining branches are singletons. We view the single vertex of a singleton branch as a leaf of this branch.

Using r , we build a collection of rooted splice diagrams $\Gamma_1, \dots, \Gamma_k$ rooted at r . Each Γ_i is obtained as the convex hull $[T_i, r]$. In particular, if $r \in [u, u']^\circ$ (as in the right of the figure), then k is the number of nodes of Γ among $\{u, u'\}$. We assume that $u \in T_1$ whenever $\delta_r = 2$.

Throughout, we consider the ideals

$$J_i := \langle \text{in}_w(f_{v,j}) : v \text{ node of } \Gamma_i, j = 1, \dots, \delta_v - 2 \rangle \quad \text{for } i \in \{1, \dots, k\}. \quad (7.6)$$

The generators of J_i are among the series of the system (7.5) per our assumption on the tails $g_{v,i}$. Since r is not a node of Γ_i , Lemma 2.19 confirms that J_i is generated by

polynomials in $\mathbb{C}[z_{\lambda'} : \lambda' \in \partial T_i]$. Viewed in this ring, the ideal J_i defines an end-curve $\mathcal{C}_i$ in $\mathbb{C}^{|\partial T_i|}$ associated to the rooted diagram Γ_i (see Remark 7.2).

Next, we consider a solution $\underline{p} \in \mathbb{C}^n$ of the system (7.5). Our assertion that $\underline{p}$ is the origin will follow by combining the next two claims:

Claim A. Fix $i \in \{1, \dots, \delta_r\}$ and assume that $p_{\lambda'} = 0$ for some leaf λ' of T_i . Then, $p_{\lambda} = 0$ for all leaves λ of T_i .

Proof If $i > k$, then $T_i = \{\lambda'\}$ and the statement is tautological. On the contrary, if $i \leq k$, then the projection of $\underline{p}$ to $\mathbb{C}^{|\partial T_i|}$ lies in the end-curve $\mathcal{C}_i$. Since $p_{\lambda'} = 0$, Theorem 2.21 guarantees that the projection of $\underline{p}$ is the origin of $\mathbb{C}^{|\partial T_i|}$, as we wanted. $\diamond$

Claim B. Assume that all coordinates of $\underline{p}$ indexed by two different branches (say, T_{i_1} and T_{i_2}) adjacent to r vanish. Then, the same is true for each remaining branch.

Proof If $\delta_r = 2$, there is nothing to show. On the contrary, if $\delta_r \geq 3$, we consider the splice type polynomials $f_{r,j}$ for $j \in \{1, \dots, \delta_r - 2\}$. By Proposition 7.1, $\text{in}_w(f_{r,j}) = f_{r,j}$ for each j .

Using the Hamm conditions, we may assume that each $f_{r,j}$ involves exactly three admissible monomials, two of which are $\underline{z}^{m_{r,T_{i_1}}}$ and $\underline{z}^{m_{r,T_{i_2}}}$. The third one equals $\underline{z}^{m_{r,T_{i_j}}}$. Since the first two monomials vanish along $\underline{p}$ by construction, we deduce that the same is true for the remaining monomial in $f_{r,j}$. Thus, for each of the remaining branches T_{i_j} adjacent to r we have $p_{\lambda'_j} = 0$ for some leaf λ'_j in T_{i_j} . Claim A ensures that the same is true for all leaves of T_{i_j} . This concludes our proof. $\diamond$

To finish our proof, it is enough to notice that the hypothesis of Claim B holds by combining Claim A with the fact that the leaves λ and μ belong to distinct branches of Γ adjacent to r . $\square$

Our second statement gives a stronger version of Corollary 2.23:

Corollary 7.12 Assume that the conditions of Lemma 7.11 hold and consider the map $F_w : \mathbb{C}^n \rightarrow \mathbb{C}^{n-2}$ obtained from the collection $\{\text{in}_w(F_{v,i})\}_{v,i}$, ordered appropriately. Then, the restriction of this map to the codimension-2 subspace $L = \{z_{\lambda} = z_{\mu} = 0\}$ of $\mathbb{C}^n$ is surjective.

Proof By Lemma 7.11, we see that the fiber of the restriction of F_w to L over the origin of $\mathbb{C}^{n-2}$ is 0-dimensional. By upper semicontinuity of fiber dimensions, the generic fiber of $F_{w|L}$ is also 0-dimensional. Since $\dim L = n - 2$, the map $F_{w|L}$ is dominant.

Since $F_{w|L}$ is defined by weighted homogeneous functions, it admits a projectivization as a map between weighted projective spaces. But a dominant projective map must be surjective. Thus, as an affine map, $F_{w|L}$ is surjective as well. $\square$

Next, we recover Theorem 2.16 (originally due to Neumann and Wahl) by combining Corollary 6.18 with the following result:

Corollary 7.13 The singularity defined by the splice type system $\mathcal{S}(\Gamma)$ is isolated. In particular, it is also irreducible.

Proof Let $(X, 0) \hookrightarrow \mathbb{C}^n$ be the germ defined by $\mathcal{S}(\Gamma)$. By Corollary 6.20, we know that $X \cap (\mathbb{C}^*)^n$ is dense in X . Since $\mathcal{S}(\Gamma)$ is a Newton non-degenerate complete intersection system by Theorem 7.3, we conclude that $X \subset \mathbb{C}^n$ admits an embedded toric resolution (see, e.g., Khovanskii [23, Sect. 2.7] and Oka [41, Chapter III, Theorem (3.4)]). In particular, for a suitable subdivision Σ of the splice fan of X , the corresponding toric morphism $\pi: \mathcal{X}_\Sigma \rightarrow \mathbb{C}^n$ induces an embedded resolution of the pair $(\mathbb{C}^n, X)$. But as we saw in Theorem 6.2, the local tropicalization of $\mathcal{S}(\Gamma)$ intersects the boundary of the non-negative orthant only along the canonical basis elements. It follows from this that the morphism π is an isomorphism outside the origin, i.e., π is a resolution of $(X, 0)$. We conclude that the singularity at the origin is isolated. As $(X, 0)$ is moreover a complete intersection of dimension two by Corollary 6.20, it is automatically irreducible by Hartshorne's connectedness theorem [21, Theorem 2.2] (see also [9, Theorem 18.12]), which states that a complete intersection singularity cannot be disconnected by removing a closed subgerm of codimension at least two. $\square$

We end this section by showing how to use our results to get embedded resolutions of complex plane curve singularities by composing re-embeddings of $\mathbb{C}^2$ into higher-dimensional smooth spaces $\mathbb{C}^n$ with toric modifications of $\mathbb{C}^n$. The fact that such resolutions are possible was proven by Goldin and Teissier [17] in the irreducible case and recently by de Felipe, González Pérez and Mourtada [7, Theorem 2.27] in full generality.

Corollary 7.14 *Let $(X, 0) \hookrightarrow \mathbb{C}^2$ be the germ of a reduced complex analytic plane curve. Then, the ambient germ $(\mathbb{C}^2, 0)$ can be holomorphically re-embedded into a suitable higher-dimensional germ $(\mathbb{C}^n, 0)$ in such a way that the induced germ $(X, 0) \hookrightarrow \mathbb{C}^n$ can be resolved by a single toric modification of $\mathbb{C}^n$.*

Proof This result is a consequence of Theorem 7.3, as we now explain. Indeed, consider a completion $(\hat{X}, 0) \hookrightarrow \mathbb{C}^2$ of the input germ $(X, 0) \hookrightarrow \mathbb{C}^2$ in the sense of [15, Definition 1.4.15]. By construction, $(\hat{X}, 0)$ is also the germ of a reduced plane curve, it contains $(X, 0)$ as a subgerm and it admits an embedded resolution (that is, a modification $\pi: S \rightarrow \mathbb{C}^2$ where S is smooth and the total transform of $\hat{X}$ on S has normal crossings) such that the strict transform of $\hat{X}$ intersects all the leaf components of the exceptional divisor. Here, leaf components correspond to leaves of the dual graph of the exceptional divisor.

Moreover, possibly after extra blowups at points, the modification $\pi: S \rightarrow \mathbb{C}^2$ can be chosen to ensure that each leaf component is intersected by exactly one irreducible component of the strict transform of $\hat{X}$. Since all the irreducible components of $(\hat{X}, 0)$ are principal divisors on $(\mathbb{C}^2, 0)$, and $(\mathbb{C}^2, 0)$ has an integral homology sphere link, Neumann and Wahl's *end-curve theorem* [39, Theorem 4.1 (3)] guarantees the existence of a holomorphic embedding $\phi: (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^n, 0)$ such that $(\phi(\mathbb{C}^2), 0)$ is a splice type singularity. Furthermore, the irreducible components of $(\phi(\hat{X}), 0)$ are exactly the intersections of $(\phi(\mathbb{C}^2), 0)$ with the coordinate hyperplanes of $(\mathbb{C}^n, 0)$.

Let Γ be the splice diagram of the splice type singularity $(\phi(\mathbb{C}^2), 0) \hookrightarrow (\mathbb{C}^n, 0)$. Consider the splice fan $\mathcal{F}$ of Γ , in the sense of Definition 6.1. By Corollary 7.10, $\mathcal{F}$ is a standard tropicalizing fan of $(\phi(\mathbb{C}^2), 0)$, in the sense of Definition 3.17. Consider a regular subdivision $\mathcal{F}'$ of $\mathcal{F}$. It is also a standard tropicalizing fan of $(\phi(\mathbb{C}^2), 0)$.



Fig. 8 Two splice diagrams used in Example 7.16: the right one is obtained from the left one by attaching an extra leaf through a weight one edge

Let $\mathcal{X}_{\mathcal{F}'}$ be the toric variety defined by the fan $\mathcal{F}'$ and let $\pi_{\mathcal{F}'}: \mathcal{X}_{\mathcal{F}'} \rightarrow \mathbb{C}^n$ be the birational toric morphism defined by the inclusion of the support of the fan $\mathcal{F}'$ in the non-negative orthant $(\mathbb{R}_{\geq 0})^n$ of the weight lattice $\mathbb{Z}^n$ of $\mathbb{C}^n$.

Consider the strict transform S' of $(\phi(\mathbb{C}^2), 0)$ by $\pi_{\mathcal{F}'}$ and denote by

$$\pi': S' \rightarrow (\phi(\mathbb{C}^2), 0)$$

the restriction of $\pi_{\mathcal{F}'}$ to S' . By construction, the support of $\mathcal{F}'$ equals the local tropicalization of $(\phi(\mathbb{C}^2), 0)$ in $\mathbb{C}^n$. Thus, Proposition 3.19 (1) implies that the bimeromorphic morphism π' is proper.

By Theorem 7.3, $(\phi(\mathbb{C}^2), 0)$ is defined by a Newton non-degenerate complete intersection system inside $\mathbb{C}^n$. Therefore, the scheme-theoretic intersections of S' with the orbits of the toric variety $\mathcal{X}_{\mathcal{F}'}$ are either empty or smooth (see [28, Remark 6.4.18]). As a consequence, S' intersects transversally the toric boundary of $\mathcal{X}_{\mathcal{F}'}$ (see [6, Proposition 3.9]). As $\mathcal{F}'$ is a regular fan, the toric variety $\mathcal{X}_{\mathcal{F}'}$ is smooth. Therefore, S' is also smooth and the total transform of $(\phi(\hat{X}), 0)$ in S' is a normal crossings curve.

This fact and the properness of π' imply that π' is an embedded resolution of $(\phi(\hat{X}), 0) \hookrightarrow (\phi(\mathbb{C}^2), 0)$. As $(X, 0) \hookrightarrow (\hat{X}, 0)$, the morphism π' is also an embedded resolution of $(\phi(X), 0) \hookrightarrow (\phi(\mathbb{C}^2), 0)$. Therefore, $\phi^{-1} \circ \pi': S' \rightarrow (\mathbb{C}^2, 0)$ is an embedded resolution of $(X, 0) \hookrightarrow (\mathbb{C}^2, 0)$, as we wanted to show. $\square$

Remark 7.15 Results from [14, Sect. 5] allow to describe the splice diagram associated to a plane curve singularity in terms of the Newton–Puiseux series of its branches. Applying Corollary 7.10 to this splice diagram yields a concrete description of the local tropicalization of the embedding ϕ from the proof of Corollary 7.14 in terms of standard combinatorial invariants of the given plane curve singularity. This method is similar to the one used by de Felipe, González Pérez and Mourtada in [7, Sect. 3] to characterize such local tropicalizations.

The construction of splice type systems by Neumann and Wahl implies that the embedding dimension of a splice type singularity is bounded from above by the number of leaves of the associated splice diagram. If an edge ending in a leaf has weight one, then it can be removed by simple elimination to produce a germ in lower dimension. As the following example illustrates, this operation need not necessarily produce a splice type singularity.

Example 7.16 Consider the splice diagram Γ on the right of Fig. 8 satisfying the edge determinant and semigroup conditions. We consider two strict splice type systems for

Γ :

$$\begin{cases} z_1^2 + z_2^3 + z_3 z_4^3 = 0, \\ z_1 z_2^4 + z_3^5 + z_4^2 + z_5 = 0, \\ 2 z_1 z_2^4 + 3 z_3^5 - 2 z_4^2 + z_5 = 0, \end{cases} \quad \begin{cases} z_1^2 + z_2^3 + z_3 z_4 z_5 = 0, \\ z_1 z_2^4 + z_3^5 + z_4^2 + z_5 = 0, \\ 2 z_1 z_2^4 + 3 z_3^5 - 2 z_4^2 + z_5 = 0. \end{cases} \quad (7.7)$$

Eliminating the z_5 variable from the left system yields a splice system associated to the splice diagram on the left of the figure, namely $z_1^2 + z_2^3 + z_3 z_4^3 = -z_1 z_2^4 - 2 z_3^5 + 3 z_4^2 = 0$. On the other hand, elimination on the rightmost system in (7.7) produces a system of two equations in four unknowns that is not of splice type:

$$\begin{cases} z_1^2 + z_2^3 + z_3^6 z_4 - 4 z_3 z_4^3 = 0, \\ -z_1 z_2^4 - 2 z_3^5 + 3 z_4^2 = 0. \end{cases}$$

□

But even if no such terminal edge with weight one exists, the embedding dimension of a splice type singularity can still be smaller than the number of leaves in the splice diagram. In particular, [39, Example 3] exhibits a hypersurface singularity $Z(f)$ in $\mathbb{C}^3$ realizing a splice type surface singularity $(X, 0)$ whose associate splice diagram Γ has six leaves.

A simple calculation reveals that the standard local tropicalization of this hypersurface is the cone over a star-shaped tree with three leaves (corresponding to the standard coordinate basis vectors) and a single node w . The w -initial form of f is non-reduced so the Newton non-degeneracy condition fails for this hypersurface presentation, despite the fact that the splice type system $\mathcal{S}(\Gamma)$ defining $(X, 0)$ is a Newton non-degenerate complete intersection system by Theorem 7.3.

Motivated by this example, we define the *Newton non-degenerate embedding dimension* $\text{edim}_{\text{NND}}(X, 0)$ of an abstract germ $(X, 0)$ as the smallest $n \geq 0$ such that $(X, 0)$ may be embedded in $\mathbb{C}^n$ as a Newton non-degenerate subgerm. In this context, the following question arises naturally:

Question 7.17 *Fix a splice diagram with n leaves satisfying the edge determinant and semigroup conditions. Assume that no edge of Γ ending in a leaf has weight one. Let $(X, 0)$ be a splice type singularity associated to Γ . Is it true that $\text{edim}_{\text{NND}}(X, 0) = |\partial \Gamma|$?*

8 Recovering splice diagrams from splice fans

Throughout this section we assume that the splice type diagrams satisfy the edge determinant condition of Definition 2.9 and the semigroup condition of Definition 2.13. The construction of splice fans from splice diagrams introduced in Definition 6.1 raises a natural question: how much data about Γ can be recovered from its splice fan, decorated with the tropical multiplicities? The main result of this section answers this question under the following coprimality restrictions, which are central to [39]:




Fig. 9 A splice diagram Γ which cannot be recovered uniquely from its splice fan, as in Remark 8.4. Here, d_1 may take the values 1, 2 or 4 and all tropical multiplicities of the splice fan equal 4

Definition 8.1 We say that a splice diagram Γ satisfying the edge determinant condition and the semigroup condition is *coprime* if the weights around each node of Γ are pairwise coprime.

As mentioned in Theorem 2.10, if Γ is coprime, then there exists a unique integral homology sphere 3-manifold $\Sigma(\Gamma)$ associated to Γ . However, there may be also non-integral rational homology sphere links with the same splice diagram, since Γ only determines the topological types of their universal abelian covers (see [38, p. 2]).

Our next result highlights the restrictions on the tropically weighted splice fan imposed by a coprime splice diagram. Its proof will be postponed until the end of this section. Precise formulas for the tropical multiplicities are given in Remark 7.9.

Theorem 8.2 *Let Γ be a coprime splice diagram. Then:*

1. *for each node v of Γ the vector $w_v \in N(\partial\Gamma) \simeq \mathbb{Z}^n$ is primitive;*
2. *all tropical multiplicities of $\text{Trop}_{>0}\langle S(\Gamma) \rangle$ equal one.*

The following is the main result in this section. It ensures that coprime splice diagrams can be recovered from their tropically weighted splice fans:

Theorem 8.3 *Assume that all tropical multiplicities on the splice fan equal one. Then, there is a unique coprime splice diagram Γ yielding the given weighted splice fan.*

The rest of the section is devoted to the proof of these two results. A series of lemmas and propositions will simplify the exposition.

Remark 8.4 Notice that the analog of Theorem 8.3 may fail if we drop the tropical multiplicity one restrictions. This can be seen by looking at the example in Fig. 9. For each choice of edge weight $d_1 = 1, 2$ or 4, the diagram Γ satisfies the semigroup and edge determinant conditions. Furthermore, all tropical multiplicities on the 2-dimensional cones of the splice fan equal four. For each value of d_1 we can choose systems $S(\Gamma)$ defining a germ in $\mathbb{C}^4$, whose local tropicalization is supported on the input splice fan.

Our first technical result will allow us to employ a pruning argument to prove Theorem 8.3. To this end, we use superscripts to indicate the underlying splice diagram considered for the computation of each linking number. The absence of a superscript refers to Γ . The same notation will be used for weight vectors.

Proposition 8.5 *Let $[u, v]$ be an internal edge of a splice diagram Γ . Let T be the branch of Γ adjacent to u and containing v . Consider $\Gamma' = [u, T]$ with weights around its nodes inherited from Γ . Then, the weighted tree Γ' also satisfies the semigroup and the edge determinant conditions.*

Proof We only need to check that the semigroup conditions hold for Γ' . The linking numbers for Γ involving a vertex v' of Γ' with $v' \neq u$ and a leaf λ of Γ can be obtained from those in Γ' via:

$$\ell_{v',\lambda} = \begin{cases} \ell_{v',\lambda}^{\Gamma'} & \text{if } \lambda \in \partial \Gamma' \cap \partial \Gamma, \\ \ell_{v',u}^{\Gamma'} \frac{\ell_{u,\lambda}}{d_{u,v}} & \text{otherwise.} \end{cases} \quad (8.1)$$

Since the semigroup condition at each v' holds for Γ , expression (8.1) implies the same is true for Γ' . $\square$

Assume that u and v are adjacent nodes of Γ and let Γ' be the associated splice diagram introduced in Proposition 8.5. Up to relabeling, we write $\partial \Gamma \setminus \partial \Gamma' = \{\lambda_1, \dots, \lambda_s\}$ for some s . Consider the following $n \times (n - s + 1)$ matrix with integer entries in block form obtained from (8.1):

$$A := \left(\begin{array}{c|c} \ell_{u,\lambda_1}/d_{u,v} & \\ \vdots & \\ \ell_{u,\lambda_s}/d_{u,v} & \mathbf{0} \\ \hline \mathbf{0} & \text{Id}_{n-s} \end{array} \right). \quad (8.2)$$

A direct computation yields the following identity for the tree Γ' :

Lemma 8.6 *For each vertex v' of Γ' with $v' \neq u$ we have $A w_{v'}^{\Gamma'} = w_{v'}$.*

Lemma 8.7 *Fix a coprime splice diagram Γ . Let u, v be two adjacent nodes of Γ , and let Γ' be the diagram from Proposition 8.5. Then, we have*

$$d_{v,u} = \gcd(\ell_{v,\lambda} : \lambda \in \partial \Gamma' \cap \partial \Gamma).$$

Proof The result follows by an easy induction on the number of nodes of Γ' . If Γ' has a single node, the identity holds by the coprimality of the weights around v . For the inductive step, we assume that v is adjacent to q leaves and k nodes other than u , denoted by $\{\mu_1, \dots, \mu_q\}$ and $\{v_1, \dots, v_k\}$, respectively. Then,

$$\gcd(\ell_{v,\mu_1}, \dots, \ell_{v,\mu_q}) = d_{v,u} \prod_{j=1}^k d_{v,v_j}. \quad (8.3)$$

For each $j \in \{1, \dots, k\}$ we let T_j be the branch of Γ' adjacent to v and containing v_j . Let $\Gamma'_j = [v, T_j]$ be the corresponding splice diagram with inherited weights. The inductive hypothesis on each Γ'_j yields $d_{v_j,v} = \gcd(\ell_{v_j,\mu} : \mu \in \partial \Gamma'_j \setminus \{v\})$. The identity $\ell_{v,\mu} = (\ell_{v_j,\mu}/d_{v_j,v}) (d_v/d_{v,v_j})$ where $\mu \in \partial \Gamma'_j \setminus \{v\}$ gives

$$\gcd(\ell_{v,\mu} : \mu \in \partial \Gamma'_j \setminus \{v\}) = \frac{d_v}{d_{v,v_j}}. \quad (8.4)$$

The result follows by combining (8.3) and (8.4) with the coprimality of the weights at v . $\square$

Proof of Theorem 8.2 We prove the statement by induction on the number of nodes of Γ , which we denote by p . If $p = 1$, we let u be the unique node of Γ . Then, the coprimality condition

$$\gcd(d_{u,\lambda_i}, d_{u,\lambda_j}) = 1 \quad \text{for } i \neq j$$

implies that w_u is a primitive vector. Furthermore, the formula in Remark 7.9 confirms that the tropical multiplicity associated to the edge $[u, \lambda]$ equals one since

$$\frac{1}{d_{u,\lambda}} \gcd \left(d_{u,\lambda} \prod_{\gamma \neq \lambda, \mu} d_{u,\gamma} : \mu \in \partial \Gamma \setminus \{\lambda\} \right) = 1.$$

Next, assume $p > 1$ and let u be an end-node of Γ . Let $\{\lambda_1, \dots, \lambda_s\}$ be the leaves adjacent to u and let v be the unique node of Γ adjacent to u . We let Γ' be the splice subdiagram of Γ obtained from u and v , as in Proposition 8.5. The coprimality condition for Γ' and our inductive hypothesis confirm that for each vertex v' of Γ' , $w_{v'}^{\Gamma'}$ is a primitive vector in $N(\partial \Gamma')$, and all tropical multiplicities of $\text{Trop}_{>0} \mathcal{S}(\Gamma')$ are one.

Since the gcd of all maximal minors of the matrix A equals 1 by the coprimality condition around u , it follows that A maps primitive vectors in $N(\partial \Gamma')$ to primitive vectors in $N(\partial \Gamma)$. This fact together with Lemma 8.6 ensures that the vector $w_{v'}$ is primitive whenever $v' \neq u$ is a node of Γ' . If $v' = u$ we have

$$w_u = \sum_{i=1}^s \frac{d_u}{d_{u,\lambda_i}} w_{\lambda_i} + \frac{d_u}{d_{u,v}} \sum_{\mu \in \partial_v \Gamma} \frac{\ell_{v,\mu}}{d_{v,u}} w_\mu,$$

where $\partial_v \Gamma$ is the set $\partial_{u,[u,v]} \Gamma$ from Definition 2.13. Since $\gcd(\ell_{v,\mu} : \mu \in \partial_v \Gamma) = d_{v,u}$ by Lemma 8.7, and $\gcd(d_u/d_{u,\lambda_i} : i = 1, \dots, s) = d_{u,v}$, the pairwise coprimality of weights around u ensures that w_u is a primitive vector in $N(\partial \Gamma)$.

To finish, we compute the tropical multiplicities. Remark 7.9, Lemma 8.7 and the coprimality of weights around u implies that the multiplicity corresponding to the edge $[u, \lambda_i]$ of Γ is one. Indeed, we have

$$\begin{aligned} & \gcd \left(\frac{\ell_{u,\mu}}{d_{u,\lambda_i}} : \mu \in \partial \Gamma \setminus \{\lambda_i\} \right) \\ &= \gcd \left(\gcd \left(\frac{d_u}{d_{u,\lambda_i} d_{u,\lambda_j}} : j = 1, \dots, s, j \neq i \right), \gcd \left(\frac{\ell_{v,\mu}}{d_{v,u}} \frac{d_u}{d_{u,\lambda_i} d_{u,v}} : \mu \in \partial_v \Gamma \right) \right) \\ &= \gcd \left(\gcd \left(\frac{d_u}{d_{u,\lambda_i} d_{u,\lambda_j}} : j = 1, \dots, s, j \neq i \right), \frac{d_u}{d_{u,\lambda_i} d_{u,v}} \right) = 1. \end{aligned}$$

If we pick an edge $[v', \lambda_j]$ with $j > s$ we get multiplicity one by the inductive hypothesis applied to Γ' combined with (8.1) and the coprimality of the weights around u . More precisely,

$$\begin{aligned} & \gcd \left(\gcd \left(\frac{\ell_{v', \lambda_k}}{d_{v', \lambda_j}} : k = s+1, \dots, n, k \neq j \right), \gcd \left(\frac{\ell_{v', \lambda_i}}{d_{v', \lambda_j}} : i = 1, \dots, s \right) \right) \\ &= \gcd \left(\gcd \left(\frac{\ell_{v', \lambda_k}^{\Gamma'}}{d_{v', \lambda_j}} : k = s+1, \dots, n, k \neq j \right), \frac{\ell_{v', u}^{\Gamma'}}{d_{v', \lambda_j}} \underbrace{\gcd \left(\frac{\ell_{u, \lambda_i}}{d_{u, v}} : i = 1, \dots, s \right)}_{=1} \right) = 1. \end{aligned}$$

Finally, the cone associated to an edge between two adjacent nodes u', v' of Γ will have tropical multiplicity one by Lemma 8.7 since $\gcd(\ell_{u', \lambda}/d_{u', v'} : \lambda \in \partial_{v', [u', v']}\Gamma) = \gcd(\ell_{v', \mu}/d_{v', u'} : \mu \in \partial_{u', [u', v']}\Gamma) = 1$. $\square$

Proof of Theorem 8.3 The combinatorial type of Γ is completely determined by intersecting Δ_{n-1} and the splice fan. In turn, Theorem 8.2 (1) allows us to characterize the vector w_u as the primitive vector associated to the corresponding ray of the fan $\mathbb{R}_{\geq 0}^t(\Gamma)$. All that remains is to determine the weights around each node of Γ from this data. We do so by induction on the number of nodes of Γ , which we denote by p .

If $p = 1$, then the coprimality of the weights around the single node u of Γ determines each d_{u, λ_i} uniquely as follows. By construction, the entries of w_u are coprime and we have

$$d_{u, \lambda} = \gcd((w_u)_\mu : \mu \in \partial \Gamma \setminus \{\lambda\}).$$

Next, assume $p > 1$ and fix an end-node u of Γ . Let $\lambda_1, \dots, \lambda_s$ be the leaves of Γ adjacent to u , and v be the unique node of Γ adjacent to u . Let Γ' be the tree obtained by pruning Γ from u , as in Proposition 8.5. The weights around u can be recovered uniquely from the splice fan of Γ . Indeed, write

$$w_u = \sum_{i=1}^s \frac{d_u}{d_{u, \lambda_i}} w_{\lambda_i} + \frac{d_u}{d_{u, v}} \sum_{j=s+1}^n \frac{\ell_{v, \lambda_j}^{\Gamma'}}{d_{v, u}} w_{\lambda_j}. \quad (8.5)$$

Notice that the coprimality condition gives $d_{u, v} = \gcd(d_u/d_{u, \lambda_i} : i = 1, \dots, s)$ and $d_u = \text{lcm}(d_u/d_{u, \lambda_i} : i = 1, \dots, s)$. From this we recover all remaining s weights at u since $d_{u, \lambda_i} = d_u/(d_u/d_{u, \lambda_i})$ for every $i \in \{1, \dots, s\}$.

Next, for each node v' of Γ with $v' \neq u$, we use the full-rank matrix A from (8.2) to recover $w_{v'}^{\Gamma'} \in \mathbb{Z}^{n-s+1}$ uniquely from $w_{v'}$. Since $w_u^{\Gamma'}$ is a prescribed canonical basis element of $N(\partial \Gamma') \simeq \mathbb{Z}^{n-s+1}$, the set of vectors $\{w_{v'}^{\Gamma'} : v' \text{ is a node of } \Gamma'\}$ allows us to determine the splice fan of Γ' . The inductive hypothesis then uniquely recovers the splice diagram Γ' , and hence Γ has been fully determined. $\square$

9 Intrinsic nature of splice type singularities

Theorem 6.2 shows that the local tropicalization of the germ defined by a given splice type system $\mathcal{S}(\Gamma)$ is independent of the choice of admissible monomials and higher order terms used to define it. In fact, Neumann and Wahl proved that the set of splice type singularities defined by splice type systems with fixed admissible monomials associated to a given splice diagram is independent of these choices, both in the coprime setting and in the general case under a suitable equivariant hypothesis on the series $g_{v,i}$ collecting the higher order terms of each series $F_{v,i}$. In this section we give a variant of their proof in the coprime case and we show by an example that without the equivariance hypothesis, the result no longer holds.

Here is the precise statement for coprime diagrams, which can be deduced from the equivariant case [38, Theorem 10.1]. For completeness, we include a direct proof:

Theorem 9.1 *Let Γ be a coprime splice diagram with n leaves, and let*

$$\mathcal{M} := \{\underline{z}^{m_{v,e}} : v \text{ is a node in } \Gamma, e \in \text{Star}_\Gamma(v)\}$$

be a complete set of admissible monomials for Γ . Let $\mathcal{S}(\Gamma)_\mathcal{M}$ be the set of all splice type systems that can be constructed using the set $\mathcal{M}$. Then, the set $X_\mathcal{M}$ of all germs $(X, 0) \hookrightarrow \mathbb{C}^n$ defined by the vanishing of a system in $\mathcal{S}(\Gamma)_\mathcal{M}$ is independent of $\mathcal{M}$.

Proof Let $\mathcal{M}'$ be a second complete set of admissible monomials. We proceed by induction on the size p of the set $\mathcal{M} \setminus \mathcal{M}'$. If $p = 0$, there is nothing to show. For the inductive step, it suffices to analyze the case when $p = 1$, i.e. when $\mathcal{M}$ and $\mathcal{M}'$ differ by exactly one admissible monomial. Fix a pair (v_0, e_0) for which the corresponding monomials in $\mathcal{M}$ and $\mathcal{M}'$ differ, and let m_{v_0, e_0} and m'_{v_0, e_0} be the associated admissible exponents. To prove the statement, we use the w_{v_0} -filtration $\mathcal{I}_\bullet$: $I_0 \supseteq I_1 \supseteq I_2 \supseteq \dots$ of ideals of the local ring $\mathcal{O} := \mathbb{C}[z_1, \dots, z_n]$, where

$$I_d := \{f \in \mathcal{O} : w_{v_0}(f) \geq d\}. \quad (9.1)$$

Let $\{F_{v,i} : v, i\}$ be a splice type system in $\mathcal{S}(\Gamma)_\mathcal{M}$. It determines a germ $(X, 0) \in X_\mathcal{M}$ together with an embedding $\varphi : X \hookrightarrow \mathbb{C}^n$. We write $F_{v,i} = f_{v,i} + g_{v,i}$ for each node v of Γ and $i \in \{0, \dots, \delta_v - 2\}$ as in (2.8) and (2.9). The w_{v_0} -filtration $\mathcal{I}_\bullet$ restricted to X yields a filtration $\mathcal{J}_\bullet$ under the corresponding surjective map of local rings $\varphi^\sharp : \mathcal{O} \twoheadrightarrow \mathcal{O}_{(X,0)}$, i.e.,

$$J_k := \varphi^\sharp(I_k) \quad \text{for all } k \geq 0. \quad (9.2)$$

By Lemma 9.2 below, there exist $a \in \mathbb{C}^*$, $g \in I_{d_{v_0}+1}$ and h in the ideal of $\mathcal{O}$ spanned by $\{F_{v,i} : v, i\}$, satisfying the equality

$$\underline{z}^{m_{v_0, e_0}} - a \underline{z}^{m'_{v_0, e_0}} = h + g. \quad (9.3)$$

Furthermore, up to moving to g all higher-order contributions of terms in h coming from each $F_{v_0,i}$, we may assume that $h = \sum_{v,i} a_{v,i} F_{v,i}$ with $a_{v_0,i} \in \mathbb{C}$ for each $i \in \{1, \dots, \delta_{v_0}\}$. This can be achieved thanks to (2.6) and (2.10).

We prove the inclusion $X_{\mathcal{M}} \subseteq X_{\mathcal{M}'}$ by constructing an explicit system $\{F'_{v,i} : v, i\}$ in $\mathcal{S}(\Gamma)_{\mathcal{M}'}$ whose vanishing set equals X . The reverse inclusion then follows by exploiting the symmetry between $\mathcal{M}$ and $\mathcal{M}'$. We consider the set $\{F'_{v,i} : v, i\}$ with $F'_{v,i} := F_{v,i}$ for each $v \neq v_0$ and $i \in \{1, \dots, \delta_v - 2\}$, whereas for $v = v_0$ and $i \in \{1, \dots, \delta_{v_0} - 2\}$ we pick $F'_{v_0,i} := f'_{v_0,i} + g'_{v_0,i}$ with

$$f'_{v_0,i} := (f_{v_0,i} - c_{v_0,e_0,i} \underline{z}^{m_{v_0,e_0}}) + (c_{v_0,e_0,i} a) \underline{z}^{m'_{v_0,e_0}} \quad \text{and} \quad g'_{v_0,i} := g_{v_0,i} + c_{v_0,e_0,i} g. \quad (9.4)$$

The scalar a and the series $g \in I_{d_{v_0}+1}$ are those from (9.3).

We claim that $\{F'_{v,i}\} \in \mathcal{S}(\Gamma)_{\mathcal{M}'}$. Indeed, by construction, each series $g'_{v_0,i}$ lies in $I_{d_{v_0}+1}$, as required by (2.10) (see Remark 2.15). In addition, the matrix of coefficients for the polynomials $\{F'_{v_0,i}\}_i$ is obtained from the matrix $(c_{v_0,e,i})_{i,e}$ after rescaling by a the column labeled with e_0 . Thus, the Hamm determinant conditions of Definition 2.14 are satisfied.

Combining (9.3) and (9.4) yields

$$F'_{v_0,i} = F_{v_0,i} - c_{v_0,e_0,i} h \in \langle F_{v,j} : v \text{ node of } \Gamma, j = 1, \dots, \delta_v - 2 \rangle.$$

We use the expression of h given above to replace $\{F'_{v,i} : v, i\}$ by a set generating the same ideal, i.e.,

$$\{F_{v,i} : v \neq v_0, i = 1, \dots, \delta_v - 2\} \cup \left\{ F_{v_0,i} - c_{v_0,e_0,i} \sum_{j=1}^{\delta_{v_0}-2} a_{v_0,j} F_{v_0,j} : i = 1, \dots, \delta_{v_0} - 2 \right\}. \quad (9.5)$$

Since both $\{F_{v,i} : v, i\} \in \mathcal{S}(\Gamma)_{\mathcal{M}}$ and $\{F'_{v,i} : v, i\} \in \mathcal{S}(\Gamma)_{\mathcal{M}'}$ determine complete intersection systems of equations by Theorem 2.16, the $(\delta_{v_0} - 2) \times (\delta_{v_0} - 2)$ -matrix of scalars $\text{Id} - (c_{v_0,e_0,i} a_{v_0,j})_{i,j}$ associated to the second set in (9.5) must be invertible. From here it follows that the vanishing sets of both collections $\{F'_{v,i} : v, i\}$ and $\{F_{v,i} : v, i\}$ agree. Thus, the germ X lies in $X_{\mathcal{M}'}$, as we wanted to show. $\square$

The following technical lemma gives a more precise version of the first half of the statement of [38, Theorem 10.1]. Its proof follows the same reasoning, so we omit it here:

Lemma 9.2 *Fix two collections of admissible monomials $\mathcal{M}, \mathcal{M}'$ with $|\mathcal{M} \setminus \mathcal{M}'| = 1$. Assume $X \in X_{\mathcal{M}}$, and let $\underline{z}^{m_{v_0,e_0}} \in \mathcal{M} \setminus \mathcal{M}'$ and $\underline{z}^{m'_{v_0,e_0}} \in \mathcal{M}' \setminus \mathcal{M}$. Then, there exists $a \in \mathbb{C}^*$ such that the restriction $(\underline{z}^{m_{v_0,e_0}} - a \underline{z}^{m'_{v_0,e_0}})|_X$ belongs to $J_{d_{v_0}+1}$, where $J_{d_{v_0}+1}$ is the ideal from (9.2).*

As we mentioned earlier, if Γ is not coprime, analogous results to Theorem 9.1 and Lemma 9.2 can be proved under the condition that the higher order terms of




Fig. 10 An example confirming the dependency of splice-type systems on the choice of admissible monomials

each system (i.e., the terms in each $g_{v,i}$) satisfy an equivariant condition under the action of a suitable finite abelian group, namely, the discriminant group of a given plumbing graph with associated splice diagram Γ . For a precise statement, we refer to [39, Theorem 10.1].

It is natural to ask whether this equivariant condition can be weakened. The next example shows that *this is not the case*.

Example 9.3 Consider the (non-coprime) splice diagram Γ from Fig. 10 and pick two sets of admissible monomials for Γ that differ only in the choice of exponent vectors for the pair $(v, [v, u])$:

$$\mathcal{M} := \{z_1^2, z_2^2, z_3 z_4, z_3^2, z_4^2\} \cup \{z_1^3\}, \quad \mathcal{M}' := \{z_1^2, z_2^2, z_3 z_4, z_3^2, z_4^2\} \cup \{z_1^2 z_2\}.$$

We claim that $X_{\mathcal{M}} \neq X_{\mathcal{M}'}$, i.e., the elements of $\mathcal{S}(\Gamma)_{\mathcal{M}}$ and $\mathcal{S}(\Gamma)_{\mathcal{M}'}$ determine different sets of subgerms of $(\mathbb{C}^n, 0)$. More precisely, we show that the germ in $X_{\mathcal{M}}$ defined by the system

$$\begin{cases} f_{u,1} := z_1^2 + z_2^2 + z_3 z_4, \\ f_{v,1} := z_1^3 + z_3^2 + z_4^2, \end{cases} \quad (9.6)$$

in $\mathcal{S}(\Gamma)_{\mathcal{M}}$ cannot be a member of $X_{\mathcal{M}'}$. To do so, it suffices to show that no power series associated to the node v of Γ , that is, no power series of the form

$$F'_{v,1} := b_1 z_1^2 z_2 + b_2 z_3^2 + b_3 z_4^2 + g_{v,1},$$

with $b_1, b_2, b_3 \in \mathbb{C}^*$ and $g_{v,1}$ satisfying (2.10) can be an element of the ideal $(f_{u,1}, f_{v,1})$ of the power series ring $\mathbb{C}\{z_1, \dots, z_4\}$.

We argue by contradiction and pick elements $A_1, A_2 \in \mathbb{C}\{z_1, \dots, z_4\}$ with

$$b_1 z_1^2 z_2 + b_2 z_3^2 + b_3 z_4^2 + g_{v,1} = A_1 f_{u,1} + A_2 f_{v,1}. \quad (9.7)$$

By construction, the w_v -initial form on the left-hand side is $b_1 z_1^2 z_2 + b_2 z_3^2 + b_3 z_4^2$ and its w_v -weight is 24. We claim that the w_v -initial form on the right-hand side of (9.7) may be written as

$$\alpha_1(\underline{z})(z_1^2 + z_2^2) + \alpha_2(\underline{z})(z_1^3 + z_3^2 + z_4^2),$$

for two w_v -homogeneous polynomials $\alpha_1(\underline{z}), \alpha_2(\underline{z})$.

We prove this claim by explicit computation, comparing the w_v -weights of both summands and noticing that $\text{in}_{w_v}(f_{u,1}) = z_1^2 + z_2^2$, $\text{in}_{w_v}(f_{v,1}) = f_{v,1}$. Three situations can occur. First, if $w_v(A_1 f_{u,1}) < w_v(A_2 f_{v,1})$, then the w_v -initial form on

the right-hand side of (9.7) comes from the first summand, i.e., $\alpha_1(\underline{z}) = \text{in}_{w_v}(A_1)$ and $\alpha_2(\underline{z}) = 0$. Similarly, if $w_v(A_1 f_{u,1}) > w_v(A_2 f_{v,1})$, then the second summand determines the w_v -initial form on the right-hand side of (9.7), so $\alpha_1(\underline{z}) = 0$ and $\alpha_2(\underline{z}) = \text{in}_{w_v}(A_2)$. Finally, if $w_v(A_1 f_{u,1}) = w_v(A_2 f_{v,1})$, the condition that the total w_v -weight of $A_1 f_{u,1} + A_2 f_{v,1}$ and $f_{v,1}$ agrees with the w_v -weight of $F'_{v,1}$ confirms that both terms contribute to the w_v -initial form, with $\alpha_1(\underline{z}) = \text{in}_{w_v}(A_1)$ and $\alpha_2(\underline{z}) = \text{in}_{w_v}(A_2)$.

Comparing the w_v -initial forms on both sides of (9.7) yields an identity of w_v -homogeneous polynomials:

$$b_1 z_1^2 z_2 + b_2 z_3^2 + b_3 z_4^2 = \alpha_1(\underline{z}) (z_1^2 + z_2^2) + \alpha_2(\underline{z}) (z_1^3 + z_3^2 + z_4^2).$$

Since the w_v -weight on both sides equals 24, we conclude that $\alpha_2(\underline{z})$ must be a constant. Evaluating both sides at $z_1 = z_2 = 0$ forces $b_2 = b_3 = \alpha_2(\underline{z})$, so in particular $\alpha_2 := \alpha_2(\underline{z}) \in \mathbb{C}^*$. We conclude from this that

$$b_1 z_1^2 z_2 = \alpha_1(\underline{z}) (z_1^2 + z_2^2) + \alpha_2 z_1^3.$$

This identity implies that the function $z_1^3/(z_1^2 z_2) = b_1 \alpha_2^{-1}$ is constant on $Z(z_1^2 + z_2^2) \cap (\mathbb{C}^*)^2$, which is false. This contradiction confirms that $X_{\mathcal{M}} \neq X_{\mathcal{M}'}$, as we wanted to show.

Acknowledgements Maria Angelica Cueto was supported by an NSF postdoctoral fellowship DMS-1103857 and NSF Standard Grants DMS-1700194 and DMS-1954163 (USA). Patrick Popescu-Pampu was supported by French grants ANR-12-JS01-0002-01 SUSI, ANR-17-CE40-0023-02 LISA and Labex CEMPI (ANR-11-LABX-0007-01). Labex CEMPI also financed a one month research stay of the first author in Lille during Summer 2018. Part of this project was carried out during two Research in triples programs, one at the Centre International de Rencontres Mathématiques (Marseille, France, Award No. 1173/2014) and one at the Center International Bernoulli (Lausanne, Switzerland). The authors would like to thank both institutes for their hospitality, and for providing excellent working conditions. The authors are very grateful to Jonathan Wahl for contributing the proof presented in the appendix of this paper. Finally, we express our warm thanks to the referee for comments and suggestions that helped us improve the exposition.

Appendix A. Initial ideals and local regular sequences

In [38], the authors invoke a folklore lemma in commutative algebra in order to prove several of their main theorems. This result involves regular sequences in a polynomial ring and their initial forms with respect to integer weight vectors. As originally stated, [38, Lemma 3.3] is not quite-correct: the global setting must be replaced by a local one. This appendix provides a complete proof of this result in the local setting of convergent power series near the origin, a result we could not locate in the literature. This local version agrees with the general framework of [38]. Throughout, we let n be a positive integer and let $(\mathcal{O}, \mathfrak{m})$ denote the local ring of convergent power series $\mathbb{C}\{z_1, \dots, z_n\}$ near the origin.

We start by stating our main result, namely, a reformulation of [38, Lemma 3.3] in the local setting. Its proof will be given at the end of this appendix, after discussing

a series of preliminary technical results. Note that the same statement and proof will hold if $\mathcal{O}$ denotes the localization of the polynomial ring $\mathbb{C}[z_1, \dots, z_n]$ at the maximal ideal of the origin of $\mathbb{C}^n$.

Theorem A.1 *Let $(f_1, \dots, f_s)$ be a finite sequence of elements in the maximal ideal $\mathfrak{m}$ of $\mathcal{O}$, and let J be the ideal generated by them. Fix a positive weight vector $w \in (\mathbb{Z}_{>0})^n$. Assume that $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is a regular sequence in $\mathcal{O}$. Then:*

1. *the sequence $(f_1, \dots, f_s)$ is also regular; and*
2. *the w -initial ideal $\text{in}_w(J)\mathcal{O}$ is generated by $\{\text{in}_w(f_1), \dots, \text{in}_w(f_s)\}$.*

Remark A.2 As mentioned earlier, Theorem A.1 does not hold in the polynomial setting. For instance, $(z_1(1 - z_1), z_2(1 - z_1))$ is a regular sequence in the local ring $\mathbb{C}\{z_1, z_2\}$ but not in the polynomial ring $\mathbb{C}[z_1, z_2]$. However, the sequence (z_1, z_2) of initial forms with respect to any weight vector $w \in (\mathbb{Z}_{>0})^2$ is regular in both rings.

Remark A.3 The regularity of the sequence of w -initial forms is needed in Theorem A.1. As an example, fix $n = 4$, $w = (1, 1, 1, 1)$, and consider the sequence (f_1, f_2) with

$$f_1 := z_1^2 + z_2^4 - z_3^3 \quad \text{and} \quad f_2 := z_1 z_2 - z_4^3.$$

By construction, (f_1, f_2) is a regular sequence in $\mathcal{O}$ defining an isolated complete intersection surface singularity. The sequence of initial forms $(\text{in}_w(f_1), \text{in}_w(f_2)) = (z_1^2, z_1 z_2)$ is not regular, and the w -initial ideal of $(f_1, f_2)\mathcal{O}$ is generated by $\text{in}_w(f_1)$, $\text{in}_w(f_2)$ and $\text{in}_w(z_2 f_1 - z_1 f_2) = -z_2 z_3^3 + z_1 z_4^3$.

Throughout, we fix $w \in (\mathbb{Z}_{>0})^n$ and an arbitrary sequence $(f_1, \dots, f_s)$ of elements of the maximal ideal $\mathfrak{m}$. We let J be the ideal generated by the f_i 's. Consider the first few steps in the Koszul complex of $\mathcal{O}$ -modules determined by it (see, e.g., [51, Sect. IV.A]):

$$\boxed{F} := \bigoplus_{1 \leq i < j \leq s} \mathcal{O} \cdot e_{ij} \xrightarrow{d_2} \boxed{E} := \bigoplus_{1 \leq i \leq s} \mathcal{O} \cdot e_i \xrightarrow{d_1} \mathcal{O} \longrightarrow \mathcal{O}/J. \quad (\text{A.1})$$

The map $d_1: E \rightarrow \mathcal{O}$ sends e_i to f_i for each $i = 1, \dots, s$ and the kernel $\boxed{R}$ of d_1 is the *module of relations* between the given generators of J . The morphism $d_2: F \rightarrow E$ sends e_{ij} to $f_j e_i - f_i e_j$, and its image is the submodule of “trivial relations” between $(f_1, \dots, f_s)$. By definition, the image of d_2 lies in R , so we view d_2 also as a map $d_2: F \rightarrow R$.

By a standard result in commutative algebra (see, e.g., [51, Proposition 3, Chapter IV.A.2]) we have:

Proposition A.4 *The sequence $(f_1, \dots, f_s)$ of elements in $\mathfrak{m}$ is regular in $\mathcal{O}$ if and only if the Koszul complex (A.1) is exact at E .*

Since the definition of E does not depend on the order of the sequence $(f_1, \dots, f_s)$, the following consequence arises naturally:

Corollary A.5 *If $(f_1, \dots, f_s)$ is a regular sequence in $\mathcal{O}$, any reordering of it is also a regular sequence.*

The weight vector w inducing the w -weight valuation (3.1) on $\mathcal{O}$ endows this ring with a weight filtration by ideals $(I_p)_{p \geq 0}$, where $\boxed{I_p} := \{g \in \mathcal{O} : w(g) \geq p\}$. Similarly, we can filter $\mathcal{O}$ via the ideals $(\mathfrak{m}^p)_{p \geq 0}$. Both filtrations are cofinal since

$$I_{dp} \subseteq \mathfrak{m}^p \subseteq I_p \quad \text{for all } p \geq 0, \quad (\text{A.2})$$

where $\boxed{d}$ is the maximum among all coordinates of w . It follows from this that the completions of $\mathcal{O}$ with respect to both filtrations are canonically isomorphic. The completion induced by the $\mathfrak{m}$ -adic filtration $(\mathfrak{m}^p)_{p \geq 0}$ is the ring of formal power series in n variables.

In a similar fashion, we can filter the modules E and F appearing in (A.1) via $(E_p)_{p \geq 0}$ and $(F_p)_{p \geq 0}$, respectively, by assigning the weights $w(f_i)$ and $w(f_i) + w(f_j)$ to e_i and e_{ij} , respectively. More precisely,

$$\begin{aligned} \boxed{E_p} &:= \left\{ \sum_{i=1}^s a_i e_i : w(a_i) \geq p - w(f_i) \ \forall i \right\} \text{ and} \\ \boxed{F_p} &:= \left\{ \sum_{i < j} b_{ij} e_{ij} : w(b_{ij}) \geq p - w(f_i) - w(f_j) \ \forall i, j \right\}. \end{aligned} \quad (\text{A.3})$$

These choices ensure that the maps d_1 and d_2 from the Koszul complex (A.1) preserve the filtration. In addition, the module R of relations is filtered as well, via

$$\boxed{R_p} := E_p \cap R. \quad (\text{A.4})$$

We use these filtrations to define the w -initial forms on E and F . We state the definition for E , since the one for F is analogous. The definition for R is given by restriction.

Definition A.6 Given any $g \in E$ with $g \neq 0$, we let p be the unique integer such that $g \in E_p \setminus E_{p+1}$. An element $g := \sum_{i=1}^s r_i e_i \in E_p \setminus E_{p+1}$ satisfies $w(r_i) + w(f_i) \geq p$ for all $i \in \{1, \dots, s\}$ and equality must hold for some index i . Let I be the set of indices where equality is achieved. The w -initial form of g is $\boxed{\text{in}_w(g)} := \sum_{i \in I} \text{in}_w(r_i) e_i$. We set $\text{in}_w(0) = 0$.

By Proposition A.4, the regularity of the sequence $(f_1, \dots, f_s)$ is equivalent to the surjectivity of the map $d_2: F \rightarrow R$ induced by (A.1). We prove the latter in Lemma A.11, assuming the regularity of the sequence of w -initial forms of all f_i 's.

Our first two lemmas use the regularity assumptions for the sequence of w -initial forms to prove the surjectivity of $d_2: F \rightarrow R$ by working with the filtrations of F and R described above.

Lemma A.7 *Assume that the sequence $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is regular in $\mathcal{O}$. Then, the morphism of $\mathbb{C}$ -vector spaces $\varphi_p: F_p/F_{p+1} \rightarrow R_p/R_{p+1}$ induced by the morphism of $\mathcal{O}$ -modules $d_2: F \rightarrow R$ is surjective for all integers $p \geq 0$.*

Proof We must show that modulo R_{p+1} , every element g of R_p is the image of an element of F_p/F_{p+1} under the map φ_p . If $g = 0$, there is nothing to show, so we assume $g \neq 0$. In particular, g lifts to an element in $R_p \setminus R_{p+1}$, which we denote by g as well. We write $g = \sum_{j=1}^s r_j e_j$.

Assume that $\text{in}_w(g)$ has k many terms, with $k \in \{1, \dots, s\}$ (see Definition A.6). By Corollary A.5, we can reorder the original sequence while preserving its regularity, and write $\text{in}_w(g)$ as

$$\text{in}_w(g) = \sum_{j=1}^k \text{in}_w(r_j) e_j \quad \text{with} \quad w(r_j) + w(f_j) = p \quad \text{for all } j \in \{1, \dots, k\}.$$

We claim that g is congruent, modulo the image of φ_p , to an element of R_p whose w -initial form lies in the ideal generated by $\{\text{in}_w(f_1), \dots, \text{in}_w(f_{k-1})\}$. The original statement will follow by induction on $k \leq s$.

Since $\sum_{j=1}^s r_j f_j = 0$ by definition of R and $w(r_j) + w(f_j) > p$ for $j \in \{k+1, \dots, s\}$, we conclude that the expected w -initial form of $\sum_{j=1}^s r_j f_j$ must vanish, i.e.,

$$\sum_{j=1}^k \text{in}_w(r_j) \text{in}_w(f_j) = 0.$$

Therefore, $\text{in}_w(r_k) \text{in}_w(f_k)$ is zero modulo the ideal $I := \langle \text{in}_w(f_1), \dots, \text{in}_w(f_{k-1}) \rangle \mathcal{O}$. Since the sequence $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is regular, we conclude that $\text{in}_w(r_k)$ must lie in I .

Taking the w -weight value of r_k and each f_j into account we write $\text{in}_w(r_k)$ as

$$\text{in}_w(r_k) = \sum_{j=1}^{k-1} a_j \text{in}_w(f_j),$$

where a_j is either 0 or a non-zero w -weighted homogeneous polynomial with $w(a_j) = p - w(f_k) - w(f_j) \geq 0$ for all $j \in \{1, \dots, k-1\}$. It follows from this that the element

$$r'_k := r_k - \sum_{j=1}^{k-1} a_j f_j$$

satisfies $w(r'_k) > p - w(f_k)$, so $r'_k e_k \in E_{p+1}$. Simple arithmetic manipulations give a new formula for g , i.e.,

$$g = \sum_{j=1}^s r_j e_j = \sum_{j=1}^{k-1} (r_j + a_j f_k) e_j + \underbrace{\sum_{j=1}^{k-1} a_j (f_j e_k - f_k e_j)}_{=: h} + r'_k e_k + \sum_{j=k+1}^s r_j e_j. \quad (\text{A.5})$$

By construction, it follows that $h = \varphi_p(\sum_{j=1}^{k-1} a_j e_{jk}) \in \varphi_p(F_p/F_{p+1})$. Furthermore, $\text{in}_w(g - h)$ only involves terms in the first of the four summands on the right-hand side of (A.5) since the last two summands lie in E_{p+1} . This establishes the claim. $\square$

Lemma A.8 *Let J be the ideal of $\mathcal{O}$ generated by $\{f_1, \dots, f_s\}$ and assume that $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is a regular sequence in $\mathcal{O}$. If $g \in J$ has w -weight equal to $p \in \mathbb{N}$, then g admits an expression of the form $g = \sum_{i=1}^s a_i f_i$, where $w(a_i f_i) \geq p$ for all i . In particular, $\text{in}_w(g)$ belongs to the ideal of $\mathcal{O}$ generated by $\{\text{in}_w(f_1), \dots, \text{in}_w(f_s)\}$.*

Proof Since $g \in J$, we may write g as $g = \sum_{i=1}^s b_i f_i$ with $b_i \in \mathcal{O}$ for each i . Consider

$$p' = \min\{w(b_i f_i) : i = 1, \dots, s\}.$$

Assume that this weight is achieved at k many terms, which we can fix to be $\{b_1 f_1, \dots, b_k f_k\}$ upon reordering. If $p' \geq p$ we have $w(a_i f_i) \geq p$ for all i and equality must hold for some i by definition of p . From here it follows that $p' = p$, so $\text{in}_w(g) = \sum_{j=1}^k \text{in}_w(a_j) \text{in}_w(f_j)$, as we wanted to show.

On the contrary assume that $p' < p$. We claim that we can find an alternative expression $g = \sum_{j=1}^s b'_j f_j$ where the corresponding minimum weight $p'' := \min\{w(b'_j f_j)\}$ satisfies $p'' \geq p'$ and the number of summands realizing p'' is strictly smaller than k . An easy induction combined with the fact that $p', p \in \mathbb{Z}_{\geq 0}$ will then yield a new expression for g with $p' \geq p$, as in our previous case.

It remains to prove the claim. Since $p' < p$, the terms in g with w -weight p' must cancel out, i.e., $\sum_{j=1}^k \text{in}_w(b_j) \text{in}_w(f_j) = 0$. As in the proof of Lemma A.7, the fact that $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is a regular sequence in $\mathcal{O}$ ensures that

$$\text{in}_w(b_k) = \sum_{j=1}^{k-1} c_j \text{in}_w(f_j),$$

where c_j is either zero or a w -homogeneous polynomial with $w(c_j) = p' - w(f_j) - w(f_k) \geq 0$. It follows from here that the element $b'_k := b_k - \sum_{j=1}^{k-1} c_j f_j$ has weight $w(b'_k) > w(b_k)$, so $w(b'_k f_k) > p'$.

An arithmetic manipulation allows us to rewrite g as follows:

$$g = \sum_{j=1}^{k-1} (b_j + c_j f_k) f_j + b'_k f_k + \sum_{j=k+1}^s b_j f_j \quad (\text{A.6})$$

By construction, the terms with minimum w -weight only appear in the first of the three summands on the right-hand side of (A.6). Furthermore, the corresponding minimum weight p'' satisfies $p'' \geq p'$ since $w(b_j f_j) \geq p'$ for all j and $w(c_j f_k f_j) \geq p'$ for $j < k$. This confirms the validity of our claim. $\square$

A standard commutative algebra result (see, e.g., [4, Lemma 10.23]) combined with Lemma A.8 yields:

Lemma A.9 *Assume that the sequence $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is regular in $\mathcal{O}$. Then, the map $d_2 : F \rightarrow R$ of filtered modules induces a surjection between their completions relative to the filtrations $(F_p)_{p \geq 0}$ and $(R_p)_{p \geq 0}$ respectively. More precisely, $\varprojlim F/F_p \twoheadrightarrow \varprojlim R/R_p$.*

We let $\hat{F}$ and $\hat{R}$ be the $\mathfrak{m}$ -adic completions of F and R respectively, which can be computed with standard methods. Indeed, by [4, Theorem 10.13], we have

$$\hat{F} \simeq F \otimes_{\mathcal{O}} \hat{\mathcal{O}} \quad \text{and} \quad \hat{R} \simeq R \otimes_{\mathcal{O}} \hat{\mathcal{O}}. \quad (\text{A.7})$$

The double inclusions in (A.2) allow us to compare the completions in Lemma A.9 induced by $(F_p)_{p \geq 0}$ and $(R_p)_{p \geq 0}$, with $\hat{F}$ and $\hat{R}$, respectively. More precisely,

Lemma A.10 *Assume that the sequence $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is regular in $\mathcal{O}$. Then, the completions appearing in Lemma A.9 agree with the $\mathfrak{m}$ -adic ones, i.e.*

$$\varprojlim F/F_p \simeq \varprojlim F/\mathfrak{m}^p F \simeq F \otimes_{\mathcal{O}} \hat{\mathcal{O}} \quad \text{and} \quad \varprojlim R/R_p \simeq \varprojlim R/\mathfrak{m}^p R \simeq R \otimes_{\mathcal{O}} \hat{\mathcal{O}}. \quad (\text{A.8})$$

Proof We let $\ell := \max\{w(f_j) : j = 1, \dots, s\}$. It suffices to prove the first isomorphism on each side of (A.8), since the remaining ones appear in (A.7). By (A.3), we have

$$E_p := \bigoplus_i I_{p-w(f_i)} e_i \quad \text{and} \quad F_p := \bigoplus_{i < j} I_{p-w(f_i)-w(f_j)} e_{ij} \quad \text{for each } p \geq 0.$$

It follows from here that $I_p E \subseteq E_p \subseteq I_{p-\ell} E$ and $I_p F \subseteq F_p \subseteq I_{p-2\ell} F$ for each $p \geq 0$. Combining these inclusions with (A.2) yields:

$$E_{dp+\ell} \subseteq I_{dp} E \subseteq \mathfrak{m}^p E \subseteq E_p \quad \text{and} \quad F_{dp+2\ell} \subseteq I_{dp} F \subseteq \mathfrak{m}^p F \subseteq F_p \quad \text{for each } p \geq 0. \quad (\text{A.9})$$

The inclusions appearing on the right of (A.9) ensure that the filtrations $(\mathfrak{m}^p F)_{p \geq 0}$ and $(F_p)_{p \geq 0}$ are cofinal in F . Thus, they yield isomorphic completions. This proves the first isomorphism in (A.8).

Next, consider the filtration R_p from (A.4). First, notice that $\mathfrak{m}^p R \subseteq R_p$ by (A.9). To finish, we claim the existence of some $k \geq 0$ for which $R_{dp+(dk+\ell)} \subseteq \mathfrak{m}^p R$ for all $p \gg 0$. Indeed, by the Artin–Rees Lemma (see, e.g., [4, Theorem 10.10]), there exists an integer $k \geq 0$ satisfying

$$R \cap \mathfrak{m}^p E = \mathfrak{m}^{p-k}(R \cap \mathfrak{m}^k E) \quad \text{for all } p \geq k.$$

Therefore, combining this fact with property (A.9) we obtained the desired inclusion:

$$R_{dp+(dk+\ell)} = R \cap E_{d(p+k)+\ell} \subset R \cap \mathfrak{m}^{p+k} E = \mathfrak{m}^p(R \cap \mathfrak{m}^k E) \subset \mathfrak{m}^p R.$$

We conclude that $(\mathfrak{m}^p R)_{p \geq 0}$ and $(R_p)_{p \geq 0}$ are cofinal filtrations in R , so they yield isomorphic completions. $\square$

We let M be the cokernel of the map $d_2: F \rightarrow R$ given by (A.1), and we let $\hat{M}$ be its $\mathfrak{m}$ -adic completion. Lemma A.10 yields the following result:

Lemma A.11 *Assume that the sequence $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is regular in $\mathcal{O}$. Then, $\hat{M} = 0$ and $M = 0$. In particular, the Koszul complex (A.1) is exact at E .*

Proof By standard commutative algebra (see, e.g., [51, Corollaire 2, Chap. II.A.5]) we know that $\hat{\mathcal{O}}$ is a flat $\mathcal{O}$ -module. Therefore, taking $\mathfrak{m}$ -adic completion is an exact functor. Since $\hat{F} \rightarrow \hat{R}$ is surjective (by combining Lemmas A.9 and A.10) it follows that $\hat{M} = 0$.

By [4, Theorem 10.17], the kernel of the canonical morphism $M \rightarrow \hat{M}$ is annihilated by an element of the form $(1+z)$ where $z \in \mathfrak{m}$. As $\mathcal{O}$ is a local ring, the element $(1+z)$ must be a unit of $\mathcal{O}$, thus $M = 0$ as claim. The exactness of the Koszul complex at E follows immediately, as it is equivalent to the surjectivity of the morphism $d_2: F \rightarrow R$. $\square$

We end this appendix by proving its main result:

Proof of Theorem A.1 Since $(\text{in}_w(f_1), \dots, \text{in}_w(f_s))$ is regular in $\mathcal{O}$, Lemma A.11 ensures that the Koszul complex (A.1) is exact at E . In turn, Proposition A.4 implies that $(f_1, \dots, f_s)$ is a regular sequence in $\mathcal{O}$. This proves item (1) of the statement.

To finish, we must show that the w -initial forms $\{\text{in}_w(f_1), \dots, \text{in}_w(f_s)\}$ generate the w -initial ideal $\text{in}_w(J)\mathcal{O}$. By definition, the ideal generated by these forms is contained in $\text{in}_w(J)\mathcal{O}$. As $\text{in}_w(J)\mathcal{O}$ is generated over $\mathcal{O}$ by all elements $\text{in}_w(g)$ with $g \in J$, the reverse inclusion will follow immediately if we show that $\text{in}_w(g) \in (\text{in}_w(f_1), \dots, \text{in}_w(f_s))\mathcal{O}$. This identity is a direct consequence of Lemma A.8. Therefore, item (2) holds. This concludes our proof. $\square$

References

- Améndola, C., Kohn, K., Lamboglia, S., Maclagan, D., Smith, B., Sommars, J., Tripoli, P., Zajackowska, M.: Tropical: a package for doing computations in tropical geometry. Version 0.1. Available at <https://www.github.com/Macaulay2/M2/tree/master/M2/Macaulay2/packages>
- Aroca, F., Gómez-Morales, M., Shabbir, K.: Torical modification of Newton non-degenerate ideals. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM* **107**(1), 221–239 (2013)
- Aroca, F., Gómez-Morales, M., Mourtada, H.: Groebner fan and embedded resolutions of ideals on toric varieties. *Beitr. Algebra Geom.* (2023). <https://doi.org/10.1007/s13366-023-00684-0>
- Atiyah, M., Macdonald, I.G.: *Commutative Algebra*. Addison Wesley, Berlin (1969)
- Bivià-Ausina, C.: Mixed Newton numbers and isolated complete intersection singularities. *Proc. Lond. Math. Soc.* (3) **94**(3), 749–771 (2007)
- Cueto, M.A., Popescu-Pampu, P., Stepanov, D.: The Milnor fiber conjecture of Neumann and Wahl, and an overview of its proof. In: *Essays in Geometry, Dedicated to Norbert A'Campo*, pp. 629–710. EMS Publishing House, Berlin (2023)
- de Felipe, A.B., González Pérez, P., Mourtada, H.: Resolving singularities of curves with one toric morphism. *Math. Ann.* (2022). <https://doi.org/10.1007/s00208-022-02504-7>
- de Jong, T., Pfister, G.: Local analytic geometry. Basic theory and applications. In: *Advanced Lectures in Mathematics*. Friedr. Vieweg & Sohn, Braunschweig (2000). xii+382 pp. ISBN: 3-528-03137-9
- Eisenbud, D.: *Commutative algebra. With a view toward algebraic geometry*. In: *Graduate Texts in Mathematics*, vol. 150. Springer, New York (1995). xvi+785 pp. ISBN: 0-387-94268-8
- Eisenbud, D., Neumann, W.: *Three-Dimensional Link Theory and Invariants of Plane Curve Singularities*. Princeton University Press, Princeton (1985)
- Ewald, G.: Combinatorial convexity and algebraic geometry. In: *Graduate Texts in Mathematics*, vol. 168. Springer, New York (1996). xiv+372 pp. ISBN: 0-387-94755-8
- Fulton, W.: *Introduction to Toric Varieties*. *Annals of Mathematics Studies*, vol. 131. Princeton University Press, Princeton (1993)
- García Barroso, E., González Pérez, P., Popescu-Pampu, P.: Ultrametric spaces of branches on arborescent singularities. In: *Singularities, Algebraic Geometry, Commutative Algebra, and Related Topics*, pp. 55–117. Springer, Berlin (2018)
- García Barroso, E., González Pérez, P., Popescu-Pampu, P.: The valuative tree is the projective limit of Eggers–Wall trees. *RACSAM* **113**, 4051–4105 (2019)
- García Barroso, E., González Pérez, P., Popescu-Pampu, P.: The combinatorics of plane curve singularities. In: *Handbook of Geometry and Topology of Singularities I*, pp. 1–150. Springer, Cham (2020)
- García Barroso, E., González Pérez, P., Popescu-Pampu, P., Ruggiero, M.: Ultrametric properties for valuation spaces of normal surface singularities. *Trans. Am. Math. Soc.* **372**(12), 8423–8475 (2019)
- Goldin, R., Teissier, B.: Resolving singularities of plane analytic branches with one toric morphism. In: *Resolution of Singularities (Oberurgel, 1997)*, *Progr. Math.*, vol. 181, pp. 315–340. Birkhäuser, Basel (2000)
- Grayson, D., Stillman, M.E.: Macaulay2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>
- Hamm, H.: Die Topologie isolierter Singularitäten von vollständigen Durchschnitten komplexer Hyperflächen. *Dissertation Bonn* (1969)
- Hamm, H.: Exotische Sphären als Umgebungsränder in speziellen komplexen Räumen. *Math. Ann.* **197**, 44–56 (1972)
- Hartshorne, R.: Complete intersections and connectedness. *Am. J. Math.* **84**, 497–508 (1962)
- Khovanskii, A.G.: Newton polyhedra and toroidal varieties. *Funktsional. Anal. i Prilozhen.* **11**, 56–64 (1977); English translation, *Functional Anal. Appl.* **11**, 289–296 (1977)
- Khovanskii, A.G.: Newton polyhedra (resolution of singularities). *J. Sov. Math.* **27**, 2811–2830 (1984)
- Kouchnirenko, A.G.: Newton polyhedron and Milnor numbers. *Funct. Anal. Appl.* **9**(1), 71–72 (1975)
- Kouchnirenko, A.G.: Polyèdres de Newton et nombres de Milnor. *Invent. Math.* **32**, 1–31 (1976)
- Küronya, A., Souza, P., Ulirsch, M.: Tropicalization of toric prevarieties (2021). [arXiv:2107.03139](https://arxiv.org/abs/2107.03139)
- Lamberson, P.J.: The Milnor fiber conjecture and iterated branched cyclic covers. *Trans. Am. Math. Soc.* **361**, 4653–4681 (2009)
- Maclagan, D., Sturmfels, B.: *Introduction to tropical geometry*. In: *Graduate Studies in Mathematics*, vol. 161. American Mathematical Society, Providence (2015)

29. Maurer, J.: Puiseux expansion for space curves. *Manuscr. Math.* **32**, 91–100 (1980)
30. Munkres, J.R.: *Topology, a First Course*, 2nd edn. Prentice Hall Inc., Upper Saddle River (2000). xvi+537 pp. ISBN: 0-13-181629-2
31. Némethi, A.: The cohomology of line bundles of splice-quotient singularities. *Adv. Math.* **229**(4), 2503–2524 (2012)
32. Némethi, A., Okuma, T.: The Seiberg–Witten invariant conjecture for splice-quotients. *J. Lond. Math. Soc.* (2) **78**(1), 143–154 (2008)
33. Némethi, A., Okuma, T.: On the Casson invariant conjecture of Neumann–Wahl. *J. Algebraic Geom.* **18**(1), 135–149 (2009)
34. Neumann, W.: Brieskorn complete intersections and automorphic forms. *Invent. Math.* **42**, 285–293 (1977)
35. Neumann, W.: Abelian covers of quasihomogeneous surface singularities. In: *Singularities, Part 2* (Arcata, Calif., 1981). *Proc. Sympos. Pure Math.*, vol. 40, pp. 233–243. Amer. Math. Soc., Providence (1983)
36. Neumann, W.: Graph 3-manifolds, splice diagrams, singularities. In: *Singularity Theory*, pp. 787–817, World Sci. Publ., Hackensack (2007)
37. Neumann, W., Wahl, J.: Universal abelian covers of surface singularities. In: *Trends in Singularities*, pp. 181–190. Birkhäuser, Basel (2002)
38. Neumann, W., Wahl, J.: Complete intersection singularities of splice type as universal abelian covers. *Geom. Topol.* **9**, 699–755 (2005)
39. Neumann, W., Wahl, J.: Complex surface singularities with integral homology sphere links. *Geom. Topol.* **9**, 757–811 (2005)
40. Neumann, W., Wahl, J.: The end curve theorem for normal complex surface singularities. *J. Eur. Math. Soc. (JEMS)* **12**(2), 471–503 (2010)
41. Oka, M.: Non-degenerate complete intersection singularity. *Actualités Mathématiques [Current Mathematical Topics]*. Hermann, Paris (1997)
42. Okuma, T.: Universal abelian covers of certain surface singularities. *Math. Ann.* **334**(4), 753–773 (2006)
43. Okuma, T.: The geometric genus of splice-quotient singularities. *Trans. Am. Math. Soc.* **360**(12), 6643–6659 (2008)
44. Okuma, T.: Another proof of the end curve theorem for normal surface singularities. *J. Math. Soc. Jpn.* **62**(1), 1–11 (2010)
45. Okuma, T.: Invariants of splice quotient singularities. In: *Singularities in Geometry and Topology*, IRMA Lect. Math. Theor. Phys., vol. 20, pp. 149–159. Eur. Math. Soc., Zürich (2012)
46. Okuma, T.: The multiplicity of abelian covers of splice quotient singularities. *Math. Nachr.* **288**(2–3), 343–352 (2015)
47. Pedersen, H.: Splice diagram determining singularity links and universal abelian covers. *Geom. Dedic.* **150**, 75–104 (2011)
48. Pedersen, H.: Splice diagram singularities and the universal abelian cover of graph orbifolds. *Geom. Dedic.* **195**, 283–305 (2018)
49. Popescu-Pampu, P., Stepanov, D.: Local tropicalization. In: *Algebraic and Combinatorial Aspects of Tropical Geometry*, *Contemp. Math.*, vol. 589, pp. 253–316. Amer. Math. Soc., Providence (2013)
50. Saia, M.J.: The integral closure of ideals and the Newton filtration. *J. Algebra Geom.* **5**, 1–11 (1996)
51. Serre, J.-P.: *Algèbre locale. Multiplicités*. *Lecture Notes in Maths.*, vol. 11. Springer, London (1965)
52. Siebenmann, L.C.: On vanishing of the Rohlin invariant and nonfinitely amphicheiral homology 3-spheres. In: *Topology Symposium, Siegen 1979* (*Proc. Sympos., Univ. Siegen, Siegen, 1979*). *Lecture Notes in Math.*, vol. 788, pp. 172–222. Springer, Berlin (1980)
53. Steenbrink, J.H.M.: Motivic Milnor fibre for nondegenerate function germs on toric singularities. In: *Bridging Algebra, Geometry, and Topology*. *Springer Proc. Math. Stat.*, vol. 96, pp. 255–267. Springer, Cham (2014)
54. Stepanov, D.: Universal valued fields and lifting points in local tropical varieties. *Commun. Algebra* **45**(2), 469–480 (2017)
55. Teissier, B.: Monomial ideals, binomial ideals, polynomial ideals. In: *Trends in Commutative Algebra*, *Math. Sci. Res. Inst. Publ.*, vol. 51, pp. 211–246. Cambridge Univ. Press, Cambridge (2004)
56. Tevelev, J.: Compactifications of subvarieties of tori. *Am. J. Math.* **129**, 1087–1104 (2007)
57. Varchenko, A.N.: Zeta-function of monodromy and Newton’s diagram. *Invent. Math.* **37**, 253–262 (1976)

- 58. Wahl, J.: Topology, geometry, and equations of normal surface singularities. In: Singularities and Computer Algebra, London Math. Soc. Lecture Note Ser., vol. 324, pp. 351–371. Cambridge University Press, Cambridge (2006)
- 59. Wahl, J.: Splice diagrams and splice quotient surface singularities. In: *Celebratio Mathematica*, Volume in Honor of Walter D. Neumann, article 1030 (2022)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.