



## OPEN ACCESS

## EDITED BY

Diana Sofia Madeira,  
University of Aveiro, Portugal

## REVIEWED BY

Renato Crespo Pereira,  
Fluminense Federal University, Brazil  
Masahiro Nakaoka,  
Hokkaido University, Japan

## \*CORRESPONDENCE

Colleen A. Burge  
✉ Colleen.Burge@wildlife.ca.gov;  
✉ caburge@ucdavis.edu

## †PRESENT ADDRESSES

Sukanya Dayal,  
Department of Biology and Marine Biology,  
University of North Carolina Wilmington,  
Wilmington, NC, United States  
James Sanghyun Lee,  
Front and Centered, Seattle, WA,  
United States  
Colleen A. Burge,  
California Department of Fish & Wildlife, UC  
Davis Bodega Marine Laboratory, Bodega  
Bay, CA, United States

<sup>†</sup>These authors have contributed  
equally to this work and share  
first authorship

<sup>‡</sup>These authors have contributed  
equally to this work and share  
last authorship

RECEIVED 28 January 2023

ACCEPTED 23 June 2023

PUBLISHED 08 August 2023

## CITATION

Venkataraman YR, Shore A, Dayal S,  
Lee JS, Alidoost Salimi M, Crandall G,  
Loeher MM, Stoops M, Swanger M,  
Eisenlord ME, Van Alstyne KL, Fast MD,  
Burge CA and Groner ML (2023)  
Characterizing host-pathogen  
interactions between *Zostera marina*  
and *Labyrinthula zosterae*.  
*Front. Mar. Sci.* 10:1152647.  
doi: 10.3389/fmars.2023.1152647

## COPYRIGHT

© 2023 Venkataraman, Shore, Dayal, Lee,  
Alidoost Salimi, Crandall, Loeher, Stoops,  
Swanger, Eisenlord, Van Alstyne, Fast, Burge  
and Groner. This is an open-access article  
distributed under the terms of the [Creative  
Commons Attribution License \(CC BY\)](#). The  
use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in  
this journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted  
which does not comply with these terms.

# Characterizing host-pathogen interactions between *Zostera marina* and *Labyrinthula zosterae*

Yaamini R. Venkataraman<sup>1†</sup>, Amanda Shore<sup>2‡</sup>, Sukanya Dayal<sup>3†</sup>,  
James Sanghyun Lee<sup>4†</sup>, Mahsa Alidoost Salimi<sup>5</sup>,  
Grace Crandall<sup>6</sup>, Malina M. Loeher<sup>7</sup>, Mark Stoops<sup>8</sup>,  
Megan Swanger<sup>6</sup>, Morgan E. Eisenlord<sup>9</sup>,  
Kathryn L. Van Alstyne<sup>10</sup>, Mark D. Fast<sup>11</sup>, Colleen A. Burge<sup>3,12\*†§</sup>  
and Maya L. Groner<sup>11,13§</sup>

<sup>1</sup>Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA, United States

<sup>2</sup>Department of Biology, Farmingdale State College, Farmingdale, NY, United States, <sup>3</sup>Institute of  
Marine & Environmental Technology, University of Maryland Baltimore County, University of Maryland  
Baltimore, Baltimore, MD, United States, <sup>4</sup>School of Marine and Environmental Affairs, University of  
Washington, Seattle, WA, United States, <sup>5</sup>School of Geography, Earth and Atmospheric Sciences,  
University of Melbourne, Victoria, VIC, Australia, <sup>6</sup>School of Aquatic and Fishery Sciences, University of  
Washington, Seattle, WA, United States, <sup>7</sup>Department of Aquatic Health Sciences, Virginia Institute of  
Marine Science, Gloucester Point, VA, United States, <sup>8</sup>University of North Carolina (UNC) Chapel Hill  
Institute of Marine Sciences, Morehead City, NC, United States, <sup>9</sup>Department of Ecology and  
Evolutionary Biology, Cornell University Corson Hall, Ithaca, NY, United States, <sup>10</sup>Shannon Point  
Marine Center, Western Washington University, Anacortes, WA, United States, <sup>11</sup>Hoplite Lab,  
Department of Pathology and Microbiology, Atlantic Veterinary College-University of Prince Edward  
Island (AVC-UPEI), Charlottetown, PE, Canada, <sup>12</sup>Department of Microbiology, University of Maryland  
Baltimore, Baltimore, MD, United States, <sup>13</sup>Bigelow Lab for Ocean Sciences, East Boothbay,  
ME, United States

**Introduction:** Seagrass meadows serve as an integral component of coastal  
ecosystems but are declining rapidly due to numerous anthropogenic stressors  
including climate change. Eelgrass wasting disease, caused by opportunistic  
*Labyrinthula* spp., is an increasing concern with rising seawater temperature. To  
better understand the host-pathogen interaction, we paired whole organism  
physiological assays with dual transcriptomic analysis of the infected host  
and parasite.

**Methods:** Eelgrass (*Zostera marina*) shoots were placed in one of two  
temperature treatments, 11° C or 18° C, acclimated for 10 days, and exposed  
to a waterborne inoculation containing infectious *Labyrinthula zosterae* (*Lz*) or  
sterile seawater. At two- and five-days post-exposure, pathogen load, visible  
disease signs, whole leaf phenolic content, and both host- and pathogen-  
transcriptomes were characterized.

**Results:** Two days after exposure, more than 90% of plants had visible lesions and  
*Lz* DNA was detectable in 100% percent of sampled plants in the *Lz* exposed  
treatment. Concentrations of total phenolic compounds were lower after 5 days  
of combined exposure to warmer temperatures and *Lz*, but were unaffected in  
other treatments. Concentrations of condensed tannins were not affected by *Lz*  
or temperature, and did not change over time. Analysis of the eelgrass  
transcriptome revealed 540 differentially expressed genes in response to *Lz*  
exposure, but not temperature. *Lz*-exposed plants had gene expression patterns

consistent with increased defense responses through altered regulation of phytohormone biosynthesis, stress response, and immune function pathways. Analysis of the pathogen transcriptome revealed up-regulation of genes potentially involved in breakdown of host defense, chemotaxis, phagocytosis, and metabolism.

**Discussion:** The lack of a significant temperature signal was unexpected but suggests a more pronounced physiological response to *Lz* infection as compared to temperature. Pre-acclimation of eelgrass plants to the temperature treatments may have contributed to the limited physiological responses to temperature. Collectively, these data characterize a widespread physiological response to pathogen attack and demonstrate the value of paired transcriptomics to understand infections in a host-pathogen system.

#### KEYWORDS

*Zostera marina*, *Labyrinthula zosterae*, eelgrass wasting disease, transcriptomics, host-pathogen interactions, marine disease

## Introduction

Despite occupying less than 0.1% of the ocean floor (primarily in estuaries and embayments), seagrasses provide numerous ecosystem services, including shoreline stabilization and protection from erosion, nutrient cycling, sediment retention, reduction of pathogens in the water column, and provision of critical nursery habitat for numerous invertebrate and fish species, many of which are of economic importance to the seafood industry (Nordlund et al., 2016; Lamb et al., 2017). Seagrass meadows are threatened by a wide range of global and local stressors associated with anthropogenic activity such as coastal development, increasing sea surface temperature, rising sea levels, sediment and nutrient runoff and disease (Orth et al., 2006; Sullivan et al., 2013). Consequently, over the past century, global seagrass beds have experienced approximately 19% loss in cover (Dunic et al., 2021).

*Zostera marina*, the most widely distributed seagrass species in the northern hemisphere (Short et al., 2007), is susceptible to infection with the protist *Labyrinthula zosterae* (*Lz*), which causes eelgrass wasting disease (EWD). While chronic *Lz* infections are nearly ubiquitous in distribution, acute epidemics of wasting disease can cause declines in eelgrass populations, most notably in the 1930s when Atlantic eelgrass populations were nearly eliminated along the coasts of North America and Europe (Short et al., 1987). *Lz* is a pathogenic Labyrinthulomycete and can be transmitted through direct contact between healthy and infected blades within an eelgrass bed (Muehlstein, 1992) or through the water column (Groner et al., 2014; Groner et al., 2018). Upon infection, *Lz* targets chloroplasts of eelgrass leaves, compromising photosynthesis, and ultimately causing necrosis of eelgrass leaves, and in some cases, entire plants (Muehlstein, 1992). Non-lethal impacts of EWD include reductions in growth and below-ground storage of carbohydrates, which may impact seasonal resilience of eelgrass

meadows (Graham et al., 2021). Strains of *Lz* vary in virulence, raising questions about mechanisms of virulence employed by this pathogen and potential costs associated with those mechanisms (Martin et al., 2016).

Eelgrass has a robust defense system which includes proteins and surface-associated metabolites like fatty acids and phenolics, *p*-coumaric acid, rosmarinic acid, and zosteric acid (Papazian et al., 2019). These compounds play a wide-range of roles including but not limited to signaling molecules, antioxidant activity, free radical scavenging activity, and regulators of auxin transport (Ma et al., 2016). Phenolic compounds which are produced in response to stress and pathogen presence can be also considered as defense-related secondary metabolites. Indeed, some phenolic compounds have been shown to inhibit growth of *Lz* *in vitro* (Vergeer and Develi, 1997) and concentrations of total phenols (% dry mass) are increased in diseased plants in the field (Groner et al., 2016). In terrestrial plants, phytohormones including abscisic acid (ABA), jasmonic acid (JA), and salicylic acid (SA) play significant roles in regulating defense responses against a great number of biotic and abiotic factors (Thaler et al., 2004; Tamaoki et al., 2013), however these hormones have not been well explored in marine plants such as eelgrass.

As with many marine diseases, EWD is sensitive to temperature (Tracy et al., 2019). Both EWD prevalence and severity are correlated with sea surface temperatures in the Salish Sea (WA, USA), and in the Isles of Scilly (UK), (Bull et al., 2012; Groner et al., 2021). Increased temperatures enhance the growth-reducing effects of EWD on eelgrass (Bull et al., 2012). These changes are likely driven by both the host and pathogen. *Lz* is also sensitive to temperature, with faster *in vitro* growth documented at 18°C compared to 11°C for a strain isolated from the Salish Sea (WA, USA) (Dawkins et al., 2018); similarly, increased disease severity was noted in eelgrass exposed to *Lz* and held at 18°C compared to 11°C (Agnew et al., 2022). Recent studies have correlated heatwaves

to substantial seagrass die-offs (Strydom et al., 2020; Groner et al., 2021), reduced restoration success (Aoki et al., 2020), and changes in fatty acid composition (Beca-Carretero et al., 2018), showing that seagrasses are sensitive to warming conditions.

To better understand the factors that contribute to *Lz* infection and provide insight on mechanisms of virulence and host defense, we characterized the production of defensive compounds in *Z. marina* and the transcriptional responses of both *Z. marina* and *Lz* following a controlled *Lz* challenge conducted at ambient and warm seawater temperatures. We hypothesize that both increased temperature and *Lz* infection will alter expression of genes related to immune function and stress response in *Z. marina*. More specifically, we hypothesize that phenolic and hormonal gene pathways will be enriched in response to presence of *Lz*, but we expected these responses may be reduced at warmer temperatures that could be stressful to *Z. marina*. Finally, we hypothesize that *Lz* will exhibit a change in gene expression in response to temperature, reflecting the association between EWD and warmer seawater temperatures found in nature (Brakel et al., 2019; Groner et al., 2021).

## Materials and methods

### Experimental design

Similarly sized eelgrass ramets were collected from False Bay, San Juan Island, WA (48.550° N, 123.008° W) on 29 July 2015. Epiphytes, grazers and damaged leaves were removed from ramets and they were held in flow-through 14°C, 0.2 µm filter-sterilized seawater at Friday Harbor Labs, WA, USA.

Eelgrass ramets were placed in one of two temperature treatments (target temperatures 11° or 18°C, actual temperatures ranged  $\pm 1^{\circ}\text{C}$  from target) on 3 Aug 2015 and allowed to acclimate to these treatments for 10 days. These temperatures represent typical summer sea surface temperatures found locally along coastal eelgrass beds at high and low tides, respectively. A constant exposure to 18°C would be unusually warm in this region. Each treatment was replicated eight times for a total of 16

experimental units. Each replicate unit contained four ramets. Experimental units were 4-L containers with 3500 mL of 0.2 µm filter-sterilized seawater. Experimental units were split into groups of eight that were clustered in a common cooler. Each cooler received flow-thru 0.2 µm filtered and UV-irradiated seawater. Seawater temperatures were continuously monitored and adjusted using Honeywell UDA2182 pH controller and Honeywell Durafet III electrodes. After flowing into the cooler, water was then piped into the 8 experimental units. Water temperatures were maintained by circulating it through chilled water set 1°C below target temperatures and, if necessary, heating it with aquatic heaters placed in the coolers (i.e. O'Donnell et al., 2013). Two temperature loggers (iButtons, Whitewater, WI) set to record every 30 min were placed in each cooler for independent confirmation of temperature. Full-spectrum lights were kept to a 12: 12 hr light: dark schedule with a mean light output of  $161 \pm 3$  (mean  $\pm$  SE) µmol/m<sup>2</sup>/sec PAR below the water surface. To reduce algal blooms, which could clog plumbing and block light, tanks were treated daily with 0.67 ppm (final concentration) of Germanium dioxide (GeO<sub>2</sub>; Markham and Hagmeier, 1982).

A 36-hr incubation with the pathogen inoculant was conducted after 10 days of temperature acclimation (Figure 1). Immediately prior to inoculation, all ramets were photographed, weighed, labelled with uniquely colored zip-ties, and pin-pricked at the sheath for later measurements of growth. All ramets within a replicate were placed in Whirlpaks with 98 mL of sterile seawater. Control treatments were inoculated with 2 mL of sterile seawater. *Lz* treatments were inoculated with 2 mL of inoculant for a final concentration of  $2.5 \times 10^5$  cells per mL (additional details in Supplementary Material). After adding the inoculate, the Whirlpaks were closed and floated in treatment containers to maintain target temperatures. 36 hours later, ramets were removed from Whirlpaks and placed back into treatment containers. The *Lz* strain used in this experiment, 8.16.D, has been used in several previous experiments, (e.g., Groner et al., 2014; Groner et al., 2018) and was isolated in 2011 from non-flowering adult *Zostera marina* shoots that were collected at Picnic Cove, Shaw Island, 48° 33.942' N, 122° 55.448' W).




FIGURE 1

Timeline of experimental treatments and sampling. After collection and cleaning of plants to remove grazers, epiphytes and diseased tissue, eelgrass ramets were held at 14°C for 1d and then transferred in groups of 4 ramets to 11°C or 18°C for a 10d acclimation period. Each group of 4 ramets was replicated 8 times during the acclimation period. After that, the groups were equally divided into control and *Lz* – exposed treatments at each of the two temperatures. *Lz* or sham exposures occurred for 36 hr. Subsampling of plants occurred at days 2 and 5 post-*Lz* exposure.

Forty-eight hours after inoculation (experiment day 13), a random selection of half of the individuals (48 total) from each experimental unit were removed for sampling (Figure 1). Each of those individuals was photographed and weighed. Then the roots and shoots were frozen in liquid nitrogen for transcriptomic sampling. The oldest leaf was split lengthwise into two pieces. One half was frozen in liquid nitrogen for transcriptomic analysis, and the other was stored in 70% molecular grade ethanol at room temperature for subsequent measurement of *Lz* load using quantitative PCR (qPCR) as described in Groner et al. (2018). The second oldest leaf was frozen in liquid nitrogen for measurement of total phenolic compounds and condensed tannins. Five days after inoculation, the experiment was ended because the plants had severe disease signs. At this point, the remaining ramets were photographed and sampled (as described above) for measurement of visual disease signs (e.g., lesions with a distinct black border, *sensu* Groner et al., 2014) as well as phenol and tannin concentrations (Figure 1).

## Measurement of total phenolic compounds and condensed tannins

Total phenolic compounds were measured in 96 well microplates using the methods described in Groner et al. (2018). Briefly, sampled eelgrass leaves were frozen (-20°C) and sent on dry ice to the Shannon Point Marine Center for analysis. Frozen tissues were lyophilized and ground to a fine powder in an SPEX mixer/mill (SPEX, Metuchen, NJ). Ground tissue (9–10 mg) was extracted overnight in 1 mL of HPLC-grade methanol, then diluted in ANSI Type I water (19 parts water to 1 part extract). Forty  $\mu$ L of 40% Folin-Ciocalteu's Reagent (Sigma F9252) was added to each 100  $\mu$ L aliquot (n=3 per sample) of each of the diluted extracts, mixed for 5 min prior to the addition of 100  $\mu$ L of 2N sodium carbonate. Samples were shaken for another 30 min in a 50°C chamber and then the absorbance of each cell was read at 765 nm. Concentrations were standardized using native standards from *Z. marina* collected from Ship Harbor WA, using caffeic acid as a secondary standard (Groner et al., 2016). The remaining methanolic extracts were used for measurements of condensed tannins, which were conducted with a sulfuric acid method (Bate-Smith and Rasper, 1969; Nitao et al., 2001) that was modified for use in a microplate reader. Condensed tannins (proanthocyanidins) were cleaved by the addition of 43% sulfuric acid in methanol (250  $\mu$ L/well containing 100  $\mu$ L of a methanol extract) for 30 min at 50°C for. Tannin cleavage produces red-colored chromophores (anthocyanins), which were then quantified spectrophotometrically at 550 nm using a Biotek Synergy multiplate reader. Cyanidin (Cayman Chemical, purity >98%) was used as a standard. All assessments were run in triplicate.

## Quantification of *Lz* abundance

Samples (mean dry weight of leaf material = 7 mg) were preserved in 70% ethanol until DNA extractions were performed

using a Qiagen DNeasy Plant Mini Kit following the manufacturer's instructions. Prior to DNA extractions, samples were removed from ethanol, patted dry, weighed and then transferred into a clean 1.5 mL micro-centrifuge tube with one 3 mm tungsten carbide bead (Qiagen) per tube. Samples were then placed on dry ice for 10–15 minutes and lysed using a Qiagen TissueLyser at a speed of 25 Hz for 5 minutes.

Presence of *Lz* DNA was detected and quantified using qPCR detection using primers and probe targeting the *Lz* ITS region (Bockelmann et al., 2013); Laby\_ITS\_Taq\_f (5'- TTG AAC GTA ACA TTC GAC TTT CGT- 3') and the reverse primer Laby\_ITS\_Taq\_r (5' -ACG CAT GAA GCG GTC TTC TT -3') were used, in addition to the TaqMan probe Laby\_ITS\_probe (5' FAM- TGG ACG AGT GTG TTT TG -MGB-NFQ 3'). qPCR reaction conditions and the standard curve (serial dilution between 1.37–1.37\*10<sup>4</sup> cells) were as described by Groner et al., 2018 on run on the Applied Biosystems 7500 Fast Real-Time PCR System. For each qPCR plate, we aimed for R2 = 0.999 and an efficiency of 90–110%; the mean reaction efficiency was 105.5% +/- 1.5% (cross three total plates). We used a cut-off of 1.37 cells as our limit of detection (amplification of 1.37 cells was achieved in all reactions in each cell curve).

## Statistical analyses of treatment effects on phenolic compounds, tannins, disease signs and *Lz* DNA

Independent and interactive effects of temperature and *Lz* exposure on concentrations of phenols, concentrations of tannins, presence of suspected disease signs and  $\log_{10}$  transformed *Lz* DNA concentrations in leaf tissues were quantified for the time points when these measurements were taken (R v. 4.1.1). Linear regression was used to quantify treatment effects on total phenolic compounds, condensed tannins, and *Lz* DNA concentrations, while logistic regression was used to quantify treatment effects on the presence of disease signs (package lme4, Bates et al., 2015). In all models, the 4L container replicate was included as a random effect.

## RNA extractions and transcriptome sequencing

Total RNA from four samples per treatment was extracted following the Qiagen RNeasy plant kit protocol; samples were chosen based on both quality of RNA and qPCR results. DNA was removed from extracted RNA using the Turbo DNA-free treatment according to the manufacturer's instructions (Ambion Inc., The Life Technologies Corporation<sup>TM</sup>, Grand Island, NY). Removal of DNA was confirmed by using RNA (1  $\mu$ L) as template in a qPCR targeting 18s ribosomal DNA as previously described (Burge and Friedman, 2012). RNA concentrations were quantified using the NanoDrop<sup>®</sup> ND-1000 (NanoDrop Technologies, Wilmington, DE). Samples were shipped to McGill University and Genome Quebec for library preparation and sequencing.

RNA quality was assessed prior to library preparation using an Agilent Bioanalyzer. The Illumina TruSeq stranded cDNA kit was used for library preparation and barcoding; quality assessment of libraries indicated that one sample was not of high enough quality for sequencing. The remaining samples were distributed across two sequencing lanes on an Illumina HiSeq 4000 platform (2x150 bp paired end reads).

## Transcriptome assembly and annotation

Prior to *de novo* assembly, reads were quality screened (FastQC Version 0.11.4, Babraham Bioinformatics), and sequences were screened for remaining Illumina adapter sequences (Trimmomatic, Bolger et al., 2014). *De novo* assembly was carried out using reads from all fifteen samples using Trinity (Grabherr et al., 2011; Haas et al., 2013, Trinity RNASeq Github). Following transcriptome assembly, reads for each sample were matched back to gene isoforms.

A multi-step annotation process was used to annotate the transcriptome in order to separate isoform consensus sequences (or contigs) matching to *Z. marina* and those non-matching contigs (“non-host”). First, a database of annotations was created from the *Z. marina* genome (GCA\_001185155.1) using a BLASTx search of the UniProtKB/Swiss-Prot database. The BLASTx algorithm (Altschul et al., 1990) was used to search the *Z. marina* database with a threshold E-value of 0.00001; each contig matching as *Z. marina* was filtered out. Similarly, the blastx algorithm was used to search the Swiss-Prot database with a threshold E-value of 0.00001 to form the non-*Zostera* annotation for subsequent analysis. Finally, to determine which non-host contigs have significant similarity to *Labyrinthula* and other closely related ‘slime-mold’ species, the blast algorithm was used to match non-host contigs against the recently added *Labyrinthula* sp. Ha genome (NCBI Accession # GCA\_015227615.1) and also against the genomes of two other, better annotated genomes, a thraustochytrid, *Hondaea fermentalgiana* (NCBI Accession # GCA\_002897355.1) and the amoeba model organism, *Dictyostelium discoideum* (NCBI Accession # GCF\_000004695.1).

Gene ontology (GO) information was used to annotate both *Z. marina* and non-*Zostera* transcriptome data separately. Gene ontology IDs and associated GO Slim terms for biological processes, cellular components, and molecular function categories were downloaded from the UniProtKB/Swiss-Prot database. Swiss-Prot identifiers from BLASTx output were joined to gene ontology terms. Since assembly and annotation were conducted at the isoform level, transcriptome data was streamlined to conduct differential expression analysis on genes. Annotated sequences were sorted by contig name and e-value; contig names included a contig, gene, and isoform identifier. The isoform entry with the lowest e-value was retained for each gene. The final annotated dataset for both *Z. marina* and non-*Zostera* transcriptomes included only one entry per gene. Annotated genes were merged with count data to prepare for differential expression analysis. The data is deposited in the NCBI database in the SRA as Bioproject (PRJNA990835).

## Analysis of *Zostera* and non-*Zostera* transcriptomes

Statistical differences in count data were conducted for *Z. marina* and non-*Zostera* annotated genes using EdgeR (Robinson et al., 2010; McCarthy et al., 2012; Chen et al., 2016). Briefly, we utilized EdgeR to filter and normalize the count data based on the library size and examine samples for outliers using MDS plots. Next, a multi-factorial model including estimated dispersion (i.e. estimate of biological variation) was used with multiple contrasts to calculate fold change, log CPM (counts per million), and raw and Benjamini-Hochberg adjusted p-values to correct for False Discovery Rate.

We tested for statistical differences using several contrasts followed by the primary questions of interest, respectively examining the relationships of the pathogen-by-temperature interaction, temperature alone, and pathogen exposure alone:

- 1) What genes are differentially expressed between Lz-exposed shoots at elevated temperatures? At ambient temperatures?
- 2) What genes are differentially expressed between elevated and ambient temperatures, irrespective of Lz-exposure?
- 3) What genes are differentially expressed between Lz-exposed and control shoots, irrespective of temperature?


Very few genes were differentially expressed in questions 1 and 2 (25 genes and 0 genes, respectively). Thus, for the remainder of the analysis we focused on the effect of Lz exposure alone. Differentially expressed genes were pooled into host and pathogen genes. Genes were considered differentially expressed based on FDR adjusted p-values from edgeR ( $p < 0.05$ ).

To identify groups of *Z. marina* genes with similar expression patterns (eigengenes) and associate these patterns with experimental conditions, a Weighted Gene Co-expression Network Analysis (WGCNA) was conducted using the WGCNA package in R (version 1.69; Langfelder and Horvath, 2008). The WGCNA was only conducted for *Z. marina* genes, as transcriptome sequencing produced a high magnitude of information for *Z. marina* that was difficult to parse without additional analysis. After identifying significant modules, overrepresented biological process and molecular function GOterms were identified using the GO-MWU enrichment method (Wright et al., 2015). More details on WGCNA and GO-MWU analyses are available in the [Supplemental Material](#).

## Results

### Disease progression and pathogen load

The inoculations were successful in transmitting Lz to the exposed plants. Two days after exposure, Lz concentrations in eelgrass tissue were 1.1 orders of magnitude greater in the 18°C treatment as compared to the 11°C treatment (with concentrations of  $4.2 \pm 4.3 \times 10^4$  and  $3.7 \pm 3.6 \times 10^3$  (mean  $\pm$  SE) Lz cells/mg of eelgrass tissue, respectively) (Figure 2). These pathogen concentrations were



**FIGURE 2**  
Lz cell equivalents (per mg eelgrass tissue) from qPCR analysis of eelgrass leaf tissue 2 days post-inoculation. Bars indicate standard errors. Letters and bars indicate significantly different treatments.

significantly higher than the control treatments, where no Lz was detected using qPCR ( $t$  value = 12.3,  $p < 0.0001$ ). No other factors affected Lz abundance in our linear regression ( $\text{all } p > 0.05$ ).

Visible disease signs were recorded in all treatments (Figure 3). Pathogen exposure was a significant predictor of disease signs on day 2 and a marginally significant predictor of disease signs on day 5. On day 2, the disease signs were recorded in  $93.7 \pm 8.8\%$  (mean  $\pm$  SE) of the exposed plants and  $28.8 \pm 19.7\%$  of the control plants ( $t = 2.1$ ,  $p = 0.04$ ). Neither temperature nor the temperature by exposure interaction were predictors of visual disease signs. These values increased on day 5 where disease signs were recorded in 100

$\pm 0\%$  (mean  $\pm$  SE) of the exposed plants and  $37.5 \pm 17.7\%$  of the control plants ( $t = 1.99$ ,  $p = 0.07$ ).


The concentrations of total phenolic compounds and condensed tannins were not impacted by temperature or Lz treatments on day 2 (Figure 4). By day 5, total phenolic compounds were significantly lower in the warmer exposed treatment, where they were  $0.70 \pm 0.17\%$  of dry mass as opposed to ranging between  $1.15$  and  $1.39\%$  of dry mass in the other three treatments ( $t$  value =  $-3.298$ ,  $p < 0.001$  for the temperature by exposed interaction). There was no significant impact of temperature or Lz exposure on the concentration of condensed tannins in eelgrass leaves on day 5.

## Transcriptome assembly and annotation

Trinity *de novo* assembly yielded 1,065,804 isoforms (767,296 potential genes) with an average read length of 654.71 bp (median of 349) and an N50 value of 1075. BLAST searches of the *de novo* assembly yielded 281,600 contigs matching the *Z. marina* genome. 784,204 contigs did not match the *Z. marina* genome and thus were categorized as ‘non-host’ contigs.

## Differential expression analysis - host contigs

Of the 3,096 contigs identified as host genes and annotated, 540 genes were differentially expressed, with 338 genes with significantly higher expression levels in exposed blades and 202 genes with lower expression levels (Supplementary Table 1). The WGCNA identified thirteen module eigengenes comprised of genes with similar expression patterns for 3,096 *Z. marina* contigs (Figure 5A; Supplementary Table 2). Of these eigengenes, four - named “brown”, “yellow”, “blue”, and “magenta” - were significantly



**FIGURE 3**  
Prevalence of observed disease signs on eelgrass blades 2 and 5 days post-inoculation. Error bars indicate  $\pm 1$  standard error of the replicate mean. Letters and bars indicate significantly different treatments.




FIGURE 4

Effects of temperature and *Lz* treatment on condensed tannins and total phenolic compounds (as a percentage of eelgrass dry mass) 2 and 5 days post-inoculation. Data are means ( $\pm 1$  SE). Letters and bars indicate significantly different treatments.

associated with *Lz* exposure (Figure 5B). These modules, labeled as color categories, are described in detail below.

The module eigengene “brown” was negatively correlated with *Lz* exposure, indicating lower module expression when eelgrass was exposed to *Lz*. ( $r^2 = -0.86$ ,  $p = 3.3 \times 10^{-5}$ ) (Figure 5B) The module contained 194 differentially expressed genes (Figure 5C; Supplementary Table 2). Seven biological processes were significantly enriched, including response to fungus (GO:0009620), response to salt stress (GO:0009651), transposition (GO:0032197;GO:0032196), and cell wall organization or biogenesis (GO:0071555;GO:0045229;GO:0071554) (Table 1). Genes were also involved in responses to abscisic acid, jasmonic acid, and wounding (Supplementary Table 2). Additionally, 17 molecular functions were enriched in this module, including transferase activity, transferring glycosyl groups (GO:0016757), xenobiotic transmembrane transporter activity (GO:0042910), and transcription regulator activity (GO:0003700;GO:0140110) (Table 1).

The module eigengene “yellow” was positively correlated with *Lz* exposure ( $r^2 = 0.52$ ,  $p = 0.049$ ) and contained five differentially expressed genes (Figures 5B, C; Supplementary Table 2). While there were no significantly enriched GO terms in this module, common biological processes included proteolysis, cellular response

to oxidative stress, response to osmotic stress, regulation of jasmonic acid metabolic process, and response to abscisic acid, and common molecular functions were binding of ATP, RNA, and metal ions (Supplementary Table 2).

The module eigengene “blue” was positively correlated with *Lz* exposure ( $r^2 = 0.99$ ,  $p = 2.0 \times 10^{-11}$ ) (Figure 5B). The “blue” module contained 263 differentially expressed genes (Figure 5C; Supplementary Table 2). There were no enriched biological processes, but genes were involved in immune response, protein transport and ubiquitination, response to salt stress and oxidative stress, cell cycle, and abscisic acid-activated signaling pathway (Supplementary Table 2). Enriched molecular functions were double-stranded DNA binding (GO:0003690) and cyclic nucleotide-dependent protein kinase activity (GO:0004691; GO:0004690) (Table 1).

The module eigengene “magenta” was positively correlated with *Lz* exposure ( $r^2 = 0.52$ ,  $p = 0.046$ ) (Figure 5B). This module eigengene had the most variable expression of all significant modules (Figure 5C). The “magenta” module contained 60 differentially expressed genes (Figure 5C; Supplementary Table 2). Peptide biosynthetic process (GO:0006412;GO:0043043) and determination of adult lifespan (GO:0008340), and structural constituent of




FIGURE 5

Results of *Z. marina* WCGNA analysis. (A) Cluster dendrogram of 3,096 genes used in WCGNA analysis. Modules were identified (Dynamic Tree Cut), then merged based on a cut height of 0.34 (Merged dynamic). All modules contain at least 30 genes. (B) Significant correlations between module eigengenes (rows) and *Lz* infection status (column). The  $R^2$  value is provided for each significant correlation, with  $P$ -value in parentheses. Positive correlations are represented by red boxes, while negative correlations are represented by blue boxes, with the intensity of the color increasing with correlation strength. (C) Expression of module eigengenes significantly correlated *Lz* exposure for exposed and control eelgrass.

TABLE 1 Biological process and molecular function GOterms enriched within significant module eigengenes for *Z. marina*.

Module	GO ID	GOterm Name	GO Category	FDR	Number of Sequences
brown	GO:0006970	response to osmotic stress	Biological Process	0.07	74
brown	GO:0009620	response to fungus	Biological Process	0	17
brown	GO:0009651	response to salt stress	Biological Process	0	53
brown	GO:0015074	DNA integration	Biological Process	0.0916667	24
brown	GO:0032197; GO:0032196	transposition	Biological Process	0.0857143	11
brown	GO:0042538	hyperosmotic salinity response	Biological Process	0.0625	8
brown	GO:0071555; GO:0045229; GO:0071554	cell wall organization or biogenesis	Biological Process	0	35
brown	GO:0003677	DNA binding	Molecular Function	0.04	190
brown	GO:0004519; GO:0004518	nuclease activity	Molecular Function	0	52
brown	GO:0008170; GO:0008276	N-methyltransferase activity	Molecular Function	0.0766667	23
brown	GO:0008270	zinc ion binding	Molecular Function	0.0636364	110
brown	GO:0008757	S-adenosylmethionine-dependent methyltransferase activity	Molecular Function	0.0833333	36

(Continued)

TABLE 1 Continued

Module	GO ID	GOterm Name	GO Category	FDR	Number of Sequences
brown	GO:0016279; GO:0016278	protein-lysine N-methyltransferase activity	Molecular Function	0.0714286	15
brown	GO:0016757	transferase activity transferring glycosyl groups	Molecular Function	0	62
brown	GO:0016763	transferase activity transferring pentosyl groups	Molecular Function	0.0941176	12
brown	GO:0016779	nucleotidyltransferase activity	Molecular Function	0.090625	73
brown	GO:0016788	hydrolase activity acting on ester bonds	Molecular Function	0	139
brown	GO:0016891; GO:0004521; GO:0016893	endonuclease activity active with either ribo- or deoxyribonucleic acids and producing 5'-phosphomonoesters	Molecular Function	0	17
brown	GO:0018024	histone-lysine N-methyltransferase activity	Molecular Function	0.0125	11
brown	GO:0034061; GO:0003964	DNA polymerase activity	Molecular Function	0	34
brown	GO:0042054	histone methyltransferase activity	Molecular Function	0.0333333	16
brown	GO:0042910	xenobiotic transmembrane transporter activity	Molecular Function	0.0769231	13
brown	GO:0140097	catalytic activity acting on DNA	Molecular Function	0	56
brown	GO:0140110; GO:0003700	transcription regulator activity	Molecular Function	0	58
blue	GO:0003690	double-stranded DNA binding	Molecular Function	0.025	32
blue	GO:0004691; GO:0004690	cyclic nucleotide-dependent protein kinase activity	Molecular Function	0	9
magenta	GO:0006412; GO:0043043	peptide biosynthetic process	Biological Process	0	259
magenta	GO:0008340	determination of adult lifespan	Biological Process	0	10
magenta	GO:0003735	structural constituent of ribosome	Molecular Function	0	248
magenta	GO:0019843	rRNA binding	Molecular Function	0.05	64

ribosome (GO:0003735) and rRNA binding (GO:0019843) were significantly enriched biological process and molecular function terms in this module, respectively (Table 1). Other prominent biological processes included translation, cell redox homeostasis, and microtubule-based process (Supplementary Table 2).

### Differential expression analysis - non-host contigs

Of the non-host contigs, 11.1% (87,550 out of 784,204) had significant matches to the Swiss-Prot database. Only 0.77% (5,959 out of 784,204) of non-host contigs had significant matches to the

*Labyrinthula* sp. Ha genome. However, 42,601 non-host contigs (5.4% of total non-host contigs) had significant matches to a closely related thraustochytrid, *Hondaea fermentalgiana* (NCBI Accession # GCA\_002897355.1). Thirty-two non-host contigs were differentially expressed; 30 up-regulated whereas two were down-regulated (Supplementary Table 3). Of the 32 differentially expressed non-host contigs, only 5 had significant similarity to the *Labyrinthula* sp. Ha genome; including arylsulfatase (SP\_P08842), patatin-like phospholipase domain-containing protein (SP\_Q8N8W4), 5-aminolevulinate synthase (SP\_P08262), dynein heavy chain 10 (*dhcA*) (SP\_Q8IVF4), and calponin-1 (SP\_P4318). In contrast, 20 differentially expressed non-host contigs had significant hits to the *H. fermentalgiana* genome

(Supplementary Table 3). Seven non-host contigs also had homology to genes found in the well-annotated genome of the slime-mold model organism, *D. discoideum*, including the dynein heavy chain *dhcA* (DDB\_G0276355), dual specificity protein kinase *shkB* (DDB\_G0288617), adenylate cyclase *sgcA* (DDB\_G0276269), calponin homology containing protein *mp20* (DDB\_G0292664), serine/threonine-protein kinase *roco5* (DDB\_G0294533), beta-ketoadipyl-CoA thiolase (DDB\_G0269588), and regulator of G-protein signaling 21 (DDB\_G0273033) (Supplementary Table 3).

Of the differentially expressed genes that displayed significant similarity to either *Lz* or to related species *H. fermentalgiana* or *D. discoideum*, ten up-regulated non-host contigs were linked to chemotaxis and motility (i.e. actin binding proteins, tubulin beta-chain, and microtubule formation), 11 contigs to amino acid and lipid metabolism (i.e. peptidases and fatty acid synthesis enzymes), 4 contigs to transcription/translation, 3 contigs to transporter activity (i.e. ion channel and transporter subunits), and 4 contigs to other processes including respiration (i.e. cytochrome c oxidase subunits) and production of secondary metabolites (i.e. phytoene synthase) (Supplementary Table 3). Several non-host contigs were related to phagocytosis, such as patatin-like phospholipase domain containing protein (SP\_Q8N8W4), beta-ketoadipyl-CoA thiolase (SP\_O26884), leucine-rich repeat serine-threonine protein kinase 2 (*roco5*) (SP\_Q1ZXD6), ATP-dependant RNA helicase (SP\_Q0DB53), hypoxanthine-guanine phosphoribosyltransferase (SP\_Q72454), pepsin A-1 (SP\_Q03168), and dynein heavy chain 10 (*dhcA*) (SP\_Q8IVF4).

## Discussion

Exposure of eelgrass ramets to *Lz* resulted in rapid progression of EWD, with necrotic lesions forming within two days post-exposure and plant death occurring after five days. The temperature did not alter the pathogen load or progression of EWD in the *Lz* exposed plants. This may be due to the high concentration of pathogen in the exposure, which was higher than in previous studies with this *Lz* strain (Groner et al., 2014; Groner et al., 2018) and/or pre-acclimation of the plants to the temperature treatments prior to exposure. Analysis of the eelgrass transcriptome revealed changes in gene expression, with patterns consistent with increased defensive responses through altered regulation of genes associated with phytohormone biosynthesis, stress response, and immune function. Analysis of non-*Zostera* genes (likely *Lz* genes), revealed expression patterns suggestive of *Lz* disrupting host immune responses and undergoing phagocytosis. This study reveals the importance of specific pathways related to plant defense against pathogen presence, and provides molecular evidence to support previous phenotypic observations of host-pathogen interactions of eelgrass wasting disease.

## Host responses

The genome of *Z. marina* allowed us to identify 3,096 host contigs, and we detected 540 differentially expressed host genes responding to *Lz* infection as opposed to increased temperature

(Supplementary Table 1). We found expression changes in genes and/or pathways involved with immunity such as pathogen detection, defense-related cell-cell signaling, defense-related metabolite production, and apoptosis. We also detected genes potentially involved in salt stress. The occurrence of many differentially expressed defense-response related GO terms, including genes associated with microbial detection and defense response to bacterial or fungal pathogens, indicates that *Z. marina* is responding to *Lz* infection.

Plant innate immune system pathways are highly conserved and are initiated by recognition of pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs) by membrane localized receptors (Choi and Klessig, 2016). Calcium signaling is a major pathway in pattern-triggered immunity, where binding of PAMPs and DAMPs to membrane receptors initiates the rapid release of calcium ions ( $Ca^{2+}$ ) by receptor like kinases and their  $Ca^{2+}$  dependent protein kinases and mitogen activated protein kinases (Hou et al., 2018). For example, calcium/calmodulin-dependent protein kinase (CcaMK) genes have been found to play a role in maintaining endosymbionts and in pathogen resistance in tomato, by promoting accumulation of hydrogen peroxide (Wang et al., 2015). Genes related to calcium- and protein-kinases show up eight and 43 times, respectively, in the blue module. In particular, upregulation of the calcium/calmodulin-dependent protein kinase (CcaMK) type 1 (CaM kinase I) suggests the role of *Lz*-related PAMP/DAMP detection and may warrant further exploration.

Pathogen detection leads to the induction of signaling pathways to induce immune and defense responses. We found that the cAMP-dependent protein kinase catalytic subunit gamma is significantly upregulated with *Lz*-exposure. Changes in the abundance of cAMPs can affect the activity of other signaling pathways and cellular processes, such as protein kinases and transcription factors (Świeżawska et al., 2018), demonstrating that *Z. marina* may respond to *Lz* infection by altering upstream signal transduction.

Phytohormone production in plants is important for defense-related signaling and is turned on after PAMP/DAMP detection. For example, compounds such as jasmonic acid (JA), salicylic acid (SA), and abscisic acid (ABA) are involved in general activation of plant defense systems against pathogens (Thaler et al., 2004; Tamaoki et al., 2013). In particular, SA is used in defense against biotic factors such as insects, mites, fungi, and bacteria and JA is used in defense pathways against saprophytic microbes like *Lz* (Tamaoki et al., 2013; Zhang et al., 2020). ABA is triggered in response to both abiotic and biotic stressors, and has complex interactions with both JA and SA responses to infection (Fan et al., 2009). We found GO terms for SA, JA, and ABA phytohormones [GO:0080140, GO:0009695, GO:0009753, and GO:0009751] were highly represented in the blades experimentally exposed to *Lz*, although these gene ontologies were not significantly enriched. Gene expression patterns in this study were indicative of phytohormone signaling via enrichment of serine/threonine-protein kinase PCRK2 gene which is involved in the resistance to bacterial pathogens and salicylate biosynthesis during pathogenic infection (Sreekanta et al., 2015). Methylesterase 1 gene, which is

required during the conversion of methyl salicylate into SA was downregulated in exposed blades. These data suggest that *Z. marina* is altering production of these defense-related compounds. We found gene expression changes in two genes related to JA biosynthesis: chloroplastic rhomboid-like protein 11 (regulation of jasmonic acid metabolic process [GO:0080140]) and phospholipase A (defense response to fungus [GO:0050832]; jasmonic acid biosynthetic process [GO:0009695]). Phospholipase A is responsible for the release of linolenic acid from the chloroplast membrane during defensive signaling, which induces JA production (Yang et al., 2007; Mata-Pérez et al., 2015). The detection of differentially regulated genes in the JA pathway is particularly interesting, as the role of JA in seagrasses is unclear. A congener of *Z. marina*, *Z. muelleri*, lost genes encoding jasmonate methyltransferase, which converts JA into a volatile compound, methyl jasmonate, but maintain other genes along the JA pathway, including for jasmonate synthesis, and signaling (Lee et al., 2016). Although the JA-associated genes were not differentially expressed in our study, changes to expression in these genes suggests a role for the jasmonic acid pathway in responding to *Lz* infection.

Gene expression changes also indicated down-stream effects of phytohormone production. In maize roots, expression of CHC1 gene increases in response to SA or ABA, suggesting involvement of CHC1 in the SA signaling pathway in maize defense responses (Zeng et al., 2013). Upregulation of the probable clathrin heavy chain 1 gene (CHC1) ([GO:0005198]) in *Lz*-infected blades suggests a similar response in eelgrass. We detected gene expression changes in factors involved in stress signaling which either require ABA or interact with ABA to trigger defense responses. For example, the S-type anion channel SLAH3 is an anion channel that can be triggered by ABA during a stress response (Roelfsema et al., 2012), and was downregulated in response to *Lz* infection. The B3 domain-containing protein was also downregulated. This protein is a transcription factor for ABA and inhibits the production of ABA (Brady et al., 2003), so a decrease in expression of B3 could result in an increase in ABA production, indicating a greater stress response to *Lz* infection. One specific pathway in which ABA affects plant defense is in the callose pathway, which is involved in defense of fungal pathogens in plants. Specifically, ABA mutants displayed impaired resistance to necrotrophic fungi, and ABA addition to the infection site mimicked callose deposition and increased resistance to necrotrophic fungi (Mauch-Mani and Mauch, 2005). Therefore, it is not surprising to observe altered ABA expression in *Z. marina* infected with *Lz*, which has necrotic capabilities.

Additional immune responses in plants include the production of antimicrobial compounds and apoptosis of infected cells, the latter of which would be important for intracellular infections, such as with *Lz*. Upregulation of peroxisomal targeting signal 1 receptor indicates potential defenses at sites of pathogen entry. This gene is associated with concentrating antifungal glucosinolate derivatives in the peroxisome for eventual transport to sites of fungal infection entry (Bednarek et al., 2009). It would be valuable to assess whether these compounds can inhibit *Lz* as well. Differential expression of GDP-L-fucose synthase supports the role of apoptosis as an immune response to *Lz*. GDP-L-fucose plays a role in stomatal and apoptosis-related defense in *Arabidopsis*, indicating a similar

role in *Z. marina* immunity (Zhang et al., 2019). Differential expression of EKC/KEOPS subunit genes, which are involved in apoptotic processes via telomere capping and elongation (Downey et al., 2006), also support the hypothesis that apoptosis is an important defense against *Lz*.

The production of phenolic compounds is another major immune response of plants. Many of phenolic metabolites detected in *Z. marina* have been implicated in *Z. marina* defense. These include flavonoids in sulfated and unsulfated form, as well as acids like caffeic, *p*-coumaric, ferulic, and zosteric, and rosmarinic acids, all of which are hydroxy, sulfoxyl, or ester forms of cinnamic acid (Papazian et al., 2019). Resistance of *Z. marina* to *Lz* is hypothesized to depend upon phenolic acids, especially caffeic acid which has demonstrated activity against *Lz* in bioassays (Vergeer and Develi, 1997; Trevathan-Tackett et al., 2015). *Lz* has also been shown to induce production of total phenols in some cases (McKone and Tanner, 2009), and inhibit them in others, potentially as a mechanism to circumvent this defense. In addition, production of phenols in seagrasses are reduced at warmer temperatures (Vergeer et al., 1995). This is reflected in the reduced concentrations of phenols found five days post *Lz*-exposure in the warmer temperature treatment.

The shikimate pathway is the mechanism through which plants produce aromatic phenolic compounds in response to stress (Santos-Sánchez et al., 2019). Four genes - caffeic acid 3-O-methyltransferase, caffeoylshikimate esterase, 4-coumarate-CoA ligase-like 7, and cinnamoyl-CoA reductase - responsible for coding enzymes involved in the biosynthesis of different forms of cinnamic acids were differentially expressed in the exposed treatment, indicating a role for the shikimate pathway in *Lz* infection response. Caffeoylshikimate esterase (CSE), converts caffeoyl shikimate into caffeic acid, which has been shown to inhibit *Lz* growth *in vitro* (Vergeer and Develi, 1997; Vanholme et al., 2013). Another differentially expressed gene, caffeic acid 3-O-methyltransferase, is associated with converting caffeic acid into products associated with the lipid biosynthesis pathway. 4-coumarate-CoA is also associated with production of additional secondary compounds including isoflavonoids and furanocoumarins as phytoalexins (i.e., inhibitors of pathogen growth) (Hamberger and Hahlbrock, 2004). Cinnamoyl-CoA reductase is involved in the formation of phenolic compounds associated with the hypersensitive response (Lauvergeat et al., 2001). The hypersensitive response in plants is akin to the innate immune response in animals and is responsible for isolated apoptosis surrounding an infection in order to prevent further spread (Morel and Dangl, 1997).

Apart from individual roles in the shikimate pathway, all four differentially expressed genes in this pathway are also associated with lignin biosynthesis. The role of lignin is unclear in aquatic plants such as *Z. marina*. While it may be protective against microbial attack, it is typically concentrated in the slower growing rhizomes, and not in the blades (Klap et al., 2000), where gene expression was measured in this study.

In addition to genes involved in immune activation, hormone pathways, and phenolic production, the transcriptional patterns of *Lz* exposed plants and gene enrichment results suggest responses to environmental stress, including salt stress. Stress-related genes were significantly upregulated in the exposed treatment, indicating that

*Lz* infection may cause damage to cells in a way that triggers this response. The mitochondrial phosphate carrier protein 3 (MPT3) is induced under high salinity conditions (Zhu et al., 2012). Increased expression of the MPT3 gene suggests that *Z. marina* may be experiencing salt stress as *Lz* breaks through host physical barriers. Similarly, the sorting nexin 1 protein regulates the process of accumulating nitrous oxide, which is an important signaling molecule in responses to abiotic stressors, including salt stress (Li et al., 2018). Higher sorting nexin 1 protein gene expression implies that *Z. marina* may accumulate more nitrous oxide in an effort to respond to abiotic stress. Finally, the upregulation of the calcium/calmodulin-dependent protein kinase type 1 (CaM kinase I) gene suggests changing cytosol calcium concentrations, which is a known plant response to stressors such as anoxia, increases temperature, and increased salinity (White and Broadley, 2003). Changes in expression of genes associated with stress response in *Z. marina* may inform our understanding of how molecular functioning is altered upon exposure to environmental changes. This information can also shed light on how *Lz* infection cause stressful conditions for *Z. marina*.

## Pathogen responses

By separating host from non-host contigs, we aimed to identify genes or pathways that may contribute to the virulence of *L. zosterae* during experimental infection of *Z. marina*. We found up-regulation of genes potentially involved in breakdown of host defense, chemotaxis and phagocytosis, and metabolism. Although an unannotated genome of another species of *Labyrinthula* (*Labyrinthula* sp. Ha) is newly available, very few contigs had significant matches to the current version of the genome, and only five of the differentially expressed genes from the non-host dataset had significant BLAST hits to this genome. However, the majority of the non-host contigs had significant matches to related slime-mold species including *Dictyostelium* species and *Hondaea fermentalgiana* (a thraustochytrid closely related to the genus *Labyrinthula*), suggesting that the identified non-host genes were likely from *Lz* inoculated in the infection treatment.

*Zostera* species (including *Z. marina*) produce high concentrations of a sulfate ester, zosteric acid, as a defense mechanism that protects blades from microbial biofouling (Grignon-Dubois et al., 2012; Vilas-Boas et al., 2017). We found a 24-fold up-regulation of a non-host contig that matched to the arylsulfatase gene (GBG30795.1) of *H. fermentalgiana*. Arylsulfatases are enzymes that break down sulfatides and organic sulfate esters. In other protists, arylsulfatases are produced in response to sulfate deprivation (Niedermeyer et al., 1987). However, for the fungal plant pathogen, *Colletotrichum gloeosporioides*, arylsulfatase expression increases during penetration of host leaf tissue after experimental inoculation (Goodwin et al., 2000). In *Lz*, an increase in arylsulfatase expression may facilitate attachment by breaking down the anti-biofouling zosteric acid produced by the host. However, the potential role of arylsulfatases to facilitate protist pathogen attachment and colonization has yet to be investigated.

Initial studies describing EWD observed *Lz* directly penetrating mesophyll cell walls, damage suggestive of feeding on plant cell organelles such as chloroplasts, and moving rapidly through the host tissues (Muehlstein, 1992; Raghukumar, 2002). In support of these microscopic observations and supporting the role of plant consumption as a mechanism of pathogenesis, we identified several up-regulated genes that are likely involved in movement, i.e. chemotaxis and cytoskeleton rearrangement and organization, as well as phagocytosis. For example, contigs homologous to *shkB*, *sgcA* and *roco5* genes were all up-regulated after experimental inoculation of *Lz*. In *D. discoideum*, null mutants for *shkB* and *sgcA* have reduced chemotaxis and phagocytosis ability (Moniakis et al., 2001; Veltman and Van Haastert, 2006; Veltman et al., 2008), and null mutants for the *roco5* genes have slow or no motility (Sawai et al., 2007). Six up-regulated non-host contigs were homologous to genes that are up-regulated during phagocytosis and/or have gene products which are a part of the 'macropinocytosis proteome' of *D. discoideum* (Sillo et al., 2008; Journet et al., 2012). Additionally, in *D. discoideum*, *dhcA* proteins cluster to phagosomes and promote phagolysosome fusion (Rai et al., 2016), and lysosomal aspartic protease CatD, a ubiquitous hydrolytic enzyme with a protein of homology in *D. discoideum*, is known to be a lysosomal pepsin in the macropinosome and is involved with phagosome maturation (Vines and King, 2019).

Genes involved in amino acid and lipid metabolism were up-regulated among the non-host contigs. M42 peptidase, a co-catalytic metallopeptidase (cytosolic enzyme) described in bacteria and archaea (reviewed by Appolaire et al., 2016), was the most expressed non-host contig. M42 may play a role in pathogenesis in bacteria by using or modifying exogenous proteins including immunological antibacterial peptides. Other peptidases were up-regulated, such as Carboxypeptidase Y and an otubain-like ubiquitin thioesterase, and may be involved with the catabolism of both large and small peptides in *Lz*. Ethylmalony-CoA decarboxylase, polyketide synthase, beta-ketoacyl-CoA thiolase, and patatin-like phosphatase domain-containing protein are all involved with fatty acid synthesis and may support membrane formation and increase membrane fluidity during cell division or for the extension of ectoplasmic nets of *Lz*. Up-regulation of these metabolism related contigs along with the up-regulation of genes associated with transcription and translation support increased metabolism and/or cellular proliferation.

Surprisingly, no genes potentially linked to the production of zoospores were up-regulated in this study. Other mechanisms of microbial virulence, such as toxin production, iron sequestration, and host immune invasion (Finlay and Falkow, 1997), were also not up-regulated in this study. Together the non-host transcriptome suggests the possible enzymatic degradation of plant cell defenses, supports the rapid dissemination of *Lz* cells within host tissue, and supports the voracious consumption of plant tissue by invading *Lz* cells as mechanisms of virulence in seagrass wasting disease. Time course experiments, with lower exposure concentrations (*sensu* Bower et al., 1989) would be useful in elucidating both the genes and virulence mechanisms involved in penetration vs. pathogenesis during this disease process.

## Conclusion

With increased outbreaks of marine disease, continued influence of anthropogenic stressors, and rapidly changing oceans contributing to significant declines in seagrass meadows across the globe, there is an increasing need to understand host-pathogen dynamics across environmental scales. Given the rapid decline of *Z. marina* globally coupled with active restoration, we have increased need for understanding of host-pathogen dynamics and how this might impact future restoration. This study demonstrates the value of dual host-pathogen transcriptomics for understanding physiological consequences of disease across environmental gradients. While the *Lz* pathogen exhibited a limited repertoire of virulence approaches, including degradation of cell walls, consumption of intracellular materials and movement within the host, the eelgrass host exhibited a wide range of responses to infection, from cascading phytohormone signals, to altered production of phenols, increased PAMP/DAMP signaling and apoptosis. As our knowledge of *Lz* increases, biologically realistic inoculation experiments can be used to further understand the role of changing nearshore conditions in facilitating or repressing disease in this important coastal habitat-forming species.

## Data availability statement

The RNAseq data presented in the study are deposited in the NCBI Short Read Archive repository, accession number PRJNA990835. Additional data presented in this study are available at FigShare (<http://doi.org/10.6084/m9.figshare.22564528>).

## Author contributions

MG and CB obtained financial support, conceived of the experimental design, and implemented the experiment. MG led statistical analyses of phenolic and diagnostic data. YV, CB, and AS led the transcriptomics analysis, with help from all authors, and KA analyzed phenols and tannins in the samples. All authors contributed to the article and approved the submitted version.

## References

Świeżawska, B., Duszyn, M., Jaworski, K., and Szmidt-Jaworska, A. (2018). Downstream targets of cyclic nucleotides in plants. *Front. Plant Sci.* 9, 1428. doi: 10.3389/fpls.2018.01428

Agnew, M. V., Groner, M. L., Eisenlord, M. E., Friedman, C. S., and Burge, C. A. (2022). Pacific oysters are a sink and a potential source of the eelgrass pathogen, *Labyrinthula zosterae*. *Aquacult Environ. Interact.* 14, 295–307. doi: 10.3354/aei00446

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment search tool. *J. Mol. Biol.* 215, 403–410. doi: 10.1016/S0022-2836(05)80360-2

Aoki, L. R., McGlathery, K. J., Wiberg, P. L., and Al-Haj, A. (2020). Depth affects seagrass restoration success and resilience to marine heat wave disturbance. *Estuaries Coast.* 43, 316–328. doi: 10.1007/s12237-019-00685-0

Appolaire, A., Colombo, M., Basbous, H., Gabel, F., Girard, E., and Franzetti, B. (2016). TET peptidases: a family of tetrahedral complexes conserved in prokaryotes. *Biochimie* 122, 188–196. doi: 10.1016/j.biochi.2015.11.001

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *J. Statistical Software*. 67 (1), 1–48. doi: 10.18637/jss.v067.i01

Bate-Smith, E. C., and Rasper, V. (1969). Tannins of grain sorghum: luteoforol (leucoleutoolinidin) 3',4',5,7-pentahydroxyflavan. *J. Food Sci.* 34, 203–209. doi: 10.1111/j.1365-2621.1969.tb00919.x

Beca-Carretero, P., Guihéneuf, F., Marín-Guirao, L., Bernardeau-Esteller, J., García-Muñoz, R., Stengel, D. B., et al. (2018). Effects of an experimental heat wave on fatty acid composition in two Mediterranean seagrass species. *Mar. Pollut. Bull.* 134, 27–37. doi: 10.1016/j.marpolbul.2017.12.057

Bednarek, P., Piślewska-Bednarek, M., Svatoš, A., Schneider, B., Doubský, J., Mansurova, M., Schulze-lefer, A., et al. (2009). A glucosinolate metabolism pathway in living plant cells mediates broad-spectrum antifungal defense. *Science* 323, 101–106. doi: 10.1126/science.1163732

## Funding

This work was funded by a seed grant from the Canadian excellence research chair in support of aquatic epidemiology to MG, start-up funds provided to CB from the University of Maryland Baltimore County and the University of Maryland Baltimore, funding from the National Science Foundation (2109607; 1215977), as well as the University of Washington Friday Harbor Labs.

## Acknowledgments

Michelle Herko, Rebecca Guenther and Connie Sullivan assisted in the execution of this experiment. N. Rivlin provided technical assistance and T. Bachvaroff bioinformatics assistance.

## Conflict of interest

All authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmars.2023.1152647/full#supplementary-material>

Bockelmann, A. C., Tams, V., Ploog, J., Schubert, P. R., and Reusch, T. B. (2013). Quantitative PCR reveals strong spatial and temporal variation of the wasting disease pathogen, *Labyrinthula zosterae* in northern European eelgrass (*Zostera marina*) beds. *PLoS One* 8, e62169. doi: 10.1371/journal.pone.0062169

Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics* 30, 2114–2120. doi: 10.1093/bioinformatics/btu170

Bower, S. M., McLean, N., and Whitaker, D. J. (1989). Mechanism of infection by *Labyrinthuloides haliotidis* (Protozoa: labyrinthomorpha), a parasite of abalone (*Haliotis kamtschatkana*) (Mollusca: Gastropoda). *J. Invert Pathol.* 53, 401–409. doi: 10.1016/0022-2011(89)90106-7

Brady, S. M., Sarkar, S. F., Bonetta, D., and McCourt, P. (2003). The abscisic acid insensitive 3 (ABI3) gene is modulated by farnesylation and is involved in auxin signaling and lateral root development in *Arabidopsis*. *Plant J.* 34, 67–75. doi: 10.1046/j.1365-313X.2003.01707.x

Brakel, J., Jakobsson-Thor, S., Bockelmann, A. C., and Reusch, T. B. (2019). Modulation of the eelgrass–*Labyrinthula zosterae* interaction under predicted ocean warming, salinity change and light limitation. *Front. Mar. Sci.* 6, 268. doi: 10.3389/fmars.2019.00268

Bull, J. C., Kenyon, E. J., and Cook, K. J. (2012). Wasting disease regulates long-term population dynamics in a threatened seagrass. *Oecologia* 169, 135–142. doi: 10.1007/s00442-011-2187-6

Burge, C. A., and Friedman, C. S. (2012). Quantifying Ostreid herpesvirus (OsHV-1) copies and expression during transmission. *Microb. Ecol.* 63 (3), 596–604. doi: 10.1007/s00248-011-9937-1

Chen, Y., Lun, A. A. T., and Smyth, G. K. (2016). “From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline.” *F1000Research* 5, 1438. doi: 10.12688/f1000research.8987.2

Choi, H. W., and Klessig, D. F. (2016). DAMPs, MAMPs, and NAMPs in plant innate immunity. *BMC Plant Biol.* 16, 232. doi: 10.1186/s12870-016-0921-2

Dawkins, P. D., Eisenlord, M. E., Yoshioka, R. M., Fiorenza, E., Fruchter, S., Giannonna, F., et al. (2018). Environment, dosage, and pathogen isolate moderate virulence in eelgrass wasting disease. *Dis. Aquat. Org.* 130, 51–63. doi: 10.3354/dao03263

Downey, M., Houlsworth, R., Maringele, L., Rollie, A., Brehme, M., Galicia, S., et al. (2006). A genome-wide screen identifies the evolutionarily conserved KEOPS complex as a telomere regulator. *Cell* 124, 1155–1168. doi: 10.1016/j.cell.2005.12.04

Dunic, J. C., Brown, C. J., Connolly, R. M., Turschwell, M. P., and Cote, I. M. (2021). Long-term declines and recovery of meadow area across the world’s seagrass bioregions. *Global Change Biol.* 27, 4096–4109. doi: 10.1111/gcb.15684

Fan, J., Hill, L., Crooks, C., Doerner, P., and Lamb, C. (2009). Abscisic acid has a key role in modulating diverse plant-pathogen interactions. *Plant Physiol.* 150, 1750–1761. doi: 10.1104/pp.109.137943

Finlay, B. B., and Falkow, S. (1997). Common themes in microbial pathogenicity revisited. *Microbiol. Mol. Biol. Rev.* 61, 136–169. doi: 10.1128/mmbr.61.2.136-169.1997

Goodwin, P. H., Li, J., and Jin, S. (2000). Evidence for sulfate derepression of an arylsulfatase gene of *Colletotrichum gloeosporioides* f. sp. *malvae* during infection of round-leaved mallow, *Malva pusilla*. *Physiol. Mol. Plant Pathol.* 57, 169–176. doi: 10.1006/pmpp.2000.0295

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., et al. (2011). Full-length transcriptome assembly from RNA-seq data without a reference genome. *Nat. Biotechnol.* 29, 644. doi: 10.1038/nbt.1883

Graham, O. J., Aoki, L. R., Stephens, T., Stokes, J., Dayal, S., Rappazzo, B., et al. (2021). Effects of seagrass wasting disease on eelgrass growth and belowground sugar in natural meadows. *Front. Mar. Sci.* 8, 768668. doi: 10.3389/fmars.2021.768668

Grignon-Dubois, M., Rezzonico, B., and Alcoverro, T. (2012). Regional scale patterns in seagrass defences: phenolic acid content in *Zostera noltii*. *Estuar. Coast. Shelf Sci.* 114, 18–22. doi: 10.1016/j.ecss.2011.09.010

Groner, M. L., Burge, C. A., Couch, C. S., Kim, C. J., Siegmund, G. F., Singhal, S., et al. (2014). Host demography influences the prevalence and severity of eelgrass wasting disease. *Dis. aquat. org.* 108, 165–175. doi: 10.3354/dao02709

Groner, M. L., Burge, C. A., Cox, R., Rivlin, N. D., Turner, M., Van Alstyne, K. L., et al. (2018). Oysters and eelgrass: potential partners in a high pCO<sub>2</sub> ocean. *Ecology* 99, 1802–1814. doi: 10.1002/ecy.2393

Groner, M. L., Burge, C. A., Kim, C. J., Rees, E., Van Alstyne, K. L., Yang, S., et al. (2016). Plant characteristics associated with widespread variation in eelgrass wasting disease. *Dis. aquat. org.* 118, 159–168. doi: 10.3354/dao02962

Groner, M. L., Eisenlord, M. E., Yoshioka, R. M., Fiorenza, E. A., Dawkins, P. D., Graham, O. J., et al. (2021). Warming sea surface temperatures fuel summer epidemics of eelgrass wasting disease. *Mar. Ecol. Prog. Ser.* 679, 47–58. doi: 10.3354/meps13902

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., et al. (2013). *De novo* transcript sequence reconstruction from RNA-seq using the trinity platform for reference generation and analysis. *Nat. Protoc.* 8, 1494. doi: 10.1038/nprot.2013.084

Hamberger, B., and Hahlbrock, K. (2004). The 4-coumarate: CoA ligase gene family in *Arabidopsis thaliana* comprises one rare, sinapate-activating and three commonly occurring isoenzymes. *PNAS* 101, 2209–2214. doi: 10.1073/pnas.0307307101

Hou, S., Jamieson, P., and He, P. (2018). The cloak, dagger, and shield: proteases in plant-pathogen interactions. *Biochem. J.* 475, 2491–2509. doi: 10.1042/bcj20170781

Journet, A., Klein, G., Brugiére, S., Vandenbrouck, Y., Chapel, A., Kieffer, S., et al. (2012). Investigating the macropinocytic proteome of *Dictyostelium* amoebae by high-resolution mass spectrometry. *Proteomics* 12, 241–245. doi: 10.1002/pmic.201100313

Klap, V. A., Hemminga, M. A., and Boon, J. J. (2000). Retention of lignin in seagrasses: angiosperms that returned to the sea. *Mar. Ecol. Prog. Ser.* 194, 1–11. doi: 10.3354/meps194001

Lamb, J. B., Van De Water, J. A., Bourne, D. G., Altier, C., Hein, M. Y., Fiorenza, E. A., et al. (2017). Seagrass ecosystems reduce exposure to bacterial pathogens of humans, fishes, and invertebrates. *Science* 355, 731–733. doi: 10.1126/science.aa11956

Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation network analysis. *BMC Bioinf.* 9, 559. doi: 10.1186/1471-2105-9-559

Lauvergeat, V., Lacomme, C., Lacombe, E., Lasserre, E., Roby, D., and Grima-Pettenati, J. (2001). Two cinnamoyl-CoA reductase (CCR) genes from *Arabidopsis thaliana* are differentially expressed during development and in response to infection with pathogenic bacteria. *Phytochemistry* 57, 1187–1195. doi: 10.1016/S0031-9422(01)00053-X

Lee, H., Golicz, A. A., Bayer, P. E., Jiao, Y., Tang, H., Paterson, A. H., et al. (2016). The genome of a southern hemisphere seagrass species (*Zostera muelleri*). *Plant Physiol.* 172, 272–283. doi: 10.1104/pp.16.00868

Li, T. T., Liu, W. C., Wang, F. F., Ma, Q. B., Lu, Y. T., and Yuan, T. T. (2018). SORTING NEXIN 1 functions in plant salt stress tolerance through changes of NO accumulation by regulating NO synthase-like activity. *Front. Plant Sci.* 9, 1634. doi: 10.3389/fpls.2018.01634

Ma, D., Li, Y., Zhang, J., Wang, C., Qin, H., Ding, H., et al. (2016). Accumulation of phenolic compounds and expression profiles of phenolic acid biosynthesis-related genes in developing grains of white, purple, and red wheat. *Front. Plant Sci.* 7, 528. doi: 10.3389/fpls.2016.00528

Markham, J. W., and Hagmeier, E. (1982). Observations on the effects of germanium dioxide on the growth of macro-algae and diatoms. *Phycologia* 21, 125–130. doi: 10.2216/0031-8884-21-2-125

Martin, D. L., Chiari, Y., Boone, E., Sherman, T. D., Ross, C., Wyllie-Echeverria, S., et al. (2016). Functional, phylogenetic and host-geographic signatures of *Labyrinthula* spp. provide for putative species delimitation and a global-scale view of seagrass wasting disease. *Estuaries Coasts* 39, 1403–1421. doi: 10.1007/s12237-016-0087-z

Mata-Pérez, C., Sánchez-Calvo, B., Begara-Morales, J. C., Luque, F., Jiménez-Ruiz, J., Padilla, M. N., et al. (2015). Transcriptomic profiling of linolenic acid-responsive genes in ROS signaling from RNA-seq data in *Arabidopsis*. *Front. Plant Sci.* 6, 122. doi: 10.3389/fpls.2015.00122

Mauch-Mani, B., and Mauch, F. (2005). The role of abscisic acid in plant-pathogen interactions. *Curr. Opin. Plant Biol.* 8, 409–414. doi: 10.1016/j.pbi.2005.05.015

McCarthy, D. J., Chen, Y., and Smyth, G. K. (2012). Differential expression analysis of multifactor RNA-seq experiments with respect to biological variation. *Nucleic Acids Res.* 40, 4288–4297. doi: 10.1093/nar/gks042

McKone, K. L., and Tanner, C. E. (2009). Role of salinity in the susceptibility of eelgrass *Zostera marina* to the wasting disease pathogen *Labyrinthula zosterae*. *Mar. Ecol. Prog. Ser.* 377, 123–130. doi: 10.3354/meps07860

Moniakis, J., Funamoto, S., Fukuzawa, M., Meisenhelder, J., Araki, T., Abe, T., et al. (2001). An SH2-domain-containing kinase negatively regulates the phosphatidylinositol-3 kinase pathway. *Genes Dev.* 15, 687–698. doi: 10.1101/gad.871001

Morel, J. B., and Dangl, J. L. (1997). The hypersensitive response and the induction of cell death in plants. *Cell Death Differ* 4, 671–683. doi: 10.1038/sj.cdd.4400309

Muehlstein, L. K. (1992). The host-pathogen interaction in the wasting disease of eelgrass, *Zostera marina*. *Can. J. Bot.* 70, 2081–2088. doi: 10.1139/b92-258

Niedermeier, I., Biedlingmaier, S., and Schmidt, A. (1987). Derepression of arylsulfatase activity by sulfate starvation in *Chlorella fusca*. *Z. für Naturforschung C*, 42 (5), 530–536, 42. doi: 10.1515/znc-1987-0507

Nitao, J. K., Birr, B. A., Nair, M. G., Herms, D. A., and Mattson, W. J. (2001). Rapid quantification of proanthocyanidins (condensed tannins) with a continuous flow analyzer. *J. Agric. Food Chem.* 49, 2207–2214. doi: 10.1021/jf001183b

Nordlund, L. M., Koch, E. W., Barbier, E. B., and Creed, J. C. (2016). Seagrass ecosystem services and their variability across genera and geographical regions. *Plos One* 11 (10), e0163091. doi: 10.1371/journal.pone.0169942

O’Donnell, M. J., George, M. N., and Carrington, E. (2013). Mussel byssus attachment weakened by ocean acidification. *Nat. Clim. Change* 3, 587–590. doi: 10.1038/nclimate1846

Orth, R. J., Carruthers, T. J., Dennison, W. C., Duarte, C. M., Fourqurean, J. W., Heck, K. L., et al. (2006). A global crisis for seagrass ecosystems. *Bioscience* 56, 987–996. doi: 10.1641/0006-3568(2006)56[987:AGCSE]2.0.CO;2

Papazian, S., Parrot, D., Burýšková, B., Weinberger, F., and Tasdemir, D. (2019). Surface chemical defence of the eelgrass *Zostera marina* against microbial foulers. *Sci. Rep.* 9, 3323. doi: 10.1038/s41598-019-39212-3

Raghukumar, S. (2002). Ecology of the marine protists, the labyrinthulomycetes (Thraustochytrids and labyrinthulids). *Eur. J. Protistol.* 38, 127–145. doi: 10.1078/0932-4739-00832

Rai, A., Pathak, D., Thakur, S., Singh, S., Dubey, A. K., and Mallik, R. (2016). Dynein clusters into lipid microdomains on phagosomes to drive rapid transport toward lysosomes. *Cell* 164, 722–734. doi: 10.1016/j.cell.2015.12.054

Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2010). edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinform* 26, 139–140. doi: 10.1093/bioinformatics/btp616

Roelfsema, M. R. G., Hedrich, R., and Geiger, D. (2012). Anion channels: master switches of stress responses. *Trends Plant Sci.* 17, 221–229. doi: 10.1016/j.tplants.2012.01.009

Santos-Sánchez, N. F., Salas-Coronado, R., Hernández-Carlos, B., and Villanueva-Cañongo, C. (2019). Shikimic acid pathway in biosynthesis of phenolic compounds. *Plant Physiol. aspects phenolic compounds* 1, 1–15. doi: 10.5772/intechopen.83815

Sawai, S., Guan, X. J., Kuspa, A., and Cox, E. C. (2007). High-throughput analysis of spatio-temporal dynamics in *Dictyostelium*. *Genome Biol.* 8, 1–15. doi: 10.1186/gb-2007-8-7-r144

Short, F., Carruthers, T., Dennison, W., and Waycott, M. (2007). Global seagrass distribution and diversity: a bioregional model. *J. Exp. Mar. Biol. Ecol.* 350, 3–20. doi: 10.1016/j.jembe.2007.06.012

Short, F. T., Muehlstein, L. K., and Porter, D. (1987). Elggrass wasting disease: cause and recurrence of a marine epidemic. *Biol. Bull.* 173, 557–562. doi: 10.2307/1541701

Sillo, A., Bloomfield, G., Balest, A., Balbo, A., Pergolizzi, B., Peracino, B., et al. (2008). Genome-wide transcriptional changes induced by phagocytosis or growth on bacteria in *Dictyostelium*. *BMC Genomics* 9, 1–22. doi: 10.1186/1471-2164-9-291

Sreekanta, S., Bethke, G., Hatsugai, N., Tsuda, K., Thao, A., Wang, L., et al. (2015). The receptor-like cytoplasmic kinase PCRK 1 contributes to pattern-triggered immunity against *Pseudomonas syringae* in *Arabidopsis thaliana*. *New Phytol.* 207, 78–90. doi: 10.1111/nph.13345

Strydom, S., Murray, K., Wilson, S., Huntley, B., Rule, M., Heithaus, M., et al. (2020). Too hot to handle: unprecedented seagrass death driven by marine heatwave in a world heritage area. *Glob. Change Biol.* 26, 3525–3538. doi: 10.1111/gcb.15065

Sullivan, B. K., Sherman, T. D., Damare, V. S., Lilje, O., and Gleason, F. H. (2013). Potential roles of *Labyrinthula* spp. in global seagrass population declines. *Fungal Ecol.* 6, 328–338. doi: 10.1016/j.funeco.2013.06.004

Tamaoki, D., Seo, S., Yamada, S., Kano, A., Miyamoto, A., Shishido, H., et al. (2013). Jasmonic acid and salicylic acid activate a common defense system in rice. *Plant Signal Behav.* 8, e24260. doi: 10.4161/psb.24260

Thaler, J. S., Owen, B., and Higgins, V. J. (2004). The role of the jasmonate response in plant susceptibility to diverse pathogens with a range of lifestyles. *Plant Physiol.* 135, 530–538. doi: 10.1104/pp.104.041566

Tracy, A. M., Pielmeyer, M. L., Yoshioka, R. M., Heron, S. F., and Harvell, C. D. (2019). Increases and decreases in marine disease reports in an era of global change. *Proc. R. Soc. B* 286, 20191718. doi: 10.1098/rspb.2019.1718

Trevathan-Tackett, S. M., Lane, A. L., Bishop, N., and Ross, C. (2015). Metabolites derived from the tropical seagrass *Thalassia testudinum* are bioactive against pathogenic *Labyrinthula* sp. *Aquat. Bot.* 122, 1–8. doi: 10.1016/j.aquabot.2014.12.005

Vanholme, R., Cesarino, I., Rataj, K., Xiao, Y., Sundin, L., Goeminne, G., et al. (2013). Caffeoyl shikimate esterase (CSE) is an enzyme in the lignin biosynthetic pathway in *Arabidopsis*. *Science* 341, 1103–1106. doi: 10.1126/science.1241602

Veltman, D. M., Keizer-Gunnik, I., and Van Haastert, P. J. (2008). Four key signaling pathways mediating chemotaxis in *Dictyostelium discoideum*. *J. Cell Biol.* 180, 747–753. doi: 10.1083/jcb.200709180

Veltman, D. M., and Van Haastert, P. J. (2006). Guanylyl cyclase protein and cGMP product independently control front and back of chemotaxing *Dictyostelium* cells. *Mol. Biol. Cell* 17, 3921–3929. doi: 10.1091/mbc.e06-05-0381

Vergeer, L. H. T., Aarts, T. L., and De Groot, J. D. (1995). The ‘wasting disease’ and the effect of abiotic factors (light intensity, temperature, salinity) and infection with *Labyrinthula zosterae* on the phenolic content of *Zostera marina* shoots. *Aquat. Bot.* 52, 35–44. doi: 10.1016/0304-3770(95)00480-N

Vergeer, L. H., and Develi, A. (1997). Phenolic acids in healthy and infected leaves of *Zostera marina* and their growth-limiting properties towards *Labyrinthula zosterae*. *Aquat. Bot.* 58, 65–72. doi: 10.1016/S0304-3770(96)01115-1

Vilas-Boas, C., Sousa, E., Pinto, M., and Correia-da-Silva, M. (2017). An antifouling model from the sea: a review of 25 years of zosteric acid studies. *Biofouling* 33, 927–942. doi: 10.1080/08927014.2017.1391951

Vines, J. H., and King, J. S. (2019). The endocytic pathways of *Dictyostelium discoideum*. *Int. J. Dev. Biol.* 63, 461–471. doi: 10.1387/ijdb.190236jk

Wang, J. P., Munyampundu, J. P., Xu, Y. P., and Cai, X. Z. (2015). Phylogeny of plant calcium and calmodulin-dependent protein kinases (CCaMKs) and functional analyses of tomato CCaMK in disease resistance. *Front. Plant Sci.* 6, 1075. doi: 10.3389/fpls.2015.01075

White, P. J., and Broadley, M. R. (2003). Calcium in plants. *Ann. Bot.* 9, 487–511. doi: 10.1093/aob/mcg164

Wright, R. M., Aglyamova, G. V., Meyer, E., and Matz, M. V. (2015). Gene expression associated with white syndromes in a reef-building coral, *Acropora hyacinthus*. *BMC Genomics* 16, 371. doi: 10.1186/s12864-015-1540-2

Yang, W., Devaiah, S. P., Pan, X., Isaac, G., Welti, R., and Wang, X. (2007). AtPLAI is an acyl hydrolase involved in basal jasmonic acid production and *Arabidopsis* resistance to *Botrytis cinerea*. *J. Biol. Chem.* 282, 18116–18128. doi: 10.1074/jbc.M700405200

Zeng, M. H., Liu, S. H., Yang, M. X., Zhang, Y. J., Liang, J. Y., Wan, X. R., et al. (2013). Characterization of a gene encoding clathrin heavy chain in maize up-regulated by salicylic acid, abscisic acid and high boron supply. *Int. J. Mol. Sci.* 14, 15179–15198. doi: 10.3390/ijms140715179

Zhang, L., Paasch, B. C., Chen, J., Day, B., and He, S. Y. (2019). An important role of L-fucose biosynthesis and protein fucosylation genes in *Arabidopsis* immunity. *New Phytol.* 222, 981–994. doi: 10.1111/nph.15639

Zhang, N., Zhou, S., Yang, D., and Fan, Z. (2020). Revealing shared and distinct genes responding to JA and SA signaling in *arabidopsis* by meta-analysis. *Front. Plant Sci.* 11, 908. doi: 10.3389/fpls.2020.00908

Zhu, W., Miao, Q., Sun, D., Yang, G., Wu, C., Huang, J., et al. (2012). The mitochondrial phosphate transporters modulate plant responses to salt stress via affecting ATP and gibberellin metabolism in *Arabidopsis thaliana*. *PLoS One* 7, e43530. doi: 10.1371/journal.pone.0043530