

Vibrational Spectroscopy and Dissociation Dynamics of Cyclohexyl Hydroperoxide

Tarun Kumar Roy^a, Yujie Qian^a, Elizabeth Karlsson^a, Rawan Rabayah^a, Christopher A. Sojda^a, Marisa C. Kozlowski^a, Tolga N. V. Karsili^b, and Marsha I. Lester^{a,*}

^a Department of Chemistry, University of Pennsylvania,
Philadelphia, PA 19104-6323 USA

^b Department of Chemistry, University of Louisiana,
Lafayette, LA USA

Abstract

Organic hydroperoxide (ROOH) are ubiquitous in the atmospheric oxidation of volatile organic compounds (VOCs) as well as in low-temperature oxidation of hydrocarbon fuels. The present work focuses on a prototypical cyclic hydroperoxide, cyclohexyl hydroperoxide (CHHP). The overtone OH stretch ($2\nu_{\text{OH}}$) spectrum of jet-cooled CHHP is recorded by IR multiphoton excitation with UV laser-induced fluorescence detection of the resulting OH products. A distinctive IR feature is observed at 7012.5 cm^{-1} . Two conformers of CHHP are predicted to have similar stabilities (within 0.2 kcal mol^{-1}) and overtone OH stretch transitions ($2\nu_{\text{OH}}$), yet are separated by a significant interconversion barrier. The IR power dependence indicates that absorption of three or more IR photons is required for dissociation of CHHP to cyclohexoxy (RO) and OH radical products. Accompanying high-level single- and multi-reference electronic structure calculations quantitatively support the experimental results. Calculations are extended to a range of organic hydroperoxides to examine trends in bond dissociation energies associated with $\text{RO} + \text{OH}$ formation and compared with prior theoretical results.

* Corresponding author email milester@sas.upenn.edu

1. Introduction

Organic hydroperoxides (ROOH) are formed in the oxidation of volatile organic compounds (VOCs) in the atmosphere or during combustion.¹⁻² The primary VOC generally reacts with hydroxyl (OH) radicals to form an alkyl radical that promptly reacts with O₂ to generate an alkylperoxy (RO₂) radical.³⁻⁴ The fate of the nascent RO₂ is heavily dependent on its molecular structure as well as the temperature and pressure of the surroundings.⁵⁻⁶

In the atmosphere, the dominant removal of nascent RO₂ radicals in pristine environments is reaction with HO₂, which forms lower volatility ROOH compounds.⁷⁻¹⁰ Alternatively, the RO₂ radical may undergo hydrogen atom transfer (HAT) to form a carbon-centered hydroperoxyalkyl radical (•QOOH),^{5-6, 11-13} which generally undergoes unimolecular decay or reacts with O₂ to form •OOQOOH. Sequential repetition of HAT and O₂ addition, termed autoxidation, may eventually form a highly oxidized ROOH compound, which has lower volatility than the precursor VOC.¹⁴⁻¹⁹ Such compounds are implicated in the formation of secondary organic aerosols, which affect radiative forcing in the atmosphere and impact public health.²⁰⁻²² In contrast, reaction with HO₂ represents a minor sink for RO₂ radicals formed during low temperature combustion (such as in modern-day vehicle engines).²³⁻²⁶ Under these conditions, the higher temperatures and pressures usually promote intramolecular HAT to form •QOOH radicals which play a vital role in fuel autoignition.^{14, 23-36}

Given their significance in atmospheric and combustion chemistry, a thorough understanding of the spectroscopy and dissociation dynamics of ROOH compounds is needed. Prior studies have focused on the vibrationally mediated photodissociation dynamics of hydrogen peroxide and simple organic peroxides.³⁷⁻⁴¹ The fundamental OH stretch and various overtone OH stretch transitions of H₂O₂ and its partially deuterated analogue HOOD have been characterized with a variety of techniques.⁴² Excitation at 6ν_{OH} provides sufficient energy to promote O-O bond dissociation to form OH X²Π fragments. These studies reveal that ca. 11% of the resulting OH fragments are formed vibrationally excited. This is in stark contrast to near UV photodissociation studies of H₂O₂ and organic hydroperoxides, which predominantly yield rovibrationally cold OH products.⁴³⁻⁴⁸

Studies have been extended to *tert*-butyl hydroperoxide (TBHP), where a sequence of OH-stretch transitions from the fundamental through the fifth overtone have been characterized.⁴⁰⁻⁴¹ Recent work revealed a strong feature at ~7017 cm⁻¹ arising from the first overtone OH stretch (2ν_{OH}), which is heavily coupled to COOH torsion.⁴⁹⁻⁵⁰ The ν_{OH} = 0 and ν_{OH} = 2 levels undergo tunneling splitting due to a double-minimum potential energy well along the COOH torsional coordinate. OH radical products are detected following ν_{OH} = 2 excitation of TBHP, arising from the absorption of three or more photons of the same wavelength to promote dissociation, while ν_{OH} = 6 excitation results in single photon dissociation. The prevailing mechanism for dissociation involves rapid intramolecular vibrational energy redistribution


(IVR) followed by O-O bond fission to form OH + RO radical products. The timescale for dissociation is controlled by unimolecular decay along a barrierless potential. Analogous vibrationally activated dissociation is expected for other ROOH compounds with greater molecular complexity. Bach and Schlegel have undertaken a systematic computational study of various organic hydroperoxides, revealing bond dissociation energies on the order of 45 kcal mol⁻¹, which are consistent with experimental values for a range of hydroperoxides.⁵¹⁻⁵²

This study characterizes the first overtone OH stretch (2ν_{OH}) of cyclohexyl hydroperoxide (cyc-C₆H₁₁O-OH, CHHP) and its multiphoton dissociation dynamics to OH products. A new synthetic approach is introduced to safely generate CHHP from cyclohexene. The present work builds on a recently reported high-resolution rotational spectroscopy study of the equatorial and axial conformers of CHHP in a Ne expansion (CHHP 1 and 2, respectively), which determined their structures and mapped out the conformational landscape involving torsional motions of the -OOH group.⁵³ Another recent study focused on the synthesis of a wide range of organic hydroperoxides, including CHHP, and determined their absolute photoionization cross sections and resultant fragmentation (-OOH loss) channels.⁵⁴ The present experimental study is complemented by high-level single- and multi-reference electronic structure calculations for CHHP and a range of other hydroperoxides.

2. Methods

2.1 Synthesis

A new synthetic approach was chosen for preparing cyclohexyl hydroperoxide (CHHP) to avoid the storage of large quantities of CHHP due to the explosive nature of organic hydroperoxides. It was hypothesized that protecting the peroxide with an appropriate silane group would increase the molecular mass and greatly decrease the risk of unexpected detonation of material.⁵⁵ Specifically, as shown in Scheme 1, cobalt-catalyzed silylperoxidation of cyclohexene directly yielded the triethylsilyl protected peroxide, (cyclohexylperoxy) triethylsilane, in excellent yields on a multigram scale.⁵⁶ From (cyclohexylperoxy) triethylsilane, conventional deprotection with pyridine•HF yielded very pure cyclohexyl hydroperoxide in poor yields. However, the poor yields of this transformation were vastly offset by rapid reaction times and more importantly the avoidance of storing bulk quantities of organic hydroperoxide, thus improving the safety of this approach. Extensive details on the synthetic preparation and analytical characterization of CHHP are presented in Electronic Supplementary Information (ESI, see Section S1 and Figures S1-S4).



Scheme 1. Synthetic route to generate cyclohexyl hydroperoxide (CHHP).

2.2 Spectroscopic Methods

The experimental method is similar to that used in a recent IR spectroscopic study of TBHP.⁴⁹ Freshly synthesized CHHP is seeded in Ar carrier gas at a pressure of 25 psig. The gas mixture is pulsed through a solenoid valve and an affixed quartz capillary tube (~25 mm length and 1 mm ID) into a vacuum chamber. The resultant supersonic expansion cools CHHP to a rotational temperature of ca. 10 K. Counter propagating IR and UV beams are spatially overlapped and intersect the gas mixture in the collision-free region ca. 1 cm downstream from the exit of the capillary. Multiphoton IR excitation of jet-cooled CHHP in the overtone OH stretch region ($2\nu_{\text{OH}}$) results in dissociation to OH $X^2\Pi_{3/2}$ ($v=0$) radical products that are detected. The tunable IR beam (with powers up to ca. 28 mJ/pulse) is the signal output of an optical parametric oscillator/amplifier (OPO/OPA, LaserVision; 0.9 cm^{-1} bandwidth) pumped by an unseeded Nd:YAG laser (1064 nm, Continuum Surelite EX, 5 Hz). The OH radical products are probed via laser-induced fluorescence (LIF) on the $A^2\Sigma^+ - X^2\Pi_{3/2}$ (1,0) $Q_1(3.5)$ transition at 282 nm utilizing UV radiation (ca. 1 mJ/pulse) generated with the frequency-doubled output of a Nd:YAG (EKSPLA NL300; 532 nm, 10 Hz) pumped dye laser (Continuum ND6000, Rhodamine 590 dye).

The fluorescence emitted on the OH $A^2\Sigma^+ - X^2\Pi_{3/2}$ (1,1) transition is detected using a gated photomultiplier tube (Electron Tubes 9813QB) and preamplified. Data acquisition involves an active background subtraction scheme (IR on – IR off) to remove a weak OH LIF background signal that arises from UV photodissociation of CHHP at 282 nm.

2.3 Computational Methods

The ground state minimum energy geometry of a range of ROOH compounds and their associated anharmonic normal mode wavenumbers were computed at the B2PLYP-D3/cc-pVTZ level of theory. Equivalent calculations were carried out for the OH and RO radicals formed upon O-O bond fission. Single point energies for the resultant optimized geometries were calculated at the CCSD(T)-F12/cc-pVTZ-F12 level of theory. Bond dissociation energies associated with RO + OH formation ($D_0[\text{RO-OH}]$) as well as the energy profile along the equatorial to axial isomerization pathway were calculated using the CCSD(T)-F12 energies and harmonic zero-point energies (ZPE) from the B2PLYP-D3/cc-pVTZ

calculations described above. Additional bond dissociation energies for CHHP and other organic hydroperoxides were computed using the W1BD composite method,⁵⁷ which is known to provide highly accurate thermochemical data within chemical accuracy (ca. 1 kcal mol⁻¹).

The ground state relaxed potential energy (PE) profile of CHHP along the O-O stretch coordinate (R_{O-O}) was calculated with single-state complete active space second-order perturbation theory (CASPT2) coupled to Dunning's aug-cc-pVTZ basis set (henceforth CASPT2/AVTZ). This was achieved by fixing R_{O-O} at various values across the range 1.0-5.6 Å and allowing the remainder of the nuclear framework to relax to their respective minimum energy configurations. These were based on a 4-state averaged complete active space self-consistent field (SA4-CASSCF) wavefunction employing the same basis set. An active space of 6 electrons in 4 orbitals was utilized to describe all significance short- and long-range interactions of the system while maintaining an appropriate computational expense. The active orbitals are displayed in Figure S5, while the orbital occupancies of the ground and excited electronic states of CHHP are shown in Figure S6 of ESI. An imaginary level shift of 0.4 E_H was used to aid convergence and to mitigate the involvement of intruder states.

The B2PLYP and W1BD calculations were performed using Gaussian 16 software suite⁵⁸ while the CCSD(T)-F12 and CASPT2 calculations were performed in the Molpro computation package.⁵⁹⁻⁶¹

3. Results and Discussion

Cyclohexyl hydroperoxide (CHHP) is predicted to have two stable chair conformations with the -OOH group in an equatorial or axial position of the ring. The present calculations indicate that the equatorial conformer is more stable by 0.2 kcal mol⁻¹. A prior investigation of the conformational landscape of CHHP (B3LYP-D3(BJ)/def2-TZVP) also found the equatorial conformer to be lower in energy, although by ca. 1 kcal mol⁻¹.⁵³ As illustrated in Figure 1, the interconversion between equatorial and axial configurations involves a sequence of geometry changes. The equatorial conformer initially transforms into a twist-boat configuration through a half-chair transition state (TS) barrier of 11.5 kcal mol⁻¹ (Table S1). Subsequently, interconversion between the two twist-boat configurations occurs via a TS (ca. 1.5 kcal mol⁻¹) with a boat geometry. The second twist-boat configuration converts into the axial conformer through another half-chair TS.

The relative stability of the equatorial and axial conformers suggests that both conformers will be populated with an estimated 60:40 ratio in favor of equatorial based on a room temperature Boltzmann population distribution. The significant TS barriers separating the equatorial and axial conformations indicates that both conformers will be populated under jet-cooled conditions, as found in a recent rotational spectroscopy study.⁵³ In addition, the earlier study anticipated a second equatorial conformer (0.67 kcal mol⁻¹ higher in energy) associated with internal rotation about the C-O bond (not shown in

Figure 1) that was not observed, presumably due to relaxation via a low torsional barrier of only 4 kcal mol⁻¹.⁵³




Figure 1. Energy profile connecting the equatorial and axial conformers of CHHP calculated at the CCSD(T)-F12/cc-pVTZ-F12//B2PLYPD3/cc-pVTZ level of theory.

In this study, we examine the IR spectrum of CHHP in the first overtone OH stretch region ($2\nu_{\text{OH}}$), guided by electronic structure calculations performed at the B2PLYP-D3/cc-pVTZ level of theory/basis. The equatorial and axial structures of CHHP are optimized, and the anharmonic vibrational frequencies and IR absorption intensities (Table S2) are calculated at these geometries using second-order vibrational perturbation theory (VPT2). The first overtone OH stretch for the equatorial and axial conformers of CHHP are predicted at 7006 and 6999 cm⁻¹, respectively, as shown in Figure 2. IR transitions associated with both equatorial and axial conformers of CHHP are anticipated based on the significant TS barrier separating the two conformers (Figure 1) and the observation of both conformers in the recent rotational spectroscopy study.⁵³

The IR action spectrum of CHHP was recorded by scanning the OPO from 6990 to 7040 cm⁻¹ at a scan speed of 0.1 cm⁻¹/s in the first overtone OH stretch ($2\nu_{\text{OH}}$) region under jet-cooled conditions, as shown in Figure 2. The IR action spectrum is obtained with LIF detection of OH products on the A-X (1,0) Q₁(3.5) transition at an IR-UV delay time of 700 ns (see Section S3 and Figure S7 of ESI). This requires IR multiphoton excitation, which induces dissociation of CHHP and leads to formation of cyclohexoxy (RO) and OH products. The IR action spectrum exhibits a single prominent band at 7012.5 cm⁻¹ with a full width at half maximum (FWHM) of ca. 7.5 cm⁻¹. The breadth of the observed feature is

noticeably greater than the ca. 5 cm^{-1} FWHM band contour expected for an isolated vibrational overtone transition of CHHP under jet-cooled conditions. The anticipated breadth is based on simulations using experimental rotational constants,⁵³ a rotational temperature of 10 K, the OPO bandwidth (0.9 cm^{-1}), and typical homogeneous broadening (1.7 cm^{-1}) associated with rapid (ps) intramolecular vibrational energy redistribution.^{12, 62}

The experimental observation of a single broadened feature at 7012.5 cm^{-1} can result from different population distributions for the equatorial and axial conformers of CHHP. One limiting case is a 60:40 population ratio of equatorial to axial conformers estimated for a Boltzmann distribution at 300 K and assumed to be unchanged upon supersonic expansion. In this case, the $2\nu_{\text{OH}}$ transitions for the equatorial and axial conformers would need to be more closely spaced, e.g. ca. 3 cm^{-1} apart, than predicted theoretically such that the two band contours merge into a single feature as shown in Figure 2. An alternative limiting scenario is that the observed $2\nu_{\text{OH}}$ feature originates primarily from the lower energy equatorial conformer with IR power broadening increasing the homogeneous linewidth to 6.0 cm^{-1} . The two limiting cases yield nearly indistinguishable spectra as illustrated in Figure S8. The observed feature appears at higher energy (ca. $10 \pm 2\text{ cm}^{-1}$) than predicted for the equatorial and axial conformers (7006 and 6999 cm^{-1} , respectively), which is well within the accuracy of the theoretical predictions ($10\text{-}20\text{ cm}^{-1}$).




Figure 2. Theoretically predicted $2\nu_{\text{OH}}$ transitions for the equatorial (red) and axial (blue) conformers of CHHP at 7006 and 6999 cm^{-1} , respectively, shown with relative intensities of 60:40 for a room temperature Boltzmann distribution. Experimental IR action spectrum (purple points) is overlaid with a simulated rotational band contour (black) for a scenario of closely spaced $2\nu_{\text{OH}}$ transitions (ca. 3 cm^{-1} apart) for equatorial and axial conformers. A nearly indistinguishable simulation results from an alternate limiting case with IR power broadening of the $2\nu_{\text{OH}}$ transition for the equatorial conformer that increases the homogeneous linewidth to 6.0 cm^{-1} (see Figure S8). The spectral shift of the observed feature (ca. $10 \pm 2\text{ cm}^{-1}$) is within the accuracy of the theoretical predictions ($10\text{-}20\text{ cm}^{-1}$).

A previous study reported an extensively broadened $2\nu_{\text{OH}}$ feature (12.2 cm^{-1} FWHM) for TBHP under jet-cooled conditions, which was attributed to tunneling across the symmetric double-well potential along the OH torsion coordinate.⁴⁹⁻⁵⁰ This resulted in multiple transitions and associated inhomogeneous broadening. Such broadening is not anticipated in CHHP since the double-well potential associated with OH torsion is asymmetric at the global minimum energy geometry of CHHP.

The IR power dependence of the $2\nu_{\text{OH}}$ transition of CHHP at 7012.5 cm^{-1} is obtained by varying the output power of the OPO from 13 to 28 mJ/pulse, as shown in Figure 3 with both axes on a $\log_{10}$ scale. The variation of the OH LIF signal with IR laser power reveals a slope of 2.3 ± 0.3 . The nonlinear dependence on IR power implies that the dissociation process requires multiphoton absorption of IR photons in $2\nu_{\text{OH}}$ region. The slope indicates that more than two photons are required to promote CHHP above the dissociation limit, yielding the OH radicals that are detected. The energies of the OH fundamental and first overtone transitions are obtained from the VPT2 calculation, while higher OH overtone levels ($3\nu_{\text{OH}}$ and $4\nu_{\text{OH}}$) illustrated in Figure 3 are estimated from an anharmonic vibrational analysis.




Figure 3. (Left) Energy diagram showing the multiphoton dissociation of CHHP. IR excitation of the $2\nu_{\text{OH}}$ transition (black arrow) followed by absorption of additional photon(s) provides sufficient energy to exceed the O-O bond dissociation limit of CHHP (dashed line), resulting in cyclohexoxy and OH radicals. The OH radicals are detected by LIF on the $A^2\Sigma^+ - X^2\Pi_{3/2}(1,0) Q_1(3.5)$ transition at 282 nm. (Right) IR-induced OH LIF signal intensity at 7012.5 cm^{-1} as a function of IR power (mJ/pulse). Both axes are plotted on a $\log_{10}$ scale, yielding a slope of 2.3 ± 0.3 . The nonlinear IR power dependence indicates multiphoton excitation of CHHP.

Figure 4 presents the CASPT2 relaxed potential energy (PE) profile along the O-O stretch coordinate (R_{OO}) for the equatorial conformer of CHHP. The PE profile shows a weak saddle point at $R_{\text{OO}} = 2.78\text{ Å}$ followed by a shallow minimum at $R_{\text{OO}} = 2.8\text{ Å}$, the latter of which arises from a hydrogen bonding interaction between the O lone pair of the cyclohexoxy radical (RO) and the H-atom of OH unit. The

energy profile along the O–O coordinate beyond $R_{OO} > 3.4$ Å proceeds smoothly to the RO + OH asymptotic limit, which is consistent with earlier studies of other hydroperoxides.^{63–64} The corresponding PE profile for the axial conformer of CHHP is shown in Figure S9. Using the asymptotic energies, bond dissociation energies (D_e) of ca. 46 kcal mol^{−1} are determined for both equatorial and axial conformers. Upon ZPE correction, the resulting D_0 values for both conformers are ca. 42 kcal mol^{−1}. The W1BD method predicts a D_0 value of 43 kcal mol^{−1} (Table 1), which agrees well with the ZPE-corrected CASPT2 energies. The W1BD method is known to give thermochemical quantities that are within chemical accuracy (1 kcal mol^{−1}).




Figure 4. Energy diagram for the equatorial conformer of CHHP (ROOH) along the O–O stretch coordinate (R_{OO}) to form cyclohexoxy (RO) + OH products. The potential is calculated at the CASPT2/aug-cc-pVTZ level of theory. A bond dissociation energy (D_e) of 46 kcal mol^{−1} is obtained for the equatorial conformer of CHHP.

Given that the IR excitation is tuned to 7012.5 cm^{−1} (ca. 20 kcal mol^{−1}), which corresponds to the $2\nu_{OH}$ transition of CHHP, multiphoton dissociation is required to initiate O–O bond fission. With a W1BD derived D_0 value of 43 kcal mol^{−1}, at least three photons of the OH first overtone transition would be required to promote O–O bond fission. The third photon would impart an excess energy of ca. 17 kcal mol^{−1} into CHHP, which upon intramolecular vibrational energy redistribution, would provide sufficient energy in O–O stretch to promote O–O bond fission.

The O–O bond dissociation energy (D_0) for CHHP (cyc-C₆H₁₁O–OH) is compared with those computed for four simpler hydroperoxides (ROOH) using the same W1BD method in Table 1. The D_0 values of the representative hydroperoxides are also evaluated with the CCSD(T)-F12/cc-pVTZ-F12 method (Table 1). The calculated dissociation energies using the W1BD and CCSD(T)-F12 methods compare favorably with those previously reported with the CBS-APNO and G4 methods.⁵² Prior

theoretical studies of O-O bond dissociation energies for ROOH with various substituents range from ca. 35 to 50 kcal mol⁻¹ with typical O-O bond dissociation energy of ca. 45 kcal mol⁻¹.⁵¹⁻⁵²

Table 1. Computed O-O dissociation energies (D_0 , kcal mol⁻¹) for several hydroperoxides evaluated at various levels of theory and comparison with experiment.

	This work		Bach <i>et al.</i> ^a		Expt.
	W1BD	CCSD(T)-F12/ cc-pVTZ-F12 ^b	CBS-APNO	G4	
HO-OH	48.4	48.1	50.2	49.0	50.30 ^c
CH ₃ O-OH	41.9	42.4	44.6	43.2	44.67 ^c
(CH ₃) ₃ CO-OH	44.0	43.5	45.8	43.9	44.1 ^d
(CH ₃) ₂ CHO-OH	43.9	44.1	-	-	-
cyc-C ₆ H ₁₁ O-OH	43.4	-	-	-	-

a. Ref. 52.

b. ZPE corrected with vibrational wavenumbers derived at the B2PLYP-D3/cc-pVTZ level.

c. Ref. 65.

d. Ref. 66.

The present calculations, along with those conducted previously,⁵¹⁻⁵² show that the organic hydroperoxides have $D_0(\text{O-O})$ values that are ca. 4-7 kcal mol⁻¹ lower than hydrogen peroxide. This can be qualitatively understood by the relative stabilities of the OH vs. OR radical products. Alkyl groups are well-known σ -electron donors and the H-atoms of the CH₃ moiety participate in hyperconjugation. Both effects are expected to stabilize the oxygen-centered odd-electron (the electron deficient p -orbital) of the OR radical. Equivalent stabilizations are not available for OH, resulting in more stable OR radicals. Interestingly, the organic hydroperoxide with a primary carbon-center (CH₃O-OH) has a D_0 value that is ca. 1-2 kcal mol⁻¹ lower than secondary [(CH₃)₂CHO-OH and cyc-C₆H₁₁O-OH] and tertiary (CH₃)₃CO-OH carbon-centered hydroperoxides. The secondary and tertiary carbon-centers are more electron-rich than primary carbon-centers due to the +I inductive effects imparted by the alkyl-substituents; this is expected to increase the nucleophilicity of the carbon-center. Additionally, hyperconjugation is not available in the secondary and tertiary carbon centers. Both effects are expected to destabilize secondary and tertiary carbon-centered organic hydroperoxides, consistent with their higher bond dissociation energies.

Conclusions

A new synthetic approach for generating CHHP is utilized in the present study of its IR spectroscopy and multiphoton dissociation dynamics. The IR spectrum of jet-cooled CHHP has been recorded in the

first overtone OH stretch ($2\nu_{\text{OH}}$) region. Quantum chemical studies reveal that the equatorial conformer of CHHP is slightly more stable than the axial conformer (ca. 0.2 kcal mol⁻¹), and separated by a substantial TS barrier for isomerization (ca. 11.5 kcal mol⁻¹). As a result, both conformers may be populated under jet-cooled conditions, in accord with a recent rotational spectroscopy investigation.⁵³ IR multiphoton excitation promotes CHHP above the O-O dissociation limit, resulting in OH X²Π ($v=0$) radical products that are detected via UV-LIF. IR action spectroscopy yields a distinctive, yet broadened, vibrational overtone feature for CHHP at 7012.5 cm⁻¹. This can be ascribed to various scenarios ranging from a single homogeneously broadened feature dominated by the equatorial conformer to overlapping band contours for the equatorial and axial conformers that merge into a single feature. The nonlinear IR power dependence of this spectral feature confirms that a minimum of three photons with energies corresponding to $2\nu_{\text{OH}}$ are required to promote O-O bond fission. This observation is confirmed by computation of relaxed CASPT2 potential energy profiles along the O-O bond stretch coordinate for the equatorial and axial conformers, and determination of the O–O bond dissociation energy using the W1BD method.

The present study adds to the small but growing body of work on larger organic hydroperoxides. In the future, we anticipate the emergence of synthetic procedures for laboratory investigation of the spectroscopy and dissociation dynamics of more complex and atmospherically relevant organic hydroperoxides.

Data availability

All supporting data and methods are available in the ESI.

Author contributions

M.I.L. and T.N.V.K. designed the experimental and theoretical aspects of the research project, respectively. C.S. and M.C.K. synthesized cyclohexyl hydroperoxide (CHHP). T.K.R and Y.Q. conducted the experiments and analyzed the experimental data. L.K., R.R., and T.N.V.K. carried out the theoretical calculations. All authors discussed the results and contributed to writing the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgments

Research in the Lester laboratory was supported by the National Science Foundation under grants CHE-1955068 and CHE-2301298. M.C.K. is grateful to the NSF (CHE-2102626) for financial support of this research. Partial instrumentation support (NMR spectrometer) was provided by NSF CHE-1827457 and the Vagelos Institute for Energy Science and Technology. T.K.R. was primarily supported by a Walter-Benjamin Scholarship funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project number 508074809. T.N.V.K is grateful to the NSF (OIA- 2229695) for financial support. This work used the Advanced Cyberinfrastructure Coordination Ecosystem: Service & Support (ACCESS) program, which is supported by National Science Foundation grants #2138259, #2138286, #2138307, #2137603, and #2138296, through the allocation TG-CHE190088.

References

1. S. Enami, Fates of Organic Hydroperoxides in Atmospheric Condensed Phases. *J. Phys. Chem. A* **2021**, *125*, 4513-4523.
2. F. Bianchi; O. Garmash; X. He; C. Yan; S. Iyer; I. Rosendahl; Z. Xu; M. P. Rissanen; M. Riva; R. Taipale; N. Sarnela; T. Petäjä; D. R. Worsnop; M. Kulmala; M. Ehn; H. Junninen, The role of highly oxygenated molecules (HOMs) in determining the composition of ambient ions in the boreal forest. *Atmos. Chem. Phys.* **2017**, *17*, 13819-13831.
3. L. Vereecken; B. Nozière, H-migration in peroxy radicals under atmospheric conditions. *Atmos. Chem. Phys.* **2020**, *20*, 7429-7458.
4. J. Lelieveld; T. M. Butler; J. N. Crowley; T. J. Dillon; H. Fischer; L. Ganzeveld; H. Harder; M. G. Lawrence; M. Martinez; D. Taraborrelli; J. Williams, Atmospheric oxidation capacity sustained by a tropical forest. *Nature* **2008**, *452*, 737-740.
5. D. L. Osborn, Reaction mechanisms on multiwell potential energy surfaces in combustion (and atmospheric) chemistry. *Annu. Rev. Phys. Chem.* **2017**, *68*, 233-260.
6. J. D. Savee; E. Papajak; B. Rotavera; H. Huang; A. J. Eskola; O. Welz; L. Sheps; C. A. Taatjes; J. Zádor; D. L. Osborn, Direct observation and kinetics of a hydroperoxyalkyl radical (QOOH). *Science* **2015**, *347*, 643-646.
7. L. Vereecken; J. S. Francisco, Theoretical studies of atmospheric reaction mechanisms in the troposphere. *Chem. Soc. Rev.* **2012**, *41*, 6259-6293.
8. G. Meloni; P. Zou; S. J. Klippenstein; M. Ahmed; S. R. Leone; C. A. Taatjes; D. L. Osborn, Energy-Resolved Photoionization of Alkylperoxy Radicals and the Stability of Their Cations. *J. Am. Chem. Soc.* **2006**, *128*, 13559-13567.
9. W. P. L. Carter, A detailed mechanism for the gas-phase atmospheric reactions of organic compounds. *Atmos. Environ. Part A. General Topics* **1990**, *24*, 481-518.
10. R. Atkinson; D. L. Baulch; R. A. Cox; R. F. Hampson, Jr.; J. A. Kerr; M. J. Rossi; J. Troe, Evaluated Kinetic, Photochemical and Heterogeneous Data for Atmospheric Chemistry: Supplement V. IUPAC Subcommittee on Gas Kinetic Data Evaluation for Atmospheric Chemistry. *J. Phys. Chem. Ref. Data* **1997**, *26*, 521-1011.
11. A. S. Hansen; T. Bhagde; K. B. Moore; D. R. Moberg; A. W. Jasper; Y. Georgievskii; M. F. Vansco; S. J. Klippenstein; M. I. Lester, Watching a hydroperoxyalkyl radical ($\bullet$ QOOH) dissociate. *Science* **2021**, *373*, 679-682.
12. A. S. Hansen; T. Bhagde; Y. Qian; A. Cavazos; R. M. Huchmala; M. A. Boyer; C. F. Gavin-Hanner; S. J. Klippenstein; A. B. McCoy; M. I. Lester, Infrared spectroscopic signature of a hydroperoxyalkyl radical ($\bullet$ QOOH). *J. Chem. Phys.* **2022**, *156*, 014301.
13. T. Bhagde; A. S. Hansen; S. Chen; P. J. Walsh; S. J. Klippenstein; M. I. Lester, Energy-resolved and time-dependent unimolecular dissociation of hydroperoxyalkyl radicals ($\bullet$ QOOH). *Faraday Discuss.* **2022**, *238*, 575-588.
14. J. D. Crounse; L. B. Nielsen; S. Jørgensen; H. G. Kjaergaard; P. O. Wennberg, Autoxidation of Organic Compounds in the Atmosphere. *J. Phys. Chem. Lett.* **2013**, *4*, 3513-3520.
15. M. Ehn; J. A. Thornton; E. Kleist; M. Sipilä; H. Junninen; I. Pullinen; M. Springer; F. Rubach; R. Tillmann; B. Lee; F. Lopez-Hilfiker; S. Andres; I.-H. Acir; M. Rissanen; T. Jokinen; S. Schobesberger; J. Kangasluoma; J. Kontkanen; T. Nieminen; T. Kurtén; L. B. Nielsen; S. Jørgensen; H. G. Kjaergaard; M. Canagaratna; M. D. Maso; T. Berndt; T. Petäjä; A. Wahner; V.-M. Kerminen; M. Kulmala; D. R. Worsnop; J. Wildt; T. F. Mentel, A large source of low-volatility secondary organic aerosol. *Nature* **2014**, *506*, 476-479.
16. T. Jokinen; M. Sipilä; S. Richters; V.-M. Kerminen; P. Paasonen; F. Stratmann; D. Worsnop; M. Kulmala; M. Ehn; H. Herrmann; T. Berndt, Rapid Autoxidation Forms Highly Oxidized RO₂ Radicals in the Atmosphere. *Angew. Chem. Int. Ed.* **2014**, *53*, 14596-14600.

17. T. Berndt; S. Richters; T. Jokinen; N. Hyttinen; T. Kurtén; R. V. Otkjær; H. G. Kjaergaard; F. Stratmann; H. Herrmann; M. Sipilä; M. Kulmala; M. Ehn, Hydroxyl radical-induced formation of highly oxidized organic compounds. *Nat. Commun.* **2016**, *7*, 13677.
18. E. Praske; R. V. Otkjær; J. D. Crounse; J. C. Hethcox; B. M. Stoltz; H. G. Kjaergaard; P. O. Wennberg, Atmospheric autoxidation is increasingly important in urban and suburban North America. *Proc. Natl. Acad. Sci. U.S.A.* **2018**, *115*, 64-69.
19. D. J. Medeiros; M. A. Blitz; P. W. Seakins; L. K. Whalley, Direct Measurements of Isoprene Autoxidation: Pinpointing Atmospheric Oxidation in Tropical Forests. *J. Am. Chem. Soc. Au* **2022**, *2*, 809-818.
20. M. Ge; S. Tong; W. Wang; W. Zhang; M. Chen; C. Peng; J. Li; L. Zhou; Y. Chen; M. Liu, Important Oxidants and Their Impact on the Environmental Effects of Aerosols. *J. Phys. Chem. A* **2021**, *125*, 3813-3825.
21. C. Li; R. Signorell, Understanding vapor nucleation on the molecular level: A review. *J. Aerosol Sci.* **2021**, *153*, 105676.
22. V. F. McNeill, Virtual Special Issue: New Advances in Organic Aerosol Chemistry. *ACS Earth Space Chem.* **2020**, *4*, 491-494.
23. E. W. Kaiser, Temperature and Pressure Dependence of the C_2H_4 Yield from the Reaction $C_2H_5 + O_2$. *J. Phys. Chem.* **1995**, *99*, 707-711.
24. E. W. Kaiser, Formation of C_3H_6 from the Reaction $C_3H_7 + O_2$ between 450 and 550 K. *J. Phys. Chem. A* **1998**, *102*, 5903-5906.
25. E. W. Kaiser; T. J. Wallington, Formation of C_3H_6 from the Reaction $C_3H_7 + O_2$ and C_2H_3Cl from $C_2H_4Cl + O_2$ at 297 K. *J. Phys. Chem.* **1996**, *100*, 18770-18774.
26. I. C. Plumb; K. R. Ryan, Kinetic studies of the reaction of C_2H_5 with O_2 at 295 K. *Int. J. Chem. Kinet.* **1981**, *13*, 1011-1028.
27. E. P. Clifford; J. T. Farrell; J. D. DeSain; C. A. Taatjes, Infrared Frequency-Modulation Probing of Product Formation in Alkyl + O_2 Reactions: I. The Reaction of C_2H_5 with O_2 between 295 and 698 K. *J. Phys. Chem. A* **2000**, *104*, 11549-11560.
28. J. D. DeSain; E. P. Clifford; C. A. Taatjes, Infrared Frequency-Modulation Probing of Product Formation in Alkyl + O_2 Reactions: II. The Reaction of C_3H_7 with O_2 between 296 and 683 K. *J. Phys. Chem. A* **2001**, *105*, 3205-3213.
29. J. D. DeSain; C. A. Taatjes; J. A. Miller; S. J. Klippenstein; D. K. Hahn, Infrared frequency-modulation probing of product formation in alkyl + O_2 reactions. Part IV. Reactions of propyl and butyl radicals with O_2 . *Faraday Discuss.* **2002**, *119*, 101-120.
30. E. W. Kaiser; I. M. Lorkovic; T. J. Wallington, Pressure dependence of the ethene yield from the reaction ethyl radical + oxygen. *J. Phys. Chem.* **1990**, *94*, 3352-3354.
31. E. G. Estupiñán; J. D. Smith; A. Tezaki; S. J. Klippenstein; C. A. Taatjes, Measurements and Modeling of DO_2 Formation in the Reactions of C_2D_5 and C_3D_7 Radicals with O_2 . *J. Phys. Chem. A* **2007**, *111*, 4015-4030.
32. J. Zádor; C. A. Taatjes; R. X. Fernandes, Kinetics of elementary reactions in low-temperature autoignition chemistry. *Prog. Energy Combust. Sci.* **2011**, *37*, 371-421.
33. J. J. Orlando; G. S. Tyndall, Laboratory studies of organic peroxy radical chemistry: an overview with emphasis on recent issues of atmospheric significance. *Chem. Soc. Rev.* **2012**, *41*, 6294-6317.
34. R. W. Walker; C. Morley, Chapter 1 Basic chemistry of combustion. In *Comprehensive Chemical Kinetics*, Pilling, M. J., Ed. Elsevier: 1997; Vol. 35, pp 1-124.
35. S. Sharma; S. Raman; W. H. Green, Intramolecular Hydrogen Migration in Alkylperoxy and Hydroperoxyalkylperoxy Radicals: Accurate Treatment of Hindered Rotors. *J. Phys. Chem. A* **2010**, *114*, 5689-5701.
36. J. D. DeSain; C. A. Taatjes, Infrared Frequency-Modulation Probing of Product Formation in Alkyl + O_2 Reactions: III. The Reaction of Cyclopentyl Radical ($c-C_5H_9$) with O_2 between 296 and 723 K. *J. Phys. Chem. A* **2001**, *105*, 6646-6654.

37. H. R. Dübal; F. F. Crim, Vibrational overtone predissociation spectroscopy of hydrogen peroxide. *J. Chem. Phys.* **1985**, *83*, 3863-3872.
38. T. M. Ticich; M. D. Likar; H. R. Dübal; L. J. Butler; F. F. Crim, Vibrationally mediated photodissociation of hydrogen peroxide. *J. Chem. Phys.* **1987**, *87*, 5820-5829.
39. D. T. Colbert; E. L. Sibert, III, Theory of vibrationally mediated photodissociation of HOOH: Delocalized tails in a localized wave function. *J. Chem. Phys.* **1991**, *94*, 6519-6545.
40. M. D. Likar; J. E. Baggott; F. F. Crim, Vibrationally mediated photodissociation of t-butyl hydroperoxide: Vibrational overtone spectroscopy and photodissociation dynamics. *J. Chem. Phys.* **1989**, *90*, 6266-6274.
41. M. D. Likar; J. E. Baggott; A. Sinha; T. M. Ticich; R. L. V. Wal; F. F. Crim, Vibrationally mediated photodissociation. *J. Chem. Soc., Faraday Trans. 2* **1988**, *84*, 1483-1497.
42. M. Brouard; M. T. Martinez; J. O'Mahony; J. P. Simons, Energy and angular momentum disposals in the vibrationally mediated photodissociation of HOOH and HOOD. *Mol. Phys.* **1990**, *69*, 65-84.
43. J. August; M. Brouard; M. P. Docker; C. J. Milne; J. P. Simons; R. Lavi; S. Rosenwaks; D. Schwartz-Lavi, Photodissociation dynamics of tert-butyl nitrite (S₂) and tert-butyl hydroperoxide at 248-250 nm. *J. Phys. Chem.* **1988**, *92*, 5485-5491.
44. Y. Inagaki; Y. Matsumi; M. Kawasaki, Formation of a Hydrogen Atom from the Photodissociation of Hydrogen Peroxide at 193 nm. *Bull. Chem. Soc. Jpn.* **1993**, *66*, 3166-3170.
45. S. K. Shin; S. O. Park; Y. S. Choi; H. L. Kim; C. R. Park, Photodissociation Dynamics of Cumene Hydroperoxide at 248 and 193 nm. *J. Phys. Chem. A* **2001**, *105*, 10018-10024.
46. B.-G. Ryu; C. R. Park; Y. Lee; S. K. Shin; H. L. Kim, Photodissociation dynamics of tert-butyl hydroperoxide at 266 nm. *J. Photochem. Photobiol., A* **2002**, *149*, 15-21.
47. S. Hsieh; R. Vushe; Y. T. Tun; J. L. Vallejo, Trends in organic hydroperoxide photodissociation and absorption cross sections between 266 and 377nm. *Chem. Phys. Lett.* **2014**, *591*, 99-102.
48. M. D. Likar; J. E. Baggott; F. F. Crim, Vibrationally mediated photodissociation of t-butyl hydroperoxide: Vibrational overtone spectroscopy and photodissociation dynamics. *J. Chem. Phys.* **1989**, *90*, 6266-6274.
49. A. S. Hansen; R. M. Huchmala; E. Vogt; M. A. Boyer; T. Bhagde; M. F. Vansco; C. V. Jensen; A. Kjærsgaard; H. G. Kjaergaard; A. B. McCoy; M. I. Lester, Coupling of torsion and OH-stretching in tert-butyl hydroperoxide. I. The cold and warm first OH-stretching overtone spectrum. *J. Chem. Phys.* **2021**, *154*, 164306.
50. E. Vogt; R. M. Huchmala; C. V. Jensen; M. A. Boyer; J. Wallberg; A. S. Hansen; A. Kjærsgaard; M. I. Lester; A. B. McCoy; H. G. Kjaergaard, Coupling of torsion and OH-stretching in tert-butyl hydroperoxide. II. The OH-stretching fundamental and overtone spectra. *J. Chem. Phys.* **2021**, *154*, 164307.
51. R. D. Bach; P. Y. Ayala; H. B. Schlegel, A Reassessment of the Bond Dissociation Energies of Peroxides. An ab Initio Study. *J. Am. Chem. Soc.* **1996**, *118*, 12758-12765.
52. R. D. Bach; H. B. Schlegel, Bond Dissociation Energy of Peroxides Revisited. *J. Phys. Chem. A* **2020**, *124*, 4742-4751.
53. P. Pinacho; W. Sun; D. A. Obenchain; M. Schnell, Conformational analysis of cyclohexyl hydroperoxide by rotational spectroscopy. *J. Mol. Spectrosc.* **2023**, *392*, 111758.
54. Z. Hu; Q. Di; B. Liu; Y. Li; Y. He; Q. Zhu; Q. Xu; P. Dagaut; N. Hansen; S. M. Sarathy; L. Xing; D. G. Truhlar; Z. Wang, Elucidating the photodissociation fingerprint and quantifying the determination of organic hydroperoxides in gas-phase autoxidation. *Proc. Natl. Acad. Sci. U.S.A.* **2023**, *120*, e2220131120.
55. J. B. Sperry; M. Azuma; S. Stone, Explosive Hazard Identification in Pharmaceutical Process Development: A Novel Screening Method and Workflow for Shipping Potentially Explosive Materials. *Org. Process Res. Dev.* **2021**, *25*, 212-224.
56. T. Tokuyasu; S. Kunikawa; A. Masuyama; M. Nojima, Co(III)-Alkyl Complex- and Co(III)-Alkylperoxo Complex-Catalyzed Triethylsilylperoxidation of Alkenes with Molecular Oxygen and Triethylsilane. *Org. Lett.* **2002**, *4*, 3595-3598.

57. E. C. Barnes; G. A. Petersson; J. A. Montgomery, Jr.; M. J. Frisch; J. M. L. Martin, Unrestricted Coupled Cluster and Brueckner Doubles Variations of W1 Theory. *J. Chem. Theory Comput.* **2009**, *5*, 2687-2693.
58. M. J. Frisch; G. W. Trucks; H. B. Schlegel; G. E. Scuseria; M. A. Robb; J. R. Cheeseman, G. Scalmani; V. Barone; G. A. Petersson; H. Nakatsuji, *et al.* *Gaussian 16*, Rev. C.01; Gaussian, Inc.: Wallingford, CT, 2016.
59. P. J. K. H.-J. Werner, P. Celani, W. Györfy, A. Hesselmann, D. Kats, G. Knizia, A. Köhn, T. Korona, D. Kreplin, R. Lindh, *et. al.*, *MOLPRO 2020.1, a package of ab initio programs*, see www.molpro.net, 2020.
60. H.-J. Werner; P. J. Knowles; G. Knizia; F. R. Manby; M. Schütz, Molpro: a general-purpose quantum chemistry program package. *WIREs Computational Molecular Science* **2012**, *2*, 242-253.
61. H.-J. Werner; P. J. Knowles; F. R. Manby; J. A. Black; K. Doll; A. Heßelmann; D. Kats; A. Köhn; T. Korona; D. A. Kreplin; Q. Ma; T. F. Miller; A. Mitrushchenkov; K. A. Peterson; I. Polyak; G. Rauhut; M. Sibae, The Molpro quantum chemistry package. *J. Chem. Phys.* **2020**, *152*, 144107.
62. A. S. Hansen; Y. Qian; C. A. Sojda; M. C. Kozlowski; V. J. Esposito; J. S. Francisco; S. J. Klippenstein; M. I. Lester, Rapid Allylic 1,6 H-Atom Transfer in an Unsaturated Criegee Intermediate. *J. Am. Chem. Soc.* **2022**, *144*, 5945-5955.
63. N. M. Kidwell; H. Li; X. Wang; J. M. Bowman; M. I. Lester, Unimolecular dissociation dynamics of vibrationally activated CH₃CHOO Criegee intermediates to OH radical products. *Nat. Chem.* **2016**, *8*, 509-514.
64. B. Marchetti; V. J. Esposito; R. E. Bush; T. N. V. Karsili, The states that hide in the shadows: the potential role of conical intersections in the ground state unimolecular decay of a Criegee intermediate. *Phys. Chem. Chem. Phys.* **2022**, *24*, 532-540.
65. Active Thermochemical Tables. <https://atct.anl.gov/Thermochemical%20Data/version%201.130/index.php>. (accessed February 2024).
66. W. Reints; D. A. Pratt; H.-G. Korth; P. Mulder, O–O Bond Dissociation Enthalpy in Di(trifluoromethyl) Peroxide (CF₃OO CF₃) as Determined by Very Low Pressure Pyrolysis. Density Functional Theory Computations on O–O and O–H Bonds in (Fluorinated) Derivatives. *J. Phys. Chem. A* **2000**, *104*, 10713-10720.

TOC

