



PAPER

Existence and uniqueness of solutions to the Peierls–Nabarro model in anisotropic media

To cite this article: Yuan Gao and James M Scott 2024 *Nonlinearity* **37** 025010

View the [article online](#) for updates and enhancements.

You may also like

- [Constrictions and Jogs in Extended Dislocations](#)
A N Stroh
- [Modelling \$\[1\ 0\ 0\]\$ and \$\[0\ 1\ 0\]\$ screw dislocations in \$\text{MgSiO}_3\$ perovskite based on the Peierls–Nabarro–Galerkin model](#)
K Gouriet, Ph Carrez and P Cordier
- [Modification of the Peierls–Nabarro Model for Edge Dislocation Core](#)
H B Huntington

Existence and uniqueness of solutions to the Peierls–Nabarro model in anisotropic media

Yuan Gao¹  and James M Scott^{2,*} 

¹ Department of Mathematics, Purdue University, West Lafayette, IN, United States of America

² Department of Applied Physics and Applied Mathematics, Columbia University, New York, NY, United States of America

E-mail: jms2555@columbia.edu

Received 11 April 2023; revised 29 August 2023

Accepted for publication 20 December 2023

Published 12 January 2024

Recommended by Dr Theodore Dimitrios Drivas



CrossMark

Abstract

We study the existence and uniqueness of solutions to the vector field Peierls–Nabarro (PN) model for curved dislocations in a transversely isotropic medium. Under suitable assumptions for the misfit potential on the slip plane, we reduce the 3D PN model to a nonlocal scalar Ginzburg–Landau equation. For a particular range of elastic coefficients, the nonlocal scalar equation with explicit nonlocal positive kernel is derived. We prove that any stable steady solution has a one-dimensional profile. As a result, we obtain that solutions to the scalar equation, as well as the original 3D system, are characterized as a one-parameter family of straight dislocations. This paper generalizes results found previously for the full isotropic case to an anisotropic setting.

Keywords: nonlocal equations, Peierls–Nabarro Model, Ginzburg–Landau equation, anisotropic elasticity, fractional Laplacian, Dirichlet-to-Neumann map, dimension reduction

Mathematics Subject Classification numbers: 35A02, 35J50, 35Q74, 35R09, 35J60

* Author to whom any correspondence should be addressed.

1. Introduction

The Peierls–Nabarro (PN) model plays a fundamental role in describing dislocations or line defects in materials. Analysis of the model provides further insight on the design of new materials with desirable atomistic properties. The vectorial PN model is a nonlinear model that describes the core structure of the dislocation by incorporating the atomistic effect in the dislocation core into the continuum elastic model [14, 18]. As the most common line defects in materials [12, 21], the core structures of dislocations are fundamental problems for studying the stability of material structure and the minimum energy barrier for plastic deformations. Particularly for general anisotropic materials, the stationary dislocation core profile and its rigidity is a central problem. The existence and rigidity properties of dislocation profiles to the vectorial PN model for fully isotropic elastic media has been explored recently in [4, 6, 7]. In this paper, we investigate analogous questions for an anisotropic constitutive medium. In particular, we find the range of parameters in the anisotropic constitutive relation for which the stationary solution to the vectorial PN model is a one-dimensional profile given by a proper one-dimensional scalar nonlocal Ginzburg–Landau equation with an explicitly computed kernel.

In the three-dimensional PN model for the displacement vector $\mathbf{u} = (u_1, u_2, u_3)$, two half-spaces separated by the slip plane of a dislocation are assumed to be linear elastic continua described by Hooke’s law. Here the slip plane is assumed to be a fixed plane Γ , where the horizontal displacement discontinuity (known as disregistry) happens. The two elastic continua are connected by a nonlinear atomistic potential force across the slip plane, which results in a boundary stress nonlinearly depending on the disregistry across the slip plane. The direction and magnitude of the above disregistry due to the dislocation are characterized by the Burgers vector $\mathbf{b}$. The magnitude of the Burgers vector represents the typical length to observe a heavily distorted region in the dislocation core. Hence it is natural to rescale all the quantities including spatial variables x_1, x_2, x_3 , the displacement vector $\mathbf{u} = (u_1, u_2, u_3)$ and $\mathbf{b}$ with respect to the magnitude of the Burgers vector. After rescaling, we regard all these quantities (with the same notations) as dimensionless and regard the magnitude of the Burgers vector as 4. This means, after some symmetric assumption on the upper/lower elastic bulk, the disregistry at far field in the shear direction yields the bi-states far field condition ± 1 ; see [4, 6, 7, 21]

In this work the elastic media are transversely isotropic; the stress–strain response of the material is isotropic about an axis normal to the transverse, or basal, plane. The stiffness tensor for transversely isotropic media is uniquely determined by five independent elastic coefficients, see (2.3) below. We compute the Dirichlet-to-Neumann map associated to the bulk constitutive law in the two half-spaces separated by the slip plane in order to reduce the 3D model to a strongly-coupled 2D nonlocal system. Because of the anisotropy of the media, different slip planes Γ result in different forms of reduced models. In this work we treat two different orientations of the slip plane. The first case is when Γ is oriented perpendicular to the plane of isotropy, and the second case is when Γ and the plane of isotropy are parallel; see the illustrations in figures 1 and 2.

The nonlinear misfit potential W , defined on the slip plane, creates the boundary stress due to the dislocation disregistry across the slip plane. In each case, we assume that W depends only on the displacement disregistry in a single direction, and that W is a periodic/double well potential in that one variable. This assumption permits further reduction of the 2D system to a nonlocal scalar Ginzburg–Landau equation. We compute the exact expression for this integro-differential operator, and find the range of elastic coefficients for which the kernel of the integro-differential operator is positive. The positivity of the kernel allows us to apply the program that was carried out in [4] to establish the existence and rigidity of minimizers to the PN model in the case of full isotropy. We conclude that the scalar Ginzburg–Landau equation




Figure 1. The slip plane oriented perpendicular to the plane of isotropy.




Figure 2. The slip plane oriented parallel to the plane of isotropy.

has a unique (up to translation and rotation) solution with a 1D profile. We then follow the elastic extension procedures in [6, 7] to conclude the well-posedness of the full 3D PN system.

Again because of the anisotropy of the full 3D media, assuming different directions for the dependence on W leads to different reduced 1D equations. In case I, when Γ is perpendicular to the plane of isotropy and $W = W(u_1)$ depends only on the in-plane-of-isotropy direction of

the displacement, the reduced 2D operator is highly-anisotropic, in the sense that its Fourier symbol is characterized by more than three distinct frequency magnitudes. In case II, when Γ is perpendicular to the plane of isotropy and $W = W(u_3)$ depends only on the out-of-plane-of-isotropy direction of the displacement, the reduced 2D operator is anisotropic but depends on three frequency magnitudes. In case III, when Γ is parallel to the plane of isotropy, the resulting disregistry depends only on the reference configuration in the plane of isotropy, i.e. the reduced 2D model closely resembles its analogue in the fully isotropic case. This 2D model is itself isotropic, and so choosing W to depend on either direction of the displacement results in equivalent models. Our results in these three cases are summarized in three theorems respectively: propositions 2.4, 2.5 and 5.2.

In the context of materials science, our results say that, for a certain set of elastic stiffness tensors, and if the misfit potential depends only on the shear displacement relative to Γ , then the equilibrium dislocation profile depends only on shear displacements in the slip plane. Moreover, this unique (up to translations) shear displacement is a strictly monotonic 1D profile that connects two stable states at far fields.

The proofs rely on the local BV estimates originally developed in [2] to study the quantitative flatness of nonlocal minimal surfaces. In [4, theorem 4.6], the authors use the flatness estimate for the scalar solution of the nonlocal differential equation by combining the interior BV estimate and a sharp interpolation inequality in [2, 5, 11, 19]. The crucial ingredients are the explicit properties for the kernel representation of the nonlocal operator. To be precise, the nonlocal kernel needs to be homogeneous with respect to dilations, strictly positive, and have good decay estimates; see [2, 3, 6, 11]. Taking into account the transverse anisotropy of the materials, for each of the three cases described above, we will take a specific range of the five elastic constants so that we have the desired properties of the kernel K that describes the scalar nonlocal operator. We use these properties along with [4, theorem 4.6] to obtain that any bounded stable solution to the reduced scalar nonlocal equation in two dimensions has a 1D profile. For other methods on the rigidity property for nonlocal Allen–Cahn equations, we refer to [6, theorem 2] for using estimates of the spectrum of a linear operator and refer to [1] for using a Liouville-type theorem for the original local equation. With the obtained rigidity result and the explicit kernel representation for the one-dimensional nonlocal Allen–Cahn equation reduced from the 3D vectorial dislocation model, one can further explore the long time behaviour of dislocations using maximum principles for the nonlocal operator. We leave this for a future work and refer to the methods in [8, 9, 15, 16] in the case of isotropy.

The PN model for transversely anisotropic media is a generalization of the PN model for full isotropic media. Indeed, each result in this paper has a corresponding result for the PN model in the full isotropic setting by making a particular choice of elastic coefficients; we illustrate this in remarks throughout the paper. In particular, many of the results characterizing the 2D and 1D nonlocal operators studied in [4, 6, 7] can be recovered from their analogous statements in this work. At the same time, in cases I and II we make some special choices of elastic coefficients so that the reduced model can be analysed relatively easily, but the question of whether the results in this work hold for a general choice of coefficients that, for instance, do not satisfy (2.11) below, is open. Our techniques become more difficult to use in the general case; in fact we have not been able to find an expression for the kernel K in the reduced scalar nonlocal operator for cases beyond those considered in this work.

The paper is organized as follows: in the next section we specify notation and necessary technical lemmas. In section 2 we define the 3D PN model in the case that Γ is perpendicular to the plane of isotropy and calculate the reduced 2D model via the Dirichlet-to-Neumann map. Section 3 contains the computation and properties of the further reduced 1D nonlocal model in the case that W depends only on u_1 , and section 4 contains the same in the case that W

depends only on u_3 . In section 5 the 3D PN model for the parallel setting is derived, and also contains the dimension reductions, computation and properties of the scalar equation. Finally in section 6 we conclude that, for each of the three reduced problems, bounded stable solutions have 1D profiles, and thus complete the proof of propositions 2.4, 2.5 and 5.2.

1.1. Notation and preliminaries

For $d \geq 2$, denote the Schwarz class as $\mathcal{S}(\mathbb{R}^d)$ and its dual as $\mathcal{S}'(\mathbb{R}^d)$, and denote the homogeneous Sobolev spaces (see [10]) as

$$H^s(\mathbb{R}^d) := \left\{ u \in \mathcal{S}'(\mathbb{R}^d) : \int_{\mathbb{R}^d} |\xi|^{2s} |\hat{u}(\xi)|^2 d\xi < \infty \right\}.$$

The Fourier transform for $L^1(\mathbb{R}^2)$ functions v is written as

$$\hat{v}(k) = \mathcal{F}v(k) = \int_{\mathbb{R}^2} e^{-ix \cdot k} v(x) dx,$$

with inverse denoted by

$$v^\vee(k) = \mathcal{F}^{-1}v(k) = (2\pi)^{-2} \int_{\mathbb{R}^2} e^{ix \cdot k} v(x) dx.$$

Lemma 1.1. *Let $\rho > 0$. The integral representation of the square root of the operator $(-\Delta)_\rho := (-\partial_{x_1}^2 - \rho \partial_{x_2}^2)$ acting on sufficiently smooth $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ is*

$$(-\Delta_\rho)^{1/2} u(x) := -\frac{1}{4\pi\sqrt{\rho}} \int_{\mathbb{R}^2} \frac{u(x-y) + u(x+y) - 2u(x)}{\left(y_1^2 + \frac{y_2^2}{\rho}\right)^{3/2}} dy,$$

and the associated Fourier symbol is $\sqrt{k_1^2 + \rho k_2^2}$.

See for instance [10] for the case $\rho = 1$; the proof for general $\rho > 0$ is similar.

2. Slip plane perpendicular to the plane of isotropy: the 3D model

For this section we introduce the 3D PN model for transversely anisotropic elasticity, and then in section 2.1 we reduce it to a 2D nonlocal model via the Dirichlet-to-Neumann map. The first two main results (propositions 2.4 and 2.5) for the case of the slip plane perpendicular to the plane of isotropy are summarized in this section. The proofs of these two results will be completed in section 6.

In the PN model, the two half spaces of the elastic medium are separated by the slip plane Γ . Let $\mathbf{u} = (u_1, u_2, u_3)$ be the displacement vector. The total energy of the system is

$$E(\mathbf{u}) := E_{\text{els}}(\mathbf{u}) + E_{\text{mis}}(\mathbf{u}). \quad (2.1)$$

The material properties of transversely isotropic materials are determined by five independent elastic constants C_{11} , C_{13} , C_{33} , C_{44} , and C_{66} . These constants determine the stress–strain relation comprising the total elastic energy $E_{\text{els}}(\mathbf{u})$ for the two half-spaces. To be precise, $E_{\text{els}}(\mathbf{u})$ is defined as

$$E_{\text{els}}(\mathbf{u}) = \frac{1}{2} \int_{\mathbb{R}^3 \setminus \Gamma} \sigma : \varepsilon dx = \frac{1}{2} \int_{\mathbb{R}^3 \setminus \Gamma} \sigma_{ij} \varepsilon_{ij} dx, \quad (2.2)$$

where ε_{ij} is the strain tensor

$$\varepsilon_{ij} = \frac{1}{2} (\partial_j u_i + \partial_i u_j), \quad \text{for } i, j = 1, 2, 3, \quad \partial_i := \frac{\partial}{\partial x_i},$$

and σ is the stress tensor in $\mathbb{R}^{3 \times 3}$ given by

$$\begin{aligned} \sigma_{11} &= C_{11}\varepsilon_{11} + (C_{11} - 2C_{66})\varepsilon_{22} + C_{13}\varepsilon_{33}, \\ \sigma_{22} &= (C_{11} - 2C_{66})\varepsilon_{11} + C_{11}\varepsilon_{22} + C_{13}\varepsilon_{33}, \\ \sigma_{33} &= C_{13}\varepsilon_{11} + C_{13}\varepsilon_{22} + C_{33}\varepsilon_{33}, \\ \sigma_{23} &= \sigma_{32} = C_{44}\varepsilon_{23}, \\ \sigma_{13} &= \sigma_{31} = C_{44}\varepsilon_{13}, \\ \sigma_{12} &= \sigma_{21} = C_{66}\varepsilon_{12}. \end{aligned} \quad (2.3)$$

We are using Einstein summation notation in (2.2):

$$\sigma_{ij}\varepsilon_{ij} = \sum_{i,j=1}^3 \sigma_{ij}\varepsilon_{ij}.$$

The five elastic moduli defining σ are assumed to satisfy the uniform ellipticity conditions (see [13, 17])

$$0 < C_{66} < C_{11}, \quad C_{13}^2 < C_{33}(C_{11} - C_{66}), \quad C_{44} > 0. \quad (2.4)$$

Note that one can recover the PN model for fully isotropic media by setting

$$C_{11} = C_{33} = 2\mu \frac{1-\nu}{1-2\nu}, \quad C_{13} = 2\mu \frac{\nu}{1-2\nu}, \quad C_{44} = C_{66} = \mu, \quad (2.5)$$

where μ is the shear modulus and ν is the Poisson ratio.

In this first setting, Γ is taken perpendicular to the plane of isotropy, that is, $\Gamma = \{(x_1, x_2, x_3) : x_2 = 0\}$; see figure 1. The misfit energy $E_{\text{mis}}(\mathbf{u})$ across the slip plane due to atomistic reactions is therefore given by

$$E_{\text{mis}}(\mathbf{u}) := \int_{\Gamma} \gamma(u_1^+ - u_1^-, u_3^+ - u_3^-) dS = \int_{\Gamma} W(u_1^+, u_3^+) dS,$$

where

$$u_i^{\pm}(x_1, x_3) := u_i(x_1, 0^{\pm}, x_3) \text{ for } i = 1, 3.$$

We assume that the nonlinear potential W is twice differentiable with Hölder-continuous derivatives, i.e. $W \in C_b^{2,a}(\mathbb{R}^2; \mathbb{R})$ for some $a \in (0, 1)$.

The equilibrium structure of a dislocation is obtained by minimizing (2.2) subject to the following conditions at the slip plane

$$\begin{aligned} u_1^+(x_1, x_3) &= -u_1^-(x_1, x_3), \\ u_2^+(x_1, x_3) &= u_2^-(x_1, x_3), \\ u_3^+(x_1, x_3) &= -u_3^-(x_1, x_3). \end{aligned} \quad (2.6)$$

We will also consider two different scenarios for the form of W ; we will treat the cases $\partial_{u_1} W = 0$ and $\partial_{u_3} W = 0$ separately in sections 3 and 4. In the former case we assume the bi-states far field boundary condition for u_1

$$u_1^{\pm}(\pm\infty, x_3) = \pm 1 \quad \text{for any } x_3 \in \mathbb{R}, \quad (2.7)$$

and in the latter case we assume a similar condition for u_3

$$u_3^+(x_1, \pm\infty) = \pm 1 \quad \text{for any } x_1 \in \mathbb{R}. \quad (2.8)$$

We remark that the Dirichlet-type boundary condition, for instance $u_1(x_1, x_3) = \pm 1$ at $x_1 = \pm r$ for some fixed r , can also be considered. However, the truncated dislocation profile can only be regarded as an approximation for the displacement profile in the entire space, since after constant extension outside a given bounded domain, the total energy for such a truncated dislocation profile is larger than the true minimizer defined via definition 2.2.

Formally, the elastic energy is infinite due to the slow decay rate of ε . We therefore define a minimizer via perturbation. To be precise, for any $\varphi \in C^\infty(\mathbb{R}^3 \setminus \Gamma; \mathbb{R}^3)$ with compact support in a Euclidean ball $B_R(0) \subset \mathbb{R}^3$ we define the perturbed elastic energy

$$\begin{aligned} \hat{E}_{\text{els}}(\varphi|\mathbf{u}) &:= \int_{\mathbb{R}^3 \setminus \Gamma} \frac{1}{2} (\sigma_{\mathbf{u}} + \sigma_{\varphi}) : (\varepsilon_{\mathbf{u}} + \varepsilon_{\varphi}) - \frac{1}{2} \sigma_{\mathbf{u}} : \varepsilon_{\mathbf{u}} \, dx \\ &= \int_{\mathbb{R}^3 \setminus \Gamma} \frac{1}{2} \left((\sigma_{\varphi})_{ij} (\varepsilon_{\varphi})_{ij} + (\sigma_{\mathbf{u}})_{ij} (\varepsilon_{\varphi})_{ij} + (\sigma_{\varphi})_{ij} (\varepsilon_{\mathbf{u}})_{ij} \right) \, dx \\ &:= E_{\text{els}}(\varphi) + \mathcal{C}_{\text{els}}(\mathbf{u}, \varphi), \end{aligned}$$

where the cross term is defined as

$$\mathcal{C}_{\text{els}}(\mathbf{u}, \varphi) := \int_{\mathbb{R}^3 \setminus \Gamma} \frac{1}{2} (\sigma_{\mathbf{u}} : \varepsilon_{\varphi} + \sigma_{\varphi} : \varepsilon_{\mathbf{u}}) \, dx = \int_{\mathbb{R}^3 \setminus \Gamma} \frac{1}{2} \left((\sigma_{\mathbf{u}})_{ij} (\varepsilon_{\varphi})_{ij} + (\sigma_{\varphi})_{ij} (\varepsilon_{\mathbf{u}})_{ij} \right) \, dx$$

and $\sigma_{\mathbf{u}}, \varepsilon_{\mathbf{u}}$ and $\sigma_{\varphi}, \varepsilon_{\varphi}$ are the stress and strain tensors corresponding to $\mathbf{u}$ and φ , respectively. We also define the perturbed misfit energy as

$$\hat{E}_{\text{mis}}(\varphi|\mathbf{u}) := \int_{\Gamma} W(u_1^+ + \varphi_1^+, u_3^+ + \varphi_3^+) - W(u_1^+, u_3^+) \, dS.$$

The perturbed total energy is then defined as

$$\hat{E}(\varphi|\mathbf{u}) := \hat{E}_{\text{els}}(\varphi|\mathbf{u}) + \hat{E}_{\text{mis}}(\varphi|\mathbf{u}).$$

Remark 2.1. Since $\mathbf{u}$ and $\mathbf{u} + \varphi$ coincide outside of B_R , the perturbed elastic energy $\hat{E}(\varphi|\mathbf{u})$ is equivalent to the local perturbed elastic energy

$$\hat{E}_{\text{els}}(\varphi|\mathbf{u}; B_R) := \int_{B_R \setminus \Gamma} \frac{1}{2} (\sigma_{\mathbf{u}} + \sigma_{\varphi}) : (\varepsilon_{\mathbf{u}} + \varepsilon_{\varphi}) - \frac{1}{2} \sigma_{\mathbf{u}} : \varepsilon_{\mathbf{u}} \, dx := E_{\text{els}}(\mathbf{u} + \varphi; B_R) - E_{\text{els}}(\mathbf{u}; B_R),$$

and similarly for the misfit energy

$$\hat{E}_{\text{mis}}(\varphi|\mathbf{u}; B_R) := \int_{B_R \cap \Gamma} W(u_1^+ + \varphi_1^+, u_3^+ + \varphi_3^+) - W(u_1^+, u_3^+) \, dS := E_{\text{mis}}(\mathbf{u} + \varphi; B_R) - E_{\text{mis}}(\mathbf{u}; B_R),$$

so that $\hat{E}(\varphi|\mathbf{u})$ is equivalent to

$$\hat{E}(\varphi|\mathbf{u}; B_R) := \hat{E}_{\text{els}}(\varphi|\mathbf{u}; B_R) + \hat{E}_{\text{mis}}(\varphi|\mathbf{u}; B_R) := E(\mathbf{u} + \varphi; B_R) - E(\mathbf{u}; B_R).$$

Hence, we say that $\mathbf{u}$ is a local minimizer, following the convention in [7].

Definition 2.2. We call $\mathbf{u}$ a local minimizer of E if $\mathbf{u}$ satisfies

$$E(\mathbf{u} + \varphi; B_R) - E(\mathbf{u}; B_R) \geq 0$$

for any $\varphi \in C^\infty(\mathbb{R}^3 \setminus \Gamma; \mathbb{R}^3)$ with compact support in a Euclidean ball $B_R(0) \subset \mathbb{R}^3$ satisfying the conditions

$$\begin{aligned}\varphi_1^+(x_1, x_3) &= -\varphi_1^-(x_1, x_3), \\ \varphi_2^+(x_1, x_3) &= \varphi_2^-(x_1, x_3), \\ \varphi_3^+(x_1, x_3) &= -\varphi_3^-(x_1, x_3).\end{aligned}\tag{2.9}$$

In order to define the Euler–Lagrange equation associated to the total energy E we need to set some notation. We define $\mathbb{L}\mathbf{u}$ to be the second-order partial differential system

$$\mathbb{L}\mathbf{u} := \operatorname{div} \sigma, \quad (\mathbb{L}\mathbf{u})_i = L_{ij}u_j, \quad i = 1, 2, 3,$$

where L_{ij} with $i, j = 1, 2, 3$, is the matrix of partial derivatives

$$\begin{aligned}L_{11} &= C_{11}\partial_{11} + C_{66}\partial_{22} + C_{44}\partial_{33}, \\ L_{22} &= C_{66}\partial_{11} + C_{11}\partial_{22} + C_{44}\partial_{33}, \\ L_{33} &= C_{44}\partial_{11} + C_{44}\partial_{22} + C_{33}\partial_{33}, \\ L_{23} &= L_{32} = (C_{13} + C_{44})\partial_{23}, \\ L_{13} &= L_{31} = (C_{13} + C_{44})\partial_{13}, \\ L_{12} &= L_{21} = (C_{11} - C_{66})\partial_{12}.\end{aligned}$$

Lemma 2.3. Assume that $\mathbf{u} \in C^2(\mathbb{R}^3 \setminus \Gamma; \mathbb{R}^3)$ satisfying (2.6) and either (2.7) or (2.8) is a local minimizer of the total energy E in the sense of definition 2.2. Then $\mathbf{u}$ satisfies the Euler–Lagrange equations

$$\begin{aligned}\mathbb{L}\mathbf{u} &= 0 \text{ in } \mathbb{R}^3 \setminus \Gamma, \\ \sigma_{12}^+ + \sigma_{12}^- &= \partial_{u_1} W(u_1^+, u_3^+) \text{ on } \Gamma, \\ \sigma_{32}^+ + \sigma_{32}^- &= \partial_{u_3} W(u_1^+, u_3^+) \text{ on } \Gamma, \\ \sigma_{22}^+ - \sigma_{22}^- &= 0 \text{ on } \Gamma.\end{aligned}\tag{2.10}$$

Proof. The computation is similar to its analogue in the case of fully isotropic elasticity; see [4, appendix A] for the isotropic case. $\square$

The analysis of the reduced models changes significantly depending on the choice of the elastic coefficients. We leave more general cases to future works, but here we make the choices

$$\sqrt{C_{11}C_{33}} - C_{13} - 2C_{44} = 0, \quad \text{and} \quad C_{11} = C_{33}.\tag{2.11}$$

The first relation greatly simplifies the analysis, while the second relation further simplifies the presentation of the results for the case of Γ perpendicular to the plane of isotropy.

Our first two main results are the following:

Proposition 2.4 (case I: Γ perpendicular to the plane of isotropy, $\partial_{u_3} W = 0$). Suppose that the five elastic constants satisfy (2.4) and (2.11), and assume that the quantities

$$\nu := \frac{C_{13}}{2(C_{13} + C_{44})}, \quad \delta := \frac{C_{66}}{C_{44}}, \quad \mu := C_{44}\tag{2.12}$$

satisfy

$$\delta > 0 \quad \text{and} \quad \max \left\{ 1 - \frac{2}{\delta}, \frac{\sqrt{\delta}}{2} \frac{2\delta - 3}{2\delta^{3/2} - 2\delta^{1/2} + 1} \right\} < \nu < \frac{2}{4 - \delta + \sqrt{\delta^2 + 8}}.\tag{2.13}$$

Assume that $\partial_{u_3} W = 0$, i.e. $W(u_1, u_3) = W(u_1)$, and that

$$\begin{aligned} &W \text{ is a periodic / double-well potential in } C_b^{2,a}(\mathbb{R}) \text{ satisfying} \\ &W(v) > W(\pm 1) \text{ for } v \in (-1, 1), \quad W''(\pm 1) > 0. \end{aligned} \quad (2.14)$$

Then the system (2.10) has a unique (up to translation in the x_1 -direction and rotation about the x_2 -axis) classical solution $\mathbf{u} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ belonging to $C^\infty(\mathbb{R}^3 \setminus \Gamma)$ with $\mathbf{u}(\cdot, x_2, \cdot) \in \dot{H}^s(\mathbb{R}^2)$ for all $s \geq 1$ and for all fixed $x_2 \neq 0$. Moreover, the solution is a local minimizer of E in the sense of definition 2.2. The first component u_1 is the unique stable solution to (3.3) with specific nonlocal kernel defined in (3.14) and u_1 has a 1D profile.

Proposition 2.5 (case II: Γ perpendicular to the plane of isotropy, $\partial_{u_1} W = 0$). Suppose that the five elastic constants satisfy (2.4) and (2.11). Define the quantities μ , ν and δ as in (2.12), and define

$$p := \delta(2(1 - \nu) - \delta(1 - 2\nu)) \quad \text{and} \quad q := 1 - \nu. \quad (2.15)$$

Suppose that

$$(\nu, \delta) \in \mathcal{U}^+, \quad (2.16)$$

where $\mathcal{U}^+ \subset \mathbb{R}^2$ is the open region enclosed by the curves $\nu = \frac{1}{2}$, $\nu = \frac{\delta(\delta-2)}{2(\delta^2-\delta+1)}$, $\nu = 1 - \frac{2}{\delta}$, $\nu = \frac{2}{1+4\delta-2\delta^2+\sqrt{1-8\delta+28\delta^2-16\delta^3+4\delta^4}}$, and $p = \tilde{r}$, where $\tilde{r}(q)$ is the smallest positive root of the polynomial

$$\begin{aligned} x \mapsto &-8(-1+q)x^4 + 8(-1+q)q^3 + q^2(13+14q-11q^2)x \\ &+ (-11+14q+13q^2)x^3 + 2q(1-18q+q^2)x^2. \end{aligned}$$

Assume that $\partial_{u_1} W = 0$, i.e. $W(u_1, u_3) = W(u_3)$ and that W satisfies (2.14). Then the system (2.10) has a unique (up to translation in the x_3 -direction and rotation about the x_2 -axis) classical solution $\mathbf{u} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ belonging to $C^\infty(\mathbb{R}^3 \setminus \Gamma)$ with $\mathbf{u}(\cdot, x_2, \cdot) \in \dot{H}^s(\mathbb{R}^2)$ for all $s \geq 1$ and for all fixed $x_2 \neq 0$. Moreover, the solution is a local minimizer of E in the sense of definition 2.2. The third component u_3 is the unique stable solution to (4.2) with specific nonlocal kernel defined in (4.7) and u_3 has a 1D profile.

Note that (2.11) and (2.12) imply that

$$C_{11} = C_{33} = 2\mu \frac{1-\nu}{1-2\nu}, \quad C_{13} = 2\mu \frac{\nu}{1-2\nu}, \quad C_{44} = \mu, \quad C_{66} = \delta\mu.$$

The case $\delta = 1$ corresponds to the case of isotropic elasticity, and μ and ν respectively correspond to the shear modulus and Poisson's ratio of an isotropic solid.

The conditions in (2.4) for uniform ellipticity are equivalent to the conditions

$$0 < \mu, \quad 0 < \delta < 4, \quad 1 - \frac{2}{\delta} < \nu < \frac{1}{2}. \quad (2.17)$$

Note that δ can be arbitrarily close to 0, so there is no lower bound on ν ; this is expected behaviour of Poisson's ratios in anisotropic media [20]. It is straightforward to check that the regions in the (ν, δ) -plane described by (2.13) and $\mathcal{U}^+$ are subsets of the region defined by (2.17).

2.1. Reduction of the 3D system to a nonlocal 2D system

We can reduce to a nonlocal 2D system that depends only on u_1^+ and u_3^+ by using the relations in (2.10) satisfied by the Dirichlet-to-Neumann maps $(-\sigma_{12}^+, -\sigma_{22}^+, -\sigma_{32}^+)$ and $(\sigma_{12}^-, \sigma_{22}^-, \sigma_{32}^-)$

for the upper and lower half-spaces respectively. We calculate this map using the Fourier transform. Here, $x = (x_1, x_3)$ and $k = (k_1, k_3)$, so that statements for the slip plane inherit the conventions used for the full 3D system. We state the dimension reduction in the following lemma.

Lemma 2.6. *Let C_{11} , C_{13} , C_{33} , C_{44} and C_{66} satisfy (2.4) and (2.11), and define μ , ν and δ as in (2.12). Suppose that $\mathbf{u}$ satisfies (2.10) and remains bounded as $|x_2| \rightarrow \infty$, and assume that u_1^+ and u_3^+ belong to the fractional Sobolev space $H^s(\mathbb{R}^2)$ for some $s \geq 1/2$. Then $\mathbf{u}$ can be expressed entirely in terms of u_1^+ and u_3^+ . In particular, (u_1^+, u_3^+) satisfies the nonlocal system*

$$\begin{bmatrix} \mathcal{F}(\sigma_{12}^+(k) + \sigma_{12}^-(k)) \\ \mathcal{F}(\sigma_{32}^+(k) + \sigma_{32}^-(k)) \end{bmatrix} := -\mathbb{A}(k) \begin{bmatrix} \hat{u}_1^+(k) \\ \hat{u}_3^+(k) \end{bmatrix} = \mathcal{F} \begin{bmatrix} \partial_{u_1} W(u_1^+, u_3^+) \\ \partial_{u_3} W(u_1^+, u_3^+) \end{bmatrix} \quad \text{on } \Gamma, \quad (2.18)$$

where the 2×2 matrix $\mathbb{A}(k)$ is given by

$$\mathbb{A}(k) := \frac{2\mu}{1-\nu} r_1 \cdot \begin{bmatrix} \frac{\delta(2(1-\nu) - \delta(1-2\nu))k_1^2 + (1-\nu)k_3^2}{r_1^2} & \frac{\delta\nu k_1 k_3}{r_1^2} \\ \frac{k_1 k_3}{\delta r_1^2} & \frac{(\nu\delta + (1-\nu)(1-\delta))r_2 + (\nu\delta^2 - (1-\nu)(1-\delta)^2)r_1}{r_1 + r_2} \end{bmatrix}, \quad (2.19)$$

and

$$r_1 := r_1(k) := \sqrt{k_1^2 + k_3^2/\delta} \quad \text{and} \quad r_2 := r_2(k) := \sqrt{k_1^2 + k_3^2}. \quad (2.20)$$

Note that when $\delta = 1$ the matrix $\mathbb{A}$ is consistent with the Fourier transform of the Dirichlet-to-Neumann map in the setting of dislocations in fully isotropic media, see [4].

Proof. Dividing the system by C_{44} , taking the Fourier transform of $\mathbb{L}\mathbf{u} = 0$ in the (x_1, x_3) variables, and using (2.11) and (2.12) gives the set of three equations

$$\begin{aligned} -\left(\frac{2(1-\nu)}{1-2\nu}k_1^2 + k_3^2\right)\hat{u}_1 + \delta\partial_{22}\hat{u}_1 + \left(\frac{2(1-\nu)}{1-2\nu} - \delta\right)(ik_1)\partial_2\hat{u}_2 - \frac{1}{1-2\nu}(k_1k_3)\hat{u}_3 &= 0, \\ \left(\frac{2(1-\nu)}{1-2\nu} - \delta\right)(ik_1)\partial_2\hat{u}_1 - (\delta k_1^2 + k_3^2)\hat{u}_2 + \frac{2(1-\nu)}{1-2\nu}\partial_{22}\hat{u}_2 + \frac{1}{1-2\nu}(ik_3)\partial_2\hat{u}_3 &= 0, \\ -\frac{1}{1-2\nu}(k_1k_3)\hat{u}_1 + \frac{1}{1-2\nu}(ik_3)\partial_2\hat{u}_2 - \left(k_1^2 + \frac{2(1-\nu)}{1-2\nu}k_3^2\right)\hat{u}_3 + \partial_{22}\hat{u}_3 &= 0. \end{aligned}$$

This is a 3×3 system of second-order ODEs in x_2 . We solve this system in the upper and lower half-spaces separately.

To solve this system in the upper half-space, we write it as a 6×6 first-order system

$$\partial_2 \begin{bmatrix} \hat{\mathbf{u}}(k, x_2) \\ \partial_2 \hat{\mathbf{u}}(k, x_2) \end{bmatrix} = \mathfrak{A}(k) \begin{bmatrix} \hat{\mathbf{u}}(k, x_2) \\ \partial_2 \hat{\mathbf{u}}(k, x_2) \end{bmatrix} \quad (2.21)$$

where the 6×6 matrix $\mathfrak{A}$ depends on k , μ , ν and δ , with boundary conditions

$$\hat{\mathbf{u}}(k_1, k_3, 0) = \hat{\mathbf{u}}^+(k_1, k_3), \quad \hat{\mathbf{u}}(k, +\infty) = 0.$$

Thanks to the conditions (2.4) and (2.11), the characteristic polynomial of $\mathfrak{A}(k)$ has six real nonzero roots $\pm r_1(k)$ and $\pm r_2(k)$ as in (2.20), and $\pm r_2$ both have multiplicity 2. So the general solution to (2.21) is an arbitrary linear combination of six terms, each of which is a product of a (generalized) eigenvector and $e^{\pm r_i(k)x_2}$, where $i = 1, 2$. The computation is straightforward.

Since $\mathbf{u}$ remains bounded as $|x_2| \rightarrow \infty$, we exclude the solutions with positive roots from the final solution, i.e. three of the coefficients in the linear combination are zero. The remaining three coefficients are completely determined by the boundary conditions at $x_2 = 0$, and so the unique solution to (2.21) has the form $\hat{\mathbf{u}}(k, x_2) = \mathbb{B}^+(k, x_2)\hat{\mathbf{u}}^+(k)$ for $x_2 > 0$, for a 3×3 matrix

$\mathbb{B}^+$. The form of the matrix $\mathbb{B}^+$ is determined only from the properties of the eigenvalues and eigenvectors of $\mathfrak{A}$.

In exactly a similar way, the unique solution to the system (2.21) for the lower half-space with boundary conditions

$$\hat{\mathbf{u}}(k_1, k_3, 0) = \hat{\mathbf{u}}^-(k_1, k_3), \quad \hat{\mathbf{u}}(k, -\infty) = 0,$$

has the form $\hat{\mathbf{u}}(k, x_2) = \mathbb{B}^-(k, x_2)\hat{\mathbf{u}}^-$ for $x_2 < 0$, for some 3×3 matrix $\mathbb{B}^-$. In fact, $\mathbb{B}^-(k, x_2) = \overline{\mathbb{B}^+(k, -x_2)}$, where $\bar{z}$ is the complex conjugate of z .

Next, we use the slip plane boundary conditions (2.6) to see that $\hat{\mathbf{u}}(k, t)$ depends only on $\hat{u}_1^+$, $\hat{u}_2^+$, and $\hat{u}_3^+$ for all $k \in \mathbb{R}^2$ and all $x_2 \in \mathbb{R}$. Finally, the equation $\mathcal{F}(\sigma_{22}^+ - \sigma_{22}^-) = 0$ in (2.10) allows us to write $\hat{u}_2^+$ in terms of $\hat{u}_1^+$ and $\hat{u}_3^+$, and inserting this relation into the other two equations involving the Dirichlet-to-Neumann map in (2.10) gives the expression (2.19) for the matrix $\mathbb{A}(k)$. □

The matrix $\mathbb{A}$ is positive-definite for $k \neq 0$ thanks to the uniform ellipticity conditions (2.4). Thus by Plancherel's identity

$$C^{-1} \|(u_1, u_3)\|_{\dot{H}^{\frac{1}{2}}(\Gamma)}^2 \leq \int_{\Gamma} \langle \mathbb{A}(k) (\hat{u}_1, \hat{u}_3), (\hat{u}_1, \hat{u}_3) \rangle dk_1 dk_3 \leq C \|(u_1, u_3)\|_{\dot{H}^{\frac{1}{2}}(\Gamma)}^2.$$

In the next two sections we treat Cases I and II as described in the introduction. In case I, W depends only on u_1^+ , and in case II, W depends only on u_3^+ . In both cases the simplification allows us to further reduce the 2D equation to a one-dimensional problem.

3. The 1D reduced equation in the perpendicular case: $\partial_{u_3} W = 0$

Under the assumption that $W(u_1, u_3) = W(u_1)$, we derive the reduced scalar equation for u_1^+ , which is an anisotropic nonlocal Ginzburg-Landau equation. We will derive the nonlocal kernel representation for this anisotropic nonlocal equation in proposition 3.1. Then we study the explicit formula for the nonlocal kernel and prove the positivity of this kernel in propositions 3.4 and 3.6.

At this point, we drop the superscripts on u_1^+ , u_3^+ for simplicity. However, we retain the index convention for the spatial and frequency variables $x = (x_1, x_3) \in \mathbb{R}^2$ and $k = (k_1, k_3) \in \mathbb{R}^2$. Our reduced nonlocal system reads

$$\mathbb{A}(k) \begin{bmatrix} \hat{u}_1(k) \\ \hat{u}_3(k) \end{bmatrix} = -\mathcal{F} \begin{bmatrix} \partial_{u_1} W(u_1, u_3) \\ \partial_{u_3} W(u_1, u_3) \end{bmatrix} \quad \text{on } \Gamma. \quad (3.1)$$

Since $\partial_{u_3} W = 0$, we can solve for $\hat{u}_3$ in terms of $\hat{u}_1$ and substitute into the remaining equation. We obtain the one-dimensional scalar anisotropic nonlocal equation for $\hat{u}_1$ in Fourier space

$$\mathcal{F}(\mathcal{L}u_1)(k_1, k_3) := \frac{\det \mathbb{A}(k)}{a_{22}(k)} \hat{u}_1(k) = -\mathcal{F}(W'(u_1)). \quad (3.2)$$

Therefore, we invert the Fourier transform and obtain a new 1D nonlocal equation

$$\begin{aligned} \mathcal{L}u_1(x_1, x_3) &= -W'(u_1(x_1, x_3)), \quad (x_1, x_3) \in \Gamma, \\ \lim_{x_1 \rightarrow \pm\infty} u_1(x_1, x_3) &= \pm 1, \quad x_3 \in \mathbb{R}, \end{aligned} \quad (3.3)$$

where the nonlocal operator $\mathcal{L}$ has the Fourier symbol

$$\tilde{m}(k) := \frac{2\mu r_2(pk_1^2 + k_3^2)}{r_1 r_2 - \nu k_1^2} \quad (3.4)$$

with p defined in (2.15). This symbol has a ‘doubly nonlocal’ nature, consisting of products of square roots of anisotropic symbols. Since $\nu < 1/2$ we have

$$2\mu \min\{1, p\} r_2(k) \leq \tilde{m}(k) \leq \mu C(\delta, \nu) r_2(k) \quad \text{for all } k \in \mathbb{R}^2.$$

We now derive the integral formulation of $\mathcal{L}$ in $\mathcal{S}'(\mathbb{R}^2)$. The integral formulation for $\mathcal{L}$ agrees with the Fourier inversion formula whenever the function it acts on is sufficiently smooth, for instance $u \in \dot{H}^s(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ for $s \geq 1$.

Proposition 3.1. *Suppose that K_1 and K_2 are the unique C^∞ solutions in $\mathbb{R}^2 \setminus \{0\}$ of the partial differential equations*

$$(\delta(1-\nu^2)\partial_1^4 + (1+\delta)\partial_1^2\partial_3^2 + \partial_3^4)K_1(x) = (\partial_1^2 + \partial_3^2)(p\partial_1^2 + \partial_3^2)\left(\frac{1}{(x_1^2 + \delta x_3^2)^{3/2}}\right) \quad (3.5)$$

and

$$(\delta(1-\nu^2)\partial_1^4 + (1+\delta)\partial_1^2\partial_3^2 + \partial_3^4)K_2(x) = (p\partial_1^4 + \partial_1^2\partial_3^2)\left(\frac{1}{(x_1^2 + x_3^2)^{3/2}}\right). \quad (3.6)$$

Then the integral form of the operator $\mathcal{L}$ defined in (3.2) is

$$\mathcal{L}w(x) = -\frac{1}{4\pi} \int_{\mathbb{R}^2} (w(x-y) + w(x+y) - 2w(x))K(y) dy, \quad (3.7)$$

where $K(x) := 2\mu\delta(\sqrt{\delta}K_1(x) + \nu K_2(x))$.

Proof. We want the denominator of $\tilde{m}$ to be a local operator so that we can integrate by parts in the half-Laplacian definition. If we rationalize the denominator, then $\tilde{m}$ remains in $\mathcal{S}'(\mathbb{R}^2)$, and the Fourier symbol becomes

$$\tilde{m}(k) = 2\mu\delta \frac{m_n(k)}{m_d(k)},$$

where the symbols $m_n(k)$ and $m_d(k)$ are defined as

$$m_n(k) := r_2(pk_1^2 + k_3^2)(r_1r_2 + \nu k_1^2)$$

and

$$m_d(k) := ck_1^4 + bk_1^2k_3^2 + k_3^4 \quad \text{with} \quad c := \delta(1-\nu^2), \quad b := (1+\delta). \quad (3.8)$$

Here m_d is a local operator, and m_n is a sum of compositions of local operators with square-root operators.

Separate m_n into two pieces: $m_n = m_{n,1} + \nu m_{n,2}$ where

$$\begin{aligned} m_{n,1} &:= r_2^2(pk_1^2 + k_3^2)r_1, \\ m_{n,2} &:= k_1^2(pk_1^2 + k_3^2)r_2. \end{aligned}$$

Note that, as functions of variables k_1^2 and k_3^2 , both $\frac{m_{n,1}}{r_1}$ and $\frac{m_{n,2}}{r_2}$ are homogeneous second-order polynomials, hence define anisotropic biharmonic-type differential operators.

Define L_n , $L_{n,1}$ and $L_{n,2}$ to be the operators whose Fourier symbols are m_n , $\frac{m_{n,1}}{r_1}$ and $\frac{m_{n,2}}{r_2}$ respectively. Therefore for any $w \in \mathcal{S}(\mathbb{R}^2)$

$$\mathcal{F}(L_n w) = \mathcal{F}\left(L_{n,1} \circ \left(-\Delta_{\frac{1}{\delta}}\right)^{1/2} w + \nu L_{n,2} \circ (-\Delta)^{1/2} w\right) = (m_{n,1}(k) + \nu m_{n,2}(k))\hat{w} = m_n(k)\hat{w}$$

in $\mathcal{S}'(\mathbb{R}^2)$, where the anisotropic fractional Laplacian has been defined in lemma 1.1.

Define L_d to be the operator whose symbol is m_d . By using a cutoff function near the origin and using the dominated convergence theorem, we may assume that $\hat{w}$ vanishes near the origin. Now define $v \in \mathcal{S}'(\mathbb{R}^2)$ by $w = L_d v$. Thus $\hat{v} = \frac{\hat{w}}{m_d(k)}$. So, we have

$$\begin{aligned}\mathcal{L}w &:= 2\mu\delta(\mathcal{L}_1 w + \nu\mathcal{L}_2 w) \\ &= 2\mu\delta\left(\left(-\Delta_{\frac{1}{\delta}}\right)^{1/2} L_{n,1} L_d^{-1} w + \nu(-\Delta)^{1/2} L_{n,2} L_d^{-1} w\right) \\ &= 2\mu\delta\left(\left(-\Delta_{\frac{1}{\delta}}\right)^{1/2} L_{n,1} v + \nu(-\Delta)^{1/2} L_{n,2} v\right).\end{aligned}$$

Treat $\mathcal{L}_1$ and $\mathcal{L}_2$ separately. By the representation formula in lemma 1.1

$$\begin{aligned}\mathcal{L}_1 w(x) &= -\frac{\sqrt{\delta}}{4\pi} \int_{\mathbb{R}^2} \frac{(L_{n,1})_x v(x-y) + (L_{n,1})_x v(x+y) - 2(L_{n,1})_x v(x)}{(y_1^2 + \delta y_3^2)^{3/2}} dy \\ &= -\frac{\sqrt{\delta}}{4\pi} \int_{\mathbb{R}^2} \frac{(L_{n,1})_y \left(v(x-y) + v(x+y) - \left(\sum_{|a| \leq 4} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) \right)}{(y_1^2 + \delta y_3^2)^{3/2}} dy.\end{aligned}$$

Here we are using multi-index notation: $a = (a_1, a_3) \in \mathbb{N}_0^2$, $|a| = a_1 + a_3$, $a! = a_1! a_3!$, $y^a = y_1^{a_1} y_3^{a_3}$, and $D^a v(x) = \frac{\partial^{|a|}}{\partial x_1^{a_1} \partial x_3^{a_3}} v(x)$. Now integrating by parts (we can use integration by parts here without difficulty by using the P.V. definition of the integral operator and observe that the boundary term $\rightarrow 0$ as the P.V. limit is taken),

$$\begin{aligned}\mathcal{L}_1 w(x) &= -\frac{\sqrt{\delta}}{4\pi} \int_{\mathbb{R}^2} \left(v(x-y) + v(x+y) - \sum_{|a| \leq 4} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) (L_{n,1})_y \left(\frac{1}{(y_1^2 + \delta y_3^2)^{3/2}} \right) dy.\end{aligned}$$

Since $K_1(y)$ satisfies

$$L_d K_1(y) = L_{n,1} \left(\frac{1}{(y_1^2 + \delta y_3^2)^{3/2}} \right),$$

integrating by parts again gives

$$\begin{aligned}\mathcal{L}_1 w(x) &= -\frac{\sqrt{\delta}}{4\pi} \int_{\mathbb{R}^2} \left(v(x-y) + v(x+y) - \sum_{|a| \leq 4} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) (L_d)_y K_1(y) dy \\ &= -\frac{\sqrt{\delta}}{4\pi} \int_{\mathbb{R}^2} (L_d)_y \left(v(x-y) + v(x+y) - \sum_{|a| \leq 4} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) K_1(y) dy \\ &= -\frac{\sqrt{\delta}}{4\pi} \int_{\mathbb{R}^2} ((L_d)_x v(x-y) + (L_d)_x v(x+y) - 2(L_d)_x v(x)) K_1(y) dy \\ &= -\frac{\sqrt{\delta}}{4\pi} \int_{\mathbb{R}^2} (w(x-y) + w(x+y) - 2w(x)) K_1(y) dy.\end{aligned}$$

We treat $\mathcal{L}_2$ the same way:

$$\begin{aligned}\mathcal{L}_2 w(x) &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} \frac{(L_{n,2})_x v(x-y) + (L_{n,2})_x v(x+y) - 2(L_{n,2})_x v(x)}{(y_1^2 + y_3^2)^{3/2}} dy \\ &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} \frac{(L_{n,2})_y \left(v(x-y) + v(x+y) - \sum_{|a| \leq 4} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right)}{(y_1^2 + y_3^2)^{3/2}} dy.\end{aligned}\quad (3.9)$$

Integrating by parts,

$$\begin{aligned}\mathcal{L}_2 w(x) &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} \left(v(x-y) + v(x+y) - \sum_{|a| \leq 4} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) (L_{n,2})_y \left(\frac{1}{(y_1^2 + y_3^2)^{3/2}} \right) dy.\end{aligned}$$

Since $K_2(y)$ satisfies

$$L_d K_2(y) = L_{n,2} \left(\frac{1}{(y_1^2 + y_3^2)^{3/2}} \right),$$

integrating by parts again results in

$$\begin{aligned}\mathcal{L}_2 w(x) &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} \left(v(x-y) + v(x+y) - \sum_{|a| \leq 6} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) (L_d)_y K_2(y) dy \\ &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} (L_d)_y \left(v(x-y) + v(x+y) - \sum_{|a| \leq 6} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) K_2(y) dy \\ &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} ((L_d)_x v(x-y) + (L_d)_x v(x+y) - 2(L_d)_x v(x)) K_2(y) dy \\ &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} (w(x-y) + w(x+y) - 2w(x)) K_2(y) dy.\end{aligned}$$

□

3.1. The formula for the kernel

We now find the formula for the nonlocal kernel K in (3.7) by solving the PDEs for $K_1(x)$ and $K_2(x)$.

Lemma 3.2. *Let μ , δ and ν satisfy (2.17), define p and q as in (2.15), and define b and c as in (3.8). Then there exists a solution K_1 to the equation (3.5) given by*

$$K_1(x_1, x_3) = \frac{Ax_1^{12} + Bx_1^{10}x_3^2 + Cx_1^8x_3^4 + Dx_1^6x_3^6 + Ex_1^4x_3^8 + Fx_1^2x_3^{10} + Gx_3^{12}}{(x_1^2 + \delta x_3^2)^{\frac{3}{2}} (x_1^4 + bx_1^2x_3^2 + cx_3^2)^3}, \quad (3.10)$$

where

$$\begin{aligned}
 A &= \frac{-2b + \delta + 2p + 2}{\delta}, \\
 B &= \frac{3(2b^2 - 2b(\delta + p + 1) - 4c + 3\delta + 3\delta p + 4p)}{\delta}, \\
 C &= \frac{9b^2\delta + 6c(3b - 5\delta - 4p - 4) - 6b(\delta^2 + \delta + \delta p - p) + 3\delta(2\delta + 2\delta p + 11p)}{\delta}, \\
 D &= \frac{b^2(\delta(2\delta + 1) + (\delta + 2)p) - 2b(c(-13\delta + p + 1) + \delta(\delta + (\delta - 13)p))}{\delta} \\
 &\quad + \frac{20c^2 - 2c(\delta(10\delta + 23) + (23\delta + 10)p) + 20\delta^2 p}{\delta}, \\
 E &= \frac{-6c(b(-\delta^2 + \delta + \delta p + p) + \delta(4\delta + 4\delta p + 5p)) + 9b\delta p(b + 2\delta) + c^2(33\delta + 6p + 6)}{\delta}, \\
 F &= 6b^2\delta p - 6c(b(\delta + \delta p + p) + 2\delta p) + 3c^2(4\delta + 3p + 3), \\
 G &= c(c(2\delta + 2\delta p + p) - 2b\delta p).
 \end{aligned} \tag{3.11}$$

K_1 is the unique solution of (3.5) in $C^\infty(\mathbb{R}^2 \setminus \{0\})$, and K_1 satisfies the following:

- i) $K_1(x) = K_1(-x)$, $K_1(\rho x) = \rho^{-3}K_1(x)$ for $\rho > 0$.
- ii) $|x^a D^a K_1(x)| < \frac{C}{|x|^3}$ for any multi-index $a \in \mathbb{N}_0^2$ and any $x \neq 0$.

Proof. Recall the definitions of the operators L_d and $L_{n,1}$ from the proof of proposition 3.1. The right-hand side of (3.5) is equal to

$$G_1(x_1, x_3) := \frac{P_1(x_1, x_3)}{(x_1^2 + \delta x_3^2)^{\frac{11}{2}}},$$

where P_1 is the fourth-degree polynomial

$$\begin{aligned}
 P_1(x_1, x_3) &:= 45(d_1(\nu, \delta)x_1^4 + d_2(\nu, \delta)x_1^2x_3^2 + d_3(\nu, \delta)x_3^4), \text{ with} \\
 d_1(\nu, \delta) &:= 8p - 2\delta(1 + p) + \delta^2, \\
 d_2(\nu, \delta) &:= \delta(-12p + 17(1 + p)\delta - 12\delta^2), \\
 d_3(\nu, \delta) &:= \delta^2(p - \delta^2(1 + p) + 8\delta^2).
 \end{aligned}$$

Our guess for a solution of (3.5) is (3.10), with constants A through G to be determined. Applying L_d to this ansatz, we obtain a function of the form $K_1(x_1, x_3) = \frac{\Pi(x_1, x_3)}{(x_1^2 + \delta x_3^2)^{\frac{11}{2}}}$, where Π is a polynomial with terms $x_1^{2i}x_3^{2(12-i)}$, where $i \in \{0, 1, \dots, 12\}$. Multiplying the equation $L_d K_1(x) = G_1(x)$ by the denominator of this expression, we obtain a linear system of thirteen equations for the seven unknown coefficients A through G . This linear system has a unique solution and we obtain that K_1 is of the form (3.10) with coefficients given by (3.11).

Properties i) and ii) for $K_1(x)$ are now apparent. Converting to polar coordinates $(x_1, x_3) = (r \cos \theta, r \sin \theta)$, the equation (3.5) becomes a fourth-order ODE in θ after multiplying by r^{11} , with coefficients that are polynomials in $\sin(\theta)$ and $\cos(\theta)$ with bounded right-hand side. We have the existence and uniqueness of a solution to this ODE with π -periodic boundary conditions on its derivatives up to order 3, and since we have also shown that K is smooth away from the origin, we deduce that K is unique. $\square$

Lemma 3.3. Let μ , δ and ν satisfy (2.17). Define p and q as in (2.15), and define b and c as in (3.8). Then there exists a solution K_2 to the equation (3.6) given by

$$K_2(x_1, x_3) = \frac{Ax_1^{12} + Bx_1^{10}x_3^2 + Cx_1^8x_3^4 + Dx_1^6x_3^6 + Ex_1^4x_3^8 + Fx_1^2x_3^{10} + Gx_3^{12}}{(x_1^2 + x_3^2)^{\frac{3}{2}}(x_1^4 + bx_1^2x_3^2 + cx_3^4)^3}. \quad (3.12)$$

where

$$\begin{aligned} A &= 2, \\ B &= -6b + 12p + 9, \\ C &= 6b(p - 1) - 24c + 33p + 6, \\ D &= b^2 - 2b(c + 1) + 2(b + 13)bp - 20cp - 46c + 20p, \\ E &= -6c((b + 5)p + b + 4) + 9b(b + 2)p + 6c^2, \\ F &= 6b^2p - 6bc(p + 1) + 3c(3c - 4p), \\ G &= c(c(p + 2) - 2bp). \end{aligned} \quad (3.13)$$

K_2 is the unique solution of (3.6) in $C^\infty(\mathbb{R}^2 \setminus \{0\})$, and K_2 satisfies the following:

- i) $K_2(x) = K_2(-x)$, $K_2(\rho x) = \rho^{-3}K_2(x)$ for $\rho > 0$.
- ii) $|x^a D^a K_2(x)| < \frac{C}{|x|^3}$ for any multi-index $a \in \mathbb{N}_0^2$ and any $x \neq 0$.

Proof. The right-hand side of (3.6) is equal to

$$G_2(x_1, x_3) := \frac{P_2(x_1, x_3)}{(x_1^2 + x_3^2)^{\frac{11}{2}}},$$

where P_2 is a fourth-degree polynomial defined as

$$P_2(x_1, x_3) := 45((8p - 2)x_1^4 + (17 - 12p)x_1^2x_3^2 + (p - 2)x_3^4).$$

Our guess for a solution of (3.6) is (3.12). In exactly the same way, we apply L_d to this ansatz and obtain a function of the form $K_2(x_1, x_3) = \frac{\Pi(x_1, x_3)}{(x_1^2 + x_3^2)^{\frac{11}{2}}(x_1^4 + bx_1^2x_3^2 + cx_3^4)^5}$, where Π is a polynomial with terms $x_1^{2i}x_3^{2(12-i)}$, where $i \in \{0, 1, \dots, 12\}$. Multiplying the equation $L_d K_2(x) = G_2(x)$ by the denominator of this expression, we obtain a linear system of thirteen equations for the seven unknown coefficients A through G . This linear system also has a unique solution, and we obtain that K_2 is of the form (3.12) with A through G given by (3.13). $\square$

In summary, we have

Proposition 3.4. The kernel function $K: \mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{R}$ of the operator (3.7) is defined by

$$K(x) := 2\mu\delta \left(\sqrt{\delta}K_1(x) + \nu K_2(x) \right). \quad (3.14)$$

Here the functions $K_1(x)$ and $K_2(x)$ are defined via (3.10)–(3.11) and (3.12)–(3.13), respectively.

Remark 3.5. When $\delta = 1$, then $p = 1$, $b = 2$ and $c = (1 + \nu)(1 - \nu)$, and so the formula for K simplifies to

$$K(z) = 2\mu \frac{Az_1^4 + Bz_1^2z_3^2 + Cz_3^4}{|z|(z_1^2 + qz_3^2)^3} \quad (3.15)$$

with

$$A = 3 - 2q, \quad B = 2(3q^2 - 5q + 3), \quad C = q(3q - 2), \quad (3.16)$$

and $q = 1 - \nu$. This is exactly the kernel associated to the 1D reduced operator in the case of isotropic elasticity [4]; see also section 5 below.

3.2. The region where K is positive

A crucial component of our analysis requires that K is positive everywhere in $\mathbb{R}^2 \setminus \{0\}$. Below, for case I, we compute the range of elastic coefficients for which this holds.

Proposition 3.6 (Positivity of kernel for case I). *Let μ , δ and ν satisfy (2.17). Define p and q as in (2.15), and define b and c as in (3.8). Suppose also that ν and δ satisfy (2.13). Let $K(x) = 2\mu\delta(\sqrt{\delta}K_1(x) + \nu K_2(x))$ be the kernel for the nonlocal operator in (3.3), where functions $K_1(x)$ and $K_2(x)$ are defined via (3.10)–(3.11) and (3.12)–(3.13), respectively. Then there exist positive constants $c_{\delta,\nu}$ and $C_{\delta,\nu}$ depending only on δ and ν such that $\frac{\mu c_{\delta,\nu}}{|x|^3} \leq K(x) \leq \frac{\mu C_{\delta,\nu}}{|x|^3}$.*

Proof. Since $K(\rho x) = \rho^{-3}K(x)$ for any $\rho > 0$, we need only be concerned with the minimum values of K on $|x| = 1$, i.e. the minimum values of $K(\cos \theta, \sin \theta)$ for $\theta \in [0, \pi]$.

Based on the plots of the exact expression for K for varying ν and δ , the most likely candidates for $\arg \min_{\theta \in [0, \pi]} K(\cos \theta, \sin \theta)$ are $\theta = 0$ and $\theta = \pi/2$. Therefore, the values of (ν, δ) for which $K(x) > 0$ is at most those for which both $K(1, 0) > 0$ and $K(0, 1) > 0$. Plugging these two values in gives

$$K(1, 0) = \delta^{3/2}(3 - 4\nu) + \delta^{5/2}(4\nu - 2) + 2\delta\nu, \quad K(0, 1) = \frac{\delta(\nu(-2\delta\nu + \delta + 2\nu - 4) + 1)}{(\nu - 1)^2}.$$

Both of these quantities are positive exactly when ν and δ satisfy (2.13). Because of the explicit formula for K , we claim that (2.13) precisely describes the range of coefficients for which K is positive. It can be checked that the boundary of figure 3 coincides numerically with the boundary of the set $\{(\nu, \delta) : (2.13) \text{ is satisfied}\}$. $\square$

Remark 3.7. When $\delta = 1$ and K is given by (3.15) and (3.16), the first derivative test applied to $\theta \mapsto K(\cos \theta, \sin \theta)$ reveals that

$$K(z) > 0 \text{ if and only if } \frac{2}{3} < q < \frac{3}{2},$$

i.e. if and only if $\nu \in (-\frac{1}{2}, \frac{1}{3})$. This is the same range for ν found in [4] via variation of parameters.

The energy corresponding to (3.3) is

$$E_\Gamma(u_1) := \frac{1}{2} \int_{\mathbb{R}^2} (\mathcal{L}u_1)u_1 \, dx + \int_{\mathbb{R}^2} W(u_1) \, dx, \quad (3.17)$$

with $E_\Gamma(u_1; B_R)$ defining the localized version. The goal now is to show existence, uniqueness, and properties of solutions to (3.3). Thanks to the properties of K , solutions to (3.3) have a 1D profile, which we will prove in section 6.




Figure 3. Plot of the numerical minima of $K(\cos \theta, \sin \theta)$ in the (ν, δ) -plane. The outer region consists of (ν, δ) that satisfy (2.17). The inner region consists of the (ν, δ) for which the numerical minimum of $K(\cos \theta, \sin \theta)$ is positive.

4. The 1D reduced equation in the perpendicular case: $\partial_{u_1} W = 0$

Assuming that $W(u_1, u_3) = W(u_3)$, we now derive analogously to section 3 the reduced scalar equation for case II and then characterize the explicit form of the nonlocal kernel in proposition 4.1. Since the material is anisotropic, making the assumption that the misfit potential depends only on $\hat{u}_3$ gives us a different reduced equation. If one makes this assumption, then proceeding the same way as we did in section 3 we get an equation for $\hat{u}_3$:

$$\mathcal{F}(\mathcal{L}u_3)(k_1, k_3) := \frac{\det \mathbb{A}(k)}{a_{11}(k)} \hat{u}_3(k) = -\mathcal{F}(W'(u_3)). \quad (4.1)$$

Inverting the Fourier transform, we obtain the 1D nonlocal equation along with (2.8):

$$\begin{aligned} \mathcal{L}u_3(x_1, x_3) &= -W'(u_3(x_1, x_3)), \quad (x_1, x_3) \in \Gamma, \\ \lim_{x_3 \rightarrow \pm\infty} u_3(x_1, x_3) &= \pm 1, \quad x_1 \in \mathbb{R}, \end{aligned} \quad (4.2)$$

where

$$\mathcal{F}(\mathcal{L}u_3)(k) = \frac{2\mu r_2 (\delta(2(1-\nu) - \delta(1-2\nu))k_1^2 + k_3^2)}{\delta(2(1-\nu) - \delta(1-2\nu))k_1^2 + (1-\nu)k_3^2} \hat{u}_3(k). \quad (4.3)$$

This equation is similar to the 1D reduced equation in the case of isotropic elasticity. However, neither the numerator nor denominator are Laplacian operators. Recalling the definition of p and q from (2.15), we can rewrite (4.3) as

$$\mathcal{F}(\mathcal{L}u_3)(k) = \frac{2\mu r_2(p k_1^2 + k_3^2)}{p k_1^2 + q k_3^2} \hat{u}_3(k). \quad (4.4)$$

Note that from definition in (2.15), $0 < p \leq 4$ and $1/2 < q < \infty$ for any ν and δ satisfying (2.17), so we have

$$C(\mu, p, q)^{-1} r_2(k) \leq \mathcal{F}(\mathcal{L}u_3)(k) \leq C(\mu, p, q) r_2(k) \text{ for all } k \in \mathbb{R}^2.$$

We use the notational conventions from section 3 (e.g. $\mathcal{L}$ has nonlocal kernel K , etc) but the explicit definitions (i.e. the expressions for $\mathcal{L}$ and K) are distinct from those in section 3 unless indicated otherwise.

Proposition 4.1. *The integral form of $\mathcal{L}$ defined in (4.1) is*

$$\mathcal{L}w = -\frac{1}{4\pi} \int_{\mathbb{R}^2} (w(x-y) + w(x+y) - 2w(x)) K(y) dy,$$

where K is the unique solution in $C^\infty(\mathbb{R}^2 \setminus \{0\})$ to the PDE

$$(p\partial_1^2 + q\partial_3^2) K(z) = 2\mu(p\partial_1^2 + \partial_3^2) \left[\frac{1}{|z|^3} \right]. \quad (4.5)$$

Proof. Define L_n and L_d to be the operators whose symbols are $m_n(k) := p k_1^2 + k_3^2$ and $m_d(k) := p k_1^2 + q k_3^2$ respectively. Let $w \in \mathcal{S}(\mathbb{R}^2)$. By using a cutoff function near the origin and using the dominated convergence theorem, we may assume that $\hat{w}$ vanishes near the origin. Define $v \in \mathcal{S}(\mathbb{R}^2)$ by $L_d v = w$, so that $\hat{v} = \frac{\hat{w}}{m_d(k)}$. Therefore we have

$$\begin{aligned} \mathcal{L}w &= 2\mu \left((-\Delta)^{1/2} L_n L_d^{-1} w \right) \\ &= 2\mu \left((-\Delta)^{1/2} L_n v \right). \end{aligned}$$

By the representation formula for $(-\Delta)^{1/2}$ in lemma 1.1

$$\begin{aligned} \mathcal{L}w(x) &= -\frac{2\mu}{4\pi} \int_{\mathbb{R}^2} \frac{(L_n)_x v(x-y) + (L_n)_x v(x+y) - 2(L_n)_x v(x)}{(y_1^2 + y_3^2)^{3/2}} dy \\ &= -\frac{2\mu}{4\pi} \int_{\mathbb{R}^2} \frac{(L_n)_y \left(v(x-y) + v(x+y) - \left(\sum_{|a| \leq 2} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) \right)}{(y_1^2 + y_3^2)^{3/2}} dy, \end{aligned}$$

where we retain the multi-index convention of proposition 3.1. Now integrating by parts,

$$\begin{aligned} \mathcal{L}w(x) &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} \left(v(x-y) + v(x+y) - \sum_{|a| \leq 2} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) (L_n)_y \left(\frac{2\mu}{(y_1^2 + y_3^2)^{3/2}} \right) dy. \end{aligned}$$

Now, since $K(y)$ satisfies

$$L_d K(y) = L_n \left(\frac{2\mu}{|y|^3} \right) \quad (4.6)$$

integrating by parts again gives

$$\begin{aligned}
 \mathcal{L}_1 w(x) &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} \left(v(x-y) + v(x+y) - \left(\sum_{|a| \leq 2} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) \right) (L_d)_y K(y) \, dy \\
 &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} (L_d)_y \left(v(x-y) + v(x+y) - \left(\sum_{|a| \leq 2} \frac{D^a v(x)}{a!} (y^a + (-y)^a) \right) \right) K(y) \, dy \\
 &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} ((L_d)_x v(x-y) + (L_d)_x v(x+y) - 2(L_d)_x v(x)) K(y) \, dy \\
 &= -\frac{1}{4\pi} \int_{\mathbb{R}^2} (w(x-y) + w(x+y) - 2w(x)) K(y) \, dy.
 \end{aligned}$$

□

Proposition 4.2 (Homogeneity and positivity of kernel for case II). *Let δ and ν satisfy (2.17). Define p and q as in (2.15). Let the kernel $K : \mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{R}$ be defined as*

$$K(z) := 2\mu \frac{Az_1^6 + Bz_1^4 z_3^2 + Cz_1^2 z_3^4 + Dz_3^6}{|z|^3 (qz_1^2 + pz_3^2)^3} \quad (4.7)$$

with

$$\begin{aligned}
 A &= 2pq^2 - 2pq + q^2, & B &= -6p^2q + 6p^2 + 9pq^2 - 6pq, \\
 C &= -6p^2q + 9p^2 + 6pq^2 - 6pq, & D &= p^3 - 2p^2q + 2p^2.
 \end{aligned} \quad (4.8)$$

Then K satisfies the following:

- (i) $K(x) = K(-x)$, $K(\rho x) = \rho^{-3}K(x)$ for $\rho > 0$.
- (ii) $|x^a D^a K(x)| < \frac{C}{|x|^3}$ for any multi-index $a \in \mathbb{N}_0^2$ and any $x \neq 0$.
- (iii) K is the unique solution in $C^\infty(\mathbb{R}^2 \setminus \{0\})$ to the equation (4.5) that is homogeneous of order -3 .
- (iv) Suppose that ν and δ satisfy (2.16). Then for each $(\nu, \delta) \in \mathcal{U}^+$ there exist positive constants $c_{\delta, \nu}$ and $C_{\delta, \nu}$ depending only on δ and ν such that $\frac{\mu c_{\delta, \nu}}{|x|^3} \leq K(x) \leq \frac{\mu C_{\delta, \nu}}{|x|^3}$.

Proof. Property (i) is obvious, and (ii) is easily derived from (i). Property (iii) can be seen by converting to polar coordinates $(x_1, x_3) = (r \cos \theta, r \sin \theta)$; after multiplying by r^3 the equation (4.5) becomes a second-order ODE with bounded C^∞ coefficients in θ with π -periodic boundary conditions on $\theta \mapsto K(\cos \theta, \sin \theta)$. Thus there exists a unique twice-differentiable solution, and that the solution must be $K(\cos \theta, \sin \theta)$ follows from property (ii) and from the fact that $K(x)$ solves (4.5) in $\mathbb{R}^2 \setminus \{0\}$.

It remains to prove property iv). Using the previous discussion, it will suffice to examine the critical points of the function $\bar{K}(\theta) := K(\cos \theta, \sin \theta)$ in the interval $[0, \pi/2]$. Using elementary algebra, the roots of the function $\bar{K}'(\theta)$ in the interval $[0, \pi/2]$ are 0 , $\pi/2$, and $\arccos(\zeta^\pm)$, where

$$\zeta^\pm := \sqrt{\frac{2q^2 - pq - p^2 \pm \sqrt{(p-q)^2(p^2 + q^2)}}{(p-q)^2}}.$$

In the range $0 < p \leq 4$ and $\frac{1}{2} < q < \infty$, $\zeta^+ \in [-1, 1]$ only for $p \geq \frac{4}{3}q$, and $\zeta^- \in [-1, 1]$ is a real number only for $p \leq \frac{3}{4}q$. Therefore, the equation $\bar{K}'(\theta) = 0$ has exactly two solutions for $\frac{3}{4}q \leq p \leq \frac{4}{3}q$, and exactly three solutions otherwise.

Therefore $K(x_1, x_3) > 0$ if and only if $\bar{K}(\theta) > 0$ if and only if

$$\bar{K}(0) > 0, \quad \bar{K}(\pi/2) > 0, \quad \text{and} \quad \bar{K}(\arccos(\zeta^\pm)) > 0.$$

Substituting directly gives the relations

$$\begin{aligned} \frac{2(2pq - 2p + q)}{q^2} &> 0, \\ \frac{2(p - 2q + 2)}{p} &> 0, \\ \frac{(q-1)(p^3 + q^3 + (p^2 + q^2)\sqrt{p^2 + q^2}) + 4(1-p)pq^2}{2p(q-p)q^2} &> 0 \text{ when } p \leq \frac{3}{4}q \text{ or } p \geq \frac{4}{3}q. \end{aligned} \quad (4.9)$$

The region $\mathcal{U}^+$ is then defined by these three conditions along with the ellipticity conditions (2.17). It can be seen after elementary algebraic manipulations that the first condition gives $\nu > \frac{\delta(\delta-2)}{2(\delta^2-\delta+1)}$ and the second gives $\nu < \frac{2}{1+4\delta-2\delta^2+\sqrt{1-8\delta+28\delta^2-16\delta^3+4\delta^4}}$. To see that the third condition leads to $p > \tilde{r}$, we use a combination of analytic and numeric justification. First, when $p \geq \frac{4}{3}q$ it is easy to see that the expression $\bar{K}(\arccos(\zeta^\pm))$ is always positive. Second, we note that $\bar{K}(\arccos(\zeta^\pm)) \leq \min\{\bar{K}(0), \bar{K}(\pi/2)\}$ in the region U defined by

$$U := \{(\nu, \delta) : (2.17) \text{ is satisfied and } 1/2 < q < 1, p \leq 3q/4\},$$

and $\bar{K}(\arccos(\zeta^\pm)) \geq \min\{\bar{K}(0), \bar{K}(\pi/2)\}$ in the region $\{(\nu, \delta) : (2.17) \text{ is satisfied and } 1 \geq q, p \leq 3q/4\}$. Third, the conjugate

$$\frac{(q-1)(p^3 + q^3 - (p^2 + q^2)\sqrt{p^2 + q^2}) + 4(1-p)pq^2}{2p(q-p)q^2}$$

is positive for all $(p, q) \in U$.

Multiplying the third expression in (4.9) by the conjugate, we obtain a rational expression whose denominator is positive and whose numerator is the polynomial

$$\begin{aligned} x \mapsto & -8(-1+q)x^4 + 8(-1+q)q^3 + q^2(13+14q-11q^2)x \\ & + (-11+14q+13q^2)x^3 + 2q(1-18q+q^2)x^2. \end{aligned}$$

The roots of this polynomial can be found exactly (although their analytic expressions are cumbersome to work with) and it can be verified numerically that all four roots are real for all $(p, q) \in U$. Putting all these observations together, we see that $\bar{K}(\arccos(\zeta^\pm)) > 0$ for $(p, q) \in U$ exactly when p is larger than either two or all of the roots of the polynomial. The following facts can be verified numerically: For $(p, q) \in U$, one root is always negative. The smallest positive root $\tilde{r}$, as a function of q , defines a curve $p = \tilde{r}$ that splits the region U into two sets (note that $\tilde{r} = 0$ for $q = 1$). The other two positive roots are always outside the region U . In conclusion $\bar{K}(\arccos(\zeta^\pm)) > 0$ for $(p, q) \in U$ only if $p > \tilde{r}$, and so iv) is proved. $\square$

Remark 4.3. When $\delta = 1$ the formula for K coincides with that of (3.15) and (3.16) for fully isotropic elasticity.

Analogous to (3.17), the energy corresponding to (4.2) is

$$E_{\Gamma}(u_3) := \frac{1}{2} \int_{\mathbb{R}^2} (\mathcal{L}u_3) u_3 \, dx + \int_{\mathbb{R}^2} W(u_3) \, dx, \quad (4.10)$$

with $E_{\Gamma}(u_3; B_R)$ defining the localized version. Just as for (3.3), analogous existence, uniqueness, and properties of solutions to (4.2) are established in section 6.

5. Slip plane parallel to the plane of isotropy

In this section, we study case III; Γ is taken parallel to the plane of isotropy, that is, $\Gamma = \{(x_1, x_2, x_3) : x_3 = 0\}$. We first introduce the 3D model and reduce to the 2D nonlocal system, and then in section 5.1 we derive the scalar reduced equation in this setting. We summarize our third result on the existence and uniqueness of the solution for case III in proposition 5.2. We characterize the explicit form and the positivity of the corresponding nonlocal kernel in section 5.1.

The Dirichlet-to-Neumann map and the subsequent 1D reduced equation turn out to have forms identical to their analogues in the setting of full isotropic elasticity. The analysis of [4] then readily applies, with one exception; the analogue of the Poisson's ratio ν here is instead a function of the five elastic constants. The constraint on the Poisson's ratio in the isotropic case from [4] here describes a region in $\mathbb{R}^5$ for which the integro-differential kernel is positive. The notation for the full 3D system and the perturbed and localized energies is retained.

Lemma 5.1. Assume that $\mathbf{u} \in C^2(\mathbb{R}^3 \setminus \Gamma; \mathbb{R}^3)$, where $\Gamma = \{x_3 = 0\}$ satisfies

$$\begin{aligned} u_1^+(x_1, x_2) &= -u_1^-(x_1, x_2), \\ u_2^+(x_1, x_2) &= -u_2^-(x_1, x_2), \\ u_3^+(x_1, x_2) &= u_3^-(x_1, x_2), \end{aligned} \quad (5.1)$$

and satisfies either

$$u_1^+(\pm\infty, x_2) = \pm 1 \quad \text{for any } x_2 \in \mathbb{R} \quad (5.2)$$

or

$$u_2^+(x_1, \pm\infty) = \pm 1 \quad \text{for any } x_1 \in \mathbb{R}. \quad (5.3)$$

Suppose additionally that $\mathbf{u}$ is a local minimizer of the total energy

$$E(\mathbf{u}) = E_{\text{els}}(\mathbf{u}) + \int_{\Gamma} W(u_1^+, u_2^+) \, dS$$

in the sense of definition 2.2. Then $\mathbf{u}$ satisfies the Euler–Lagrange equations

$$\begin{aligned} \mathbb{L}\mathbf{u} &= 0 \text{ in } \mathbb{R}^3 \setminus \Gamma, \\ \sigma_{13}^+ + \sigma_{13}^- &= \partial_{u_1} W(u_1^+, u_2^+) \text{ on } \Gamma, \\ \sigma_{23}^+ + \sigma_{23}^- &= \partial_{u_2} W(u_1^+, u_2^+) \text{ on } \Gamma, \\ \sigma_{33}^+ - \sigma_{33}^- &= 0 \text{ on } \Gamma. \end{aligned} \quad (5.4)$$

Proposition 5.2 (case III: Γ parallel to the plane of isotropy, $\partial_{u_1} W = 0$). Suppose that the five elastic constants satisfy (2.4), and suppose further that $(C_{11}, C_{33}, C_{13}, C_{44}, C_{66}) \in \mathcal{V}^+$, where $\mathcal{V}^+ \subset \mathbb{R}^5$ is the region

$$\mathcal{V}^+ := \left\{ (C_{11}, C_{33}, C_{13}, C_{44}, C_{66}) : \frac{2}{3} < \frac{\eta_1}{\eta_2} < \frac{3}{2} \right\}, \quad (5.5)$$

where η_1 and η_2 are the real numbers defined as

$$\eta_1 := 2\sqrt{C_{44}C_{66}} \quad (5.6)$$

and

$$\eta_2 := \frac{1}{\tau} \left(C_{11} - C_{13} - C_{44} + \frac{C_{13}C_{44}}{C_{33}} + \frac{\sqrt{C_{11}C_{33}}(C_{33} - C_{13})(C_{44} + C_{13})}{C_{33}^2} \right), \quad (5.7)$$

and where

$$\tau := \frac{\sqrt{\sqrt{C_{11}C_{33}} - C_{13}}\sqrt{\sqrt{C_{11}C_{33}} + C_{13} + 2C_{44}}}{2\sqrt{C_{33}C_{44}}}. \quad (5.8)$$

Assume that $\partial_{u_1} W = 0$, i.e. $W(u_1, u_2) = W(u_2)$, and that W satisfies (2.14). Then the system (5.4) has a unique (up to translation in the x_1 -direction and rotation about the x_3 -axis) classical solution $\mathbf{u} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ belonging to $C^\infty(\mathbb{R}^3 \setminus \Gamma; \mathbb{R}^3)$ with $\mathbf{u}(\cdot, \cdot, x_3) \in H^s(\mathbb{R}^2)$ for all $s \geq 1$ and for all fixed $x_3 \neq 0$. Moreover, the solution is a local minimizer of E in the sense of definition 2.2. The second component u_2 is the unique stable solution to (5.13) with specific nonlocal kernel defined in (5.15) and u_2 has a 1D profile.

Now we reduce the 3D system to a 2D system just as in section 2. Here $x = (x_1, x_2) \in \mathbb{R}^2$ and $k = (k_1, k_2) \in \mathbb{R}^2$.

Lemma 5.3. Suppose that the five elastic constants satisfy (2.4). Suppose that $\mathbf{u}$ satisfies (5.4) and remains bounded as $|x_3| \rightarrow \infty$, and assume that u_1^+ and u_2^+ belong to $H^s(\mathbb{R}^2)$ for some $s \geq 1/2$. Then $\mathbf{u}$ can be expressed entirely in terms of u_1^+ and u_2^+ . In particular, (u_1^+, u_2^+) satisfies the nonlocal system

$$\begin{bmatrix} \mathcal{F}(\sigma_{13}^+(k) + \sigma_{13}^-(k)) \\ \mathcal{F}(\sigma_{23}^+(k) + \sigma_{23}^-(k)) \end{bmatrix} := -\mathbb{A}(k) \begin{bmatrix} \hat{u}_1^+(k) \\ \hat{u}_2^+(k) \end{bmatrix} = \mathcal{F} \begin{bmatrix} \partial_{u_1} W(u_1^+, u_2^+) \\ \partial_{u_2} W(u_1^+, u_2^+) \end{bmatrix} \quad \text{on } \Gamma, \quad (5.9)$$

where the 2×2 matrix $\mathbb{A}(k)$ is given by

$$\mathbb{A}(k) = \eta_1 |k| \mathbb{I} + (\eta_2 - \eta_1) |k| \frac{k \otimes k}{|k|^2}, \quad (5.10)$$

and where $\mathbb{I}$ denotes the 2×2 identity matrix and $(k \otimes k)_{ij} = k_i k_j$ for $i, j = 1, 2$.

Proof. The proof is a straightforward computation using the process outlined in lemma 2.6. The 2×2 matrix $\mathbb{A}$ is determined by the 6×6 matrix $\mathfrak{A}$. The characteristic polynomial of $\mathfrak{A}(k)$ is given by

$$\tilde{p}(\lambda) := (\lambda^2 - \delta |k|^2) (\alpha \lambda^4 - \gamma |k|^2 \lambda^2 + \beta |k|^4),$$

where the four dimensionless parameters α, β, γ and δ are given by

$$\alpha = \frac{C_{33}}{C_{44}}, \quad \beta = \frac{C_{11}}{C_{44}}, \quad \gamma = 1 + \alpha\beta - \left(\frac{C_{13}}{C_{44}} + 1 \right)^2, \quad \delta = \frac{C_{66}}{C_{44}}.$$

From the uniform ellipticity conditions (2.4), we see that the constants satisfy at least $\alpha > 0$, $\beta > 0$, $\delta > 0$ and $-2\sqrt{\alpha\beta} < \gamma \leq \alpha\beta + 1$. See also [17] for a detailed discussion on the ellipticity properties of $\mathbb{L}$. The six roots of $\tilde{p}(\lambda)$ are

$$\pm r_i(k) = \pm \vartheta_i |k|, \quad i = 1, 2, 3,$$

where the (possibly complex) constants ϑ_i are given by

$$\pm \vartheta_1 = \pm \sqrt{\delta}, \quad \pm \vartheta_2 = \pm \sqrt{\frac{\gamma + \sqrt{\gamma^2 - 4\alpha\beta}}{2\alpha}}, \quad \pm \vartheta_3 = \pm \sqrt{\frac{\gamma - \sqrt{\gamma^2 - 4\alpha\beta}}{2\alpha}}.$$

In terms of the original elastic constants, $\vartheta_2 = \tau + \tilde{\tau}$ and $\vartheta_3 = \tau - \tilde{\tau}$, where τ is defined in (5.8) and $\tilde{\tau}$ is defined as

$$\tilde{\tau} := \frac{\sqrt{\sqrt{C_{11}C_{33}} + C_{13}} \sqrt{\sqrt{C_{11}C_{33}} - C_{13} - 2C_{44}}}{2\sqrt{C_{33}C_{44}}}.$$

The exact form of $\mathfrak{A}$ will vary depending on whether the roots repeat or are complex, but the formula (5.10) for the reduced matrix $\mathbb{A}$ holds for all cases. $\square$

The matrix $\mathbb{A}(k)$ is positive definite for all $k \neq 0$ so long as

$$\eta_1 > 0 \text{ and } \eta_2 > 0.$$

Clearly $\eta_1 > 0$ by the ellipticity conditions on C_{44} and C_{66} , but the region in $\mathbb{R}^4$ describing the range of C_{11} , C_{13} , C_{33} and C_{44} for which $\eta_2 > 0$ is more difficult to describe. However, it is a nonempty set; in fact it can be checked that η_2 is positive for the materials whose elastic constants have been determined in [17, pg. 3]. A special case is when $\sqrt{C_{11}C_{33}} - C_{13} - 2C_{44} = 0$; it is straightforward to check that $\eta_2 > 0$ exactly when

$$\begin{aligned} & \frac{1}{\alpha} < \beta < 6 + \frac{2}{\alpha} + \alpha + \frac{2(1+\alpha)}{\alpha} \sqrt{1+4\alpha} \text{ when } 0 < \alpha \leq 2 + \sqrt{5}, \\ & 6 + \frac{2}{\alpha} + \alpha - \frac{2(1+\alpha)}{\alpha} \sqrt{1+4\alpha} < \beta < 6 + \frac{2}{\alpha} + \alpha + \frac{2(1+\alpha)}{\alpha} \sqrt{1+4\alpha} \text{ when } 2 + \sqrt{5} < \alpha. \end{aligned}$$

5.1. The 1D equation

Clearly the reduced 2D operator is isotropic, and so it will suffice to consider the case

$$W(u_1^+, u_2^+) = W(u_2^+);$$

the other case

$$W(u_1^+, u_2^+) = W(u_1^+)$$

is equivalent to the first case via a rotation of coordinates. Similar to the previous sections, we drop the superscripts on u_1^+ , u_2^+ for simplicity. Our reduced nonlocal system reads

$$\left(\eta_1 |k| \mathbb{I} + (\eta_2 - \eta_1) |k| \frac{k \otimes k}{|k|^2} \right) \begin{bmatrix} \hat{u}_1(k) \\ \hat{u}_2(k) \end{bmatrix} = -\mathcal{F} \begin{bmatrix} \partial_{u_1} W(u_1, u_2) \\ \partial_{u_2} W(u_1, u_2) \end{bmatrix} \quad \text{on } \Gamma. \quad (5.11)$$

If we assume additionally that $\partial_{u_1} W = 0$, i.e. $W(u_1, u_2) = W(u_2)$, then we can solve for $\hat{u}_1$ in terms of $\hat{u}_2$ and substitute into the remaining equation. We obtain the one-dimensional scalar anisotropic nonlocal equation in Fourier space

$$\mathcal{F}(\mathcal{L}u_2)(k_1, k_2) := \frac{\det \mathbb{A}(k)}{a_{11}(k)} \hat{u}_2(k) = -\mathcal{F}(W'(u_2)). \quad (5.12)$$

Inverting the Fourier transform and using (5.2), we obtain the scalar nonlocal equation

$$\begin{aligned}\mathcal{L}u_2(x_1, x_2) &= -W'(u_2(x_1, x_2)), \quad (x_1, x_2) \in \Gamma, \\ \lim_{x_2 \rightarrow \pm\infty} u_2(x_1, x_2) &= \pm 1, \quad x_1 \in \mathbb{R},\end{aligned}\quad (5.13)$$

where

$$\mathcal{F}(\mathcal{L}u_2)(k) = \frac{\eta_1 \eta_2 |k|^3}{\eta_2 k_1^2 + \eta_1 k_2^2} \hat{u}_2(k). \quad (5.14)$$

If $\eta_i > 0$ for $i = 1, 2$, this equation has the same form as the 1D reduced equation in the case of isotropic elasticity [4]. Setting $\mu = \eta_1/2$, $p = 1$ and $q = \eta_1/\eta_2$ with k_2 in place of k_3 in proposition 4.1, we obtain the following:

Proposition 5.4. *The integral form of $\mathcal{L}$ defined in (5.12) is*

$$\mathcal{L}w(x) = -\frac{1}{4\pi} \int_{\mathbb{R}^2} (w(x-y) + w(x+y) - 2w(x)) K(y) dy,$$

where

$$K(z) = \eta_1 \eta_2 \frac{Az_1^4 + Bz_1^2 z_2^2 + Cz_2^4}{|z|(\eta_1 z_1^2 + \eta_2 z_2^2)^3} \quad (5.15)$$

with

$$A = 3\eta_1^2 - 2\eta_1 \eta_2, \quad B = 2(3\eta_1^2 - 5\eta_1 \eta_2 + 3\eta_2^2), \quad C = 3\eta_2^2 - 2\eta_1 \eta_2 \quad (5.16)$$

is the unique solution in $C^\infty(\mathbb{R}^2 \setminus \{0\})$ to the PDE

$$(\eta_2 \partial_1 + \eta_1 \partial_2) K(z) = \eta_1 \eta_2 (\partial_1^2 + \partial_2^2) \left[\frac{1}{|z|^3} \right]. \quad (5.17)$$

Proposition 5.5 (Homogeneity and positivity of kernel for case III). *Define η_1 and η_2 as in (5.6) and (5.7). Let the kernel $K: \mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{R}$ be defined as in (5.15) and (5.16). Then K satisfies the following:*

- (i) $K(x) = K(-x)$, $K(\rho x) = \rho^{-3} K(x)$ for $\rho > 0$.
- (ii) $|x^a D^a K(x)| < \frac{C}{|x|^3}$ for any multi-index $a \in \mathbb{N}_0^2$ and any $x \neq 0$.
- (iii) K is the unique solution to the equation (5.17) in $C^\infty(\mathbb{R}^2 \setminus \{0\})$ that is homogeneous of order -3 .
- (iv) Suppose that the five elastic constants are in $\mathcal{V}^+$. Then there exist positive constants c and C depending only on the five elastic constants such that $\frac{c}{|x|^3} \leq K(x) \leq \frac{C}{|x|^3}$.

Proof. We proceed identically to proposition 4.2. Set $\ell = \eta_1/\eta_2$. The roots of $\bar{K}'(\theta) = \frac{d}{d\theta} K(\cos(\theta), \sin(\theta))$ for $\theta \in [0, \pi/2]$ are $0, \pi/2$, and $\arccos(\zeta)$, where

$$\zeta := \sqrt{\frac{1 + 2\ell - 2\sqrt{1 + \ell^2}}{\ell - 1}} \quad \text{for } \ell \in (1/2, 3/4) \cup (4/3, 2).$$

In the range $\frac{1}{2} < \ell < 2$, $\zeta \in [0, 1]$ only for $\ell \in (1/2, 3/4) \cup (4/3, 2)$. Therefore, the equation $\bar{K}'(\theta) = 0$ has exactly three solutions in $[0, \pi/2]$ for $\ell \in (1/2, 3/4) \cup (4/3, 2)$, and exactly two solutions otherwise. Therefore $K(x_1, x_2) > 0$ if and only if $\bar{K}(\theta) > 0$ if and only if

$$\bar{K}(0) > 0, \quad \bar{K}(\pi/2) > 0, \quad \text{and} \quad \bar{K}(\arccos(\zeta)) > 0 \quad \text{when } \ell \in (1/2, 3/4) \cup (4/3, 2).$$

Substituting directly gives the relations

$$\begin{aligned} \frac{2(3\ell-2)}{\ell^2} &> 0, \\ 2(3-2\ell) &> 0, \\ \frac{\ell^3 + \ell^2\sqrt{\ell^2+1} + \sqrt{\ell^2+1} + 1}{2\ell^2} &> 0 \text{ when } \frac{4}{3} < \ell < 2 \text{ or } \frac{1}{2} < \ell < \frac{3}{4}. \end{aligned} \quad (5.18)$$

Clearly the third relation in (5.18) is always satisfied, and the other two relations give the condition $2/3 < \ell < 3/2$. $\square$

Note that taking the elastic constants as in (2.5) gives $\frac{\eta_1}{\eta_2} = 1 - \nu$.

6. Bounded stable solutions have 1D profiles

We now prove well-posedness and properties of solutions to the reduced scalar equations. We simultaneously consider the following three settings:

- (I) $W(u_1, u_3) = W(u_1)$, $u = u_1$ satisfies (3.3), the kernel K is defined in proposition 3.4, and the elastic constants satisfy the assumptions of proposition 2.4.
- (II) $W(u_1, u_3) = W(u_3)$, $u = u_3$ satisfies (4.2), the kernel K is defined in proposition 4.2, and the elastic constants satisfy the assumptions of proposition 2.5.
- (III) $W(u_1, u_2) = W(u_2)$, $u = u_2$ satisfies (5.13), the kernel K is defined in proposition 5.4, and the elastic constants satisfy the assumptions of proposition 5.2.

Thanks to the work in the previous sections, each of these settings can be cast in the mold of the following problem: find u satisfying

$$\begin{aligned} \mathcal{L}u(x_1, x_2) &= -W'(u(x_1, x_2)), \quad (x_1, x_2) \in \mathbb{R}^2, \\ \lim_{x_1 \rightarrow \pm\infty} u(x_1, x_2) &= \pm 1, \quad x_2 \in \mathbb{R}, \end{aligned} \quad (6.1)$$

where $x = (x_1, x_2)$,

$$\mathcal{L}w(x) = -\frac{1}{4\pi} \int_{\mathbb{R}^2} (w(x-y) + w(x+y) - 2w(x)) K(y) dy, \quad (6.2)$$

and where the nonlocal kernel $K \in C^\infty(\mathbb{R}^2 \setminus \{0\})$ satisfies

- (i) $K(x) = K(-x)$, $K(\rho x) = \rho^{-3}K(x)$ for any $\rho > 0$,
- (ii) $|x^a D^a K(x)| < \frac{C}{|x|^3}$ for any multi-index $a \in \mathbb{N}_0^2$ and any $x \neq 0$,
- (iii) there exist positive constants ℓ and L such that $\frac{\ell}{|x|^3} \leq K(x) \leq \frac{L}{|x|^3}$ for all $x \in \mathbb{R}^2 \setminus \{0\}$.

In this section, we show that any bounded stable solution in each setting (I), (II) and (III) has a 1D profile, i.e. $u(x) = \psi(e \cdot x)$ for some $e \in \mathbb{S}^1$, where ψ is the unique (up to translations) solution to a 1D scalar problem.

Let us first clarify the definition of stable solutions.

Definition 6.1. We say that u is a stable solution to (6.1) if

$$\int_{\mathbb{R}^2} (\mathcal{L}v + W''(u)v) v dx \geq 0 \quad \text{for any } v \in C_c^2(\mathbb{R}^2).$$

Define the total energy of u in any Euclidean ball $B_R \subset \mathbb{R}^2$ as

$$\begin{aligned} E_\Gamma^0(u; B_R) &:= \frac{C_d}{4} \iint_{\mathbb{R}^2 \times \mathbb{R}^2 \setminus B_R^c \times B_R^c} |u(x) - u(y)|^2 K(x-y) \, dx \, dy + \int_{B_R} W(u(x)) \, dx \\ &:= \frac{C_d}{4} \mathcal{E}(u; B_R) + F(u; B_R), \end{aligned} \quad (6.3)$$

where K satisfies properties (i)-(ii)-(iii) above. In this general setting we can obtain the interior BV estimate for stable solutions.

Lemma 6.2. *Let $|u| \leq M$ be a bounded stable solution to (6.1) in the sense of definition 6.1. Assume that W satisfies (2.14) and $L_* := \max\{2, \|W\|_{C_b^{2,a}(\mathbb{R})}\}$. Then there exists a constant $C = C(\ell, L, M, L_*)$ such that for any $B_R \subset \mathbb{R}^2$ and $R \geq 1$,*

$$\int_{B_R} |\nabla u| \, dx \leq C(\ell, L, M, L_*) R \log(L_* R), \quad \mathcal{E}(u, B_R) \leq C(\ell, L, M, L_*) R \log^2(L_* R). \quad (6.4)$$

Sketch of the proof of lemma 6.2. We only sketch the proof of this lemma because it follows the same procedures as [4, proposition 4.5].

The first key ingredient is the interior BV estimate for any direction $e \in \mathbb{S}^1$

$$\begin{aligned} \left(\int_{B_{\frac{1}{2}}} (\partial_e u(x))_+ \, dx \right) \left(\int_{B_{\frac{1}{2}}} (\partial_e u(y))_- \, dy \right) &\leq C(\delta, \nu) \frac{\mathcal{E}(u, B_R)}{R^2}, \\ \int_{B_{\frac{1}{2}}} |\nabla u(x)| \, dx &\leq C(\delta, \nu) \left(1 + \sqrt{\mathcal{E}(u, B_1)} \right). \end{aligned} \quad (6.5)$$

Indeed, let $P_{t,e}u$ be a local perturbation of u such that $P_{t,e}u(x) = u(x - te)$ for $x \in B_{\frac{1}{2}}$ and $|t|$ small enough. Then using the identity for the nonlocal energy $\mathcal{E}$, cf [4, lemma 4.1], we have

$$\begin{aligned} \mathcal{E}(u_m, B_R) + \mathcal{E}(u_M, B_R) + 2 \int_{B_{\frac{1}{2}}} \int_{B_{\frac{1}{2}}} (u(x - te) - u(x))_+ (u(y - te) - u(y))_- K(x-y) \, dx \, dy \\ \leq \mathcal{E}(u, B_R) + \mathcal{E}(P_{t,e}u, B_R), \end{aligned}$$

where $u_M := \max\{P_{t,e}u, u\}$, $u_m := \min\{P_{t,e}u, u\}$. This, together with the equality for the local energy $F(u_m, B_R) + F(u_M, B_R) = F(P_{t,e}u, B_R) + F(u, B_R)$ and the property (iii) for the kernel $0 < \frac{\ell}{|x-y|^3} \leq K(x-y)$, we have

$$\begin{aligned} E_\Gamma^0(u_m, B_R) + E_\Gamma^0(u_M, B_R) + C(\beta) \int_{B_{\frac{1}{2}}} \int_{B_{\frac{1}{2}}} (u(x - te) - u(x))_+ (u(y - te) - u(y))_- \, dx \, dy \\ \leq E_\Gamma^0(u, B_R) + E_\Gamma^0(P_{t,e}u, B_R). \end{aligned}$$

Then by the stability of the solution u , we conclude

$$\int_{B_{\frac{1}{2}}} \int_{B_{\frac{1}{2}}} (u(x - te) - u(x))_+ (u(y - te) - u(y))_- \, dx \, dy \leq C(\ell) \frac{t^2}{R^2} \mathcal{E}(u, B_R).$$

Dividing both sides by t^2 and taking $t \rightarrow 0$, we conclude the first equation (6.5). The second equation is a direct consequence of the first equation by noticing $|\nabla u| \leq |\partial_{e_1} u| + |\partial_{e_2} u|$ and the minimum of $\int_{B_{\frac{1}{2}}} (\partial_e u(x))_\pm \, dx \leq \frac{C(\ell)}{R} \sqrt{\mathcal{E}(u, B_1)}$.

The second key ingredient is a sharp interpolation inequality proved in [5, lemma 3.1]

$$\mathcal{E}(u, B_1) \leq \iint_{\mathbb{R}^2 \times \mathbb{R}^2 \setminus B_1^c \times B_1^c} \frac{|u(x) - u(y)|^2}{|x - y|^3} dx dy \leq C(M) \log L_0 \left(1 + \int_{B_2} |\nabla u| dx \right),$$

where $L_0 := \max\{2, \|\nabla u\|_{L^\infty(B_2)}\}$. Using this and (6.5), we have

$$\begin{aligned} \int_{B_{\frac{1}{2}}} |\nabla u(x)| dx &\leq C(\beta, M) \left(1 + \sqrt{C(M) \log L_2 \left(1 + \int_{B_2} |\nabla u| dx \right)} \right) \\ &\leq \frac{C(\beta, M) \log L_*}{\delta} + \delta \int_{B_2} |\nabla u| dx, \end{aligned}$$

where L_* is a constant depending on $\|W\|_{C_b^{2,\alpha}}$ and $\delta > 0$. Then using the property (i) for the kernel K , by a standard scaling argument and an iteration procedure, one can complete the proof. $\square$

Thus for each setting (I), (II), and (III), we conclude that

Theorem 6.3. Assume that $|u| \leq M$ is a bounded stable solution to (6.1) and W satisfies (2.14). Suppose the kernel K satisfies conditions (i)-(ii)-(iii) above. Then u has a 1D monotone profile and $|u| \leq 1$. Moreover, the solution to (6.1) can be characterized as $u(x) = \psi(e \cdot x)$ for any $e := (\cos \theta, \sin \theta) \in \mathbb{S}^1$ with fixed $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, where ψ is the unique solution (up to translations) to the 1D scalar problem

$$\begin{aligned} (-\Delta)^{\frac{1}{2}} \psi(x_1) &= \frac{-1}{m(\cos \theta, \sin \theta)} W'(\psi(x_1)), \quad x_1 \in \mathbb{R}, \\ \lim_{x_1 \rightarrow \pm\infty} \psi(x_1) &= \pm 1, \end{aligned}$$

and where m is the symbol associated to $\mathcal{L}$ that is defined in (3.4), (4.4), or (5.14).

Sketch of proofs for theorem 6.3. The proof is same as that of [4, theorem 4.6], so we only sketch it.

First, combining the energy estimate (6.4) with the interior BV estimate (6.5), we obtain that as $R \rightarrow +\infty$, for any direction $e \in \mathbb{S}^1$ and any half ball in $\mathbb{R}^2$ we have

$$\partial_e u \geq 0 \text{ in } \mathbb{R}^2 \quad \text{or} \quad \partial_e u \leq 0 \text{ in } \mathbb{R}^2 \quad \text{for any } e \in \mathbb{S}^1,$$

which yields the conclusion that u has a 1D monotone profile.

Second, we need to find ψ such that $u(x) = \psi(e \cdot x)$ satisfies (6.1), i.e.

$$-\mathcal{F}(W'(u))(k) = -\mathcal{F}(W'(\psi(e \cdot x)))(k) = m(k_1, k_2) \mathcal{F}(u)(k) = m(k_1, k_2) \mathcal{F}(\psi(e \cdot x))(k).$$

Then using elementary calculations, and the property that the Fourier transform commutes with rotations, one has

$$|k_1| \hat{\psi}(k_1) = -\frac{1}{m(\cos \theta, \sin \theta)} \mathcal{F}(W'(\phi))(k_1),$$

which concludes the AD scalar problem for ψ . $\square$

Proofs of propositions 2.4, 2.5, and 5.2. From theorem 6.3, the solution to (6.1) is unique and has a 1D profile. That means the solutions to (3.3), (4.2) and (5.13) is unique and has a 1D profile respectively. From obtained one component of the 3D solution, we can further solve for the other two components in the 3D system by using the elastic extension based on the relevant Dirichlet-to-Neumann map in section 2 (see [7] for the details in the fully isotropic case). Finally, the unique stable solution to the full 3D system corresponding to each of the three settings is obtained. $\square$

Remark 6.4. For more general anisotropic materials, or for elastic coefficients not satisfying (2.11), the expression of the matrix $\mathbb{A}$ associated to the Dirichlet-to-Neumann map is more complicated, making the program followed in this work less tractable. The results in this paper remain open in those more general settings.

Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

Acknowledgments

Y G is supported by NSF grant DMS-2204288. J S is supported in part by NSF DMS-1937254 and DMS-2012562.

ORCID iDs

Yuan Gao  <https://orcid.org/0000-0002-7231-5672>

James M Scott  <https://orcid.org/0009-0007-0778-4177>

References

- [1] Cabré X and Solá-Morales J 2005 Layer solutions in a half-space for boundary reactions *Commun. Pure Appl. Math.* **58** 1678–732
- [2] Cinti E, Serra J and Valdinoci E 2019 Quantitative flatness results and BV -estimates for stable nonlocal minimal surfaces *J. Differ. Geom.* **112** 447–504
- [3] Dipierro S, Serra J and Valdinoci E 2020 Improvement of flatness for nonlocal phase transitions *Am. J. Math.* **142** 1083–160
- [4] Dong H and Gao Y 2021 Existence and uniqueness of bounded stable solutions to the peierls–nabarro model for curved dislocations *Calc. Var. Partial Differ. Equ.* **60** 1–26
- [5] Figalli A and Serra J 2020 On stable solutions for boundary reactions: a De Giorgi-type result in dimension $4 + 1$ *Invent. Math.* **219** 153–77
- [6] Gao Y, Liu J-G and Liu Z 2021 Existence and rigidity of the vectorial Peierls–Nabarro model for dislocations in high dimensions *Nonlinearity* **34** 7778
- [7] Gao Y, Liu J-G, Luo T and Xiang Y 2021 Revisit of the Peierls–Nabarro model for edge dislocations in Hilbert space *Discrete Contin. Dyn. Syst. B* **26** 3177
- [8] Gao Y and Roquejoffre J-M 2022 Asymptotic stability for diffusion with dynamic boundary reaction from Ginzburg–Landau energy (arXiv:2201.02105)
- [9] González M and Monneau R 2012 Slow motion of particle systems as a limit of a reaction–diffusion equation with half-Laplacian in dimension one *Discrete Contin. Dyn. Syst.* **32** 1255–86
- [10] Grafakos L 2009 *Modern Fourier Analysis* vol 250 (Springer)
- [11] Gui C and Li Q 2020 Some energy estimates for stable solutions to fractional Allen–Cahn equations *Calc. Var. Partial Differ. Equ.* **59** 49

- [12] Hirth J P and Lothe J 1982 *Theory of Dislocations* 2nd edn (Wiley)
- [13] Merodio J and Ogden R W 2003 A note on strong ellipticity for transversely isotropic linearly elastic solids *Q. J. Mech. Appl. Math.* **56** 589–91
- [14] Nabarro F R N 1947 Dislocations in a simple cubic lattice *Proc. Phys. Soc.* **59** 256–72
- [15] Patrizi S and Valdinoci E 2016 Relaxation times for atom dislocations in crystals *Calc. Var. Partial Differ. Equ.* **55** 1–44
- [16] Patrizi S and Valdinoci E 2017 Long-time behavior for crystal dislocation dynamics *Math. Models Methods Appl. Sci.* **27** 2185–228
- [17] Payton R C 2012 *Elastic Wave Propagation in Transversely Isotropic Media* vol 4 (Springer)
- [18] Peierls R 1940 The size of a dislocation *Proc. Phys. Soc.* **52** 34
- [19] Savin O 2018 Rigidity of minimizers in nonlocal phase transitions *Anal. PDE* **11** 1881–900
- [20] Ting T C T and Chen T 2005 Poisson’s ratio for anisotropic elastic materials can have no bounds *Q. J. Mech. Appl. Math.* **58** 73–82
- [21] Xiang Y 2006 Modeling dislocations at different scales *Commun. Comput. Phys.* **1** 383–424 (available at: http://global-sci.org/intro/article_detail/cicp/7962.html)