



Profinite properties of algebraically clean graphs of free groups

Kasia Jankiewicz ^{a,*}, Kevin Schreve ^b



^a Department of Mathematics, University of California, Santa Cruz, CA 95064, United States of America

^b Department of Mathematics, Louisiana State University, Baton Rouge, LA 70806, United States of America

ARTICLE INFO

Article history:

Received 22 December 2023

Received in revised form 30 June 2024

Available online 14 July 2024

Communicated by P. Tip

MSC:

20F65; 20E26; 20F36

Keywords:

Algebraically clean graphs of groups

Residual p -finiteness

Cohomological p -completeness

Cohomological goodness

Artin groups

ABSTRACT

We prove that for every prime p algebraically clean graphs of groups are virtually residually p -finite and cohomologically p -complete. We also prove that they are cohomologically good. We apply this to certain 2-dimensional Artin groups.

© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

An *algebraically clean graph of free groups* is a graph of groups where each vertex group and edge group are finite rank free groups, and all the inclusion maps are inclusions of free factors. Examples of the fundamental groups of algebraically clean graph groups include free-by-cyclic groups, the fundamental groups of clean 2-complexes in the sense of Wise [36], and certain 2-dimensional Artin groups [22,23]. We note that (many among) the former examples are known to not admit (virtual) cocompact actions on CAT(0) cube complexes, so they are not virtually cocompactly special. In particular, the family of algebraically clean graphs of groups is strictly larger than the family of the fundamental groups of finite clean 2-complexes, which all are virtually special.

* Corresponding author.

E-mail addresses: kasia@ucsc.edu (K. Jankiewicz), kschreve@lsu.edu (K. Schreve).

1.1. Virtual residual p -finiteness

A group G is *residually finite* if for every $g \in G - \{1\}$ there exists a quotient $\phi : G \rightarrow K$ where K is a finite group and $\phi(g) \neq 1$. The fundamental groups of algebraically clean graph of free groups are known to be residually finite [37, Thm 3.4].

Let p be a prime number. A group G is *residually p -finite* if for every $g \in G - \{1\}$ there exists a quotient $\phi : G \rightarrow K$ where K is a finite p -group and $\phi(g) \neq 1$. Clearly, every residually p -finite group is residually finite, but the converse does not hold.

Theorem 1.1. *For every prime p , the fundamental group of an algebraically clean graph of free groups has a finite index subgroup that is residually p -finite.*

We do not know whether algebraically clean graphs of free groups are linear. We note that linear groups are known to be virtually residually p -finite [28] for all but finitely many primes p . There have been previous combination theorems concerning residual p -finiteness, originating in the work of Higman [17], see e.g. [34,2,33] and references therein.

1.2. Cohomology of profinite and pro- p completions

For a residually finite group G , the *profinite completion* \hat{G} of G is defined as

$$\hat{G} = \varprojlim_{[G:H] < \infty} G/H,$$

where the inverse limit is taken over the system of finite quotients of G . For every G , there is a canonical homomorphism $i : G \rightarrow \hat{G}$ which sends $g \in G$ to the cosets gH . A group G is called *cohomologically good* (also known as *good in the sense of Serre*) if for every finite G -module M the induced homomorphism

$$H_{cont}^*(\hat{G}, M) = \varinjlim_{[G:H] < \infty} H^*(G/H, M) \xrightarrow{i^*} H^*(G, M)$$

is an isomorphism. We always take the cohomology of a profinite group to be its continuous cohomology. Goodness was introduced in [32, Exercises 2.6].

We can analogously define *cohomological p -completeness* for a residually p -finite group. In this case, the *pro- p completion* \hat{G}_p of G is given by

$$\hat{G}_p = \varprojlim G/H$$

where H varies over all the normal subgroups of G whose index is a power of p . Then G is *cohomologically p -complete* if the homomorphism $G \rightarrow \hat{G}_p$ induces an isomorphism

$$H_{cont}^*(\hat{G}_p, \mathbb{F}_p) \rightarrow H^*(G, \mathbb{F}_p)$$

where we assume the G -action on \mathbb{F}_p is trivial.

Theorem 1.2. *The fundamental group of an algebraically clean graph of free groups is*

- (1) *cohomologically good,*
- (2) *for every prime p , virtually cohomologically p -complete.*

For each p , the cohomologically complete finite index subgroup is a priori different. General graphs of free groups do not always satisfy the above theorem. Indeed, there exist examples of amalgamated products of free groups that are not residually finite [7,35]. There are even examples of simple groups that split as amalgamated products of free groups [10].

1.3. Virtual poly-freeness

A group G is *poly-free* if it admits a chain of subgroups $1 = G_0 \trianglelefteq G_1 \trianglelefteq \cdots \trianglelefteq G_n = G$ such that G_i/G_{i-1} is a free group (of possibly infinite rank). We say G is *normally poly-free* if additionally all subgroups G_i are normal in G .

Theorem 1.3. *Algebraically clean graphs of free groups are normally poly-free.*

This has a number of consequences; for instance it implies these groups are locally indicable, hence left-orderable [31], and satisfy the $K-$ and $L-$ theoretic Farrell-Jones Conjecture [4,9].

1.4. Applications to Artin groups

An *Artin group* is given by a presentation

$$A = \langle s_1, \dots, s_k \mid \underbrace{s_i s_j s_i \cdots}_{m_{ij} \text{ terms}} = \underbrace{s_j s_i s_j \cdots}_{m_{ij} \text{ terms}} \rangle$$

where $m_{ij} \in \{2, 3, \dots\} \cup \{\infty\}$. We understand $m_{ij} = \infty$ as no relation involving s_i and s_j . A *triangle Artin group* $A_{\ell mn}$ is an Artin group where $k = 3$, and $m_{12} = \ell$, $m_{23} = m$, and $m_{13} = n$.

Since finite type Artin groups are linear ([24], [8] for braid groups, and [12], [13] in general), it follows that they are also virtually residually p -finite. Moreover, pure Artin groups of type A_n, C_n, G_2 and $I_2(n)$ are residually p and cohomologically p -complete for all p [1], and cohomologically good [32].

With the next corollary in mind, we note that the only spherical triangle Artin groups are the $A_{22n} = A(I_2(n)) \times \mathbb{Z}$ for $n \geq 2$, and A_{23n} where $n \in \{3, 4, 5\}$. Among those, the even ones, A_{22n} for even n , all are known to be cohomologically good and virtually residually p -finite and cohomologically p -complete for all p .

Corollary 1.4. *A triangle Artin $A_{\ell mn}$ where $\ell \leq m \leq n$ is*

- *residually finite and cohomologically good,*
- *for each prime p , virtually residually p -finite and cohomologically p -complete,*
- *virtually normally poly-free,*

provided that

- $\ell = 2$, and $m, n \geq 4$ and at least one of them is even, or
- $\ell, m, n \geq 4$ except for the case where $\ell = m = 4$ and n is odd.

In particular, all even triangle Artin groups and all extra-extra-large triangle Artin groups (i.e. where $\ell, m, n \geq 5$) satisfy the above.

Moreover, there are many more 2-dimensional Artin groups that have the above properties. See [22] for a combinatorial criterion on the defining graph, which ensure that the associated Artin group is virtually algebraically clean graph of free groups.

The Artin groups above were shown to virtually split as algebraically clean graphs of free groups in [22,23]. We note that “virtual” in the above statement is necessary. Indeed, a group G that is residually p -finite for all primes p is bi-orderable [29] (see also [25]), but the only bi-orderable Artin groups are right-angled Artin groups. However, it is possible that each Artin group listed above contains a finite index subgroup that is residually p -finite for all primes p .

Artin groups that are known to be poly-free are right-angled Artin groups [14,19,20], even FC-type Artin groups [6], and even large type Artin groups [5]. Artin groups of types A_n , $B_n = C_n$, D_n , F_4 , G_2 and $I_2(n)$ [11], as well as \tilde{A}_n , \tilde{B}_n , \tilde{C}_n , \tilde{D}_n [30] are known to be virtually poly-free. Independently, Wu-Ye proved that all triangle Artin groups except A_{23n} where n is odd, are virtually poly-free [39]. Wu-Ye also show that some triangle Artin groups are not poly-free.

Finally, we also establish residual finiteness and cohomological goodness for all even Artin groups whose defining graphs contains no 4-cliques.

Theorem 1.5. *Let Γ be a finite labeled graph with all even labels that does not contain a 4-clique. Then A_Γ is residually finite and cohomologically good.*

Such Artin groups are also poly-free by [5,38].

Acknowledgments

We thank the anonymous referee for their useful comments. The first author was supported by the NSF grant DMS-2203307 and DMS-2238198. The second author was supported by the NSF grant DMS-2203325.

2. Graphs of groups

2.1. Graph of groups notation

We recall the basic definitions and set the notation.

A graph Y consists of a set $V(Y)$ of its *vertices*, and a set $E(Y)$ of its *edges*, and two maps:

- (1) $\bar{\cdot} : E(Y) \rightarrow E(Y)$ satisfying $\bar{\bar{e}} = e$, where we think of \bar{e} as the edge e with the orientation reversed,
- (2) $\tau : E(Y) \rightarrow V(Y)$, which we think of as taking the endpoint of an edge.

A graph of groups \mathcal{G} with underlying graph Y consists of a family of vertex groups $\{G_v\}_{v \in V(Y)}$ and edge groups $\{G_e\}_{e \in E(Y)}$ where $G_e = G_{\bar{e}}$ together with maps $\{f_e : G_e \rightarrow G_{\tau(e)}\}_{e \in E(Y)}$.

Let $T \subseteq E(Y)$ be a set of edges of a spanning tree of Y . The *fundamental group* $\pi_1 \mathcal{G}$ of the graph of groups \mathcal{G} is constructed as the quotient

$$\pi_1 \mathcal{G} = (*_{v \in V(Y)} G_v * F(E(Y))) / K$$

where K is a set of the following relations

- (1) $ef_e(g)\bar{e} = f_{\bar{e}}(g)$ for all $e \in E(Y)$ and $g \in G_e$, and
- (2) $\bar{e} = e^{-1}$, and $e = 1$ if and only if $e \in T$.

2.2. Algebraically clean graph of groups

An *algebraically clean graph of free groups* is a graph of groups \mathcal{G} with finite underlying graph Y , where G_v is a finite rank free group for all $v \in V(Y)$, G_e is finitely generated for all $e \in E(Y)$, and the maps $f_e : G_e \rightarrow G_{\tau(e)}$ are injective maps onto free factors.

Let G be a group, and $N, M \subseteq G$ be two subgroups. We say that an isomorphism $\phi : N \rightarrow M$ is a *partial automorphism*, if there exists an automorphism $\phi^{ext} : G \rightarrow G$ such that $\phi|_N = \phi$. A *partial identity* is a partial automorphism that can be extended to the identity.

Proposition 2.1. *Every algebraically clean graph of free groups \mathcal{G} admits a splitting as an algebraically clean graph of groups \mathcal{G}' where the underlying graph Y' has a unique vertex, and up to renaming e and \bar{e} , $G_e \subseteq G_{\tau(e)}$ is a free factor, f_e is the inclusion map, and $f_{\bar{e}}$ is a partial automorphism of $G_{\tau(e)}$.*

Proof. Consider a spanning tree T in the underlying graph Y of \mathcal{G} . We define a new graph Y' to have the vertex set $V(Y') = \{T\}$ and edge set $E(Y') = \{e \in E(Y) \mid e \notin E(T)\}$. Let \mathcal{G}_T denote the graph of groups with the underlying graph T , and the vertex and edge groups and maps as in \mathcal{G} . Then $\pi_1 \mathcal{G}_T = *_{v \in T} G_v / \{f_e(g) = f_{\bar{e}}(g)\}$. By induction on the number of vertices in T , we can argue that $\pi_1 \mathcal{G}_T$ is a finite rank free group, where each G_v for $v \in T$ embeds as a direct factor. That is clearly the case when $|V(T)| = 1$. Let $V(T) \geq 1$, and let $v \in T$ be a leaf, and T' be the subtree of T excluding v , and let e be the unique edge in T such that $\tau(e) = v$. Then $\pi_1 \mathcal{G}_T = \mathcal{G}_{T'} *_{f_e(g) = f_{\bar{e}}(g)} G_v$, which is an amalgamated product of two finite rank free groups along their free factors, so it is a finite rank free group. By construction, the group G_v and every free factor of $\mathcal{G}_{T'}$ embed in \mathcal{G}_T as free factors.

By “collapsing” T in Y , we can identify $\pi_1 \mathcal{G}$ with the fundamental group of a graph of groups \mathcal{G}' with underlying graph Y' , where

- the unique vertex group G_T is $\pi_1 \mathcal{G}_T$,
- for each edge $e \in E(Y)$, i.e. $e \notin E(T)$, G_e becomes identified with $f_e(G_e) \subseteq G_{\tau(e)} \subseteq G_T$ which is a free factor in $G_{\tau(e)}$ and therefore also in G_T , and the map $f_{\bar{e}} : G_{\bar{e}} = G_e \rightarrow G_{\tau(\bar{e})} \subseteq G_T$ is an embedding onto some free factor of $G_{\tau(\bar{e})}$ and again also a free factor of G_T . We can thus think of that map $f_{\bar{e}}$ as a partial automorphism of G_T . \square

3. Residual p -finiteness

Throughout this section p is a fixed prime.

3.1. Well-known basics on residual p -finiteness

We start with stating some easy facts that we will use later.

Lemma 3.1. *Let G be a finitely generated group.*

- (1) *Let $N \trianglelefteq G$ be a subgroup whose index is a power of p . Then there exists a characteristic subgroup $K \trianglelefteq G$ whose index is a power of p , such that $K \subseteq N$.*
- (2) *Suppose G fits in a short exact sequence*

$$1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$$

where Q is a finite p -group, and N is residually p -finite. Then G is residually p -finite.

Proof. (1) Let K be the intersection $\bigcap H$ of all the normal subgroups H of G of index $[G : N]$. Note that K is also the kernel of a homomorphism $G \rightarrow \prod_H G/H$, since the order of each G/H is a power of p , so is the order of $\prod_H G/H$. In particular, the index $[G : K]$ is a power of p .

(2) Let $g \in G$. If g survives in Q , then Q is the required finite p -quotient of G . Suppose $g \in N$. Since N is residually p -finite, then using (1) we know that there exists a characteristic subgroup $K \subseteq N$ such that $g \notin K$ and whose index is a power of p . Since K is characteristic in N , it is normal in G , and $[G : K]$ is a power of p . \square

3.2. Basics on lower central p -series

Let G be a finitely generated group. For subgroup $H, K \subseteq G$ we denote:

- $H^p = \langle h^p \mid h \in H \rangle$,
- $[H, K] = \langle [h, k] \mid h \in H, k \in K \rangle$, and we use the convention that $[h, k] = hkh^{-1}k^{-1}$,
- $HK = \langle hk \mid h \in H, k \in K \rangle$.

Let G be a finitely generated group. A *filtration* of G is a collection $(G_n)_{n \in \mathbb{N}}$ of subgroups of G where $G_1 = G$, and $G_{n+1} \subseteq G_n$ for each $n \in \mathbb{N}$. A filtration $(G_n)_{n \in \mathbb{N}}$ is *normal* if $G_n \trianglelefteq G$ is normal for each $n \in \mathbb{N}$, and it is *separating* if $\bigcap_{n \in \mathbb{N}} G_n = \{1\}$.

The *lower p -central filtration* $\{\gamma_n^p(G)\}_n$ of G is defined as:

$$\gamma_1^p(G) := G, \quad \gamma_{n+1}^p(G) := (\gamma_n^p(G))^p [G, \gamma_n^p(G)].$$

We also denote $L_n^p(G) = \gamma_n^p(G)/\gamma_{n+1}^p(G)$. In particular, $L_1^p(G) = H_1(G, \mathbb{F}_p)$. The lower p -central filtration of G is a normal filtration, and it is separating if and only if G is residually p -finite. We note a couple of basic well-known properties of the lower p -central series. For completeness, we provide proofs.

Lemma 3.2.

- (1) For each n we have $(\gamma_n^p(G))^p \subseteq \gamma_{n+1}^p(G)$.
- (2) For each n, m we have $[\gamma_m^p(G), \gamma_n^p(G)] \subseteq \gamma_{n+m}^p(G)$.
- (3) Each $\gamma_n^p(G)$ is a characteristic subgroup of G . In particular, for each i there are natural homomorphisms $\theta_n : \text{Aut}(G) \rightarrow \text{Aut}(L_n^p(G))$ and $\sigma_n : \text{Aut}(G) \rightarrow \text{Aut}(G/\gamma_n^p(G))$.

Proof. (1) Follows immediately from the definition.

(2) We induct on m . For $m = 1$ the statement follows directly from the definition for every n . Suppose that $[\gamma_{m-1}^p(G), \gamma_n^p(G)] \subseteq \gamma_{n+m-1}^p(G)$ for every n .

First we claim that $[(\gamma_{m-1}^p(G))^p, \gamma_n^p(G)] \subseteq \gamma_{n+m}^p(G)$. Given $k \in \gamma_n^p(G)$ and $h \in \gamma_{m-1}^p(G)$ we need to show that $[h^p, k] \in \gamma_{n+m}^p(G)$. First note that $[h^p, k] = h^p u^p$ where $u = kh^{-1}k^{-1}$. By the inductive assumption $u = h^{-1}\ell$ for some $\ell \in \gamma_{n+m-1}^p(G)$.

We have

$$\begin{aligned} [h, k]^p &= (hkh^{-1}k^{-1})^p \\ &= h^p(h^{-(p-1)}uh^{(p-1)})(h^{-(p-2)}uh^{(p-2)}) \dots (h^{-1}uh)u. \end{aligned}$$

By substituting $u = h^{-1}\ell$ we get

$$h^{-(p-i)}uh^{(p-i)} = h^{-(p-i)}h^{-1}\ell h^{(p-i)} = h^{-1}h^{-(p-i)}\ell h^{(p-i)} = h^{-1}\ell\ell_i = u\ell_i$$

for some $\ell_i \in \gamma_{n+m}$ where the equality $h^{-(p-i)}\ell h^{(p-i)} = \ell\ell_i$ follows from $[G, \gamma_{n+m-1}] \subseteq \gamma_{n+m}$. Thus we have

$$[h, k]^p = h^p u \ell_{p-1} u \ell_{p-2} \dots, u \ell_1 u \in (h^p u^p) \gamma_{n+m}^p(G),$$

and in particular $[h^p, k] \gamma_{n+m}^p(G) = (h^p u^p) \gamma_{n+m}^p(G) = [h, k]^p \gamma_{n+m}^p(G)$. Since $[h, k] \in \gamma_{n+m-1}^p(G)$ by induction, we have $[h, k]^p \in \gamma_{n+m}^p(G)$ by Lemma 3.2(1). We conclude that $[h^p, k] \in \gamma_{n+m}^p(G)$, as claimed.

Now we claim that $[[G, \gamma_{m-1}^p(G)], \gamma_n^p(G)] \subseteq \gamma_{n+m}^p(G)$. By the three subgroup lemma (see e.g. [21, Cor 8.28])

$$\begin{aligned} [[G, \gamma_{m-1}^p(G)], \gamma_n^p(G)] &\subseteq [[\gamma_{m-1}^p(G), \gamma_n^p(G)], G] \cdot [[G, \gamma_n^p(G)], \gamma_{m-1}^p(G)] \\ &\subseteq [\gamma_{n+m-1}^p(G), G] \cdot [\gamma_{n+1}^p(G), \gamma_{m-1}^p(G)] \\ &\subseteq \gamma_{n+m}^p(G) \end{aligned}$$

and the second and third line follow from the inductive hypothesis. Thus we conclude that $[\gamma_m^p(G), \gamma_n^p(G)] = [(\gamma_{m-1}^p(G))^p, \gamma_{m-1}^p(G)] \subseteq \gamma_{n+m}^p(G)$, as desired.

(3) We now induct on n . For $n = 1$, clearly $\gamma_1^p(G) = G$ is characteristic in G . We assume that the statements are true for $n - 1$ and prove it for n . Let $h \in \gamma_n^p(G) = \gamma_{n-1}^p(G)[G, \gamma_{n-1}^p(G)]$, i.e. $h = h_1^p \cdot [k, h_2]$ where $h_1, h_2 \in \gamma_{n-1}^p(G)$ and $k \in G$. Let $\phi \in \text{Aut}(G)$. Then

$$\phi(h) = \phi(h_1^p \cdot [k, h_2]) = \phi(h_1)^p \cdot [\phi(k), \phi(h_2)].$$

Since $\gamma_{n-1}^p(G)$ is characteristic, $\phi(h_1), \phi(h_2) \in \gamma_{n-1}^p(G)$, so $\phi(h) \in \gamma_{n-1}^p(G)[G, \gamma_{n-1}^p(G)] = \gamma_n^p(G)$. Thus $\gamma_n^p(G)$ is characteristic.

Since $\gamma_n^p(G)$ is characteristic in G , every automorphism $\phi : G \rightarrow G$ preserves $\gamma_n^p(G)$, and therefore $\sigma_n(\phi) : G/\gamma_n^p(G) \rightarrow G/\gamma_n^p(G)$ is well-defined. It is clear that σ_n is a homomorphism. The automorphism ϕ restricts to $\phi|_{\gamma_n^p(G)} : \gamma_n^p(G) \rightarrow \gamma_n^p(G)$, and to $\phi|_{\gamma_{n+1}^p(G)} : \gamma_{n+1}^p(G) \rightarrow \gamma_{n+1}^p(G)$. Thus ϕ descends to a well-defined automorphism of $L_n^p(G)$. The map θ_n is clearly a homomorphism. \square

Proposition 3.3 ([16, Chap VIII.1]). *Let $\phi \in \text{Aut}(G)$ such that $\theta_1(\phi) = \text{Id}_{L_1^p(G)}$.*

- (1) *We have $\theta_n(\phi) = \text{Id}_{L_n^p(G)}$ for all n .*
- (2) *The order of $\sigma_n(\phi)$ is a power of p .*

Proof. (1) We induct on n . The case of $n = 1$ is immediate. We assume that the statement holds for $n - 1$.

Let first $h \in \gamma_{n-1}^p(G)$. Then by assumption $\phi(h) = hk_n$ where $k_n \in \gamma_n^p(G)$. We have

$$\begin{aligned} \phi(h^p \gamma_{n+1}^p(G)) &= (hk_n)^p \gamma_{n+1}^p(G) \\ &= h^p (h^{-(p-1)} k_n h^{p-1}) (h^{-(p-2)} k_n h^{p-2}) \cdots (h^{-1} k_n h) k_i \gamma_{n+1}^p(G) \\ &= h^p k_n \ell_{p-1} k_n \ell_{p-2} \cdots k_n \ell_1 k_n \gamma_{n+1}^p(G) \\ &= h^p k_n^p (k_n^{-(p-1)} \ell_{p-1} k_n^{p-1}) (k_n^{-(p-2)} \ell_{p-2} k_n^{p-2}) \cdots (k_n^{-1} \ell_1 k_n) \gamma_{n+1}^p(G) \\ &= h^p k_n^p \gamma_{n+1}^p(G) \\ &= h^p \gamma_{n+1}^p(G) \end{aligned}$$

where $\ell_1, \dots, \ell_{p-1}, \ell, \ell'$ are some elements of $\gamma_{n+1}^p(G)$. Indeed, the fact that $h^{-j}k_nh^j = k_n\ell_j$ follows from the fact that $[\gamma_{n-1}^p(G), \gamma_n^p(G)] \subseteq [G, \gamma_n^p(G)] \subseteq \gamma_{n+1}^p(G)$ (Proposition 3.2(2)). Finally, the fact that $k_n^p \in \gamma_{n+1}^p$ follows from Proposition 3.2(1).

Now let $g \in G$. Then $\phi(g) = g\ell_2$ for some $\ell_2 \in \gamma_2^p(G)$. We have

$$\begin{aligned}\phi([g, h]\gamma_{n+1}^p(G)) &= [g\ell_2, hk_n]\gamma_{n+1}^p(G) \\ &= g(\ell_2hk_n\ell_2^{-1})g^{-1}k_n^{-1}h^{-1}\gamma_{n+1}^p(G) \\ &= gh(k_nd_{n+1}g^{-1}k_n^{-1})h^{-1}\gamma_{n+1}^p(G) \\ &= ghd_{n+1}g^{-1}d'_{n+1}h^{-1}\gamma_{n+1}^p(G) \\ &= ghg^{-1}h^{-1}\gamma_{n+1}^p(G) \\ &= [g, h]\gamma_{n+1}^p(G)\end{aligned}$$

where we used Proposition 3.2(2) to write $\ell_2hk_n\ell_2^{-1} = hk_nd_{n+1}$ for some $d_{n+1} \in \gamma_{n+1}^p(G)$ since $[\gamma_2^p(G), \gamma_{n-1}^p(G)] \subseteq \gamma_{n+1}^p(G)$, and $k_nd_{n+1}g^{-1}k_n^{-1} = d_{n+1}g^{-1}d'_{n+1}$ for some $d'_{n+1} \in \gamma_{n+1}^p(G)$ since $[\gamma_n^p(G), G] \subseteq \gamma_{n+1}^p(G)$.

Finally, every generator h of $\gamma_n^p(G)$ is of the form $h = h_1^p[g, h_2]$ for some $h_1, h_2 \in \gamma_{n-1}^p(G)$. We have

$$\phi(h\gamma_{n+1}^p(G)) = \phi(h_1^p)\phi([g, h_2])\gamma_{n+1}^p(G) = h_1^p[g, h_2]\gamma_{n+1}^p(G),$$

for some $\ell \in \gamma_{i+1}^p(G)$. This proves that $\theta_i(\phi) = \text{Id}_{L_i^p(G)}$ as claimed.

(2) We prove by induction on n that $\sigma_n(\phi^{p^{n-1}}) = \text{Id}_{G/\gamma_n^p(G)}$. The case of $n = 1$ is immediate as $\sigma_1 = \theta_1$. We assume that the statement holds for $n - 1$, in particular for every $h \in G$ we have $\phi^{p^{n-2}}(h\gamma_{n-1}^p(G)) = h\gamma_{n-1}^p(G)$, i.e. there exists $k \in \gamma_{n-1}^p(G)$ such that $\phi^{p^{n-2}}(h) = hk$. We have

$$\phi^{p^{n-1}}(h) = (\phi^{p^{n-2}})^p(h) = (\phi^{p^{n-2}})^{p-1}(\phi^{p^{n-2}}(h)) = (\phi^{p^{n-2}})^{p-1}(hk).$$

Since $k \in \gamma_{n-1}^p(G)$ and $\theta_n(\phi^{p^{n-2}}) = \theta_n(\phi)^{p^{n-2}} = \text{Id}_{L_n^p(G)}$, using Proposition 3.3(1), we get

$$\begin{aligned}(\phi^{p^{n-2}})^{p-1}(hk) &= (\phi^{p^{n-2}})^{p-2}(\phi^{p^{n-2}}(h)\phi^{p^{n-2}}(k)) \\ &\in (\phi^{p^{n-2}})^{p-2}(hk \cdot k\gamma_n^p(G)) \\ &= (\phi^{p^{n-2}})^{p-3}(hk^3\gamma_n^p(G)) = \dots = hk^p\gamma_n^p(G).\end{aligned}$$

In particular, since $k^p \in \gamma_n^p(G)$ by Proposition 3.2(1), we conclude $\phi^{p^{n-1}}(h) \in h\gamma_n^p(G)$ as required. \square

Corollary 3.4. *Let $A \subseteq \text{Aut}(G)$ be a subgroup.*

- (1) *The image $\theta_1(A) \subseteq \text{Aut}(L_1^p(G))$ is a finite p -group if and only if $\theta_n(A) \subseteq \text{Aut}(L_n^p(G))$ is a finite p -group for every $n \geq 1$.*
- (2) *The image $\theta_1(A) \subseteq \text{Aut}(G/\gamma_1^p(G))$ is a finite p -group if and only if $\sigma_n(A) \subseteq \text{Aut}(G/\gamma_n^p(G))$ is a finite p -group for every $n \geq 1$.*

Proof. (1) Fix $n \geq 1$, and suppose that $\theta_1(A)$ is a finite p -group. By Proposition 3.3(1) for every $\phi \in A$, if $\theta_1(\phi) = \text{Id}_{L_1^p(G)}$, then $\theta_n(\phi) = \text{Id}_{L_n^p(G)}$. Thus $\theta_i(A)$ is a quotient of $\theta_1(A)$. In particular, $\theta_i(A)$ is a finite p -group as required.

(2) Fix $n \geq 1$, and suppose that $\sigma_1(A) = \theta_1(A)$ is a finite p -group. For every $\phi \in A$ there exists $k \geq 1$ such that $\theta_1(\phi^{p^k}) = \theta_1(\phi)^{p^k} = \text{Id}_{L_1^p(G)}$. By Proposition 3.3(2) the order of $\sigma_n(\phi^{p^k})$ is a power of p , and therefore the order of $\sigma_n(\phi)$ is a power of p . We conclude that $\sigma_n(A)$ is a p -group. As a subgroup of the automorphism group of a finite group $\sigma_n(A)$ is also finite. \square

The above corollary implies that for every subgroup K of $\ker(\text{Aut}(G) \rightarrow \text{Aut}(L_1^p(G)))$ (which has finite index in $\text{Aut}(G)$), all the images $\sigma_i(K)$ are finite p -groups. This observation is crucial in the proof of Theorem 1.1.

We note the following observation that the operators θ_i and σ_i can be extended to partial automorphisms.

Lemma 3.5. *A partial automorphism $\phi : N \rightarrow M$ induces a partial automorphism $N \cap \gamma_n^p(G) \rightarrow M \cap \gamma_n^p(G)$ of $\gamma_n^p(G)$. In particular, it descends to the following partial automorphisms*

- (1) $\sigma_n(\phi) : N/N \cap \gamma_n^p(G) \rightarrow M/M \cap \gamma_n^p(G)$, and
- (2) $\theta_n(\phi) : N \cap \gamma_n^p(G)/N \cap \gamma_{n+1}^p(G) \rightarrow M \cap \gamma_n^p(G)/M \cap \gamma_{n+1}^p(G)$.

Proof. Since ϕ is a partial automorphism of G , there exists $\phi^{\text{ext}} \in \text{Aut}(G)$ such that $\phi|_N^{\text{ext}} = \phi$. By Proposition 3.2(3), $\gamma_n^p(G)$ is a characteristic subgroup of G , so $\phi^{\text{ext}}(\gamma_n^p(G)) = \gamma_n^p(G)$ for every n . Thus $\phi(N \cap \gamma_n^p(G)) = \phi^{\text{ext}}(N \cap \gamma_n^p(G)) = M \cap \gamma_n^p(G)$ for every n . The lemma follows. \square

3.3. Residual p -finiteness criterion for graphs of groups

We generalize a theorem of [1] that every graph of virtually residually p -groups, where edge group inclusions are isomorphisms, are virtually residually p -finite. Their result, in particular, applies to free-by-cyclic groups. We will use a criterion for residual p -finiteness of graphs of groups stated therein.

A *filtration* \mathbf{G} of a graph of groups \mathcal{G} is a collection $\{\mathbf{G}_v\}_v$ of *compatible* filtrations $\mathbf{G}_v = \{G_{v,n}\}_n$ of G_v for each $v \in V(\Gamma)$, in the sense that for all n

$$f_e^{-1}(G_{\tau(e),n}) = f_{\bar{e}}^{-1}(G_{\tau(\bar{e}),n}).$$

For a given property X (e.g. normal, separating), we say that \mathbf{G} is X if for every $v \in V(Y)$ the filtration \mathbf{G}_v is X . We say that a filtration \mathbf{G} of a graph of groups \mathcal{G} *separates edge groups* if $f_e(G_e) = \bigcap_n G_{\tau(e),n} \cdot f_e(G_e)$ for all edges e .

Let \mathbf{G}_n be the n -th depth subgroups of the filtration \mathbf{G} of \mathcal{G} , i.e. $\mathbf{G}_n = (G_{v,n})_{v \in V(Y)}$. Since the filtrations of the vertex groups are compatible, there exists a natural graph of groups quotient \mathcal{G}/\mathbf{G}_n which has Y as its underlying graph and vertex groups $G_v/G_{v,n}$.

The following will be used to prove Theorem 1.1.

Theorem 3.6 ([1, Cor 3.14]). *Let \mathbf{G} be a normal separating filtration of \mathcal{G} which separates edge groups of \mathcal{G} , such that $\pi_1(\mathcal{G}/\mathbf{G}_n)$ is residually p -finite for every $n \geq 1$. Then $\pi_1\mathcal{G}$ is residually p -finite.*

3.4. Main proof

The following lemma can be deduced from [34] but in this case it is easy to prove it directly.

Lemma 3.7. *Let P be a finite p -group. Let \mathcal{G} be a graph of p -groups where each vertex group comes with an injective homomorphism $\psi_v : G_v \rightarrow P$, and each edge group comes with an inclusion $\psi_e : G_e \rightarrow P$ such that $\psi_e = \psi_{\bar{e}}$. Moreover, assume that for each edge e the composition $\psi_{\tau(e)} \cdot f_e = \psi_e$. Then $\pi_1\mathcal{G}$ is residually p -finite.*

Proof. The assumption on \mathcal{G} implies that there exists an epimorphism $\psi : \pi_1 \mathcal{G} \rightarrow P$ which is an isomorphism on each vertex group. Indeed, ψ is defined as ψ_v on each vertex group G_v , and sending all edge generators (not edge groups) to the identity. The kernel $\ker \psi$ is thus a finite index subgroup of $\pi_1 \mathcal{G}$ and splits as a finite graph of trivial groups, i.e. $\ker \psi$ is a finite rank free group. By Lemma 3.1(2) $\pi_1 \mathcal{G}$ is residually p -finite. \square

We are now ready to prove the main theorem.

Theorem 3.8. *For every prime p , the fundamental group of an algebraically clean graph of free groups has a finite index subgroup that is residually p -finite.*

To illustrate the proof, we first consider the special case of free-by-free groups.

Proof for $F \rtimes Q$. Let F, Q be finite rank free groups. In particular, F and Q are residually p -finite. Let $\alpha : Q \rightarrow \text{Aut}(F)$ be a homomorphism associated to the semi-direct product. By Lemma 3.2(3) $\gamma_2^p(F)$ is characteristic in F , so every automorphism $\phi \in \text{Aut}(F)$ descends to an automorphism of $L_1^p(F)$, i.e. there is a well-defined homomorphism $\beta : \text{Aut}(F) \rightarrow \text{Aut}(L_1^p(F))$. By composing α with β , we obtain a homomorphism to a finite group $\beta \cdot \alpha : Q \rightarrow \text{Aut}(L_1^p(F))$. Let Q' be its kernel. Then $F \rtimes Q'$ is a finite index subgroup of $F \rtimes Q$, which we claim is residually p -finite.

By Corollary 3.4(2) the image $\sigma_i(Q')$ is a p -group for every i . In particular $(F/\gamma_i^p) \rtimes Q'$ is residually p -finite. Indeed if $g \in (F/\gamma_i^p) \rtimes Q'$ survives in Q' , then we can use the fact that Q' is residually p -finite. Otherwise, when $g \in \ker((F/\gamma_i^p) \rtimes Q' \rightarrow Q')$, then g must survive in the quotient $(F/\gamma_i^p) \rtimes \sigma_i(Q')$, which is a p group as its order is a power of p . Since every element $g \in F \rtimes Q'$ survives in $F/\gamma_i^p(F) \rtimes Q'$ for some i , we conclude that $F \rtimes Q'$ is residually p -finite. \square

We now move to the general case. Let F be a finite rank free group, and let N, M be two subgroups of the same rank, each being a free factor of F . Every isomorphism $\phi : N \rightarrow M$ can be extended to an automorphism $\phi^{ext} : F \rightarrow F$, i.e. $\phi^{ext}|_N = \phi$, which we call an *extension of ϕ to F* . Note that an extension of ϕ is far from being unique. Indeed, it is only unique if $N = M = F$.

Proof in general case. Let G' be a fundamental group of an algebraically clean graph of free groups. By Proposition 2.1 we can think of G' as the fundamental group of a graph of groups \mathcal{G}' where the underlying graph Y' of \mathcal{G}' is a wedge of k oriented circles $\{e_1, \dots, e_n\}$. The unique vertex group of \mathcal{G}' is identified with a finite rank free group F , and for each $1 \leq i \leq k$ the edge groups G_{e_i} can be identified with a free factor N_i so that $f_{e_i} = \text{Id}_{N_i}$ and $f_{\bar{e}_i} = \phi_i : N_i \rightarrow M_i$ is a partial automorphism onto a free factor M_i of F . Let Q' be a free group freely generated by $\{\phi_1, \dots, \phi_k\}$, which can be naturally identified with $\pi_1(Y)$.

Step 1. *We construct a finite index normal subgroup Q of Q' such that $\theta_n(Q)$ is trivial.*

Let $\{\phi_1^{ext}, \dots, \phi_k^{ext}\}$ be a choice of extensions of $\{\phi_1, \dots, \phi_k\}$, i.e. for each $1 \leq i \leq k$ $\phi_i^{ext} \in \text{Aut}(F)$ such that $\phi_i^{ext}|_{N_i} = \phi_i$. Recall the homomorphism $\theta_1 : \text{Aut}(F) \rightarrow \text{Aut}(L_1^p(F))$ from Proposition 3.2(3). We construct a subgroup Q of Q' as

$$Q = \ker(Q' \rightarrow \text{Aut}(F) \rightarrow \text{Aut}(L_1^p(F)))$$

where the first map sends ϕ_i to $\phi_i^{ext} \in \text{Aut}(F)$, and the second map is θ_1 . Since $\text{Aut}(L_1^p(F))$ is a finite group, the index $[Q' : Q]$ is finite.

By Corollary 3.4(1), the image of Q in $\text{Aut}(L_1^p(F))$ is trivial for every n .

Step 2. *We construct the corresponding finite index normal subgroup G of G' and realize it as the fundamental group of a graph of groups \mathcal{G} covering the graph of groups for \mathcal{G}' .*

Consider the finite index subgroup G of G' corresponding to Q , i.e. $G = \ker(G' \rightarrow Q' \rightarrow Q'/Q)$. The group G is the fundamental group of the following graph of groups \mathcal{G} . The underlying graph Y of \mathcal{G} is the finite covering space of Y' corresponding to $Q \subseteq Q'$. Each vertex group of \mathcal{G} is a copy of F . The edge groups of edges labeled with e_i are copies of N_i with the maps Id_{N_i} and ϕ_i into the respective vertex groups. We note that \mathcal{G} is still an algebraically clean graph of finite rank free groups, with a natural quotient Q .

Step 3. *There is a natural filtration \mathbf{G} of \mathcal{G} where $\mathbf{G}_v = \{\gamma_n^p(G_v)\}_n$ of G_v for each $v \in V(Y)$. The filtration is normal, separating, and it separates the edge groups.*

Lemma 3.5 implies that the filtrations \mathbf{G}_v on the individual vertex groups are compatible and indeed define a filtration on \mathcal{G} . It is immediate that \mathbf{G} is normal. Since all the vertex groups are free groups, hence residually p -finite, their lower p -central series are separating. Since the edge groups are retracts of the vertex groups, and vertex groups are residually p -finite, [1, Lem 1.6] implies that the filtration separates the edge groups.

Step 4. *For each n , $\pi_1(\mathcal{G}/\mathbf{G}_n)$ is residually p -finite.*

The graph of groups \mathcal{G}/\mathbf{G}_n has all the vertex groups naturally isomorphic to $F/\gamma_n^p(F)$ and the edge groups are $(N_i/(N_i \cap \gamma_n^p(F)))$ for respective i , with the respective edge maps being partial identities and partial automorphisms ϕ_i . We construct a further subgroup Q_n of Q as

$$Q_n = \ker(Q \rightarrow \text{Aut}(F) \rightarrow \text{Aut}(F/\gamma_n^p(F)))$$

the first map sends ϕ_i to $\phi_i^{\text{ext}} \in \text{Aut}(F)$, and the second map is the map σ_n defined in Lemma 3.2(3). By Corollary 3.4(2), the image of Q in $\text{Aut}(F/\gamma_n^p(F))$ is a finite p -group, and therefore $[Q : Q_n]$ is a p -power. We now claim that the kernel $K_n = \ker(\pi_1(\mathcal{G}/\mathbf{G}_n) \rightarrow Q \rightarrow Q/Q_n)$ is the fundamental group of a graph of groups satisfying the assumptions of Lemma 3.7, and therefore is residually p -finite. Indeed, K_n is a finite cover \mathcal{G}_n of the graph of groups \mathcal{G}/\mathbf{G}_n whose all the vertex groups are still naturally isomorphic to $F/\gamma_n^p(F)$, and edge groups are $(N_i/(N_i \cap \gamma_n^p(F)))$ for respective i , with the respective edge maps being partial identities and partial automorphisms ϕ_i . We fix a vertex group G_{v_0} of \mathcal{G}_n and for each $v \in V(Y)$ we construct a map $\phi_v : G_v \rightarrow G_{v_0}$. We describe each map as an automorphism $\psi_v \in \text{Aut}(F/\gamma_n^p(F))$ using the natural identification of each G_v with $F/\gamma_n^p(F)$. First, $\psi_{v_0} = \text{Id}_{F/\gamma_n^p(F)}$. For v such that there is a path from v to v_0 is labeled by edges $e_{i_1} \dots e_{i_k}$ we define $\psi_v = \sigma_n(\phi_{i_k}^{\text{ext}} \dots \phi_{i_1}^{\text{ext}})$.

We claim that ψ_v does not depend on the choice of the path from v to v_0 . Indeed, given some other path with labels $e_{j_1} \dots e_{j_{k'}}$ we get that

$$\theta_n(\phi_{i_k}^{\text{ext}} \dots \phi_{i_1}^{\text{ext}}) \theta_n(\phi_{j_{k'}}^{\text{ext}} \dots \phi_{j_1}^{\text{ext}})^{-1} = \theta_n(\phi_{i_k}^{\text{ext}} \dots \phi_{i_1}^{\text{ext}} (\phi_{j_1}^{\text{ext}})^{-1} \dots (\phi_{j_{k'}}^{\text{ext}})^{-1}) = \text{Id}_{G_{v_0}}$$

by our choice of Q_n . This proves that $\theta_n(\phi_{i_k}^{\text{ext}} \dots \phi_{i_1}^{\text{ext}}) = \theta_n(\phi_{j_{k'}}^{\text{ext}} \dots \phi_{j_1}^{\text{ext}})$. For any edge e with label i , we set $\psi_e = \psi_{\bar{e}} = \psi_{\tau(e)|G_e}$, and easily verify that this choice is compatible with both $\psi_{\tau(e)}$ and $\psi_{\tau(\bar{e})}$. By Lemma 3.7 \mathcal{G}_n is residually p -finite. By Lemma 3.1, so is \mathcal{G}/\mathbf{G}_n since it has a power- p subgroup that is residually p -finite.

Step 5. *The group $\pi_1(\mathcal{G})$ is residually p -finite.*

Residual p -finiteness of $\pi_1(\mathcal{G})$ follows from Theorem 3.6 and Steps 3 and 4. \square

4. Cohomological p -completeness and goodness

In this section we prove Theorem 1.1. The proofs are nearly identical for goodness/ p -completeness. Therefore, we will just prove the p -completeness statements, and mention in the last subsection how the same arguments work for goodness.

4.1. Cohomological p -completeness

Recall from the introduction that a discrete group G is *cohomologically p -complete* if the canonical homomorphism $G \rightarrow \widehat{G}_p$ to the pro- p completion induces an isomorphism

$$H_{cont}^*(\widehat{G}_p, \mathbb{F}_p) \rightarrow H^*(G, \mathbb{F}_p).$$

Theorem 4.1. *The following groups are cohomologically p -complete for all p :*

- (1) *Free groups* [27].
- (2) *Finitely generated nilpotent groups* [26].
- (3) *Right-angled Artin groups* [26].
- (4) *Free products of cohomologically p -complete groups* [27].
- (5) *Direct products of cohomologically p -complete groups where at least one factors has a finite classifying space* [27].
- (6) *Retracts of cohomologically p -complete groups* [27].

We refer to [27] for further details on cohomologically p -complete groups. The idea behind our proof is simple. We are considering multiple HNN extensions of a free group F , and the cohomology of these can be computed by a Mayer-Vietoris sequence. It is well-known that free groups are cohomologically p -complete for every p , hence four out of every five terms in the Mayer-Vietoris sequence are cohomology groups of cohomologically p -complete groups, and the remaining term is $H^*(G, \mathbb{F}_p)$. If there was a similar exact sequence for the cohomology of the pro- p completion \widehat{G}^p , then we would be done by the Five Lemma (this is essentially the argument for right-angled Artin groups used in [26]). The following property of a graph of groups is a sufficient condition for this pro- p Mayer-Vietoris sequence [1, Lem 5.11]. A profinite version can be found in [40, Prop 4.3].

Definition 4.2. Let G be the fundamental group of a graph of groups, and suppose G is residually p -finite. The pro- p topology on G is *p -efficient* if the vertex and edge groups of G are closed in the pro- p topology of G and if the pro- p topology on G induces the full pro- p topologies on the vertex and edge groups of G .

In general, if $H < G$ and G is residually p -finite, the pro- p topology on G induces the full pro- p topology on H if and only if for every p^n -index subgroup $K < H$, there is a p^m -index subgroup $J < G$ with $J \cap H \subset K$. A subgroup $H < G$ is closed in the pro- p topology if it is the intersection of p^n -index subgroups.

Lemma 4.3. *Let \mathcal{G} be a graph of groups where the edge groups are retracts of the vertex group. Then $\pi_1 \mathcal{G}$ is p -efficient if and only if*

- (1) $G = \pi_1 \mathcal{G}$ is residually p -finite,
- (2) the pro- p topology on G induces the pro- p topology on G_v for all vertices v , and
- (3) every vertex group G_v is closed in the pro- p topology of G .

Proof. Every homomorphism $\phi : G_e \rightarrow P$ to a finite p -group P extends to a homomorphism from G_v . This proves that the pro- p topology on G_v induces to the pro- p topology on G_e , and it follows that the pro- p topology on G induces the pro- p topology on G_v . When the edge group G_e is a retract of a vertex group G_v , then it is closed in the pro- p topology of G_v by [1, Lem 1.6]. Thus if G_v is closed in the pro- p topology of G , then so is G_e . \square

We state the criterion.

Theorem 4.4 ([1, Cor 5.12]). *Let \mathcal{G} be a p -efficient graph of finitely generated groups, where all vertex and edge groups are cohomologically p -complete. Then $\pi_1\mathcal{G}$ is cohomologically p -complete.*

Theorem 4.5. *For every prime p , the fundamental group of an algebraically clean graph of free groups has a finite index subgroup which is cohomologically p -complete*

Proof. We have already constructed in the proof of Theorem 3.8 a finite index subgroup $\pi_1(\mathcal{G})$ of $\pi_1(\mathcal{G}')$ which is residually p -finite. We claim that the corresponding decomposition as a graph of free groups is efficient. Recall that the pro- p topology on vertex groups is generated by the filtration $\gamma_n^p(G_v)$. By construction, for every n , $\pi_1(\mathcal{G})$ admits a homomorphism to a finite p -group which restricts to $G_v \rightarrow G_n/\gamma_n^p(G_v)$. This combined with Lemma 4.3 shows that the pro- p topology on $\pi_1(\mathcal{G})$ is efficient, so we are done by Theorem 4.4. \square

4.2. Goodness

Cohomological goodness is a bit easier to establish; we will give a more straightforward proof without restating the relevant definitions (which essentially involves replacing the pro- p completion/topology everywhere with the profinite completion/topology).

Theorem 4.6. *The fundamental group G of an algebraically clean graph of free groups is cohomologically good.*

Proof. We know G decomposes as an iterated HNN extension of a free group F , where the edge maps extend to automorphisms of F . We claim these decompositions are efficient. To see this, take a finite index characteristic subgroup C of F . There is an induced homomorphism from G to an iterated HNN extension of F/C , denoted by G' . Since F/C is finite, G' is virtually free, so let H'' be any finite index free subgroup which intersects F/C trivially. The preimage H of H'' hence intersects F inside of C . This shows the HNN extension is efficient, so we are done by the profinite Mayer-Vietoris sequence [40, Prop 4.3]. \square

Since finite extensions of good groups are good, this implies that any Artin group satisfying the conditions of Corollary 1.4 is good as well.

5. Virtual poly-freeness

Proof of Theorem 1.3. Let \mathcal{G} be an algebraically clean graph of finite rank free groups over a finite graph Γ . By Proposition 2.1, we can assume that Γ has a unique vertex v , and some finite number of loops. Then the fundamental group of \mathcal{G} fits in the following short exact sequence

$$1 \rightarrow \langle\langle G_v \rangle\rangle \rightarrow \pi_1(\mathcal{G}) \rightarrow \pi_1\Gamma \rightarrow 1.$$

We claim that $\langle\langle G_v \rangle\rangle$ is a (possibly infinite rank) free group. Indeed, the induced graph of groups decomposition of $\langle\langle G_v \rangle\rangle$ is an infinite tree of G_v , amalgamated along free factors. The chain $1 \trianglelefteq \langle\langle G_v \rangle\rangle \trianglelefteq \pi_1(\mathcal{G})$ is a chain witnessing the normal poly-freeness of $\pi_1(G(\Gamma))$. \square

6. Even Artin groups

Lemma 6.1. *Let Γ be a graph labeled by even numbers ≥ 2 , and let $\Lambda \subseteq \Gamma$ be any induced subgraph. Then A_Γ retracts onto A_Λ .*

Proof. The retraction is obtained by mapping each generator $s \in V(\Lambda)$ to itself, and each generator $s \in V(\Gamma) - V(\Lambda)$ to 1. \square

Proof of Theorem 1.5. The proof is an induction on the number of non-edges in the defining graph Γ of A_Γ . If Γ is a full graph, then by the assumption on no 4-cliques Γ has at most three vertices. If Γ has one vertex, then $A_\Gamma = \mathbb{Z}$ is residually finite and cohomologically good. If Γ has two vertices, then A_Γ is virtually $F \times \mathbb{Z}$ (see e.g. [18, Lem 4.3]), which is residually finite and cohomologically good. Since residual finiteness and cohomological goodness pass to finite index supergroups [15, Lem 3.2], A_Γ has those properties. Finally, when Γ has three vertices then A_Γ is residually finite and cohomologically good by Corollary 1.4.

We now prove the inductive step. Let Γ have a non-edge $\{u, v\}$. Then A_Γ splits as an amalgamated product $A_{\Gamma - \{u\}} *_{A_{\Gamma - \{u, v\}}} A_{\Gamma - \{v\}}$. By the inductive assumption $A_{\Gamma - \{u\}}$ and $A_{\Gamma - \{v\}}$ are residually finite and cohomologically good. By Lemma 6.1, A_Γ is an amalgamated product along retracts, so A_Γ is residually finite by [3], and cohomologically good by [15, Prop 3.5]. \square

More generally, the argument in the proof above shows that for even Artin groups being residually finite and cohomologically good are “free-of-infinity properties”, in the sense that if we prove that all even Artin groups whose defining graphs are cliques are residually finite and cohomologically good, then we will be able to deduce that the same holds for all even Artin groups.

CRediT authorship contribution statement

Kasia Jankiewicz: Writing – original draft. **Kevin Schreve:** Writing – original draft.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Matthias Aschenbrenner, Stefan Friedl, 3-manifold groups are virtually residually p , Mem. Am. Math. Soc. 225 (1058) (2013) viii+100.
- [2] D.N. Arzakov, Residual p -finiteness of generalized free products of groups, Russ. Math. 61 (2017) 1–6.
- [3] James Boler, Benny Evans, The free product of residually finite groups amalgamated along retracts is residually finite, Proc. Am. Math. Soc. 37 (1973) 50–52.
- [4] Mladen Bestvina, Koji Fujiwara, Derrick Wigglesworth, The Farrell-Jones conjecture for hyperbolic-by-cyclic groups, Int. Math. Res. Not. 2023 (7) (2023) 5887–5904.
- [5] Rubén Blasco-García, Poly-freeness in large even Artin groups, J. Pure Appl. Algebra 225 (4) (2021) 106551.
- [6] Rubén Blasco-García, Conchita Martínez-Pérez, Luis Paris, Poly-freeness of even Artin groups of FC type, Groups Geom. Dyn. 13 (1) (2019) 309–325.
- [7] Meenaxi Bhattacharjee, Constructing finitely presented infinite nearly simple groups, Commun. Algebra 22 (11) (1994) 4561–4589.
- [8] Stephen J. Bigelow, Braid groups are linear, J. Am. Math. Soc. 14 (2) (2001) 471–486 (electronic).

- [9] Benjamin Brück, Dawid Kielak, Xiaolei Wu, The Farrell-Jones conjecture for normally poly-free groups, *Proc. Am. Math. Soc.* 149 (6) (2021) 2349–2356.
- [10] Marc Burger, Shahar Mozes, Finitely presented simple groups and products of trees, *C. R. Acad. Sci., Sér. I Math.* 324 (7) (1997) 747–752.
- [11] Egbert Brieskorn, Sur les groupes de tresses [d’après V. I. Arnol’d], in: *Séminaire Bourbaki*, 24ème année (1971/1972), in: *Lecture Notes in Math.*, vol. 317, Springer, Berlin-New York, 1973, pp. 21–44, pages Exp. No. 401.
- [12] Arjeh M. Cohen, David B. Wales, Linearity of Artin groups of finite type, *Isr. J. Math.* 131 (2002) 101–123.
- [13] François Digne, On the linearity of Artin braid groups, *J. Algebra* 268 (1) (2003) 39–57.
- [14] G. Duchamp, D. Krob, Free partially commutative structures, *J. Algebra* 156 (2) (1993) 318–361.
- [15] F. Grunewald, A. Jaikin-Zapirain, P.A. Zalesskii, Cohomological goodness and the profinite completion of Bianchi groups, *Duke Math. J.* 144 (1) (2008) 53–72.
- [16] Bertram Huppert, Norman Blackburn, *Finite Groups. II*, Grundlehren der Mathematischen Wissenschaften, vol. 242, Springer-Verlag, Berlin-New York, 1982, AMD, 44.
- [17] Graham Higman, Amalgams of p -groups, *J. Algebra* 1 (1964) 301–305.
- [18] Jingyin Huang, Kasia Jankiewicz, Piotr Przytycki, Cocompactly cubulated 2-dimensional Artin groups, *Comment. Math. Helv.* 91 (3) (2016) 519–542.
- [19] James Howie, Bestvina-Brady groups and the plus construction, *Math. Proc. Camb. Philos. Soc.* 127 (3) (1999) 487–493.
- [20] Susan Hermiller, Zoran Šunić, Poly-free constructions for right-angled Artin groups, *J. Group Theory* 10 (1) (2007) 117–138.
- [21] I. Martin Isaacs, *Algebra: a Graduate Course*, Graduate Studies in Mathematics, vol. 100, American Mathematical Society, Providence, RI, 2009, Reprint of the 1994 original.
- [22] Kasia Jankiewicz, Residual finiteness of certain 2-dimensional Artin groups, *Adv. Math.* 405 (2022) 108487.
- [23] Kasia Jankiewicz, Splittings of triangle Artin groups, *Groups Geom. Dyn.* 18 (1) (2022) 91–108.
- [24] Daan Krammer, Braid groups are linear, *Ann. Math.* (2) 155 (1) (2002) 131–156.
- [25] Thomas Koberda, Alexander I. Suciu, Residually finite rationally p groups, *Commun. Contemp. Math.* 22 (3) (2020) 1950016.
- [26] Karl Lorensen, Groups with the same cohomology as their profinite completions, *J. Algebra* 320 (4) (2008) 1704–1722.
- [27] Peter Linnell, Thomas Schick, Finite group extensions and the Atiyah conjecture, *J. Am. Math. Soc.* 20 (4) (2007) 1003–1051.
- [28] V.P. Platonov, A certain problem for finitely generated groups, *Dokl. Akad. Nauk Belarusi* 12 (1968) 492–494.
- [29] A.H. Rhemtulla, Residually F_p -groups, for many primes p , are orderable, *Proc. Am. Math. Soc.* 41 (1973) 31–33.
- [30] S.K. Roushon, Some virtually poly-free Artin groups, 2020.
- [31] Akbar Rhemtulla, Dale Rolfsen, Local indicability in ordered groups: braids and elementary amenable groups, *Proc. Am. Math. Soc.* 130 (9) (2002) 2569–2577.
- [32] Jean-Pierre Serre, *Galois Cohomology*, Springer-Verlag, Berlin, 1997, Translated from the French by Patrick Ion and revised by the author.
- [33] E.V. Sokolov, On the residual nilpotence of generalized free products of groups, *J. Algebra* 657 (2024) 292–326.
- [34] Gareth Wilkes, A criterion for residual p -finiteness of arbitrary graphs of finite p -groups, *J. Group Theory* 22 (5) (2019) 837–844.
- [35] Daniel T. Wise, Non-positively curved squared complexes, aperiodic tilings, and non-residually finite groups, PhD thesis, Princeton University, 1996.
- [36] Daniel T. Wise, Subgroup separability of graphs of free groups with cyclic edge groups, *Q. J. Math.* 51 (1) (2000) 107–129.
- [37] Daniel T. Wise, The residual finiteness of negatively curved polygons of finite groups, *Invent. Math.* 149 (3) (2002) 579–617.
- [38] Xiaolei Wu, Poly-freeness of Artin groups and the Farrell-Jones conjecture, *J. Group Theory* 25 (1) (2022) 11–24.
- [39] Xiaolei Wu, Shengkui Ye, Splittings and poly-freeness of triangle Artin groups, arXiv:2312.08681 [abs], 2023.
- [40] Henry Wilton, Pavel Zalesskii, Profinite properties of graph manifolds, *Geom. Dedic.* 147 (2010) 29–45.