

Field-controlled dynamics of skyrmions and monopoles

Jung-Shen B. Tai^{1†}, Andrew J. Hess^{1‡}, Jin-Sheng Wu¹ and Ivan I. Smalyukh^{1,2,3,4*}

¹*Department of Physics and Chemical Physics Program, University of Colorado, Boulder, CO 80309, USA*

²*Department of Electrical, Computer, and Energy Engineering, Materials Science and Engineering Program and Soft Materials Research Center, University of Colorado, Boulder, CO 80309, USA*

³*Renewable and Sustainable Energy Institute, National Renewable Energy Laboratory and University of Colorado, Boulder, CO 80309, USA*

⁴*International Institute for Sustainability with Knotted Chiral Meta Matter (WPI-SKCM²), Hiroshima University, Higashihiroshima, Hiroshima, 739-8526, Japan*

* Correspondence to: ivan.smalyukh@colorado.edu

† Present address: Molecular, Cellular, and Developmental Biology, Yale University, New Haven, CT 06511, USA

‡ Present address: National Renewable Energy Laboratory, Golden, CO 80401, USA

Magnetic monopoles, despite their ongoing experimental search as elementary particles, have inspired the discovery of analogous excitations in condensed matter systems. In chiral condensed matter systems, emergent monopoles are responsible for the onset of transitions between topologically distinct states and phases, like in the case of transitions from helical and conical phase to A-phase comprising periodic arrays of skyrmions. By combining numerical modeling and optical characterizations, we describe how different geometrical configurations of skyrmions terminating at monopoles can be realized in liquid crystals and liquid crystal ferromagnets. We demonstrate how such complex structures can be effectively manipulated by external magnetic and electric fields. Furthermore, we discuss how our findings may hint at similar dynamics in other physical systems, and their potential

28 **applications.**

29 **Teaser**

30 Skyrmions and emergent monopoles in liquid-crystal ferromagnets manipulated by external

31 electric and magnetic fields.

Introduction

Magnetic monopoles, hypothetical elementary particles, which are the point sources of magnetic fields, are the holy grail of particle physics. Despite promising restoration of duality symmetry in the electromagnetic theory and various theoretical predictions, the experimental discovery of magnetic monopoles has remained elusive(1, 2). Like in the case of macroscopic magnets, the sources and sinks of magnetic field could not be found separated from each other at the level of elementary particles. However, structures analogous to monopole-like excitations have been recently discovered in various condensed matter systems, including spin ices(3), Bose-Einstein condensates(4) and magnetic solids(5). Such excitations are particularly useful in understanding emergent behavior of these condensed matter systems and physically resemble the elementary-particle counterparts of magnetic monopoles beyond topological classification (as a matter of fact, monopoles as topological objects are rather common). In magnets, $\pi_2(\mathbb{S}^2)$ topological point defects are monopoles not just in the magnetization field $\mathbf{m}(\mathbf{r})$, but also in the emergent magnetic field, $(B_{\text{em}})_i \equiv (\hbar/2e)\epsilon^{ijk}\mathbf{m} \cdot (\partial_j\mathbf{m} \times \partial_k\mathbf{m})$, a fictitious field describing the interaction between conduction electrons and the underlying spin texture, where $\hbar$ and e are the reduced Planck constant and the elementary charge, respectively(6–8). This is because the emergent magnetic flux from a $\pi_2(\mathbb{S}^2)$ topological point defect is directly proportional to its topological charge as both measure the solid angle subtended by $\mathbf{m}(\mathbf{r})$ in a confining surface. Here, $\pi_n(\mathbb{S}^m)$ refers to the n^{th} homotopy group of m -sphere $\mathbb{S}^m$, which is used to classify topological defects or solitons as maps from the configuration space to the order-parameter space $\mathbb{S}^n \mapsto \mathbb{S}^m$; the case of monopoles corresponds to 2-spheres representing both spaces, $n=m=2$ (9, 10). Emergent monopoles in chiral ferromagnets play an important role during a phase transition between topologically distinct phases, serving as nucleation points for the inception of topological transformation(5). In liquid crystals

(LCs) and LC ferromagnets, two- and three-dimensional topological point defects are ubiquitous(11, 12). Topological defects in LCs also serve as a means to embed topological solitons of lower dimensions in the topologically trivial background of a higher dimensional space(13). A few examples include embedding $\pi_2(\mathbb{S}^2)$ skyrmions in the three-dimensional space as torons and $\pi_1(\mathbb{S}^1)$ twist walls in the two-dimensional space as one of the common cholesteric finger structures and recently observed configurations of Möbiusons(13, 14).

In this article, using externally applied magnetic and electric fields, we demonstrate the dynamical control of $\pi_2(\mathbb{S}^2)$ skyrmions and the accompanying $\pi_2(\mathbb{S}^2)$ topological defects, monopoles in the emergent field. We show that, much like the skyrmion A-phase, periodic lattices of skyrmions can also form in LCs when embedded within either a uniform or a helical background. Moreover, we find that skyrmions and monopoles can be effectively controlled by external fields while the motions of the monopoles are directly observed with an optical microscope with which observations benefit particularly from in-plane configurations of skyrmions. Furthermore, we demonstrate that in a chiral LC material, the vertical skyrmion terminating at monopoles (the so-called torons) can be hosted in a helical or uniform field background with controllable inter-soliton interaction, modulated by the applied voltage. While the analogy of our skyrmions and monopoles to their high-energy-physics counterparts like Dirac monopoles(15) and $\pi_3(\mathbb{S}^3)$ Skyrme solitons(2), which in particle-physics models describe subatomic particles with different baryon numbers, is distant, the observed findings closely mimic what were found in chiral magnets (5) where even its Hamiltonian can have a form similar to that of a chiral LC, so that LCs could potentially serve as model systems and provide useful insights. The diversity of observed geometrically distinct but topologically equivalent torons, or monopole-terminating skyrmions, points to possible topology-preserving switching between them, in analogy to the recently studied

inter-transformation between geometrically distinct states of hopfions(16). Finally, we discuss the fundamental implications of monopole control in LC systems and their potential technological applications.

Results

$\pi_2(\mathbb{S}^2)$ point defects are monopoles in emergent field that embed $\pi_2(\mathbb{S}^2)$ skyrmions in topologically trivial backgrounds

Ubiquitously observed in the order-parameter field $\mathbf{n}(\mathbf{r})$ of condensed matter systems, a hedgehog point defect and a hyperbolic point defect are both topological point defects classified by the $\pi_2(\mathbb{S}^2)$ homotopy group and have a topological charge of unity (Fig. 1, A and B). In the emergent magnetic field, despite the distinct difference in their geometrical field configurations, both hedgehog and hyperbolic defects are monopoles – radial sources of emergent magnetic fields (Fig. 1, A-C). In LCs, LC ferromagnets, and solid-state magnets, torons are $\pi_2(\mathbb{S}^2)$ skyrmions terminating at hyperbolic point defects of the same $\pi_2(\mathbb{S}^2)$ field topology, often observed in a uniform field background with a perpendicular boundary condition (Fig. 1D)(12, 17–19). Note that since all $\mathbf{n}(\mathbf{r})$ structures discussed in this work can be consistently vectorized even when the order parameter of the system has nonpolar head-tail symmetry, namely $\mathbf{n}(\mathbf{r}) \equiv -\mathbf{n}(\mathbf{r})$ and the corresponding order-parameter space and homotopy group are $\mathbb{S}^2/\mathbb{Z}_2$ and $\pi_2(\mathbb{S}^2/\mathbb{Z}_2)$, respectively, we focus on vectorized $\mathbf{n}(\mathbf{r})$ exclusively for simplicity(20). The two chiral hyperbolic point defects – deformable from the hyperbolic defect in Fig. 1B (or its oppositely-charged counterpart by reversing $\mathbf{n}(\mathbf{r})$ direction) and thus topologically equivalent – act as a topological switch that transitions between regions of trivial ($N_{\text{sk}} = 0$) and non-trivial topology ($|N_{\text{sk}}| = 1$) and are required to embed skyrmions in the trivial uniform background (Fig. 1, D and

E). Here N_{sk} is the skyrmion number of the 2D $\mathbf{n}(\mathbf{r})$ field with the definition, $N_{\text{sk}} \equiv \frac{1}{4\pi} \int d^2\mathbf{r} \mathbf{n} \cdot (\partial_x \mathbf{n} \times \partial_y \mathbf{n})$. The emergent field $\mathbf{B}_{\text{em}}$ derived from $\mathbf{n}(\mathbf{r})$ of a toron shows the corresponding two defects in $\mathbf{B}_{\text{em}}$ are a pair of a monopole and an anti-monopole, acting as the source and sink of the emergent field (Fig. 1F). The $\mathbf{B}_{\text{em}}$ flux streaming from the monopole to the anti-monopole is reminiscent of the Dirac string in Dirac monopole(15). In Dirac monopoles, the Dirac string is an infinitesimally thin solenoid attached to the monopole with quantized magnetic flux. Here, the distribution of $\mathbf{B}_{\text{em}}$ flux is determined by $\mathbf{n}(\mathbf{r})$ configurations, though the total flux is also quantized due to topology. Therefore, the emergence of a skyrmion with non-trivial $\pi_2(\mathbb{S}^2)$ field topology in a trivial field background is mediated by the $\pi_2(\mathbb{S}^2)$ defects, which are a pair of monopoles in $\mathbf{B}_{\text{em}}$.

In-plane skyrmions and monopoles can be stabilized in a uniform or helical background

In chiral materials such as chiral LCs and chiral magnets, skyrmions are often recognized as axisymmetric whirling configurations that smoothly transition into a uniform far-field $\mathbf{n}_0$. Such uniform-far-field skyrmions can be approximated analytically by the ansatz $\mathbf{n}(\mathbf{r}) = (\sin f(r) \cos(\theta + \gamma), \sin f(r) \sin(\theta + \gamma), \cos f(r))$, where $f(r)$ is a monotonic function with $f(0) = \pi$ and $f(\infty) = 0$, θ is the polar angle of the 2D spatial coordinate $\mathbf{r}$, and $\gamma = \pi/2$ for right-handed chiral material. The corresponding $\mathbf{B}_{\text{em}}$ (or equivalently, skyrmion number density) derived from the ansatz also possesses the axial symmetry (Fig. 2, A and C). Such skyrmions in a uniform background are often found in the horizontal mid-plane perpendicular to $\mathbf{n}_0$ in chiral LCs and chiral LC ferromagnets between substrates with a perpendicular (homeotropic) boundary condition, where torons emerge as a result of geometrical frustration between the material's chirality and the substrate's surface alignment (Fig. 1, D-F)(13). A somewhat less commonly

familiar configuration of skyrmions is that with a helical field background; in this case, the corresponding $\mathbf{n}(\mathbf{r})$ can be thought of as splitting and splicing the uniform-background skyrmion while extending the region of helical configuration into a uniformly helical background with constant helical axis χ_0 , which defines the axis that $\mathbf{n}(\mathbf{r})$ twists around in the far field (Fig. 2, B and D)(10). Helical skyrmions constructed as such show distinct symmetry in the distribution of skyrmion number density where it is localized at two lobes around the so-called λ disclination lines (often also referred to as fractional skyrmions and merons), the singular line defects in the helical-axis field that remain nonsingular in $\mathbf{n}(\mathbf{r})$ (21). Note in the case of helical skyrmion configurations, which in LC literature are sometimes referred to as Lehman clusters, the helical background is also topologically trivial and allows for compactification of the configuration space $\mathbb{R}^2$ to $\mathbb{S}^2$ and thus the classification of the observed configuration within $\pi_2(\mathbb{S}^2)$ topology.

The richness in geometrical configurations of skyrmions expands the confinement and alignment conditions that can host the solitons and associated monopoles. In particular, torons and skyrmions can also emerge in cells with planar alignment, where their stability could be established by frustration induced by cholesteric compression/dilation or by coupling to externally applied electric and magnetic fields. Computer simulations based on Frank-Oseen free-energy minimization show skyrmions stabilized both in a uniform-field background and in a helical-field background, respectively (Fig. 2, E and H). The $\pi_2(\mathbb{S}^2)$ topological defects in $\mathbf{n}(\mathbf{r})$ mediate the embedding of skyrmions into the field backgrounds (Fig. 2, F and I), and are sources and sinks of $\mathbf{B}_{\text{em}}$ where the streamlines of $\mathbf{B}_{\text{em}}$ run along the in-plane direction parallel to the substrates, representing monopoles of opposite charges (Fig. 2, G and J). Despite the dissimilar $\mathbf{n}(\mathbf{r})$ configurations around the defects, interestingly, they are classified by the $\pi_2(\mathbb{S}^2)$ homotopy group and correspond to consistent charges of monopoles in $\mathbf{B}_{\text{em}}$. Compared to the more commonly

realized skyrmions in perpendicular-alignment cells, in-plane skyrmions and monopoles allow the same topology to be realized in a geometry where they could reconfigure in the direction parallel to the confining substrates; this allows better accessibility for optical characterization and freedom of dynamics for monopoles.

In-plane skyrmion emergence under applied electric field and their 1D lattices

To observe in-plane skyrmions experimentally, we confined chiral LCs between substrates with planar anchoring in LC cells with a thickness-to-pitch ratio of $d/p = 1$. Under such conditions, the uniformly helical (cholesteric) state with far-field helical axis χ_0 perpendicular to the substrates is the global ground state and no stable skyrmions or monopoles are observed. Upon heating and quenching from the isotropic state to the cholesteric state, dark fragments of stripe-like patterns seen under a polarizing optical microscope (POM) between crossed polarizers can be stabilized when a voltage across the substrates of $U \approx 1.4$ V is applied to the LC with positive dielectric anisotropy (Fig. 3A). To reveal unambiguously the $\mathbf{n}(\mathbf{r})$ of these fragments, we employed a nonlinear fluorescence polarizing imaging (3PEF-PM, *Methods*) where the generated patterns can be compared with those from $\mathbf{n}(\mathbf{r})$ simulations. The results obtained using different polarizations of excitation are consistent with $\mathbf{n}(\mathbf{r})$ of in-plane skyrmions terminating at monopoles in a helical background (Fig. 3B). The in-plane helical skyrmions are stabilized by an electric field $\mathbf{E}$ because of the skyrmion's energetically favored vertical $\mathbf{n}(\mathbf{r})$ along $\mathbf{E}$, which is not present in the uniformly helical background (Fig. 2H).

Besides spontaneous emergence after quenching, in-plane skyrmions can also nucleate from sample edges or be controllably generated by local melting and reorienting of LCs using laser tweezers when an electric field is applied (*Methods*). As skyrmions increase in number and fill the

sample, they form a 1D lattice in the direction perpendicular to the direction of planar alignment (Fig. 3, C and D). Defects of the skyrmion lattice as quasi-dislocations can be observed readily in a large lattice, where skyrmions terminating at monopoles locally disrupt the periodic arrangement (Fig. 3E). 3PEF-PM images show changes in N_{sk} in the vertical cross-sections mediated by monopoles, where the presence of a monopole is associated with a change in N_{sk} by unity. Specifically, Fig. 3G shows the effect of topological switching mediated by monopoles: N_{sk} increases from -3 to -2 by a monopole, and decreases from -2 to -3 by an anti-monopole (Fig. 3, E-G). The 1D skyrmion lattice in chiral LCs is analogous to the 2D skyrmion A-phase in chiral magnets, where both form as a result of energy competition between material's free energy minimization and coupling to externally applied fields. Notably, the unwinding of the chiral magnetic A-phase into ferromagnetic phase is also mediated by monopoles(5). The energetics of the emergence of skyrmions/torons mediated by monopole-antimonopole nucleations in this geometry competes with that of the formation of undulations of chiral LC's quasilayers, where they, for certain material parameters in a similar experimental geometry, can emerge as a transformation that preserves the trivial topology of the helical background (22).

In-plane skyrmions can also be realized in a uniform-field background (Figs. 2E and 3H,I). For a confined chiral LC with planar anchoring and $d/p \approx 1$, an in-plane electric field $\sim 1 \text{ V}/\mu\text{m}$ unwinds the helical $\mathbf{n}(\mathbf{r})$ configuration and makes the uniform state $\mathbf{n}_0 \parallel \mathbf{E}$ the ground state. Uniform-background in-plane skyrmions and monopoles can thus be created spontaneously through quenching or controllably by laser tweezers. Compared to the case of helical-background skyrmions, the stabilization of uniform-background skyrmions with an electric field requires high voltages (at least $\sim 1000 \text{ V}$) because of the large lateral dimensions compared to the thickness in our LC samples. However, more sophisticated sample geometry such as those with patterned

electrodes that reduce the gap size should alleviate the need for high voltages.

External fields control dynamics of skyrmions and monopoles in LCs and LC ferromagnets

Stabilization of in-plane skyrmions by an electric field $\mathbf{E}$ is enabled by the balance between elastic energy and dielectric coupling energy, where dielectric coupling energetically favors $\mathbf{n}(\mathbf{r}) \parallel \mathbf{E}$ in a material with positive dielectric anisotropy. In the case of helical-background skyrmions, $\mathbf{E} \parallel \hat{\mathbf{z}}$ promotes vertical $\mathbf{n}(\mathbf{r})$ orientations and the expansion of skyrmions, which counters the shrinking tendency due to mismatch between twist deformation in the skyrmion and molecular pitch p . Beyond simply stabilizing $\mathbf{n}(\mathbf{r})$ configurations, modulation of $\mathbf{E}$ can also be used to control the dynamics of monopoles on which the skyrmions terminate. Here we focus on in-plane helical-background skyrmions because of their ease of manipulation and observation, though the same field-controlled dynamics can be straightforwardly applied to other geometrical configurations.

For a pair of a monopole and an anti-monopole at the ends of a skyrmion fragment, we define the separation velocity as the rate in the change of their distance of separation. Away from the equilibrium voltage at $U < 1.4$ V, the dielectric coupling is not strong enough to compensate for the cost in elastic energy in the skyrmion, and the skyrmion shrinks (Fig. 4A and movie S1). Whereas at $U > 1.4$ V, it becomes favorable in the overall free energy for the skyrmion to grow and acquire larger regions of vertical orientations in the system (Fig. 4B and movie S1). Thus, the separation velocity becomes positive and the width of the skyrmion, as determined in POM by the region deviating away from the helical background, is also visibly larger. The dependence of separation velocity on U is plotted in Fig. 4C. The magnitude of this separation velocity ≈ -60 $\mu\text{m/s}$ is largest when the skyrmion shrinks at $U = 0$ V, while at $U = 1.6$ V, the monopole pair separate at 20 $\mu\text{m/s}$. In the range $U < 1.6$ V, the skyrmions connecting monopole pairs remain

approximately linear in shape. Above $U = 1.6$ V, distortion in the helical background takes place in the form of undulations(22), resulting in deformation of the linear shape of the skyrmion and obstruction in the separation of monopoles. We note that when $U < 1.4$ V, the skyrmion shrinks and the monopole pair come together and annihilate, indicating the absence of an energy barrier between the monopole pairs. This contrasts with topological solitons, where $\mathbf{n}(\mathbf{r})$ is continuous and topologically nontrivial; erasing them requires generation and annihilation of singular defects, leading to an energy barrier and the so-called topological protection(21).

The dynamics and stability of skyrmions and monopoles can also be controlled by magnetic fields in chiral LC ferromagnets where magnetically monodomain nanoplates are uniformly dispersed at high concentrations in the hosting chiral LC (23). By forming a monodomain in the magnetization $\mathbf{m}(\mathbf{r})$ of ferromagnetic nanoplates and introducing strong homeotropic surface anchoring of LC molecules to the nanoplates (*Methods*), $\mathbf{m}(\mathbf{r})$ follows $\mathbf{n}(\mathbf{r})$ and physically vectorizes $\mathbf{n}(\mathbf{r})$ from a director field into a vector field. When an external magnetic field $\mathbf{H}$ is applied, $\mathbf{H}$ and $\mathbf{m}(\mathbf{r})$ couple linearly and compete with free-energy contributions from elastic and dielectric interactions. Figure 4, D and E, show the magnetic control of skyrmions and monopoles with $\mathbf{H}$ applied in different directions. When $\mathbf{H}$ is parallel to the average $\mathbf{m}(\mathbf{r})$ within a skyrmion, $\mathbf{m}_s$, which is also the magnetization at the top and bottom substrates (Fig. 2H), the linear magnetic coupling promotes the expansion of skyrmions and separation of monopoles (Fig. 4D). The width of the skyrmions also becomes visibly wider. When $\mathbf{H}$ is antiparallel to $\mathbf{m}_s$, skyrmions contract and eventually monopole pairs annihilate (Fig. 4E). Compared to electric-field control of monopoles, the coupling between $\mathbf{H}$ and $\mathbf{m}(\mathbf{r})$ is linear and polar, and can be applied with less constraint pertaining to sample geometry, enabling an orthogonal dimension in controlling

monopole dynamics.

Electric field controls interaction and assembly of vertical torons by modulating energetic landscape around solitons

Thus far, we have focused on in-plane skyrmions and monopoles because of their accessibility for manipulation and observation. However, vertical skyrmions along the normal of substrates and terminating at monopoles in a cell with homeotropic boundary conditions, also known as torons, also display manipulable interaction and assembly modulated by an electric field. In a homeotropic cell, when the confined chiral LC has an elastic anisotropy such that $K_{33}/K_{22} \leq 1$, i.e., the energetic penalty of bend deformation is smaller than that of twist deformation, the background $\mathbf{n}(\mathbf{r})$ can be switched between the helical state and the uniform state by an electric field. For a material with positive dielectric anisotropy, the background is helical at no field and can be unwound to the uniform state when an electric field is applied (16, 24). Vertical skyrmions stabilized in a helical background with no field display attractive pair-wise interaction and form a close-packed hexagonal lattice when several of them are in close proximity (Fig. 5A). This is in stark contrast to the repulsive interaction between vertical skyrmions stabilized in a uniform background(17, 20). When an electric field across the thickness of an LC cell was applied to unwind the helical background into a uniform background, the skyrmion-skyrmion interaction switched from attractive to repulsive (Fig. 5B). Computer simulations based on free-energy minimization reveal the detailed $\mathbf{n}(\mathbf{r})$ transformation when the background state changes between the helical state and the uniform state (Fig. 5, C and D). Before and after the transformation of the background state, the monopole pairs and skyrmion number in the horizontal cross-sections are conserved, preserving their respective $\pi_2(\mathbb{S}^2)$ topology. However, the energetic landscape in the

lateral dimensions changes. When the background is helical at no electric field, the free energy density at the periphery of the skyrmion is higher than that of the background; when the background is unwound by an electric field, the free energy density at the periphery of the skyrmion becomes lower than that of the background (Fig. 5E). To minimize total free energy, skyrmions (in a helical background) with a high-energy periphery comes closer together to share and reduce the volume with a high energy density, leading to an attractive interaction, whereas skyrmions (in the uniform background) with a low-energy periphery stay away from each other to increase the space with low energy density, leading to repulsion. We note that, in noncentrosymmetric chiral magnets, skyrmions in a uniformly polarized background are commonly observed, and skyrmions with attractive interaction have also been demonstrated in a conical background(25). In LCs, the switching of pair-wise interactions between skyrmions through background modulation can be recorded in real time due to their unparalleled experimental accessibility.

Discussion

In this work, we have presented the magnetic and electric control of the equilibrium configuration of $\pi_2(\mathbb{S}^2)$ topological configurations, namely skyrmions and monopoles (with their composite structures referred to as torons), and their dynamics in chiral LC materials. We found that $\pi_2(\mathbb{S}^2)$ point defects at the termini of skyrmions are indeed monopoles in the emergent magnetic field. We further demonstrated that LC skyrmions can take on versatile geometrical configurations under different boundary conditions and applied fields. Notably, in-plane skyrmions in planar-confined cells offer exceptional accessibility for observation and manipulation by external fields. Compared to vertical skyrmions in a homeotropically aligned cell (torons), in-plane skyrmions offer the

opportunity of controlling the separation of monopoles in the lateral dimensions, whereas the separation between the monopoles in vertical skyrmions is limited due to confinement by the substrates, though their lateral spatial extent can be controlled by fields(12, 26). To enhance the dynamical range of the electric control of skyrmions and monopoles, dual-frequency LCs may be used to allow the sign of LC dielectric anisotropy to be adjustable, such that positive (negative) dielectric anisotropy that promotes expansion (contraction) can be used in the same material(27). The linear coupling enabled by magnetic fields, distinct from the quadratic nature of dielectric coupling to electric fields, could also potentially allow for orientational control of solitons and monopoles.

Additionally, we showed that the pair-wise interaction between vertical skyrmions can be switched between attractive and repulsive, by altering the background configuration that embeds them through applying electric fields. We expect that similar switching can be achieved as well in LC ferromagnets using magnetic fields (as well as in magnetic solids), where the far-field $\mathbf{m}(\mathbf{r})$ can be polarized by $\mathbf{H}$ applied to the sample. Crossover between different pair-wise interactions of LC skyrmions has been observed between vertical and in-plane skyrmions in homeotropically aligned cells, albeit it was done differently through changing the dimension of the confined LC or anchoring(28). The field-controlled pair-wise interaction of LC skyrmions could potentially further couple with their recently observed squirming motion, activated by a modulated electric field, to facilitate versatile control and manipulation of LC topological solitons and defects(29).

It is informative to compare the emergent magnetic monopoles studied here to Dirac monopoles(15), as well as emergent monopoles in other condensed-matter systems such as those in spin ice and artificial spin ice(3, 30). Dirac showed magnetic monopoles can be consistent with quantum mechanics by considering a monopole as the end of a semi-infinitely long and intestinally

thin solenoid. The solenoid, also called the Dirac string, has no observable quantum mechanical Aharonov-Bohm effect when satisfying Dirac's quantization condition. The Dirac string is thus infinitesimally thin and undetectable. In contrast, emergent magnetic monopoles are sources or sinks of the emergent magnetic field with finite-sized "strings" in between. In $\mathbf{n}(\mathbf{r})$, emergent monopoles are $\pi_2(\mathbb{S}^2)$ point defects connected by $\pi_2(\mathbb{S}^2)$ skyrmions. Since the monopoles and skyrmions are imbedded in a topologically trivial background, the emergent flux is also quantized as a result of the $\pi_2(\mathbb{S}^2)$ field topology. The interaction between emergent magnetic monopoles depends on the energetics of the material field $\mathbf{n}(\mathbf{r})$. Notably, as shown above, the tension in the skyrmion string that connects monopoles can be controlled between positive and negative values by external electric fields or magnetic fields that couple to $\mathbf{n}(\mathbf{r})$. This is analogous to their counterparts in solid-state chiral magnets, where the transition into and out of the skyrmion A-phase is mediated by nucleation and annihilation of emergent magnetic monopole-anti-monopole pairs(5). Emergent monopoles in spin ice or artificial spin ice, on the other hand, are sources of the actual magnetic field(3, 30). When the local "ice rule" is violated in geometrically frustrated spin ice materials, monopoles emerge with their charge determined by the dipole moment and lattice parameters. Since skyrmion spin ice could be potentially formed in chiral molecular and colloidal LCs(31) or even magnetic solids, future explorations of the interplay between different types of monopoles may be possible too.

On the application side, the high degree of distortion in the alignment field around the emergent magnetic monopoles in this study and the central isotropic core are elastic-energy-enabled traps for micro- or nanoparticles(32, 33). Thus, potential technological applications could arise from dynamical control of skyrmions and monopoles, such as electro/magneto-optic devices, microfluidic devices, optical logic devices, etc. The widespread electro-optic applications of LCs

typically rely on switching between topologically trivial states that are homeomorphic one to another, but our findings point to the possibility of realizing controlled conditions for achieving novel electro-optic effects either by morphing topologically nontrivial configurations or even by changing topology, for example, through the monopole-mediated skyrmion nucleation and reconfiguration. Lastly, the study of dynamics and control of LC skyrmions and monopoles could shed light on the physics of topologically equivalent solitonic structures in other ordered systems, such as in the solid-state magnetic(5), ferroelectric(34), or optical fields(35).

Materials & Methods

Sample preparation

Chiral LC mixtures used in this work comprise 4-Cyano-4'-pentylbiphenyl (5CB, from EM Chemicals) added with the right-handed chiral dopant CB-15 (EM Chemicals). The helical pitch p of the mixture was measured in a wedge cell and ranged from 2.33 to 60 μm (36). For the LC mixture with reduced bend elasticity, 4',4''-(heptane-1,7-diyl)-dibiphenyl-4-carbonitrile (CB7CB; from SYNTHON Chemicals, Germany) was additionally added to the mixture at 40 wt%(37).

LC cells were assembled from substrates of glass slides or coverslips coated with indium tin oxide (ITO) and treated with polyimide PI2555 and SE5661 (both from Nissan Chemicals) for planar and homeotropic boundary conditions (anchoring) for LC director, respectively. The treatment involves spin-coating polyimide onto ITO glasses, followed by baking at 180°C for 1h. PI2555-coated glass slides or coverslips were additionally rubbed to impose unidirectional planar anchoring. Silica microbeads or microcylinders were used as spacers to assemble treated ITO glasses into cells with well-defined thicknesses ranging from 7 to 60 μm by UV-activated glue. Finally, metal wires were soldered to ITO glasses as electrodes. For LC cells where an in-plane electric field parallel to the confining substrates was applied, glasses without ITO coating were polyimide-treated and used as confining substrates of LCs, and a pair of ITO glasses with a gap of ~ 3 mm were assembled perpendicular to the substrates and soldered with wires. LC mixtures were introduced into cells via capillary force.

Electric control of LCs was achieved by connecting the electrodes of LC cells to a function generator (DS345; Stanford Research Systems) operating at a 1 kHz carrier frequency to preclude complex hydrodynamic effects. To achieve an ~ 1 V/ μm electric field in the in-plane direction, a high-voltage amplifier (Model 10/10B, Trek) was used to raise the voltage to ~ 3000 V. Magnetic

control of LC ferromagnets was achieved by permanent neodymium magnets (K&J Magnetics) or a previously reported home-built electromagnet system containing solenoids with machined cast-iron cores driven by power supplies (BOP20-5M, Kepco)(38, 39). The solenoids were arranged in Helmholtz coils to produce uniform magnetic fields up to ~ 30 mT. Magnetic fields used to control the dynamics of skyrmions and monopoles in LC ferromagnets were typically ~ 10 mT.

Laser generation and imaging of skyrmions and monopoles

Skyrmions and monopoles were controllably generated by laser tweezers that locally melt and align LC directors. The tweezers setup is based on an ytterbium-doped fiber laser (YLR-10-1064, IPG Photonics) and a phase-only spatial light modulator (P512-1064, Boulder Nonlinear Systems) integrated with an inverted optical microscope (IX81, Olympus)(17). Upon focusing the 1,064 nm infrared laser into the LC sample, local heating and optical realignment created initial $\mathbf{n}(\mathbf{r})$ that eventually relaxed into skyrmions and monopoles. In a planar-alignment cell, at $d = p = 10$ μm and using a 5CB-based chiral LC mixture, in-plane skyrmions in a helical background can be reliably generated at $U \approx 1.27$ V upon laser-induced melting, while in-plane skyrmions in a uniform background can be generated at an in-plane electric field of ~ 1 V/ μm . In a homeotropic-alignment cell, vertical torons in a uniform background can be generated without the applied electric field.

Bright-field microscopy and polarizing optical microscopy were performed using an Olympus IX-81 inverted microscope and a charge-coupled device camera (Flea-COL, from PointGrey Research)(17). Video-microscopy was conducted at a frame rate of 15 fps. Image analysis of monopole separation velocity was performed using the ridge detection plugin in ImageJ (National Institute of Health). Under different applied voltages, the skyrmion and monopoles as its end points in each POM video were identified. The separation velocity of the monopole-

antimonopole pair at different voltages, which varied only modestly with the length of the skyrmion, was extracted as the average velocity.

Nonlinear optical imaging via three-photon excitation fluorescence polarizing microscopy (3PEF-PM) was performed using a previously reported setup to unambiguously characterize 3D $\mathbf{n}(\mathbf{r})$ configurations in LCs(17, 21). Briefly, 3PEF-PM was performed using a setup built around the same IX-81 microscope with a Ti:sapphire laser (Chameleon Ultra II; Coherent) operating at 900 nm wavelength, 140 fs pulse duration, and 80 MHz repetition rate. 5CB and CB-15 molecules in the LC mixture were excited by the ultrashort pulses and the fluorescence was epi-collected by a 100× oil-immersion objective (NA = 1.44) and detected by a photomultiplier tube (H5784-20, Hamamatsu) after a 417/60-nm bandpass filter. The polarization state of the excitation was controlled by a pair of half-wave and quarter-wave retardation plates. In 3PEF-PM imaging, the image intensity scales as $\cos^6 \beta$, where β is the angle between the dipole moment of the LC molecule, orientating along $\mathbf{n}(\mathbf{r})$, and the polarization of the excitation light(40). LC mixtures used in samples for 3PEF-PM imaging comprise additionally 15% of diacrylate nematic reactive mesogen RM 257 (Merck) and 1% of UV-sensitive photoinitiator Irgacure 369 (Sigma-Aldrich), making them partially polymerizable. Before imaging, samples were polymerized by UV illumination and the unpolymerized components were replaced with immersion oil. This process preserves the $\mathbf{n}(\mathbf{r})$ configuration of the sample, while reducing the material's birefringence by an order of magnitude, thus minimizing imaging artifacts(41).

Synthesis and dispersion of magnetic nanoplates in LCs

Barium hexaferrite $\text{BaFe}_{11}\text{CrO}_{19}$ ferromagnetic nanoplates were synthesized by the hydrothermal method described in(42). Briefly, 0.01 M of $\text{Ba}(\text{NO}_3)_2$, 0.04 M of $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$, and 0.01 M of $\text{Cr}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ (all from Alfa Aesar), according to the nominal stoichiometry, were dissolved in

deionized (DI) water and co-precipitated by 2.72 M of NaOH (Alfa Aesar) aqueous solution with a final volume of 20 mL in a 25 mL Teflon-lined autoclave. The resulting material was hydrothermally heated to 220°C at a rate of 3°C/min, held at 220 °C for 1 h, and then cooled down to room temperature. The precipitated nanoplates were then washed with 10 wt% nitric acid and acetone and redispersed in 1 mL of DI water. The thickness of the as-synthesized nanoplates was determined to be ~10 nm and the average diameter ~105 nm using transmission electron microscopy. The nanoplates are magnetically monodomain with an magnetic moment approximately 1×10^{-17} Am², orthogonal to their large-area faces(12). The magnetic nanoplates were then surface-functionalized by 5 kDa silane-terminated polyethylene glycol (JemKem Technology) to increase their stability in LCs and introduce strong homeotropic anchoring to LC molecules.

To obtain LC ferromagnets, 15 µL of LC mixtures were mixed with 15 µL of ethanol and 15 µL of 1 wt% magnetic nanoplates dispersed in ethanol. The mixture was kept at 90°C for 3 h for the ethanol to fully evaporate before being cooled rapidly to nematic phase while vigorously disturbed to reduce formation of aggregates at domain boundaries. The ensuing mixture was then centrifuged at 2,200 rpm for 5 min to remove any residual aggregates. To achieve monodomain LC ferromagnets, an unwinding electric field (~0.5 V/µm) and a colinear magnetic field perpendicular to the substrates (~15 mT) were applied while filling the LC cell(43). The samples were confirmed to be magnetically monodomain by the response of $\mathbf{n}(\mathbf{r})$ to externally applied magnetic fields(38).

Numerical modeling

Numerical modeling of configurations and energetics of skyrmions and monopoles was based on energy minimization of Frank-Oseen free energy that describes energetic penalties of different

elastic deformations modes in $\mathbf{n}(\mathbf{r})$, supplemented by dielectric and magnetic (for LC ferromagnets) coupling terms(11, 16, 21). For a chiral LC, the elastic and dielectric terms read

$$F_{\text{elastic}} = \int d^3\mathbf{r} \left\{ \frac{K_{11}}{2} (\nabla \cdot \mathbf{n}(\mathbf{r}))^2 + \frac{K_{22}}{2} [\mathbf{n}(\mathbf{r}) \cdot (\nabla \times \mathbf{n}(\mathbf{r}))]^2 + \frac{K_{33}}{2} [\mathbf{n}(\mathbf{r}) \times (\nabla \times \mathbf{n}(\mathbf{r}))]^2 + \frac{2\pi K_{22}}{p} \mathbf{n}(\mathbf{r}) \cdot (\nabla \times \mathbf{n}(\mathbf{r})) \right\} \quad (1),$$

$$F_{\text{dielectric}} = -\frac{\epsilon_0 \epsilon_a}{2} \int d^3\mathbf{r} (\mathbf{E} \cdot \mathbf{n}(\mathbf{r}))^2 \quad (2).$$

Here, $\mathbf{n}(\mathbf{r})$ is the LC molecular alignment field, K_{11} , K_{22} , and K_{33} are the Frank elastic constants describing the energetic costs of splay, twist, and bend deformations, respectively, and p is the helical pitch of the material. Surface energy and saddle-splay deformation were not included by assuming strong boundary conditions on the surfaces, consistent with conditions in experiments. In the dielectric coupling term, ϵ_0 is vacuum permittivity, ϵ_a is the dielectric anisotropy of the LC and $\mathbf{E}$ is the applied electric field. In a chiral LC ferromagnet, the magnetization unit vector field $\mathbf{m}(\mathbf{r}) \equiv \mathbf{M}(\mathbf{r})/|\mathbf{M}(\mathbf{r})|$, where $\mathbf{M}(\mathbf{r})$ is the local magnetization, is assumed to be collinear with $\mathbf{n}(\mathbf{r})$ due to strong homeotropic surface anchoring of LC molecules to the magnetic nanoplates. Therefore, the elastic free energy and dielectric coupling of a chiral LC ferromagnet are identical to Eqs. (1) and (2), with $\mathbf{n}(\mathbf{r})$ substituted by $\mathbf{m}(\mathbf{r})$. An additional magnetic coupling term between $\mathbf{m}(\mathbf{r})$ and an applied magnetic field $\mathbf{H}$ reads

$$F_{\text{magnetic}} = -\mu_0 |\mathbf{M}(\mathbf{r})| \int d^3\mathbf{r} \mathbf{H} \cdot \mathbf{m}(\mathbf{r}). \quad (3)$$

Here, μ_0 is vacuum permeability and $|\mathbf{M}(\mathbf{r})| = \rho m_p$ is the magnetization of the material (approximated to be uniform) as the product of the nanoplate density ρ ($10/\mu\text{m}^3$) and the average magnetic moment of a nanoplate m_p ($1 \times 10^{-17} \text{ Am}^2$), both adopted from experiments (38, 39, 44).

Minimization of free energy can be performed iteratively using standard numerical methods of finite difference discretization in space and forward Euler method in time and has been implemented in commercial software and custom energy-minimization routines (16, 17, 21, 45). Previously relaxed configurations or those obtained from experimental 3PEF-PM images were used as initial conditions for $\mathbf{n}(\mathbf{r})$ or $\mathbf{m}(\mathbf{r})$. Briefly, $\mathbf{n}(\mathbf{r})$ or $\mathbf{m}(\mathbf{r})$ was updated iteratively from an initial configuration using Euler-Lagrange equation until the change in the spatial average of functional derivatives converged at an energy minimum. In all simulations, the computational volume was sampled by isotropic voxels on a cubic grid at 24 grid points per p and periodic boundary conditions were applied at the lateral faces of the volume.

Material parameters in simulations were chosen to match those of 5CB (Table S1) (21). To model the elasticity of the bend-reduced LC mixture containing 5CB and CB7CB, material parameters were set to be the same as those of 5CB, except that K_{33} was reduced to a smaller value such that $K_{33}/K_{22} \leq 1$. In the limit where $K \equiv K_{11} = K_{22} = K_{33}$, Eq. (1) can be rewritten into the following form,

$$F_{\text{elastic,1-const}} = \int d^3\mathbf{r} \left\{ \frac{K}{2} (\nabla \mathbf{n}(\mathbf{r}))^2 + \frac{2\pi K}{p} \mathbf{n}(\mathbf{r}) \cdot (\nabla \times \mathbf{n}(\mathbf{r})) \right\}. \quad (4)$$

Equation (4), or Frank-Oseen free energy with the so-called one-constant approximation, has been frequently adopted to simplify calculations of LC energetics, and takes an identical functional form to the micromagnetic Hamiltonian density of solid-state chiral magnets (with the nonlocal dipole-dipole interactions neglected)(46–48). The connection between the continuum energy functionals of LCs and solid-state magnets suggests that similar configurations and phenomena can be anticipated in these distinct physical systems.

The numerical calculation of $\mathbf{B}_{\text{em}}$ from $\mathbf{m}(\mathbf{r})$ configurations is based on the definition $(B_{\text{em}})_i \equiv (\hbar/2e)\epsilon^{ijk} \mathbf{m} \cdot (\partial_j \mathbf{m} \times \partial_k \mathbf{m})$. The numerical simulation of 3PEF-PM images is based

474 on the $\cos^6 \beta$ dependence of the angle between $\mathbf{n}(\mathbf{r})$ and the different polarization directions of
475 the excitation light. Both numerical functions implemented in MATLAB are available on Zenodo:
476 [10.5281/zenodo.10092838](https://doi.org/10.5281/zenodo.10092838).
477

References

1. A. Rajantie, The search for magnetic monopoles. *Phys. Today*. **69**, 40–46 (2016).
2. Y. M. Shnir, *Magnetic Monopoles* (Springer Berlin Heidelberg, 2006).
3. C. Castelnovo, R. Moessner, S. L. Sondhi, Magnetic monopoles in spin ice. *Nature*. **451**, 42–45 (2008).
4. M. W. Ray, E. Ruokokoski, S. Kandel, M. Möttönen, D. S. Hall, Observation of Dirac monopoles in a synthetic magnetic field. *Nature*. **505**, 657–660 (2014).
5. P. Milde, D. Köhler, J. Seidel, L. M. Eng, A. Bauer, A. Chacon, J. Kindervater, S. Mühlbauer, C. Pfleiderer, S. Buhrandt, C. Schütte, A. Rosch, Unwinding of a skyrmion lattice by magnetic monopoles. *Science* **340**, 1076–1080 (2013).
6. G. E. Volovik, Linear momentum in ferromagnets. *J. Phys. C: Solid State Phys.* **20**, L83 (1987).
7. S. Zhang, S. S.-L. Zhang, Generalization of the Landau-Lifshitz-Gilbert equation for conducting ferromagnets. *Phys. Rev. Lett.* **102**, 086601 (2009).
8. J. Zang, M. Mostovoy, J. H. Han, N. Nagaosa, Dynamics of skyrmion crystals in metallic thin films. *Phys. Rev. Lett.* **107**, 136804 (2011).
9. A. Hatcher, *Algebraic Topology* (Cambridge University Press, 2002).
10. J.-S. Wu, I. I. Smalyukh, Hopfions, heliknotons, skyrmions, torons and both abelian and nonabelian vortices in chiral liquid crystals. *Liq. Cryst. Rev.* 1–35 (2022).
11. P. M. Chaikin, T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, 1995).
12. Q. Zhang, P. J. Ackerman, Q. Liu, I. I. Smalyukh, Ferromagnetic switching of knotted vector fields in liquid crystal colloids. *Phys. Rev. Lett.* **115**, 097802 (2015).
13. J.-S. B. Tai, I. I. Smalyukh, Surface anchoring as a control parameter for stabilizing torons, skyrmions, twisted walls, fingers, and their hybrids in chiral nematics. *Phys. Rev. E*. **101**, 042702 (2020).
14. H. Zhao, J.-S. B. Tai, J.-S. Wu, I. I. Smalyukh, Liquid crystal defect structures with Möbius strip topology. *Nat. Phys.* **19**, 451–459 (2023).
15. P. A. M. Dirac, Quantised singularities in the electromagnetic field. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*. **133**, 60–72 (1931).

16. J.-S. B. Tai, J.-S. Wu, I. I. Smalyukh, Geometric transformation and three-dimensional hopping of Hopf solitons. *Nat. Commun.* **13**, 2986 (2022).
17. I. I. Smalyukh, Y. Lansac, N. A. Clark, R. P. Trivedi, Three-dimensional structure and multistable optical switching of triple-twisted particle-like excitations in anisotropic fluids. *Nat. Mater.* **9**, 139–145 (2010).
18. A. O. Leonov, K. Inoue, Homogeneous and heterogeneous nucleation of skyrmions in thin layers of cubic helimagnets. *Phys. Rev. B.* **98**, 054404 (2018).
19. Y. Liu, R. K. Lake, J. Zang, Binding a hopfion in a chiral magnet nanodisk. *Phys. Rev. B.* **98**, 174437 (2018).
20. D. Foster, C. Kind, P. J. Ackerman, J.-S. B. Tai, M. R. Dennis, I. I. Smalyukh, Two-dimensional skyrmion bags in liquid crystals and ferromagnets. *Nat. Phys.* **15**, 655–659 (2019).
21. J.-S. B. Tai, I. I. Smalyukh, Three-dimensional crystals of adaptive knots. *Science* **365**, 1449–1453 (2019).
22. B. I. Senyuk, I. I. Smalyukh, O. D. Lavrentovich, Undulations of lamellar liquid crystals in cells with finite surface anchoring near and well above the threshold. *Phys. Rev. E.* **74**, 011712 (2006).
23. A. Mertelj, D. Lisjak, M. Drofenik, M. Čopič, Ferromagnetism in suspensions of magnetic platelets in liquid crystal. *Nature* **504**, 237–241 (2013).
24. R. B. Meyer, Effects of electric and magnetic fields on the structure of cholesteric liquid crystals. *App. Phys. Lett.* **12**, 281–282 (2003).
25. J. C. Loudon, A. O. Leonov, A. N. Bogdanov, M. C. Hatnean, G. Balakrishnan, Direct observation of attractive skyrmions and skyrmion clusters in the cubic helimagnet Cu₂OSeO₃. *Phys. Rev. B.* **97**, 134403 (2018).
26. P. J. Ackerman, J. van de Lagemaat, I. I. Smalyukh, Self-assembly and electrostriction of arrays and chains of hopfion particles in chiral liquid crystals. *Nat. Commun.* **6**, 6012 (2015).
27. H. Xianyu, S.-T. Wu, C.-L. Lin, Dual frequency liquid crystals: a review. *Liquid Crystals* **36**, 717–726 (2009).
28. H. R. O. Sohn, S. M. Vlasov, V. M. Uzdin, A. O. Leonov, I. I. Smalyukh, Real-space observation of skyrmion clusters with mutually orthogonal skyrmion tubes. *Phys. Rev. B.* **100**, 104401 (2019).
29. P. J. Ackerman, T. Boyle, I. I. Smalyukh, Squirmer motion of baby skyrmions in nematic fluids. *Nat. Commun.* **8**, 673 (2017).

30. S. H. Skjærvø, C. H. Marrows, R. L. Stamps, L. J. Heyderman, Advances in artificial spin ice. *Nat. Rev. Phys.* **2**, 13–28 (2020).
31. A. Duzgun, C. Nisoli, Skyrmion Spin Ice in Liquid Crystals. *Phys. Rev. Lett.* **126**, 047801 (2021).
32. P. J. Ackerman, H. Mundoor, I. I. Smalyukh, J. van de Lagemaat, Plasmon–exciton interactions probed using spatial coentrapment of nanoparticles by topological singularities. *ACS Nano*. **9**, 12392–12400 (2015).
33. H. R. O. Sohn, P. J. Ackerman, T. J. Boyle, G. H. Sheeta, B. Fornberg, I. I. Smalyukh, Dynamics of topological solitons, knotted streamlines, and transport of cargo in liquid crystals. *Phys. Rev. E*. **97**, 052701 (2018).
34. S. Das, Y. L. Tang, Z. Hong, M. a. P. Gonçalves, M. R. McCarter, C. Klewe, K. X. Nguyen, F. Gómez-Ortiz, P. Shafer, E. Arenholz, V. A. Stoica, S.-L. Hsu, B. Wang, C. Ophus, J. F. Liu, C. T. Nelson, S. Saremi, B. Prasad, A. B. Mei, D. G. Schlom, J. Iñiguez, P. García-Fernández, D. A. Muller, L. Q. Chen, J. Junquera, L. W. Martin, R. Ramesh, Observation of room-temperature polar skyrmions. *Nature*. **568**, 368–372 (2019).
35. S. Tsesses, E. Ostrovsky, K. Cohen, B. Gjonaj, N. H. Lindner, G. Bartal, Optical skyrmion lattice in evanescent electromagnetic fields. *Science*. **361**, 993–996 (2018).
36. P. R. Gerber, On the determination of the cholesteric screw sense by the grandjean-cano-method. *Zeitschrift für Naturforschung A*. **35**, 619–622 (1980).
37. M. R. Tuchband, M. Shuai, K. A. Graber, D. Chen, L. Radzihovsky, A. Klitnick, L. Foley, A. Scarbrough, J. H. Porada, M. Moran, E. Korblova, D. M. Walba, M. A. Glaser, J. E. MacLennan, N. A. Clark, The twist-bend nematic phase of bent mesogenic dimer CB7CB and its mixtures. arXiv:1511.07523 (2015).
38. A. J. Hess, Q. Liu, I. I. Smalyukh, Optical patterning of magnetic domains and defects in ferromagnetic liquid crystal colloids. *App. Phys. Lett.* **107**, 071906 (2015).
39. Q. Liu, P. J. Ackerman, T. C. Lubensky, I. I. Smalyukh, Biaxial ferromagnetic liquid crystal colloids. *Proceedings of the National Academy of Sciences*. **113**, 10479–10484 (2016).
40. T. Lee, R. P. Trivedi, I. I. Smalyukh, Multimodal nonlinear optical polarizing microscopy of long-range molecular order in liquid crystals. *Opt. Lett.* **35**, 3447–3449 (2010).
41. J. S. Evans, P. J. Ackerman, D. J. Broer, J. van de Lagemaat, I. I. Smalyukh, Optical generation, templating, and polymerization of three-dimensional arrays of liquid-crystal defects decorated by plasmonic nanoparticles. *Phys. Rev. E*. **87**, 032503 (2013).
42. D. Lisjak, M. Drofenik, Chemical substitution—an alternative strategy for controlling the particle size of barium ferrite. *Crystal Growth & Design*. **12**, 5174–5179 (2012).

- 578 43. P. Medle Rupnik, D. Lisjak, M. Čopič, S. Čopar, A. Mertelj, Field-controlled structures
579 in ferromagnetic cholesteric liquid crystals. *Science Advances* **3**, e1701336 (2017).
- 580 44. J.-S. B. Tai, P. J. Ackerman, I. I. Smalyukh, Topological transformations of Hopf
581 solitons in chiral ferromagnets and liquid crystals. *Proceedings of the National Academy*
582 *of Sciences*. **115**, 921–926 (2018).
- 583 45. J. E. Anderson, P. J. Bos, P. E. Watson, *LC3D: Liquid Crystal Display 3-D Director*
584 *Simulator : Software and Technology Guide* (Artech House, Boston, 2001).
- 585 46. A. O. Leonov, Surface anchoring as a control parameter for shaping skyrmion or toron
586 properties in thin layers of chiral nematic liquid crystals and noncentrosymmetric
587 magnets. *Phys. Rev. E*. **104**, 044701 (2021).
- 588 47. J.-S. B. Tai, I. I. Smalyukh, Static Hopf solitons and knotted emergent fields in solid-
589 state noncentrosymmetric magnetic nanostructures. *Phys. Rev. Lett.* **121**, 187201 (2018).
- 590 48. R. Voinescu, J.-S. B. Tai, I. I. Smalyukh, Hopf solitons in helical and conical
591 backgrounds of chiral magnetic solids. *Phys. Rev. Lett.* **125**, 057201 (2020).
- 592

593 **Acknowledgements:** We acknowledge discussions and technical assistance of Qingkun Liu and
594 Haridas Mundoor. We are grateful to Patrick Davidson for providing CB7CB used at the initial
595 stages of this study.

596 **Funding:** This research was supported by the U.S. National Science Foundation grant DMR-
597 1810513.

598 **Author Contributions:** J.-S.B.T. performed experiments. A.J.H. synthesized magnetic
599 nanoplates. J.-S.B. T. and J.-S.W. performed numerical modelling. I.I.S. directed research and
600 provided funding. All authors contributed to the writing of the manuscript.

601 **Competing Interests:** The authors have no competing interests.

602 **Data Availability:** All data generated or analyzed during this study are included in the published
603 article and its Supplementary Information.

604 **Code Availability:** The code used for numerical calculations are available on Zenodo:
605 10.5281/zenodo.10092838.

606




Fig. 1. Monopoles and point defects. (A,B) A hedgehog (A) and a hyperbolic (B) point defect in the material alignment field $\mathbf{n}(\mathbf{r})$. (C) Emergent magnetic field $\mathbf{B}_{\text{em}}$ derived from the ansatzes of the hedgehog (A) and the hyperbolic point defect (B) in $\mathbf{n}(\mathbf{r})$ both shows radial monopole-like field configuration. (D,E) Computer-simulated $\mathbf{n}(\mathbf{r})$ configuration of a toron in a cell with a perpendicular boundary condition shown in the vertical cross-section parallel to the uniform far-field $\mathbf{n}_0 \parallel \hat{z}$ (D) and horizontal cross-sections at different positions along z indicated in (D) and their respective skyrmion number N_{sk} . Point defects (monopoles) in (D) are circled in red with their charges labeled. Here $\mathbf{n}(\mathbf{r})$ is visualized by colored arrows using the color scheme in (G). The blue slabs indicate the substrates confining LCs. (F) $\mathbf{B}_{\text{em}}$ or equivalently the 3D skyrmion number density derived from the $\mathbf{n}(\mathbf{r})$ configuration of the toron in (D) shown by the isosurfaces of the scalar magnitude of $\mathbf{B}_{\text{em}}$ and streamlines of $\mathbf{B}_{\text{em}}$ connecting the positively-charged (top) and the negatively-charged (bottom) monopoles. (G) Order-parameter space $\mathbb{S}^2$ colored according to the vectorized $\mathbf{n}(\mathbf{r})$ orientation.




Fig. 2. Toron in a uniform or helical background and the associated monopoles in emergent magnetic field. (A,B) Torons are formed by monopole-associated skyrmions which can be embedded in a uniform far-field $\mathbf{n}_0$ (A) and a helical field background with constant helical axis χ_0 (B), respectively. (C,D) Skyrmion number density derived from $\mathbf{n}(\mathbf{r})$ in (A) and (B), respectively. (E) Computer-simulated cross-sections of $\mathbf{n}(\mathbf{r})$ of an in-plane skyrmion terminating at two monopoles (circled in red with the sign of the charge labeled) and embedded in a uniform far-field $\mathbf{n}_0 \parallel \hat{y}$. (F) Detailed $\mathbf{n}(\mathbf{r})$ around each monopole in (E). (G) Emergent field $\mathbf{B}_{em}$ derived from $\mathbf{n}(\mathbf{r})$ in (E) shown by streamlines of $\mathbf{B}_{em}$ colored by its magnitude. (H) Computer-simulated cross-sections of $\mathbf{n}(\mathbf{r})$ of a skyrmion terminating at two monopoles (circled in red with the sign of

633 the charge labeled) and embedded in a helical far-field with helical axis $\chi_0 \parallel \hat{z}$. **(I)** Detailed $\mathbf{n}(\mathbf{r})$
634 around each monopole in (H). **(J)** Emergent field $\mathbf{B}_{\text{em}}$ derived from $\mathbf{n}(\mathbf{r})$ in (H) shown by
635 streamlines of $\mathbf{B}_{\text{em}}$, colored by its magnitude. In simulations, $d/p = 0.8$ in (E-G), and $d/p = 1$
636 and $U = 1.5$ V in (H-J), respectively.
637




Fig. 3. Torons comprising skyrmion and monopoles in confined chiral LCs. (A) POM images of an individual toron with a skyrmion (A) in a helical far-field background (constant helical axis χ_0) terminating at two monopoles (B) Experimental (exp.) and computer-simulated (sim.) 3PEF-PM images in the vertical cross-section through the skyrmion. (C) 1D lattice of skyrmions in a helical background. (D) Experimental (exp.) and computer-simulated (sim.) 3PEF-PM images in the vertical cross-section through an 1D lattice of helical-background skyrmions. (E) POM image of a skyrmion lattice with defects. The sample in (E) was polymerized and washed before imaging, reducing the birefringence. (F,G) 3PEF-PM images of skyrmions and monopoles in horizontal and vertical cross-sections indicated in (E). The skyrmion number N_{sk} of each vertical cross-section in the region defined by the gray box are shown. (H,I) POM images of an individual skyrmion in a uniform far-field background n_0 terminating at two monopoles with variable length under a smaller (H) or a larger (I) in-plane electric field. Scale bars are 10 μm in A, B, D, F and 30 μm in C, E, H. Sample thicknesses are 30 μm in (C) and (E) and 10 μm otherwise. $d/p = 1$ in all samples. The polarization states of the linearly polarized excitation light for 3PEF-PM in B,D,F,G are marked in red.




Fig. 4. Field-controlled dynamics of monopoles at skyrmion ends. (A,B) Snapshots of POM images of two monopoles connected by a skyrmion in a helical background moving close to (A) and away from (B) each other at $U = 0$ V and $U = 1.5$ V, respectively. (C) Separation velocity of the monopole-antimonopole pair vs. the applied voltage U . The error in measuring velocity is ± 2 $\mu\text{m/s}$. (D,E) Snapshots of POM images of two monopoles connected by a skyrmion in a helical background moving close to (D) and away from (E) each other when an external magnetic field $\mathbf{H}$ is applied parallel or antiparallel to magnetization $\mathbf{m}_s$, the average $\mathbf{m}(\mathbf{r})$ within the skyrmion, respectively. Scale bars are 20 μm in (A,B) and 50 μm in (D,E). Sample thicknesses are 10 μm in (A) and (B) and 60 μm in (D) and (E). $d/p = 1$ in all samples.




Fig. 5 | Controlling pair-wise interaction between vertical skyrmions. (A) POM images of a vertical skyrmion in a helical background with perpendicular boundary conditions individually (left) and aggregating into a close-packed crystallite (right). (B) The pair-wise interaction between skyrmions is altered by changing the embedding far-field between helical (up) and uniform (bottom) through modulating the applied voltage U . (C) Preimages of $\mathbf{n}(\mathbf{r})$ orientations along $\pm\hat{z}$ and $\pm\hat{y}$ colored according to order-parameter space S^2 shown in Fig. 1G for computer-simulated skyrmions in a helical background (left) and a uniform background (right), respectively. (D) Computer-simulated vertical cross-section of $\mathbf{n}(\mathbf{r})$ of a vertical skyrmion terminating at monopoles in a helical (left) and a uniform (right) background, respectively. Monopoles are circled in red with their charges labeled. (E) Free-energy density of the computer-simulated skyrmion field configurations shown in (C) and (D) in a helical (left) and a uniform (right) background, respectively. The energy density is averaged over thickness (z direction) and excludes regions containing singular monopoles, shown in units of Kp , the product of the average elastic constant and pitch. In experiments and simulations, $d/p = 3$ and $d = 10 \mu\text{m}$. $\sqrt{K_{33}/K_{22}} = 0.8$ and $U = 4.1 \text{ V}$ were used in simulations. Scale bars are $5 \mu\text{m}$.

- 685 1. A. Rajantie, The search for magnetic monopoles. *Physics Today* **69**, 40–46 (2016).
- 686 2. Y. M. Shnir, *Magnetic Monopoles* (Springer Berlin Heidelberg, 2006;
687 <https://books.google.com/books?id=G5huCG0k2dAC>) *Theoretical and Mathematical*
688 *Physics*.
- 689 3. C. Castelnovo, R. Moessner, S. L. Sondhi, Magnetic monopoles in spin ice. *Nature* **451**, 42–
690 45 (2008).
- 691 4. M. W. Ray, E. Ruokokoski, S. Kandel, M. Möttönen, D. S. Hall, Observation of Dirac
692 monopoles in a synthetic magnetic field. *Nature* **505**, 657–660 (2014).
- 693 5. P. Milde, D. Köhler, J. Seidel, L. M. Eng, A. Bauer, A. Chacon, J. Kindervater, S.
694 Mühlbauer, C. Pfleiderer, S. Buhrandt, C. Schütte, A. Rosch, Unwinding of a Skyrmion
695 Lattice by Magnetic Monopoles. *Science* **340**, 1076–1080 (2013).
- 696 6. G. E. Volovik, Linear momentum in ferromagnets. *J. Phys. C: Solid State Phys.* **20**, L83
697 (1987).
- 698 7. S. Zhang, S. S.-L. Zhang, Generalization of the Landau-Lifshitz-Gilbert Equation for
699 Conducting Ferromagnets. *Phys. Rev. Lett.* **102**, 086601 (2009).
- 700 8. J. Zang, M. Mostovoy, J. H. Han, N. Nagaosa, Dynamics of Skyrmion Crystals in Metallic
701 Thin Films. *Phys. Rev. Lett.* **107**, 136804 (2011).
- 702 9. A. Hatcher, *Algebraic Topology* (Cambridge University Press, 2002).
- 703 10. J.-S. Wu, I. I. Smalyukh, Hopfions, heliknotons, skyrmions, torons and both abelian and
704 nonabelian vortices in chiral liquid crystals. *Liquid Crystals Reviews* **0**, 1–35 (2022).
- 705 11. P. M. Chaikin, T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge
706 University Press, 1995; https://books.google.com/books?id=Bw_ooQEACAAJ).
- 707 12. Q. Zhang, P. J. Ackerman, Q. Liu, I. I. Smalyukh, Ferromagnetic Switching of Knotted
708 Vector Fields in Liquid Crystal Colloids. *Phys. Rev. Lett.* **115**, 097802 (2015).
- 709 13. J.-S. B. Tai, I. I. Smalyukh, Surface anchoring as a control parameter for stabilizing torons,
710 skyrmions, twisted walls, fingers, and their hybrids in chiral nematics. *Phys. Rev. E* **101**,
711 042702 (2020).
- 712 14. H. Zhao, J.-S. B. Tai, J.-S. Wu, I. I. Smalyukh, Liquid crystal defect structures with Möbius
713 strip topology. *Nat. Phys.* **19**, 451–459 (2023).
- 714 15. P. A. M. Dirac, Quantised singularities in the electromagnetic field,. *Proceedings of the*
715 *Royal Society of London. Series A, Containing Papers of a Mathematical and Physical*
716 *Character* **133**, 60–72 (1931).

- 717 16. J.-S. B. Tai, J.-S. Wu, I. I. Smalyukh, Geometric transformation and three-dimensional
718 hopping of Hopf solitons. *Nat Commun* **13**, 2986 (2022).
- 719 17. I. I. Smalyukh, Y. Lansac, N. A. Clark, R. P. Trivedi, Three-dimensional structure and
720 multistable optical switching of triple-twisted particle-like excitations in anisotropic fluids.
721 *Nature Mater* **9**, 139–145 (2010).
- 722 18. A. O. Leonov, K. Inoue, Homogeneous and heterogeneous nucleation of skyrmions in thin
723 layers of cubic helimagnets. *Phys. Rev. B* **98**, 054404 (2018).
- 724 19. Y. Liu, R. K. Lake, J. Zang, Binding a hopfion in a chiral magnet nanodisk. *Phys. Rev. B* **98**,
725 174437 (2018).
- 726 20. D. Foster, C. Kind, P. J. Ackerman, J.-S. B. Tai, M. R. Dennis, I. I. Smalyukh, Two-
727 dimensional skyrmion bags in liquid crystals and ferromagnets. *Nat. Phys.* **15**, 655–659
728 (2019).
- 729 21. J.-S. B. Tai, I. I. Smalyukh, Three-dimensional crystals of adaptive knots. *Science* **365**,
730 1449–1453 (2019).
- 731 22. B. I. Senyuk, I. I. Smalyukh, O. D. Lavrentovich, Undulations of lamellar liquid crystals in
732 cells with finite surface anchoring near and well above the threshold. *Phys. Rev. E* **74**,
733 011712 (2006).
- 734 23. A. Mertelj, D. Lisjak, M. Drofenik, M. Čopič, Ferromagnetism in suspensions of magnetic
735 platelets in liquid crystal. *Nature* **504**, 237–241 (2013).
- 736 24. R. B. Meyer, Effects of electric and magnetic fields on the structure of cholesteric liquid
737 crystals. *Applied Physics Letters* **12**, 281–282 (2003).
- 738 25. J. C. Loudon, A. O. Leonov, A. N. Bogdanov, M. C. Hatnean, G. Balakrishnan, Direct
739 observation of attractive skyrmions and skyrmion clusters in the cubic helimagnet
740 $\text{Cu}_{1-x}\text{Fe}_x\text{SeO}_3$. *Phys. Rev. B* **97**, 134403 (2018).
- 741 26. P. J. Ackerman, J. van de Lagemaat, I. I. Smalyukh, Self-assembly and electrostriction of
742 arrays and chains of hopfion particles in chiral liquid crystals. *Nat Commun* **6**, 6012 (2015).
- 743 27. H. Xianyu, S.-T. Wu, C.-L. Lin, Dual frequency liquid crystals: a review. *Liquid Crystals* **36**,
744 717–726 (2009).
- 745 28. H. R. O. Sohn, S. M. Vlasov, V. M. Uzdin, A. O. Leonov, I. I. Smalyukh, Real-space
746 observation of skyrmion clusters with mutually orthogonal skyrmion tubes. *Phys. Rev. B*
747 **100**, 104401 (2019).
- 748 29. P. J. Ackerman, T. Boyle, I. I. Smalyukh, Squirming motion of baby skyrmions in nematic
749 fluids. *Nat Commun* **8**, 673 (2017).

- 750 30. S. H. Skjærvø, C. H. Marrows, R. L. Stamps, L. J. Heyderman, Advances in artificial spin
751 ice. *Nat Rev Phys* **2**, 13–28 (2020).
- 752 31. A. Duzgun, C. Nisoli, Skyrmion Spin Ice in Liquid Crystals. *Phys. Rev. Lett.* **126**, 047801
753 (2021).
- 754 32. P. J. Ackerman, H. Mundoor, I. I. Smalyukh, J. van de Lagemaat, Plasmon–Exciton
755 Interactions Probed Using Spatial Coentrapment of Nanoparticles by Topological
756 Singularities. *ACS Nano* **9**, 12392–12400 (2015).
- 757 33. H. R. O. Sohn, P. J. Ackerman, T. J. Boyle, G. H. Sheetah, B. Fornberg, I. I. Smalyukh,
758 Dynamics of topological solitons, knotted streamlines, and transport of cargo in liquid
759 crystals. *Phys. Rev. E* **97**, 052701 (2018).
- 760 34. S. Das, Y. L. Tang, Z. Hong, M. a. P. Gonçalves, M. R. McCarter, C. Klewe, K. X. Nguyen,
761 F. Gómez-Ortiz, P. Shafer, E. Arenholz, V. A. Stoica, S.-L. Hsu, B. Wang, C. Ophus, J. F.
762 Liu, C. T. Nelson, S. Saremi, B. Prasad, A. B. Mei, D. G. Schlom, J. Íñiguez, P. García-
763 Fernández, D. A. Muller, L. Q. Chen, J. Junquera, L. W. Martin, R. Ramesh, Observation
764 of room-temperature polar skyrmions. *Nature* **568**, 368–372 (2019).
- 765 35. S. Tsesses, E. Ostrovsky, K. Cohen, B. Gjonaj, N. H. Lindner, G. Bartal, Optical skyrmion
766 lattice in evanescent electromagnetic fields. *Science* **361**, 993–996 (2018).
- 767 36. P. R. Gerber, On the Determination of the Cholesteric Screw Sense by the Grandjean-Cano-
768 Method. *Zeitschrift für Naturforschung A* **35**, 619–622 (1980).
- 769 37. M. R. Tuchband, M. Shuai, K. A. Graber, D. Chen, L. Radzihovsky, A. Klitnick, L. Foley,
770 A. Scarbrough, J. H. Porada, M. Moran, E. Korblova, D. M. Walba, M. A. Glaser, J. E.
771 MacLennan, N. A. Clark, The twist-bend nematic phase of bent mesogenic dimer CB7CB
772 and its mixtures. arXiv arXiv:1511.07523 [Preprint] (2015).
773 <https://doi.org/10.48550/arXiv.1511.07523>.
- 774 38. A. J. Hess, Q. Liu, I. I. Smalyukh, Optical patterning of magnetic domains and defects in
775 ferromagnetic liquid crystal colloids. *Applied Physics Letters* **107**, 071906 (2015).
- 776 39. Q. Liu, P. J. Ackerman, T. C. Lubensky, I. I. Smalyukh, Biaxial ferromagnetic liquid crystal
777 colloids. *Proceedings of the National Academy of Sciences* **113**, 10479–10484 (2016).
- 778 40. T. Lee, R. P. Trivedi, I. I. Smalyukh, Multimodal nonlinear optical polarizing microscopy of
779 long-range molecular order in liquid crystals. *Opt. Lett., OL* **35**, 3447–3449 (2010).
- 780 41. J. S. Evans, P. J. Ackerman, D. J. Broer, J. van de Lagemaat, I. I. Smalyukh, Optical
781 generation, templating, and polymerization of three-dimensional arrays of liquid-crystal
782 defects decorated by plasmonic nanoparticles. *Phys. Rev. E* **87**, 032503 (2013).
- 783 42. D. Lisjak, M. Drofenik, Chemical Substitution—An Alternative Strategy for Controlling the
784 Particle Size of Barium Ferrite. *Crystal Growth & Design* **12**, 5174–5179 (2012).

- 785 43. P. Medle Rupnik, D. Lisjak, M. Čopič, S. Čopar, A. Mertelj, Field-controlled structures in
786 ferromagnetic cholesteric liquid crystals. *Science Advances* **3**, e1701336 (2017).
- 787 44. J.-S. B. Tai, P. J. Ackerman, I. I. Smalyukh, Topological transformations of Hopf solitons in
788 chiral ferromagnets and liquid crystals. *Proceedings of the National Academy of Sciences*
789 **115**, 921–926 (2018).
- 790 45. J. E. Anderson, P. J. Bos, P. E. Watson, *LC3D: Liquid Crystal Display 3-D Director*
791 *Simulator : Software and Technology Guide* (Artech House, Boston, 2001;
792 <http://catdir.loc.gov/catdir/toc/fy0904/2009286012.html>) *Artech House optoelectronics*
793 *library*.
- 794 46. A. O. Leonov, Surface anchoring as a control parameter for shaping skyrmion or toron
795 properties in thin layers of chiral nematic liquid crystals and noncentrosymmetric magnets.
796 *Phys. Rev. E* **104**, 044701 (2021).
- 797 47. J.-S. B. Tai, I. I. Smalyukh, Static Hopf Solitons and Knotted Emergent Fields in Solid-State
798 Noncentrosymmetric Magnetic Nanostructures. *Phys. Rev. Lett.* **121**, 187201 (2018).
- 799 48. R. Voinescu, J.-S. B. Tai, I. I. Smalyukh, Hopf Solitons in Helical and Conical Backgrounds
800 of Chiral Magnetic Solids. *Phys. Rev. Lett.* **125**, 057201 (2020).

801