

## Visible Light-Induced Reactions of Diazo Compounds and Their Precursors

Ziyan Zhang and Vladimir Gevorgyan\*



Cite This: *Chem. Rev.* 2024, 124, 7214–7261




Read Online

ACCESS |

Metrics & More

Article Recommendations

**ABSTRACT:** In recent years, visible light-induced reactions of diazo compounds have attracted increasing attention in organic synthesis, leading to improvement of existing reactions, as well as to the discovery of unprecedented transformations. Thus, photochemical or photocatalytic generation of both carbenes and radicals provide milder tools toward these key intermediates for many valuable transformations. However, the vast majority of the transformations represent new reactivity modes of diazo compounds, which are achieved by the photochemical decomposition of diazo compounds and photoredox catalysis. In particular, the use of a redox-active photocatalysts opens the avenue to a plethora of radical reactions. The application of these methods to diazo compounds led to discovery of transformations inaccessible by the classical reactivity associated with carbenes and metal carbenes. In most cases, diazo compounds act as radical sources but can also serve as radical acceptors. Importantly, the described processes operate under mild, practical conditions. This Review describes this subfield of diazo compound chemistry, particularly focusing on recent advancements.



### CONTENTS


1. Introduction	7215	4.1. Introduction	7236
2. Diazo Compounds As Singlet Carbene Precursors	7216	4.2. Radical Addition to Alkenes	7236
2.1. Introduction	7216	4.2.1. Giese- or Heck-Type Reactions	7236
2.2. The Wolff Rearrangement	7216	4.2.2. 1,2-Difunctionalization of Alkenes	7238
2.2.1. Reactions with Nucleophiles	7216	4.3. Radical Addition to Heteroarenes	7239
2.2.2. Formal Cycloadditions	7218	4.4. Radical–Radical Recombinations	7240
2.3. [2 + 1] Cycloadditions	7219	4.5. Other Radical Reactions	7240
2.3.1. Cyclopropanation	7219	5. Diazo Compounds as Diazomethyl Radical Precursors	7242
2.3.2. Cyclopropenation	7221	5.1. Introduction	7242
2.4. C–H Insertions	7222	5.2. Radical Addition to Arenes	7242
2.5. X–H Insertions	7223	5.3. Heterocyclization	7242
2.5.1. B–H Insertions	7223	6. Diazo Compounds As Radical Acceptors	7244
2.5.2. N–H Insertions	7223	7. Diazo Compounds as Nucleophiles	7246
2.5.3. O–H Insertions	7224	8. Miscellaneous Reactions of Diazo Compounds	7248
2.5.4. Si–H Insertions	7224	9. Precursors of Diazo Compounds	7251
2.5.5. S–H Insertions	7225	9.1. Diazirines	7251
2.6. Ylides Formation	7225	9.2. <i>N</i> -Tosylhydrazones	7251
2.6.1. Nitrogen Ylides	7226	9.3. Pyridotriazoles	7253
2.6.2. Oxygen Ylides	7227		
2.6.3. Sulfur Ylides	7229		
2.7. Miscellaneous Reactions	7231		
3. Diazo Compounds As Triplet Carbene Precursors	7234		
4. Diazo Compounds as Alkyl or Vinyl Radical Precursors	7236		

Received: November 26, 2023

Revised: April 20, 2024

Accepted: April 30, 2024

Published: May 16, 2024



9.4. 1,3,4-Oxadiazolines	7253
9.5. ( $\beta$ -Amino- $\alpha,\alpha$ -difluoroethyl)phosphonates	7253
10. Conclusions	7254
Author Information	7254
Corresponding Author	7254
Author	7254
Notes	7254
Biographies	7254
Acknowledgments	7255
References	7255

## 1. INTRODUCTION


Diazo compounds are versatile reagents in organic synthesis. In particular, they are extensively utilized as precursors for generating free carbene or metal carbenoid intermediates. These reactive species can undergo a variety of potent synthetic transformations, including cyclopropanations, X–H insertions, and ylide formations. The earliest developed techniques employ thermal conditions, although their applicability is limited due to the harsh reaction conditions and low selectivity.<sup>1</sup> Consequently, transition metal catalysis has emerged as a powerful tool, providing access to metal carbenes whose electronic properties and stabilities can be easily tuned. In this scenario, transformations frequently occur at room temperature. Importantly, the careful design of catalysts and ligands enables exceptional control over selectivity and reactivity, offering opportunities for asymmetric reactions. The topic of transition metal-catalyzed reactions of diazo compounds has been extensively reviewed.<sup>2</sup>

In contrast to the numerous protocols reported on thermal processes throughout the years, photochemical studies with diazo compounds have been performed mostly several decades ago.<sup>3,4</sup> Contemporary applications were relatively scarce until recently. Photolysis is a critical tool in investigating denitrogenative reactions of diazo compounds within the realm of physical organic chemistry. Thus, it has played an important role in establishing the basis for contemporary carbene chemistry. Matrix isolation techniques<sup>5</sup> and ultrafast spectroscopy studies significantly contributed to the experimental determination of the structural and electronic properties of carbene intermediates. Platz and co-workers conducted a series of studies<sup>6–10</sup> on ultrafast photolysis of diazo compounds to carbenes and their subsequent reactions, such as cyclopropanation.<sup>9</sup> They observed a singlet carbene produced directly from the excited state of a diazo precursor. Additionally, they studied dynamics of solvation of the singlet carbene.<sup>10</sup> The initial absorbance of the carbene is solvent dependent and is red-shifted in coordinating solvents, such as alcohols, indicating the formation of hydrogen bonds. However, it is important to note that these approaches produced carbene species from diazo compounds using high-energy UV irradiation. This typically required special medium- and high-pressure xenon and mercury lamps in combination with monochromators or filters. Given that the majority of organic molecules are capable of absorbing UV light, exposing a reaction to UV radiation can induce simultaneous and competitive excitation of multiple compounds, such as reagents, intermediates, products, and even solvents. These harsh reaction conditions led to numerous side reactions, thus limiting the synthetic applications of the methods.

Recently, the application of visible light in organic synthesis has garnered tremendous attention and evolved into a powerful

tool for developing a wide range of transformations. Consequently, interests in the chemistry of diazo compounds under visible light-induced conditions have been reignited, as evidenced by several review articles.<sup>11</sup> Diazo compounds are often colored molecules with their absorption strongly dependent on the substituents of the carbon bearing the diazo functional group. Notably, the presence of a donor group, often an aryl ring, induces a bathochromic shift in the absorption band. Thus, donor–acceptor diazo compounds, such as aryl diazoacetates, exhibit strong absorption in the visible light region (Scheme 1A). Furthermore, the donor–

### Scheme 1. Diazo Compounds under Visible Light Irradiation



acceptor carbenes are capable of very efficient reactions without the need for stabilization by a transition metal, as was seen in thermal studies conducted a few years earlier.<sup>12–15</sup> In 2018, Jurberg and Davies pioneered a photochemical method to generate carbenes from aryl diazoacetates using blue light.<sup>16</sup> Since then, various transformations of aryl diazoacetates have been developed under mild conditions, utilizing visible light as the sole energy source at ambient temperature, and frequently without the need for a catalyst. Theoretical studies indicate that the preferred electronic configuration of carbenes originating from aryl diazoacetates varies based on the aryl substituent. For instance, a singlet ground state is preferred in the presence of a donor substituent like a *para*-methoxy group, whereas a triplet ground state is favored with an acceptor substituent such as a *para*-nitro group.<sup>17</sup> While earlier reports

primarily focused on aryl diazoacetate substrates, recent developments in the field have demonstrated the possibilities of using donor–donor and acceptor diazo compounds for carbene generation.

Generally, when exposed to light, diazo compounds in the ground state undergo photoexcitation to singlet state (Scheme 1B).<sup>6,7,9</sup> Subsequently, they could follow different decomposition pathways, leading to the formation of carbenes in either the singlet or triplet states, which exhibit orthogonal reactivity. In the former case, the singlet diazo compound undergoes denitrogenation, producing a singlet carbene with two nonbonding electrons at the divalent carbon. The two electrons have antiparallel spins, occupying a single hybridized orbital while leaving a p orbital vacant. Thus, determined by the electronic properties of nearby substituents, singlet carbenes exhibit ambiphilic behavior, acting as both electrophiles and nucleophiles. On the other hand, triplet carbenes can be formed through two mechanisms: one involves the singlet carbene converting into a triplet carbene via intersystem crossing (ISC), while the other entails the singlet diazo compound to its triplet form through intersystem crossing, followed by decomposing into the triplet carbene. These pathways to triplet carbenes are typically facilitated by energy transfer from a photosensitizer. Unlike the singlet state, triplet carbenes have unpaired electrons with parallel spins distributed across two different orbitals. As a result, they are commonly classified as diradicals and typically participate in stepwise reactions, including hydrogen atom abstraction or nonstereospecific addition to olefins.

Importantly, in combination with photoredox catalysis,<sup>18–20</sup> diazo compounds can serve as precursors for various radical species under visible light irradiation (Scheme 1C). This has led to the discovery of transformations that were previously inaccessible through classical free carbene and metal carbene reactivity. Thus, diazo compounds undergo efficient single-electron reduction by a photocatalyst to produce C-centered radicals with the extrusion of dinitrogen, even in cases where diazo compounds are transparent in the visible light region. Furthermore, specifically designed diazoalkanes that possess a redox-active leaving group, such as a hypervalent iodine or sulfonium functionalities, yield diazomethyl radicals under visible light irradiation, thus preserving the valuable diazo functionality. Additionally, other photocatalytic transformations, in which diazo compounds can act as radical acceptors or nucleophiles have also attracted increasing attention in recent years.


This review intends to offer readers an overview of this emerging field, spanning from its early development to recent advances. As each reactive intermediate produced from diazo compounds exhibits unique reactivity, we have organized this review based on these reactive intermediates. Photoinduced reactions of diazo compound precursors are also covered. We have restricted our discussion to photochemical or photocatalytic methods that utilize visible light or near-visible light (>390 nm). Therefore, UV-induced transformations, which requires the use of specialized glassware, are not included in this discussion. It is worth noting that, given the typical focus of newly developed synthetic methodologies on expanding the scope of reactions, certain mechanistic aspects of these protocols may remain unclear. Therefore, the reaction mechanisms presented herein represent reasonable proposals by the authors and may include speculative steps.

## 2. DIAZO COMPOUNDS AS SINGLET CARBENE PRECURSORS

### 2.1. Introduction

The majority of the reactions covered in this review utilize diazo compounds as precursors for singlet carbenes, which serve as crucial intermediates in various transformations. These transformations include the Wolff rearrangement, [2 + 1] cycloadditions, insertions into C–H and X–H bonds, and the formation of ylides (Scheme 2). While many of these reactions

**Scheme 2. Diverse Transformations of Singlet Carbenes**



are well-established in classical metal carbene chemistry, the photochemical method for generating free carbenes has proven to be a valuable alternative, at times demonstrating complementary reactivities.


### 2.2. The Wolff Rearrangement

The Wolff rearrangement is one of the most extensively studied reactions of  $\alpha$ -diazoketones.<sup>21</sup> Initially, this rearrangement was primarily employed to prepare homologues of carboxylic acids, serving as a key step in Arndt–Eistert reaction. The process starts with the extrusion of dinitrogen, followed by a 1,2-shift to afford highly reactive ketenes,<sup>22</sup> which can occur either through a concerted mechanism or in a stepwise manner involving a carbene intermediate. Many mechanistic investigations have been conducted by aiming at determining whether migration occurs in a concerted fashion with the loss of dinitrogen.<sup>21</sup> The conclusion that has emerged is that a carbene is formed in photochemical reactions,<sup>23</sup> whereas the reaction is concerted under thermal conditions.

Traditionally, the Wolff rearrangement of  $\alpha$ -diazoketones is promoted by thermolysis, photolysis, or metal ions, with the latter two methods being more widely employed.<sup>21</sup> However, these approaches often require high-energy UV lights or metal salts. Recent advances have shown that Wolff rearrangements can be efficiently conducted under visible light irradiation. The resulting ketene then reacts with various nucleophiles to yield carboxylic acid derivatives in both intermolecular and intramolecular fashion (Scheme 3). Additionally, they can participate in (formal) cycloaddition reactions with various unsaturated systems. These new protocols offer milder and more sustainable reaction conditions, enabling practical operations in the laboratory setting.

**2.2.1. Reactions with Nucleophiles.** Early reports in this field involved intramolecular reactions with nitrogen nucleo-


## Scheme 3. Wolff Rearrangement and Its Applications



philes. In 2010, Konopelski and co-workers reported synthesis of  $\beta$ -lactams 4.3 from  $\alpha$ -diazo- $\beta$ -ketoamide 4.1. This transformation proceeds through a sequence of photochemical Wolff rearrangement and intramolecular nucleophilic attack of ketene 4.2 by amine (Scheme 4).<sup>24</sup> White compact fluorescent

Scheme 4. Synthesis of  $\beta$ -Lactams via Wolff Rearrangement

Konopelski 2010




lamps (CFLs) were utilized as the light source, proving to be equally efficient as the commonly employed high-energy UV lamps in these processes. Additionally, flow conditions were implemented to accommodate the reaction at larger scales.

Very recently, Zhou's group achieved an asymmetric synthesis of chiral amides 5.3 using  $\alpha$ -diazo ketones 5.1 and amines 5.2 (Scheme 5).<sup>25</sup> This transformation is catalyzed by a

## Scheme 5. Chiral Phosphoric Acid-Catalyzed Enantioselective Synthesis of Chiral Amides via Wolff Rearrangement

Zhou 2023




chiral phosphoric acid, which not only accelerates the reaction between ketene 5.4 and amines to form an *E*-configured amino enol species A but also enables the enantioselective proton transfer process to yield chiral amides.


Furthermore, it was found that oxygen nucleophiles can participate in reactions with ketenes. In 2015, Burtoloso's group efficiently synthesized allylic esters and carboxylic acids 6.2 using alcohols, phenols, and water as nucleophiles under white LEDs (Scheme 6a).<sup>26</sup> Later, Han's group achieved

## Scheme 6. Esterification via Wolff Rearrangement

a) Burtoloso 2015



b) Han 2019: asymmetric esterification




asymmetric esterification toward  $\alpha,\alpha$ -disubstituted carboxylic esters 6.8 (Scheme 6b).<sup>27</sup> They employed a chiral isothiourea catalyst 6.9 to facilitate the formation of an ammonium enolate intermediate B from ketene A. The enolate then underwent an asymmetric proton-transfer process with phenol, followed by nucleophilic attack by a phenoxide anion (C) to yield the products. A wide range of chiral esters (6.10–6.14) were efficiently prepared with high enantioselectivity.

Carbon nucleophiles have also been employed in reactions with ketene. However, in certain cases, the addition of a suitable catalyst may be necessary. In 2018, Lu's and Xiao's group documented a regioselective acylation method for  $\beta$ -ketoesters 7.1. The selectivity was controlled by the use of either a Lewis acid or base catalyst (Scheme 7).<sup>28</sup> The presence of the Lewis acidic  $\text{NiCl}_2\text{-glyme}$  facilitated the generation of an LA-chelated enolate intermediate 7.6, which subsequently reacted with ketene 7.5 to form the C-acylation product 7.3. In contrast, the use of a Lewis base catalyst resulted in *O*-acylation of the carbonyl group at the  $\beta$ -ketoester. Nucleophilic addition of the Lewis base to ketene produced a zwitterionic intermediate 7.7 capable of abstracting a proton from the  $\beta$ -ketoester. The resulting transient acylated species 7.8 then reacted with enolates, furnishing enol esters 7.4.

In 2021, Basso's group employed isocyanides 8.2 as C-nucleophiles in a three-component reaction with  $\alpha$ -diazo ketones 8.1 (Scheme 8).<sup>29</sup> The reaction is proposed to proceed through the nucleophilic addition of isocyanides to ketenes,


**Scheme 7. Regioselective Acylation of  $\beta$ -Ketoesters with  $\alpha$ -Diazo Ketones**

Lu & Xiao 2018



**Scheme 8. Isocyanides as C-Nucleophiles**

Basso 2021




producing a nitrilium ion **8.6**. The latter was captured by either a carboxylic acid or a silanol. Finally, an intramolecular rearrangement of intermediate **8.7** occurs to afford the corresponding  $\alpha$ -acyloxy- or  $\alpha$ -silyloxyacrylamides **8.4**.

Sulfoxonium ylides have been shown to engage in nucleophilic addition to ketenes, as evidenced by two simultaneous reports. The groups of Oliveira and Burtoloso disclosed an acylation reaction of monocarbonyl sulfoxonium ylides **9.2** and  $\alpha$ -diazo ketones **9.1** (Scheme 9).<sup>30</sup> This method enables the synthesis of dicarbonylated products **9.3** under 400 nm irradiation. Notably, the transformation can be carried out using a continuous flow setup, leading to improved yields and shorter reaction times. Independently, Wang's group reported a similar approach using 450 nm LEDs.<sup>31</sup> The proposed mechanism involves the nucleophilic addition of sulfoxonium ylides **9.2** to ketenes **9.4** and a subsequent proton transfer event, delivering 1,3-dicarbonyl sulfur ylides **9.3**.

**2.2.2. Formal Cycloadditions.** Apart from directly coupling with nucleophiles, ketenes, which are formed through the visible light-induced Wolff rearrangement of  $\alpha$ -diazo ketones, can also participate in formal cycloadditions with

**Scheme 9. Synthesis of Dicarbonyl Sulfoxonium Ylides**

Oliveira & Burtoloso 2022, Wang 2022




different unsaturated  $\pi$ -systems. These cyclization reactions often follow a stepwise mechanism and demonstrate excellent stereocontrol when chiral ligand or organocatalyst is present.

In 2017, Lu and Xiao's groups achieved Pd-catalyzed asymmetric formal [4 + 2] cycloadditions between vinyl benzoxazinanes **10.1** and  $\alpha$ -diazo ketones **10.2**, utilizing a chiral P,S ligand **10.4** (Scheme 10a).<sup>32</sup> Various aryl  $\alpha$ -diazoketones proved to be suitable substrates, providing the corresponding quinolinones **10.5–10.9** in excellent yields with high degrees of enantioselectivity and diastereoselectivity. The reaction mechanism begins with the oxidative addition of Pd<sup>0</sup> to the allyl ester, affording a  $\pi$ -allyl palladium complex **A**. A subsequent decarboxylation of the latter yields tosyl amide anion **B**, which can undergo nucleophilic addition to ketene, producing a zwitterionic intermediate **C**. Finally, an intramolecular asymmetric allylic alkylation generates quinolinones. Following this, the same groups employed vinylethylene carbonates **10.10** in a formal [5 + 2] asymmetric cycloaddition reaction, enabling the efficient synthesis of 7-membered chiral lactones **10.11** via a similar mechanism (Scheme 10b).<sup>33</sup> Furthermore, it was found that vinyl cyclopropane **11.1** acts as an efficient reaction partner for a formal [3 + 2] cycloaddition with ketenes, as reported by Lu's group in 2019 (Scheme 11).<sup>34</sup> This approach allows access to a wide range of highly functionalized tetrahydrofurans **11.5–11.9**. Under similar Pd-catalyzed reaction conditions, vinyl cyclopropanes undergo a ring-opening process in the presence of Pd(0), producing a 1,3-dipolar intermediate **A**. The nucleophilic addition of the latter to ketene and a subsequent intramolecular allylic alkylation event **B** furnished tetrahydrofurans **11.3**.


In addition to Pd catalysis, the asymmetric formal cycloaddition of ketenes can also be achieved through organocatalysis. Chiral organocatalysts efficiently react with photochemically generated ketenes to form chiral enolates **A**, enabling precise control over the stereoselectivity of the subsequent cyclization processes with unsaturated  $\pi$ -systems (Scheme 12a). Thus, in 2019, Song and co-workers reported a chiral isothiourea-catalyzed approach for the formal [4 + 2] cycloaddition of azadienes **12.1** with ketenes (Scheme 12b).<sup>35</sup> This method allows for a highly enantioselective synthesis of benzofuran-fused heterocycles **12.5–12.9** with all-carbon quaternary stereocenters. Similarly, Hui's group employed a chiral N-heterocyclic carbene (NHC) catalyst **12.12** for

**Scheme 10. Pd-Catalyzed Asymmetric Formal [4 + 2] and [5 + 2] Cycloadditions**

a) Lu & Xiao 2017: formal [4+2] cycloaddition



b) Lu & Xiao 2019: formal [5+2] cycloaddition




preparation of tetrahydropyranoindoles 12.11 from enamides 12.10 and  $\alpha$ -diazo ketones 12.2 (Scheme 12c).<sup>36</sup> This visible light-induced protocol yielded the formal [4 + 2] cycloaddition products 12.13–12.17 in high yields and good stereoselectivities.

Imines (13.1) represent a class of substrates suitable for efficient cycloaddition reactions with ketenes. In 2021, Jurberg's group introduced a photochemical method for the formal [2 + 2] cycloaddition to synthesize  $\beta$ -lactams 13.3 (Scheme 13a).<sup>37</sup> They also reported a formal [3 + 2] cycloaddition reaction employing sulfoxonium ylides 13.5 for

**Scheme 11. Pd-Catalyzed Formal [3 + 2] Cycloaddition Employing Vinyl Cyclopropanes**

Lu 2019: formal [3+2] cycloaddition




synthesis of furanones 13.6 (Scheme 13b).<sup>37</sup> The proposed mechanism involves the nucleophilic attack by the sulfoxonium ylide onto the ketene intermediate. Subsequently, the elimination of dimethyl sulfoxide (DMSO) takes place, followed by a Nazarov-type cyclization to give the desired product.

### 2.3. [2 + 1] Cycloadditions


**2.3.1. Cyclopropanation.** Cyclopropanation of C–C double bonds is one of the most well-developed, valuable, and characteristic transformations of diazo compounds.<sup>38,39</sup> The cyclopropane ring itself is widely present in various natural products<sup>40,41</sup> and serves as a versatile building block in organic synthesis.<sup>42,43</sup> Although approaches involving various transition metals are well-established for cyclopropanation reactions, visible light-induced metal-free photolysis of aryl diazoacetates has recently emerged as a more straightforward and milder method for accessing the corresponding cyclopropanes.<sup>44</sup> In recent years, significant progress has been made in the field of metal-free photochemical cyclopropanation reactions. Scheme 14 summarizes the contributions made by various research groups in employing different alkenes or arenes for these transformations. Many of the cyclopropanes were formed with high diastereoselectivity. However, enantioselective cyclopropanation has not been achieved, as typically no catalysts or ligands were employed in these photochemical reactions.

In 2018, the pioneering work by the groups of Jurberg and Davies revealed that aryl diazoacetates absorb in the blue light region (460–490 nm) to undergo photolysis at room

Scheme 12. Asymmetric Formal [4 + 2] Cycloaddition via Organocatalysis



Scheme 13. Photochemical Formal [2 + 2] and [3 + 2] Cycloaddition



temperature toward carbenes.<sup>16</sup> In the presence of styrenes **14.4**, arenes **14.10**, and heteroarenes **14.7**, the formed carbene species efficiently participated in cycloaddition reactions to produce the corresponding cyclopropanes in good yields with high diastereoselectivity (Scheme 14a–c). Following this work, Niu and Song's groups reported cyclopropanation of aryl diazoacetates with *N*-protected indoles and pyrroles (Scheme 14b).<sup>45</sup> However, it was observed that unprotected indoles did not display any reactivity under these conditions. Later, the groups of Reiser and Barham adapted the cyclopropanation reaction to be carried out in continuous flow with various

heteroarenes, further improving the safety, sustainability, and scalability of this transformation.<sup>46</sup>


In 2019, the groups of Nguyen and Koenigs made further advancements by expanding the scope of arenes to include different types of methyl-substituted arenes, such as toluene **14.11**, xylenes, and *p*-cymene (Scheme 14c).<sup>47</sup> These arenes were used in excess as solvents to achieve efficient cyclopropanation with diazo compounds. Furthermore, for polycyclic aromatic compounds, such as naphthalenes **14.12**, anthracene and triphenylene, reactions can be conducted in DCM using 10 equiv of these reagents. Additionally, Chen and co-workers contributed to extending this reaction to novel carbene precursors, 3-diazooxindoles, enabling synthesis of spiro-fused norcaradiene indolinones **14.15**.<sup>48</sup>

In 2020, the groups of Fallon and Koenigs revealed that (poly-)unsaturated carbocycles **14.16** can also serve as suitable reaction partners for cyclopropanation with diazo compounds (Scheme 14d).<sup>49</sup> Despite the presence of multiple double bonds in the substrates, the exclusive formation of monocyclopropanated products **14.17** and **14.18** was observed. This selectivity was achieved by employing large excess of alkenes.

In 2022, Koenigs and Xuan's groups employed NHC boranes **14.19** in cyclopropanation reactions under photochemical conditions (Scheme 14e).<sup>50</sup> This unusual cyclopropanation reaction chemoselectively occurred at the imidazolium backbone of the NHC boranes (**14.20** and **14.21**).


Photochemical intramolecular cyclopropanation of diazo compounds with *N*-protected indoles **15.1** was achieved by Sen and co-workers (Scheme 15).<sup>51</sup> By modifying the substitution pattern at the nitrogen atom of the indole, they were able to access different products. When employing *N*-acetyl and *N*-Boc protected indoles, cyclopropane-fused polycyclic products **15.2** were obtained, whereas *N*-methyl and *N*-H substrates delivered cyclopropyl ring expansion products. Subsequently, the same research group further expanded this photochemical methodology to an intermolecular cyclopropanation/ring expansion cascade.<sup>52</sup>

Scheme 14. Intermolecular Cyclopropanation of Diazo Compounds with Alkenes and Arenes

Scheme 15. Intramolecular Cyclopropanation of Diazo Compounds with *N*-Protected Indoles

Very recently, Koenigs and co-workers introduced a one-pot procedure that combines cyclopropanation of diazo compounds **16.1** with transesterification using allylic alcohols **16.2** (Scheme 16). This method allows for synthesis of various


Scheme 16. One-Pot Cyclopropanation and Transesterification Approach Toward Cyclopropane-Fused Lactones



cyclopropane-fused bicyclic lactones **16.3** with high diastereoselectivity.<sup>53</sup> Notably, this photochemical approach demonstrates excellent chemoselectivity toward cyclopropanation of allylic alcohols, without the formation of O–H insertion or ylide rearrangement products.

**2.3.2. Cyclopropanation.** In 2019, Koenigs and co-workers developed a photochemical catalyst-free approach for cyclopropanation of aryl diazoacetate **17.1** and alkyne **17.2** (Scheme 17).<sup>54</sup> Various phenylacetylene derivatives, as well as internal alkynes, efficiently participated in this transformation. The protocol was further examined in a continuous-flow setup, which increased productivity by 36-fold compared to the

Scheme 17. Cyclopropanation of Diazo Compounds with Alkynes




conventional batch cyclopropanation reactions. Following this success, the same research group extended this photochemical cyclopropanation method to propargylic alcohols.<sup>55</sup> Important differences in reactivity were observed between photochemical and metal-catalyzed reaction of carbenes. In the metal-free transformation, a highly chemoselective cyclopropanation reaction (17.8–17.11) of the tertiary alcohol occurred. This is in a contrast with the reactions involving transition metals, where rearrangements of an ylide intermediate are more common due to the increased reactivity of more electrophilic metal carbene complexes toward nucleophilic alcohols.<sup>56,57</sup> However, when switching to primary alcohols, exclusive formation of the O–H insertion products, propargylic ethers, were observed, without the generation of cyclopropene or rearrangement products.

In 2021, they further investigated this photochemical cyclopropanation reaction by employing diaryl diazo compounds **18.1** with alkynes **18.2** (Scheme 18).<sup>58</sup> Upon

**Scheme 18. Cyclopropanation of Diaryl Diazo Compounds with Alkynes**

Koenigs 2021



photolysis, the formed singlet diaryl carbenes are stabilized by the introduction of electron-withdrawing nitro group at the aryl rings. Consequently, these carbenes exhibit similar characteristics to the ones substituted with an ester group. Thus, these can engage in reactions that are common for singlet carbenes, such as cyclopropanation of alkynes. Unlike the reactions involving aryl diazoacetates, the photolysis of diaryl diazo compounds, in order to achieve high yields, requires an anaerobic and anhydrous atmosphere.


**2.4. C–H Insertions**

Given the ubiquitous presence of C–H bonds in organic molecules, C–H insertion reactions of diazo compounds represent a powerful strategy for forging new carbon–carbon bonds.<sup>2b</sup> Over the last few decades, this field has undergone substantial development, notably in the realm of enantioselective intermolecular reactions catalyzed by transition metals.<sup>2a</sup> Recently, visible light offered new opportunities for C–H functionalizations under milder and metal-free conditions.

Pioneering investigations by Jurberg and Davies demonstrated the use of cyclic alkanes **19.2** and electron-rich arenes **19.6** in photochemical intermolecular C–H insertion reactions with diazo compounds (Scheme 19).<sup>16</sup> Various electron-rich

**Scheme 19. Intermolecular C–H Insertions of Diazo Compounds**

Jurberg & Davies 2018




arenes, such as *N*-methylindole (**19.3**), *N*-methyl pyrrole (**19.4**), or 1,3,5-trimethoxybenzene (**19.5**), efficiently participated in this transformation. However, reactions of cyclic alkanes need to be conducted in neat or with a large excess of the substrate to compensate their lower reactivity. In the same report, the authors also found that this reaction can be performed intramolecularly with C(sp<sup>3</sup>)–H to afford 5-membered lactone **20.2** (Scheme 20a). Later, Sun and co-workers extended this intramolecular strategy to C(sp<sup>2</sup>)–H bonds, which allowed for preparation of fluorene derivatives **20.4** (Scheme 20b).<sup>59</sup> Very recently, Hari's group demon-

**Scheme 20. Intramolecular C–H Insertions of Diazo Compounds**

a) Jurberg & Davies 2018



b) Sun 2023




c) Hari 2023



strated the efficient synthesis of various spiro- $\beta$ -lactones and lactams **20.6** using this photochemical approach (Scheme 20c).<sup>60</sup>

In 2021, Koenigs and co-workers employed diaryl diazo compounds **21.1** in photochemical intermolecular C–H insertion reactions with alkynes **21.2** (Scheme 21).<sup>58</sup> In a

**Scheme 21. Intermolecular C–H Insertions of Diaryl Diazo Compounds with Terminal Alkynes**




combined experimental and computational study, the authors proposed that incorporating two electron-donating methoxy groups onto aryl rings leads to the nearly degenerated singlet–triplet carbene intermediates. Therefore, the reaction outcome strongly depends on substrates. For instance, terminal aromatic alkynes or TMS-acetylene reacted exclusively at the C(sp)–H site (**21.3**) via a low-lying transition state involving a singlet carbene, whereas aliphatic alkynes produced C–H insertion products **21.4** at the propargylic C(sp<sup>3</sup>)–H site via the triplet carbene intermediates.

Diazo compounds containing an *N*-(acyloxy)phthalimide (NHPI) ester moiety are attractive reagents in organic synthesis. They have previously been employed by Mendoza's group in Rh-catalyzed reactions with alkenes for the enantioselective synthesis of highly substituted cyclopropanes.<sup>61,62</sup> On the other hand, because redox-active esters can produce C-centered radicals through the photoexcitation of electron donor–acceptor (EDA) complexes,<sup>63</sup> such diazo compounds hold the potential for orthogonal functionalizations. Very recently, Gryko and co-workers investigated the

photochemical behavior of these two functionalities coexisting within a single molecule, a diazo compound containing a redox-active ester group 22.1.<sup>64</sup> UV/vis spectra analysis of the diazo compounds with the NHPI moiety indicates that they absorb in the visible light region, albeit with a hypsochromic shift of their  $\lambda_{\text{max}}$ . Nevertheless, further investigation demonstrated that the reactions exclusively occur at the diazo group, leaving the NHPI ester functionality intact. For example, it was found that under photochemical conditions, the diazo compounds undergo efficient C–H insertions with both linear and cyclic alkanes (22.4–22.6) (Scheme 22).

**Scheme 22. C–H Insertions of Aryl Redox-Active Diazo Compounds**



## 2.5. X–H Insertions


X–H (X = B, N, O, Si, S) insertion by carbenes is a potent strategy for constructing diverse C–heteroatom bonds. This field is primarily dominated by transition metal-catalyzed reactions,<sup>65,66</sup> with only a few instances of metal-free transformations at elevated temperatures.<sup>13,15</sup> Recent advancements in the direct photolysis of diazo compounds demonstrate that visible light-induced approaches offer a valuable alternative and, in some cases, are highly complementary to the traditional transition metal-catalyzed pathways. While the product resulting from X–H insertion can be formally considered as the outcome of a carbene inserting into the X–H bond, the actual mechanistic processes involved may vary widely. Under photolytic conditions, free carbenes are generated from diazo compounds, which can either undergo nucleophilic attack by various heteroatoms, followed by protonation, or proceed via a concerted process.<sup>67,68</sup>

**2.5.1. B–H Insertions.** In the previously mentioned study on the cyclopropanation of NHC-boranes using aryl diazoacetates conducted by Koenigs and Xuan,<sup>50</sup> it was observed that when substitutions blocked the cyclopropanation site on imidazole, B–H insertion products 23.4 were formed (Scheme 23). Other Lewis base borane complexes, such as pyridine borane (23.5) and phosphine borane (23.6), also efficiently participate in this reaction. Furthermore, cyclopropanation and B–H insertion can occur simultaneously on NHC-boranes when both reaction sites are available.

**2.5.2. N–H Insertions.** In the previously referenced report by Jurberg and Davies, they demonstrated N–H insertion of aryl diazoacetates into amines and anilines 24.4 under blue light irradiation (Scheme 24a).<sup>16</sup> Various N-alkylated products 24.5 and 24.6 were obtained in great efficiency. Following this report, several research groups have further advanced and expanded upon this photochemical strategy for constructing C–N bonds. The scope of this transformation is summarized in Scheme 24.

**Scheme 23. B–H Insertions of Diazo Compounds with NHC Boranes**

Koenigs & Xuan 2022




Aza-arenes are capable reaction partners in photochemical N–H insertion protocols with aryl diazoacetates. In 2019, the groups of Patureau and Koenigs employed carbazoles 24.7 to construct N-alkylated products (Scheme 24b).<sup>69</sup> Later, Jurberg and co-workers also reported the N–H insertion of aryl diazoacetates into carbazoles, as well as pyrazoles 24.10 and triazoles 24.13 (Scheme 24b,c).<sup>70</sup> However, nonsymmetrical pyrazoles and 1,2,3-triazoles produced different N-alkylated regioisomers 24.14. In 2021, Sen's group further expanded the scope of N-heterocycles to include indoles 24.28, under both batch and flow conditions (Scheme 24e).<sup>71</sup> Notably, other competing reactions of carbenes, such as cyclopropanation and C–H insertion at the C3 position, were not observed in this case.

Apart from N-heterocycles, Jurberg and co-worker demonstrated that amides 24.16, isatins 24.19, sulfonamides 24.22, and imides 24.25 can effectively engage in N–H insertion reactions with aryl diazoacetates (Scheme 24d).<sup>72</sup> It was observed that substrates with less acidic N–H bonds, such as amides 24.16 and isatins 24.19, required the addition of a catalytic amount of DBU to improve yields.

As summarized above, while racemic photochemical N–H insertion of free carbenes has been extensively studied, there are limited examples of its enantioselective counterparts. A recent development in this area involves the use of a chiral phosphoric acid catalyst, as reported by the groups of Guo and Sun. (Scheme 25).<sup>73</sup> They accomplished asymmetric N–H insertion reactions of aryl diazoacetates 25.1 for the synthesis of enantioenriched  $\alpha$ -amino esters 25.3. Various amines were found to be efficient reaction partners (25.5–25.8). However, it was observed that aliphatic amines lead to racemic products. This was attributed to the increased basicity of aliphatic amines, which may deactivate the catalyst. The proposed mechanism begins with the rapid capture of free carbene A by DMSO to produce a sulfoxonium ylide B. A subsequent protonation of the latter by a chiral acid catalyst affords sulfoxonium ions C and D. These chiral anions then guide the amine to preferentially react with one of the two diastereomers, delivering the enantioenriched product 25.3 through dynamic kinetic resolution.

Scheme 24. N–H Insertions of Diazo Compounds



Thereafter, the groups of Li and Zhou designed spiro phosphamide catalysts **26.4** to achieve enantioselective N–H insertion reactions (Scheme 26).<sup>74</sup> Differently from the previous case, photochemically generated free carbene species were directly captured by amine substrates **26.2** through a nucleophilic attack. The resulting ammonium ylide intermediate then engaged in an enantioselective proton transfer process with the chiral phosphamide catalyst, affording the enantioenriched product **26.3**. Various aromatic amines proved to be suitable reaction partners (**26.5–26.7**). Moreover, challenging substrates, such as benzylamine (**26.8**) and ammonia (**26.9**), also participated in the reaction, albeit with lower enantioselectivities. Soon after, Guo's group reported a similar N–H insertion approach for the asymmetric synthesis of  $\alpha$ -amino esters, enabled by an enantioselective proton transfer process using a chiral phosphoric acid catalyst.<sup>75</sup>


**2.5.3. O–H Insertions.** In 2018, Jurberg and Davies reported that carboxylic acids **27.4** and alcohols **27.7** efficiently undergo O–H insertions with aryl diazoacetates under blue light irradiation (Scheme 27a,b).<sup>16</sup> Additionally, control experiments revealed that reactions involving stronger carboxylic acids can proceed in dark, which presumably involved protonation of the diazo compound followed by  $S_N2$  displacement of  $N_2$ . Later, Rostovskii and co-workers employed carboxylic acids containing an azirine moiety to synthesize the corresponding esters **27.6**.<sup>76</sup> They observed that photochemically generated free carbene chemoselectively reacted at the carboxyl group rather than at the azirine ring. In contrast, the metal carbene underwent nucleophilic attack

by the azirine nitrogen in the presence of a gold catalyst. In 2022, the groups of Yang and Xia further extended the scope of alcohols (Scheme 27b).<sup>77</sup> In the same report, the authors also utilized water for O–H insertions targeting  $\alpha$ -hydroxy esters **27.11** and **27.12** (Scheme 27c). The use of oximes **27.13** in photochemical O–H insertion reactions with aryl diazoacetates was independently developed by the groups of Xuan,<sup>78</sup> Wang,<sup>79</sup> and Jurberg<sup>80</sup> (Scheme 27d). Various oxime esters were obtained in good yields from both ketoximes **27.14** and aldoximes **27.15**. It was also shown that donor–donor type diazo compound, such as diphenyl diazomethane, can be utilized in this reaction.<sup>78</sup> Furthermore, the groups of Zhang, Li, and Xia demonstrated the O–H insertion of hydroxylamides **27.16** formed *in situ* through nitroso aldol reaction using  $\beta$ -keto esters and nitrosopyridines (Scheme 27e).<sup>81</sup> This approach provided an efficient pathway toward trisubstituted hydroxylamines **27.17** and **27.18**. Moreover, Yang, Zhou, Huo, and co-workers demonstrated that 2-hydroxypyridines **27.19**, formed *in situ* from 2-pyridones, also participated in O–H insertion reactions (Scheme 27f).<sup>82</sup> The proposed mechanism suggests the initial formation of free carbene from diazo compounds, which then engage in radical pathways.

**2.5.4. Si–H Insertions.** Si–H insertions are convenient and effective methods for constructing C–Si bonds. In 2020, Koenigs and co-workers reported a photochemical insertion reaction of silanes **28.2** with aryl diazoacetates **28.1** (Scheme 28).<sup>83</sup> This approach enabled synthesis of a wide range of silanes with moderate to good yields (**28.4–28.7**), including sterically hindered substrates such as triphenyl silane (**28.6**).


**Scheme 25. Enantioselective N–H Insertion Catalyzed by Chiral Phosphoric Acid**

Guo & Sun 2023




**Scheme 26. Enantioselective N–H Insertion Catalyzed by Chiral Phosphamide**

Li & Zhou 2023



**2.5.5. S–H Insertions.** In 2020, Qiu and Guo's groups reported a photochemical S–H insertion reaction of aryl diazoacetates **29.1** with cysteine residues **29.4** (Scheme 29a). This reaction was also found to be applicable to common thiols (29.8) and thiophenol (29.9), resulting in various S-alkylated products. Additionally, during the same year, the groups of Yang, Ma, and Huo also disclosed a S–H insertion reaction of thiols **29.10** and **29.13** (Scheme 29b). Both

**Scheme 27. O–H Insertions of Diazo Compounds**



donor–acceptor and acceptor-type diazo compounds were found to be suitable reagents. However, it was observed that ethyl diazoacetate failed reacted with alkyl thiols. The proposed mechanism suggests the formation of thiyl radicals from thiols, followed by their subsequent addition to carbenes.


**2.6. Ylides Formation**

Free singlet carbenes derived from  $\alpha$ -diazocarbonyl compounds are highly reactive electrophiles and thus can readily react with various heteroatoms possessing lone pairs such as

## Scheme 28. Si–H Insertions of Diazo Compounds




## Scheme 29. S–H Insertions of Diazo Compounds



ethers, sulfides, and amines (Scheme 30).<sup>86</sup> These reactions lead to the formation of “free ylides,” which are versatile

## Scheme 30. Formation of Ylides




intermediates capable of undergoing various reactions, including [2,3]-sigmatropic rearrangements of allyl-substituted species, [1,2]-rearrangements, and cycloadditions.


**2.6.1. Nitrogen Ylides.** In 2019, Koenigs and co-workers reported a photochemical [2,3]-sigmatropic rearrangement reaction of ammonium ylides A (Scheme 31).<sup>87</sup> Different allylic amines 31.2 participate in reactions with aryl diazoacetates 31.1, delivering various homoallylic products 31.4–31.7.

Subsequently, Xu and Hu’s groups developed a photochemical annulation reaction of ammonium ylides A for synthesis of polysubstituted indolines 32.3 from aryl

## Scheme 31. [2,3]-Sigmatropic Rearrangement of Ammonium Ylides




diazoacetates 32.1 and *o*-aminoacetophenones 32.2 (Scheme 32).<sup>88</sup> Various aryl diazoacetates smoothly reacted in this

Scheme 32. Formal [4 + 1]-Annulation of Diazoacetates with *o*-Aminoacetophenones

transformation, affording the corresponding products 32.4–32.6 with high diastereoselectivity. DFT calculations suggested the formation of (E)-enol intermediates B through the H-shift of ylide A. The high diastereoselectivity observed in the subsequent intramolecular Aldol-type addition is facilitated by H-bonding interactions between the enol species and the tethered carbonyl group.

In 2021, Wang and Xuan’s groups reported a photochemical protocol for the chemodivergent synthesis of aziridine 33.3 and imidazolidine 33.4 frameworks using diazo compounds 33.1 and 1,3,5-triazines 33.2 (Scheme 33).<sup>89</sup> Interestingly,

## Scheme 33. Divergent Synthesis of Aziridines and Imidazolidines




they were able to access different products by altering the solvents used. Mechanistic studies and DFT calculations indicated the involvement of key ammonium ylide intermediates, which formed through the trapping of free carbenes by 1,3,5-triazines.

In the same year, Xuan and Xiao’s groups described a highly efficient photochemical method for constructing amides 34.3

from aryl diazoacetates 34.1 and nitrosoarenes 34.2 (Scheme 34).<sup>90</sup> Mechanistically, free carbenes were trapped by the

**Scheme 34. Synthesis of Amide from Diazo Compounds and Nitrosoarenes**

Xuan & Xiao 2021




nucleophilic *N*-center of nitrosoarenes, producing ylides **A** that predominantly exist in their resonance form as nitrones (**B**). Under blue light irradiation, the latter gets converted to an oxaziridine **C**, which subsequently is rearranged into an amide product.


Unlike the generation of ammonium ylides using donor-acceptor aryl diazoacetates, nitrile ylides were formed using acceptor-only diazo compounds. In 2022, Zhao and Li reported a three-component photochemical approach for constructing diacylglycine esters 35.6 and *N*-imidoylbenzotriazoles 35.5 from  $\alpha$ -diazo esters 35.1 and nitriles 35.2.<sup>91</sup> Carboxylic acids 35.4 and benzotriazoles 35.3 were used as nucleophiles, respectively (Scheme 35A). The reaction

**Scheme 35. Formation of Nitrile and Pyridinium Ylides**

*A. Formation of nitrile ylide*



*B. Formation of pyridinium ylide*




mechanism involves the capture of free carbenes by nitriles to generate nitrile ylides **A**, followed by protonation to afford a nitrilium ion. A nucleophilic attack of the latter by carboxylate, followed by the Mumm rearrangement, delivered imides 35.6. Alternatively, the nitrilium ion can undergo nucleophilic attack by benzotriazole to produce various *N*-imidoylbenzotriazoles 35.5. Independently, Xuan's group later reported a similar approach using carboxylic acids.<sup>92</sup> Pyridinium ylides can be formed through reactions of free carbenes and pyridines. Thus, in 2021, Beeler and co-workers demonstrated the dearomatized ring expansion of pyridinium ylides to produce azepines.<sup>93</sup> However, this reaction has a low yield of 25% due to

competitive carbene side reactions. Subsequently, Sen's group utilized pyridinium ylides **B** in a [3 + 2] cycloaddition with maleimides 35.9 (Scheme 35B).<sup>94</sup> The targeted itaconimide derivatives 35.10 were obtained through the sequential oxidation and ring-opening of the adducts, followed by the addition of another equivalent of maleimide.

**2.6.2. Oxygen Ylides.** In 2019, the groups of Xu and Koenigs reported photochemical ring expansion of oxonium ylides **A**, allowing for synthesis of various tetrahydrofurans 36.3 from oxetanes 36.2 and aryl diazoacetates 36.1 under mild and operationally simple reaction conditions (Scheme 36A).<sup>95</sup> DFT calculations suggested a diradical pathway (**B**)

**Scheme 36. Ring Expansion and Contraction of Oxonium Ylides**




for the ring expansion of oxonium ylides. Later, Sen's group reported similar ring expansion of dioxanes 36.7 to afford a wide range of dioxepanes 36.8 and 36.9.<sup>96</sup> However, unlike the previous report, the proposed mechanism involved a [1,2]-rearrangement of oxonium ylides **C**.

Apart from the ring expansion reactions, oxonium ylides can also participate in ring contraction processes. In 2022, the research groups of Guo, Song, and Xia reported a photochemical approach for synthesis of vinyl oxetanes 36.12 from 2,5-dihydrofurans 36.11 and diazo compounds 36.10 (Scheme 36B).<sup>97</sup> The DFT calculations indicated a mechanistic sequence similar to that proposed in the work of Xu and Koenigs, suggesting the formation of an oxonium ylide intermediate **D**, followed by a diradical rearrangement pathway.

In recent years, significant progress has been made in the development of photochemical cascade ring-opening reactions

## Scheme 37. Photochemical Cascade Ring-Opening Reactions of Oxonium Ylides



of oxonium ylides. The mechanism of these multicomponent transformations begins with the nucleophilic addition of cyclic ethereal solvents **37.2**, such as THF or dioxane, to free carbenes producing the key oxonium ylides. A subsequent ring-opening of the ylides in the presence of various nucleophiles **37.3** furnishes products **37.4** with a short polyether side chain. Scheme 37 summarizes the contributions from different research groups regarding the employment of a diverse range of nucleophiles.

The initial discovery was made by Koenig's group in 2020. They reported a photochemical fluoro-amino etherification reaction of aryl diazoacetates **37.1** with *N*-fluorobenzolsulfonamide (NFSI) **37.5** in 1,4-dioxane (Scheme 37a).<sup>98</sup> In this reaction, the formed oxonium ylide undergoes electrophilic

fluorination with NFSI, followed by ring-opening reaction of resulting oxonium ion to afford the final products **37.6** with newly constructed C–O, C–F, and C–N bonds.


Other groups have adopted this photochemical platform to install polyether side chains with diverse functional groups. In the same year, Jiang and co-workers reported synthesis of carbamates **37.8** by employing carbamate ions formed *in situ* from CO<sub>2</sub> and amines as nucleophiles. (Scheme 37b).<sup>99</sup> Later, the groups of Wei, Zhao, and He developed an analogous reaction using CS<sub>2</sub> to prepare dithiocarbamates **37.16** (Scheme 37e).<sup>100</sup> A further advance was achieved by Xuan's group, using oximes **37.9** to produce various oxime ethers **37.10** (Scheme 37c).<sup>78</sup> In this process, the ylide intermediate was protonated by oximes, followed by a ring-opening reaction to

yield the desired products. Independently, Jurberg and co-workers reported a similar approach using oximes.<sup>80</sup> Sen and co-workers further explored this cascade ring opening approach of oxonium ylides, incorporating various *N*-heterocycles such as phthalimides 37.11 and indoles 37.13 (Scheme 37d).<sup>96</sup> In 2022, Jurberg systematically tested a variety of nucleophiles, expanding the scope to include carboxylic acids, water, alcohols, HCl, sulfonamides 37.25, and isatins 37.27 (Scheme 37f).<sup>101</sup> During the same year, the groups of Yue and Wei disclosed a four-component photochemical reactions of aryl diazoesters, cyclic ethers, S<sub>8</sub>, and *H*-phosphonates 37.29 (Scheme 37g).<sup>102</sup> In this reaction, a nucleophilic phosphorothioic anion was formed *in situ*, which subsequently participated in the ring-opening process of oxonium ylides, affording a broad range of *S*-alkyl phosphorothioates 37.30. Later, Wei's group reported an analogous approach employing O<sub>2</sub> from the air for the synthesis of various phosphonates 37.32 and phosphinates (Scheme 37h).<sup>103</sup> Moreover, the same group demonstrated the utilization of TMSN<sub>3</sub>, TMSCN, pyrazolones, and dicarbonyl compounds as nucleophiles in three-component cascade reactions involving diazo compounds and cyclic ethers (Scheme 37i,j).<sup>104,105</sup> More recently, An and Wang's groups employed the *in situ* formed O-substituted hydroxamic acids 37.38 toward various *N*-alkylated acrylamides 37.39 (Scheme 37k).<sup>106</sup>

The photochemical cascade ring-opening reaction of oxonium ylides can be synergistically combined with NHC catalysis, thus further expanding the range of suitable substrates for this transformation. In 2022, Xuan and Koenigs introduced a four-component approach for efficient synthesis of a wide range of *O*-protected hydroxylamines 38.4 using aryl diazoacetates 38.1, aldehydes 38.2, and nitroso arenes 38.3 (Scheme 38).<sup>107</sup> The NHC catalyst was used to promote the *in*

### Scheme 38. NHC-Catalyzed Approach for Synthesis of *O*-Protected Hydroxylamines

Xuan & Koenigs 2022




*in situ* formation of hydroxamic acids **A** from aldehydes and nitroso arenes, which then acted as nucleophiles in ring-opening reactions of oxonium ylides.

The vast majority of published studies of cascade ring-opening reactions of oxonium ylides employ donor–acceptor diazo compounds, particularly aryl diazoacetates. Much less attention has been given to other diazo substrates, such as acceptor diazo compounds, mainly due to their limited absorption in visible light region. However, a recent report by Zhao and Li demonstrate the successful employment of acceptor-type diazo compounds 39.1 in cascade ring-opening reactions of oxonium ylides under blue light (Scheme 39).<sup>108</sup> A variety of *N*-, *O*-, and *S*-based nucleophiles (39.5–39.8) efficiently participated in this transformation. Competition

### Scheme 39. Cascade Ring-Opening Reactions of Oxonium Ylides Using Acceptor-Type Diazo Compounds

Zhao & Li 2023




experiments indicated that, although *α*-diazoesters are viable substrates, aryl diazoacetates still more readily generate free carbenes, owing to their relatively strong absorption in the visible light region.

In addition to cyclic ethers, Chatterjee and co-workers have demonstrated that nitroso arenes 40.2 can also react with free carbenes to form oxygen ylides **A** (Scheme 40).<sup>109</sup> They developed a photochemical oxygenation process for the preparation of a series of ketones 40.3.

### Scheme 40. Synthesis of Ketones from Nitroso Arenes


Chatterjee 2021



Very recently, the groups of Cheng, Xu, and Li reported a photochemical oxy-alkynylation reaction of diazo compounds 41.1 with ethynylbenziodoxolones 41.2 (Scheme 41).<sup>110</sup> This

### Scheme 41. Synthesis of Propargylic Esters via Oxonium Ylide Intermediates

Cheng, Xu & Li 2023




method efficiently delivers a wide range of propargylic esters 41.3 under metal-free conditions. In comparison to the previously developed Cu-catalyzed version of this transformation by Waser and co-workers,<sup>111</sup> this reaction proceeds via an oxonium ylide **B**. Subsequently, a 1,2-rearrangement occurs, followed by a reductive elimination process of intermediate **C**, yielding a propargylic ester. Both acceptor– and donor–acceptor-type diazo compounds are suitable substrates in this reaction.

**2.6.3. Sulfur Ylides.** The [2,3]-sigmatropic rearrangement of sulfonium ylides in the presence of a transition metal, known

as the Doyle–Kirmse reaction, was first reported by Kirmse in 1968<sup>112</sup> and later modified by Doyle in 1981.<sup>113</sup> However, the metal-free, photochemical process was only recently reported by Koenigs and co-workers in 2019 (Scheme 42a).<sup>54</sup> Under

**Scheme 42. [2,3]-Sigmatropic Rearrangement of Sulfonium Ylides**




blue light irradiation, various allylic sulfides 42.5 smoothly reacted with aryl diazo acetates 42.1 in a Doyle–Kirmse reaction. This method provides access to a wide range of homoallylic sulfides 42.6–42.10, including both aliphatic and heteroaromatic thioethers, in high yields. Later, the groups of Jiang and Xiao employed *gem*-difluoroallyl sulfides 42.11 in Doyle–Kirmse reaction for synthesis of corresponding homoallylic sulfides 42.12–42.16 (Scheme 42b).<sup>114</sup> The obtained products possess a vinyl bromide moiety that can be further functionalized. The scope of the photochemical [2,3]-sigmatropic rearrangement reaction extends beyond allyl compounds. In 2020, Gryko's group demonstrated the use of propargyl sulfides 42.17 for synthesis of allenes 42.18–42.22 (Scheme 42c).<sup>115</sup> While these studies mainly focus on employing aryl diazoacetates, in 2021, Lu and Wang expanded the scope of diazo compounds to 4-diazo-1,4-dihydroisoquinolin-3-ones, which participate in reactions with both allyl and propargyl sulfides.<sup>116</sup>

The Sommelet–Hauser rearrangement was observed in sulfonium ylides obtained from the reaction of  $\alpha$ -mercapto esters and free carbenes. In 2019, Koenigs and co-workers

revealed the photochemical synthesis of various alkylated arenes 43.3 from aryl diazoacetates 43.1 and  $\alpha$ -mercapto esters 43.2 (Scheme 43A).<sup>117</sup> Thus, the initially formed sulfur ylide A

**Scheme 43. Sommelet–Hauser and [1,2]-Rearrangement of Sulfonium Ylides**




tautomerizes to generate the more reactive ylide B, which subsequently engages in consecutive [2,3]-sigmatropic dearomatization and [1,3]-shift rearomatization to yield the *ortho*-alkylated arenes. In the same report, the authors also explored the reactivity of aryl diazoacetates in the presence of sulfenyl phthalimides 43.7 and benzylic sulfides 43.10 (Scheme 43B).<sup>117</sup> Thus, it was found that the resulting sulfonium ylides underwent [1,2]-rearrangement. The scope mainly involved aryl diazoesters and sulfides with different substitution patterns at the aromatic rings.

Sulfonium ylides can participate in ring expansion reactions. Thus, in 2019, the groups of Xu and Koenigs disclosed a ring expansion reaction of thietanes 44.2 with donor–acceptor diazo compounds 44.1 toward thiolanes 44.3 (Scheme 44).<sup>95</sup> The DFT calculations revealed that, similarly to the aforementioned mechanism of oxygen ylides, the subsequent ring expansion of sulfonium ylides likely proceeds through a diradical pathway.

Very recently, Muthusamy's group introduced a photochemical protocol for synthesis of arylidene oxindoles 45.3 from diazoindolones 45.1 and thioketones 45.2 (Scheme 45).<sup>118</sup> The proposed mechanism suggests the formation of sulfonium ylide A, followed by an electrocyclization reaction to produce episulfide B. Subsequently, elemental sulfur is eliminated, furnishing various tetrasubstituted alkenes 45.3.


**Scheme 44. Ring Expansion of Sulfonium Ylides**

XII & Koenigs 2019



**Scheme 45. Synthesis of 3-Arylidene Oxindoles**


Muthusamy 2023



In an effort to achieve high levels of enantiocontrol for formal [4 + 1] cycloadditions of sulfonium ylides **A** and ethynylbenzoxazinones **46.2**, Lu and co-workers examined a copper-catalyzed approach employing chiral bisoxazoline ligand **L\*** (Scheme 46).<sup>119</sup> Sulfur ylides were generated from

**Scheme 46. Copper-Catalyzed Asymmetric Formal [4 + 1] Cycloadditions**

Lu 2023




diazocompounds and dimethyl sulfide under blue light irradiation. This method allowed for preparation of a wide range of chiral indolines **46.3**, bearing C2-quaternary stereocenters with good degrees of enantio- and diastereoselectivity.

## 2.7. Miscellaneous Reactions

Shortly after the work by Jurberg and Davies,<sup>16</sup> in 2018, another seminal publication on photochemical reactions of diazo compounds was reported by He and Zhou (Scheme 47).<sup>120</sup> In this study, a mixture of donor–acceptor 47.1 and acceptor diazoalkane 47.2 was subjected to a reaction under blue light irradiation. Only the latter absorbed a sufficient amount of visible light for photolysis to generate singlet carbenes. The acceptor-type diazoalkanes, however, remained untouched during this photochemical process. As a result, it can subsequently attack the electrophilic singlet carbene to afford trisubstituted alkenes 47.3 as the final products. Various

**Scheme 47. Selective Visible Light Photolysis of Donor–Acceptor Diazoalkanes in the Presence of Acceptor-Type Diazo Compound**

He & Zhou 2018




$\alpha$ -diazo ketones, amides, and esters successfully participated in the reaction, producing a series of  $\alpha,\beta$ -unsaturated *E*-alkenes (47.4–47.13). This report highlighted the difference in reactivity between the two types of diazo compounds when subjected to visible light irradiation.

In 2020, Singh and co-workers disclosed a photochemical heterocyclization reaction between  $\alpha$ -diazo 1,3-diketones **48.1** and  $\beta$ -ketothiamides **48.2** toward highly functionalized thiadiazoles **48.3** (Scheme 48).<sup>121</sup> The reaction mechanism involves a

**Scheme 48. Synthesis of Thiazolines from  $\alpha$ -Diazo 1,3-Diketones and  $\beta$ -Ketothiamides**


*Sinah 2020*



nucleophilic attack by the sulfur atom of the  $\beta$ -ketothiamides at carbene to afford intermediate A. An intramolecular cyclization of the latter led to intermediate B, which upon elimination of water produced thiazolines. It should be noted that Wolff rearrangements of  $\alpha$ -diazo 1,3-diketones were not observed in this case.

In the same year, Jurberg's group reported a photochemical arylation/alkylation protocol using aryl diazoacetates **49.1** and boronic acids **49.2** (Scheme 49). This method enables preparation of various  $\alpha$ -branched esters **49.3** in moderate to good yields.<sup>122</sup> The proposed mechanism suggests that free singlet carbenes **A** can react with boronic acids to form ylide intermediates **B**. The ylides then undergo a [1,2]-sigmatropic shift, which upon protodeboronation leads to the formation of


**Scheme 49. Arylation and Alkylation of Diazo Compounds with Boronic Acids**



C–C coupled products **49.3**. Furthermore, it was also found that the reaction could proceed in the dark under thermal conditions, albeit with much more limited scope. The reaction presumably proceeded through the coordination of the aryl diazoacetate with boronic acid in the absence of light, which can stabilize the resulting carbene intermediate through hydrogen bonding, thus following the elimination of  $N_2$ .

In 2021, the groups of Yang and Ma demonstrated a photocatalytic strategy for the site-selective  $N^1$ -alkylation of benzotriazoles **50.2** with  $\alpha$ -diazoacetates **50.1** (Scheme 50).<sup>123</sup> The  $N$ -alkylation of benzotriazoles usually results in a mixture


**Scheme 50. Site-Selective  $N^1$ -Alkylation of *N*-heterocycles with  $\alpha$ -Diazoacetates**



of  $N^1$ - and  $N^2$ -substituted products, given that benzotriazole exists in equilibrium between its  $1H$  and  $2H$  tautomers.<sup>124,125</sup> In this work, the authors addressed this synthetic problem by generating a  $N^1$ -centered radical from benzotriazole. The reaction mechanism commences with the photoexcitation of 1,4-benzoquinone (PBQ) catalyst, which produces radical intermediate A from benzotriazole via either HAT or PCET. A subsequent capture of radical species A by a carbene leads to the formation of carbon-centered radical B, which upon a HAT event with intermediate C affords  $N^1$ -alkylated product **50.3**. Various aryl diazo esters worked well in this reaction (**50.4**–**50.8**). It was also found that other *N*-heterocycles, such as indazole (**50.7**) and pyrazole (**50.8**), are also capable substrates. This work demonstrated complementary selectivity compared to that by previously reported Rh-catalyzed  $N^2$ -selective N–H insertion of diazo compounds into benzotriazoles<sup>126</sup> and visible light-induced methods involving 1,2,3-triazoles.<sup>70</sup>

In the same year, the groups of Li, Xu, and Xuan reported a photochemical protocol for preparation of sulfoxonium ylides **51.3** using diazo compounds **51.1** in combination with sulfoxides **51.2** (Scheme 51).<sup>127</sup> Shortly after, similar


**Scheme 51. Synthesis of Imines and Sulfoxonium Ylides Using Diazo Compounds**



approaches toward sulfoxonium ylides were independently disclosed by Bhat<sup>128</sup> and Jurberg.<sup>37</sup> Furthermore, in Jurberg's report, they also explored the reactivity of aryl diazoacetates in the presence of azides **51.4**, which, upon light irradiation, produced a diverse array of imines **51.5**.


A year later, Jurberg's group developed a formal C–C bond insertion protocol of aryl diazoacetates **52.2** with 1,3-diketones **52.1** (Scheme 52).<sup>129</sup> This method allowed for efficient

**Scheme 52. Racemic Approach for Formal C–C Bond Insertion**



synthesis of various 1,4-dicarbonyl compounds **52.3**. The proposed mechanism begins with the formation of a free carbene **A** under blue light irradiation, which then undergoes cyclopropanation with the enol form **B** of 1,3-diketone. The resulting cyclopropanol intermediates then rearrange into formal C–C bond insertion products **52.3**. Later, a similar racemic approach using enaminones was demonstrated by the groups of Lin and Jin.<sup>130</sup> Furthermore, an analogous asymmetric version of this formal C–C bond insertion reaction of aryl diazoacetates was achieved by Hui's group (Scheme 53).<sup>131</sup> A wide range of enantioenriched 1,4-

**Scheme 53. Asymmetric Formal C–C Bond Insertion Using Chiral Phosphoric Acid Catalysts**




dicarbonyl compounds **53.5–53.14** bearing a quaternary carbon center were efficiently prepared. Asymmetric induction was accomplished by employing a chiral phosphoric acid catalyst **53.4**. Dual hydrogen-bonding interactions between the catalyst and the enol tautomer of 1,3-diketone ensured the

stereoselectivity of the following cyclopropanation step. In 2023, Lui and co-workers employed chiral guanidine amide catalysts **54.4** in the asymmetric formal C–C bond insertion reaction (Scheme 54).<sup>132</sup> Interestingly, in the cases of *ortho*-

**Scheme 54. Asymmetric Formal C–C and C–H Bond Insertion Using Chiral Guanidine Amide Catalysts**

Lui 2023

a) formal C–C bonds insertion




substituted aryl diazo esters **54.13**, formal C–H bond insertion products, 1,3-diketones **54.14**, were obtained in moderate yields. This chemoselectivity was not observed in the previous catalyst-free racemic approach and chiral phosphoric acid-catalyzed process. Control experiments and DFT calculations indicated a stepwise reaction mechanism rather than a direct cyclopropanation process, which could explain the influence of substitution at the aryl group of  $\alpha$ -diazoesters and the hydrogen-bonding ability of the catalyst on both the chemoselectivity and enantioselectivity observed in the reaction.


In 2022, Rostovskii and co-workers reported a photochemical approach for construction of various ketenimines **55.3** from donor–acceptor type diazo compounds **55.1** and alkyl isocyanides **55.2** (Scheme 55a).<sup>133</sup> The DFT calculations and mechanistic studies showed that carbenes in both singlet and triplet states can react with isocyanides; however, only the reaction of the former leads to the smooth formation of ketenimines. In the same year, the groups of Xu and Xuan reported preparation of tetrasubstituted alkenes **55.6** from diazo compounds **55.4** and iodonium ylides **55.5** (Scheme 55b).<sup>134</sup> Mechanistically, the nucleophilic attack of carbenes by iodonium ylide afforded intermediate A. A subsequent elimination of iodobenzene provided the final all-carbon tetrasubstituted olefins. However, it should be mentioned that an alternative mechanism involving the photolysis of

**Scheme 55. Synthesis of Ketenimines and Tetrasubstituted Alkenes**

a) Rostovskii 2022




b) Xu & Xuan 2022



iodonium ylide to generate diradical species cannot be ruled out.<sup>135</sup>

An example of free singlet carbenes functioning as hydride acceptors came from Jin's group, who reported the synthesis of functionalized isoxazolidines 56.3 using tertiary amines 56.1 and *N*-hydroximoyl chloride 56.2 (Scheme 56). In this work,

**Scheme 56. Free Carbenes as Hydride Acceptor**



the highly reactive electrophilic carbene species were observed to abstract hydrides adjacent to the nitrogen in tertiary amines, resulting in the formation of ion pairs A and B. The deprotonation occurring at the  $\beta$ -position of the iminium ion afforded the reduced product C, along with an enamine intermediate D. A subsequent 1,3-dipolar cycloaddition of the latter with the *in situ* formed nitrile oxide 56.4 delivered isoxazoline derivatives 56.3.


In 2023, the groups of Guo, Song, and Xia reported a Mn-catalyzed trifluoromethylsulfonylation reaction of aryl diazoesters 57.1 to prepare  $\alpha$ -trifluoromethyl sulfone esters 57.2 (Scheme 57).<sup>136</sup> This transformation features the generation of trifluoromethyl sulfone radicals A through coordination to the Mn(acac)<sub>3</sub> catalyst, followed by their combination with a carbene, producing a stable radical intermediate B.

Very recently, the groups of Cheng and Liu disclosed a Pd-catalyzed strategy for the directed C–H alkylation of 1-naphthylamines 58.2 with  $\alpha$ -diazoesters 58.1 (Scheme 58).<sup>137</sup> The proposed mechanism suggests the formation of carbene species under blue light irradiation, which undergo addition to Pd<sup>II</sup> complexes. However, the reactions required a considerable excess of diazo substrates, which served as the solvent.

As previously discussed, aryl-substituted NHPI diazoacetates were involved in C–H insertions with alkanes. In the same report, Gryko and co-workers observed that NHPI diazo-


**Scheme 57. Trifluoromethylsulfonylation Reaction of Aryl Diazoesters Enabled by Manganese Catalysis**

Guo, Song & Xia 2023




**Scheme 58. Pd-Catalyzed C–H Alkylation of 1-Naphthylamines with  $\alpha$ -Diazoester**

Cheng & Liu 2023



acetates with alkyl substitutions 59.1 can also undergo photolysis to produce singlet carbenes (Scheme 59).<sup>64</sup> However, in this scenario, either a 1,2-H shift (59.3 and 59.4) or an alkyl shift (59.5) occurs, producing vinyl NHPI esters 59.2.

**Scheme 59. 1,2-Rearrangement of Alkyl Redox-Active Diazo Compounds**




### 3. DIAZO COMPOUNDS AS TRIPLET CARBENE PRECURSORS

Compared with extensive studies on transformations via singlet carbenes, reactions involving triplet carbenes are relatively underdeveloped.<sup>138</sup> A direct photolysis of diazo compounds can lead to triplet carbenes, although singlet carbenes are typically formed through this process, as the nitrogen dissociation of the excited diazoalkene is usually faster than intersystem crossing to the triplet state. The electronic properties of substituents attached to carbenes greatly affect the spin state and the energy difference between their singlet and triplet states. Thus, it is possible to switch the spin state by modifying substitutions on them, for example, by employing donor–donor type diazo compounds with tunable aryl rings. Diphenyl carbene favors the triplet state over the singlet state

by 4 kcal/mol,<sup>139</sup> while the energy difference between the singlet and triplet states of the electron-rich bis(4-methoxyphenyl)carbene is nearly negligible, with the singlet state being only 0.3 kcal/mol energetically favored, as was reported by Sander and co-workers.<sup>140</sup> They reported that the spin state of bis(4-methoxyphenyl)carbene can be reversibly switched between the singlet and triplet states by utilizing different wavelengths (Scheme 60).<sup>140</sup> Furthermore, Sander

### Scheme 60. Spin State of Bis(4-methoxyphenyl)carbene

Savitsky, Sanchez-Garcia & Sander 2016



reported the switching of the spin ground state of diphenylcarbene from triplet to singlet when the carbene interacts with methanol.<sup>141</sup> The singlet carbene becomes more stable compared to its triplet state by forming hydrogen bonds with an alcohol, which represents a unique way to control chemical reactivity. This effect was further analyzed in-depth by Sanchez-Garcia and Nuernberger, using broadband femtosecond transient absorption and quantum mechanics/molecular mechanics simulations.<sup>142</sup>


Another important method to access triplet carbenes is the denitrogenative decomposition of diazo compounds through energy transfer with a photosensitizer. The employment of a photosensitizer allows for mild, efficient activation of diazo compounds that may not have sufficient absorption in the visible light region.

In 2016, the groups of Zawada, Kadish, and Gryko reported a photosensitized approach for  $\alpha$ -alkylation of aldehydes **61.2** using diazo compounds **61.1** (Scheme 61). In this transformation, acceptor-type diazo compounds underwent energy transfer with a free base tetraphenylporphyrin ( $H_2TPP$ ) catalyst **61.4**, leading to the formation of triplet carbenes. Various diazoacetates were found to be competent substrates (**61.5–61.7**). Notably, aldehydes bearing an alkene moiety furnished C–H alkylated compound **61.8** exclusively, without the formation of cyclopropane. The reaction mechanism begins with the photoexcitation of  $H_2TPP$  from the singlet ground state to the excited state. The energy transfer from the catalyst to diazoacetates produced triplet carbene with simultaneous loss of dinitrogen. In organocatalytic cycle, enamines **A** were formed from aldehyde substrate **61.2** in the presence of an amine catalyst. Subsequently, the photo-excited  $H_2TPP$  oxidizes enamine **A** to form a porphyrin radical anion and an active cation radical **B**. The latter reacts with triplet carbene, furnishing a radical intermediate **C**. A following single electron transfer from the porphyrin radical anion and a subsequent protonation led to the final alkylated products **61.3**. Notably, the porphyrin serves both as a photoredox catalyst and as a photosensitizer in this transformation. However, it should be noted that chain propagation processes may also be involved.<sup>143</sup>

In 2021, Koenigs and co-workers conducted a systematic study on the photochemical generation of free carbenes from diaryl diazo compounds **62.1**, along with their subsequent chemoselective reactions with terminal alkynes **62.2** (Scheme 62).<sup>144</sup> The introduction of electronically diverse substituents at


### Scheme 61. Generation of Triplet Carbene via Energy Transfer

Zawada, Kadish & Gryko 2016



### Scheme 62. Reaction of Aryl–Aryl Diazo Compounds with Terminal Alkynes via Triplet Carbene Intermediates


Koenigs 2021



the aryl rings effectively controlled the reactivity of free diaryl carbenes. This led to a significant perturbation of the singlet–triplet energy splitting of the diaryl carbene intermediates, as well as the activation energies of their subsequent reactions. For instance, the reaction of an electron-neutral diazo compound with terminal alkynes afforded indenes **62.3**, following a stepwise process involving a triplet carbene intermediate formed through intersystem crossing from a singlet carbene state.

A year later, the same research group developed a photosensitized method for generating triplet carbenes from acceptor-type diazo compounds **63.1** using an Ir complex as a photosensitizer (Scheme 63A).<sup>145</sup> This approach allows for efficient *gem*-difluoroalkylation reactions of  $\alpha$ -trifluorome-


**Scheme 63. Photocatalytic *gem*-Difluoroalkenylation, Stereoconvergent Cyclopropanation Reactions and C3-Alkylation of Indoles by Diazo Acetates via Triplet Carbene Intermediates**



thylstyrenes **63.2**. The produced triplet carbene undergoes efficient addition to  $\alpha$ -trifluoromethylstyrenes, followed by the  $\beta$ -F elimination to furnish the products **63.3**. It was also found that the use of a tertiary amine is crucial for preventing the formation of cyclopropanes. Building on this success, the authors reported a photocatalytic strategy that allows the employment of a mixture of *E*- and *Z*-isomers of olefins **63.5** to give access to a single isomer of cyclopropane products **63.6** in a stereoconvergent reaction (Scheme 63B).<sup>146</sup> Both experimental and theoretical evidence indicate the presence of a triplet carbene intermediate, which readily reacts with either isomer of the olefin. The intersystem crossing from triplet to singlet state proceeds in a diastereoselective fashion, offering an explanation for this highly stereoselective reaction. Additionally, the authors adapted this photosensitization strategy for synthesis of multisubstituted furans.<sup>147</sup> Very recently, Gryko and Koenigs further investigated reactivity of triplet carbenes in a photosensitized alkylation reaction of indoles **63.7** using diazo acetates **63.4** (Scheme 63C).<sup>148</sup> While the C2-alkylation occurs in protic solvent, the employment of aprotic solvents leads to exclusive C3-functionalization (**63.8**). Mechanistic investigations and DFT calculations revealed that the unusual selectivity arises from the generation of triplet carbenes in aprotic solvents. Although the triplet carbene initially adds to the C2 position of indoles, a subsequent 1,2-alkyl radical shift and 1,2-hydrogen shift led to the observed C3-alkylated products in aprotic solvents.

In 2022, the groups of Zhou and Yang reported a visible light-induced aerobic oxyphosphorylation reaction of  $\alpha$ -diazoesters **64.1** and H-phosphine oxides **64.2** (Scheme 64).<sup>149</sup> In this transformation, the authors proposed that the triplet carbene **B** serves as a HAT reagent, which upon reaction with phosphine oxides produce phosphinoyl radicals **C**. The latter is then promptly intercepted by molecular oxygen to

**Scheme 64. Triplet Carbene as an HAT Reagent**



deliver a peroxy radical intermediate **D**, which then abstracts a hydrogen atom to furnish phosphoryl hydroperoxide **E**. A subsequent homolysis and radical–radical coupling afforded the desired product **64.3**.

## 4. DIAZO COMPOUNDS AS ALKYL OR VINYL RADICAL PRECURSORS

### 4.1. Introduction


Diazo compounds have traditionally served as precursors for carbenes, and significant advancements have been made in this field. In contrast, radical reactions involving diazo compounds have received less attention. Nevertheless, with the recent emergence of photocatalysis<sup>18–20</sup> and visible-light induced transition metal catalysis,<sup>150,151</sup> new possibilities have arisen. The combination of these photocatalytic methods with diazo compounds has led to the discovery of novel radical reactivities that differ from classical carbene transformations. Typically, a photocatalyst reduces diazo compounds through a single electron transfer process. After extrusion of dinitrogen and protonation by an acid or a protic solvent, radical intermediates are formed. Acceptor-type diazo compounds, such as  $\alpha$ -diazo esters, are more commonly used in these processes, possibly due to their limited absorption in visible light region, thus making them less prone to decomposition or side reactions. The resulting radicals can then participate in various transformations, such as addition to alkenes or heteroarenes, as well as recombination with other radical species. These breakthroughs have significantly expanded synthetic applications of diazo compounds, offering new opportunities in organic synthesis.

### 4.2. Radical Addition to Alkenes

**4.2.1. Giese- or Heck-Type Reactions.** In 2016, Gryko and co-workers reported cobaltested-catalyzed method for the generation of alkyl radicals from diazoacetates **65.1** under white LEDs, allowing for alkylation reaction of alkenes **65.2** (Scheme 65).<sup>152</sup> The resulting alkyl radicals undergo an intermolecular Heck-type reaction. However, the authors observed a mixture of Heck-type (oxidative) and Giese-type (reductive) products. Thus, a subsequent hydrogenation step was performed to obtain reduced products **65.3**.  $\alpha$ -Diazo esters reacted well with aliphatic or disubstituted olefins (**65.3–65.5**). However, transformations with styrene derivatives

**Scheme 65. Generation of Alkyl Radicals from Diazo Compounds via Cobalt Catalysis**

Gryko 2016




(65.6) were found to be less efficient. The proposed mechanism begins with the reduction of cobalester to form the catalytically active Co(I) species. Upon reaction with diazo compounds, alkyl-Co(III) intermediates A are formed. A subsequent homolytic cleavage of the Co–C bonds produces electrophilic radical B, which upon addition to the electron-rich alkene affords radical C, that can either recombine with Co(II) species or abstract a proton, leading to reduced products. Dehydrocobalatation of intermediate D yields the Heck-type product. Alternatively, the Giese-type products can be formed through reduction by hydridocobaltester. It is important to note that, in all cases, the typical cobalt-catalyzed cyclopropanation process<sup>153,154</sup> was not observed, highlighting the crucial role of visible light irradiation in this reaction.

In 2020, the groups of Qiu and Doyle employed diazo compounds 66.1 to access primary and secondary alkyl radicals via reductive proton coupled electron transfer (PCET) with a Ru<sup>II</sup> photocatalyst (Scheme 66).<sup>155</sup> The formed C-centered radicals efficiently added across both aliphatic alkenes and styrene derivatives, allowing for synthesis of various anti-Markovnikov hydroalkylated products (66.4–66.12). The proposed mechanism begins with photoexcitation of the ruthenium complex Ru(bpy)<sub>3</sub>Cl<sub>2</sub>·6H<sub>2</sub>O to form Ru<sup>II\*</sup>, which is able to oxidize Hantzsch ester to generate a radical cation

**Scheme 66. Generation of Alkyl Radicals from Diazo Compounds via PCET**


Qiu & Doyle 2020



and Ru<sup>I</sup> species. The newly formed Ru<sup>I</sup> species participates in an SET process with the diazo compound, while regenerating Ru<sup>II</sup> catalyst. A sequential extrusion of N<sub>2</sub> with protonation delivers C-centered radical A, which upon radical addition to alkene B, followed by HAT, delivers Giese-type product 66.3. A thiol catalyst was added for promoting the HAT process. This methodology also marks a distinct difference from the traditional reactivity of diazo compounds in the presence of alkenes, where cyclopropanation via the intermediary of carbenes are dominant. Inspired by this work, many other groups have adapted this photocatalytic PCET strategy for the generation of alkyl radicals from diazo compounds to develop many other transformations. For example, Qiu and Hu's groups successfully developed a hydroalkylation reaction of DNA-tagged alkenes using diazo compounds toward the construction of DNA-encoded libraries using this platform.<sup>156</sup>

In 2022, Gevorgyan's group reported a photoinduced alkyl-Heck reaction of diazo compounds 67.1 via hybrid alkyl-Pd radical species (Scheme 67).<sup>157</sup> A diverse range of acceptor and donor–acceptor type diazo compounds were found to efficiently participate in reactions with styrene derivatives,

**Scheme 67. Heck Reaction of Diazo Compounds via Photoinduced Palladium Catalysis**




leading to the exclusive formation of allylic compounds **67.4–67.8**. Three possible mechanistic pathways were proposed for this reaction. Path **a** begins with oxidative addition of Pd(0) into HX to afford an active palladium hydride species. A denitrogenative reaction between H–Pd<sup>II</sup>X with diazo compound takes place, producing a Pd–carbene intermediate **A**. A subsequent hydride shift leads to the formation of palladium species **B**. Upon blue light irradiation, the C–Pd bond undergoes homolysis, producing a hybrid alkyl Pd–radical species **C**. Alternatively, the formation of the latter can occur through two additional pathways. In path **b**, it is generated by a single electron transfer (SET) to diazonium ion, which is the protonated form of the diazo compound. In path **c**, a PCET takes place directly from the diazo compounds, furnishing species **C**. Subsequently, the hybrid alkyl Pd–radical species **C** undergoes an addition reaction with styrene to deliver a hybrid benzylic radical **D**. The rest of the catalytic cycle follows the classical Heck reaction mechanism, which includes a  $\beta$ -H-elimination step to afford the desired Heck products **67.3**.

**4.2.2. 1,2-Difunctionalization of Alkenes.** Upon the addition of radicals to alkenes, the newly formed radical species can be effectively captured by various radical traps. Two C–C or C–heteroatom bonds are formed efficiently in these

multicomponent photocatalytic reactions of diazo compounds to deliver a wide range of 1,2-difunctionalized products under mild conditions.

In 2021, the groups of Yang and Xia employed the photocatalytic PCET strategy to generate radicals from ethyl diazoacetate and to engage **68.1** in a 1,2-carboamination reaction of alkenes **68.2** with diarylamines **68.3** (Scheme 68).<sup>158</sup> In this transformation, alkyl radicals **A** efficiently added

**Scheme 68. Photocatalytic 1,2-Amino-alkylation of Styrenes with Diazoacetates and Diarylamines**



to various styrene derivatives, affording relatively persistent benzylic radical intermediates **B**. The final products were formed through a radical recombination between the latter and transient N-centered radicals **C** derived from diarylamines.


In 2022, Wang and co-workers disclosed a method for 1,2-alkyl-alkylation of alkenes **69.2** by employing diazo esters **69.1** and aldehydes **69.3**, featuring cooperative photoredox and NHC catalysis (Scheme 69).<sup>159</sup> A wide range of functionalities on benzaldehydes were well tolerated (**69.6–69.11**). Vinyl heteroarenes also proven to be competent substrates (**69.12** and **69.13**). The proposed mechanism commences with the generation of an active NHC catalyst from thiiazolium in the presence of a base. The produced NHC catalyst then reacts with aldehydes to deliver Breslow intermediate **A**, which undergoes a single electron oxidation by excited photocatalyst to produce ketyl radical **B**. Similarly to the previous approaches, alkyl radicals are produced from diazo compounds through a PCET process. Radical addition to alkenes and a subsequent radical recombination with the ketyl radical **B** afford tetrahedral intermediates **D**. Finally, ketones were formed with the release of the NHC catalyst.

In the same year, Koenigs and co-workers developed a photocatalytic approach to achieve 1,2-oxo-alkylation of vinyl arenes **70.2** employing diazoacetates **70.1** and hydroperoxide **70.3** (Scheme 70).<sup>160</sup> Alkyl radicals efficiently added to alkenes, followed by the capture of the resulting benzylic radical by peroxy radical, resulting in the formation of peroxy ethers. Under basic conditions, Kornblum–DeLaMare rearrangement of peroxy ethers occurred, affording 1,2-difunctionalized products **70.4**.

The groups of Qiu and Doyle reported a three-component photocatalytic Minisci reactions of *N*-heterocycles **71.3**, diazo compounds **71.1**, and alkenes **71.2** (Scheme 71).<sup>161</sup> Various aliphatic alkenes and acceptor-type diazo compounds under-


**Scheme 69. Difunctionalization of Alkenes via Cooperative Photoredox and NHC Catalysts**

Wang 2022




**Scheme 70. Photocatalytic 1,2-Oxo-Alkylation of Styrenes with Diazoacetates and Hydroperoxides**

Koenigs 2022



**Scheme 71. Photocatalytic Minisci Reactions**

Qiu & Doyle 2022




went smooth transformation with electron-deficient *N*-heteroarenes in good yields (71.4–71.8). An example of benzimidazole was also demonstrated, albeit with diminished efficiency (71.9). In this transformation, electrophilic C-centered radicals **A** generated from acceptor diazo compounds **71.1** undergo selective addition to alkenes, affording nucleophilic radicals, which readily react with various electron-deficient *N*-heteroarenes. Trifluoroacetic acid (TFA) was employed to protonate both the heteroarene and the diazo compound in this reaction.

Zhao, Li, and co-workers reported a photocatalytic difunctionalization reaction of alkenes **72.2** with diazo compounds **72.1** and *tert*-butyl nitrites **72.3** (Scheme 72).<sup>162</sup> In this case, *tert*-butyl nitrites were used as radical traps, which allowed for synthesis of various  $\gamma$ -oximino esters **72.4**.

**Scheme 72. Photocatalytic Difunctionalization of Alkenes with Diazo Compounds and *tert*-Butyl Nitrite**


Zhao & Li 2022



### 4.3. Radical Addition to Heteroarenes

Apart from alkenes, the electrophilic radicals produced from acceptor type diazo compounds undergo efficient addition to electron-rich heteroarenes. In 2019, Gryko and co-workers reported a photocatalytic method for alkylation of pyrroles and indoles **73.2** using  $\alpha$ -diazo esters **73.1** (Scheme 73a).<sup>163</sup> This approach allows for synthesis of diverse C2-alkylated indole and pyrrole derivatives **73.3**. In the case of **73.6**, C3-alkylation

**Scheme 73. Photocatalytic Alkylation of Pyrroles and Indoles with  $\alpha$ -Diazo Esters**




product prevails, possibly as a result of a parallel carbene C–H insertion pathway.<sup>16</sup> N-methyl protected and unprotected indoles, as well as pyrroles, delivered the respective products in good yields (73.4–73.9). When electron-deficient substrates are used, the addition of a sacrificial reductant, such as amine, is required. In protic solvent, diazo compounds undergo protonation to form diazonium ions A, which can be reduced to alkyl radicals B via a SET by excited Ru photocatalyst. Subsequently, the alkyl radicals undergo addition to N-heterocycle to give radical C. After oxidation and deprotonation, the desired alkylated product is formed. A year later, Hajra and co-workers reported a similar radical approach using diazo compounds for C3-alkylation of imidazopyridines under the same reaction conditions (Scheme 73b).<sup>164</sup> In addition to aforementioned Ru complex, graphitic carbon nitride ( $\text{g-C}_3\text{N}_4$ ) also could serve as a photocatalyst in alkylation of indoles with diazo compounds, as shown by Zhou and Guo.<sup>165</sup>

#### 4.4. Radical–Radical Recombinations

Besides radical additions to alkenes and heteroarenes, as mentioned above, alkyl radicals derived from diazo compounds can also participate in radical–radical couplings. In 2022, Cho and co-workers successfully developed a photocatalytic alkylation reaction of anilines 74.1 with acceptor–acceptor type diazo compounds 74.2 (Scheme 74A).<sup>166</sup> This protocol showed no sensitivity to electronics, as both electron-deficient and electron-rich anilines performed equally well. The proposed mechanism involves radical recombination between persistent radical A and transient radical B, followed by rearomatization, affording *para*-C–C coupled aniline derivatives. In the same year, the groups of Hu and Qiu developed a

**Scheme 74. Photocatalytic Transformations Involving Radical–Radical Recombinations**




photocatalytic, three-component reaction of secondary amines, aldehydes, and diazo compounds (Scheme 74B).<sup>166</sup> This protocol afforded various  $\gamma$ -amino acid derivatives in good yields. In this reaction, enamines were formed *in situ* via condensation between aldehyde 74.6 and secondary amine 74.5, which can be oxidized to radical intermediate C by an excited photocatalyst. Radical recombination between intermediate C and a transient  $\alpha$ -ester radical D occurs, followed by reduction using a Hantzsch ester, delivering the desired  $\gamma$ -amino acid derivatives. Very recently, the groups of Hou and Li reported a photocatalytic approach for synthesis of various indole-fused tricyclic compounds 74.9 using diazo compounds 74.4 and N-phenylpiperidines 74.8 (Scheme 74C).<sup>167</sup> The mechanism of this protocol features the recombination between nucleophilic radical E and electrophilic radical D.

#### 4.5. Other Radical Reactions

In 2016, Meggers and co-workers pioneered a photocatalytic approach employing  $\alpha$ -diazo esters 75.2 for asymmetric alkylation of 2-acyl imidazoles 75.1 (Scheme 75).<sup>168</sup> This method was enabled by the combination of a chiral Lewis acidic  $\Delta$ -RhS complex with a Ru photocatalyst. A variety of acceptor-type diazo compounds were compatible with this transformation, delivering the alkylated products in good yields and high enantioselectivities (75.5–75.8). The mechanism of this reaction features dual photoredox and asymmetric catalytic cycles. Upon coordination of the substrate with the chiral rhodium complex A and deprotonation, rhodium enolate B is formed. This electron-rich enolate acts as a single electron

## Scheme 75. Asymmetric Alkylation with Diazo Compounds

Meggers 2016




donor to the photoexcited ruthenium complex, leading to the formation of strongly reducing  $[\text{Ru}(\text{bpy})_3]^{+}$ , which subsequently transfers a single electron to the diazo esters. Extrusion of  $\text{N}_2$  and protonation produces alkyl radicals, which stereoselectively add to the rhodium enolate to form the Rh-coordinated ketyl radical C. This highly reducing agent can either directly reduce the diazo reagent for chain propagation or quench the photoexcited  $[\text{Ru}(\text{bpy})_3]^{2+}$ . Ultimately, the alkylated products are obtained through the oxidation of radical D and the recoordination of the substrate with  $\Delta\text{-RhS}$ .

In 2019, Rastogi and co-workers reported a visible light-induced benzannulation reaction of diazo ketones 76.1 and alkynes 76.2 (Scheme 76).<sup>169</sup> This reaction proceeds through single-electron reduction of the diazo enolate by Hantzsch ester to form vinyl radical A. Then, this radical is subsequently

## Scheme 76. Visible Light-Induced Benzannulation Reaction of Diazo Ketones and Alkynes

Rastogi 2019




trapped by alkyne, delivering functionalized naphthalenol 76.3 in good efficiency.

In 2022, Li, Xuan and co-workers disclosed an organophotocatalytic protocol for synthesis of phenanthridines 77.3 using acceptor-type diazo compounds 77.1 and biphenyl isocyanides 77.2 (Scheme 77).<sup>170</sup> Similarly to the previous

## Scheme 77. Organophotocatalytic Approach for Synthesis of Phenanthridines

Li &amp; Xuan 2022



report, alkyl radical A is generated via PCET mechanism. The formed C-centered radical A then engages in a radical addition/intramolecular radical cyclization cascade with biphenyl isocyanates. The resulting 2-alkylated phenanthridine B exists in equilibrium with its tautomer C, which can capture another equivalent of an alkyl radical to afford phenanthridine product bearing a succinate moiety.

In the same year, the groups of Kessler, Röschenthaler, and Han introduced a photocatalytic approach for synthesizing  $\alpha$ -fluorovinylphosphonates 78.2 from difluoroalkyl diazo compounds 78.1 (Scheme 78).<sup>171</sup> Differently from the previously

Scheme 78. Synthesis of  $\alpha$ -Fluorovinylphosphonates

Kessler, Röschenthaler &amp; Han 2022




reported PCET mechanism, the authors proposed that the alkyl radical A is produced through a Hantzsch ester-assisted HAT process, as supported by mechanistic experiments and DFT calculations.

More recently, Jiang and co-workers reported a cooperative photoredox and NHC-catalyzed protocol for 1,3-difunctionalization of [1.1.1]propellane 79.2 using  $\alpha$ -diazo esters 79.1 and benzaldehydes 79.3 (Scheme 79a).<sup>172</sup> Benzaldehydes bearing various functional groups efficiently participated in this transformation, producing various bicyclo[1.1.1]pentanes

**Scheme 79. Synthesis of Disubstituted Bicyclo[1.1.1]pentanes**

a) Jiang 2023



b) Jiang 2023



**79.6–79.14.** Heteroaryl aldehydes (79.12 and 79.13) were also capable substrates. The proposed mechanism begins with the generation of NHC A in the presence of a base, which then reacts with aldehydes to form the Breslow intermediates B. Subsequently, the photoexcited Ir<sup>III</sup> complex oxidizes the intermediate B, leading to the formation of radical species C. The resulting Ir<sup>II</sup> catalyst then reduces diazo ester, generating an alkyl radical through a PCET process. This transient radical rapidly adds to [1.1.1]propellane to afford the radical species D. The recombination of two radical intermediates eventually leads to the formation of difunctionalized product 79.4, along with the regeneration of the NHC catalyst. Next, the same group expanded the scope of radical acceptors to quinoxalinones 79.15 in analogous difunctionalization of [1.1.1]-propellanes (Scheme 79b).<sup>173</sup>

**5. DIAZO COMPOUNDS AS DIAZOMETHYL RADICAL PRECURSORS**

**5.1. Introduction**

In above-mentioned reports, the generation of reactive intermediates require the release of dinitrogen from diazo compounds. However, by utilizing specifically designed substrates, such as  $\alpha$ -diazo onium salts, the valuable diazo functional group can be retained.<sup>174</sup> These diazo species possess a redox-active leaving group, such as a hypervalent iodine<sup>175,176</sup> or sulfonium<sup>177</sup> group. These substrates could be converted into diazomethyl radicals, thus unlocking new reactivities that can be accessed from diazo compounds using visible light. Initial reports focused on the employment of a photocatalyst, which produces the key diazomethyl radical species through a single-electron reduction event. However, later studies demonstrated the possibilities of visible light-induced homolytic cleavage of the weak C–I bond in hypervalent iodine reagents, without the need of a catalyst, to arrive at the same reactive intermediates. The produced diazomethyl radical is highly electrophilic, which undergoes efficient addition to arenes, delivering C–H functionalized products, or participates in a stepwise cyclization with other unsaturated systems, affording various heterocycles.

**5.2. Radical Addition to Arenes**

In 2018, Suero and co-workers pioneered a photocatalytic approach for the generation of diazomethyl radicals from  $\alpha$ -diazo iodonium triflates 80.1 (Scheme 80).<sup>178</sup> This mild protocol allowed for efficient C–H diazomethylation of arenes 80.2, affording various aryl diazo esters 80.3. This method is also amenable to the functionalization of complex molecules 80.10 and 80.11. Under white light irradiation, hypervalent iodine reagents undergo single-electron reduction by a photoexcited Ru complex, leading to the formation of key diazomethyl radical intermediates A. This electrophilic radical undergoes efficient addition to arenes, affording cyclohexadienyl radical B. Subsequently, oxidation with Ru<sup>III</sup> complex and proton elimination deliver the desired diazomethylation product. Furthermore, it was demonstrated that the obtained products can undergo various carbene transformations, enabling the efficient synthesis of complex chiral molecules from readily available chemical feedstocks.


In 2023, Li and co-workers developed a photocatalytic cascade reaction for the preparation of acridines 81.3 via double sp<sup>2</sup> C–H bond activation of diarylamines 81.2 by using hypervalent iodine reagents 81.1 (Scheme 81).<sup>179</sup> Similarly to the Suero's method, the mechanism commences with an SET from the photoexcited Ru complex to hypervalent iodine substrate, producing the key diazomethyl radical intermediate A. A subsequent radical addition of the latter to the aryl ring of diphenylamine affords radical intermediate B. Upon oxidation and proton loss, diazo compound C is formed, which produces free carbene species D under blue light irradiation with the extrusion of dinitrogen. Finally, the intramolecular cyclization of D and oxidation leads to the formation of acridine 81.3. Notably, this reaction highlights the dual reactivity of hypervalent iodine diazo reagents, producing both radical and carbene species.

**5.3. Heterocyclization**

Besides radical addition to arenes, diazomethyl radicals can also undergo cyclization reactions with different unsaturated  $\pi$ -systems toward various N-heterocycles. In 2020, the groups of

**Scheme 80. Generating Diazomethyl Radicals via Photoredox Catalysis**


Suero 2018



Liu and Li reported a photocatalytic approach for synthesizing oxadiazoles **82.3** from hypervalent iodine reagents **82.1** and benzoyl formic acids **82.2** (Scheme 82a).<sup>180</sup> The proposed mechanism begins with the photoexcitation of 4CzPN catalyst, followed by its reductive quenching by benzoyl formic acid, producing a benzoyl cation radical **A** and 4CzPN<sup>•-</sup>. Subsequent deprotonation and decarboxylation afford benzoyl radical **B**. Upon single electron reduction by 4CzPN<sup>•-</sup>, a diazomethyl radical is formed, which is a resonance form of intermediate **C**. The recombination of **C** with benzoyl radical **B**, followed by cyclization, furnish the targeted oxadiazole. A year later, the groups of Chen and Han revealed an analogous catalyst-free method toward oxadiazoles **82.6** (Scheme 82b).<sup>181</sup> In this case, the proposed mechanism suggests that diazotrifluoroethyl radical is formed through visible-light-induced homolysis of the C—I bond in hypervalent iodine reagent with the concurrent generation of iodanyl radical **A**. Subsequently, a HAT process between iodanyl radical **A** and  $\alpha$ -ketoacid **82.5**, followed by decarboxylation, affords the benzoyl radical **D**. Similarly to the previous method, radical–radical coupling and ring-closure furnished the heterocyclic product. In the same year, the groups of Li and Li reported a similar approach employing benzaldehydes and hypervalent iodine

**Scheme 81. Photocatalytic Synthesis of Acridines**

Li 2023





reagents for synthesizing oxadiazoles under catalyst-free conditions.<sup>182</sup>

In 2021, Alcarazo and co-workers employed  $\alpha$ -diazo sulfonium triflates **83.1** for the generation of diazomethyl radicals under photocatalytic conditions, allowing for efficient synthesis of *N*-amino triazoles **83.3** with hydrazones **83.2** (Scheme 83).<sup>183</sup> Mechanistically, the photoexcited Ru complex is capable of reducing  $\alpha$ -diazo sulfonium triflates via a single-electron transfer process, producing the key diazomethyl radical **A**. Addition of the latter to the azomethine carbon of *N,N*-dialkyl hydrazones affords an aminyl radical species **B**, which is subsequently oxidized by Ru<sup>III</sup> complex. Upon deprotonation, intermediate **C** is formed, which cyclizes into triazoles. Simultaneously, Li's group reported a similar photocatalytic method employing hypervalent iodine reagents **84.1** for synthesis of triazoles **84.3** (Scheme 84a).<sup>184</sup> In 2022, the groups of Liu and Li disclosed a related photocatalytic strategy for synthesizing triazoloquinoxalinones **84.5** from hypervalent iodine reagents **84.1** and quinoxalinones **84.4** (Scheme 84b).<sup>185</sup> This reaction proceeds via a similar mechanism involving diazomethyl radical intermediates. Subsequently, Pla-Quintana, Poater, and co-workers conducted DFT calculations of this transformation, unveiling that the rate-determining step is the C–C bond formation step through the attack of diazomethyl radicals onto the C–N double bond of the quinoxalinone.<sup>186</sup>



In 2022, the group of Yuan and Zhao reported a photochemical catalyst-free annulation reaction of  $\alpha$ -diazo sulfonium triflates **85.1** with  $\alpha$ -oxocarboxylic acids **85.2** and alkynes, producing oxadiazoles **85.3** and pyrazoles **85.4**, respectively (Scheme 85).<sup>187</sup> In this catalyst-free reaction, the authors proposed that diazomethyl radicals were formed through the homolytic cleavage of S–C bond in  $\alpha$ -diazosulfonium triflates upon exposure to blue light irradiation.

## Scheme 82. Synthesis of Oxadiazoles via Diazomethyl Radical Intermediates



a) Liu &amp; Li 2020: photocatalytic approach



b) Chen &amp; Han 2021: photochemical catalyst-free approach

Scheme 83. Synthesis of N-Amino Triazoles from  $\alpha$ -Diazo Sulfonium Triflates

Alcarazo 2021




## 6. DIAZO COMPOUNDS AS RADICAL ACCEPTORS


Diazo compounds can act as acceptors for radicals, generating an additional free radical by releasing dinitrogen, enabling cascade transformations. In 2018, Wan and co-workers reported a photocatalytic approach to access oxalate esters 86.3 from diazo compounds 86.2 and  $\alpha$ -bromo-ketones 86.1 in the presence of molecular oxygen (Scheme 86).<sup>188</sup> The mechanism of this reaction begins with the single-electron reduction of  $\alpha$ -bromo-ketones by photoexcited eosin Y to generate alkyl radical species A, which is then trapped by molecular oxygen, producing superoxide radical B. Upon HAT and a subsequent homolysis of the weak O–O bond, the oxygen-centered radical C is formed. Radical addition to diazo compound and a subsequent extrusion of dinitrogen produces radical intermediate D. Similarly, successive reactions of radical

## Scheme 84. Synthesis of Triazole Derivatives from Hypervalent Iodine Reagents


a) Li 2021



b) Liu &amp; Li 2022

Scheme 85. Synthesis of Oxadiazoles and Pyrazoles from  $\alpha$ -Diazo Sulfonium Triflates

Yuan &amp; Zhao 2022




D with molecular oxygen produce oxygen-centered radical G, which is then oxidized to furnish oxalate ester.

One year later, Yu and co-workers disclosed a photocatalytic method for synthesizing N-alkyl hydrazones 87.3 from alkyl N-hydroxyphthalimide esters (NHPI esters) 87.2 and diazo compounds 87.1 (Scheme 87).<sup>189</sup> Donor–acceptor type diazo compounds were found to be suitable substrates. Additionally, primary, secondary, and tertiary alkyl radicals all effectively participated in this transformation 87.4–87.9. A plausible mechanism for this reaction begins with the single-electron

**Scheme 86. Photocatalytic Method for Accessing Oxalate Esters from Diazo Compounds and  $\alpha$ -Bromo-ketones**

Wan 2018





reduction of the redox-active ester **87.2** by a photoexcited rose bengal catalyst. A subsequent decarboxylation leads to alkyl radical **A**, which undergoes addition to the terminal nitrogen of the diazo compound. The resulting radical species **B** is a resonance form of intermediate **C**, which, upon HAT from the Hantzsch radical cation, produces *N*-alkyl hydrazone as the final product.

In 2021, Zhou and co-workers utilized vinyl diazo compounds **88.1** as radical acceptors to construct fluorinated acridines **88.3** in a cascade radical cyclization reaction (Scheme 88).<sup>190</sup> In this transformation, chlorodifluoromethyl alkynyl ketoimine **88.2** serves as a radical precursor, which upon single-electron reduction by a photoexcited Ru complex produces difluoromethyl radical **A**. A subsequent trapping of the latter by vinyl diazo compounds generates a vinyl radical **B**, which intramolecularly adds to C–C triplet bond of ketoimine to deliver vinyl radical **C**. Then, intramolecular homolytic aromatic substitution (HAS) and oxidation provides vinyl cation **D**, which upon elimination of HF furnishes 4-fluoroacridine as the final product. In addition to chlorides, perfluoroalkyl iodides were also employed as radical sources in this denitrogenative reaction with vinyl diazo compounds. Thus, the groups of Ke and Zhou documented an atom transfer radical addition (ATRA) reaction targeting fluorinated vinyl iodides.<sup>191</sup> It was found that the reaction does not require external photocatalysts or additives.

In the following studies, the same group developed a photochemical 1,3-selenosulfonylation reaction that also employs vinyl diazo compounds **89.1** (Scheme 89A).<sup>192</sup> In this transformation, selenosulfonates **89.2** serve as radical precursors, which allows for the efficient preparation of various  $\gamma$ -seleno allylic sulfones **89.3**. A plausible mechanism was proposed that begins with the visible light-induced homolysis of Se–S bonds, producing sulfonyl **A** and selenyl radicals. The addition of an electrophilic tosyl radical **A** to the C–C double bond of vinyl diazo compounds results in the formation of radical **B**, which has a resonance structure represented by

**Scheme 87. Synthesis of *N*-Alkyl Hydrazones**

Yu 2019




intermediate **C**. Following a 1,2-radical shift and the extrusion of  $\text{N}_2$ , vinyl radical **D** is formed, which affords 1,3-selenosulfonylation product upon reaction with selenosulfonates **89.2**. A year later, they expanded this photochemical strategy to include alkynyl diazo compounds **89.4**, thus enabling synthesis of tetrasubstituted allenes **89.6** (Scheme 89B).<sup>193</sup> The reaction mechanism closely resembles the one described earlier. Moreover, apart from selenosulfonates, it was observed that this method can be applied to thiosulfonates and sulfonyl iodides, delivering 1,3-thiosulfonylation and iodosulfonylation products, respectively.

In the same year, the groups of Koenigs and Xuan reported a photochemical reaction between diazo compounds **90.1**, azodicarboxylates **90.2**, and nitriles **90.3** for preparation of dihydrotriazole **90.4** (Scheme 90).<sup>194</sup> Control experiments and computational studies suggested the formation of a triplet species **A** from azodicarboxylates upon light irradiation. A subsequent radical-type addition to diazo compound produces diradical intermediate **B**. Upon extrusion of dinitrogen, azomethine ylide intermediate **C** is formed, which has a resonance form of intermediate **D**. Finally, dipolar cycloaddition of the latter with acetonitrile **90.3** delivered the desired dihydrotriazole **90.4**.


**Scheme 88. Synthesis of 4-Fluoroacridines via Cascade Radical Cyclization**

Zhou 2021




**Scheme 89. 1,3-Functionalization of Vinyl and Alkynyl Diazo Compounds**

Zhou 2021 — A. 1,3-Difunctionalization of vinyl diazo compounds




Zhou 2022 — B. 1,3-Difunctionalization of alkynyl diazo compounds



Very recently, Zhao, Ma, and co-workers reported a photocatalytic three-component reaction between  $\alpha$ -diazo ketones **91.1**, perfluoroalkyl iodides **91.2** and amino *N*-heterocycles **91.3** (Scheme 91).<sup>195</sup> This method allows for efficient synthesis of *N*-heterocycles containing valuable CF<sub>3</sub> and perfluoroalkyl groups. Mechanistically, the produced perfluoroalkyl radical **A** undergoes efficient addition to the diazo compound, resulting in the formation of radical intermediate **B**. A subsequent single-electron reduction with a phosphoranyl radical delivers anion **C**. Upon  $\beta$ -F elimination, alkenyl fluoride **D** is formed, which cyclizes with amino benzoimidazole, delivering fluorinated tricyclic heterocycle **91.4**.


**Scheme 90. Photochemical Synthesis of Dihydrotriazole from Diazo Compounds, Azodicarboxylates, and Nitriles**

Koenigs & Xuan 2022



**Scheme 91. Synthesis of Fluorinated N-Heterocycles via Multicomponent Cyclization Reactions**


Zhao & Ma 2023



## 7. DIAZO COMPOUNDS AS NUCLEOPHILES

Diazo compounds exhibit ambiphilic reactivity. This property is evident from the resonance structure of the diazo group, which indicates a pronounced negative charge at the carbon to which the diazo group is attached, thus rendering it nucleophilic (Scheme 92).<sup>196</sup> Conversely, the terminal nitrogen of the diazo group serves as a preferred electrophilic site. Among various diazo reagents, vinyl diazo compounds have garnered attention as versatile and valuable substrates in recent

## Scheme 92. Nucleophilic Sites in Diazo Compounds



years.<sup>197</sup> Apart from the above-mentioned nucleophilic carbon to which the diazo group is attached, vinyl diazo compounds possess an extra nucleophilic site at the terminal alkene carbon. Furthermore, the vinylic moiety can be considered an activated olefin capable of engaging in cyclization reactions.

Many transformations featuring diazo compounds as nucleophiles have been documented in the past.<sup>198</sup> More recently, photoredox catalysis has allowed for mild generation of diverse radical species from a wide range of radical precursors. The process of single-electron oxidation produces various radical cations, which can either function as electrophiles themselves or undergo transformations into electrophilic species before reacting with diazo compounds. In certain instances, the valuable diazo functionality remains intact during the nucleophilic addition, thus enabling subsequent functionalizations.


In 2014, Zhou's group reported an organic dye-catalyzed coupling reaction involving tertiary amines **93.1** and  $\alpha$ -diazocarbonyl compounds **93.2** (Scheme 93a).<sup>199</sup> In this

## Scheme 93. Organic Dye-Catalyzed Coupling Reaction Between Tertiary Amines and Diazo Compounds


a) Zhou 2014



mechanism



b) Rastogi 2017




transformation, the diazo functionality is preserved, affording donor–acceptor-type diazo compounds **93.3** as final products. The mechanism begins with the oxidation of the tertiary amine by the photoexcited rose bengal catalyst, yielding radical cation **A** and the reduced form of the catalyst  $\text{RB}^{\bullet-}$ . A subsequent oxidation of the latter by oxygen regenerates the catalyst and produces a superoxide radical anion capable of HAT from radical cation **A**, delivering iminium ion **B**. A nucleophilic attack of the latter by a diazo compound generates intermediate **C**, which is then deprotonated by a hydroperoxide anion to furnish the final product **93.3**. It is important

to note that photolysis of diazo compounds was not observed under the given conditions, likely due to the relatively low energy of the green light utilized in this reaction. Later, Rastogi and co-workers published a follow-up work employing silyl diazoenolate **93.5** as a nucleophile under similar reaction conditions (Scheme 93b).<sup>200</sup>


Building on the efficient nucleophilic addition of diazo compounds to the *in situ* formed iminium ions, in 2016, Zhou's group introduced an additional photocatalytic approach that enables the efficient synthesis of aziridines **94.3** from *N*-aryl glycines **94.1** (Scheme 94).<sup>201</sup> Mechanistically, the decarbox-

## Scheme 94. Synthesis of Aziridines via Organophotoredox Catalysis

Zhou 2016



mechanism




ylation of the radical cation **A** derived from *N*-aryl glycine produces an  $\alpha$ -amino radical **B**, which, following oxidation yields an iminium cation **C**. Subsequently, the nucleophilic addition of a diazo compound to imine **D**, followed by a denitrogenative 3-*exo*-*tert* cyclization, furnishes aziridine.

In 2017, Ferreira and co-workers reported a Cr-catalyzed cyclopropanation reaction of diazo compounds **95.2** and alkenes **95.1** by using CFL as a light source (Scheme 95).<sup>202</sup> It was found that this reaction does not involve a carbene intermediate. Instead, the diazo reagent serves as a nucleophile. The mechanism commences with the single-electron oxidation of electron-rich alkene by a photoexcited Cr(III) catalyst. The resulting radical cation **A** undergoes nucleophilic attack by the diazo compound, which, upon loss of dinitrogen, delivers a radical cation intermediate **C**. A subsequent reduction and intramolecular cyclization furnish the cyclopropane. The reaction proceeds well for alkenes whose reduction potentials fall within the range of +1.11 V to +1.80 V (vs SCE). Different diazo compounds also underwent this transformation smoothly **95.5–95.9**.

Following the successful development of radical cyclopropanation, the same research group has further achieved a formal [3 + 2] cycloaddition reaction of alkenes **96.1** using vinyl diazo compounds **96.2** (Scheme 96a).<sup>203</sup> It was discovered that both chromium and ruthenium complexes are capable of catalyzing this transformation. Although similar yields were obtained for both catalysts, the reaction time was significantly shorter for the Ru catalyst. Mechanistically, the nucleophilic attack of vinyl diazo compound at radical cation **A**

**Scheme 95. Radical Cation Cyclopropanations via Chromium Photooxidative Catalysis**




produces a stabilized benzylic radical **B**. This offers an explanation for the observed high regioselectivity. A rare *5-endo-trig* radical cyclization generates radical species **C**, which upon reduction and loss of  $\text{N}_2$ , forms cyclopentene **96.3**. The authors also proposed an alternative route that involves *4-exo-trig* cyclization of **B**, followed by the ring expansion. Mechanistic studies supported the radical cyclization pathway instead of a cyclopropanation and isomerization process. Later, the authors expanded this formal  $[3 + 2]$  cycloaddition protocol to include indoles **96.7**, which allowed for synthesis of a wide variety of indoline-fused tricyclic compounds **96.9** (Scheme 96b).<sup>204</sup>

In 2020, Nicewicz and co-workers reported an organophotocatalytic approach for C–H alkylation of arenes **97.2** (Scheme 97).<sup>205</sup> It was shown that acceptor-type diazo compounds **97.1** are capable substrates for this transformation. Simple aromatics and electron-rich anisole-derived substrates also reacted well (97.4–97.7); however, regioisomers were obtained in some cases 97.4. Generally, functionalization at *ortho*-position to electron-donating groups is observed, while the *para*-selectivity is favored in more sterically hindered cases. The reaction mechanism begins with the single-electron oxidation of the arene by the photoexcited acridinium catalyst, producing a radical cation **A**. A subsequent nucleophilic addition by diazo compounds leads to the formation of distonic cation radical **B**, which is then reduced by an acridine radical ( $\text{Mes-Acr}\bullet$ ) to form a norcaradiene intermediate **C**. Following a second SET process, intermediate **C** undergoes a ring-opening process to afford distonic benzylic radical cation **D**, which upon reduction and deprotonation, produces alkylated arenes.

Very recently, Koenigs, Zhou, and co-workers developed a photocatalytic method for the self-cyclization of vinyl diazo

**Scheme 96. Formal  $[3 + 2]$  Cycloadditions of Vinyl Diazo Compounds with Alkenes or Indoles**





compounds **98.1** (Scheme 98).<sup>206</sup> This transformation preserves the valuable diazo functional group, allowing for the efficient synthesis of cyclopentenyl  $\alpha$ -diazo compounds **98.2**. The reaction mechanism commences with the single-electron oxidation of the vinyl diazo substrate by the photo-excited  $\text{Ir}^{\text{III}}$  complex, yielding a radical cation intermediate. A subsequent nucleophilic attack of radical cation **A** by another vinyl diazo molecule generates radical intermediate **B**. Following this addition step, a *5-endo* radical cyclization of **B**, with the extrusion of dinitrogen, produces intermediate **C**. Finally, single-electron reduction of **C** by  $\text{Ir}^{\text{II}}$  provides the cyclopentenyl  $\alpha$ -diazo compounds **98.2**.

## 8. MISCELLANEOUS REACTIONS OF DIAZO COMPOUNDS

In 2018, the groups of Shi and Li introduced an iodine-catalyzed approach for activating ethyl diazoacetate **99.1** to access radical intermediates (Scheme 99a).<sup>207</sup> This photocatalytic protocol allows for synthesis of a wide variety of cyclopropanes **99.3**. Upon reaction of diazo compounds with iodine, a diiodoalkyl intermediate **A** is generated, which is reduced by a photoexcited  $\text{Ru}^{\text{II}}$  complex to produce the carbon-centered radical **B**. A subsequent radical addition to an alkene gives a radical intermediate **C**. Intramolecular


## Scheme 97. Regioselective C–H Alkylation of Arenes Enabled by Organic Photoredox Catalysis

Nicewicz 2020



## Scheme 98. Photocatalytic Self-Cyclization of Vinyldiazo Compounds

Koenigs &amp; Zhou 2023




cyclization of the latter affords a cyclopropane **99.3** with the release of an iodine radical. Following this, the produced iodine radical reacts with iodide, forming an anionic radical  $I_2^{-\bullet}$ , which is further oxidized to regenerate iodine and the photocatalyst. Inspired by this work, Chen and co-workers later reported a photochemical iodine-mediated method for homocoupling of diazo substrates, enabling synthesis of tetrasubstituted alkenes.<sup>208</sup> In 2023, Wu's group developed an iodine-mediated multicomponent coupling between trifluorodiazoethane **99.4** and aryl alkynes **99.5** (Scheme 99b).<sup>209</sup> This reaction involves two diazo and two alkyne components, providing a convenient access to bis trifluoromethylated conjugated trienes **99.6**. Shortly after, the same research group expanded this strategy to allyl sulfones **99.7** as radical acceptors under photocatalytic conditions (Scheme 99c).<sup>210</sup> This transformation features a  $\beta$ -sulfonyl elimination process, leading to the formation of  $\alpha$ -CF<sub>3</sub>-substituted homoallylic iodides **99.8**.


In 2020, Koenigs and co-workers reported a photoexcited proton transfer strategy for O–H functionalization of weakly acidic alcohols **100.2** using diazo compounds **100.1** (Scheme 100).<sup>211</sup> The reaction initiates with the formation of a hydrogen bonding complex **A**. Upon photoirradiation, the photoexcited complex readily deprotonates the mildly acidic substrates. The resulting ion pair undergoes rapid nucleophilic substitution, extruding dinitrogen in a concerted or stepwise

## Scheme 99. Iodine-Enabled Diazo Activation to Access Radical Reactivity


a) Shi &amp; Li 2018




mechanism



b) Wu 2023





c) Wu 2023



manner, leading to O–H functionalized products **100.3**. It was found that various fluorinated alcohols efficiently participated in this reaction **100.4–100.7**. In same year, the authors expanded the scope of the developed photochemical platform to include cyclic diazo amides<sup>212</sup> and phenols.<sup>213</sup> Later on, both the same research group<sup>214,215</sup> and Petit's group<sup>216</sup> conducted computational studies, which suggested a different mechanism: the O–H insertion mechanism in several acidic alcohols includes free carbenes as essential reactive intermediates.


In 2021, Yue, Wei, and co-workers established a photocatalytic method for synthesis of  $\alpha$ -ketoesters **101.2** through aerobic oxidation of  $\alpha$ -diazoesters **101.1** (Scheme 101).<sup>217</sup> The

## Scheme 100. Photoexcited Proton Transfer Reaction

Scheme 101. Synthesis of  $\alpha$ -Ketoesters via Aerobic Oxidation of  $\alpha$ -Diazoesters

proposed mechanism entails the formation of singlet oxygen via an energy transfer process with the photoexcited eosin Y catalyst. Following this, a concerted 1,3-dipolar cycloaddition of  $\alpha$ -diazoesters **101.1** with  $^1\text{O}_2$  generates the dioxadiazole intermediate, which swiftly decomposes to yield the desired product **101.2** while releasing  $\text{N}_2\text{O}$ .


A year later, the groups of Qu and Wang reported a rare example of photocatalytic generation of free carbyne radical intermediates **A** from  $\alpha$ -diazo sulfonium triflates **102.1** (Scheme 102).<sup>218</sup> This three-component coupling reaction between internal alkyne **102.2**,  $\alpha$ -diazo sulfonium triflate **102.1**, and water allows for synthesis of various 1,4-dicarbonyl Z-alkenes **102.3**. The isotope labeling experiments indicated that both the oxygen and hydrogen atoms in the product originated from water. Furthermore, it was discovered that dibenzothiophene-substituted  $\alpha$ -diazo sulfonium salts failed to react. However, the addition of  $\text{SMe}_2$  promoted the reaction, albeit with low efficiency. This observation, coupled with cyclic voltammetry analysis and fluorescent quenching experiments, suggests that the *in situ* formed  $\text{Me}_2\text{S}$  is necessary for the initial single-electron reduction of the photocatalyst, which initiates the catalytic cycle. The reduced photocatalyst is excited once more and undergoes single-electron transfer with  $\alpha$ -diazo sulfonium triflate, which leads to the formation of a radical carbyne intermediate. Addition of the latter to alkyne produces radical intermediate **B**, which is oxidized by the excited photocatalyst to the corresponding cyclopropenium cation **C**. A subsequent capture of the latter by water delivers cyclopropeneol **D**, followed by a  $\text{KH}_2\text{PO}_4$ -assisted ring-

Scheme 102. Generation of Free Carbyne Radical Intermediates from  $\alpha$ -Diazo Sulfonium Triflates

opening process that furnishes the dicarbonyl alkenes with Z-configuration.

Very recently, the groups of Koenigs and Chauhan observed that cyclic diazo imides **103.1** are capable of engaging in two different reaction pathways in the presence of thiols under light-induced conditions (Scheme 103).<sup>219</sup> In acetonitrile solvent, an unusual reduction of the diazo compound to a hydrazone **103.2** was observed, with no carbene intermediate formed (Scheme 103a). However, in dichloromethane, a triplet carbene is generated under 365 nm light irradiation, which then participates in a cascade cyclization/thiolation reaction,

## Scheme 103. Solvent-Dependent Selectivity of Cyclic Diazo Imides



ultimately yielding indane-fused pyrrolidines **103.3** (Scheme 103b).

## 9. PRECURSORS OF DIAZO COMPOUNDS

### 9.1. Diazirines


Diazirines are three-membered cyclic isomers of diazo compounds. These carbene precursors are more stable compared to diazo compounds, without the need for an electron-withdrawing group to ensure their safe handling. However, harsh conditions, such as elevated temperatures or UV photolysis, are required for their activation.<sup>220</sup>

The photochemical transformations of diazirines are well-documented.<sup>220</sup> Moreover, exciting biological applications have been extensively reported, wherein they serve as probes in photoaffinity labeling<sup>221–223</sup> and chemical proteomics.<sup>224</sup> However, these reactions utilize high energy UV light, as diazirines exhibit little to no direct absorption of blue light (410–490 nm).<sup>225</sup> Recently, Fadeyi, Oslund, and MacMillan introduced a microenvironment-mapping ( $\mu$ Map) platform that exploits photocatalytic carbene generation from diazirines to selectively identify protein–protein interactions on cell membranes.<sup>226</sup> Iridium (Ir) photocatalysts facilitate the conversion of nearby diazirines into carbenes through Dexter energy transfer and biotinylate proteins within a  $\sim$ 4 nm radius, allowing for high spatial and temporal control over labeling. Importantly, this technology has been employed to rapidly profile immunological synapse interactions,<sup>226</sup> identify small molecule targets,<sup>227</sup> and track chromatin reorganization events,<sup>228</sup> and map small molecule binding site mapping.<sup>229</sup>

### 9.2. N-Tosylhydrazones

N-Tosylhydrazones are useful synthons in organic synthesis,<sup>230,231</sup> notably via the *in situ* generation of diazo compounds through the Bamford–Stevens reaction (Scheme 104).<sup>232</sup> In turn, N-tosylhydrazones can easily be synthesized


#### Scheme 104. *In Situ* Formation of Diazo Compounds from N-Tosylhydrazones



through the condensation of ketones or aldehydes with  $\text{TsNNH}_2$ . They are generally stable compounds, often crystalline solids that are easy to purify. Particularly, N-tosylhydrazones serve as safe precursors for diazo compounds without electron-withdrawing groups at the diazo carbon, such as alkyl- and aryl-substituted diazomethane derivatives, which are typically unstable and challenging to handle.

In 2020, Koenigs and co-workers achieved the photochemical generation of carbenes from N-tosylhydrazones **105.1** (Scheme 105).<sup>233</sup> Under basic conditions, N-tosylhydrazones readily participated in reactions with indoles **105.2**, yielding either cyclopropanation **105.3** or C–H insertion products **105.4**, depending on the electronic properties of the substrates. Additionally, the authors tested unprotected N-heterocycles **105.5**, including indoles, carbazoles, and piperidine derivatives. In this scenario, N–H insertion products **105.6** were observed. Moreover, the employment of N-tosylhydrazones under photochemical conditions for X–H (X = C, N, O, S)


#### Scheme 105. Photochemical Generation of Carbenes from N-Tosylhydrazones



insertions,<sup>234</sup> [2 + 1] cycloadditions,<sup>235</sup> Doyle–Kirmse and Sommelet–Hauser reactions,<sup>236</sup> was also reported.

N-Sulfonylhydrazones can be applied in denitrogenative sulfonylation reactions. In 2021, Xie, Zeng, and co-workers introduced a photocatalytic method for synthesizing aryl or alkyl sulfones **106.2** (Scheme 106).<sup>237</sup> It was found that aryl or

#### Scheme 106. Denitrogenative Sulfonylation of N-Sulfonylhydrazones




heteroaryl-substituted sulfonylhydrazones **106.1** worked well in this transformation. Similarly, sulfonylhydrazones derived from acetophenone, benzophenone, and benzocycloketone were also proven to be capable substrates (**106.3–106.8**). The reaction mechanism begins with the formation of a diazo compound from N-sulfonylhydrazones, accompanied by desulfonylation. A subsequent denitrogenation generates a carbene intermediate **A**, which is oxidized by the photoexcited  $\text{Ir}^{\text{II}}$  complex, yielding radical cation **B**. The nucleophilic attack of the latter by tosylate results in the formation of radicals **C**, followed by reduction and protonation to give the desired sulfone product.

In 2022, Gevorgyan's group reported an alkyl-Heck reaction of N-tosylhydrazones **107.1** with vinyl arenes **107.2** (Scheme 107).<sup>157</sup> Similarly to the previously mentioned method for

**Scheme 107. Heck Reaction of *N*-Tosylhydrazones via Photoinduced Palladium Catalysis**

Gevorgyan 2022




Heck reaction of diazo compounds by the same group, this reaction proceeds via a hybrid alkyl palladium radical intermediate **A**. *N*-Tosylhydrazones bearing different primary and secondary alkyl chains are capable substrates for this transformation (107.4–107.6). Various vinyl arenes and heteroarenes also reacted well (107.6–107.10).

In the same year, König and co-workers developed a photochemical method for the *in situ* formation of highly reactive alkyl diazo compounds (Scheme 108).<sup>238</sup> This

**Scheme 108. Synthesis of Ketones and Aldehydes Enabled by Photoexcitation of *N*-Tosylhydrazone Anions**

König 2022




approach proceeds via the photoexcitation of *N*-tosylhydrazone anions **A**, enabling the efficient preparation of complex carbonyl compounds **108.2** from the corresponding aldehydes. The UV-vis absorption spectra demonstrate that *N*-tosylhydrazone derived from cyclohexanone absorbs exclusively in the UV region (<340 nm). However, upon the addition of one equivalent of  $\text{Cs}_2\text{CO}_3$ , a notable bathochromic shift of 50 nm

occurs, extending the absorption band into the near-visible light region (390 nm). Thus, upon the deprotonation of *N*-tosylhydrazone, anion **A** undergoes photoexcitation under 385 nm light irradiation. Subsequently, the extrusion of a tosylate delivers the diazo intermediate, which participates in nucleophilic attack on the aldehyde, producing a diazonium alkoxide **B**. Finally, a 1,2-hydride shift of intermediate **B** occurs, which, upon extrusion of dinitrogen, furnishes the product.

In contrast to König's method, which relies on UVA light, the groups of Hong, Zhang, and Zhang recently introduced a visible light-mediated energy transfer approach by using a photosensitizer for the *in situ* formation of donor–donor-type diazo compounds from *N*-tosylhydrazones **109.1** (Scheme 109).<sup>239</sup> Notably, this method enables access to tetrasub-

**Scheme 109. Synthesis of Tetrasubstituted Cyclopropanes Enabled by Energy Transfer**

Hong, Zhang & Zhang 2023




stituted cyclopropanes **109.3** featuring two all-carbon quaternary centers. In terms of the reaction mechanism, the process begins with the deprotonation of *N*-tosylhydrazone by a base, resulting in the formation of the *N*-tosylhydrazone anion **A**. This anion then engages in an energy transfer process with the photoexcited 4CzIPN, affording intermediate in triplet state **B**. Subsequently, the triplet intermediates readily undergo desulfonylation to generate diazo compounds in a triplet state. After the release of dinitrogen, a triplet carbene species **C** is formed, which can then add to alkenes, ultimately yielding the desired cyclopropanes.

Recently, Wang, Das, and Zhang's groups have introduced a photochemical method for the [3 + 2] cycloaddition of alkenes **110.2** and *in situ* formed donor–donor-type diazo compounds from *N*-tosylhydrazones **110.1** (Scheme 110). This method proves highly effective in synthesizing a diverse range of pyrazolines **110.3**. Moreover, by applying modified reaction conditions with additional bases or oxidants, the obtained pyrazolines can be easily transformed into pyrazoles.

**Scheme 110. [3 + 2] Cycloaddition of *in Situ* Formed Diazo Compounds and Alkenes**


Wang, Das & Zhang 2023



### 9.3. Pyridotriazoles

1,2,3-Triazoles can serve as viable precursors for diazo compounds. These heterocyclic compounds are robust, stable, and easily available.<sup>240</sup> Recently, transition metal-catalyzed denitrogenative transformations of pyridotriazoles have emerged as a powerful tool for synthesis of diverse *N*-heterocyclic scaffolds.<sup>241,242</sup> These transformations take advantage of the ring-chain tautomerization of pyridotriazoles in solution into the corresponding diazo tautomer. Although the generation of carbene species from pyridotriazoles under photochemical conditions has been reported, it is important to note that these photolysis approaches involved the use of harmful, high-energy UVC irradiation, leading to many side reactions, thus limiting the synthetic applications of these methods.<sup>243</sup> In 2019, Gevorgyan group observed that pyridotriazoles **111.1** bearing aromatic substitutions at C3 position exhibited good absorption around 390 nm region.<sup>244</sup> This observation inspired the authors to develop light-induced metal-free approach for arylation, X–H insertion, and cyclopropanation reactions of pyridotriazoles (Scheme 111). Under 390 nm light irradiation, the C–C bond forming

**Scheme 111.** Light-Induced and Metal-Free Transformations of Pyridotriazoles



reaction of pyridotriazoles with boronic acids **111.2** proceeded efficiently in the presence of  $K_2CO_3$ , producing a series of triarylmethanes **111.4**–**111.7**. Moreover, 3-aryl pyridotriazoles underwent X–H insertion reactions with alcohols, amines, and carboxylic acids delivering benzylpyridine derivatives without any additives or catalysts. In addition, it was also found that upon photoirradiation, the pyridotriazoles participated in efficient [2 + 1] cyclopropanation with a variety of alkenes. However, this approach, is limited to pyridotriazoles with aryl substituents at C3 position. Other derivatives, which are transparent in that area, such as 3-methyl and ester pyridotriazoles, remained intact and did not show reactivity under these conditions.

### 9.4. 1,3,4-Oxadiazolines

1,3,4-Oxadiazolines are reasonably stable compounds that can be easily prepared, making them valuable substrates for organic synthesis. Depending on the reaction conditions used, these heterocyclic compounds can undergo decomposition to produce various reactive intermediates. When subjected to thermolysis, bis(heteroatom)carbenes can be generated via ylide intermediates.<sup>245</sup> Extensive research by Warkentin and co-workers has demonstrated their utility as dimethoxycarbene precursors in synthesis of structurally diverse heterocycles.<sup>246</sup> Conversely, exposure to UV light leads to the formation of nonstabilized alkyl diazo compounds from these substrates. Notably, although the initial report on photolysis dates to 1968,<sup>247</sup> it was only recently demonstrated by Ley's group that 1,3,4-oxadiazolines efficiently serve as diazo precursors under UV light irradiation. Utilizing this strategy, they accomplished aryl–alkyl cross-coupling reactions of *in situ*-formed nonstabilized diazo compounds with boronic acids via a flow method.<sup>248</sup> Additionally, they showcased the synthesis of various unsymmetrical ketones by adding the *in situ*-formed diazo compound to an aldehyde, followed by a subsequent 1,2-hydride shift.<sup>249</sup>


Very recently, Chaladaj, Gryko, and co-workers developed a visible light-mediated strategy for activating oxadiazolines **112.1** through triplet energy transfer (Scheme 112).<sup>250</sup> This method enables the efficient generation of dialkyl-substituted carbenes, which, upon reacting with Michael acceptors, produce a wide range of spirocyclopropanes **112.3**. Mechanistically, oxadiazolines engage in energy transfer with a photosensitizer, leading to their triplet state. Subsequently, the cleavage of the C–N bond in oxadiazoline produces a diazenyl radical, which decomposes into triplet carbene A and diazoalkane B. In the presence of an alkene, triplet carbene A delivers cyclopropanes in a stepwise manner through a diradical intermediate. Simultaneously, diazoalkane participates in 1,3-dipolar cycloaddition with the alkene, producing pyrazoline C, which upon photosensitization, delivers spirocyclopropane via a diradical intermediate D with the extrusion of dinitrogen. Mechanistic experiments support both carbene and diazoalkane-mediated pathways. However, at this point, there is insufficient evidence to determine which of these pathways predominates.

### 9.5. ( $\beta$ -Amino- $\alpha,\alpha$ -difluoroethyl)phosphonates


In 2021, Röschenharter and Han's groups disclosed the *in situ* formation of (2-diazo-1,1-difluoroethyl)phosphonates **113.1** from corresponding amine precursors by using *tert*-butyl nitrite (Scheme 113).<sup>251</sup> Upon photoirradiation, a wide range of sulfonate esters **113.3** were efficiently synthesized from sulfonic acid. The proposed reaction mechanism involves a

## Scheme 112. Cyclopropanation of Oxadiazolines

Chaladaj &amp; Gryko 2023

Scheme 113. *In Situ* Formation of Diazo Species from ( $\beta$ -Amino- $\alpha,\alpha$ -difluoroethyl)phosphonates

Röschenthaler &amp; Han 2023



nucleophilic substitution of diazonium by sulfonate. It is worth noting that the transformation also proceeded in the absence of blue light, albeit with lower efficiency.

## 10. CONCLUSIONS

Visible light-induced reactions of diazo compounds are a rapidly emerging area in organic synthesis. Despite being a relatively recent field of study, these methods have provided access to a variety of reactive intermediates from diazo compounds, as demonstrated by the wide array of transformations discussed in this review. It is anticipated that this subfield will continue to grow in the future.

Many of the photochemical, metal-free reactions of diazo compounds were performed at room temperature, frequently in air, and without requisite removal of moisture from the solvent. The ability to conduct transformations under such mild conditions enhances selectivity and prevents side reactions commonly found in thermal reactions or those induced by higher-energy UV radiation. Importantly, the reactivity or selectivity of these photochemical reactions can vary significantly from that of metal-catalyzed reactions. In the case of photocatalytic reactions, utilizing a photoredox catalyst enables the generation of radical intermediates from diazo

compounds. This offers new reactivity modes for diazo compounds, which were previously regarded as carbene precursors only.

Although a substantial progress has been made in the field of visible light-induced transformations of diazo compounds, this area is still in its infancy. Furthermore, certain limitations are yet to be overcome. Within the context of photochemical reactions, a majority of transformations presented herein employ donor–acceptor or donor–donor type diazo compounds that have good absorption in the visible light region. Thus, one of the remaining challenges is to engage diazo compounds that are transparent in this region, such as acceptor types. Moreover, although several enantioselective methods have been reported by employing chiral catalyst or ligand, achieving stereoselective transformations in photochemical reactions remains majorly underexplored. Additionally, many of the reactions require the excess of one of the reacting partners. Besides these limitations, in-depth mechanistic investigations are essential to provide a more comprehensive understanding of the reaction mechanisms. Thus, it is expected that development toward asymmetric transformations and the engagement of novel diazo substrates or precursors will constitute an important future direction in this field.

We anticipated that visible light-induced reactions of diazo compounds will become a versatile strategy with broad applications in organic synthesis, pharmaceutical chemistry, and material science.

## AUTHOR INFORMATION

## Corresponding Author

Vladimir Gevorgyan – Department of Chemistry and Biochemistry, The University of Texas at Dallas, Richardson, Texas 75080-3021, United States; [orcid.org/0000-0002-7836-7596](https://orcid.org/0000-0002-7836-7596); Email: [vlad@utdallas.edu](mailto:vlad@utdallas.edu)

## Author

Ziyan Zhang – Department of Chemistry and Biochemistry, The University of Texas at Dallas, Richardson, Texas 75080-3021, United States

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.chemrev.3c00869>

## Notes

The authors declare no competing financial interest.

## Biographies

Ziyan Zhang received her B.Sc. from Beijing Technology and Business University. In 2017, she joined the Gevorgyan group at the University of Illinois at Chicago as a Ph.D. student and later moved to the University of Texas at Dallas. Her research interests are focused on the development of novel photoinduced methodologies.

Vladimir Gevorgyan received his Ph.D. from the Latvian Institute of Organic Synthesis. After two years of postdoctoral research (1992–1994, JSPS- and CibaGeigy International Postdoctoral Fellowships) at Tohoku University, Japan, and a visiting professorship (1995) at CNR, Bologna, Italy, he joined the faculty at Tohoku University (Assistant Professor, 1996; Associate Professor, 1997–1999). In 1999, Vladimir Gevorgyan moved to USA to join UIC (Associate Professor, 1999; Professor, 2003; LAS Distinguished Professor, 2012). In 2019, he joined the University of Texas at Dallas to become a Robert A. Welch Distinguished Chair in Chemistry. Vladimir also holds a Professor position at the University of Texas Southwestern

Medical Center. His group is interested in the development of novel synthetic methodology, particularly toward biologically relevant molecules.

## ACKNOWLEDGMENTS

We thank the National Institutes of Health (GM120281), National Science Foundation (CHE-2246075), and Welch Foundation (Chair, AT-0041) for financial support.

## REFERENCES

- (1) Ye, T.; McKervey, M. A. Organic Synthesis with  $\alpha$ -Diazo Carbonyl Compounds. *Chem. Rev.* **1994**, *94*, 1091–1160.
- (2) (a) Davies, H. M. L.; Morton, D. Guiding Principles for Site Selective and Stereoselective Intermolecular C–H Functionalization by Donor/acceptor Rhodium Carbene. *Chem. Soc. Rev.* **2011**, *40*, 1857–1869. (b) Davies, H. M. L.; Manning, J. R. Catalytic C–H Functionalization by Metal Carbenoid and Nitrenoid Insertion. *Nature* **2008**, *451*, 417–424. (c) Gillingham, D.; Fei, N. Catalytic X–H Insertion Reactions Based on Carbenoids. *Chem. Soc. Rev.* **2013**, *42*, 4918–4931. (d) Ford, A.; Miel, H.; Ring, A.; Slattery, C. N.; Maguire, A. R.; McKervey, M. A. Modern Organic Synthesis with  $\alpha$ -Diazocarbonyl Compounds. *Chem. Rev.* **2015**, *115*, 9981–10080. (e) Candeias, N. R.; Paterna, R.; Gois, P. M. P. Homologation Reaction of Ketones with Diazo Compounds. *Chem. Rev.* **2016**, *116*, 2937–2981. (f) Cheng, Q.-Q.; Deng, Y.; Lankelma, M.; Doyle, M. P. Cycloaddition Reactions of Enoldiazo Compounds. *Chem. Soc. Rev.* **2017**, *46*, 5425–5443. (g) Davies, H. M. L.; Denton, J. R. Application of Donor/acceptor-Carbenoids to the Synthesis of Natural Products. *Chem. Soc. Rev.* **2009**, *38*, 3061–3071.
- (3) Candeias, N. R.; Afonso, C. A. M. Developments in the Photochemistry of Diazo Compounds. *Curr. Org. Chem.* **2009**, *13*, 763–787.
- (4) Galkina, O. S.; Rodina, L. L. Photochemical Transformations of Diazocarbonyl Compounds: Expected and Novel Reactions. *Russ. Chem. Rev.* **2016**, *85*, 537.
- (5) Sander, W.; Bucher, G.; Wierlacher, S. Carbenes in Matrixes: Spectroscopy, Structure, and Reactivity. *Chem. Rev.* **1993**, *93*, 1583–1621.
- (6) Wang, J.-L.; Likhotvorik, I.; Platz, M. S. A Laser Flash Photolysis Study of 2-Naphthyl(carbomethoxy)carbene. *J. Am. Chem. Soc.* **1999**, *121*, 2883–2890.
- (7) Wang, J.; Burdzinski, G.; Gustafson, T. L.; Platz, M. S. Ultrafast Study of p-Biphenylyldiazoethane. The Chemistry of the Diazo Excited State and the Relaxed Carbene. *J. Am. Chem. Soc.* **2007**, *129*, 2597–2606.
- (8) Zhang, Y.; Kubicki, J.; Wang, J.; Platz, M. S. 2-Naphthyl(carbomethoxy)carbene Revisited: Combination of Ultrafast UV–vis and Infrared Spectroscopic Study. *J. Phys. Chem. A* **2008**, *112*, 11093–11098.
- (9) Zhang, Y.; Kubicki, J.; Platz, M. S. Ultrafast UV–Visible and Infrared Spectroscopic Observation of a Singlet Vinylcarbene and the Intramolecular Cyclopropanation Reaction. *J. Am. Chem. Soc.* **2009**, *131*, 13602–13603.
- (10) Wang, J.; Kubicki, J.; Gustafson, T. L.; Platz, M. S. The Dynamics of Carbene Solvation: An Ultrafast Study of p-Biphenylyl trifluoromethylcarbene. *J. Am. Chem. Soc.* **2008**, *130*, 2304–2313.
- (11) (a) Ciszewski, L. W.; Rybicka-Jasińska, K.; Gryko, D. Recent Developments in Photochemical Reactions of Diazo Compounds. *Org. Biomol. Chem.* **2019**, *17*, 432–448. (b) Yang, Z.; Stivannin, M. L.; Jurberg, I. D.; Koenigs, R. M. Visible Light-Promoted Reactions with Diazo Compounds: a Mild and Practical Strategy towards Free Carbene Intermediates. *Chem. Soc. Rev.* **2020**, *49*, 6833–6847. (c) Durka, J.; Turkowska, J.; Gryko, D. Lightening Diazo Compounds? *ACS Sustain. Chem. Eng.* **2021**, *9*, 8895–8918. (d) Empel, C.; Pei, C.; Koenigs, R. M. Unlocking Novel Reaction Pathways of Diazoalkanes with Visible Light. *Chem. Commun.* **2022**, *58*, 2788–2798. (e) Gallo, R. D. C.; Cariello, G.; Goulart, T. A. C.; Jurberg, I. D. Visible Light-Mediated Photolysis of Organic Molecules: the Case Study of Diazo Compounds. *Chem. Commun.* **2023**, *59*, 7346–7360.
- (12) Ovalles, S. R.; Hansen, J. H.; Davies, H. M. L. Thermally Induced Cycloadditions of Donor/Acceptor Carbenes. *Org. Lett.* **2011**, *13*, 4284–4287.
- (13) Hansen, S. R.; Spangler, J. E.; Hansen, J. H.; Davies, H. M. L. Metal-Free N–H Insertions of Donor/Acceptor Carbenes. *Org. Lett.* **2012**, *14*, 4626–4629.
- (14) Alford, J. S.; Davies, H. M. L. Expanding the Scope of Donor/Acceptor Carbenes to N-Phthalimido Donor Groups: Diastereoselective Synthesis of 1-Cyclopropane  $\alpha$ -Amino Acids. *Org. Lett.* **2012**, *14*, 6020–6023.
- (15) Tortoreto, C.; Rackl, D.; Davies, H. M. L. Metal-Free C–H Functionalization of Alkanes by Aryldiazoacetates. *Org. Lett.* **2017**, *19*, 770–773.
- (16) Jurberg, I. D.; Davies, H. M. L. Blue Light-Promoted Photolysis of Aryldiazoacetates. *Chem. Sci.* **2018**, *9*, 5112–5118.
- (17) Geise, C. M.; Wang, Y.; Mykhaylova, O.; Frink, B. T.; Toscano, J. P.; Hadad, C. M. Computational and Experimental Studies of the Effect of Substituents on the Singlet–Triplet Energy Gap in Phenyl(carbomethoxy)carbene. *J. Org. Chem.* **2002**, *67*, 3079–3088.
- (18) Romero, N. A.; Nicewicz, D. A. Organic Photoredox Catalysis. *Chem. Rev.* **2016**, *116*, 10075–10166.
- (19) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. Photoredox Catalysis in Organic Chemistry. *J. Org. Chem.* **2016**, *81*, 6898–6926.
- (20) Chan, A. Y.; Perry, I. B.; Bissonnette, N. B.; Buksh, B. F.; Edwards, G. A.; Frye, L. I.; Garry, O. L.; Lavagnino, M. N.; Li, B. X.; Liang, Y.; Mao, E.; Millet, A.; Oakley, J. V.; Reed, N. L.; Sakai, H. A.; Seath, C. P.; MacMillan, D. W. C. Metallaphotoredox: The Merger of Photoredox and Transition Metal Catalysis. *Chem. Rev.* **2022**, *122*, 1485–1542.
- (21) Kirmse, W. 100 Years of the Wolff Rearrangement. *Eur. J. Org. Chem.* **2002**, *2002*, 2193–2256.
- (22) Allen, A. D.; Tidwell, T. T. Ketenes and Other Cumulenes as Reactive Intermediates. *Chem. Rev.* **2013**, *113*, 7287–7342.
- (23) (a) Chang, S. J.; Ravi Shankar, B. K.; Shechter, H. Chemistry of Diazoacenaphthenones and Diazoacenaphthenes. *J. Org. Chem.* **1982**, *47*, 4226–4234. (b) Trost, B. M.; Kinson, P. L. On the Applicability of the Wolff Rearrangement to Strained Hydrocarbons. *Tetrahedron Lett.* **1973**, *14*, 2675–2678. (c) Trost, B. M.; Kinson, P. L. Perturbed [12]annulenes. Derivatives of Dibenzo[cd,gh]pentalene. *J. Am. Chem. Soc.* **1975**, *97*, 2438–2449.
- (24) Vaske, Y. S. M.; Mahoney, M. E.; Konopelski, J. P.; Rogow, D. L.; McDonald, W. J. Enantiomerically Pure trans- $\beta$ -Lactams from  $\alpha$ -Amino Acids via Compact Fluorescent Light (CFL) Continuous-Flow Photolysis. *J. Am. Chem. Soc.* **2010**, *132*, 11379–11385.
- (25) Pan, J.-B.; Yang, Z.-C.; Zhang, X.-G.; Li, M.-L.; Zhou, Q.-L. Enantioselective Synthesis of Chiral Amides by a Phosphoric Acid Catalyzed Asymmetric Wolff Rearrangement\*\*. *Angew. Chem., Int. Ed.* **2023**, *62*, No. e202308122.
- (26) Bernardim, B.; Hardman-Baldwin, A. M.; Burtoloso, A. C. B. LED lighting as a simple, inexpensive, and sustainable alternative for Wolff rearrangements. *RSC Adv.* **2015**, *5*, 13311–13314.
- (27) Meng, J.; Ding, W.-W.; Han, Z.-Y. Synthesis of Chiral Esters via Asymmetric Wolff Rearrangement Reaction. *Org. Lett.* **2019**, *21*, 9801–9805.
- (28) Liu, D.; Ding, W.; Zhou, Q.-Q.; Wei, Y.; Lu, L.-Q.; Xiao, W.-J. Catalyst-Controlled Regioselective Acylation of  $\beta$ -Ketoesters with  $\alpha$ -Diazo Ketones Induced by Visible Light. *Org. Lett.* **2018**, *20*, 7278–7282.
- (29) Capurro, P.; Lambruschini, C.; Lova, P.; Moni, L.; Basso, A. Into the Blue: Ketene Multicomponent Reactions under Visible Light. *J. Org. Chem.* **2021**, *86*, 5845–5851.
- (30) Echemendia, R.; de Oliveira, K. T.; Burtoloso, A. C. B. Visible-Light-Promoted Synthesis of 1,3-Dicarbonyl Sulfoxonium Ylides. *Org. Lett.* **2022**, *24*, 6386–6390.
- (31) Dong, Y.; Tian, Y.; Zhang, Z.; Wang, T. Blue Light-Promoted Reaction of  $\alpha$ -Diazoketones and Sulfoxonium Ylides: Synthesis of 1,3-

Dicarbonyl Sulfoxonium Ylides. *Adv. Synth. Catal.* **2022**, *364*, 4026–4030.

(32) Li, M.-M.; Wei, Y.; Liu, J.; Chen, H.-W.; Lu, L.-Q.; Xiao, W.-J. Sequential Visible-Light Photoactivation and Palladium Catalysis Enabling Enantioselective [4 + 2] Cycloadditions. *J. Am. Chem. Soc.* **2017**, *139*, 14707–14713.

(33) Wei, Y.; Liu, S.; Li, M.-M.; Li, Y.; Lan, Y.; Lu, L.-Q.; Xiao, W.-J. Enantioselective Trapping of Pd-Containing 1,5-Dipoles by Photo-generated Ketenes: Access to 7-Membered Lactones Bearing Chiral Quaternary Stereocenters. *J. Am. Chem. Soc.* **2019**, *141*, 133–137.

(34) Liu, J.; Li, M.-M.; Qu, B.-L.; Lu, L.-Q.; Xiao, W.-J. A photoinduced Wolff rearrangement/Pd-catalyzed [3 + 2] cyclo-addition sequence: an unexpected route to tetrahydrofurans. *Chem. Commun.* **2019**, *55*, 2031–2034.

(35) Fan, T.; Zhang, Z.-J.; Zhang, Y.-C.; Song, J. Construction of All-Carbon Quaternary Stereocenters via Sequential Photoactivation/Isothiourea Catalysis. *Org. Lett.* **2019**, *21*, 7897–7901.

(36) Wang, C.; Wang, Z.; Yang, J.; Shi, S.-H.; Hui, X.-P. Sequential Visible-Light and N-Heterocyclic Carbene Catalysis: Stereoselective Synthesis of Tetrahydropyrano[2,3-b]indoles. *Org. Lett.* **2020**, *22*, 4440–4443.

(37) Munaretto, L. S.; dos Santos, C. Y.; Gallo, R. D. C.; Okada, C. Y., Jr.; Deflon, V. M.; Jurberg, I. D. Visible-Light-Mediated Strategies to Assemble Alkyl 2-Carboxylate-2,3,3-Trisubstituted  $\beta$ -Lactams and S-Alkoxy-2,2,4-Trisubstituted Furan-3(2H)-ones Using Aryldiazoacetates and Aryldiazoketones. *Org. Lett.* **2021**, *23*, 9292–9296.

(38) Ebner, C.; Carreira, E. M. Cyclopropanation Strategies in Recent Total Syntheses. *Chem. Rev.* **2017**, *117*, 11651–11679.

(39) Qian, D.; Zhang, J. Gold-Catalyzed Cyclopropanation Reactions using a Carbenoid Precursor Toolbox. *Chem. Soc. Rev.* **2015**, *44*, 677–698.

(40) Tang, P.; Qin, Y. Recent Applications of Cyclopropane-Based Strategies to Natural Product Synthesis. *Synthesis* **2012**, *44*, 2969–2984.

(41) Chen, D. Y. K.; Pouwer, R. H.; Richard, J.-A. Recent Advances in the Total Synthesis of Cyclopropane-Containing Natural Products. *Chem. Soc. Rev.* **2012**, *41*, 4631–4642.

(42) Ghosh, K.; Das, S. Recent Advances in Ring-Opening of Donor Acceptor Cyclopropanes using C-Nucleophiles. *Org. Biomol. Chem.* **2021**, *19*, 965–982.

(43) Pirenne, V.; Muriel, B.; Waser, J. Catalytic Enantioselective Ring-Opening Reactions of Cyclopropanes. *Chem. Rev.* **2021**, *121*, 227–263.

(44) Chen, Z.-L.; Xie, Y.; Xuan, J. Visible Light-Mediated Cyclopropanation: Recent Progress. *Eur. J. Org. Chem.* **2022**, *2022*, No. e202201066.

(45) Zhang, X.; Du, C.; Zhang, H.; Li, X.-C.; Wang, Y.-L.; Niu, J.-L.; Song, M.-P. Metal-Free Blue-Light-Mediated Cyclopropanation of Indoles by Aryl(diazo)acetates. *Synthesis* **2019**, *51*, 889–898.

(46) Klöpfer, V.; Eckl, R.; Floß, J.; Roth, P. M. C.; Reiser, O.; Barham, J. P. Catalyst-free, scalable heterocyclic flow photocyclopropanation. *Green Chem.* **2021**, *23*, 6366–6372.

(47) Guo, Y.; Nguyen, T. V.; Koenigs, R. M. Norcaradiene Synthesis via Visible-Light-Mediated Cyclopropanation Reactions of Arenes. *Org. Lett.* **2019**, *21*, 8814–8818.

(48) Zhao, S.; Chen, X.-X.; Gao, N.; Qian, M.; Chen, X. Visible-Light-Mediated Cyclopropanation Reactions of 3-Diazoxyindoles with Arenes. *J. Org. Chem.* **2021**, *86*, 7131–7140.

(49) Guo, Y.; Empel, C.; Pei, C.; Atodiresei, I.; Fallon, T.; Koenigs, R. M. Photochemical Cyclopropanation of Cyclooctatetraene and (Poly-)unsaturated Carbocycles. *Org. Lett.* **2020**, *22*, 5126–5130.

(50) Chen, Z.-L.; Empel, C.; Wang, K.; Wu, P.-P.; Cai, B.-G.; Li, L.; Koenigs, R. M.; Xuan, J. Enabling Cyclopropanation Reactions of Imidazole Heterocycles via Chemoselective Photochemical Carbene Transfer Reactions of NHC-Boranes. *Org. Lett.* **2022**, *24*, 2232–2237.

(51) Chauhan, J.; Ravva, M. K.; Gremaud, L.; Sen, S. Blue LED Mediated Intramolecular C–H Functionalization and Cyclopropanation of Tryptamines: Synthesis of Azepino[4, 5-b]indoles and Natural Product Inspired Polycyclic Indoles. *Org. Lett.* **2020**, *22*, 4537–4541.

(52) Guha, S.; Gadde, S.; Kumar, N.; Black, D. S.; Sen, S. Orthogonal Syntheses of  $\gamma$ -Carbolinone and Spiro[pyrrolidinone-3,3']indole Derivatives in One Pot through Reaction Telescopin. *J. Org. Chem.* **2021**, *86*, 5234–5244.

(53) Pei, C.; Empel, C.; Koenigs, R. M. Photochemical Intermolecular Cyclopropanation Reactions of Allylic Alcohols for the Synthesis of [3.1.0]-Bicyclohexanes. *Org. Lett.* **2023**, *25*, 169–173.

(54) Hommelsheim, R.; Guo, Y.; Yang, Z.; Empel, C.; Koenigs, R. M. Blue-Light-Induced Carbene-Transfer Reactions of Diazoalkanes. *Angew. Chem., Int. Ed.* **2019**, *58*, 1203–1207.

(55) He, F.; Koenigs, R. M. Visible light mediated, metal-free carbene transfer reactions of diazoalkanes with propargylic alcohols. *Chem. Commun.* **2019**, *55*, 4881–4884.

(56) Li, Z.; Boyarskikh, V.; Hansen, J. H.; Autschbach, J.; Musaev, D. G.; Davies, H. M. L. Scope and Mechanistic Analysis of the Enantioselective Synthesis of Allenes by Rhodium-Catalyzed Tandem Ylide Formation/[2,3]-Sigmatropic Rearrangement between Donor/Acceptor Carbeneoids and Propargylic Alcohols. *J. Am. Chem. Soc.* **2012**, *134*, 15497–15504.

(57) Wang, J.; Yao, X.; Wang, T.; Han, J.; Zhang, J.; Zhang, X.; Wang, P.; Zhang, Z. Synthesis of 2,5-Dihydrofurans via a Gold(I)-Catalyzed Formal [4 + 1] Cycloaddition of  $\alpha$ -Diazoesters and Propargyl Alcohols. *Org. Lett.* **2015**, *17*, 5124–5127.

(58) Jana, S.; Pei, C.; Empel, C.; Koenigs, R. M. Photochemical Carbene Transfer Reactions of Aryl/Aryl Diazoalkanes—Experiment and Theory\*\*. *Angew. Chem., Int. Ed.* **2021**, *60*, 13271–13279.

(59) Zeng, T.-T.; Zhang, H.-Y.; Xie, Z.-B.; Dong, Y.-Y.; Gong, S.-S.; Sun, Q. Facile and Efficient Synthesis of Fluorene and Indeno[1,2-*e*]Carboxylates from Biaryldiazoacetates via BlueLight-promoted Intramolecular CarbeneC–H Insertion. *Asian J. Org. Chem.* **2023**, *12*, No. e202200638.

(60) Samim Mondal, A. R.; Ghorai, B.; Hari, D. P. Photoinduced Temperature-Regulated Selective Carbene C–H Insertion for the Synthesis of Functionalized Spiro- $\beta$ -lactones and -lactams. *Org. Lett.* **2023**, *25*, 4974–4979.

(61) Montesinos-Magraner, M.; Costantini, M.; Ramírez-Contreras, R.; Muratore, M. E.; Johansson, M. J.; Mendoza, A. General Cyclopropane Assembly by Enantioselective Transfer of a Redox-Active Carbene to Aliphatic Olefins. *Angew. Chem., Int. Ed.* **2019**, *58*, 5930–5935.

(62) Yu, Z.; Mendoza, A. Enantioselective Assembly of Congested Cyclopropanes using Redox-Active Aryldiazoacetates. *ACS Catal.* **2019**, *9*, 7870–7875.

(63) Costantini, M.; Mendoza, A. Modular Enantioselective Synthesis of cis-Cyclopropanes through Self-Sensitized Stereoselective Photodecarboxylation with Benzothiazolines. *ACS Catal.* **2021**, *11*, 13312–13319.

(64) Santiago, J. V.; Orlowska, K.; Ociepa, M.; Gryko, D. Aryl versus Alkyl Redox-Active Diazoacetates — Light-Induced C–H Insertion or 1,2-Rearrangement. *Org. Lett.* **2023**, *25*, 6267–6271.

(65) Zhu, S.-F.; Zhou, Q.-L. Transition-Metal-Catalyzed Enantioselective Heteroatom–Hydrogen Bond Insertion Reactions. *Acc. Chem. Res.* **2012**, *45*, 1365–1377.

(66) Bergstrom, B. D.; Nickerson, L. A.; Shaw, J. T.; Souza, L. W. Transition Metal Catalyzed Insertion Reactions with Donor/Donor Carbenes. *Angew. Chem., Int. Ed.* **2021**, *60*, 6864–6878.

(67) Toscano, J. P.; Platz, M. S.; Nikolaev, V.; Popic, V. Carboethoxycarbene. A Laser Flash Photolysis Study. *J. Am. Chem. Soc.* **1994**, *116*, 8146–8151.

(68) Candeias, N. R.; Afonso, C. A. M. Developments in the Photochemistry of Diazo Compounds. *Curr. Org. Chem.* **2009**, *13*, 763–787.

(69) Empel, C.; Patureau, F. W.; Koenigs, R. M. Visible Light Induced Metal-Free Carbene N-Carbazolation. *J. Org. Chem.* **2019**, *84*, 11316–11322.

(70) Stivanin, M. L.; Fernandes, A. A. G.; da Silva, A. F.; Okada, C. Y., Jr.; Jurberg, I. D. Blue Light-Promoted N–H Insertion of Carbazoles, Pyrazoles and 1,2,3-Triazoles into Aryldiazoacetates. *Adv. Synth. Catal.* **2020**, *362*, 1106–1111.

(71) Maiti, D.; Das, R.; Sen, S. Blue LED-Mediated N–H Insertion of Indoles into Aryldiazoesters at Room Temperature in Batch and Flow: Reaction Kinetics, Density Functional Theory, and Mechanistic Study. *J. Org. Chem.* **2021**, *86*, 2522–2533.

(72) Okada, C. Y.; dos Santos, C. Y.; Jurberg, I. D. Blue Light-Promoted N–H Insertion of Amides, Isatins, Sulfonamides and Imides into Aryldiazoacetates: Synthesis of Unnatural  $\alpha$ -Aryl Amino Acid Derivatives. *Tetrahedron* **2020**, *76*, 131316.

(73) Guo, W.; Zhou, Y.; Xie, H.; Yue, X.; Jiang, F.; Huang, H.; Han, Z.; Sun, J. Visible-Light-Induced Organocatalytic Enantioselective N–H Insertion of  $\alpha$ -Diazoesters Enabled by Indirect Free Carbene Capture. *Chem. Sci.* **2023**, *14*, 843–848.

(74) Pan, J.-B.; Zhang, X.-G.; Shi, Y.-F.; Han, A.-C.; Chen, Y.-J.; Ouyang, J.; Li, M.-L.; Zhou, Q.-L. A Spiro Phosphamide Catalyzed Enantioselective Proton Transfer of Ylides in a Free Carbene Insertion into N–H Bonds. *Angew. Chem., Int. Ed.* **2023**, *62*, No. e202300691.

(75) Zhou, Y.; Zhang, Y.; Yu, C.; Yue, X.; Jiang, F.; Guo, W. Visible-light-mediated catalytic asymmetric synthesis of  $\alpha$ -amino esters via free carbene insertion into NH bond. *Tetrahedron Lett.* **2023**, *122*, 154496.

(76) Titov, G. D.; Antonychev, G. I.; Novikov, M. S.; Khlebnikov, A. F.; Rogacheva, E. V.; Kraeva, L. A.; Rostovskii, N. V. Gold vs Light: Chemodivergent Reactivity of Diazoesters toward 2H-Azirine-2-carboxylic Acids. *Org. Lett.* **2023**, *25*, 2707–2712.

(77) Bai, J.; Qi, D.; Song, Z.; Li, B.; Guo, L.; Yang, C.; Xia, W. Synthesis of  $\alpha$ -Hydroxy and  $\alpha$ -Alkoxy Esters Enabled by a Visible-Light-Induced O–H Insertion Reaction of Diazo Compounds. *Synlett* **2022**, *33*, 2048–2052.

(78) Li, Q.; Cai, B.-G.; Li, L.; Xuan, J. Oxime Ether Synthesis through O–H Functionalization of Oximes with Diazo Esters under Blue LED Irradiation. *Org. Lett.* **2021**, *23*, 6951–6955.

(79) Qi, Z.; Wang, S. Chemodivergent Synthesis of Oxazoles and Oxime Ethers Initiated by Selective C–N/C–O Formation of Oximes and Diazo Esters. *Org. Lett.* **2021**, *23*, 8549–8553.

(80) Stivanin, M. L.; Duarte, M.; Leão, L. P. M. O.; Saito, F. A.; Jurberg, I. D. Visible-Light-Mediated Strategies for the Preparation of Oxime Ethers Derived from O–H Insertions of Oximes into Aryldiazoacetates. *J. Org. Chem.* **2021**, *86*, 17528–17532.

(81) Cai, B.-G.; Li, Q.; Zhang, Q.; Li, L.; Xuan, J. Synthesis of Trisubstituted Hydroxylamines by a Visible Light-Promoted Multi-component Reaction. *Org. Chem. Front.* **2021**, *8*, 5982–5987.

(82) Yang, J.; Wang, G.; Zhou, H.; Li, Z.; Ma, B.; Song, M.; Sun, R.; Huo, C. Visible-Light-Promoted Selective O-Alkylation of 2-Pyridones with  $\alpha$ -Aryldiazoacetates. *Org. Biomol. Chem.* **2021**, *19*, 394–398.

(83) He, F.; Li, F.; Koenigs, R. M. Metal-Free Insertion Reactions of Silanes with Aryldiazoacetates. *J. Org. Chem.* **2020**, *85*, 1240–1246.

(84) Qin, L.-Z.; Yuan, X.; Cui, Y.-S.; Sun, Q.; Duan, X.; Zhuang, K.-Q.; Chen, L.; Qiu, J.-K.; Guo, K. Visible-Light-Mediated S–H Bond Insertion Reactions of Diazoalkanes with Cysteine Residues in Batch and Flow. *Adv. Synth. Catal.* **2020**, *362*, 5093–5104.

(85) Yang, J.; Wang, G.; Chen, S.; Ma, B.; Zhou, H.; Song, M.; Liu, C.; Huo, C. Catalyst-Free, Visible-Light-Promoted S–H Insertion Reaction between Thiols and  $\alpha$ -Diazoesters. *Org. Biomol. Chem.* **2020**, *18*, 9494–9498.

(86) Jana, S.; Guo, Y.; Koenigs, R. M. Recent Perspectives on Rearrangement Reactions of Ylides via Carbene Transfer Reactions. *Chem.—Eur. J.* **2021**, *27*, 1270–1281.

(87) Li, F.; He, F.; Koenigs, R. M. Catalyst-Free [2,3]-Sigmatropic Rearrangement Reactions of Photochemically Generated Ammonium Ylides. *Synthesis* **2019**, *51*, 4348–4358.

(88) Chen, J.; Liu, S.; Lv, X.; Hong, K.; Lei, J.; Xu, X.; Hu, W. Blue Light-Promoted Formal [4 + 1]-Annulation of Diazoacetates with  $\alpha$ -Aminoacetophenones: Synthesis of Polysubstituted Indolines and Computational Study. *J. Org. Chem.* **2020**, *85*, 13920–13928.

(89) Cheng, X.; Cai, B.-G.; Mao, H.; Lu, J.; Li, L.; Wang, K.; Xuan, J. Divergent Synthesis of Aziridine and Imidazolidine Frameworks under Blue LED Irradiation. *Org. Lett.* **2021**, *23*, 4109–4114.

(90) Cai, B.-G.; Luo, S.-S.; Li, L.; Li, L.; Xuan, J.; Xiao, W.-J. Visible Light-Promoted Amide Bond Formation via One-Pot Nitrone in Situ Formation/Rearrangement Cascade. *CCS Chemistry* **2021**, *3*, 2764–2771.

(91) Zhu, K.; Cao, M.; Zhao, G.; Zhao, J.; Li, P. Visible Light-Promoted Diazoacetates and Nitriles Generating Nitrilium Ions Trapped by Benzotriazoles and Carboxylic Acids. *Org. Lett.* **2022**, *24*, 5855–5859.

(92) Cai, B.-G.; Yao, W.-Z.; Li, L.; Xuan, J. Visible-Light-Induced Imide Synthesis through a Nitrile Ylide Formation/Trapping Cascade. *Org. Lett.* **2022**, *24*, 6647–6652.

(93) Mailloux, M. J.; Fleming, G. S.; Kumta, S. S.; Beeler, A. B. Unified Synthesis of Azepines by Visible-Light-Mediated Dearomatic Ring Expansion of Aromatic N-Ylides. *Org. Lett.* **2021**, *23*, 525–529.

(94) Maiti, D.; Das, R.; Prabakar, T.; Sen, S. Blue LED Induced Solvent-Free Multicomponent Reactions Among Aryl Diazoacetates, Pyridine Derivatives and Maleimides: Direct Eco-Friendly Synthesis of Densely Functionalized Itaconimides. *Green Chem.* **2022**, *24*, 3001–3008.

(95) Jana, S.; Yang, Z.; Pei, C.; Xu, X.; Koenigs, R. M. Photochemical ring expansion reactions: synthesis of tetrahydrofuran derivatives and mechanism studies. *Chem. Sci.* **2019**, *10*, 10129–10134.

(96) Maiti, D.; Das, R.; Sen, S. Photolytic Amino Etherification Reactions of Aryl Diazoacetates with N-Heterocycles and a Stoichiometric Amount of Dioxane/tetrahydropyran in Aqueous Medium: Synthesis of 1,4-Dioxepane/1,4,7-dioxazan-6-one Systems. *Green Chem.* **2021**, *23*, 8533–8544.

(97) Qi, D.; Bai, J.; Zhang, H.; Li, B.; Song, Z.; Ma, N.; Guo, L.; Song, L.; Xia, W. Photoinduced Synthesis of Functionalized Oxetanes via Diradical-mediated Ring Contraction. *Green Chem.* **2022**, *24*, 5046–5051.

(98) He, F.; Pei, C.; Koenigs, R. M. Photochemical fluoro-amino etherification reactions of aryldiazoacetates with NFSI under stoichiometric conditions. *Chem. Commun.* **2020**, *56*, 599–602.

(99) Cheng, R.; Qi, C.; Wang, L.; Xiong, W.; Liu, H.; Jiang, H. Visible Light-Promoted Synthesis of Organic Carbamates from Carbon Dioxide under Catalyst- and Additive-Free Conditions. *Green Chem.* **2020**, *22*, 4890–4895.

(100) Lv, Y.; Liu, R.; Ding, H.; Wei, W.; Zhao, X.; He, L. Metal-free Visible-light-induced Multi-component Reactions of  $\alpha$ -Diazoesters Leading to S-Alkyl Dithiocarbamates. *Org. Chem. Front.* **2022**, *9*, 3486–3492.

(101) Stivanin, M. L.; Gallo, R. D. C.; Spadeto, J. P. M.; Cormanich, R. A.; Jurberg, I. D. A Visible Light-mediated Three-component Strategy Based on the Ring-opening of Cyclic Ethers with Aryldiazoacetates and Nucleophiles. *Org. Chem. Front.* **2022**, *9*, 1321–1326.

(102) Qu, C.; Liu, R.; Wang, Z.; Lv, Y.; Yue, H.; Wei, W. Visible-light-driven Multicomponent Reactions to Access S-Alkyl Phosphorothioates Using Elemental Sulfur as the Sulfur Source. *Green Chem.* **2022**, *24*, 4915–4920.

(103) Qu, C.; Hao, J.; Ding, H.; Lv, Y.; Zhao, X.-E.; Zhao, X.; Wei, W. Visible-Light-Initiated Multicomponent Reactions of  $\alpha$ -Diazoesters to Access Organophosphorus Compounds. *J. Org. Chem.* **2022**, *87*, 12921–12931.

(104) Wang, Z.; Hao, J.; Lv, Y.; Qu, C.; Yue, H.; Wei, W. Additive-Free Visible-Light-Initiated Three-Component Cyanation and Azidation. *Asian J. Org. Chem.* **2022**, *11*, No. e202200484.

(105) Ding, H.; Wang, Z.; Qu, C.; Lv, Y.; Zhao, X.; Wei, W. Visible-light-mediated Multi-component Carbene Transfer Reactions of  $\alpha$ -Diazoesters to Construct Multisubstituted Pyrazoles and 1,3-Dicarbonyl Derivatives. *Org. Chem. Front.* **2022**, *9*, 5530–5535.

(106) Xia, S.; Jian, Y.; Zhang, L.; Zhang, C.; An, Y.; Wang, Y. Visible-light-promoted N–H Functionalization of O-Substituted Hydroxamic Acid with Diazo Esters. *RSC Adv.* **2023**, *13*, 14501–14505.

(107) Cai, B.-G.; Li, Q.; Empel, C.; Li, L.; Koenigs, R. M.; Xuan, J. Dark and Light Reactions of Carbenes—Merging Carbene Transfer

Reactions with N-Heterocyclic Carbene Catalysis for the Synthesis of Hydroxamic Acid Esters. *ACS Catal.* **2022**, *12*, 11129–11136.

(108) Zhu, K.; Zhou, X.; Ren, Y.; Dong, L.; Zhao, G.; Zhao, J.; Li, P. Visible Light-Induced Carbene Reactivity of Acceptor Diazoalkanes: Deconstructive Difunctionalizations of Cyclic Ethers with Nucleophiles. *Chem. Commun.* **2023**, *59*, 631–634.

(109) Roy, S.; Kumar, G.; Chatterjee, I. Photoinduced Diverse Reactivity of Diazo Compounds with Nitrosoarenes. *Org. Lett.* **2021**, *23*, 6709–6713.

(110) Cao, M.; Ren, Y.; Zhang, R.; Xu, H.; Cheng, P.; Xu, H.; Xu, Y.; Li, P. Photochemical “Cut and Sew” Transformations of Ethynylbenziodoxolone Reagents and Diazo Compounds. *Org. Lett.* **2023**, *25*, 6300–6304.

(111) Hari, D. P.; Waser, J. Copper-Catalyzed Oxy-Alkynylation of Diazo Compounds with Hypervalent Iodine Reagents. *J. Am. Chem. Soc.* **2016**, *138*, 2190–2193.

(112) Kirmse, W.; Kapps, M. Reaktionen des Diazomethans mit Diallylsulfid und Allylättern unter Kupfersalz-Katalyse. *Chem. Ber.* **1968**, *101*, 994–1003.

(113) Doyle, M. P.; Tamblyn, W. H.; Bagheri, V. Highly Effective Catalytic Methods for Ylide Generation from Diazo Compounds. Mechanism of the Rhodium- and Copper-Catalyzed Reactions with Allylic Compounds. *J. Org. Chem.* **1981**, *46*, 5094–5102.

(114) Yang, J.; Wang, J.; Huang, H.; Qin, G.; Jiang, Y.; Xiao, T. gem-Difluoroallylation of Aryl Diazoesters via Catalyst-Free, Blue-Light-Mediated Formal Doyle–Kirmse Reaction. *Org. Lett.* **2019**, *21*, 2654–2657.

(115) Orlowska, K.; Rybicka-Jasińska, K.; Krajewski, P.; Gryko, D. Photochemical Doyle–Kirmse Reaction: A Route to Allenes. *Org. Lett.* **2020**, *22*, 1018–1021.

(116) Xie, J.; Suleiman, M.; Wang, Z.; Mao, X.; Mao, B.; Fan, J.; Lu, P.; Wang, Y. Syntheses of 4-Allyl-/4-allenyl-4-(arylthio)-1,4-dihydroisoquinolin-3-ones via the Photochemical Doyle–Kirmse Reaction. *Org. Biomol. Chem.* **2021**, *19*, 6341–6345.

(117) Yang, Z.; Guo, Y.; Koenigs, R. M. Photochemical, Metal-Free Sigmatropic Rearrangement Reactions of Sulfur Ylides. *Chem.—Eur. J.* **2019**, *25*, 6703–6706.

(118) Muthusamy, S.; Ramesh, C. Blue LED-Mediated Synthesis of 3-Arylidene Oxindoles. *J. Org. Chem.* **2023**, *88*, 5609–5621.

(119) Qu, B.-L.; Shi, B.; He, L.; Shi, J.-W.; Xiao, W.-J.; Lu, L.-Q. Photoinduced Carbene Transfer for Copper-Catalyzed Asymmetric [4 + 1] Cycloadditions: an Entry to Chiral Indolines Bearing Quaternary Stereocenters. *Org. Chem. Front.* **2023**, *10*, 3498–3503.

(120) Xiao, T.; Mei, M.; He, Y.; Zhou, L. Blue light-promoted cross-coupling of aryl diazoacetates and diazocarbonyl compounds. *Chem. Commun.* **2018**, *54*, 8865–8868.

(121) Ansari, M. A.; Yadav, D.; Singh, M. S. Visible-Light-Driven Photocatalyst- and Additive-Free Cross-Coupling of  $\beta$ -Ketothioamides with  $\alpha$ -Diazo 1,3-Diketones: Access to Highly Functionalized Thiazolines. *Chem.—Eur. J.* **2020**, *26*, 8083–8089.

(122) da Silva, A. F.; Afonso, M. A. S.; Cormanich, R. A.; Jurberg, I. D. Room Temperature Coupling of Aryldiazoacetates with Boronic Acids Enhanced by Blue Light Irradiation. *Chem.—Eur. J.* **2020**, *26*, 5648–5653.

(123) Yang, J.; Duan, J.; Wang, G.; Zhou, H.; Ma, B.; Wu, C.; Xiao, J. Visible-Light-Promoted Site-Selective N1-Alkylation of Benzotriazoles with  $\alpha$ -Diazoacetates. *Org. Lett.* **2020**, *22*, 7284–7289.

(124) Song, C.; Dong, X.; Yi, H.; Chiang, C.-W.; Lei, A. DDQ-Catalyzed Direct C(sp<sup>3</sup>)–H Amination of Alkylheteroarenes: Synthesis of Biheteroarenes under Aerobic and Metal-Free Conditions. *ACS Catal.* **2018**, *8*, 2195–2199.

(125) Singh, M. K.; Akula, H. K.; Satishkumar, S.; Stahl, L.; Lakshman, M. K. Ruthenium-Catalyzed C–H Bond Activation Approach to Azolyl Aminals and Hemiaminal Ethers, Mechanistic Evaluations, and Isomer Interconversion. *ACS Catal.* **2016**, *6*, 1921–1928.

(126) Wang, K.; Chen, P.; Ji, D.; Zhang, X.; Xu, G.; Sun, J. Rhodium-Catalyzed Regioselective N2-Alkylation of Benzotriazoles with Diazo Compounds/Enynones via a Nonclassical Pathway. *Angew. Chem., Int. Ed.* **2018**, *57*, 12489–12493.

(127) Lu, J.; Li, L.; He, X.-K.; Xu, G.-Y.; Xuan, J. Visible Light-Promoted Sulfoxonium Ylides Synthesis from Aryl Diazoacetates and Sulfoxides. *Chin. J. Chem.* **2021**, *39*, 1646–1650.

(128) Khade, V. V.; Thube, A. S.; Warghude, P. K.; Bhat, R. G. DABCO Mediated One Pot Synthesis of Sulfoxonium Ylides under Blue LED. *Tetrahedron Lett.* **2021**, *77*, 153258.

(129) Gallo, R. D. C.; Duarte, M.; da Silva, A. F.; Okada, C. Y., Jr.; Deflon, V. M.; Jurberg, I. D. A Selective C–C Bond Cleavage Strategy Promoted by Visible Light. *Org. Lett.* **2021**, *23*, 8916–8920.

(130) Li, W.; Yang, Y.; Tang, Z.; Yu, X.; Lin, J.; Jin, Y. Visible-Light-Promoted Carbene Insertion and Decarbonylation for the Synthesis of  $\alpha$ -Substituted  $\gamma$ -Ketoesters. *J. Org. Chem.* **2022**, *87*, 13352–13362.

(131) Zhang, H.; Wang, Z.; Wang, Z.; Chu, Y.; Wang, S.; Hui, X.-P. Visible-Light-Mediated Formal Carbene Insertion Reaction: Enantioselective Synthesis of 1,4-Dicarbonyl Compounds Containing All-Carbon Quaternary Stereocenter. *ACS Catal.* **2022**, *12*, 5510–5516.

(132) Li, S.; Zhang, C.; Pan, G.; Yang, L.; Su, Z.; Feng, X.; Liu, X. Enantioselective Photochemical Carbene Insertion into C–C and C–H Bonds of 1,3-Diketones by a Guanidine-Amide Organocatalyst. *ACS Catal.* **2023**, *13*, 4656–4666.

(133) Sakharov, P. A.; Novikov, M. S.; Nguyen, T. K.; Kinzhakov, M. A.; Khlebnikov, A. F.; Rostovskii, N. V. Blue Light-Promoted Cross-Coupling of  $\alpha$ -Diazo Esters with Isocyanides: Synthesis of Ester-Functionalized Ketenimines. *ACS Omega* **2022**, *7*, 9071–9079.

(134) Zhao, Y.-R.; Li, L.; Xu, G.-Y.; Xuan, J. All-Carbon Tetrasubstituted Olefins Synthesis from Diazo Compounds and Iodonium Ylides under Blue LED Irradiation. *Adv. Synth. Catal.* **2022**, *364*, 506–511.

(135) Chidley, T.; Jameel, I.; Rizwan, S.; Peixoto, P. A.; Pouységu, L.; Quideau, S.; Hopkins, W. S.; Murphy, G. K. Blue LED Irradiation of Iodonium Ylides Gives Diradical Intermediates for Efficient Metal-free Cyclopropanation with Alkenes. *Angew. Chem., Int. Ed.* **2019**, *58*, 16959–16965.

(136) Bai, J.; Li, S.; Qi, D.; Song, Z.; Li, B.; Guo, L.; Song, L.; Xia, W. Visible-Light-Induced Trifluoromethylsulfonylation Reaction of Diazo Compounds Enabled by Manganese Catalysis. *Org. Lett.* **2023**, *25*, 2410–2414.

(137) Zhao, B.; Li, H.; Jiang, F.; Wan, J.-P.; Cheng, K.; Liu, Y. Synergistic Visible Light and Pd-Catalyzed C–H Alkylation of 1-Naphthylamines with  $\alpha$ -Diazoesters. *J. Org. Chem.* **2023**, *88*, 640–646.

(138) Hirai, K.; Itoh, T.; Tomioka, H. Persistent Triplet Carbenes. *Chem. Rev.* **2009**, *109*, 3275–3332.

(139) Eisenthal, K. B.; Turro, N. J.; Sitzmann, E. V.; Gould, I. R.; Hefferon, G.; Langan, J.; Cha, Y. Singlet-Triplet Interconversion of Diphenylmethylene. Energetics, Dynamics and Reactivities of Different Spin States. *Tetrahedron* **1985**, *41*, 1543–1554.

(140) Costa, P.; Lohmiller, T.; Trosien, I.; Savitsky, A.; Lubitz, W.; Fernandez-Oliva, M.; Sanchez-Garcia, E.; Sander, W. Light and Temperature Control of the Spin State of Bis(p-methoxyphenyl)-carbene: A Magnetically Bistable Carbene. *J. Am. Chem. Soc.* **2016**, *138*, 1622–1629.

(141) Costa, P.; Sander, W. Hydrogen Bonding Switches the Spin State of Diphenylcarbene from Triplet to Singlet. *Angew. Chem., Int. Ed.* **2014**, *53*, 5122–5125.

(142) Knorr, J.; Sokkar, P.; Schott, S.; Costa, P.; Thiel, W.; Sander, W.; Sanchez-Garcia, E.; Nuernberger, P. Competitive solvent-molecule interactions govern primary processes of diphenylcarbene in solvent mixtures. *Nat. Commun.* **2016**, *7*, 12968.

(143) Rybicka-Jasińska, K.; Shan, W.; Zawada, K.; Kadish, K. M.; Gryko, D. Porphyrins as Photoredox Catalysts: Experimental and Theoretical Studies. *J. Am. Chem. Soc.* **2016**, *138*, 15451–15458.

(144) Cismesia, M. A.; Yoon, T. P. Characterizing Chain Processes in Visible Light Photoredox Catalysis. *Chem. Sci.* **2015**, *6*, 5426–5434.

(145) Li, F.; Pei, C.; Koenigs, R. M. Photocatalytic gem-Difluoroolefination Reactions by a Formal C–C Coupling/Defluor

nation Reaction with Diazoacetates. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202111892.

(146) Langletz, T.; Empel, C.; Jana, S.; Koenigs, R. M. Stereoconvergent, Photocatalytic Cyclopropanation Reactions of  $\beta$ -Substituted Styrenes with Ethyl Diazoacetate. *Tetrahedron Chem.* **2022**, *3*, 100024.

(147) Zhu, S.; Li, F.; Empel, C.; Jana, S.; Pei, C.; Koenigs, R. M. Furan Synthesis via Triplet Sensitization of Acceptor/Acceptor Diazoalkanes. *Adv. Synth. Catal.* **2022**, *364*, 3149–3154.

(148) Empel, C.; Jana, S.; Ciszewski, Ł. W.; Zawada, K.; Pei, C.; Gryko, D.; Koenigs, R. M. C–H Functionalization of Heterocycles with Triplet Carbenes by means of an Unexpected 1,2-Alkyl Radical Migration\*\*. *Chem.—Eur. J.* **2023**, *29*, No. e202300214.

(149) Zhou, H.; Wang, G.; Wang, C.; Yang, J. Visible-Light-Promoted Aerobic Oxyphosphorylation of  $\alpha$ -Diazoesters with H-Phosphine Oxides. *Org. Lett.* **2022**, *24*, 1530–1535.

(150) Parasram, M.; Gevorgyan, V. Visible Light-Induced Transition Metal-Catalyzed Transformations: Beyond Conventional Photosensitizers. *Chem. Soc. Rev.* **2017**, *46*, 6227–6240.

(151) Cheung, K. P. S.; Sarkar, S.; Gevorgyan, V. Visible Light-Induced Transition Metal Catalysis. *Chem. Rev.* **2022**, *122*, 1543–1625.

(152) Giedyk, M.; Goliszewska, K.; ó Proinsias, K.; Gryko, D. Cobalt(i)-catalysed CH-alkylation of terminal olefins, and beyond. *Chem. Commun.* **2016**, *52*, 1389–1392.

(153) Chen, Y.; Zhang, X. P. Vitamin B12 Derivatives as Natural Asymmetric Catalysts: Enantioselective Cyclopropanation of Alkenes. *J. Org. Chem.* **2004**, *69*, 2431–2435.

(154) Epping, R. F. J.; Vesseur, D.; Zhou, M.; de Bruin, B. Carbene Radicals in Transition-Metal-Catalyzed Reactions. *ACS Catal.* **2023**, *13*, 5428–5448.

(155) Su, Y.-L.; Liu, G.-X.; Liu, J.-W.; Tram, L.; Qiu, H.; Doyle, M. P. Radical-Mediated Strategies for the Functionalization of Alkenes with Diazo Compounds. *J. Am. Chem. Soc.* **2020**, *142*, 13846–13855.

(156) Fu, X.; Tang, J.; Hua, R.; Li, X.; Kang, Z.; Qiu, H.; Hu, W. Functionalization of DNA-Tagged Alkenes with Diazo Compounds via Photocatalysis. *Org. Lett.* **2022**, *24*, 2208–2213.

(157) Zhang, Z.; Kvasovs, N.; Dubrovina, A.; Gevorgyan, V. Visible Light Induced Brønsted Acid Assisted Pd-Catalyzed Alkyl Heck Reaction of Diazo Compounds and N-Tosylhydrazones. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202110924.

(158) Ma, N.; Guo, L.; Qi, D.; Gao, F.; Yang, C.; Xia, W. Visible-Light-Induced Multicomponent Synthesis of  $\gamma$ -Amino Esters with Diazo Compounds. *Org. Lett.* **2021**, *23*, 6278–6282.

(159) Zhang, B.; Qi, J.-Q.; Liu, Y.; Li, Z.; Wang, J. Visible-Light-Driven Bisfunctionalization of Unactivated Olefins via the Merger of Proton-Coupled Electron Transfer and Carbene Catalysis. *Org. Lett.* **2022**, *24*, 279–283.

(160) Li, F.; Zhu, S.; Koenigs, R. M. Photocatalytic 1,2-oxo-alkylation reaction of styrenes with diazoacetates. *Chem. Commun.* **2022**, *58*, 7526–7529.

(161) Su, Y.-L.; Liu, G.-X.; De Angelis, L.; He, R.; Al-Sayyed, A.; Schanze, K. S.; Hu, W.-H.; Qiu, H.; Doyle, M. P. Radical Cascade Multicomponent Minisci Reactions with Diazo Compounds. *ACS Catal.* **2022**, *12*, 1357–1363.

(162) Liu, Y.; Zhu, K.; Zhao, J.; Li, P. Photocatalytic Regioselective Difunctionalization of Alkenes with Diazo Compounds and tert-Butyl Nitrite: Access to  $\gamma$ -Oximino Esters. *Org. Lett.* **2022**, *24*, 6834–6838.

(163) Ciszewski, Ł. W.; Durka, J.; Gryko, D. Photocatalytic Alkylation of Pyrroles and Indoles with  $\alpha$ -Diazo Esters. *Org. Lett.* **2019**, *21*, 7028–7032.

(164) Bhattacharjee, S.; Laru, S.; Samanta, S.; Singsardar, M.; Hajra, A. Visible light-induced photocatalytic C–H ethoxycarbonylmethylation of imidazoheterocycles with ethyl diazoacetate. *RSC Adv.* **2020**, *10*, 27984–27988.

(165) Li, P.; Chang, R.; Guan, F.; Yang, W.; Zhou, C.-Y.; Guo, Z. Heterogeneous Photocatalytic C–H Functionalization of Indoles with Diazo Compounds Enabled by Carbon Nitride. *J. Org. Chem.* **2023**, *88*, 8703–8708.

(166) Karmakar, U.; Hwang, H. S.; Lee, Y.; Cho, E. J. Photocatalytic para-Selective C–H Functionalization of Anilines with Diazomalonates. *Org. Lett.* **2022**, *24*, 6137–6141.

(167) Li, W.; Qiu, J.; Li, H.; Chen, W.; Hou, C.; Li, H. Indolization of N-Aryl Tertiary Amines with Diazoacetates by a Single Organophotocatalyst. *Org. Lett.* **2023**, *25*, 3778–3783.

(168) Huang, X.; Webster, R. D.; Harms, K.; Meggers, E. Asymmetric Catalysis with Organic Azides and Diazo Compounds Initiated by Photoinduced Electron Transfer. *J. Am. Chem. Soc.* **2016**, *138*, 12636–12642.

(169) Nagode, S. B.; Kant, R.; Rastogi, N. Hantzsch Ester-Mediated Benzannulation of Diazo Compounds under Visible Light Irradiation. *Org. Lett.* **2019**, *21*, 6249–6254.

(170) Ye, H.-B.; Zhou, X.-Y.; Li, L.; He, X.-K.; Xuan, J. Photochemical Synthesis of Succinic Ester-Containing Phenanthridines from Diazo Compounds as 1,4-Dicarbonyl Precursors. *Org. Lett.* **2022**, *24*, 6018–6023.

(171) Wang, Q.; Liu, J.; Mei, H.; Pajkert, R.; Kessler, M.; Röschenthaler, G.-V.; Han, J. Ru-Catalyzed Hydrogen Atom Transfer/C–F Bond Cleavage of Difluoroalkyl Diazo Compounds with Hantzsch Ester via a Photocatalytic Radical Process. *Org. Lett.* **2022**, *24*, 8036–8040.

(172) Gao, Y.; Zheng, Z.; Zhu, Y.; Xu, W.; Zhou, Y.; Yu, C.; Jiang, X. Visible Light-Induced Synthesis of 1,3-Disubstituted Bicyclo[1.1.1]-pentane Ketones via Cooperative Photoredox and N-Heterocyclic Carbene Catalysis. *Green Chem.* **2023**, *25*, 3909–3915.

(173) Xu, W.; Zheng, Z.; Bao, G.; Wang, Y.; Gao, Y.; Zhu, H.; Xu, G.; Zhu, Y.; Yu, C.; Jiang, X. Photocatalytic Three-Component Synthesis of 3-Heteroaryl bicyclo[1.1.1]-pentane-1-acetates. *Org. Lett.* **2023**, *25*, 4050–4055.

(174) Timmann, S.; Alcarazo, M.  $\alpha$ -Diazo- $\lambda^3$ -Iodanes and  $\alpha$ -Diazo Sulfonium Salts: the Umpolung of Diazo Compounds. *Chem. Commun.* **2023**, *59*, 8032–8042.

(175) Yoshimura, A.; Zhdankin, V. V. Advances in Synthetic Applications of Hypervalent Iodine Compounds. *Chem. Rev.* **2016**, *116*, 3328–3435.

(176) Wang, X.; Studer, A. Iodine(III) Reagents in Radical Chemistry. *Acc. Chem. Res.* **2017**, *50*, 1712–1724.

(177) Kaiser, D.; Klose, I.; Oost, R.; Neuhaus, J.; Maulide, N. Bond-Forming and -Breaking Reactions at Sulfur(IV): Sulfoxides, Sulfonium Salts, Sulfur Ylides, and Sulfinate Salts. *Chem. Rev.* **2019**, *119*, 8701–8780.

(178) Wang, Z.; Herraiz, A. G.; del Hoyo, A. M.; Suero, M. G. Generating Carbyne Equivalents with Photoredox Catalysis. *Nature* **2018**, *554*, 86–91.

(179) Liu, L.; Zhang, Y.; Zhao, W.; Wen, J.; Dong, C.; Hu, C.; Li, J. Photoredox-Catalyzed Cascade sp<sup>2</sup> C–H Bond Functionalization to Construct Substituted Acridine with Diarylamine and Hypervalent Iodine(III) Reagents. *Org. Lett.* **2023**, *25*, 592–596.

(180) Li, J.; Lu, X.-C.; Xu, Y.; Wen, J.-X.; Hou, G.-Q.; Liu, L. Photoredox Catalysis Enables Decarboxylative Cyclization with Hypervalent Iodine(III) Reagents: Access to 2,5-Disubstituted 1,3,4-Oxadiazoles. *Org. Lett.* **2020**, *22*, 9621–9626.

(181) Zhao, W.-W.; Shao, Y.-C.; Wang, A.-N.; Huang, J.-L.; He, C.-Y.; Cui, B.-D.; Wan, N.-W.; Chen, Y.-Z.; Han, W.-Y. Diazotrifluoroethyl Radical: A CF<sub>3</sub>-Containing Building Block in [3 + 2] Cycloaddition. *Org. Lett.* **2021**, *23*, 9256–9261.

(182) Li, J.; Wen, J.-X.; Lu, X.-C.; Hou, G.-Q.; Gao, X.; Li, Y.; Liu, L. Catalyst-Free Visible-Light-Promoted Cyclization of Aldehydes: Access to 2,5-Disubstituted 1,3,4-Oxadiazole Derivatives. *ACS Omega* **2021**, *6*, 26699–26706.

(183) Li, X.; Golz, C.; Alcarazo, M.  $\alpha$ -Diazo Sulfonium Triflates: Synthesis, Structure, and Application to the Synthesis of 1-(Dialkylamino)-1,2,3-triazoles. *Angew. Chem., Int. Ed.* **2021**, *60*, 6943–6948.

(184) Dong, J.-Y.; Wang, H.; Mao, S.; Wang, X.; Zhou, M.-D.; Li, L. Visible Light-Induced [3 + 2] Cyclization Reactions of Hydrazones with Hypervalent Iodine Diazo Reagents for the Synthesis of 1-Amino-1,2,3-Triazoles. *Adv. Synth. Catal.* **2021**, *363*, 2133–2139.

(185) Wen, J.; Zhao, W.; Gao, X.; Ren, X.; Dong, C.; Wang, C.; Liu, L.; Li, J. Synthesis of [1,2,3]Triazolo-[1,5-a]quinoxalin-4(SH)-ones through Photoredox-Catalyzed [3 + 2] Cyclization Reactions with Hypervalent Iodine(III) Reagents. *J. Org. Chem.* **2022**, *87*, 4415–4423.

(186) Monreal-Corona, R.; Besalú, E.; Pla-Quintana, A.; Poater, A. Photoredox catalysis leading to triazolo-quinoxalinones at room temperature: selectivity of the rate determining step. *Org. Biomol. Chem.* **2022**, *20*, 9330–9336.

(187) Xu, X.-C.; Wu, D.-N.; Liang, Y.-X.; Yang, M.; Yuan, H.-Y.; Zhao, Y.-L. Visible Light-Induced Coupling Cyclization Reaction of  $\alpha$ -Diazosulfonium Triflates with  $\alpha$ -Oxocarboxylic Acids or Alkynes. *J. Org. Chem.* **2022**, *87*, 16604–16616.

(188) Ma, M.; Hao, W.; Ma, L.; Zheng, Y.; Lian, P.; Wan, X. Interception of Radicals by Molecular Oxygen and Diazo Compounds: Direct Synthesis of Oxalate Esters Using Visible-Light Catalysis. *Org. Lett.* **2018**, *20*, 5799–5802.

(189) Chan, C.-M.; Xing, Q.; Chow, Y.-C.; Hung, S.-F.; Yu, W.-Y. Photoredox Decarboxylative C(sp<sup>3</sup>)-N Coupling of  $\alpha$ -Diazoacetates with Alkyl N-Hydroxyphthalimide Esters for Diversified Synthesis of Functionalized N-Alkyl Hydrazones. *Org. Lett.* **2019**, *21*, 8037–8043.

(190) Li, W.; Zhou, L. Vinyldiazo Compounds as 3-Carbon Radical Acceptors: Synthesis of 4-Fluoroacridines via Visible-Light-Promoted Cascade Radical Cyclization. *Org. Lett.* **2021**, *23*, 4279–4283.

(191) Li, W.; Zhou, X.; Xiao, T.; Ke, Z.; Zhou, L. External Photocatalyst-Free Visible Light-Promoted 1,3-Addition of Perfluoroalkyl Iodides to Vinyldiazoacetates. *CCS Chemistry* **2022**, *4*, 638–649.

(192) Li, W.; Zhou, L. Visible-Light-Driven Radical 1,3-Addition of Selenosulfonates to Vinyldiazo Compounds. *Green Chem.* **2021**, *23*, 6652–6658.

(193) Li, W.; Zhou, L. Synthesis of Tetrasubstituted Allenes via Visible-Light-Promoted Radical 1,3-Difunctionalization of Alkynyl Diazo Compounds. *Org. Lett.* **2022**, *24*, 3976–3981.

(194) Cai, B.-G.; Bao, Y.-P.; Pei, C.; Li, Q.; Li, L.; Koenigs, R. M.; Xuan, J. Photochemical Synthesis of 1,2,4-Triazoles via Addition Reaction of Triplet Intermediates to Diazoalkanes and Azomethine Ylide Intermediates. *Chem. Sci.* **2022**, *13*, 13141–13146.

(195) Zhao, P.; Wang, L.; Guo, X.; Chen, J.; Liu, Y.; Wang, L.; Ma, Y. Visible Light-Driven  $\alpha$ -Diazoketones as Denitrogenated Synthons: Synthesis of Fluorinated N-Heterocycles via Multicomponent Cyclization Reactions. *Org. Lett.* **2023**, *25*, 3314–3318.

(196) Bug, T.; Hartnagel, M.; Schlierf, C.; Mayr, H. How Nucleophilic Are Diazo Compounds? *Chem.—Eur. J.* **2003**, *9*, 4068–4076.

(197) López, E.; González-Pelayo, S.; López, L. A. Recent Developments in Coinage Metal Catalyzed Transformations of Stabilized Vinyldiazo Compounds: Beyond Carbenic Pathways. *Chem. Rec.* **2017**, *17*, 312–325.

(198) Zhang, Y.; Wang, J. Recent Development of Reactions with  $\alpha$ -Diazocarbonyl Compounds as Nucleophiles. *Chem. Commun.* **2009**, 5350–5361.

(199) Xiao, T.; Li, L.; Lin, G.; Mao, Z.-w.; Zhou, L. Metal-Free Visible-Light Induced Cross-Dehydrogenative Coupling of Tertiary Amines with Diazo Compounds. *Org. Lett.* **2014**, *16*, 4232–4235.

(200) Pramanik, M. M. D.; Nagode, S. B.; Kant, R.; Rastogi, N. Visible light catalyzed Mannich reaction between tert-amines and silyl diazoenolates. *Org. Biomol. Chem.* **2017**, *15*, 7369–7373.

(201) Liu, Y.; Dong, X.; Deng, G.; Zhou, L. Synthesis of Aziridines by Visible-Light Induced Decarboxylative Cyclization of N-Aryl Glycines and Diazo Compounds. *Sci. China Chem.* **2016**, *59*, 199–202.

(202) Sarabia, F. J.; Ferreira, E. M. Radical Cation Cyclopropanations via Chromium Photooxidative Catalysis. *Org. Lett.* **2017**, *19*, 2865–2868.

(203) Sarabia, F. J.; Li, Q.; Ferreira, E. M. Cyclopentene Annulations of Alkene Radical Cations with Vinyl Diazo Species Using Photocatalysis. *Angew. Chem., Int. Ed.* **2018**, *57*, 11015–11019.

(204) Gall, B. K.; Smith, A. K.; Ferreira, E. M. Dearomatic (3 + 2) Cycloadditions between Indoles and Vinyldiazo Species Enabled by a Red-Shifted Chromium Photocatalyst. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202212187.

(205) Holmberg-Douglas, N.; Onuska, N. P. R.; Nicewicz, D. A. Regioselective Arene C–H Alkylation Enabled by Organic Photoredox Catalysis. *Angew. Chem., Int. Ed.* **2020**, *59*, 7425–7429.

(206) Li, W.; Li, S.; Empel, C.; Koenigs, R. M.; Zhou, L. Photoredox-Enabled Self-(3 + 2) Cyclization of Vinyldiazo Reagents: Synthesis of Cyclopentenyl  $\alpha$ -Diazo Compounds. *Angew. Chem., Int. Ed.* **2023**, *62*, No. e202309947.

(207) Li, P.; Zhao, J.; Shi, L.; Wang, J.; Shi, X.; Li, F. Iodine-catalyzed diazo activation to access radical reactivity. *Nat. Commun.* **2018**, *9*, 1972.

(208) Zhao, S.; Gao, N.; Bao, N.; Qian, M.; Chen, Z.-Y.; Zhang, M.-J.; Chen, X. Stereoselective Synthesis of Tetrasubstituted Olefins via Visible-Light-Promoted Iodine-Mediated Homo-Coupling of Diazo. *J. Org. Chem.* **2022**, *87*, 11826–11837.

(209) Zhou, Y.; Pang, T.; Liu, Y.; Zhong, F.; Wu, G. Iodine-Mediated Coupling of 2,2,2-Trifluorodiazooethane and Alkynes To Access Bistrifluoromethylated 1,3,5-Trienes. *Org. Lett.* **2023**, *25*, 538–542.

(210) Liu, Y.; Pang, T.; Yao, W.; Zhong, F.; Wu, G. Visible-Light-Induced Radical gem-Iodoallylation of 2,2,2-Trifluorodiazooethane. *Org. Lett.* **2023**, *25*, 1958–1962.

(211) Jana, S.; Yang, Z.; Li, F.; Empel, C.; Ho, J.; Koenigs, R. M. Photoinduced Proton-Transfer Reactions for Mild O–H Functionalization of Unreactive Alcohols. *Angew. Chem., Int. Ed.* **2020**, *59*, 5562–5566.

(212) Empel, C.; Jana, S.; Pei, C.; Nguyen, T. V.; Koenigs, R. M. Photochemical O–H Functionalization of Aryldiazoacetates with Phenols via Proton Transfer. *Org. Lett.* **2020**, *22*, 7225–7229.

(213) Empel, C.; Jana, S.; Pei, C.; Nguyen, T. V.; Koenigs, R. M. Photochemical O–H Functionalization of Aryldiazoacetates with Phenols via Proton Transfer. *Org. Lett.* **2020**, *22*, 7225–7229.

(214) Pei, C.; Koenigs, R. M. A Computational Study on the Photochemical O–H Functionalization of Alcohols with Diazoacetates. *J. Org. Chem.* **2022**, *87*, 6832–6837.

(215) Empel, C.; Pei, C.; He, F.; Jana, S.; Koenigs, R. M. Proton or Carbene Transfer? On the Dark and Light Reaction of Diazoalkanes with Alcohols. *Chem.—Eur. J.* **2022**, *28*, No. e202104397.

(216) Gallardo, G. M.; Ventura, D. J.; Petit, A. S. Computationally Probing the Mechanism of the Blue-Light-Driven O–H Functionalization of Alcohols by Aryldiazoacetates: Photobasicity or Carbene Chemistry. *J. Org. Chem.* **2022**, *87*, 6212–6223.

(217) Liu, R.; Liu, Q.; Meng, H.; Ding, H.; Hao, J.; Ji, Z.; Yue, H.; Wei, W. Metal-Free Visible-Light-Induced Aerobic Oxidation of  $\alpha$ -Diazooesters Leading to  $\alpha$ -Ketoesters in Air. *Org. Chem. Front.* **2021**, *8*, 1970–1975.

(218) Wang, X.; Tong, W.-Y.; Huang, B.; Cao, S.; Li, Y.; Jiao, J.; Huang, H.; Yi, Q.; Qu, S.; Wang, X. Convergent Synthesis of 1,4-Dicarbonyl Z-Alkenes through Three-Component Coupling of Alkynes,  $\alpha$ -Diazooesters, and Water. *J. Am. Chem. Soc.* **2022**, *144*, 4952–4965.

(219) Hussain, Y.; Empel, C.; Koenigs, R. M.; Chauhan, P. Carbene Formation or Reduction of the Diazo Functional Group? An Unexpected Solvent-Dependent Reactivity of Cyclic Diazo Imides. *Angew. Chem., Int. Ed.* **2023**, *62*, No. e202309184.

(220) Liu, M. T. H. The Thermolysis and Photolysis of Diazirines. *Chem. Soc. Rev.* **1982**, *11*, 127–140.

(221) Das, J. Aliphatic Diazirines as Photoaffinity Probes for Proteins: Recent Developments. *Chem. Rev.* **2011**, *111*, 4405–4417.

(222) Ge, S.-S.; Chen, B.; Wu, Y.-Y.; Long, Q.-S.; Zhao, Y.-L.; Wang, P.-Y.; Yang, S. Current advances of carbene-mediated photoaffinity labeling in medicinal chemistry. *RSC Adv.* **2018**, *8*, 29428–29454.

(223) Dubinsky, L.; Krom, B. P.; Meijler, M. M. Diazirine based photoaffinity labeling. *Biorg. Med. Chem.* **2012**, *20*, 554–570.

(224) Halloran, M. W.; Lumb, J.-P. Recent Applications of Diazirines in Chemical Proteomics. *Chem.—Eur. J.* **2019**, *25*, 4885–4898.

(225) Brunner, J.; Senn, H.; Richards, F. M. 3-Trifluoromethyl-3-phenyldiazirine. A new carbene generating group for photolabeling reagents. *J. Biol. Chem.* **1980**, *255*, 3313–3318.

(226) Geri, J. B.; Oakley, J. V.; Reyes-Robles, T.; Wang, T.; McCarver, S. J.; White, C. H.; Rodriguez-Rivera, F. P.; Parker, D. L.; Hett, E. C.; Fadeyi, O. O.; Oslund, R. C.; MacMillan, D. W. C. Microenvironment Mapping via Dexter Energy Transfer on Immune Cells. *Science* **2020**, *367*, 1091–1097.

(227) Trowbridge, A. D.; Seath, C. P.; Rodriguez-Rivera, F. P.; Li, B. X.; Dul, B. E.; Schwaid, A. G.; Buksh, B. F.; Geri, J. B.; Oakley, J. V.; Fadeyi, O. O.; Oslund, R. C.; Ryu, K. A.; White, C.; Reyes-Robles, T.; Tawa, P.; Parker, D. L.; MacMillan, D. W. C. Small Molecule Photocatalysis Enables Drug Target Identification via Energy Transfer. *Proc. Natl. Acad. Sci. U. S. A.* **2022**, *119*, No. e2208077119.

(228) Seath, C. P.; Burton, A. J.; Sun, X.; Lee, G.; Kleiner, R. E.; MacMillan, D. W. C.; Muir, T. W. Tracking Chromatin State Changes Using Nanoscale Photo-proximity Labelling. *Nature* **2023**, *616*, 574–580.

(229) Huth, S. W.; Oakley, J. V.; Seath, C. P.; Geri, J. B.; Trowbridge, A. D.; Parker, D. L., Jr.; Rodriguez-Rivera, F. P.; Schwaid, A. G.; Ramlil, C.; Ryu, K. A.; White, C. H.; Fadeyi, O. O.; Oslund, R. C.; MacMillan, D. W. C.  $\mu$ Map Photoproximity Labeling Enables Small Molecule Binding Site Mapping. *J. Am. Chem. Soc.* **2023**, *145*, 16289–16296.

(230) Xia, Y.; Wang, J. N-Tosylhydrazone: Versatile Synthons in the Construction of Cyclic Compounds. *Chem. Soc. Rev.* **2017**, *46*, 2306–2362.

(231) Xiao, Q.; Zhang, Y.; Wang, J. Diazo Compounds and N-Tosylhydrazone: Novel Cross-Coupling Partners in Transition-Metal-Catalyzed Reactions. *Acc. Chem. Res.* **2013**, *46*, 236–247.

(232) Bamford, W. R.; Stevens, T. S. 924. The Decomposition of Toluene-p-Sulphonylhydrazone by Alkali. *J. Chem. Soc.* **1952**, *4735*–4740.

(233) Jana, S.; Li, F.; Empel, C.; Verspeek, D.; Aseeva, P.; Koenigs, R. M. Stoichiometric Photochemical Carbene Transfer by Bamford–Stevens Reaction. *Chem.—Eur. J.* **2020**, *26*, 2586–2591.

(234) Xu, Y.; Lv, G.; Yan, K.; He, H.; Li, J.; Luo, Y.; Lai, R.; Hai, L.; Wu, Y. Blue Light-promoted Carbene Transfer Reactions of Tosylhydrazone. *Chemistry – An Asian Journal* **2020**, *15*, 1945–1947.

(235) Yan, K.; He, H.; Li, J.; Luo, Y.; Lai, R.; Guo, L.; Wu, Y. Blue Light-Promoted Cyclopropenizations of N-Tosylhydrazone in Water. *Chin. Chem. Lett.* **2021**, *32*, 3984–3987.

(236) Wang, Y.; Jia, P.; Hao, Y.; Li, J.; Lai, R.; Guo, L.; Wu, Y. Blue Light Induced [2,3]-Sigmatropic Rearrangement Reactions of Tosylhydrazone. *Tetrahedron Lett.* **2022**, *107*, 154098.

(237) Huang, X.; Chen, X.; Xie, H.; Tan, Z.; Jiang, H.; Zeng, W. Visible-Light-Catalyzed In Situ Denitrogenative Sulfenylation of Sulfonylhydrazone. *Org. Lett.* **2021**, *23*, 6784–6788.

(238) Wang, H.; Wang, S.; George, V.; Llorente, G.; König, B. Photo-Induced Homologation of Carbonyl Compounds for Iterative Syntheses. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202211578.

(239) Xia, D.; Wu, R.; Wang, J.; Han, X.; Li, Y.; Li, Q.; Luan, X.; Hong, X.; Zhang, Y.; Zhang, W.-D. Visible-Light-Mediated Energy Transfer Enables Cyclopropanes Bearing Contiguous All-Carbon Quaternary Centers. *ACS Catal.* **2023**, *13*, 9806–9816.

(240) Akter, M.; Rupa, K.; Anbarasan, P. 1,2,3-Triazole and Its Analogue: New Surrogates for Diazo Compounds. *Chem. Rev.* **2022**, *122*, 13108–13205.

(241) Anbarasan, P.; Yadagiri, D.; Rajasekar, S. Recent Advances in Transition-Metal-Catalyzed Denitrogenative Transformations of 1,2,3-Triazoles and Related Compounds. *Synthesis* **2014**, *46*, 3004–3023.

(242) Shen, R.; Yin, K.; Yu, M.; Li, X. Progress on Pyridotriazole Chemistry: Synthesis and Transformations. *Asian J. Org. Chem.* **2023**, *12*, No. e202200631.

(243) (a) Kuzaj, M.; Lüerssen, H.; Wentrup, C. ESR Observation of Thermally Produced Triplet Nitrenes and Photochemically Produced Triplet Cycloheptatrienylidene. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 480–482. (b) Tomioka, H.; Ichikawa, N.; Komatsu, K. Photochemistry of 2-(Methoxycarbonyl)phenyl Azide Studied by Matrix-Isolation Spectroscopy. A New Slippery Energy Surface for Phenyl-nitrene. *J. Am. Chem. Soc.* **1993**, *115*, 8621–8626. (c) Kuhn, A.; Plüg, C.; Wentrup, C. Direct Observation of a Carbene–Pyridine Ylide by Matrix IR Spectroscopy. Rearrangements of 2-Pyridylacylcarbenes. *J. Am. Chem. Soc.* **2000**, *122*, 1945–1948. (d) Plüg, C.; Kuhn, A.; Wentrup, C. Quinolizine-2,4-Diones by Reversible Dimerisation 2-Pyridylketenes. *J. Chem. Soc., Perkin Trans. 2* **2002**, *1*, 1366–1368. (e) Andersen, H. G.; Bednarek, P.; Wentrup, C. Methoxy(2-pyridyl)ketene. *J. Phys. Org. Chem.* **2003**, *16*, 519–524. (f) Fu, N.; Allen, A. D.; Kobayashi, S.; Sequeira, P. A.; Shang, M.; Tidwell, T. T.; Mishima, M. Observable Azacyclobutene Ylides with Antiromatic Character from 2-Diazoacetyl-azaaromatics. *J. Am. Chem. Soc.* **2007**, *129*, 6210–6215. (g) Zimmerman, H. E.; Ignatchenko, A. Control of the Stereochemistry of Kinetic Protonation: Intramolecular Proton Delivery 1,2. *J. Org. Chem.* **1999**, *64*, 6635–6645.

(244) Zhang, Z.; Yadagiri, D.; Gevorgyan, V. Light-induced metal-free transformations of unactivated pyridotriazoles. *Chem. Sci.* **2019**, *10*, 8399–8404.

(245) El-Saidi, M.; Kassam, K.; Pole, D. L.; Tadey, T.; Warkentin, J. 2,2-Dialkoxy- $\Delta$ 3-1,3,4-Oxadiazolines: Convenient Thermal Sources of Dialkoxy Carbene. *J. Am. Chem. Soc.* **1992**, *114*, 8751–8752.

(246) Warkentin, J. 2,5-Dihydro-1,3,4-oxadiazoles and Bis(heteroatom-substituted)carbenes. *Acc. Chem. Res.* **2009**, *42*, 205–212.

(247) Hoffmann, R. W.; Luthardt, H. J. Thermolyse und Photolyse von  $\Delta$ 3-1,3,4-Oxadiazolinen. *Chem. Ber.* **1968**, *101*, 3861–3871.

(248) Greb, A.; Poh, J.-S.; Greed, S.; Battilocchio, C.; Pasau, P.; Blakemore, D. C.; Ley, S. V. A Versatile Route to Unstable Diazo Compounds via Oxadiazolines and their Use in Aryl–Alkyl Cross-Coupling Reactions. *Angew. Chem., Int. Ed.* **2017**, *56*, 16602–16605.

(249) Dingwall, P.; Greb, A.; Crespin, L. N. S.; Labes, R.; Musio, B.; Poh, J.-S.; Pasau, P.; Blakemore, D. C.; Ley, S. V. C–H Functionalisation of Aldehydes Using Light Generated, Non-Stabilised Diazo Compounds in Flow. *Chem. Commun.* **2018**, *54*, 11685–11688.

(250) Orlowska, K.; Santiago, J. V.; Krajewski, P.; Kisiel, K.; Deperasińska, I.; Zawada, K.; Chaładaj, W.; Gryko, D. UV Light Is No Longer Required for the Photoactivation of 1,3,4-Oxadiazolines. *ACS Catal.* **2023**, *13*, 1964–1973.

(251) Mei, H.; Liu, J.; Pajkert, R.; Wang, L.; Röschenthaler, G.-V.; Han, J. Design of ( $\beta$ -Diazo- $\alpha$ , $\alpha$ -difluoroethyl)phosphonates and Their Application as Masked Carbenes in Visible Light-Promoted Coupling Reactions with Sulfonic Acids. *Org. Chem. Front.* **2021**, *8*, 767–772.