

Chiral fermion anomaly as a memory effect

Adrián del Río^{1,2,*} and Ivan Agullo^{3,4,†}

¹*Departamento de Física Teórica and IFIC, Universitat de Valencia - CSIC,
Dr. Moliner 50, 46100, Burjassot (Valencia), Spain*

²*Institute for Gravitation and the Cosmos & Physics Department, The Pennsylvania State University,
University Park, Pennsylvania 16802, USA*

³*Department of Physics and Astronomy, Louisiana State University, Baton Rouge,
Louisiana 70803-4001, USA*

⁴*Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo,
Ontario, Canada N2L 2Y5*



(Received 18 August 2023; accepted 30 October 2023; published 30 November 2023)

We study the nonconservation of the chiral charge of Dirac fields between past and future null infinity due to the Adler-Bell-Jackiw chiral anomaly. In previous investigations [A. del Rio, *Phys. Rev. D* **104**, 065012 (2021)], we found that this charge fails to be conserved if electromagnetic sources in the bulk emit circularly polarized radiation. In this article, we unravel yet another contribution coming from the nonzero, infrared “soft” charges of the external, electromagnetic field. This new contribution can be interpreted as another manifestation of the ordinary memory effect produced by transitions between different infrared sectors of Maxwell theory, but now on test quantum fields rather than on test classical particles. In other words, a flux of electromagnetic waves can leave a memory on quantum fermion states in the form of a permanent, net helicity. We elaborate this idea in both $1+1$ and $3+1$ dimensions. We also show that, in sharp contrast, gravitational infrared charges do not contribute to the fermion chiral anomaly.

DOI: 10.1103/PhysRevD.108.105025

I. INTRODUCTION

Not all symmetries of a classical theory remain exact after quantization. When this occurs, i.e., when a symmetry of the action is broken by quantum effects, one speaks about anomalies [1]. Anomalies were first discovered in the late 1960s, in the seminal works by Adler, Bell, and Jackiw, as an attempt of solving the pion decay puzzle [2,3]. They found that the chiral symmetry of the action of a massless Dirac field $\Psi(x)$ that interacts with an electromagnetic background is broken in quantum field theory. Mathematically, this outstanding result is beautifully encoded in the nonconservation equation of the fermionic chiral current $j_A^a(x) = \bar{\Psi}(x)\gamma^a\gamma^5\Psi(x)$, which on a $3+1$ dimensional Minkowski spacetime takes the form

$$\langle \nabla_a j_A^a \rangle = -\frac{\hbar q^2}{8\pi^2} F_{ab} \star F^{ab}, \quad (1)$$

where F_{ab} is the field strength of the background electromagnetic field, $\star F^{ab}$ its Hodge dual, and q the charge of the fermion. This is the celebrated chiral or axial anomaly.

Besides electromagnetic fields, gravitational backgrounds have also the ability of triggering a chiral anomaly, as it was soon after found in [4–6]. Mathematically, this contribution generalizes the previous equation by adding a new term proportional to the pseudoscalar curvature invariant $R_{abcd} \star R^{abcd}$, where R_{abcd} is the Riemann tensor and $\star R_{abcd}$ its Hodge dual with respect to the first two indices:

$$\langle \nabla_a j_A^a \rangle = -\frac{\hbar q^2}{8\pi^2} F_{ab} \star F^{ab} + \frac{\hbar}{192\pi^2} R_{abcd} \star R^{abcd}. \quad (2)$$

The following years experienced an outbreak of fascinating results involving anomalies, both regarding physics and mathematics. Examples include, besides the prediction of the neutral pion decay rate to two photons, applications to the matter-antimatter asymmetry of the universe, the U(1) and strong CP problems in QCD, implications for anomaly cancellation in the Standard Model (see [7] for a nice

*adrian.rio@uv.es
†agullo@lsu.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

summary of all these applications), and connections with the index theorems in geometric analysis [8–10]. The notion of chiral anomalies has also been extended to other fields, including integer spin fields [11–14].

In this article, we investigate yet another aspect of chiral anomalies, related to *global* properties of both the fermionic and the background fields. These global properties appear when discussing the Noether charge associated with the chiral current, namely $Q_A = \int_{\Sigma} d\Sigma j_A^0$, where the integral is computed on any constant-time Cauchy hypersurface Σ . For the sake of clarity, let us focus on electromagnetic backgrounds, although similar arguments apply also for gravitational backgrounds—except for some important differences that we unravel in this article. Classically, the chiral charge Q_A of a Dirac field measures the difference in the amplitude of the two helical components of the field. Quantum mechanically, this quantity translates to the difference in the number of positive and negative helicity particles, together with possible contributions from “vacuum polarization.” The charge Q_A is strictly conserved in the classical theory, but it is not quantum mechanically due to nonconservation of the current (2). This can be easily shown by considering any two Cauchy hypersurfaces, Σ_{in} and Σ_{out} , and noticing that the change in the vacuum expectation value of Q_A between Σ_{in} and Σ_{out} is equal to the integral of $\langle \nabla_a j_A^a \rangle$ in the four-dimensional region R bounded by Σ_{in} and Σ_{out} :

$$\begin{aligned} \int_R d^4x \langle \nabla_\mu j_A^\mu \rangle &= \int_R d^4x (\langle \partial_t j_A^0 \rangle + \langle \vec{\nabla} \cdot \vec{j}_A \rangle) \\ &= \int_R d^4x \langle \partial_t j_A^0 \rangle = \langle Q_A \rangle_{\Sigma_{\text{out}}} - \langle Q_A \rangle_{\Sigma_{\text{in}}}, \end{aligned} \quad (3)$$

where, in the second equality, the second term vanishes due to Stoke’s theorem and standard falloff conditions of the fields at spatial infinity. Hence, a nonzero value of the integral $\int_R d^4x \langle \nabla_a j_A^a \rangle$ implies the nonconservation of the chiral charge of the quantum fermionic field.

Our goal is to understand what characteristics the electromagnetic backgrounds must have to produce a nonzero value of this integral. It is easy to check that the pseudoscalar $F_{ab} \star F^{ab}$ appearing in (1) can be written as the divergence of a vector, $j_{\text{CS}}^a = 2A_b \star F^{ab}$, where A_a is the electromagnetic potential—CS stands for Chern-Simons. Repeating the steps used to produce Eq. (3), Eqs. (1) and (3) automatically imply that the fermionic chiral charge $\langle Q_A \rangle$ fails to be conserved if and only if the scalar $Q_{\text{CS}} = \int_{\Sigma} d\Sigma j_{\text{CS}}^0$ associated with the vector j_{CS}^a changes between Σ_{in} and Σ_{out} :

$$\langle Q_A \rangle_{\Sigma_{\text{out}}} - \langle Q_A \rangle_{\Sigma_{\text{in}}} = -\frac{q^2 \hbar}{8\pi^2} [Q_{\text{CS}}|_{\Sigma_{\text{out}}} - Q_{\text{CS}}|_{\Sigma_{\text{in}}}] . \quad (4)$$

This simple observation provides an interesting strategy to classify electromagnetic backgrounds that are able to trigger an anomalous nonconservation of the chiral charge

of fermionic quantum fields propagating thereon.¹ This strategy was initiated in [15,16] for both electromagnetic and gravitational backgrounds, where some aspects of the scalar Q_{CS} were analyzed for asymptotically flat spacetimes, in which the hypersurfaces Σ_{in} and Σ_{out} can be chosen to be past (\mathcal{I}^-) and future null infinity (\mathcal{I}^+), respectively. This is a natural choice when studying massless quantum fields. It was shown that, at these limiting surfaces, the scalar Q_{CS} receives a contribution from the net helicity of the radiative content of the electromagnetic and gravitational fields. This implies that, if there are sources in the bulk emitting helical or circularly polarized radiation—in gravity, this happens, for instance, in the coalescence of a large family of binary black hole mergers [15,17]—there is a net change of Q_{CS} between \mathcal{I}^- and \mathcal{I}^+ , which induces a change in the chiral charge $\langle Q_A \rangle$ of fermionic quantum fields propagating thereon. This is a profound relation between the radiative content of the background field and the chiral charge of quantum fields. We emphasize that this is a quantum effect; classically, $\langle Q_A \rangle$ is strictly conserved for massless fields, regardless of the properties of the electromagnetic and gravitational backgrounds.

This article unravels another contribution to Q_{CS} —which, consequently, also acts as a source of fermionic helicity $\langle Q_A \rangle$ —originated in the existence of certain electromagnetic infrared or “soft” charges. Infrared charges have received a good deal of attention in the recent past, due to their theoretical importance in the study of the S-matrix in quantum electrodynamics and quantum gravity, and due to their connection with soft theorems (see the reviews [18,19] and references therein). On the other hand, nonzero infrared charges indicate the generation of “memory effects” in physical systems. To give an example, test charged particles can experience a permanent change in their velocity (a “kick”) after the passage of electromagnetic waves [20]. In electrodynamics, this was the first example of memory reported in the literature. Other memory effects have been identified in recent years (see for instance [21,22] for an effect related with the helicity of radiation, and references therein).

Therefore, the results of this article can be interpreted as another type of memory effect produced by infrared charges, now on *test quantum fields* rather than on classical test particles. Quite interestingly, we find that this new manifestation of electromagnetic memory effect does not occur for the gravitationally-induced chiral anomaly.

¹For Yang-Mills fields, the standard strategy consists in looking for instanton solutions in Euclidean space. However, there are no instanton solutions of Maxwell (Abelian) equations in four dimensions. Furthermore, it is useful to work directly within the framework of asymptotically flat spacetimes, since it captures the full causal structure of physical (Lorentzian) spacetimes.

The rest of this article is organized as follows. Section II introduces a simple example of pedagogical value: a massless Dirac field in $1+1$ -dimensional flat spacetime coupled to an electromagnetic background. Section III contains a brief summary of the asymptotic properties of the electromagnetic field at past and future null infinity, including the notion of soft charges and memory effect in this framework; readers already familiar with the notation can skip this section. Section IV contains the main analysis of this article, where the contribution of soft electromagnetic charges for the Adler-Bell-Jackiw anomaly is derived. This section also includes a simple example of an electromagnetic configuration for which the relevant infrared charges are different from zero. The gravitational case is discussed in Sec. V, and Sec. VI closes the paper with a few conclusions and remarks.

Throughout this paper, we use geometric units in which $G = c = 1$, and we keep \hbar explicit in our equations to emphasize quantum effects. The metric signature is chosen to be $(-, +, +, +)$; ∇_a represents the Levi-Civita connection; the Riemann tensor is defined by $2\nabla_{[a}\nabla_{b]}v_c = R_{abc}^{\quad d}v_d$ for any covector v_d . Unless otherwise stated, all tensors will be assumed to be smooth.

II. CHIRAL ANOMALY IN TWO DIMENSIONS

This section discusses a massless, charged Dirac field in a $1+1$ -dimensional flat spacetime, propagating on a homogeneous electric background with finite support in time. To make our arguments simpler, we assume that the spacetime manifold is $\mathbb{R} \times \mathbb{S}^1$, i.e., the spatial dimension has been compactified to the circle. The electric field is assumed to be strong enough to make the backreaction of the quantum field negligibly small. This setup has great pedagogical value to illustrate some of the main messages of this paper, particularly the relation between the chiral fermion anomaly and the memory effect. We also discuss the relation of these concepts with spontaneous quantum particle-pair creation.

In a $1+1$ -dimensional spacetime, F_{ab} has only one independent component—the electric field— $F_{ab} = E\epsilon_{ab}$, where ϵ_{ab} is the totally antisymmetric tensor. Among the two Maxwell equations, $dF = 0$ is a trivial identity in $1+1$ dimensions, since it involves antisymmetrizing three indices which can only take two different values. The other set of Maxwell equations, $d^*F = {}^*j$, lead to $\epsilon^{ab}\partial_b E = -j^a$. These equations imply that the electric field cannot vary out of the support of the sources j , neither in space nor in time. Hence, in $1+1$ dimensions there are neither magnetic fields nor electric waves.

Let us consider a fixed, time-dependent electric field that is uniform in space. This can be generated by a time-dependent current of the form $j^a = (0, j(t))$. We further assume that the electric field is different from zero only during a finite interval, $E(t) \neq 0$ for $t_{\text{in}} < t < t_{\text{out}}$. Despite

the fact that $dF = 0$ is an identity and there is no magnetic field, it is still useful to introduce a vector potential, $F = dA$, in terms of which the electric field reads $E = \partial_t A_\theta - \partial_\theta A_t$. A gauge transformation changes $A_a \rightarrow A_a + \partial_a \alpha$, with α a continuous function in the spacetime manifold. We can always use this freedom to make $A_t = 0$, so that, under this gauge choice, $E = \partial_t A_\theta$.

In $1+1$ dimensions, the expression (1) for the chiral anomaly is replaced by [23]

$$\langle \nabla_a j_A^a \rangle = \frac{q\hbar}{2\pi} \epsilon^{ab} F_{ab}. \quad (5)$$

The right-hand side can be written as $\frac{q\hbar}{2\pi} \nabla_a j_{\text{CS}}^a$, where $j_{\text{CS}}^a = 2\epsilon^{ab} A_b$. Although this vector is not gauge invariant, its divergence, as well as the scalar $Q_{\text{CS}}(t) \equiv \int_t dx j_{\text{CS}}^0(t, x)$, are both gauge invariant.² Following the argument described in the Introduction, Eq. (5) implies that the change of the chiral charge $\langle Q_A \rangle(t) = \int_0^L d\theta \langle j_A^0 \rangle(t, \theta)$ from t_{in} to t_{out} can be written, for any quantum state, as

$$\langle Q_A \rangle(t_{\text{out}}) - \langle Q_A \rangle(t_{\text{in}}) = \frac{q\hbar}{2\pi} [Q_{\text{CS}}(t_{\text{out}}) - Q_{\text{CS}}(t_{\text{in}})], \quad (6)$$

where $Q_{\text{CS}} \equiv \int_0^L d\theta j_{\text{CS}}^0 = 2 \int_{\mathbb{S}^1} A_a d\ell^a$. As mentioned above, this quantity is manifestly gauge invariant. Recall also that this scalar is purely electric, i.e., it does not know anything about the Dirac field. From this, we have

$$\begin{aligned} Q_{\text{CS}}(t_{\text{out}}) - Q_{\text{CS}}(t_{\text{in}}) &= 2 \int_0^L d\theta (A_\theta(t_{\text{out}}) - A_\theta(t_{\text{in}})) \\ &= 2L \int_{t_{\text{in}}}^{t_{\text{out}}} dt E(t), \end{aligned} \quad (7)$$

where in the last equality we have used that $E = \partial_t A_\theta$, that the electric field is homogeneous, and that L is the length of the spatial sections. Hence, Eq. (6) tells us that the anomalous nonconservation of the chiral charge $\langle Q_A \rangle$ is dictated by the value of the time integral of the electric field

$$\langle Q_A \rangle(t_{\text{out}}) - \langle Q_A \rangle(t_{\text{in}}) = \frac{q\hbar}{\pi} L \int_{t_{\text{in}}}^{t_{\text{out}}} dt E(t). \quad (8)$$

This result shows that the vacuum expectation value $\langle Q_A \rangle$ “keeps memory” of the past history of the electric field. In particular, the effect of switching on an electric field for some period of time $t_{\text{in}} < t < t_{\text{out}}$ can leave a residual, permanent helicity contribution on the vacuum state of the quantum field (quantified by the value of $\langle Q_A \rangle$ at late

²A more rigorous derivation of this vector gives $j_{\text{CS}}^a = 2\epsilon^{ab}(A_b - A_b^0)$ with $\nabla_{[a} A_{b]}^0 = 0$, which includes an auxiliary potential A_b^0 that makes j_{CS}^a gauge invariant. Since this extra term does not affect physical quantities, we can simply take $A_b^0 = 0$, as customary in the literature.

times), *even* after switching off completely the external field. One can think about this residual helicity as the way the quantum field retains information about the past influence of the electric background.

The integral on the right-hand side above also features in the memory effect found for classical particles [20]. A test charged particle in our background would suffer a permanent change in its velocity after the passage of this electromagnetic pulse if and only if $\int_{t_{\text{in}}}^{t_{\text{out}}} dt E(t) \neq 0$. From this point of view, the permanent change in the vacuum expectation value $\langle Q_A \rangle$ found above can be thought of as another manifestation of the electromagnetic memory effect, but now on quantum fields.

It may be surprising that, despite the fact that the external electric field vanishes, the quantum system does not return to its original configuration. To better understand this effect, one has to resort to the electromagnetic potential, which makes manifest that the memory actually originates from transitions between inequivalent vacua of the electric field (hence, the background does not really return to the same exact configuration either). To see this, recall that our electric background evolves from a vacuum configuration, $E(t_{\text{in}}) = 0$, to another vacuum configuration $E(t_{\text{out}}) = 0$. Classically, the two electric vacuum states are equivalent, but quantum mechanically they may not be.³ In particular, note that the change in potential from t_{in} to t_{out} , $A_a(t_{\text{out}}) - A_a(t_{\text{in}})$, is nontrivial. This can be seen from the fact that the loop integral $\int_{\mathbb{S}^1} d\ell^a (A_a(t_{\text{out}}) - A_a(t_{\text{in}}))$, which is gauge invariant, is different from zero if and only if $\int_{t_{\text{in}}}^{t_{\text{out}}} dt E(t) \neq 0$. But since the electric field vanishes at early and late times, $A_a(t_{\text{out}})$ can only differ from the initial potential $A_a(t_{\text{in}})$ by a residual gauge transformation, left by the dynamical evolution of the electric field. A straightforward calculation shows that this gauge transformation is given by $A_a(t_{\text{out}}) = A_a(t_{\text{in}}) + \partial_a \alpha$ with $\alpha(\theta) = \theta \int_{t_{\text{in}}}^{t_{\text{out}}} dt E(t) + \alpha_0$, for some constant α_0 . However, this is not an ordinary gauge transformation because $\alpha(\theta)$ is not a continuous function on \mathbb{S}^1 (because $\alpha(L) \neq \alpha(0)$). Instead, it belongs to the family of so-called “large” gauge transformations [23], which carry physical implications, and which can be used to label inequivalent notions of vacuum states of the quantum electromagnetic theory.⁴

In summary, the passage of an electric pulse with $\int_{t_{\text{in}}}^{t_{\text{out}}} dt E(t) \neq 0$ induces a large gauge transformation in

the vector potential, which produces a memory effect, not only on classical charged particles but also on the states of quantum fermion fields. As will be discussed below, in $3 + 1$ dimensions there is another contribution to the chiral anomaly coming from the radiative content of the electromagnetic field; such contribution does not arise in $1 + 1$ due to the absence of electromagnetic radiation.

The permanent change in the vacuum expectation value of the chiral charge $\langle Q_A \rangle$, described on the left-hand side of (8), can also be understood in terms of the standard notion of electromagnetic memory for particles. Heuristically, virtual charged particles populating the quantum vacuum would suffer a permanent change in its velocity after switching on this electric pulse, provided the integral $\int_{t_{\text{in}}}^{t_{\text{out}}} dt E(t)$ does not vanish. In $1 + 1$ dimensions these charges can only propagate in two directions, left or right. Therefore, positive charges suffer a “kick” in the direction of the electric field, while negative charges are kicked in the opposite direction. Both particles in the pair have the same helicity. If the kick is strong enough, it will turn virtual charges into physical excitations out of the quantum vacuum. This results in a net creation of helicity, which explains the permanent change of the quantum state or of the chiral charge $\langle Q_A \rangle$.

This heuristic picture can be made rigorous through a calculation of particle-pair creation using Bogoliubov coefficients. We finish this section with a brief allusion to this. If the electric field is nonzero only during a finite interval, $t_{\text{in}} < t < t_{\text{out}}$, we can define natural “in” and “out” notions of vacua and particles. The question of interest is, if the field is prepared in the “in” vacuum before t_{in} , and evolved until a time after t_{out} , what is the number of “out” quanta in the final state?

This question can be answered without much difficulty in the case in which the electric field is uniform at all times (Appendix A contains a detailed derivation of this and of the general case of an electric field that varies both in space and time). As before, let us work in a gauge in which the vector potential is purely spatial, $A_t = 0$. Without loss of generality, we can also consider $A_\theta(t) = 0$ for $t < t_{\text{in}}$. Let $A_0 = \int_{t_{\text{in}}}^{t_{\text{out}}} E(t) dt$ denote then the value of A_θ at late times, after t_{out} . In short, a nonzero value of A_0 induces a permanent frequency shift between the “in” and “out” basis of solutions of the field equations, which define the “in” and “out” vacua, respectively. Namely, for modes with

³The most prominent example of this is the Aharonov-Bohm effect [24].

⁴More precisely, when the potential A_a is viewed as a gauge connection on a $U(1)$ principle bundle over $\mathbb{R} \times \mathbb{S}^1$, we can speak of infinitesimal gauge transformations, as well as of global or finite gauge transformations. In the temporal gauge fixing $A_t = 0$, a finite gauge transformation, $A_a \rightarrow A_a - ig^{-1} \nabla_a g$, is determined by a continuous map $g: \mathbb{S}^1 \rightarrow \mathbb{S}^1$, given in a local coordinate system by $g(\theta) = e^{i\alpha(\theta)}$. Continuous maps on \mathbb{S}^1 can be divided in different (homotopy) classes, where two elements of the same class can be deformed continuously into each other. The classification of continuous maps is determined by the first homotopy group, $\Pi(\mathbb{S}^1) \simeq \mathbb{Z}$, which shows that each class of gauge functions is labeled by an integer. This is easy to infer from the requirement that g is continuous, because it demands $\alpha(L) - \alpha(0) = 2\pi n$, for $n \in \mathbb{Z}$. Two gauge functions g, g' belonging to different classes cannot be deformed continuously into each other. An ordinary gauge transformation is a gauge transformation g that belongs to the trivial class or $n = 0$, while gauge functions with $n \neq 0$ lead to “large” gauge transformations [23].

spatial dependence $e^{ik\theta}$, with $k \in (2\pi/L)n$ and $n \in \mathbb{Z}$, the “in” modes oscillate with frequency $\omega_{\text{in}} = k$, while “out” modes oscillate with frequency $\omega_{\text{out}} = k + qA_0$. Because k is discretized (due to the compactness of the spatial sections) there is a finite number of modes within the frequency interval $(0, q|A_0|)$. Namely, there are $[q|A_0| \frac{L}{2\pi}]$ modes within this interval, where the square brackets denote integer part.

This shift in frequency automatically implies that the evolution creates a number $[q|A_0| \frac{L}{2\pi}]$ of fermion-antifermion pairs. Because of linear momentum conservation (note that the background is homogeneous), antifermions (positive charges) move in the direction of $E(t)$, while fermions (negative charges) move in the opposite direction. However, helicity is not conserved in this process. Fermions moving to the right (left), and antifermions moving to the left (right), both have negative (positive) helicity. As a result, both members of the pair have positive \hbar (negative $-\hbar$) helicity if $A_0 > 0$ ($A_0 < 0$). The total helicity carried by the excited pairs is $2\hbar[qA_0 \frac{L}{2\pi}] = 2\hbar[q \int_{t_{\text{in}}}^{t_{\text{out}}} E(t) dt \frac{L}{2\pi}]$. This agrees, except for the noninteger part, with the prediction for $\langle Q_A \rangle(t_{\text{out}}) - \langle Q_A \rangle(t_{\text{in}})$ given in Eq. (8). The difference is due to the “vacuum polarization,” i.e. the helicity leftover in the “out” vacuum, which did not reach the threshold to excite another pair.

Although, for the sake of pedagogy, in this section we have restricted to uniform, background electric fields, all the arguments generalize to arbitrary functions $E(t, \theta)$. Appendix A contains information about this generalization and further details which have been omitted in this section.

III. ASYMPTOTIC STRUCTURE OF THE ELECTROMAGNETIC FIELD AND INFRARED CHARGES: A BRIEF REVIEW

The rest of this paper will focus on the chiral anomaly in asymptotically flat spacetimes in $3 + 1$ dimensions. The presence of electromagnetic radiation, not present in $1 + 1$ dimensions, makes it more convenient to use past and future null infinity for the initial and final Cauchy hypersurface of zero-rest mass fields. This section contains a brief summary of tools concerning the asymptotic structure of the electromagnetic field at null infinity and infrared charges. These tools are well known [18,25–27], and the reader familiar with them can skip this section.

A. Review on the asymptotic structure of the electromagnetic field

The electromagnetic radiation generated by charges and currents can be rigorously studied within the framework of asymptotically flat spacetimes [25–27]. This framework makes use of the notions of conformally compactified spacetimes introduced by Penrose in the 1960s [28].

Let $(\mathbb{R}^4, \hat{\eta}_{ab})$ represent the physical, Minkowski spacetime, and let (M, η_{ab}) denote an extended (unphysical)

spacetime obtained from $(\mathbb{R}^4, \hat{\eta}_{ab})$ by an ordinary conformal compactification, i.e. by the addition of “points at infinity.”⁵ More precisely, the new metric is obtained from the physical one by a conformal transformation $\eta_{ab} = \Omega^2(x)\hat{\eta}_{ab}$, while the new manifold is constructed by attaching smoothly a null boundary \mathcal{J} to the physical manifold, $M = \mathbb{R}^4 \cup \mathcal{J}$. Locally, \mathcal{J} corresponds to the hypersurface $\Omega = 0$ and has null normal $\eta^{ab}\nabla_b\Omega \neq 0$. From a physical viewpoint, the elements of \mathcal{J} represent the “points of (null) infinity,” i.e. the points that can be asymptotically reached by following radial, null geodesics in the physical spacetime. The boundary \mathcal{J} is made of two portions, past (\mathcal{J}^-) and future (\mathcal{J}^+) null infinity. In the following, we will focus on \mathcal{J}^+ . The construction is similar for \mathcal{J}^- .

For example, in a Bondi-Sachs coordinate system $\{u, r, \theta, \phi\}$, where $u = t - r$ is the standard retarded time, the Minkowski metric reads $d\hat{s}^2 = -du^2 + 2dudr + r^2d\omega^2$ and one uses $\Omega = 1/r$ to obtain $ds^2 = -\Omega^2 du^2 + 2dud\Omega + d\omega^2$ after the conformal transformation mentioned above. The restriction of this line element to the $\Omega = 0$ hypersurface gives a well-defined (although degenerate) metric. The limit $r \rightarrow \infty$ keeping u, θ, ϕ constant follows the geodesics of outgoing radiation propagating to future null infinity, getting to $\Omega = 0$ in finite time as measured by the unphysical metric. The extended manifold is obtained then by including all these limiting points $\{u, \Omega = 0, \theta, \phi\}$ to the original manifold, and future null infinity is described then by the submanifold $\mathbb{R} \times \mathbb{S}^2$.

This framework makes it possible to study the behavior of the electromagnetic field in a neighborhood of infinity (which in this case is simply a boundary of the spacetime manifold) using standard techniques in differential geometry. To see this, let us first note that the electromagnetic field tensors are conformal invariant, $\hat{F}_{ab} = F_{ab}$, $\hat{A}_a = A_a$. These tensors are well defined in the entire extended spacetime, including at the boundary \mathcal{J}^+ . The electromagnetic field F_{ab} has six independent components. In a Newman-Penrose basis $\{n^a, \ell^a, m^a, \bar{m}^a\}$ [29], where typically one takes $\ell_a = -\nabla_a u$ as the vector tangent to outgoing null geodesics, the six electric and magnetic components of F_{ab} can be captured in the following three complex scalars

$$\Phi_2 := F_{ab}n^a\bar{m}^b, \quad (9)$$

$$\Phi_1 := \frac{1}{2}[F_{ab}n^a\ell^b + F_{ab}m^a\bar{m}^b], \quad (10)$$

$$\Phi_0 := F_{ab}m^a\ell^b. \quad (11)$$

If we assume smooth fields, the Peeling theorem guarantees that these scalars admit the following Taylor expansion in Ω in a neighborhood of future null infinity [30]:

⁵We shall use the hat symbol for any tensor and quantity intrinsic to the physical spacetime.

$$\Phi_2(u, \Omega, \theta, \phi) = \Phi_2^0(u, \theta, \phi) + \Omega \Phi_2^1(u, \theta, \phi) + \dots, \quad (12)$$

$$\Phi_1(u, \Omega, \theta, \phi) = \Phi_1^0(u, \theta, \phi) + \Omega \Phi_1^1(u, \theta, \phi) + \dots, \quad (13)$$

$$\Phi_0(u, \Omega, \theta, \phi) = \Phi_0^0(u, \theta, \phi) + \Omega \Phi_0^1(u, \theta, \phi) + \dots, \quad (14)$$

where we denote $\Phi_2^0(u, \theta, \phi) \equiv \Phi_2(u, \Omega = 0, \theta, \phi)$ and similarly for $\Phi_1^0(u, \theta, \phi)$ and $\Phi_0^0(u, \theta, \phi)$. These fields encode all the information about the electromagnetic field at \mathcal{J}^+ . They are, however, not independent. Using Maxwell's equations, one finds

$$\partial_u \Phi_1^0 = \eth \Phi_2^0, \quad (15)$$

$$\partial_u \Phi_0^0 = \eth \Phi_1^0, \quad (16)$$

where $\partial_u f = n^a \nabla_a f$ for any function f (this is a consequence of the Newman-Penrose normalization $n^a \ell_a = -1$), and where \eth is a spin-weighted derivative operator [30], defined by $V_1^b V_2^c \dots m^a \nabla_a T_{bc\dots} = \eth(V_1^b V_2^c \dots T_{bc\dots})$, for arbitrary tensors V_i^a and $T_{bc\dots}$. These equations determine the evolution of the scalars Φ_0^0 and Φ_1^0 along the retarded time u in \mathcal{J}^+ , upon giving initial conditions, and also some input for Φ_2^0 . In contrast, the dynamics of Φ_2^0 along u is not determined by Maxwell equations. This scalar serves as the free data for a characteristic value formulation of Maxwell theory at \mathcal{J}^+ .

By switching back to the original physical spacetime, with the appropriate conformal rescaling of the Newman-Penrose vectors, one can see that $\hat{\Phi}_2 \sim O(r^{-1})$ and $\hat{\Phi}_1 \sim O(r^{-2})$. From this, one identifies the scalar Φ_2^0 as describing the two radiative degrees of freedom of the electromagnetic field, while Φ_1^0 represents the Coulombic part of the field. In fact, the total energy flux radiated to \mathcal{J}^+ is given by $F = \int dud\mathbb{S}^2 T_{ab} n^a n^b = \int dud\mathbb{S}^2 |\Phi_2^0|^2$, where T_{ab} is the energy-momentum tensor, and therefore is entirely determined from Φ_2^0 . Similarly, the electric charge of sources in the bulk can be determined from \mathcal{J}^+ using Gauss's law as $Q = \frac{1}{2\pi} \int d\mathbb{S}^2 \text{Re} \Phi_1^0(u, \theta, \phi)$, and is completely determined from Φ_1^0 . Using Maxwell equations in \mathcal{J}^+ , it is straightforward to check that $\partial_u Q = 0$, reflecting the conservation of the electric charge. Note also that the requirement of finite energy flux at \mathcal{J}^+ , $F < \infty$, requires $\Phi_2^0(u, \theta, \phi) \rightarrow 0$ as $u \rightarrow \pm\infty$.

In terms of an electromagnetic potential, one introduces the scalars $A_2 := A_a \bar{m}^a$, $A_1 = A_a n^a$, $A_0 = A_a \ell^a$. The falloff conditions of the potential for large r is not given beforehand from the theory and requires some input. Physical considerations require that these components admit an asymptotic series with leading behavior $O(r^{-1})$ [31,32]. Using $F = dA$, one can obtain the following formulas,

valid at future null infinity \mathcal{J}^+ :

$$\Phi_2^0 = \dot{A}_2^0 - \bar{\eth} A_1^0, \quad (17)$$

$$\text{Im} \Phi_1^0 = \text{Im} \bar{\eth} A_2^0, \quad (18)$$

where dot denotes derivative with respect to retarded time u . Furthermore, by integrating Maxwell equations at \mathcal{J}^+ , one can further obtain

$$\text{Re} \Phi_1^0 = \text{Re} \bar{\eth} A_2^0 - \int_{u_0}^u du' \bar{\eth} \bar{\eth} A_1^0 + G(u_0, \theta, \phi), \quad (19)$$

where $G(u_0, \theta, \phi)$ arises as an integration factor. From (9) we see that the two electromagnetic radiative degrees of freedom are distributed between A_2^0 and A_1^0 . But these are three real-valued scalars, so there is, as expected, some gauge redundancy in the description. A useful gauge fixing is $A_1^0 \equiv A_a n^a = 0$. With this gauge choice, the real and imaginary parts of A_2^0 represent the two radiative degrees of freedom, electric and magnetic respectively. The Coulombic aspects of the field are all encoded in $G(u_0, \theta, \phi)$, in particular $Q = \frac{1}{2\pi} \int_{\mathbb{S}^2} G(u_0, \theta, \phi)$.

B. Electromagnetic soft charges and the memory effect

The phenomenon of memory effect is well known in Maxwell's theory [20]. The most prominent example is a charged pointlike particle of initial velocity \vec{v}_1 that suffers a “kick” and changes its direction of propagation to \vec{v}_2 after the passage of an electromagnetic pulse. This is an example of electric-type memory. In the intermediate process, the charged particle emits radiation by Bremsstrahlung; the properties of the emitted radiation carry information about this effect.

At future null infinity, the phenomenon of electromagnetic memory is encoded in the following quantities:

$$q_\alpha = \int d\mathbb{S}^2 \alpha(\theta, \phi) (\Phi_1^0(\infty, \theta, \phi) - \Phi_1^0(-\infty, \theta, \phi)), \quad (20)$$

where α is a smooth real function on the sphere \mathbb{S}^2 . The complex numbers q_α are called soft charges, and they measure permanent changes in the multipolar structure of the Coulombic part of the electromagnetic field after some process. From (15), one infers that $q_\alpha \neq 0$ only if $\Phi_2(u, \theta, \phi) \neq 0$, i.e. only, if there is a flux of electromagnetic radiation reaching infinity. When this happens, one says that the electromagnetic field keeps memory on the radiation flux emitted to infinity in the past.

The relation of the charges q_α and the radiation reaching \mathcal{J}^+ can be explicitly shown by using Maxwell equations to

replace Φ_1^0 by Φ_2^0 in (20):

$$\begin{aligned} q_\alpha &= \int d\mathbb{S}^2 du \alpha(\theta, \phi) \dot{\Phi}_1^0(u, \theta, \phi) \\ &= \int d\mathbb{S}^2 du \alpha \delta \Phi_2^0(u, \theta, \phi) \\ &= - \int d\mathbb{S}^2 du \delta \alpha(\theta, \phi) \Phi_2^0(u, \theta, \phi). \end{aligned} \quad (21)$$

Expanding in spin-weighted spherical harmonics, this can be further simplified as

$$\begin{aligned} q_\alpha &= \sum_{\ell m} (-1)^m \sqrt{\ell(\ell+1)} \alpha_{\ell m} \int_{-\infty}^{+\infty} du \Phi_{2\ell m}^0(u) \\ &\equiv \sum_{\ell m} \tilde{\alpha}_{\ell m} q_{\ell m}. \end{aligned} \quad (22)$$

The problem is reduced to study the basis $q_{\ell m} = \int_{-\infty}^{+\infty} du \Phi_{2\ell m}^0(u)$ of charges. Note that, for $\ell = 0$, i.e. when $\alpha(\theta, \phi) = \text{const}$, the soft charge is identically zero. This is consistent with the fact that the monopole of the electromagnetic field (the electric charge) is conserved and cannot be radiated away. In contrast, dipolar and higher order structure ($\ell \geq 1$) can be radiated away. That phenomena is encoded in $q_{\ell m} \neq 0$ for $\ell > 0$.

As argued in the previous subsection, finiteness of energy fluxes require that $\Phi_{2\ell m}^0(u)$ belongs to $L^2(\mathbb{R}, \mathbb{C})$, which implies that it admits a Fourier transform on \mathcal{J}^+ :

$$\tilde{\Phi}_{2\ell m}^0(\omega) = \int_{-\infty}^{+\infty} du \Phi_{2\ell m}^0(u) e^{-i\omega u}. \quad (23)$$

This automatically implies that the charges $q_{\ell m}$ are simply the “zero mode” of $\tilde{\Phi}_{2\ell m}^0$

$$q_{\ell m} = \tilde{\Phi}_{2\ell m}^0(0). \quad (24)$$

Therefore, only the zero-frequency modes of the emitted electromagnetic radiation leave a memory on the multipolar structure of the field. This is the reason why these charges are called “soft,” as they are associated with “soft photons” [19,33].

These charges are intrinsically associated to asymptotic symmetries of Maxwell theory. One way of looking into this is by considering the phase space Γ of the electromagnetic degrees of freedom at \mathcal{J}^+ . This phase space is made of pairs of canonically conjugate fields⁶ (A_a, \mathcal{E}^a) , where $A_a = -A_1^0 \ell_a + A_2^0 m_a + \bar{A}_2^0 \bar{m}_a$ and $\mathcal{E}^a = -2\text{Re}\Phi_1^0 n^a + \Phi_2^0 m^a + \bar{\Phi}_2^0 \bar{m}^a$.⁷ Γ can be endowed

⁶In the Newman-Penrose basis introduced above, notice that the tangent space at each point of future null infinity is spanned by the three vectors $\{n^a, m^a, \bar{m}^a\}$, while the cotangent space is spanned by the three covectors $\{\ell_a, m_a, \bar{m}_a\}$.

⁷No gauge condition on A_a has been imposed at this level.

with the structure of an infinite-dimensional Banach manifold. The usual symplectic structure for Maxwell theory can be written on future null infinity as

$$\begin{aligned} \Omega((A_a^{(1)}, \mathcal{E}^{(1)a}), (A_a^{(2)}, \mathcal{E}^{(2)a})) \\ = \int_{\mathbb{R} \times \mathbb{S}^2} du d\mathbb{S}^2 \left(A_a^{(1)} \mathcal{E}^{(2)a} - A_a^{(2)} \mathcal{E}^{(1)a} \right). \end{aligned} \quad (25)$$

Together with suitable falloff conditions at $u \rightarrow \pm\infty$ required to make this integral well defined, the pair (Γ, Ω) defines the phase space for the radiative degrees of freedom of Maxwell theory.

Now, consider the (restricted) family of gauge transformations $A_a \rightarrow A_a + D_a \alpha$, with $\alpha = \alpha(\theta, \phi)$. This transformation is generated in phase space by the quantity

$$\begin{aligned} Q_\alpha &\equiv \frac{1}{2} \Omega((A_a, \mathcal{E}^a), (D_a \alpha, 0)) \\ &= -\frac{1}{2} \int_{\mathbb{R} \times \mathbb{S}^2} du d\mathbb{S}^2 (D_a \alpha) \mathcal{E}^a \\ &= -\frac{1}{2} \int_{\mathbb{R} \times \mathbb{S}^2} du d\mathbb{S}^2 D_a(\alpha \mathcal{E}^a) \\ &= \int_{\mathbb{R} \times \mathbb{S}^2} du d\mathbb{S}^2 n^a D_a(\alpha \text{Re}\Phi_1^0), \end{aligned}$$

where in the second equality we have used Maxwell equations (15) to write $D_a \mathcal{E}^a \propto \text{Re}\dot{\Phi}_1^0 - \text{Re}\delta\Phi_2^0 = 0$. Recalling that $\partial_u = n^a D_a$, the right-hand side of this equation happens to be equal to the real part of the soft charges defined in (20). (A similar analysis using a “dual” potential Z_a produces the imaginary part of the soft charges.)

This observation tells us that soft charges q_α can be identified with the generators of gauge transformations in the radiative phase space. Since soft charges can be different from zero, one concludes that transformations $A_a \rightarrow A_a + D_a \alpha$ in \mathcal{J}^+ are actual symmetries of our phase space (Γ, Ω) , rather than mere gauge transformations. Therefore, they have physical significance (which is, precisely, the electromagnetic memory). From the viewpoint of the bulk, these are gauge transformations that do not vanish at infinity. To distinguish them from ordinary gauge transformations, they are called “large” gauge transformations. The set of all large gauge transformations constitutes the infinite-dimensional, asymptotic symmetry group of Maxwell theory.

IV. THE CHIRAL ANOMALY INDUCED BY AN ELECTROMAGNETIC BACKGROUND

This section contains the main results of this article. We consider a quantum, massless Dirac field propagating in

Minkowski spacetime in $3 + 1$ dimensions with metric η_{ab} , coupled to an electromagnetic field F_{ab} . The spin $1/2$ field is treated as a test field, i.e. we neglect its backreaction on the electromagnetic and spacetime backgrounds. This external electromagnetic field is assumed to be generated by some distribution of electric charges and currents, that are smooth and confined in space, but otherwise arbitrary. To keep the parallelism with the $1 + 1$ -dimensional chiral anomaly discussed in the previous section, the sources will be “switched on” only for a finite amount of time, in the sense that they become stationary at sufficiently late and early times. All possible electromagnetic waves are radiated during a finite period of time.

As discussed above, the electromagnetic field can induce a change in the helicity of the fermionic field due to the chiral anomaly. Our starting point is expression (3) for the change of the chiral charge of the quantum field

$$\begin{aligned} \langle Q_A \rangle_{\Sigma_{\text{out}}} - \langle Q_A \rangle_{\Sigma_{\text{in}}} &= \int_R d^4x \sqrt{-\eta} \langle \nabla_a j_A^a \rangle \\ &= -\frac{\hbar q^2}{8\pi^2} \int_R d^4x \sqrt{-\eta} F_{ab} \star F^{ab}. \end{aligned} \quad (26)$$

If we integrate over the entire spacetime manifold, $R = M \simeq \mathbb{R}^4$, then Σ_{in} and Σ_{out} correspond to past and future null infinity, respectively. This choice makes it possible to use the machinery summarized in the previous section to disentangle the properties of the electromagnetic field that can make the right-hand side different from zero. This problem was worked out in [16], where it was found that, assuming no incoming electromagnetic radiation from past null infinity, the right-hand side of (26) can be written in terms of boundary data on future null infinity as

$$\langle Q_A \rangle_{\mathcal{J}^+} - \langle Q_A \rangle_{\mathcal{J}^-} = \frac{\hbar q^2}{4\pi^2} \int_{-\infty}^{\infty} du \int d\mathbb{S}^2 \text{Im}\{(A_2^0 - \eth\alpha_0) \tilde{\Phi}_2^0\}. \quad (27)$$

Here, α_0 is a smooth real-valued function on the sphere and $\eth\alpha_0$ is a pure gauge potential (i.e. it produces no electromagnetic field, $\Phi_2^0 = 0$). This expression was derived in the gauge $A_1^0 = 0$. Notice that a nonzero value is obtained in the integral (27) because of the weak decay behavior of the radiative solutions of Maxwell equations in a neighborhood of future null infinity: $A_2 \sim 1/r$, $\Phi_2 \sim 1/r$ (recall the discussion earlier). These two radial factors compensate the r^2 factor in the integral measure.⁸

To analyze the physical interpretation of the right-hand side of the previous equation, it is convenient to work with a compactified retarded coordinate u . We will consider $u \in [-L/2, L/2]$ and let $L \rightarrow \infty$ at the end of the

⁸There are no contributions to (26) from spatial infinity, nor from future or past timelike infinities, because the product of the electromagnetic field and its potential decays too fast [as $O(r^{-3})$] in those directions; see [16] for details.

calculation. As explained above, if Φ_2^0 is the electromagnetic radiation field, the requirement that the energy flux across \mathcal{J}^+ is finite implies $\Phi_2^0(\cdot, \theta, \phi) \in L^2(\mathbb{R})$ for all $(\theta, \phi) \in \mathbb{S}^2$, and in particular $\Phi_2^0 \rightarrow 0$ as $u \rightarrow \pm\infty$. Therefore, we will consider functions $\Phi_2^0(\cdot, \theta, \phi) \in L^2((-L/2, L/2))$, for all $(\theta, \phi) \in \mathbb{S}^2$, with boundary conditions given by $\Phi_2^0(\pm \frac{L}{2}, \theta, \phi) = 0$. This will guarantee that $\Phi_2^0 \rightarrow 0$ as $u \rightarrow \pm\infty$ at the end of the calculation. Since the functions $\Phi_2^0(\cdot, \theta, \phi)$ happen to be periodic with period L , an orthonormal basis for $L^2((-L/2, L/2))$ is given by $\{\frac{e^{-i\omega_n L/2}}{\sqrt{L}} e^{-i\omega_n u}\}_{n \in \mathbb{Z}}$, where $\omega_n = \frac{2\pi}{L} n$, so one can expand in Fourier series:

$$\Phi_2^0(u, \theta, \phi) = \sum_{n=-\infty}^{+\infty} \tilde{\Phi}_2^0(\omega_n, \theta, \phi) e^{-i\omega_n \frac{L}{2}} \frac{e^{-i\omega_n u}}{\sqrt{L}}. \quad (28)$$

The inverse Fourier series is

$$\tilde{\Phi}_2^0(\omega_n, \theta, \phi) = e^{i\omega_n \frac{L}{2}} \frac{1}{\sqrt{L}} \int_{-L/2}^{L/2} \Phi_2^0(u, \theta, \phi) e^{+i\omega_n u} du. \quad (29)$$

The basis modes are orthonormal with respect to the L^2 norm:

$$\int_{-L/2}^{L/2} du \frac{1}{\sqrt{L}} e^{-i\omega_n u} (-1)^n \frac{1}{\sqrt{L}} e^{+i\omega_n u} (-1)^n = \delta_{nn'}. \quad (30)$$

The continuous limit will be recovered using the formula $\lim_{L \rightarrow \infty} \frac{1}{L} \sum_{n \in \mathbb{Z}} f(\frac{n}{L}) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} dw f(w)$.

In order to disentangle the potential contribution of IR charges to Eq. (27) we will make an explicit distinction between the zero frequency mode $\tilde{\Phi}_2^0(0, \theta, \phi) \neq 0$ and the rest of the modes. Let us then write the field and potential as

$$\Phi_2^0(u, \theta, \phi) = \sum_{n \neq 0} \tilde{\Phi}_2^0(\omega_n, \theta, \phi) (-1)^n \frac{e^{-i\omega_n u}}{\sqrt{L}} + \frac{1}{\sqrt{L}} \Phi_2^0(0, \theta, \phi), \quad (31)$$

$$\begin{aligned} A_2^0(u, \theta, \beta) &= \frac{1}{\sqrt{L}} \sum_{n \neq 0} \tilde{\Phi}_2^0(u, \theta, \phi) \frac{e^{-i\omega_n u} - e^{-i\omega_n u_0}}{-i\omega_n} (-1)^n \\ &+ \frac{u - u_0}{\sqrt{L}} \tilde{\Phi}_2^0(0, \theta, \phi) + A_2^0(u_0, \theta, \phi), \end{aligned} \quad (32)$$

where the second line is derived using $\Phi_2^0 = \dot{A}_2^0$, which is valid in the gauge $A_1^0 = A_a n^a = 0$. Next, we substitute this expansion in (27) and keep track of the contribution of the zero mode. The calculation is tedious, and is written in detail in Appendix B. We focus here in the result and its

physical meaning:

$$\begin{aligned} \langle Q_A \rangle_{\mathcal{J}^+} - \langle Q_A \rangle_{\mathcal{J}^-} &= \frac{\hbar q^2}{4\pi^2} \left[\frac{1}{2\pi} \int d\mathbb{S}^2 \int_{-\infty}^{\infty} d\omega \frac{|\tilde{\Phi}_2^0(\omega, \theta, \phi) - \tilde{\Phi}_2^0(0, \theta, \phi)|^2}{\omega} \right. \\ &\quad \left. + \text{Im}q_\alpha - \text{Re}q_\beta \right]. \end{aligned} \quad (33)$$

The first term in the right-hand side of this equation contains the contribution from electromagnetic radiation with nonzero frequencies reaching \mathcal{J}^+ ; the subtraction of $\tilde{\Phi}_2^0(0, \theta, \phi)$ removes the zero mode from the integral. This, in turn, makes the integrand finite in the limit $\omega \rightarrow 0$ (the integral is well defined for $\omega \rightarrow \pm\infty$ by Plancherel theorem). This term can be further expressed as [16]

$$\begin{aligned} &\int_{-\infty}^{\infty} d\omega \frac{|\tilde{\Phi}_2^0(\omega, \theta, \phi) - \tilde{\Phi}_2^0(0, \theta, \phi)|^2}{\omega} \\ &= \int_0^{\infty} \frac{d\omega}{\omega} \left(|\tilde{\Phi}_R(\omega, \theta, \phi)|^2 - |\tilde{\Phi}_L(\omega, \theta, \phi)|^2 \right), \end{aligned} \quad (34)$$

where $\Phi_R(\omega, \theta, \phi) := \tilde{\Phi}_2^0(\omega, \theta, \phi) - \tilde{\Phi}_2^0(0, \theta, \phi)$ defined for $\omega > 0$, and $\tilde{\Phi}_L(\omega, \theta, \phi) := \tilde{\Phi}_2^0(-\omega, \theta, \phi) - \tilde{\Phi}_2^0(0, \theta, \phi)$ defined for $\omega < 0$, describe right- and left-handed circularly polarized radiation, respectively. Expression (34) has a neat physical interpretation: it measures the net electromagnetic helicity radiated to \mathcal{J}^+ .

The second and third terms in (33) come entirely from the zero mode of the electromagnetic field, and correspond to two infrared charges of magnetic and electric type, $\text{Im}q_\alpha$ and $\text{Re}q_\beta$, respectively:

$$\begin{aligned} \text{Im}q_\alpha &= \int d\mathbb{S}^2 \alpha(\infty, \theta, \phi) \\ &\quad \times \left(\text{Im}\Phi_1^0(\infty, \theta, \phi) - \text{Im}\Phi_1^0(-\infty, \theta, \phi) \right), \end{aligned} \quad (35)$$

$$\begin{aligned} \text{Re}q_\beta &= \int d\mathbb{S}^2 \beta(\infty, \theta, \phi) \\ &\quad \times \left(\text{Re}\Phi_1^0(\infty, \theta, \phi) - \text{Re}\Phi_1^0(-\infty, \theta, \phi) \right). \end{aligned} \quad (36)$$

In these equations the real-valued functions α and β are defined from the longitudinal and transverse part of the electromagnetic potential at future timelike infinity, as follows. In the gauge we are using, in which $A_1^0 = A_a n^a = 0$, the one-form A_a^0 lives on the cotangent space of each cross section \mathbb{S}^2 of \mathcal{J}^+ . Therefore, it can be expressed as the sum of a gradient and a curl: $A_a^0 = D_a \alpha + e_a^b D_b \beta$, where D_a is the covariant derivative on \mathbb{S}^2 . This equation defines $\alpha(u, \theta, \phi)$ and $\beta(u, \theta, \phi)$.

Using (18) and (19), they can be solved from $\text{Re}\Phi_1^0(u, \theta, \phi) = \Delta\alpha(u, \theta, \phi) + G(\theta, \phi)$ and $\text{Im}\Phi_1^0(u, \theta, \phi) = \Delta\beta(u, \theta, \phi)$, where Δ denotes the two-dimensional Laplacian. Since the soft charges $\text{Im}q_\alpha$ and $\text{Re}q_\beta$ are specified from functions α and β , which depend on the electromagnetic potential, these are *field-dependent* soft charges. Since α, β originate from a gradient and a curl, respectively, the function α can be thought of as the electric degree of freedom of the emitted waves, while β is the magnetic one.

In summary, the change of the chiral charge of a quantum, massless, Dirac field between past to future null infinity, resulting from its coupling to an electromagnetic background, yields

$$\begin{aligned} \langle Q_A \rangle_{\mathcal{J}^+} - \langle Q_A \rangle_{\mathcal{J}^-} &= \frac{\hbar q^2}{4\pi^2} \left[\frac{1}{2\pi} \int d\mathbb{S}^2 \int_0^{\infty} \frac{d\omega}{\omega} \right. \\ &\quad \times (|\tilde{\Phi}_R(\omega, \theta, \phi)|^2 - |\tilde{\Phi}_L(\omega, \theta, \phi)|^2) \\ &\quad \left. + \text{Im}q_\alpha - \text{Re}q_\beta \right]. \end{aligned} \quad (37)$$

This is the main result of this paper. It shows that the anomalous nonconservation of fermionic helicity receives two types of contributions from an external electromagnetic field. Namely, $\langle Q_A \rangle$ can change in time if (i) a distribution of electric currents and charges in the bulk are able to radiate chiral electromagnetic waves, and (ii) there is a change in the infrared sector of the external electromagnetic field, such that the two soft charges (35) and (36) are different from zero. The presence of \hbar emphasizes that this is a quantum effect with no classical analog; it originates from the chiral anomaly.

We finish this section with a few remarks.

Remark 1. We have assumed no incoming radiation from \mathcal{J}^- . If the electromagnetic field is not trivial at past null infinity, we just need to replace quantities at \mathcal{J}^+ above with differences between \mathcal{J}^+ and \mathcal{J}^- .

Remark 2. The contribution from soft charges bears some similarity with the rationale behind the theory of instantons [34,35], in which quantum-mechanical transitions between “topologically inequivalent” vacuum states of the Hilbert space underlying a non-Abelian gauge theory induces an anomaly. In the quantum theory of the electromagnetic field, for each nontrivial IR sector one has a representation of the canonical commutation relations which is unitarily inequivalent to the usual Fock representation. So, just like with the interpretation of the instantons, we can say here that tunneling transitions between the different IR vacuum states of the electromagnetic field induces the fermionic chiral anomaly.⁹ In contrast, in this approach there is no need to work with

⁹Notice that this was precisely the origin of the chiral anomaly in 1 + 1 dimensions discussed in Sec. II.

Euclidean field equations. In fact, by working with solutions of the Lorentzian Maxwell equations we also get a radiative contribution, in addition to the contribution from soft charges. This radiative contribution is not predicted in the Euclidean case, where everything is “instantaneous.”

A. Examples

We discuss now examples of electromagnetic sources that are able to trigger the chiral anomaly obtained in (37). A physical configuration of electric charges and currents that can radiate circularly polarized electromagnetic waves was described in [16], namely, an electric-magnetic oscillating dipole. In this subsection we focus on examples that produce nonzero values of the infrared charges (36) and (35).

Soft charges of electric type are determined by the Coulombic contribution (i.e. $\sim 1/r^2$) of the radial component of the electric field:

$$\text{Re}q_{\ell m} = \oint d\mathbb{S}^2 \text{Re}\Phi_1^0(u, \theta, \phi) Y_{\ell m}(\theta, \phi) \Big|_{-\infty}^{+\infty},$$

$$\text{Re}\Phi_1^0(u, \theta, \phi) = \lim_{r \rightarrow \infty} [r^2 E_r(u, r, \theta, \phi)]. \quad (38)$$

The canonical example where these soft charges are not zero is a charged particle with some initial velocity that interacts with an external source (a nucleus, another charged particle, etc.) and changes its velocity [20]. In this process, the charged particle emits Bremstrahlung, which is known to possess zero-frequency photons. We shall review this example here for completeness.

The electromagnetic field generated by a moving charged particle can be obtained in closed form from the Lienard-Wiechart potentials [36]:

$$\vec{E}(t, \vec{r}) = \frac{q}{4\pi(1 - \vec{n}_s \cdot \vec{v}_s)^3} \left[\frac{\vec{n}_s - \vec{v}_s}{(1 - v_s^2) \|\vec{r} - \vec{r}_s\|^2} + \frac{\vec{n}_s \times [(\vec{n}_s - \vec{v}_s) \times \dot{\vec{v}}_s]}{\|\vec{r} - \vec{r}_s\|} \right] \Big|_{t=t_r}, \quad (39)$$

$$\vec{B}(t, \vec{r}) = \vec{v}_s(t_s) \times \vec{E}(t, \vec{r}), \quad (40)$$

where $\vec{r}_s(t)$ is the location of the charge, $\vec{v}_s = \dot{\vec{r}}_s$ its velocity, $\vec{n}_s = \frac{\vec{r} - \vec{r}_s(t)}{\|\vec{r} - \vec{r}_s(t)\|}$, and $t_r = t - \|\vec{r} - \vec{r}_s(t_r)\|$ is the retarded time [which is a function of (t, \vec{r})]. For a particle with constant velocity, $\dot{\vec{v}}_s = 0$, and the electric field above can be rewritten as [36]

$$\vec{E}(t, \vec{r}) = \frac{q}{4\pi} \frac{1 - v^2}{\left[1 - v^2 + \left(\frac{(\vec{r} - \vec{v}t) \cdot \vec{v}}{\|\vec{r} - \vec{v}t\|} \right)^2 \right]^{3/2}} \times \frac{\vec{r} - \vec{v}t}{\|\vec{r} - \vec{v}t\|^3}, \quad (\vec{r}_s(t) = \vec{v}_s \cdot t). \quad (41)$$

Changing to Bondi-Sachs coordinates $\{u, r, \theta, \phi\}$, and taking the limit $r \rightarrow \infty$ keeping $\{u, \theta, \phi\}$ constant, one obtains the following expression for the radial component of the electric field

$$E^a(u, r, \theta, \phi) \nabla_a r = \frac{q}{4\pi r^2} \frac{1 - v^2}{(1 - v^a \nabla_a r)^2} = \frac{q}{4\pi r^2} \frac{1 - v^2}{(1 - v \cos \theta)^2}, \quad (42)$$

where in the last equality we have chosen \hat{z} in the direction of \vec{v}_s . From this expression and (38) one readily obtains

$$\text{Re}\Phi_1^0(u, \theta, \phi) = \frac{q}{4\pi} \frac{1 - v^2}{(1 - v \cos \theta)^2}. \quad (43)$$

For a particle that always moves with the same constant velocity, $\text{Re}q_{\ell m} = 0$ for all ℓ . This is easy to see if we choose the reference system comoving with the particle, so that $\vec{v}_s = 0$ and $\text{Re}\Phi_1^0(u, \theta, \phi) = \frac{q}{4\pi}$. However, if the particle interacts with some external potential and changes its velocity, then we can no longer choose an inertial reference system attached to the particle at all times. While at early times we may have $\text{Re}\Phi_1^0(u \rightarrow -\infty, \theta, \phi) = \frac{q}{4\pi}$, at late times we will have $\text{Re}\Phi_1^0(u \rightarrow +\infty, \theta, \phi) = \frac{q}{4\pi} \frac{1 - v^2}{(1 - v \cos \theta)^2}$. The electric soft charges can be now computed:

$$\text{Re}q_{\ell m} = \frac{q}{4\pi} (1 - v^2) \oint_{t^+} d\mathbb{S}^2 \frac{Y_{\ell m}(\theta, \phi)}{(1 - v \cos \theta)^2} - q\delta_{\ell 0}$$

$$= \frac{q(1 - v^2)}{2} \delta_{m0} \int_{-1}^1 dx \frac{P_{\ell}(x)}{(1 - vx)^2} - q\delta_{\ell 0}. \quad (44)$$

In particular, $\text{Re}q_{00} = 0$ due to electric charge conservation, as expected. By taking different values of ℓ one can check that this expression is indeed different from zero. In summary, soft charges of electric type can be generated by Lorentz boosting an electric charge.

Soft charges of magnetic type are determined by the Coulombic contribution (i.e. $\sim 1/r^2$) of the radial component of the magnetic field:

$$\text{Im}q_{\ell m} = \oint d\mathbb{S}^2 \text{Im}\Phi_1^0(u, \theta, \phi) Y_{\ell m}(\theta, \phi) \Big|_{-\infty}^{+\infty},$$

$$\text{Im}\Phi_1^0(u, \theta, \phi) = \lim_{r \rightarrow \infty} [r^2 B_r(u, r, \theta, \phi)]. \quad (45)$$

Because magnetic charges have not been observed, we do not have, in principle, a magnetic analog of a boosted charge. To the best of our knowledge, there are no examples of magnetic memory reported in the literature. We discuss here one such example which, although it could be challenging to materialize physically [37], it certainly contains pedagogical value.

To think in potential situations that exhibit magnetic memory it may be useful to rewrite the radial component of the magnetic field of a moving particle (40) as

$$B^a(t, \vec{r}) \nabla_a r = \frac{q/m}{4\pi(1 - \vec{n}_s \cdot \vec{v}_s)^3 \|\vec{r} - \vec{r}_s(t_r)\|^2} \times \left\{ (\vec{n}_s \cdot \vec{a}_s) L_r + (1 - \vec{n}_s \cdot \vec{v}_s) \dot{L}_r + \frac{(1 - v_s^2) L_r}{\|\vec{r} - \vec{r}_s(t_r)\|} \right\} \Big|_{t_r}, \quad (46)$$

where $\vec{a}_s = \dot{\vec{v}}_s$ is the acceleration of the charge, and L_r is the radial component of the particle's angular momentum $\vec{L} = m\vec{r}_s \times \vec{v}_s$. Equivalently, these terms are related to the magnetic dipole moment of the moving charge:

$$\vec{m}(t) = \int_{\mathbb{R}^3} \vec{j}(t) \times \vec{r} d^3\vec{r} = q \int_{\mathbb{R}^3} d^3\vec{r} \delta^{(3)}(\vec{r} - \vec{r}_s(t)) \vec{v}_s(t) \times \vec{r} = q \vec{v}_s(t) \times \vec{r}_s(t) = -\frac{q}{m} \vec{L}(t). \quad (47)$$

Equation (46) shows, in particular, that for a charged particle moving with constant velocity, $\text{Im}\Phi_1^0 = 0$, and all soft charges of magnetic type are zero.

Only the first two terms in (46) may lead to $\text{Im}\Phi_1^0 \neq 0$, as the third term decays as $O(r^{-3})$. Furthermore, *a priori* one would expect that physically reasonable situations demand $a_s(u \rightarrow \pm\infty) \rightarrow 0$. So, for sources consisting of a moving charged particle, to get magnetic memory one needs a situation in which the particle acquires a permanent rate of change for the radial angular momentum at infinity, $\dot{L}_r \neq 0$.

One can imagine a situation in which this occurs. Consider polar coordinates $\{t, \rho, \phi, z\}$ and suppose there exists a nonvanishing magnetic field in $z \in (0, z_0)$ that has

the profile $\vec{B}(t, \rho, \phi, z) = B_0/\rho\hat{z}$ for $\rho > \rho_0$, for some constants B_0 and $\rho_0 > 0$. Due to the inhomogeneous magnetic field, a charged particle that initially is at rest at some point in the interval $(0, z_0)$ and $\rho > \rho_0$, and suffers a “kick” at some instant of time, will start spiraling outwards. It can be expected then that $\dot{L}_r \neq 0$ at infinity.

This problem can be solved in closed form as follows. If the kinematical variables of the charged particle are

$$\vec{r}_s = \rho_s \vec{u}_\rho, \quad (48)$$

$$\vec{v}_s = \dot{\rho}_s \vec{u}_\rho + \rho_s \dot{\phi}_s \vec{u}_\phi, \quad (49)$$

$$\vec{a}_s = (\ddot{\rho}_s - \rho_s \dot{\phi}_s^2) \vec{u}_\rho + (\rho_s \ddot{\phi}_s + 2\dot{\rho}_s \dot{\phi}_s) \vec{u}_\phi, \quad (50)$$

then one can solve

$$\dot{\rho}_s(t) = +\sqrt{v^2 - \left(\frac{qB_0}{m} + \frac{c}{\rho_s(t)}\right)^2}, \quad (51)$$

$$\dot{\phi}_s(t) = \mp \frac{qB_0}{m\rho_s(t)} + \frac{c}{\rho_s^2(t)}, \quad (52)$$

$$\frac{d}{dt}(\rho_s^2(t) \dot{\phi}_s(t)) = \frac{qB_0}{m} \dot{\rho}_s(t), \quad (53)$$

where c is a constant of integration. Since $\dot{\rho}_s(t) > 0$, $\rho_s(t)$ is monotonically increasing, therefore $\rho_s(t \rightarrow \infty) \rightarrow +\infty$ and consequently $\dot{\phi}_s \rightarrow 0$. The angular momentum is simply given by $\vec{L} = \rho_s^2(t) \dot{\phi}_s(t) \vec{u}_z$, so

$$\dot{L}_r(t) = \frac{qB_0}{m} \dot{\rho}_s(t) \cos \theta. \quad (54)$$

One can further check that $\vec{a}_s(t \rightarrow \infty) \rightarrow \vec{0}$.

Taking into account that $t_r(t, r \rightarrow \infty) \rightarrow u$, and also $\hat{r} \cdot \vec{v}_s(t) = \dot{\rho}_s(t) \sin \theta \cos(\phi - \phi_s(t)) + \rho_s(t) \dot{\phi}_s(t) \sin(\phi - \phi_s(t))$, the limit $r \rightarrow \infty$ keeping $\{u, \theta, \phi\}$ constant gives $\|\vec{r} - \vec{r}_s(t_r)\| \sim \|\vec{r} - \vec{r}_s(u)\| \sim r$ and $\vec{n}_s(t_r) \cdot \vec{v}_s(t_r) \sim \vec{n}_s(u) \cdot \vec{v}_s(u) \sim \hat{r} \cdot \vec{v}_s(u)$. From (46) we get

$$B_r(u, r, \theta, \phi) = \frac{q/m}{4\pi r^2 (1 - \dot{\rho}_s(u) \sin \theta \cos(\phi - \phi_s(u)) - \rho_s(u) \dot{\phi}_s(u) \sin(\phi - \phi_s(u)))^2} \frac{qB_0}{m} \dot{\rho}_s(u) \cos \theta + O(r^{-3}). \quad (55)$$

At future timelike infinity we have $\dot{\phi}_s(u \rightarrow \infty) \rightarrow 0$, $\phi_s(u \rightarrow \infty) \rightarrow \phi_0$, $\dot{\rho}_s(u \rightarrow \infty) \rightarrow \sqrt{v^2 - \frac{q^2 B_0^2}{m^2}}$. Taking into account all of this,

$$\text{Im}\Phi_1^0(u \rightarrow \infty) = \frac{q/m}{4\pi (1 - \sqrt{v^2 - \frac{q^2 B_0^2}{m^2}} \sin \theta \cos(\phi - \phi_0))^2} \frac{qB_0}{m} \sqrt{v^2 - \frac{q^2 B_0^2}{m^2}} \cos \theta. \quad (56)$$

To finish this section, recall from the discussion below (36) that $\text{Re}\Phi_1^0(u, \theta, \phi) = \Delta\alpha(u, \theta, \phi) + G(\theta, \phi)$ and $\text{Im}\Phi_1^0(u, \theta, \phi) = \Delta\beta(u, \theta, \phi)$. Therefore, from (43), obtained in the first example, it is possible to obtain a nonzero value of $\alpha(u, \theta, \phi)$ for $u \rightarrow \infty$, as well as $\text{Re}\Phi_1^0(\infty, \theta, \phi) - \text{Re}\Phi_1^0(-\infty, \theta, \phi) \neq 0$; while from (56), obtained in the second example, it is possible to obtain a nonzero value of $\beta(u, \theta, \phi)$ when $u \rightarrow +\infty$, and $\text{Im}\Phi_1^0(\infty, \theta, \phi) - \text{Im}\Phi_1^0(-\infty, \theta, \phi) \neq 0$. Combining the two examples, it is not difficult to check that (35) and (36) are both nonvanishing.

V. THE CHIRAL ANOMALY INDUCED BY A GRAVITATIONAL BACKGROUND

As remarked in the Introduction, if instead of an electromagnetic field we consider an external gravitational background, described by a curved spacetime (M, g_{ab}) , Dirac fields (as well as the electromagnetic field itself [11–14]) experience a gravitationally induced chiral anomaly [see Eq. (2)]. Similarly to what we did for electromagnetic backgrounds, we explore here global properties of this chiral anomaly by studying the change of the chiral charge Q_A .

As usual in general relativity, we restrict our study to globally hyperbolic spacetimes to ensure the well posedness of the Cauchy problem. This allows us to foliate the manifold in the form $M \simeq \mathbb{R} \times \Sigma$. We will further assume that the spatial slices are $\Sigma \simeq \mathbb{R}^3$. Performing a similar analysis as in (3), the permanent change in the chiral charge predicted by the chiral anomaly is dictated now by the Chern-Pontryagin integral

$$\langle Q_A \rangle_{\mathcal{J}^+} - \langle Q_A \rangle_{\mathcal{J}^-} = \frac{\hbar}{192\pi^2} \int_{\mathbb{R}^4} d^4x \sqrt{-g} R_{abcd} \star R^{abcd}, \quad (57)$$

where $\{x^a\}$ is a global coordinate system for $M \simeq \mathbb{R}^4$. The right-hand side of this equation was investigated in [15,16]. Although it may appear intractable from an analytical viewpoint, it is actually possible to rewrite it in a form that allows us to extract information of physical value without having to resort to numerical techniques. More precisely, assuming no incoming gravitational waves from past null infinity \mathcal{J}^- , it is possible to rewrite it as an integral over future null infinity only:

$$\langle Q_A \rangle_{\mathcal{J}^+} - \langle Q_A \rangle_{\mathcal{J}^-} = -\frac{\hbar}{96\pi^2} \int_{-\infty}^{\infty} du \int d\mathbb{S}^2 \text{Im}(N\bar{\Psi}_4). \quad (58)$$

In this expression, $\Psi_4(u, \theta, \phi) = -\lim_{r \rightarrow \infty} r C_{abcd} \bar{m}^a \times n^b \bar{m}^c n^d$ is a complex scalar constructed from the Weyl tensor C_{abcd} , which carries the two radiative degrees of freedom of gravitational waves; it is the gravitational analog of the complex scalar $\Phi_2^0(u, \theta, \phi)$ in electrodynamics [compare with equations (9) and (12)]. On the other

hand, $N(u, \theta, \phi) = N_{ab}(u, \theta, \phi) m^a m^b$ is the relevant component of the Bondi news tensor N_{ab} [25], which measures the time evolution of the asymptotic shear of outgoing null geodesics at \mathcal{J}^+ . It is a symmetric, transverse ($N_{ab} n^b = 0$), and traceless tensor on \mathcal{J}^+ that, just like Ψ_4 , captures the two gravitational degrees of freedom at future null infinity. The two quantities are related by $\Psi_4 = -\frac{1}{2}N$, so N can be thought of as the gravitational analog of the electromagnetic potential A_2^0 [compare with equation (17) with the gauge choice $A_1^0 = 0$]. The total amount of energy carried away by the gravitational waves across \mathcal{J}^+ is proportional to $\int_{-\infty}^{\infty} d\mathbb{S}^2 du |N(u, \theta, \phi)|^2$. Because of this, the Bondi news indicates unambiguously if a system is radiating gravitational waves. If $N = 0$ then the sources do not emit radiation, while $N \neq 0$ indicates the presence of radiation. Finiteness of this energy flux requires $N(\cdot, \theta, \phi) \in L^2(\mathbb{R}, \mathbb{C})$ for all $(\theta, \phi) \in \mathbb{S}^2$, and in particular $N \rightarrow 0$ as $u \rightarrow \pm\infty$. These properties carry over to Ψ_4 .

In view of the results found in Sec. IV, it is natural to ask if gravitational soft charges, or gravitational memory, may also contribute to the fermion chiral anomaly (58). The gravitational memory effect [38–44] consists in the permanent relative displacement that a set of free test masses may experience after the passage of a gravitational wave burst. The deformation of a congruence of free observers or curves is controlled by the shear. If $\sigma(u, \theta, \phi)$ denotes the asymptotic shear of outgoing null geodesics at future null infinity, a flux of gravitational radiation will make $\sigma(u, \theta, \phi)$ evolve with time u , while it remains constant otherwise. As commented above, this effect is captured precisely in the Bondi news, which is related to the shear via the equation $N = 2\dot{\sigma}$. Because $N \rightarrow 0$ as $u \rightarrow \pm\infty$, $\sigma(u, \theta, \phi)$ reaches constant values at early and late times. However, $\sigma(-\infty, \theta, \phi) \neq \sigma(\infty, \theta, \phi)$ in general, and there can remain a permanent distortion in the shear. The amount of gravitational memory encoded in free test masses is quantified then by the overall change in the asymptotic shear $\sigma(u, \theta, \phi)$ of outgoing null geodesics between early and late times:

$$q_\alpha = \frac{1}{8\pi} \int d\mathbb{S}^2 \bar{\delta}^2 \alpha(\theta, \phi) (\sigma(\infty, \theta, \phi) - \sigma(-\infty, \theta, \phi)), \quad (59)$$

where α is an arbitrary real-valued function on the sphere. These quantities are called gravitational infrared charges [27] [compare this definition with the electromagnetic analog (20)]. Following the analogy with the electromagnetic case, it can also be proven that these charges can be identified with the Hamiltonian generating Bondi-Metzner-Sachs (BMS) supertranslations in the radiative phase space of general relativity [45,46]. From the point of view of the bulk, supertranslations are diffeomorphisms (the gauge transformations in general relativity) that do not vanish at infinity, as a result of which they are called “large.”

Notice that the physical manifestations of the gravitational and electromagnetic memory effects are qualitatively different. An electromagnetic field does not generate a permanent, relative displacement of electrically charged particles; instead, it generates a permanent, relative velocity between the charges.

Using the relation between the shear and the Bondi news, we can formulate the gravitational infrared charges in terms of the radiative degrees of freedom,

$$q_\alpha = \frac{1}{16\pi} \int_{-\infty}^{+\infty} du d\theta d\phi S^2 N(u, \theta, \phi) \bar{\partial}^2 \alpha(\theta, \phi). \quad (60)$$

Expanding in a basis of spin-weighted spherical harmonics, this expression reduces to

$$q_\alpha = \frac{1}{16\pi} \int_{-\infty}^{+\infty} du \sum_{\ell m} \alpha_{\ell m} N_{\ell m}(u) \equiv \frac{1}{16\pi} \sum_{\ell m} \alpha_{\ell m} q_{\ell m}, \quad (61)$$

for real-valued coefficients $\alpha_{\ell m}$. In the second equality, we have defined the parameters $q_{\ell m} = \int_{-\infty}^{+\infty} du N_{\ell m}(u)$. Now, because $N(\cdot, \theta, \phi) \in L^2(\mathbb{R}, \mathbb{C})$, each of its harmonic modes admits a Fourier transform on \mathcal{J}^+

$$\tilde{N}_{\ell m}(\omega) = \int_{-\infty}^{+\infty} du N_{\ell m}(u) e^{-i\omega u}. \quad (62)$$

Therefore, just like in the electromagnetic case, we conclude that the infrared charges are determined by the zero-frequency mode of the gravitational radiation *as described by the Bondi news* N . Namely,

$$q_\alpha = \frac{1}{16\pi} \sum_{\ell m} \alpha_{\ell m} \tilde{N}_{\ell m}(0). \quad (63)$$

Notice, however, that, in sharp contrast with the electromagnetic case (20), the infrared charges are determined by the zero modes of the “potential” $N(u, \theta, \phi)$ and not by the zero modes of the “field” $\Psi_4(u, \theta, \phi)$. While this may look like an irrelevant comment, it is an important point in our analysis. The calculation of the right-hand side of (58) is formally equal to the electromagnetic case (27) if we identify A_2^0 with N , and Φ_2^0 with Ψ_4 . In the previous section we found that the electromagnetic infrared charges contribute to the chiral anomaly through the zero modes of the electromagnetic field $\Phi_2^0(u, \theta, \phi)$. Similarly, in the gravitational case, (58) only receives contributions from the zero modes of $\Psi_4^0(u, \theta, \phi)$, while the zero mode of

$N(u, \theta, \phi)$ never appears. However, $\Psi_4^0(u, \theta, \phi)$ has no zero mode:

$$\begin{aligned} \tilde{\Psi}_4(0, \theta, \phi) &= \int_{-\infty}^{+\infty} du \Psi_4(u, \theta, \phi) \\ &= -\frac{1}{2} [N(+\infty, \theta, \phi) - N(-\infty, \theta, \phi)] = 0, \end{aligned} \quad (64)$$

where in the second equality we made use of $\Psi_4 = -\frac{1}{2} \dot{N}$, and the last equality follows from $N(\pm\infty, \theta, \phi) = 0$. This is in sharp contrast with electrodynamics, where $\Phi_2^0(u, \theta, \phi)$ —the electromagnetic analog of Ψ_4 —does have a zero mode. As a consequence, only the radiative part of the gravitational field contributes to the chiral fermion anomaly in (58). There is no gravitational memory contributing to the change of the chiral charge $\langle Q_A \rangle$, and the total change from \mathcal{J}^- to \mathcal{J}^+ is determined by the helicity carried away by gravitational waves generated in the bulk [15,16].

VI. CONCLUSIONS

Chiral fermion anomalies have been extensively studied in the literature for several decades and from multiple viewpoints. Despite that, this topic is sufficiently rich to allow for yet another intriguing insight. We have found one such new aspect by studying global aspects of the chiral anomaly, related to the failure of the chiral charge Q_A of a massless Dirac field to be conserved. This charge is strictly conserved classically, as well as in quantum field theory for free Dirac fields. However, the presence of background fields, either electromagnetic or gravitational, may induce a local nonconservation of the chiral current j_A^a by quantum fluctuations, which can potentially produce a time evolution in the vacuum expectation value $\langle Q_A \rangle$.

The identification of external fields that can or cannot trigger a change of $\langle Q_A \rangle$ is a nontrivial problem. For non-Abelian gauge fields, a traditional approach is to look for instanton solutions in a Euclidean spacetime, which display a complex topological/global structure. To address this question, we have evaluated instead the change in $\langle Q_A \rangle$ between past and future null infinity using familiar, global techniques within the framework of asymptotically flat spacetimes. For an external electromagnetic field, our results are neatly summarized in Eq. (37). This equation tells us that $\langle Q_A \rangle$ can change between past and future null infinity if (i) electromagnetic sources in the bulk emit circularly polarized electromagnetic waves (i.e. radiation with net helicity) and/or (ii) if electromagnetic sources in the bulk produce transitions between certain infrared sectors of Maxwell theory. The relevant transitions are determined by a concrete pair of infrared charges of electric and magnetic type, respectively, written in Eqs. (35) and (36). To gain physical intuition, we have devised an academic example where the required soft charges are different from zero.

Physically, nonzero infrared charges are known to produce memory effects on physical systems. This is

how the transitions between the infrared quantum vacua can leave observable imprints. To the best of our knowledge, the only electromagnetic memory effects known to date involve classical systems. Here, we have shown that quantum states of a field theory can also keep memory of the past influence of electromagnetic radiation, by storing a certain amount of helicity.

The connection of electromagnetic memory and the change of $\langle Q_A \rangle$ has also been worked out in $1+1$ dimensions, which is cleaner because there are no electromagnetic waves. This example also allowed us to interpret this new memory effect in terms of “kicks” of virtual charges and excitation of particle pairs out of the quantum vacuum.

Overall, the results in this paper, together with our previous analysis [15,16], open up an unforeseen connection between chiral anomalies, the radiative content of the electromagnetic field, infrared charges, and the memory effect.

Although our approach is qualitatively different, the contribution from soft charges to the chiral anomaly bears some similarity with the rationale underlying instantons in Euclidean gauge-field theories. According to the usual interpretation [34,35], instantons mediate quantum-mechanical transitions between inequivalent vacuum states of the Hilbert space of the background (non-Abelian) gauge field. These transitions, which are labeled by the instanton charge, are able to induce the chiral anomaly [47]. On the other hand, the quantization of the electromagnetic field at future null infinity leads naturally to a Hilbert space that can be divided in different, disjoint infrared sectors [26,48], which represent inequivalent notions of quantum vacua. The infrared charges label transitions between the different infrared sectors, and therefore play the same role of the instanton charge. We have shown in this article that these transitions contribute to the chiral anomaly in a specific manner.

To finish, we have also checked that, quite interestingly, gravitational infrared charges do not contribute to the fermion chiral anomaly.

ACKNOWLEDGMENTS

I. A. thanks Miguel Campiglia for numerous illuminating discussions on the content of this article. A. D. R. is grateful to A. Ashtekar for a deep training of many of the techniques employed in this article. A. D. R. is supported through a M. Zambrano grant (ZA21-048) with reference UP2021-044 from the Spanish Ministerio de Universidades, funded within the European Union-Next Generation EU. This work is also supported by the Spanish Grant PID2020-116567GB-C21 funded by MCIN/AEI/10.13039/501100011033 and the project PROMETEO/2020/079 (Generalitat Valenciana); as well as by the NSF Grant PHY-1806356, and the Eberly Chair funds of Penn State. I. A. is supported by the NSF Grant PHY-

2110273, by the RCS program of Louisiana Boards of Regents through the Grant LEQSF(2023-25)-RD-A-04, by the Hearne Institute for Theoretical Physics, and in part by Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science, and Economic Development, and by the Province of Ontario through the Ministry of Colleges and Universities.

APPENDIX A: CHIRAL FERMION ANOMALY AND SPONTANEOUS PARTICLE CREATION IN $1+1$ DIMENSIONS

In Sec. II we argued that the permanent change in the vacuum expectation value $\langle Q_A \rangle$ produced by the chiral anomaly (2) can be understood as a net creation of helicity resulting from virtual particles that are “kicked” out of the vacuum state. In $1+1$ dimensions it is relatively easy to see the connection between the chiral fermion anomaly and particle pair creation. In this appendix we show this connection and deduce the anomaly from the analysis of Bogoliubov transformations between canonical in and out vacua. We perform the analysis for a general, nonuniform background electric field $E(t, x)$.

When the external electric field is uniform, $E = E(t)$, a heuristic argument involving the Dirac “sea” has been given in several occasions in the literature (see [23,49–51]). However, to the best of our knowledge, an explicit and/or rigorous, complete calculation is still lacking. In particular, this heuristic picture always misses a contribution from vacuum polarization, which we provide here.

Let us consider a massless, quantum Dirac field $\Psi(t, x)$ living in a two-dimensional flat spacetime, $(\mathbb{R} \times \mathbb{S}^1, \eta_{ab})$, and coupled to an external electric field $E(t, x)$. The spatial sections will have length equal to L . The electric field departs from an initial “vacuum” configuration at early times, $E(t_{\text{in}}, x) = 0$; it is switched on for a finite amount of time, and eventually returns to another “vacuum” state, $E(t_{\text{out}}, x) = 0$. At early and late times, in which the field strength vanishes, one can introduce canonical “in” and “out” vacuum states for the fermion field. As remarked in Sec. II, the two electric vacua are equivalent in the classical theory, but they may differ in the quantum theory if the potential $A_a(t, x)$ changes nontrivially, as a result of which the “out” state of the fermion field will potentially differ from the “in” state.

As in Sec. II, we work with the temporal gauge fixing: $A_t(t, x) = 0$. This can always be obtained by a suitable gauge transformation. There is still a residual gauge freedom, which consists in $A_a(t, x) \rightarrow A_a(t, x) + \nabla_a \alpha$, for $\alpha = \alpha(x)$. This residual freedom can be fixed by demanding $A_a(t \rightarrow -\infty, x) \rightarrow 0$, which we will adopt here onwards. At early and late times, where $E(t, x) = 0$, we have $\partial_t A_x = \partial_x A_t$, and the electromagnetic connection one-form takes the form $A(t \rightarrow -\infty, x) = 0$ and $A(t \rightarrow \infty, x) = A(x)dx$.

Massless Dirac fields $\Psi(t, x)$ split into two decoupled, left- and right-handed Weyl spinors that will be denoted by u_+ , u_- , respectively. In $1 + 1$ dimensions these spinors are represented by ordinary functions on the spacetime. For $t < t_{\text{in}}$, the two Weyl equations read

$$\begin{aligned} (i\partial_t - i\partial_x)u_+ &= 0, \\ (i\partial_t + i\partial_x)u_- &= 0, \end{aligned} \quad (\text{A1})$$

while for $t > t_{\text{out}}$ we have

$$(i\partial_t - i\partial_x - qA(x))u_+ = 0, \quad (\text{A2})$$

$$(i\partial_t + i\partial_x + qA(x))u_- = 0, \quad (\text{A3})$$

where the evolution of the electric field has left a residual gauge potential in the field equations. To study the effects of the external, dynamical electric field on the fermion modes, it is useful to analyze the Bogoliubov transformations between the “in” and “out” vacuum states. To specify these states we have to provide first a basis of positive-frequency solutions for the in and out Hilbert spaces.

Let us focus on Eqs. (A2) and (A3) (the solutions at early times can be simply recovered by taking $A = 0$). These equations admit separable solutions of the form $u_{\pm}(t, x) = f_{\pm}(t)g_{\pm}(x)$, which produce

$$\frac{i\partial_t f_{\pm}(t)}{f_{\pm}(t)} = \pm \frac{(\partial_x + qA(x))g_{\pm}(x)}{g_{\pm}(x)} \equiv \omega_{\pm} = \text{const}, \quad (\text{A4})$$

for some separation constants ω_{\pm} . The left-hand side of this equation can be solved to give $f_{\pm}(t) = e^{-i\omega_{\pm}t}$ and are of positive frequency with respect to the operator $H = i\partial_t$ if $\omega_{\pm} > 0$. To solve the spatial dependence of the field modes, note that the potential must be pure gauge (because $E = 0$), so let us write $qA = \partial_x\phi(x)$ for some function $\phi(x)$. Then $(i\partial_x + qA(x))g_{\pm}(x) = e^{+i\phi(x)}i\partial_x(e^{-i\phi(x)}g_{\pm})$ and the equation above yields $g_{\pm}(x) = e^{\mp i\omega_{\pm}x}e^{i\phi(x)}$. In conclusion, a basis of positive-frequency solutions to the Weyl equations above consist of $\{u_{\pm,\omega}(t, x)\}_{\omega > 0}$, where

$$u_{\pm,\omega}(t, x) = e^{-i\omega_{\pm}(t\pm x)}e^{i\phi(x)}. \quad (\text{A5})$$

These functions represent left- and right-moving modes, respectively, and they define the two chiral sectors of the theory. In $1 + 1$ dimensions handedness is nothing but the direction of propagation in the spatial dimension.

At early times the connection is identically zero, $A = 0$, so $\phi(x) = 0$ for the in modes: $u_{\pm,\omega}^{\text{in}}(t, x) = e^{-i\omega_{\pm}(t\pm x)}$. However, the dynamics of the electric field could produce $\phi(x) \neq 0$ at late times. This is the electric memory mentioned in the main text. The out modes can differ by a position-dependent, global phase with respect to the in modes: $u_{\pm,\omega}^{\text{out}}(t, x) = e^{-i\omega_{\pm}(t\pm x)}e^{i\phi(x)}$.

The compactness of the spatial spacetime dimension imposes severe constraints on the field modes. On \mathbb{S}^1 there are two inequivalent spin structures: the trivial one (that corresponds to imposing periodic boundary conditions on the field modes), and the nontrivial one (that corresponds to imposing antiperiodic boundary conditions) [52]. That is, $u_{\pm,\omega}(t, x + L) = e^{i2\pi\delta}u_{\pm,\omega}(t, x)$, where $\delta = 0$ or $1/2$ for periodic and antiperiodic boundary conditions, respectively. We shall assume here periodic boundary conditions, for simplicity. This implies $e^{\mp i\omega_{\pm}L}e^{i\phi(x+L)} = e^{i\phi(x)}$, which produces $\mp\omega_{\pm,n}L + \phi(x + L) - \phi(x) = -n2\pi, n \in \mathbb{Z}$, or

$$\omega_{\pm,n} = \pm n \frac{2\pi}{L} \pm \frac{(\phi(x + L) - \phi(x))}{L}. \quad (\text{A6})$$

In other words, the allowed frequencies of the field modes take only discrete values. To convert the right-hand side in terms of the electromagnetic potential, recall that $qA_x = \partial_x\phi(x)$, so $\phi(x) = q \int^x A(x')dx'$, and $\phi(x + L) - \phi(x) = q \int_x^{x+L} A(x')dx'$. This integral is well defined, so we can make a change of variable $x' \rightarrow x' - x$ to rewrite it as $\phi(x + L) - \phi(x) = q \int_0^L A(x')dx' =: 2\pi qCS[A]$, where $CS[A]$ is the Chern-Simons [9]. Therefore,

$$\omega_{\pm,n} = \pm \frac{2\pi}{L}(n + qCS[A]). \quad (\text{A7})$$

From this result we can infer

$$\omega_{\pm,n} > 0 \text{ implies } \begin{cases} n > -qCS[A] & \text{for positive chirality,} \\ n < -qCS[A] & \text{for negative chirality,} \end{cases} \quad (\text{A8})$$

$$\omega_{\pm,n} < 0 \text{ implies } \begin{cases} n < -qCS[A] & \text{for positive chirality,} \\ n > -qCS[A] & \text{for negative chirality.} \end{cases} \quad (\text{A9})$$

The field equations (A2) and (A3) are linear, and therefore the space of solutions of each chiral sector has the structure of a vector space. As usual, we endow these vector spaces with the Dirac inner product $(u_1, u_2) = \int_0^L dx \bar{u}_1(t, x)u_2(t, x)$, which is preserved in time by the Weyl equations and by the periodic boundary conditions. Then, an orthonormal basis $f_{\pm,n}^{\text{in}}(t, x)$ of periodic, positive-frequency modes for the “in” Hilbert space $L^2((0, L)) \oplus L^2((0, L))$ is given by $\{\frac{e^{-i\frac{2\pi}{L}(t+x)}}{\sqrt{L}}\}_{n \in \mathbb{Z}^+}$ for left-moving spinors and $\{\frac{e^{i\frac{2\pi}{L}(t-x)}}{\sqrt{L}}\}_{n \in \mathbb{Z}^-}$ for right-moving spinors. If we define the helicity as $h := \hbar \frac{\omega_{\pm,n}}{\frac{2\pi}{L}} = \pm \hbar$, we see that for left-moving spinors, u_+ , positive-frequency modes have positive helicity, while for right-moving spinors u_- positive-frequency modes have negative helicity ($\frac{2\pi}{L}n$ can be thought of as the wave number of the mode). On the other hand, an orthonormal basis $f_{\pm,n}^{\text{out}}(t, x)$ of periodic, positive-frequency modes for the out Hilbert

space $L^2((0, L)) \oplus L^2((0, L))$ is given, at late times, by

$$\left\{ \frac{e^{-i\frac{2\pi}{L}n(t+x)}}{\sqrt{L}} e^{i\frac{2\pi}{L}qCS[A](-t-x)+iq\int^x A(x')dx'} \right\}_{n+qCS[A] \in \mathbb{Z}^+} \quad (\text{A10})$$

for left-moving spinors, and

$$\left\{ \frac{e^{i\frac{2\pi}{L}n(t-x)}}{\sqrt{L}} e^{i\frac{2\pi}{L}qCS[A](t-x)+iq\int^x A(x')dx'} \right\}_{n+qCS[A] \in \mathbb{Z}^-} \quad (\text{A11})$$

for right-moving spinors.

In the basis of in modes the quantum fields can be expanded as

$$u_+(t, x) = \sum_{n>0}^{+\infty} a_{+,n}^{\text{in}} f_{+,n}^{\text{in}}(t, x) + \sum_{n<0}^{-\infty} b_{-,n}^{\text{in}\dagger} f_{+,n}^{\text{in}}(t, x), \quad (\text{A12})$$

$$u_-(t, x) = \sum_{n<0}^{-\infty} a_{-,n}^{\text{in}} f_{-,n}^{\text{in}}(t, x) + \sum_{n>0}^{+\infty} b_{+,n}^{\text{in}\dagger} f_{-,n}^{\text{in}}(t, x), \quad (\text{A13})$$

where $f_{\pm,n}^{\text{in}}(t, x) \rightarrow \frac{e^{\mp i\frac{2\pi}{L}n(t\pm x)}}{\sqrt{L}}$ at early times. In this expression $a_{\pm,n}^{\text{in}}$ annihilate fermions (negative charges) with positive (+) and negative (−) helicity, while $b_{\mp,n}^{\text{in}\dagger}$ create

antifermions (positive charges) with negative (−) and positive (+) helicity. The ± signs on u_{\pm} and $f_{\pm,n}^{\text{in}}$ simply denote the sign of chirality (i.e. the direction of propagation). On the other hand, in the basis of out modes we have instead

$$u_+(t, x) = \sum_{n>-qCS[A]}^{+\infty} a_{+,n}^{\text{out}} f_{+,n}^{\text{out}}(t, x) + \sum_{n<-qCS[A]}^{-\infty} b_{-,n}^{\text{out}\dagger} f_{+,n}^{\text{out}}(t, x), \quad (\text{A14})$$

$$u_-(t, x) = \sum_{n<-qCS[A]}^{-\infty} a_{-,n}^{\text{out}} f_{-,n}^{\text{out}}(t, x) + \sum_{n>-qCS[A]}^{+\infty} b_{+,n}^{\text{out}\dagger} f_{-,n}^{\text{out}}(t, x), \quad (\text{A15})$$

where $f_{\pm,n}^{\text{out}}(t, x) \rightarrow \frac{e^{\mp i\frac{2\pi}{L}n(t\pm x)}}{\sqrt{L}} e^{i\frac{2\pi}{L}qCS[A](\mp t-x)+iq\int^x A(x')dx'}$ at late times. The Bogoliubov coefficients that relate the in and out representations are now easy to obtain. Since $f_{\pm,n}^{\text{out}}(t, x)$ form a complete basis of the Hilbert space of solutions, $f_{\pm,n}^{\text{in}}(t, x)$ can be expanded as

$$n > 0 \text{ (positive energy): } f_{+,n}^{\text{in}}(t, x) = \sum_{n'>-qCS[A]}^{+\infty} \alpha_{nn'}^+ f_{+,n'}^{\text{out}}(t, x) + \sum_{n'<-qCS[A]}^{-\infty} \beta_{nn'}^+ f_{+,n'}^{\text{out}}(t, x), \quad (\text{A16})$$

$$n < 0 \text{ (negative energy): } f_{+,n}^{\text{in}}(t, x) = \sum_{n'<-qCS[A]}^{-\infty} \tilde{\alpha}_{nn'}^+ f_{+,n'}^{\text{out}}(t, x) + \sum_{n'>-qCS[A]}^{+\infty} \tilde{\beta}_{nn'}^+ f_{+,n'}^{\text{out}}(t, x), \quad (\text{A17})$$

$$n < 0 \text{ (positive energy): } f_{-,n}^{\text{in}}(t, x) = \sum_{n'<-qCS[A]}^{-\infty} \alpha_{nn'}^- f_{-,n'}^{\text{out}}(t, x) + \sum_{n'>-qCS[A]}^{+\infty} \beta_{nn'}^- f_{-,n'}^{\text{out}}(t, x), \quad (\text{A18})$$

$$n > 0 \text{ (negative energy): } f_{-,n}^{\text{in}}(t, x) = \sum_{n'>-qCS[A]}^{+\infty} \tilde{\alpha}_{nn'}^- f_{-,n'}^{\text{out}}(t, x) + \sum_{n'<-qCS[A]}^{-\infty} \tilde{\beta}_{nn'}^- f_{-,n'}^{\text{out}}(t, x). \quad (\text{A19})$$

Using the normalization conditions $(f_{\pm,n}, f_{\pm,n'}) = \delta_{n,n'}$ one can get several useful identities. For instance:

$$\delta_{nn'} = (f_{+,n}^{\text{in}}, f_{+,n'}^{\text{in}}) = \sum_{n''>-qCS[A]}^{+\infty} \tilde{\alpha}_{nn''}^+ \alpha_{n'n''}^+ + \sum_{n''<-qCS[A]}^{-\infty} \tilde{\beta}_{nn''}^+ \beta_{n'n''}^+, \quad (\text{A20})$$

$$\delta_{nn'} = (f_{-,n}^{\text{in}}, f_{-,n'}^{\text{in}}) = \sum_{n''<-qCS[A]}^{-\infty} \tilde{\alpha}_{nn''}^- \alpha_{n'n''}^- + \sum_{n''>-qCS[A]}^{+\infty} \tilde{\beta}_{nn''}^- \beta_{n'n''}^-. \quad (\text{A21})$$

Setting $n = n'$, the convergence of the integrals imply that $\lim_{n'' \rightarrow \pm\infty} |\beta_{nn''}^{\pm}| = 0$. In what follows we will also need the inverse Bogoliubov transformations. Since $f_{\pm,n}^{\text{in}}(t, x)$ also form a complete basis of the Hilbert space of solutions, the elements $f_{\pm,n}^{\text{out}}(t, x)$ can equivalently be expanded as

$$n > -qCS[A] \text{ (positive energy): } f_{+,n}^{\text{out}}(t, x) = \sum_{n'>0}^{+\infty} \gamma_{nn'}^+ f_{+,n'}^{\text{in}}(t, x) + \sum_{n'<0}^{-\infty} \delta_{nn'}^+ f_{+,n'}^{\text{in}}(t, x), \quad (\text{A22})$$

$$n < -qCS[A] \text{ (positive energy): } f_{-,n}^{\text{out}}(t, x) = \sum_{n'<0}^{-\infty} \gamma_{nn'}^- f_{-,n'}^{\text{in}}(t, x) + \sum_{n'>0}^{+\infty} \delta_{nn'}^- f_{-,n'}^{\text{in}}(t, x). \quad (\text{A23})$$

For $n'' > 0$ we have $\gamma_{nn''}^+ = (f_{+,n''}^{\text{in}}, f_{+,n''}^{\text{out}}) = \overline{\alpha_{n''n}^+}$ and for $n'' < 0$ we have $\delta_{nn''}^+ = (f_{+,n''}^{\text{in}}, f_{+,n''}^{\text{out}}) = \overline{\tilde{\beta}_{n''n}^+}$. Similarly, for $n'' < 0$ we obtain $\gamma_{nn''}^- = (f_{-,n''}^{\text{in}}, f_{-,n''}^{\text{out}}) = \overline{\alpha_{n''n}^-}$, while for $n'' > 0$ we get $\delta_{nn''}^- = (f_{-,n''}^{\text{in}}, f_{-,n''}^{\text{out}}) = \overline{\tilde{\beta}_{n''n}^-}$. Using $(f_{\pm,n}, f_{\pm,n'}) = \delta_{n,n'}$ we further obtain the following identities:

$$\delta_{nn'} = (f_{+,n}^{\text{out}}, f_{+,n'}^{\text{out}}) = \sum_{n'' > 0}^{+\infty} \overline{\alpha_{n''n}^+} \alpha_{n''n'}^+ + \sum_{n'' < 0}^{-\infty} \overline{\tilde{\beta}_{n''n}^+} \tilde{\beta}_{n''n'}^+, \quad (\text{A24})$$

$$\delta_{nn'} = (f_{-,n}^{\text{out}}, f_{-,n'}^{\text{out}}) = \sum_{n'' < 0}^{-\infty} \overline{\alpha_{n''n}^-} \alpha_{n''n'}^- + \sum_{n'' > 0}^{+\infty} \overline{\tilde{\beta}_{n''n}^-} \tilde{\beta}_{n''n'}^-. \quad (\text{A25})$$

Setting $n = n'$, the convergence of the integrals imply that $\lim_{|n''| \rightarrow +\infty} |\tilde{\beta}_{n''n}^{\pm}| = 0$.

Plugging (A16) and (A17) into (A12), and rewriting as in (A14), we can read off the Bogoliubov transformation of the creation and annihilation operators:

$$a_{+,n'}^{\text{out}} = \sum_{n>0}^{+\infty} \alpha_{nn'}^+ a_{+,n}^{\text{in}} + \sum_{n<0}^{-\infty} \tilde{\beta}_{nn'}^+ b_{-,n}^{\text{in}\dagger}, \quad (\text{A26})$$

$$b_{-,n'}^{\text{out}\dagger} = \sum_{n<0}^{-\infty} \tilde{\alpha}_{nn'}^+ b_{-,n}^{\text{in}\dagger} + \sum_{n>0}^{+\infty} \beta_{nn'}^+ a_{+,n}^{\text{in}}. \quad (\text{A27})$$

Similarly, plugging (A18) and (A19) in (A13), and rewriting as in (A15), we can read off the Bogoliubov transformation for the remaining creation and annihilation operators:

$$a_{-,n'}^{\text{out}} = \sum_{n<0}^{-\infty} \alpha_{nn'}^- a_{-,n}^{\text{in}} + \sum_{n>0}^{+\infty} \tilde{\beta}_{nn'}^- b_{+,n}^{\text{in}\dagger}, \quad (\text{A28})$$

$$b_{+,n'}^{\text{out}\dagger} = \sum_{n>0}^{+\infty} \tilde{\alpha}_{nn'}^- b_{+,n}^{\text{in}\dagger} + \sum_{n<0}^{-\infty} \beta_{nn'}^- a_{-,n}^{\text{in}}. \quad (\text{A29})$$

We have now all of the required ingredients to see the connection between the chiral anomaly and particle pair production. Using the normalization equations $(f_{\pm,n}, f_{\pm,n'}) = \delta_{n,n'}$, the chiral charge can be formally evaluated as:

$$\begin{aligned} Q_5(t) &= \int_0^L dx \bar{\Psi}(t, x) \gamma^0 \gamma_5 \Psi(t, x) = \int_0^L dx \Psi^\dagger(t, x) \gamma_5 \Psi(t, x) = \int_0^L dx (\bar{u}_+ u_+ - \bar{u}_- u_-) = (u_+, u_+) - (u_-, u_-) \\ &= \sum_{n>-qCS[A]} a_{+,n}^{\text{out}\dagger} a_{+,n}^{\text{out}} + \sum_{n<-qCS[A]} b_{-,n}^{\text{out}\dagger} b_{-,n}^{\text{out}} - \sum_{n<-qCS[A]} a_{-,n}^{\text{out}\dagger} a_{-,n}^{\text{out}} - \sum_{n>-qCS[A]} b_{+,n}^{\text{out}\dagger} b_{+,n}^{\text{out}}. \end{aligned}$$

However, this operator is not well defined in our Fock space; its expectation values produce divergent sums. This is because $Q_5(t)$ is quadratic in the quantum fields, and, consequently, the evaluation of expectation values requires renormalization. The quantity of interest is $\langle \text{in} | Q_5(t) | \text{in} \rangle_{\text{ren}}$, whose time evolution tells us whether there exists an anomaly or not. To obtain this result one can apply renormalization directly. However, there is an alternative, indirect procedure which, as we shall see, provides useful insights on the physical interpretation of $\langle \text{in} | Q_5(t) | \text{in} \rangle_{\text{ren}}$. Let us introduce the following fiducial (“normal-ordered”) operator:

$$\begin{aligned} :Q_5: (t_{\text{out}}) &:= \int_0^L dx \lim_{y \rightarrow x} [\bar{\Psi}(t_{\text{out}}, x) \gamma^0 \gamma_5 \Psi(t_{\text{out}}, y) - \mathbb{I} \langle \text{out} | \bar{\Psi}(t_{\text{out}}, x) \gamma^0 \gamma_5 \Psi(t_{\text{out}}, y) | \text{out} \rangle] \\ &= \sum_{n>-qCS[A]} a_{+,n}^{\text{out}\dagger} a_{+,n}^{\text{out}} - \sum_{n<-qCS[A]} b_{-,n}^{\text{out}\dagger} b_{-,n}^{\text{out}} - \sum_{n<-qCS[A]} a_{-,n}^{\text{out}\dagger} a_{-,n}^{\text{out}} + \sum_{n>-qCS[A]} b_{+,n}^{\text{out}\dagger} b_{+,n}^{\text{out}}. \end{aligned} \quad (\text{A30})$$

This operator, which is given in terms of particle number operators of the out vacuum state, is now well defined on the Fock space. In particular, the expectation value $\langle \text{in} | :Q_5:(t_{\text{out}}) | \text{in} \rangle$ exists. However, keep in mind that this is just an auxiliary operator that we introduced for convenience. What truly determines the quantum anomaly is the time evolution of the charge Q_5 , *not* of the fiducial operator $:Q_5:$. Using the definition above we can obtain the relation between the two:

$$\langle \text{in} | :Q_5:(t_{\text{out}}) | \text{in} \rangle = \langle \text{in} | Q_5(t_{\text{out}}) | \text{in} \rangle_{\text{ren}} - \langle \text{out} | Q_5(t_{\text{out}}) | \text{out} \rangle_{\text{ren}}. \quad (\text{A31})$$

We can now invert this expression to finally get the result of interest:

$$\langle \text{in} | Q_5(t_{\text{out}}) | \text{in} \rangle_{\text{ren}} = \langle \text{in} | :Q_5:(t_{\text{out}}) | \text{in} \rangle + \langle \text{out} | Q_5(t_{\text{out}}) | \text{out} \rangle_{\text{ren}}. \quad (\text{A32})$$

The first contribution on the right-hand side depends only on the Bogoliubov coefficients and does not require renormalization. Then, it can be understood in terms of particle pairs created with net helicity by the external, electric background. The second term on the right-hand side requires renormalization, and it represents a vacuum polarization effect. To the quantum anomaly, *both* effects contribute.¹⁰

As a side remark, it is interesting to note the similarity of this result with the Hawking effect for black holes. In the formation of a black hole by gravitational collapse, one can compute the expectation value of the particle number operator using Bogoliubov transformations between past and future null infinity (with in and out states, respectively), as Hawking originally did. This calculation is well defined and does not require renormalization. In our problem, this would be analogous to the calculation of $\langle \text{in} | :Q_5: (t_{\text{out}}) | \text{in} \rangle$, which is related to the particle number operators. On the other hand, one can also study the Hawking effect by computing the nondiagonal, flux component of the expectation value of the stress-energy tensor across future null infinity. This calculation, based only on the in state, does require renormalization. This is because, apart from the particle pair creation, there is yet another contribution coming from vacuum polarization effects. In our case, since Q_5 is a quadratic operator, the evaluation of $\langle \text{in} | Q_5(t_{\text{out}}) | \text{in} \rangle_{\text{ren}}$ is analogous to the calculation of the Hawking effect using the stress-energy tensor and not via the particle number operator.

The evaluation of $\langle \text{in} | :Q_5: (t_{\text{out}}) | \text{in} \rangle$ in (A32) is straightforward from the expressions (A26)–(A29) above and the canonical commutation relations. It produces

$$\begin{aligned} \langle \text{in} | :Q_5: (t_{\text{out}}) | \text{in} \rangle / \hbar = & \sum_{\substack{n > -qCS[A] \\ n' < 0}} |\tilde{\beta}_{n'n}^+|^2 - \sum_{\substack{n < -qCS[A] \\ n' > 0}} |\beta_{n'n}^+|^2 \\ & - \sum_{\substack{n < -qCS[A] \\ n' > 0}} |\tilde{\beta}_{n'n}^-|^2 + \sum_{\substack{n > -qCS[A] \\ n' < 0}} |\beta_{n'n}^-|^2. \end{aligned}$$

Note that each sum is convergent because each summand decays in both indices as $n \rightarrow \infty$. Using (A20)–(A21),

¹⁰Mathematically, $:Q_5:$ (and not Q_5) is the relevant operator that is related with the index theorem in geometric analysis [53], and, because of this, one may be tempted to identify it with the chiral anomaly. Historically, chiral anomalies were studied on compact manifolds without boundary, that arise naturally using Euclidean techniques, and in these cases the chiral fermion anomaly was found to match the predictions from the Atiyah-Singer index theorem. As a result, the statement that chiral anomalies are predicted by index theorems became a standard lore. However, this is not true in more general cases. In particular, for manifolds with boundary, extra contributions arise in the index theorem, like the APS eta index η_{APS} [53], and the agreement with the anomaly fails. Physically, these extra boundary terms are represented by the vacuum polarization effects $\langle \text{out} | Q_5(t_{\text{out}}) | \text{out} \rangle_{\text{ren}}$ pointed out in this appendix.

(A24)–(A25) one can write:

$$\sum_{n>0}^k 1 = \sum_{n>0}^k \left(\sum_{n'' > -qCS[A]}^{+\infty} |\alpha_{nn''}^+|^2 + \sum_{n'' < -qCS[A]}^{-\infty} |\beta_{nn''}^+|^2 \right), \quad (\text{A33})$$

$$\sum_{n<0}^{-k} 1 = \sum_{n<0}^{-k} \left(\sum_{n'' < -qCS[A]}^{-\infty} |\alpha_{nn''}^-|^2 + \sum_{n'' > -qCS[A]}^{+\infty} |\beta_{nn''}^-|^2 \right), \quad (\text{A34})$$

$$\sum_{n>-qCS[A]}^k 1 = \sum_{n>-qCS[A]}^k \left(\sum_{n'' > 0}^{+\infty} |\alpha_{n''n}^+|^2 + \sum_{n'' < 0}^{-\infty} |\tilde{\beta}_{n''n}^+|^2 \right), \quad (\text{A35})$$

$$\sum_{n<-qCS[A]}^{-k} 1 = \sum_{n<-qCS[A]}^{-k} \left(\sum_{n'' < 0}^{-\infty} |\alpha_{n''n}^-|^2 + \sum_{n'' > 0}^{+\infty} |\tilde{\beta}_{n''n}^-|^2 \right), \quad (\text{A36})$$

for some positive integer k . Subtracting (A33) from (A35), and then taking the limit $k \rightarrow \infty$, we obtain $\sum_{\substack{n > -qCS[A] \\ n' < 0}} |\tilde{\beta}_{n'n}^+|^2 - \sum_{\substack{n < -qCS[A] \\ n' > 0}} |\beta_{n'n}^+|^2 = [qCS[A]]$, where $[\cdot]$ indicates the integer part. Subtracting now (A34) from (A36), and taking again the limit $k \rightarrow \infty$, we get $\sum_{\substack{n < -qCS[A] \\ n' > 0}} |\tilde{\beta}_{n'n}^-|^2 - \sum_{\substack{n > -qCS[A] \\ n' < 0}} |\beta_{n'n}^-|^2 = -[qCS[A]]$. In conclusion,

$$\langle \text{in} | :Q_5: (t_{\text{out}}) | \text{in} \rangle = 2\hbar[qCS[A]]. \quad (\text{A37})$$

This is the main result of this appendix. It makes manifest that the vacuum expectation value of the Dirac chiral charge Q_5 at late times, (A32), receives an important contribution from particle pair creation.

The evaluation of $\langle \text{out} | Q_5(t_{\text{out}}) | \text{out} \rangle_{\text{ren}}$ in (A32), on the other hand, is technically more involved and requires renormalization. This can be done using the adiabatic method [54–56]. Since the main purpose of this appendix was to show the connection with particle pair creation, we just give the final answer without entering into the details, which is

$$\langle \text{out} | Q_5(t_{\text{out}}) | \text{out} \rangle_{\text{ren}} = 2\hbar(qCS[A] - [qCS[A]]). \quad (\text{A38})$$

The final result reads $\langle \text{in} | Q_5(t_{\text{out}}) | \text{in} \rangle_{\text{ren}} = 2\hbar qCS[A]$, and $\langle \text{in} | Q_5(t_{\text{in}}) | \text{in} \rangle_{\text{ren}} = 0$. This agrees, precisely, with the prediction of the Adler-Bell-Jackiw anomaly. In other words, there is a nontrivial evolution of the Noether charge in the quantum theory, which violates the classical symmetry.

As a final remark, notice that, since $(qCS[A] - [qCS[A]]) \in [0, 1[$, unless $[qCS[A]]$ is very small, the particle-creation contribution dominates against the vacuum polarization effect.

Example: Uniform electric field. We address now the problem described at the end of Sec. II. When the electric background field is homogeneous, $E = E(t)$, the Weyl equations read, for any time t ,

$$\begin{aligned} (i\partial_t - i\partial_x - qA_x(t))u_+ &= 0, \\ (i\partial_t + i\partial_x + qA_x(t))u_- &= 0. \end{aligned} \quad (\text{A39})$$

In our gauge choice we have $E(t) = \partial_t A_x(t)$, with $A_x(t \rightarrow -\infty) \rightarrow 0$ and $A_x(t \rightarrow +\infty) \rightarrow A$, for some real-valued constant A . The Weyl equations can be solved in full closed form. The properly normalized in modes defined above are

$$f_{\pm,n}^{\text{in}}(t, x) = \frac{1}{\sqrt{L}} e^{\mp i \frac{2\pi n}{L} (t \pm x) + q \int_{-\infty}^t A_x(t') dt'}, \quad (\text{A40})$$

while the out modes are

$$f_{\pm,n}^{\text{out}}(t, x) = \frac{1}{\sqrt{L}} e^{\mp i \frac{2\pi n}{L} (t \pm x) - qAt_{\text{out}} + q \int_{t_{\text{out}}}^t A_x(t') dt'}. \quad (\text{A41})$$

Notice that the in modes satisfy the initial condition $f_{\pm,n}^{\text{in}}(t, x) \sim \frac{1}{\sqrt{L}} e^{\mp i \frac{2\pi n}{L} (t \pm x)}$ at early times, while the out

modes satisfy the required final condition $f_{\pm,n}^{\text{out}}(t, x) \sim \frac{1}{\sqrt{L}} e^{\mp i [t(\frac{2\pi n}{L} + qA) \pm \frac{2\pi n}{L} x]}$ at late times.

Using these explicit solutions, we can now calculate the relevant Bogoliubov coefficients in full closed form. It is straightforward to get

$$\begin{aligned} (f_{\pm,n'}^{\text{out}}, f_{\pm,n}^{\text{in}}) &= \frac{1}{L} e^{\mp iq(At_{\text{out}} + \int_{-\infty}^{t_{\text{out}}} dt' A_x(t'))} \int_0^L dx e^{\mp i \frac{2\pi}{L} (t \pm x)(n - n')} \\ &= e^{\mp iq(At_{\text{out}} + \int_{-\infty}^{t_{\text{out}}} dt' A_x(t'))} \delta_{nn'}. \end{aligned} \quad (\text{A42})$$

Note that (i) the result is time independent, as expected from the Dirac inner product, (ii) $|(f_{\pm,n'}^{\text{out}}, f_{\pm,n}^{\text{in}})| = 1$, and (iii) the emergence of $\delta_{nn'}$ is a direct consequence of the homogeneity of the background field.

Let us assume that $qA > 0$ (similar results hold when $qA < 0$). Then from (A16)–(A19) we can simplify [we neglect the irrelevant phase factor of (A42)]

$$n > 0 \text{ (positive energy): } f_{+,n}^{\text{in}}(t, x) = f_{+,n}^{\text{out}}(t, x), \quad (\text{A43})$$

$$n < 0 \text{ (negative energy): } f_{+,n}^{\text{in}}(t, x) = \Theta(-n - qAL/2\pi) f_{+,n}^{\text{out}}(t, x) + \Theta(n + qAL/2\pi) f_{+,n}^{\text{out}}(t, x), \quad (\text{A44})$$

$$n < 0 \text{ (positive energy): } f_{-,n}^{\text{in}}(t, x) = \Theta(-n - qAL/2\pi) f_{-,n}^{\text{out}}(t, x) + \Theta(n + qAL/2\pi) f_{-,n}^{\text{out}}(t, x), \quad (\text{A45})$$

$$n > 0 \text{ (negative energy): } f_{-,n}^{\text{in}}(t, x) = f_{-,n}^{\text{out}}(t, x). \quad (\text{A46})$$

This result shows that there is a number $[qAL/2\pi]$ of negative-frequency, left-moving in modes that transform into positive-frequency, left-moving out modes. This means that the electric field has created $[qAL/2\pi]$ left-moving fermions (with negative electric charge) out of the in vacuum. Similarly, there is a number $[qAL/2\pi]$ of positive-frequency, right-moving in modes that are measured as negative-frequency, right-moving modes by out observers. In other words, the electric field has excited $[qAL/2\pi]$ right-moving antifermions (with positive electric charge). All of these particles have positive helicity \hbar . This explains the net helicity found in the general result (A37).

APPENDIX B: PROOF OF (33)

We include here the technical details and computations of Sec. IV. The starting point is Eqs. (31) and (32). Our task is to plug these expressions in (27) and get (33).

Let us denote by $A_2^{\text{aux}}(u, \theta, \phi)$ the first term on the right-hand side of (32). Using (27), we divide the calculation in three terms. First,

$$\begin{aligned} \int_{-L/2}^{L/2} du \int d\mathbb{S}^2 \text{Im}(A_2^{\text{aux}} \tilde{\Phi}_2^0) &= \int d\mathbb{S}^2 \text{Im} \frac{1}{L} \left\{ \sum_{n,n' \neq 0} \tilde{\Phi}_2^0(n, \theta, \phi) \overline{\tilde{\Phi}_2^0(n', \theta, \phi)} \frac{e^{-i(\omega_n - \omega_{n'})u} - e^{-i\omega_n u_0} e^{i\omega_{n'} u}}{-i\omega_n} (-1)^{n+n'} \right. \\ &\quad \left. + \sum_{n \neq 0} \tilde{\Phi}_2^0(n, \theta, \phi) \frac{e^{-i\omega_n u} - e^{-i\omega_n u_0}}{-i\omega_n} \overline{\tilde{\Phi}_2^0(0, \theta, \phi)} (-1)^n \right\} \\ &= \int d\mathbb{S}^2 \text{Im} \frac{1}{L} \left\{ \sum_{n \neq 0} L \frac{|\tilde{\Phi}_2^0(n, \theta, \phi)|^2}{-i\omega_n} + \frac{1}{i} L \overline{\tilde{\Phi}_2^0(0, \theta, \phi)} \cdot \sum_{n \neq 0} \tilde{\Phi}_2^0(n, \theta, \phi) \frac{e^{-i\omega_n u_0}}{\omega_n} (-1)^n \right\} \\ &= \int d\mathbb{S}^2 \sum_{n \neq 0} \frac{|\tilde{\Phi}_2^0(n, \theta, \phi)|^2}{\omega_n} - \text{Re} \left(\overline{\tilde{\Phi}_2^0(0, \theta, \phi)} \sum_{n \neq 0} \tilde{\Phi}_2^0(n, \theta, \phi) \frac{e^{-i\omega_n u_0}}{\omega_n} (-1)^n \right), \end{aligned} \quad (\text{B1})$$

where in the second equality we used the orthonormal properties of the basis modes (30), and the identity $\int_{-L/2}^{L/2} du e^{i2\pi n u/L} = 0$ to get rid of some terms. Second,

$$\begin{aligned}
& \int_{-L/2}^{L/2} du \int d\mathbb{S}^2 \text{Im} \left\{ \left(\frac{-u_0}{\sqrt{L}} \tilde{\Phi}_2(0, \theta, \phi) + A_2^0(u_0, \theta, \phi) \right) \left(\sum_{n \neq 0} \overline{\tilde{\Phi}_2}(n, \theta, \phi) \frac{e^{i\omega_n u} (-1)^n}{\sqrt{L}} + \frac{1}{\sqrt{L}} \overline{\tilde{\Phi}_2(0, \theta, \phi)} \right) \right\} \\
&= L \int d\mathbb{S}^2 \text{Im} \left\{ \left(\frac{-u_0}{\sqrt{L}} \tilde{\Phi}_2(0, \theta, \phi) + A_2^0(u_0, \theta, \phi) \right) \frac{1}{\sqrt{L}} \overline{\tilde{\Phi}_2(0, \theta, \phi)} \right\} \\
&= \sqrt{L} \int d\mathbb{S}^2 \text{Im}(A_2^0(u_0, \theta, \phi) \overline{\tilde{\Phi}_2(0, \theta, \phi)}) \\
&= \int_{-L/2}^{L/2} du d\mathbb{S}^2 \text{Im}(A_2^0(u_0, \theta, \phi) \overline{\tilde{\Phi}_2(0, \theta, \phi)}), \tag{B2}
\end{aligned}$$

where in the first equality we used the identity $\int_{-L/2}^{L/2} du e^{i2\pi n u/L} = 0$; in the second equality we noticed that one term was real, so its imaginary part vanishes; and in the last equality we recalled the definition of Fourier transform. Finally,

$$\begin{aligned}
& \int_{-L/2}^{L/2} du \int d\mathbb{S}^2 \text{Im} \left\{ \frac{u}{\sqrt{L}} \tilde{\Phi}_2(0, \theta, \phi) \left(\sum_{n \neq 0} \overline{\tilde{\Phi}_2(\omega_n, \theta, \phi)} \frac{e^{+i\omega_n u}}{\sqrt{L}} (-1)^n + \frac{1}{\sqrt{L}} \overline{\tilde{\Phi}_2(0, \theta, \phi)} \right) \right\} \\
&= \int_{-L/2}^{L(2)} du \int d\mathbb{S}^2 \text{Im} \left\{ \frac{u}{\sqrt{L}} \tilde{\Phi}_2(0, \theta, \phi) \sum_{n \neq 0} \overline{\tilde{\Phi}_2(\omega_n \theta, \phi)} \frac{e^{i\omega_n u}}{\sqrt{L}} (-1)^n \right\} \\
&= \int_{-L/2}^{L(2)} du \int d\mathbb{S}^2 \text{Im} \left\{ \frac{1}{L} \tilde{\Phi}_2(0, \theta, \phi) \sum_{n \neq 0} \overline{\tilde{\Phi}_2(\omega_n \theta, \phi)} e^{i\omega_n u} (-1)^n \frac{1}{i} \frac{d}{d\epsilon} \Big|_{\epsilon=0} e^{i\epsilon u} \right\} \\
&= \int d\mathbb{S}^2 \text{Im} \left\{ \frac{1}{L} \tilde{\Phi}_2(0, \theta, \phi) \sum_{n \neq 0} \overline{\tilde{\Phi}_2(\omega_n \theta, \phi)} (-1)^n \frac{1}{i} \frac{d}{d\epsilon} \Big|_{\epsilon=0} \int_{-L/2}^{L(2)} du e^{i(\epsilon+\omega_n)u} \right\} \\
&= \int d\mathbb{S}^2 \text{Im} \left\{ \frac{1}{L} \tilde{\Phi}_2(0, \theta, \phi) \sum_{n \neq 0} \overline{\tilde{\Phi}_2(\omega_n \theta, \phi)} \frac{L}{i\omega_n} \right\} \\
&= - \int d\mathbb{S}^2 \text{Re} \left\{ \tilde{\Phi}_2(0, \theta, \phi) \sum_{n \neq 0} \frac{\overline{\tilde{\Phi}_2(\omega_n \theta, \phi)}}{\omega_n} \right\}, \tag{B3}
\end{aligned}$$

where in the first equality we noted that one term has vanishing imaginary part; and in the fourth equality we used $\int_{-L/2}^{L(2)} du e^{i(\epsilon+\omega_n)u} = (-1)^n \frac{e^{i\epsilon L/2} - e^{-i\epsilon L/2}}{i(\omega_n + \epsilon)}$.

Combining all three terms above, and for $u_0 = -L/2$, which allows to further simplify some terms, we end up with

$$\begin{aligned}
& \int_{-L/2}^{L/2} du \int d\mathbb{S}^2 \text{Im} \{ (A_2^0 - \bar{\delta}\alpha_0) \tilde{\Phi}_2 \} \\
&= \int d\mathbb{S}^2 \left\{ \sum_{n \neq 0} \frac{|\tilde{\Phi}_2(n, \theta, \phi)|^2}{\omega_n} - 2 \text{Re} \left(\overline{\tilde{\Phi}_2(0, \theta, \phi)} \frac{\tilde{\Phi}_2(n, \theta, \phi)}{\omega_n} \right) \right\} + \int_{-L/2}^{L/2} du d\mathbb{S}^2 \text{Im}(A_2^0(-L/2, \theta, \phi) - \bar{\delta}\alpha_0) \overline{\tilde{\Phi}_2(u, \theta, \phi)}. \tag{B4}
\end{aligned}$$

Note that $\sum_{n \neq 0} \frac{|\tilde{\Phi}_2^0(0, \theta, \phi)|^2}{\omega_n} \sim \sum_{n \neq 0} \frac{1}{n} = 0$. Therefore, we can rewrite the first term above as $\int d\mathbb{S}^2 \sum_{n \neq 0} \frac{|\tilde{\Phi}_2^0(n, \theta, \phi) - \tilde{\Phi}_2^0(0, \theta, \phi)|^2}{\omega_n}$. The second term above can also be greatly simplified and has a nice physical interpretation. To see this, note first that, in the gauge $A_1^0 = A_a n^a = 0$, the one-form A_a^0 lives on the tangent space of \mathbb{S}^2 , so it can be expressed as the sum of a gradient and a curl: $A_a^0 = D_a \alpha + \epsilon_a^b D_b \beta$, where D_a is the connection on \mathbb{S}^2 . Thus, $A_2^0 = \bar{\delta}(\alpha + i\beta)$, with $\alpha, \beta \in C^\infty(\mathbb{R} \times \mathbb{S}^2, \mathbb{R})$. There still exists a residual gauge freedom represented by $A_a^0 \rightarrow A_a^0 + D_a \Lambda$, with $\Lambda = 0$. Under this transformation β remains invariant, and although α does not remain invariant, the combination $\alpha - \alpha_0$ does. With this new terminology we can reexpress the second term on the right-hand side above as

$$\begin{aligned}
\int_{-L/2}^{L/2} du d\mathbb{S}^2 \text{Im}(A_2^0(-L/2, \theta, \phi) - \delta\alpha_0) \overline{\Phi_2^0(u, \theta, \phi)} &= \int_{-L/2}^{L/2} du d\mathbb{S}^2 \text{Im}(i\bar{\delta}\beta + \bar{\delta}(\alpha - \alpha_0)) \overline{\Phi_2^0(u, \theta, \phi)} \\
&= - \int_{-L/2}^{L/2} du d\mathbb{S}^2 \beta \text{Re} \overline{\delta\Phi_2^0(u, \theta, \phi)} + (\alpha - \alpha_0)\beta \text{Im} \overline{\delta\Phi_2^0(u, \theta, \phi)} \\
&= -\text{Re}q_\beta + \text{Im}q_{\alpha-\alpha_0},
\end{aligned} \tag{B5}$$

where in the last step we made use of Maxwell equations (15) and the definition of infrared charges (20).

As we can see this contribution in the chiral anomaly is related to memory of the electromagnetic background. Despite $\Phi_2^0(\pm\frac{L}{2}, \theta, \phi) = 0$, if $\Phi_2^0(u, \theta, \phi) \neq 0$, then $\beta(u, \theta, \phi)$ and $\alpha(u, \theta, \phi) - \alpha_0(u, \theta, \phi)$ evolve in time u , leading to nonzero soft charges. Using (18) and (19), these functions can be solved from $\text{Re}\Phi_1^0(u, \theta, \phi) = \Delta\alpha(u, \theta, \phi) + G(\theta, \phi)$ and $\text{Im}\Phi_1^0(u, \theta, \phi) = \Delta\beta(u, \theta, \phi)$, where Δ denotes the two-dimensional Laplacian. The parameter $\beta(-L/2, \theta, \phi)$ tells us about the magnetic field at spatial infinity i^0 when we take $L \rightarrow \infty$, while the combination $\alpha(-L/2, \theta, \phi) - \alpha_0(-L/2, \theta, \phi)$ tells us about the electric field. The residual gauge freedom mentioned above can be fully fixed by choosing G such that $G(\theta, \phi) = -\frac{Q}{2}$, where $Q = \frac{1}{2\pi} \int_{\mathbb{S}^2} d\mathbb{S}^2 \text{Re}\Phi_1^0$ is the (constant) electric charge of the sources. Because α_0 was chosen such that $A_a^0 = D_a\alpha_0$ produces $\Phi_2^0 = \Phi_1^0 = 0$, in this fully fixed gauge choice α_0 satisfies $\Delta\alpha_0 = 0$, which for smooth functions on the sphere is equivalent to $\alpha_0 = 0$. In summary, for a fully gauge-fixed theory the final result

reads

$$\begin{aligned}
&\int_{-\infty}^{\infty} du \int d\mathbb{S}^2 \text{Im}\{(A_2^0 - \bar{\delta}\alpha_0)\bar{\Phi}_2^0\} \\
&= \frac{1}{2\pi} \int d\mathbb{S}^2 \int_{-\infty}^{\infty} d\omega \frac{|\tilde{\Phi}_2^0(\omega, \theta, \phi) - \tilde{\Phi}_2^0(0, \theta, \phi)|^2}{\omega} \\
&\quad + \text{Im}q_\alpha - \text{Re}q_\beta,
\end{aligned} \tag{B6}$$

where the infrared charges are evaluated for $\alpha(-\infty, \theta, \phi)$, $\beta(-\infty, \theta, \phi)$.

Note that the integral is well defined in the infrared limit $\omega \rightarrow 0$. On the other hand, when the soft charges are all zero, we also have $\tilde{\Phi}_2^0(0, \theta, \phi) = 0$, and this expression reduces to the result obtained in [16]. As shown in [16], the first contribution on the right-hand side above represents the difference between right-handed and left-handed circularly polarized radiation reaching future null infinity. This is a purely radiative contribution.

The contribution from the IR charges in (B6) can be rewritten in a different form:

$$\begin{aligned}
\text{Im}q_\alpha - \text{Re}q_\beta &= \int d\mathbb{S}^2 \alpha(-\infty, \theta, \phi) (\text{Im}\Phi_1^0(\infty, \theta, \phi) - \text{Im}\Phi_1^0(-\infty, \theta, \phi)) \\
&\quad - \int d\mathbb{S}^2 \beta(-\infty, \theta, \phi) (\text{Re}\Phi_1^0(\infty, \theta, \phi) - \text{Re}\Phi_1^0(-\infty, \theta, \phi)) \\
&= \int d\mathbb{S}^2 \alpha(-\infty, \theta, \phi) \Delta(\beta(\infty, \theta, \phi) - \beta(-\infty, \theta, \phi)) - \int d\mathbb{S}^2 \beta(-\infty, \theta, \phi) \Delta(\alpha(\infty, \theta, \phi) - \alpha(-\infty, \theta, \phi)) \\
&= \int d\mathbb{S}^2 [\alpha(-\infty, \theta, \phi) \Delta\beta(\infty, \theta, \phi) - \beta(-\infty, \theta, \phi) \Delta\alpha(\infty, \theta, \phi)].
\end{aligned} \tag{B7}$$

As a final remark, note that Eq. (B6) can also be obtained if we had chosen $u_0 = +L/2$ to write (B4) instead of $u_0 = -L/2$. The only difference is that α and β in that equation would be evaluated at $u = +L/2$ and not at $u = -L/2$. Although apparently different, the two expressions are actually equivalent. The equivalence is manifest from (B7), which implies $\text{Im}q_{\alpha(\infty)} - \text{Re}q_{\beta(\infty)} = \text{Im}q_{\alpha(-\infty)} - \text{Re}q_{\beta(-\infty)}$ after integration by parts. Equation (B6) with this last choice is the result of Eq. (33) of the main text.

[1] R. Bertlmann, *Anomalies in Quantum Field Theory*, International Series of Monographs on Physics Vol. 91 (Clarendon Press, Oxford, England, 1996).

[2] S. L. Adler, *Phys. Rev.* **177**, 2426 (1969).

[3] J. S. Bell and R. Jackiw, *Il Nuovo Cimento A (1965–1970)* **60**, 47 (1969).

[4] T. Kimura, *Prog. Theor. Phys.* **42**, 1191 (1969).

[5] R. Delbourgo and A. Salam, *Phys. Lett.* **40B**, 381 (1972).

[6] T. Eguchi and P. G. O. Freund, *Phys. Rev. Lett.* **37**, 1251 (1976).

[7] M. D. Schwartz, *Quantum Field Theory and the Standard Model* (Cambridge University Press, Cambridge, England, 2014).

[8] T. Eguchi, P. B. Gilkey, and A. J. Hanson, *Phys. Rep.* **66**, 213 (1980).

[9] M. Nakahara, *Geometry, Topology and Physics*, Graduate Student Series in Physics (Hilger, Bristol, England, 1990).

[10] F. Bastianelli and P. van Nieuwenhuizen, *Path Integrals and Anomalies in Curved Space*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 2006).

[11] I. Agullo, A. del Rio, and J. Navarro-Salas, *Phys. Rev. Lett.* **118**, 111301 (2017).

[12] I. Agullo, A. del Rio, and J. Navarro-Salas, *Int. J. Mod. Phys. D* **26**, 1742001 (2017).

[13] I. Agullo, A. del Rio, and J. Navarro-Salas, *Phys. Rev. D* **98**, 125001 (2018).

[14] I. Agullo, A. del Rio, and J. Navarro-Salas, *Symmetry* **10**, 763 (2018).

[15] A. del Rio, N. Sanchis-Gual, V. Mewes, I. Agullo, J. A. Font, and J. Navarro-Salas, *Phys. Rev. Lett.* **124**, 211301 (2020).

[16] A. del Rio, *Phys. Rev. D* **104**, 065012 (2021).

[17] N. Sanchis-Gual and A. del Rio, *Phys. Rev. D* **108**, 044052 (2023).

[18] A. Ashtekar, M. Campiglia, and A. Laddha, *Gen. Relativ. Gravit.* **50**, 140 (2018).

[19] A. Strominger, *Lectures on the Infrared Structure of Gravity and Gauge Theory* (Princeton University Press, Princeton, NJ, 2017).

[20] L. Bieri and D. Garfinkle, *Classical Quantum Gravity* **30**, 195009 (2013).

[21] B. Oblak and A. Seraj, [arXiv:2304.12348](https://arxiv.org/abs/2304.12348).

[22] A. Seraj and B. Oblak, [arXiv:2112.04535](https://arxiv.org/abs/2112.04535).

[23] R. Jackiw, in *Current Algebra and Anomalies*, edited by S. B. Treiman *et al.* (World Scientific, Singapore, 1985), p. 211.

[24] Y. Aharonov and D. Bohm, *Phys. Rev.* **115**, 485 (1959).

[25] R. Geroch, in *Proceedings of a Symposium on Asymptotic Structure of Space-Time*, edited by F. Esposito and L. Witten (University of Cincinnati, Cincinnati, OH, 1977), pp. 1–105.

[26] A. Ashtekar, *Asymptotic Quantization*, based on 1984 Naples Lectures (Bibliopolis, Naples, 1987).

[27] A. Ashtekar, in *Surveys in Differential Geometry*, edited by L. Bieri and S. T. Yau (International Press, Boston, MA, 2015), p. 99.

[28] R. Penrose, *Phys. Rev. Lett.* **10**, 66 (1963).

[29] E. Newman and R. Penrose, *J. Math. Phys. (N.Y.)* **3**, 566 (1962).

[30] J. Stewart, *Advanced General Relativity*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 1991).

[31] A. Ashtekar and B. Bonga, *Classical Quantum Gravity* **34**, 20LT01 (2017).

[32] A. Ashtekar and B. Bonga, *Gen. Relativ. Gravit.* **49**, 122 (2017).

[33] S. Weinberg, *Phys. Rev.* **140**, B516 (1965).

[34] R. Jackiw, *Rev. Mod. Phys.* **49**, 681 (1977).

[35] M. A. Shifman, *Instantons in Gauge Theories* (World Scientific, Singapore, 1994).

[36] D. J. Griffiths, *Introduction to Electrodynamics*, 4th ed. (Pearson, Boston, MA, 2013).

[37] J. Winicour, *Classical Quantum Gravity* **31**, 205003 (2014).

[38] B. Zel'dovich and A. G. Polnarev, *Sov. Astron.* **18**, 17 (1974).

[39] V. B. Braginsky and L. P. Grishchuk, *Sov. Phys. JETP* **62**, 427 (1985).

[40] V. B. Braginskii and K. S. Thorne, *Nature (London)* **327**, 123 (1987).

[41] D. C. Christodoulou, *Phys. Rev. Lett.* **67**, 1486 (1991).

[42] K. S. Thorne, *Phys. Rev. D* **45**, 520 (1992).

[43] J. Frauendiener, *Classical Quantum Gravity* **9**, 1639 (1992).

[44] L. Bieri and D. Garfinkle, *Phys. Rev. D* **89**, 084039 (2014).

[45] A. Ashtekar, *Phys. Rev. Lett.* **46**, 573 (1981).

[46] A. Ashtekar, *J. Math. Phys. (N.Y.)* **22**, 2885 (1981).

[47] G. 't Hooft, *Phys. Rev. Lett.* **37**, 8 (1976).

[48] A. Ashtekar and K. S. Narain, *Presented at the VI International Conference on Mathematical Physics* (Syracuse University, Berlin, 1981).

[49] H. B. Nielsen and M. Ninomiya, *Phys. Lett.* **130B**, 389 (1983).

[50] J. Ambjorn, J. Greensite, and C. Peterson, *Nucl. Phys. B* **221**, 381 (1983).

[51] L. Alvarez-Gaume and M. A. Vazquez-Mozo, *An Invitation to Quantum Field Theory*, Lecture Notes in Physics (Springer, Berlin, Heidelberg, 2012).

[52] L. Parker and D. J. Toms, *Quantum Field Theory in Curved Spacetime: Quantized Fields and Gravity* (Cambridge University Press, Cambridge, England, 2009).

[53] C. Baer and A. Strohmaier, *Commun. Math. Phys.* **347**, 703 (2016).

[54] A. Ferreiro and J. Navarro-Salas, *Phys. Rev. D* **97**, 125012 (2018).

[55] J. F. Barbero, A. Ferreiro, J. Navarro-Salas, and E. J. S. Villasenor, *Phys. Rev. D* **98**, 025016 (2018).

[56] A. Ferreiro, J. Navarro-Salas, and S. Pla, *Phys. Rev. D* **98**, 045015 (2018).