



ASYMPTOTIC SPREADING FOR FISHER-KPP REACTION-DIFFUSION EQUATIONS WITH HETEROGENEOUS SHIFTING DIFFUSIVITY

GRÉGORY FAYE*

Institut de Mathématiques de Toulouse
UMR 5219, Université de Toulouse
UPS IMT, F-31062 Toulouse Cedex 9, France

THOMAS GILETTI

IECL ; UMR 7502, University of Lorraine
B.P. 70239, F-54506 Vandoeuvre-lès-Nancy Cedex, France

MATT HOLZER

Department of Mathematical Sciences
Center for Mathematics and Artificial Intelligence (CMAI)
George Mason University, Fairfax, VA, USA

ABSTRACT. We determine the asymptotic spreading speed of the solutions of a Fisher-KPP reaction-diffusion equation, starting from compactly supported initial data, when the diffusion coefficient is a fixed bounded monotone profile that is shifted at a given forcing speed and satisfies a general uniform ellipticity condition. Depending on the monotonicity of the profile, we are able to characterize this spreading speed as a function of the forcing speed and the two linear spreading speeds associated to the asymptotic problems at $x = \pm\infty$. Most notably, when the profile of the diffusion coefficient is increasing we show that there is an intermediate range for the forcing speed where spreading actually occurs at a speed which is larger than the linear speed associated with the homogeneous state around the position of the front. We complement our study with the construction of strictly monotone traveling front solutions with strong exponential decay near the unstable state when the profile of the diffusion coefficient is decreasing and in the regime where the forcing speed is precisely the selected spreading speed.

1. Introduction. In this work we aim at understanding the influence of a heterogeneous diffusion on the spreading speed of solutions of a scalar reaction-diffusion equation of the Fisher-KPP type [15, 21]. More precisely, we consider the situation of a shifting environment, i.e. the diffusion coefficient depends on a shifting variable

2020 *Mathematics Subject Classification.* Primary: 35K57, 35B40, 35K45, 35C07.

Key words and phrases. reaction-diffusion, spreading speed, heterogeneous diffusion, traveling waves.

GF acknowledges support from an ANITI (Artificial and Natural Intelligence Toulouse Institute) Research Chair and from Labex CIMI under grant agreement ANR-11-LABX-0040. GF and TG acknowledge support from the ANR project Indyana under grant agreement ANR-21-CE40-0008-01. The research of MH was partially supported by the National Science Foundation (DMS-2007759).

*Corresponding author: Grégory Faye.

$x - c_{het}t$ where the parameter $c_{het} \in \mathbb{R}$. The equation we consider can be written as follows:

$$\partial_t u = \chi(x - c_{het}t) \partial_x^2 u + \alpha u(1 - u), \quad t > 0, x \in \mathbb{R}, \quad (1)$$

where $\alpha > 0$. Throughout this paper we assume that the function $\chi : \mathbb{R} \mapsto \mathbb{R}$ is positive and smooth; further assumptions will be added below.

Such a reaction-diffusion equation is often used in ecology and population dynamics to model spatial propagation phenomena [1]. In this context, the unknown function u denotes the density of some species, and the parameter α stands for its per capita growth rate. The function χ stands for the motility of individuals, and therefore the heterogeneity means that the motility depends on local environmental conditions. In the past decade, there has been several series of works dedicated to the study of a variation of (1) with constant motility (say $\chi \equiv 1$ to fix the ideas) but with shifting growth rates, that typically reads

$$\partial_t u = \partial_x^2 u + u(r(x - c_{het}t) - u),$$


where the function r is not necessarily monotone and can take negative values [8, 2, 10, 25, 3]. One of the main motivations behind the introduction of a shifting growth rate came from modeling climate change and its impact on the survival of species [2, 12]. Some spreading-vanishing dichotomy properties were typically found, depending on either the size of a favorable patch [2] or of the initial population [10].

Here our choice of a shifting heterogeneity (i.e. depending only on a shifting variable $x - c_{het}t$) comes from systems coupling several species densities. Most of the mathematical literature on spreading speeds for systems has been concerned with situations where these species only interact through their growth rates, e.g. prey-predator and competition systems [13, 18, 19, 23]. Yet we would like to understand the influence that such inter-species interactions may play on the respective motilities of each species. For instance, it seems natural that the motility of predators should be higher when the density of the prey is low, while the motility of preys should be higher when the density of predators is high.

To illustrate this, let us consider the reaction-diffusion system,

$$\begin{cases} \partial_t u = d(v) \partial_x^2 u + \alpha u(1 - u), \\ \partial_t v = \partial_x^2 v + \beta v(1 - v), \end{cases} \quad (2)$$

where $\beta > 0$ and $d : \mathbb{R}^+ \mapsto \mathbb{R}$ is a positive function. Here u and v denote two distinct species, which are coupled through the motility of individuals of the former species. For simplicity we have excluded other coupling terms, and in particular there are no predation nor competition terms in the growth rates; this allows us to focus purely on the cross-diffusion phenomenon. As suggested above, the function d may be taken as a monotonic function depending on whichever species acts as the predator. More precisely, if u is the prey then the motility function d should be increasing as the number of predators v grows; conversely, if u is the predator then its motility d should be decreasing as the number of preys v grows. Other forms of cross-diffusion have been proposed and studied in the literature, typically when the diffusion term is either of the form $\partial_x^2(d(v)u)$ or $\partial_x(d(v)\partial_x u)$ [27, 24, 17]. While such systems have been considered for instance in the context of chemotaxis, little is known from the point of view of propagation phenomena, apart from perturbative or singular limit results that reduce the system to a weakly coupled system which allow the construction of traveling front solutions (see for example [16] in the context of cancer modeling).

FIGURE 1. Illustration of χ in cases (I) (left) and (II) (right).

It is well-known that, for a large class of initial data, the solution of the Fisher-KPP equation

$$\partial_t v = \partial_x^2 v + \beta v(1 - v),$$

converges to a traveling front solution, i.e. an entire in time solution $v(t, x) = V(x - ct)$ where V is a decreasing function and satisfies

$$V(-\infty) = 1, \quad V(+\infty) = 0.$$

More precisely, such a traveling front solution exists if and only if $c \geq 2\sqrt{\beta}$, and the traveling front with minimal speed is typically the most biologically meaningful because it is attractive with respect to compactly supported initial conditions (up to some drift phenomenon) [15, 21, 1]. Due to the unilateral coupling in system (2), it is thus natural to assume that v is identical to the traveling front with minimal speed, and then the equation for u reduces to (1) with $c_{het} = 2\sqrt{\beta}$. In other words, the shifting heterogeneity arises as a result of the propagation of another surrounding species, which individuals of the species u may either chase or flee from.

Remark 1. From the above discussion, it may seem natural to consider a slightly more general heterogeneous diffusion $\tilde{\chi}(t, x)$, which converges as $t \rightarrow +\infty$ to $\chi(x - c_{het}t)$, or even to $\chi(x - c_{het}t + m(t))$ where $m(t) = o(t)$ as $t \rightarrow +\infty$ accounts for the drift phenomenon. One may check that most of our analysis applies to such situations with straightforward adaptations. We choose to only consider (1) since it makes the presentation simpler and captures the same phenomena.

Following this motivation, we will study the large-time behavior of solutions of (1) with $c_{het} \geq 0$ and under either assumptions

$$\begin{cases} \chi > 0, & \chi' \leq 0, \\ \chi(-\infty) = d_+, & \chi(+\infty) = d_-, \end{cases} \quad (\text{I})$$

or

$$\begin{cases} \chi > 0, & \chi' \geq 0, \\ \chi(-\infty) = d_-, & \chi(+\infty) = d_+, \end{cases} \quad (\text{II})$$

where

$$0 < d_- < d_+ < +\infty.$$

Let us point out that, equivalently, one may fix the function χ satisfying either assumption, allow c_{het} to vary on the whole real line $\mathbb{R}$ and investigate spreading in corresponding directions. Here we will instead pick c_{het} nonnegative and study spreading only to the right, i.e. in the direction of the moving heterogeneity. While spreading in the opposite direction may also be investigated, there the solution is expected to behave just like the solution of the homogeneous problem with diffusion

coefficient equal to $\chi(-\infty)$. This can easily be proved by similar and even simpler arguments than those we develop here, hence we will omit it.

Finally, equation (1) will be supplemented together with some initial condition u_0 such that

$$0 \leq u_0 \leq 1 \text{ and it is nontrivial and compactly supported.}$$

2. Main results. In the first case when (I) holds, then the ‘high diffusion’ part of the environment where individuals diffuse faster is growing with speed c_{het} , and therefore one may expect that the solution will be spreading with positive speed. On the other hand, in the second case when (II) holds, then the ‘high diffusion’ part of the environment is actually receding and a key point will be whether the solution is able to ‘keep up’ with the shifting speed c_{het} of the heterogeneity.

This leads us to introduce, before we state our main results, the speeds

$$c_+ := 2\sqrt{d_+\alpha}, \quad c_- := 2\sqrt{d_-\alpha},$$

which correspond respectively to the speeds of the homogeneous Fisher-KPP equation

$$\partial_t u = d\partial_x^2 u + \alpha u(1-u),$$

with either $d = d_+$ or $d = d_-$. In other words, these are the speeds respectively in the ‘high diffusion’ environment where the motility function χ is at its maximal value, and in the ‘low diffusion’ environment where χ is at its minimal value. The large time behavior of solutions of (1) will largely depend on how the shifting speed c_{het} of the heterogeneity compares with the values of c_- and c_+ .

2.1. Case (I). We first consider the situation when the diffusivity is ‘high’ (resp. ‘low’) behind (resp. beyond) the moving frame with speed c_{het} . In this case, we will prove the following result:

Theorem 2.1. *Assume that χ satisfies case (I). Consider a compactly supported and nontrivial initial datum u_0 such that $0 \leq u_0 \leq 1$. Then the solution of (1) spreads to the right with some positive speed c_u^* in the sense that*

$$\begin{aligned} \forall 0 < c < c_u^*, \quad \limsup_{t \rightarrow +\infty} \sup_{0 \leq x \leq ct} |1 - u(t, x)| &= 0, \\ \forall c > c_u^*, \quad \limsup_{t \rightarrow +\infty} \sup_{x \geq ct} u(t, x) &= 0. \end{aligned}$$

Moreover, we have either:

$$c_u^* = c_- \quad \text{if} \quad c_{het} < c_-,$$

or

$$c_u^* = c_{het} \quad \text{if} \quad c_- \leq c_{het} \leq c_+,$$

or

$$c_u^* = c_+ \quad \text{if} \quad c_+ < c_{het}.$$

It is indeed natural that the spreading speed c_u^* (if it exists) should be less than both c_+ (which is the speed when diffusivity is maximal) and the maximum of c_{het} and c_- (since the solution may not spread faster than c_- in the part where diffusivity is minimal beyond $x \approx c_{het}t$). This theorem states that these intuitive upper bounds give precisely the spreading speed. We refer to Figure 2 (a) for a numerical illustration.

We also note that when $c_{het} \in [c_-, c_+]$, the selected spreading speed c_u^* is precisely the speed c_{het} of the heterogeneity and the invasion process for the u component is




FIGURE 2. Numerically computed spreading speed c_u^* (pink circles) as a function of c_{het} for Case (I) (left) and Case (II) (right). The purple plain line is the theoretical spreading speed provided by Theorem 2.1 and Theorem 2.3. In both cases parameters are fixed with $\alpha = 1$, $d_+ = 1$ and $d_- = 1/4$, such that the corresponding linear speeds are $c_+ = 2$ and $c_- = 1$. The function χ was set to $\chi(x) = \frac{d_+ e^{-\lambda x} + d_-}{1 + e^{-\lambda x}}$ in Case (I) and to $\chi(x) = \frac{d_- e^{-\lambda x} + d_+}{1 + e^{-\lambda x}}$ in Case (II) with $\lambda = 2$. Numerical simulations were performed by discretizing equation (1) via finite differences in space and a semi-implicit scheme in time. Typical discretization step sizes were set to $\delta t = 0.02$ in time and $\delta x = 0.02$ in space.

locked to the heterogeneity; see also [14]. We expect that in that case, the dynamics is dictated by traveling wave solutions U of

$$0 = \chi(x)U''(x) + c_{het}U'(x) + \alpha U(x)(1 - U(x)), \quad x \in \mathbb{R}, \quad (3)$$

that satisfy the conditions

$$U(-\infty) = 1, \quad U(+\infty) = 0, \quad \text{with} \quad 0 < U < 1. \quad (4)$$

In order to state our second main result, we need to further assume the following assumption

$$|\chi(x) - d_{\pm}| = \mathcal{O}(e^{-\nu|x|}), \quad |\chi'(x)| = \mathcal{O}(e^{-\nu|x|}), \quad \text{as } x \rightarrow \pm\infty, \quad (5)$$

for some $\nu > 0$. The above assumption is natural if we come back to the reaction-diffusion system (2) for which the traveling front solutions $v(t, x) = V(x - ct)$ for $c \geq 2\sqrt{\beta}$ are known to converge at an exponential rate towards their asymptotic limit states.

Theorem 2.2. *Assume that χ satisfies case (I) and that (5) is verified. For each $c_{het} \in (c_-, c_+)$, there exists a unique strictly monotone traveling wave solution U of (3)-(4) with strong exponential decay at $+\infty$:*

$$U(x) \underset{x \rightarrow +\infty}{\sim} \gamma_s e^{-\lambda_s x}, \quad \lambda_s := \frac{c_{het} + \sqrt{c_{het}^2 - c_-^2}}{2d_-},$$

for some $\gamma_s > 0$. For $c_{het} \in [0, c_-)$ or $c_{het} > c_+$ no such traveling wave can exist.

While there may exist other traveling wave solutions with a different (slower) exponential decay at $+\infty$, we conjecture that, for compactly supported initial data, the solution $u(t, x)$ of (1) converges as $t \rightarrow +\infty$, in the moving frame with speed c_{het} , to the traveling wave U provided by Theorem 2.2. However, we do not address this issue here.

2.2. Case (II). The other situation turns out to be slightly more intricate. Our main result writes as follows:

Theorem 2.3. *Assume that χ satisfies case (II) and that (5) is verified. Consider a compactly supported and nontrivial initial datum u_0 such that $0 \leq u_0 \leq 1$. Then the solution of (1) spreads to the right with some speed c_u^* in the sense that*

$$\forall 0 < c < c_u^*, \quad \limsup_{t \rightarrow +\infty} \sup_{0 \leq x \leq ct} |1 - u(t, x)| = 0,$$

$$\forall c > c_u^*, \quad \limsup_{t \rightarrow +\infty} \sup_{x \geq ct} u(t, x) = 0.$$

Moreover, we have

$$c_u^* = c_+ \quad \text{if} \quad c_{het} < c_+,$$

while

$$c_u^* = \frac{c_{het}}{2} \left(1 - \sqrt{1 - \frac{d_-}{d_+}} \right) + \frac{c_-^2}{2c_{het} \left(1 - \sqrt{1 - \frac{d_-}{d_+}} \right)} \in (c_-, c_+] \quad \text{if} \quad c_+ \leq c_{het} < c_{int},$$

and

$$c_u^* = c_- \quad \text{if} \quad c_{int} \leq c_{het}.$$

Here

$$c_{int} := c_+ \left(\sqrt{\frac{d_+}{d_-}} + \sqrt{\frac{d_+}{d_-} - 1} \right).$$

Again we find several subcases depending on the value of c_{het} . When $c_+ > c_{het}$, then individuals move fast enough to keep up with the ‘high diffusion’ zone beyond the front of the heterogeneity, so that propagation reaches it full speed c_+ . In particular, the population spreads as if there was no heterogeneity.

The case when $c_+ < c_{het}$ is less intuitive. Because the solution cannot spread faster than c_+ and therefore it vanishes in the ‘high diffusion’ zone as time goes to infinity, one may have expected that the solution behaves as in the equation with lower diffusivity d_- and thus spreads with speed c_- . According to Theorem 2.3, it turns out that this intuition is true if c_{het} is large enough. Yet there exists some intermediate range where spreading actually occurs with speed c_u^* which is strictly larger than c_- . This means that the far away ‘high diffusion’ zone still plays a role in the propagation, which can be related to the phenomena of accelerated fronts or nonlocal pulling [18, 19]. We refer to Figure 2 (b) for a numerical illustration.

This observation is especially striking when d_- is small. Indeed, let us take the formal limit $d_- = 0$ in Theorem 2.3. Then $c_{int} = c_+ \left(\sqrt{\frac{d_+}{d_-}} + \sqrt{\frac{d_+}{d_-} - 1} \right) \rightarrow +\infty$ and it can also be checked that $c_u^* \rightarrow \frac{4d_+ \alpha}{c_{het}} > 0$ for any $c_{het} > c_+$ as $d_- \rightarrow 0$. This strikingly suggests that the spreading speed is bounded by a positive constant uniformly with respect to d_- , and even as the diffusivity vanishes around any positive level set of the solution. We refer to Figure 3 for a numerical illustration.




FIGURE 3. Numerically computed spreading speed c_u^* (pink circles) as a function of c_{het} for Case (II) in the degenerate case where $d_- = 0$. The purple plain line is the curve $c_{het} \mapsto \frac{4d_+\alpha}{c_{het}}$ obtained by formally taking the limit $d_- = 0$ in Theorem 2.3. Other parameters are fixed with $\alpha = 1$ and $d_+ = 1$, such that the corresponding linear speed is $c_+ = 2$. The function χ was set to $\chi(x) = \frac{d_+}{1+e^{-\lambda x}}$ with $\lambda = 2$.

Before we proceed to the proofs, let us give some formal computation to explain the appearance of an anomalous speed. Assume that $c_+ < c_{het}$, and that

$$\chi(z) := \begin{cases} d_- & \text{if } z \leq 0, \\ d_+ & \text{if } z > 0. \end{cases}$$

Notice that such χ is no longer smooth but it allows us to perform an explicit computation. Since the reaction term $\alpha u(1-u)$ is concave, it is of the Fisher-KPP type and it is reasonable to expect that the spreading speed is dictated by the linearized equation

$$\partial_t u = \chi(x - c_{het}t) \partial_x^2 u + \alpha u.$$

By analogy with the usual Fisher-KPP equation, it is also natural to look for an exponential ansatz. However, if one tries an ansatz of the type

$$e^{-\lambda(x-ct)},$$

one immediately sees that both resulting dispersion equations (for $x \leq c_{het}t$ and $x > c_{het}t$) cannot be satisfied simultaneously. Therefore, we instead look for an ansatz of the type

$$\begin{cases} e^{-\lambda(x-ct)} & \text{if } x \leq c_{het}t, \\ e^{-\lambda(c_{het}-c)t} \times e^{-\mu(x-c_{het}t)} & \text{if } x \geq c_{het}t, \end{cases}$$

where $\lambda, \mu > 0$ and $c \in \mathbb{R}$. Note that the resulting function is continuous. Plugging this in the linearized equation, we find that

$$d_- \lambda^2 - c\lambda + \alpha = 0 \quad \text{and} \quad d_+ \mu^2 - c_{het}\mu + \lambda(c_{het} - c) + \alpha = 0.$$

On the one hand, it is easy to check that both conditions can be satisfied when $c \geq c_{het}$, by taking λ as any of the two roots of the first equation, and noticing that

the discriminant of the second equation is then positive (recall that $c_{het} > c_+ > c_-$ here). On the other hand, when $c < c_{het}$, the resulting conditions for finding such an ansatz are

$$c \geq c_-, \quad g(c) \geq 0, \quad (6)$$

where

$$g(c) := c_{het}^2 - 4d_+ \left[\alpha + \frac{c - \sqrt{c^2 - 4d_- \alpha}}{2d_-} (c_{het} - c) \right].$$

Notice that in the formula for $g(c)$ we used the smaller positive root for λ , as the larger one would only lead to a more restrictive condition. Now it is straightforward to compute that

$$\forall c \in [c_-, c_{het}], \quad g'(c) > 0.$$

In particular, if $g(c_-) \geq 0$, then we conclude that there is an exponential ansatz for any $c \geq c_-$. This occurs when

$$c_{het}^2 - 4d_+ \sqrt{\frac{\alpha}{d_-}} c_{het} + 4d_+ \alpha \geq 0,$$

which is equivalent to

$$c_{het} \geq c_+ \left(\sqrt{\frac{d_+}{d_-}} + \sqrt{\frac{d_+}{d_-} - 1} \right) = c_{int}.$$

By analogy with the homogeneous case, it is reasonable to expect that under the previous condition the solution spreads with speed c_- which is precisely what is stated in Theorem 2.3.

It remains to consider the case when $g(c_-) < 0$. Then we claim that

$$g(c_+) > 0.$$

To check this, first compute

$$g(c_+) = c_{het}^2 - 4d_+ \left[\alpha + \frac{c_+ - \sqrt{c_+^2 - 4d_- \alpha}}{2d_-} (c_{het} - c_+) \right]. \quad (7)$$

Notice that $g(c_+) = 0$ when $c_{het} = c_+$. Hence it is enough to show that the derivative with respect to c_{het} is positive, which is indeed the case as

$$\begin{aligned} 2c_{het} - 2\frac{d_+}{d_-} \left[c_+ - \sqrt{c_+^2 - 4d_- \alpha} \right] &\geq 2c_+ - 2\frac{d_+}{d_-} \left[c_+ - \sqrt{c_+^2 - 4d_- \alpha} \right] \\ &= 4\sqrt{d_+ \alpha} \left[1 - \frac{d_+}{d_-} + \frac{d_+}{d_-} \sqrt{1 - \frac{d_-}{d_+}} \right] \\ &= 4\sqrt{d_+ \alpha} \frac{d_+}{d_-} \left[-\left(1 - \frac{d_-}{d_+}\right) + \sqrt{1 - \frac{d_-}{d_+}} \right] \\ &> 0. \end{aligned}$$

The claim that $g(c_+) > 0$ is proved and it follows that $g^{-1}(0) \in (c_-, c_+)$ (here g is understood as a function on the interval $[c_-, c_+]$ where it is invertible), and that an exponential ansatz exists if and only if $c \geq g^{-1}(0)$. Furthermore, upon denoting $\lambda_* := \alpha - \frac{c_{het}^2}{4d_+} < 0$ (as $c_+ < c_{het}$) and letting

$$c = c_{het} - \frac{1}{\zeta},$$

one may check that $g(c) = 0$ is equivalent to

$$2d_- \lambda_* \zeta^2 + c_{het} \zeta - 1 = \sqrt{(c_{het} \zeta - 1)^2 - 4d_- \alpha \zeta^2}.$$

The part inside the square root is positive, so that one eventually reaches

$$d_- \lambda_* \zeta^2 + c_{het} \zeta - 1 + \frac{\alpha}{\lambda_*} = 0.$$

Since ζ must be positive in order for c to belong to the interval (c_-, c_+) , we conclude that

$$g^{-1}(0) = c_{het} - \frac{2d_- \lambda_*}{-c_{het} + \sqrt{c_{het}^2 - 4d_- (\alpha - \lambda_*)}},$$

and we recover the formula for the spreading speed in Theorem 2.3 by using the expression for λ_* .

Building upon the argument above, we can further motivate the emergence of the accelerated front with speed c_u^* as a matching condition in the following sense. We attempt to build a solution of (1) which resembles a traveling front of the Fisher-KPP equation $0 = d_- U'' + cU' + \alpha U(1 - U)$ on the left concatenated on the right with a solution of the linearized equation near zero in a frame moving with the heterogeneity. If $c > c_-$ then there exists a family of traveling fronts traveling with speed c and whose profile resembles $e^{-\lambda(c)(x-ct)}$ as $x - ct \rightarrow +\infty$. In a frame of reference moving with speed c_{het} this front can be viewed as forming an effective boundary condition for the PDE (1) linearized near zero. Rescaling by $u(t, x) := e^{\eta(c)t} v(t, x)$ this PDE satisfies the boundary value problem

$$v_t = \chi(x)v_{xx} + c_{het}v_x + (\alpha - \eta(c))v, \quad v_x(-L) = -\lambda(c)v(-L), \quad v_x(L) = -\gamma(c)v(L) \quad (8)$$

for some L sufficiently large, $\eta(c) = -\lambda(c)(c_{het} - c)$ and $\gamma(c)$ is the largest positive root of $d_+ \gamma^2 - c_{het} \gamma + \alpha - \eta(c) = 0$. For generic separated boundary conditions, it is known that the point spectrum of the linear operator appearing on the right hand side of (8) will accumulate on the absolute spectrum in the limit as $L \rightarrow \infty$; see [26, 20]. The absolute spectrum in this case is

$$\Sigma_{abs} = \left\{ \lambda \leq \alpha - \eta(c) - \frac{c_{het}^2}{4d_+} \right\}.$$

Thus, if c is selected so that $\eta(c) < \alpha - \frac{c_{het}^2}{4d_+}$ then one would expect that the constructed solution would be pointwise unstable due to unstable point spectrum of (8). Conversely, if $\eta(c) > \alpha - \frac{c_{het}^2}{4d_+}$ then solutions of (8) will decay pointwise and matching with the front will fail. Therefore, we select c_u^* so that

$$\eta(c_u^*) = \alpha - \frac{c_{het}^2}{4d_+} = \lambda_*,$$

which upon inspection is equivalent to the condition that $g(c_u^*) = 0$.

Outline of the paper. In Section 3, we provide general bounds on the spreading speed, and prove that if it exists it should be bounded respectively from above and below by c_+ and c_- . Sections 4 and 5 are dedicated to the proofs of our main theorems on spreading speeds in cases (I) and (II) respectively. For each case, we construct sub and/or super-solutions to bound adequately the spreading speed. Finally, in the last Section 6, we prove the existence and uniqueness of traveling front solutions with strong exponential decay at $+\infty$ in case (I) and for each $c_{het} \in (c_-, c_+)$.

3. General bounds on the spreading speed. In this section, we aim at confirming rigorously the natural intuition that the spreading speed should always be bounded from respectively above and below by $c_+ = 2\sqrt{d_+\alpha}$ and $c_- = 2\sqrt{d_-\alpha}$. We will do this by constructing some super and sub-solutions which are valid in all the cases considered in our main theorems. Actually, our bounds on the spreading speed remain true for more general form of diffusivity and therefore our results here can be of independent interest.

Throughout this section, we consider

$$\partial_t u = \chi(t, x) \partial_x^2 u + \alpha u(1 - u), \quad (9)$$

with $\alpha > 0$ and some smooth function χ that only satisfies

$$d_- \leq \chi(t, x) \leq d_+ \quad \text{for all } t \geq 0 \text{ and } x \in \mathbb{R},$$

where

$$0 < d_- < d_+.$$

We will prove the following result.

Proposition 1. *Consider a compactly supported and nontrivial initial datum u_0 such that $0 \leq u_0 \leq 1$. Then the solution u of (9) satisfies that*

$$\begin{aligned} \forall 0 < c < c_-, \quad \limsup_{t \rightarrow +\infty} \sup_{0 \leq x \leq ct} |1 - u(t, x)| &= 0, \\ \forall c > c_+, \quad \limsup_{t \rightarrow +\infty} \sup_{x \geq ct} u(t, x) &= 0. \end{aligned}$$

In particular, the rightward spreading speed c_u^* (if it exists) must satisfy $c_- \leq c_u^* \leq c_+$. We refer to [6, 5] for other general results on monostable equations with spatio-temporal heterogeneities.

3.1. A general super-solution. We start the proof of Proposition 1 with the construction of a super-solution.

Lemma 3.1. *For any $C > 0$, the function $w(t, x) = \min \left\{ 1, C e^{-\frac{c_+}{2d_+}(x-c_+t)} \right\}$ is a super-solution of (9).*

Proof. The proof is a simple calculation. Let

$$N(v) := \partial_t v - \chi(t, x) \partial_x^2 v - \alpha v(1 - v).$$

We compute $N(1) = 0$ and, for any $C > 0$,

$$\begin{aligned} N(C e^{-\lambda(x-ct)}) &= C \left(\lambda c - d_+ \lambda^2 - \alpha \right) e^{-\lambda(x-ct)} + C \lambda^2 (d_+ - \chi(t, x)) e^{-\lambda(x-ct)} + \alpha C^2 e^{-2\lambda(x-ct)} \\ &> C \left(\lambda c - d_+ \lambda^2 - \alpha \right) e^{-\lambda(x-ct)}, \end{aligned}$$

as $d_+ - \chi(t, x) \geq 0$ for all $x \in \mathbb{R}$ and $t \geq 0$. Then letting $\lambda = \frac{c_+}{2d_+}$ and $c = c_+$ we get that $w(t, x)$ is a super-solution. $\square$

Now, for any $0 \leq u_0 \leq 1$ a compactly supported and nontrivial initial datum, one can find $C > 0$ large enough so that $u_0 \leq \min \{1, C e^{-\frac{c_+}{2d_+}x}\}$. A consequence of the above lemma and the comparison principle is that we have

$$\forall c > c_+, \quad \limsup_{t \rightarrow +\infty} \sup_{x \geq ct} u(t, x) = 0,$$

for $u(t, x)$ solution of (9) from a compactly supported, nontrivial initial datum $0 \leq u_0 \leq 1$. This proves the second assertion of Proposition 1.




FIGURE 4. Illustration of the building block of the general sub-solution (10) (before its scaling by ϵ) which is composed of two parts $\rho\Psi_+$ (pink curve) and Ψ_- (blue curve) in the moving frame $z = x - ct$. It is of class $\mathcal{C}^2$ and compactly supported on $\left[-\frac{\pi}{2\omega} - z_+^*, \frac{\pi}{2\beta} - z_-^*\right]$.

3.2. A general sub-solution. We now construct a sub-solution and conclude the proof of Proposition 1. For each $c \in (0, c_-)$, there exist $\delta > 0$ small enough and $\eta > 0$ such that

$$0 < c < 2\sqrt{d_-(\alpha - \delta)} < c_- < 2\sqrt{d_+(\alpha - \delta)} < c_+,$$

and

$$(\alpha - \delta)u < \alpha u(1 - u), \quad \forall u \in (0, \eta).$$

We first set

$$\lambda := \frac{c}{2d_-}, \quad \text{and} \quad \beta := \frac{1}{2d_-}\sqrt{4d_-(\alpha - \delta) - c^2} > 0,$$

together with

$$\gamma := \frac{c}{2d_+}, \quad \text{and} \quad \omega := \frac{1}{2d_+}\sqrt{4d_+(\alpha - \delta) - c^2} > 0.$$

Then, the functions $\Psi_-(z) := e^{-\lambda z} \cos(\beta z)$ and $\Psi_+(z) := e^{-\gamma z} \cos(\omega z)$ are respectively solutions of

$$d_- \Psi'' + c \Psi' + (\alpha - \delta) \Psi = 0,$$

and

$$d_+ \Psi'' + c \Psi' + (\alpha - \delta) \Psi = 0.$$

We consider the intervals $\Omega_- = \left(-\frac{\pi}{2\beta}, \frac{\pi}{2\beta}\right)$ and $\Omega_+ = \left(-\frac{\pi}{2\omega}, \frac{\pi}{2\omega}\right)$, and we note that $\Psi_{\pm} > 0$ on $\Omega_{\pm}$. We denote $z_{\pm}^* \in \Omega_{\pm}$ the unique values where $\Psi_{\pm}''(z_{\pm}^*) = 0$, which are given by

$$z_-^* = -\frac{1}{\beta} \arctan\left(\frac{\lambda^2 - \beta^2}{2\lambda\beta}\right), \quad \text{and} \quad z_+^* = -\frac{1}{\omega} \arctan\left(\frac{\gamma^2 - \omega^2}{2\gamma\omega}\right).$$

Finally, we introduce the function

$$\underline{u}_{c,\delta,\epsilon}(t,x) = \begin{cases} 0, & x - ct \leq -\frac{\pi}{2\omega} - z_+^*, \\ \epsilon\rho\Psi_+(x - ct + z_+^*), & -\frac{\pi}{2\omega} - z_+^* < x - ct \leq 0, \\ \epsilon\Psi_-(x - ct + z_-^*), & 0 < x - ct < \frac{\pi}{2\beta} - z_-^*, \\ 0, & x - ct \geq \frac{\pi}{2\beta} - z_-^*. \end{cases} \quad (10)$$

Here, $\rho > 0$ is chosen so as to ensure continuity at $x - ct = 0$, that is

$$\rho := \frac{\Psi_-(z_-^*)}{\Psi_+(z_+^*)}.$$

We fix $\epsilon > 0$ small enough such that

$$0 \leq \underline{u}_{c,\delta,\epsilon}(t,x) < \eta, \quad \forall t \geq 0, x \in \mathbb{R}.$$

We now check that $\underline{u}_{c,\delta,\epsilon}(t,x)$ is a generalized sub-solution.

- For $z = x - ct \in (-\frac{\pi}{2\omega} - z_+^*, 0]$, one has $0 < \epsilon\rho\Psi_+(z + z_+^*) < \eta$, and so

$$N(\epsilon\rho\Psi_+(z + z_+^*)) < (d_+ - \chi(t,x))\epsilon\rho\Psi_+''(z + z_+^*) \leq 0,$$

as $\Psi_+''(z + z_+^*) \leq 0$ on $(-\frac{\pi}{2\omega} - z_+^*, 0]$ and $\chi(t,x) \leq d_+$.

- For $z = x - ct \in (0, \frac{\pi}{2\beta} - z_-^*)$, one has $0 < \epsilon\Psi_-(z + z_-^*) < \eta$, and so

$$N(\epsilon\Psi_-(z + z_-^*)) < (d_- - \chi(t,x))\epsilon\rho\Psi_-''(z + z_-^*) \leq 0,$$

as $\Psi_-''(z + z_-^*) \geq 0$ on $(0, \frac{\pi}{2\beta} - z_-^*)$ and $d_- \leq \chi(t,x)$.

- At $z = x - ct = 0$, we already have continuity since $\rho\Psi_+(z_+^*) = \Psi_-(z_-^*)$, and we also have that $\Psi_+''(z_+^*) = \Psi_-''(z_-^*) = 0$ by definition of $z_\pm^*$. Now using the equations satisfied by $\Psi_\pm$ evaluated at $z_\pm^*$, we also obtain

$$c(\Psi_-'(z_-^*) - \rho\Psi_+'(z_+^*)) = -(\alpha - \delta)(\Psi_-(z_-^*) - \rho\Psi_+(z_+^*)) - d_- \Psi_-''(z_-^*) + d_+ \rho\Psi_+''(z_+^*) = 0.$$

As a consequence, we have $\Psi_-'(z_-^*) = \rho\Psi_+'(z_+^*)$ and the sub-solution is of class

$$\mathcal{C}^2 \text{ for } x - ct \in \left(-\frac{\pi}{2\omega} - z_+^*, \frac{\pi}{2\beta} - z_-^*\right).$$

We have now reached the next result.

Lemma 3.2. *Let $c \in (0, c_-)$. Then there exists $\delta_0(c) > 0$ such that for each $0 < \delta < \delta_0(c)$ one can find $\epsilon_0(c, \delta) > 0$ such that for all $0 < \epsilon < \epsilon_0(c, \delta)$ the function*

$$\underline{u}_{c,\delta,\epsilon}(t,x) = \begin{cases} 0, & x - ct \leq -\frac{\pi}{2\omega} - z_+^*, \\ \epsilon\rho\Psi_+(x - ct + z_+^*), & -\frac{\pi}{2\omega} - z_+^* < x - ct \leq 0, \\ \epsilon\Psi_-(x - ct + z_-^*), & 0 < x - ct < \frac{\pi}{2\beta} - z_-^*, \\ 0, & x - ct \geq \frac{\pi}{2\beta} - z_-^*, \end{cases}$$

is a generalized sub-solution where $\Psi_\pm$, $z_\pm^$ and ρ are defined as above. We refer to Figure 4 for an illustration.*

With the above lemma, we can now conclude the proof of Proposition 1. Let u be the solution of the Cauchy problem starting from a compactly supported initial condition $0 \leq u_0 \leq 1$, and fix $0 < c < c' < c_-$. By the strong maximum principle, it satisfies $u(1, \cdot) > 0$ and, upon diminishing further ϵ in the previous lemma, one can ensure that

$$\underline{u}_{c',\delta,\epsilon}(1, \cdot) \leq u(1, \cdot).$$

By the comparison principle, we infer that

$$\inf_{t \geq 1} u(t, c't) \geq \nu > 0,$$

for some $\nu \in (0, 1)$ which depends on c' . By a straightforward symmetry argument, one may check that $\underline{u}_{c', \delta, \epsilon}(t, -x)$ is also a sub-solution of (9) and therefore we also get that

$$\inf_{t \geq 1} u(t, -c't) \geq \nu > 0.$$

Next, up to reducing ν we can also assume that

$$u(1, x) \geq \nu,$$

for all $x \in [-c', c']$. Therefore, applying a comparison principle on $\{(t, x) \mid t \geq 1 \text{ and } x \in [-c't, c't]\}$, with the constant ν as a sub-solution, we infer that actually

$$\liminf_{t \rightarrow +\infty} \inf_{-c't \leq x \leq c't} u(t, x) \geq \nu.$$

Now we prove the first assertion of Proposition 1. We proceed by contradiction and, since $u \leq 1$, we assume that there exist sequences $t_n \rightarrow +\infty$ and $x_n \in [0, ct_n]$ such that

$$\limsup_{n \rightarrow +\infty} u(t_n, x_n) < 1.$$

By looking at the reaction as a L^∞ -bounded source term $(t, x) \mapsto f(u(t, x))$, one can apply standard parabolic estimates to find that, for any compact set $D \subset \mathbb{R}^2$ and p arbitrarily large, the sequence $u(t_n + \cdot, x_n + \cdot)$ is uniformly bounded in the parabolic Sobolev spaces $W_p^{1,2}(D)$, hence in some Holder space $C^{\alpha, 1+\alpha}(D)$ with $\alpha \in (0, 1)$. One can then extract a subsequence such that $u(t_n + t, x_n + x)$ converges (locally uniformly and weakly in $W_{p,loc}^{1,2}(\mathbb{R}^2)$ for any p) as $n \rightarrow +\infty$ to an entire in time solution u_∞ of

$$\partial_t u_\infty = \tilde{\chi}(t, x) \partial_x^2 u_\infty + \alpha u_\infty (1 - u_\infty),$$

for some $\tilde{\chi}$ which also satisfies $d_- \leq \tilde{\chi} \leq d_+$. Moreover, it follows from the above where $c' > c$ that $u_\infty(t, x) \geq \nu > 0$ for all $(t, x) \in \mathbb{R}^2$. By a straightforward comparison with the ODE and regardless of the actual function $\tilde{\chi}$, the function u_∞ must be identical to 1, which is a contradiction with the fact that $u_\infty(0, 0) < 1$ by construction. This proves the last statement of Proposition 1.

4. Construction of super and sub-solutions for case (I) and proof of Theorem 2.1. In this section, we construct super and sub-solutions in case where χ satisfies assumption (I). We proceed step by step and consider each case depicted in Theorem 2.1.

4.1. Subcase $c_{het} < c_-$. When $c_{het} < c_-$, we need to prove that the rightward spreading speed is $c_u^* = c_-$. From Proposition 1, we already have proved that

$$\forall 0 < c < c_-, \quad \limsup_{t \rightarrow +\infty} \sup_{0 \leq x \leq ct} |1 - u(t, x)| = 0,$$

and thus it remains to provide a super-solution in this case.

We let $c > c_-$. Then, one can find $\epsilon > 0$ small enough such that we have $c_- = 2\sqrt{d_- \alpha} < 2\sqrt{(d_- + \epsilon)\alpha} < \min(c, c_+)$. Next we fix $\tau_\epsilon > 0$ to be large enough such that for any $\tau \geq \tau_\epsilon$, we have

$$d_- \leq \chi(\tau) \leq d_- + \epsilon.$$

Lemma 4.1. *Let $c_\epsilon := 2\sqrt{(d_- + \epsilon)\alpha}$. For all $\tau \geq \tau_\epsilon$, the following function*

$$u_\tau(t, x) = \min \left\{ 1, e^{-\frac{c_\epsilon}{2\chi(\tau)}(x - c_\epsilon t - \tau)} \right\}, \quad t \geq 0, \quad x \in \mathbb{R},$$

is a super-solution for (1).

Proof. We recall the functional N defined as

$$N(u) = \partial_t u - \chi(x - c_{het}t) \partial_x^2 u - \alpha u(1 - u).$$

We readily have that $N(1) = 0$ and that

$$N\left(e^{-\lambda(x-c_\epsilon t-\tau)}\right) \geq (\lambda c_\epsilon - \chi(x - c_{het}t) \lambda^2 - \alpha) e^{-\lambda(x-c_\epsilon t-\tau)},$$

for all $\lambda > 0$. Next, we note that for each $x \geq c_\epsilon t + \tau$, we have

$$x - c_{het}t \geq \underbrace{(c_\epsilon - c_{het})}_{>0} t + \tau \geq \tau.$$

As a consequence, for all $x \geq c_\epsilon t + \tau$, using the monotonicity of χ , we get

$$N\left(e^{-\lambda(x-c_\epsilon t-\tau)}\right) \geq (\lambda c_\epsilon - \chi(\tau) \lambda^2 - \alpha) e^{-\lambda(x-c_\epsilon t-\tau)}.$$

Now evaluating at $\lambda = \frac{c_\epsilon}{2\chi(\tau)}$, we obtain

$$N\left(e^{-\frac{c_\epsilon}{2\chi(\tau)}(x-c_\epsilon t-\tau)}\right) \geq \frac{c_\epsilon^2 - 4\chi(\tau)\alpha}{4\chi(\tau)} e^{-\frac{c_\epsilon}{2\chi(\tau)}(x-c_\epsilon t-\tau)}, \quad x \geq c_\epsilon t + \tau.$$

Finally, as $\tau \geq \tau_\epsilon$, we get

$$c_\epsilon^2 - 4\chi(\tau)\alpha \geq c_\epsilon^2 - 4(d_- + \epsilon)\alpha = 0,$$

which concludes the proof. $\square$

As a consequence of the above lemma and the comparison principle, we have that

$$\forall c > c_- > c_{het}, \quad \limsup_{t \rightarrow +\infty} \sup_{x \geq ct} u(t, x) = 0,$$

for the solution $u(t, x)$ of (1) from a compactly supported, nontrivial initial datum $0 \leq u_0 \leq 1$. And thus, the rightward spreading speed of (1) is less than or equal to c_- when $c_{het} < c_-$ which concludes the proof of Theorem 2.1 in this case.

4.2. Subcase $c_+ < c_{het}$. When $c_+ < c_{het}$, we need to prove that the rightward spreading speed is $c_u^* = c_+$. From Proposition 1, we already have that the spreading speed is less than or equal to c_+ , in the sense that

$$\forall c > c_+, \quad \limsup_{t \rightarrow +\infty} \sup_{x \geq ct} u(t, x) = 0,$$

and thus it remains to provide a sub-solution in this case.

As a matter of fact this sub-solution is valid in all subcases of (I), and hereafter we simply let $0 < c < \min(c_+, c_{het})$. Then, one can find $\epsilon > 0$ such that $c < 2\sqrt{d_+(\alpha - 2\epsilon)} < c_+$ together with $\eta_\epsilon > 0$ such that

$$(\alpha - \epsilon)u \leq \alpha u(1 - u), \quad u \in [0, \eta_\epsilon].$$

We introduce two positive real numbers

$$\beta_+(\epsilon, c) := \frac{\sqrt{4d_+(\alpha - \epsilon) - c^2}}{2d_+} > 0, \text{ and } \beta_+(c) := \beta_+(0, c) = \frac{\sqrt{4d_+\alpha - c^2}}{2d_+} > 0,$$

together with the following family of functions

$$u_{\tau, \epsilon}^c(t, x) := \begin{cases} \delta_\epsilon \left[e^{-\frac{c}{2d_+}(x-ct+\tau)} \cos(\beta_+(\epsilon, c)(x - ct + \tau)) + \epsilon \right], & x - ct + \tau \in \Omega_\epsilon(c), \\ 0, & \text{otherwise,} \end{cases}$$

with $\Omega_\epsilon(c) = \left[-\frac{\pi}{2\beta_+(\epsilon, c)} - z_\epsilon^-(c), \frac{\pi}{2\beta_+(\epsilon, c)} + z_\epsilon^+(c)\right]$. Here $z_\epsilon^\pm(c)$ are defined through

$$z_\epsilon^\pm(c) \in \left(0, \frac{\pi}{2\beta_+(\epsilon, c)}\right),$$

and

$$e^{\mp \frac{c}{2d_+} \left(\frac{\pi}{2\beta_+(\epsilon, c)} + z_\epsilon^\pm(c)\right)} \sin(\beta_+(\epsilon, c) z_\epsilon^\pm(c)) = \epsilon,$$

so that the functions $u_{\tau, \epsilon}^c$ are nonnegative and continuous. Then we have the asymptotics

$$z_\epsilon^\pm(c) = \frac{e^{\mp \frac{c\pi}{4d_+\beta_+(c)}}}{\beta_+(c)} \epsilon + o(\epsilon), \quad \text{as } \epsilon \rightarrow 0.$$

Furthermore, $\delta_\epsilon > 0$ is fixed such that $0 \leq u_{\tau, \epsilon}^c(t, x) \leq \eta_\epsilon$ for all $x - ct + \tau \in \Omega_\epsilon(c)$ and can be chosen independent of $c < 2\sqrt{d_+(\alpha - 2\epsilon)}$.

We first remark that when $x - ct + \tau \in \Omega_\epsilon(c)$, we have

$$x - c_{het}t \in \left[-\frac{\pi}{2\beta_+(\epsilon, c)} - z_\epsilon^-(c) + (c - c_{het})t - \tau, \frac{\pi}{2\beta_+(\epsilon, c)} + z_\epsilon^+(c) + (c - c_{het})t - \tau\right].$$

Next, as $\chi(-\infty) = d_+$, there exists $A > 0$ such that for all $\xi \leq -A$, we get

$$|\chi(\xi) - d_+| \leq \epsilon^2.$$

As a consequence, for all

$$\tau > \tau_\epsilon(c) := A + \frac{\pi}{2\beta_+(\epsilon, c)} + z_\epsilon^+(c),$$

we have $x - c_{het}t \leq -A$ and

$$\left| (d_+ - \chi(x - c_{het}t)) \partial_x^2 u_{\tau, \epsilon}^c(t, x) \right| \leq \epsilon^2 \delta_\epsilon \left(\frac{c}{2d_+} + \beta_+(\epsilon, c) \right)^2 e^{\frac{c}{2d_+} \left(\frac{\pi}{2\beta_+(\epsilon, c)} + z_\epsilon^-(c) \right)} := \epsilon^2 \delta_\epsilon K_\epsilon(c),$$

for all $x - ct + \tau \in \Omega_\epsilon(c)$. This implies that for all $\tau > \tau_\epsilon(c)$

$$\begin{aligned} N(u_{\tau, \epsilon}^c(t, x)) &\leq (d_+ - \chi(x - c_{het}t)) \partial_x^2 u_{\tau, \epsilon}^c(t, x) - \delta_\epsilon \epsilon (\alpha - \epsilon) \\ &\leq \epsilon \delta_\epsilon [(K_\epsilon(c) + 1)\epsilon - \alpha], \quad x - ct + \tau \in \Omega_\epsilon(c). \end{aligned}$$

Notice that

$$\liminf_{\epsilon \rightarrow 0} K_\epsilon(c) > 0.$$

As a consequence, we obtain the following lemma.

Lemma 4.2. *Let $0 < c < \min(c_+, c_{het})$. There is $\epsilon_0(c) > 0$ such that for all $\epsilon \in (0, \epsilon_0(c))$, the function $u_{\tau, \epsilon}^c$ is a sub-solution for all $\tau > \tau_\epsilon(c)$.*

Now, using similar arguments as in the proof of Proposition 1 we deduce that the solutions spread at least with speed $\min(c_+, c_{het})$, which concludes the proof of Theorem 2.1 in the subcase when $c_+ < c_{het}$.

4.3. Subcase $c_- \leq c_{het} \leq c_+$. When $c_- \leq c_{het} \leq c_+$, our main Theorem 2.1 asserts that the selected rightward spreading speed is precisely the speed of the heterogeneity, that is $c_u^* = c_{het}$. In that case, we cannot rely on Proposition 1 to obtain either inequality, and we need to refine our analysis. Nevertheless, we can use Lemma 4.2 to obtain in this subcase that the spreading speed is larger than or equal to $c_{het} = \min(c_+, c_{het})$, and it only remains to construct a super-solution which spreads at speed c_{het} . This is precisely the result of the following lemma.

Lemma 4.3. *Let $c_- \leq c_{het} \leq c_+$. Then, for any $\epsilon > 0$ there is $\tau_0 > 0$ such that for all $\tau \geq \tau_0$ the function $\bar{u}_\tau(t, x) = \min \left\{ 1, e^{-\frac{c_{het}}{2\chi(\tau)}(x - (c_{het} + \epsilon)t - \tau)} \right\}$ is a super-solution of (1).*

Proof. Let ϵ be any positive number. As $c_- \leq c_{het}$ and $\chi(x) \rightarrow d_-$ as $x \rightarrow +\infty$, there exists $\tau_0 > 0$ such that for all $\tau \geq \tau_0$ one has

$$c_{het}^2 + 2c_{het}\epsilon - 4\alpha\chi(\tau) \geq 0.$$

Next, we have for $\tau \leq x - c_{het}t$ that $\chi(x - c_{het}t) \leq \chi(\tau)$ and

$$N(\bar{u}_\tau(t, x)) \geq \frac{c_{het}^2 + 2c_{het}\epsilon - 4\alpha\chi(\tau)}{4\chi(\tau)} e^{-\frac{c_{het}}{2\chi(\tau)}(x - (c_{het} + \epsilon)t - \tau)} \geq 0.$$

This already concludes the proof. $\square$

5. Construction of super and sub-solutions in case (II) and proof of Theorem 2.3. In this section, we construct super and sub-solutions in the case where χ satisfies assumption (II). We first treat the most difficult case when $c_{het} \in (c_+, c_{int})$ and then explain how to deal with the remaining two cases.

5.1. Case $c_{het} \in [c_+, c_{int})$. Recall from the discussion in Section 2.2 that

$$g(c) = c_{het}^2 - 4d_+ \left[\alpha + \frac{c - \sqrt{c^2 - 4d_- \alpha}}{2d_-} (c_{het} - c) \right],$$

and since $c_{het} \in [c_+, c_{int})$, there exists a unique $c \in (c_-, c_+]$ such that $g(c) = 0$. For convenience, throughout Section 5.1 we will denote it by c_u^* . Our goal is indeed to prove that this c_u^* , whose explicit formula is given in Theorem 2.3, is the spreading speed of the solution in that case.

5.1.1. Super-solution. According to Proposition 1, we already know that the spreading speed is less than or equal to c_+ . Thus to construct a super-solution here we only need to consider the case when $c_{het} \in (c_+, c_{int})$. Then recall that $c_u^* \in (c_-, c_+)$ is such that $g(c_u^*) = 0$, and let any $c \in (c_u^*, c_{het})$. We introduce the family of (continuous) functions

$$u_\tau(t, x) = C \times \begin{cases} 1, & x \leq ct - \tau, \\ e^{-\lambda(x - ct + \tau)}, & ct - \tau < x < c_{het}t - \tau, \\ e^{-\lambda(c_{het} - c)t} e^{-\mu(x - c_{het}t + \tau)}, & x \geq c_{het}t - \tau, \end{cases}$$

where λ , μ and τ are positive and will be adjusted as follows, and $C \geq 1$ is also positive but arbitrary.

- For $x \in (ct - \tau, c_{het}t - \tau)$, we compute

$$\begin{aligned} N(Ce^{-\lambda(x - ct + \tau)}) &= C(\lambda c - \lambda^2 \chi(x - c_{het}t) - \alpha) e^{-\lambda(x - ct + \tau)} + C^2 \alpha e^{-2\lambda(x - ct + \tau)} \\ &\geq C(\lambda c_u^* - \lambda^2 d_- - \alpha) e^{-\lambda(x - ct + \tau)} \\ &\quad + C[\lambda^2(d_- - \chi(-\tau)) + \lambda(c - c_u^*)] e^{-\lambda(x - ct + \tau)}, \end{aligned}$$

where we used the fact that χ is nondecreasing and $x - c_{het}t \leq -\tau$. We now select

$$\lambda = \lambda(c_u^*),$$

where $\lambda(c)$ denotes the smaller positive (thanks to $c > c_u^* > c_-$) solution of

$$\lambda c - \lambda^2 d_- - \alpha = 0, \tag{11}$$

that is

$$\lambda(c) := \frac{c - \sqrt{c^2 - 4d_- \alpha}}{2d_-}.$$

There exists $\tau_0 > 0$ such that for all $\tau \geq \tau_0$ we have

$$\lambda(c_u^*)^2(d_- - \chi(-\tau)) + \lambda(c_u^*)(c - c_u^*) > 0,$$

hence $N(u_\tau(t, x)) > 0$ for $x \in (ct - \tau, c_{het}t - \tau)$.

- For $x > c_{het}t - \tau$, we have that

$$\begin{aligned} N(Ce^{-\lambda(c_{het}-c)t}e^{-\mu(x-c_{het}t+\tau)}) &= C \left(-\lambda(c_{het} - c_u^*) + \mu c_{het} - \alpha - d_+ \mu^2 \right) e^{-\lambda(c_{het}-c)t} e^{-\mu(x-c_{het}t+\tau)} \\ &\quad + C \left[\mu^2(d_+ - \chi(x - c_{het}t)) + \lambda(c - c_u^*) \right] e^{-\lambda(c_{het}-c)t} e^{-\mu(x-c_{het}t+\tau)} \\ &\quad + C^2 \alpha e^{-2\lambda(c_{het}-c)t} e^{-2\mu(x-c_{het}t+\tau)}. \end{aligned}$$

The last two terms are nonnegative as $\chi \leq d_+$ and we finally select $\mu > 0$ such that

$$-\lambda(c_{het} - c_u^*) + \mu c_{het} - \alpha - d_+ \mu^2 = 0, \text{ with } \lambda = \lambda(c_u^*).$$

That is, we let

$$\mu := \frac{c_{het} - \sqrt{g(c_u^*)}}{2d_+} = \frac{c_{het}}{2d_+},$$

as $g(c_u^*) = 0$, and we get that $N(u_\tau(t, x)) \geq 0$ for $x > c_{het}t - \tau$. We note that the above formula with $\lambda = \lambda(c_u^*)$ and $\mu = \frac{c_{het}}{2d_+}$ writes

$$-\lambda(c_u^*)(c_{het} - c_u^*) = \alpha - \frac{c_{het}^2}{4d_+} = \lambda_\star.$$

- To be consistent with the definition of the generalized super-solution, it remains to prove that there is a negative jump in the derivative at $x = ct - \tau$ and $x = c_{het}t - \tau$. This is trivial at the former point, and concerning the latter we will show that

$$0 > \partial_x u_\tau(t, (c_{het}t - \tau)^-) > \partial_x u_\tau(t, (c_{het}t - \tau)^+).$$

This is equivalent to show that $\lambda = \lambda(c_u^*) < \mu = \frac{c_{het}}{2d_+}$, where c_u^* is such that $g(c_u^*) = 0$. We claim that we have

$$\lambda(c_u^*) = \frac{c_{het}}{2d_-} - \frac{1}{2d_-} \sqrt{c_{het}^2 - 4d_- \alpha + 4d_- \lambda_\star}, \quad (12)$$

where $\lambda_\star = \alpha - \frac{c_{het}^2}{4d_+}$. Let us assume that the claim (12) holds. Then, we obtain

$$\lambda = \lambda(c_u^*) = \frac{c_{het}}{2d_-} \left(1 - \sqrt{1 - \frac{d_-}{d_+}} \right) < \frac{c_{het}}{2d_-} \left(1 - \left(1 - \frac{d_-}{d_+} \right) \right) = \frac{c_{het}}{2d_+} = \mu,$$

since $0 < 1 - \frac{d_-}{d_+} < 1$ and we have verified the fact that there is a negative jump in the derivative at $x = c_{het}t - \tau$. Coming back to the formula (12), we first note that by definition of $\lambda(c_u^*)$ it solves

$$d_- \lambda(c_u^*)^2 - c_u^* \lambda(c_u^*) + \alpha = 0,$$

and thus it is also solution of

$$d_- \lambda(c_u^*)^2 - c_{het} \lambda(c_u^*) + \alpha = \lambda_\star,$$

since

$$-\lambda(c_u^*)(c_{het} - c_u^*) = \lambda_\star.$$




FIGURE 5. Sketch of the super-solution $u_\tau(t, x)$ given in Lemma 5.1 with $C = 1$ which is composed of three parts: it is constant and equal to 1 for $x \leq ct - \tau$ (gray curve), and then it is the concatenation of two exponentials (blue and pink curves) for $x \geq ct - \tau$ which are glued at $x = c_{het}t - \tau$. Note that the factor $\rho(t)$ is to ensure continuity between the two exponentials.

Finally, the fact that $\lambda(c_u^*)$ is given by (12) (and not the other positive root) can be deduced by recalling that it was defined as the smaller positive root of (11) and comparing the formulas in the limiting case $c_{het} = c_+$.

Lemma 5.1. *Let $c_+ < c_{het} < c_{int}$. Let also c_u^* be such that $g(c_u^*) = 0$ and $\lambda(c) := \frac{c - \sqrt{c^2 - 4d_- - \alpha}}{2d_-}$. Then for any $c > c_u^*$, there exists $\tau_0 > 0$ such that, for each $\tau \geq \tau_0$ and $C \geq 1$,*

$$u_\tau(t, x) = C \times \begin{cases} 1, & x \leq ct - \tau, \\ e^{-\lambda(c_u^*)(x-ct+\tau)}, & ct - \tau < x < c_{het}t - \tau, \\ e^{-\lambda(c_u^*)(c_{het}-c)t} e^{-\frac{c_{het}}{2d_+}(x-c_{het}t+\tau)}, & x \geq c_{het}t - \tau, \end{cases}$$

is a super-solution of (1).

We refer to Figure 5 for an illustration.

Choosing C large enough so that $u_\tau(0, \cdot) \geq u_0$, we find that

$$\forall c > c_u^*, \quad \limsup_{t \rightarrow +\infty} \sup_{x \geq ct} u(t, x) = 0.$$

In other words, the solution of (1) with compactly supported initial datum spreads at speed less than or equal to c_u^* .

5.1.2. Sub-solution. Here we assume that $c_{het} \in [c_+, c_{int})$. We let $c \in (c_-, c_+)$ be such that $c_- < c < c_u^* = g^{-1}(0) \leq c_+$. We also let $\epsilon > 0$ (to be made arbitrarily small) and $\eta > 0$ be such that

$$(\alpha - \epsilon)u < \alpha u(1 - u), \quad \text{for } u \in [0, \eta].$$

We are going to construct a suitable sub-solution of

$$\partial_t u - \chi(x - c_{het}t) \partial_x^2 u - (\alpha - \epsilon)u \leq 0, \quad t \geq 0, \quad x \in \mathbb{R}, \quad (13)$$

moving with speed c . Provided that this sub-solution is smaller than η , then clearly it is also a sub-solution of (1). For convenience, we introduce the linear operator

$$M(u) := \partial_t u - \chi(x - c_{het}t) \partial_x^2 u - (\alpha - \epsilon)u.$$

We give a first sub-solution of (13), which has compact support to the left, and that writes

$$\underline{u}_{1,\tau}(t, x) = \max \left\{ 0, e^{-\lambda(x-ct+\tau)} - e^{-(\lambda+\gamma)(x-ct+\tau)} \right\},$$

where λ , γ and τ are positive constants which we adjust below.

- First, we select λ as a root to the equation

$$\lambda^2 d_- - \lambda c + \alpha - \epsilon = 0, \quad \lambda = \frac{c - \sqrt{c^2 - 4d_-(\alpha - \epsilon)}}{2d_-} > 0.$$

By our choice of λ we will always have

$$M \left(e^{-\lambda(x-ct+\tau)} \right) = \lambda^2 (d_- - \chi(x - c_{het}t)) e^{-\lambda(x-ct+\tau)} \leq 0.$$

- Next we pick γ such that

$$0 < d_- \gamma < c - 2\lambda d_- = \sqrt{c^2 - 4d_-(\alpha - \epsilon)}.$$

Then, on the support of $\underline{u}_{1,\tau}$ we have

$$M(\underline{u}_{1,\tau}(t, x)) = \partial_t \underline{u}_{1,\tau}(t, x) - \chi(x - c_{het}t) \partial_x^2 \underline{u}_{1,\tau}(t, x) - (\alpha - \epsilon) \underline{u}_{1,\tau}(t, x),$$

and by linearity we get

$$M(\underline{u}_{1,\tau}(t, x)) = M(e^{-\lambda(x-ct+\tau)}) - M(e^{-(\lambda+\gamma)(x-ct+\tau)}) \leq -M(e^{-(\lambda+\gamma)(x-ct+\tau)}).$$

Let us also note that

$$M \left(e^{-(\lambda+\gamma)(x-ct+\tau)} \right) = ((\lambda + \gamma)c - (\lambda + \gamma)^2 \chi(x - c_{het}t) - (\alpha - \epsilon)) e^{-(\lambda+\gamma)(x-ct+\tau)},$$

and using the fact that $\lambda > 0$ is such that $\lambda^2 d_- - \lambda c + \alpha - \epsilon = 0$, we can simplify the above expression to

$$M \left(e^{-(\lambda+\gamma)(x-ct+\tau)} \right) = (\gamma[c - 2\lambda d_- - \gamma d_-] + (d_- - \chi(x - c_{het}t))(\lambda + \gamma)^2) e^{-(\lambda+\gamma)(x-ct+\tau)}.$$

From our choice of γ , one can find $x_* \in \mathbb{R}$ such that for $x - c_{het}t \leq x_*$ we have

$$\chi(x - c_{het}t) - d_- \leq \frac{\gamma[c - 2\lambda d_- - \gamma d_-]}{(\lambda + \gamma)^2}.$$

As a consequence, for $x - c_{het}t \leq x_*$, we have $M(e^{-(\lambda+\gamma)(x-ct+\tau)}) \geq 0$ and so

$$M(\underline{u}_{1,\tau}(t, x)) \leq 0$$

on $x - c_{het}t \leq x_*$.

- We now assume that $x - c_{het}t \geq x_*$. From the previous computations we obtain that

$$M(\underline{u}_{1,\tau}(t, x)) \leq (d_- - \chi(x - c_{het}t)) \left(\lambda^2 - (\lambda + \gamma)^2 e^{-\gamma(x-ct+\tau)} \right) e^{-\lambda(x-ct+\tau)}.$$

Recall that $c < c_+ \leq c_{het}$. It follows that there exists $\tau_* > 0$ such that for all $\tau \geq \tau_*$ we have

$$\frac{\lambda^2}{(\lambda + \gamma)^2} > e^{-\gamma(x-c_{het}t+\tau)} > e^{-\gamma(x-ct+\tau)}, \quad x - c_{het}t \geq x_*,$$

hence $M(\underline{u}_{1,\tau}(t, x)) < 0$.

As a conclusion, we have obtained the following lemma.

Lemma 5.2. *Let $c \in (c_-, c_+)$ be such that $c_- < c < c_u^* = g^{-1}(0) \leq c_+$. Then there exists $\tau_* > 0$ such that*

$$\underline{u}_{1,\tau}(t, x) = \max \left\{ 0, e^{-\lambda(x-ct+\tau)} - e^{-(\lambda+\gamma)(x-ct+\tau)} \right\},$$

is a sub-solution of (13) for all $\tau \geq \tau_*$, with

$$\lambda = \lambda_\epsilon(c) := \frac{c - \sqrt{c^2 - 4d_-(\alpha - \epsilon)}}{2d_-} > 0,$$

and $0 < \gamma = \gamma_\epsilon(c) < \frac{1}{d_-} \sqrt{c^2 - 4d_-(\alpha - \epsilon)}$ for some $\epsilon > 0$ small enough.

We now use the above sub-solution to construct another one which will be compactly supported. Up to reducing $\epsilon > 0$ we assume that $\alpha - \frac{c_{het}^2}{4d_-} < \lambda_* - \epsilon < \lambda_* = \alpha - \frac{c_{het}^2}{4d_+}$ and denote $\varphi_{\lambda_*-\epsilon}$ the solution to

$$\mathcal{L}\varphi = (\lambda_* - \epsilon)\varphi, \quad \text{with} \quad \mathcal{L} := \chi(x)\partial_x^2 + c_{het}\partial_x + \alpha,$$

with prescribed asymptotic expansion

$$\varphi_{\lambda_*-\epsilon}(x) = e^{\left(-\frac{c_{het}}{2d_-} + \frac{1}{2d_-} \sqrt{c_{het}^2 - 4d_- \alpha + 4d_- (\lambda_* - \epsilon)}\right)x} \left(1 + \mathcal{O}\left(e^{\nu'x}\right)\right), \text{ as } x \rightarrow -\infty,$$

for some $0 < \nu' < \nu$, and with damped oscillations at $+\infty$ due to $c_{het}^2 < 4d_+(\alpha - \lambda_* + \epsilon)$. The existence of such a solution follows from the robustness of exponential dichotomies following [11] after noting the exponential convergence of $\chi(x)$ to its asymptotic limits; see (5). We cut-off $\varphi_{\lambda_*-\epsilon}$ to the right at the smallest point $x_\epsilon \in \mathbb{R}$ where it vanishes and denote

$$\tilde{\varphi}_{\lambda_*-\epsilon}(x) = \begin{cases} \varphi_{\lambda_*-\epsilon}(x), & x \leq x_\epsilon, \\ 0, & x > x_\epsilon. \end{cases}$$

We define, for any $\tau > -2x_\epsilon$, the following sub-solution

$$\underline{u}_{2,\tau}(t, x) = \begin{cases} \underline{u}_{1,\tau}(t, x), & x - c_{het}t \leq -\tau/2, \\ c_\tau \underline{u}_{1,\tau}(t, c_{het}t - \tau/2) \tilde{\varphi}_{\lambda_*-\epsilon}(x - c_{het}t), & x - c_{het}t > -\tau/2, \end{cases}$$

where $c_\tau = 1/\varphi_{\lambda_*-\epsilon}(-\tau/2) > 0$. With Lemma 5.2 and the definition of $\tilde{\varphi}_{\lambda_*-\epsilon}$, we only need to verify that $\underline{u}_{2,\tau}(t, x)$ is a sub-solution of (13) for $-\tau/2 \leq x - c_{het}t \leq x_\epsilon$, and also that the jump of the spatial derivative at $x - c_{het}t$ has the correct sign.

For $-\tau/2 < x - c_{het}t < x_\epsilon$, we have

$$\begin{aligned} M(\underline{u}_{2,\tau}(t, x)) &= -\lambda_\epsilon(c)(c_{het} - c)c_\tau e^{-\lambda_\epsilon(c)((c_{het}-c)t+\tau/2)} \varphi_{\lambda_*-\epsilon}(x - c_{het}t) \\ &\quad + (\lambda_\epsilon(c) + \gamma_\epsilon(c))(c_{het} - c)c_\tau e^{-(\lambda_\epsilon(c)+\gamma_\epsilon(c))((c_{het}-c)t+\tau/2)} \varphi_{\lambda_*-\epsilon}(x - c_{het}t) \\ &\quad - c_{het}c_\tau \underline{u}_{1,\tau}(t, c_{het}t - \tau)\varphi'_{\lambda_*-\epsilon}(x - c_{het}t) \\ &\quad - \chi(x - c_{het}t)c_\tau \underline{u}_{1,\tau}(t, c_{het}t - \tau)\varphi''_{\lambda_*-\epsilon}(x - c_{het}t) \\ &\quad - (\alpha - \epsilon)c_\tau \underline{u}_{1,\tau}(t, c_{het}t - \tau)\varphi_{\lambda_*-\epsilon}(x - c_{het}t). \end{aligned}$$

As $\mathcal{L}\varphi = (\lambda_* - \epsilon)\varphi$, we have $\mathcal{L}\varphi - \epsilon\varphi = (\lambda_* - 2\epsilon)\varphi$ and

$$\begin{aligned} M(\underline{u}_{2,\tau}(t, x)) &= -\lambda_\epsilon(c)(c_{het} - c)c_\tau e^{-\lambda_\epsilon(c)((c_{het}-c)t+\tau/2)} \varphi_{\lambda_*-\epsilon}(x - c_{het}t) \\ &\quad + (\lambda_\epsilon(c) + \gamma_\epsilon(c))(c_{het} - c)c_\tau e^{-(\lambda_\epsilon(c)+\gamma_\epsilon(c))((c_{het}-c)t+\tau/2)} \varphi_{\lambda_*-\epsilon}(x - c_{het}t) \\ &\quad - (\lambda_* - 2\epsilon)c_\tau \underline{u}_{1,\tau}(t, c_{het}t - \tau)\varphi_{\lambda_*-\epsilon}(x - c_{het}t) := \mathcal{R}(t, x). \end{aligned}$$

Next, we recall that

$$\lambda_* = -\lambda(c_u^*)(c_{het} - c_u^*),$$




FIGURE 6. Sketch of the sub-solution given in Proposition 2 which is the concatenation of the sub-solution $\underline{u}_{1,\tau}(x-ct)$ given in Lemma 5.2 (composed of the difference of two exponentials) and the function $\varphi_{\lambda_\star-\epsilon}$ which solves $\mathcal{L}\varphi = (\lambda_\star - \epsilon)\varphi$ with prescribed asymptotic behavior at $-\infty$. Note that the factor $\rho(t)$ is to ensure continuity at the matching point $x = c_{het}t - \tau/2$.

such that the right-hand side of the previous inequality can be written as

$$\mathcal{R}(t, x) = \mathcal{D}(t, x) c_\tau e^{-\lambda_\epsilon(c)((c_{het}-c)t+\tau/2)} \varphi_{\lambda_\star-\epsilon}(x - c_{het}t),$$

with

$$\mathcal{D}(t, x) = -\lambda_\epsilon(c)(c_{het}-c) + \lambda(c_u^*)(c_{het}-c_u^*) + 2\epsilon + [(\lambda_\epsilon(c) + \gamma_\epsilon(c))(c_{het}-c) + \lambda_\star - 2\epsilon] e^{-\gamma_\epsilon(c)((c_{het}-c)t+\tau/2)}.$$

First, we can pick $\tau \geq \tau_\epsilon$ really large such that

$$[(\lambda_\epsilon(c) + \gamma_\epsilon(c))(c_{het}-c) + \lambda_\star - 2\epsilon] e^{-\gamma_\epsilon(c)((c_{het}-c)t+\tau/2)} < \epsilon.$$

Then we note that as $c < c_u^* \leq c_{het}$ we have $\lambda(c_u^*) < \lambda(c)$ and

$$-\lambda(c)(c_{het}-c) + \lambda(c_u^*)(c_{het}-c_u^*) < 0;$$

here we recall that $\lambda(c)$ denotes the smallest of the two solutions of $d_- \lambda^2 - c\lambda + \alpha = 0$. Similarly, $\lambda_\epsilon(c)$ denoted the smallest solution of the same equation with α replaced by $\alpha - \epsilon$. As a consequence, we can chose $\epsilon > 0$ small enough such that

$$-\lambda_\epsilon(c)(c_{het}-c) + \lambda(c_u^*)(c_{het}-c_u^*) + 3\epsilon < 0.$$

It follows that $M(\underline{u}_{2,\tau}(t, x)) \leq 0$ for $-\tau/2 < x - c_{het}t < x_\epsilon$. It remains to deal with the jump of the spatial derivative at $x - c_{het}t$, which we adress after the statement for the resulting sub-solution.

Proposition 2. *Let $c \in (c_-, c_+)$ be such that $c_- < c < c_u^* = g^{-1}(0) \leq c_+$. Then, there is $\epsilon_0(c) > 0$ and $\tau_0(\epsilon, c) > 0$ such that for all $0 < \epsilon < \epsilon_0(c)$ and all $\tau \geq \tau_0(\epsilon, c)$, we have that*

$$\underline{u}_{c,\epsilon,\tau}(t, x) = \begin{cases} \underline{v}_{c,\epsilon,\tau}(t, x), & x - c_{het}t \leq -\tau/2, \\ c_{\epsilon,\tau} \underline{v}_{c,\epsilon,\tau}(t, c_{het}t - \tau/2) \tilde{\varphi}_{\lambda_\star-\epsilon}(x - c_{het}t), & x - c_{het}t > -\tau/2, \end{cases}$$

is a sub-solution of (13) for all $(t, x) \in \mathbb{R}^+ \times \mathbb{R}$. Here, we have set

$$\underline{v}_{c,\epsilon,\tau}(t, x) = \max \left\{ 0, e^{-\lambda_\epsilon(c)(x-ct+\tau)} - e^{-(\lambda_\epsilon(c)+\gamma_\epsilon(c))(x-ct+\tau)} \right\},$$

with $\lambda_\epsilon(c) = \frac{c - \sqrt{c^2 - 4d_-(\alpha - \epsilon)}}{2d_-} > 0$ and $0 < \gamma_\epsilon(c) < \frac{1}{d_-} \sqrt{c^2 - 4d_-(\alpha - \epsilon)}$. Moreover, the function $\tilde{\varphi}_{\lambda_\star - \epsilon}(x - c_{het}t)$ is nonnegative and defined from $\varphi_{\lambda_\star - \epsilon}$ which solves $\mathcal{L}\varphi = (\lambda_\star - \epsilon)\varphi$ with prescribed asymptotic behavior at $-\infty$:

$$\varphi_{\lambda_\star - \epsilon}(x) = e^{\left(-\frac{c_{het}}{2d_-} + \frac{1}{2d_-} \sqrt{c_{het}^2 - 4d_-\alpha + 4d_-(\lambda_\star - \epsilon)}\right)x} \left(1 + \mathcal{O}\left(e^{\nu'x}\right)\right), \text{ as } x \rightarrow -\infty,$$

for some $0 < \nu' < \nu$. Finally, the normalizing constant $c_{\epsilon, \tau} > 0$ is given by $c_{\epsilon, \tau} = 1/\varphi_{\lambda_\star - \epsilon}(-\tau/2)$.

We refer to Figure 6 for an illustration.

Proof. Let $c \in (c_-, c_+)$ be such that $c_- < c < c_u^* = g^{-1}(0) \leq c_+$. Then, there exists $\epsilon_0(c) > 0$ such that for all $0 < \epsilon < \epsilon_0(c)$, we have

$$\begin{aligned} \alpha - \frac{c_{het}^2}{4d_-} &< \lambda_\star - \epsilon < \lambda_\star, \\ \lambda_\epsilon(c) &= \frac{c - \sqrt{c^2 - 4d_-(\alpha - \epsilon)}}{2d_-} > 0, \\ -\lambda_\epsilon(c)(c_{het} - c) + \lambda(c_u^*)(c_{het} - c_u^*) + 3\epsilon &< 0. \end{aligned}$$

Then, there exists $\tau_0(\epsilon, c)$ such that for all $\tau \geq \tau_0(\epsilon, c)$ one has

$$\begin{aligned} \frac{\lambda_\epsilon(c)^2}{(\lambda_\epsilon(c) + \gamma_\epsilon(c))^2} &> e^{-\gamma_\epsilon(c)(x - ct + \tau)}, \quad x - c_{het}t \geq x_*(\epsilon, c), \\ [(\lambda_\epsilon(c) + \gamma_\epsilon(c))(c_{het} - c) + \lambda_\star - 2\epsilon] &e^{-\gamma_\epsilon(c)((c_{het} - c)t + \tau/2)} < \epsilon, \end{aligned}$$

where $x_*(\epsilon, c) \in \mathbb{R}$ is defined such that

$$\chi(x - c_{het}t) - d_- \leq \frac{\gamma_\epsilon(c)[c - 2\lambda_\epsilon(c)d_- - \gamma_\epsilon(c)]}{(\lambda_\epsilon(c) + \gamma_\epsilon(c))^2}, \quad x - c_{het}t \leq x_*(\epsilon, c).$$

According to the above discussions, we already have that $\underline{u}_{c, \epsilon, \tau}$ is a sub-solution on each subdomains $x - c_{het}t < -\tau/2$ and $x - c_{het}t > -\tau/2$.

It remains to check that there is a positive jump in the spatial derivative of the sub-solution at $x = c_{het}t - \tau/2$, that is

$$\partial_x \underline{u}_{c, \epsilon, \tau}(t, (c_{het}t - \tau/2)^-) < \partial_x \underline{u}_{c, \epsilon, \tau}(t, (c_{het}t - \tau/2)^+) < 0.$$

First, we compute:

$$\partial_x \underline{u}_{c, \epsilon, \tau}(t, (c_{het}t - \tau/2)^-) = -\lambda_\epsilon(c)e^{-\lambda_\epsilon(c)[(c_{het} - c)t + \tau/2]} \left[1 - \frac{\lambda_\epsilon(c) + \gamma_\epsilon(c)}{\lambda_\epsilon(c)} e^{-\gamma_\epsilon(c)[(c_{het} - c)t + \tau/2]}\right],$$

and

$$\partial_x \underline{u}_{c, \epsilon, \tau}(t, (c_{het}t - \tau/2)^+) = \frac{\varphi'_{\lambda_\star - \epsilon}(-\tau/2)}{\varphi_{\lambda_\star - \epsilon}(-\tau/2)} e^{-\lambda_\epsilon(c)[(c_{het} - c)t + \tau/2]} \left[1 - e^{-\gamma_\epsilon(c)[(c_{het} - c)t + \tau/2]}\right].$$

We now use the prescribed asymptotic behavior at $-\infty$ of $\varphi_{\lambda_\star - \epsilon}$ to get that

$$\frac{\varphi'_{\lambda_\star - \epsilon}(-\tau/2)}{\varphi_{\lambda_\star - \epsilon}(-\tau/2)} \longrightarrow -\frac{c_{het}}{2d_-} + \frac{1}{2d_-} \sqrt{c_{het}^2 - 4d_-\alpha + 4d_-(\lambda_\star - \epsilon)}, \quad \text{as } \tau \rightarrow +\infty.$$

Recall (12), that is

$$\lambda(c_u^*) = \frac{c_{het}}{2d_-} - \frac{1}{2d_-} \sqrt{c_{het}^2 - 4d_-\alpha + 4d_-\lambda_\star},$$

and also that $\lambda(c) > \lambda(c_u^*)$ due to $c < c_u^*$. Thus we can select $\epsilon > 0$ even smaller to have

$$-\lambda_\epsilon(c) < -\frac{c_{het}}{2d_-} + \frac{1}{2d_-} \sqrt{c_{het}^2 - 4d_- \alpha + 4d_- (\lambda_* - \epsilon)}.$$

This implies that upon taking τ larger we can always ensure that

$$\partial_x \underline{u}_{c,\epsilon,\tau,\eta}(t, (c_{het}t - \tau/2)^-) < \partial_x \underline{u}_{c,\epsilon,\tau,\eta}(t, (c_{het}t - \tau/2)^+) < 0.$$

This concludes the proof of Proposition 2. $\square$

We are now in a position to prove that the solution u of (1) with compactly supported initial datum spreads with speed larger than or equal to c_u^* . Take any $c \in (c_-, c_u^*)$ arbitrarily close to c_u^* , and notice that $\underline{u}_{c,\epsilon,\tau}$ from Proposition 2 is uniformly bounded from above since it is compactly supported and continuous. In particular, we can find $\delta_0 > 0$ small enough so that, for any $0 < \delta \leq \delta_0$,

$$0 \leq \delta \underline{u}_{c,\epsilon,\tau} \leq \eta,$$

where η is such that

$$(\alpha - \epsilon)u < \alpha u(1 - u), \quad \text{for } u \in [0, \eta].$$

It is then straightforward to check that, using the linearity of M ,

$$N(\delta \underline{u}_{c,\epsilon,\tau}) < \delta M(\underline{u}_{c,\epsilon,\tau}) \leq 0,$$

i.e. $\delta \underline{u}_{c,\epsilon,\tau}$ is a sub-solution of (1) for any $\delta \leq \delta_0$. Proceeding as in the proof of Proposition 1, we can infer that the solution spreads with speed larger than or equal to c_u^* . We omit the details and Theorem 2.3 is proved in the case when $c_{het} \in [c_+, c_{int})$.

5.2. Case $c_{het} \geq c_{int}$. When $c_{het} \geq c_{int}$, we need to prove that the rightward spreading speed is $c_u^* = c_-$. In this case, using the second statement of Proposition 1 we have already proved that the solution spreads with speed larger than or equal to c_- , and it only remains to construct a super-solution to conclude the proof of Theorem 2.1 in that case. This is precisely the purpose of the next lemma.

Lemma 5.3. *Assume that $c_- < c_+ < c_{int} \leq c_{het}$. There exists $\tau_0 > 0$ such that for each $\tau \geq \tau_0$,*

$$u_\tau(t, x) = \begin{cases} 1, & x \leq c_-t + \tau, \\ e^{-\frac{c_-}{2d_-}(x - c_-t - \tau)}, & c_-t + \tau < x < c_{het}t + \tau, \\ e^{-\frac{c_-}{2d_-}(c_{het} - c_-)t} e^{-\mu_-(x - c_{het}t - \tau)}, & x \geq c_{het}t + \tau, \end{cases}$$

is a super-solution of (1), with $\mu_- := \frac{c_{het} + \sqrt{g(c_-)}}{2d_+} > 0$.

Proof. As $c_{het} \geq c_{int}$, we have that

$$g(c_-) = c_{het}^2 - 4d_+ \left(\alpha + \frac{c_-}{2d_-} (c_{het} - c_-) \right) \geq 0,$$

and μ_- is well-defined. For $x \in (c_-t + \tau, c_{het}t + \tau)$, we have

$$N \left(e^{-\frac{c_-}{2d_-}(x - c_-t - \tau)} \right) \geq \left[\frac{c_-^2}{4d_-^2} (d_- - \chi(\tau)) + \alpha e^{\frac{c_-}{2d_-}\tau} \right] e^{-\frac{c_-}{2d_-}(x - c_-t - \tau)}.$$

Thus, we fix $\tau_0 > 0$ such that for all $\tau \geq \tau_0$

$$\frac{c_-^2}{4d_-^2} (d_- - \chi(\tau)) + \alpha e^{\frac{c_-}{2d_-}\tau} > 0.$$

Next, for $x > c_{het}t + \tau$, we have that

$$N(u_\tau(t, x)) > \left(-\frac{c_-}{2d_-}(c_{het} - c_-) + \mu_- c_{het} - \alpha - d_+ \mu_-^2\right) e^{-\frac{c_-}{2d_-}(c_{het} - c_-)t} e^{-\mu_-(x - c_{het}t - \tau)} = 0,$$

as $\mu_- > 0$ is the largest positive root of

$$-\frac{c_-}{2d_-}(c_{het} - c_-) + \mu c_{het} - \alpha - d_+ \mu^2 = 0.$$

Finally, we have that

$$\mu_- > \frac{c_-}{2d_-},$$

since

$$\mu_- \geq \frac{c_{het}}{2d_+} \geq \frac{c_{int}}{2d_+} > \sqrt{\frac{\alpha}{d_-}} = \frac{c_-}{2d_-}.$$

This insures that the jump of the spatial derivative at $x = c_{het}t + \tau$ has the correct sign. It is also obvious that the jump of the spatial derivative is negative at $x = c_-t + \tau$ and thus u_τ is a super-solution of (1). $\square$

5.3. Case $c_{het} < c_+$. When $c_{het} < c_+$, we need to prove that the rightward spreading speed is $c_u^* = c_+$. In this case, using the first statement of Proposition 1 we have already proved that the spreading speed is less than or equal to c_+ and it only remains to construct a sub-solution to conclude the proof of Theorem 2.3.

We proceed as in the proof of Lemma 4.2. First we let $c_{het} < c < c_+$. Then, one can find $\epsilon > 0$ such that $c < 2\sqrt{d_+}(\alpha - 2\epsilon) < c_+$ together with $\eta_\epsilon > 0$ such that

$$(\alpha - \epsilon)u \leq \alpha u(1 - u), \quad u \in [0, \eta_\epsilon].$$

We again define

$$\beta_+(\epsilon, c) := \frac{\sqrt{4d_+(\alpha - \epsilon) - c^2}}{2d_+} > 0, \text{ and } \beta_+(c) := \beta_+(0, c) = \frac{\sqrt{4d_+\alpha - c^2}}{2d_+} > 0,$$

together with the following family of functions

$$u_{\tau, \epsilon}^c(t, x) := \begin{cases} \delta_\epsilon \left[e^{-\frac{c}{2d_+}(x - ct - \tau)} \cos(\beta_+(\epsilon, c)(x - ct - \tau)) + \epsilon \right], & x - ct - \tau \in \Omega_\epsilon(c), \\ 0, & \text{otherwise,} \end{cases}$$

with $\Omega_\epsilon(c) = \left[-\frac{\pi}{2\beta_+(\epsilon, c)} - z_\epsilon^-(c), \frac{\pi}{2\beta_+(\epsilon, c)} + z_\epsilon^+(c)\right]$, $\delta_\epsilon > 0$ is fixed such that $0 \leq u_{\tau, \epsilon}^c(t, x) \leq \eta_\epsilon$, and

$$z_\epsilon^\pm(c) \in \left(0, \frac{\pi}{2\beta_+(\epsilon, c)}\right),$$

together with

$$e^{\mp \frac{c}{2d_+} \left(\frac{\pi}{2\beta_+(\epsilon, c)} + z_\epsilon^\pm(c) \right)} \sin(\beta_+(\epsilon, c) z_\epsilon^\pm(c)) = \epsilon,$$

which ensure the nonnegativity and continuity of $u_{\tau, \epsilon}^c$. Notice the asymptotics

$$z_\epsilon^\pm(c) = \frac{e^{\mp \frac{c\pi}{4d_+\beta_+(c)}}}{\beta_+(c)} \epsilon + o(\epsilon), \quad \text{as } \epsilon \rightarrow 0.$$

Next, as $\chi(+\infty) = d_+$, there exists $A > 0$ such that for all $\xi \geq A$, we get

$$|\chi(\xi) - d_+| \leq \epsilon^2.$$

Proceeding as for Lemma 4.2, we then find that for any c , this is a sub-solution provided that ϵ is small enough. As in the proof of Proposition 1, one can deduce

that the solution of (1) spreads with speed larger than or equal to c_+ , which ends the proof of Theorem 2.3.

6. Traveling fronts in case (I). Throughout this section we assume that $c_{het} \in (c_-, c_+)$. Before proceeding with the proof of Theorem 2.2, we introduce the following notion of generalized principal eigenvalue, which can be found in [7, 4, 9], for the elliptic operator $\mathcal{L}$ defined as

$$\mathcal{L} = \chi(x)\partial_x^2 + c_{het}\partial_x + \alpha, \quad x \in \mathbb{R}.$$

We define $\mu^* \in \mathbb{R}$ to be

$$\mu^* := \sup \{ \mu \mid \exists \varphi \in \mathcal{C}^2(\mathbb{R}), \varphi > 0, (\mathcal{L} + \mu) \leq 0 \}.$$

One of the key property of the generalized principal eigenvalue μ^* is that it can be obtained as the limit of the Dirichlet principal eigenvalue. More precisely, consider the following Dirichlet problem:

$$\begin{cases} \mathcal{L}\varphi = -\mu\varphi, & |x| < r, \\ \varphi(\pm r) = 0, \end{cases}$$

for each $r > 0$ and denote $\mu_d(r)$ the principal eigenvalue given by Krein-Rutman theory [22]. Then [4, Proposition 4.2] ensures that $r \mapsto \mu_d(r) \in \mathbb{R}$ decreases and

$$\mu_d(r) \longrightarrow \mu^*, \text{ as } r \rightarrow +\infty.$$

We claim that we have the following result.

Lemma 6.1. *When $c_{het} \in (c_-, c_+)$ and χ satisfies (I), then the principal eigenvalue of $\mathcal{L}$ satisfies $\mu^* < 0$.*

Proof. We check that the conditions of [4, Theorem 4.3] are satisfied in our case which translate into our setting by checking that the function $q(x) := 4\chi(x)\alpha - c_{het}^2$ is above a fixed positive constant on some (large) interval. As $c_{het} \in (c_-, c_+)$, we have that $q(-\infty) = 4d_+\alpha - c_{het}^2 > 0$. As a consequence, there exists $\epsilon > 0$ and $x_0 < 0$ such that for all $x \leq x_0$, we have

$$q(x) = 4\chi(x)\alpha - c_{het}^2 \geq \epsilon,$$

which implies that $\mu^* < 0$ from [4, Theorem 4.3]. $\square$

Existence. In order to prove the existence of traveling front solutions, we are first going to construct generalized sub and super-solutions for (3). The construction of the sub-solution relies on the aforementioned properties of the generalized principal eigenvalue μ^* . For each $r > 0$, we denote by φ_r the corresponding eigenfunction to the Dirichlet principal eigenvalue $\mu_d(r)$ which satisfies $\varphi_r > 0$ and normalized with $\varphi_r(0) = 1$. As $\mu^* < 0$ from Lemma 6.1, there exists some $R > 0$ such that for any $r \geq R$ we also have $\mu_d(r) < 0$. As a consequence, there exists $0 < \kappa_r < 1$ small enough such that

$$(\alpha + \mu_d(r))(\kappa_r \varphi_r) \leq \alpha \kappa_r \varphi_r (1 - \kappa_r \varphi_r).$$

Thus, if one defines $\underline{U}_r$ the following family of functions

$$\underline{U}_r(x) = \begin{cases} \kappa_r \varphi_r(x), & |x| \leq r, \\ 0, & \text{otherwise,} \end{cases}$$

then for all $r \geq R$, the function $\underline{U}_r$ is a generalized sub-solution to (3).

Next, as $c_{het} \in (c_-, c_+)$, there exists $\epsilon_0 > 0$ small enough such that for each $0 < \epsilon < \epsilon_0$ one has $c_- < 2\sqrt{(d_- + \epsilon)\alpha} < c_{het}$. For such an ϵ , one can find $\tau_\epsilon > 0$ such that for any $\tau \geq \tau_\epsilon$ we have

$$d_- \leq \chi(x) \leq d_- + \epsilon, \quad x \geq \tau.$$

We now introduce $\overline{U}_{\epsilon, \tau}$ defined as

$$\overline{U}_{\epsilon, \tau}(x) = \max \left\{ 1, e^{-\lambda_\epsilon(x-\tau)} \right\}, \quad \lambda_\epsilon = \frac{c_{het} + \sqrt{c_{het}^2 - 4(d_- + \epsilon)\alpha}}{2(d_- + \epsilon)} > 0.$$

For all $\tau \geq \tau_\epsilon$, one can check that $\overline{U}_{\epsilon, \tau}$ is a generalized super-solution to (3). Up to further reducing κ_r (or taking τ larger), we can always ensure that

$$0 \leq \underline{U}_r \leq \overline{U}_{\epsilon, \tau} \leq 1,$$

for some $r \geq R$, $\epsilon \in (0, \epsilon_0)$ and $\tau \geq \tau_\epsilon$.

Now denote by u the solution of

$$\partial_t u = \chi(x) \partial_x^2 u + c_{het} \partial_x u + \alpha u(1 - u), \quad (14)$$

with the initial condition

$$u(t = 0, \cdot) \equiv \overline{U}_{\epsilon, \tau}.$$

Since $\overline{U}_{\epsilon, \tau}$ is a super-solution of (3), hence also of (14), it follows from parabolic comparison principles that u is nonincreasing in the time variable. Therefore it converges to some function U as $t \rightarrow +\infty$, and by parabolic estimates we get that U is a solution of (3).

Moreover, by construction and another use of the comparison principle, we get that

$$0 \leq \underline{U}_r \leq U \leq \overline{U}_{\epsilon, \tau} \leq 1, \quad \text{on } \mathbb{R}.$$

Using the strong maximum principle, we actually get that $0 < U < 1$. Indeed, assume there is $x_0 \in \mathbb{R}$ for which $U(x_0) = 0$, then the strong maximum principle implies that $0 \leq \underline{U}_r \leq U \equiv 0$, which is impossible. A similar argument holds for the other inequality.

We also remark that U must be nonincreasing. Indeed, notice that

$$\partial_x u(t = 0) = \overline{U}'_{\epsilon, \tau} \leq 0.$$

Moreover, derivating (14), we get that $v = \partial_x u$ solves

$$\partial_t v = \chi(x) \partial_x^2 v + (c_{het} + \chi'(x)) \partial_x v + \alpha(1 - 2u)v.$$

By another comparison principle, we find that

$$v(t, x) \leq 0,$$

for all $t > 0$ and $x \in \mathbb{R}$, hence $U' \leq 0$.

This latter fact combined with $0 \leq \underline{U}_r \leq U$ ensures that $U(-\infty) > 0$, and one necessarily gets that $U(-\infty) = 1$ as U is solution of the ODE (3). Next, using the fact that $\overline{U}_{\epsilon, \tau}(x) \rightarrow 0$ as $x \rightarrow +\infty$, we get that $U(+\infty) = 0$ by comparison. We claim that we actually have $U' < 0$ on $\mathbb{R}$. This property is satisfied near $-\infty$ as there exists a unique stable direction. We let $x_* \in \mathbb{R}$ be such that $U'(x_*) = 0$ and $U'(x) < 0$ for all $x < x_*$. Then, we have

$$\chi(x_*) U''(x_*) = -\alpha U(x_*) (1 - U(x_*)) < 0,$$

which is a contradiction.

Asymptotic behavior at $+\infty$. As $c_{het} > c_-$ we have that $\frac{\sqrt{c_{het}^2 - 4d_- \alpha}}{2d_-} > 0$, and upon eventually reducing $\epsilon_0 > 0$, we can ensure that for all $\epsilon \in (0, \epsilon_0)$,

$$\lambda_\epsilon > \frac{c_{het}}{2d_-}.$$

As a consequence, we have that for all $x \geq \tau$

$$0 < e^{\frac{c_{het}}{2d_-}x} U(x) \leq e^{\frac{c_{het}}{2d_-}x} \overline{U}_{\epsilon, \tau}(x) = e^{-(\lambda_\epsilon - \frac{c_{het}}{2d_-})x + \lambda_\epsilon \tau}.$$

We now prove that U has the strong exponential decay given in Theorem 2.2. Near $+\infty$, system (3) can be written in condensed form

$$\mathbf{U}'(x) = A(x)\mathbf{U}(x) + N(x, \mathbf{U}(x)), \quad x \in \mathbb{R}, \quad (15)$$

with $\mathbf{U}(x) = (U(x), V(x))^t$ and

$$A(x) := \begin{pmatrix} 0 & 1 \\ -\frac{\alpha}{\chi(x)} & -\frac{c_{het}}{\chi(x)} \end{pmatrix}, \quad N(x, \mathbf{U}) := \begin{pmatrix} 0 \\ \frac{\alpha}{\chi(x)} U^2 \end{pmatrix}.$$

As χ converges at an exponential rate at $+\infty$, so does $A(x)$, and since there is a gap between the strong stable and weak stable eigenvalues of $A_\infty := \lim_{x \rightarrow +\infty} A(x)$, the constant coefficient asymptotic system has an exponential dichotomy. By classical arguments on the robustness of exponential dichotomies [11], the non-autonomous system inherits one with the same decay rates as χ converges at an exponential rate at $+\infty$. Note that the strong stable eigenvalue is precisely given by $-\lambda_s$ defined in Theorem 2.2 while the weak stable eigenvalue is $-\lambda_w$ given by

$$0 < \lambda_w := \frac{c_{het} - \sqrt{c_{het}^2 - c_-^2}}{2d_-} < \lambda_s.$$

As a consequence, since we have $0 < e^{\frac{c_{het}}{2d_-}x} U(x) \leq e^{-(\lambda_\epsilon - \frac{c_{het}}{2d_-})x + \lambda_\epsilon \tau}$ for all $x \geq \tau$, we deduce that necessarily

$$U(x) \underset{x \rightarrow +\infty}{\sim} \gamma_s e^{-\lambda_s x},$$

for some $\gamma_s > 0$.

Uniqueness. We first prove that when $c_{het} > c_- = 2\sqrt{d_- \alpha}$ solutions of (3)-(4) are unique in $H_{\frac{c_{het}}{2d_-}}^1(\mathbb{R}) := \left\{ U \mid e^{\frac{c_{het}}{2d_-}x} U \in H^1(\mathbb{R}) \right\}$. Let $U \in H_{\frac{c_{het}}{2d_-}}^1(\mathbb{R})$ and $V \in H_{\frac{c_{het}}{2d_-}}^1(\mathbb{R})$ be two solutions of (3)-(4) and assume by contradiction that $U \not\equiv V$.

Without loss of generality, we may assume that $U(x) > V(x)$ on some interval $(a, b) \subset \mathbb{R}$ with $U(a) = V(a)$ and $U(b) = V(b)$. Note that $a, b \in \mathbb{R} \cup \{\pm\infty\}$. Multiplying the equation for U with $e^{\beta(x)} \frac{V(x)}{\chi(x)}$ and the equation for V with $e^{\beta(x)} \frac{U(x)}{\chi(x)}$, where we set $\beta(x) := \int_0^x \frac{c_{het}}{\chi(y)} dy$, we obtain

$$\begin{cases} 0 = V(x) \frac{d}{dx} \left(e^{\beta(x)} U'(x) \right) + \frac{\alpha}{\chi(x)} U(x) V(x) (1 - U(x)) e^{\beta(x)}, \\ 0 = U(x) \frac{d}{dx} \left(e^{\beta(x)} V'(x) \right) + \frac{\alpha}{\chi(x)} U(x) V(x) (1 - V(x)) e^{\beta(x)}. \end{cases}$$

As a consequence, we have

$$\int_a^b V(x) \frac{d}{dx} \left(e^{\beta(x)} U'(x) \right) - U(x) \frac{d}{dx} \left(e^{\beta(x)} V'(x) \right) dx = \alpha \int_a^b \frac{U(x) V(x)}{\chi(x)} (U(x) - V(x)) e^{\beta(x)} dx.$$

Note that the above integrals are convergent as on the one hand $U, V \in H^1_{\frac{c_{het}}{2d}}(\mathbb{R})$ which ensures integrability when $b = +\infty$, and on the other hand $\beta(x) \sim \frac{c_{het}}{d_+}x$ as $x \rightarrow -\infty$ which ensures integrability when $a = -\infty$ (recall that $U(-\infty) = V(-\infty) = 1$, and the convergence must be exponential by classical arguments on (15)). Integrating by parts the integral on the left-hand side of the equality, we obtain

$$e^{\beta(x)} (V(x)U'(x) - U(x)V'(x)) \Big|_a^b = \alpha \int_a^b \frac{U(x)V(x)}{\chi(x)} (U(x) - V(x)) e^{\beta(x)} dx.$$

When both $a, b \in \mathbb{R}$, we have that the right-hand side is strictly positive while the left-hand side is negative using that $U(x) = V(x)$ at $x \in \{a, b\}$ and $U(x) > V(x)$ for $x \in (a, b)$. If $a = -\infty$, we have that

$$e^{\beta(x)} (V(x)U'(x) - U(x)V'(x)) \rightarrow 0, \quad x \rightarrow -\infty,$$

and the left-hand side is negative. If $b = +\infty$, we have that

$$e^{\beta(x)} |V(x)U'(x)| = e^{\int_0^x c_{het} \left(\frac{1}{\chi(y)} - \frac{1}{d_-} \right) dy} e^{\frac{c_{het}}{d_-} x} |V(x)U'(x)| \rightarrow 0, \quad x \rightarrow +\infty,$$

$$e^{\beta(x)} |V'(x)U(x)| = e^{\int_0^x c_{het} \left(\frac{1}{\chi(y)} - \frac{1}{d_-} \right) dy} e^{\frac{c_{het}}{d_-} x} |V'(x)U(x)| \rightarrow 0, \quad x \rightarrow +\infty,$$

and the left-hand side is again negative. Thus, we have reached a contradiction, and the solution of (3)-(4) with strong exponential decay at $+\infty$, if it exists, is unique. Non existence of solutions of (3)-(4) when $c_{het} < c_-$. Using the fact that $\chi(+\infty) = d_- > 0$, we readily obtain the necessary condition that $c_{het} \geq 2\sqrt{d_- \alpha} = c_-$ for the corresponding solution to remain positive near the equilibrium $u = 0$.

Non existence of solutions of (3)-(4) when $c_{het} > c_+$ with strong exponential decay. Let us assume that $c_{het} > c_+ = 2\sqrt{d_+ \alpha}$. We are going to prove that any solution satisfies

$$U'(x) > -\frac{c_{het}}{2d_+} U(x), \quad x \in \mathbb{R}. \quad (16)$$

We know that the above inequality holds true near $-\infty$, and suppose by contradiction that $x_* \in \mathbb{R}$ is the leftmost point where

$$\begin{cases} 0 = \chi(x_*)U''(x_*) + c_{het}U'(x_*) + \alpha U(x_*)(1 - U(x_*)), \\ U'(x_*) = -\frac{c_{het}}{2d_+} U(x_*). \end{cases}$$

Using the fact that $\alpha U(1 - U) \leq \alpha U < \frac{c_{het}^2}{4d_+} U$, we have that

$$\begin{aligned} \chi(x_*)U''(x_*) &= -c_{het}U'(x_*) - \alpha U(x_*)(1 - U(x_*)) \\ &> -c_{het}U'(x_*) - \frac{c_{het}^2}{4d_+} U(x_*), \\ &= -\frac{c_{het}}{2} U'(x_*), \end{aligned}$$

from which we deduce that $U''(x_*) > -\frac{c_{het}}{2d_+} U'(x_*)$ as $\chi(x_*) \leq d_+$ and $U'(x_*) < 0$. This is a contradiction since by construction $U' > -\frac{c_{het}}{2d_+} U$ on $(-\infty, x_*)$ and $U'(x_*) = -\frac{c_{het}}{2d_+} U(x_*)$. Therefore (16) must hold for all x . Integrating (16) we obtain that

$$U(x) > e^{-\frac{c_{het}}{2d_+} x} U(0), \quad x > 0,$$

and thus the solution U , if it exists, has weak exponential decay near $+\infty$. As a consequence, as $0 < \frac{c_{het}}{2d_+} < \frac{c_{het}}{2d_-} < \lambda_s$, when $c_{het} > 2\sqrt{d_+\alpha}$ there cannot exist a solution of (3)-(4) with strong exponential decay given by λ_s .

REFERENCES

- [1] D. G. Aronson and H. F. Weinberger, [Multidimensional nonlinear diffusion arising in population genetics](#), *Adv. in Math.*, **30** (1978), 33–76.
- [2] H. Berestycki, O. Diekmann, C. J. Nagelkerke and P. A. Zegeling, [Can a species keep pace with a shifting climate?](#), *Bull. Math. Biol.*, **71** (2009), 399–429.
- [3] H. Berestycki and J. Fang, [Forced waves of the Fisher–KPP equation in a shifting environment](#), *J. Differential Equations*, **264** (2018), 2157–2183.
- [4] H. Berestycki, F. Hamel and L. Rossi, [Liouville-type results for semilinear elliptic equations in unbounded domains](#), *Ann. Mat. Pura Appl.*, **186** (2007), 469–507.
- [5] H. Berestycki and G. Nadin, Asymptotic spreading for general heterogeneous equations, *Memoirs of the American Mathematical Society*.
- [6] H. Berestycki and G. Nadin, [Spreading speeds for one-dimensional monostable reaction-diffusion equations](#), *J. Math. Phys.*, **53** (2012), 115619, 23pp.
- [7] H. Berestycki and L. Rossi, [On the principal eigenvalue of elliptic operators in \$\mathbb{R}^n\$ and applications](#), *J. Eur. Math. Soc. (JEMS)*, **8** (2006), 195–215.
- [8] H. Berestycki and L. Rossi, [Reaction-diffusion equations for population dynamics with forced speed I—the case of the whole space](#), *Discrete Contin. Dyn. Syst.*, **21** (2008), 41–67.
- [9] H. Berestycki and L. Rossi, [Generalizations and properties of the principal eigenvalue of elliptic operators in unbounded domains](#), *Comm. Pure Appl. Math.*, **68** (2015), 1014–1065.
- [10] J. Bouhours and T. Giletti, [Spreading and vanishing for a monostable reaction–diffusion equation with forced speed](#), *J. Dynam. Differential Equations*, **31** (2019), 247–286.
- [11] W. A. Coppel, *Dichotomies in Stability Theory*, Vol. 629. Springer-Verlag, Berlin-New York, 1978.
- [12] C. Cosner, [Challenges in modeling biological invasions and population distributions in a changing climate](#), *Ecological Complexity*, **20** (2014), 258–263.
- [13] A. Ducrot, T. Giletti and H. Matano, [Spreading speeds for multidimensional reaction-diffusion systems of the prey-predator type](#), *Calc. Var. Partial Differential Equations*, **58** (2019), 137, 34pp.
- [14] G. Faye and M. Holzer, [Bifurcation to locked fronts in two component reaction-diffusion systems](#), *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **36** (2019), 545–584.
- [15] R. A. Fisher, [The wave of advance of advantageous genes](#), *Annals of Eugenics*, **7** (1937), 355–369.
- [16] T. Gallay and C. Mascia, [Propagation fronts in a simplified model of tumor growth with degenerate cross-dependent self-diffusivity](#), *Nonlinear Anal. Real World Appl.*, **63** (2022), 103387, 28pp.
- [17] R. A. Gatenby and E. T. Gawlinski, A reaction-diffusion model of cancer invasion, *Cancer Research*, **56** (1996), 5745–5753.
- [18] L. Girardin and K.-Y. Lam, [Invasion of open space by two competitors: Spreading properties of monostable two-species competition-diffusion systems](#), *Proc. Lond. Math. Soc.*, **119** (2019), 1279–1335.
- [19] M. Holzer and A. Scheel, [Accelerated fronts in a two-stage invasion process](#), *SIAM J. Math. Anal.*, **46** (2014), 397–427.
- [20] T. Kapitula and K. Promislow, *Spectral and Dynamical Stability of Nonlinear Waves*, Applied Mathematical Sciences, 185. Springer, New York, 2013.
- [21] A. N. Kolmogorov, Étude de l’équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique, *Bull. Univ. Moskow, Ser. Internat., Sec. A*, **1** (1937), 1–25.
- [22] M. G. Krein and M. A. Rutman, Linear operators leaving invariant a cone in a banach space, *Uspehi Matem. Nauk (N. S.)*, **3** (1948), 3–95.
- [23] K.-Y. Lam and X. Yu, Asymptotic spreading of kpp reactive fronts in heterogeneous shifting environments, preprint, [arXiv:2101.06698](#), 2021.
- [24] M. Lewis and J. Murray, [Modelling territoriality and wolf–deer interactions](#), *Nature*, **366** (1993), 738–740.

- [25] B. Li, S. Bewick, J. Shang and W. F. Fagan, [Persistence and spread of a species with a shifting habitat edge](#), *SIAM J. Appl. Math.*, **74** (2014), 1397–1417.
- [26] B. Sandstede and A. Scheel, [Absolute and convective instabilities of waves on unbounded and large bounded domains](#), *Phys. D*, **145** (2000), 233–277.
- [27] N. Shigesada, K. Kawasaki and E. Teramoto, [Spatial segregation of interacting species](#), *J. Theoret. Biol.*, **79** (1979), 83–99.

Received March 2021; revised September 2021; early access November 2021.

E-mail address: gregory.faye@math.univ-toulouse.fr

E-mail address: thomas.giletti@univ-lorraine.fr

E-mail address: mholzer@gmu.edu