

## RESEARCH ARTICLE

# Population structure and connectivity among coastal and freshwater Kelp Gull (*Larus dominicanus*) populations from Patagonia

Tatiana Kasinsky  <sup>1</sup>\*, Natalia Rosciano  <sup>2</sup>‡, Juliana A. Vianna <sup>3,4</sup>‡, Pablo Yorio <sup>1,5</sup>‡, Leonardo Campagna  <sup>6,7</sup>¶

**1** Centro Para El Estudio de Sistemas Marinos, CONICET, Puerto Madryn, Chubut, Argentina, **2** Instituto de Investigaciones en Biodiversidad y Medio Ambiente, CONICET, Universidad Nacional del Comahue, San Carlos de Bariloche, Río Negro, Argentina, **3** Departamento de Ecología, Facultad de Ciencias Biológicas, Instituto para el Desarrollo Sustentable, Pontificia Universidad Católica de Chile, Santiago, Chile, **4** Millennium Institute Center for Genome Regulation (CRG), Millennium Institute of Biodiversity of Antarctic and Subantarctic Ecosystems (BASE), Millennium Nucleus of Patagonian Limit of Life (LiLi), Valdivia, Chile, **5** Wildlife Conservation Society Argentina, Ciudad Autónoma de Buenos Aires, Argentina, **6** Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY, United States of America, **7** Fuller Evolutionary Biology Program, Cornell Lab of Ornithology, Ithaca, NY, United States of America

¶ These authors contributed equally to this work.  
 ‡ NR, JAV and PY also contributed equally to this work.  
 \* [kasinsky@cenpat-conicet.gob.ar](mailto:kasinsky@cenpat-conicet.gob.ar)



## OPEN ACCESS

**Citation:** Kasinsky T, Rosciano N, A. Vianna J, Yorio P, Campagna L (2024) Population structure and connectivity among coastal and freshwater Kelp Gull (*Larus dominicanus*) populations from Patagonia. PLoS ONE 19(4): e0301004. <https://doi.org/10.1371/journal.pone.0301004>

**Editor:** Vitor Hugo Rodrigues Paiva, MARE – Marine and Environmental Sciences Centre, PORTUGAL

**Received:** September 26, 2023

**Accepted:** March 9, 2024

**Published:** April 18, 2024

**Copyright:** © 2024 Kasinsky et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its [Supporting Information files](#).

**Funding:** This project was funded by NSF DEB-2232929 (to L.C.), Wildlife Conservation Society Argentina (to P.Y.) and ANID – Programa Iniciativa Milenio – ICN2021\_044 (CGR), ICN2021\_002 (BASE) and NCN2021-050 (LiLi) to J.A.V.

**Competing interests:** The authors have declared that no competing interests exist.

## Abstract

The genetic identification of evolutionary significant units and information on their connectivity can be used to design effective management and conservation plans for species of concern. Despite having high dispersal capacity, several seabird species show population structure due to both abiotic and biotic barriers to gene flow. The Kelp Gull is the most abundant species of gull in the southern hemisphere. In Argentina it reproduces in both marine and freshwater environments, with more than 100,000 breeding pairs following a metapopulation dynamic across 140 colonies in the Atlantic coast of Patagonia. However, little is known about the demography and connectivity of inland populations. We aim to provide information on the connectivity of the largest freshwater colonies (those from Nahuel Huapi Lake) with the closest Pacific and Atlantic populations to evaluate if these freshwater colonies are receiving immigrants from the larger coastal populations. We sampled three geographic regions (Nahuel Huapi Lake and the Atlantic and Pacific coasts) and employed a reduced-representation genomic approach to genotype individuals for single-nucleotide polymorphisms (SNPs). Using clustering and phylogenetic analyses we found three genetic groups, each corresponding to one of our sampled regions. Individuals from marine environments are more closely related to each other than to those from Nahuel Huapi Lake, indicating that the latter population constitutes the first freshwater Kelp Gull colony to be identified as an evolutionary significant unit in Patagonia.


## Introduction

Uncovering the mechanisms behind population genetic divergence is relevant for understanding both evolutionary processes and identifying evolutionary significant units for species in need of management and conservation strategies [1–4]. Moreover, the implementation of genomic techniques in the fields of ecology and conservation genetics has resulted in an increased ability to delimit populations, generate more complex models of demographic histories, and identify regions of the genome that confer important ecological functions and contribute to adaptation [4, 5].

Seabirds reproduce in discrete spatial populations (colonies); however, their ability to disperse over long distances [6] can lead to varying degrees of connectivity, resulting in metapopulation dynamics. However, several studies show some species have strong population structure, resulting from factors such as their non-breeding distribution, physical barriers to dispersal, the geographic distance between colonies, philopatry, and individual dispersal patterns [7].

The Kelp Gull (*Larus dominicanus*) is the most abundant and widely distributed gull species in the southern hemisphere, nesting in South America, Africa, New Zealand, Australia, sub-Antarctic islands, and the Antarctic Peninsula [8]. In Chile, they nest in at least 26 locations along the Pacific coast [9]. The estimated population in Chile comprises approximately ~8,200 breeding pairs, and colony sizes vary, ranging from just a few pairs to as many as 3,000, though due to the difficulty in surveying the geographically complex southern region, the actual numbers may potentially be higher [see 9]. Information regarding population trends in Chile is relatively scarce; however, preliminary data suggest an upward trend in the number of breeding pairs nesting on the rooftops of coastal cities [9]. On the other hand, in Argentina, the Kelp Gull breeds in both marine and freshwater environments [10], and its reproductive population in marine environments of Atlantic Patagonia was estimated at 106,200 breeding pairs distributed in 140 colonies [see 9]. Long-term monitoring of more than sixty colonies in northern Patagonia and the development of demographic models have shown high population connectivity and increases in numbers in recent decades, with demographic behaviors that vary depending on the breeding location [11, 12]. Reproductive Kelp Gull colonies in freshwater environments are limited to a few locations in Patagonia and Uruguay [9, 13, 14]. Little is known about demographic trends from freshwater populations in Argentinian Patagonia, however in the case of the colonies from Lake Nahuel Huapi, there has been an increase in the number of individuals breeding at one of the most important nesting sites, archipelago Islas de los Fósiles, within less than a decade (from 350 nests in 2012 to 575 nests in 2019).

Studies conducted on gulls belonging to the genus *Larus* in various parts of the world have revealed genetic structure among populations, often following a pattern of isolation by distance [15–18]. Moreover, Herring Gull (*Larus argentatus*) populations from the Laurentian Great Lakes were found to be genetically distinct from their marine counterparts [19, 20], supporting the idea that reproductive Herring Gulls from the Great Lakes region constitute a closed and isolated system [21]. Genetic studies in the Kelp Gull show different degrees of population structure depending on their geographic scale, over a background of demographic expansion during the Holocene [16, 22–24]. However, the resolution provided by the genetic tools employed (mitochondrial DNA and microsatellite markers) and the scope of the sampling, particularly in the Patagonian region, may be underpowered to detect fine-scale population differences and adequately understand population connectivity. Studies carried out on other highly mobile seabirds have shown that the higher resolution of genomic data can reveal genetic structure that was previously undetected with mtDNA and microsatellite markers [25, 26]. In addition, Kelp Gull population studies in Argentina have focused on coastal regions,



**Fig 1. Map of the study area showing the Kelp Gull colonies from three regions.** Pacific coast: Maiquillahue (n = 7), Isla Conejo (n = 3), Puñihuil (n = 4). Nahuel Huapi Lake: islets Gaviotero 1 (n = 36) and Gaviotero 2 (n = 32) (archipelago Islas de los Fósiles), Isla el Roble (n = 4). Atlantic coast: Isla de Los Pájaros (n = 15), Punta León (n = 16).

<https://doi.org/10.1371/journal.pone.0301004.g001>

leaving an information gap regarding freshwater colonies. This study aims to provide information on the connectivity of the Lake Nahuel Huapi freshwater population with the closest coastal populations of the Pacific and Atlantic and to evaluate whether this freshwater population is an evolutionary significant unit. If the genetic structure in this species follows a pattern of isolation by distance, we would expect the Nahuel Huapi Lake individuals to show a stronger connection to those found along the Pacific coast, owing to the significantly shorter geographic distance than with respect to the Atlantic coast (see Fig 1). Despite the Andes Mountains potentially acting as a barrier between Nahuel Huapi Lake and the Pacific coast, it is conceivable that Kelp Gulls could be utilizing the interconnected network of lakes and rivers as natural conduits for genetic exchange between the two regions.

## Methods

### Fieldwork and sampling

We collected a total of 117 Kelp Gull samples (110 from blood and 7 from eggshells) across three different regions during the reproductive season (October to January of 2021 and 2022). The first region was located on the Pacific coast in Chile (PC, n = 14), where we sampled the colonies of Maiquillahue ( $39^{\circ} 27'S$ ,  $73^{\circ} 16'W$ ), Puñihuil ( $41^{\circ} 55'S$ ,  $74^{\circ} 2'W$ ), and Isla Conejo ( $42^{\circ} 54'S$ ,  $73^{\circ} 35'W$ ). The second area was Nahuel Huapi Lake (NHL, n = 72), Argentina, where we collected samples from three colonies: the islets Gaviotero 1 and Gaviotero 2 ( $41^{\circ} 0'S$ ,  $71^{\circ} 32'W$ ) within the archipelago Islas de los Fósiles, and Isla el Roble ( $41^{\circ} 00'S$ ,  $71^{\circ} 27'W$ ). Finally, the third region was the Atlantic coast in Argentina (AC, n = 31), where we sampled the colonies of Punta León ( $43^{\circ} 04'S$ ,  $64^{\circ} 29'W$ ) and Isla de Los Pájaros ( $42^{\circ} 25'S$ ,  $64^{\circ} 31'W$ ) (Fig 1). We captured the incubating adult from nests using a leg-noose trap or incubation traps [27] and subsequently collected a blood sample. In cases where it was impossible to capture adults, membranes were collected from newly hatched eggs, or blood (0.3–0.5 mL) was taken from non-related chicks (belonging to different nests and sectors of the colony). We conserved all samples in 96% ethanol until DNA extraction. We obtained the relevant permits for collecting and transporting samples from the corresponding national authorities in Argentina (Administración de Parques Nacionales DRPN N°1814, Dirección de Flora y Fauna Silvestre del Chubut Disp. N° 27/2022-DFyFS, Ministerio de Ambiente y Desarrollo Sostenible de la Argentina CE-2022-47128129-APN-DNBI#MAD, CE-2022-45430867-APN-DNBI#MAD) and Chile (Servicio Agrícola y Ganadero, La Corporación Nacional Forestal N° 5446/2022).

### Marker discovery and genotyping

We generated double digest restriction-site associated DNA markers (ddRAD tags) by first extracting genomic DNA using the DNeasy blood and tissue kit (Qiagen, Valencia, CA, USA). The DNA of each individual was digested with two restriction enzymes (*SbfI* and *MspI*; New England Biolabs; Ipswich, MA) and then ligated to specific adapters at the 5' and 3' ends for downstream bioinformatic processing. Adapter-ligated DNA from different individuals was combined into nine index groups and DNA fragments of between 400 and 700 base pairs were selected using the BluePippin size selection system (Sage Science, Beverly, MA, USA). The ddRAD protocol is described in detail in [28]. Index groups were combined into a single genomic library and sequencing took place on a lane of Illumina's NextSeq500 platform at the Cornell Institute of Biotechnology (single-end x 150 bp mode), together with samples from another project.

We obtained an average of  $453,047 \pm 219,091$  raw Illumina reads per individual. We evaluated read quality using FastQC version 0.11.6 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). All sequences were trimmed to 112 bp using fastX trimmer [29] to remove lower-quality base pairs towards the 3' end. Subsequently, reads that had any base calls with a quality score below Phred 10 (90% call accuracy) or with more than 20% of their base calls between 10 and 20 (90–99% call accuracy) were filtered using fastq\_quality\_filter (fastx-Toolkit). The reads were then demultiplexed and adapters were removed using the process\_ddrtags module of the Stacks pipeline version 2.3e [30] to obtain files containing sequences specific to each individual. All sequences were trimmed to 105 bp after 5–7 bp inline barcodes were removed. Individual sequences were assembled using two methods implemented in the Stacks bioinformatics pipeline, a *de novo* pipeline, and a reference-based pipeline using the Herring Gull (*Larus smithsonianus*) reference genome obtained from [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) (GCA\_013400295.1\_ASM1340029v1).

We aligned sequences to the Herring Gull reference genome using Bowtie 2 [31] with an average alignment rate of 89.17%. We subsequently assembled reads into markers using Stack's gstacks module. This pipeline generated a catalog of 148,795 RAD loci with an effective per-sample coverage of  $19.5 \pm 9.0$ . After filtering out markers absent in more than 15% of individuals and with a minor allele count lower than 5 with the *populations* module, we retained and exported 6,569 SNPs. The *de novo* assembly was performed using 'ustacks/cstacks/sstacks' as executed by the "denovo\_map" pipeline and produced 2,735 SNPs after filtering. However, due to the similarity in the results from both assembly strategies, we decided to present those obtained from the reference-based assembly. We discarded five individuals with missing data higher than 25% (three from Gaviotero 1 and one from Gaviotero 2 in NHL, plus one from Isla de Los Pájaros in the Atlantic coast), retaining 112 individuals for all downstream analyses.

### Mitochondrial DNA amplification and sequencing

We obtained mitochondrial DNA (cytochrome c oxidase I, COI) from a subsample of five individuals from each population. COI amplification was conducted using polymerase chain reactions (PCRs) carried out as described in [32]. PCR products were cleaned by performing an Exo-SAP treatment (exonuclease -Exo- and shrimp alkaline phosphatase—SAP) before sequencing by incubating 10  $\mu$ l of PCR product with 5 U of Exonuclease I and 0.5 U of SAP for 30 min at 37 C, followed by an inactivation step at 90 C for 10 min. Finally, the samples were Sanger sequenced bidirectionally at the Cornell Institute of Biotechnology with the same primers used for amplification (BirdF1 and COIBirdR2, [32]). Sequences were aligned using BioEdit v.7.0.5.3 [33], and alignments were checked manually. Relationships between haplotypes (maternally inherited) were estimated based on the Median-Joining method using PopART 1.7 [34].


### Population genomic analyses

We measured the degree of genetic differentiation among populations by conducting analyses of population genetics and landscape genetics based on allelic and haplotype frequencies. We evaluated genetic structure using a principal component analysis (PCA) generated with the SNPRelate package version 3.3 [35] in R 3.5.2 [36] and with the STRUCTURE 2.3.4 [37] and Admixture 1.3.0 [38] programs, exploring values of K from 1 to 6. For the Admixture analysis we first thinned our dataset to avoid including linked markers by retaining 3,958 SNPs that were at least 10 kb apart, using VCFtools version 0.1.16 [39]. We compared models with different K values using the cross-validation method with 10 iterations. We chose to plot the results for K = 3 in the main manuscript as these are concordant with our other analyses. K values beyond this point did not show further structure and are displayed in the Supplementary Figures in [S1 File](#). For the Structure analysis, we exported a single SNP per locus (retaining 4,527 SNPs), selected randomly using the populations Stacks module, to avoid including tightly linked SNPs. Each run consisted of 500,000 generations after 250,000 iterations were discarded as burn-in. We ran 10 iterations per value of K and did not incorporate sample location in the model. We implemented the admixture ancestry model with correlated allele frequencies and  $\lambda = 1$  as the allele frequency prior. The level of genomic differentiation (between sectors: Pacific coast, Nahuel Huapi Lake, and Atlantic coast) was determined by calculating the  $F_{ST}$  values for each SNP, and by performing an AMOVA (analysis of molecular variance) with the *population* module in Stacks. We also used the *population* module to calculate heterozygosity, nucleotide diversity and inbreeding coefficients. We ran fineRADstructure v0.3 [40], which clusters individuals based on

haplotype frequencies to determine population structure, conducting 100,000 iterations following 100,000 burn in repetitions (sampling every 1,000 iterations), and inferring a tree. We built a Maximum Likelihood phylogenetic tree in RaxML version 8.2.4 [41] after implementing the ASC\_GTRGAMMA model and the Lewis correction for ascertainment bias and conducting 200 bootstrap replicates to assess node support. We used VCFtools to calculate relatedness values among individuals from the Nahuel Huapi Lake colonies. Finally, we estimated migration among geographic regions using BayesAss-SNPs 3.0.5.6 [42], running the program on the same SNP dataset used for the Admixture analysis, for 10 million generations following a burn in of four million generations. We fine-tuned the MCMC mixing parameters by using the following commands: -a0.23 -f0.01. We assessed convergence by inspecting traces and effective sample sizes in Tracer 1.7.1 [43].

## Results

We found individuals from the Pacific coast, Nahuel Huapi Lake, and the Atlantic coast belong to three genetically differentiated populations ( $F_{ST}$  and  $\Phi_{ST}$ : NHL vs. PC = 0.018 and 0.04, respectively; NHL vs. AC = 0.023 and 0.04; PC vs. AC = 0.027 and 0.03). The PCA (based on 6,569 SNPs) shows three distinct clusters corresponding to each one of the three geographic regions (Fig 2A). The first principal component (PC1) explains 4.14% of the total variance, separating the freshwater and coastal populations. PC2 explains 2.3% of the variance and separates the Pacific colonies from the remaining individuals, as well as resolving differences among these colonies (which did not occur for colonies in the remaining areas). In addition, the fineRADstructure results are consistent with the PCA, showing that individuals within the three populations cluster together (Fig 2B), as shown by the dendrogram above the inter-individual co-ancestry matrix. Within Nahuel Huapi Lake, we found groups of individuals that cluster together despite being sampled at different locations. While the average relatedness among NHL individuals is 0.01 (and, for comparison, 0.06 and 0.083 in AC and PC, respectively), the smaller clusters average 0.07 and 0.28, with pairs of individuals reaching relatedness values of up to 0.72 (Fig 2B). This suggests we sampled related individuals in these smaller breeding populations. The fineRADstructure and particularly the Admixture analysis (Fig 2C), show admixture among all three regions, with a lesser degree of Nahuel Huapi Lake ancestry in the Pacific coast. We obtained a similar result using Structure, and for both analyses values of K above 3 did not uncover additional population structure (S1 and S2 Fig in S1 File). The fineRADstructure analysis revealed an individual sampled on the Atlantic coast with Atlantic coast and Nahuel Huapi Lake ancestry, that clusters with other freshwater individuals, which is evidence of connectivity among colonies. Our phylogenetic analysis as well as the clustering analyses show that the three populations are similarly differentiated from each other, with populations from marine environments being slightly more closely related to each other than to the Nahuel Huapi Lake population (e.g., Fig 2B and 2D). We note that in agreement with the relatively low levels of differentiation among populations, our phylogenetic analysis produced generally low node support (Fig 2D). Overall, different measures of diversity were similar across the three regions, with the Nahuel Huapi Lake population showing the lowest heterozygosity and nucleotide diversity ( $\pi$ ) (Table 1). Finally, the highest proportion of migrants was estimated from NHL into PC, followed by NHL into AC (Table 2). Gene flow into NHL from either PC or AC was comparatively much lower. Our mtDNA analysis did not provide sufficient resolution to differentiate populations, as we found two COI haplotypes among our samples, one of which was present in both the NHL and PC colonies, while the second was found in the three regions (Fig 2E).



vertical bar broken into different colored genetic clusters showing the proportion of genetic ancestry assigned to each cluster. The orange arrow indicates the individual mentioned in (B). (D) Maximum Likelihood tree showing phylogenetic relationships between individuals from the three regions represented with three different colors as in (B). Node support for each of the main clades was 87 (PC), 61 (AC), 50 (NHL), and 50 (AC, PC). (E) Haplotype network generated from COI mitochondrial sequences.

<https://doi.org/10.1371/journal.pone.0301004.g002>

## Discussion

This is the first genetic study to include freshwater Kelp Gull colonies from Patagonia, and we find that those from Nahuel Huapi Lake are genetically differentiated from their closest coastal Atlantic and Pacific counterparts. Previous studies on Kelp Gulls have found genetic structure both at a South American continental scale [22–24] and among populations along the Argentine coast [16]. Particularly, Lyons et al. [16] found some level of differentiation among the most geographically distant colonies in their study, as expected from a pattern of isolation by distance with gradually diminishing connectivity with distance along the Argentine coast. Our findings also indicate closer affinities among colonies within each specific region, as opposed to the relationships observed between colonies situated in distinct geographic regions. In this context, we initially expected higher connectivity between Nahuel Huapi Lake populations and those from the Pacific coast, given their proximity and the existence of potential ecological corridors across the Andes Mountains, compared to the Atlantic coast. Instead, our results show that coastal colonies are more closely related to each other than to those from freshwater, despite their considerable geographic separation. This generally low level of differentiation between Atlantic and Pacific colonies may be related to the continuous presence of numerous colonies along these coasts, with a gap in the southern region of Chile which may be due to a lack of information resulting from the difficulty of accessing and surveying this area [9]. However, this does not imply a lack of connectivity between coastal colonies and those from Nahuel Huapi Lake. First, our results show admixture between freshwater and coastal populations, primarily between NHL and AC. Moreover, we observed an adult individual sampled in the Isla de Los Pájaros colony (on the Atlantic coast) with a considerable proportion of genetic ancestry from NHL and the remaining ancestry from AC. This bird is likely the offspring of a Nahuel Huapi Lake migrant which bred with an Atlantic coast individual (although more complex scenarios of dispersal are also compatible with this finding). Additionally, during the course of this study, two records of juvenile individuals banded in Nahuel Huapi Lake were recovered in coastal localities, one was found on the Pacific coast (Osorno, Chile) and the other on the Atlantic coast (Rio Negro Province, Argentina), indicating gulls are capable of dispersing among geographic regions, although this is not necessarily indicative of breeding. Concordantly, our estimates of gene flow suggest migration from NHL into both PC and AC. We did not find direct records of individuals captured in the lake that originated from coastal colonies. Taken together, the generally low levels of differentiation observed in our ddRAD data, and the lack of differentiation in mtDNA, suggests a recent history of divergence, with ongoing gene flow among regions.

An evolutionary significant unit refers to a population or group of individuals that share genetic characteristics [1], and can evolve even in metapopulations of species due to factors

**Table 1. Summary statistics for different Kelp Gull (*Larus dominicanus*) populations.** Observed ( $H_{obs}$ ) and expected ( $H_{exp}$ ) heterozygosity, nucleotide diversity ( $\pi$ ), and inbreeding coefficient ( $F_{IS}$ ).

	$H_{obs}$	$H_{exp}$	$\pi$	$F_{IS}$
Nahuel Huapi Lake	0.248	0.233	0.235	-0.025
Pacific coast	0.267	0.252	0.262	-0.005
Atlantic coast	0.272	0.260	0.265	-0.012

<https://doi.org/10.1371/journal.pone.0301004.t001>

**Table 2. Estimates of gene flow among populations.** Donating populations are shown in the column on the left while receiving populations are in the top row. Values represent the fraction of individuals in the receiving population that are migrants derived from the donating population (per generation). For example, ~30% of individuals from PC are estimated to be derived from NHL in every generation.

	NHL	PC	AC
NHL	0.983	0.297	0.056
PC	0.008	0.683	0.017
AC	0.008	0.015	0.924

<https://doi.org/10.1371/journal.pone.0301004.t002>

such as philopatry [25, 26]. In this regard, our findings imply that the colonies in Lake Nahuel Huapi should be considered an evolutionary significant unit, separate from the coastal colonies. The growth of Kelp Gull populations in many of their coastal colonies is influenced by anthropogenic food subsidies [11]. The available information from colonies situated in Nahuel Huapi Lake shows that Kelp Gulls also use landfill sites as a food source both during and after the breeding season [13, 44]. There is a strong relationship between the areas utilized by seabirds during the winter and their dispersal movements, and this is associated with the existence of population structure [7]. As a consequence, the use of local anthropogenic food sources during the winter might explain the patterns of genetic clustering observed in our study. Furthermore, our results also suggest that the demography of these freshwater colonies is not directly dependent on the movement of individuals from nearby colonies, and thus the likely population growth of Kelp Gulls in Nahuel Huapi Lake is self-sustaining.

Another Patagonian seabird, the Imperial Cormorant (*Leucocarbo atriceps*) also has freshwater populations that are genetically differentiated, both in Lake Yehuin (Tierra del Fuego, Argentina) and in Nahuel Huapi Lake [45]. While the NHL population is likely derived from Pacific coast individuals, those from Lake Yehuin show stronger genetic differentiation [45]. The NHL population shows ecological differences with respect to other marine Imperial Cormorants [46], and those from Lake Yehuin are morphologically differentiated (showing smaller body size and nasal-glands), perhaps as a consequence of local adaptation [47]. It remains to be determined whether the genetically differentiated freshwater population of Kelp Gulls also shows specific adaptations to these environments that differentiate them from those found breeding on the coast.

Understanding the dynamics of Kelp Gull populations, which may function as sources, sinks, or occur in relative isolation from other colonies, has implications for conservation and disease ecology. The Kelp Gull, like other gull species [48–50], is a known carrier of pathogens and parasites in several areas along its distribution, some of which can cause diseases in humans. For instance, Kelp Gulls can be carriers of enterobacteria such as *Salmonella* [51, 52], influenza virus [53, 54], and helminths of the trematoda class (ex *Schistosoma*) that can cause dermatitis in humans. Recent reports have confirmed the presence of cestodes from the genus *Diphyllobothrium* in Kelp Gulls nesting in Nahuel Huapi Lake [55]. The eggs of these cestodes are released in the aquatic environments together with the feces of the definitive host (ichthyophagous birds and mammals, including humans), and can cause *Diphyllobothriasis* [56]. Furthermore, considering that this species can be problematic, negatively affecting other birds with which it usually nests [56], knowledge of the connectivity of its populations and the identification of management units can be fundamental tools when applying management and conservation measures. Our study is the first to identify a freshwater Kelp Gull colony from Patagonia as an evolutionary significant unit. However, to achieve a more complete understanding of the metapopulation dynamics and connectivity of this species as a whole, as well as the origin of freshwater colonies, it is necessary to expand the geographic scope of sampling and include more colonies across the species' distribution.

## Supporting information

### S1 File.

(PDF)

## Acknowledgments

We thank Bronwyn Butcher from the Cornell Lab of Ornithology's Fuller Evolutionary Biology Lab for help with marker discovery and genotyping. We thank the Directorate of Wild Fauna and Flora of the Province of Chubut for administrative advice in the processing of permits, and the Center for the Study of Marine Systems (CONICET) for institutional support. We also thank Juan Cortes, Martin Frixione, Fabiola León, Josefina Jorquera, and Elisa González for their assistance with fieldwork. We thank Lucila Amione and SAG (Servicio Agrícola Ganadero, Chile), for providing the resighting observations of banded individuals. The data from this study are deposited in Dryad (DOI <https://doi.org/10.5061/dryad.dr7sqvb5d>).

## Author Contributions

**Conceptualization:** Tatiana Kasinsky.

**Formal analysis:** Tatiana Kasinsky, Leonardo Campagna.

**Funding acquisition:** Juliana A. Vianna, Pablo Yorio, Leonardo Campagna.

**Resources:** Tatiana Kasinsky, Natalia Rosciano, Juliana A. Vianna, Leonardo Campagna.

**Writing – original draft:** Tatiana Kasinsky.

**Writing – review & editing:** Tatiana Kasinsky, Natalia Rosciano, Juliana A. Vianna, Pablo Yorio, Leonardo Campagna.

## References

1. Moritz C. Applications of mitochondrial DNA analysis in conservation: a critical review. *Mol Ecol*. 1994; 3(4): 401–411.
2. Palsbøll PJ, Bérubé M, Allendorf FW. Identification of management units using population genetic data. *Trends Ecol Evol*. 2006; 22: 11–2. <https://doi.org/10.1016/j.tree.2006.09.003> PMID: 16982114
3. Friesen VL. New roles for molecular genetics in understanding seabird evolution, ecology and conservation. *Mar Ornithol*. 2007; 35: 89–96.
4. Toews DPL, Walsh J, Campagna L. Population genomics of birds: evolutionary history and conservation. In: Hohenlohe PA, Rajora P, editors. *Population Genomics: Wildlife*; 2018. pp. 255–294.
5. Campagna L, Toews DP. The genomics of adaptation in birds. *Current Biology*. 2022; 32(20): R1173–R1186. <https://doi.org/10.1016/j.cub.2022.07.076> PMID: 36283387
6. Oro D. Managing seabird metapopulations in the Mediterranean: constraints and challenges. *Sci Mar*. 2003; 67(S2): 13–22.
7. Friesen VL, Burg TM, McCoy KD. Mechanisms of population differentiation in seabirds. *Mol Ecol*. 2007; 16(9): 1765–1785. <https://doi.org/10.1111/j.1365-294X.2006.03197.x> PMID: 17444891
8. Burger J, Gochfeld M. Family Laridae (Gulls). In: del Hoyo J, Elliott A, Sargatal J, editors. *Handbook of the Birds of the World*. Vol. 3: Hoatzin to Auks. Barcelona; 1996. pp. 572–623.
9. Yorio P, Branco JO, Lenzi J, Luna-Jorquera G, Zavalaga C. Distribution and trends in Kelp Gull (*Larus dominicanus*) coastal breeding populations in South America. *Waterbirds*. 2016; 39(1): 114–135.
10. Yorio P, Bertellotti M, García-Borboroglu P. Estado poblacional y de conservación de gaviotas que se reproducen en el litoral marítimo argentino. *Hornero*. 2005; 20: 53–74.
11. Lislizer N, García-Borboroglu P, Yorio P. Spatial and temporal variations in Kelp Gull population trends in northern Patagonia, Argentina. *Emu*. 2011; 111: 259–267.

12. Lisnizer L, García-Borboroglu P, Pascual M, Yorio P. Transfer processes drive population dynamics of Kelp Gull colonies in Patagonia: implications for management strategies. *Mar Biol Res.* 2015; 11(7): 738–746.
13. Frixione MG, Casaux R, Villanueva C, Alarcón PAE. A recently established Kelp Gull colony in a freshwater environment supported by an inland refuse dump in Patagonia. *Emu.* 2012; 112(2): 174–178.
14. Roesler CI. Conservación del Macá Tobiano (*Podiceps gallardo*): factores que afectan la viabilidad de sus poblaciones. PhD Thesis, Universidad de Buenos Aires. Facultad de Ciencias Exactas y Naturales. 2016. Available from: [http://hdl.handle.net/20.500.12110/tesis\\_n5958\\_Roesler](http://hdl.handle.net/20.500.12110/tesis_n5958_Roesler).
15. Given AD. Phylogenetics and population genetics of the Australasian silver gull (*Larus novaehollandiae*). PhD Thesis, University of Toronto. 2004. Available from: <https://hdl.handle.net/1807/121318>
16. Lyons AC, Lisnizer N, García-Borboroglu P, Yorio P, Bouzat JL. Genetics and demography of kelp gulls (*Larus dominicanus*) in Patagonia. *Ornitol Neotrop.* 2015; 26: 349–362.
17. Mischler C, Veale A, Van Stijn T, Brauning R, McEwan JC, Maloney R, et al. Population connectivity and traces of mitochondrial introgression in New Zealand black-billed gulls (*Larus bulleri*). *Genes.* 2018; 9(11): 544. <https://doi.org/10.3390/genes9110544> PMID: 30424003
18. Mancilla-Morales MD., Velarde E, Aguilar A, Contreras-Rodríguez A, Ezcurra E, Rosas-Rodríguez JA, et al. Strong philopatry, isolation by distance, and local habitat have promoted genetic structure in Heermann's Gull. *Diversity.* 2022; 14(2): 108.
19. Yauk CL, Quinn JS. Genetic structure among breeding herring gulls (*Larus argentatus*) from the Great Lakes and Eastern Canada. *J Great Lakes Res.* 1999; 25(4): 856–864.
20. Chen JZ, Yauk CL, Hebert C, Hebert PD. Genetic variation in mitochondrial DNA of North American herring gulls, *Larus argentatus*. *J Great Lakes Res.* 2001; 27(2): 199–209.
21. Weseloh DVC. The origins of banded herring gulls recovered in the Great Lakes regions. *J Field Ornithol.* 1994; 55(2): 190–195.
22. Dantas GPDM, Meyer D, Godinho R, Ferrand N, Morgante JS. Genetic variability in mitochondrial and nuclear genes of *Larus dominicanus* (Charadriiformes, Laridae) from the Brazilian coast. *Genet Mol Biol.* 2012; 35: 847–885.
23. Santos AF, Morgante JS, Frere E, Millones A, Sander M, de Abreu Vianna J, et al. Evolutionary history of the Kelp Gull (*Larus dominicanus*) in the Southern Hemisphere supported by multilocus evidence. *J Ornithol.* 2016; 157: 1103–1113.
24. Linhares HH, Frere E, Millions A, Pires de Mendonça Dantas G. Evolutionary history of Kelp Gulls at the South Hemisphere. *J Ornithol.* 2023; 1–9.
25. Rexer-Huber K, Veale AJ, Catry P, Cherel Y, Dutoit L, Foster Y, et al. Genomics detects population structure within and between ocean basins in a circumpolar seabird: the white-chinned petrel. *Mol Ecol.* 2019; 28(20): 4552–4572. <https://doi.org/10.1111/mec.15248> PMID: 31541577
26. Frugone MJ, Cole TL, López ME, Clucas G, Matos-Maraví P, Lois NA, et al. Taxonomy based on limited genomic markers may underestimate species diversity of rockhopper penguins and threaten their conservation. *Divers Distrib.* 2021; 27(11): 2277–2296.
27. Suárez N, Retana V, Yorio P. Spatial patterns in the use of foraging areas and its relationship with prey resources in the threatened Olrog's gull (*Larus atlanticus*). *J Ornithol.* 2012; 153: 861–871.
28. Thrasher DJ, Butcher BG, Campagna L, Webster MS, Lovette IJ. Double-digest RAD sequencing outperforms microsatellite loci at assigning paternity and estimating relatedness: A proof of concept in a highly promiscuous bird. *Mol Ecol Res.* 2018; 18(5): 953–965.
29. Gordon A, Hannon G. Fastx-toolkit. FASTQ/A short-reads preprocessing tools. 2010; 5. [http://hannonlab.cshl.edu/fastx\\_toolkit/](http://hannonlab.cshl.edu/fastx_toolkit/).
30. Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. Stacks: an analysis tool set for population genomics. *Mol Ecol.* 2013; 22(11): 3124–3140. <https://doi.org/10.1111/mec.12354> PMID: 23701397
31. Langmead B & Salzberg S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 2012; 9: 357–359. <https://doi.org/10.1038/nmeth.1923> PMID: 22388286
32. KERR KCR, Lijtmaer DA, Barreira AS, Hebert PD, Tubaro PL. Probing evolutionary patterns in Neotropical birds through DNA barcodes. *PLoS One.* 2009; 4(2): e4379. <https://doi.org/10.1371/journal.pone.0004379> PMID: 19194495
33. Hall T, Biosciences I, Carlsbad CJGBB. BioEdit: an important software for molecular biology. *GERF Bull Biosci.* 2011; 2(1): 60–61.
34. Leigh JW, Bryant D. POPART: Full-feature software for haplotype network construction. *Methods Ecol Evol.* 2015; 6: 1110–1116.

35. Zheng X. SNPRelate: Parallel computing toolset for genome-wide association studies. R package version. 2012; 95.
36. R Development Core Team. R: A language and environment for statistical computing. 2015. In: R Foundation for Statistical Computing, Vienna, Austria, Available at. <http://www.R-project.org>.
37. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multi-locus genotype data. *Genetics*. 2000; 155: 945–959. <https://doi.org/10.1093/genetics/155.2.945> PMID: 10835412
38. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res*. 2009; 19: 1655–1664, <https://doi.org/10.1101/gr.094052.109> PMID: 19648217
39. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format and VCFtools. *Bioinformatics*. 2011; 27: 2156–2158. <https://doi.org/10.1093/bioinformatics/btr330> PMID: 21653522
40. Malinsky M, Trucchi E, Lawson DJ, Falush D. RADpainter and fine RADstructure: population inference from RADseq data. *Mol Biol Evol*. 2018; 35(5): 1284–90. <https://doi.org/10.1093/molbev/msy023> PMID: 29474601
41. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 2014; 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033> PMID: 24451623
42. Mussmann M, Douglas MR, Chafin TK, Douglas ME. BA3-SNPs: Contemporary migration reconfigured in BayesAss for next-generation sequence data. *Methods Ecol Evol*. 2019; 10(10): 1808–1813.
43. Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Syst Biol*. 2018; 67(5): 901–904. <https://doi.org/10.1093/sysbio/syy032> PMID: 29718447
44. Frixione MG, Alarcón PAE. Composición de la dieta post-reproductiva de la Gaviota Cocinera (*Larus dominicanus*) en el Lago Nahuel Huapi, Patagonia Argentina. *Ornitol Neotrop*. 2016; 27: 217–222.
45. Calderón L, Quintana F, Cabanne GS, Lougheed SC, Tubaro PL. Phylogeography and genetic structure of two Patagonian shag species (Aves: Phalacrocoracidae). *Mol Phylogen Evol*. 2014; 72: 42–53. <https://doi.org/10.1016/j.ympev.2013.12.011> PMID: 24418531
46. Casaux R, Bertolin ML, Tartara MA, Alarcón P, Porro G. The unexpected diet of breeding imperial shags (*Phalacrocorax atriceps*) at the Nahuel Huapi Lake, Patagonia: implications on population trends *Ornitol Neotrop*. 2010; 21: 457–462
47. Rasmussen PC. Geographic variation in morphology and allozymes of South American imperial shags. *Auk*. 1994; 111: 143–161
48. Savić V, Labrović A, Zelenika TA, Balenović M, Šeparović S, Jurinović L. Multiple introduction of Asian H5N1 avian influenza virus in Croatia by wild birds during 2005–2006 and isolation of the virus from apparently healthy black-headed gulls (*Larus ridibundus*). *Vector Borne Zoonotic Dis*. 2010; 10(9): 915–920. <https://doi.org/10.1089/vbz.2009.0107> PMID: 20370426
49. Wille M, Robertson GJ, Whitney H, Ojkic D, Lang AS. Reassortment of American and Eurasian genes in an influenza A virus isolated from a great black-backed gull (*Larus marinus*), a species demonstrated to move between these regions. *Arch Virol*. 2011; 156: 107–115. <https://doi.org/10.1007/s00705-010-0839-1> PMID: 21053031
50. Tarasiuk K, Kycko A, Knitter M, Świętoń E, Wyrostek K, Domańska-Blicharz K, et al. Pathogenicity of highly pathogenic avian influenza H5N8 subtype for herring gulls (*Larus argentatus*): impact of homo- and heterosubtypic immunity on the outcome of infection. *Vet Res*. 2022; 53(1): 1–11.
51. Frere E, Gandini P, Martinez P. Gaviota Cocinera (*Larus dominicanus*) como vector potencial de patógenos, en la costa patagónica. *Hornero*. 2000; 15(2): 93–97.
52. Barrera V, Retamal P. Detección de *Salmonella* enterica, desde aves silvestres acuáticas en Chile. *Avances en Ciencias Veterinarias*. 2012; 27(2).
53. Pereda AJ, Uhart M, Perez AA, Zaccagnini ME, La Sala L, Decarre J, et al. Avian influenza virus isolated in wild waterfowl in Argentina: evidence of a potentially unique phylogenetic lineage in South America. *Virology*. 2008; 378(2): 363–370. <https://doi.org/10.1016/j.virol.2008.06.010> PMID: 18632129
54. Ghersi BM, Blazes DL, Icochea E, Gonzalez RI, Kochel T, Tinoco Y, et al. Avian influenza in wild birds, central coast of Peru. *Emerg Infect Dis*. 2009; 15(6): 935. <https://doi.org/10.3201/eid1506.080981> PMID: 19523296
55. Casalins LM. La parasitofauna de *Laurus dominicanus* en el Parque Nacional Nahuel Huapi. PhD. Thesis, Universidad Nacional del Comahue. 2023. Available from: <http://rdi.uncoma.edu.ar/handle/uncoma/17095>.
56. Casalins L, Arbetman M, Semenás L, Velezán AA, Flores VR, Viozzi GP. Difilobotrosis en gaviota. Pasado y presente de esta zoonosis en Parque Nacional Nahuel Huapi. *Revista Argentina de Zoonosis y Enfermedades Infecciosas Emergentes*. 2015; 1(10): 38–40.