



OPEN

DATA DESCRIPTOR

The Caribbean and Mesoamerica Biogeochemical Isotope Overview (CAMBIO)

Claire E. Ebert¹✉, Sean W. Hixon^{2,3}, Gina M. Buckley⁴, Richard J. George⁵, Sofía I. Pacheco-Fores⁶, Juan Manuel Palomo⁷, Ashley E. Sharpe⁸, Óscar R. Solís-Torres^{2,3,9}, J. Britt Davis¹⁰, Ricardo Fernandes^{3,11,12,13} & Douglas J. Kennett⁵✉

The Caribbean & Mesoamerica Biogeochemical Isotope Overview (CAMBIO) is an archaeological data community designed to integrate published biogeochemical data from the Caribbean, Mesoamerica, and southern Central America to address questions about dynamic interactions among humans, animals, and the environment in the region over the past 10,000 years. Here we present the CAMBIO human dataset, which consists of more than 16,000 isotopic measurements from human skeletal tissue samples ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$, $\delta^{18}\text{O}$, $^{87}\text{Sr}/^{86}\text{Sr}$, $^{206}/^{204}\text{Pb}$, $^{207}/^{204}\text{Pb}$, $^{208}/^{204}\text{Pb}$, $^{207}/^{206}\text{Pb}$) from 290 archaeological sites dating between 7000 BC to modern times. The open-access dataset also includes detailed chronological, contextual, and laboratory/sample preparation information for each measurement. The collated data are deposited on the open-access CAMBIO data community via the Pandora Initiative data platform (<https://pandoradata.earth/organization/cambio>).

Background & Summary

The Caribbean, Mesoamerica, and southern Central America are extremely diverse in their geology, ecology, and climate. Archaeological research has nevertheless demonstrated millennia of inter-regional interaction resulting in shared cultural trajectories and the development of common economic, social, and political practices^{1–11}. Over the past five decades, isotopic reconstructions of past human diets and mobility in these regions have increased dramatically, providing key insights into the origins of maize agriculture and its intensification, the rise of social complexity and urbanism, and the impacts of European colonization^{8,12–20}.

Despite decades of extensive biogeochemical research in Caribbean, Mesoamerican, and Central American archaeology^{13,15,17,20}, comparative regional and diachronic syntheses of human diet and mobility have been limited by several factors. First, preservation of skeletal remains is variable in the humid, tropical environments of the region. Laboratory observations estimate that between 30–50% of human skeletal samples are too poorly preserved for many types of biogeochemical analysis (e.g., collagen extraction)^{21–23}. Second, sampling bias has resulted in uneven geographic representation throughout the study area (Fig. 1). The highest coverage of isotopic research is in the southern Maya lowlands of Mexico, Guatemala, Belize, and western Honduras^{14,18,24}. Substantial research has also focused on several large (100+ individuals), well-studied skeletal assemblages

¹Department of Anthropology, University of Pittsburgh, 3302 WPH, Pittsburgh, PA, 15260, USA. ²Department of Integrative Biology, Oregon State University, 4575 SW Research Way, Corvallis, OR, 97331, USA. ³Max Planck Institute of Geoanthropology, Kahlaische Strasse 10, D-07745, Jena, Germany. ⁴Interdisciplinary Center for Archaeology and the Evolution of Human Behaviour (ICArEHB), Faculdade das Ciências Humanas e Sociais, Universidade do Algarve, Campus de Gambelas, 8005-139, Faro, Portugal. ⁵Department of Anthropology, University of California, Santa Barbara, Santa Barbara, CA, 93106, USA. ⁶Anthropology Department, Hamline University, 1536 Hewitt Avenue, Saint Paul, MN, 55104, USA. ⁷Department of Anthropology, University of Arizona, 1009 E South Campus Dr, Tucson, AZ, 85721, USA. ⁸Center for Tropical Paleocology and Archaeology, Smithsonian Tropical Research Institute, Luis Clement Avenue, Bldg. 401 Tupper, Ancon, Panama, Republic of Panama. ⁹Instituto Nacional de Antropología e Historia (INAH), Moneda 16, Col. Centro, Alcaldía Cuauhtémoc, 06060, Ciudad de México, México. ¹⁰School of Human Evolution and Social Change, Arizona State University, 900 S. Cady Mall, Tempe, AZ, 85281, USA. ¹¹Department of Bioarchaeology, Faculty of Archaeology, University of Warsaw, Krakowskie Przedmieście 26/28, 00-927, Warsaw, Poland. ¹²Climate Change and History Research Initiative, Princeton University, 129 Dickinson Hall, Princeton, NJ, 08544-1017, USA. ¹³Arne Faculty of Arts, Masaryk University, Nováka 1, 602 00, Brno, Czech Republic. ✉e-mail: c.ebert@pitt.edu; kennett@anth.ucsb.edu




Fig. 1 Spatial distributions of human site locations compiled in CAMBIO. Heatmap visualization shows sample size densities of isotopic measurements (>20 measurements). Sites and geographic regions with >200 measurements are labelled. Base map images are the intellectual property of Esri and are used herein under license. Copyright 2023 Esri and its licensors. All rights reserved.

from the Caribbean islands of Cuba, Puerto Rico, and Guadeloupe^{13,23,25–27}, and in central Mexico at the city of Teotihuacan^{28–33}. Third, a lack of standardization has created inconsistencies in reporting and publication of isotopic datasets. Early studies frequently presented only average isotope values for multiple individuals and infrequently reported sample quality control parameters (e.g., collagen yields and/or atomic C:N ratios^{34,35}). Other inconsistencies are related to reporting (or lack thereof) of chronological information and age/sex estimates. For example, scholars have applied different methods for reporting age estimates that are not easily comparable¹⁵. In other instances, some studies include previously published data without citing the original publication or reporting which variables come from these sources. Finally, and perhaps most importantly, large-scale synthetic isotopic studies from the Caribbean, Mesoamerica, and southern Central America have been prohibited by a lack of open access to published datasets. Most archaeological biogeochemical studies are published in English-language journals and are often inaccessible to our colleagues and communities based in Latin America and the Caribbean due to pay-wall restrictions. Limited data access highlights the academic exclusivity of researchers at universities in the United States, Canada, and Europe^{36,37}. The recent availability of multiple large biogeochemical datasets from around the globe demonstrates that isotopic analysis is a powerful tool for understanding aspects of past human lifeways including diet, nutrition, and population movement^{38–41}.

The Caribbean & Mesoamerica Biogeochemical Isotope Overview (CAMBIO) is a collaborative effort led by early career researchers in archaeology based in the US, Latin America, and Europe to systematically compile published biogeochemical data from the Caribbean, Mesoamerica, and southern Central America in an open-access and multilingual format (Spanish, French, and English). Importantly, our efforts focus not only on collecting isotopic datasets, but also the inclusion of provenance and chronological information to accompany datasets, facilitating multiple archaeological, bioarchaeological, and paleoenvironmental applications. The total number of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$, $\delta^{18}\text{O}$, $^{87}\text{Sr}/^{86}\text{Sr}$, and Pb measurements included is currently 16,512 from at least 5,353 individuals recovered from 290 archaeological sites. The temporal coverage of the dataset represents a broad time span, from the Archaic (~7000 BC) to the Colonial/Historic period (~AD 1500–1800).

Most recorded data (~81%) are stable isotope measurements ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$, $\delta^{18}\text{O}$) from bone and dentine collagen (~41.6%), bone bioapatite (~13.1%), enamel bioapatite (22.4%), and bone and enamel phosphate (~3.7%) (Table 1). The remainder of the data consists of radiogenic isotopic measurements, mostly $^{87}\text{Sr}/^{86}\text{Sr}$ measurements from enamel and bone bioapatite (~16.6%). Approximately 1.5% of the dataset includes Pb measurements, reflecting the exploratory nature of Pb analysis to track the origin and movement of human populations in the CAMBIO region⁴². For both $^{87}\text{Sr}/^{86}\text{Sr}$ and Pb, a local or non-local origin was designated based on reporting by the original authors.

Methods

Data collection for the CAMBIO human dataset began in May 2021, focusing on three major geographic regions: 1) the Caribbean (islands of the Bahamian Archipelago and the Greater and Lesser Antilles), 2) Mesoamerica (modern countries of Mexico, Guatemala, Belize, Honduras, and El Salvador), and 3) southern Central America (modern countries of Nicaragua, Costa Rica, and Panama). A very small number of measurements were also included from northern South America (coastal Venezuela) based on reported cultural

Analyzed Component	Stable Isotopes				Radiogenic Isotopes					Total Measurements	Total Database %
	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	$\delta^{34}\text{S}$	$\delta^{18}\text{O}$	$^{87}\text{Sr}/^{86}\text{Sr}$	$^{206}/^{204}\text{Pb}$	$^{207}/^{204}\text{Pb}$	$^{208}/^{204}\text{Pb}$	$^{207}/^{206}\text{Pb}$		
Bone collagen	3165	3067	138	0	0	0	0	0	0	6370	38.6%
Dentine collagen	260	228	0	0	0	0	0	0	0	488	3.0%
Dentine carbonate	0	0	0	0	5	0	0	0	0	5	0.0%
Bone bioapatite	1,391	0	0	771	288	0	0	0	0	2450	14.8%
Enamel bioapatite	1492	0	0	2208	2450	117	117	117	88	6589	39.9%
Bone phosphate	0	0	0	346	0	0	0	0	0	346	2.1%
Enamel phosphate	0	0	0	264	0	0	0	0	0	264	1.6%
Total Measurements	6308	3295	138	3589	2743	117	117	117	88	16512	100.0%
Total Database %	38.2%	20.0%	0.8%	21.7%	16.6%	0.7%	0.7%	0.7%	0.5%	100.0%	

Table 1. Number of measurements in CAMBIO human database listed by skeletal tissue (“Analyzed Component”) and isotopic measurement type.

affiliation with pre-contact Caribbean groups. The dataset was compiled from published sources including journal articles, books and book chapters, conference proceedings, publicly available academic theses, and archaeological reports in both English and Spanish. Studies were obtained through scientific search engines (e.g., Web of Science, Google Scholar) and online library searches. In cases where datasets could not be obtained online, we contacted the original authors who generously shared their publications. Resources reporting human stable isotope data as of March 2024 are included in the inaugural version of CAMBIO. The dataset in its current form, however, is not completely exhaustive, and will be regularly updated following publication of new studies or the location of previously published studies not yet integrated into the dataset.

The structure of the CAMBIO human dataset is depicted in Figure 2 and described in detail in the dataset’s accompanying metadata file. The CAMBIO dataset is organized according to a series of nested descriptive fields. Each entry has a unique numerical Entry ID. Entries are listed by skeletal tissue type (“Analyzed component”): bone and dentine collagen; dentine carbonate; bone and enamel bioapatite; and bone and enamel phosphate. Isotopic measurements ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$, $\delta^{18}\text{O}$, $^{87}\text{Sr}/^{86}\text{Sr}$, $^{206}/^{204}\text{Pb}$, $^{207}/^{204}\text{Pb}$, $^{208}/^{204}\text{Pb}$, $^{207}/^{206}\text{Pb}$) are then reported based on these tissue types. A single individual or burial, therefore, may have multiple entries in CAMBIO if multiple tissues were analyzed (e.g., separate entries from bone collagen and bone bioapatite isotope values). They may also have multiple entries if different elements were sampled (e.g., multiple bones and/or teeth from the same individual), or if samples were incremental (e.g., tooth increments) or duplicated. This data structure was chosen to more easily address research questions that can be evaluated by specific skeletal tissue types or isotopic systems. The approach also facilitates identification of measurements from different elements that might signal changes to diet or movement throughout a single individual’s life. Entries are tied together by burial identification numbers, if assigned by the original investigators, or sample numbers when burial numbers were not reported. Variations of burial/sample numbers were standardized so that entries for the same individual could be correlated, and to eliminate duplicate entries with identical elemental data.

Burial and site information in CAMBIO comprise both spatial and descriptive attributes. Site locations for each entry are reported as decimal degrees (“Latitude”; “Longitude”) relative to the WGS84 spatial reference system. Coordinates are designated as either “reported” or “estimated”. Reported coordinates are those listed in the original publication. When not available, coordinates were estimated by either locating sites or georeferencing maps from the original publications in Google Earth. Spatial coordinates for Maya lowland sites were also derived from the Electronic Atlas of Ancient Maya Sites⁴³. Local regions (e.g., Maya Lowlands, Greater Antilles) and cultural affiliations (e.g., Mixtec, Aztec, Lucayan Taino) are also included for each entry when described in the source publication. Other burial information recorded in CAMBIO includes provenience data including a description of the site, structure, excavation unit, level, and/or other information about the location of the burial. A description of pathology was also included if reported in the source publication along with isotopic data. Because researchers use several different methods for age estimation to answer different research questions^{15,44}, age estimates are listed as those reported in the source publication, with minimum and maximum age in years, if reported. A second age category (“Age 2”) with adult (18+ yrs.) and sub-adult (0–18 yrs.) designations was also assigned to facilitate data comparisons between broader age cohorts.

The CAMBIO database also includes laboratory and sample preparation information for each measurement. This includes the name of the lab where samples were processed and/or measured, a description of pretreatment protocols, and mass spectrometry instrumentation information. Isotopists are continuously investigating the impacts of different pretreatment protocols on isotope values⁴⁵. For collagen extraction, laboratories use various chemical protocols (e.g., NaOH rinse) and mechanical techniques (e.g., ultrafiltration, or the modified version of the Longin method⁴⁶) to remove exogenous contaminants, with differing impacts on collagen yields and quality^{47–51}. More recently, amino acids have been extracted and purified using XAD resin column chromatography^{52,53}. This has greatly improved the success rate in obtaining carbon and nitrogen isotopic data from degraded bone samples^{21,52,54,55}. Potential complications also exist in the removal of organics from bone and enamel bioapatite before carbonate and phosphate analysis^{56–62}. Following best practice recommendations for reporting in archaeological isotopic studies⁶³, CAMBIO lists a short description of each sample pretreatment protocol and citation for pretreatment protocols when reported in the original publication.

Temporal assignments (“Min date”; “Max date” in years BC/AD) for each measurement were based on radiocarbon dates when available, but were primarily assigned using contextual, epigraphic, and historic documents




Fig. 2 CAMBIO human database structure.




Fig. 3 Summed probability distribution of all human individuals ($n = 5,353$) in the CAMBIO dataset represented by major geographic region. AMS ^{14}C dates were calibrated using methods described in text. Contextually dated samples were modelled as a uniform distribution based on minimum and maximum assigned chronological values. Graph was produced using OxCal v. 4.4.3. and edited in Adobe Illustrator v27.9.

as listed in the original publications. Relative time periods and site-specific archaeological phase designations are also listed when available. All reported radiocarbon measurements in the database are listed by “ ^{14}C date lab number” (if reported) along with the conventional ^{14}C date (in cal yr BP) and error. Radiocarbon dates listed in CAMBIO were calibrated in the calibration software OxCal v.4.4⁶⁴ primarily using the IntCal20 calibration curve⁶⁵. Temporal assignments, however, followed calibration procedures reported in the original study. For example, a mixed IntCal20/Marine20⁶⁶ or local freshwater or marine reservoir correction (ΔR) was used if reported, and is listed in the “Curve used for Calibration” and “ ^{14}C reservoir effect” categories in the dataset, respectively. We report all calibrated ^{14}C dates at the 2-sigma range. Figure 3 shows an estimate for the chronological distribution of the individuals with isotopic measurements based on both ^{14}C and relative temporal assignments (i.e., a summed probability distribution). Samples primarily fall between ~AD 100–1000, reflecting a bias in isotopic data collection, which often focuses on the Classic/Epiclassic urban and elite contexts. The dataset includes a small number of modern measurements from forensic case studies to further contextualize archaeological isotopic values.

We also collected information about the accessibility of each study in the dataset. This information is useful for researchers working outside of the United States, Canada, and Europe or those unaffiliated with an academic institution attempting to locate source publications. Accessibility is listed under the “Open Access” category. We also listed the fees associated with accessing the study if it is not open access. Approximately 59% of studies in the CAMBIO human dataset are behind a paywall with an average of \$37.45 USD per publication for unaffiliated researchers. A list of funding sources reported in the original publication is included. Most publications (~82%) in the CAMBIO human database received partial or complete financial support from federal (US), state, and/or other national sources (e.g., National Science Foundation (USA), Consejo Nacional de Ciencia y Tecnología (Mexico), Social Sciences and Humanities Research Council of Canada).

Data Records

The complete CAMBIO human isotopic dataset is available via a single table in.xlsx format ('cambio_humans_v.1.xlsx') through the CAMBIO data community within the Pandora Initiative data platform (<https://doi.org/10.48493/6c2a-r758>)⁶⁷. Metadata descriptions and the current dataset bibliography are provided on separate sheets within the file.

Technical Validation

CAMBIO includes information about quality control parameters in bioarchaeological isotopic studies when available in the source publication. For bone and dentine, these metrics can be used to assess collagen preservation ("‰C"; "‰N"; "Atomic C:N ratio"; "Atomic C:S ratio"; "Atomic N:S ratio"). Measured values that fall outside the recommended ranges for collagen^{34,68} are included in the database since they are useful for assessing patterns of sample preservation. Other quality control parameters for bone and enamel apatite can be used to document diagenesis and possibly chemical alteration. For ⁸⁷Sr/⁸⁶Sr, this includes Sr ppm, crystallinity index ("CI") values, Ca/P ratios, U/Ca ratios, and Nd/Ca ratios^{69–72}. Studies that lack quality control criteria, or that employ different reference values, are included in CAMBIO and can be filtered by researchers prior to analyses, if desired.

CAMBIO differentiates between $\delta^{18}\text{O}$ measurements from phosphate and carbonates based on the "analyzed component" (bone/enamel phosphate vs bone/enamel bioapatite). Phosphate $\delta^{18}\text{O}$ values are expressed relative to the VSMOW standard⁷³. Most bioapatite carbonate $\delta^{18}\text{O}$ measurements are expressed relative to VPDB in the source publication. Therefore, bioapatite carbonate values originally reported relative to VSMOW were converted to VPDB to standardize the dataset^{74–76}. Computational details are listed in the "Notes" section of the database. Some studies also applied conversions of bioapatite carbonate VPDB values to compare with drinking water VSMOW baseline values. In these instances, if VPDB values were not reported, our conversions may introduce some error by passing observations through equations multiple times^{77,78}. In cases of conversion, we refer researchers to the original publications before conducting additional analyses.

Usage Notes

The CAMBIO human isotopic data compilation combines isotopic data informative of diet and place of origin with chronological, bioarchaeological, and archaeological and historical information. This dataset provides facilitated investigations of significant archaeological questions about human behavior and cultural developments in the Caribbean, Mesoamerica, and southern Central America over the past 10,000 years. Potential use includes inter-regional, regional, and site-specific syntheses of paleodietary isotopic data ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$) to assess broad patterns of human resource acquisition and subsistence change. For example, the broad spatial scope of the database facilitates studies about the timing for the spread of domesticates like maize (*Zea mays*) from Mesoamerica to the Caribbean and North and South America. Analyses of other isotopic systems ($\delta^{18}\text{C}$, ⁸⁷Sr/⁸⁶Sr, Pb) can also be used to document large-scale population movement. Potential questions of interests include immigration to Classic period cities, including Teotihuacan and lowland Maya kingdoms, and the forced movement of indigenous populations and enslaved Africans during the colonial era.

Code availability

No custom code was used to generate or process the data.

Received: 4 November 2023; Accepted: 20 March 2024;

Published online: 08 April 2024

References

1. Cooke, R. Prehistory of Native Americans on the Central American Land Bridge: Colonization, Dispersal, and Divergence. *J. Archaeol. Res.* **13**, 129–187 (2005).
2. Englehardt, J. D. & Carrasco, M. D. *Interregional Interaction in Ancient Mesoamerica*. (University Press of Colorado, 2019).
3. Hofman, C. L. *et al.* Island networks: Transformations of inter-community social relationships in the Lesser Antilles at the advent of European colonialism. *J. Isl. Coast. Archaeol.* **16**, 290–316 (2021).
4. Keegan, W. F., Hofman, C. L. & Rodríguez Ramos, R. *The Oxford Handbook of Caribbean Archaeology*. (Oxford University Press, 2013).
5. Kennett, D. J. *et al.* South-to-north migration preceded the advent of intensive farming in the Maya region. *Nat. Commun.* **13**, 1530 (2022).
6. Kistler, L. *et al.* Archaeological Central American maize genomes suggest ancient gene flow from South America. *Proc. Natl. Acad. Sci.* **117**, 33124–33129 (2020).
7. McCafferty, G., Amador, F. E., Salgado Gonzalez, S. & Dennett, C. Archaeology on Mesoamerica's Southern Frontier. in *The Oxford Handbook of Mesoamerican Archaeology* (eds. Nichols, D. L. & Pool, C. A.) 83–105 (Oxford University Press, 2012).
8. Reid, B. A. *The Archaeology of Caribbean and Circum-Caribbean Farmers (6000 BC - AD 1500)*. (Routledge, 2018).
9. Ranere, A. J. & Cooke, R. G. Late glacial and Early Holocene migrations, and Middle Holocene settlement on the lower isthmian land-bridge. *Quat. Int.* **578**, 20–34 (2021).
10. Robinson, E. & Davies, G. *Routes, Interaction and Exchange in the Southern Maya Area*. (Routledge, 2023).
11. Rodríguez Ramos, R. Isthmo-Antillean Engagements. in *The Oxford Handbook of Caribbean Archaeology* (eds. Keegan, W. F., Hofman, C. L. & Rodríguez Ramos, R.) 155–170 (Oxford University Press, 2013).
12. Laffoon, J. Paleomobility Research in Caribbean Contexts: New Perspectives from Isotope Analysis. in *The Oxford Handbook of Caribbean Archaeology* (eds. Keegan, W. F., Hofman, C. L. & Rodríguez Ramos, R.) 418–435 (Oxford University Press, 2013).
13. Pestle, W. J. Stable Isotopes Analysis of Paleodiet in the Caribbean. in *The Oxford Handbook of Caribbean Archaeology* (eds. Keegan, W. F., Hofman, C. L. & Rodríguez Ramos, R.) 407–417 (Oxford University Press, 2013).
14. Price, T. D. *et al.* Strontium Isotopes and the Study of Human Mobility in Ancient Mesoamerica. *Lat. Am. Antiq.* **19**, 167–180 (2008).
15. Spence, M. W. & White, C. D. Mesoamerican Bioarchaeology: Past and Future. *Anc. Mesoam.* **20**, 233–240 (2009).
16. Scherer, A. K. Bioarchaeology and the Skeletons of the Pre-Columbian Maya. *J. Archaeol. Res.* **25**, 133–184 (2017).
17. Tiesler, V. *The Routledge Handbook of Mesoamerican Bioarchaeology*. (Routledge, 2022).
18. Tykot, R. H. Contribution of Stable Isotope Analysis to Understanding Dietary Variation among the Maya. in *Archaeological Chemistry: Materials, Methods, and Meaning* (ed. Jakes, K. A.) 214–230 (American Chemical Society, 2002).

19. Sharpe, A. E. *et al.* A preliminary multi-isotope assessment of human mobility and diet in pre-Columbian Panama. *J. Archaeol. Sci. Rep.* **36**, 102876 (2021).
20. Willermet, C. & Cucina, A. *Bioarchaeology of Pre-Columbian Mesoamerica: And Interdisciplinary Approach*. (University Press of Florida, 2018).
21. Ebert, C. E., Hoggarth, J. A., Awe, J. J., Culleton, B. J. & Kennett, D. J. The Role of Diet in Resilience and Vulnerability to Climate Change among Early Agricultural Communities in the Maya Lowlands. *Curr. Anthropol.* **60**, 589–601 (2019).
22. Freiwald, C. Excavation and Curation Strategies for Complex Burials in Tropical Environments. *Adv. Archaeol. Pract.* **7**, 10–22 (2019).
23. Pestle, W. J. & Colvard, M. Bone collagen preservation in the tropics: a case study from ancient Puerto Rico. *J. Archaeol. Sci.* **39**, 2079–2090 (2012).
24. Izzo, V. S. R., Wright, L. E. & Canterbury, A. Geographic Variation in Mesoamerican Paleodiets: A Review of Recent Stable Carbon and Nitrogen Isotopic Analyses. in *The Routledge Handbook of Mesoamerican Bioarchaeology* (ed. Tiesler, V.) 441–465 (Routledge, 2022).
25. Pestle, W. J. Fishing Down a Prehistoric Caribbean Marine Food Web: Isotopic Evidence from Punta Candeler, Puerto Rico. *J. Isl. Coast. Archaeol.* **8**, 228–254 (2013).
26. Laffoon, J. E., Hoogland, M. L. P., Davies, G. R. & Hofman, C. L. A Multi-Isotope Investigation of Human and Dog Mobility and Diet in the Pre-Colonial Antilles. *Environ. Archaeol.* **24**, 132–148 (2019).
27. Laffoon, J. E. *et al.* Diverse and Dynamic Dietary Patterns in Early Colonial Cuba: New Insights from Multiple Isotope Analyses. *Lat. Am. Antiq.* **31**, 103–121 (2020).
28. Buckley, G. M. *et al.* New Perspectives on Migration into the Tlajinga District of Teotihuacan: A Dual-Isotope Approach. *Lat. Am. Antiq.* **32**, 536–556 (2021).
29. Casar, I., Morales, P., Manzanilla, L. R., Cienfuegos, E. & Otero, F. Dietary differences in individuals buried in a multiethnic neighborhood in Teotihuacan: stable dental isotopes from Teopancazco. *Archaeol. Anthropol. Sci.* **9**, 99–115 (2017).
30. Nado, K. L., Zolotova, N. & Knudson, K. J. Paleodietary analysis of the sacrificial victims from the Feathered Serpent Pyramid, Teotihuacan. *Archaeol. Anthropol. Sci.* **9**, 117–132 (2017).
31. Price, T. D., Manzanilla, L. & Middleton, W. D. Immigration and the Ancient City of Teotihuacan in Mexico: a Study Using Strontium Isotope Ratios in Human Bone and Teeth. *J. Archaeol. Sci.* **27**, 903–913 (2000).
32. White, C. D., Spence, M. W., Longstaffe, F. J., Stuart-Williams, H. & Law, K. R. Geographic Identities of the Sacrificial Victims from the Feathered Serpent Pyramid, Teotihuacan: Implications for the Nature of State Power. *Lat. Am. Antiq.* **13**, 217–236 (2002).
33. White, C. D., Price, T. D. & Longstaffe, F. J. Residential Histories of the Human Sacrifices at the Moon Pyramid, Teotihuacan: Evidence from Oxygen and Strontium Isotopes. *Anc. Mesoam.* **18**, 159–172 (2007).
34. DeNiro, M. J. Postmortem preservation and alteration of *in vivo* bone collagen isotope ratios in relation to palaeodietary reconstruction. *Nature* **317**, 806–809 (1985).
35. van Klinken, G. J. Bone Collagen Quality Indicators for Palaeodietary and Radiocarbon Measurements. *J. Archaeol. Sci.* **26**, 687–695 (1999).
36. Marwick, B. *et al.* Open Science in Archaeology. *SAA Archaeol. Rec.* **17**, 8–14 (2017).
37. McKiernan, E. C. *et al.* How open science helps researchers succeed. *eLife* **5**, e16800 (2016).
38. Coccozza, C., Cirelli, E., Groß, M., Teegen, W.-R. & Fernandes, R. Presenting the Compendium Isotoporum Medii Aevi, a Multi-Isotope Database for Medieval Europe. *Sci. Data* **9**, 354 (2022).
39. Fernandes, R. *et al.* The ARCHIPELAGO Archaeological Isotope Database for the Japanese Islands. *J. Open Archaeol. Data*, <https://doi.org/10.5334/joad.73> (2021).
40. Goldstein, S. *et al.* Presenting the AfriArch Isotopic Database. *J. Open Archaeol. Data*, <https://doi.org/10.5334/joad.94> (2022).
41. Plomp, E. *et al.* The IsoArch initiative: Working towards an open and collaborative isotope data culture in bioarchaeology. *Data Brief* **45**, 108595 (2022).
42. Sharpe, A. E. *et al.* Lead (Pb) Isotope Baselines for Studies of Ancient Human Migration and Trade in the Maya Region. *PLOS ONE* **11**, e0164871 (2016).
43. Witschey, W. R. T. & Brown, C. Electronic Atlas of Ancient Maya Sites. (2010).
44. Boldsen, J. L., Milner, G. R. & Ousley, S. D. Paleodemography: From archaeology and skeletal age estimation to life in the past. *Am. J. Biol. Anthropol.* **178**, 115–150 (2022).
45. Pestle, W. J., Crowley, B. E. & Weirauch, M. T. Quantifying Inter-Laboratory Variability in Stable Isotope Analysis of Ancient Skeletal Remains. *PLoS ONE* **9**, e102844 (2014).
46. Longin, R. New Method of Collagen Extraction for Radiocarbon Dating. *Nature* **230**, 241–242 (1971).
47. Guiry, E. J., Szpak, P. & Richards, M. P. Effects of lipid extraction and ultrafiltration on stable carbon and nitrogen isotopic compositions of fish bone collagen. *Rapid Commun. Mass Spectrom.* **30**, 1591–1600 (2016).
48. Higham, T. F. G., Jacobi, R. M. & Ramsey, C. B. AMS Radiocarbon Dating of Ancient Bone Using Ultrafiltration. *Radiocarbon* **48**, 179–195 (2006).
49. Sealy, J., Johnson, M., Richards, M. & Nehlich, O. Comparison of two methods of extracting bone collagen for stable carbon and nitrogen isotope analysis: comparing whole bone demineralization with gelatinization and ultrafiltration. *J. Archaeol. Sci.* **47**, 64–69 (2014).
50. Szpak, P., Krippner, K. & Richards, M. P. Effects of Sodium Hydroxide Treatment and Ultrafiltration on the Removal of Humic Contaminants from Archaeological Bone. *Int. J. Osteoarchaeol.* **27**, 1070–1077 (2017).
51. Talamo, S., Fewlass, H., Maria, R. & Jaouen, K. "Here we go again": the inspection of collagen extraction protocols for ¹⁴C dating and palaeodietary analysis. *STAR. Sci. Technol. Archaeol. Res.* **7**, 62–77 (2021).
52. Kennett, D. J. *et al.* Early isotopic evidence for maize as a staple grain in the Americas. *Sci. Adv.* **6**, eaba3245 (2020).
53. Stafford, T. W., Hare, P. E., Currie, L., Jull, A. J. T. & Donahue, D. J. Accelerator radiocarbon dating at the molecular level. *J. Archaeol. Sci.* **18**, 35–72 (1991).
54. Hoggarth, J. A. *et al.* Building High-Precision AMS ¹⁴C Bayesian Models for the Formation of Peri-Abandonment Deposits at Baking Pot, Belize. *Radiocarbon* **63**, 977–1002 (2021).
55. Hoggarth, J. A., Culleton, B. J., Awe, J. J. & Kennett, D. J. Questioning Postclassic Continuity at Baking Pot, Belize, Using Direct AMS ¹⁴C Dating of Human Burials. *Radiocarbon* **56**, 1057–1075 (2014).
56. Crowley, B. E. & Wheatley, P. V. To bleach or not to bleach? Comparing treatment methods for isolating biogenic carbonate. *Chem. Geol.* **381**, 234–242 (2014).
57. Freiwald, C. Oxygen Isotope Values in the Maya Region. in *Isotopic Proveniening and Mobility: The Current State of Research* (ed. Price, T. D.) 67–84 (Springer Nature, 2023).
58. Grimes, V. & Pellegrini, M. A comparison of pretreatment methods for the analysis of phosphate oxygen isotope ratios in bioapatite. *Rapid Commun. Mass Spectrom.* **27**, 375–390 (2013).
59. Pederzani, S. & Britton, K. Oxygen isotopes in bioarchaeology: Principles and applications, challenges and opportunities. *Earth-Sci. Rev.* **188**, 77–107 (2019).
60. Pellegrini, M. & Snoeck, C. Comparing bioapatite carbonate pre-treatments for isotopic measurements: Part 2 — Impact on carbon and oxygen isotope compositions. *Chem. Geol.* **420**, 88–96 (2016).

61. Snoeck, C. & Pellegrini, M. Comparing bioapatite carbonate pre-treatments for isotopic measurements: Part 1—Impact on structure and chemical composition. *Chem. Geol.* **417**, 394–403 (2015).
62. Ventresca Miller, A. *et al.* Sampling and Pretreatment of Tooth Enamel Carbonate for Stable Carbon and Oxygen Isotope Analysis. *J. Vis. Exp.* e58002, <https://doi.org/10.3791/58002> (2018).
63. Roberts, P. *et al.* Calling all archaeologists: guidelines for terminology, methodology, data handling, and reporting when undertaking and reviewing stable isotope applications in archaeology. *Rapid Commun. Mass Spectrom.* **32**, 361–372 (2018).
64. Reimer, P. J. *et al.* The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon* **62**, 725–757 (2020).
65. Bronk Ramsey, C. Bayesian Analysis of Radiocarbon Dates. *Radiocarbon* **51**, 337–360 (2009).
66. Heaton, T. J. *et al.* Marine20—The Marine Radiocarbon Age Calibration Curve (0–55,000 cal BP). *Radiocarbon* **62**, 779–820 (2020).
67. Ebert, C. E. *et al.* CAMBIO Human Isotopic Dataset. <https://doi.org/10.48493/6c2a-r758> (2024).
68. Nehlich, O. & Richards, M. P. Establishing collagen quality criteria for sulphur isotope analysis of archaeological bone collagen. *Archaeol. Anthropol. Sci.* **1**, 59–75 (2009).
69. Budd, P., Montgomery, J., Barreiro, B. & Thomas, R. G. Differential diagenesis of strontium in archaeological human dental tissues. *Appl. Geochem.* **15**, 687–694 (2000).
70. Hedges, R. E. M. Bone diagenesis: an overview of processes. *Archaeometry* **44**, 319–328 (2002).
71. Price, T. D., Blitz, J., Burton, J. & Ezzo, J. A. Diagenesis in prehistoric bone: Problems and solutions. *J. Archaeol. Sci.* **19**, 513–529 (1992).
72. Reynard, B., Lécuyer, C. & Grandjean, P. Crystal-chemical controls on rare-earth element concentrations in fossil biogenic apatites and implications for paleoenvironmental reconstructions. *Chem. Geol.* **155**, 233–241 (1999).
73. Chenery, C. A., Pashley, V., Lamb, A. L., Sloane, H. J. & Evans, J. A. The oxygen isotope relationship between the phosphate and structural carbonate fractions of human bioapatite. *Rapid Commun. Mass Spectrom.* **26**, 309–319 (2012).
74. Brand, W. A., Coplen, T. B., Vogl, J., Rosner, M. & Prohaska, T. Assessment of international reference materials for isotope-ratio analysis (IUPAC Technical Report). **86**, 425–467 (2014).
75. Coplen, T. B. Normalization of oxygen and hydrogen isotope data. *Chem. Geol. Isot. Geosci. Sect.* **72**, 293–297 (1988).
76. Coplen, T. B. Guidelines and recommended terms for expression of stable-isotope-ratio and gas-ratio measurement results. *Rapid Commun. Mass Spectrom.* **25**, 2538–2560 (2011).
77. Pellegrini, M., Lee-Thorp, J. A. & Donahue, R. E. Exploring the variation of the $\delta^{18}\text{O}_\text{p}$ and $\delta^{18}\text{O}_\text{c}$ relationship in enamel increments. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **310**, 71–83 (2011).
78. Pollard, A. M., Pellegrini, M. & Lee-Thorp, J. A. Technical note: Some observations on the conversion of dental enamel $\delta^{18}\text{O}_\text{p}$ values to $\delta^{18}\text{O}_\text{v}$ to determine human mobility. *Am. J. Phys. Anthropol.* **145**, 499–504 (2011).

Acknowledgements

The Pandora and IsoMemo initiatives are supported by Max Planck Institute for Geoanthropology. We are grateful to the researchers who have published stable isotope measurements from the Caribbean, Mesoamerica, and Lower Central America, therefore contributing to this dataset. We would also like to thank student volunteers who helped with data compilation (University of Pittsburgh – Lydia Bailey and Megan Myers; Hamline University – Kashama Bhakta, Kathleen Kelley; University of California, Santa Barbara – Louisa Crouthamel, Jonathan Chastain, Rosie Morales). We also thank two anonymous reviewers who provided helpful feedback and pointed out sources missing in the CAMBIO human dataset. Funding to support data collection at the University of California, Santa Barbara provided by the National Science Foundation (DISES-2307175 awarded to D.J.K.). Open access funding provided by the Department of Anthropology, University of Pittsburgh (C.E.E.).

Author contributions

C.E.E., S.W.H., G.M.B., R.J.G., S.I.P.-F., J.M.P., A.E.S., and O.R.S. co-designed meta-data structure. C.E.E., S.W.H., G.M.B., R.J.G., S.I.P.-F., J.M.P., and A.E.S. oversaw data collection of human stable isotope datasets. C.E.E. and S.W.H. designed the study. C.E.E. created the figures, assisted by J.B.D. C.E.E., D.J.K., and R.F. supervised the study. C.E.E. and S.W.H. wrote the paper with contributions from all authors.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to C.E.E. or D.J.K.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024