

SYNTHESIS

The Relationship Between Maturation Size and Maximum Tree Size From Tropical to Boreal Climates

Valentin Journé¹  | Michał Bogdziewicz²  | Benoit Courbaud¹ | Georges Kunstler¹  | Tong Qiu³  | Marie-Claire Aravena Acuña⁴ | Davide Ascoli⁵  | Yves Bergeron⁶ | Daniel Berveiller⁷ | Thomas Boivin⁸ | Raul Bonal⁹ | Thomas Caignard¹⁰ | Maxime Cailleret¹¹ | Rafael Calama¹² | J. Julio Camarero¹³  | Chia-Hao Chang-Yang¹⁴  | Jerome Chave¹⁵ | Francesco Chianucci¹⁶ | Thomas Curt¹⁷ | Andrea Cutini¹⁸ | Adrian Das¹⁹ | Evangelia Daskalakou²⁰ | Hendrik Davi⁸ | Nicolas Delpierre⁷ | Sylvain Delzon¹⁰ | Michael Dietze²¹  | Sergio Donoso Calderon²² | Laurent Dormont²³ | Josep Maria Espelta²⁴ | William Farfan-Rios²⁵ | Michael Fenner²⁶ | Jerry Franklin²⁷ | Catherine Gehring²⁸ | Gregory Gilbert²⁹ | Georg Gratzner³⁰ | Cathryn H. Greenberg³¹ | Arthur Guignabert³²  | Qinfeng Guo³³ | Andrew Hacket-Pain³⁴ | Arndt Hampe³⁵ | Qingmin Han³⁶ | Mick E. Hanley³⁷ | Janneke Hille Ris Lambers³⁸ | Jan Holík³⁹ | Kazuhiko Hoshizaki⁴⁰ | Ines Ibanez⁴¹ | Jill F. Johnstone⁴² | Johannes M. H. Knops⁴³  | Richard K. Kobe⁴⁴ | Hiroko Kurokawa⁴⁵  | Jonathan Lageard⁴⁶ | Jalene LaMontagne⁴⁷  | Mateusz Ledwon⁴⁸ | François Lefèvre⁸ | Theodor Leininger⁴⁹ | Jean-Marc Limousin⁵⁰ | James Lutz⁵¹  | Diana Macias⁵² | Anders Mårell⁵³  | Eliot McIntire⁵⁴  | Emily V. Moran⁵⁵ | Renzo Motta⁵ | Jonathan Myers⁵⁶ | Thomas A. Nagel⁵⁷ | Shoji Naoe⁵⁸  | Mahoko Noguchi⁵⁸ | Julian Norghauer⁵⁹ | Michio Oguro⁴⁵ | Jean-Marc Ourcival⁵⁰ | Robert Parmenter⁶⁰ | Ian Pearse⁶¹ | Ignacio M. Pérez-Ramos⁶² | Łukasz Piechnik⁶³ | Tomasz Podgórski⁶⁴ | John Poulsen⁶⁵ | Miranda D. Redmond⁶⁶ | Chantal D. Reid⁶⁷ | Pavel Samonil³⁹ | C. Lane Scher⁶⁷  | William H. Schlesinger⁶⁷ | Barbara Seget⁶³ | Shubhi Sharma⁶⁸ | Mitsue Shibata⁴⁵ | Miles Silman⁶⁹ | Michael Steele⁷⁰ | Nathan Stephenson¹⁹ | Jacob Straub⁷¹ | Samantha Sutton⁶⁷ | Jennifer J. Swenson⁷² | Margaret Swift⁶⁷ | Peter A. Thomas⁷³ | Maria Uriarte⁷⁴  | Giorgio Vacchiano⁷⁵ | Amy Whipple⁷⁶ | Thomas Whitham⁷⁶ | S. Joseph Wright⁷⁷  | Kai Zhu⁴¹  | Jess Zimmerman⁷⁸ | Magdalena Żywiec⁶³  | James S. Clark^{1,67}

Correspondence: Valentin Journé (journe.valentin@gmail.com) | James S. Clark (jimclark@duke.edu)

Received: 19 September 2023 | **Revised:** 12 August 2024 | **Accepted:** 13 August 2024

Funding: This work was supported by Belmont Forum, (1854976). Agence Nationale de la Recherche, (18-MPGA-0004). Advanced Exploration Systems, (AIST16-0052, AIST18-0063). Directorate for Biological Sciences, (DEB-1754443).

Keywords: allometry | life history | seed production | size | tree fecundity | tree maturation

ABSTRACT

The fundamental trade-off between current and future reproduction has long been considered to result in a tendency for species that can grow large to begin reproduction at a larger size. Due to the prolonged time required to reach maturity, estimates of tree maturation size remain very rare and we lack a global view on the generality and the shape of this trade-off. Using seed production from five continents, we estimate tree maturation sizes for 486 tree species spanning tropical to boreal climates. Results show that a species' maturation size increases with maximum size, but in a non-proportional way: the largest species begin reproduction at smaller sizes than would be expected if maturation were simply proportional to maximum size. Furthermore, the decrease in relative maturation size is steepest in cold climates. These findings on maturation size drivers are key to accurately represent forests' responses to disturbance and climate change.

1 | Introduction

The size or age at maturity is critical for tree population fitness and forest regeneration because recruitment opportunities can

occur when trees are any size or age (Dietze and Clark 2008; McDowell et al. 2020; Qiu et al. 2021). In trees, reproduction follows an extended maturation phase (Thomas 1996; Clark, LaDeau, and Ibanez 2004). Juvenile allocation to leaves and

the roots and architecture that supports them build the large light- and water-harvesting capacity characteristic of the tree life form. Allocation can then shift to include reproduction. This delayed maturation of trees is linked to the fundamental trade-off between current and future reproduction (Stearns 1989).

On one hand, delayed reproduction sacrifices early seed production to reap future benefits. In environments marked by intense competition and a reliable future, delayed reproduction benefits from large size and the resources that accumulate if juveniles allocate to growth and survival (Falster and Westoby 2003; Wenk and Falster 2015). On the other hand, the advantages of large size can come with costs, including water transport high into the crown and biomechanical risk of bole fracture or windthrow (Niklas 1994; Koch et al. 2004; Dietze and Clark 2008; Lines et al. 2012). Current reproduction avoids the risks of an uncertain future, and it contributes most to fitness in non-competitive or frequently disturbed environments (Charlesworth 2000).

Among tree species, the potential trade-off between current and future reproduction might require a balance of benefits and risks and result in a positive association across species in maturation and maximum size or age (Loehle 1988; Thomas 1996; Davies and Ashton 1999; Westoby et al. 2002; Falster and Westoby 2003; Wenk and Falster 2015; Visser et al. 2016). Our understanding of the variation of maturation size among tree species is, however, extremely limited and there are no large-scale studies on this topic. We thus have a poor understanding of how maturation size varies with species maximum size and the relative importance of other factors such as species climate niche and functional traits.

In the absence of maturation estimates, earth system models (ESMs) incorporate assumptions that are expected to bias lifetime reproduction. Many ESMs omit impacts of life history on disturbance response entirely (see McDowell et al. 2020 for a review). In models that do accommodate life history, maturation size— d_{mat} (for diameter)—is independent of maximum size— d_{max} (Kohler and Huth 2004; Wallentin et al. 2008; Yang et al. 2022) (Figure 1, black dotted line). An alternative hypothesis is that maturation size is proportional to maximum size. It is consistent with a study at Barro Colorado Island in Panama; where, Visser et al. (2016) reported that $d_{\text{mat}} = d_{\text{max}} / 2$ (red dashed line in Figure 1). This is related to the classical prediction of a proportionate increase in maturation age with increased maximum age from simple optimisation models in animals (Charnov and Berrigan 1990, 1991; Jensen 1996; Thorson et al. 2017) and trees (Clark 1991). Indeed, if the predictions for age also apply to size, then this *proportionate risk model* means that the maturation delay incurred for increased maximum size is the same for species large and small, represented by the red line in Figure 1.

In fact, it would be remarkable if this relationship was the same for species of all sizes because the constraints on the large size and the relative contribution to fitness of early seed production could vary widely depending on the species' maximum size. If mortality risks and allocation demands change with age and size (Charnov and Berrigan 1990), then the strictly proportional relationship between maturation size (d_{mat}) and maximum size (d_{max}) can be generalised to a power relationship,




FIGURE 1 | Hypothesised association between maturation size (d_{mat}) and maximum size (d_{max}) (a) and the relative size at maturation ($d_{\text{rel}} = d_{\text{mat}} / d_{\text{max}}$) (b) (Equation (1)). To highlight the effects of size (parameter β_d), values of parameter α are selected to yield an equivalent diameter at $d_{\text{max}} = 60$ cm. Two ‘baseline’ hypotheses (dashed lines) are independence between d_{mat} and d_{max} (black dotted) and proportionate delay (red dotted), the latter is expected if increased size incurs the same maturation delay at all size classes. Two alternative hypotheses are increasing (purple—accelerating risk model) or decreasing (blue—diminishing risk model) maturation delays in the largest size classes.

$$d_{\text{mat}} = \alpha \times d_{\text{max}}^{\beta_d} \quad (1)$$

The proportionate model means that $\beta_d = 1$ in Equation (1) (Figure 1a). An *accelerating risk model* refers to the case where the maturation delay required for large species to increase maximum size is greater than for small species. If $\beta_d > 1$ (purple in Figure 1), the relative size at maturation ($d_{\text{rel}} = d_{\text{mat}} / d_{\text{max}}$) continues to increase for species in the largest size classes (Figure 1b). Consider, for example, an expected fitness gain from extending end-of-life reproduction that comes with increased size and age. The early-life investments in structural support or defences needed for an incremental increase in maximum size might be especially high for the largest species.

Alternatively, a *diminishing risk model* refers to the case where the need to further delay maturation size declines for species already at large size. If $0 < \beta_d < 1$, then species that

reach large size do not incur the same proportionate delay risk as small species. As β_d approaches zero, the largest species mature at nearly the same size as the small species. Then, the relative size at maturation d_{rel} declines with maximum size (Figure 1b).

Improved understanding of maturation size confronts challenges posed by slow dynamics and limited observation. Where there is a dense canopy, the reproductive status of individual trees is often not detected, and crop failures (little or no seed production in trees that have reached maturity) are common. Likewise, seed traps often fail to recover seeds from nearby reproductive trees, especially the low seed production of newly mature individuals (LaDeau and Clark 2001). Detection error can be minimised where observations come from above the canopy or open settings like savannas, seed orchards or common gardens (LaDeau and Clark 2001; Caignard et al. 2021). Still, a time series of observations is needed because an individual reproducing this year will have matured at some time in the past. Estimates of maturation status from time series data allow for detection error similar to the way it is used in capture–recapture models. Maturation can be treated as a hidden Markov process (see detailed Supplements to Clark, LaDeau, and Ibanez 2004; Clark, Nuñez, and Tomasek 2019). As in capture–recapture models, the probability that an individual is mature in a given year depends not only on failure to detect in the current year but also on the history and future of observations on the same individual. The more times that reproduction is not detected in the past (or future), the lower the probability that a tree is mature now. In tree fecundity studies, the complexity is compounded by the ‘masting’ phenomenon, where quasi-synchronous, quasi-periodic crops require observations over several years, making a large number of observations in a single year insufficient. Most studies where maturation size has been estimated focus on open-grown trees and/or have limited taxonomic breadth, habitat variation or both (Wenk et al. 2018; Thomas 1996; Davies and Ashton 1999; Kohyama et al. 2003; Wright et al. 2005; Visser et al. 2016; Minor and Kobe 2019).

Inferring the relationship between maturation and maximum size has also to control for the environment (Wenk and Falster 2015) and species characteristics (Visser et al. 2016). While the effects of climate on maturation size are unknown, tree fecundity responds to seasonal temperature and moisture, soils and light availability, which depends on the local competitive environment (Clark et al. 2014; Caignard et al. 2017; Minor and Kobe 2019; Le Roncé et al. 2021; Qiu et al. 2022; Journé et al. 2022). Also, fast growth and accelerated competition that comes from long growing seasons in the wet tropics do not necessarily imply small or large maturation sizes.

The relationship between maturation and maximum size could be associated with other plant functional traits, that would reflect diverse plant strategies. Fast growth in open environments is often associated with low wood density and high specific leaf area (SLA) (Moles et al. 2004, 2006; Thomas, Martin, and Mycroft 2015; Visser et al. 2016; Wenk et al. 2018). However, it is unclear whether the same traits that are involved in fast growth are also associated with maturation at a small size. Furthermore, if large-seeded species need to accumulate resource reserves, then there could be a positive association between seed size and

maturation size (Moles et al. 2004). Relatedly, high reproductive expenditures, measured as seed size $\times$ seed number (Qiu et al. 2022), might be associated with delayed maturation size. Due to their co-dependence, it is necessary to model all of these traits jointly, while accounting for the effects of habitat and phylogenetic groups (Clark 2016; Seyednasrollah and Clark 2020; Bogdziewicz et al. 2023; Qiu et al. 2023).

In this study, we provide the first comprehensive estimates of tree maturation size, obtained for 486 tree species on five continents, incorporating effects of the environment over a large range of tree diameters and habitats. We use the Masting Inference and Forecasting (MASTIF) network and modelling framework to accommodate the dependence between observations between trees and within trees over time (Clark et al. 2021; Sharma et al. 2022; Qiu et al. 2022; Journé et al. 2022). Based on MASTIF estimates we derive maturation size as tree diameter at the onset of female reproductive function allowing us to compare maturation sizes across species that vary in reproductive biology (e.g., *Pinaceae* commonly produce male cones earlier than female cones; many species have no such separation) and where pollen production can be hard to quantify. We first evaluate how maturation varies with species' maximum size and test the three alternative models of Figure 1 with our estimates of maturation size and estimates of species' maximum size. Then, we evaluate how the relationship between maturation size and maximum size is influenced by climate and its association with other plant functional traits.

2 | Materials and Methods

Our analysis includes three elements (Figure 2). We first parameterise a model for individual maturation status and fecundity based on diameter, shade conditions and environmental variables (Figure 2a). The year in which an individual achieves maturity is almost never observed. Instead, seeds counted in traps or in crowns vary from year to year. Successive observations represent a time series for every tree. This first step estimates maturation status and conditional fecundity (seeds per tree per year given that it is mature) for all trees in the network. From this fitted model, we generate predictive distributions of maturation status across diameter with other variables held at intermediate values to estimate d_{mat} from the model. Again, this prediction from the model is necessitated by the fact that true maturation status is an estimate, not a state that is directly observed. Second, we estimated the model of Figure 1 to obtain estimates of α and β_d (Figure 2b), while controlling for other variables that could affect their relationship. Finally, we evaluate the species-level trait relationship that includes maturation size (Figure 2c). The following section describes these elements of the analysis.

2.1 | MASTIF Data and Model

The MASTIF model and data summarised here are detailed in Clark, Nuñez, and Tomasek (2019) and its extended Supplement (see also Qiu et al. 2021, 2022; Journé et al. 2022). Data are of two types, crop counts on trees and seed traps in mapped inventory plots (MASTIF) (Clark, Nuñez, and Tomasek 2019)




FIGURE 2 | Three elements of the analysis include (a) an individual-scale analysis (blue) to estimate maturation status each year and to parameterise relationships that control maturation. This fitted model is the basis for species-level prediction of maturation size (red). (b) Species-level expected maturation size based on the proportionate risk model, controlling for species' differences in their climate domains. (c) Analysis of species-level trait relationships with maturation size.

(Figure S1). The initial sample size is approximately 12 million tree-years from five continents on 898 species and 112 families. The majority of observations (99%) are derived from longitudinal studies, involving repeated observations of all trees on a plot or individual trees. The remaining crop count observations (1%) are collected opportunistically through the iNaturalist project MASTIF (Clark, Nuñez, and Tomasek 2019). The number of species observed per plot ranges from 1 to 221 species. The number of species observations is larger for seed trap monitoring (476 species in total, 22,929 tree-year observations on average) than for crop count monitoring (130 species in total, 1058 tree-year observations on average). Most plots are localised in North America and central Europe (97%), whereas most species observations are coming from South America (54%). On average, 75% of individual tree-year observation are coming from the tropics. Additional information is provided in Table S1 and Supplementary Files S1 and S2. For both data types, observations include species, diameter, shade class (ranging from 'full sun', Class 1, to 'full shade', Class 5), number of fruiting structures, and an estimate of the fraction of the total crop represented by the count. For crop counts, the data model is beta-binomial, with binomial uncertainty for the counts given crop fraction, and beta uncertainty for crop fraction. The seed traps data additionally include mapped locations of trees and seed traps, which is used to jointly estimate fecundity, dispersal, and, for seeds identified only to genus, species identity. For seed traps, the data model is Poisson for counts given dispersal and species, a bivariate Student's t (i.e., 2Dt) redistribution kernel for dispersal (Clark et al. 1999; Clark, Nuñez, and Tomasek 2019), and a multinomial species probability (many seeds are identified only to genus level).

The MASTIF model is a dynamic model for year-to-year and tree-to-tree seed production. The model allows for conditional independence in crop counts and seed traps data through latent states. It estimates maturation state and conditional fecundity (seed production given the individual is mature), which depend on tree size, shading, local climate and soil conditions. Random

effects on individuals and years allow for wide variation between trees and over time. The posterior distribution includes the parameters and latent states presented in Clark, Nuñez, and Tomasek (2019), and summarised in Qiu et al. (2022) and Journé et al. (2022). Model fitting was accomplished with Gibbs sampling, a Markov chain Monte Carlo technique based on sampling from conditional distributions. Model structure and methodology are implemented with the R package Mast Inference and Forecasting (MASTIF, v1.0.1) (Clark, Nuñez, and Tomasek 2019).

2.2 | Derivation of Tree Maturation Size From Fitted MASTIF Model

Tree maturation size (d_{mat}) is derived from an individual-scale model fitted to each species with MASTIF. We define tree maturation size (d_{mat}) as the diameter when a tree is mature and has the capacity to produce enough seed to construct one fruiting structure, f_{min} . For species that produce one-seeded fruits (e.g., *Quercus*, *Juglandaceae*), $f_{min} = 1$. For species that produce cones (e.g., *Pinaceae*, *Cupressaceae*), pods (e.g., *Fabaceae*, *Bignoniaceae*) or other capsules that house multiple seeds (e.g., *Fagus* capsules), f_{min} is the number of seeds contained in that structure. The data and definitions we use to determine d_{mat} differ from those employed in previous studies (e.g., Visser et al. 2016), as we use both crop count and seed trap observation and not only maturation status. The estimation of individual fecundities, obtained through MASTIF model, is also included, taking into account tree characteristics and environment (Clark et al. 2021; Qiu et al. 2021; Journé et al. 2022). MASTIF models the effects of environmental predictors on conditional fecundity (given mature status), because immature trees do not respond to predictors (it is always zero) [Modelling environmental effects on (unconditional) fecundity would make no more sense than including immature individuals in studies of masting intervals or synchronicity]. Conditional fecundity ψ is represented by a log-normal distribution, which allows for the effects of the environment. The log-normal is undefined for zero seeds. Zeros

are accommodated by the fact that trees can be in the immature state ($\rho = 0$), or conditional fecundity can be below the threshold $f_{\min}$, as in a failed seed crop (Clark, LaDeau, and Ibanez 2004; Clark, Nuñez, and Tomasek 2019),

$$f = \begin{cases} \psi & \rho = 1 \\ 0 & (\rho = 0) + (\rho = 1)(\psi < f_{\min}) \end{cases} \quad (2)$$

An individual is immature until the first time fecundity rises above the threshold for producing fruit, that is, $(\rho_{i,t} = 1)(\psi_{i,t} > f_{\min})$. Specifically for tree i in year t ,

$$\begin{aligned} f_{i,t} &= \psi_{i,t} \times \rho_{i,t} \\ \rho_{i,t} | \rho_{i,t-1}, \rho_{i,t+1} &\sim \text{Bernoulli}(\rho_{i,t-1} + (1 - \rho_{i,t-1})\rho_{i,t+1}\Phi(\beta_0^p + \beta_1^p d_{i,t})) \\ \log \psi_{i,t} &\sim N(\mathbf{x}'_{i,t}\beta^x + \dots, \sigma^2) \end{aligned} \quad (3)$$

where $\Phi(\cdot)$ is the standard normal cumulative distribution function for the probit probability of transitioning to the mature state, depending on tree diameter $d_{i,t}$. Importantly, $\Phi(\beta_0^p + \beta_1^p d_{i,t})$ it is the probability of making the transition for an individual that is now in the immature state. For this reason, the coefficients β_0^p, β_1^p for maturation in the second line of Equation (3) engage only for the transition tree-years, $[\rho_{i,t} | \rho_{i,t-1} = 0, \rho_{i,t+1} = 1]$. Predictors in the design vector for conditional fecundity $\mathbf{x}'_{i,t}$ include the mean climate variables (defined at the species level) tested here and competition by neighbours, and β^x is the estimated parameter vector. The ellipses (...) in Equation (3) includes individual effects (subscript i) and year effects (subscript t) (Clark, Nuñez, and Tomasek 2019). The variance not assigned to predictors is $s^2 = \sigma^2 + \text{Var}(\text{individuals}) + \text{Var}(\text{years})$.

Setting all other fitted variables at their mean values and intermediate shade (shade class 3 on the scale from 1 to 5), we obtained (unconditional) fecundity f from the fitted model. We first factored the joint distribution of conditional fecundity and maturation,

$$[\psi > f_{\min}, \rho = 1] = [\psi > f_{\min} | \rho = 1][\rho = 1] \quad (4)$$

Using Bayes' theorem, the cumulative distribution function for maturation diameter is

$$\begin{aligned} [d_{\text{matr}} > d | \psi > f_{\min}, \rho = 1] &\propto [\psi > f_{\min} | \rho = 1][d] \\ &= [\psi > f_{\min} | \rho = 1, d][\rho = 1][d] \\ &= \Phi(z_1)\Phi(z_2)[d] \end{aligned} \quad (5)$$

where $z_1 = \frac{\log f_{\min} - \mathbf{x}'\beta^y}{s}$ (log normal fecundity), $z_2 = \beta_0^p + \beta_1^p d$ (probit maturation) and again, s^2 is the marginal variance for conditional fecundity. We have taken the diameter distribution $[d]$ to be uniform. The distribution of maturation size is obtained using inverse distribution sampling from Equation (5), and we estimated d_{mat} as the mean of this distribution. We selected species for which maturation and fecundity schedules could be estimated with confidence. The selection was based on estimates of maturation status from the MASTIF model, and we retained species with at least 10 immature and 10 mature individuals. This included 486 species observed over a range of values for diameters.

2.3 | Trait and Climate Data

Like maximum tree height or age, maximum tree diameter is a useful concept, despite the fact that it cannot be known. To incorporate the concept of size differences, we use extreme sizes available from literature and our inventory data, recognising that the concept of a maximum becomes most meaningful with large sample sizes, which are not available for all species. For the final analysis here, we kept the highest estimates of d_{max} . Sources in Table S3 include tropical species from large plots in central Panama (189 species) and French Guyana (33 species), which together represent 45.7% of values. Estimates extracted from the internet (e.g., encyclopaedia, online flora) include 173 species (35.6%) (Table S3). For species not estimated in other sources, we used forest inventory data, evaluated by two approaches, both based on order statistics. We avoided using the absolute largest reported value in forest/MASTIF inventories due to the high noise levels associated with extremes. Order statistics were preferred over quantiles, the latter being determined by whether there are huge numbers of small trees in the data set; quantiles are based on the entire stand structure, whereas here the goal is to estimate the largest sizes, regardless of whether there are few or many small trees. For species present in national forest inventories we estimated d_{max} following Qiu et al. (2021) by using the tenth largest order statistic (38 species, 7.8%). For the remaining species present in MASTIF inventories, and with at least more than 90 unique individuals, we used the fifth largest order statistics (representing in total 8.4%). For species having only maximal plant height (Liu et al. 2019), but no d_{max} , we converted them to d_{max} using allometric equations of Feldpausch et al. (2011) (12 species, <2.5%). Observations of d_{max} coming from the internet are usually higher than data from National Forest Inventories, allometric predictions and MASTIF inventories (Figure S2). Seed size estimates came from measurements in our lab (Clark et al. 2021), the primary literature and the TRY Plant Trait Database (Kattge et al. 2011). Wood density and SLA are from the compilation of Carmona et al. (2021). We used genus- or family-level means for seed size, SLA and wood density values that were missing at the species level (15%, 28% and 26%, respectively). We defined a species' seed productivity as (mass per seed) $\times$ (mean seeds per tree basal area) (Qiu et al. 2022).

For species' climate, we extracted average temperature (in °C) and moisture deficit (evapotranspiration minus precipitation, in mm) for each species based on all occurrences in the Global Biodiversity Information Facility (GBIF) through the R package rgbif (Chamberlain and Boettiger 2017). The GBIF request is available from reference GBIF.org (2022). For species that are absent from GBIF, we extracted temperature and deficit from the MASTIF sites where those species were reported (162 species, 33%). Climate variables were obtained from CHELSA (Karger et al. 2017).

2.4 | Maturation and Maximum Size

To test the alternative hypotheses that the maturation diameter decreases ($\beta_d < 1$) or increases ($\beta_d > 1$) with maximum species size (Figure 2b), we estimated parameters in Equation (1) with the model

$$\begin{aligned} \log_{10}(d_{\text{mat}_s}) &= \log_{10}(\alpha) + \beta_d \times \log_{10}(d_{\text{max}_s}) + \dots + \varepsilon_s \\ \varepsilon &\sim N(0, \sigma^2) \end{aligned} \quad (6)$$

where for species s , where the ellipsis includes climatic variables (moisture deficit and temperature) and their interactions with $d_{\max}$. We tested alternative models including independence between maturation and maximum size (fitted α with β_d fixed at zero), proportionate increase (fitted α with $\beta_d = 1$) and changing relationship with size (both α and β_d estimated). Models were fitted with regression by using species average estimates of $d_{\max}$ as a response, and we included the inverse of the standard error of $d_{\max}$ as weights. Model selection and fit were evaluated with AIC and root mean square error (RMSE). Regression dilution could cause underestimation of the strength between here $d_{\max}$ and $d_{\max}$ when a predictor (i.e., $d_{\max}$) contains errors (Frost and Thompson 2000; Detto et al. 2019). We thus ran additional analyses to test the robustness of our results to the regression dilution effect (see Supplementary Material A2). First, we corrected the parameter $\hat{\beta}_d$ from measurement error by using the R package mecor (Nab 2021) (v1.0). Secondly, we tested if the relationship between $d_{\max}$ and $d_{\max}$ varies depending on the origin of $d_{\max}$.

2.5 | Joint Trait Analysis

We evaluated the association between maturation size and other species' traits from the ability of $d_{\max}$ to predict other trait values while allowing for climate and phylogeny effects (Figure 2c). The marginal correlations that are commonly used for this purpose do not account for the many ways that traits can be related to one another. For instance, maturation size might be associated with maximum size because both tend to be high in warm climates, or in the phylogenetic groups that tend to occur in warm climates. To accommodate co-dependence between trait values we used generalised joint attribute modelling (GJAM) with traits as responses (Clark 2016). To account for phylogeny in the joint traits model, we diverged from traditional assumptions concerning residual covariance. Instead, we adopted a direct inference of the effects of phylogenetic groups. Traditional approaches of phylogenetic correction build on highly specific assumptions for the residual variance (random walk, or more complex models representing stabilising selection such as the Ornstein–Uhlenbeck model). Our departure from these assumptions stems from the recognition that natural selection does not operate uniformly, neither within a given species pair nor across a broad spectrum of species. Our GJAM analysis explored phylogenetic contributions, with species groups treated as random effects and covariance that is unconstrained by assumptions on divergence rates (Qiu et al. 2023). Explanatory variables included temperature, moisture deficit and their interaction. Traits included wood density (g m^{-3}), SLA ($\text{mm}^2 \text{mg}^{-1}$), species seed productivity (kg m^{-2} basal area), seed size (g), maximum diameter ($d_{\max}$) (cm) and maturation diameter ($d_{\max}$) (cm). All traits were log-transformed. We included a random phylogenetic group effect in the joint trait analysis (Qiu et al. 2022, 2023; Bogdziewicz et al. 2023). For species in speciose genera (more than 10 species), genus was used as the phylogenetic group. For species in less speciose genera but belonging to families with more than five species, family was used as the phylogenetic group. For the remaining species (<25% of the total), an 'other' category was used. To estimate the direct effect of traits (i.e., SLA, wood density,

species seed productivity and seed size) and climatic variables on $d_{\max}$, we report conditional parameters from GJAM. Conditional parameters are estimated by extracting the parameters of the conditional distribution of traits conditioned on $d_{\max}$. Conditional parameters estimate the direct associations between traits while accounting for climate and phylogeny. Conditional parameters were obtained with the gjam R package (v2.6.2) (Supplementary Material, Section A1).

2.6 | Relation of $d_{\max}$ Along the Phylogeny

We visualised how $d_{\max}$ varies across species phylogeny by making a phylogenetic tree plot. We used the phylogeny from Zanne et al. (2014), and retrieved phylogenetic information for 400 out of the 486 studied species. Of the species missing from the phylogeny (i.e., 86 species), the relative proportion of missing phylogenetic information is about 13.2% for temperate species and about 19.4% for tropical species. We then tested for a phylogenetic signal in $d_{\max}$ and $d_{\max}$ using Pagel's λ (Pagel 1999) (which test for a Brownian motion evolutionary signal), with values close to 0 indicating low phylogenetic signal and values close to 1 suggesting a phylogenetic correlation. We plotted the phylogenetic tree with ggtree R package (v3.8) (Yu et al. 2017). We estimated the Pagel's λ by using the phylosig function from phytools (v1.5) (Revell 2012).

3 | Results

Maturation size is associated with maximum size, but not proportionately so (Figure 3a). Large inter-specific variation in $d_{\max}$ estimates had 95% quantiles that ranged from 4.0 to 51 cm, with relative maturation size ($d_{\max} = d_{\max}/d_{\max}$) quantiles of (0.07, 0.65). Contrary to the baseline independence model ($\beta_d = 0$), trees did not start to reproduce at a constant size (dashed black line in Figure 3a). If we force proportionality (fix β_d at 1), the estimate of $\hat{\alpha} = 0.24(0.23, 0.26)$ (Line 3 of Table 1) is consistent with Loehle's (1988) range for hardwoods (1/5–1/4), but far outside his range for conifers (1/15–1/10). The 95% CI that is well below 0.5. This differs from the Visser et al. (2016)'s estimate of 1/2 for Barro Colorado Island ($N = 60$ species), Panama and with Minor and Kobe (2019) La Selva, Costa Rica ($N = 16$ species). It is crucial to acknowledge that the aforementioned authors employed a distinct definition of $d_{\max}$ and estimated larger $d_{\max}$ (Figure S3). Moreover, this proportional cost model ($\beta_d = 1$) fits poorly, with twice the RMSE and a higher AIC than the best-fitting model (Table 1).

Fitting both α and β_d (Line 2 of Table 1) shows strong support for the diminishing risk model ($0 < \hat{\beta}_d < 1$). Allowing for environmental predictors further decreases the estimate to $\hat{\beta}_d = 0.30(0.15, 0.46)$. The exponent $0 < \beta_d < 1$ means that relative size at maturation ($d_{\max}$) decreases in large species (blue in Figure 3a).

The best-fitting model (lowest AIC and RMSE) includes a negative effect of temperature (maturation at small size for species most common in cold climates) and a positive interaction between temperature T and $d_{\max}$ (Table 1). This positive interaction means that the relationship between maturation and maximum




FIGURE 3 | Tree maturation size (a, b), and relative size at maturation (c, d) for 486 species. Each dot represents one species. Alternative models are dashed lines, black for independence between maturation size and maximum size ($\beta_d = 0$), and red for the proportional cost model ($\beta_d = 1$). The best fitting model (blue with 95%CI) supports the diminishing risk model ($\beta_d < 1$, Table 1). Panels b and d are predictions from the fitted model with an interaction between continuous d_{max} and temperature (Line 1 of Table 1). This model gives a continuous surface plot of maturation size as a function of maximum size and temperature (see Figure S4). However, for clarity, we represent only the prediction at cold (8°C, purple) and warm temperatures (25°C, green) spanning observed diameter ranges.

TABLE 1 | Coefficient estimates and fit to Equation (6). The selected model with the lowest AIC (bold font at top) includes temperature (β_T) and the interaction between d_{max} and temperature (β_{dT}). The proportional cost model has β_d fixed at 1. The independence model has β_d fixed at 0. Additional models that include moisture deficit and temperature have higher AIC values (Table S4).

α	β_d	β_T	β_{dT}	σ	AIC	RMSE
3.71 [1.94, 7.07]	0.30 [0.15, 0.46]	-0.023 [-0.035, -0.011]	0.012 [0.0058, 0.019]	0.089	-62	10.2
1.08 [0.93, 1.25]	0.59 [0.55, 0.63]	—	—	0.090	-52	10.0
0.24 [0.23, 0.26]	1	—	—	0.12	248	18.8
9.25 [8.69, 9.85]	0	—	—	0.15	447	15.4

Abbreviation: RMSE, root mean square error.

size tends to steepen for species in warm climates (Figure 3b,d), approaching the proportionate risk model (Table 1); the rise in d_{mat} with d_{max} increases with temperature. However, the main plus interaction effect remains below 1 even in warm climates showing that the diminishing risk model is supported across this temperature range. The β_d remained below 1 even when we restricted the analysis to a single source of d_{max} (Table S1 and Figure S1, see Supplementary Section A2). After correcting for risks of regression dilution, the average value of β_d remained below 1, with corrected $\beta_d = 0.73$; however, the confidence interval is between 0.03 and 1.43 (Table S2, see Supplementary Section A2).

The joint trait model incorporating random phylogenetic group and climate exhibits a root mean square prediction error

1.17 units smaller compared to the model that includes only climate. Conditional parameter estimates from the joint trait analysis show that d_{max} has a stronger effect on d_{mat} than other traits. There is a weak positive association with seed size, and a negative association with species fecundity (see Methods, Trait and Climate Data section) (Figure 4). There are no meaningful associations with wood density or SLA. The joint trait analysis also confirms the absence of a direct climate effect on d_{mat} after accounting for d_{max} in the conditional trait analysis (see Table S5 for joint trait model and Table S6 for conditional parameters). Previous linear models showed that temperature was significant only in the interaction with d_{max} (Table S4). Joint trait analysis indicates that the temperature effect on maturation size in Figure 2c could be due to the abundance of small species (small d_{max}) in warm climates (Figure S5). The trait relationships do




FIGURE 4 | Conditional parameter estimates for the direct effect of traits on tree size at maturation diameter (d_{mat}) while accounting for trait covariance, climate, and phylogeny. Conditional parameters are evaluated on a standardised scale (predictors are centred and standardised) making trait effects on d_{mat} respective to their variation in the data set. Shown are posterior means and 95% credible intervals. Blue and red represent positive and negative associations where 95% of the posterior does not include zero. SLA = specific leaf area.

not depend on the source of climatic data: (i) GBIF species occurrence in Figure 4 or (ii) more narrowly, where they occur in the MASTIF data network (Figure S6).

Both maturation (d_{mat}) and relative size (d_{rel}) show evidence of phylogenetic conservation ($\lambda_{mat} = 0.83$, $p < 0.0001$; $\lambda_{rel} = 0.51$, $p < 0.0001$, $n = 400$, Figure S7a), yet with substantial variation within some groups. The two-sample t-test for unequal variances shows differences between gymnosperms and angiosperms. Gymnosperms have high mean values for both d_{mat} and d_{max} and low mean values for d_{rel} (all $p < 0.0001$). Within gymnosperms, *Pinales* and *Cupressaceae* mature at large size, but large d_{max} gives them lower d_{rel} than most angiosperms (Figures S8 and S9). Most *Pinaceae* (*Picea*, *Pinus*) and *Cupressaceae* (*Thuja*, *Sequoia*), *Fagaceae* (*Quercus* and *Fagus*) and *Juglandaceae* have low d_{rel} (Figure S10). Plant groups with both tree and shrub habits, such as *Rosales*, *Magnoliales*, *Rubiaceae* and *Fabaceae*, have mixed d_{rel} . However, we did not find a significant effect of tree versus shrub habit on d_{rel} , possibly due to high variation in the data (Figure S10). Shrubs may tend to have high d_{rel} , but higher d_{rel} is also observed in trees genera like *Magnolia* and *Poulsenia*.

4 | Discussion

Our analysis suggests a diminishing risk model for the relationship between maturation and maximum size (i.e., d_{mat} and d_{max}). The novelty here comes from the low coupling we find. Indeed, the coefficient $\hat{\beta}_d = 0.30$ in Table 1 that we found is closer to zero (no relationship) than one. In contrast, the estimated exponent values fitted to vertebrates are greater than 1/2 (Prothero 1993; Herculano-Houzel 2019), twice the value

of $\hat{\beta}_d$ we find for trees. Nevertheless, the comparison across groups is complex due to the uncertainty on maximum size (d_{max}), which could influence the value of the exponent $\hat{\beta}_d$, but this issue remained unexplored in other taxa. The biological difference of $\hat{\beta}_d$ may arise because trees differ from other species groups in the gains that come from allocation to growth, as the gain is due to the relative difference in height with other competitive individuals. In most tree species, individuals in the understory produce no seed at all, while dominant stature can yield multi-order-of-magnitude gains in fecundity over crowded neighbours (Clark, LaDeau, and Ibanez 2004). In contrast, in vertebrates, improved parental condition and size can translate to incremental increases in clutch size or survival of well-provisioned offspring. Gestation times and physical limits on clutch size (e.g., one offspring) may allow only muted near-term benefits of reproductive delay.

In trees, large size comes with uncertainty that could weaken the potential benefit of delaying maturation. Wind exposure and risk of hydraulic failure both increase with size (Bennett et al. 2015; Jackson et al. 2021; Gardiner 2021; Barrere et al. 2023). The vanishing probability that a seed survives to large size, combined with the fact that fecundity can plateau and even decline late in life (Qiu et al. 2021) means that the competitive advantages of extremely large size can rarely make up for lost benefits of early reproduction. At a stand scale, the risk of stand-replacing disturbances can increase with stand age and development (e.g., accumulated fuels increase fire risk), such that species that fail to reach minimum reproductive size before the next disturbance can be excluded from communities (Clark 1991; McDowell et al. 2020). The fact that maturation size increases with maximum size means that the two are not independent. But the cost is not proportionate (Figure 3).

The fact that some correlation exists does not conflict with a disproportionate importance of near-term gains that can follow delayed maturation. Instead, it suggests that the benefits of large size probably do not come at the end of life. The capacity to reach a large size pays benefits throughout life, contributing with many other variables to current size and fecundity, not just as a tree approaches the maximum.

Both climate and species traits contribute to the relationships between maturation and maximum size. The negative main effect of temperature and its positive interaction steepens the relationship with maximum size in warm climates (Figure 3), where growth and mortality rates are generally higher than in temperate forests (Stephenson and Van Mantgem 2005; Locosselli et al. 2020). Abundant resources may offer a disproportionate advantage to early maturation (van Noordwijk and de Jong 1986; Kozłowski 1992; Wenk and Falster 2015). Long growing seasons in warm climates might have similar effects. However, intense competition on nutrient-rich sites might also favour delayed reproduction as trees compete for canopy access. Theoretical studies (Falster et al. 2017; Detto, Levine, and Pacala 2022) have shown that a trade-off between maximum size and maturation size can promote niche diversification and maintain species coexistence, and can be typically observed in tropical where there is a wide range of maximum sizes forests (Falster et al. 2017).

We did not find that high SLA is associated with maturation at small size (Visser et al. 2016) or early age (Wenk et al. 2018). The relationship reported in Wenk et al. (2018) includes leaf area from 1 year and one site, and the correlation estimated in that study does not appear to control for phylogeny. Similarly, lack of association with wood density in our study does not agree with suggestions that shade-tolerant species with high wood density mature at small size (Thomas, Martin, and Mycroft 2015). The inclusion of a wider range of plant species may reveal a different pattern of traits. For example, a comparison over a large number of perennial plant species such as herbs, graminoids, shrubs, and trees shows that traits that promote longevity are associated with greater variability in seed production (Journé, Hacket-Pain, and Bogdziewicz 2023). The fact that species that produce large seeds also allocate more to reproductive effort (Qiu et al. 2022) could contribute in a small way to delayed maturation. The differences between our result and previous work may be due to the larger species coverage, and to the control of the effect of climate and phylogeny in our joint analysis.

Results highlight the importance of large data sets and how they are modelled. This first compilation of tree maturation size for hundreds of species on five continents shows strong support for a diminishing risk model—trees that can get big can still mature at relatively small sizes. The result is a decline in the relative size of maturation for large trees (Figure 3d). The benefits of extensive data here parallel the shift from early theory that argued for a constant relative maturation size (d_{rel}) in fish (Charnov and Berrigan 1990), followed by studies showing an exponent that is less than one (Froese and Binohlan 2000; Tsikliras and Stergiou 2014; Thorson et al. 2017). However, it is important to acknowledge that our coverage of tree species diversity is still patchy, with most data coming from Europe and North America and limited spatial coverage in Africa, South America, Asia and Oceania as data are concentrated in a few large plots in these areas (Daru and Rodriguez 2023).

New insight from this analysis comes first from extending observations beyond a small number of tropical sites, few species or limited sample size (Thomas 1996, 2011; Wright et al. 2005; Visser et al. 2016; Minor and Kobe 2019). The expanded coverage of species and sites permitted the incorporation of climatic drivers into the analysis of d_{mat} in relation to d_{max} , which influenced the estimation of α and β_d . Second, this study also benefited from accommodating detection and temporal dependence to infer maturation. The estimation of d_{mat} was possible by combining diverse datasets, either based from direct crop measurement and seed trap monitoring and by the use of MASTIF model which could estimate jointly a probability of maturation and individual fecundities. For instance, estimates of d_{mat} from Visser et al. (2016) are, on average, approximately 1.8 times larger than our estimates for the species in common in the two studies (Figure S3). This discrepancy could be attributed to a different definition of size at maturation and methods of analysis, as our method also includes the number of seeds produced.

Due to the high juvenile mortality, the maturation sizes quantified here are expected to impact predictions from demographic vegetation models, including earth system models (ESMs) that include effects of maturation size. In one ESM

study that considered the effects of maturation height, variation in a single value applied to all species did not have a large impact on simulated stand productivity (Raczka et al. 2018). However, when differences in species maturation size are accounted for in models, the effect can be larger. Few individuals survive to large size and, thus, their ability to reproduce early can be important. The fact that species capable of large size tend to retain this capacity to reproduce while still small highlights the importance of understanding maturation size. Accurate estimation of maturation size (d_{mat}) is likewise important for assessing response to disturbance regimes, especially as the time to maturity begins to exceed the interval between disturbances. For instance, several species that compared pairs of species found that species with smaller size at maturation can have better post-disturbance dynamics than species with larger maturation size (Alfaro-Sánchez et al. 2022; Andrus et al. 2020). Our results provided the data to test such hypotheses at a much larger scale. This effect can even scale up at the ecosystem scale. In boreal habitats, exposure to more frequent disturbances that exceed the tree maturation time can completely change a tree community to a grass-dominated community without a return to a forest stand within centuries (Buma et al. 2013).

Current ESMs suffer from limited information on allocation to reproduction (Wenk and Falster 2015), including empirical data (Hanbury-Brown, Ward, and Kueppers 2022b). Similarly, management actions intended to assure regeneration from seed also need to consider if the minimum harvest diameter is smaller than maturation size (Ouédraogo et al. 2018). Maturation size may play an important role in the ability of species to respond to disturbance and climate change (McDowell et al. 2020) when tree maturation can be reached faster under elevated CO₂ exposure (LaDeau and Clark 2001). Considering the difference in maturation size between species may be crucial, although it can be more challenging to comprehend due to the impact of CO₂ on maturation size. Understanding how fecundity strategies differ between species and phylogenetic groups, such as lower relative size at maturation (d_{rel}) for gymnosperms than angiosperms, may open a new avenue to better understand species diversification and responses to disturbances (Bond 1989; Verdu 2002; Qiu et al. 2022). Developing a model that represents the size at which a species begins to produce seeds could improve the representation of the regeneration of each functional type (Hanbury-Brown et al. 2022a) and colonisation rates (Snell 2014) and improve our understanding of species coexistence. Our study contributes to the maturation sizes needed for each of these objectives.

Author Contributions

V.J., G.K. and J.S.C. performed analyses, led the paper and designed the study. V.J., M.B., B.C., G.K., T.Q. and J.S.C. co-wrote the paper. J.S.C. compiled the MASTIF network, and wrote the MASTIF model and software. All authors contributed data and revised the paper.

Affiliations

¹Université Grenoble Alpes, Institut National de Recherche Pour Agriculture, Alimentation et Environnement (INRAE), Laboratoire EcoSystèmes et Sociétés En Montagne (LESSEM), Grenoble,

France | ²Forest Biology Center, Institute of Environmental Biology, Adam Mickiewicz University, Poznań, Poland | ³Department of Ecosystem Science and Management, Pennsylvania State University, University Park, Pennsylvania, USA | ⁴Centro Austral de Investigaciones Científicas (CADIC), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), B. Houssay 200 (9410), Ushuaia, Tierra del Fuego, Argentina | ⁵Department of Agriculture, Forest and Food Sciences, University of Torino, Turin, Italy | ⁶Forest Research Institute, University of Quebec in Abitibi-Temiscamingue, Rouyn-Noranda, Quebec, Canada | ⁷Université Paris-Saclay, CNRS, AgroParisTech, Ecologie Systématique et Evolution, Gif-sur-Yvette, France | ⁸Institut National de Recherche Pour Agriculture, Alimentation et Environnement (INRAE), Ecologie des Forêts Méditerranéennes, Avignon, France | ⁹Université Bordeaux, Institut National de Recherche Pour Agriculture, Alimentation et Environnement (INRAE), biodiversity, Genes, and Communities (BIOGECO), Pessac, France | ¹⁰Department of Biodiversity, Ecology and Evolution, Complutense University of Madrid, Madrid, Spain | ¹¹INRAE, Aix-Marseille University, UMR RECOVER, Aix-En-Provence, France | ¹²ICIFOR (Forest Research Institute), INIA-CSIC, Madrid, Spain | ¹³Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas (IPE-CSIC), Zaragoza, Spain | ¹⁴Department of Biological Sciences, National Sun Yat-Sen University, Kaohsiung, Taiwan | ¹⁵Unité Evolution et Diversité Biologique (EDB), CNRS, IRD, UPS, Toulouse, France | ¹⁶CREA-Research Centre for Forestry and Wood, Arezzo, Italy | ¹⁷Aix Marseille Université, Institut National de Recherche Pour Agriculture, Alimentation et Environnement (INRAE), Aix-en-Provence, France | ¹⁸Research Centre for Forestry and Wood, Arezzo, Italy | ¹⁹USGS Western Ecological Research Center, Three Rivers, California, USA | ²⁰Institute of Mediterranean and Forest Ecosystems, Hellenic Agricultural Organization, Athens, Greece | ²¹Earth and Environment, Boston University, Boston, Massachusetts, USA | ²²Facultad de Ciencias Forestales y de la Conservación de la Naturaleza (FCFCN), Universidad de Chile, Santiago, Chile | ²³Centre d'Ecologie Fonctionnelle et Evolutive (CEFE), Centre National de la Recherche Scientifique (CNRS), Montpellier, France | ²⁴Centre de Recerca Ecològica i Aplicacions Forestals (CREAF), Bellaterra, Catalunya, Spain | ²⁵Biology Department, Center for Energy, Environment, and Sustainability, Wake Forest University, Winston Salem, North Carolina, USA | ²⁶Biology Department, University of Southampton, Southampton, UK | ²⁷Forest Resources, University of Washington, Seattle, Washington, USA | ²⁸Department of Biological Sciences, Center for Adaptive Western Landscapes, Flagstaff, Arizona, USA | ²⁹Department of Environmental Studies, University of California, Santa Cruz, California, USA | ³⁰Department of Forest and Soil Sciences, Institute of Forest Ecology, University of Natural Resources and Life Sciences, Vienna, Austria | ³¹Bent Creek Experimental Forest, USDA Forest Service, Asheville, North Carolina, USA | ³²INRAE, Bordeaux Sciences Agro, UMR 1391 ISPA, Villenave d'Ornon, France | ³³Eastern Forest Environmental Threat Assessment Center, USDA Forest Service, Southern Research Station, Research Triangle Park, Durham, North Carolina, USA | ³⁴Department of Geography and Planning, School of Environmental Sciences, University of Liverpool, Liverpool, UK | ³⁵BIOGECO, INRAE, University of Bordeaux, Cestas, France | ³⁶Department of Plant Ecology, Forestry and Forest Products Research Institute (FFPRI), Tsukuba, Japan | ³⁷School of Biological and Marine Sciences, University of Plymouth, Plymouth, UK | ³⁸Plant Ecology, Institute of Integrative Biology, D-USYS, ETH Zürich, Zürich, Switzerland | ³⁹Department of Forest Ecology, Silva Tarouca Research Institute, Brno, Czech Republic | ⁴⁰Department of Biological Environment, Akita Prefectural University, Akita, Japan | ⁴¹School for Environment and Sustainability, University of Michigan, Ann Arbor, Michigan, USA | ⁴²Institute of Arctic Biology, University of Alaska, Fairbanks, Alaska, USA | ⁴³Health and Environmental Sciences Department, Xian Jiaotong-Liverpool University, Suzhou, China | ⁴⁴Department of Plant Biology, Program in Ecology, Evolutionary Biology, and Behavior, Michigan State

University, East Lansing, Michigan, USA | ⁴⁵Department of Forest Vegetation, Forestry and Forest Products Research Institute, Tsukuba, Ibaraki, Japan | ⁴⁶Department of Natural Sciences, Manchester Metropolitan University, Manchester, UK | ⁴⁷Department of Biological Sciences, DePaul University, Chicago, Illinois, USA | ⁴⁸Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Krakow, Poland | ⁴⁹USDA, Forest Service, Stoneville, Mississippi, USA | ⁵⁰CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France | ⁵¹Department of Wildland Resources, and The Ecology Center, Utah State University, Logan, Utah, USA | ⁵²Department of Biology, University of New Mexico, Albuquerque, New Mexico, USA | ⁵³INRAE, UR EFNO, Nogent-Sur-Vernisson, France | ⁵⁴Pacific Forestry Centre, Victoria, British Columbia, Canada | ⁵⁵School of Natural Sciences, UC Merced, Merced, California, USA | ⁵⁶Department of Biology, Washington University in St. Louis, St. Louis, Missouri, USA | ⁵⁷Department of Forestry and Renewable Forest Resources, Biotechnical Faculty, University of Ljubljana, Ljubljana, Slovenia | ⁵⁸Tohoku Research Center, Forestry and Forest Products Research Institute, Morioka, Iwate, Japan | ⁵⁹Institute of Plant Sciences, University of Bern, Bern, Switzerland | ⁶⁰Valles Caldera National Preserve, National Park Service, Jemez Springs, Jemez Springs, New Mexico, USA | ⁶¹Fort Collins Science Center, Fort Collins, Colorado, USA | ⁶²Inst. de Recursos Naturales y Agrobiología de Sevilla, Consejo Superior de Investigaciones Científicas (IRNAS-CSIC), Andalucía, Spain | ⁶³W. Szafer Institute of Botany, Polish Academy of Sciences, Krakow, Poland | ⁶⁴Department of Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic | ⁶⁵The Nature Conservancy, Boulder, Colorado, USA | ⁶⁶Department of Environmental Science Policy and Management, University of California Berkeley, Berkeley, California, USA | ⁶⁷Nicholas School of the Environment, Duke University, Durham, North Carolina, USA | ⁶⁸Ecology and Evolutionary Biology Department, Yale University, New Haven, Connecticut, USA | ⁶⁹Department of Biology, Wake Forest University, Winston-Salem, North Carolina, USA | ⁷⁰Department of Biology, Wilkes University, Wilkes-Barre, Pennsylvania, USA | ⁷¹Department of Environmental Science and Ecology, State University of New York-Brockport, Brockport, New York, USA | ⁷²Data Science, College of William and Mary, Williamsburg, Virginia, USA | ⁷³School of Life Sciences, Keele University, Staffordshire, UK | ⁷⁴Department of Ecology, Evolution and Environmental Biology, Columbia University, New York, New York, USA | ⁷⁵Department of Agricultural and Environmental Sciences-Production, Territory, Agroenergy (DISAA), University of Milan, Milano, Italy | ⁷⁶Department of Biological Sciences, Northern Arizona University, Flagstaff, Arizona, USA | ⁷⁷Smithsonian Tropical Research Institute, Panama, Republic of Panama | ⁷⁸Department of Environmental Sciences, University of Puerto Rico, Rio Piedras, Puerto Rico, USA

Acknowledgements

We thank the National Ecological Observatory Network (NEON) for access to sites and vegetation structure data. The project has been funded by grants to JSC from the National Science Foundation, most recently DEB-1754443, and by the Belmont Forum (1854976), NASA (AIST16-0052, AIST18-0063) and the Programme d'Investissement d'Avenir under project FORBIC (18-MPGA-0004) (*Make Our Planet Great Again*). Jerry Franklin's data remain accessible through NSF LTER DEB-1440409. Puerto Rico data were funded by NSF grants, most recently, DEB 0963447 and LTREB 11222325. Data from the Andes Biodiversity and Ecosystem Research Group were funded by the Gordon and Betty Moore Foundation and NSF LTREB 1754647. V.J. was supported by project FORBIC (18-MPGA-0004) and project No. 2021/43/P/NZ8/01209 co-funded by the Polish National Science Centre and the EU H2020 research and innovation programme under the MSCA GA No. 945339. For the purpose of Open Access, the authors has applied a CC-BY public copyright licence to any Author Accepted Manuscript (AAM) version arising

from this submission. M.B. was supported by grant no. 2019/35/D/NZ8/00050 from the Polish National Science Centre, and Polish National Agency for Academic Exchange Bekker programme PPN/BEK/2020/1/00009/U/00001. Research by the USDA Forest Service and the USGS was funded by these agencies. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the US Government.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

Data and code supporting our results are archived on the Open Science Framework (OSF) Repository: <https://doi.org/10.17605/OSF.IO/U23VY>. All analyses used R Core Team (2023) (v4.3.0) and published R packages.

Peer Review

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/10.1111/ele.14500>.

References

- Alfaro-Sánchez, R., J. F. Johnstone, S. G. Cumming, et al. 2022. "What Drives Reproductive Maturity and Efficiency in Serotinous Boreal Conifers?" *Frontiers in Ecology and Evolution* 10: 1–12.
- Andrus, R. A., B. J. Harvey, A. Hoffman, and T. T. Veblen. 2020. "Reproductive Maturity and Cone Abundance Vary With Tree Size and Stand Basal Area for Two Widely Distributed Conifers." *Ecosphere* 11: 3092.
- Barrere, J., B. Reineking, T. Cordonnier, et al. 2023. "Functional Traits and Climate Drive Interspecific Differences in Disturbance-Induced Tree Mortality." *Global Change Biology* 29: 2836–2851.
- Bennett, A. C., N. G. McDowell, C. D. Allen, and K. J. Anderson-Teixeira. 2015. "Larger Trees Suffer Most During Drought in Forests Worldwide." *Nature Plants* 1: 15139.
- Bogdziewicz, M., M. C. A. Acuña, R. Andrus, et al. 2023. "Linking Seed Size and Number to Trait Syndromes in Trees." *Global Ecology and Biogeography* 32: 683–694.
- Bond, W. J. 1989. "The Tortoise and the Hare: Ecology of Angiosperm Dominance and Gymnosperm Persistence." *Biological Journal of the Linnean Society* 36: 227–249.
- Buma, B., C. D. Brown, D. C. Donato, J. B. Fontaine, and J. F. Johnstone. 2013. "The Impacts of Changing Disturbance Regimes on Serotinous Plant Populations and Communities." *Bioscience* 63: 866–876.
- Caignard, T., A. Kremer, X. P. Bouteiller, et al. 2021. "Counter-Gradient Variation of Reproductive Effort in a Widely Distributed Temperate Oak." *Functional Ecology* 35: 1745–1755.
- Caignard, T., A. Kremer, C. Firmat, M. Nicolas, S. Venner, and S. Delzon. 2017. "Increasing Spring Temperatures Favor Oak Seed Production in Temperate Areas." *Scientific Reports* 7: 1–8.
- Carmona, C. P., C. G. Bueno, A. Toussaint, et al. 2021. "Fine-Root Traits in the Global Spectrum of Plant Form and Function." *Nature* 597: 683–687.
- Chamberlain, S., and C. Boettiger. 2017. "R Python, and Ruby Clients for GBIF Species Occurrence Data." *PeerJ PrePrints* 5: e3304v1.
- Charlesworth, B. 2000. "Fisher, Medawar, Hamilton and the Evolution of Aging." *Genetics* 156: 927–931.
- Charnov, E. L., and D. Berrigan. 1990. "Age of Maturity Versus the Adult Lifespan." *Evolutionary Ecology* 4: 273–275.
- Charnov, E. L., and D. Berrigan. 1991. "Evolution of Life History Parameters in Animals With Indeterminate Growth, Particularly Fish." *Evolutionary Ecology* 5: 63–68.
- Clark, J. S. 1991. "Disturbance and Tree Life History on the Shifting Mosaic Landscape." *Ecology* 72: 1102–1118.
- Clark, J. S. 2016. "Why Species Tell More About Traits Than Traits About Species: Predictive Analysis." *Ecology* 97: 1979–1993.
- Clark, J. S., R. Andrus, M. Aubry-Kientz, et al. 2021. "Continent-Wide Tree Fecundity Driven by Indirect Climate Effects." *Nature Communications* 12: 1–11.
- Clark, J. S., D. M. Bell, M. C. Kwit, and K. Zhu. 2014. "Competition-Interaction Landscapes for the Joint Response of Forests to Climate Change." *Global Change Biology* 20: 1979–1991.
- Clark, J. S., S. LaDeau, and I. Ibanez. 2004. "Fecundity of Trees and the Colonization–Competition Hypothesis." *Ecological Monographs* 74: 415–442.
- Clark, J. S., C. L. Nuñez, and B. Tomasek. 2019. "Foodwebs Based on Unreliable Foundations: Spatiotemporal Masting Merged With Consumer Movement, Storage, and Diet." *Ecological Monographs* 89: 1–24.
- Clark, J. S., M. Silman, R. Kern, E. Macklin, and J. HilleRisLambers. 1999. "Seed Dispersal Near and Far: Patterns Across Temperate and Tropical Forests." *Ecology* 80: 1475–1494.
- Daru, B. H., and J. Rodriguez. 2023. "Mass Production of Unvouchered Records Fails to Represent Global Biodiversity Patterns." *Nature Ecology & Evolution* 7: 816–831.
- Davies, S. J., and P. S. Ashton. 1999. "Phenology and Fecundity in Pioneer Species of Euphorbiaceae." *American Journal of Botany* 86: 1786–1795.
- Detto, M., J. M. Levine, and S. W. Pacala. 2022. "Maintenance of High Diversity in Mechanistic Forest Dynamics Models of Competition for Light." *Ecological Monographs* 92: e1500.
- Detto, M., M. D. Visser, S. J. Wright, and S. W. Pacala. 2019. "Bias in the Detection of Negative Density Dependence in Plant Communities." *Ecology Letters* 22: 1923–1939.
- Dietze, M. C., and J. S. Clark. 2008. "Changing the Gap Dynamics Paradigm: Vegetative Regeneration Control on Forest Response to Disturbance." *Ecological Monographs* 78: 331–347.
- Falster, D. S., Å. Brännström, M. Westoby, and U. Dieckmann. 2017. "Multitrait Successional Forest Dynamics Enable Diverse Competitive Coexistence." *Proceedings of the National Academy of Sciences of the United States of America* 114: E2719–E2728.
- Falster, D. S., and M. Westoby. 2003. "Plant Height and Evolutionary Games." *Trends in Ecology & Evolution* 18: 337–343.
- Feldpausch, T. R., L. Banin, O. L. Phillips, et al. 2011. "Height-Diameter Allometry of Tropical Forest Trees." *Biogeosciences* 8: 1081–1106.
- Froese, R., and C. Binohlan. 2000. "Empirical Relationships to Estimate Asymptotic Length, Length at First Maturity and Length at Maximum Yield Per Recruit in Fishes, With a Simple Method to Evaluate Length Frequency Data." *Journal of Fish Biology* 56: 758–773.
- Frost, C., and S. G. Thompson. 2000. "Correcting for Regression Dilution Bias: Comparison of Methods for a Single Predictor Variable." *Journal of the Royal Statistical Society. Series A, Statistics in Society* 163: 173–189.
- Gardiner, B. 2021. "Wind Damage to Forests and Trees: A Review With an Emphasis on Planted and Managed Forests." *Journal of Forest Research* 26: 248–266.
- GBIF.org. 2022. "GBIF Occurrence Download." <https://doi.org/10.15468/dl.wevh3v>.

- Hanbury-Brown, A. R., T. L. Powell, H. C. Muller-Landau, S. J. Wright, and L. M. Kueppers. 2022a. "Simulating Environmentally-Sensitive Tree Recruitment in Vegetation Demographic Models." *New Phytologist* 235: 78–93.
- Hanbury-Brown, A. R., R. E. Ward, and L. M. Kueppers. 2022b. "Forest Regeneration Within Earth System Models: Current Process Representations and Ways Forward." *New Phytologist* 235: 20–40.
- Herculano-Houzel, S. 2019. "Longevity and Sexual Maturity Vary Across Species With Number of Cortical Neurons, and Humans Are no Exception." *Journal of Comparative Neurology* 527: 1689–1705.
- Jackson, T. D., A. F. Shenkin, N. Majalap, et al. 2021. "The Mechanical Stability of the World's Tallest Broadleaf Trees." *Biotropica* 53: 110–120.
- Jensen, A. L. 1996. "Beverton and Holt Life History Invariants Result From Optimal Trade-Off of Reproduction and Survival." *Canadian Journal of Fisheries and Aquatic Sciences* 822: 820–822.
- Journé, V., R. Andrus, M. C. Aravena, et al. 2022. "Globally, Tree Fecundity Exceeds Productivity Gradients." *Ecology Letters* 25: 1471–1482.
- Journé, V., A. Hacket-Pain, and M. Bogdziewicz. 2023. "Evolution of Masting in Plants Is Linked to Investment in Low Tissue Mortality." *Nature Communications* 14: 7998.
- Karger, D. N., O. Conrad, J. Böhner, et al. 2017. "Climatologies at High Resolution for the Earth's Land Surface Areas." *Scientific Data* 4: 1–20.
- Kattge, J., S. Díaz, S. Lavorel, et al. 2011. "TRY—A Global Database of Plant Traits." *Global Change Biology* 17: 2905–2935.
- Koch, G. W., S. C. Stillet, G. M. Jennings, and S. D. Davis. 2004. "The Limits to Tree Height." *Nature* 428: 851–854.
- Kohler, P. K., and A. Huth. 2004. "Simulating Growth Dynamics in a South-East Asian Rainforest Threatened by Recruitment Shortage and Tree Harvesting." *Climatic Change* 67: 95–117.
- Kohyama, T., E. Suzuki, T. Partomihardjo, T. Yamada, and T. Kubo. 2003. "Tree Species Differentiation in Growth, Recruitment and Allometry in Relation to Maximum Height in a Bornean Mixed Dipterocarp Forest." *Journal of Ecology* 91: 797–806.
- Kozłowski, J. 1992. "Optimal Allocation of Resources to Growth and Reproduction: Implications for Age and Size at Maturity." *Trends in Ecology & Evolution* 7: 15–19.
- LaDeau, S. L., and J. S. Clark. 2001. "Rising CO₂ Levels and the Fecundity of Forest Trees." *Science* 292: 95–98.
- Le Roncé, I., J. Gavinet, J. M. Ourcival, F. Mouillot, I. Chuine, and J. M. Limousin. 2021. "Holm oak Fecundity Does Not Acclimate to a Drier World." *New Phytologist* 231: 631–645.
- Lines, E. R., M. A. Zavala, D. W. Purves, and D. A. Coomes. 2012. "Predictable Changes in Aboveground Allometry of Trees Along Gradients of Temperature, Aridity and Competition." *Global Ecology and Biogeography* 21: 1017–1028.
- Liu, H., S. M. Gleason, G. Hao, et al. 2019. "Hydraulic Traits Are Coordinated With Maximum Plant Height at the Global Scale." *Science Advances* 5: eaav1332.
- Locosselli, G. M., R. J. W. Brien, M. S. Leite, et al. 2020. "Global Tree-Ring Analysis Reveals Rapid Decrease in Tropical Tree Longevity With Temperature." *Proceedings of the National Academy of Sciences* 117: 33358–33364.
- Loehle, C. 1988. "Tree Life History Strategies: The Role of Defenses." *Canadian Journal of Forest Research* 18: 209–222.
- McDowell, N., C. Allen, K. Anderson-Teixeira, et al. 2020. "Pervasive Shifts in Forest Dynamics in a Changing World." *Science* 368: eaaz9463.
- Minor, D. M., and R. K. Kobe. 2019. "Fruit Production Is Influenced by Tree Size and Size-Asymmetric Crowding in a Wet Tropical Forest." *Ecology and Evolution* 9: 1458–1472.
- Moles, A. T., D. D. Ackerly, J. C. Tweddle, et al. 2006. "Global Patterns in Seed Size." *Global Ecology and Biogeography* 16: 109–116.
- Moles, A. T., D. S. Falster, M. R. Leishman, and M. Westoby. 2004. "Small-Seeded Species Produce More Seeds Per Square Metre of Canopy Per Year, but Not Per Individual Per Lifetime." *Journal of Ecology* 92: 384–396.
- Nab, L. 2021. "Mecor: Measurement Error Correction in Linear Models With a Continuous Outcome." R Package Version 1.0.0.
- Niklas, K. J. 1994. *Plant Allometry. The Scaling of Form and Process*. Chicago: The University of Chicago Press.
- van Noordwijk, A., and G. de Jong. 1986. "Acquisition and Allocation of Resources: Their Influence on Variation in Life History Tactics." *American Naturalist* 128: 137–142.
- Ouédraogo, D. Y., J. L. Doucet, K. Dainou, et al. 2018. "The Size at Reproduction of Canopy Tree Species in Central Africa." *Biotropica* 50: 465–476.
- Pagel, M. 1999. "Inferring the Historical Patterns of Biological Evolution." *Nature* 401: 877–884.
- Prothero, J. 1993. "Adult Life Span as a Function of Age at Maturity." *Experimental Gerontology* 28: 529–536.
- Qiu, T., R. Andrus, M. C. Aravena, et al. 2022. "Limits to Reproduction and Seed Size-Number Trade-Offs That Shape Forest Dominance and Future Recovery." *Nature Communications* 13: 2381.
- Qiu, T., M. C. Aravena, R. Andrus, et al. 2021. "Is There Tree Senescence? The Fecundity Evidence." *Proceedings of the National Academy of Sciences of the United States of America* 118: 1–10.
- Qiu, T., M. C. Aravena, D. Ascoli, et al. 2023. "Masting Is Uncommon in Trees That Depend on Mutualist Dispersers in the Context of Global Climate and Fertility Gradients." *Nature Plants* 9: 1044–1056.
- R Core Team. 2023. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Raczka, B., M. C. Dietze, S. P. Serbin, and K. J. Davis. 2018. "What Limits Predictive Certainty of Long-Term Carbon Uptake?" *Journal of Geophysical Research: Biogeosciences* 123: 3570–3588.
- Revell, L. J. 2012. "Phytools: An R Package for Phylogenetic Comparative Biology (And Other Things)." *Methods in Ecology and Evolution* 3: 217–223.
- Seyednasrollah, B., and J. S. Clark. 2020. "Where Resource-Acquisitive Species Are Located: The Role of Habitat Heterogeneity." *Geophysical Research Letters* 47: 1–12.
- Sharma, S., R. Andrus, Y. Bergeron, et al. 2022. "North American Tree Migration Paced by Climate in the West, Lagging in the East." *Proceedings of the National Academy of Sciences of the United States of America* 119: e2116691118.
- Snell, R. S. 2014. "Simulating Long-Distance Seed Dispersal in a Dynamic Vegetation Model." *Global Ecology and Biogeography* 23: 89–98.
- Stearns, S. C. 1989. "Trade-Offs in Life-History Evolution." *Functional Ecology* 3: 259–268.
- Stephenson, N. L., and P. J. Van Mantgem. 2005. "Forest Turnover Rates Follow Global and Regional Patterns of Productivity." *Ecology Letters* 8: 524–531.
- Thomas, S. C. 1996. "Relative Size at Onset of Maturity in Rain Forest Trees: A Comparative Analysis of 37 Malaysian Species." *Oikos* 76: 1450154.
- Thomas, S. C. 2011. "Age-Related Changes in Tree Growth and Functional Biology: The Role of Reproduction." In *Size- and Age-Related Changes in Tree Structure and Function*, edited by F. C. Meinzer, B. Lachenbruch, and T. E. Dawson, vol. 4, 33–64. Dordrecht: Springer Netherlands.

- Thomas, S. C., A. R. Martin, and E. E. Mycroft. 2015. "Tropical Trees in a Wind-Exposed Island Ecosystem: Height-Diameter Allometry and Size at Onset of Maturity." *Journal of Ecology* 103: 594–605.
- Thorson, J. T., S. B. Munch, J. M. Cope, and J. Gao. 2017. "Predicting Life History Parameters for all Fishes Worldwide." *Ecological Applications* 27: 2262–2276.
- Tsikliras, A. C., and K. I. Stergiou. 2014. "Size at Maturity of Mediterranean Marine Fishes." *Reviews in Fish Biology and Fisheries* 24: 219–268.
- Verdu, M. 2002. "Age at Maturity and Diversification in Woody Angiosperms." *Evolution* 56: 1352–1361.
- Visser, M. D., M. Bruijning, S. J. Wright, et al. 2016. "Functional Traits as Predictors of Vital Rates Across the Life Cycle of Tropical Trees." *Functional Ecology* 30: 168–180.
- Wallentin, G., U. Tappeiner, J. Strobl, and E. Tasser. 2008. "Understanding Alpine Tree Line Dynamics: An Individual-Based Model." *Ecological Modelling* 218: 235–246.
- Wenk, E. H., K. Abramowicz, M. Westoby, and D. S. Falster. 2018. "Investment in Reproduction for 14 Iteroparous Perennials Is Large and Associated With Other Life-History and Functional Traits." *Journal of Ecology* 106: 1338–1348.
- Wenk, E. H., and D. S. Falster. 2015. "Quantifying and Understanding Reproductive Allocation Schedules in Plants." *Ecology and Evolution* 5: 5521–5538.
- Westoby, M., D. S. Falster, A. T. Moles, P. A. Vesk, and I. J. Wright. 2002. "Plant Ecological Strategies: Some Leading Dimensions of Variation Between Species." *Annual Review of Ecology and Systematics* 33: 125–159.
- Wright, S. J., M. A. Jaramillo, J. Pavon, R. Condit, S. P. Hubbell, and R. B. Foster. 2005. "Reproductive Size Thresholds in Tropical Trees: Variation Among Individuals, Species and Forests." *Journal of Tropical Ecology* 21: 307–315.
- Yang, X., A. L. Angert, P. A. Zuidema, et al. 2022. "The Role of Demographic Compensation in Stabilising Marginal Tree Populations in North America." *Ecology Letters* 25: 1676–1689.
- Yu, G., D. K. Smith, H. Zhu, Y. Guan, and T. T. Y. Lam. 2017. "Ggtree: An R Package for Visualization and Annotation of Phylogenetic Trees With Their Covariates and Other Associated Data." *Methods in Ecology and Evolution* 8: 28–36.
- Zanne, A. E., D. C. Tank, W. K. Cornwell, et al. 2014. "Three Keys to the Radiation of Angiosperms Into Freezing Environments." *Nature* 506: 89–92.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.