

EVALUATIONS OF LINK POLYNOMIALS AND RECENT CONSTRUCTIONS IN HEEGAARD FLOER THEORY

LARRY GU AND ANDREW MANION

ABSTRACT. Using a definition of Euler characteristic for fractionally-graded complexes based on roots of unity, we show that the Euler characteristics of Dowlin’s “ $\mathfrak{sl}(n)$ -like” Heegaard Floer knot invariants HFK_n recover both Alexander polynomial evaluations and $\mathfrak{sl}(n)$ polynomial evaluations at certain roots of unity for links in S^3 . We show that the equality of these evaluations can be viewed as the decategorified content of the conjectured spectral sequences relating $\mathfrak{sl}(n)$ homology and HFK_n .

1. INTRODUCTION

Ozsváth–Szabó’s theory of Heegaard Floer homology [OSz04b] is a flexible set of constructions yielding many types of invariants for low-dimensional manifolds. Even for knots and links in S^3 , the ideas of Heegaard Floer homology can be applied in several ways to produce a family of related invariants known collectively as knot Floer homology or HFK [OSz04a, Ras03]. The simplest variant, \widehat{HFK} as applied to (single-component) knots in S^3 , assigns to a knot K a bigraded vector space $\widehat{HFK}(K)$ (say over \mathbb{Q}) whose graded Euler characteristic is the Alexander polynomial $\Delta_K(t)$. Here we focus on more recent variants \widehat{HFK}_n (which we will call \overline{HFK}_n) and HFK_n , due to Dowlin [Dow18a], with relationships to Khovanov homology and $\mathfrak{sl}(n)$ homology more generally.

Reduced and unreduced $\mathfrak{sl}(n)$ homology [KR08a], like \widehat{HFK} , also assign bigraded vector spaces to knots K . Their graded Euler characteristics are the reduced and unreduced $\mathfrak{sl}(n)$ polynomials of K . The $\mathfrak{sl}(n)$ polynomials and the Alexander polynomial are all specializations of the two-variable HOMFLY-PT polynomial of K , leading to various relationships between the $\mathfrak{sl}(n)$ and Alexander polynomials at special values.

For the homology theories at the categorified level, one can often think of these relationships between knot polynomial evaluations as being categorified by certain spectral sequences that are known or conjectured to exist. For instance, the appearance of the $\mathfrak{sl}(n)$ polynomial as an evaluation of the HOMFLY-PT polynomial is categorified by Rasmussen’s spectral sequences [Ras15] from triply graded HOMFLY-PT homology [KR08b] to $\mathfrak{sl}(n)$ homology; the appearance of the Alexander polynomial as a HOMFLY-PT evaluation should be categorified by the conjectured spectral sequence from HOMFLY-PT homology to HFK [DGR06].

In general, given some construction or conjecture in the realm of $\mathfrak{sl}(n)$ homology or HFK , it is natural to ask “what does it categorify, if anything?”; in other words, “what is its decategorified content?”. Often this is something simpler than what one started with; for example, the identities relating $\mathfrak{sl}(n)$ polynomials and Alexander polynomials with the HOMFLY-PT polynomial are simpler than the known and conjectured spectral sequences from HOMFLY-PT homology to $\mathfrak{sl}(n)$ homology and HFK . Investigating the decategorified level can be an easy way to gain valuable information about the structure one expects at the categorified level.

For this reason, it is natural to ask about the decategorified content of Dowlin's conjectured spectral sequences [Dow18a, Conjecture 1.6] from reduced and unreduced $\mathfrak{sl}(n)$ homology to HFK (generalizing Rasmussen's conjecture for $\mathfrak{sl}(2)$, proved by Dowlin in [Dow18c]). The specific variants of HFK appearing in the conjectured spectral sequences are Dowlin's singly-graded variants \overline{HFK}_n and HFK_n for links (the first of these agrees with a grading collapse of \widehat{HFK} when applied to knots).

One complication is that as defined, \overline{HFK}_n and HFK_n are the homology of complexes whose differentials increase the single grading gr_n by n . The usual Euler characteristic formula, applied to such a complex, will not always be homotopy invariant. Instead, we divide the grading on \overline{HFK}_n and HFK_n by n , producing $\frac{1}{n}\mathbb{Z}$ -graded complexes whose differentials increase the grading by one. We introduce a natural generalization of the Euler characteristic to this setting (based on roots of unity and admitting an interpretation in terms of Grothendieck groups of triangulated categories); while it is plausible that our definition has appeared previously in the literature, we have not seen it elsewhere.

Theorem 1.1. *Let L be an ℓ -component link in S^3 and let $n \geq 2$. The Euler characteristic of $\frac{\text{gr}_n}{n}$ -graded $\overline{HFK}_n(L)$, in our sense, equals $e^{\pi i(1-\ell)/n} \Delta_L(t)|_{t^{1/2} = -e^{-\pi i/n}}$ where $\Delta_L(t)$ is the symmetric single-variable Alexander polynomial of L (a Laurent polynomial in $t^{1/2}$), and the Euler characteristic of $HFK_n(L)$ is zero. For $n = 1$, the Euler characteristics of both $\overline{HFK}_1(L)$ and $HFK_1(L)$ are 1 for all links L .*

Theorem 1.1 is new even for knots ($\ell = 1$), although the proof is relatively straightforward in this case (see Remark 6.5); when $\ell > 1$, a more involved spectral-sequence argument is required.

We introduce grading-modified versions $\overline{HFK}'_n(L)$ and $HFK'_n(L)$ of Dowlin's invariants such that the Euler characteristic of $\overline{HFK}'_n(L)$ equals $\Delta_L(t)|_{t^{1/2} = -e^{\pi i/n}}$. These n -dependent invariants are related to bigraded versions $\overline{HFK}'(L)$ and $HFK'(L)$ categorifying $\Delta_L(t)$ and zero respectively.¹

We propose $\overline{HFK}'_n(L)$ and $HFK'_n(L)$ as the E_∞ pages of Dowlin's conjectured spectral sequences, and support our proposal with Euler characteristic evidence. The E_2 pages of these conjectured spectral sequences should be reduced and unreduced $\mathfrak{sl}(n)$ homology with the bigrading collapsed to a single grading and divided by n as above. By analogy with HFK_n , we will refer² to the single collapsed grading as gr_n and its quotient by n as $\frac{\text{gr}_n}{n}$.

Theorem 1.2. *Let L be a link in S^3 . The Euler characteristic of the $\frac{\text{gr}_n}{n}$ -graded reduced $\mathfrak{sl}(n)$ homology of L , in our sense, equals the reduced $\mathfrak{sl}(n)$ polynomial of L evaluated at $q = e^{\pi i/n}$. The Euler characteristic of the $\frac{\text{gr}_n}{n}$ -graded unreduced $\mathfrak{sl}(n)$ homology of L equals the unreduced $\mathfrak{sl}(n)$ polynomial of L evaluated at $q = e^{\pi i/n}$.*

As with Theorem 1.1 in the case of knots, Theorem 1.2 is new, although the proof is straightforward given our definition of fractionally-graded Euler characteristic and known results about the bigraded Euler characteristic of $\mathfrak{sl}(n)$ homology.

¹From the representation-theoretic perspective, $\overline{HFK}'(L)$ categorifies the $U_q(\mathfrak{gl}(1|1))$ invariant of a link with one component cut open to form a $(1, 1)$ -tangle, while $HFK'(L)$ categorifies the $U_q(\mathfrak{gl}(1|1))$ invariant of a closed link which is zero.

²This usage of gr_n conflicts with the notation in [Ras15], where gr_n is used for what we call the quantum grading on $\mathfrak{sl}(n)$ homology (itself related in an n -dependent way to the quantum and horizontal gradings on HOMFLY-PT homology; see [Ras15]).

For $n \geq 2$ the polynomial evaluations appearing in Theorem 1.2 equal $\Delta_L(t)|_{t^{1/2} = -e^{\pi i/n}}$ and zero respectively; indeed, in the reduced case both evaluations are equal to $\overline{P}_L(-1, e^{\pi i/n})$ where $\overline{P}_L(a, q)$ is the reduced HOMFLY-PT polynomial of L , and similarly in the unreduced case. When $n = 1$ both evaluations are equal to 1. As we discuss below, we can view Dowlin's conjectured spectral sequences as categorifications of these equalities.

We situate these results in the context of Rasmussen's spectral sequences from HOMFLY-PT homology to $\mathfrak{sl}(n)$ homology and the conjectured spectral sequences from HOMFLY-PT homology to HFK , which fit with Dowlin's conjectured spectral sequences into a square as shown in [Dow18a, Figure 1]. We review the decategorified content of the known and conjectured spectral sequences starting at HOMFLY-PT homology, which we generalize to links in terms of our shifted gradings, and we add to Dowlin's square by labeling the edges with their decategorified content (see Figure 2). Examining the decategorified content along the possible paths in the square, in terms of link polynomial evaluations, reveals a compatibility that could be a sign of a more structured relationship between these spectral sequences at the categorified level.

Example 1.3. Dowlin computes HFK_n of the unknot in [Dow18a, Example 2.10]; the result is $\mathbb{Q}[U]/(U^n)$, and the generator $1 \in \mathbb{Q}[U]$ has gr_n equal to $1 - n$ (so it has $\frac{\text{gr}_n}{n}$ equal to $\frac{1}{n} - 1$). Dowlin writes that the graded Euler characteristic of this homology is $\frac{q^n - q^{-n}}{q - q^{-1}}$, agreeing with the $\mathfrak{sl}(n)$ polynomial of the unknot.

We propose that $\frac{q^n - q^{-n}}{q - q^{-1}}$ is the gr_n -graded Poincaré polynomial of this homology group, where the coefficient of q^i in the polynomial is the dimension of the homology in $\text{gr}_n = i$. In fact, HFK_n of the unknot is isomorphic to unreduced $\mathfrak{sl}(n)$ homology of the unknot, which is naturally bigraded; in this example, the homological component of the bigrading is zero and the intrinsic component agrees with gr_n . With respect to this bigrading, which makes sense on HFK_n of the unknot but not on HFK_n in general, it is indeed true that the graded Euler characteristic of HFK_n of the unknot is $\frac{q^n - q^{-n}}{q - q^{-1}}$.

However, here we are considering gr_n (divided by n), which makes sense on HFK_n of arbitrary links, as a homological grading. Since it is only a single grading, its Euler characteristic in our sense will be a single complex number (not necessarily an integer because the grading is fractional). A generator of the homology in $\text{gr}_n = k$ will contribute a term $e^{k\pi i/n}$ to the Euler characteristic by our definitions; the Euler characteristic for the unknot homology is thus

$$e^{\pi i(1/n-1)} + e^{\pi i(3/n-1)} + \cdots + e^{\pi i(1-3/n)} + e^{\pi i(1-1/n)}.$$

For $n = 1$ we get $e^{\pi i(0)} = 1$; for $n \geq 2$ the sum is zero, since the roots of unity are distributed symmetrically around the origin.

Remark 1.4. Here we see chain complexes with gradings by $\frac{1}{n}\mathbb{Z}$ (and $d^2 = 0$) categorifying evaluations of $\mathfrak{sl}(n)$ polynomials at $2n^{\text{th}}$ roots of unity. For categorification of these polynomials at roots of unity $e^{2\pi i/p}$ for p prime, complexes with $d^2 = 0$ are no longer suitable, and one often works with p -complexes satisfying $d^p = 0$ (see e.g. [Kho16, Qi14]). Combining these ideas, one could look for p -complexes with gradings by $\frac{1}{n}\mathbb{Z}$ categorifying evaluations of $\mathfrak{sl}(n)$ polynomials at pn^{th} roots of unity, although we are not aware of such complexes in the literature.

Remark 1.5. If one is only interested in categorifying e.g. $\overline{P}_{n,L}(e^{\pi i/n})$ using the ideas of this paper where $\overline{P}_{n,L}(q)$ is the $\mathfrak{sl}(n)$ polynomial, one does not need to use \overline{HFK}_n ; it suffices

to take a grading-collapse of reduced $\mathfrak{sl}(n)$ homology. However, \overline{HFK}_n is a more natural or minimal categorification of this evaluation; analogously, to categorify the Alexander polynomial one can take a grading collapse of HOMFLY-PT homology, but \overline{HFK} is a more minimal way to do it.

Remark 1.6. Let K be a knot. For $n = 2$ where a spectral sequence from Khovanov homology to HFK has been constructed by Dowlin [Dow18c], the equality $\overline{P}_{2,K}(i) = \Delta_K(-1)$ is familiar (both evaluations give the knot determinant) and is a sign of a deeper relationship between the representation theory of $U_q(\mathfrak{gl}(1|1))$ and $U_q(\mathfrak{gl}(2))$ at $q = i$; see [KS91, Section 1]. We do not know whether there is any similar story one can tell about the analogous equalities for $n > 2$, although the $n = 2$ case is special at least in that both $\mathfrak{gl}(1|1)$ and $\mathfrak{gl}(2)$ are defined using 2×2 matrices.

Organization. In Section 2 we define Euler characteristics for fractionally graded complexes and discuss spectral sequences. In Section 3 we review what we need about HOMFLY-PT polynomials, $\mathfrak{sl}(n)$ polynomials, and Alexander polynomials as well as HOMFLY-PT homology and $\mathfrak{sl}(n)$ homology (focusing on the gradings). In Section 4 we do the same for HFK while introducing bigrading-shifted versions of HFK theories adapted to the three variants of HOMFLY-PT homology. In Section 5 we recall the definitions of Dowlin's HFK_n invariants; in Section 6 we compute their fractionally-graded Euler characteristics, introduce grading-shifted variants of HFK_n , and prove Theorem 1.1. In Section 7 we prove Theorem 1.2. We also state a version of Dowlin's spectral sequence conjectures involving grading-shifted HFK_n , compute its decategorified content, and place it in the context of spectral sequences from HOMFLY-PT homology to $\mathfrak{sl}(n)$ homology and HFK .

Acknowledgments. We would like to thank Aaron Lauda for useful conversations and suggestions. A.M. was partially supported by NSF grant DMS-1902092 and Army Research Office W911NF2010075.

2. ALGEBRAIC PRELIMINARIES

2.1. Euler characteristics of fractionally-graded complexes. Following [Dow18a], we will work over \mathbb{Q} .

Definition 2.1. For $n \geq 1$, a $\frac{1}{n}\mathbb{Z}$ -graded complex C of \mathbb{Q} -vector spaces (or just a $\frac{1}{n}\mathbb{Z}$ -graded complex for short) is a $\frac{1}{n}\mathbb{Z}$ -graded \mathbb{Q} -vector space

$$C = \bigoplus_{\alpha \in \frac{1}{n}\mathbb{Z}} C_\alpha$$

equipped with a \mathbb{Q} -linear endomorphism d of degree $+1$ satisfying $d^2 = 0$.

The above definition is a special case of the definition of complexes with group-valued gradings. By contrast, Definitions 2.3 and 2.4 below are new as far as we are aware.

Remark 2.2. A $\frac{1}{n}\mathbb{Z}$ -graded complex is the same data as n ordinary complexes, one for each element of $(\frac{1}{n}\mathbb{Z})/\mathbb{Z}$. However, the examples of interest here more naturally give a $\frac{1}{n}\mathbb{Z}$ -graded complex than n ordinary complexes.

The category of $\frac{1}{n}\mathbb{Z}$ -graded complexes and homotopy classes of degree-zero chain maps is triangulated; the translation functor is degree shift downward by one. Furthermore, degree

shift downward by $\frac{1}{n}$ equips this triangulated category with an n^{th} root of its translation functor [1], that is, a functor $[\frac{1}{n}]$ such that $[\frac{1}{n}]$ composed n times equals [1]. We let $C[\alpha]$ denote C with its degrees shifted downward by $\alpha \in \frac{1}{n}\mathbb{Z}$, so that $(C[\alpha])_{\alpha'} = C_{\alpha'+\alpha}$.

Definition 2.3. Let \mathcal{C} be an (essentially small) triangulated category equipped with an n^{th} root $[\frac{1}{n}]$ of its translation functor [1]. Let $\zeta_n = e^{\pi i/n}$. We define the Grothendieck group $K_0(\mathcal{C})$ to be the quotient of the free $\mathbb{Z}[\zeta_n]$ -module spanned by isomorphism classes of objects of \mathcal{C} by the relations $X - Y + Z = 0$ for every distinguished triangle $X \rightarrow Y \rightarrow Z \rightarrow X[1]$ in \mathcal{C} , as well as

$$\left[X \left[\frac{1}{n} \right] \right] = \zeta_n^{-1} [X]$$

for all objects X of \mathcal{C} .

We can apply Definition 2.3 to the homotopy category H of finite-dimensional $\frac{1}{n}\mathbb{Z}$ -graded complexes (a full triangulated subcategory of the homotopy category of all $\frac{1}{n}\mathbb{Z}$ -graded complexes, preserved by the n^{th} root of the translation functor). The result is a free $\mathbb{Z}[\zeta_n]$ -module $K_0(H)$ of rank 1 spanned by $[\mathbb{Q}]$, where $[\mathbb{Q}]$ denotes the class of the complex that has \mathbb{Q} in degree zero and zero in all other degrees.

Definition 2.4. Let C be a finite-dimensional $\frac{1}{n}\mathbb{Z}$ -graded complex. The Euler characteristic $\chi(C)$ of C is the unique element of $\mathbb{Z}[\zeta_n]$ such that $[C] = \chi(C)[\mathbb{Q}]$ in $K_0(H)$. Explicitly,

$$\chi(C) = \sum_{\alpha \in \frac{1}{n}\mathbb{Z}} e^{\pi i \alpha} \dim_{\mathbb{Q}} C_{\alpha}.$$

Just as for ordinary Euler characteristics, we have $\chi(C) = \chi(H_*(C))$. We also recall the usual graded Euler characteristics for bigraded and triply-graded complexes.

Definition 2.5. Let $C = (\{C_{I,J} : I, J \in \mathbb{Z}\}, d)$ be a bigraded chain complex which is finite-dimensional in each I -degree, such that d has degree $(0, -1)$ or $(0, 1)$. The graded Euler characteristic of C is defined to be

$$\chi_u(C) = \sum_{I, J \in \mathbb{Z}} (-1)^J u^I \dim_{\mathbb{Q}} (C_{I,J}),$$

a formal Laurent series in a variable u . If the I -grading is valued in $\frac{1}{2}\mathbb{Z}$ rather than \mathbb{Z} , the same definition gives a formal Laurent series in $u^{1/2}$.

Similarly, let $C = (\{C_{I,J,K} : I, J, K \in \mathbb{Z}\}, d)$ be a triply graded chain complex which is finite-dimensional in each (I, J) -bidegree, such that d has degree $(0, 0, -1)$ or $(0, 0, 1)$. The graded Euler characteristic of C is defined to be

$$\chi_{u,v}(C) = \sum_{I, J, K \in \mathbb{Z}} (-1)^K u^I v^J \dim_{\mathbb{Q}} (C_{I,J,K}),$$

a formal Laurent series in variables u and v .

Remark 2.6. Rather than u and v , we will often use variable names corresponding to the gradings in question.




FIGURE 1. Links appearing in the HOMFLY-PT skein relation.

2.2. Spectral sequences.

Definition 2.7. A spectral sequence of $\frac{1}{n}\mathbb{Z}$ -graded complexes is a sequence of $\frac{1}{n}\mathbb{Z}$ -graded complexes $(E_r, d_r)_{r \geq 0}$ together with isomorphisms $H_*(E_r, d_r) \cong E_{r+1}$ for $r \geq 0$.

All spectral sequences in this paper have $d_r = 0$ for large enough r , so that for some $\frac{1}{n}\mathbb{Z}$ -graded vector space E_∞ we have $(E_r, d_r) = (E_\infty, 0)$ for large enough r .

Remark 2.8. It is built into the above definition that the differential d_r on each page of the spectral sequence has degree $+1$ with respect to the $\frac{1}{n}\mathbb{Z}$ grading.

Since $\chi(C) = \chi(H_*(C))$ for finite-dimensional $\frac{1}{n}\mathbb{Z}$ -graded chain complexes, if some page E_r of a spectral sequence as in Definition 2.7 is finite-dimensional then for any $r' \geq r$ we have $\chi(E_r) = \chi(E_{r'})$; in particular, $\chi(E_r) = \chi(E_\infty)$.

Remark 2.9. Suppose one has a spectral sequence of bigraded complexes as in Definition 2.5, such that each d_r has bidegree $(0, -1)$ or $(0, 1)$. Suppose that some page E_r is finite-dimensional in each j -degree; the same is then true for each page $E_{r'}$ for $r \geq r$, and it follows for the same reason as above that $\chi_u(E_r) = \chi_u(E_{r'})$. In particular, $\chi_u(E_r) = \chi_u(E_\infty)$. A similar equality for $\chi_{u,v}$ holds in the triply graded case, assuming each d_r has bidegree $(0, 0, -1)$ or $(0, 0, 1)$.

3. LINK POLYNOMIALS AND KHOVANOV–ROZANSKY HOMOLOGY

All links below are assumed to be oriented.

3.1. Link polynomials. The HOMFLY-PT polynomial $P_L(a, q)$ of a link L in S^3 [FYH⁺85, PT87] is defined by the skein relation

$$aP_{L_+}(a, q) - a^{-1}P_{L_-}(a, q) = (q - q^{-1})P_{L_0}(a, q)$$

(where L_+ , L_- , and L_0 are related near a crossing as in Figure 1) together with the HOMFLY-PT polynomial of the unknot as a normalization. We consider three variants:

- The reduced HOMFLY-PT polynomial $\overline{P}_L(a, q)$ has $\overline{P}_{\text{unknot}}(a, q) = 1$.
- The middle HOMFLY-PT polynomial $P_L^-(a, q)$ has $P_{\text{unknot}}^-(a, q) = \frac{-1}{q - q^{-1}}$.
- The unreduced HOMFLY-PT polynomial $P_L(a, q)$ has $P_{\text{unknot}}(a, q) = \frac{a - a^{-1}}{q - q^{-1}}$.

Remark 3.1. In these variables the middle and unreduced HOMFLY-PT “polynomials” are rational functions in general, although they are Laurent polynomials in a and $z = q - q^{-1}$.

We also consider two variants of the $\mathfrak{sl}(n)$ polynomial:

- The reduced $\mathfrak{sl}(n)$ polynomial is $\overline{P}_{n,L}(q) := \overline{P}_L(q^n, q)$.
- The unreduced $\mathfrak{sl}(n)$ polynomial is $P_{n,L}(q) := P_L(q^n, q)$.

Finally, if we let $\Delta_L(t)$ denote the (symmetric single-variable) Alexander polynomial of L , a Laurent polynomial in $t^{1/2}$, then we have

- $\overline{P}_L(1, t^{1/2}) = \Delta_L(t)$,
- $P_L^-(1, t^{1/2}) = \frac{\Delta_L(t)}{t^{-1/2} - t^{1/2}}$,
- $P_L(1, t^{1/2}) = 0$.

More relevant for us will be the following identities, which are consequences of the symmetries $\overline{P}_L(a, q) = \overline{P}(-a, -q)$, $P_L^-(a, q) = -P_L^-(a, -q)$, and $P_L(-a, -q) = P_L(a, q)$ of the HOMFLY-PT polynomials:

- $\overline{P}_L(-1, -t^{1/2}) = \Delta_L(t)$,
- $P_L^-(1, -t^{1/2}) = \frac{\Delta_L(t)}{t^{1/2} - t^{-1/2}}$,
- $P_L(-1, -t^{-1/2}) = 0$.

3.2. Khovanov–Rozansky homology.

3.2.1. *Gradings and Euler characteristics.* We briefly establish notation for the $\mathfrak{sl}(n)$ homology and HOMFLY-PT homology of Khovanov–Rozansky [KR08a, KR08b]; see also [Ras15]. Let $\overline{H}(L)$, $H^-(L)$, and $H(L)$ be the reduced, middle, and unreduced HOMFLY-PT homology of a link L in S^3 . In the notation of [Ras15], these variants of HOMFLY-PT homology (denoted there by $\overline{H}(L)$, $H(L)$, and $\widetilde{H}(L)$ respectively) have a \mathbb{Z} -grading gr_q (or just q), a $\frac{1}{2}\mathbb{Z}$ -grading gr_h , and a $\frac{1}{2}\mathbb{Z}$ -grading gr_v . Rasmussen also writes $i = \text{gr}_q$, $j = 2\text{gr}_h$, and $k = 2\text{gr}_v$; the value of $j - k$ is always even. We let

- $\text{gr}_A = 2\text{gr}_h = j$,
- $\text{gr}_Q = \text{gr}_q = i$,
- $\text{gr}_H = \text{gr}_v - \text{gr}_h = \frac{k-j}{2}$,

each of which is a grading by \mathbb{Z} on the above three variants of HOMFLY-PT homology.

Remark 3.2. While h in gr_h stands for horizontal, H in gr_H stands for homological.

Each variant of HOMFLY-PT homology is finite-dimensional in each $(\text{gr}_A, \text{gr}_Q)$ -bidegree, so the following proposition makes sense.

Proposition 3.3 (cf. Theorem 2.11, Section 2.8 of [Ras15]). *For a link L in S^3 , we have:*

- $\chi_{a,q}(\overline{H}(L)) = \overline{P}_L(a, q)$,
- $\chi_{a,q}(H^-(L)) = P_L^-(a, q)$,
- $\chi_{a,q}(H(L)) = P_L(a, q)$.

Now let $\overline{H}_n(L)$ and $H_n(L)$ be the reduced and unreduced $\mathfrak{sl}(n)$ homology of a link L in S^3 ; the reduced homology $H_n(L)$ also depends on a choice of component of L . Both variants of $\mathfrak{sl}(n)$ homology have \mathbb{Z} -gradings $\text{gr}_{Q,n}$ and gr_H ; in [Ras15, Section 2.9] these gradings are called gr_n and gr_v respectively, while in [Ras15, Section 5] they are called gr'_n and gr_- (see [Ras15, Proposition 5.14]).

Proposition 3.4 (cf. Theorem 2.16 of [Ras15]). *Write χ_q for the $\text{gr}_{Q,n}$ -graded Euler characteristic, with gr_H treated as the homological grading. For a link L in S^3 , we have:*

- $\chi_q(\overline{H}_n(L)) = \overline{P}_{n,L}(q)$,
- $\chi_q(H_n(L)) = P_{n,L}(q)$.

3.2.2. *Rasmussen's spectral sequences.* In [Ras15], Rasmussen constructs spectral sequences with E_2 page $\overline{H}(L)$ (respectively $H(L)$) and E_∞ page $\overline{H}_n(L)$ (respectively $H_n(L)$) for $n \geq 1$. As discussed in [Ras15, beginning of Section 5], the differentials on each page have $\text{gr}_{Q,n} = 0$ and $\text{gr}_H = 1$ where $\text{gr}_{Q,n}$ on HOMFLY-PT homology is defined by $\text{gr}_{Q,n} = \text{gr}_Q + n \text{gr}_A$. Thus, each page gets a bigrading as the homology of the previous page, and the induced bigrading on the E_∞ page agrees with $(\text{gr}_{Q,n}, \text{gr}_H)$ on $\mathfrak{sl}(n)$ homology.

These spectral sequences give equalities of Euler characteristics $\chi_q(\overline{H}(L)) = \chi_q(\overline{H}_n(L))$ and $\chi_q(H(L)) = \chi_q(H_n(L))$, where we are viewing $\overline{H}(L)$ and $H(L)$ as bigraded by $(\text{gr}_{Q,n}, \text{gr}_H)$ and χ_q denotes the $\text{gr}_{Q,n}$ -graded Euler characteristic. As in the proof of [Ras15, Lemma 5.4], we have $\chi_q(\overline{H}(L)) = \overline{P}_L(q^n, q)$; similarly, we have $\chi_q(H(L)) = P_L(q^n, q)$. Thus, applying Euler characteristics to these spectral sequences recovers the usual identity of the $\mathfrak{sl}(n)$ polynomial with an evaluation of the HOMFLY-PT polynomial. We view these identities as the “decategorified content” of Rasmussen's spectral sequences; in other words, we view the spectral sequences as categorifications of these identities.

4. KNOT FLOER HOMOLOGY

4.1. **The master complex.** Let L be a link in S^3 ; let \mathcal{H} be a multi-pointed Heegaard diagram for L (versions of the below theories can be defined for links in more general 3-manifolds but we restrict attention to links in S^3 here). Write

$$\{z_1, w_1, \dots, z_m, w_m\}$$

for the set of basepoints in \mathcal{H} . We assume that \mathcal{H} is equipped with the appropriate analytic data such that the knot Floer homology “master complex” $CFK_{U,V}(L)$, a finitely generated bigraded free module over $\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]$ with an endomorphism $\partial_{U,V}$ satisfying

$$\partial_{U,V}^2 = \sum_{i=1}^m (U_{a(i)} - U_{b(i)}) V_i,$$

is defined (here $a(i)$ denotes the index of the unique w basepoint in the same component as z_i of the Heegaard surface with alpha curves removed, and similarly for $b(i)$ and beta curves). See [Zem19, Dow18a] for more details on the master complex; the differential counts holomorphic disks whose domains are allowed to have nonzero multiplicity at any type of basepoint (w and/or z), and the contribution of a domain to the differential is weighted by powers of U_i determined by the multiplicities at w_i basepoints as well as powers of V_i determined by the multiplicities at z_i basepoints.

The two gradings on $CFK_{U,V}(L)$ are called the Alexander grading gr_T (a grading by $\frac{1}{2}\mathbb{Z}$ in general) and the Maslov grading gr_M (a grading by \mathbb{Z}); our conventions for these gradings follow [OSz04a, OSz08]. The variables U_i have $\text{gr}_T = -1$ and $\text{gr}_M = -2$, the variables V_i have $\text{gr}_T = 1$ and $\text{gr}_M = 0$, and $\partial_{U,V}$ has $\text{gr}_T = 0$ and $\text{gr}_M = -1$.

In particular, to fix the absolute Maslov grading, one can work with a Heegaard diagram \mathcal{H} whose number m of z and w basepoints is equal to the number ℓ of components of L . Then the homology of

$$CFK_{U,V}(L) \otimes_{\mathbb{Q}[U_1, V_1, \dots, U_\ell, V_\ell]} \frac{\mathbb{Q}[U_1, V_1, \dots, U_\ell, V_\ell]}{(U_1, V_1 - 1, \dots, V_\ell - 1)},$$

³While it would be more accurate to write $CFK_{U,V}(\mathcal{H})$, it will help avoid confusion with reduced and unreduced versions of knot Floer homology below to write $CFK_{U,V}(L)$, with the \mathcal{H} dependence left implicit.

which is a complex ($\partial^2 = 0$) with a single grading by gr_M , computes $\widehat{HF}(S^3) \cong \mathbb{Q}$ (see the discussion after Theorem 4.4 of [OSz08]), while the homology of

$$CFK_{U,V}(L) \otimes_{\mathbb{Q}[U_1, V_1, \dots, U_\ell, V_\ell]} \frac{\mathbb{Q}[U_1, V_1, \dots, U_\ell, V_\ell]}{(U_1, \dots, U_\ell, V_1 - 1, \dots, V_\ell - 1)}$$

computes $\widehat{HF}(\#^{\ell-1}(S^2 \times S^1)) \cong \wedge^* V$ where V is a vector space of dimension $\ell - 1$. As mentioned in [OSz08, proof of Theorem 1.1], in the grading conventions of that paper the top-dimensional generator of $\wedge^* V$ corresponds to the generator of $\widehat{HF}(S^3) \cong \mathbb{Q}$. By the discussion after [OSz08, Theorem 1.2], the absolute Maslov grading on CFK is fixed so that the top-dimensional generator of $\wedge^* V$ has Maslov degree zero, so we can equivalently say that the generator of $\widehat{HF}(S^3) \cong \mathbb{Q}$ has degree zero. The absolute Alexander grading is fixed by symmetry. That is, fix the absolute Alexander grading gr_T such that for all $n \in \mathbb{Z}$ and

$$\widehat{CFK}(L) := CFK_{U,V}(L) \otimes_{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]} \frac{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]}{(V_1, \dots, V_m, U_{i_1}, \dots, U_{i_\ell})},$$

$\widehat{CFK}(L)$ satisfies $\dim_{\mathbb{Q}}(\widehat{CFK}(L)_{\text{gr}_T=n}) = \dim_{\mathbb{Q}}(\widehat{CFK}(L)_{\text{gr}_T=-n})$.

4.2. Other bigraded variants of HFK . The following complexes are all derived from the master complex and satisfy $\partial^2 = 0$. We focus on bigraded versions of HFK ; there are also multi-graded versions as in [OSz08]. Let L be an ℓ -component link in S^3 ; when we mention knot Floer complexes for links, the dependence on a choice of Heegaard diagram for L (say with basepoints $\{z_1, w_1, \dots, z_m, w_m\}$) is implicit.

Definition 4.1. The bigraded complex $\widetilde{CFK}(L)$ is

$$CFK_{U,V}(L) \otimes_{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]} \mathbb{Q}.$$

Definition 4.2. Assume that, in our Heegaard diagram \mathcal{H} representing L , we are given some choice of basepoints (z_{i_j}, w_{i_j}) on each component L_j of L . The bigraded complex $\widehat{CFK}(L)$ is

$$CFK_{U,V}(L) \otimes_{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]} \frac{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]}{(V_1, \dots, V_m, U_{i_1}, \dots, U_{i_\ell})}.$$

Definition 4.3. Assume that L is equipped with a distinguished component and that the basepoints z_m, w_m of the Heegaard diagram \mathcal{H} representing L lie on the distinguished component of L . The bigraded complex $\overline{CFK}(L)$ is

$$CFK_{U,V}(L) \otimes_{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]} \frac{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]}{(V_1, \dots, V_m, U_m)}.$$

In general, the homotopy type of $\overline{CFK}(L)$ depends on the choice of distinguished component. However, the choice of basepoints for the same component give homotopy-equivalent complexes (see Dowlin [Dow18a, Corollary 2.15]). Following [Dow18a, Definition 2.19] and in analogy to $\mathfrak{sl}(n)$ homology, we will refer to $\overline{CFK}(L)$ as the reduced knot Floer complex of L (with its distinguished component) and its homology as the reduced knot Floer homology of L .

Definition 4.4. The bigraded complex $CFK^-(L)$ is

$$CFK_{U,V}(L) \otimes_{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]} \frac{\mathbb{Q}[U_1, V_1, \dots, U_m, V_m]}{(V_1, \dots, V_m)}.$$

Definition 4.5. [cf. Section 2.6 of [Dow18a]] Let L' be the disjoint union of L with a split unknot; we choose the unknot component as a distinguished component for L' , and we assume that the only basepoints of the diagram \mathcal{H}' we choose to represent L' that lie on the distinguished component of L' are the final pair $(z_{m'}, w_{m'})$ of basepoints. We define

$$CFK(L) := t^{-1/2} \overline{CFK}(L')$$

where $t^{-1/2}$ denotes a downward shift by $\frac{1}{2}$ in the Alexander grading gr_T . Following [Dow18a], we will refer to $CFK(L)$ as the unreduced knot Floer complex of L and its homology as the unreduced knot Floer homology of L .

Remark 4.6. The use of a split unknot to define unreduced HFK appears in Baldwin–Levine–Sarkar [BLS17], although these authors use the term “unreduced HFK ” for a slightly different theory.

The homology of each of these complexes is an invariant of L (equipped with a distinguished component in Definition 4.3) and will be denoted by $\widehat{HFK}(L)$, $\widehat{HFK}(L)$, etc. Each of the above bigraded versions of HFK is finite-dimensional in each Alexander degree.

Remark 4.7. The complex $\overline{CFK}(L)$ and its homology appear to be less common in the literature; we use $\overline{(\cdot)}$ to match the notation of HOMFLY-PT homology and $\mathfrak{sl}(n)$ homology, although it is possible that our use of the notation $\overline{HFK}(L)$ conflicts with uses of this notation elsewhere.

4.3. Graded Euler characteristics.

Proposition 4.8. *For a link L in S^3 , we have:*

- $\chi_t(\widehat{HFK}(L)) = (t^{-1/2} - t^{1/2})^{\ell-1} \Delta_L(t) = (-1)^{\ell-1} t^{\frac{\ell-1}{2}} (1 - t^{-1})^{\ell-1} \Delta_L(t)$,
- $\chi_t(\overline{HFK}(L)) = (-1)^{\ell-1} t^{\frac{\ell-1}{2}} \Delta_L(t)$,
- $\chi_t(HFK^-(L)) = (-1)^{\ell-1} t^{\frac{\ell-1}{2}} \frac{\Delta_L(t)}{1-t^{-1}}$,
- $\chi_t(HFK(L)) = 0$.

Proof. The first claim follows from [OSz04a, equation (1)] and [OSz08, Theorem 1.1].⁴ The second claim follows from the first because the chain groups in $\overline{CFK}(L)$ are free modules over polynomial rings in $\ell - 1$ more variables than the corresponding chain groups in $\widehat{CFK}(L)$; the graded Euler characteristic of a polynomial ring in one of these variables (with degrees $A = -1$ and $M = -2$) is $\frac{1}{1-t^{-1}}$. The third claim follows similarly; the fourth claim follows from the second claim along with the fact that $\Delta_L(t)$ vanishes on split links. \square

Remark 4.9. For an ℓ -component link L in S^3 , the single-variable and multi-variable

Alexander polynomials of L satisfy the relation $\Delta_L(t) \doteq \begin{cases} \Delta_L^{\text{multi}}(t) & \ell = 1 \\ \Delta_L^{\text{multi}}(t, \dots, t)(1-t) & \ell > 1 \end{cases}$,

where \doteq means equality up to multiplication by a unit in $\mathbb{Z}[t, t^{-1}]$ (see [Kaw96, Proposition 7.3.10(1)]). This relation explains why, unlike in [OSz08, equations (1) and (2)], we do not need to treat $\ell = 1$ and $\ell > 1$ separately in Proposition 4.8.

Definition 4.10. To more closely match the three variants of HOMFLY-PT homology, we introduce grading-shifted variants $\overline{HFK}'(L)$, $(HFK^-)'(L)$, and $HFK'(L)$ of knot Floer

⁴Note that [OSz08, equation (1)], when proved in [OSz08, Proposition 9.1], is stated with a \pm sign.

homology. We first replace the Alexander and Maslov degrees of $\overline{HFK}(L)$, $HFK^-(L)$, and $HFK(L)$ by their negatives, so that the variable U_i now has $\text{gr}_A = 1$ and $\text{gr}_M = 2$ and the differentials on CFK complexes now have Maslov degree $+1$. In the Euler characteristic computations of Proposition 4.8, t gets replaced by t^{-1} ; note that $\Delta_L(t^{-1}) = (-1)^{\ell-1} \Delta_L(t)$. We then make the following shifts:

- For $\overline{HFK}'(L)$, we shift the Alexander grading on grading-reversed $\overline{HFK}(L)$ upward by $\frac{\ell-1}{2}$. We have

$$\chi_t(\overline{HFK}'(t)) = \Delta_L(t).$$

- For $(HFK^-)'(L)$, we shift the Alexander grading on grading-reversed $HFK^-(L)$ upward by $\frac{\ell}{2}$; we also shift the Maslov grading upward by 1. We have

$$\chi_t((HFK^-)'(L)) = \frac{\Delta_L(t)}{t^{1/2} - t^{-1/2}}.$$

- For $HFK'(L)$, we shift the Alexander grading on grading-reversed $HFK(L)$ upward by $\frac{\ell-1}{2}$ (recall that $HFK(L)$ already had an Alexander grading shift in Definition 4.5). We have $\chi_t(HFK'(L)) = 0$.

4.4. Conjectured spectral sequences from HOMFLY-PT homology to HFK . In [DGR06], Dunfield–Gukov–Rasmussen conjectured the existence of spectral sequences from $\overline{H}(K)$ to $\widehat{HFK}(K)$ for knots K in S^3 . Manolescu [Man14] gives a similar conjecture for HFK^- and the middle HOMFLY-PT homology. Dowlin [Dow18a] conjectures spectral sequences from $\overline{H}(L)$ to $\widehat{HFK}(L)$ and from $H(L)$ to $HFK(L)$ for all links in S^3 . We believe a spectral sequence from $\overline{H}(L)$ to $\overline{HFK}(L)$ is more plausible for links, so we will state the following version of these spectral sequence conjectures.

Conjecture 4.11. *Let L be a link in S^3 . Ignoring gradings at first, there are spectral sequences with:*

- E_2 page $\overline{H}(L)$ and E_∞ page $\overline{HFK}(L)$;
- E_2 page $H^-(L)$ and E_∞ page $HFK^-(L)$;
- E_2 page $H(L)$ and E_∞ page $HFK(L)$.

Moreover, such sequences are given by the construction of Manolescu [Man14, Theorem 1.1], which is known to give E_∞ pages recovering HFK .

Remark 4.12. In [Dow18b], Dowlin identifies the E_1 page of the spectral sequence from [Man14, Theorem 1.1] with the appropriate sum of HOMFLY-PT complexes for singular resolutions of L ; it remains to identify the E_2 page with HOMFLY-PT homology for links with nonsingular crossings.

Manolescu [Man14, Section 4] discusses the grading properties of his conjectured spectral sequences from HOMFLY-PT homology to HFK in detail; we will rephrase some of his discussion in terms of the grading-shifted variants of HFK from Definition 4.10. We define a $\frac{1}{2}\mathbb{Z}$ -grading gr_T and a \mathbb{Z} -grading gr_M on each of the variants of HOMFLY-PT homology by

- $\text{gr}_T = \frac{\text{gr}_Q}{2}$,
- $\text{gr}_M = \text{gr}_A + \text{gr}_Q + \text{gr}_H$

(Manolescu includes constant grading-shift terms in the above formulas but here we incorporate the grading shifts into HFK ; he also has negative signs since, unlike us, he has not multiplied the Alexander and Maslov gradings on HFK by -1). The differential d_r on the E_r page of Manolescu's conjectured sequences has $\text{gr}_A = 2 - 2r$, $\text{gr}_Q = 0$, and $\text{gr}_H = 2r - 1$. Thus, d_r has $\text{gr}_T = 0$ and $\text{gr}_M = 1$. Writing $\overline{H}(L) = \bigoplus_{i,j,k \in \mathbb{Z}} \overline{H}^{i,j,k}(L)$ as in [Ras15] (and similarly for the other versions), we equivalently have $\text{gr}_T = i/2$ and $\text{gr}_M = i + j/2 + k/2$.

Conjecture 4.13. *Let L be a link in S^3 . There are spectral sequences with each page bigraded by $(\text{gr}_T, \text{gr}_M)$, such that the differential on each page has $(\text{gr}_T, \text{gr}_M) = (0, 1)$ and each page is the bigraded homology of the previous page, and with*

- E_2 page $\overline{H}(L)$ and E_∞ page $\overline{HFK}'(L)$;
- E_2 page $H^-(L)$ and E_∞ page $(HFK^-)'(L)$;
- E_2 page $H(L)$ and E_∞ page $HFK'(L)$

as bigraded vector spaces.

These spectral sequences would give equalities of Euler characteristics

$$\chi_t(\overline{H}(L)) = \chi_t(\overline{HFK}'(L)), \quad \chi_t(H^-(L)) = \chi_t((HFK^-)'(L)), \quad \chi_t(H(L)) = \chi_t(HFK'(L))$$

where χ_T denotes the gr_T -graded Euler characteristic. We have

$$\begin{aligned} \chi_t(\overline{H}(L)) &= \sum_{I \in \frac{1}{2}\mathbb{Z}, J \in \mathbb{Z}} (-1)^J t^I \dim_{\mathbb{Q}} (\overline{H}(L)_{\text{gr}_T=I, \text{gr}_M=J}) \\ &= \sum_{i,j,k \in \mathbb{Z}} (-1)^{i+j/2+k/2} t^{i/2} \dim_{\mathbb{Q}} (\overline{H}^{i,j,k}(L)) \\ &= \left(\sum_{i,j,k \in \mathbb{Z}} a^j q^i (-1)^{(k-j)/2} \dim_{\mathbb{Q}} (\overline{H}^{i,j,k}(L)) \right) \Big|_{a=-1, q=-t^{1/2}} \\ &= \overline{P}_L(-1, -t^{1/2}). \end{aligned}$$

Similarly,

$$\chi_t(H^-(L)) = P_L^(-1, -t^{1/2}), \quad \chi_t(H(L)) = P_L(-1, -t^{1/2}) = 0.$$

Thus, these conjectured spectral sequences can be viewed as categorifications of the three equalities involving Alexander polynomials and HOMFLY-PT polynomial evaluations (with $a = -1$) at the end of Section 3.1.

5. DOWLIN'S HFK_n INVARIANTS

We now consider two versions of HFK defined by Dowlin [Dow18a], applied to links in S^3 rather than more general 3-manifolds. Rather than bigradings, these versions will have single gradings by $\frac{1}{n}\mathbb{Z}$ in our conventions.

Let L be a link in S^3 , represented by a Heegaard diagram \mathcal{H} as in Section 4.1 with basepoints $\{z_1, w_1, \dots, z_m, w_m\}$. Following Dowlin, for $n \geq 1$ we consider a collapse gr_n of the bigrading on $CFK_{U,V}(L)$ defined by

$$\text{gr}_n = -n \text{gr}_M + 2(n-1) \text{gr}_T.$$

We divide gr_n by n to get

$$\frac{\text{gr}_n}{n} = -\text{gr}_M + 2 \left(1 - \frac{1}{n}\right) \text{gr}_T$$

which is valued in $\frac{1}{n}\mathbb{Z}$ even for half-integral values of gr_T . The variables U_i have $\frac{\text{gr}_n}{n} = \frac{2}{n}$, the variables V_i have $\frac{\text{gr}_n}{n} = 2 - \frac{2}{n}$, and $\partial_{U,V}$ has $\frac{\text{gr}_n}{n} = 1$.

Definition 5.1 (cf. Definition 2.19 of [Dow18a]). Assume that L is equipped with a distinguished component and that the basepoints z_m, w_m of the Heegaard diagram \mathcal{H} representing L lie on the distinguished component. The $\frac{1}{n}\mathbb{Z}$ -graded complex $\overline{CFK}_n(L)$ is

$$CFK_{U,V}(L) \otimes_{\mathbb{Q}[U_1, \dots, U_m, V_1, \dots, V_m]} \frac{\mathbb{Q}[U_1, \dots, U_m, V_1, \dots, V_m]}{\left(V_i - \frac{U_{a(i)}^n - U_{b(i)}^n}{U_{a(i)} - U_{b(i)}} : 1 \leq i \leq m-1\right) + (U_m, V_m)}$$

where $a(i)$ and $b(i)$ are defined as in Section 4.1. The grading is given by $\frac{\text{gr}_n}{n}$; note that $\frac{U_{a(i)}^n - U_{b(i)}^n}{U_{a(i)} - U_{b(i)}}$ equals the telescoping sum $U_{a(i)}^{n-1} + U_{a(i)}^{n-2}U_{b(i)} + \dots + U_{b(i)}^{n-1}$, which (like V_i) has $\frac{\text{gr}_n}{n} = 2 - \frac{2}{n}$.

Definition 5.2 (cf. Definition 2.5 of [Dow18a]). Let L' be the disjoint union of L with a split unknot, and choose the unknot component to be distinguished. As in Definition 4.5, we assume that the only basepoints of the diagram \mathcal{H}' we choose to represent L' that lie on the distinguished component of L' are the final pair $(z_{m'}, w_{m'})$ of basepoints. We define

$$CFK_n(L) := \overline{CFK}_n(L')[1 - 1/n];$$

note that a downward shift by $\frac{1}{2}$ in gr_T as in [Dow18a, Section 2.2] produces a downward shift by $1 - \frac{1}{n}$ in $\frac{\text{gr}_n}{n} = -\text{gr}_M + 2(1 - 1/n) \text{gr}_T$.

We write ∂_n for the differential on either variant of CFK_n ; it satisfies $\partial_n^2 = 0$. When $n = 1$, the complex $\overline{CFK}_1(L)$ computes $\widehat{HF}(S^3)$ (see Section 4.1), so its homology is \mathbb{Q} in $\text{gr}_M = 0$ (and thus $\frac{\text{gr}_n}{n} = 0$) and zero in other degrees; see [Dow18a, Lemma 5.2]. It follows that $HFK_1(L)$ is also \mathbb{Q} in degree 0 and zero in other degrees.

Since the tensor product (after annihilating the final pair of variables) sets each V_i variable equal to a polynomial in the U_i variables while imposing no further relations on the U_i variables, the complexes $\overline{CFK}_n(L)$ and $CFK_n(L)$ are free over $\mathbb{Q}[U_1, \dots, U_{m-1}]$ and $\mathbb{Q}[U_1, \dots, U_{m'-1}]$ respectively. Their homology groups $\overline{HFK}_n(L)$ (respectively $HFK_n(L)$) depend only on L with its distinguished component (respectively, L) and are finite-dimensional over \mathbb{Q} as shown in [Dow18a].

Remark 5.3. In [Dow18a], Dowlin uses the notation $\widehat{HFK}_n(L)$ to refer to what we call $\overline{HFK}_n(L)$; however, in [Dow18c], \widehat{HFK}_n (at least for $n = 2$) is given a different definition which is closer to Definition 4.2 for $\widehat{HFK}(L)$.

6. THE EULER CHARACTERISTIC OF HFK_n

Let L be an ℓ -component link in S^3 equipped with a distinguished component; in this section we compute the Euler characteristics of $\overline{HFK}_n(L)$ and $HFK_n(L)$.

For simplicity, assume we are working with a Heegaard diagram \mathcal{H} for L that has exactly 2ℓ basepoints. Let $R = \mathbb{Q}[U_1, \dots, U_{\ell-1}]$, a $\frac{1}{n}\mathbb{Z}$ -graded ring where U_i has degree $\frac{2}{n}$ as in Section 5,

and let K be the $\frac{1}{n}\mathbb{Z}$ -graded Koszul complex

$$K = \bigotimes_{i=1}^{\ell-1} \left(R \left[1 - \frac{2}{n} \right] \xrightarrow{U_i} R \right)$$

where the tensor products are over R .

Lemma 6.1. *Let $n \geq 2$. There exists a spectral sequence with each page graded by $\frac{1}{n}\mathbb{Z}$, with differentials of degree $+1$ such that each page is the $\frac{1}{n}\mathbb{Z}$ -graded homology of the previous page, and with E_1 page $\widehat{HFK}_n(L) \otimes_R K$ and E_∞ page $\widehat{HFK}(L)$ as $\frac{1}{n}\mathbb{Z}$ -graded vector spaces.*

Proof. The complex $\widehat{CFK}_n(L) \otimes_R K$ can be viewed as a cube of dimension $\ell - 1$ in which each vertex is a copy of $\widehat{CFK}_n(L)$. We equip $\widehat{CFK}_n(L) \otimes_R K$ with a filtration such that every oriented edge of this cube increases the filtration level by 1. Then the differential d on $\widehat{CFK}_n(L) \otimes_R K$ can be decomposed as $d = d_0 + d_1$, where d_0 is the differential on each copy of $\widehat{CFK}_n(L)$ and d_1 comes from the differential on K .

From this filtration, we get a spectral sequence whose E_1 page is $(\widehat{HFK}_n(L) \otimes_R K, (d_1)_*)$. The spectral sequence converges because there are only finitely many nontrivial filtration levels, and the E_∞ page is $\widehat{HFK}(L)$; indeed, we claim that the total complex $\widehat{CFK}_n(L) \otimes_R K$ has a contractible subcomplex such that the quotient by this subcomplex is $\widehat{CFK}(L)$.

To see this, note that R has a basis of monomials and we get a corresponding basis for K (which is a direct sum of copies of R). Let K' denote the span of all basis elements of K except for the monomial 1 in the unique R -summand of K having no outward-pointing arrows in the differential on K (this is the R -summand with no degree shift). As a complex, we have $K \cong K' \oplus \mathbb{Q}$, and since the homology of K is the homology of the \mathbb{Q} summand in this decomposition (namely \mathbb{Q}), the complex K' is contractible.

The subcomplex of $\widehat{CFK}_n(L) \otimes_R K$ we consider is $\widehat{CFK}_n(L) \otimes_R K'$; by construction we have

$$\frac{\widehat{CFK}_n(L) \otimes_R K}{\widehat{CFK}_n(L) \otimes_R K'} \cong \widehat{CFK}(L).$$

To see that $\widehat{CFK}_n(L) \otimes_R K'$ is contractible, define a filtration on $\widehat{CFK}_n(L) \otimes_R K'$ as the Maslov grading plus the above-defined filtration on $\widehat{CFK}_n(L) \otimes_R K$. Then components of the differential internal to $\widehat{CFK}_n(L)$ decrease filtration level by 1 while components of the differential coming from K' preserve the filtration level. The associated graded complex to the filtered complex $\widehat{CFK}_n(L) \otimes_R K'$ is isomorphic to a direct sum of copies of K' , so it is contractible; it follows that $\widehat{CFK}_n(L) \otimes_R K'$ is contractible. \square

Proof of Theorem 1.1. Since $\widehat{HFK}_n(L)$ is finitely generated over \mathbb{Q} , the same is true for $\widehat{HFK}_n(L) \otimes_R K$. Thus, the spectral sequence of Lemma 6.1 gives an equality between the $\frac{1}{n}\mathbb{Z}$ -graded Euler characteristics of $\widehat{HFK}_n(L) \otimes_R K$ and $\widehat{HFK}(L)$. The Euler characteristics of $\widehat{HFK}_n(L) \otimes_R K$ and $\widehat{HFK}_n(L)$ are related by

$$\chi(\widehat{HFK}_n(L) \otimes_R K) = (1 - e^{2\pi i/n})^{\ell-1} \chi(\widehat{HFK}_n(L)),$$

so if $n \geq 2$ we get

$$\chi(\widehat{HFK}_n(L)) = (1 - e^{2\pi i/n})^{1-\ell} \chi(\widehat{HFK}(L)).$$

Using Proposition 4.8, we can compute the $\frac{\text{gr}_n}{n}$ -graded Euler characteristic of $\widehat{HFK}(L)$ as follows:

$$\begin{aligned}
\chi(\widehat{HFK}(L)) &= \sum_{\alpha \in \frac{1}{n}\mathbb{Z}} e^{\pi i \alpha} \dim_{\mathbb{Q}} \left(\widehat{HFK}(L)_{\frac{\text{gr}_n}{n}=\alpha} \right) \\
&= \sum_{I \in \frac{1}{2}\mathbb{Z}, J \in \mathbb{Z}} e^{\pi i (-J+2(1-1/n)I)} \dim_{\mathbb{Q}} \left(\widehat{HFK}(L)_{\text{gr}_T=I, \text{gr}_M=J} \right) \\
&= \left(\sum_{I \in \frac{1}{2}\mathbb{Z}, J \in \mathbb{Z}} (-1)^J t^I \dim_{\mathbb{Q}} \left(\widehat{HFK}(L)_{\text{gr}_T=I, \text{gr}_M=J} \right) \right) \Big|_{t^{1/2}=e^{\pi i (1-1/n)}=-e^{-\pi i /n}} \\
&= \left((-1)^{\ell-1} t^{\frac{\ell-1}{2}} (1-t^{-1})^{\ell-1} \Delta_L(t) \right) \Big|_{t^{1/2}=-e^{-\pi i /n}} \\
&= e^{\pi i (1-\ell)/n} (1-e^{2\pi i /n})^{\ell-1} \Delta_L(t) \Big|_{t^{1/2}=-e^{-\pi i /n}}.
\end{aligned}$$

It follows that for $n \geq 2$, the $\frac{\text{gr}_n}{n}$ -graded Euler characteristic of $\overline{HFK}_n(L)$ is

$$e^{\pi i (1-\ell)/n} \Delta_L(t) \Big|_{t^{1/2}=-e^{-\pi i /n}}.$$

Since $HFK_n(L)$ is defined as a grading shift of \overline{HFK}_n of the disjoint union of L with a split unknot, and the Alexander polynomial vanishes on split links, we see that if $n \geq 2$, the $\frac{\text{gr}_n}{n}$ -graded Euler characteristic of $HFK_n(L)$ is zero for all links L . When $n = 1$, we have

$$\chi(\overline{HFK}_1(L)) = \chi(HFK_1(L)) = 1$$

for all links L . \square

Definition 6.2. As in Definition 4.10, we define grading-shifted variants $\overline{HFK}'_n(L)$ and $HFK'_n(L)$ of $\overline{HFK}_n(L)$ and $HFK_n(L)$. Starting with bigradings on $\overline{CFK}_n(L)$ and $CFK_n(L)$ corresponding to the bigradings on $\overline{CFK}'(L)$ and $CFK'(L)$, the differentials ∂_n have degree $+1$ with respect to

$$\frac{\text{gr}_n}{n} := \text{gr}_M - 2 \left(1 - \frac{1}{n} \right) \text{gr}_T$$

(note that since we still want $+1$ differentials on $\frac{1}{n}\mathbb{Z}$ -graded complexes, this is the negative of the earlier definition of $\frac{\text{gr}_n}{n}$ in terms of gr_T and gr_M). We define $\overline{CFK}'_n(L)$ to be $\overline{CFK}'(L)$ with grading given by $\frac{\text{gr}_n}{n}$ and differential given by ∂_n ; we define $CFK'_n(L)$ similarly.

Remark 6.3. Starting from $\overline{HFK}(L)$, we negated both gradings and shifted the Alexander grading upward by $\frac{\ell-1}{2}$ to get $\overline{HFK}'(L)$, then applied the collapse $\text{gr}_M - 2(1 - 1/n) \text{gr}_T$ to get the grading on $\overline{HFK}'_n(L)$. Equivalently, we could first apply the collapse $-\text{gr}_M + 2(1 - 1/n) \text{gr}_T$ on $\overline{HFK}(L)$ to get the grading on $\overline{HFK}_n(L)$, then shift this $\frac{1}{n}\mathbb{Z}$ grading upward by $(1-\ell)(1-1/n)$. In other words, $\overline{HFK}'_n(L)$ is $\overline{HFK}_n(L)$ with its $\frac{1}{n}\mathbb{Z}$ -grading shifted upward by $(1-\ell)(1-1/n)$; similarly, $HFK'_n(L)$ is $HFK_n(L)$ with its $\frac{1}{n}\mathbb{Z}$ -grading shifted upward by $(1-\ell)(1-1/n)$.

Corollary 6.4. For $n \geq 2$, the $\frac{\text{gr}_n}{n}$ -graded Euler characteristic of $\overline{HFK}'_n(L)$ is

$$\Delta_L(t) \Big|_{t^{1/2}=-e^{\pi i /n}},$$

and the $\frac{\text{gr}_n}{n}$ -graded Euler characteristic of $HFK'_n(L)$ is zero.

Proof. For the reduced case, we have

$$e^{\pi i(1-\ell)(1-1/n)} e^{\pi i(1-\ell)/n} \Delta_L(t)|_{t^{1/2}=-e^{-\pi i/n}} = (-1)^{1-\ell} \Delta_L(t)|_{t^{1/2}=-e^{-\pi i/n}} = \Delta_L(t)|_{t^{1/2}=-e^{\pi i/n}}$$

(using that $\Delta_L(t^{-1}) = (-1)^{\ell-1} \Delta_L(t)$); for the unreduced case, we have $e^{\pi i(1-\ell)(1-1/n)} \cdot 0 = 0$. \square

The proof of [Dow18a, Lemma 2.23] gives us $\frac{1}{n}\mathbb{Z}$ -graded spectral sequences from $\overline{HFK}'(L)$ to $\overline{HFK}'_n(L)$ and from $HFK'(L)$ to $HFK'_n(L)$, where $\frac{\text{gr}_n}{n}$ on $\overline{HFK}'(L)$ and $HFK'(L)$ is defined to be $\text{gr}_M - 2(1 - 1/n) \text{gr}_T$.

When $n = 1$, the shifted homology groups $\overline{HFK}'_1(L)$ and $HFK'_1(L)$ agree with $\overline{HFK}_1(L)$ and $HFK_1(L)$ respectively, so their Euler characteristics are both 1.

Remark 6.5. The arguments in this section can be made simpler in the case of knots ($\ell = 1$), where by [Dow18a, Lemma 2.20], $\overline{HFK}_n(L)$ is isomorphic to $\frac{\text{gr}_n}{n}$ -graded $\widehat{HFK}(L) = \widehat{HFK}(L)$. In particular, Lemma 6.1 is unnecessary in this case.

7. EULER CHARACTERISTICS AND SPECTRAL SEQUENCES

Dowlin [Dow18a, Conjecture 1.6] conjectures the existence of spectral sequences from $\overline{H}_n(L)$ to $\overline{HFK}_n(L)$ and from $H_n(L)$ to $HFK_n(L)$. These sequences are conjectured to respect the $\frac{1}{n}\mathbb{Z}$ -gradings, where the $\frac{1}{n}\mathbb{Z}$ -grading $\frac{\text{gr}_n}{n}$ on reduced and unreduced $\mathfrak{sl}(n)$ homology is defined by

$$\frac{\text{gr}_n}{n} = \frac{1}{n} \text{gr}_{Q,n} + \text{gr}_H.$$

Dowlin works with n times this grading, which we would write as $\text{gr}_{Q,n} + n \text{gr}_H$.⁵ We state the following version of Dowlin's conjectures in terms of the grading-shifted theories \overline{HFK}'_n and HFK'_n .

Conjecture 7.1. *Let L be a link in S^3 . There exist spectral sequences with each page graded by $\frac{1}{n}\mathbb{Z}$, with differentials of degree +1 such that each page is the $\frac{1}{n}\mathbb{Z}$ -graded homology of the previous page, and with*

- E_2 page $\overline{H}_n(L)$ and E_∞ page $\overline{HFK}'_n(L)$;
- E_2 page $H_n(L)$ and E_∞ page $HFK'_n(L)$

as $\frac{1}{n}\mathbb{Z}$ -graded vector spaces.

These spectral sequences would give equalities of $\frac{1}{n}\mathbb{Z}$ -graded Euler characteristics

$$\chi(\overline{H}_n(L)) = \chi(\overline{HFK}'_n(L)), \quad \chi(H_n(L)) = \chi(HFK'_n(L)).$$

⁵In Dowlin's notation this grading is called $\text{gr}_n + \frac{n}{2}\text{gr}_v$ (where gr_n corresponds to our $\text{gr}_{Q,n}$). As specified in [Dow18a, Section 4.2], the grading gr_v here is $k = 2\text{gr}_v$ in the notation of [Ras15], where gr_v corresponds to our gr_H . This accounts for the factor of $\frac{1}{2}$ in Dowlin's formula.

Proof of Theorem 1.2. We have

$$\begin{aligned}
\chi(\overline{H}_n(L)) &= \sum_{\alpha \in \frac{1}{n}\mathbb{Z}} e^{\pi i \alpha} \dim_{\mathbb{Q}} \left(\overline{H}_n(L)_{\frac{1}{n} \text{gr}_{Q,n} + \text{gr}_H = \alpha} \right) \\
&= \sum_{I, J \in \mathbb{Z}} e^{\pi i (I/n + J)} \dim_{\mathbb{Q}} \left(\overline{H}_n(L)_{\text{gr}_{Q,n} = I, \text{gr}_H = J} \right) \\
&= \left(\sum_{I, J \in \mathbb{Z}} (-1)^J q^I \dim_{\mathbb{Q}} \left(\overline{H}_n(L)_{\text{gr}_{Q,n} = I, \text{gr}_H = J} \right) \right) \Big|_{q = e^{\pi i/n}} \\
&= \overline{P}_{n,L}(e^{\pi i/n});
\end{aligned}$$

similarly, $\chi(H_n(L)) = P_{n,L}(e^{\pi i/n})$, which is 0 for $n \geq 2$ and 1 for $n = 1$. Thus, for $n \geq 2$ these conjectured spectral sequences can be viewed as categorifications of the equalities

$$\overline{P}_{n,L}(e^{\pi i/n}) = \Delta_L(t)|_{t^{1/2} = -e^{\pi i/n}}, \quad P_{n,L}(e^{\pi i/n}) = 0.$$

For $n = 1$, the equalities are $\overline{P}_{1,L}(-1) = 1$ and $P_{1,L}(-1) = 1$ (note that $\overline{P}_{1,L}(q) = P_{1,L}(q) = 1$ in general). \square

Remark 7.2. Let $n \geq 2$ for simplicity. The spectral sequences of Conjecture 7.1, together with the ones from Sections 3.2.2 and 4.4 and the spectral sequences from HFK to HFK_n , can be organized as shown in Figure 2, following [Dow18a, Figure 1]. The arrows in this figure represent spectral sequences (solid for known, dotted for conjectural); we augment Dowlin's figure by labeled the arrows with their decategorified content.⁶

It is interesting to look at the square formed by the reduced theories; traveling along the left edge and then the bottom edge amounts to starting with $\overline{P}_L(a, q)$, evaluating at $a = q^n$, and then evaluating the result at $q = e^{\pi i/n}$ to get $\overline{P}_L(-1, e^{\pi i/n})$. On the other hand, traveling along the top edge and then the right edge amounts to starting with $\overline{P}_L(a, q)$, evaluating at $a = -1$ and $q = -t^{1/2}$, and then evaluating the result at $t^{1/2} = -e^{\pi i/n}$ to get $\overline{P}_L(-1, e^{\pi i/n})$. This compatibility at the level of Euler characteristics could be a sign of a more elaborate compatibility relationship between the conjectured spectral sequences at the categorified level.

REFERENCES

- [BLS17] J. A. Baldwin, A. S. Levine, and S. Sarkar. Khovanov homology and knot Floer homology for pointed links. *J. Knot Theory Ramifications*, 26(2):1740004, 49, 2017. arXiv:1512.05422.
- [DGR06] N. M. Dunfield, S. Gukov, and J. Rasmussen. The superpolynomial for knot homologies. *Experiment. Math.*, 15(2):129–159, 2006. arXiv:math/0505662.
- [Dow18a] N. Dowlin. A family of sl_n -like invariants in knot Floer homology, 2018. arXiv:1804.03165.
- [Dow18b] N. Dowlin. Knot Floer homology and Khovanov-Rozansky homology for singular links. *Algebr. Geom. Topol.*, 18(7):3839–3885, 2018. arXiv:1511.08953.
- [Dow18c] N. Dowlin. A spectral sequence from Khovanov homology to knot Floer homology, 2018. arXiv:1811.07848.

⁶In fact, “decategorified content” is not immediately defined for the arrows on the right edge of the square; the grading-collapsed versions of $\overline{HFK}(n)$ and $HFK(n)$ are not finite-dimensional over \mathbb{Q} so their Euler characteristics as $\frac{1}{n}\mathbb{Z}$ -graded complexes are not defined. It is reasonable to guess that if a natural definition of these Euler characteristics did exist, it would give the answer in the figure.




FIGURE 2. Spectral sequences from [Dow18a, Figure 1], with arrows labeled by decategorified content.

- [FYH⁺85] P. Freyd, D. Yetter, J. Hoste, W. B. R. Lickorish, K. Millett, and A. Ocneanu. A new polynomial invariant of knots and links. *Bull. Amer. Math. Soc. (N.S.)*, 12(2):239–246, 1985. projecteuclid/1183552531.
- [Kaw96] A. Kawauchi. *A survey of knot theory*. Birkhäuser Verlag, Basel, 1996. Translated and revised from the 1990 Japanese original by the author.
- [Kho16] M. Khovanov. Hopfological algebra and categorification at a root of unity: the first steps. *J. Knot Theory Ramifications*, 25(3):1640006, 26, 2016. arXiv:math/0509083.
- [KR08a] M. Khovanov and L. Rozansky. Matrix factorizations and link homology. *Fund. Math.*, 199(1):1–91, 2008. arXiv:math/0401268.
- [KR08b] M. Khovanov and L. Rozansky. Matrix factorizations and link homology. II. *Geom. Topol.*, 12(3):1387–1425, 2008. arXiv:math/0505056.
- [KS91] L. H. Kauffman and H. Saleur. Free fermions and the Alexander-Conway polynomial. *Comm. Math. Phys.*, 141(2):293–327, 1991. projecteuclid/1104248302.
- [Man14] C. Manolescu. An untwisted cube of resolutions for knot Floer homology. *Quantum Topol.*, 5(2):185–223, 2014. arXiv:1108.0032.
- [OSz04a] P. S. Ozsváth and Z. Szabó. Holomorphic disks and knot invariants. *Adv. Math.*, 186(1):58–116, 2004. arXiv:math/0209056.
- [OSz04b] P. S. Ozsváth and Z. Szabó. Holomorphic disks and topological invariants for closed three-manifolds. *Ann. of Math. (2)*, 159(3):1027–1158, 2004. arXiv:math/0101206.
- [OSz08] P. S. Ozsváth and Z. Szabó. Holomorphic disks, link invariants and the multi-variable Alexander polynomial. *Algebr. Geom. Topol.*, 8(2):615–692, 2008. arXiv:math/0512286.
- [PT87] J. H. Przytycki and P. Traczyk. Conway algebras and skein equivalence of links. *Proc. Amer. Math. Soc.*, 100(4):744–748, 1987.
- [Qi14] Y. Qi. Hopfological algebra. *Compos. Math.*, 150(1):1–45, 2014. arXiv:1205.1814.
- [Ras03] J. Rasmussen. *Floer homology and knot complements*. PhD thesis, Harvard University, 2003.
- [Ras15] J. Rasmussen. Some differentials on Khovanov-Rozansky homology. *Geom. Topol.*, 19(6):3031–3104, 2015. arXiv:math/0607544.
- [Zem19] I. Zemke. Link cobordisms and functoriality in link Floer homology. *J. Topol.*, 12(1):94–220, 2019. arXiv:1610.05207.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA

Email address: ajmanion@ncsu.edu

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC