

SPECTROSCOPIC CONFIRMATION OF AN ULTRA-MASSIVE GALAXY IN A PROTOCLUSTER AT $z \sim 4.9^*$

STEPHANIE M. URBANO STAWINSKI^{1,†}, M. C. COOPER¹, BEN FORREST², ADAM MUZZIN³, DANILO MARCHESINI⁴, GILLIAN WILSON⁵, PERCY GOMEZ⁶, IAN MCCONACHIE⁷, Z. CEMILE MARSAN³, MARIANNA ANNUNZIATELLA⁸, AND WENJUN CHANG⁷

¹Department of Physics & Astronomy, University of California, Irvine, 4129 Reines Hall, Irvine, CA 92697, USA

²Department of Physics and Astronomy, University of California Davis, One Shields Avenue, Davis, CA 95616 USA

³Department of Physics and Astronomy, York University, 4700, Keele Street, Toronto, ON MJ3 1P3, Canada

⁴Department of Physics and Astronomy, Tufts University, 574 Boston Avenue, Medford, MA 02155, USA

⁵Department of Physics, University of California Merced, 5200 North Lake Rd., Merced, CA 95343, USA

⁶W.M. Keck Observatory, 65-1120 Mamalahoa Hwy., Kamuela, HI 96743, USA

⁷Department of Physics & Astronomy, University of California Riverside, 900 University Ave., Riverside, CA 92521 USA and

⁸Centro de Astrobiología (CSIC-INTA), Ctra de Torrejón a Ajalvir, km 4, E-28850 Torrejón de Ardoz, Madrid, Spain

Version June 14, 2024

ABSTRACT

We present spectroscopic confirmation of an ultra-massive galaxy (UMG) with $\log(M_\star/M_\odot) = 10.98 \pm 0.07$ at $z_{\text{spec}} = 4.8947$ in the Extended Groth Strip (EGS), based on deep observations of Ly α emission with Keck/DEIMOS. The ultra-massive galaxy (UMG-28740) is the most massive member in one of the most significant overdensities in the EGS, with four additional photometric members with $\log(M_\star/M_\odot) > 10.5$ within $R_{\text{proj}} \sim 1$ cMpc. The Ly α profile is highly asymmetric ($A_f = 3.56$), suggesting the presence of neutral gas within the interstellar medium, circumgalactic medium, or via AGN-driven outflows. Spectral energy distribution (SED) fitting using a large suite of star formation histories and two sets of high-quality photometry from ground- and space-based facilities consistently estimates the stellar mass of UMG-28740 to be $\log(M_\star/M_\odot) \sim 11$ with a small standard deviation between measurements ($\sigma = 0.07$). While the best-fit SED models agree on stellar mass, we find discrepancies in the estimated star formation rate for UMG-28740, resulting in either a star-forming or quiescent system. *JWST*/NIRCam photometry of UMG-28740 strongly favors a quiescent scenario, demonstrating the need for high-quality mid-IR observations. Assuming the galaxy to be quiescent, UMG-28740 formed the bulk of its stars at $z > 10$ and is quenching at $z \sim 8$, resulting in a high star formation efficiency at high redshift ($\epsilon \sim 0.2$ at $z \sim 5$ and $\epsilon \gtrsim 1$ at $z \gtrsim 8$). As the most massive galaxy in its protocluster environment, UMG-28740 is a unique example of the impossibly early galaxy problem.

Subject headings: galaxies:high-redshift – galaxies:evolution – galaxies:clusters

1. INTRODUCTION

The discovery of extremely massive, passive galaxies at $z \sim 3-4$ has challenged our understanding of galaxy formation and evolution at early times (Schreiber et al. 2018; Forrest et al. 2020a,b; Valentino et al. 2020). These quiescent ultra-massive galaxies (UMGs, $M_\star > 10^{11} M_\odot$) have little to no ongoing star formation, requiring a period of intense star formation followed by rapid quenching within the first 2 Gyr of the Universe. Such extreme systems are difficult to reproduce in galaxy formation models, with current simulations struggling to form UMGs so early in cosmic time and often failing to quench these systems by $z \sim 3$ (Forrest et al. 2020a; Valentino et al. 2023; Xie et al. 2024). Thus, although rare, confirmation of UMGs provide powerful constraints on theoretical models of galaxy formation and evolution (e.g. Baugh 2006; Croton et al. 2006; Nelson et al. 2018).

Intriguingly, the modeled ages and star formation histories of UMGs at $z \sim 3$ suggest that some systems assembled the majority of their mass and subsequently quenched as early as $z \sim 5$ (Forrest et al. 2020b; Schreiber et al. 2018). Overall, there appears to be rapid evolution of the UMG population at $4 < z < 6$, as some of these massive systems transition from highly star-forming to passive. Imaging campaigns that

originally uncovered UMGs at $z \sim 3$ also corroborate these findings, yielding samples of photometric UMG candidates at $z > 4$ (e.g. Muzzin et al. 2013a,b; Weaver et al. 2023). Meanwhile, measurements of the number density of UMGs find a significant increase in their abundance with cosmic time from $z \sim 6$ to $z \sim 3$ (Stefanon et al. 2015; Davidzon et al. 2017; Marsan et al. 2022; Weibel et al. 2024). Altogether, deep photometric surveys point to not only the existence of these extreme systems at $z \sim 5$, but also that this epoch is critical for understanding the rapid evolution of this extreme population.

While it is presumed that ultra-massive galaxies should exist at $z \sim 5$, spectroscopic confirmation of candidate systems has been challenging. Studies searching for UMGs at $z \sim 5$ have encountered difficulty in identifying and detecting these systems via both ground- and space-based spectroscopy (Antwi-Danso et al. 2023; Marsan et al. in prep). The few existing, spectroscopically-confirmed UMGs at $z > 4$ represent a biased population of systems with exceptionally high star formation rates ($\text{SFR} \gtrsim 500 M_\odot \text{ yr}^{-1}$) and with relatively poorly-constrained stellar masses (e.g. Xiao et al. 2023, but see also Carnall et al. 2023b).

Herein, we present the analysis surrounding the spectroscopic discovery (see Urbano Stawinski et al. 2024a)¹ of an ultra-massive ($M_\star \sim 10^{11} M_\odot$) galaxy at $z = 4.8947$ that

[†]Corresponding author: ststawinski@gmail.com

*The data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation.

¹ While finalizing this manuscript, de Graaff et al. (2024) presented *JWST*/NIRSpec observations of this object, spectroscopically confirming its quiescence. These complementary and parallel results are discussed in more detail in §4.4.

resides within an extreme overdense region in the Extended Groth Strip (EGS). In §2, we describe our target selection criterion and spectroscopic observations. We present the resulting spectral fitting and characterization of the protocluster environment in §3. We discuss the competing star formation histories (SFH), compare our object to existing simulations, and investigate the mass assembly history of this object in §4. Finally, we summarize our results in §5. Unless otherwise specified, we adopt a Λ CDM cosmology throughout with $\Omega_M = 0.3$, $\Omega_\Lambda = 0.70$, and $H_0 = 68 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (Planck Collaboration et al. 2016).

2. DATA

2.1. Target Selection and Photometric Measurements

We selected UMG candidates ($M_\star \gtrsim 10^{11} M_\odot$) at $3.5 < z_{\text{phot}} < 6.5$ in the Extended Groth Strip (EGS) using the multi-wavelength photometric catalog from the Cosmic Assembly Near-infrared Deep Extragalactic Legacy Survey (CANDELS; Grogin et al. 2011; Koekemoer et al. 2011). The CANDELS photometric catalog (Stefanon et al. 2017), which spans $0.4 - 8 \mu\text{m}$ via a suite of ground- and space-based imaging, incorporates photometric redshifts and stellar masses from seven separate photo- z codes and/or spectral energy distribution (SED) template sets. For selection purposes, we used the median z_{phot} and $M_\star$ reported in the catalog. To increase the likelihood of detection of Ly α , we prioritized objects that had an apparent CFHT r -band AB magnitude of < 28 and star formation rate (SFR) of $\gtrsim 100 M_\odot \text{ yr}^{-1}$. To increase our sample size, we included UMG candidates within the given redshift range that had lower SFRs and masses as filler targets.

UMG-28740 ($\alpha = 14^{\text{h}}19^{\text{m}}39.7^{\text{s}}$ and $\delta = +52^\circ56'56.5''$), in particular, was selected as a star-forming candidate, with an estimated SFR of $20 - 200 M_\odot \text{ yr}^{-1}$ and an apparent CFHT r -band magnitude of $27.9^{+0.6}_{-0.4}$. Within the Stefanon et al. (2017) catalog, UMG-28740 has a median photometric redshift of $z_{\text{phot}} = 4.495$ and a median stellar mass of $\log(M_\star/M_\odot) = 11.0$. In total, the CANDELS catalog reports photometric measurements of UMG-28740 at $ugriz$ and JHK_s bands from CFHT, F606W, F814W, F125W, F140W, and F160W from *HST*, as well as all four *Spitzer*/IRAC channels.

In addition to the photometry from CANDELS, this galaxy was recently observed with *JWST*/NIRCam (Rieke et al. 2003; Beichman et al. 2012) as part of the Cosmic Evolution Early Release Survey (CEERS, PID 12117; Finkelstein et al. 2023; Bagley et al. 2023). Using *JWST*/NIRCam (pointing 2), UMG-28740 was observed with the F115W, F150W, F200W, F277W, F356W, F410M, and F444W filters at a 5σ depth ranging from AB magnitudes of 28.35–29.2. While the CEERS photometry was not incorporated in the initial target selection, it is utilized in this work as a supplemental set of photometry in §3.1.

2.2. Spectroscopic Observations and Reduction

Using Keck/DEIMOS (Faber et al. 2003), we observed UMG-28740 in March and May of 2023 on three different slitmasks. For all observations, we used the 600 lines mm^{-1} grating blazed at $7500 \text{ \AA}$ and tilted to a central wavelength of $7200 \text{ \AA}$, with the GG455 order-blocking filter employed and no dithering between exposures. With slit widths of $1''$, the spectral resolution (FWHM) for the 600g grating on DEIMOS is $\sim 3.5 \text{ \AA}$ (Weiner et al. 2006), with a dispersion of $0.65 \text{ \AA}$ per pixel. Calibrations for each slitmask included three internal quartz lamp flat-field frames and an arc lamp spectrum




FIG. 1.— (top) A roughly $150 \text{ \AA}$ -wide section of the co-added 2D spectrum for UMG-28740 from our Keck/DEIMOS observations, highlighting the Lyman- α emission line. (bottom) The corresponding co-added 1D spectrum with the resulting best-fit redshift shown in red. Top x-axis shows the velocity offsets from $z_{\text{Ly}\alpha}$ while the bottom x-axis shows the observed wavelength.

(using Kr, Ar, Ne, and Xe lamps). We were able to observe UMG-28740 with the three different slitmasks for a combined exposure time of 6.9 hours. The average seeing for these observations ranged from ~ 0.8 – $1''$ with variable transparency.

The Keck/DEIMOS observations were reduced using the spec2d² DEEP2/DEEP3 DEIMOS data reduction pipeline (Newman et al. 2013; Cooper et al. 2012), yielding sky-subtracted, wavelength-calibrated 1D and 2D spectra for each target. We then combined the three 1D spectra of UMG-28740 from the reduction of each slitmask, creating a final co-added 1D spectrum as shown in Figure 1. For visualization of the line, we also co-added the corresponding 2D spectra, as shown in the top panel of Figure 1.

To measure a best-fit redshift, we utilized a custom template fitter (as used in Urbano Stawinski et al. 2024b) that incorporates both an emission-line galaxy template as well as an asymmetric Gaussian profile to probe for a single Ly α line. For UMG-28740, we found a single emission line nearby the expected wavelength for Ly α given the photometric redshift. Fitting this profile, we find the resulting heliocentric-corrected spectroscopic redshift for UMG-28740 is $z_{\text{spec}} = 4.8947$ (best-fit model is shown by the red line in Fig. 1).

While this redshift is determined by the presence of a single emission line, there are multiple arguments consolidating this feature to be Ly α . First, the asymmetric profile of the emission line follows the characteristic shape of Ly α emitters at $z > 3$ (e.g. Verhamme et al. 2008; Martin et al. 2015). Second, the measured redshift is consistent with two sets of photometry from CANDELS and CEERS, including high-quality photometric measurements from *HST*/WFC3 and *JWST*/NIRCam (see Fig. 2). Lastly, the location of this galaxy coincides with a large overdense region at $z \sim 5$ (first highlighted by Naidu et al. 2022 and later discussed in §3.2), which further solidifies $z_{\text{spec}} = 4.8947$ as the correct redshift.

The asymmetric profile of Ly α suggests the presence of neutral gas around this object. We quantify the asymmetry of

² <https://sites.uci.edu/spec2d/>

the Ly α emission by taking the ratio of the red and blue sides of the line profile following $A_f = (\int_{\lambda_{\text{peak}}}^{\infty} f_{\lambda} d\lambda) / (\int_{\lambda_{\text{trough}}}^{\lambda_{\text{peak}}} f_{\lambda} d\lambda)$ where f_{λ} is the flux, λ_{peak} is the wavelength center of the profile, and λ_{trough} is the blue edge (Kakiichi & Gronke 2021; Rhoads et al. 2003). We find the asymmetry parameter to be high, $A_f = 3.56$, potentially indicating sufficient neutral gas surrounding the galaxy within the interstellar medium, the circumgalactic medium, or possibly as part of AGN-driven outflows (Verhamme et al. 2008; Laursen et al. 2009).

3. ANALYSIS

3.1. SED Fitting

With our measured spectroscopic redshift ($z_{\text{spec}} = 4.8947$), we re-fit the SED of UMG-28740. Here, we employ both the photometry from CANDELS (Stefanon et al. 2017) and from CEERS (Bagley et al. 2023). From the CANDELS photometric catalog, we exclude the *HST*/ACS photometry at F606W and F814W due to uncertainties related to sky subtraction, such that the SED fits are made to the photometry at *ugrizJHK_s* from CFHT, F125W, F140W, and F160W from *HST*/WFC3-IR, and all four *Spitzer*/IRAC channels. Meanwhile, the CEERS catalog includes photometric measurements at F606W and F814W from *HST*/ACS, F125W, F140W, and F160W from *HST*/WFC3-IR, and finally F115W, F150W, F200W, F277W, F356W, F410M, and F444W from *JWST*/NIRCam.

The majority of our SED analysis was completed using the SED-fitting code Bayesian Analysis of Galaxies for Physical Inference and Parameter ESTimation or BAGPIPES (Carnall et al. 2018).³ BAGPIPES assumes Bruzual and Charlot stellar populations (Bruzual & Charlot 2003), a Kroupa IMF (Kroupa & Boily 2002), and a Calzetti dust law (Calzetti et al. 2000). Adopting a double-power law star formation history, we measure best-fit model SEDs for each set of photometry (from CANDELS and CEERS). We fit these datasets separately to avoid any systematic biases associated with the varying photometric methods, such as different aperture sizes and adopted PSF-matching methodology. The SEDs are fit to 10 free parameters, allowing for variation in metallicity, age, the slope of the Calzetti dust law, and the ionization parameter. The full set of free parameters used and their allowed distributions are outlined in Table 1.

As shown in Figure 2, the resulting SED fits give a stellar mass of $\log(M_{\star}/M_{\odot}) = 10.98^{+0.04}_{-0.03}$ using the CEERS photometry and $\log(M_{\star}/M_{\odot}) = 11.08^{+0.08}_{-0.1}$ using the CANDELS photometry. While the stellar mass is tightly constrained between the two sets of photometry, we find that the SFR is dependant on the photometric catalog utilized. The CANDELS photometry yields a dusty, highly star-forming galaxy ($\text{SFR} = 285.0^{+125.5}_{-79.5} M_{\odot} \text{ yr}^{-1}$) and the fit to the *JWST*/NIRCam photometry produces a very low star formation rate ($\text{SFR} = 9.5^{+4.8}_{-2.7} M_{\odot} \text{ yr}^{-1}$) and low dust attenuation. The latter, *nearly* quenched SED is shown in orange in Figure 2, and is consistent with both sets of photometry within $\pm 1\sigma$ error. On the other hand, the star-forming SED (shown in blue in Fig. 2) has insufficient emission in the rest-frame optical ($\sim 4000\text{--}6000\text{\AA}$), so as to under-produce the flux measured in the *JWST*/NIRCam F277W band. In addition, the star-forming SED overpredicts the observed flux in the *JWST*/NIRCam F356W, F410M, and F444W bands due to contribution from

TABLE 1
LIST OF FREE PARAMETERS USED IN THE SED FITS WITH BAGPIPES

Parameter	Unit	Range	Prior
General ^a			
Stellar Mass Formed	$M_{\star}/M_{\odot}$	(1, 15)	Logarithmic
Peak Star Formation	τ/Gyr	(0.3, t_{obs})	Logarithmic
Age	$t_{\text{age}}/\text{Gyr}$	(0.1, 1)	Logarithmic
Metallicity	$Z/Z_{\odot}$	(0.01, 2.5)	Logarithmic
V-band attenuation	A_V	(0, 8)	Uniform
Deviation from Calzetti slope	δ	(-0.3, 0.3)	Gaussian ^b
2175Å bump strength	B	(0, 5)	Uniform
Ionization	U	(-4, -2)	Logarithmic
Double-power law ^c			
double-power law falling slope	α	(0.01, 1000)	Logarithmic
double-power law rising slope	β	(0.01, 1000)	Logarithmic
Burst ^d			
Stellar Mass Formed	$M_{\star}/M_{\odot}$	(0, 15)	Logarithmic
Age	$t_{\text{age}}/\text{Gyr}$	(0.1, 1)	Logarithmic
Metallicity	$Z/Z_{\odot}$	(0.01, 2.5)	Logarithmic
AGN ^d			
Continuum Flux at 5100Å	f_{5100} erg/s/cm ² /Å	(0, 10^{-19})	Uniform
H α Flux	$f_{\text{H}\alpha}$ erg/s/cm ²	(0, $10^{-16.6}$)	Uniform
H α Velocity Dispersion	$\sigma_{\text{H}\alpha}/\text{km/s}$	(1000, 5000)	Logarithmic
Power law slope:			
at $\lambda < 5000\text{\AA}$	$\alpha_{\lambda < 5000}$	(-2, 2)	Gaussian ^e
at $\lambda > 5000\text{\AA}$	$\alpha_{\lambda > 5000}$	(-2, 2)	Gaussian ^f

^aParameters used in all SED fits.

^bThe Gaussian distribution for the Calzetti slope has a mean of $\mu = 0$ and a standard deviation of $\sigma = 0.1$.

^cParameters specifically used in the double-power law fits.

^dParameters used in added AGN and burst components.

^eThe Gaussian distribution for the power law slope at $\lambda < 5000\text{\AA}$ has a mean of $\mu = -1.5$ and a standard deviation of $\sigma = 0.5$.

^fThe Gaussian distribution for the power law slope at $\lambda > 5000\text{\AA}$ has a mean of $\mu = 0.5$ and a standard deviation of $\sigma = 0.5$.

strong emission lines such as H α at $\sim 3.9\mu\text{m}$. Based on the photometry alone, this favors the quenched scenario for UMG-28740 (see §4.1 for a more detailed discussion of these discrepancies).

To test the robustness of our measurements for mass and SFR, we ran BAGPIPES again using a variety of different models; varying the included free parameters, changing the assumed star formation history, restricting the assumed metallicity to solar, and adding different components such as bursts and the broad-line region of an active galactic nucleus (AGN; modeled by Gaussian fits to the H α and H β broad lines, see Carnall et al. 2023b). In summary, we highlight the results from five different models: a double-power law SFH, a double-power law SFH including an AGN, a burst, or fixing the metallicity to solar metallicity, and a delayed- τ model (see Table 2). Adding an AGN or burst component is of interest to the evolution of massive galaxies in the early Universe, as the spectra of a subset of UMGs have significant AGN contributions (Marsan et al. 2017; Forrest et al. 2020b, Forrest et al. in prep.) and current theoretical models speculate that the rapid formation of such systems require extreme bursts of star formation (Sun et al. 2023).

³ <https://github.com/ACCarnall/bagpipes>




FIG. 2.— (top) From left to right: image stamps of UMG-28740 in the *JWST*/NIRCam 1.50 μm , 2.77 μm , and 4.44 μm filters, followed by an RGB image (F150M in blue, F277M in green, and F444W in red). (bottom) Best-fit SEDs (star-forming in blue and quenched in orange) from BAGPIPES forced at the measured z_{spec} . The two sets of observed photometry are shown as dark blue (Stefanon et al. 2017) and orange (Bagley et al. 2023) markers. The width of each SED indicates the $\pm 1\sigma$ confidence intervals resulting from BAGPIPES. The open blue crosses show the predicted *JWST*/NIRCam flux at F277W, F356W, F410M, and F444W for the blue star-forming SED. Here, the predicted *JWST*/NIRCam flux for the star-forming scenario is under-estimated at 2.77 μm and over-estimated at 3.56 – 4.44 μm , suggesting that UMG-28740 is likely quenched.

We also compared the results from BAGPIPES to fits using the C++-adapted SED fitting code FAST++⁴ (Kriek et al. 2009). With FAST++, we adopt the same parameters used in the literature to model UMGs at $z \sim 3.5$ (Forrest et al. 2020a,b), assuming a Bruzual Charlot stellar population (Bruzual & Charlot 2003), the Calzetti dust law (Calzetti et al. 2000), and a Chabrier IMF (Chabrier 2003). The Chabrier IMF is related to the Kroupa IMF used in BAGPIPES by a factor of ~ 1 (Wright et al. 2017), allowing for direct comparison between results of the two SED codes. We allow FAST++ to vary the extinction from $0 < A_V < 4$ with a $\Delta A_V = 0.1$, the age from $7 < \log(\text{Age}) < \text{age of the Universe at } z_{\text{spec}}$ with $\Delta \log(\text{Age}) = 0.05$, and a metallicity of $Z/Z_\odot = 0.02$ as in Forrest et al. (2020a,b). For the assumed SFH, we adopt a double-exponential SFH as used in Forrest et al. (2020a,b).

Remarkably, regardless of the photometric dataset used, SFH assumed, underlying SED code, or range of free parameters, we find *all* best-fit SEDs result in a massive galaxy with a stellar mass of $\log(M_\star/M_\odot) \sim 11$. Across the various fits, UMG-28740 has a median mass of $\log(M_\star/M_\odot) = 10.98$

with a 1σ standard deviation of 0.07 (see Table 2 for measurements from each SED fit). Hereinafter, we conservatively use this stellar mass and error as our best estimate for UMG-28740. This SED fitting analysis also highlights a discrepancy between SFR and dust extinction – resulting in the two possible scenarios, star-forming or quenched (see §4.1 for a more detailed discussion).

3.2. An Overdense Region around UMG 28740

UMG-28740 is embedded within a significant overdensity at $z \sim 5$, as traced by the surface density of galaxies at $4.5 < z_{\text{phot}} < 5.5$ in the CANDELS photometric catalog. As shown in Figure 3, UMG-28740 resides near the center of the overdense region and is the most massive member. However, we caution that the observed overdensity is located near the edge of the CANDELS survey area within the EGS, with UMG-28740 only $\sim 16''$ from the edge of the *HST*/WFC3 imaging footprint, such that the full extent of the structure may be underestimated. Naidu et al. (2022) first identified this particular overdense region, finding that a $\sim 0.8 \text{ arcmin}^2$ area centered on a lower-mass photometric neighbor of UMG-

⁴ <https://github.com/cschreib/fastpp>

TABLE 2
STELLAR MASS AND SFR FROM ALTERNATIVE SED FITS

SFH	Code	Mass $\log(M_\star/M_\odot)$	SFR $M_\odot/\text{yr}$
CANDELS (Stefanon et al. 2017)			
Double-power law ^a	BAGPIPES	11.08 ^{0.08} _{0.1}	285.0 ^{125.5} _{79.5}
Double-power law+AGN	BAGPIPES	10.97 ^{0.04} _{0.04}	8.6 ^{3.9} _{2.4}
Double-power law+burst	BAGPIPES	10.98 ^{0.08} _{0.09}	419.6 ^{156.7} _{162.2}
Double-power law at $Z_\odot$	BAGPIPES	10.91 ^{0.1} _{0.07}	299.9 ^{99.5} _{60.1}
Delayed- τ	BAGPIPES	10.92 ^{0.11} _{0.13}	312.1 ^{159.1} _{106.2}
Double Exponential	FAST++	10.95 ^{0.00} _{0.11}	24.5 ^{16.6} _{3.3}
CEERS (Bagley et al. 2023)			
Double-power law ^a	BAGPIPES	10.98 ^{0.04} _{0.03}	9.5 ^{4.8} _{2.7}
Double-power law+AGN	BAGPIPES	10.96 ^{0.04} _{0.04}	0.0 ^{0.0} _{0.0}
Double-power law+burst	BAGPIPES	11.17 ^{0.03} _{0.04}	18.1 ^{6.2} _{4.7}
Double-power law at $Z_\odot$	BAGPIPES	10.99 ^{0.02} _{0.02}	7.4 ^{1.0} _{0.7}
Delayed- τ	BAGPIPES	11.05 ^{0.02} _{0.03}	32.2 ^{10.8} _{5.7}

^aSED fit is shown in Fig. 2.

28740 contained $4\times$ the average surface density of sources at $4.5 < z_{\text{phot}} < 5.5$. However, this lower-mass neighbor ($M_\star \sim 10^{9.6} M_\odot$, spectroscopically confirmed by Arrabal Haro et al. 2023b), is located $\geq 20''$ to the south (≥ 0.8 cMpc at $z = 5$) of UMG-28740 and outside of the apparent center of the overdense region as traced by the CANDELS photometric catalog (see the highlighted red cross in Fig. 3).

To investigate the extent of the overdense region around UMG-28740, we search for other overdensities across the entire CANDELS footprint within the EGS field by comparing the galaxy surface densities in redshift slices of $\Delta z = 1$ to the average surface density at each corresponding epoch. We use the CANDELS catalog to identify sources at $2.5 < z_{\text{phot}} < 6.5$, as this catalog covers a larger footprint within the EGS (versus CEERS). We limit this analysis to sources with $\log(M_\star/M_\odot) > 9$, so as to define a tracer population that is unbiased (complete) over the full redshift range probed. Across the entire field, we compute the galaxy surface density as a function of position on the sky within an aperture of $45'' \times 45''$ and a depth of $\Delta z = 1$. Stepping through redshift from $z = 3$ to $z = 6$ with a step size of $\Delta z = 0.1$, we identify regions that are overdense compared to the average surface density at the corresponding redshift. We find that the region hosting UMG-28740 is one of the most extreme overdense regions within the EGS at $3 < z < 6$, with a surface density that is $\sim 10\times$ higher than the average surface density of the field at $z = 4.9$.

At $z \sim 5$, the effective radii of protoclusters are estimated to extend up to $R_e = 5 - 10$ cMpc (Chiang et al. 2013). UMG-28740 is positioned nearby the apparent center of the overdense region, such that we can estimate R_{proj} from UMG-28740 to trace the effective radius of the protocluster. However, we caution this region is also near the edge of the EGS field, limiting observed galaxies past $R_{\text{proj}} \sim 0.6$ cMpc to the north-west and potentially skewing the true center of the overdensity. Nevertheless, within $R_{\text{proj}} = 3.44$ cMpc from UMG-28740, there are 26 photometric protocluster members at $4.5 < z < 5.5$ and with $M_\star > 10^9 M_\odot$ in the CANDELS photometric cata-




FIG. 3.— Overdensity hosting UMG-28740 (blue star) at $z \sim 5$. The underlying 2D histogram and contours show the galaxy surface density for objects in the CANDELS catalog at $4.5 < z_{\text{phot}} < 5.5$ and $\log(M_\star/M_\odot) > 9$. The beam size (with a diameter of $4.5''$) used to convolve the density map is shown in the upper right corner. Cyan circles denote photometric sources found in the CANDELS (Stefanon et al. 2017) and CEERS (Bagley et al. 2023) catalogs, where white circles are low-mass galaxies only found in the deeper CEERS catalog. The size of the circle indicates the estimated stellar mass of the object. Red crosses and the orange diamond are spectroscopically confirmed sources from Arrabal Haro et al. (2023a) and Urbano Stawinski et al. (2024b), respectively. The highlighted red cross indicates the location of the lower-mass member from Naidu et al. (2022).

log along with 160 members identified in the CEERS catalog, pushing down to lower stellar masses ($M_\star > 10^8 M_\odot$). Additionally, out of all the photometric members there are 4 massive neighbors with $10.5 < \log(M_\star/M_\odot) < 11$ within just $29.5''$, or ~ 1.13 cMpc, of UMG-28740. The closest of these massive members is only separated from UMG-28740 by 0.29 cMpc. This makes the protocluster surrounding UMG-28740 the most compact overdensity of massive $\log(M_\star/M_\odot) > 10.5$ galaxies within the entire EGS at $z \sim 5$ (according to the CANDELS catalog), with only two other compact regions each containing two similarly massive galaxies at a similar spatial scale ($R_{\text{proj}} < 3$ cMpc). Moreover, there are *no* other instances of > 3 massive galaxies within $R_{\text{proj}} < 3$ cMpc at this redshift in the CANDELS catalog.

This protocluster also hosts 5 spectroscopically-confirmed galaxies from Keck/DEIMOS (Urbano Stawinski et al. 2024b) and JWST/NIRSpec (Arrabal Haro et al. 2023a) spectroscopy within $R_{\text{proj}} = 3.44$ cMpc (see Fig. 3). The measured mass of these spectroscopic members range from $\log(M_\star/M_\odot) = 9.18 - 9.95$ and span 0.75 cMpc to 2.10 cMpc from UMG-28740. In summary, all members within $R_{\text{proj}} = 4$ cMpc, both spectroscopically confirmed and all photometric candidates, are estimated to be less massive than UMG-28740.

4. DISCUSSION

4.1. Investigation into Competing Star Formation Histories

As mentioned in §3.1, the two sets of photometry yield two different SFH scenarios, one with a highly star-forming and dusty SED versus one with low star formation and little




FIG. 4.— Star formation histories for the two competing scenarios, star-forming (blue) and quenched (orange), based on the best-fit SEDs from BAGPIPES and FAST++ (see Table 2). The solid lines, as well as the shaded region, show the best-fit and $\pm 1\sigma$ double-power law SFH from BAGPIPES (shown in Fig. 2). The formation time (t_{form}) corresponds to the epoch at which 50% of the stellar mass is formed for the double-power law models. The higher-quality *JWST*/NIRCam photometry favors the quenched scenario and suggests that UMG-28740 formed much of its stellar mass as early as $z \gtrsim 10$.

dust extinction. The high-quality *JWST*/NIRCam photometry favors a nearly quenched, SED. In contrast, the photometry from CANDELS favors a star-forming SED, with the exception of a quenched solution when assuming a double-exponential SFH with FAST++ (as described in §3.1). We show a variety of these SFHs, highlighting the best fit SEDs from the final BAGPIPES run, in Figure 4.

Looking at the SED fits alone, the $\pm 1\sigma$ spread of the orange, quenched SED is consistent with both sets of photometry within the errors. However, there is a clear offset between the *JWST*/NIRCam photometry and the blue, star-forming SED, with the largest discrepancy at F277W and F410M. To test the consistency between the NIRCam photometry and the star-forming SED, we predict the flux that we would expect from a mock F277W, F356W, F410M, and F444W observation of the star-forming SED fit, using the ‘predict’ function within the BAGPIPES code (see the open blue crosses in Fig. 2). For F277W, the expected flux would be elevated from the continuum due to the emission lines nearby $2.77\mu\text{m}$; however, we find this predicted flux is insufficient to reproduce the observed F277W flux even within $\pm 1\sigma$. For F356W and F410M, the predicted flux is elevated to a greater extent, due to contributions from strong emission lines such as $H\alpha$ at $\sim 3.9\mu\text{m}$. However, this added emission over-predicts the flux measured by the NIRCam photometry (also seen, to a lesser extent, in the F444W predicted flux). Taken together, this suggests that the quenched scenario is more consistent with the observed photometry from both catalogs, and thus more likely to be the true SFH of UMG-28740. Moreover, running BAGPIPES with a double-power law SFH on both sets of photometry simultaneously (i.e. the combined CANDELS and CEERS photometric dataset), we find the best-fit model to be quenched ($\text{SFR} \sim 8 M_{\odot} \text{ yr}^{-1}$), further indicating the importance of the mid-IR photometry in supporting a quenched SFH for UMG-28740.

One argument against a quenched SFH is the detection of $\text{Ly}\alpha$ emission. However, this may be explained by either AGN contribution or remnant star formation consistent with a post-starburst SED. Indeed, a recent study of a post-starburst galaxy at $z \sim 3$ has detected significant amounts of neutral gas from AGN driven outflows via ALMA (Scholtz et al. 2024). Outflowing neutral gas around UMG-28740 could also contribute to the asymmetric $\text{Ly}\alpha$ profile measured in §2.2.

To explore the consistency of the observed $\text{Ly}\alpha$ profile with a quiescent model, we compare the observed and expected $\text{Ly}\alpha$ equivalent widths (EW) from different SED fits. Assuming no AGN contamination, we again use the ‘predict’ function in BAGPIPES to estimate the EW of $\text{Ly}\alpha$ in the various SED fits. For the nearly quenched scenario, the SFR from all models without an AGN component ranges from $9.5 M_{\odot} \text{ yr}^{-1}$ to $32.2 M_{\odot} \text{ yr}^{-1}$. Measuring the EW of $\text{Ly}\alpha$ from the model SEDs (using the $\pm 1\sigma$ spread in the quenched models) yields an $\text{EW}_{\text{Ly}\alpha} = 16.5 - 35.1 \text{ \AA}$. We then estimate the EW of the observed $\text{Ly}\alpha$ line from Keck/DEIMOS. We find the continuum level just redward of the line profile and calculate the EW using a custom python code. Because the continuum level is so low, we conservatively call this measurement an upper limit on the true EW. Performing this analysis yields an observed $\text{EW}_{\text{Ly}\alpha} > 34.0 \text{ \AA}$, within the range of the predicted values for the post-starburst SED. Hence, we cannot rule out the nearly quenched scenario by detection of $\text{Ly}\alpha$ emission alone.

We perform the same analysis on the SEDs with AGN contribution. With AGN contribution we find the $\pm 1\sigma$ spread for the quenched SEDs yields $\text{EW}_{\text{Ly}\alpha} = 29.9 - 31.7 \text{ \AA}$, which is below the estimated observed $\text{Ly}\alpha$ EW. This would suggest that AGN contribution alone may not explain the presence of $\text{Ly}\alpha$ emission seen in the Keck/DEIMOS spectroscopy, and that likely there is some remnant star formation (even with the presence of AGN contamination).




FIG. 5.— SFR versus stellar mass of UMG-28740 for the two scenarios, star-forming (blue star) and quenched (orange star). The black circles show galaxies from the Illustris TNG-300 simulation at $z = 5$. The red dashed line and white squares trace the median of the galaxies in the TNG sample, while the red dotted lines show the $\pm 1\sigma$ spread. We also include other spectroscopically-confirmed massive galaxies at this epoch from the literature as purple circles (Xiao et al. 2023) and a peach cross (Carnall et al. 2023b).

The best-fit SEDs for UMG-28740 yield two different SFH scenarios, a highly star-forming galaxy or a nearly quiescent one. However, due to the flux required by the *JWST*/NIRCam photometry in the mid-IR (F277W, F356W, F410M, and F444W) and the consistency in the EW of the observed and predicted $\text{Ly}\alpha$ emission line, we argue that the quenched SED is likely the true SFH of this object.

4.2. Comparison to Simulations

The number density of UMGs at $z \sim 5$ is extremely low, especially for more quiescent systems (Stefanon et al. 2015; Davidzon et al. 2017; Marsan et al. 2022; Weaver et al. 2023). In addition, due to various observational challenges associated with observing these systems, there are only a handful of spectroscopically-confirmed UMGs at this epoch in the literature (e.g. Antwi-Danso et al. 2023; Xiao et al. 2023; Carnall et al. 2023b). To compare the SFR of UMG-28740 to a large, unbiased sample of galaxies, we plot SFR versus stellar mass for UMG-28740 with simulated galaxies. Many simulations struggle to produce massive systems at high redshift, and most fail to quench these objects at early epochs. However, the Illustris TNG-300 simulation (Nelson et al. 2019) has been able to form massive systems and subsequently suppress SFR to reproduce quiescent UMGs up to $z \sim 3.5$ (Forrest et al. 2020a; Schreiber et al. 2018; Glazebrook et al. 2017). Hence, we compare UMG-28740 to simulated galaxies in TNG-300 at $z = 5$ (see Fig. 5).

While TNG-300 reproduces massive quiescent systems at $z \sim 3.5$, as well as massive star-forming galaxies at earlier epochs (Forrest et al. 2020a), TNG-300 fails to create massive and quiescent galaxies like UMG-28740 (assuming a quenched scenario) and one other spectroscopically-confirmed quenched system in the literature to date (Carnall et al. 2023b) at $z \sim 5$. The measured SFR of UMG-28740 for the quenched

scenario is ~ 2 dex below that of any other simulated galaxy at $\log(M_\star/M_\odot) \sim 11$. On the other hand, the TNG-300 simulation does reproduce massive, star-forming galaxies at this epoch, including those from Xiao et al. (2023) and the star-forming scenario of UMG-28740. The lack of massive quenched objects is likely a result of one of two possibilities: the TNG-300 simulation fails to catch the cessation of star formation at this epoch or that the size of TNG-300 is too small ($V = 302.6^3 \text{ Mpc}^3$) to capture a system as rare as UMG-28740 and its protocluster environment.

Recently, Xie et al. (2024) used the GAEA semi-analytic model (De Lucia et al. 2024), applied to a larger box size than that of TNG-300 ($V = 685^3 \text{ Mpc}^3$), to study the formation of massive quiescent galaxies at early cosmic times. In summary, Xie et al. (2024) find a larger relative number of quenched massive galaxies at $z \sim 4.5$, with a quenched fraction for systems with $M_\star = (0.8 - 1.2) \times 10^{11} M_\odot$ of $f_q \sim 0.06$. Xie et al. (2024) suggest that the primary quenching mechanism for massive galaxies at high redshift is accretion disk feedback via quasar winds. Yet even within the GAEA simulation UMG-28740 is exceptionally rare; galaxies as quiescent as the best-fit quenched model, with a specific SFR of $\sim 10^{-10} \text{ yr}^{-1}$, have a number density of only $\sim 2 \times 10^{-7}$ per cMpc^3 at $z \sim 5$ in GAEA (De Lucia et al. 2024).

Observational studies suggest that massive quiescent galaxies at $z \sim 5$ exist at greater numbers than those predicted in these models (e.g. Carnall et al. 2023a). However, spectroscopic detection of these systems has been nearly unachievable prior to the launch of *JWST*. With *JWST* in operation, more quiescent massive systems are being uncovered (see Carnall et al. 2024 for 4 such systems at $3 < z < 5$). These discoveries suggest systems such as UMG-28740 are not as unique as current simulations project, placing valuable constraints on modern models of galaxy formation.

4.3. Mass Assembly History of UMG-28740

To investigate the potential evolution of UMG-28740, in Figure 6 we plot the mass assembly history corresponding to the best-fit star-forming and quiescent models from BAGPIPES (see Fig. 4). We compare the mass growth of UMG-28740 to that of spectroscopically-confirmed UMGs at $z \sim 3.5$ from the Massive Ancient Galaxies At $z > 3$ NEar-infrared (MAGAZ3NE) survey (Forrest et al. 2020a,b, 2022), a Keck/MOSFIRE effort that has successfully spectroscopically confirmed a large sample of star-forming and quenched UMGs at $z \sim 3.5$. We also include a sample of massive dusty star-forming galaxies (DSFGs) at $z > 5$ from the literature (Riechers et al. 2020; Williams et al. 2019; Jin et al. 2019; Zavala et al. 2018; Pavesi et al. 2018; Riechers et al. 2017; Strandet et al. 2017; Ma et al. 2015; Cooray et al. 2014; Riechers et al. 2013; Walter et al. 2012), including the dusty star-forming UMGs spectroscopically confirmed by Xiao et al. (2023). Finally, we also include the massive, spectroscopically-confirmed quiescent galaxy from Carnall et al. (2023b) and the photometric massive galaxy candidates from Labbé et al. (2023). Below we discuss the possible connections between UMG-28740 and these various populations.

For the mass assembly history predicted by the best-fit star-forming models, UMG-28740 is consistent with being the progenitor of the existing UMG population at $z \sim 3.5$, such that the ongoing star formation in the system will continue to grow its stellar mass at $3 \lesssim z \lesssim 5$, likely becoming similar to one of the more massive UMGs detected by the MAGAZ3NE




FIG. 6.— Mass assembly history of UMG-28740 for the best-fit star-forming models (blue lines) and quiescent models (orange lines). We include the double-power law fits (solid line), the fixed $Z = Z_{\odot}$ double-power law fits (dashed dotted lines), and the delayed- τ fits (dotted lines). The $\pm 1\sigma$ uncertainties for the double-power law fits are shown as the shaded blue and orange regions. The green tracks and squares show the assembly histories for spectroscopically-confirmed UMGs at $z \sim 3$ from Forrester et al. (2020b). Spectroscopically-confirmed massive galaxies at $z \sim 5$ from Xiao et al. (2023) and (Carnall et al. 2023b) are shown as purple circles and a peach cross, respectively. Finally, we include a sample of $z > 5$ DSFGs (grey open hexagons) from the literature and a sample of photometric massive galaxy candidates from Labbé et al. (2023) (cyan diamonds). The grey shaded regions show the stellar mass prohibited by Λ CDM cosmology, following $M_{\star}^{\max} = \epsilon f_b M_{\text{halo}}^{\max}$ where ϵ is the star formation efficiency and f_b is the cosmic baryon fraction. We include three different thresholds, by assuming the area of the EGS at a 20% efficiency and a 100% efficiency as well as the area of the full sky and 100% efficiency.

survey. Similarly, the DSFG population at $z \sim 5$, including the galaxies identified by Xiao et al. (2023), are also likely to evolve into members of the existing UMG population at $z \sim 3.5$ (Forrester et al. 2020b). Looking towards higher z for the best-fit star-forming models, the assembly history of UMG-28740 is consistent with the massive galaxy candidates from Labbé et al. (2023) as potential progenitors, with UMG-28740 forming much of its stellar mass at $z \lesssim 8$.

On the other hand, assuming the more-likely quenched SFH, UMG-28740 will evolve into a lower-mass UMG at $z \sim 3.5$ ($M_{\star} \sim 10^{11} M_{\odot}$). Interestingly, however, in this scenario UMG-28740 would not be included in the sample of known UMGs at $z \sim 3.5$. Due to the difficulty in detecting absorption lines in faint systems from the ground, MAGAZ3NE observations were limited to an AB magnitude of $K_s < 21.7$. In contrast, if we passively evolve the quenched model SED for UMG-28740 at $z = 4.8947$ to $z = 3.5$, we find that UMG-28740 would have an AB magnitude of $K_s = 22.7$, fainter than the selection limit for the MAGAZ3NE survey and indicating that existing samples of UMGs at intermediate redshift may be biased against the oldest systems (see Forrester et al. 2020b). Instead, UMG-28740 would more likely resemble much fainter UMGs at $z \sim 3$, such as those that have been targeted with *JWST*/NIRSpec (Nanayakkara et al. 2024; Glazebrook et al. 2024).

Towards high- z , the evolutionary history of UMG-28740 drastically changes assuming a quenched model. At $z \sim 5$, UMG-28740 is already an extreme system, more massive than the other spectroscopically-confirmed quiescent system at this epoch (Carnall et al. 2023b). Moreover, in contrast to the massive quiescent galaxy from Carnall et al. (2023b), which formed its mass at $z \sim 7$, we find UMG-28740 to already be

quenched by $z \sim 8$, with the bulk of its stellar mass formed at much earlier cosmic time ($z > 10$). Based on the quenched SED fits, the mass assembly of UMG-28740 is inconsistent with the DSFG population at $z > 5$ and the massive galaxy candidates from Labbé et al. (2023), instead forming a more extreme amount of mass at very high redshift.

The formation of such massive galaxies at high redshift is a highly active area of study thanks to a wealth of new observations from *JWST*. For example, early *JWST* imaging campaigns discovered a surplus of massive galaxy candidates at $z = 8-14$ that stand in tension with predictions from Λ CDM (Casey et al. 2023; Labbé et al. 2023; Boylan-Kolchin 2023). These results have sparked a myriad of questions and scrutiny in our current understanding of the physical processes driving galaxy evolution, including those that underpin SED fitting codes. The extreme build-up of mass required to assemble massive galaxies suggests intense bursts of star formation that convert $> 20\%$ of the available baryons into stars (Steinhardt et al. 2016; Boylan-Kolchin 2023; Sun et al. 2023; Casey et al. 2023; Xiao et al. 2023; Antwi-Danso et al. 2023).

To investigate the star formation efficiency of UMG-28740, we plot the limiting mass as imposed by a Λ CDM cosmology (shown as the grey shaded regions in Fig. 6). To do this, we follow the methodology from Boylan-Kolchin (2023), first calculating the number density of halos of a given mass as a function of redshift using *hmf*⁵, a python package that calculates the halo mass function at different redshifts (Murray 2014). We then find the maximum halo mass, given a survey volume, for which we would observe 1 such system. To convert to stellar mass, we follow $M_{\star}^{\max} = \epsilon f_b M_{\text{halo}}^{\max}$, where

⁵ <https://hmf.readthedocs.io>

f_b is the cosmic baryon fraction and ϵ is the efficiency of converting baryons into stars. We perform this calculation for the area of the CANDELS footprint within the EGS (205 arcmin², [Stefanon et al. 2017](#)) using a line-of-sight depth of $\Delta z = 2$ and an efficiency of 20% ($\epsilon = 0.2$) and 100% ($\epsilon = 1.0$). We also calculate the maximum stellar mass for observations across the full sky with an efficiency of 100%, which is the upper limit for what is allowed in a Λ CDM cosmology.

At $z \sim 5$, UMG-28740 has a stellar mass consistent with a relatively high star formation efficiency ($\sim 20\%$), comparable to that found in larger samples of massive galaxies at high redshift (e.g. [Chworowsky et al. 2023](#)). In contrast, recent analyses of massive star-forming galaxies at $z > 5$ find even more extreme star formation efficiencies ($\epsilon \sim 0.5$, [Xiao et al. 2023](#); [Casey et al. 2023](#)). If UMG-28740 is quiescent at $z \sim 5$, then its evolution is also consistent with an elevated formation efficiency at higher z . As shown in Figure 6, at $z \gtrsim 8$ where the quenched model for UMG-28740 is still star forming, the implied formation efficiency is on the order of (and potentially in excess of) unity, consistent with the expectations from models of feedback-free star formation ([Dekel et al. 2023](#)). Moreover, for the quenched model, UMG-28740 reaches significant tension with Λ CDM at even earlier times, exceeding $\epsilon = 1$ across the area of the full sky at $z \sim 15$. We tested this tension by fitting the CEERS photometry with a model set to high metallicity ($Z = 2Z_\odot$), to further explore the potential degeneracy between stellar age and metallicity. Even at super-solar abundances, however, we find that the formation time is only reduced by 0.15 Gyr, still in conflict with Λ CDM at high redshifts. This suggests UMG-28740 formed its stars at nearly impossible rates as currently predicted by theoretical models.

4.4. RUBIES-EGS-QG-1

While we finalized this manuscript, [de Graaff et al. \(2024\)](#) presented a *JWST*/NIRSpec PRISM and G395M spectrum of UMG-28740 (identified as RUBIES-EGS-QG-1 in [de Graaff et al. 2024](#)). The spectroscopic observations were collected in March 2024, with coverage of the Balmer break and the H α emission line. This effort was part of the RUBIES survey (GO-4233; PIs A. de Graaff and G. Brammer), which uses *JWST*/NIRSpec to observe galaxies in the EGS and UDS fields. In this section, we discuss the results of this independent work and compare those to our work.

The physical properties of RUBIES-EGS-QG-1 were measured from SED fitting of the *JWST*/NIRSpec PRISM spectrum, as well as photometry from *HST*/WFC3 (F125W, F140W, and F160W), *HST*/ACS (F814W and F606W), and *JWST*/NIRCam (F115W, F150W, F200W, F277W, F356W, F444W, and F410M). The spectrum and photometry were fit simultaneously via Prospector ([Leja et al. 2017](#)), assuming a Chabrier IMF ([Chabrier 2003](#)), stellar population models from Flexible Stellar Population Synthesis (FSPS, [Conroy & Gunn 2010](#); [Conroy et al. 2009](#)), the MILES spectral library ([Sánchez-Blázquez et al. 2006](#)), and MESA Isochrones and stellar tracks (MIST, [Choi et al. 2016](#); [Dotter 2016](#)). With Prospector, [de Graaff et al. \(2024\)](#) explored four different star formation history models – a delayed- τ , rising, and fixed $Z = Z_\odot$ model as well as a SFH comprised of 14 bins of constant star formation. The resulting best-fit stellar mass is in agreement with our results, yielding $M_\star = 8.7^{+0.5}_{-0.4} \times 10^{10} M_\odot$. With inclusion of mid-IR spectroscopy from *JWST*/NIRSpec, [de Graaff et al. \(2024\)](#) pin down the SFR in the last 100 Myr

as $\text{SFR}_{100} = 1.9^{+1.5}_{-1.0} M_\odot \text{ yr}^{-1}$, as the first work to spectroscopically confirm its quiescence. This independent work uses a separate SED modeling code and includes a high-quality spectrum that probes key spectral features, yet the final results are in excellent agreement with our quenched model.

A remarkable note between the results presented in [de Graaff et al. \(2024\)](#) and this manuscript is the detection of Ly α emission in a high-redshift quiescent galaxy. The observed spectrum and SED fit from [de Graaff et al. \(2024\)](#) did not include a Ly α emission line, potentially due to poor resolution at the blue end of the *JWST*/NIRSpec PRISM spectrum and the low SFR in the SED models. We investigate the potential origins of this Ly α emission by estimating the EW of Ly α for a best-fit model SED to the observed photometry given $\text{SFR} = 6 M_\odot \text{ yr}^{-1}$ (the highest SFR in the last 10 Myr for any model in [de Graaff et al. 2024](#)). We find the resulting Ly α EW to be $18.4^{+4.6}_{-2.5} \text{ \AA}$, well below the limiting EW given by our observations ($\text{EW}_{\text{Ly}\alpha} > 34.0 \text{ \AA}$, see §4.1). This suggests that star formation from this galaxy alone may not be the cause of this Ly α emission, but rather there could be additional contribution from an AGN or the surrounding circumgalactic medium. In addition, [de Graaff et al. \(2024\)](#) measured a slightly lower redshift via emission line modelling of [OIII], [NII], [SII], and H α ($z_{\text{spec}} = 4.8963$) such that the peak Ly α emission is redshifted by 479.67 km s^{-1} . Taken together with the asymmetric profile discussed in §2.2, this could be a sign of a significant amount of neutral gas surrounding the galaxy or AGN-driven galactic outflows.

The complementary analysis from [de Graaff et al. \(2024\)](#), taken together with the conclusions of our SED fitting, highlights the importance of high-quality mid-IR photometry – currently only accessible via *JWST*/NIRCam – in understanding systems like UMG-28740. Only with such data can we accurately measure key physical properties of $z \gtrsim 4$ galaxies. This example (UMG-28740) demonstrates the importance of *JWST* as we continue to search and study the rarest and most massive systems in the Universe.

5. CONCLUSIONS

In this work, we present spectroscopic confirmation of an ultra-massive galaxy (UMG, ID 28740) with $\log(M_\star/M_\odot) = 10.98 \pm 0.07$ at $z_{\text{spec}} = 4.8947$ based on deep spectroscopy with Keck/DEIMOS. UMG-28740 is one of the most massive spectroscopically-confirmed galaxies at $z \sim 5$ and lies within one of the most extreme overdensities in the EGS. Here, we summarize the main results of our analysis:

- We observe an asymmetric Ly α emission line profile with a relatively high asymmetry parameter of $A_f = 3.56$, suggesting the presence of neutral gas within or surrounding the galaxy. This neutral component could be due to gas in the interstellar medium, circumgalactic medium, or potentially associated with AGN-driven outflows.
- Using the spectroscopic redshift, we run a large suite of SED modeling (assuming various SFHs and using two SED fitting codes, BAGPIPES and FAST++) on two sets of photometric data from CANDELS ([Stefanon et al. 2017](#)) and CEERS ([Bagley et al. 2023](#)). Regardless of the SED code, set of free parameters, or SFH assumed, the stellar mass of UMG-28740 is tightly constrained, with a median mass of $\log(M_\star/M_\odot) = 10.98$ and a standard deviation of ± 0.07 between all runs.

- We find a discrepancy between the measured SFR and photometric dataset assumed, with multiple SED fits to the CANDELS photometry yielding a highly star-forming object. However, when fitting to the *JWST*/NIRCam photometry from CEERS, UMG-28740 is nearly quiescent, with $\text{SFR} < 32.2 \text{ M}_{\odot} \text{ yr}^{-1}$. Furthermore, while the quenched SED is consistent with both sets of photometry, the predicted flux from the best-fit star-forming SED under-predicts the measured flux at F277W and over-predicts the measured flux at F356W, F410M, and F444W; suggesting a quenched scenario is more likely. Finally, we show the presence of $\text{Ly}\alpha$ does not rule out a post-starburst, nearly quenched scenario.
- We find UMG-28740 to be the most massive member of a protocluster environment at $z \sim 5$, with the surface density of galaxies in the region around UMG-28740 roughly $10\times$ higher than the average coeval field surface density. Within $R_{\text{proj}} = 3.44 \text{ cMpc}$ of UMG-28740, we find 26 photometric members with $M_{\star} > 10^9 \text{ M}_{\odot}$ and 160 photometric members with $M_{\star} > 10^8 \text{ M}_{\odot}$ at $4.5 < z < 5.5$, using the CANDELS and CEERS photometric catalogs, respectively. Finally, we find four additional massive galaxies ($M_{\star} > 10^{10.5} \text{ M}_{\odot}$) within $0.29 - 1.13 \text{ cMpc}$ of UMG-28740 in the CANDELS photometric catalog.
- Comparing the SFR of UMG-28740 to the Illustris TNG-300 simulation box at $z = 5$ shows that, if quenched, UMG-28740 is extremely rare – with a $\text{SFR} \sim 2$ dex lower than any simulated galaxy at $\log(M_{\star}/\text{M}_{\odot}) \gtrsim 11$.
- We find UMG-28740 has a high star formation efficiency ($\sim 20\%$) at $z \sim 5$. Assuming a quenched model, this efficiency exceeds $\epsilon = 1$ for the area of the full sky,

suggesting UMG-28740 would need to form its stars at faster rates than predicted by theoretical models.

ACKNOWLEDGEMENTS

The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain. This work is based in part on observations taken by the CANDELS Multi-Cycle Treasury Program with the NASA/ESA *Hubble Space Telescope*, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. This work is also based in part on observations taken by the Cosmic Evolution Early Release Science Survey (CEERS) team with the NASA/ESA/CSA *JWST* from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127. SMUS and MCC acknowledge support from the National Science Foundation through grant AST-1815475. BF acknowledges support from *JWST*-GO-02913.001-A. GW gratefully acknowledges support from the National Science Foundation through grant AST-2205189 and from HST program number GO-16300. This research was supported by the International Space Science Institute (ISSI) in Bern, through ISSI International Team project “Understanding the evolution and transitioning of distant proto-clusters into clusters”. We also gratefully acknowledge the Lorentz Center in Leiden (NL) for facilitating discussions related to this project. MCC thanks Michael Boylan-Kolchin, Kristin Turney, and Lucas Cooper for helpful discussions related to this work. SMUS thanks Adam Carnall for additional support with BAGPIPES functionality. Finally, we thank the anonymous referee for their insightful comments and suggestions that improved this work.

REFERENCES

- Antwi-Danso J., et al., 2023, *arXiv e-prints*, p. [arXiv:2307.09590](#)
- Arrabal Haro P., et al., 2023a, *Nature*, **622**, 707
- Arrabal Haro P., et al., 2023b, *ApJ*, **951**, L22
- Bagley M. B., et al., 2023, *ApJ*, **946**, L12
- Baugh C. M., 2006, *Reports on Progress in Physics*, **69**, 3101
- Beichman C. A., Rieke M., Eisenstein D., Greene T. P., Krist J., McCarthy D., Meyer M., Stansberry J., 2012, in Clampin M. C., Fazio G. G., MacEwen H. A., Oeschmann Jacobus M. J., eds, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 8442*. Space Telescopes and Instrumentation 2012: Optical, Infrared, and Millimeter Wave, p. 84422N, doi:[10.1117/12.925447](#)
- Boylan-Kolchin M., 2023, *Nature Astronomy*, **7**, 731
- Bruzual G., Charlot S., 2003, *MNRAS*, **344**, 1000
- Calzetti D., Armus L., Bohlin R. C., Kinney A. L., Koornneef J., Storch-Bergmann T., 2000, *ApJ*, **533**, 682
- Carnall A. C., McLure R. J., Dunlop J. S., Davé R., 2018, *MNRAS*, **480**, 4379
- Carnall A. C., et al., 2023a, *Monthly Notices of the Royal Astronomical Society*, **520**, 3974
- Carnall A. C., et al., 2023b, *Nature*, **619**, 716
- Carnall A. C., et al., 2024, *arXiv e-prints*, p. [arXiv:2405.02242](#)
- Casey C. M., et al., 2023, *arXiv e-prints*, p. [arXiv:2308.10932](#)
- Chabrier G., 2003, *Publications of the Astronomical Society of the Pacific*, **115**, 763
- Chiang Y.-K., Overzier R., Gebhardt K., 2013, *ApJ*, **779**, 127
- Choi J., Dotter A., Conroy C., Cantiello M., Paxton B., Johnson B. D., 2016, *The Astrophysical Journal*, **823**, 102
- Chworowsky K., et al., 2023, *arXiv e-prints*, p. [arXiv:2311.14804](#)
- Conroy C., Gunn J. E., 2010, *ApJ*, **712**, 833
- Conroy C., Gunn J. E., White M., 2009, *ApJ*, **699**, 486
- Cooper M. C., Newman J. A., Davis M., Finkbeiner D. P., Gerke B. F., 2012, *spec2d: DEEP2 DEIMOS Spectral Pipeline*, Astrophysics Source Code Library, record ascl:1203.003 (ascl:1203.003)
- Cooray A., et al., 2014, *ApJ*, **790**, 40
- Croton D. J., et al., 2006, *MNRAS*, **365**, 11
- Davidzon I., et al., 2017, *A&A*, **605**, A70
- De Lucia G., Fontanot F., Xie L., Hirschmann M., 2024, *arXiv e-prints*, p. [arXiv:2401.06211](#)
- Dekel A., Sarkar K. C., Birnboim Y., Mandelker N., Li Z., 2023, *MNRAS*, **523**, 3201
- Dotter A., 2016, *The Astrophysical Journal Supplement Series*, **222**, 8
- Faber S. M., et al., 2003, in Iye M., Moorwood A. F. M., eds, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 4841*. Instrument Design and Performance for Optical/Infrared Ground-based Telescopes, pp 1657–1669, doi:[10.1117/12.460346](#)
- Finkelstein S. L., et al., 2023, *ApJ*, **946**, L13
- Forrest B., et al., 2020a, *ApJ*, **890**, L1
- Forrest B., et al., 2020b, *ApJ*, **903**, 47
- Forrest B., et al., 2022, *ApJ*, **938**, 109
- Glazebrook K., et al., 2017, *Nature*, **544**, 71
- Glazebrook K., et al., 2024, *Nature*, **628**, 277
- Grogin N. A., et al., 2011, *ApJS*, **197**, 35
- Jin S., et al., 2019, *ApJ*, **887**, 144
- Kakiichi K., Gronke M., 2021, *ApJ*, **908**, 30
- Koekemoer A. M., et al., 2011, *ApJS*, **197**, 36
- Kriek M., van Dokkum P. G., Labbé I., Franx M., Illingworth G. D., Marchesini D., Quadri R. F., 2009, *ApJ*, **700**, 221
- Kroupa P., Boily C. M., 2002, *MNRAS*, **336**, 1188
- Labbé I., et al., 2023, *Nature*, **616**, 266
- Laursen P., Razoumov A. O., Sommer-Larsen J., 2009, *ApJ*, **696**, 853
- Leja J., Johnson B. D., Conroy C., van Dokkum P. G., Byler N., 2017, *ApJ*, **837**, 170
- Ma J., et al., 2015, *ApJ*, **812**, 88
- Marsan Z. C., Marchesini D., Brammer G. B., Geier S., Kado-Fong E., Labbé I., Muzzin A., Stefanon M., 2017, *The Astrophysical Journal*, **842**, 21
- Marsan Z. C., et al., 2022, *The Astrophysical Journal*, **924**, 25
- Martin C. L., Dijkstra M., Henry A., Soto K. T., Danforth C. W., Wong J., 2015, *ApJ*, **803**, 6
- Murray S., 2014, HMF: Halo Mass Function calculator, Astrophysics Source Code Library, record ascl:1412.006

- Muzzin A., et al., 2013a, *ApJS*, **206**, 8
Muzzin A., et al., 2013b, *ApJ*, **777**, 18
Naidu R. P., et al., 2022, *arXiv e-prints*, p. [arXiv:2208.02794](https://arxiv.org/abs/2208.02794)
Nanayakkara T., et al., 2024, *Scientific Reports*, **14**, 3724
Nelson D., et al., 2018, *MNRAS*, **475**, 624
Nelson D., et al., 2019, *Computational Astrophysics and Cosmology*, **6**, 2
Newman J. A., et al., 2013, *ApJS*, **208**, 5
Pavesi R., et al., 2018, *ApJ*, **861**, 43
Planck Collaboration et al., 2016, *A&A*, **594**, A13
Rhoads J. E., et al., 2003, *AJ*, **125**, 1006
Riechers D. A., et al., 2013, *Nature*, **496**, 329
Riechers D. A., et al., 2017, *ApJ*, **850**, 1
Riechers D. A., et al., 2020, *ApJ*, **895**, 81
Rieke M. J., et al., 2003, in Mather J. C., ed., *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 4850, IR Space Telescopes and Instruments*, pp 478–485, doi:[10.1117/12.489103](https://doi.org/10.1117/12.489103)
Sánchez-Blázquez P., et al., 2006, *MNRAS*, **371**, 703
Scholtz J., et al., 2024, *arXiv e-prints*, p. [arXiv:2405.19401](https://arxiv.org/abs/2405.19401)
Schreiber C., et al., 2018, *A&A*, **618**, A85
Stefanon M., et al., 2015, *ApJ*, **803**, 11
Stefanon M., et al., 2017, *ApJS*, **229**, 32
Steinhardt C. L., Capak P., Masters D., Speagle J. S., 2016, *ApJ*, **824**, 21
Strandet M. L., et al., 2017, *ApJ*, **842**, L15
Sun G., Faucher-Giguère C.-A., Hayward C. C., Shen X., Wetzel A., Cochrane R. K., 2023, *The Astrophysical Journal Letters*, **955**, L35
Urbano Stawinski S., Cooper M., Magaz3ne Team 2024a, in *American Astronomical Society Meeting Abstracts*, p. 306.31
Urbano Stawinski S. M., et al., 2024b, *MNRAS*, **528**, 5624
Valentino F., et al., 2020, *ApJ*, **889**, 93
Valentino F., et al., 2023, *ApJ*, **947**, 20
Verhamme A., Schaerer D., Atek H., Tapken C., 2008, *A&A*, **491**, 89
Walter F., et al., 2012, *Nature*, **486**, 233
Weaver J. R., et al., 2023, *A&A*, **677**, A184
Weibel A., et al., 2024, *arXiv e-prints*, p. [arXiv:2403.08872](https://arxiv.org/abs/2403.08872)
Weiner B. J., et al., 2006, *ApJ*, **653**, 1027
Williams C. C., et al., 2019, *ApJ*, **884**, 154
Wright A. H., et al., 2017, *Monthly Notices of the Royal Astronomical Society*, **470**, 283
Xiao M., et al., 2023, *arXiv e-prints*, p. [arXiv:2309.02492](https://arxiv.org/abs/2309.02492)
Xie L., et al., 2024, *arXiv e-prints*, p. [arXiv:2402.01314](https://arxiv.org/abs/2402.01314)
Zavala J. A., et al., 2018, *Nature Astronomy*, **2**, 56
de Graaff A., et al., 2024, *arXiv e-prints*, p. [arXiv:2404.05683](https://arxiv.org/abs/2404.05683)

This paper was built using the Open Journal of Astrophysics $\LaTeX$ template. The OJA is a journal which provides fast and easy peer review for new papers in the *astro-ph* section of the

arXiv, making the reviewing process simpler for authors and referees alike. Learn more at <http://astro.theoj.org>.